

SEPARATING G_2 -INVARIANTS OF SEVERAL OCTONIONS

ARTEM LOPATIN AND ALEXANDR N. ZUBKOV

ABSTRACT. We describe separating G_2 -invariants of several copies of the algebra of octonions over an algebraically closed field of characteristic two. We also obtain a minimal separating and a minimal generating sets for G_2 -invariants of several copies of the algebra of octonions in case of a field of odd characteristic.

Keywords: polynomial invariants, exceptional groups, separating set, generating set, matrix invariants, positive characteristic.

2020 MSC: 20G05, 14L30, 17A75, 17D05, 20F29.

1. INTRODUCTION

Assume that \mathbb{F} is an algebraically closed field of arbitrary characteristic $p = \text{char } \mathbb{F} \geq 0$. All vector spaces and algebras are over \mathbb{F} and all algebras are associative, unless stated otherwise.

We continue the study of the invariants of the diagonal action of the exceptional simple group G_2 on the space of several octonions, over a field of positive characteristic. Over the field of complex numbers, this was done in [20]. This result has been generalized to an arbitrary infinite field of odd characteristic in [21], using a much finer technique of modules with good filtration, together with some results from the theory of groups with triality.

Unfortunately, the technique of modules with good filtration no longer works over a field of characteristic two and the complete description of the generating invariants in this case seems to be extremely difficult problem. Thus, it makes sense to describe separating invariants, since they satisfy the most important property of ordinary invariants to separate closed orbits in Zariski topology. The latter problem is usually more accessible and it does not require extremely technical methods. We describe the separating invariants over an algebraically closed field of characteristic two, using a detailed description of the subalgebras of the octonion algebra (up to the action of G_2) and the Hilbert-Mumford criterion.

The article is organized as follows. In Sections 2.1 and 2.2 we define the octonion algebra \mathbf{O} , the group G_2 and the algebra of G_2 -invariants $\mathbb{F}[\mathbf{O}^n]^{G_2}$ of n copies of the algebra of octonions \mathbf{O} . We use notations from [21]. Generators and relations between generators for $\mathbb{F}[\mathbf{O}^n]^{G_2}$ were described by Schwarz [20] over $\mathbb{F} = \mathbb{C}$. Zubkov and Shestakov described generators for $\mathbb{F}[\mathbf{O}^n]^{G_2}$ over an arbitrary field with $\text{char } \mathbb{F} \neq 2$ (see Section 2.4), but generators for the algebra $\mathbb{F}[\mathbf{O}^n]^{G_2}$ are still not known in case $p = 2$. Note that the invariants for the action of F_4 on several copies of the split Albert algebra were studied in [11]. Our results are formulated in Section 2.5. In Section 3 some definitions and notations are given. In Section 4 we describe a minimal generating and a minimal separating sets for $\mathbb{F}[\mathbf{O}^n]^{G_2}$ in

The work was supported by UAEU grant G00003324 and partially supported in accordance with the state task of the IM SB RAS, project FWNF-2022-0003.

case $p \neq 2$. In Section 5 a minimal generating set is constructed for the subalgebra $T_n \subset \mathbb{F}[\mathbf{O}^n]^{\mathbf{G}_2}$ of trace invariants in case $p = 2$. In Section 6 subalgebras of \mathbf{O} of dimension ≤ 3 are described modulo \mathbf{G}_2 -action in case $p = 2$. This result is applied in Section 7 to obtain our main result which is the description of a separating set for $\mathbb{F}[\mathbf{O}^n]^{\mathbf{G}_2}$ in case $p = 2$.

2. INVARIANTS OF OCTONIONS

2.1. Octonions. The *octonion algebra* $\mathbf{O} = \mathbf{O}(\mathbb{F})$, also known as the *split Cayley algebra*, is the vector space of all matrices

$$a = \begin{pmatrix} \alpha & \mathbf{u} \\ \mathbf{v} & \beta \end{pmatrix} \text{ with } \alpha, \beta \in \mathbb{F} \text{ and } \mathbf{u}, \mathbf{v} \in \mathbb{F}^3,$$

endowed with the following multiplication:

$$aa' = \begin{pmatrix} \alpha\alpha' + \mathbf{u} \cdot \mathbf{v}' & \alpha\mathbf{u}' + \beta'\mathbf{u} - \mathbf{v} \times \mathbf{v}' \\ \alpha'\mathbf{v} + \beta\mathbf{v}' + \mathbf{u} \times \mathbf{u}' & \beta\beta' + \mathbf{v} \cdot \mathbf{u}' \end{pmatrix}, \text{ where } a' = \begin{pmatrix} \alpha' & \mathbf{u}' \\ \mathbf{v}' & \beta' \end{pmatrix},$$

$\mathbf{u} \cdot \mathbf{v} = u_1v_1 + u_2v_2 + u_3v_3$ and $\mathbf{u} \times \mathbf{v} = (u_2v_3 - u_3v_2, u_3v_1 - u_1v_3, u_1v_2 - u_2v_1)$. For short, denote $\mathbf{c}_1 = (1, 0, 0)$, $\mathbf{c}_2 = (0, 1, 0)$, $\mathbf{c}_3 = (0, 0, 1)$, $\mathbf{0} = (0, 0, 0)$ from \mathbb{F}^3 . Consider the following basis of \mathbf{O} :

$$e_1 = \begin{pmatrix} 1 & \mathbf{0} \\ \mathbf{0} & 0 \end{pmatrix}, e_2 = \begin{pmatrix} 0 & \mathbf{0} \\ \mathbf{0} & 1 \end{pmatrix}, \mathbf{u}_i = \begin{pmatrix} 0 & \mathbf{c}_i \\ \mathbf{0} & 0 \end{pmatrix}, \mathbf{v}_i = \begin{pmatrix} 0 & \mathbf{0} \\ \mathbf{c}_i & 0 \end{pmatrix}$$

for $i = 1, 2, 3$. The unity of \mathbf{O} is denoted by $1_{\mathbf{O}} = e_1 + e_2$. We identify octonions

$$\alpha 1_{\mathbf{O}}, \begin{pmatrix} 0 & \mathbf{u} \\ \mathbf{0} & 0 \end{pmatrix}, \begin{pmatrix} 0 & \mathbf{0} \\ \mathbf{v} & 0 \end{pmatrix}$$

with $\alpha \in \mathbb{F}$, $\mathbf{u}, \mathbf{v} \in \mathbb{F}^3$, respectively. Similarly to $\mathbf{O}(\mathbb{F})$ we define the algebra of octonions $\mathbf{O}(\mathcal{A})$ over any commutative associative \mathbb{F} -algebra \mathcal{A} .

The algebra \mathbf{O} has a linear involution

$$\bar{a} = \begin{pmatrix} \beta & -\mathbf{u} \\ -\mathbf{v} & \alpha \end{pmatrix}, \text{ satisfying } \overline{aa'} = \bar{a}'\bar{a},$$

a *norm* $n(a) = a\bar{a} = \alpha\beta - \mathbf{u} \cdot \mathbf{v}$, and a non-degenerate symmetric bilinear *form* $q(a, a') = n(a + a') - n(a) - n(a') = \alpha\beta' + \alpha'\beta - \mathbf{u} \cdot \mathbf{v}' - \mathbf{u}' \cdot \mathbf{v}$. Define the linear function *trace* by $\text{tr}(a) = a + \bar{a} = \alpha + \beta$. The subspace $\{a \in \mathbf{O} \mid \text{tr}(a) = 0\}$ of traceless octonions is denoted by \mathbf{O}_0 . Notice that

$$\text{tr}(aa') = \text{tr}(a'a) \text{ and } n(aa') = n(a)n(a'). \quad (2.1)$$

The following quadratic equation holds:

$$a^2 - \text{tr}(a)a + n(a) = 0. \quad (2.2)$$

Since

$$n(a + a') = n(a) + n(a') - \text{tr}(aa') + \text{tr}(a)\text{tr}(a'), \quad (2.3)$$

the linearization of equation (2.2) implies

$$aa' + a'a - \text{tr}(a)a' - \text{tr}(a')a - \text{tr}(aa') + \text{tr}(a)\text{tr}(a') = 0. \quad (2.4)$$

The algebra \mathbf{O} is a simple *alternative* algebra, i.e., the following identities hold for $a, b \in \mathbf{O}$:

$$a(ab) = (aa)b, \quad (ba)a = b(aa). \quad (2.5)$$

The linearization implies that

$$a(a'b) + a'(ab) = (aa' + a'a)b, \quad (ba)a' + (ba')a = b(aa' + a'a). \quad (2.6)$$

The trace is associative, i.e., for all $a, b, c \in \mathbf{O}$ we have

$$\mathrm{tr}((ab)c) = \mathrm{tr}(a(bc)). \quad (2.7)$$

Note that

$$2n(a) = -\mathrm{tr}(a^2) + \mathrm{tr}^2(a) \text{ for each } a \in \mathbf{O}. \quad (2.8)$$

More details on \mathbf{O} can be found in Sections 1 and 3 of [21].

2.2. The group G_2 . The group $G_2 = G_2(\mathbb{F})$ is known to be the group $\mathrm{Aut}(\mathbf{O})$ of all automorphisms of the algebra \mathbf{O} . The group G_2 contain a Zariski closed subgroup $\mathrm{SL}_3 = \mathrm{SL}_3(\mathbb{F})$. Namely, every $g \in \mathrm{SL}_3$ defines the following automorphism of \mathbf{O} :

$$a \rightarrow \begin{pmatrix} \alpha & \mathbf{u}g \\ \mathbf{v}g^{-T} & \beta \end{pmatrix},$$

where g^{-T} stands for $(g^{-1})^T$ and $\mathbf{u}, \mathbf{v} \in \mathbb{F}^3$ are considered as row vectors. In what follows SL_3 is regarded as this subgroup of G_2 only. For every $\mathbf{u}, \mathbf{v} \in \mathbf{O}$ define $\delta_1(\mathbf{u}), \delta_2(\mathbf{v})$ from $\mathrm{Aut}(\mathbf{O})$ as follows:

$$\delta_1(\mathbf{u})(a') = \begin{pmatrix} \alpha' - \mathbf{u} \cdot \mathbf{v}' & (\alpha' - \beta' - \mathbf{u} \cdot \mathbf{v}')\mathbf{u} + \mathbf{u}' \\ \mathbf{v}' - \mathbf{u}' \times \mathbf{u} & \beta' + \mathbf{u} \cdot \mathbf{v}' \end{pmatrix},$$

$$\delta_2(\mathbf{v})(a') = \begin{pmatrix} \alpha' + \mathbf{u}' \cdot \mathbf{v} & \mathbf{u}' + \mathbf{v}' \times \mathbf{v} \\ (-\alpha' + \beta' - \mathbf{u}' \cdot \mathbf{v})\mathbf{v} + \mathbf{v}' & \beta' - \mathbf{u}' \cdot \mathbf{v} \end{pmatrix}.$$

The group G_2 is generated by SL_3 and $\delta_1(t\mathbf{u}_i), \delta_2(t\mathbf{v}_i)$ for all $t \in \mathbb{F}$ and $i = 1, 2, 3$. As an example, by straightforward calculations we can see that

$$h : \mathbf{O} \rightarrow \mathbf{O}, \text{ defined by } a \rightarrow \begin{pmatrix} \beta & -\mathbf{v} \\ -\mathbf{u} & \alpha \end{pmatrix}, \quad (2.9)$$

belongs to G_2 (see also the proof of Lemma 1 of [21]).

The action of G_2 on \mathbf{O} satisfies the following properties:

$$\overline{ga} = g\overline{a}, \quad \mathrm{tr}(ga) = \mathrm{tr}(a), \quad n(ga) = n(a), \quad q(ga, ga') = q(a, a').$$

Thus, \mathbf{O}_0 is a G_2 -submodule of \mathbf{O} .

Consider the diagonal action of G_2 on the vector space $\mathbf{O}^n = \mathbf{O} \oplus \cdots \oplus \mathbf{O}$ (n copies), i.e., $g(a_1, \dots, a_n) = (ga_1, \dots, ga_n)$ for all $g \in G_2$ and $a_1, \dots, a_n \in \mathbf{O}$. The coordinate ring of the affine variety \mathbf{O}^n is the polynomial \mathbb{F} -algebra $K_n = \mathbb{F}[\mathbf{O}^n] = \mathbb{F}[z_{ij} \mid 1 \leq i \leq n, 1 \leq j \leq 8]$, where $z_{ij} : \mathbf{O}^n \rightarrow \mathbb{F}$ is defined by $(a_1, \dots, a_n) \rightarrow \alpha_{ij}$ for

$$a_i = \begin{pmatrix} \alpha_{i1} & (\alpha_{i2}, \alpha_{i3}, \alpha_{i4}) \\ (\alpha_{i5}, \alpha_{i6}, \alpha_{i7}) & \alpha_{i8} \end{pmatrix} \in \mathbf{O}. \quad (2.10)$$

The action of $\mathrm{GL}(\mathbf{O})$ on \mathbf{O} induces the action on K_n by $(gf)(\underline{a}) = f(g^{-1}\underline{a})$ for all $g \in \mathrm{GL}(\mathbf{O})$, $f \in K_n$, $\underline{a} \in \mathbf{O}^n$.

To explicitly describe the action of G_2 on K_n consider the *generic octonions*

$$Z_i = \begin{pmatrix} z_{i1} & (z_{i2}, z_{i3}, z_{i4}) \\ (z_{i5}, z_{i6}, z_{i7}) & z_{i8} \end{pmatrix} \in \mathbf{O}(K_n)$$

for $1 \leq i \leq n$. Given $g \in G_2$, denote by $g \bullet Z_i$ the octonion

$$\begin{pmatrix} gz_{i1} & (gz_{i2}, gz_{i3}, gz_{i4}) \\ (gz_{i5}, gz_{i6}, gz_{i7}) & gz_{i8} \end{pmatrix} \in \mathbf{O}(K_n).$$

For any commutative algebra \mathcal{A} , the action of G_2 on \mathbf{O} extends for $\mathbf{O}(\mathcal{A})$ by \mathcal{A} -linearity. In particular, G_2 acts on $\mathbf{O}(K_n)$. It is easy to see that

$$g \bullet Z_i = g^{-1} Z_i, \quad (2.11)$$

where $g^{-1} Z_i$ stands for the action of g^{-1} on the octonion $Z_i \in \mathbf{O}(K_n)$.

The algebra of G_2 -invariants of several octonions (octonion G_2 -invariants, for short) is

$$K_n^{G_2} = \mathbb{F}[\mathbf{O}^n]^{G_2} = \{f \in \mathbb{F}[\mathbf{O}^n] \mid gf = f \text{ for all } g \in G_2\}.$$

In other words,

$$K_n^{G_2} = \{f \in \mathbb{F}[\mathbf{O}^n] \mid f(g\underline{a}) = f(\underline{a}) \text{ for all } g \in G_2, \underline{a} \in \mathbf{O}^n\}.$$

Similarly we can define $\mathbb{F}[\mathbf{O}_0^n]^{G_2}$, since $\mathbf{O}_0 \subset \mathbf{O}$ is invariant with respect to G_2 -action. Namely, the coordinate ring of the affine variety \mathbf{O}_0^n is $K_{0,n} = \mathbb{F}[\mathbf{O}_0^n] = \mathbb{F}[z_{ij} \mid 1 \leq i \leq n, 1 \leq j \leq 7]$. The *generic traceless octonions* are

$$X_i = \begin{pmatrix} z_{i1} & (z_{i2}, z_{i3}, z_{i4}) \\ (z_{i5}, z_{i6}, z_{i7}) & -z_{i1} \end{pmatrix}.$$

The analogue of formula (2.11) also holds for the generic traceless octonions, namely, $g \bullet X_i = g^{-1} X_i$ for all $g \in G_2$ and $1 \leq i \leq n$. The algebra of G_2 -invariants of several traceless octonions is $K_{0,n}^{G_2}$.

2.3. Separating sets. Consider a finite dimensional vector space \mathcal{V} and a linear group $G < \text{GL}(\mathcal{V})$. In 2002 Derksen and Kemper [2] (see [3] for the second edition) introduced the notion of separating invariants as a weaker concept than generating invariants. Given a subset S of $\mathbb{F}[\mathcal{V}]^G$, we write $S(u) = S(v)$ in case for every invariant $f \in S$ we have $f(u) = f(v)$. Elements u, v of \mathcal{V} are *separated by S* if $S(u) \neq S(v)$. If $u, v \in \mathcal{V}$ are separated by $\mathbb{F}[\mathcal{V}]^G$, then we say that they are *separated*. A subset $S \subset \mathbb{F}[\mathcal{V}]^G$ of the invariant ring is called *separating* if for any u, v from \mathcal{V} that are separated we have that they are separated by S . It is well-known that there always exists a finite separating set (see [2], Theorem 2.3.15). We say that a separating set is *minimal* if it is minimal w.r.t. inclusion. Obviously, any generating set is also separating. Minimal separating sets and upper bounds on degrees of elements of a separating set for different actions were constructed in [1, 4, 5, 6, 7, 9, 12, 13, 15, 17].

2.4. Known results. Denote by $\text{alg}_{\mathbb{F}}\{Z\}_n$ the non-associative \mathbb{F} -algebra generated by the generic octonions Z_1, \dots, Z_n and $1_{\mathbf{O}}$. Any product of the generic octonions is called a word of $\text{alg}_{\mathbb{F}}\{Z\}_n$. The unit $1_{\mathbf{O}} \in \text{alg}_{\mathbb{F}}\{Z\}_n$ is called the empty word. Note that for every $A, B \in \text{alg}_{\mathbb{F}}\{Z\}_n$ we have

$$\text{tr}(gA) = \text{tr}(A), \quad n(gA) = n(A), \quad g(AB) = (gA)(gB). \quad (2.12)$$

- Lemma 2.1.**
- (a) The trace of any (non-associative) product of X_1, \dots, X_n and $n(X_i)$ belong to $K_{0,n}^{G_2}$.
 - (b) The trace of any (non-associative) product of Z_1, \dots, Z_n and $n(Z_i)$ belong to $K_n^{G_2}$.
 - (c) The trace of any (non-associative) product of $Z_1, \dots, Z_n, \overline{Z_1}, \dots, \overline{Z_n}$ belong to $K_n^{G_2}$.

Proof. Let $w = w(Z_1, \dots, Z_n)$ be some (non-associative) product of Z_1, \dots, Z_n . Given $g \in G_2$, equalities (2.11), (2.12) imply that $g \operatorname{tr}(w) = \operatorname{tr}(w(g \bullet Z_1, \dots, g \bullet Z_n)) = \operatorname{tr}(w(g^{-1}Z_1, \dots, g^{-1}Z_n)) = \operatorname{tr}(g^{-1}w) = \operatorname{tr}(w)$. The case of $n(Z_i)$ is considered similarly. Part (b) is proven. The proof of part (a) is the same. Part (c) follows from part (b) and formulas

$$\operatorname{tr}(\bar{a}) = \operatorname{tr}(a), \quad n(\bar{a}) = n(a), \quad \operatorname{tr}(\bar{a}b) = \operatorname{tr}(a) \operatorname{tr}(b) - \operatorname{tr}(ab)$$

for all $a, b \in \mathbf{O}$. □

In case $\mathbb{F} = \mathbb{Q}$ for every $A_1, \dots, A_4 \in \operatorname{alg}_{\mathbb{F}}\{Z\}_n$ denote by $Q'(A_1, A_2, A_3, A_4)$ the complete skew symmetrization of $\operatorname{tr}(((A_1 A_2) A_3) A_4)$ with respect to its arguments, i.e.,

$$Q'(A_1, A_2, A_3, A_4) = \frac{1}{24} \sum_{\sigma \in S_4} (-1)^\sigma \operatorname{tr}(((A_{\sigma(1)} A_{\sigma(2)}) A_{\sigma(3)}) A_{\sigma(4)}).$$

In [21] it was shown that all coefficients of $Q'(X_1, X_2, X_3, X_4)$ belong to $\mathbb{Z}[\frac{1}{2}]$. Lemma 4.1 (see below) implies that all coefficients of $Q'(Z_1, Z_2, Z_3, Z_4)$ also belong to $\mathbb{Z}[\frac{1}{2}]$. Thus $Q'(A_1, A_2, A_3, A_4)$ is well-defined over an arbitrary field of odd characteristic.

In case $\operatorname{char} \mathbb{F} \neq 2$ the algebra of invariants

- $K_{0,n}^{G_2}$ is generated by $\operatorname{tr}(X_i X_j)$, $\operatorname{tr}((X_i X_j) X_k)$, $Q'(X_i, X_j, X_k, X_l)$;
- $K_n^{G_2}$ is generated by $\operatorname{tr}(Z_i)$, $\operatorname{tr}(Z_i Z_j)$, $\operatorname{tr}((Z_i Z_j) Z_k)$, $Q'(Z_i, Z_j, Z_k, Z_l)$

for all $1 \leq i, j, k, l \leq n$ (see Corollary 9 of [21], Section 1 of [21] and Lemma 3.1 below).

2.5. New results. Denote by $S_{0,n}$ the set

$$n(X_i); \operatorname{tr}(\cdots((X_{i_1} X_{i_2}) X_{i_3}) \cdots) X_{i_k}), \quad i_1 < \cdots < i_k, \quad k > 1,$$

for all $1 \leq i, i_1, \dots, i_k \leq n$, and by S_n the set

$$n(Z_i); \operatorname{tr}(\cdots((Z_{i_1} Z_{i_2}) Z_{i_3}) \cdots) Z_{i_k}), \quad i_1 < \cdots < i_k, \quad k > 0,$$

for all $1 \leq i, i_1, \dots, i_k \leq n$. Given $1 \leq k \leq n$, denote by $S_{0,n}^{(k)}$ and $S_n^{(k)}$ (respectively) the subset of $S_{0,n}$ and S_n (respectively) of elements of degree less or equal to k .

In case $\operatorname{char} \mathbb{F} = 2$ generators for the algebras $K_{0,n}^{G_2}$ and $K_n^{G_2}$ are not known. We introduce the algebra of *trace G_2 -invariants of octonions* $T_n \subset K_n^{G_2}$, i.e., the algebra T_n is generated by $n(Z_1), \dots, n(Z_n)$ and the traces of all (non-associative) products of Z_1, \dots, Z_n (see Lemma 2.1). In case $\operatorname{char} \mathbb{F} \neq 2$ we obviously have that $T_n = K_n^{G_2}$. We obtain the following results:

- $S_n^{(4)}$ is a minimal (w.r.t. inclusion) generating set for $K_n^{G_2}$ in case $\operatorname{char} \mathbb{F} \neq 2$ (see Proposition 4.4);
- $S_n^{(4)}$ is a minimal (w.r.t. inclusion) separating set for $K_n^{G_2}$ in case $\operatorname{char} \mathbb{F} \neq 2$ (see Proposition 4.6);
- T_n is minimally generated by S_n in case $\operatorname{char} \mathbb{F} = 2$ (see Theorem 5.2);
- $S_n^{(8)}$ is a separating set for $K_n^{G_2}$ in case $\operatorname{char} \mathbb{F} = 2$ (see Theorem 7.12).

3. AUXILIARIES

3.1. Symmetric algebra. A basis of \mathbf{O}^n is the set of all e_{ij} for $1 \leq i \leq n$, $1 \leq j \leq 8$, where by definition for $(a_1, \dots, a_n) \in \mathbf{O}^n$ with a_i as in (2.10) we have $a_i = \sum_{j=1}^8 \alpha_{ij} e_{ij}$. The symmetric algebra $S(\mathbf{O}^n)$ is a polynomial \mathbb{F} -algebra generated by all $\{e_{ij}\}$. Denote $O(n) = \{g \in \text{GL}(\mathbf{O}) \mid n(ga) = n(a) \text{ for all } a \in \mathbf{O}\}$. Obviously, $G_2 < O(n)$ and $q(ga, gb) = q(a, b)$ for all $a, b \in \mathbf{O}$.

Lemma 3.1. *Define the isomorphism of algebras $\psi : S(\mathbf{O}^n) \rightarrow K_n$ by*

$$\begin{pmatrix} e_{i1} & (e_{i2}, e_{i3}, e_{i4}) \\ (e_{i5}, e_{i6}, e_{i7}) & e_{i8} \end{pmatrix} \rightarrow Z_i,$$

$1 \leq i \leq n$, where the image of an element in a certain position of the left-hand side matrix is the element in the same position of the right-hand side matrix. Then ψ is an isomorphism of $O(n)$ -modules.

Proof. By Example 1 of [21] the isomorphism of algebras $\psi_0 : S(\mathbf{O}^n) \rightarrow K_n$ defined by

$$\begin{pmatrix} e_{i1} & (e_{i2}, e_{i3}, e_{i4}) \\ (e_{i5}, e_{i6}, e_{i7}) & e_{i8} \end{pmatrix} \rightarrow \begin{pmatrix} z_{i8} & (-z_{i5}, -z_{i6}, -z_{i7}) \\ (-z_{i2}, -z_{i3}, -z_{i4}) & z_{i1} \end{pmatrix}$$

is also an isomorphism of $O(n)$ -modules. Since $\psi = \hbar \psi_0$ for $\hbar \in G_2$ from (2.9), the required is proven. \square

3.2. Indecomposable invariants. Denote by $\mathbb{F}\{\mathbb{X}\}_n$ the free non-associative and non-commutative unital \mathbb{F} -algebra with free generators x_1, \dots, x_n , which are called letters. A word w is a non-empty product of letters. The number of a letters in w is the degree $\deg(w)$ of w . The degree of w in x_i is denoted by $\deg_{x_i}(w)$ and the total degree of w is denoted by $\deg(w)$. The multidegree of a word w is $\text{mdeg}(w) = (\deg_{x_1}(w), \dots, \deg_{x_n}(w))$. A word w with $\deg_{x_i}(w) \leq 1$ for all i is called multilinear. An element $f = \alpha_i w_i$ of $\mathbb{F}\{\mathbb{X}\}_n$, where $\alpha_i \in \mathbb{F}$ and w_i is a word, is \mathbb{N} -homogeneous (\mathbb{N}^n -homogeneous, respectively) if there exists d ($\Delta \in \mathbb{N}^n$, respectively) such that $\deg(w_i) = d$ ($\text{mdeg}(w_i) = \Delta$, respectively) for all i , where \mathbb{N} stands for non-negative integers. Define a homomorphisms of \mathbb{F} -algebras $\phi_0 : \mathbb{F}\{\mathbb{X}\}_n \rightarrow \text{alg}_{\mathbb{F}}\{X\}_n$ and $\phi : \mathbb{F}\{\mathbb{X}\}_n \rightarrow \text{alg}_{\mathbb{F}}\{Z\}_n$ by $x_i \rightarrow X_i$ and $x_i \rightarrow Z_i$ (respectively) for all i . In other words, for $f = f(x_1, \dots, x_n) \in \mathbb{F}\{\mathbb{X}\}_n$ we have $\phi(f) = f(Z_1, \dots, Z_n) \in \text{alg}_{\mathbb{F}}\{Z\}_n$. We write $x_{i_1} \circ \dots \circ x_{i_k}$ for some non-associative product of x_{i_1}, \dots, x_{i_k} . Similar notation we use for non-associative product in $\text{alg}_{\mathbb{F}}\{Z\}_n$.

For $f \in K_n$ denote by $\deg(f)$ its degree and by $\text{mdeg}(f)$ its multidegree, i.e., $\text{mdeg}(f) = (t_1, \dots, t_n)$, where t_i is the total degree of the polynomial f in z_{ij} , $1 \leq j \leq 8$, and $\deg(f) = t_1 + \dots + t_n$. For $f \in K_{0,n}$ the degree and multidegree are defined as above. It is well-known that the algebras $K_{0,n}^{G_2}$ and $K_n^{G_2}$ have \mathbb{N} -gradings by degrees and \mathbb{N}^n -gradings by multidegrees.

Consider an \mathbb{N}^n -graded unital (possibly, non-associative) algebra \mathcal{A} with the component of degree zero equal to \mathbb{F} . Denote by \mathcal{A}^+ the subalgebra generated by homogeneous elements of positive degree. A set $\{a_i\} \subseteq \mathcal{A}$ is a minimal (by inclusion) \mathbb{N}^n -homogeneous generating set (m.h.g.s.) of \mathcal{A} as a unital algebra if and only if the a_i 's are \mathbb{N}^n -homogeneous and $\{\bar{a}_i\} \cup \{1\}$ is a basis of the vector space $\bar{\mathcal{A}} = \mathcal{A}/(\mathcal{A}^+)^2$. We say that an element $a \in \mathcal{A}$ is *decomposable* and we write $a \equiv 0$ if $\bar{a} = 0$ in $\bar{\mathcal{A}}$. In other words, a decomposable element is equal to a polynomial

in elements of strictly less degree. Therefore the largest degree of indecomposable elements of \mathcal{A} is equal to the least upper bound for the degrees of elements of a m.h.g.s. for \mathcal{A} .

3.3. One-parameter subgroups of G_2 . Consider a finite dimensional vector space \mathcal{V} and a linear reductive group $G < \text{GL}(\mathcal{V})$. For a point $v \in \mathcal{V}$ and for a one-parameter subgroup $\theta : \mathbb{F}^\times \rightarrow G$ we have $\theta(t)v = \sum_{i \in I(v)} t^i v^{(i)}$ for all $t \in \mathbb{F}^\times$, where $I(v) = \{i \in \mathbb{Z} \mid v^{(i)} \neq 0\}$. Following [14] we say that $\lim_{t \rightarrow 0} \theta(t)v$ exists if and only if $I(v)$ consists of non-negative integers. Then $\lim_{t \rightarrow 0} \theta(t)v = 0$ if and only if $I(v)$ consists of positive integers only, otherwise $\lim_{t \rightarrow 0} \theta(t)v = v^{(0)}$. It is clear that if $\lim_{t \rightarrow 0} \theta(t)v$ exists, then it is contained in \overline{Gv} .

Given $\underline{\lambda} \in \mathbb{Z}^3$ with $\lambda_1 + \lambda_2 + \lambda_3 = 0$, the *standard* one-parameter subgroup $\theta_{\underline{\lambda}}$ of G_2 is defined by

$$\theta_{\underline{\lambda}}(t)e_i = e_i, \quad \theta_{\underline{\lambda}}(t)\mathbf{u}_j = t^{\lambda_j}\mathbf{u}_j, \quad \theta_{\underline{\lambda}}(t)\mathbf{v}_j = t^{-\lambda_j}\mathbf{v}_j,$$

for all $i = 1, 2$ and $1 \leq j \leq 3$.

4. MINIMAL GENERATING AND SEPARATING SETS

In this section we write $\text{tr}(i_1, \dots, i_k)$ for $\text{tr}((\dots((Z_{i_1}Z_{i_2})Z_{i_3})\dots)Z_{i_k})$, where $1 \leq i_1, \dots, i_k \leq n$. The following lemma can be proven by straightforward calculations.

Lemma 4.1. *Assume that $\text{char } \mathbb{F} \neq 2$. Then*

$$\begin{aligned} Q'(Z_1, Z_2, Z_3, Z_4) &= \text{tr}(1234) + \frac{1}{2}(-\text{tr}(1)\text{tr}(2)\text{tr}(3)\text{tr}(4) + \\ &- \text{tr}(1)\text{tr}(234) - \text{tr}(2)\text{tr}(134) - \text{tr}(3)\text{tr}(124) - \text{tr}(4)\text{tr}(123) + \\ &- \text{tr}(12)\text{tr}(34) + \text{tr}(13)\text{tr}(24) - \text{tr}(14)\text{tr}(23) + \\ &\text{tr}(1)\text{tr}(2)\text{tr}(34) + \text{tr}(1)\text{tr}(4)\text{tr}(23) + \text{tr}(2)\text{tr}(3)\text{tr}(14) + \text{tr}(3)\text{tr}(4)\text{tr}(12)). \end{aligned}$$

Recall that the definition of T_n was given in Section 2.5.

Lemma 4.2. *Let $w \in \mathbb{F}\{\mathbb{X}\}_n$ be a word.*

1. *If w is not multilinear and $\deg(w) > 2$, then $\text{tr}(w(Z_1, \dots, Z_n)) \equiv 0$ in T_n .*
2. *If w is multilinear and w is a product of letters x_{i_1}, \dots, x_{i_k} for $1 \leq i_1 < \dots < i_k \leq n$, then*

$$\text{tr}(w(Z_1, \dots, Z_n)) \equiv \pm \text{tr}(i_1, \dots, i_k) \text{ in } T_n.$$

3. *For all $1 \leq i_1 < \dots < i_k \leq n$ with $k \geq 3$ and every permutation $\sigma \in \mathcal{S}_k$ we have*

$$\text{tr}(i_{\sigma(1)}, \dots, i_{\sigma(k)}) \equiv (-1)^\sigma \text{tr}(i_1, \dots, i_k) \text{ in } T_n.$$

Proof. Combining (2.4) and (2.6) we obtain that

$$a(a'b) + a'(ab) = (\text{tr}(a)a' + \text{tr}(a')a + \text{tr}(aa') - \text{tr}(a)\text{tr}(a'))b$$

for all $a, a', b \in \mathbf{O}$. Since \mathbb{F} is infinite, the same equality holds for the generic octonions. We multiply it from the left and from the right by the generic octonions and then apply the trace. Since the trace is a linear function, we obtain that

$$\text{tr}(C_1 \circ \dots \circ C_r \circ (A(A'B)) \circ C_{r+1} \circ \dots \circ C_s) \equiv -\text{tr}(C_1 \circ \dots \circ C_r \circ (A'(AB)) \circ C_{r+1} \circ \dots \circ C_s) \quad (4.1)$$

for all products of the generic octonions $A, A', B, C_1, \dots, C_s$ with $0 \leq r \leq s$ and $s \geq 0$. Similarly we obtain that

$$\mathrm{tr}(C_1 \circ \dots \circ C_r \circ ((BA)A') \circ C_{r+1} \circ \dots \circ C_s) \equiv -\mathrm{tr}(C_1 \circ \dots \circ C_r \circ ((BA')A) \circ C_{r+1} \circ \dots \circ C_s) \quad (4.2)$$

In the same manner as above, (2.2) and (2.4) imply that

$$\mathrm{tr}(C_1 \circ \dots \circ C_r \circ (A^2) \circ C_{r+1} \circ \dots \circ C_s) \equiv 0, \quad (4.3)$$

$$\mathrm{tr}(C_1 \circ \dots \circ C_r \circ (AA') \circ C_{r+1} \circ \dots \circ C_s) \equiv -\mathrm{tr}(C_1 \circ \dots \circ C_r \circ (A'A) \circ C_{r+1} \circ \dots \circ C_s), \quad (4.4)$$

where in both cases $0 \leq r \leq s$ and $s > 0$. We claim that

$$\begin{aligned} &\text{If } W = Z_{i_1} \circ \dots \circ Z_{i_k} \text{ is a product of generic octonions,} \\ &\text{then } \mathrm{tr}(W) \equiv \pm \mathrm{tr}(i_{\sigma(1)}, \dots, i_{\sigma(k)}) \text{ for some } \sigma \in \mathcal{S}_k. \end{aligned} \quad (4.5)$$

Assume that claim (4.5) does not hold. Then there exists $\tau \in \mathcal{S}_k$ such that some product $W' = Z_{i_{\tau(1)}} \circ \dots \circ Z_{i_{\tau(k)}}$ satisfy the following conditions: $\mathrm{tr}(W) \equiv \pm \mathrm{tr}(W')$ and

- $W' = C_1 \circ \dots \circ (U(V_1V_2)) \circ \dots \circ C_s$ or $W' = C_1 \circ \dots \circ (VU) \circ \dots \circ C_s$, where
- $U = (\dots((Z_{j_1}Z_{j_2})Z_{j_3})\dots)Z_{j_r}$ for some $1 \leq j_1, \dots, j_r \leq n$ and $r \geq 2$ is maximal,
 - V, V_1, V_2 are some products of generic octonions,
 - C_1, \dots, C_s are generic octonions with $s \geq 0$.

By (2.1) and (4.4), we can assume that $W' = C_1 \circ \dots \circ (U(V_1V_2)) \circ \dots \circ C_s$. Consequently applying equivalence (4.1) and equivalence (2.1) or (4.4), we obtain that

$$\begin{aligned} \mathrm{tr}(C_1 \circ \dots \circ (U(V_1V_2)) \circ \dots \circ C_s) &\equiv -\mathrm{tr}(C_1 \circ \dots \circ (V_1(UV_2)) \circ \dots \circ C_s) \equiv \\ &\pm \mathrm{tr}(C_1 \circ \dots \circ ((UV_2)V_1) \circ \dots \circ C_s). \end{aligned}$$

If V_2 is a product of more than one generic octonions, then $V_2 = V_2'V_2''$ for some products V_2', V_2'' of generic octonions and we repeat the reasoning for $C_1 \circ \dots \circ (U(V_2'V_2'')) \circ \dots \circ C_s$. Therefore, we can assume that $V_2 = Z_j$ for some j ; a contradiction to the definition of W' .

Part 2 is a partial case of claim (4.5). Equivalences (4.2) and (4.4) imply that part 3 is valid for $1 \leq i_1 \leq \dots \leq i_k \leq n$, where $k \geq 3$. This fact together with claim (4.5) and formula (4.3) implies part 1. \square

As an immediate corollary of Lemma 4.2 and formula (2.8) we obtain the following.

Corollary 4.3. *Let $w \in \mathbb{F}\{\mathbb{X}\}_n$ be a word. Then $\mathrm{tr}(w(Z_1, \dots, Z_n))$ is a polynomial in $f \in S_n$ with $\mathrm{mdeg}(f) = (\delta_1, \dots, \delta_n)$ for some $\delta_1 \leq \deg_{x_1}(w), \dots, \delta_n \leq \deg_{x_n}(w)$.*

Proposition 4.4. *In case $\mathrm{char} \mathbb{F} \neq 2$ the algebra of invariants $K_n^{\mathrm{G}_2}$ is minimally generated by $S_n^{(4)}$.*

Proof. The description of generators for $K_n^{\mathrm{G}_2}$ from [21] (see Section 2.4 for the details) together with by Lemma 4.1 and Corollary 4.3 implies that the set $S_n^{(4)}$ generates the algebra $K_n^{\mathrm{G}_2}$. By Corollary 1 of [21] and formula (2.8), the invariants

$$\mathrm{tr}(i), n(Z_i), \mathrm{tr}(12), \mathrm{tr}(13), \mathrm{tr}(23), \mathrm{tr}(123),$$

where $1 \leq i \leq 3$, are algebraically independent over \mathbb{F} . Thus the required statement is proven for $n \leq 3$.

Assume $n \geq 4$. Thus $S_n^{(4)} \setminus \{f\}$ is not a generating set for any $f \in S_n^{(4)}$ with $\deg(f) \neq 4$.

Assume that $S_n^{(4)} \setminus \{\text{tr}(1234)\}$ is a generating set. Then $\text{tr}(1234)$ is a linear combination of $\text{tr}(12)\text{tr}(34)$, $\text{tr}(13)\text{tr}(24)$, $\text{tr}(14)\text{tr}(23)$ and products containing $\text{tr}(i)$ for some $1 \leq i \leq 4$. Considering substitutions $Z_1 \rightarrow \mathbf{v}_1$, $Z_2 \rightarrow \mathbf{v}_2$, $Z_3 \rightarrow \mathbf{v}_3$, $Z_4 \rightarrow e_1 - e_2$ and using equalities $\text{tr}(((\mathbf{v}_1\mathbf{v}_2)\mathbf{v}_3)(e_1 - e_2)) = -1$ and $\text{tr}(\mathbf{v}_i(e_1 - e_2)) = 0$ for $1 \leq i \leq 3$, we obtain a contradiction. The proposition is proven. \square

Remark 4.5. 1. By (2.8), in the formulation of Proposition 4.4 we can replace $n(Z_i)$ by $\text{tr}(Z_i^2)$ for all $1 \leq i \leq n$.
2. It easily follows from Proposition 4.4 (see also Section 1 of [21]) that $K_{0,n}^{G_2}$ is minimally generated by $S_{0,n}^{(4)}$.

Proposition 4.6. Assume $\text{char } \mathbb{F} \neq 2$. Then $S_{0,n}^{(4)}$ and $S_n^{(4)}$ are minimal separating sets for $K_{0,n}^{G_2}$ and $K_n^{G_2}$ (respectively) for all $n > 0$;

Proof. By Proposition 4.4 and Remark 4.5, the sets $S_{0,n}^{(4)}$ and $S_n^{(4)}$ are separating for $K_{0,n}^{G_2}$ and $K_n^{G_2}$ (respectively). For $a = 0$, $b = \mathbf{u}_1 + \mathbf{v}_1$ we have $\text{tr}(a) = n(a) = \text{tr}(b) = 0$, but $n(b) = 1$. For $a = 0$, $b = e_1$ we have $\text{tr}(a) = n(a) = n(b) = 0$, but $\text{tr}(b) = 1$. Hence, S_1 is a minimal separating set for $K_1^{G_2}$. Claims 1, 2, 3 (see below) imply that $S_{0,n}^{(4)}$ is a *minimal* separating set for $K_{0,n}^{G_2}$. Therefore, $S_n^{(4)}$ is also a minimal separating set for $K_n^{G_2}$.

Claim 1. Let $n = 2$. Then $S_{0,2} \setminus \{\text{tr}(X_1X_2)\}$ is not separating $K_{0,2}^{G_2}$.

To prove this claim consider $\underline{a} = (0, 0)$ and $\underline{b} = (\mathbf{u}_1, \mathbf{v}_1)$ from \mathbf{O}_0^2 . Then $\text{tr}(a_1a_2) \neq \text{tr}(b_1b_2)$.

Claim 2. Let $n = 3$. Then $S_{0,3} \setminus \{\text{tr}((X_1X_2)X_3)\}$ is not separating for $K_{0,3}^{G_2}$.

To prove this claim we consider $\underline{a} = (0, 0, 0)$ and $\underline{b} = (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$ from \mathbf{O}_0^3 . Then $\text{tr}(a_i a_j) = \text{tr}(b_i b_j) = 0$ for all $1 \leq i < j \leq 3$, but $\text{tr}(a_1 a_2 a_3) \neq \text{tr}(b_1 b_2 b_3)$.

Claim 3. Let $n = 4$. Then $S_{0,4} \setminus \{\text{tr}(((X_1X_2)X_3)X_4)\}$ is not separating for $K_{0,4}^{G_2}$.

To prove this claim we consider $\underline{a} = (\mathbf{u}_1, \mathbf{v}_1, c, \mathbf{u}_2)$ and $\underline{b} = (\mathbf{u}_1, \mathbf{v}_1, c, -\mathbf{v}_2)$ from \mathbf{O}_0^4 , where $c = e_1 + \mathbf{u}_2 - \mathbf{v}_2 - e_2$. Then $\text{tr}(a_i a_4) = \text{tr}(b_i b_4)$ for $1 \leq i \leq 3$ and $\text{tr}((a_i a_j) a_4) = \text{tr}((b_i b_j) b_4)$ for $1 \leq i < j \leq 3$, but $\text{tr}(((a_1 a_2) a_3) a_4) = 0$ and $\text{tr}(((b_1 b_2) b_3) b_4) = -1$. \square

5. TRACE INVARIANTS

The group $\text{GL}_2 = \text{GL}_2(\mathbb{F})$ acts on $M_2^n = M_2(\mathbb{F})^{\oplus n}$ diagonally by conjugation. The coordinate ring $\mathbb{F}[M_2^n] = \mathbb{F}[z_{i1}, z_{i2}, z_{i5}, z_{i8} \mid 1 \leq i \leq n]$ is also a GL_2 -module, where the *generic matrices* are

$$\widehat{Z}_i = \begin{pmatrix} z_{i1} & z_{i2} \\ z_{i5} & z_{i8} \end{pmatrix}.$$

Note that we consider $\mathbb{F}[M_2^n]$ as a subalgebra of K_n . In [8] it was shown that

$$\det(\widehat{Z}_i), \quad 1 \leq i \leq n; \quad \text{tr}(\widehat{Z}_{i_1} \cdots \widehat{Z}_{i_k}), \quad 1 \leq i_1 < \cdots < i_k \leq n, \quad k > 0, \quad (5.1)$$

is a minimal generating set for $\mathbb{F}[M_2^n]^{\text{GL}_2}$, where $k \leq 3$ in case $\text{char } \mathbb{F} \neq 2$. In particular, all elements from set (5.1) are indecomposable. Note that a minimal separating set for $\mathbb{F}[M_2^n]^{\text{GL}_2}$ was obtained in [12].

Define a surjective homomorphism of \mathbb{F} -algebras $\Psi : K_n \rightarrow \mathbb{F}[M_2^n]$ as follows: $z_{i3} \rightarrow 0, z_{i4} \rightarrow 0, z_{i6} \rightarrow 0, z_{i7} \rightarrow 0$ for all i . We can naturally extend Ψ to the linear map $\widehat{\Psi} : \mathbf{O}(K_n) \rightarrow \mathbf{O}(\mathbb{F}[M_2^n])$ by

$$\widehat{\Psi} \begin{pmatrix} f_1 & (f_2, f_3, f_4) \\ (f_5, f_6, f_7) & f_8 \end{pmatrix} = \begin{pmatrix} \Psi(f_1) & (\Psi(f_2), \Psi(f_3), \Psi(f_4)) \\ (\Psi(f_5), \Psi(f_6), \Psi(f_7)) & \Psi(f_8) \end{pmatrix}$$

for $f_1, \dots, f_8 \in K_n$.

For an associative commutative \mathbb{F} -algebra \mathcal{A} define a map $\mathcal{F} : M_2(\mathcal{A}) \rightarrow \mathbf{O}(\mathcal{A})$ by

$$\begin{pmatrix} a_1 & a_2 \\ a_3 & a_4 \end{pmatrix} \rightarrow \begin{pmatrix} a_1 & (a_2, 0, 0) \\ (a_3, 0, 0) & a_4 \end{pmatrix}$$

for $a_1, \dots, a_4 \in \mathcal{A}$. It is easy to see that \mathcal{F} is an injective homomorphism of algebras, since $(a, 0, 0) \times (b, 0, 0) = 0$ for all $a, b \in \mathcal{A}$. In what follows, we consider $\mathcal{A} = K_n$.

Lemma 5.1. *For all $1 \leq i, i_1, \dots, i_k \leq n$ we have*

- (a) $\widehat{\Psi}(\cdots((Z_{i_1} Z_{i_2}) Z_{i_3}) \cdots) Z_{i_k}) = \mathcal{F}(\widehat{Z}_{i_1} \cdots \widehat{Z}_{i_k});$
- (b) $\Psi(\text{tr}(\cdots((Z_{i_1} Z_{i_2}) Z_{i_3}) \cdots) Z_{i_k})) = \text{tr}(\widehat{Z}_{i_1} \cdots \widehat{Z}_{i_k});$
- (c) $\Psi(n(Z_i)) = \det(\widehat{Z}_i).$

Proof. (a) We prove by induction on $k \geq 1$. If $k = 1$, then the equality $\widehat{\Psi}(Z_{i_1}) = \mathcal{F}(\widehat{Z}_{i_1})$ is trivial.

Assume $\widehat{\Psi}(\cdots((Z_{i_1} Z_{i_2}) Z_{i_3}) \cdots) Z_{i_{k-1}}) = \mathcal{F}(\widehat{Z}_{i_1} \cdots \widehat{Z}_{i_{k-1}})$. Then

$$\begin{aligned} \widehat{\Psi}(\cdots((Z_{i_1} Z_{i_2}) Z_{i_3}) \cdots) Z_{i_k}) &= \widehat{\Psi}(\cdots((Z_{i_1} Z_{i_2}) Z_{i_3}) \cdots) Z_{i_{k-1}}) \widehat{\Psi}(Z_{i_k}) = \\ &= \mathcal{F}(\widehat{Z}_{i_1} \cdots \widehat{Z}_{i_{k-1}}) \mathcal{F}(\widehat{Z}_{i_k}). \end{aligned}$$

We conclude the proof by the fact that \mathcal{F} is a homomorphism of algebras.

(b) Part (b) follows from the equalities $\Psi(\text{tr}(U)) = \text{tr}(\widehat{\Psi}(U))$ for all $U \in \mathbf{O}(K_n)$ and $\text{tr}(\mathcal{F}(A)) = \text{tr}(A)$ for all $A \in M_2(K_n)$ together with part (a).

(c) Since

$$\Psi(n(Z_i)) = z_{i1} z_{i8} - z_{i2} z_{i5} = \det(\widehat{Z}_i),$$

part (c) is proven. \square

Lemma 2.1, Lemma 5.1 and the description of generators of $\mathbb{F}[M_2^n]^{\text{GL}_2}$ imply that

$$\mathbb{F}[M_2^n]^{\text{GL}_2} \subset \Psi(K_n^{\text{G}_2}), \quad (5.2)$$

where we have the equality in case $\text{char } \mathbb{F} \neq 2$.

Theorem 5.2. *In case $\text{char } \mathbb{F} = 2$ the algebra of trace G_2 -invariants T_n is minimally generated by S_n .*

Proof. By Corollary 4.3, the algebra T_n is generated by S_n . To show that S_n is a minimal generating set, it is enough to prove that every element $f \in S_n$ is indecomposable in T_n . Assume the contrary. If $f = \text{tr}(\cdots((Z_{i_1} Z_{i_2}) Z_{i_3}) \cdots) Z_{i_k})$, then by parts (b) and (c) of Lemma 5.1, $\Psi(f) = \text{tr}(\widehat{Z}_{i_1} \cdots \widehat{Z}_{i_k})$ is decomposable in $\mathbb{F}[M_2^n]^{\text{GL}_2}$; a contradiction. Similarly, if $f = n(Z_i)$, then $\Psi(f) = \det(\widehat{Z}_i)$ is decomposable in $\mathbb{F}[M_2^n]^{\text{GL}_2}$; a contradiction. \square

6. SUBALGEBRAS OF \mathbf{O} OF LOW DIMENSION

Given a subalgebra \mathcal{A} of \mathbf{O} , the *equivalence class* $[\mathcal{A}] = G_2 \cdot \mathcal{A}$ of \mathcal{A} is the family of all subalgebras of \mathbf{O} isomorphic to \mathcal{A} with respect to the action of G_2 . The family of all equivalence classes of all subalgebras of \mathbf{O} is denoted by $\Omega(\mathbf{O})$. Since all algebras from a given class $\mathfrak{A} \in \Omega(\mathbf{O})$ have the same dimension, we call it the *dimension* of \mathfrak{A} . A set of (linearly independent) octonions is said to be a *basis* of \mathfrak{A} , provided they form a basis of at least one algebra from \mathfrak{A} . An equivalence class $\mathfrak{A} \in \Omega(\mathbf{O})$ is called *closed* if there exists a subalgebra \mathcal{A} of \mathbf{O} with $[\mathcal{A}] = \mathfrak{A}$ and there is an \mathbb{F} -basis a_1, \dots, a_n of \mathcal{A} such that the G_2 -orbit of (a_1, \dots, a_n) is closed in \mathbf{O}^n . Note that the given definition is not equivalent to the trivial claim that \mathcal{A} is closed in \mathbf{O} . More details on the definition of a closed equivalence class can be found in Remark 7.3 (see below).

The main result of this section is the following statement.

Proposition 6.1. *Assume $\text{char } \mathbb{F} = 2$ and an equivalence class $\mathfrak{A} \in \Omega(\mathbf{O})$ has the dimension $d \leq 3$. Then one of the following sets is a basis for \mathfrak{A} :*

$$\begin{aligned} d = 1: & \{1_{\mathbf{O}}\}, \{\mathbf{u}_1\}, \{e_1\}; \\ d = 2: & \{1_{\mathbf{O}}, \mathbf{u}_1\}, \{\mathbf{u}_1, \mathbf{v}_2\}, \{e_1, \mathbf{u}_1\}, \{e_1, \mathbf{v}_1\}, \{e_1, e_2\}; \\ d = 3: & \{1_{\mathbf{O}}, \mathbf{u}_1, \mathbf{v}_2\}, \{e_1, e_2, \mathbf{u}_1\}, \{e_1, \mathbf{u}_1, \mathbf{v}_2\}, \{\mathbf{u}_1, \mathbf{v}_2, a\} \text{ for non-zero } a = \alpha 1_{\mathbf{O}} + \\ & \beta \mathbf{u}_3 + \gamma \mathbf{v}_3 \text{ with } \alpha^2 = \beta\gamma \text{ for some } \alpha, \beta, \gamma \in \mathbb{F}. \end{aligned}$$

Note that we do not require for a subalgebra of \mathbf{O} to be unital. The proof of Proposition 6.1 will be given in the series of propositions and lemmas, which are interesting on their own.

Proposition 6.2 (Proposition 3.4 of [18]). *For each $a \in \mathbf{O}$ there exists $g \in G_2$ such that ga is a canonical octonion of one of the following types:*

$$\begin{aligned} \text{(D)} & \begin{pmatrix} \alpha_1 & \mathbf{0} \\ \mathbf{0} & \alpha_8 \end{pmatrix}, \\ \text{(K}_1\text{)} & \begin{pmatrix} \alpha_1 & (1, 0, 0) \\ \mathbf{0} & \alpha_1 \end{pmatrix}, \end{aligned}$$

for some $\alpha_1, \alpha_8 \in \mathbb{F}$. Moreover, these canonical octonions are unique modulo permutation $\alpha_1 \leftrightarrow \alpha_8$ for type (D).

Proposition 6.3 (Corollary 5.2 of [18]). *Assume $\text{char } \mathbb{F} = 2$. For each $(a, b) \in \mathbf{O}_0^2$ there exists $g \in G_2$ such that $g(a, b)$ is a pair of one of the following types:*

$$\begin{aligned} \text{(EE)} & (\alpha_1 1_{\mathbf{O}}, \beta_1 1_{\mathbf{O}}), \\ \text{(EK}_1\text{)} & \left(\alpha_1 1_{\mathbf{O}}, \begin{pmatrix} \beta_1 & (1, 0, 0) \\ \mathbf{0} & \beta_1 \end{pmatrix} \right), \\ \text{(K}_1\text{E)} & \left(\begin{pmatrix} \alpha_1 & (1, 0, 0) \\ \mathbf{0} & \alpha_1 \end{pmatrix}, \beta_1 1_{\mathbf{O}} \right), \\ \text{(K}_1\text{L}_1\text{)} & \left(\begin{pmatrix} \alpha_1 & (1, 0, 0) \\ \mathbf{0} & \alpha_1 \end{pmatrix}, \begin{pmatrix} \beta_1 & (\beta_2, 0, 0) \\ \mathbf{0} & \beta_1 \end{pmatrix} \right) \text{ with } \beta_2 \neq 0, \\ \text{(K}_1\text{L}_1^\top\text{)} & \left(\begin{pmatrix} \alpha_1 & (1, 0, 0) \\ \mathbf{0} & \alpha_1 \end{pmatrix}, \begin{pmatrix} \beta_1 & \mathbf{0} \\ (\beta_5, 0, 0) & \beta_1 \end{pmatrix} \right) \text{ with } \beta_5 \neq 0, \\ \text{(K}_1\text{M}_1\text{)} & \left(\begin{pmatrix} \alpha_1 & (1, 0, 0) \\ \mathbf{0} & \alpha_1 \end{pmatrix}, \begin{pmatrix} \beta_1 & (0, 1, 0) \\ \mathbf{0} & \beta_1 \end{pmatrix} \right), \\ \text{(K}_1\text{M}_1^\top\text{)} & \left(\begin{pmatrix} \alpha_1 & (1, 0, 0) \\ \mathbf{0} & \alpha_1 \end{pmatrix}, \begin{pmatrix} \beta_1 & \mathbf{0} \\ (0, 1, 0) & \beta_1 \end{pmatrix} \right). \end{aligned}$$

for all $\alpha_1, \beta_1, \beta_2, \beta_5 \in \mathbb{F}$.

For a vector $\mathbf{u} = (u_1, u_2, u_3)$ of \mathbb{F}^3 and a permutation $\sigma \in \mathcal{S}_3$ we write $\mathbf{u} \diamond \sigma$ for

- $(u_{\sigma(1)}, u_{\sigma(2)}, u_{\sigma(3)})$ in case $\sigma \in \{(1, 2, 3), (1, 3, 2)\}$ is a cycle or σ is the identity permutation;
- $(u_2, u_1, -u_3)$ if $\sigma = (1, 2)$;
- $(u_3, -u_2, u_1)$ if $\sigma = (1, 3)$;
- $(-u_1, u_3, u_2)$ if $\sigma = (2, 3)$.

Remark 6.4 (Remarks 3.2 and 3.3 of [18]). Assume $a = \begin{pmatrix} \alpha & \mathbf{u} \\ \mathbf{v} & \beta \end{pmatrix} \in \mathbf{O}$. Then

- (a) if $\mathbf{u} \neq 0$, then there exists $g \in \mathrm{SL}_3$ such that $ga = \begin{pmatrix} \alpha & (1, 0, 0) \\ \mathbf{v}' & \beta \end{pmatrix}$, where $\mathbf{v}' = (*, 0, 0)$ or $\mathbf{v}' = (0, 1, 0)$;
- (b) if $\mathbf{v} \neq 0$, then there exists $g \in \mathrm{SL}_3$ such that $ga = \begin{pmatrix} \alpha & \mathbf{u}' \\ (1, 0, 0) & \beta \end{pmatrix}$, where $\mathbf{u}' = (*, 0, 0)$ or $\mathbf{u}' = (0, 1, 0)$.
- (c) for a permutation $\sigma \in \mathcal{S}_3$ exists $g \in \mathrm{SL}_3$ such that $ga = \begin{pmatrix} \alpha & \mathbf{u} \diamond \sigma \\ \mathbf{v} \diamond \sigma & \beta \end{pmatrix}$.
- (d) if $\mathbf{u} = (\gamma_1, \gamma_2, \gamma_3)$ with $\gamma_2 \neq 0$ or $\gamma_3 \neq 0$ and $\mathbf{v} = (\delta, 0, 0)$, then there exists $g \in \mathrm{SL}_3$ such that $ga = \begin{pmatrix} \alpha & (\gamma_1, 0, 1) \\ (\delta, 0, 0) & \beta \end{pmatrix}$.

Lemma 6.5. *If $\mathcal{A} \subset \mathbf{O}$ is a non-zero subalgebra, then there exists $g \in \mathrm{G}_2$ such that $1_{\mathbf{O}} \in g\mathcal{A}$ or $\mathbf{u}_1 \in g\mathcal{A}$ or $e_1 \in g\mathcal{A}$. In particular, if $\mathrm{char} \mathbb{F} = 2$ and $\mathcal{A} \not\subset \mathbf{O}_0$ is a non-zero subalgebra of \mathbf{O} , then there exists $g \in \mathrm{G}_2$ such that $e_1 \in g\mathcal{A}$.*

Proof. Consider some non-zero octonion $a \in \mathcal{A}$. Proposition 6.2 implies that up to the action of G_2 one has to consider one of the following cases:

1. $a = \alpha e_1 + \beta e_2$ for some $\alpha, \beta \in \mathbb{F}$. In case $\alpha \neq \beta$ are both non-zero, the octonions a and $a^2 = \alpha^2 e_1 + \beta^2 e_2$ are linearly independent; in particular, $e_1, e_2 \in \mathcal{A}$. In case $\alpha = 0$ we have $\beta \neq 0$ and $e_2 \in \mathcal{A}$; thus, acting by \tilde{h} on \mathcal{A} we obtain that $e_1 \in \mathcal{A}$. In case $\beta = 0$, we have $\alpha \neq 0$ and $e_1 \in \mathcal{A}$. Finally, in case $\alpha = \beta$ we have $\alpha \neq 0$ and $1_{\mathbf{O}} \in \mathcal{A}$.

2. $a = \alpha 1_{\mathbf{O}} + \mathbf{u}_1$ for some $\alpha \in \mathbb{F}$. Then $a^2 = \alpha^2 1_{\mathbf{O}} + 2\alpha \mathbf{u}_1$ lies in \mathcal{A} . If $\alpha = 0$, then $\mathbf{u}_1 \in \mathcal{A}$. Otherwise, $\alpha 1_{\mathbf{O}} + 2\mathbf{u}_1 \in \mathcal{A}$. Hence, $\mathbf{u}_1 \in \mathcal{A}$.

In case $\mathrm{char} \mathbb{F} = 2$ and $\mathcal{A} \not\subset \mathbf{O}_0$ we repeat the proof for $a \in \mathcal{A}$ with $\mathrm{tr}(a) \neq 0$. Hence we have only case 1 for a , where $\alpha \neq \beta$. \square

6.1. The case of traceless subalgebra. In this section we assume that $\mathrm{char} \mathbb{F} = 2$ and $\mathcal{A} \subset \mathbf{O}$ is a subalgebra of traceless octonions, i.e., $\mathcal{A} \subset \mathbf{O}_0$.

Remark 6.6. If $a = \begin{pmatrix} \alpha & \mathbf{u} \\ \mathbf{v} & \beta \end{pmatrix} \in \mathcal{A}$ is *triangular* (i.e., $\mathbf{u} = \mathbf{0}$ or $\mathbf{v} = \mathbf{0}$), and $\alpha \neq 0$ or $\beta \neq 0$, then $1_{\mathbf{O}} \in \mathcal{A}$.

Proof. Since $\alpha = \beta$ is non-zero, considering $a^2 = \alpha^2 1_{\mathbf{O}}$ we obtain the required. \square

Lemma 6.7. *If $\dim \mathcal{A} \geq 2$, then there exists $g \in \mathrm{G}_2$ such that one of the following possibilities holds:*

- (a) $\{1_{\mathbf{O}}, \mathbf{u}_1\} \subset g\mathcal{A}$;

(b) $\{\mathbf{u}_1, \mathbf{v}_2\} \subset g\mathcal{A}$ and $1_{\mathbf{O}} \notin g\mathcal{A}$.

Proof. By Lemma 6.5, we assume that one of the following alternatives holds:

1. $1_{\mathbf{O}} \in \mathcal{A}$. There exists $a \in \mathcal{A}$ such that $\{1_{\mathbf{O}}, a\}$ are linearly independent. Since $G_2 1_{\mathbf{O}} = 1_{\mathbf{O}}$, by Proposition 6.2, we can assume that $a = \alpha e_1 + \beta e_2$ or $a = \alpha 1_{\mathbf{O}} + \mathbf{u}_1$ for some $\alpha, \beta \in \mathbb{F}$. In the first case we have $\alpha = \beta$ and $\{1_{\mathbf{O}}, a\}$ are linearly dependent; a contradiction. In the second case we obtain that $\mathbf{u}_1 = a - \alpha 1_{\mathbf{O}}$ lies in \mathcal{A} .

2. There is $\mathbf{u}_1 \in \mathcal{A}$ and $1_{\mathbf{O}} \notin \mathcal{A}$. There exists $b \in \mathcal{A}$ such that $\{\mathbf{u}_1, b\}$ are linearly independent. Consider $g \in G_2$ such that $g(\mathbf{u}_1, b) = (a', b')$ is one of the pairs from Proposition 6.3. Since $\text{tr}(a') = n(a') = 0$ and $a' \neq 0$, one easily sees that $a' = \mathbf{u}_1$. By Remark 6.6 and the fact that $1_{\mathbf{O}} \notin \mathcal{A}$ one sees that both diagonal entries of b' are equal to zero. Using the fact that $\{\mathbf{u}_1, b'\}$ are linearly independent, we obtain that the pair (\mathbf{u}_1, b') has one of the following types:

- ($K_1 L_1^\top$) $b' = \beta \mathbf{v}_1$, where $\beta \in \mathbb{F} \setminus \{0\}$. Since $\mathbf{u}_1 b' = \beta e_1$ and $\text{tr}(e_1) \neq 0$, we obtain a contradiction.
 ($K_1 M_1$) $b' = \mathbf{u}_2$. Since $\mathbf{u}_1 b' = \mathbf{v}_3$, acting by a suitable element of SL_3 and using part (c) of Remark 6.4, we obtain case (b).
 ($K_1 M_1^\top$) $b' = \mathbf{v}_2$, i.e., we have case (b).

□

Lemma 6.8. *If $\dim \mathcal{A} \geq 3$, then there exists $g \in G_2$ such that one of the following possibilities holds:*

- (a) $\{1_{\mathbf{O}}, \mathbf{u}_1, \mathbf{v}_2\} \subset g\mathcal{A}$;
 (b) $\{\mathbf{u}_1, \mathbf{v}_2, a\} \subset g\mathcal{A}$ and $1_{\mathbf{O}} \notin g\mathcal{A}$, where the element $a = \alpha 1_{\mathbf{O}} + \beta \mathbf{u}_3 + \gamma \mathbf{v}_3$ is non-zero and $\alpha, \beta, \gamma \in \mathbb{F}$ satisfy $\alpha^2 = \beta\gamma$.

Proof. By Lemma 6.7, one can assume that one of the following possibilities holds:

1. $\{1_{\mathbf{O}}, \mathbf{u}_1\} \subset g\mathcal{A}$. There exists $b \in \mathcal{A}$ such that $\{1_{\mathbf{O}}, \mathbf{u}_1, b\}$ are linearly independent. Since $G_2 1_{\mathbf{O}} = 1_{\mathbf{O}}$ and the \mathbb{F} -span of \mathbf{u}_1, b does not contain $1_{\mathbf{O}}$, applying Proposition 6.3 to the pair (\mathbf{u}_1, b) we can assume that one of the following alternatives hold (see for the details part 2 of the proof of Lemma 6.7):

- ($K_1 L_1^\top$) $\{1_{\mathbf{O}}, \mathbf{u}_1, \beta \mathbf{v}_1\} \subset \mathcal{A}$ for some non-zero $\beta \in \mathbb{F}$. Since $u_1 v_1 = e_1$, we obtain a contradiction.
 ($K_1 M_1$) $\{1_{\mathbf{O}}, \mathbf{u}_1, \mathbf{u}_2\} \subset \mathcal{A}$. Since $\mathbf{u}_1 \mathbf{u}_2 = \mathbf{v}_3$, acting by a suitable element of SL_3 from part (c) of Remark 6.4 we obtain case (a).
 ($K_1 M_1^\top$) $\{1_{\mathbf{O}}, \mathbf{u}_1, \mathbf{v}_2\} \subset \mathcal{A}$, i.e., we have case (a).

2. $\{\mathbf{u}_1, \mathbf{v}_2\} \subset \mathcal{A}$ and $1_{\mathbf{O}} \notin g\mathcal{A}$. Consider $b \in \mathcal{A}$ such that $\{\mathbf{u}_1, \mathbf{v}_2, b\}$ are linearly independent. Moreover, one can assume that $b = \begin{pmatrix} \beta_1 & (0, \beta_3, \beta_4) \\ (\beta_5, 0, \beta_7) & \beta_1 \end{pmatrix}$ for some $\beta_i \in \mathbb{F}$. Since

$$\mathbf{u}_1 b = \begin{pmatrix} \beta_5 & (\beta_1, 0, 0) \\ (0, -\beta_4, \beta_3) & 0 \end{pmatrix} \quad \text{and} \quad \mathbf{v}_2 b = \begin{pmatrix} 0 & (-\beta_7, 0, \beta_5) \\ (0, \beta_1, 0) & \beta_3 \end{pmatrix},$$

we have $\beta_3 = \beta_5 = 0$. The equality $b^2 = (\beta_1^2 + \beta_4 \beta_7) 1_{\mathbf{O}}$ implies that case (b) holds. □

6.2. The case of non-traceless subalgebra. In this section we assume that $\text{char } \mathbb{F} = 2$ and $\mathcal{A} \not\subset \mathbf{O}_0$ is a subalgebra of \mathbf{O} .

Lemma 6.9. *If $\dim \mathcal{A} \geq 2$, then there exists $g \in G_2$ such that one of the following possibilities holds:*

- (a) $\{e_1, \mathbf{u}_1\} \subset g\mathcal{A}$;
- (b) $\{e_1, \mathbf{v}_1\} \subset g\mathcal{A}$;
- (c) $\{e_1, e_2\} \subset g\mathcal{A}$.

Proof. By Lemma 6.5 we can assume that $e_1 \in \mathcal{A}$. There exists $b \in \mathcal{A}$ such that $\{e_1, b\}$ are linearly independent. Moreover, one can also assume that $b = \begin{pmatrix} 0 & \mathbf{u} \\ \mathbf{v} & \beta \end{pmatrix}$ for some $\mathbf{u}, \mathbf{v} \in \mathbb{F}^3$ and $\beta \in \mathbb{F}$.

Assume $\mathbf{u} \neq \mathbf{0}$. Since $e_1 b = \begin{pmatrix} 0 & \mathbf{u} \\ \mathbf{0} & 0 \end{pmatrix}$, by part (a) of Remark 6.4 there exists $g \in \text{SL}_3$ such that $g(e_1, e_1 b) = (e_1, \mathbf{u}_1)$, i.e., the case (a) holds.

Assume $\mathbf{v} \neq \mathbf{0}$. Since $b e_1 = \begin{pmatrix} 0 & \mathbf{0} \\ \mathbf{v} & 0 \end{pmatrix}$, by part (b) of Remark 6.4 there exists $g \in \text{SL}_3$ such that $g(e_1, b e_1) = (e_1, \mathbf{v}_1)$, i.e., the case (b) holds.

In case $\mathbf{u} = \mathbf{v} = \mathbf{0}$ we have $\beta \neq 0$, i.e., the case (c) holds. \square

Lemma 6.10. *If $\dim \mathcal{A} \geq 3$, then there exists $g \in G_2$ such that one of the following possibilities holds:*

- (a) $\{e_1, e_2, \mathbf{u}_1\} \subset g\mathcal{A}$;
- (b) $\{e_1, \mathbf{u}_1, \mathbf{v}_2\} \subset g\mathcal{A}$.

Before the proof of this lemma we formulate the following remark.

Remark 6.11. (a) $\{e_1, e_2, \mathbf{u}_1\} \subset \mathcal{A}$ if and only if $\{e_1, e_2, \mathbf{v}_1\} \subset \hbar\mathcal{A}$;
 (b) $\{e_1, \mathbf{u}_1, \mathbf{v}_2\} \subset \mathcal{A}$ if and only if $\{e_1, \mathbf{u}_2, \mathbf{v}_1\} \subset g\mathcal{A}$ for some $g \in G_2$ (see part (c) of Remark 6.4).

Proof of Lemma 6.10. By Lemma 6.9, we assume that one of the following possibilities holds:

1. $\{e_1, \mathbf{u}_1\} \subset \mathcal{A}$. There exists $b \in \mathcal{A}$ such that $\{e_1, \mathbf{u}_1, b\}$ are linearly independent.

Moreover, we can assume that $b = \begin{pmatrix} 0 & \mathbf{u} \\ \mathbf{v} & \beta \end{pmatrix}$ for some $\mathbf{u} = (0, *, *) \in \mathbb{F}^3$, $\mathbf{v} = (\gamma_1, \gamma_2, \gamma_3) \in \mathbb{F}^3$ and $\beta \in \mathbb{F}$.

Assume $\mathbf{u} \neq \mathbf{0}$. Since $e_1 b = \begin{pmatrix} 0 & \mathbf{u} \\ \mathbf{0} & 0 \end{pmatrix}$, by part (d) of Remark 6.4 there exists $g \in \text{SL}_3$ such that $g(e_1, \mathbf{u}_1, e_1 b) = (e_1, \mathbf{u}_1, \mathbf{u}_3)$. The equality $\mathbf{u}_1 \mathbf{u}_3 = -\mathbf{v}_2$ implies that the case (b) holds.

Assume $\mathbf{u} = \mathbf{0}$. Note that $b e_1 = \begin{pmatrix} 0 & \mathbf{0} \\ \mathbf{v} & 0 \end{pmatrix}$ lies in \mathcal{A} .

Let $\gamma_1 \neq 0$. Similarly to part (d) of Remark 6.4, there exists $g \in \text{SL}_3$ such that $g(e_1, \mathbf{u}_1, b e_1) = (e_1, \mathbf{u}_1, \mathbf{v}_1)$. The equality $\mathbf{v}_1 \mathbf{u}_1 = e_2$ implies that the case (a) holds.

Otherwise, $\gamma_1 = 0$. In case $\gamma_2 \neq 0$ or $\gamma_3 \neq 0$, by Remark 3.1 of [18] there exists $g \in \text{SL}_3$ such that $g(e_1, \mathbf{u}_1, b e_1) = (e_1, \mathbf{u}_1, \mathbf{v}_2)$, i.e., the case (b) holds. If $\gamma_2 = \gamma_3 = 0$, then $\beta \neq 0$ and $e_2 \in \mathcal{A}$, i.e., the case (a) holds.

2. The case $\{e_1, \mathbf{v}_1\} \subset \mathcal{A}$ is similar to case 1.

3. $\{e_1, e_2\} \subset \mathcal{A}$. There exists $b = \begin{pmatrix} \alpha & \mathbf{u} \\ \mathbf{v} & \beta \end{pmatrix}$ in \mathcal{A} such that $\{e_1, e_2, b\}$ are linearly independent. Moreover, we can assume that $\alpha = \beta = 0$.

Assume $\mathbf{u} \neq \mathbf{0}$. Since $e_1 b = \begin{pmatrix} 0 & \mathbf{u} \\ \mathbf{0} & 0 \end{pmatrix}$, by part (a) of Remark 6.4 there exists $g \in \mathrm{SL}_3$ such that $g(e_1, e_2, e_1 b) = (e_1, e_2, \mathbf{u}_1)$, i.e., the case (a) holds.

Otherwise, $\mathbf{v} \neq \mathbf{0}$. Since $b e_1 = \begin{pmatrix} 0 & \mathbf{0} \\ \mathbf{v} & 0 \end{pmatrix}$ by part (b) of Remark 6.4 there exists $g \in \mathrm{SL}_3$ such that $g(e_1, e_2, b e_1) = (e_1, e_2, \mathbf{v}_1)$. By part (a) of Remark 6.11 we obtain that case (a) holds. \square

6.3. Proof of Proposition 6.1. Assume $\mathfrak{A} = [\mathcal{A}]$ for some subalgebra \mathcal{A} of \mathbf{O} . Lemmas 6.5, 6.7, 6.8, 6.9, 6.10 imply that there exist $g \in G_2$ such that $g\mathcal{A}$ contains one of the sets from the formulation of Proposition 6.1. Since the \mathbb{F} -span of each of these sets is an algebra, the proof is completed.

7. SEPARATING INVARIANTS IN CASE $\mathrm{char} \mathbb{F} = 2$

In this section we assume that $\mathrm{char} \mathbb{F} = 2$. Introduce the following notations for $\underline{a} \in \mathbf{O}^n$:

- the *rank* $\mathrm{rk}(\underline{a})$ is the dimension of the subspace of \mathbf{O} spanned by a_1, \dots, a_n ;
- $\mathrm{alg}(\underline{a})$ is the \mathbb{F} -algebra (in general, non-unital) generated by a_1, \dots, a_n .

Obviously, $\mathrm{rk}(g\underline{a}) = \mathrm{rk}(\underline{a})$ for every $g \in G_2$. Denote by

$$\mathbb{M} = \begin{pmatrix} * & (*, 0, 0) \\ (*, 0, 0) & * \end{pmatrix} \quad \text{and} \quad \mathbb{S} = \begin{pmatrix} * & (*, *, 0) \\ (*, 0, *) & * \end{pmatrix},$$

respectively, the subalgebra of *quaternions* and *sextonions* of \mathbf{O} , respectively, where the term *sextonions* was introduced in [16]. Note that $\mathcal{F} : M_2(\mathbb{F}) \rightarrow \mathbb{M}$ is the isomorphism of \mathbb{F} -algebras (see Section 5 for the details).

The following remark is well-known (for example, see Theorem 2.3.6 of [3]).

Remark 7.1. Assume $\underline{a} \in \mathbf{O}^n$. Then there exist a unique closed G_2 -orbit $O = O_{\underline{a}}$ in the closure of $G_2 \underline{a}$. Moreover, $O_{\underline{a}}$ is the only closed orbit in the fiber

$$\{\underline{c} \in \mathbf{O}^n \mid f(\underline{a}) = f(\underline{c}) \text{ for all } f \in K_n^{G_2}\}.$$

In particular, $f(\underline{a}) = f(\underline{c})$ for every $f \in K_n^{G_2}$ and $\underline{c} \in O_{\underline{a}}$.

Observe that the group GL_n acts naturally on \mathbf{O}^n on the right as follows: for any $A = (\alpha_{ij}) \in \mathrm{GL}_n$ and $\underline{a} \in \mathbf{O}^n$ we set

$$(\underline{a}A)_i = \sum_{1 \leq k \leq n} \alpha_{ki} a_k \quad \text{for } 1 \leq i \leq n.$$

This action commutes with the left G_2 -action.

Lemma 7.2. Given $\underline{a}, \underline{b} \in \mathbf{O}^n$, define $\underline{a}' = \underline{a}A$ and $\underline{b}' = \underline{b}A$ for some $A \in \mathrm{GL}_n$. Then

- $G_2 \underline{a} = G_2 \underline{b}$ if and only if $G_2 \underline{a}' = G_2 \underline{b}'$;
- given some $d \geq 2$, we have that \underline{a} and \underline{b} are not separated by $S_n^{(d)}$ if and only if \underline{a}' and \underline{b}' are not separated by $S_n^{(d)}$;
- $G_2 \underline{a}$ is closed if and only if $G_2 \underline{a}'$ is closed; $G_2 \underline{b}$ is closed if and only if $G_2 \underline{b}'$ is closed.

Proof. Since A is invertible, for each part of this lemma it is sufficient to prove “only if” implication.

- (a) For each $g \in G_2$ the equality $g\underline{a} = \underline{b}$ implies $g\underline{a}' = \underline{b}'$, hence our claim follows.
- (b) Assume that \underline{a} and \underline{b} are not separated by $S_n^{(d)}$, i.e., $f(\underline{a}) = f(\underline{b})$ for all $f \in S_n^{(d)}$. The linearity of the trace together with Corollary 4.3 and formula (2.3) implies that $h(\underline{a}') = h(\underline{b}')$ for all $h \in S_n^{(d)}$.
- (c) The right action by A on \mathbf{O}^n gives a homeomorphism of \mathbf{O}^n with respect to Zariski topology. Hence it sends closed subsets to closed subsets. \square

The following remark is a consequence of part (c) of Lemma 7.2.

Remark 7.3. An equivalence class $\mathfrak{A} \in \Omega(\mathbf{O})$ is closed if and only if for every subalgebra \mathcal{A} of \mathbf{O} with $[\mathcal{A}] = \mathfrak{A}$ we have that \mathcal{A} is the \mathbb{F} -span of some a_1, \dots, a_n , then the G_2 -orbit of (a_1, \dots, a_n) is closed in \mathbf{O}^n .

Proposition 7.4. *The set $S_m^{(8)} \subset K_m^{G_2}$ is separating for every $m > 0$ if and only if $S_n^{(8)}$ separates different closed G_2 -orbits of $\underline{a} = (a_1, \dots, a_l, 0, \dots, 0) \in \mathbf{O}^n$ and $\underline{b} \in \mathbf{O}^n$ for all $n > 0$, where*

- a_1, \dots, a_l is a basis of some subalgebra \mathcal{A} of \mathbf{O} ,
- b_1, \dots, b_n of \mathbf{O} linearly generate some subalgebra \mathcal{B} of \mathbf{O} ,
- $\dim \mathcal{A} \geq \dim \mathcal{B}$.

Proof. We only have to prove “if” part of the statement. Assume that $\underline{a}, \underline{b} \in \mathbf{O}^n$ are not separated by $S_n^{(8)}$. To obtain the required we will show that $G_2\underline{a} = G_2\underline{b}$.

By Remark 7.1 we can assume that $G_2\underline{a}$ and $G_2\underline{b}$ are closed.

Denote by \mathcal{A} the algebra generated by a_1, \dots, a_n and by \mathcal{B} the algebra generated by b_1, \dots, b_n . Denote $l = \dim \mathcal{A}$. Without loss of generality we can assume that $l \geq \dim \mathcal{B}$.

Claim 1. Without loss of generality, we can assume that \mathbb{F} -span of a_1, \dots, a_n is \mathcal{A} and \mathbb{F} -span of b_1, \dots, b_n is \mathcal{B} .

Let us prove Claim 1. There exist words v_1, \dots, v_r of $\mathbb{F}\{\mathbb{X}\}_n$ such that the \mathbb{F} -span of the set $a_1, \dots, a_n, v_1(\underline{a}), \dots, v_r(\underline{a})$ is \mathcal{A} . Similarly, there exist words w_1, \dots, w_s of $\mathbb{F}\{\mathbb{X}\}_n$ such that the \mathbb{F} -span of the set $b_1, \dots, b_n, w_1(\underline{b}), \dots, w_s(\underline{b})$ is \mathcal{B} .

It is easy to see that the G_2 -orbits of

$$\begin{aligned} c_1 &= (a_1, \dots, a_n, v_1(\underline{a}), \dots, v_r(\underline{a}), w_1(\underline{a}), \dots, w_s(\underline{a})), \\ c_2 &= (b_1, \dots, b_n, v_1(\underline{b}), \dots, v_r(\underline{b}), w_1(\underline{b}), \dots, w_s(\underline{b})) \end{aligned}$$

are closed. Obviously, $G_2\underline{a} = G_2\underline{b}$ if and only if $G_2c_1 = G_2c_2$. By Corollary 4.3, \underline{a} and \underline{b} are not separated by $S_n^{(8)}$ if and only if c_1 and c_2 are not separated by $S_n^{(8)}$. Therefore, we can consider c_1, c_2 instead of $\underline{a}, \underline{b}$ and Claim 1 is proven.

Claim 2. Given an \mathbb{F} -basis a'_1, \dots, a'_l of \mathcal{A} , without loss of generality, we can assume that $\underline{a} = (a'_1, \dots, a'_l, 0, \dots, 0) \in \mathbf{O}^n$.

To prove Claim 2, we apply part 2 of Lemma 4.2 to see that each simultaneous permutation of entries of \underline{a} and \underline{b} preserves the property that $\underline{a}, \underline{b}$ are not separated by $S_n^{(8)}$. Therefore, without loss of generality we can assume that a_1, \dots, a_l is a basis for \mathcal{A} for some $1 \leq l \leq n$. For every $1 \leq i \leq l$ we write $a'_i = \sum_{j=1}^l \gamma_{ij} a_j$

and for every $l < i \leq n$ we write $a_i = \sum_{j=1}^l \gamma_{ij} a_j$ for some $\gamma_{ij} \in \mathbb{F}$ and denote $b'_i = b_i - \sum_{j=1}^l \gamma_{ij} b_j$ for all $1 \leq i \leq n$.

By part (c) of Lemma 7.2, the G_2 -orbits of

$$c_3 = (a'_1, \dots, a'_l, \underbrace{0, \dots, 0}_{n-l}) \text{ and } c_4 = (b'_1, \dots, b'_l, b'_{l+1}, \dots, b'_n)$$

are closed. By part (a) of Lemma 7.2, $G_2 \underline{a} = G_2 \underline{b}$ if and only if $G_2 c_3 = G_2 c_4$. By part (b) of Lemma 7.2, \underline{a} and \underline{b} are not separated by $S_n^{(8)}$ if and only if c_3 and c_4 are not separated by $S_n^{(8)}$. Therefore, we can consider c_3, c_4 instead of $\underline{a}, \underline{b}$ and Claim 2 is proven.

Since $\underline{a}, \underline{b}$ satisfy conditions (a), (b) from the formulation of the lemma and $G_2 \underline{a}, G_2 \underline{b}$ are closed, we obtain that $G_2 \underline{a} = G_2 \underline{b}$. \square

The analogue of the following lemma over a finite field is part (ii) of Lemma 3.3 from [10].

Lemma 7.5. *Every automorphism of the \mathbb{F} -algebra \mathbb{M} can be extended to an automorphism of the algebra \mathbf{O} .*

Proof. Each automorphism f of \mathbb{M} is given by the action of some $g \in \text{GL}_2$ on $\mathbb{M} \simeq M_2(\mathbb{F})$, i.e., $f(\mathcal{F}(A)) = \mathcal{F}(g^{-1}Ag)$ for all $A \in M_2(\mathbb{F})$. Moreover, we can assume that $g \in \text{SL}_2$. Note that the group SL_2 is generated by

$$g_1(t) = \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}, \quad g_2(t) = \begin{pmatrix} 1 & 0 \\ t & 1 \end{pmatrix}$$

for all $t \in \mathbb{F}$. Since

$$\delta_1(t, 0, 0)(\mathcal{F}(A)) = \mathcal{F}(g_1(t)^{-1}Ag_1(t)) \text{ and } \delta_2(t, 0, 0)(\mathcal{F}(A)) = \mathcal{F}(g_2(t)^{-1}Ag_2(t))$$

for all $t \in \mathbb{F}$ and $a \in M_2(\mathbb{F})$, the proof is concluded. \square

- Remark 7.6.**
1. For every $a \in \mathbb{M}$ with $\text{tr}(a) = 1$ and $n(a) = 0$ there exists g from the stabilizer $\text{Stab}_{G_2}(\mathbb{M}) = \{g \in G_2 \mid g\mathbb{M} \subset \mathbb{M}\}$ such that $ga = e_1$.
 2. For every $a \in \mathbb{M}$ with $\text{tr}(a) = 0$ and $n(a) = 1$ there exists $g \in \text{Stab}_{G_2}(\mathbb{M})$ such that $ga \in \{1_{\mathbf{O}}, 1_{\mathbf{O}} + \mathbf{u}_1\}$.
 3. Given non-zero $\gamma \in \mathbb{F}$, there exists $\xi_\gamma \in \text{Stab}_{G_2}(\mathbb{M})$ such that for every $\alpha_1, \dots, \alpha_4 \in \mathbb{F}$ we have

$$\xi_\gamma \begin{pmatrix} \alpha_1 & (\alpha_2, 0, 0) \\ (\alpha_3, 0, 0) & \alpha_4 \end{pmatrix} = \begin{pmatrix} \alpha_1 & (\gamma\alpha_2, 0, 0) \\ (\gamma^{-1}\alpha_3, 0, 0) & \alpha_4 \end{pmatrix}.$$

4. Assume that $\underline{a} = (e_1, e_2)$ and $\underline{b} \in \mathbb{M}^2$ satisfy $S_n^{(2)}(\underline{a}) = S_n^{(2)}(\underline{b})$. Then there exists $g \in \text{Stab}_{G_2}(\mathbb{M})$ such that $gb_1 = e_1$ and $gb_2 \in \{e_2, e_2 + \mathbf{u}_1, e_2 + \mathbf{v}_1\}$.
5. If $b \in \mathbb{M}$ satisfies $\text{tr}(b) = n(b) = \text{tr}(e_1 b) = 0$, then $b \in \mathbb{F}\mathbf{u}_1$ or $b \in \mathbb{F}\mathbf{v}_1$.

Proof. **1.** For $A = \mathcal{F}^{-1}(a)$ we have $\text{tr}(A) = 1$ and $\det(A) = 0$. Hence there exists $g \in \text{GL}_2$ such that $g^{-1}Ag = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ and Lemma 7.5 implies the required.

2. For $A = \mathcal{F}^{-1}(a)$ we have $\text{tr}(A) = 0$ and $\det(A) = 1$. Hence there exists $g \in \text{GL}_2$ such that $g^{-1}Ag = \begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}$ or $g^{-1}Ag = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}$ for some $\lambda, \lambda_1, \lambda_2$ and Lemma 7.5 implies the required.

3. Given $g = \begin{pmatrix} 1 & 0 \\ 0 & \gamma^{-1} \end{pmatrix} \in \text{GL}_2$, we have

$$g^{-1} \begin{pmatrix} \alpha_1 & \alpha_2 \\ \alpha_3 & \alpha_4 \end{pmatrix} g = \begin{pmatrix} \alpha_1 & \gamma\alpha_2 \\ \gamma^{-1}\alpha_3 & \alpha_4 \end{pmatrix}.$$

Lemma 7.5 concludes the proof.

4. By part 1 we assume that $b_1 = e_1$. Denote $\mathcal{F}^{-1}(b_2) = B_2 = \begin{pmatrix} \beta_1 & \beta_2 \\ \beta_3 & \beta_4 \end{pmatrix}$. Since $0 = \text{tr}(a_1 a_2) = \text{tr}(b_1 b_2)$, we obtain $\beta_1 = 0$. The equalities $\text{tr}(b_2) = 1$ and $n(b_2) = 0$ imply $\beta_4 = 1$ and $\beta_2 \beta_3 = 0$. Part 3 concludes the proof.

5. Denote $\mathcal{F}^{-1}(b) = B = \begin{pmatrix} \beta_1 & \beta_2 \\ \beta_3 & \beta_4 \end{pmatrix}$. Since $0 = \text{tr}(e_1 b) = \beta_1$, the equalities $\text{tr}(b) = n(b) = 0$ imply the required. \square

Lemma 7.7. *Assume that $\underline{a} = (a_1, 0, \dots, 0) \in \mathbb{M}^n$ and $\underline{b} \in \mathbb{M}^n$ are not separated by $S_n^{(2)}$, where $a_1 \in \{1_{\mathbf{O}}, e_1\}$ and $\dim(\text{alg}(\underline{b})) \leq 1$. Then $G_2 \underline{a} = G_2 \underline{b}$.*

Proof. 1. Let $a_1 = 1_{\mathbf{O}}$. Since $\text{tr}(b_1) = 0$ and $n(b_1) = 1$, by part 2 of Remark 7.6 we can assume that $b_1 = 1_{\mathbf{O}}$ or $b_1 = 1_{\mathbf{O}} + \mathbf{u}_1$.

In the first case the condition $\dim(\text{alg}(\underline{b})) \leq 1$ implies $\underline{b} = (1_{\mathbf{O}}, \beta_2 1_{\mathbf{O}}, \dots, \beta_n 1_{\mathbf{O}})$ for some $\beta_2, \dots, \beta_n \in \mathbb{F}$. Since $0 = n(b_i) = \beta_i^2$ for all $1 < i \leq n$, we have $\underline{a} = \underline{b}$.

In the second case we have that b_1 and $b_1^2 = 1_{\mathbf{O}}$ are linearly independent; a contradiction.

2. Let $a_1 = e_1$. Since $\text{tr}(b_1) = 1$ and $n(b_1) = 0$, by part 1 of Remark 7.6 we can assume that $b_1 = e_1$. Then the condition $\dim(\text{alg}(\underline{b})) \leq 1$ implies that $\underline{b} = (e_1, \beta_2 e_1, \dots, \beta_n e_1)$ for some $\beta_2, \dots, \beta_n \in \mathbb{F}$. For each $1 < i \leq n$ we have $0 = \text{tr}(a_1 0) = \text{tr}(b_1 b_i) = \beta_i$. Therefore, $\underline{a} = \underline{b}$. \square

Lemma 7.8. *Assume that $\underline{a} = (e_1, e_2, 0, \dots, 0) \in \mathbb{M}^n$ and $\underline{b} \in \mathbb{M}^n$ are not separated by $S_n^{(2)}$ and $\dim(\text{alg}(\underline{b})) \leq 2$. Then $G_2 \underline{a} = G_2 \underline{b}$.*

Proof. By part 4 of Remark 7.6 we can assume that $b_1 = e_1$ and $b_2 \in \{e_2, e_2 + \mathbf{u}_1, e_2 + \mathbf{v}_1\}$.

Let $b_2 = e_2$. For $3 \leq i \leq n$ part 5 of Remark 7.6 implies that $b_i \in \mathbb{F}\mathbf{u}_1$ or $b_i \in \mathbb{F}\mathbf{v}_1$, since $\text{tr}(b_i) = n(b_i) = \text{tr}(b_1 b_i) = 0$. It follows from $\dim(\text{alg}(\underline{b})) \leq 2$ that $b_i = 0$ for all $3 \leq i \leq n$. Therefore, $\underline{a} = \underline{b}$.

In case $b_2 = e_2 + \mathbf{u}_1$ we consider $b_1 b_2 = \mathbf{u}_1$ and obtain that $\{e_1, \mathbf{u}_1, e_2\} \subset \text{alg}(\underline{b})$; a contradiction.

In case $b_2 = e_2 + \mathbf{v}_1$ we consider $b_2 b_1 = \mathbf{v}_1$ and obtain that $\{e_1, \mathbf{v}_1, e_2\} \subset \text{alg}(\underline{b})$; a contradiction. \square

Lemma 7.9. *If $\underline{a} = (e_1, e_2, \mathbf{u}_1, \mathbf{v}_1, 0, \dots, 0) \in \mathbb{M}^n$ and $\underline{b} \in \mathbb{M}^n$ are not separated by $S_n^{(3)}$, then $G_2 \underline{a} = G_2 \underline{b}$.*

Proof. By part 4 of Remark 7.6 we can assume that $b_1 = e_1$ and $b_2 \in \{e_2, e_2 + \mathbf{u}_1, e_2 + \mathbf{v}_1\}$. Assume $3 \leq i \leq n$. We have $\text{tr}(b_i) = n(b_i) = \text{tr}(b_1 b_i) = 0$, since $\text{tr}(a_1 a_3) = \text{tr}(a_1 a_4) = 0$. Thus part 5 of Remark 7.6 implies that $b_i = \beta_i \mathbf{u}_1$ or $b_i = \beta_i \mathbf{v}_1$ for some $\beta_i \in \mathbb{F}$. Since $\text{tr}(b_3 b_4) = \text{tr}(a_3 a_4) = 1$, we obtain that $\beta_3 \beta_4 = 1$ and either $b_3 = \beta_3 \mathbf{u}_1$, $b_4 = \beta_4 \mathbf{v}_1$ or $b_3 = \beta_3 \mathbf{v}_1$, $b_4 = \beta_4 \mathbf{u}_1$ for some non-zero $\beta \in \mathbb{F}$.

1. Let $b_3 = \beta_3 \mathbf{u}_1$, $b_4 = \beta_3^{-1} \mathbf{v}_1$. By part 3 of Remark 7.6 we can assume that $\beta_3 = 1$. Consider $5 \leq i \leq n$. If $b_i = \beta_i \mathbf{u}_1$, then $0 = \text{tr}(a_4 a_i) = \text{tr}(b_4 b_i) = \beta_i$. If $b_i = \beta_i \mathbf{v}_1$, then $0 = \text{tr}(a_4 a_i) = \text{tr}(b_4 b_i) = \beta_i$. Therefore, $\underline{a} = \underline{b}$.
2. In case $b_3 = \beta_3 \mathbf{v}_1$, $b_4 = \beta_3^{-1} \mathbf{u}_1$ we have $0 = \text{tr}((b_1 b_3) b_4) = \text{tr}((a_1 a_3) a_4) = \text{tr}(\mathbf{u}_1 \mathbf{v}_1) = 1$; a contradiction. \square

Lemma 7.10. *If $\underline{a} = (e_1, e_2, \mathbf{u}_1, \mathbf{v}_1, \mathbf{u}_2, \mathbf{v}_2, \mathbf{u}_3, \mathbf{v}_3, 0, \dots, 0) \in \mathbf{O}^n$ and $\underline{b} \in \mathbf{O}^n$ are not separated by $S_n^{(3)}$, then $G_2 \underline{a} = G_2 \underline{b}$.*

Proof. Given c_1, \dots, c_8 , denote by M_{c_1, \dots, c_8} the Gram matrix $(\text{tr}(c_i c_j))_{1 \leq i, j \leq 8}$. Since the trace is a bilinear non-degenerate form on \mathbf{O} and a_1, \dots, a_8 are linearly independent, we obtain that $\det(G_{a_1, \dots, a_8}) = \det(G_{b_1, \dots, b_8})$ is non-zero. Hence, b_1, \dots, b_8 are also linearly independent. In particular, \mathbb{F} -span of b_1, \dots, b_8 is \mathbf{O} .

For every $1 \leq i \leq 8$ and $8 < j \leq n$ we have that $\text{tr}(a_i a_j) = \text{tr}(b_i b_j)$ is zero. Therefore, $\text{tr}(b b_j) = 0$ for all $b \in \mathbf{O}$. Since tr is non-degenerate on \mathbf{O} , we obtain $\underline{b} = (b_1, \dots, b_8, 0, \dots, 0)$.

For every $1 \leq i, j \leq 8$ there exists $1 \leq k_{ij} \leq 8$ and $\eta_{ij} \in \mathbb{F}$ such that $a_i a_j = \eta_{ij} a_{k_{ij}}$. Therefore, for each $1 \leq l \leq 8$ we have that $\text{tr}((a_i a_j - \eta_{ij} a_{k_{ij}}) a_l) = \text{tr}((b_i b_j - \eta_{ij} b_{k_{ij}}) b_l)$ is zero. Hence, $b_i b_j = \eta_{ij} b_{k_{ij}}$. Consider a linear map $f : \mathbf{O} \rightarrow \mathbf{O}$ defined on the basis of \mathbf{O} by $f(a_i) = b_i$ for all $1 \leq i \leq 8$. Since the multiplication table for a_1, \dots, a_8 is the same as for b_1, \dots, b_8 , we can see that $f \in G_2$. The required is proven. \square

The following statement is a corollary of Proposition 6.1.

Corollary 7.11. *Assume $\text{char } \mathbb{F} = 2$ and a closed equivalence class $\mathfrak{A} \in \Omega(\mathbf{O})$ has the dimension $d \leq 3$. Then one of the following sets is a basis for \mathfrak{A} :*

- $d = 1$: $\{1_{\mathbf{O}}\}, \{e_1\}$;
- $d = 2$: $\{e_1, e_2\}$.

Proof. We need to show that any basis $\{a_1, \dots, a_n\}$ from Proposition 6.1, different from the above bases, generates non-closed equivalence class. For each $\underline{a} = (a_1, \dots, a_n)$ the arguments are the same: we find an element \underline{a}' in the closure of $G_2 \underline{a}$ such that $\text{rk}(\underline{a}') < \text{rk}(\underline{a})$, that obviously implies that $G_2 \underline{a}$ is not closed.

- If $\underline{a} = (\mathbf{u}_1) \in \mathbf{O}^1$, then for the standard one-parameter subgroup $\theta_{\underline{\lambda}}$ with $\underline{\lambda} = (1, -1, 0)$ the element $\underline{a}' = \lim_{t \rightarrow 0} \theta_{\underline{\lambda}}(t) \underline{a} = (0)$ lies in $\overline{G_2 \underline{a}}$ (see Section 3.3 for the more details).
- If $\underline{a} = (1_{\mathbf{O}}, \mathbf{u}_1)$, then $\underline{a}' = \lim_{t \rightarrow 0} \theta_{(1, -1, 0)}(t) \underline{a} = (1_{\mathbf{O}}, 0)$ lies in $\overline{G_2 \underline{a}}$.
- If $\underline{a} = (\mathbf{u}_1, \mathbf{v}_2)$, then $\underline{a}' = \lim_{t \rightarrow 0} \theta_{(1, -1, 0)}(t) \underline{a} = (0, 0)$ lies in $\overline{G_2 \underline{a}}$.
- If $\underline{a} = (e_1, \mathbf{u}_1)$, then $\underline{a}' = \lim_{t \rightarrow 0} \theta_{(1, -1, 0)}(t) \underline{a} = (e_1, 0)$ lies in $\overline{G_2 \underline{a}}$.
- If $\underline{a} = (e_1, \mathbf{v}_1)$, then $\underline{a}' = \lim_{t \rightarrow 0} \theta_{(-1, 1, 0)}(t) \underline{a} = (e_1, 0)$ lies in $\overline{G_2 \underline{a}}$.
- If $\underline{a} = (1_{\mathbf{O}}, \mathbf{u}_1, \mathbf{v}_2)$, then $\underline{a}' = \lim_{t \rightarrow 0} \theta_{(1, -1, 0)}(t) \underline{a} = (1_{\mathbf{O}}, 0, 0)$ lies in $\overline{G_2 \underline{a}}$.
- If $\underline{a} = (e_1, e_2, \mathbf{u}_1)$, then $\underline{a}' = \lim_{t \rightarrow 0} \theta_{(1, -1, 0)}(t) \underline{a} = (e_1, e_2, 0)$ lies in $\overline{G_2 \underline{a}}$.
- If $\underline{a} = (e_1, \mathbf{u}_1, \mathbf{v}_2)$, then $\underline{a}' = \lim_{t \rightarrow 0} \theta_{(1, -1, 0)}(t) \underline{a} = (e_1, 0, 0)$ lies in $\overline{G_2 \underline{a}}$.
- If $\underline{a} = (\mathbf{u}_1, \mathbf{v}_2, a)$ for $a = \alpha 1_{\mathbf{O}} + \beta \mathbf{u}_3 + \gamma \mathbf{v}_3$ with $\alpha^2 = \beta \gamma$ for some $\alpha, \beta, \gamma \in \mathbb{F}$, then $\underline{a}' = \lim_{t \rightarrow 0} \theta_{(1, -1, 0)}(t) \underline{a} = (0, 0, a)$ lies in $\overline{G_2 \underline{a}}$. \square

Theorem 7.12. *The set $S_n^{(8)}$ is a separating set for $K_n^{G_2}$ in case $\text{char } \mathbb{F} = 2$.*

Proof. We will apply Proposition 7.4 to obtain the required statement. Assume that G_2 -orbits of $\underline{a} = (a_1, \dots, a_l, 0, \dots, 0) \in \mathbf{O}^n$, $\underline{b} \in \mathbf{O}^n$ are closed, a_1, \dots, a_l is a basis of some subalgebra \mathcal{A} of \mathbf{O} , octonions b_1, \dots, b_n linearly generate some subalgebra \mathcal{B} of \mathbf{O} , and $\dim \mathcal{A} \geq \dim \mathcal{B}$. Moreover, we assume that \underline{a} and \underline{b} are not separated by $S_n^{(8)}$. By Lemma 7.2, we can choose any basis of \mathcal{A} as a_1, \dots, a_l .

Let $\dim \mathcal{A} = 8$. We choose $\underline{a} = (e_1, e_2, \mathbf{u}_1, \mathbf{v}_1, \mathbf{u}_2, \mathbf{v}_2, \mathbf{u}_3, \mathbf{v}_3, 0, \dots, 0)$. Then Lemma 7.10 implies that $G_2 \underline{a} = G_2 \underline{b}$.

Let $\dim \mathcal{A} < 8$. Then \mathcal{A} lies in a maximal proper subalgebra of \mathbf{O} . By Theorem 5 of [19], the algebra of sextonions \mathbb{S} is the unique maximal proper subalgebra of \mathbf{O} modulo G_2 -action. Therefore, without loss of generality we can assume that $\mathcal{A} \subset \mathbb{S}$. Similarly, we can assume that $\mathcal{B} \subset \mathbb{S}$.

Therefore, without loss of generality we can assume that $\mathcal{A} \subset \mathbb{S}$, i.e., for all $1 \leq i \leq l$ we have

$$a_i = \begin{pmatrix} \alpha_{i1} & (\alpha_{i2}, \alpha_{i3}, 0) \\ (\alpha_{i4}, 0, \alpha_{i5}) & \alpha_{i6} \end{pmatrix}$$

for some $\alpha_{ij} \in \mathbb{F}$. Similarly, we can assume that $\mathcal{B} \subset \mathbb{S}$.

Since for the standard one-parameter subgroup $\theta_{\underline{\lambda}}$ with $\underline{\lambda} = (0, 1, -1)$ we have

$$\theta_{\underline{\lambda}}(t)a_i = \begin{pmatrix} \alpha_{i1} & (\alpha_{i2}, t\alpha_{i3}, 0) \\ (\alpha_{i4}, 0, t\alpha_{i5}) & \alpha_{i6} \end{pmatrix},$$

then the limit $\underline{a}' = \lim_{t \rightarrow 0} \theta_{\underline{\lambda}}(t)\underline{a}$ exists (see Section 3.3 for more details). Obviously, $\underline{a}' = (a'_1, \dots, a'_l, 0, \dots, 0)$ lies in \mathbb{M}^n . The orbit $G_2 \underline{a}$ is closed, therefore, $\underline{a}' \in G_2 \underline{a}$. Thus, without loss of generality we can replace \underline{a} by \underline{a}' . Therefore, we can assume that $\underline{a} \subset \mathbb{M}^n$. Therefore, $\mathcal{A} \subset \mathbb{M}$. In the same manner we can assume that $\mathcal{B} \subset \mathbb{M}$.

In case $\dim \mathcal{A} = 4$ Lemma 7.9 implies that $G_2 \underline{a} = G_2 \underline{b}$.

Let $\dim \mathcal{A} \leq 3$. By Corollary 7.11 we can assume that \underline{a} is one of the next elements: $(1_{\mathbf{O}})$, (e_1) , (e_1, e_2) . If $\underline{a} = (1_{\mathbf{O}})$ or $\underline{a} = (e_1)$, then Lemma 7.7 implies that $G_2 \underline{a} = G_2 \underline{b}$. If $\underline{a} = (e_1, e_2)$, then Lemma 7.8 implies that $G_2 \underline{a} = G_2 \underline{b}$.

Finally, by Proposition 7.4 the set $S_n^{(8)}$ is separating for $K_n^{G_2}$. \square

Remark 7.13. It follows from Theorem 7.12 that the set $S_{0,n}^{(8)}$ is a separating set for $K_{0,n}^{G_2}$ in case $\text{char } \mathbb{F} = 2$.

REFERENCES

- [1] F. B. CAVALCANTE AND A. LOPATIN, *Separating invariants of three nilpotent 3×3 matrices*, Linear Algebra and its Applications, 607 (2020), pp. 9–28.
- [2] H. DERKSEN AND G. KEMPER, *Computational Invariant Theory*, Springer, Heidelberg, 2002.
- [3] ———, *Computational Invariant Theory (Second Edition)*, Springer, Heidelberg, 2015.
- [4] H. DERKSEN AND V. MAKAM, *Algorithms for orbit closure separation for invariants and semi-invariants of matrices*, Algebra and Number Theory, 14 (2020), pp. 2791–2813.
- [5] M. DOMOKOS, *Degree bound for separating invariants of abelian groups*, Proc. Amer. Math. Soc., 145 (2017), pp. 3695–3708.
- [6] ———, *Addendum to “Characteristic free description of semi-invariants of 2×2 matrices”* [*J. Pure Appl. Algebra* 224 (2020), no. 5, 106220], Journal of Pure and Applied Algebra, 224 (2020), p. 106270.
- [7] ———, *Characteristic free description of semi-invariants of 2×2 matrices*, Journal of Pure and Applied Algebra, 224 (2020), p. 106220.
- [8] M. DOMOKOS, S. KUZMIN, AND A. ZUBKOV, *Rings of matrix invariants in positive characteristic*, J. Pure Appl. Algebra, 176 (2002), pp. 61–80.

- [9] E. DUFRESNE, J. ELMER, AND M. SEZER, *Separating invariants for arbitrary linear actions of the additive group*, Manuscripta Mathematica, 143 (2014), pp. 207–219.
- [10] A. N. GRISHKOV, M. DE LOURDES MERLINI GIULIANI, AND A. V. ZAVARNITSINE, *Classification of subalgebras of the cayley algebra over a finite field*, Journal of Algebra and its Applications, 9 (2010), pp. 791–808.
- [11] A. ILTYAKOV AND I. SHESTAKOV, *On invariants of F_4 and the center of the Albert algebra*, Journal of Algebra, 179 (1996), pp. 838–851.
- [12] I. KAYGORODOV, A. LOPATIN, AND Y. POPOV, *Separating invariants for 2×2 matrices*, Linear Algebra and its Applications, 559 (2018), pp. 114–124.
- [13] G. KEMPER, A. LOPATIN, AND F. REIMERS, *Separating invariants over finite fields*, Journal of Pure and Applied Algebra, 226 (2022), p. 106904.
- [14] G. R. KEMPF, *Instability in invariant theory*, Ann. of Math. (2), 108 (1978), pp. 299–316.
- [15] M. KOHLS AND M. SEZER, *Separating invariants for the Klein four group and cyclic groups*, International Journal of Mathematics, 24 (2013), p. 1350046.
- [16] J. M. LANDSBERG AND L. MANIVEL, *The sextonions and $E_{7\frac{1}{2}}$* , Advances in Mathematics, 201 (2006), pp. 143–179.
- [17] A. LOPATIN AND F. REIMERS, *Separating invariants for multisymmetric polynomials*, Proceedings of the American Mathematical Society, 149 (2021), pp. 497–508.
- [18] A. LOPATIN AND A. N. ZUBKOV, *Classification of G_2 -orbits for pairs of octonions*, arXiv: 2208.08122.
- [19] M. RACINE, *On maximal subalgebras*, J. Algebra, 30 (155–180), p. 1974.
- [20] G. SCHWARZ, *Invariant theory of G_2 and Spin_7* , Comment. Math. Helvetici, 63 (1988), pp. 624–663.
- [21] A. N. ZUBKOV AND I. SHESTAKOV, *Invariants of G_2 and $\text{Spin}(7)$ in positive characteristic*, Transformation Groups, 23 (2018), pp. 555–588.

ARTEM LOPATIN, STATE UNIVERSITY OF CAMPINAS, 651 SERGIO BUARQUE DE HOLANDA, 13083-859 CAMPINAS, SP, BRAZIL

Email address: dr.artem.lopatin@gmail.com (Artem Lopatin)

ALEXANDR N. ZUBKOV, DEPARTMENT OF MATHEMATICAL SCIENCES, UNITED ARAB EMIRATES UNIVERSITY, AL-AIN, UNITED ARAB EMIRATES; SOBOLEV INSTITUTE OF MATHEMATICS, OMSK BRANCH, PEVTZOVA 13, 644043 OMSK, RUSSIA

Email address: a.zubkov@yahoo.com (Alexandr N. Zubkov)