

REVISITING JACOBI-TRUDI IDENTITIES VIA THE BGG CATEGORY \mathcal{O}

TAO GUI AND ARTHUR L. B. YANG

ABSTRACT. Using the BGG resolution, Akin presented a realization of the Jacobi-Trudi expansion of a Schur polynomial as the Euler-Poincaré characteristic of an exact sequence in the category of polynomial representations of the general linear groups. Zelevinskii gave an independent treatment for a general skew Schur polynomial. Based on their construction of resolutions, one can obtain the Schur positivity of certain truncations of the Jacobi-Trudi expansion of a skew Schur polynomial. By interpreting Kostka numbers as tensor product multiplicities in the BGG category \mathcal{O} of the special linear Lie algebras, we establish a new relationship between the Schur positivity of these truncations of Jacobi-Trudi expansion and the BGG resolution. Specifically, we show that coefficients in the combination of the Schur polynomial of these truncations of Jacobi-Trudi expansion also appear as tensor product multiplicities in the BGG category \mathcal{O} . This point of view also enables us to derive the Jacobi-Trudi identity directly from the celebrated Weyl character formula.

Acknowledgments The first author would like to thank Ming Fang, Hongsheng Hu, and Nanhua Xi for useful comments. The second author is supported in part by the Fundamental Research Funds for the Central Universities and the National Science Foundation of China (Nos.11522110, 11971249).

1. INTRODUCTION

Many open problems in symmetric function theory are related to the *Schur positivity* questions, that is, showing that certain symmetric functions can be expressed as non-negative linear combinations of *Schur functions*; see, for example, [5, 7, 8, 11, 15]. For our purpose here, we only need to consider *Schur polynomials*, which are obtained from Schur functions by reducing the number of variables. The significance of Schur positivity stems from the fact that Schur polynomials not only have interesting combinatorial interpretations but also admit natural algebraic or geometric meanings. In combinatorics, Schur polynomials can be defined as generating functions of weighted semi-standard Young tableaux. In representation theory, they appear as characters of finite-dimensional irreducible polynomial representations (known as *Schur modules*) of the general linear groups $GL_n(\mathbb{C})$, as well as Frobenius images of irreducible representations of the symmetric group. The relationship between Schur polynomials and representation theory was discovered by Schur, hence these important polynomials are named after him. In geometry, Schur polynomials appear as the cohomology classes of the Schubert cycles of the corresponding Grassmannian. For more information on Schur polynomials, we refer the reader to [6, 13, 16]. Due to these connections, whether a particular symmetric

Key words and phrases. Jacobi-Trudi identity, Schur positivity, Kostka number, BGG category \mathcal{O} , Weyl character formula, BGG resolution.

polynomial can be expanded positively in terms of Schur polynomials is of particular interest in symmetric function theory, representation theory, and classical Schubert calculus.

The main objective of this paper is to study the Schur positivity of certain truncations of the Jacobi-Trudi expansion of skew Schur polynomials. For ordinary partitions, this positivity can be implied by the existence of a conjectural resolution (due to Lascoux [12]) of irreducible representations of $GL_n(\mathbb{C})$, which was later constructed by Akin[1] and Zelevinskii[17] independently. Zelevinskii's approach could also handle the case of skew partitions. Although the Schur positivity of these truncations of Jacobi-Trudi expansions is known, it is an open problem to give a combinatorial interpretation of these non-negative coefficients in the Schur polynomial expansion[16, p.461]. By the work of Akin and Zelevinskii, one can interpret these coefficients as multiplicities of irreducible representations of some polynomial representation of $GL_n(\mathbb{C})$. Using the classical BGG resolution, we show that these coefficients also appear as tensor product multiplicities in the BGG category \mathcal{O} of the special linear Lie algebra $\mathfrak{sl}_n(\mathbb{C})$, thus providing a new algebraic interpretation.

In order to state our results explicitly, let us now recall some related definitions and results. We will adopt the terminology and notation of symmetric functions in [16]. An integer partition λ of a positive integer d , denoted by $\lambda \vdash d$, is a weakly decreasing sequence $(\lambda_1, \lambda_2, \dots, \lambda_n)$ of non-negative integers such that $\sum_{i=1}^n \lambda_i = d$. For each partition λ , let m_λ , h_λ , s_λ denote the corresponding monomial symmetric polynomial, the complete symmetric polynomial, and the Schur polynomial, respectively. It is known that each of $\{m_\lambda \mid \lambda \vdash d\}$, $\{h_\lambda \mid \lambda \vdash d\}$, and $\{s_\lambda \mid \lambda \vdash d\}$ is a basis of the space of homogeneous symmetric polynomials of degree d . Here we adopt the definition of the Schur polynomial s_λ given by

$$(1.1) \quad s_\lambda := \sum_{\mu} K_{\lambda, \mu} m_{\mu},$$

where each $K_{\lambda, \mu}$, called the *Kostka number*, is equal to the number of semistandard Young tableaux of shape λ and type μ .

If μ, ν are two partitions such that $\mu_i \geq \nu_i$ for each i , then one can define a skew partition μ/ν . Let $s_{\mu/\nu}$ denote the corresponding skew Schur polynomial indexed by μ/ν , which can be defined by the property

$$(1.2) \quad \langle s_{\mu/\nu}, s_\lambda \rangle = \langle s_\mu, s_\lambda s_\nu \rangle,$$

where the scalar product is defined by claiming that Schur polynomials form an orthonormal basis. The *Jacobi-Trudi identity* states that

$$(1.3) \quad s_{\mu/\nu} = \det (h_{\mu_i - \nu_j - i + j})_{i, j=1}^n,$$

where we set $h_0 = 1$ and $h_k = 0$ for $k < 0$.

There are several proofs of the Jacobi-Trudi identity. The classical combinatorial proof is to relate the semi-standard Young tableaux with non-intersection lattice paths and then to use the Lindström–Gessel–Viennot lemma; see [14, Section 4.5], or the first proof of [16, Theorem 7.16.1]. A purely algebraic proof can also be given; see [13, Section 1.3], or the second proof of [16, Theorem 7.16.1]. Since Schur polynomials also appear as characters of Schur modules of the general linear groups, the alternating sum format in (1.3) resembles an *Euler-Poincaré characteristic*, so

it is natural to ask whether there is a resolution of Schur modules to realize the identity (1.3) at least for the case of $\nu = \emptyset$.

Now consider the Grothendieck ring of the category of polynomial representations of $GL_n(\mathbb{C})$. Let $V(\mu)$ denote the irreducible polynomial representation of $GL_n(\mathbb{C})$ of highest weight $\mu = (\mu_1, \mu_2, \dots, \mu_n) \in \mathbb{Z}^n$, relative to the maximal torus $T \subseteq GL_n(\mathbb{C})$ of diagonal matrices. For any weight $\tau = (\tau_1, \dots, \tau_n) \in \mathbb{Z}^n$, let $S(\tau)$ denote the representation of $GL_n(\mathbb{C})$ formed by taking the tensor product $S^{\tau_1}(V) \otimes \dots \otimes S^{\tau_n}(V)$ of symmetric powers of the standard representation $V = \mathbb{C}^n$. It is known that the character of $S(\tau)$ is the complete symmetric polynomial $h_\tau = h_{\tau_1} h_{\tau_2} \dots h_{\tau_n}$. Thus the Jacobi-Trudi identity (1.3) for the case of $\nu = \emptyset$ can be written as

$$(1.4) \quad [V(\mu)] = \sum_{w \in S_n} (-1)^{\ell(w)} [S(\mu - w \cdot \emptyset)]$$

in the Grothendieck ring, where the dot action of W on \mathbb{Z}^n is defined by the twisted rule $w \cdot \tau = w(\tau + \delta) - \delta$, and δ denotes the sequence $(n-1, n-2, \dots, 1, 0) \in \mathbb{Z}^n$. Let $Y_i(\mu)$ be the direct sum $\bigoplus S(\mu - w \cdot \emptyset)$ over all permutations w of length $\ell(w) = i$. Lascoux stated in [12] the existence of an exact sequence of the following form:

$$(1.5) \quad 0 \rightarrow Y_{\binom{n}{2}}(\mu) \rightarrow \dots \rightarrow Y_1(\mu) \rightarrow Y_0(\mu) \rightarrow V(\mu) \rightarrow 0,$$

but he did not define the boundary maps. It is clear that such an exact sequence categorifies the identity in (1.4). A rigorous treatment of Lascoux's work was subsequently given by Akin in [1–3], and independently by [17].

For general μ/ν , let $V(\mu/\nu)$ denote the $GL_n(\mathbb{C})$ -module with character $s_{\mu/\nu}$, and let $Y_i(\mu/\nu)$ be the direct sum $\bigoplus S(\mu - w \cdot \nu)$ over all permutations w of length $\ell(w) = i$. With some work (by replacing $\nu = \emptyset$ in [17, Example 1] with a general ν and doing some deductions), the approach of Zelevinskii can be used to construct the following exact sequence

$$(1.6) \quad 0 \rightarrow Y_{\binom{n}{2}}(\mu/\nu) \xrightarrow{\partial_{\binom{n}{2}}} \dots \rightarrow Y_1(\mu/\nu) \xrightarrow{\partial_1} Y_0(\mu/\nu) \xrightarrow{\partial_0} V(\mu/\nu) \rightarrow 0.$$

The existence of such a resolution has the following implication, see [16, Exercise 7.38].

Theorem 1 (Akin-Zelevinskii theorem). *For $0 \leq k \leq \binom{n}{2}$, the symmetric polynomial*

$$(1.7) \quad g_{\mu/\nu}^k = (-1)^k \sum_{w \in S_n, \ell(w) \geq k} (-1)^{\ell(w)} h_{\mu + \delta - w(\nu + \delta)},$$

a truncation of the Jacobi-Trudi expansion of $s_{\lambda/\mu}$, is Schur positive, where μ and ν are partitions of length at most n , $\ell(w)$ is the number of inversions of the permutation $w \in S_n$, and $\delta = (n-1, n-2, \dots, 1, 0)$.

The Schur positivity of $g_{\mu/\nu}^k$ follows from the existence of (1.6), since it is not hard to see that $g_{\mu/\nu}^k$ is indeed the character of the image of the boundary map ∂_k by truncating (1.6). The boundary maps were explicitly described in [3] for the case of $\nu = \emptyset$, while for general ν , they were only given implicitly in [17].

In this paper, we give a new interpretation of the coefficients in the Schur polynomial expansion of $g_{\mu/\nu}^k$ via the BGG category \mathcal{O} of $\mathfrak{sl}_n(\mathbb{C})$. We have the following result, which in turn provides a new proof of the Schur positivity of $g_{\mu/\nu}^k$.

Theorem 2. For $0 \leq k \leq \binom{n}{2}$, we have

$$(1.8) \quad g_{\mu/\nu}^k = \sum_{\lambda} [L(\lambda) \otimes V(\nu, k) : L(\mu)]_{s_{\lambda}},$$

where $L(\lambda)$ (and $L(\mu)$, respectively) denotes the simple module of highest weight λ (and μ , respectively) of the special linear Lie algebra $\mathfrak{sl}_n(\mathbb{C})$, $V(\nu, k)$ is the image of the boundary map d_k in the BGG resolution of $L(\nu)$:

$$(1.9) \quad 0 \rightarrow C_{\binom{n}{2}}^{\nu} \xrightarrow{d_{\binom{n}{2}}} C_{\binom{n}{2}-1}^{\nu} \rightarrow \cdots \rightarrow C_1^{\nu} \xrightarrow{d_1} C_0^{\nu} \xrightarrow{d_0} L(\nu) \rightarrow 0,$$

and the bracket denotes the composition factor multiplicity in the BGG category \mathcal{O} .

Our approach is different in the sense that we transform the Schur positivity problem into the positivity problem of a virtual character in the BGG category \mathcal{O} and then use classical BGG resolution to prove the positivity. It also allows us to give a new proof of the Jacobi-Trudi identity (1.3). Actually, we can derive it from the Weyl character formula, as illustrated in subsequent sections.

With Candice X. T. Zhang, we observed that truncations of the Jacobi-Trudi expansion of a skew Schur polynomial seem to share similar convexity of Schur polynomials, see [9]. Specifically, we have checked that for partitions μ with less than or equal to 9 boxes, all *normalized* truncations of the Jacobi-Trudi expansion of a Schur polynomial $g_{\mu/\emptyset}^k$ are *Lorentzian*. It would be interesting to see whether our approach can be used to understand the Lorentzian property for the skew Schur polynomials, see [9, Conjecture 19].

The remaining part of this paper is organized as follows. In Section 2, we provide preliminary results on the BGG category \mathcal{O} and the classical BGG resolution. In Section 3, we first give a new interpretation of Kostka numbers in the BGG category \mathcal{O} , and then leave the world of symmetric functions and use Verma module characters to prove the Jacobi-Trudi identity and Theorem 2.

2. PRELIMINARIES

In this section, we recall the relevant representation theory of complex semi-simple Lie algebras. For a more detailed background, we refer the reader to the comprehensive reference [10].

2.1. The BGG category \mathcal{O} . Let \mathfrak{g} be a finite-dimensional complex semi-simple Lie algebra (for our application, $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{C})$). We fix a Cartan decomposition $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$. (In the case of $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{C})$, we could choose \mathfrak{n}_+ and \mathfrak{n}_- to be the strictly upper-triangular and strictly lower-triangular matrices, respectively, and \mathfrak{h} to be the diagonal matrices.) We consider the *Bernstein-Gelfand-Gelfand category* \mathcal{O} , which is a full subcategory of $\text{Mod } U(\mathfrak{g})$ (the category of left-modules of the enveloping algebra of \mathfrak{g}). Each object of \mathcal{O} is a module M satisfying the following three conditions:

- (O1) M is a finitely generated $U(\mathfrak{g})$ -module.
- (O2) M is \mathfrak{h} -semi-simple, that is, M is a weight module: $M = \bigoplus_{\lambda \in \mathfrak{h}^*} M_{\lambda}$, where $M_{\lambda} = \{m \in M : h \cdot m = \lambda(h)m \text{ for all } h \in \mathfrak{h}\}$.
- (O3) M is locally \mathfrak{n}_+ -finite, that is, for each $v \in M$, the subspace $U(\mathfrak{n}_+) \cdot v$ of M is finite-dimensional.

For example, all finite-dimensional modules lie in \mathcal{O} . It follows from the axioms that all weight spaces of $M \in \mathcal{O}$ are finite-dimensional. Thus one could define the (formal) character of $M \in \mathcal{O}$ by

$$\text{ch}(M) = \sum_{\lambda} \dim(M_{\lambda}) x^{\lambda},$$

where the symbols x^{λ} denote formal exponentials, multiplying by the rule $x^{\lambda}x^{\mu} = x^{\lambda+\mu}$. Furthermore, if $M \in \mathcal{O}$ and L is finite-dimensional, then $L \otimes M \in \mathcal{O}$ and $\text{ch}(L \otimes M) = \text{ch } L \text{ ch } M$.

For each $\lambda \in \mathfrak{h}^*$, there is a one-dimensional representation \mathbb{C}_{λ} of $\mathfrak{b} := \mathfrak{h} \oplus \mathfrak{n}_+$, where \mathfrak{h} acts via λ and \mathfrak{n}_+ acts by zero. Now consider the Verma (or standard) module $\Delta(\lambda)$, defined as $\Delta(\lambda) := U(\mathfrak{g}) \otimes_{U(\mathfrak{b})} \mathbb{C}_{\lambda}$, where $U(\mathfrak{g})$ and $U(\mathfrak{b})$ are the universal enveloping algebras of \mathfrak{g} and \mathfrak{b} , respectively. It can be proved that simple objects $L(\lambda)$ (simple highest weight module) in \mathcal{O} are parametrized by $\lambda \in \mathfrak{h}^*$ and can be uniformly constructed as the unique simple quotient of the corresponding $\Delta(\lambda)$.

Since each $M \in \mathcal{O}$ is both artinian and noetherian, it follows that M possesses a composition series with simple quotients isomorphic to various $L(\lambda)$. The multiplicity of $L(\lambda)$ in M is independent of the choice of composition series and is denoted by $[M : L(\lambda)]$. It is known that category \mathcal{O} is an abelian category and if $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ is a short exact sequence in \mathcal{O} , then $\text{ch } M = \text{ch } M' + \text{ch } M''$. Thus any $\text{ch } M$ is determined uniquely by the formal characters of the composition factors of M , taken with multiplicity.

2.2. Weyl character formula and BGG resolution. One of the basic questions in the representation theory of the complex semi-simple Lie algebra \mathfrak{g} is to compute the character of $L(\lambda)$. Those $L(\lambda)$ with λ dominant and integral are precisely all the finite-dimensional simple modules of \mathfrak{g} , and the characters are given by the celebrated Weyl character formula ¹

$$(2.1) \quad \text{ch } L(\lambda) = \frac{\sum_{w \in W} (-1)^{\ell(w)} x^{w(\lambda + \rho)}}{\sum_{w \in W} (-1)^{\ell(w)} x^{w(\rho)}},$$

where W is the Weyl group, and ρ is the Weyl vector, defined as half the sum of positive roots. For $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{C})$, the Weyl group W is the permutation group S_n , and ρ corresponds to the partition $\delta = (n-1, n-2, \dots, 1, 0)$.

In terms of characters of Verma modules, the Weyl character formula is equivalent to

$$(2.2) \quad \text{ch } L(\lambda) = \sum_{w \in W} (-1)^{\ell(w)} \text{ch } \Delta(w \cdot \lambda),$$

where the dot action is defined as $w \cdot \lambda = w(\lambda + \rho) - \rho$, in other words, this shifts the standard action of W to have center $-\rho$, and the characters of the Verma modules are given by

$$(2.3) \quad \text{ch } (\Delta(\lambda)) = x^{\lambda} / \prod (1 - x^{-\alpha}),$$

¹It is easy to see that Cauchy's bialternant formula for the Schur polynomial $s_{\lambda}(x_1, x_2, \dots, x_n) = \det(x_j^{\lambda_i + n - i}) / \det(x_j^{n - i})$, where $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$ is a partition, is a special case of the Weyl character formula for $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{C})$ and $W = S_n$.

where α ranges over all positive roots of the complex semi-simple Lie algebra \mathfrak{g} .

In a classical paper [4], Bernstein, Gelfand and Gelfand studied the Weyl character formula and constructed a resolution of $L(\lambda)$ for λ dominant and integral to realize the alternating sum in the Weyl character formula (2.2) as an Euler-Poincaré characteristic of a complex, which is now known as the *BGG resolution*. Specifically, they proved the following theorem.

Theorem 3. [4, Theorem 10.1] *Let λ be a dominant and integral weight and $L(\lambda)$ be the irreducible finite-dimensional \mathfrak{g} -module with highest weight λ . Then there exists an exact sequence of \mathfrak{g} -modules*

$$(2.4) \quad 0 \rightarrow C_{\binom{n}{2}}^\lambda \xrightarrow{d_{\binom{n}{2}}} C_{\binom{n}{2}-1}^\lambda \rightarrow \cdots \rightarrow C_1^\lambda \xrightarrow{d_1} C_0^\lambda \xrightarrow{d_0} L(\lambda) \rightarrow 0,$$

where $C_k^\lambda = \bigoplus_{w \in W, \ell(w)=k} \Delta(w \cdot \lambda)$.

For general $\lambda \in \mathfrak{h}^*$, it is not hard ² to show that

$$(2.5) \quad \text{ch } L(\lambda) = \sum_{w \cdot \lambda \leq \lambda} b(\lambda, w) \text{ch } \Delta(w \cdot \lambda) \text{ with } b(\lambda, w) \in \mathbb{Z} \text{ and } b(\lambda, \lambda) = 1,$$

where the partial ordering is defined by $\mu \leq \lambda$ if and only if $\lambda - \mu$ is a non-negative linear combination of simple roots. That is, $b(\lambda, w)$ is uni-triangular with respect to the partial ordering. It follows that formal characters $\text{ch } L(\lambda)$ with $\lambda \in \mathfrak{h}^*$ are linearly independent and form a \mathbb{Z} -basis of the additive group generated by all $\text{ch } \Delta(\lambda)$, while formal characters of Verma modules provide an alternative \mathbb{Z} -basis. The multiplicity $[M : L(\lambda)]$ of $L(\lambda)$ in M is equal to the coefficient of $\text{ch } L(\lambda)$ in the expansion of $\text{ch } M$ in terms of irreducible characters.

3. THE MAIN RESULTS

The purpose of this section is two-fold: first, we give a new proof of the Jacobi-Trudi identity, and secondly, we use BGG resolution to prove Theorem 2.

Our approach given here relies on an interpretation of the Kostka numbers as tensor product multiplicities in the BGG category \mathcal{O} of $\mathfrak{sl}_n(\mathbb{C})$. Note that

$$(3.1) \quad h_\tau = \sum_{\lambda} K_{\lambda, \tau} s_\lambda,$$

which is equivalent to the definition (1.1) of the Schur polynomial, see [6, Section 6.1]. On the other hand, since the Schur polynomial s_λ is the character of the finite-dimensional irreducible polynomial representation V_λ (the Schur module) of $GL_n(\mathbb{C})$, by restricting to the special linear group $SL_n(\mathbb{C})$ and differentiating, it is also the formal character of the simple module $L(\lambda)$ of the special linear Lie algebra $sl_n(\mathbb{C})$, namely,

$$(3.2) \quad \text{ch } L(\lambda) = s_\lambda, \text{ for } \lambda \text{ dominant and integral, i.e., } \lambda \text{ is a partition.}$$

It follows that the Kostka number $K_{\lambda, \tau}$ is also the weight multiplicity of τ in $L(\lambda)$, namely,

$$(3.3) \quad K_{\lambda, \tau} = \dim(L(\lambda)_\tau),$$

²However, the determination of the coefficients $b(\lambda, w)$ is rather difficult, and they are given by the famous Kazhdan-Lusztig polynomials in virtue of the Kazhdan-Lusztig conjecture. We do not need the exact formula in this paper.

where λ is a partition, regarded as a dominant weight of $\mathfrak{sl}_n(\mathbb{C})$, and τ is any sequence $(\tau_1, \dots, \tau_n) \in \mathbb{Z}^n$ with $|\lambda| = |\tau|$. By the Weyl character formula (2.2) and the characters of the Verma modules (2.3), we have

$$\begin{aligned}
 (3.4) \quad K_{\lambda, \tau} &= \dim(L(\lambda)_\tau) \\
 &= \sum_{v \in S_n} (-1)^{\ell(v)} \dim(\Delta((v \cdot \lambda)_\tau)) \\
 &= \sum_{v \in S_n} (-1)^{\ell(v)} \mathfrak{p}(v \cdot \lambda - \tau),
 \end{aligned}$$

where \mathfrak{p} is the Kostant partition function. Recall that \mathfrak{p} is defined by

$$(3.5) \quad \frac{1}{\prod (1 - e^{-\alpha})} = \sum_{\mu} \mathfrak{p}(\mu) e^{-\mu},$$

where α ranges over all positive roots of $sl_n(\mathbb{C})$.

To prove Theorem 2, we use (3.4) to give an alternative interpretation of the Kostka numbers. In view of (1.7) and (3.1), we directly consider the Kostka numbers of the form $K_{\lambda, \mu + \delta - w(\nu + \delta)}$. The following result plays a key role for our proof of Theorem 2.

Proposition 4. *The Kostka number $K_{\lambda, \mu + \delta - w(\nu + \delta)}$ is equal to $[L(\lambda) \otimes \Delta(w \cdot \nu) : L(\mu)]$, the multiplicity of irreducible composition factor $L(\mu)$ in $L(\lambda) \otimes \Delta(w \cdot \nu)$, where λ , μ and ν are partitions, regarded as dominant weights for $\mathfrak{sl}_n(\mathbb{C})$, and $\delta = (n-1, n-2, \dots, 1, 0)$. In particular, $K_{\lambda, \mu} = [L(\lambda) \otimes \Delta(0) : L(\mu)]$.*

Proof. One can use the *standard filtration* (also known as the *Verma flag*) to prove this proposition; see ([10, Theorem 3.6]). Note that μ corresponds to a dominant weight, so the multiplicity with which $\Delta(\mu)$ occurs in a standard filtration of $L(\lambda) \otimes \Delta(w \cdot \nu)$ is equal to the multiplicity of $L(\mu)$ in a Jordan-Hölder series of $L(\lambda) \otimes \Delta(w \cdot \nu)$.

Here we give a direct computational proof using formal characters. If f is a formal character, let $[\text{ch } L(\lambda)]f$ denote the coefficient of $\text{ch } L(\lambda)$ in the expansion of f in terms of irreducible characters, $[\text{ch } \Delta(\lambda)]f$ denote the coefficient of $\text{ch } \Delta(\lambda)$ in its expansion by Verma module characters, and let $[x^\lambda]f$ denote the coefficient of x^λ in f . Note that

$$[L(\lambda) \otimes \Delta(w \cdot \nu) : L(\mu)] = [\text{ch } L(\mu)] \text{ch } L(\lambda) \text{ch } \Delta(w \cdot \nu).$$

Since μ is dominant, firstly we have

$$[\text{ch } L(\mu)] \text{ch } L(\lambda) \text{ch } \Delta(w \cdot \nu) = [\text{ch } \Delta(\mu)] \text{ch } L(\lambda) \text{ch } \Delta(w \cdot \nu)$$

by the uni-triangularity (2.5). Then, by the Weyl character formula (2.2), we obtain

$$[\text{ch } \Delta(\mu)] \text{ch } L(\lambda) \text{ch } \Delta(w \cdot \nu) = [\text{ch } \Delta(\mu)] \left(\sum_{v \in S_n} (-1)^{\ell(v)} \text{ch } \Delta(v \cdot \lambda) \text{ch } \Delta(w \cdot \nu) \right).$$

Furthermore, by (2.3) and (3.5), we have

$$\begin{aligned}
 [\text{ch } \Delta(\mu)] \text{ch } \Delta(v \cdot \lambda) \text{ch } \Delta(w \cdot \nu) &= [x^\mu] x^{v \cdot \lambda + w \cdot \nu} / \prod (1 - x^{-\alpha}) \\
 &= \mathfrak{p}(v \cdot \lambda + w \cdot \nu - \mu).
 \end{aligned}$$

Combining the above identities and (3.4) leads to

$$[\text{ch } L(\mu)] \text{ch } L(\lambda) \text{ch } \Delta(w \cdot \nu) = \sum_{v \in S_n} (-1)^{\ell(v)} \mathbf{p}(v \cdot \lambda + w \cdot \nu - \mu) = K_{\lambda, \mu + \delta - w(\nu + \delta)},$$

as desired. In particular, taking $w = \text{id}$ and $\nu = \emptyset$, we get

$$K_{\lambda, \mu} = [L(\lambda) \otimes \Delta(0) : L(\mu)].$$

The proof is complete. \square

The above proposition interprets Kostka numbers as tensor product multiplicities in the BGG category \mathcal{O} . With this interpretation, now we are able to give a new proof of the Jacobi-Trudi identity.

A new proof of (1.3). On the one hand, in view of $\text{ch } L(\lambda) = s_\lambda$ and (1.2), we have

$$\langle s_{\mu/\nu}, s_\lambda \rangle = \langle s_\mu, s_\lambda s_\nu \rangle = [\text{ch } L(\mu)] \text{ch } L(\lambda) \text{ch } L(\nu) = [L(\lambda) \otimes L(\nu) : L(\mu)],$$

and thus

$$s_{\mu/\nu} = \sum_{\lambda} [L(\lambda) \otimes L(\nu) : L(\mu)] s_\lambda.$$

On the other hand, by (3.1) and proposition 4, we have

$$\begin{aligned} \det (h_{\mu_i - \nu_j + j - i})_{i, j=1}^n &= \sum_{w \in S_n} (-1)^{\ell(w)} h_{\mu + \delta - w(\nu + \delta)} \\ &= \sum_{w \in S_n} \sum_{\lambda} (-1)^{\ell(w)} K_{\lambda, \mu + \delta - w(\nu + \delta)} s_\lambda \\ &= \sum_{w \in S_n} \sum_{\lambda} (-1)^{\ell(w)} [L(\lambda) \otimes \Delta(w \cdot \nu) : L(\mu)] s_\lambda \\ &= \sum_{\lambda} [L(\lambda) \otimes L(\nu) : L(\mu)] s_\lambda, \end{aligned}$$

where the last equality follows from the Weyl character formula (2.2). This completes the proof of the Jacobi-Trudi identity (1.3). \square

Remark 5. In particular, for the case $\nu = \emptyset$ of the Jacobi-Trudi identity, we have

$$\begin{aligned} \det (h_{\mu_i + j - i}) &= \sum_{w \in S_n} \sum_{\lambda} (-1)^{\ell(w)} [L(\lambda) \otimes \Delta(w \cdot 0) : L(\mu)] s_\lambda \\ &= \sum_{\lambda} [L(\lambda) \otimes L(0) : L(\mu)] s_\lambda \\ &= \sum_{\lambda} [L(\lambda) : L(\mu)] s_\lambda \\ &= s_\mu, \end{aligned}$$

which follows from the Weyl character formula (2.2) for the trivial module $L(0)$.

We proceed to prove Theorem 2.

Proof of Theorem 2. By Theorem 3, there exists an exact sequence of $\mathfrak{sl}_n(\mathbb{C})$ -modules

$$(3.6) \quad 0 \rightarrow C_{\binom{n}{2}}^\nu \xrightarrow{d_{\binom{n}{2}}} C_{\binom{n}{2}-1}^\nu \rightarrow \cdots \rightarrow C_1^\nu \xrightarrow{d_1} C_0^\nu \xrightarrow{d_0} L(\nu) \rightarrow 0,$$

where $C_k^\nu = \bigoplus_{w \in S_n, \ell(w)=k} \Delta(w \cdot \nu)$. For $0 \leq k \leq \binom{n}{2}$, let $V(\nu, k)$ be the image of the boundary map d_k in the BGG resolution of $L(\nu)$, and consider the exact sequence by truncating (3.6):

$$(3.7) \quad 0 \rightarrow C_{\binom{n}{2}}^\nu \rightarrow C_{\binom{n}{2}-1}^\nu \rightarrow \cdots \rightarrow C_k^\nu \xrightarrow{d_k} V(\nu, k) \rightarrow 0.$$

Taking the formal characters of the exact sequence (3.7), we obtain

$$(3.8) \quad \text{ch}(V(\nu, k)) = \text{ch}(\text{Im}(d_k)) = (-1)^k \sum_{w \in S_n, \ell(w) \geq k} (-1)^{\ell(w)} \text{ch}(\Delta(w \cdot \nu)).$$

Now by Proposition 4, we have

$$\begin{aligned} g_{\mu/\nu}^k &= (-1)^k \sum_{w \in S_n, \ell(w) \geq k} (-1)^{\ell(w)} h_{\mu+\delta-w(\nu+\delta)} \\ &= (-1)^k \sum_{w \in S_n, \ell(w) \geq k} \sum_{\lambda} (-1)^{\ell(w)} K_{\lambda, \mu+\delta-w(\nu+\delta)} s_\lambda \\ &= (-1)^k \sum_{w \in S_n, \ell(w) \geq k} \sum_{\lambda} (-1)^{\ell(w)} [L(\lambda) \otimes \Delta(w \cdot \nu) : L(\mu)] s_\lambda \\ &= (-1)^k \sum_{w \in S_n, \ell(w) \geq k} \sum_{\lambda} (-1)^{\ell(w)} [\text{ch } L(\mu)] \text{ch } L(\lambda) \text{ch } \Delta(w \cdot \nu) s_\lambda \\ &= \sum_{\lambda} [\text{ch } L(\mu)] \text{ch } L(\lambda) \text{ch } V(\nu, k) s_\lambda \\ &= \sum_{\lambda} [L(\lambda) \otimes V(\nu, k) : L(\mu)] s_\lambda \end{aligned}$$

The proof is complete. \square

Remark 6. The usual representation theory approach to establish Schur positivity of a given symmetric polynomial is to show that it is the characters of some finite-dimensional polynomial representation of the general linear groups or the Frobenius images of some representation of the symmetric groups. As it can be seen from the proof, our approach is to transform the Schur positivity problem into the positivity problem of a virtual character in the BGG category \mathcal{O} of $sl_n(\mathbb{C})$, see (3.8). It would be interesting to know whether there are other Schur positivity problems can be handled using BGG category \mathcal{O} techniques.

REFERENCES

- [1] Kaan Akin, *On complexes relating the Jacobi-Trudi identity with the Bernstein-Gelfand-Gelfand resolution*, J. Algebra **117** (1988), no. 2, 494–503.
- [2] ———, *Resolutions of representations*, Contemp. Math. **88** (1989), 209–217.
- [3] ———, *On complexes relating the Jacobi-Trudi identity with the Bernstein-Gelfand-Gelfand resolution, II*, J. Algebra **152** (1992), no. 2, 417–426.
- [4] IN Bernstein, Israel M Gelfand, and Sergei I Gelfand, *Differential operators on the base affine space and a study of g -modules*, Lie groups and their representations (Proc. Summer School, Bolyai János Math. Soc., Budapest, 1971) (1975), 21–64.
- [5] Vyjayanthi Chari, Ghislain Fourier, and Daisuke Sagaki, *Posets, tensor products and Schur positivity*, Algebra Number Theory **8** (2014), no. 4, 933–961.
- [6] William Fulton, *Young tableaux: With applications to representation theory and geometry*, Cambridge University Press, 1997.
- [7] Mark Haiman, *Hecke algebra characters and immanant conjectures*, J. Amer. Math. Soc. **6** (1993), no. 3, 569–595.

- [8] ———, *Hilbert schemes, polygraphs, and the Macdonald positivity conjecture*, J. Amer. Math. Soc. **14** (2001), 941–1006.
- [9] June Huh, Jacob Matherne, Karola Mészáros, and Avery St Dizier, *Logarithmic concavity of Schur and related polynomials*, Trans. Amer. Math. Soc. **375** (2022), no. 06, 4411–4427.
- [10] James E Humphreys, *Representations of semisimple lie algebras in the bgg category \mathcal{O}* , Vol. 94, American Mathematical Society, Providence, RI, 2008.
- [11] Thomas Lam, Alexander Postnikov, and Pavlo Pylyavskyy, *Schur positivity and Schur log-concavity*, Amer. J. Math **129** (2007), no. 6, 1611–1622.
- [12] Alain Lascoux, *Polynômes symétriques, foncteurs de schur et grassmanniennes*, Ph.D. Thesis, 1977.
- [13] Ian Grant Macdonald, *Symmetric functions and hall polynomials*, Oxford university press, 1998.
- [14] Bruce E Sagan, *The symmetric group: Representations, combinatorial algorithms, and symmetric functions*, Vol. 203, Springer Science & Business Media, 2001.
- [15] Richard P Stanley, *Graph colorings and related symmetric functions: ideas and applications a description of results, interesting applications, & notable open problems*, Discrete Mathematics **193** (1998), no. 1-3, 267–286.
- [16] Richard P. Stanley, *Enumerative combinatorics. Volume 2*, Camb. Stud. Adv. Math., vol. 62, Cambridge: Cambridge University Press, 1999 (English).
- [17] Andrei Vladlenovich Zelevinskii, *Resolvents, dual pairs, and character formulas*, Funct. Anal. Appl. **21** (1987), no. 2, 152–154.

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING, 100190, P.R. CHINA

Email address: `guitao18@mailsucas.ac.cn`

CENTER FOR COMBINATORICS, LPMC, NANKAI UNIVERSITY, TIANJIN 300071, P. R. CHINA

Email address: `yang@nankai.edu.cn`