

COMPACTNESS OF HAMILTONIAN STATIONARY LAGRANGIAN SUBMANIFOLDS IN SYMPLECTIC MANIFOLD

JINGYI CHEN, JOHN MAN SHUN MA

ABSTRACT. In this work, we prove a compactness theorem on the space of all Hamiltonian stationary Lagrangian submanifolds in a compact symplectic manifold with uniform bounds on area and total extrinsic curvature. This generalizes the compactness theorems in [8] and [6].

1. INTRODUCTION

Let (M, ω, h, J) be a $2n$ -dimensional symplectic manifold with a symplectic 2-form ω , an almost complex structure J and a compatible metric h . An immersed submanifold L in M is Lagrangian if $\dim L = n$ and $\omega|_L = 0$, and a Lagrangian immersion is Hamiltonian stationary (HSL) if it is a critical point of the volume functional among all Hamiltonian variations. The notion was first introduced in [15], [16]. A compact and graded Lagrangian in a Calabi-Yau manifold is Hamiltonian stationary if and only if it is special [19]. Thus HSL submanifolds are natural generalization of special Lagrangians submanifolds for general symplectic manifolds.

Examples of HSL submanifolds include the totally geodesic $\mathbb{R}\mathbb{P}^n$ in $\mathbb{C}\mathbb{P}^n$ and the flat tori $\mathbb{S}^1(a_1) \times \cdots \times \mathbb{S}^1(a_n)$ in \mathbb{C}^n [15], [16]. In two dimension, Schoen and Wolfson constructed in [17] HSL surfaces with conical singularity in any Kähler surfaces, which are also area minimizer in its Lagrangian homology class. On the other hand, using perturbation methods, Joyce, Lee and Schoen proved in [11] the existence of closed HSL submanifolds in every compact symplectic manifolds (see also the previous works [4, 13] and a family version in [14]). A large classes of examples of HSLs are constructed using techniques in integrable system. We refer the readers to the bibliography of [6] for a more comprehensive list of references.

Using the regularity theory developed in [7], Chen and Warren obtain the smoothness estimates and small Willmore energy regularity in [8] and prove a compactness theorem for HSL submanifolds in \mathbb{C}^n with uniformly bounded areas and total extrinsic curvatures in \mathbb{C}^n . The regularity and compactness results in [7, 8] rely on the assumption that the ambient space is \mathbb{C}^n since it is used, in an essential way, that the Lagrangian phase angle Θ can be written as $\arctan \lambda_1 + \cdots + \arctan \lambda_n$ for the graphic representation (x, Du) , where λ_i 's are the eigenvalues of D^2u .

Chen is partially supported by an NSERC Discovery Grant (22R80062). Ma is partially supported by DFF Sapere Aude 7027-00110B, by CPH-GEOTOP-DNRF151 and by CF21-0680 from respectively the Independent Research Fund Denmark, the Danish National Research Foundation and the Carlsberg Foundation.

In another direction, using the bubble tree convergence for conformal mappings with uniformly bounded area and Willmore energy, we proved in [6] a compactness theorem on the space of compact HSL surfaces with bounded area, genus and Willmore energy in a Kähler surfaces. A sequence of such HSL surfaces would converges smoothly away from finitely many points to a branched HSL immersion. Unlike the case for \mathbb{C}^n where one make use of the Lagrangian angle, in [6] we exploit the fact that the mean curvature one form of a HSL immersion in a Kähler manifold satisfies a first order elliptic system of "Hodge type"; this is also essential in the regularity results in [17] for a Lagrangian area minimizers in two dimensions.

Our main result is

Theorem 1.1. *Let (M, ω, h) be a compact $2n$ -dimensional symplectic manifold, where h is a Hermitian metric compatible with ω . Let C_V, C_A be positive constants and let $(L_k)_{k=1}^\infty$ be a sequence of connected compact Hamiltonian stationary Lagrangian immersion in M so that*

$$(1.1) \quad \text{Vol}(L_k) \leq C_V, \quad \|A_k\|_{L^\infty} \leq C_A, \quad \forall k \in \mathbb{N}.$$

Then either (L_k) converges to a point, or there is a finite set S so that a subsequence (L_{k_i}) of (L_k) converges smoothly locally graphically to a Hamiltonian stationary Lagrangian immersion L on $M \setminus S$ and

$$(1.2) \quad \text{Vol}(L) = \lim_{i \rightarrow \infty} \text{Vol}(L_{k_i}).$$

Also, the closure \bar{L} in M is connected and admits a structure of a Lagrangian varifold, and is Hamiltonian stationary in the sense that the generalized mean curvature vector \vec{H} of the varifold (\bar{L}, μ_L) satisfies

$$(1.3) \quad \int_M h(J\nabla^M f, \vec{H}) d\mu_L = 0, \quad \forall f \in C_c^\infty(M).$$

This generalizes the compactness theorems in [8], [6] to HSLs in any compact symplectic manifolds with metrics compacitble with the symplectic structures and the almost complex structures. The main step in proving compactness is to derive a local a-priori estimates for HSLs with small total curvature. In [8], [6], this is done by representating the Hamiltonian stationary conditions as a coupled elliptic systems of lower order. This simplification is not available for HSLs in a general symplectic manifold. Therefore, the methods in [8], [6] do not directly yield a proof of Theorem 1.1. In a Darboux coordinates, if one represents the HSL as a gradient graph $(x, Du(x))$ of a function u , then the Hamiltonian stationary condition is captured by a fourth order PDE on u . In a recent work [2], Bhattachaya, Chen and Warren have studied a fourth order elliptic PDE of double divergence form and derive a regularity results. Together with the constructions of Darboux coordinates with estimates in [11] (See also subsection 2.2), it is proved that a C^1 HSL in a general symplectic manifolds is smooth. In this work, we apply the results in [2] to derive a-priori estimates (Theorem 4.2) and a ϵ -regularity result (Theorem 4.7) for HSL submanifolds, which is essential to the proof of Theorem 1.1.

The paper is organized as follows. In section 2, we discuss some background in Lagrangian submanifolds, which includes the definition of HSL and the Darboux coordinates with estimates constructed in [11]. In section 3, we use the regularity results in [2] to derive local C^k estimates for functions which satisfy a fourth order elliptic equation in double divergence form. In section 4, we prove a ϵ -regularity theorem. Theorem 1.1 is proved in sections 5 and 6.

In the following, given any vector space W with a metric h , we denote $|w|_h = \sqrt{h(w, w)}$ for $w \in W$. When W is the Euclidean space, we write $|w| = |w|_{h_0}$, where h_0 is the Euclidean metric on W . Given any open subset Ω of \mathbb{R}^n and any function $u : \Omega \rightarrow \mathbb{R}$ on Ω , we use $\|u\|_{C^{k,\alpha}(\Omega)}$, $\|u\|_{L^p(\Omega)}$ and $\|u\|_{W^{k,p}(\Omega)}$ to denote respectively the $C^{k,\alpha}$, L^p and $W^{k,p}$ norms of u . For any $k \in \mathbb{N}$, $x \in \mathbb{R}^k$ and $r > 0$, the open ball with radius r and center x in \mathbb{R}^k is denoted $B_r^k(x)$. We also write B_r^k for $B_r^k(0)$.

2. BACKGROUND

2.1. Hamiltonian stationary Lagrangian submanifolds. Let (M, h) be a Riemannian manifold. Unless otherwise specified, throughout this paper a submanifold of M is an immersed submanifold, that is, a submanifold of M is a pair (L, ι) , where L is a subset of M and $\iota : N \rightarrow M$ is an immersion with $\iota(N) = L$. An immersed submanifold is compact (resp. connected), if N is compact (resp. connected), and is proper if ι is a proper map.

Assume that M is compact. By the Nash embedding theorem, there is an isometric embedding of (M, h) into \mathbb{R}^K for some positive integer K . Fixing such an embedding, the definition of varifolds on M is given in [1]. Let U be an open subset of M , and let L be a properly immersed k -dimensional submanifold of U given by the immersion $\iota : N \rightarrow U$. Let $g = \iota^*h$ be the induced metric on N and let dV_g be the volume form. The volume of a submanifold L is

$$(2.1) \quad \text{Vol}(L) = \int_N dV_g.$$

Let $\pi_k : G_k M \rightarrow M$ be the Grassmann bundle on M , where each fiber at $x \in M$ is the Grassmann manifold of k -dimensional subspaces of $T_x M$. Given any k -dimensional immersed submanifold L of M , let $G_\iota : N \rightarrow G_k M$ be the Gauss map given by $x \mapsto (\iota_*)_x T_x N$. Then (L, ι) is given a structure of a varifold μ_L by pushing forward: $|L| := (G_\iota)_\# dV_g$, or

$$(2.2) \quad |L|(f) := \int_N f \circ G_\iota dV_g, \quad \forall f \in C(G_k M).$$

For any integral varifold V , the weight measure μ_V on M is given by

$$(2.3) \quad \int_M \phi d\mu_V := \int_{G_k M} \phi(x) dV, \quad \forall \phi \in C(M).$$

Let $\mu_L := m\mu_{|L|}$. Clearly μ_L and $|L|$ depend not only on L but also on ι .

Let (M, ω, h, J) be a $2n$ -dimensional symplectic manifold, where ω is the symplectic form, J is an almost complex structure and h is a Riemannian metric. We assume that

ω, J, h satisfy the compatibility conditions

$$(2.4) \quad h(X, Y) = \omega(X, JY), \quad h(JX, JY) = h(X, Y)$$

for all tangent vectors X, Y at the same point.

An n -dimensional immersed submanifold L in M given by an immersion ι is called Lagrangian if $\iota^*\omega = 0$. Given an immersed Lagrangian submanifold L , (2.4) implies that J maps tangent vectors of L to normal vectors. Thus the second fundamental form of L can be written as

$$(2.5) \quad A : T_p L \times T_p L \times T_p L \rightarrow \mathbb{R}, \quad A(X, Y, Z) = h(\bar{\nabla}_X \tilde{Y}, JZ) = \omega(Z, \bar{\nabla}_X \tilde{Y}),$$

where \tilde{Y} is any local extension of Y . The *mean curvature one form* of L is a one form on N defined by

$$(2.6) \quad \alpha_x(X) = \omega_{\iota(x)}(\iota_* X, \vec{H}(x))$$

for any $x \in N$ and $X \in T_x N$. Here \vec{H} is the mean curvature vector of the immersion $\iota : N \rightarrow M$.

Next we give the definition of a Hamiltonian stationary Lagrangian immersion. As in [8], we first give the definition for general Lagrangian integral n -varifolds (An integral n -varifold V on (M, ω, h) is called *Lagrangian* if $\omega|_{T_x V} = 0$ for V -almost every x).

Definition 2.1. *A Lagrangian integral n -varifold V in U is called Hamiltonian stationary Lagrangian (HSL) in U if*

$$(2.7) \quad \int_U h(J\nabla^M f, \vec{H}) d\mu_V = 0, \quad \forall f \in C_c^\infty(U),$$

where \vec{H} is the generalized mean curvature vector of V .

Definition 2.2. *Let L be a Lagrangian immersion into an open subset U of M given by a proper immersion ι , and let $|L|$ be the integral n -varifold associated with L as in (2.2). L is called a Hamiltonian stationary Lagrangian (HSL) immersion in U if $|L|$ is HSL as in Definition 2.1.*

Let $\iota : N \rightarrow U$ be a proper Lagrangian immersion. For any $f \in C_c^\infty(U)$, let $\bar{f} = f \circ \iota$. By (2.3), (2.4) and (2.6), if the immersed submanifold (L, ι) is HSL in U , then

$$\int_N (d\bar{f}, \alpha)_g dV_g = 0,$$

where α is the mean curvature one form of L defined in (2.6). Let L be an embedded Lagrangian submanifold on M . Since every smooth function with compact support ψ on L can be extended to $f \in C_c^\infty(M)$, L is HSL if and only if

$$\int_N (d\psi, \alpha)_g dV_g = 0, \quad \forall \psi \in C_c^\infty(N).$$

This implies that α is co-closed. That is,

$$(2.8) \quad d^* \alpha = 0,$$

where $d^* = \pm * d *$ is the formal adjoint of the exterior derivative d with respect to the metric g .

Let L be an immersed submanifold in M given by an immersion $\iota : N \rightarrow M$ and let $U \subset M$ be an open set in M . Decompose $\iota^{-1}(U)$ as connected components

$$(2.9) \quad \iota^{-1}(U) = \bigsqcup_i V_i.$$

If $\iota|_{V_i}$ is an embedding for each i , we say that each $L_i = \iota(V_i)$ is an *embedded connected component* of $U \cap L$ and that ι splits into embedded connected components on U .

As in [8, Proposition 2.2], if L is a proper immersion, for each $p \in L$ there is an open neighborhood U of p so that $L \cap U$ splits into finitely many embedded connected components. We will have a more precise statement in Proposition 4.6.

Proposition 2.3. *Let L be a Lagrangian submanifold in an open subset U of M given by a proper immersion $\iota : N \rightarrow U$. Then L is HSL as in Definition 2.2 if and only if its mean curvature one form α is co-closed.*

Proof. The proof is similar to that of [8, Proposition 2.5 (1)] and we only sketch the proof for the direction (\Rightarrow). Let $p \in L$. By [8, Proposition 2.2], let W be an open neighborhood of p in U so that $\iota^{-1}(W)$ splits into finitely many connected components $\iota|_{E_i} : E_i \rightarrow \Sigma_i$, where $\Sigma_i = \iota(E_i)$. We recall that $q \in W \cap L$ is called an *embedded point* of L if there is an open neighborhood V_q of q in W so that $L \cap V_q$ is an embedded submanifold. Let $\mathcal{E} \subset L \cap W$ be the set of all embedded points of L . Using embeddedness, one can argue as in above that $d^*\alpha = 0$ on $\iota^{-1}(\mathcal{E})$ using Definition 2.2. For each i , one can prove that $\mathcal{E} \cap \Sigma_i$ is dense in Σ_i . Hence $d^*\alpha = 0$ on $\iota^{-1}(W)$. Since $p \in L$ is arbitrary, α is co-closed. \square

In particular, the Hamiltonian stationary condition holds on each embedded connected components.

Corollary 2.4. *Let L be a properly immersed HSL in an open subset U of M . Let W be an open subset of U such that $W \cap L$ splits into embedded connected components $\iota|_{E_i} : E_i \rightarrow \Sigma_i$. Then for each i ,*

$$(2.10) \quad \int_W h(J\nabla^M f, \vec{H}_i) d\mu_{\Sigma_i} = 0, \quad \forall f \in C_c^\infty(W).$$

Here \vec{H}_i is the mean curvature vector of Σ_i .

2.2. Darboux coordinates with estimates. In this subsection we recall the Darboux coordinates with estimates constructed in [11]. Let (M, ω) be a compact symplectic $2n$ -manifold, and let h be a Riemannian metric on M compatible with ω . Let \mathbb{U} be the $U(n)$ frame bundle of M : that is, for each $p \in M$, the fiber \mathbb{U}_p consists of unitary linear mappings $v : \mathbb{R}^{2n} \rightarrow T_p M$.

We recall Proposition 3.2 in [11]:

Proposition 2.5. *For small $\epsilon > 0$, we can choose a family of embeddings $\Upsilon_{p,v} : B_\epsilon^{2n} \rightarrow M$ depending smoothly on $(p, v) \in \mathbb{U}$ such that for all $(p, v) \in \mathbb{U}$, we have*

- (i) $\Upsilon_{p,v}(0) = p$ and $d\Upsilon_{p,v}|_0 = v : \mathbb{R}^{2n} \rightarrow T_p M$;
- (ii) $\Upsilon_{p,v \circ \gamma} \cong \Upsilon_{p,v} \circ \gamma$ for all $\gamma \in U(n)$;
- (iii) $\Upsilon_{p,v}^*(\omega) = \omega_0 = \frac{i}{2} \sum_{j=1}^n dz_j \wedge d\bar{z}_j$; and
- (iv) $\Upsilon_{p,v}^*(h) = h_0 + O(|z|) = \sum_{j=1}^n |dz_j|^2 + O(|z|)$.

From now on, by scaling ω, h if necessary, we assume $\epsilon = 2$ in Proposition 2.5.

Definition 2.6. Let $p \in M$ and $0 < r \leq 1$. The Darboux ball at p with radius r is

$$(2.11) \quad \mathcal{B}_r(p) := \Upsilon_{p,v}(B_r^{2n}).$$

By (ii) in Proposition 2.5, $\mathcal{B}_r(p)$ is well-defined, independent of $v \in \mathbb{U}_p$. For each $R \geq 1$ and $0 < t \leq R^{-1}$, we define $t : B_R^{2n} \rightarrow B_1^{2n}$ as $z \mapsto tz$. The following is proved in [11, Proposition 3.4].

Proposition 2.7. There exists K_0, K_1, K_2, \dots , depending only on (M, ω, h) such that for all $(p, v) \in U$ and $t \in (0, R^{-1}]$, the metric $h_{p,v}^t = t^{-2}(\Upsilon_{p,v} \circ t)^* h$ satisfies on B_R^{2n}

$$(2.12) \quad |h_{p,v}^t - h_0| \leq K_0 t \quad \text{and} \quad |\partial_j h_{p,v}^t| \leq K_j t^j \quad \text{for } j = 1, 2, \dots.$$

Again, by scaling ω, h if necessary, we assume that

$$(2.13) \quad \max\{K_0, K_1\} \leq \epsilon_1 < \max\{1/2, \epsilon_1(1/2, n)\},$$

where $\epsilon_1(1/2, n)$ is given in [2, Lemma 3.2]. By the first inequality in (2.12) and $K_0 < 1/2$,

$$(2.14) \quad B_{r/2}^M(p) \subset \mathcal{B}_r(p) \subset B_{2r}^M(p)$$

for all $p \in M$ and $r \in (0, 1)$, where $B_s^M(p)$ is the metric ball with radius s in (M, h) .

2.3. Convergence of immersed submanifolds. We recall the definition of smooth convergence as local graphs of a sequence of immersed submanifolds introduced in [8] (see also [9], [18] for similar definitions for sequences of embedded hypersurfaces).

Definition 2.8. Let $(\Sigma_k)_{k=1}^\infty$ be a sequence of properly immersed submanifolds in an open subset U of a smooth manifold M given by a proper immersion $\iota_k : N_k \rightarrow U$ for each k . Let S be an integral varifold in (M, h) and $m \in \mathbb{N} \cup \{\infty\}$. We say that $(\Sigma_k)_{k=1}^\infty$ converges graphically in U to S in C^m -topology, if there is $I \in \mathbb{N} \cup \{0\}$ such that

- Σ_k splits into I embedded connected components $\iota_k|_{E_k^i} : E_k^i \rightarrow \Sigma_k^i$, $i = 1, \dots, I$. Here E_k^i is open in N_k and $\iota_k|_{E_k^i}$ is an embedding onto Σ_k^i , and
- $S \cap U = \Sigma^1 \cup \dots \cup \Sigma^I$, where each Σ^i , $i = 1, \dots, I$, is an embedded submanifold in U .

Moreover, for each $i = 1, \dots, I$, there is $p_i \in \Sigma^i$ such that Σ_k^i , $k \in \mathbb{N}$, and Σ^i are graphs of a smooth mappings X_k^i and X^i respectively, where X_k^i and X^i are defined on some open subsets of $T_{p_i} \Sigma^i$, such that X_k^i converges in C^m norm to X^i as $k \rightarrow \infty$.

We remark when $I = 0$, it is understood that $\Sigma_k = S \cap U = \emptyset$.

Definition 2.9. Let $(\Sigma_k)_{k=1}^\infty$ be a sequence of properly immersed submanifolds in M , and let V be an open subset of M . We say that $(\Sigma_k)_{k=1}^\infty$ converges in V smoothly as local graphs to an integral varifold S , if for all $q \in V$, there is an open neighborhood U of q contained in V such that $(\Sigma_k \cap U)_{k=1}^\infty$ converges graphically to $S \cap U$ in C^∞ -topology.

3. ESTIMATES FOR DOUBLE DIVERGENCE EQUATION

In [2], the authors study the following fourth order equation in double divergence form

$$(3.1) \quad \partial_{x^i} \partial_{x^j} F^{jl}(x, Du, D^2u) = \partial_{x^k} a^k(x, Du, D^2u) - b(x, Du, D^2u)$$

where F^{ij} , a^k and b are smooth functions in x, Du, D^2u , defined in a convex neighborhood $U \subset B_r^n \times \mathbb{R}^n \times S^{n \times n}$, where $S^{n \times n}$ is the space of $n \times n$ symmetric matrices. A $W^{2,\infty}$ function u on B_r^n is called a weak solution to (3.1) if

$$(3.2) \quad \int_{B_r} [F^{ij} \eta_{ij} + a^k \eta_k + b\eta] dx = 0, \quad \text{for all } \eta \in C_c^\infty(B_r^n).$$

Definition 3.1. Let $\Lambda > 0$. We say that (3.1) is Λ -uniform on U if the standard Legendre ellipticity condition

$$(3.3) \quad \frac{\partial F^{jl}}{\partial u_{ik}}(\xi) \sigma_{ij} \sigma_{kl} \geq \Lambda |\sigma|^2, \quad \forall \sigma \in S^{n \times n}$$

is satisfied for any $\xi \in U$.

For each $p = 1, \dots, n$, let $h_p = h e_p$,

$$(3.4) \quad \eta^{-h_p}(x) := \frac{\eta(x - h_p) - \eta(x)}{h},$$

and $f := u^{h_p}$. Define

$$\begin{aligned} \xi_0 &= (x, Du(x), D^2u(x)), \\ \xi_h &= (x + h_p, Du(x + h_p), D^2u(x + h_p)), \\ \vec{V} &= \xi_h - \xi_0. \end{aligned}$$

One can check that (3.2) implies

$$(3.5) \quad \int_{B_r} \left(\beta^{ij,kl} f_{ik} \eta_{jl} + \gamma^{il} \eta_{jl} + \psi^k \eta_k^{-h_p} + \zeta \eta^{-h_p} \right) dx = 0 \quad \forall \eta \in C_c^\infty(B_{r-h}^n),$$

where

$$\begin{aligned} \beta^{ij,kl}(x) &= \int_0^1 \frac{\partial F^{jl}}{\partial u_{ik}}(\xi_0 + tV) dt, \\ \gamma^{jl}(x) &= \int_0^1 \left(\frac{\partial F^{jl}}{\partial u_k}(\xi_0 + tV) f_k + \frac{\partial F^{jl}}{\partial x_p}(\xi_0 + tV) \right) dt, \\ \psi^k(x) &= a^k(x, Du, D^2u), \\ \zeta(x) &= b(x, Du, D^2u). \end{aligned}$$

In the following, we assume that u is at least C^2 and satisfies (3.2). The following propositions are the key steps in proving the regularity results in [2].

Proposition 3.2. Assume that $u \in C^{2,\alpha}(B_r^n)$ satisfies (3.5), then for all $0 < r' < r$ and $0 < \alpha' < \alpha$ we have $u \in C^{3,\alpha'}(B_{r'}^n)$ and

$$(3.6) \quad \|D^3u\|_{C^{\alpha'}(B_{r'}^n)} \leq C(\Lambda, \alpha, \alpha', r, r', \|u\|_{W^{2,\infty}(B_r^n)}, \|\Psi\|_{C^\alpha(B_r^n)})$$

where for simplicity we write

$$\Psi = ((\beta_0^{ij,kl})_{i,j,k,l}, (\gamma_p^{jl})_{j,l,p}, (\psi^k)_k, \zeta).$$

and

$$(3.7) \quad \beta_0^{ij,kl}(x) = \frac{\partial F^{jl}}{\partial u_{ik}}(x, Du, D^2u),$$

$$(3.8) \quad \gamma_p^{jl}(x) = \frac{\partial F^{jl}}{\partial u_k}(x, Du, D^2u)u_{pk} + \frac{\partial F^{jl}}{\partial x^p}(x, Du, D^2u).$$

Proof. We first assume $r' = 1/5$ and $r = 1$. Without rewriting the whole proof again, we merely point out that in the proof of [2, Proposition 2.3], the constants used have the following dependence:

$$\begin{aligned} C_3 &= \|\beta^{ij,kl}\|_{C^\alpha(B_r^n)}, \\ C_4 &= \|\beta^{ij,kl}\|_{C^\alpha(B_r^n)}, \\ C_5 &= C(n)\|\gamma^{jl}\|_{C^\alpha(B_r^n)}, \\ C_6 &= C(n)\|\psi^k\|_{C^\alpha(B_r^n)}, \\ C_7 &= C(n)\|\zeta\|_{C^\alpha(B_r^n)}, \\ C_2 &= C(\Lambda) \\ C_1 &= C_1(n) \\ \tilde{\alpha} &= 1 - \delta, \\ \tilde{q} &= \frac{n}{2 - 2\tilde{\alpha}}, \\ \tilde{K} &= C(n)(\|\gamma^{jl}\|_{L^{2\tilde{q}}(B_1^n)}^2 + \|\psi^k\|_{L^{2\tilde{q}}(B_1^n)}^2 + \|\zeta\|_{L^{2\tilde{q}}(B_1^n)}), \\ r_0 &= r_0(C_1, \Lambda, n, \delta), \\ C_8 &= C_8(C_1, n, \delta), \\ C_9 &= C_9(C_1, C_8, r_0, \Lambda, \|D^2f\|_{L^2(B_{1/2}^n)}), \\ C_{10} &= C_{10}(C_4, C_5, C_7, \Lambda, \alpha, \delta, r_0, \|D^2f\|_{L^2(B_{1/2}^n)}, C_2) \end{aligned}$$

Thus by Proposition 2.1 and Proposition 2.3 in [2],

$$(3.9) \quad \|D^3u\|_{C^{\alpha-\delta/2}(B_{1/5})} \leq C(\Lambda, \alpha, \delta, \|u\|_{W^{2,\infty}(B_1^n)}, \|\Psi\|_{C^\alpha(B_1^n)}),$$

In general, given $0 < r' < r \leq 1$ and $x_0 \in B_{r'}^n$, we scale the ball $B_{r-r'}^n(x_0)$ to radius 1 and obtain, via (3.9),

$$\|D^3u\|_{C^{\alpha-\delta/2}(B_{(r-r')/5}^n(x_0))} \leq C(\Lambda, \alpha, \delta, r, r', \|u\|_{W^{2,\infty}(B_r^n)}, \|\Psi\|_{C^\alpha(B_r^n)}),$$

Since $x_0 \in B_{r'}^n$ is arbitrary, (3.6) is proved. \square

From the above proposition, we obtain

Proposition 3.3. *For each $0 < r' < r \leq 1$, $0 < \alpha' < \alpha < 1$ and $k \geq 3$, assume that $u \in C^{2,\alpha}(B_r^n)$ satisfies (3.2) and (F^{jl}) satisfies the Legendre ellipticity condition (3.3), then u is smooth in B_r and*

$$(3.10) \quad \|D^k u\|_{C^{\alpha'}(B_{r'}^n)} \leq C(k, \Lambda, \alpha, \alpha', r, r', \|u\|_{W^{2,\infty}(B_r^n)}, \|(\vec{F}, \vec{a}, b)\|_{C^{k-3}(B_r^n)}).$$

Proof. That u is smooth is proved in [2]. Taking the limit $h \rightarrow 0$ in (3.5), we obtain (writing $f = u_p$)

$$(3.11) \quad \int_{B_r^n} [(\beta_0^{ij,kl} f_{ij} + \gamma_p^{jl})\eta_{jl} + \partial_p \psi^k \eta_k + \partial_p \zeta \eta] dx = 0, \quad \forall \eta \in C_c^\infty(B_r^n).$$

This is an equation of the form (3.2) with the leading term

$$(3.12) \quad F_{(p)}^{ij}(x, Df, D^2 f) = \beta_0^{ij,kl} f_{ij} + \gamma_p^{jl}$$

and in particular

$$(3.13) \quad \frac{\partial F_{(p)}^{jl}}{\partial f_{ik}} = \beta_0^{ij,kl}.$$

Hence (3.11) also satisfies the Λ -uniform condition (3.3).

In general, one can inductively show that for any p_1, \dots, p_m , the function $f = u_{p_1 \dots p_m}$ satisfies

$$(3.14) \quad \int_{B_r^n} [(\beta_0^{ij,kl} f_{ik} + \gamma_{(p_1 \dots p_m)}^{jl})\eta_{jl} + \psi_{p_1 \dots p_m}^k \eta_k + \zeta_{p_1 \dots p_m} \eta] dx = 0, \quad \forall \eta \in C_c^\infty(B_r^n),$$

here $\gamma_{(p_1 \dots p_m)}^{jl}$ is defined inductively by

$$(3.15) \quad \gamma_{(p_1 \dots p_m)}^{jl} = (\gamma_{(p_1 \dots p_{m-1})}^{jl})_{x_{p_m}} + (\beta_0^{ij,kl})_{x_{p_m}} u_{ik p_1 \dots p_m}.$$

By Proposition 3.2, we obtain

$$(3.16) \quad \|D^3 f\|_{C^{\alpha'}(B_{r'}^n)} \leq C(\Lambda, \alpha, \alpha', r, r', \|f\|_{W^{2,\infty}(B_r^n)}, \|\Psi_{(p_1 \dots p_m)}\|_{C^\alpha(B_r^n)}),$$

where

$$(3.17) \quad \Psi_{(p_1 \dots p_m)} = ((\beta_0^{ij,kl})_{i,j,k,l}, (\gamma_{(p_1 \dots p_m)}^{jl})_{j,l,p_1, \dots, p_m}, (\partial_{p_1 \dots p_m} \psi^k)_k, \partial_{p_1 \dots p_m} \zeta).$$

Arguing inductively from (3.8), (3.15) that

$$\|\Psi_{p_1 \dots p_m}\|_{C^\alpha(B_r^n)} \leq C(m, \|\vec{a}\|_{C^m(B_r^n)}, \|F\|_{C^m(B_r^n)}, \|b\|_{C^m(B_r^n)}, \|u\|_{C^{m+2,\alpha}(B_r^n)})$$

Hence we have for any $0 < r' < r < 1$ and $0 < \alpha' < \alpha < 1$,

$$(3.18) \quad \|u\|_{C^{m+3,\alpha'}(B_{r'}^n)} \leq C(m, \Lambda, r, r', \alpha, \alpha', \|(\vec{F}, \vec{a}, b)\|_{C^m(B_r^n)}, \|u\|_{C^{m+2,\alpha}(B_r^n)})$$

The proposition is then proved by choosing

$$\begin{aligned} r' &= r_m < r_{m-1} < \dots < r_1 < r_0 = r, \\ \alpha' &= \alpha_m < \alpha_{m-1} < \dots < \alpha_1 < \alpha_0 = \alpha \end{aligned}$$

and applying (3.18) $m-1$ times (with r, r', α, α' replaced by $r_{j-1}, r_j, \alpha_{j-1}, \alpha_j$ respectively) and lastly Proposition 3.2. \square

4. LOCAL CALCULATIONS (SYMPLECTIC)

In this section, we represent locally a Lagrangian immersion as a gradient graph in a Darboux coordinates chart given in Proposition 2.5. We apply the estimates in the previous section. Under a smallness assumption on the L^n -norm of the second fundamental form, we derive a ϵ -regularity result (Theorem 4.7).

Let $h = h_{p,v}^t$ be a Riemannian metric on B_R^{2n} which satisfies (2.12) for some $t \in (0, R^{-1}]$ and K_0, K_1 satisfy (2.13).

Let (x^1, \dots, x^{2n}) be the standard coordinates on B_R^{2n} and e_1, \dots, e_{2n} be the standard basis of \mathbb{R}^{2n} . The Christoffel symbols for the Levi-Civita connection of h is given by

$$\nabla_{e_i} e_j = \Gamma_{ij}^k e_k + \Gamma_{ij}^{\bar{k}} e_{\bar{k}},$$

where $\bar{k} := n + k$ and the repeated indices are summed from 1 to n .

From (2.12) we have

$$(4.1) \quad |\Gamma_{ij}^k| + |\Gamma_{ij}^{\bar{k}}| \leq K_1 R^{-1}, \quad \forall i, j, k.$$

Let L be a Lagrangian submanifold in (B_R^{2n}, ω_0) given by a gradient graph

$$(4.2) \quad \Gamma(u) := \{\Gamma_u(x) : x \in U_R \subset B_R^n \subset \mathbb{R}^n\},$$

where

$$(4.3) \quad \Gamma_u(x) = (x, Du(x)).$$

for some smooth function $u : U_R \rightarrow \mathbb{R}$, where $U_R \subset B_R^n$. In this case, $L = \Gamma(u)$ satisfies (2.2) if it is a critical point of the area functional

$$(4.4) \quad \text{Vol}(u) = \int_{U_R} \sqrt{\det g_{ij}} dx,$$

where $g = \Gamma_u^* h$, among all Hamiltonian variations $t \mapsto u + t\eta$, where $\eta \in C_c^\infty(U_R)$. The Euler-Lagrange equation has been computed in [2, Lemma 3.1], which is

$$(4.5) \quad \int_{U_R} [F^{jl} \eta_{jl} + c^k \eta_k] dx = 0, \quad \forall \eta \in C_c^\infty(U_R)$$

where

$$(4.6) \quad F^{jl}(x, Du, D^2u) = \sqrt{g} g^{ij} ((\delta^{kl} + \mathcal{B}_{lk}) u_{ik} + \mathcal{C}_{li}),$$

$$(4.7) \quad c^k(x, Du, D^2u) = \frac{1}{2} \sqrt{g} g^{ij} (D_{y_k} \mathcal{A}_{ij} + 2u_{im} D_{y_k} \mathcal{C}_{mj} + u_{im} u_{jl} D_{y_k} \mathcal{B}_{kl}),$$

here \mathcal{A} , \mathcal{B} and \mathcal{C} are the components of $(h - I_{2n}) \circ \Gamma_u$ (see (3.2) in [2]). Note that \sqrt{g} , g^{ij} can also be represented by D^2u , \mathcal{A} , \mathcal{B} and \mathcal{C} . By (2.12) we obtain the following lemma.

Lemma 4.1. *For each $m \in \mathbb{N}$, there is a constant $C = C(r, m, K_0, \dots, K_m, \|D^2u\|_{L^\infty(B_r^n)})$ such that*

$$(4.8) \quad \|F^{jl}\|_{C^m(B_r^n)} \leq C,$$

$$(4.9) \quad \|c^k\|_{C^{m-1}(B_r^n)} \leq C,$$

where F^{jl} and c^k are defined in (4.6), (4.7).

By (2.13) we have $|Dh| < \epsilon_1$. By [2, Lemma 3.2], if

$$(4.10) \quad \|D^2u\|_{C^0(B_r^n)} \leq \epsilon_1,$$

then (4.5) is Λ -uniform with $\Lambda = 1/2$. Hence we can apply Proposition 3.3 and Lemma 4.1 to obtain

Theorem 4.2. *Let $u : B_r^n \rightarrow \mathbb{R}$ be a smooth function which satisfies $u(0) = Du(0) = D^2u(0) = 0$ and (4.10). If the graph $\Gamma(u)$ is Hamiltonian stationary Lagrangian in (B_r^{2n}, ω, h) , where h satisfies (2.12) and K_0, K_1 satisfy (2.13). Then for each $k \geq 3$ and $x \in B_{r/2}^n$,*

$$(4.11) \quad |D^k u(x)| \leq C(k, r, K_0, \dots, K_{k-3}).$$

4.1. Curvature estimates, ϵ -regularity. Next we use the result in the previous subsection to derive local curvature estimates for HSL immersions in (M, ω, J, h) with small L^n -norm of the second fundamental form.

Using the Darboux coordinates in subsection 2.2, one can think of $L \cap \mathcal{B}_r(p)$ as a Lagrangian immersion in (B_r^{2n}, ω_0) . We first derive an inequality comparing the second fundamental form of L computed using two different Riemannian metrics.

Proposition 4.3. *Let $k, K \in \mathbb{N}$ and $k < K$. Let L be a k -dimensional immersed submanifold in an open subset U of \mathbb{R}^K and let h_1 be a Riemannian metrics on U so that $\|h_1 - I\|_{C^0(U)} + \|Dh_1\|_{C^0(U)} < \epsilon$ for some positive number $\epsilon < 1/2$. Let A_1, A_0 be the second fundamental form of L calculated with respect to h_1 and the Euclidean metric $h_0 = I$ respectively. Then there is $C = C(K)$ so that*

$$|A_0(X, X)| \leq (1 + C\epsilon)|A_1(X, X)|_{h_1} + C\epsilon|X|_{h_1}^2$$

for all $x \in L$ and $X \in T_x L$.

Proof. Let ∇^1, ∇^0 be the Levi-Civita connections of h_1, h_0 respectively. Let \mathcal{C} be the tensor $\mathcal{C}(X, Y) = \nabla_X^1 Y - \nabla_X^0 Y$. Let $x \in L$ and let π_α^\perp be the orthogonal projection onto the orthogonal complement of $T_x L$ with respect to h_α for $\alpha = 1, 0$. Let $\{e_1, \dots, e_K\}$ be a basis of \mathbb{R}^K so that $\{e_1, \dots, e_k\}$ forms a basis for $T_x L$. Then

$$\pi_\alpha^\perp(Z) = Z - \sum_{a,b=1}^k H_\alpha^{ab} h_\alpha(Z, e_a) e_b, \quad \alpha = 1, 0.$$

Here $(H_\alpha)_{ab} := h_\alpha(e_a, e_b)$ and (H_α^{ab}) is the inverse matrix of $(H_\alpha)_{ab}$. Hence

$$\begin{aligned} (\pi_1^\perp - \pi_0^\perp)Z &= \sum_{a,b=1}^k (H_1^{ab} h_1(Z, e_a) - H_0^{ab} h_0(Z, e_a)) e_b \\ &= \sum_{a,b=1}^k ((H_1^{ab} - H_0^{ab}) h_1(Z, e_a) + H_0^{ab} (h_1(Z, e_a) - h_0(Z, e_a))) e_b. \end{aligned}$$

Write $e_a = (e_a^1, \dots, e_a^K)$ and $Z = (Z^1, \dots, Z^K)$, then

$$(H_\alpha)_{ab} = \sum_{i,j=1}^K e_a^i e_b^j (h_\alpha)_{ij}, \quad h_\alpha(Z, e_a) = \sum_{i,j=1}^K Z^i e_a^j (h_\alpha)_{ij}.$$

This implies

$$\begin{aligned} |H_1^{ab} - H_0^{ab}| &\leq C \|h_1 - h_0\|_{C^0(U)}, \\ |h_1(Z, e_a) - h_0(Z, e_a)| &\leq C \|h_1 - h_0\|_{C^0(U)} |Z| \end{aligned}$$

and

$$(4.12) \quad |(\pi_1^\perp - \pi_2^\perp)Z| \leq C \|h_1 - h_0\|_{C^0(U)} |Z|.$$

for some $C = C(K)$. Let X, Y be any tangent vector fields of L defined on some open neighborhood of x . Then

$$\begin{aligned} A_0(X, Y) &= \pi_0^\perp(\nabla_X^0 Y) \\ &= \pi_0^\perp(-\mathcal{C}(X, Y) + \nabla_X^1 Y) \\ &= -\pi_0^\perp \mathcal{C}(X, Y) + \pi_1^\perp \nabla_X^1 Y + (\pi_0^\perp - \pi_1^\perp)(\nabla_X^1 Y) \\ &= A_1(X, Y) - \pi_0^\perp \mathcal{C}(X, Y) + (\pi_0^\perp - \pi_1^\perp)(\nabla_X^1 Y). \end{aligned}$$

Now let X be chosen such that $\nabla_X^1 X = A_1(X, X)$ at x . Using (4.12), at x we have

$$\begin{aligned} |A_0(X, X)| &\leq |A_1(X, X)| + |\mathcal{C}(X, X)| + C \|h_1 - h_2\|_{C^0(U)} |A_1(X, X)| \\ &\leq (1 + C\epsilon) |A_1(X, X)|_{h_1} + |\mathcal{C}(X, X)|. \end{aligned}$$

For some $C = C(K)$. To estimate $\mathcal{C}(X, X)$, note that locally

$$\mathcal{C}_{ij}^k = (\Gamma_1)_{ij}^k - (\Gamma_0)_{ij}^k = (\Gamma_1)_{ij}^k,$$

where $(\Gamma_1)_{ij}^k$ are the Christoffel symbols of h_1 . Since $\|Dh_1\|_{C^0(U)} < \epsilon$ by assumption and $(\Gamma_1)_{ij}^k = h_1^{-1} * Dh_1$,

$$|\mathcal{C}(X, X)| \leq C\epsilon |X|_{h_1}^2$$

for some $C = C(K)$. This finishes the proof of Proposition 4.3. \square

Lemma 4.4. *Let U be an open subset in \mathbb{R}^{2n} and let L be an immersed Lagrangian submanifold in (U, ω_0, h, J) , so that h satisfies (2.12) and the C^0 -norm of the second fundamental form of L is bounded by C_A . Then there is C_1 depending on n, K_0, K_1 so that the following holds: for any $\epsilon \in (0, 1]$, write $r = \epsilon / (C_1(C_A + 1))$. For any $p \in L \cap U$ so that $B_r^{2n}(p) \subset U$, any embedded connected component L^i of $L \cap B_r^{2n}(p)$ containing p can be written as a gradient graph Γ_u , where u is a function defined on $U_i \subset T_p L^i$, $B_{r/2}^n \subset U_i$ and u satisfies*

$$(4.13) \quad u(0) = Du(0) = D^2u(0) = 0$$

and

$$(4.14) \quad \|D^2u\|_{C^0(U_i)} \leq \epsilon.$$

Proof. Apply Proposition 4.3 with $h_1 = h$, we conclude that $|A_0| \leq c_2(C_A + 1)$, where A_0 is the second fundamental form of L in (U, h_0) and c_2 depends only on n, K_0, K_1 . By [3, Theorem 2.6] (see also [12, Theorem 2.4]), there is a dimensional constant c so that if L^i is an embedded connected component of $L \cap B_{c\epsilon(C_A+1)^{-1}}^{2n}(p)$ containing p , then L^i is graphical: there is $X_i : U_i \rightarrow \mathbb{R}^n$, where $B_{c\epsilon(2c_2(C_A+1))^{-1}}^{2n}(p) \subset U_i$, so that

$$L^i = \{(x, X_i(x)) : x \in U_i\}$$

with $X_i(0) = DX_i(0) = 0$ and $|X_i(x)| \leq \epsilon$ for all $x \in U_i$. Since L^i is Lagrangian, $X_i = Du_i$ for some smooth function $u_i : U_i \rightarrow \mathbb{R}$ with $u_i(0) = 0$. This finishes the proof of the lemma, by choosing $C_1 = 2c_2/c$. \square

The following lemma can be proved by a direct computation, using the bounds on $D^2u, h - I$ and Dh .

Lemma 4.5. *With the same assumption as in Lemma 4.4, there is $C_2 > 0$ depending only on n, K_0, K_1 such that*

$$(4.15) \quad |D^3u(x)| \leq C_2(|A(x)| + 1), \quad |A(x)| \leq C_2(|D^3u(x)| + 1)$$

whenever $|x| \leq \epsilon_1(C_1(C_A + 1))^{-1}$.

From the above lemmas we obtain the following proposition.

Proposition 4.6. *Let L be a properly immersed Lagrangian submanifold in M . Let $p \in M$ and let δ, C_A be positive numbers. Assume that the C^0 -norm of the second fundamental form A of L is bounded above by C_A in the Darboux ball $\mathcal{B}_\delta(p)$. Then there is $C_2 > 0$ depending only on n, K_0, K_1 such that the following holds: for any $\epsilon \in (0, \epsilon_1]$, $L \cap \mathcal{B}_r(p)$ splits into embedded connected components, where $r = \min\{\delta/2, \epsilon(C_2(C_A + 1))^{-1}\}$. Moreover, for each $v \in \mathbb{U}_p$, each embedded connected component L_i of $L \cap \mathcal{B}_r(p)$, $z \in \tilde{L}^i := \Upsilon_{p,v}^{-1}(L_i)$, up to a unitary action, \tilde{L}^i is the gradient graph of a function u_i with $z = (0, 0)$,*

- (i) $u_i(0) = Du_i(0) = D^2u_i(0)$,
- (ii) $|D^2u_i(x)| \leq \epsilon$, and
- (iii) $|D^3u_i(x)| \leq C_1(C_A + 1)$.

Proof. For any $r < \delta$ and $v \in \mathbb{U}_p$, let $\tilde{L} := \Upsilon_{p,v}^{-1}(L \cap \mathcal{B}_p(r))$. Then \tilde{L} is a immersed Lagrangian submanifold in B_r^{2n} . For any $\epsilon \in (0, \epsilon_1]$, let $r_1(\epsilon) = \epsilon/(C_1(C_A + 1))$, where C_1 is defined in Lemma 4.4 and let $r_2 = r_1(\epsilon)/4$. Then for all $q \in B_{r_2}^{2n}$ we have $B_{r_2}^{2n} \subset B_{2r_1}^{2n}(q) \subset B_{r_1}^{2n}$. By Lemma 4.4, when $r = r_1(\epsilon)$, \tilde{L} splits into finitely many embedded connected components. Let $\bar{L}^1, \dots, \bar{L}^I$ be those components that intersects $B_{r_2}^{2n}$. For any $i = 1, \dots, I$ and $z \in \bar{L}^i \cap B_{r_2}^{2n}$, apply Lemma 4.4 to the ball $B_{2r_2}^{2n}(z)$. Thus up to a unitary action, $\bar{L}^i \cap B_{2r_2}^{2n}$ is the gradient graph of a function u_i . (i), (ii), (iii) follows from Lemma 4.4 and Lemma 4.5. Since $B_{r_2}^{2n} \subset B_{2r_2}^{2n}$, the Proposition is proved by choosing $C_2 = 4C_1$. \square

The following ϵ -regularity theorem is essential to the proof of Theorem 1.1.

Theorem 4.7. *There are positive numbers $\epsilon_0, C_{\epsilon_0}$ depending only on K_0, K_1, K_2 such that the following holds: if L is an n -dimensional properly immersed HSL submanifold in a symplectic manifold M , $p \in L$ and*

$$(4.16) \quad \int_{L \cap \mathcal{B}_{r_0}(p)} |A|^n d\mu_L \leq \epsilon_0,$$

where $r_0 \leq 1$. Then for all $0 < \sigma \leq r_0$ and $y \in \mathcal{B}_{r_0-\sigma}(p) \cap L$,

$$(4.17) \quad \sigma |A(y)| \leq C_{\epsilon_0}.$$

Proof. Let L be given by an immersion $\iota : N \rightarrow M$. Let $\Upsilon_{p,v} : B_{r_0}^{2n} \rightarrow M$ be given by Proposition 2.5, where $v \in \mathbb{U}_p$. Let z_0 be the maximum of the function

$$z \mapsto (r_0 - |z|)^2 \max_{s \in \iota^{-1}(\Upsilon_{p,v}(z))} |A(s)|^2$$

defined on $\Upsilon_{p,v}^{-1}(L \cap \mathcal{B}_{r_0}(p))$. Note that maximum exists since $\iota^{-1}(\Upsilon_{p,v}(z))$ is finite for each z . We assume this maximum is positive, or otherwise the result is trivial. So $|z_0| < r_0$. Let $s_0 \in \iota^{-1}(\Upsilon_{p,v}(z_0))$ such that

$$|A(s_0)|^2 = \max_{s \in \iota^{-1}(\Upsilon_{p,v}(z_0))} |A(s)|^2.$$

For all $z \in B_{\frac{r_0-|z_0|}{2}}^{2n}(z_0)$ and $s \in \iota^{-1}(\Upsilon_{p,v}(z))$,

$$\begin{aligned} |A(s)|^2 &\leq \frac{(r_0 - |z_0|)^2}{(r_0 - |z|)^2} |A(s_0)|^2 \\ &\leq \frac{(r_0 - |z_0|)^2}{\left(\frac{r_0 - |z_0|}{2}\right)^2} |A(s_0)|^2 \\ &\leq 4 |A(s_0)|^2. \end{aligned}$$

Let $p_0 = \Upsilon_{p,v}(z_0)$. By (2.14), $U := \mathcal{B}_{\frac{r_0-|z_0|}{8}}(p_0) \subset \Upsilon_{p,v}(B_{\frac{r_0-|z_0|}{2}}^{2n}(z_0))$. By Proposition 4.6, there is $C_1 > 0$ such that $L \cap U \cap \mathcal{B}_r(p_0)$ splits into embedded connected components, where $r = (C_1(2|A(s_0)| + 1))^{-1}\epsilon_1$ and each of the components is HSL by Corollary 2.4. Moreover, let L_i be a connected component so that $\iota^{-1}(L_i)$ contains s_0 . Choose $v_0 \in \mathbb{U}_{p_0}$ so that $v_0(\mathbb{R}^n \times \{0\}) = T_{p_0}L_i$. Then $\Upsilon_{p_0,v_0}^{-1}L_i$ is the gradient graph of a smooth function u with $u(0) = Du(0) = D^2(0) = 0$ and

$$|D^2u(x)| \leq \epsilon_1, \quad |D^3u(x)| \leq C_1(2|A(s_0)| + 1)$$

for all x in the domain of u .

Let

$$R = \frac{r_0 - |z_0|}{8} |A(s_0)|, \quad t = \frac{1}{|A(s_0)|}$$

and let

$$t : B_R^{2n} \rightarrow B_{\frac{r_0-|z_0|}{8}}^{2n}$$

be the scaling $z \mapsto tz$. Then $\tilde{L} = (\Upsilon_{p_0, v_0} \circ t)^{-1}(L_i)$ is an immersed HSL in $(B_R^{2n}, \omega_0, \tilde{h})$ given by

$$\tilde{\iota} = t^{-1} \circ \iota : V \rightarrow B_R^{2n}$$

with metric $\tilde{h} = t^{-2}(\Upsilon_{p_0, v_0} \circ t)^*h$. By the choice of R , the second fundamental form \tilde{A} of \tilde{L} satisfies $\|\tilde{A}\|_0 \leq 2$ and $|\tilde{A}(0)| = 1$. Also, \tilde{L} is a gradient graph of a smooth function \tilde{u} with

$$|D^2\tilde{u}(x)| \leq \epsilon_1, \quad |D^3\tilde{u}(x)| \leq \frac{C_1(2|A(s_0)| + 1)}{|A(s_0)|} \leq 3C_1$$

for all $x \in B_{(3C_1)^{-1}}^{2n} \cap B_R^{2n}$. Moreover,

$$(4.18) \quad \int_{\tilde{L}} |\tilde{A}|^n d\mu_{\tilde{L}} \leq \int_{B_{\frac{r_0 - |z_0|}{8}}} |A|^n d\mu < \epsilon_0$$

since the quantity is scale-invariant. Using Proposition 3.3 with $m = 4$ and (4.1)

$$|D^4\tilde{u}(x)| \leq C(K_0, K_1, K_2)$$

for all $x \in B_R^{2n}$ with $|x| \leq (4C_1)^{-1}$. This implies

$$|\tilde{\nabla}\tilde{A}| \leq C(K_0, K_1, K_2)$$

for all $x \in B_R^{2n}$ with $|x| \leq (4C_1)^{-1}$. Since $|\tilde{A}(0)| = 1$, one concludes

$$|\tilde{A}(x)| \geq \frac{1}{2}$$

for all $x \in \tilde{L}^1$ with $|x| \leq \tilde{r}$, where $\tilde{r} = \tilde{r}(K_0, K_1, K_2)$. Now choose $\epsilon_0 = \omega_n^{-1}(4\tilde{r})^n$. If $R > \tilde{r}$, then

$$\int_{B_R^{2n}} |\tilde{A}|^n d\mu_{\tilde{L}} \geq \int_{B_{\tilde{r}}^{2n}} |\tilde{A}|^n d\mu_{\tilde{L}} \geq \frac{1}{2^n} \mu_{\tilde{L}}(B_{\tilde{r}}^{2n}) \geq \frac{1}{\omega_n(4\tilde{r})^n} = \epsilon_0,$$

which is a contradiction. Thus

$$R = \frac{r_0 - |z_0|}{8} |A(s_0)| \leq \tilde{r},$$

which implies for all $z \in \mathcal{B}_{r_0}(p)$ and $s \in \iota^{-1}(\Upsilon_{p,v}(z))$,

$$|A(s)| \leq \frac{8\tilde{r}}{r_0 - |z|}.$$

□

5. PROOF OF THEOREM 1.1 FOR $n = 1$

In this section we prove Theorem 1.1 for $n = 1$. The general case is proved in the next section. We remark that it is necessary to split the proof into two cases: when $n \geq 2$ we use [8, Corollary 4.5], which says that if L is a Hamiltonian stationary Lagrangian in $M \setminus S$ and S is finite, then \bar{L} is also Hamiltonian stationary Lagrangian varifold in M . As pointed out in the introduction in [8], this does not hold when $n = 1$.

Proof of Theorem 1.1 for $n = 1$. In this case, (L_k) is a sequence of compact connected immersed curves in an oriented Riemannian surface (M, h, J) and the Lagrangian condition is automatically satisfied by any smooth curve. For each k , L_k is given by an immersion $\gamma_k : \mathbb{S}^1 \rightarrow M$. We assume each γ_k to be parameterized proportional to arc length.

Since J is compatible to h , one checks that an immersed curve $\gamma : \mathbb{S}^1 \rightarrow M$ in (M, h) is Hamiltonian stationary if and only if its curvature is constant. We remark that if h is generic then (M, h) admits infinitely many embedded circles with constant curvatures [20].

For each $k \in \mathbb{N}$, let κ_k be the (constant) curvature of γ_k . Let $\text{Length}(L)$ denotes the length of the immersed curves L defined in (2.1). The condition (1.1) implies

$$(5.1) \quad \text{Length}(\gamma_k) = \text{Length}(L_k) \leq C_V$$

and

$$(5.2) \quad |\kappa_k| \text{Length}(\gamma_k) = \int_{L_k} |\kappa_k| \leq C_A.$$

Taking a subsequence of (L_k) if necessary, we may assume that

$$(5.3) \quad \lim_{k \rightarrow \infty} \text{Length}(\gamma_k) = \ell_\infty \in [0, C_V]$$

and

$$(5.4) \quad \lim_{k \rightarrow \infty} \kappa_k = \kappa_\infty \in [-\infty, \infty].$$

By compactness of M , we may also assume that

$$(5.5) \quad \lim_{k \rightarrow \infty} \gamma_k(0) = p, \quad \lim_{k \rightarrow \infty} \gamma'_k(0) = v$$

for some $p \in M$ and $v \in T_p M$.

If $\ell_\infty = 0$, then (γ_k) converges to the point p . If $\ell_\infty > 0$, then $\kappa_\infty \neq \pm\infty$ by (5.2). Using (5.5) and the smooth dependence of ODE on initial data and parameters [10, Theorem 4.1], (γ_k) converges smoothly to an immersed curve γ_∞ with $\gamma_\infty(0) = p$, $\gamma'_\infty(0) = v$ and constant curvature κ_∞ . This finishes the proof of Theorem 1.1 when $n = 1$. □

6. PROOF OF THEOREM 1.1 FOR $n \geq 2$

First we prove a simple covering lemma by Darboux balls.

Proposition 6.1. *Let $\lambda \in (0, 1/4)$. There is b depending only on n and λ such that the following holds. For any $r < 1$, there is a finite open covering \mathcal{U} of M by Darboux balls $\mathcal{B}_r(a)$ so that (i) each member of \mathcal{U} intersect with at most $b - 1$ other members of \mathcal{U} , and (ii) the open cover $\{\mathcal{B}_{\lambda r}(p) : \mathcal{B}_r(p) \in \mathcal{U}\}$ still covers M .*

Proof. Let $\mathcal{V} = \{\mathcal{B}_{\lambda r/8}(p_1), \dots, \mathcal{B}_{\lambda r/8}(p_N)\}$ be a maximal collection of disjoint Darboux balls of radius $\lambda r/8$ in M . Then $\{\mathcal{B}_{\lambda r}(p_1), \dots, \mathcal{B}_{\lambda r}(p_N)\}$ covers M : to see this, assume

$p \in M$ and is not in one of $\mathcal{B}_{\lambda r/8}(p_i)$. Since \mathcal{V} is maximal, there is i so that $\mathcal{B}_{\lambda r/8}(p) \cap \mathcal{B}_{\lambda r/8}(p_i)$ contains an element y . By (2.14) and triangle inequality,

$$d(p, p_i) \leq d(p, y) + d(y, p_i) < \frac{\lambda r}{4} + \frac{\lambda r}{4} = \frac{\lambda r}{2}.$$

Hence $p \in B_{\lambda r/2}^M(p_i) \subset \mathcal{B}_{\lambda r}(p_i)$ by (2.14) again. Let \mathcal{U} be the open cover

$$\mathcal{U} = \{\mathcal{B}_r(p_1), \dots, \mathcal{B}_r(p_N)\}.$$

Note that \mathcal{U} satisfies (ii) by construction. To show (i), let p_i be fixed and assume that y_{i_j} lies in the intersection of $\mathcal{B}_r(p_i)$ and $\mathcal{B}_r(p_{i_j})$, for some $j = 1, \dots, k$. Then $d(p_i, p_{i_j}) \leq 4r$, or $p_{i_j} \in \mathcal{B}_{8r}(p_i)$ for all i_j . However, since \mathcal{V} is a collection of disjoint Darboux balls with radius $\lambda r/8$, one has $d(p_{i_j}, p_{i_l}) \geq \lambda r/8$. Let $v_i \in \mathbb{U}_{p_i}$ be fixed and let $z_j = \Upsilon_{p_i, v_i}^{-1}(p_{i_j})$. Then $\{z_1, \dots, z_k\}$ be a collection of points in B_{8r}^{2n} so that $|z_i - z_j| \geq \lambda r/16$ whenever $i \neq j$. This implies $k \leq b$, where b depends only on n, λ but not on r . This finishes the proof of (i). \square

Proof of Theorem 1.1 for $n \geq 2$. Choose $\lambda = \frac{\epsilon_1}{27C_2C_{\epsilon_0}}$, where ϵ_1, C_2 and C_{ϵ_0} are defined in (4.10), Proposition 4.6 and Theorem 4.7 respectively. By Proposition 6.1, there is a constant $b = b(n, \lambda)$ so that the following holds: for any $m \in \mathbb{N}$, there is a finite open cover

$$\mathcal{U}_m = \{\mathcal{B}_{2^{-m}}(\tilde{p}_1^m), \dots, \mathcal{B}_{2^{-m}}(\tilde{p}_{j(m)}^m)\}$$

of M by Darboux balls of radius 2^{-m} , so that

- (i) each elements in \mathcal{U} intersects at most $b - 1$ other members in \mathcal{U} , and
- (ii) the collection

$$\mathcal{U}_{m, \lambda} = \{\mathcal{B}_{\lambda 2^{-m}}(\tilde{p}_1^m), \dots, \mathcal{B}_{\lambda 2^{-m}}(\tilde{p}_{j(m)}^m)\}$$

still covers M .

The first condition implies that for each $k \in \mathbb{N}$,

$$(6.1) \quad \sum_{j=1}^{j(m)} \int_{\mathcal{B}_{2^{-m}}(\tilde{p}_j^m) \cap L_k} |A_k|^n d\mu_k < b \int_{L_k} |A_k|^n d\mu_k < bC_A,$$

where C_A is give in Theorem 1.1 and $\mu_k := \mu_{L_k}$. For any fixed k and m , let $J_{k,m} = \{p_{1,k}^m, \dots, p_{\ell,k}^m\}$ be the subset of $\{\tilde{p}_1^m, \dots, \tilde{p}_{j(m)}^m\}$ so that

$$\int_{\mathcal{B}_{2^{-m}}(p_{j,k}^m) \cap L_k} |A_k|^n d\mu_k \geq \epsilon_0.$$

Here ϵ_0 is given in Theorem 4.7 and $\ell = \ell(m, k)$ is less than bC_A/ϵ_0 by (6.1). Taking a diagonal subsequence of (L_k) if necessary, we may assume that $\ell(k, m) = \ell(m)$ and $J_{k,m} = J_m$ are both independent of k for all $m \in \mathbb{N}$. Write

$$J_m = \{p_1^m, \dots, p_{\ell(m)}^m\}.$$

Using $\ell(m) < bC_A/\epsilon_0$, there is a subsequence (m_i) such that $\ell(m_i) = \ell$ for all $i \in \mathbb{N}$. Using the compactness of M and taking a further subsequence of (m_i) if necessary, we may assume that

$$\lim_{i \rightarrow \infty} p_j^{m_i} = p_j$$

for all $j = 1, \dots, \ell$.

Let $S = \{p_1, \dots, p_\ell\}$ and $r_i = 2^{-m_i}$. Fix any $p \in \{\tilde{p}_1^{m_i}, \dots, \tilde{p}_{j(m)}^{m_i}\} \setminus J_{m_i}$. By definition of J_{m_i} this implies

$$\int_{\mathcal{B}_{r_i}(p) \cap L_k} |A_k|^n d\mu_k < \epsilon_0$$

for all $k \in \mathbb{N}$. By Theorem 4.7,

$$(6.2) \quad |A_k(q)| \leq 2C_{\epsilon_0}/r_i, \quad \forall k \in \mathbb{N} \text{ and } q \in \mathcal{B}_{r_i/2}(p).$$

Setting $C_A = 2C_{\epsilon_0}/r_i$ and note that

$$\frac{\epsilon_1}{C_2(C_A + 1)} = \frac{r_i \epsilon_1}{C_2(2C_{\epsilon_0} + r_i)} \geq \frac{\epsilon_1}{3C_2 C_{\epsilon_0}} r_i = 9\lambda r_i.$$

By Proposition 4.6, for each $k \in \mathbb{N}$, $\mathcal{B}_{8\lambda r_i}(p) \cap L_k$ splits into finitely many embedded connected components

$$\mathcal{B}_{8\lambda r_i}(p) \cap L_k = L_k^1 \cup \dots \cup L_k^{I_k}.$$

Let $v \in \mathbb{U}_p$ be fixed and for each $a = 1, \dots, I_k$, denote $\tilde{L}_k^a := \Upsilon_{p,v}^{-1}(L_k^a)$.

Lemma 6.2. *For each $i \in \mathbb{N}$, there is $M_i \in \mathbb{N}$ depending on i, λ, n and C_V in (1.1) only so that $\mathcal{B}_{8\lambda r_i}(p) \cap L_k$ has at most M_i embedded connected components which intersect $\mathcal{B}_{\lambda r_i}(p)$.*

Proof of Lemma 6.2. Let L_k^a be one of the components $\mathcal{B}_{8\lambda r_i}(p) \cap L_k$ which intersects $\mathcal{B}_{\lambda r_i}(p)$. Let $z_k^a \in B_{\lambda r_i}^{2n} \cap \tilde{L}_k^a$. Then there is a unitary matrix A_k^a and a smooth function u_k^a defined on U_k^a so that

$$(\Upsilon_k^a)^{-1}(\tilde{L}_k^a \cap B_{8\lambda r_i}^{2n}(z_k^a)) = \{(x, Du_k^a(x)) : x \in U_k^a\},$$

where $\Upsilon_k^a(z) = A_k^a z + z_k^a$. Using (2.12), we obtain $\mu(L_k^a) \geq C(n)(\lambda r_i)^n$ for some dimensional constant $C(n)$. Together with the assumption on the area bound in (1.1), the lemma is proved. \square

By Lemma 6.2 and taking a diagonal subsequence of (L_k) if necessary, we may assume that for any $i \in \mathbb{N}$ and $p \in \{\tilde{p}_1^{m_i}, \dots, \tilde{p}_{j(m)}^{m_i}\} \setminus J_{m_i}$, there is an integer $n_i(p)$ such that $\mathcal{B}_{8\lambda r_i}(p) \cap L_k$ has exactly $n_i(p)$ connected components which intersect $\mathcal{B}_{\lambda r_i}(p)$, where $n_i(p) \leq M_i$. Let $L_k^1, \dots, L_k^{n_i(p)}$ be such connected components.

Let $z_k^a, A_k^a, \Upsilon_k^a$ and $u_k^a : U_k^a \rightarrow \mathbb{R}$ be defined as in the proof of Lemma 6.2, where $u = u_k^a$ satisfies (4.13) and (4.14) with $\epsilon = \epsilon_1$. By Theorem 4.2, for any $m \geq 3$, there are \overline{C}_m depending on $r_i, \lambda, m, K_0, \dots, K_{m-3}$ so that

$$(6.3) \quad |D^m u_k^a(x)| \leq \overline{C}_m, \quad \text{for all } x \in B_{4\lambda r_i}^n.$$

Taking a diagonal subsequence of $(L^k)_{k=1}^\infty$ if necessary, we may assume that for each $i \in \mathbb{N}$ and p ,

- (1) the sequence of points $(z_k^a)_{k=1}^\infty$ converges to $z^a \in \overline{B_{\lambda r_i}^{2n}}$ as $k \rightarrow \infty$,
- (2) the sequence of unitary matrices $(A_k^a)_{k=1}^\infty$ converges to some $A^a \in U(n)$ as $k \rightarrow \infty$, and
- (3) the sequence of functions $(u_k^a)_{k=1}^\infty$ converges in $C^m(B_{3\lambda r_i}^n)$ for all $m \in \mathbb{N}$ to a smooth function $u^a : B_r^n \rightarrow \mathbb{R}$. This is possible by the higher order estimates (6.3).

Then the sequence of immersions $(\Upsilon_k^a \circ \Gamma_{u_k^a})_k$ converges to the immersion $\Upsilon^a \circ \Gamma_{u^a}$, where $\Upsilon^a(z) = A^a z + z^a$. In particular, there is \tilde{u}_k^a defined on $B_{2\lambda r_i}^n$ so that \tilde{L}_k^a is locally given by the immersion $\Upsilon^a \circ \Gamma_{\tilde{u}_k^a}$.

Next we define the Lagrangian integral n -varifold L on $M \setminus S$. Let

$$\mathcal{B} = \{\mathcal{B}_{\lambda r_i}(p) : i \in \mathbb{N}, p \in \{\tilde{p}_1^{m_i}, \dots, \tilde{p}_j^{m_i}\} \setminus J_{m_i}\}.$$

Clearly \mathcal{B} is an open cover of $M \setminus S$. For each $K \in \mathbb{N}$, let

$$\mathcal{B}_K = \{\mathcal{B}_{\lambda r_i}(p) \in \mathcal{B} : \mathcal{B}_{\lambda r_i}(p) \cap S_{1/K} = \emptyset\},$$

where for any $\rho > 0$ we set $S_\rho = \{q \in M : d^M(q, s) < \rho \text{ for some } s \in S\}$. Then $\tilde{\mathcal{B}} = \cup_K \mathcal{B}_K$ is an open cover of $M \setminus S$. For each $q \in M \setminus S$, then $q \in \mathcal{B}_{\lambda r_i}(p)$ for some $U = \mathcal{B}_{\lambda r_i}(p) \in \tilde{\mathcal{B}}$. Define $|L|$ on $\mathcal{B}_{\lambda r_i}(p)$ by

$$|L| = |L^1| + \dots + |L^{n_i(p)}|,$$

where for each $a = 1, \dots, n_i(p)$, L^a is the limits of $(L_k^a \cap \mathcal{B}_{\lambda r_i}(p))_{k=1}^\infty$ constructed in the previous paragraph. Since $z^a \in B_{\lambda r_i}^{2n}$ for each $a = 1, \dots, n_i(p)$, $B_{\lambda r_i}^{2n} \subset B_{2\lambda r_i}^{2n}(z^a)$. Hence $(L_k^a \cap \mathcal{B}_{\lambda r_i}(p))_{k=1}^\infty$ converges graphically in $\mathcal{B}_{\lambda r_i}(p)$ to $L \cap U$ in C^∞ -topology. Since $q \in M \setminus S$ is arbitrary, $(L_k)_{k=1}^\infty$ converges in $M \setminus S$ smoothly locally graphically to L . Since the convergence is locally smooth and each L_k is HSL, L is also HSL on $M \setminus S$. By [8, Corollary 4.4], \bar{L} admits a structure of an n -integral Lagrangian varifold so that (4.5) holds (although Theorem 4.3 and Corollary 4.4 in [8] are stated only for Lagrangian immersions in \mathbb{R}^{2n} , by isometrically embed (M, h) into some Euclidean space, the proofs work for immersions in (M, ω, J, h)). Connectedness of \bar{L} can be shown as in the proof of [8, Theorem 1.1].

Lastly, it remains to show (1.2). By the Nash embedding theorem, one can assume that (M, h) is an embedded submanifold of \mathbb{R}^{n+K} for some natural number K . Let L be an immersed submanifold M and let \tilde{A}, \tilde{H} be the second fundamental form and mean curvature vector of L respectively as a submanifold in \mathbb{R}^{n+K} . Then $\tilde{A} = A + A^M$, where A^M is the second fundamental form of M in \mathbb{R}^{n+K} . Together with the fact that $|A^M| \leq C(M)$ since M is compact, we obtain

$$\left(\int_L |\tilde{H}|^n d\mu_L \right)^{1/n} \leq \left(\int_L |H|^n d\mu_L \right)^{1/n} + C(M)\mu(L).$$

Thus we can apply [8, Proposition 4.1] to $L = L_k$ and conclude that

$$(6.4) \quad \text{Vol}(B_\rho(p) \cap L_k) \leq C(|\ln \rho| + 1)^n \rho^n$$

for all $p \in M$ and $\rho > 0$ small enough, here C depends only on M and C_V, C_A in Theorem 1.1. For any $\rho > 0$, by (6.4) we have

$$(6.5) \quad \text{Vol}(L_k) - C(|\ln \rho| + 1)^n \rho^n \leq \text{Vol}(L_k \setminus S_\rho) \leq \text{Vol}(L_k).$$

Since (L_k) converges smoothly graphically to L away from S_ρ , taking $k \rightarrow \infty$ in (6.5) gives

$$\lim_{k \rightarrow \infty} \text{Vol}(L_k) - C(|\ln \rho| + 1)^n \rho^n \leq \text{Vol}(L \setminus S_\rho) \leq \lim_{k \rightarrow \infty} \text{Vol}(L_k).$$

Thus (1.2) is shown by taking $\rho \rightarrow 0$ and this finishes the proof of Theorem 1.1. \square

REFERENCES

- [1] W.K. Allard: *On the first variation of a varifold*. Ann. Math. (2) **95**, 417-491 (1972).
- [2] A. Bhattachaya; J. Chen and M. Warren: *Regularity of Hamiltonian stationary Lagrangian equations in symplectic manifolds*, arXiv:2108.00325v1.
- [3] P. Breuning: *Immersion with Bounded Second Fundamental Form*, J. Geom. Anal. **25**, No. 2, 1344-1386 (2015).
- [4] A. Butscher and J. Corvino: *Hamiltonian stationary tori in Kähler manifolds*, Calc. Var. Partial Differ. Equ. **45**, No. 1-2 (2012), 63-100.
- [5] J. Chen and Y. Li: *Bubble tree of branched conformal immersions and applications to the Willmore functional*, Amer. J. Math. Vol. **136**, No.4 (2014).
- [6] J. Chen and J.M.S. Ma, *On the compactness of Hamiltonian stationary Lagrangian surfaces in Kahler surfaces*, Calc. Var. Partial Differ. Equ. **60**, No. 2, Paper No. 75, 23 p. (2021).
- [7] J. Chen and M. Warren: *On the regularity of Hamiltonian stationary Lagrangian submanifolds*. Adv. Math., (**343**), 316-352.
- [8] J. Chen and M. Warren: *Compactification of the space of Hamiltonian Stationary Lagrangian submanifolds with bounded total extrinsic curvature and volume*, arXiv:1901.03316v1 to appear in J. Diff. Geom.
- [9] H. Choi and R. Schoen: *The space of minimal embeddings of a surface into a three dimensional manifold of positive Ricci curvature*, Invent. Math., Vol. **81**, No. 3 , 387-394 (1985).
- [10] P. Hartman: *Ordinary Differential Equations*, second edition. Classics in Applied Mathematics, **CL38**, (2002).
- [11] D. Joyce; Y.-I., Lee and R. Schoen: *On the existence of Hamiltonian stationary Lagrangian submanifolds in symplectic manifolds*, Am. J. Math. **133**, No. 4, 1067-1092 (2011).
- [12] J. Langer: *A compactness theorem for surfaces with L_p -bounded second fundamental form*, Math. Ann. **270** (1985), no. 2, 223-234.
- [13] Y.-I. Lee: *The existence of Hamiltonian stationary Lagrangian tori in Kähler manifolds of any dimension*, Calc. Var. Partial Differ. Equ. **45**, No. 1-2 (2012), 231-251.
- [14] E. Legendre and Y. Rollin: *Hamiltonian stationary Lagrangian fibrations*, J. Symplectic Geom. **17**, No. 3, 753-791 (2019).
- [15] Y.-G. Oh: *Second variation and stabilities of minimal Lagrangian submanifolds in Kähler manifolds*, Invent. Math. **101**, No. 2 (1990), 501-519.
- [16] Y.-G. Oh: *Volume minimization of Lagrangian submanifolds under Hamiltonian deformations*, Math. Z. **212**, No. 2 (1993), 175-192.
- [17] R. Schoen and J. Wolfson: *Minimizing Area Among Lagrangian Surfaces: The Mapping Problem*, J. Differ. Geom. **58**, No. 1 (2001), 1-86.
- [18] B. Sharp: *Compactness of minimal hypersurfaces with bounded index*. J. Differ. Geom. **106** (2) 317-339, (2017).
- [19] R. Thomas and S.T. Yau: *Special Lagrangians, stable bundles and mean curvature flow*, Comm. Anal. Geom. **10**, No. 5 (2002), 1075-1113.

[20] R. Ye: *Foliation by constant mean curvature spheres*. Pac. J. Math. **147**, No. 2, 381-396 (1991).

JINGYI CHEN, DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF BRITISH COLUMBIA, 1984
MATHEMATICS RD, VANCOUVER, BC V6T 1Z2, CANADA

Email address: `jychen@math.ubc.edu.ca`

JOHN MAN SHUN MA, DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF COPENHAGEN,
UNIVERSITETSPARKEN 5, DK-2100 COPENHAGEN Ø, DENMARK

Email address: `jm@math.ku.dk`