

Riemann operators on higher K -groups

Nobushige Kurokawa* Hidekazu Tanaka†

September 19, 2022

Abstract

We introduce Riemann operators acting on Quillen’s higher K -groups $K_n(A)$ for the integer ring A of an algebraic number field K . Especially we prove that gamma factors of Dedekind zeta function of K are obtained as regularized determinants of Riemann operators.

Introduction

In this paper we introduce “Riemann operators” \mathcal{R} acting on Quillen’s higher K -groups $K_n(A)$ for the integer ring A of an algebraic number field K . We expect that such Riemann operators would supply determinant expressions for zeta allied functions. In fact, here we show that so called gamma factors of the Dedekind zeta function $\zeta_A(s)$ — the Hasse zeta function of $\text{Spec}(A)$ — are naturally obtained as the zeta regularized determinants of \mathcal{R} .

We remark that certain different “Riemann operators” so called “absolute Frobenius operators” were studied by Deninger [D1] [D2] around 30 years ago: see Manin [M] for an excellent survey lectured in 1992.

Now we define \mathcal{R} by

Definition 1. Let $\mathcal{R}|K_n(A)_{\mathbb{C}} = \frac{1-n}{2}$, where $K_n(A)_{\mathbb{C}} = K_n(A) \otimes_{\mathbb{Z}} \mathbb{C}$.

Then we prove

Theorem 1.

$$\det((sI - \mathcal{R})| \bigoplus_{n=0}^{\infty} K_n(A)_{\mathbb{C}}) = (s - \frac{1}{2})s^{-1} \left[\Gamma_{\mathbb{R}}(s)^{r_1} \Gamma_{\mathbb{C}}(s)^{r_2} (\sqrt{2\pi})^{[K:\mathbb{Q}]s} \right]^{-1} C(K),$$

*Department of Mathematics, Tokyo Institute of Technology

†6-15-11-202 Otsuka, Bunkyo-ku, Tokyo

where $C(K)$ is a constant given as

$$C(K) = (2\sqrt{\pi})^{r_1} (2\sqrt{2\pi})^{r_2}.$$

We notice that the determinant means the zeta regularized one:

$$\begin{aligned} \det((sI - \mathcal{R}) | \bigoplus_{n=0}^{\infty} K_n(A)_{\mathbb{C}}) &= \prod_{n=0}^{\infty} (s - \frac{1-n}{2})^{\dim K_n(A)_{\mathbb{C}}} \\ &= \exp\left(-\frac{\partial}{\partial w} \varphi_A(w, s) \Big|_{w=0}\right), \end{aligned}$$

where $\varphi_A(w, s) = \sum_{n=0}^{\infty} \dim K_n(A)_{\mathbb{C}} (s + \frac{n-1}{2})^{-w}$. Moreover, as usual, r_1 and r_2 are determined by the isomorphism of \mathbb{R} -algebras

$$K \otimes_{\mathbb{Q}} \mathbb{R} \cong \mathbb{R}^{r_1} \times \mathbb{C}^{r_2}.$$

We recall that the functional equation of the Dedekind zeta function $\zeta_A(s)$ is given as

$$\widehat{\zeta}_A(x) = \widehat{\zeta}_A(1-s),$$

where

$$\widehat{\zeta}_A(x) = \zeta_A(s) \Gamma_{\mathbb{R}}(s)^{r_1} \Gamma_{\mathbb{C}}(s)^{r_2} |D(K)|^{\frac{s}{2}}$$

with

$$\begin{aligned} \Gamma_{\mathbb{R}}(s) &= \Gamma\left(\frac{s}{2}\right) \pi^{-\frac{s}{2}}, \\ \Gamma_{\mathbb{C}}(s) &= \Gamma(s) 2(2\pi)^{-s} \end{aligned}$$

and $D(K)$ being the Discriminant of K over \mathbb{Q} . By letting $K = \mathbb{Q}$ in Theorem 1 we get the following

Corollary 1.

$$\det((sI - \mathcal{R}) | \bigoplus_{n=0}^{\infty} K_n(\mathbb{Z})_{\mathbb{C}}) = 2\sqrt{\pi} \left(s - \frac{1}{2}\right) s^{-1} \Gamma_{\mathbb{R}}(s)^{-1},$$

where $\Gamma_{\mathbb{R}}(s)$ is the gamma factor of the Riemann zeta function $\zeta_{\mathbb{Z}}(s)$.

1 Zeta regularized products

We recall zeta regularized products

$$\prod_{\lambda} (\lambda)^{a(\lambda)} = \exp(-\varphi'(0)),$$

where

$$\varphi(w) = \sum_{\lambda} a(\lambda) \lambda^{-w}.$$

We refer [D1] [D2] [M] [KT] for details.

Example 1.

$$\prod_{n=1}^{\infty} n = \sqrt{2\pi}.$$

Example 2 (Lerch 1894).

$$\prod_{n=0}^{\infty} (n+x) = \frac{\sqrt{2\pi}}{\Gamma(x)}.$$

2 Borel's result

Next, we recall Borel's result [B] concerning

$$\text{rank}K_n(A) = \dim K_n(A)_{\mathbb{C}}$$

for $n = 0, 1, 2, \dots$. It is given as:

$$\text{rank}K_n(A) = \begin{cases} 1 & \text{if } n = 0, \\ r_1 + r_2 - 1 & \text{if } n = 1, \\ r_1 + r_2 & \text{if } n > 1 \text{ and } n \equiv 1 \pmod{4}, \\ r_2 & \text{if } n \equiv 3 \pmod{4}, \\ 0 & \text{if otherwise.} \end{cases}$$

Thus we know that

$$\varphi_A(w, s) = (s - \frac{1}{2})^{-w} + (r_1 + r_2 - 1)s^{-w} + (r_1 + r_2)\varphi_1(w, s) + r_2\varphi_2(w, s),$$

where

$$\varphi_1(w, s) = \sum_{\substack{n>1 \\ n \equiv 1 \pmod{4}}} (\frac{n-1}{2} + s)^{-w}$$

and

$$\varphi_2(w, s) = \sum_{n \equiv 3 \pmod{4}} (\frac{n-1}{2} + s)^{-w}.$$

3 Lemmas

We prepare calculations of two regularized products.

Lemma 1.

$$\prod_{\substack{n>1 \\ n \equiv 1 \pmod{4}}} \left(\frac{n-1}{2} + s\right) = s^{-1} \Gamma_{\mathbb{R}}(s)^{-1} (\sqrt{2\pi})^{-s} 2\sqrt{\pi}.$$

Proof of Lemma 1. Since

$$\prod_{\substack{n>1 \\ n \equiv 1 \pmod{4}}} \left(\frac{n-1}{2} + s\right) = \exp\left(-\frac{\partial}{\partial w} \varphi_1(w, s) \Big|_{w=0}\right),$$

where

$$\begin{aligned} \varphi_1(w, s) &= \sum_{\substack{n>1 \\ n \equiv 1 \pmod{4}}} \left(\frac{n-1}{2} + s\right)^{-w} \\ &= \sum_{k=1}^{\infty} (2k + s)^{-w} \\ &= 2^{-w} \sum_{k=1}^{\infty} \left(k + \frac{s}{2}\right)^{-w} \\ &= 2^{-w} \zeta\left(w, 1 + \frac{s}{2}\right) \end{aligned}$$

with the standard notation of the Hurwitz zeta function

$$\zeta(w, x) = \sum_{k=0}^{\infty} (k + x)^{-w},$$

we get

$$\begin{aligned} -\frac{\partial}{\partial w} \varphi_1(w, s) \Big|_{w=0} &= (\log 2) \zeta\left(0, 1 + \frac{s}{2}\right) - \frac{\partial}{\partial w} \zeta\left(w, 1 + \frac{s}{2}\right) \Big|_{w=0} \\ &= (\log 2) \left(\frac{1}{2} - \left(1 + \frac{s}{2}\right)\right) + \log \frac{\sqrt{2\pi}}{\Gamma\left(1 + \frac{s}{2}\right)} \\ &= -\frac{s+1}{2} \log 2 + \log \frac{\sqrt{2\pi}}{\Gamma\left(1 + \frac{s}{2}\right)}. \end{aligned}$$

and

$$\begin{aligned}
\prod_{\substack{n>1 \\ n\equiv 1 \pmod{4}}} \left(\frac{n-1}{2} + s\right) &= 2^{-\frac{s+1}{2}} \frac{\sqrt{2\pi}}{\Gamma(1 + \frac{s}{2})} \\
&= 2^{-\frac{s}{2}} \frac{\sqrt{\pi}}{\frac{s}{2}\Gamma(\frac{s}{2})} \\
&= s^{-1} 2^{1-\frac{s}{2}} \frac{\sqrt{\pi}}{\Gamma_{\mathbb{R}}(s)\pi^{\frac{s}{2}}} \\
&= s^{-1} \Gamma_{\mathbb{R}}(s)^{-1} 2^{1-\frac{s}{2}} \pi^{\frac{1-s}{2}} \\
&= s^{-1} \Gamma_{\mathbb{R}}(s)^{-1} (\sqrt{2\pi})^{-s} 2\sqrt{\pi}.
\end{aligned}$$

□

Lemma 2.

$$\prod_{n\equiv 3 \pmod{4}} \left(\frac{n-1}{2} + s\right) = \Gamma_{\mathbb{R}}(s+1)^{-1} (\sqrt{2\pi})^{-s} \sqrt{2}.$$

Proof of Lemma 2. We calculate

$$\begin{aligned}
\varphi_2(w, s) &= \sum_{n\equiv 3 \pmod{4}} \left(\frac{n-1}{2} + s\right)^{-w} \\
&= \sum_{k=0}^{\infty} (2k+1+s)^{-w} \\
&= 2^{-w} \sum_{k=0}^{\infty} \left(k + \frac{s+1}{2}\right)^{-w} \\
&= 2^{-w} \zeta\left(w, \frac{s+1}{2}\right).
\end{aligned}$$

Hence, we get

$$\begin{aligned}
\prod_{n\equiv 3 \pmod{4}} \left(\frac{n-1}{2} + s\right) &= \exp\left(-\frac{\partial}{\partial w} \varphi_2(w, s) \Big|_{w=0}\right) \\
&= 2^{-\frac{s}{2}} \frac{\sqrt{2\pi}}{\Gamma(\frac{s+1}{2})} \\
&= 2^{-\frac{s}{2}} \frac{\sqrt{2\pi}}{\Gamma_{\mathbb{R}}(s+1)\pi^{\frac{s+1}{2}}} \\
&= \Gamma_{\mathbb{R}}(s+1)^{-1} (\sqrt{2\pi})^{-s} \sqrt{2}.
\end{aligned}$$

□

4 Proof of Theorem 1

Proof of Theorem 1. By using the previous calculations we obtain

$$\begin{aligned}
& \det((sI - \mathcal{R}) | \bigoplus_{n=0}^{\infty} K_n(A)_{\mathbb{C}}) \\
&= (s - \frac{1}{2}) s^{r_1+r_2-1} \left(\prod_{\substack{n>1 \\ n \equiv 1 \pmod{4}}} (\frac{n-1}{2} + s) \right)^{r_1+r_2} \left(\prod_{n \equiv 3 \pmod{4}} (\frac{n-1}{2} + s) \right)^{r_2} \\
&= (s - \frac{1}{2}) s^{r_1+r_2-1} \left(s^{-1} \Gamma_{\mathbb{R}}(s)^{-1} (\sqrt{2\pi})^{-s} 2\sqrt{\pi} \right)^{r_1+r_2} \left(\Gamma_{\mathbb{R}}(s+1)^{-1} (\sqrt{2\pi})^{-s} \sqrt{2} \right)^{r_2} \\
&= (s - \frac{1}{2}) s^{r_1+r_2-1} s^{-r_1-r_2} \Gamma_{\mathbb{R}}(s)^{-r_1-r_2} \Gamma_{\mathbb{R}}(s+1)^{-r_2} (\sqrt{2\pi})^{-(r_1+2r_2)s} (2\sqrt{\pi})^{r_1} (2\sqrt{2\pi})^{r_2} \\
&= (s - \frac{1}{2}) s^{-1} \Gamma_{\mathbb{R}}(s)^{-r_1} \left(\Gamma_{\mathbb{R}}(s) \Gamma_{\mathbb{R}}(s+1) \right)^{-r_2} (\sqrt{2\pi})^{-[K:\mathbb{Q}]s} C(K) \\
&= (s - \frac{1}{2}) s^{-1} \left(\Gamma_{\mathbb{R}}(s)^{r_1} \Gamma_{\mathbb{C}}(s)^{r_2} (\sqrt{2\pi})^{[K:\mathbb{Q}]s} \right)^{-1} C(K),
\end{aligned}$$

where we used that

$$\begin{aligned}
\Gamma_{\mathbb{C}}(s) &= \Gamma_{\mathbb{R}}(s) \Gamma_{\mathbb{R}}(s+1), \\
r_1 + 2r_2 &= [K : \mathbb{Q}]
\end{aligned}$$

and

$$C(K) = (2\sqrt{\pi})^{r_1} (2\sqrt{2\pi})^{r_2} \notin \overline{\mathbb{Q}}.$$

□

References

- [B] A. Borel “Stable real cohomology of arithmetic groups” *Ann. Sci. École Norm. Sup. (4)* **7** (1974) 235–272.
- [D1] C. Deninger “On the Γ -factors attached to motives” *Inventiones Math.* **104** (1991) 245–261.
- [D2] C. Deninger “Local L -factors of motives and regularized determinants” *Inventiones Math.* **107** (1992) 135–150.

- [KT] N. Kurokawa and H. Tanaka “Absolute zeta functions and the automorphy” *Kodai Math. J.*, **40** (2017), 584–614.
- [M] Y. Manin “Lectures on zeta functions and motives (according to Deninger and Kurokawa)” *Asterisque*, **228** (1995), 121–163.