

HIGH ORDER APPROXIMATIONS OF THE COX-INGERSOLL-ROSS PROCESS SEMIGROUP USING RANDOM GRIDS

AURÉLIEN ALFONSI AND EDOARDO LOMBARDO

ABSTRACT. We present new high order approximations schemes for the Cox-Ingersoll-Ross (CIR) process that are obtained by using a recent technique developed by Alfonsi and Bally (2021) for the approximation of semigroups. The idea consists in using a suitable combination of discretization schemes calculated on different random grids to increase the order of convergence. This technique coupled with the second order scheme proposed by Alfonsi (2010) for the CIR leads to weak approximations of order $2k$, for all $k \in \mathbb{N}^*$. Despite the singularity of the square-root volatility coefficient, we show rigorously this order of convergence under some restrictions on the volatility parameters. We illustrate numerically the convergence of these approximations for the CIR process and for the Heston stochastic volatility model and show the computational time gain they give.

1. INTRODUCTION

The present paper develops approximations, of any order, of the semigroup $P_t f(x) := \mathbb{E}[f(X_t^x)]$ associated to the following Stochastic Differential Equation (SDE) known as the Cox-Ingersoll-Ross (CIR) process

$$X_t^x = x + \int_0^t (a - kX_s^x) ds + \int_0^t \sigma \sqrt{X_s^x} dW_s, \quad t \geq 0, \quad (1.1)$$

where W is a Brownian motion, $x, a \geq 0$, $k \in \mathbb{R}$ and $\sigma > 0$. Let us recall that the process (1.1) is nonnegative and the semigroup $(P_t)_{t \geq 0}$ is well defined on the space of functions $f : \mathbb{R} \rightarrow \mathbb{R}$ with polynomial growth. The diffusion (1.1) is widely used in financial mathematics, in particular because of its simple parametrisation and the affine property that enables to use numerical methods based on Fourier techniques. We mention here the Cox-Ingersoll-Ross model [7] for the short interest rate and the Heston stochastic volatility model [8], that have been followed by many other ones. Developing efficient numerical methods for the process (1.1) is thus of practical importance.

To deal with the approximation of SDE's semigroups, a common approach is to consider stochastic approximations and the most standard one is the Euler-Maruyama scheme. The error between the approximated semigroup and the exact one is called the weak error, as opposed to the strong error that quantifies the error "omega by omega" on the probability space. The seminal work of Talay and Tubaro [14] shows, under regularity assumptions on the SDE coefficients, that the weak error given by the Euler-Maruyama scheme is of

Date: September 28, 2022.

2010 Mathematics Subject Classification. 60H35, 91G60, 65C30, G5C05.

Key words and phrases. Weak approximation schemes, random grids, Cox-Ingersoll-Ross model, Heston model.

This work benefited from the support of the "chaire Risques financiers", Fondation du Risque.

order one, i.e. is proportional to the time step. They also obtain an error expansion that enables to use Richardson-Romberg extrapolations as developed by Pagès [13]. Higher order schemes for SDEs and related extrapolations have been proposed by Kusuoka [9], Ninomiya and Victoir [11], Ninomiya and Ninomiya [10] and Oshima et al. [12] to mention a few. Recently, Alfonsi and Bally [4] have given a method to construct weak approximation of general semigroups of any order by using random time grids.

These general results on weak approximation of SDEs do not apply to the CIR (1.1) process. This is basically due to the diffusion coefficient, namely the singularity of the square-root at the origin. Besides this, classical schemes such as the plain Euler-Maruyama scheme are not well-defined for (1.1), and one has to work with dedicated schemes. Under some restrictions on the parameters, the weak convergence of order one for some discretization schemes of the CIR process has been obtained by Alfonsi [1], Bossy and Diop [5], and more recently by Briani et al. [6] who also study the weak convergence of a semigroup approximation for the Heston model. Adapting ideas from Ninomiya and Victoir [11] who developed a second order scheme for general SDEs, Alfonsi [2] has introduced second order and third order schemes for the CIR and proved their weak order of convergence, without any restriction on the parameters.

The goal of the present paper is to boost the second order scheme developed in [2] and get approximations of any order. To do so, we rely on the method developed recently by Alfonsi and Bally [4] to construct approximation of semigroups of any order. Roughly speaking, this method allows to get, from an elementary weak approximation scheme of order $\alpha > 0$, approximation schemes of any order by computing the elementary scheme on appropriate random grids. The method is illustrated in [4] on the case of the Euler-Maruyama scheme for SDEs, under regularity assumptions on the coefficients that do not hold for the CIR process (1.1). This method is presented briefly in Section 2. It relies on an appropriate choice of a function space endowed with a family of seminorms. Section 3 then presents the second order scheme that is used as an elementary scheme to get higher order approximation. It states in Theorem 3.2 the main result of this paper: we prove, when $\sigma^2 \leq 4a$, that we get weak approximations of any orders for smooth test functions f with derivatives having at most a polynomial growth. Section 4 illustrates the boosting method when considering the space of polynomials function with their usual norm. In this simple case, proofs are quite elementary so that the method can be followed easily. Section 5 is more involved: it first defines the appropriate family of seminorms on the space of smooth functions with derivative of polynomial growth and then proves Theorem 3.2. Last, we illustrate in Section 6 the convergence of the high order approximations for different parameter sets. It validates our theoretical results and shows important computational gains given by the new approximations. We also test the method on the Heston model and obtain similar convincing results.

2. HIGH ORDER SCHEMES WITH RANDOM GRIDS: THE METHOD IN A NUTSHELL

In this paragraph, we recall briefly the method developed by Alfonsi and Bally in [4] to construct approximations of any order from a family of approximation schemes. We consider F a vector space endowed with a family of seminorms $(\|\cdot\|_k)_{k \in \mathbb{N}}$ such that $\|f\|_k \leq \|f\|_{k+1}$. We consider a time horizon $T > 0$ and set, for $n \in \mathbb{N}^*$ and $l \in \mathbb{N}$,

$$h_l = \frac{T}{n^l}. \tag{2.1}$$

To achieve this goal, we consider a family of linear operators $(Q_l)_{l \in \mathbb{N}}$ and suppose that the two following conditions are satisfied. The first quantifies how Q_l approximates P_{h_l} :

$$\exists \alpha > 0, \forall l, k \in \mathbb{N}, \exists C > 0, \forall f \in F, \|(P_{h_l} - Q_l)f\|_k \leq C \|f\|_{\psi_Q(k)} h_l^{1+\alpha}, \quad (\overline{H_1})$$

where $\psi_Q : \mathbb{N} \rightarrow \mathbb{N}$ is a function¹. The second one is a uniform bound with respect to all the seminorms:

$$\forall l, k \in \mathbb{N}, \exists C > 0, \max_{0 \leq j \leq n^l} \|Q_l^{[j]} f\|_k + \sup_{t \leq T} \|P_t f\|_k \leq C \|f\|_k. \quad (\overline{H_2})$$

Then, for any $\nu \in \mathbb{N}^*$, Alfonsi and Bally [4] show how one can construct, by mixing the operators Q_l , a linear operator $\hat{P}_T^{\nu, n}$ such that

$$\exists C > 0, k \in \mathbb{N}^*, \forall f \in F, \|P_T f - \hat{P}_T^{\nu, n} f\|_0 \leq C \|f\|_k n^{-\nu\alpha}. \quad (2.2)$$

Let us explain how it works for $\nu = 1$ and $\nu = 2$. For $\nu = 1$, we mainly repeat the proof of Talay and Tubaro [14] for the weak error of the Euler scheme. For $l \in \mathbb{N}$, we note $Q_l^{[0]} = I$ the identity operator on F and, for $j \in \mathbb{N}^*$, $Q_l^{[j]} = Q_l^{[j-1]} Q_l$ the operator obtained by composition. From the semigroup property, we have

$$P_T f - Q_1^{[n]} f = P_{nh_1} f - Q_1^{[n]} f = \sum_{k=0}^{n-1} P_{(n-(k+1))h_1} [P_{h_1} - Q_1] Q_1^{[k]} f. \quad (2.3)$$

We get by using $(\overline{H_2})$, then $(\overline{H_1})$ and then again $(\overline{H_2})$

$$\begin{aligned} \|P_T f - Q_1^{[n]} f\|_0 &\leq \sum_{k=0}^{n-1} C \| [P_{h_1} - Q_1] Q_1^{[k]} f \|_0 \leq \sum_{k=0}^{n-1} C \| Q_1^{[k]} f \|_{\psi_Q(0)} h_1^{1+\alpha} \\ &\leq C \|f\|_{\psi_Q(0)} n (T/n)^{1+\alpha} = C \|f\|_{\psi_Q(0)} T^{1+\alpha} n^{-\alpha}. \end{aligned} \quad (2.4)$$

Here, and through the paper, C denotes a positive constant that may change from one line to another. So, $\hat{P}_T^{1, n} = Q_1^{[n]}$ satisfies (2.2) with $\nu = 1$, $k = \psi_Q(0)$. The approximation scheme simply consists in using n times the scheme Q_1 , which can be seen as a scheme on the regular time grid with time step h_1 .

We now present the approximation scheme (2.2) for $\nu = 2$. To do so, we use again (2.3) to get $P_{(n-(k+1))h_1} - Q_1^{[n-(k+1)]} = \sum_{k'=0}^{n-(k+2)} P_{(n-(k+k'+2))h_1} [P_{h_1} - Q_1] Q_1^{[k']}$ and then expand further (2.3):

$$\begin{aligned} P_T f - Q_1^{[n]} f &= \sum_{k=0}^{n-1} Q_1^{[n-(k+1)]} [P_{h_1} - Q_1] Q_1^{[k]} f + R_2^{h_1}(n) f, \\ \text{with } R_2^{h_1}(n) &= \sum_{k=0}^{n-1} \sum_{k'=0}^{n-(k+2)} P_{(n-(k+k'+2))h_1} [P_{h_1} - Q_1] Q_1^{[k']} [P_{h_1} - Q_1] Q_1^{[k]} \end{aligned} \quad (2.5)$$

Using $(\overline{H_1})$ three times and $(\overline{H_2})$ twice, we obtain

$$\|R_2^{h_1}(n) f\|_0 \leq C \|f\|_{\psi_Q(\psi_Q(0))} \frac{n(n-1)}{2} h_1^{2(1+\alpha)} \leq C \|f\|_{\psi_Q(\psi_Q(0))} \frac{T^{2(1+\alpha)}}{2} n^{-2\alpha}.$$

¹Note that in [4], it is taken $\psi_Q(k) = k + \beta$ for some $\beta \in \mathbb{N}$, but it can be easily generalized to any function ψ_Q . In this paper, we will work with a doubly indexed norm and take $\psi_Q(m, L) = (2(m+3), L-1)$.

Thus, $Q_1^{[n]} + \sum_{k=0}^{n-1} Q_1^{[n-(k+1)]}[P_{h_1} - Q_1]Q_1^{[k]}f$ is an approximation of order 2α , but it still involves the semigroup through P_{h_1} . To get an approximation that is obtained only with the operators Q_l , we use again (2.3) with time step h_2 and final time $h_1 = nh_2$:

$$P_{h_1}f - Q_2^{[n]}f = \sum_{k=0}^{n-1} P_{(n-(k+1))h_2}[P_{h_2} - Q_2]Q_2^{[k]}f.$$

We have $\|P_{h_1}f - Q_2^{[n]}f\|_0 \leq C\|f\|_{\psi_Q(0)}nh_2^{1+\alpha}$ by using again (\overline{H}_1) and (\overline{H}_2) . We get from (2.5)

$$P_Tf - Q_1^{[n]}f = \sum_{k=0}^{n-1} Q_1^{[n-(k+1)]}[Q_2^{[n]} - Q_1]Q_1^{[k]}f + \sum_{k=0}^{n-1} Q_1^{[n-(k+1)]}[P_{h_1} - Q_2^{[n]}]Q_1^{[k]}f + R_2^{h_1}(n)f, \quad (2.6)$$

with $\|\sum_{k=0}^{n-1} Q_1^{[n-(k+1)]}[P_{h_1} - Q_2^{[n]}]Q_1^{[k]}f\|_0 \leq C\|f\|_{\psi_Q(0)}n^2h_2^{1+\alpha} = C\|f\|_{\psi_Q(0)}T^{1+\alpha}n^{-2\alpha}$. Therefore, the approximation

$$\hat{P}^{2,n}f := Q_1^{[n]}f + \sum_{k=0}^{n-1} Q_1^{[n-(k+1)]}[Q_2^{[n]} - Q_1]Q_1^{[k]}f \quad (2.7)$$

satisfies (2.2) with $\nu = 2$ and is obtained only with the approximating operators Q_l . The first term $Q_1^{[n]}$ corresponds to apply the scheme Q_1 on the regular time grid with time step h_1 , while each term $Q_1^{[n-(k+1)]}[Q_2^{[n]} - Q_1]Q_1^{[k]}$ is the difference between this scheme and the one where $Q_2^{[n]}$ is used instead of Q_1 for the $(k+1)$ -th time step. This amounts to refine this time step and split it into n time steps of size h_2 , and to use the scheme Q_2 on this time grid.

In practice, it is inefficient to calculate one by one the terms in $\hat{P}^{2,n}f$. In fact, each term requires a number of calculations that is proportional to n , and the overall computation cost would be of the same order as n^2 . Since the convergence is in $O(n^{-2\alpha})$ it would not be better asymptotically than using $\hat{P}^{1,n^2}f$. To avoid this, we use randomization. We sample a uniform random variable κ on $\{0, \dots, n-1\}$ and calculate $n\mathbb{E}[Q_1^{[n-(\kappa+1)]}[Q_2^{[n]} - Q_1]Q_1^{[\kappa]}f] = \sum_{k=0}^{n-1} Q_1^{[n-(k+1)]}[Q_2^{[n]} - Q_1]Q_1^{[k]}f$. This amounts to consider the regular time grid with time step h_1 , to select randomly one time step and to refine it, and then to compute the difference between the schemes simulated on the refined time-grid and on the regular time-grid. This explains why we use random time grids.

We have presented here how to construct $\hat{P}^{\nu,n}$ for $\nu = 1$ and $\nu = 2$, and it is possible by repeating the same arguments to construct by induction approximations of any order. Unfortunately, the induction is quite involved. It is fully described in [4, Theorem 3.10]. We do not reproduce it in this paper because it would require much more notation, and we will mainly use the scheme (2.7). Here, we give in addition the explicit form of $\hat{P}^{3,n}$, $n \geq 2$:

$$\begin{aligned} \hat{P}^{3,n}f := & \hat{P}^{2,n} + \sum_{0 \leq k_1 < k_2 < n}^{n-1} Q_1^{[n-(k_2+1)]}[Q_2^{[n]} - Q_1]Q_1^{[k_2-k_1-1]}[Q_2^{[n]} - Q_1]Q_1^{[k_1]}f \\ & + \sum_{k=0}^{n-1} Q_1^{[n-(k+1)]} \left[\sum_{k'=0}^{n-1} Q_2^{[n-(k'+1)]}[Q_3^{[n]} - Q_2]Q_2^{[k']} \right] Q_1^{[k]}f. \end{aligned} \quad (2.8)$$

By similar arguments, it satisfies (2.2) with $\nu = 3$.

3. SECOND ORDER SCHEMES FOR THE CIR PROCESS AND MAIN RESULT

In this section, we focus on the approximation of the semigroup of the CIR process $P_t f(x) = \mathbb{E}[f(X_t^x)]$, where

$$X_t^x = x + \int_0^t (a - kX_s^x) ds + \sigma \int_0^t \sqrt{X_s^x} dW_s, \quad t \geq 0.$$

Equation (2.4) shows that, necessarily, approximating operators Q_l that satisfy both (\overline{H}_1) and (\overline{H}_2) lead to a weak error of order α . Therefore, we are naturally interested in approximation schemes of the CIR for which we know the rate of convergence α for the weak error. [1, Proposition 4.2] gives a rate $\alpha = 1$ for a family of approximation schemes that are basically obtained as a correction of the Euler scheme. Ninomiya and Victoir [11] have developed a generic method to construct second order schemes ($\alpha = 2$) for Stochastic Differential Equations with smooth coefficients. Applied to the Cox-Ingersoll-Ross process, their method leads to the following approximation scheme

$$\hat{X}_t^x = \varphi(x, t, \sqrt{t}N), \quad (3.1)$$

where $N \sim \mathcal{N}(0, 1)$ and $\varphi : \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}_+$ is defined by

$$\begin{aligned} \varphi(x, t, w) &= e^{-kt/2} \left(\sqrt{(a - \sigma^2/4)\psi_k(t/2) + e^{-kt/2}x + \sigma w/2} \right)^2 + (a - \sigma^2/4)\psi_k(t/2) \quad (3.2) \\ &= X_0(t/2, X_1(w, X_0(t/2, x))), \quad \text{with} \end{aligned}$$

$$X_0(t, x) = e^{-kt}x + \psi_k(t)(a - \sigma^2/4), \quad \psi_k(t) = \frac{1 - e^{-kt}}{k}, \quad (3.3)$$

$$X_1(t, x) = (\sqrt{x} + t\sigma/2)^2, \quad (3.4)$$

with the convention that $\psi_0(t) = t$. This scheme corresponds to approximate $P_t f(x)$ by $\hat{P}_t f(x) = \mathbb{E}[f(\hat{X}_t^x)]$ for $x, t \geq 0$, and then to set $Q_l = \hat{P}_{hl}$. Its construction comes from the splitting of the infinitesimal generator of the CIR process

$$\mathcal{L}f(x) = (a - kx)f'(x) + \frac{1}{2}\sigma^2 x f''(x), \quad f \in \mathcal{C}^2, x \geq 0, \quad (3.5)$$

as $\mathcal{L} = V_0 + \frac{1}{2}V_1^2$ with

$$V_0 f(x) = \left(a - \frac{\sigma^2}{4} - kx \right) f'(x) \quad \text{and} \quad V_1 f(x) = \sigma \sqrt{x} f'(x). \quad (3.6)$$

The function $t \mapsto X_0(t, x)$ is the solution of the ODE $X_0'(t, x) = a - \frac{\sigma^2}{4} - kX_0(t, x)$ such that $X_0(0, x) = x$, while $X_1(W_t, x)$ solves the SDE associated to the infinitesimal generator $V_1^2/2$.

The scheme (3.1) is well defined for $\sigma^2 \leq 4a$. Instead, for $\sigma^2 > 4a$, it is not well defined for any $x \geq 0$ since the argument in the square-root is negative when x is close to zero. To correct this, Alfonsi [2] has proposed the following scheme

$$\hat{X}_t^x = (\mathbb{1}_{x \geq K_2^Y(t)} \varphi(x, t, \sqrt{t}Y) + \mathbb{1}_{x < K_2^Y(t)} \hat{X}_t^{x,d}), \quad (3.7)$$

where Y is a random variable with compact support on $[-A_Y, A_Y]$ $A_Y > 0$ such that $\mathbb{E}[Y^k] = \mathbb{E}[N^k]$ for $k \leq 5$, and $\hat{X}_t^{x,d}$ is a nonnegative random variable such that $\mathbb{E}[(\hat{X}_t^{x,d})^i] = \mathbb{E}[(X_t^x)^i]$

for $i \in \{1, 2\}$ and $K_2^Y(t)$ is a nonnegative threshold defined by

$$K_2^Y(t) = \mathbf{1}_{\sigma^2 > 4a} \left[e^{\frac{kt}{2}} \left((\sigma^2/4 - a)\psi_k(t/2) + \left(\sqrt{e^{\frac{kt}{2}}(\sigma^2/4 - a)\psi_k(t/2) + \frac{\sigma}{2}A_Y\sqrt{t}} \right)^2 \right) \right]. \quad (3.8)$$

Note that when $\sigma^2 \leq 4a$, we have $K_2^Y(t) = 0$ and thus $\hat{X}_t^x = \varphi(x, t, \sqrt{t}Y)$.

We now restate [2, Theorem 2.8] that analyses the weak error.

Theorem 3.1. *Let \hat{X}_t^x be the scheme defined by (3.1) for $\sigma^2 \leq 4a$ or by (3.7) for any $\sigma > 0$. Let $f : \mathbb{R}_+ \rightarrow \mathbb{R}$ be a C^∞ function such that:*

$$\forall k \in \mathbb{N}, \exists C_k, E_k \in \mathbb{R}_+, \forall x \geq 0, |f^{(k)}(x)| \leq C_k(1 + |x|^{E_k}).$$

Then, $Q_1^{[n]}f(x) - P_T f(x) = O(1/n^2)$ where $Q_1 f(x) = \mathbb{E}[f(\hat{X}_{h_1}^x)]$.

The goal of this paper is to extend this result and prove the estimates (\overline{H}_1) and (\overline{H}_2) for a suitable space of functions and a suitable family of seminorms. We are able to prove such results only in the case $\sigma^2 \leq 4a$: the indicator function in (3.7) creates a singularity that is difficult to handle in the analysis. In Section 4, we first prove (\overline{H}_1) and (\overline{H}_2) for polynomial test functions. Then, we deal in Section 5 with the much technical case of smooth test functions with derivatives of polynomial growth. We state here our main result, the proof of which is given in Section 5.

Theorem 3.2. *Let \hat{X}_t^x be the scheme defined by (3.1) for $\sigma^2 \leq 4a$ and $Q_l f(x) = \mathbb{E}[f(\hat{X}_{h_l}^x)]$, for $l \geq 1$. Let $f : \mathbb{R}_+ \rightarrow \mathbb{R}$ be a C^{18} function such that:*

$$\forall k \leq 18, \exists C_k, E_k \in \mathbb{R}_+, \forall x \geq 0, |f^{(k)}(x)| \leq C_k(1 + |x|^{E_k}).$$

Then, $\hat{P}^{2,n}f(x) - P_T f(x) = O(1/n^4)$ as $n \rightarrow \infty$.

If $f : \mathbb{R}_+ \rightarrow \mathbb{R}$ is a C^∞ function such that:

$$\forall k \in \mathbb{N}, \exists C_k, E_k \in \mathbb{R}_+, \forall x \geq 0, |f^{(k)}(x)| \leq C_k(1 + |x|^{E_k}).$$

Then, we have $\hat{P}^{\nu,n}f(x) - P_T f(x) = O(1/n^{2\nu})$.

4. THE CASE OF POLYNOMIAL TEST FUNCTIONS

In this section, we want to illustrate the method and consider test functions that are polynomial test functions. We define for $L \in \mathbb{N}$

$$\mathfrak{P}_L(\mathbb{R}_+) = \{f : \mathbb{R}_+ \rightarrow \mathbb{R}, \exists a_0, \dots, a_L \in \mathbb{R}, f(x) = \sum_{j=0}^L a_j x^j\},$$

the vector space of polynomial functions over \mathbb{R}_+ with degree less or equal to L . We also define $\mathfrak{P}(\mathbb{R}_+) = \cup_{L \in \mathbb{N}} \mathfrak{P}_L(\mathbb{R}_+)$ the space of polynomial functions. We endow $\mathfrak{P}(\mathbb{R}_+)$ with the following norm:

$$\|f\| = \sum_{j=0}^L |a_j|, \text{ for } f(x) = \sum_{j=0}^L a_j x^j. \quad (4.1)$$

We consider the case $\sigma^2 \leq 4a$ and consider the scheme (3.7) for the CIR process with a time step $t > 0$, $\hat{X}_t^x = \varphi(x, t, \sqrt{t}Y)$. The approximation scheme Q_l is then defined by

$Q_l f = \mathbb{E}[f(\hat{X}_{h_l}^x)]$. The goal of this section is to prove $(\overline{H_1})$ and $(\overline{H_2})$ for the norm (4.1). We make the following assumption on Y .

Assumption (\mathcal{H}_Y) : $Y : \Omega \rightarrow \mathbb{R}$ is a symmetric random variable such that $\mathbb{E}[|Y|^k] < \infty$ for all $k \in \mathbb{N}$, and $\mathbb{E}[Y^k] = \mathbb{E}[N^k]$ for $k \in \{2, 4\}$ with $N \sim \mathcal{N}(0, 1)$.

We now state two lemmas that will enable us to prove that $(\overline{H_2})$ is satisfied by the scheme (3.7). Lemma 4.1 shows that polynomial functions are preserved by the approximation scheme, and gives short time estimate for the polynomial norm. Lemma 4.2 gives similar results for the CIR diffusion. The proofs of these lemmas are quite elementary and are postponed to Appendix A.

Lemma 4.1. *Let $T \geq 0$, $t \in [0, T]$, $f \in \mathfrak{P}_L(\mathbb{R}_+)$ and assume (\mathcal{H}_Y) and $\sigma^2 \leq 4a$. Then, we have $f(X_0(t, \cdot)), \mathbb{E}[f(X_1(\sqrt{t}Y, \cdot))] \in \mathfrak{P}_L(\mathbb{R}_+)$ where X_0 and X_1 are defined by (3.3) and (3.4), and*

$$\begin{aligned} (1) \quad & \|f(X_0(t, \cdot))\| \leq (1 \vee e^{-kLt})(1 + C_{X_0}^L t) \|f\|, \\ (2) \quad & \|\mathbb{E}[f(X_1(\sqrt{t}Y, \cdot))]\| \leq (1 + \mathbb{E}[Y^{2L}]C_{X_1}^L t) \|f\|, \end{aligned}$$

for some constants C_{X_0}, C_{X_1} depending only on (a, σ, T) .

Lemma 4.2. *Let $(X_t^x, t \geq 0)$ be the CIR process starting from $x \in \mathbb{R}_+$. For $m \in \mathbb{N}$, we define $\tilde{u}_m(t, x) := \mathbb{E}[(X_t^x)^m]$. There exists C^∞ functions $\tilde{u}_{j,m} : \mathbb{R}_+ \rightarrow \mathbb{R}$ that depend on (k, a, σ) such that:*

$$\tilde{u}_m(t, x) = \sum_{j=0}^m \tilde{u}_{j,m}(t) x^j. \quad (4.2)$$

If $f \in \mathfrak{P}_L(\mathbb{R}_+)$, then we have $\mathbb{E}[f(X_t)] \in \mathfrak{P}_L(\mathbb{R}_+)$ and for $t \in [0, T]$,

$$\|\mathbb{E}[f(X_t)]\| \leq C_{cir}(L, T) \|f\|, \quad (4.3)$$

with $C_{cir}(L, T) = \max_{t \in [0, T], m \in \{0, \dots, L\}} \sum_{j=0}^m |\tilde{u}_{j,m}(t)|$.

We are now in position to prove the main result of this section, which is a weaker (but easier to prove) version of our main Theorem 3.2, since it only applies to polynomial test functions.

Proposition 4.3. *Let $\sigma^2 \leq 4a$ and assume that Y satisfies (\mathcal{H}_Y) . For any $L \in \mathbb{N}$, the properties $(\overline{H_1})$ and $(\overline{H_2})$ are satisfied by the scheme (3.7) $\hat{X}_t^x = \varphi(x, t, \sqrt{t}Y)$ for $F = \mathfrak{P}_L(\mathbb{R}_+)$ and the norm (4.1). Then, we have for any $f \in \mathfrak{P}_L(\mathbb{R}_+)$,*

$$\|\mathbb{E}[f(X_T^x) - \hat{P}^{\nu, n} f]\| \leq C_L \|f\| n^{-2\nu},$$

for some constant C_L .

Proof. We first prove $(\overline{H_2})$. The property $\sup_{t \in [0, T]} \|P_t f\|$ is given by Lemma 4.2. Since $\hat{X}_t^x = X_0(t/2, X_1(\sqrt{t}Y, X_0(t/2, x)))$, we get by Lemma 4.1

$$\|\mathbb{E}[f(\hat{X}_t^x)]\| \leq [(1 \vee e^{-kLt/2})(1 + C_{X_0}^L t/2)]^2 (1 + \mathbb{E}[Y^{2L}]C_{X_1}^L t) \|f\|.$$

We now use that $1 + x \leq e^x$ to get

$$\|\mathbb{E}[f(\hat{X}_t^x)]\| \leq e^{((-k)^+ L + C_{X_0}^L + \mathbb{E}[Y^{2L}]C_{X_1}^L)t} \|f\|. \quad (4.4)$$

Since $Q_l f(x) = \mathbb{E}[f(\hat{X}_{T/n^l}^x)]$, this yields to $\max_{0 \leq j \leq n^l} \|Q_l^{[j]} f\| \leq e^{((-k)^+ L + C_{X_0}^L + \mathbb{E}[Y^{2L}]C_{X_1}^L)T} \|f\|$.

We now prove (\overline{H}_1) . Let $m \in \{0, \dots, L\}$ and $0 < x_0 < \dots < x_L$ be fixed real numbers (one may take for example $x_\ell = \ell + 1$). Lemmas 4.1 and 4.2 give that $v_m(t, x) = \mathbb{E}[(\hat{X}_t^x)^m] - \mathbb{E}[(X_t^x)^m] = \sum_{j=0}^m v_{j,m}(t)x^j$. By [2, Proposition 2.4], we know that there exists C'_m, E'_m such that $\forall t \in (0, 1), |v_m(t, x)| \leq C'_m t^3(1 + |x|^{E'_m})$. Therefore, $\exists \tilde{C}_m \in \mathbb{R}_+, \forall 0 \leq \ell \leq L, |v_m(t, x_\ell)| \leq \tilde{C}_m t^3$. By using the invertibility of the Vandermonde matrix, we get that

$$\exists C_m \in \mathbb{R}_+, \forall j \in \{0, \dots, m\}, |v_{j,m}(t)| \leq C_m t^3.$$

Therefore, we get for $f \in \mathfrak{P}_L(\mathbb{R}_+)$

$$\|\mathbb{E}[f(\hat{X}_t)] - \mathbb{E}[f(X_t)]\| \leq \sum_{m=0}^L |a_m| \sum_{j=0}^m C_m t^3 \leq L \max_{m \in \{0, \dots, L\}} C_m \|f\| t^3,$$

that gives (\overline{H}_1) . We conclude by applying [4, Theorem 3.10]. \square

5. PROOF OF THEOREM 3.2

In Section 4, we have obtained the convergence for test functions that are polynomial functions. For these test functions, the choice of the norm is straightforward and the proofs are not very technical and quite easy. However, one would like to obtain the convergence result for a much larger class of test functions. This is the goal of this section.

We consider test functions that are smooth with polynomial growth, whose derivatives have a polynomial growth. Namely, we introduce for $m, L \in \mathbb{N}$,

$$\mathcal{C}_{\text{pol}}^{m,L}(\mathbb{R}_+) = \left\{ f : \mathbb{R}_+ \rightarrow \mathbb{R} \text{ of class } \mathcal{C}^m : \forall j \in \{0, \dots, m\}, \sup_{x \geq 0} \frac{|f^{(j)}(x)|}{1 + x^L} < \infty \right\}, \quad (5.1)$$

which we endow with the norm

$$\|f\|_{m,L} = \max_{j \in \{0, \dots, m\}} \sup_{x \geq 0} \frac{|f^{(j)}(x)|}{1 + x^L}. \quad (5.2)$$

To prove Theorem 3.2, we need to prove the estimates (\overline{H}_1) and (\overline{H}_2) for this family of norms. This is the goal of the two next subsections. Before, we summarize in the next lemma some properties of the norms defined in Equation (5.2) that we will use later on.

Lemma 5.1. *Let $m, L \in \mathbb{N}$. We have the following basic properties:*

- (1) $\|f\|_{m',L} = \sup_{j \in \{0, \dots, m'\}} \|f^{(j)}\|_{0,L}$ for $f \in \mathcal{C}_{\text{pol}}^{m,L}(\mathbb{R}_+)$ and $m' \in \{0, \dots, m\}$.
- (2) $\mathcal{C}_{\text{pol}}^{m+1,L}(\mathbb{R}_+) \subset \mathcal{C}_{\text{pol}}^{m,L}(\mathbb{R}_+)$ and $\|f\|_{m,L} \leq \|f\|_{m+1,L}$ for $f \in \mathcal{C}_{\text{pol}}^{m+1,L}(\mathbb{R}_+)$.
- (3) $\|f^{(i)}\|_{m,L} \leq \|f\|_{m+i,L}$ for $i \in \mathbb{N}$ and $f \in \mathcal{C}_{\text{pol}}^{m+i,L}(\mathbb{R}_+)$.
- (4) $\mathcal{C}_{\text{pol}}^{m,L}(\mathbb{R}_+) \subset \mathcal{C}_{\text{pol}}^{m,L+1}(\mathbb{R}_+)$ and $\|f\|_{m,L+1} \leq 2\|f\|_{m,L}$ for $f \in \mathcal{C}_{\text{pol}}^{m,L}(\mathbb{R}_+)$.
- (5) Let \mathcal{M}_1 be the operator defined by $f \mapsto \mathcal{M}_1 f$, $\mathcal{M}_1 f(x) = x f(x)$. Then, $\mathcal{M}_1 f \in \mathcal{C}_{\text{pol}}^{m,L+1}(\mathbb{R}_+)$ for $f \in \mathcal{C}_{\text{pol}}^{m,L}(\mathbb{R}_+)$ and $\|\mathcal{M}_1 f\|_{m,L+1} \leq (2m + 3)\|f\|_{m,L}$.
- (6) Let $\mathcal{L}f(x) = (a - kx)f'(x) + \frac{1}{2}\sigma^2 x f''(x)$ be the infinitesimal generator of the CIR process. Then, we have for $f \in \mathcal{C}_{\text{pol}}^{m+2,L}(\mathbb{R}_+)$,

$$\|\mathcal{L}f\|_{m,L+1} \leq (2a + (2m + 3)(|k| + \sigma^2/2)) \|f\|_{m+2,L}.$$

We also have $\|(V_1^2/2)f\|_{m,L+1} \leq \sigma^2(m+2)\|f\|_{m+2,L}$ and $\|V_0f\|_{m,L+1} \leq [2|a - \sigma^2/4| + (2m+3)|k|]\|f\|_{m+1,L}$, where V_0 and V_1 are defined by (3.6).

Proof. Properties (1)–(3) are straightforward, and we prove only (4)–(6).

(4) We use the fact that $\forall x \geq 0$ $1 + x^L \leq 2(1 + x^{L+1})$, hence

$$\sup_{j \in \{0, \dots, m\}} \sup_{x \geq 0} \frac{|f^{(j)}(x)|}{1 + x^{L+1}} \leq 2 \sup_{j \in \{0, \dots, m\}} \sup_{x \geq 0} \frac{|f^{(j)}(x)|}{1 + x^L}.$$

(5) Let $f \in \mathcal{C}_{\text{pol}}^{m,L}(\mathbb{R}_+)$. We will use the fact that for all $x \geq 0$, $(1+x)(1+x^L) \leq 3(1+x^{L+1})$ so

$$\sup_{x \geq 0} \frac{x|f^{(j)}(x)|}{1 + x^{L+1}} \leq 3 \sup_{x \geq 0} \frac{x}{1+x} \sup_{x \geq 0} \frac{|f^{(j)}(x)|}{1 + x^L} = 3 \sup_{x \geq 0} \frac{|f^{(j)}(x)|}{1 + x^L}.$$

Now, we use the Leibniz rule on \mathcal{M}_1f and get $(xf(x))^{(j)} = jf^{(j-1)}(x) + xf^{(j)}(x)$, so

$$\sup_{x \geq 0} \frac{|(xf(x))^{(j)}|}{1 + x^{L+1}} \leq j \sup_{x \geq 0} \frac{|f^{(j-1)}(x)|}{1 + x^{L+1}} + \sup_{x \geq 0} \frac{x|f^{(j)}(x)|}{1 + x^{L+1}}.$$

Maximizing both sides on $j \in \{0, \dots, m\}$ and using the previous inequality gives $\|\mathcal{M}_1f\|_{m,L+1} \leq m\|f\|_{m-1,L+1} + 3\|f\|_{m,L}$. We get the bound by using properties (2) and (4).

(6) We have $\|\mathcal{L}f\|_{m,L+1} \leq a\|f'\|_{m,L+1} + (2m+3)[|k|\|f'\|_{m,L} + \frac{\sigma^2}{2}\|f''\|_{m,L}]$ by using the property (5). We get the estimate by using (3), (4) and (2). The other estimate for $V_1^2/2$ is obtained by taking $a = \sigma^2/4$ and $k = 0$, while the one for V_0 follows by using the same arguments. \square

We also state the following elementary lemma that will be useful to prove both (\overline{H}_1) and (\overline{H}_2) .

Lemma 5.2. *Let $T > 0$, $\sigma^2 \leq 4a$ and X_0 be defined by (3.3). Then, there exists a constant $K \geq 0$ such that for any function $f \in \mathcal{C}_{\text{pol}}^{m,L}(\mathbb{R}_+)$, we have*

$$\forall t \in [0, T], \|f(X_0(t, \cdot))\|_{m,L} \leq e^{Kt}\|f\|_{m,L}.$$

Proof. We first prove the following inequality

$$1 + X_0(t, x)^L \leq (1 \vee e^{-Lkt})(1 + \tilde{C}_{X_0}t)(1 + x^L),$$

for some constant \tilde{C}_{X_0} . To do so, we develop the term $X_0(t, x)^L$ and get

$$\begin{aligned} 1 + X_0(t, x)^L &= 1 + \sum_{j=0}^L \binom{L}{j} e^{-(L-j)kt} x^{(L-j)} (\psi_k(t)(a - \sigma^2/4))^j \\ &= 1 + e^{-Lkt} x^L + \psi_k(t) \sum_{j=1}^L \binom{L}{j} e^{-(L-j)kt} x^{(L-j)} \psi_k(t)^{j-1} (a - \sigma^2/4)^j. \end{aligned}$$

We remark that for $k \geq 0$, $0 \leq \psi_k(t) \leq t \leq 1 \vee T$ for all $t \in [0, T]$. For $k < 0$, we have $\psi_k(t) = e^{-kt}\psi_{-k}(t)$ and thus $\psi_k(t) \leq e^{(-k)^+t}t$ for all $t \in [0, T]$ and $k \in \mathbb{R}$. Using $x^j \leq 1 + x^L$ for all $j \in \{1, \dots, L\}$, we can rewrite the previous identity as

$$1 + X_0(t, x)^L \leq (1 \vee e^{-Lkt})(1 + x^L)$$

$$\begin{aligned}
& + te^{(-k)^+t}(1 \vee e^{-Lkt})(1+x^L) \sum_{j=0}^L \binom{L}{j} (e^{(-k)^+T}(1 \vee T)(a - \sigma^2/4))^j \\
& \leq (1 \vee e^{-Lkt})(1 + \tilde{C}_{X_0}t)(1+x^L),
\end{aligned}$$

where $\tilde{C}_{X_0} = e^{(-k)^+T}(1 + e^{(-k)^+T}(1 \vee T)(a - \sigma^2/4))^L$.

We are now in position to prove the claim. For $i \leq m$, we have:

$$\begin{aligned}
|\partial_x^i f(X_0(t, x))| & = |e^{-ikt} f^{(i)}(X_0(t, x))| \leq e^{-ikt} \|f\|_{m,L} (1 + X_0(t, x))^L \\
& \leq \|f\|_{m,L} (1 \vee e^{-mkt})(1 \vee e^{-Lkt})(1 + \tilde{C}_{X_0}t)(1 + x^L) \\
& \leq \|f\|_{m,L} e^{[\tilde{C}_{X_0} + (L+m)(-k)^+]t} (1 + x^L).
\end{aligned}$$

This gives $\|f(X_0(t, \cdot))\|_{m,L} \leq \|f\|_{m,L} e^{[\tilde{C}_{X_0} + (L+m)(-k)^+]t}$. \square

5.1. Proof of $(\overline{H_1})$. In this subsection, we prove the following result which is a direct consequence of Propositions 5.4 and 5.8 that are stated below.

Proposition 5.3. *Let Y satisfy (\mathcal{H}_Y) , $\sigma^2 \leq 4a$ and $\hat{X}_t^x = \varphi(x, t, \sqrt{t}Y)$ be the scheme (3.7). Let $m \in \mathbb{N}$, $L \in \mathbb{N}^*$ and $f \in \mathcal{C}_{\text{pol}}^{2(m+3),L}(\mathbb{R}_+)$. Then, there exists a constant $C \in \mathbb{R}_+^*$ such that for $t \in [0, T]$,*

$$\|\mathbb{E}[f(\hat{X}_t)] - \mathbb{E}[f(X_t)]\|_{m,L+3} \leq Ct^3 \|f\|_{2(m+3),L}.$$

To prove this result, we compare each term with the expansion $f(x) + t\mathcal{L}f(x) + \frac{t^2}{2}\mathcal{L}^2f(x)$ of order two. The next proposition analyses the difference between such expansion and the semigroup of the CIR process.

Proposition 5.4. *Let $m, \nu \in \mathbb{N}$, $L \in \mathbb{N}^*$, $T > 0$ and $f \in \mathcal{C}_{\text{pol}}^{m+2(\nu+1),L}(\mathbb{R}_+)$. Let X^x be the CIR process and \mathcal{L} its infinitesimal generator. Then, for $t \in [0, T]$, we have*

$$\mathbb{E}[f(X_t^x)] = \sum_{i=0}^{\nu} \frac{t^i}{i!} \mathcal{L}^i f(x) + t^{\nu+1} \int_0^1 \frac{(1-s)^\nu}{\nu!} \mathbb{E}[\mathcal{L}^{\nu+1} f(X_{ts}^x)] ds \quad (5.3)$$

where the function $x \mapsto \int_0^1 \frac{(1-s)^\nu}{\nu!} \mathbb{E}[\mathcal{L}^{\nu+1} f(X_s^x)] ds$ belongs to $\mathcal{C}_{\text{pol}}^{m,L}(\mathbb{R}_+)$ and we have the following estimate for all $t \in [0, T]$,

$$\left\| \int_0^1 \frac{(1-s)^\nu}{\nu!} \mathbb{E}[\mathcal{L}^{\nu+1} f(X_{ts}^x)] ds \right\|_{m,L+\nu+1} \leq C \|f\|_{m+2(\nu+1),L}, \quad (5.4)$$

for some constant $C \in \mathbb{R}_+$ depending on $(a, k, \sigma, \nu, m, L, T)$.

Proof. Let $f \in \mathcal{C}_{\text{pol}}^{m+2(\nu+1),L}(\mathbb{R}_+)$. Since the coefficients of the CIR SDE have sublinear growth, we have bounds on the moments of X_s^x : $\forall q \in \mathbb{N}^*$, $\exists C_q > 0$, $\forall s \in [0, T]$, $\mathbb{E}[|X_s^x|^q] \leq C_q(1+x^q)$. Using iterations of Itô's formula and a change of variable (in time), we then easily get (5.3) for $t \in [0, T]$. To get the estimate (5.4), we first use Lemma 5.1 and obtain

$$\|\mathcal{L}^{\nu+1} f\|_{m,L+\nu+1} \leq K_{\text{cir}}(m, \nu)^{\nu+1} \|f\|_{m+2(\nu+1),L},$$

with $K_{\text{cir}}(m, \nu) = 2a + (2m + 4\nu + 3)(|k| + \sigma^2/2)$. By the triangle inequality, we have

$$\left\| \int_0^1 \frac{(1-s)^\nu}{\nu!} \mathbb{E}[\mathcal{L}^{\nu+1} f(X_{ts})] ds \right\|_{m, L+\nu+1} \leq \int_0^1 \frac{(1-s)^\nu}{\nu!} \left\| \mathbb{E}[\mathcal{L}^{\nu+1} f(X_{ts})] \right\|_{m, L+\nu+1} ds.$$

Since $t \leq T$, we have $\left\| \mathbb{E}[\mathcal{L}^{\nu+1} f(X_{ts})] \right\|_{m, L+\nu+1} \leq C_{\text{cir}}(m, L + \nu + 1, T) \left\| \mathcal{L}^{\nu+1} f \right\|_{m, L+\nu+1}$ by Proposition 5.10. This gives by Lemma 5.1

$$\left\| \int_0^1 \frac{(1-s)^\nu}{\nu!} \mathbb{E}[\mathcal{L}^{\nu+1} f(X_{ts})] ds \right\|_{m, L+\nu+1} \leq \frac{C_{\text{cir}}(m, L + \nu + 1, T)}{(\nu + 1)!} K_{\text{cir}}(m, \nu)^{\nu+1} \|f\|_{m+2(\nu+1), L}.$$

□

We now focus on the approximation scheme. The main difficulty comes from the differentiation of the square-root that may lead to derivatives that blow up at the origin. Here, we exploit the fact that Y is a symmetric random variable to cancel these blowing terms. More precisely, we will then need to differentiate in x the following quantity

$$g(X_1(s\sqrt{t}, x)) + g(X_1(-s\sqrt{t}, x)) = g\left(x + \sigma s\sqrt{t}\sqrt{x} + \frac{\sigma^2}{4}ts^2\right) + g\left(x - \sigma s\sqrt{t}\sqrt{x} + \frac{\sigma^2}{4}ts^2\right),$$

and the next lemma enables us to have a sharp estimate of the derivatives.

Lemma 5.5. *Let $g : \mathbb{R}_+ \rightarrow \mathbb{R}$ be a \mathcal{C}^{2n} function, $\beta \in \mathbb{R}_+$ and $\gamma \geq \beta^2/4$. Then, the function $\psi_g(x) := g(x + \beta\sqrt{x} + \gamma) + g(x - \beta\sqrt{x} + \gamma)$, $x \geq 0$ is \mathcal{C}^n with derivatives*

$$\psi_g^{(n)}(x) = \psi_{g^{(n)}}(x) + \sum_{j=1}^n \binom{n}{j} \beta^{2j} \int_0^1 g^{(n+j)}(x + \beta(2u-1)\sqrt{x} + \gamma) \frac{(u-u^2)^{j-1}}{(j-1)!} du \quad (5.5)$$

Proof. For $x > 0$, we have

$$\begin{aligned} \psi'_g(x) &= \left(1 + \frac{\beta}{2\sqrt{x}}\right) g'(x + \beta\sqrt{x} + \gamma) + \left(1 - \frac{\beta}{2\sqrt{x}}\right) g'(x - \beta\sqrt{x} + \gamma) \\ &= \underbrace{g'(x + \beta\sqrt{x} + \gamma) + g'(x - \beta\sqrt{x} + \gamma)}_{\psi_{g'}(x)} + \beta^2 \int_0^1 g''(x + \beta(2u-1)\sqrt{x} + \gamma) du, \end{aligned}$$

since $\frac{d}{du} g'(x + \beta(2u-1)\sqrt{x} + \gamma) = 2\beta\sqrt{x} g''(x + \beta(2u-1)\sqrt{x} + \gamma)$. Clearly, this derivative is continuous at 0 which shows that ψ_g is C^1 .

We are now in position to prove (5.5) by induction on n . It is true for $n = 0, 1$. We assume that it is true for n . Then, we get by using the case $n = 1$, differentiating (5.5) and an integration by parts for the fourth term:

$$\begin{aligned} \psi_g^{(n+1)}(x) &= \psi_{g^{(n+1)}}(x) + \beta^2 \int_0^1 g^{(n+2)}(x + \beta(2u-1)\sqrt{x} + \gamma) du \\ &\quad + \sum_{j=1}^n \binom{n}{j} \beta^{2j} \left(\int_0^1 g^{(n+j+1)}(x + \beta(2u-1)\sqrt{x} + \gamma) \frac{(u-u^2)^{j-1}}{(j-1)!} du \right. \\ &\quad \left. + \beta^2 \int_0^1 g^{(n+j+2)}(x + \beta(2u-1)\sqrt{x} + \gamma) \frac{(u-u^2)^j}{j!} du \right). \end{aligned}$$

We then reorganize the terms as follows

$$\begin{aligned} \psi_g^{(n+1)}(x) &= \psi_{g^{(n+1)}}(x) + (n+1)\beta^2 \int_0^1 g^{(n+2)}(x + \beta(2u-1)\sqrt{x} + \gamma) du \\ &\quad + \beta^{2n+2} \int_0^1 g^{(2n+2)}(x + \beta(2u-1)\sqrt{x} + \gamma) \frac{(u-u^2)^n}{n!} du \\ &\quad + \sum_{j=2}^n \binom{n}{j} \beta^{2j} \left(\int_0^1 g^{(n+j+1)}(x + \beta(2u-1)\sqrt{x} + \gamma) \frac{(u-u^2)^{j-1}}{(j-1)!} du \right) \\ &\quad + \sum_{j=1}^{n-1} \binom{n}{j} \beta^{2j+2} \left(\int_0^1 g^{(n+j+2)}(x + \beta(2u-1)\sqrt{x} + \gamma) \frac{(u-u^2)^j}{j!} du \right). \end{aligned}$$

The last sum is equal to $\sum_{j=2}^n \binom{n}{j-1} \beta^{2j} \left(\int_0^1 g^{(n+j+1)}(x + \beta(2u-1)\sqrt{x} + \gamma) \frac{(u-u^2)^{j-1}}{(j-1)!} du \right)$ by changing j to $j-1$, and we conclude by using that $\binom{n}{j} + \binom{n}{j-1} = \binom{n+1}{j}$. \square

Corollary 5.6. *Let $m, L \in \mathbb{N}$, $\beta \geq 0$ and $g \in \mathcal{C}_{\text{pol}}^{2m,L}(\mathbb{R}_+)$. Then, $\psi_g(x) = g((\sqrt{x} + \beta/2)^2) + g((\sqrt{x} - \beta/2)^2)$ belongs to $\mathcal{C}_{\text{pol}}^{m,L}(\mathbb{R}_+)$, and for all $n \in \{0, \dots, m\}$ we have the following estimates*

$$\|\psi_g\|_{n,L} \leq C_{\beta,m,L} \|g\|_{2n,L}, \quad (5.6)$$

with $C_{\beta,m,L} = ((1 + \beta/2)^{2L} + (1 - \beta/2)^{2L} + 2(1 + \beta^2/2)^L(1 + \beta^2/2)^m)$.

Proof. We use (5.5) with $\gamma = \beta^2/4$. We first notice that

$$\begin{aligned} |\psi_{g^{(n)}}(x)| &\leq \|g\|_{n,L} (2 + (\sqrt{x} + \beta/2)^{2L} + (\sqrt{x} - \beta/2)^{2L}) \\ &= \|g\|_{n,L} \left(2 + 2x^L + 2 \sum_{i=1}^L \binom{2L}{2i} (\beta/2)^{2i} x^{L-i} \right). \end{aligned}$$

Using that $x^i \leq 1 + x^L$ for $0 \leq i \leq L-1$, we get

$$|\psi_{g^{(n)}}(x)| \leq 2\|g\|_{n,L} (1 + x^L) \sum_{i=0}^L \binom{2L}{2i} (\beta/2)^{2i} = \|g\|_{n,L} (1 + x^L) ((1 + \beta/2)^{2L} + (1 - \beta/2)^{2L}).$$

For the other terms, we use that for $u \in [0, 1]$, $x \geq 0$ and $j \in \{1, \dots, n\}$,

$$\begin{aligned} |g^{(n+j)}(x + \beta(2u-1)\sqrt{x} + \beta^2/4)| &\leq \|g\|_{2n,L} (1 + (x + \beta(2u-1)\sqrt{x} + \beta^2/4)^L) \\ &\leq \|g\|_{2n,L} (1 + (\sqrt{x} + \beta/2)^{2L}). \end{aligned}$$

We again expand $(\sqrt{x} + \beta/2)^L = x^L + \sum_{i=1}^{2L} \binom{2L}{i} (\beta/2)^i x^{L-i/2}$ and use that $x^{L-i/2} \leq 1 + x^L$ to get

$$|g^{(n+j)}(x + \beta(2u-1)\sqrt{x} + \beta^2/4)| \leq \|g\|_{2n,L} (1 + \beta/2)^{2L} (1 + x^L).$$

Besides, we have $u - u^2 \leq 1/4$ for $u \in [0, 1]$ and thus $\int_0^1 (u - u^2)^j du \leq \frac{1}{4^j}$, which gives $\sup_{x \geq 0} \frac{|\psi_{g^{(n)}}(x)|}{1+x^L} \leq \tilde{C}(\beta)$ with

$$\tilde{C}(\beta) = \|g\|_{n,L} ((1 + \beta/2)^{2L} + (1 - \beta/2)^{2L}) + \|g\|_{2n,L} (1 + \beta/2)^{2L} \sum_{j=1}^n \binom{n}{j} \left(\frac{\beta^2}{4} \right)^j$$

$$\begin{aligned}
&= \|g\|_{n,L}((1 + \beta/2)^{2L} + (1 - \beta/2)^{2L}) + \|g\|_{2n,L}(1 + \beta/2)^{2L}(1 + \beta^2/4)^n \\
&\leq \|g\|_{2n,L}((1 + \beta/2)^{2L} + (1 - \beta/2)^{2L} + (1 + \beta/2)^{2L}(1 + \beta^2/4)^m) = C_{\beta,m,L}\|g\|_{2n,L},
\end{aligned}$$

which gives the claim. \square

Lemma 5.7. *Let $\nu \in \mathbb{N}, L \in \mathbb{N}^*, T > 0, t \in [0, T]$. Let Y be a symmetric random variable such that $\mathbb{E}[Y^k] = \mathbb{E}[N^k]$ for $k \leq 2\nu$ and $\mathbb{E}[Y^{2k}] < \infty$ for all $k \in \mathbb{N}$. We have, for $f \in C_{\text{pol}}^{m+\nu+1,L}(\mathbb{R}_+)$,*

$$f(X_0(t, x)) = \sum_{i=0}^{\nu} \frac{t^i}{i!} V_0^i f(x) + t^{\nu+1} \int_0^t \frac{(1-u)^\nu}{\nu!} V_0^{\nu+1} f(X_0(ut, x)) du, \quad (5.7)$$

with $\| \int_0^1 \frac{(1-u)^\nu}{\nu!} V_0^{\nu+1} f(X_0(ut, x)) du \|_{m,L+\nu+1} \leq C_0 \|f\|_{m+\nu+1,L}$; and for $f \in C_{\text{pol}}^{2(m+\nu+1),L}(\mathbb{R}_+)$,

$$\begin{aligned}
\mathbb{E}[f(X_1(\sqrt{t}Y, x))] &= \sum_{i=0}^{\nu} \frac{t^i}{i!} \left(\frac{1}{2} V_1^2\right)^i f(x) \\
&\quad + t^{\nu+1} \mathbb{E} \left[Y^{2\nu+2} \int_0^1 \frac{(1-u)^{2\nu+1}}{(2\nu+1)!} V_1^{2\nu+2} f(X_1(u\sqrt{t}Y, x)) du \right],
\end{aligned} \quad (5.8)$$

with $\| \mathbb{E} \left[Y^{2\nu+2} \int_0^1 \frac{(1-u)^{2\nu+1}}{(2\nu+1)!} V_1^{2\nu+2} f(X_1(u\sqrt{t}Y, x)) du \right] \|_{m,L+\nu+1} \leq C_1 \|f\|_{2(m+\nu+1),L}$, for some constants $C_0, C_1 \in \mathbb{R}^+$ depending on $(a, k, \sigma), T, m, M$ and ν .

Proof. Equation (5.7) holds by using Taylor formula since $\frac{d}{dt} f(X_0(t, x)) = V_0 f(X_0(t, x))$. We have by Property (6) of Lemma 5.1 $\|V_0 f\|_{m,L+1} \leq |a - \frac{\sigma^2}{4}| \|f'\|_{m,L+1} + |k|(2m+3) \|f'\|_{m,L} \leq (2|a - \frac{\sigma^2}{4}| + |k|(2m+3)) \|f\|_{m+1,L}$ and thus $\|V_0^{\nu+1} f\|_{m,L+\nu+1} \leq C \|f\|_{m+\nu+1,L}$ for some constant C depending on a, σ, k, ν, m . Using the triangular inequality and Lemma 5.2, we get the result.

Equation (5.8) holds by using Taylor formula and then taking the expectation, using that $\mathbb{E}[Y^{2k+1}] = 0$. We have $\|\frac{1}{2} V_1^2 f\|_{m,L+1} \leq \sigma^2(m+2) \|f\|_{m+2,L}$ by using Lemma 5.1 (6). Then, we observe that by symmetry of Y ,

$$\begin{aligned}
&\mathbb{E} \left[Y^{2\nu+2} \int_0^1 \frac{(1-u)^{2\nu+1}}{(2\nu+1)!} V_1^{2\nu+2} f(X_1(u\sqrt{t}Y, x)) du \right] \\
&= \frac{1}{2} \mathbb{E} \left[Y^{2\nu+2} \int_0^1 \frac{(1-u)^{2\nu+1}}{(2\nu+1)!} [V_1^{2\nu+2} f(X_1(u\sqrt{t}Y, x)) + V_1^{2\nu+2} f(X_1(-u\sqrt{t}Y, x))] du \right]
\end{aligned}$$

By Corollary 5.6, we have

$$\begin{aligned}
\|V_1^{2\nu+2} f(X_1(u\sqrt{t}Y, \cdot)) + V_1^{2\nu+2} f(X_1(-u\sqrt{t}Y, \cdot))\|_{m,L+\nu+1} &\leq C_{\sigma u\sqrt{t}Y, m, L} \|V_1^{2\nu+2} f\|_{2m, L+\nu+1} \\
&\leq C' C_{\sigma u\sqrt{t}Y, m, L} \|f\|_{2(m+\nu+1), L},
\end{aligned}$$

with $C' = (4\sigma^2(m + \nu + 1))^{\nu+1}$. The conclusion follows by using the triangle inequality, the polynomial growth of the constant $C_{\sigma u\sqrt{t}Y, m, L}$ given by Corollary 5.6 and the finite moments $\mathbb{E}[Y^{2k}]$ for k sufficiently large. \square

We are now in position to prove the estimate for the approximation scheme (3.7). Since this scheme is obtained as the composition of the schemes X_0 and X_1 , the proof consists in using iteratively the estimates of Lemma 5.7.

Proposition 5.8. *Let Y be a symmetric random variable such that $\mathbb{E}[Y^k] = \mathbb{E}[N^k]$ for $k \leq 4$ and $\mathbb{E}[Y^{2k}] < \infty$ for all $k \in \mathbb{N}$. Let $\sigma^2 \leq 4a$ and \hat{X}_t^x be the scheme (3.7). Let $m \in \mathbb{N}$, $L \in \mathbb{N}^*$ and $f \in \mathcal{C}_{\text{pol}}^{2(m+3),L}(\mathbb{R}_+)$. Then, we have for $t \in [0, T]$,*

$$\mathbb{E}[f(\hat{X}_t^x)] = f(x) + t\mathcal{L}f(x) + \frac{t^2}{2}\mathcal{L}^2f(x) + \bar{R}f(t, x),$$

with $\|\bar{R}f(t, \cdot)\|_{m,L+3} \leq Ct^3\|f\|_{2(m+3),L}$.

Proof. We use $\hat{X}_t^x = X_0(t/2, X_1(\sqrt{t}Y, X_0(t/2, x)))$ and apply first (5.7):

$$\mathbb{E}[f(X_0(t/2, X_1(\sqrt{t}Y, X_0(t/2, x))))] = \mathbb{E}\left[\left(f + \frac{t}{2}V_0f + \frac{t^2}{8}V_0^2f\right)(X_1(\sqrt{t}Y, X_0(t/2, x)))\right] + R_I f(t, x),$$

$$\text{with } R_I f(t, x) = \left(\frac{t}{2}\right)^3 \int_0^1 \frac{(1-u)^2}{2} \mathbb{E}[V_0^3 f(X_0(ut/2, X_1(\sqrt{t}Y, X_0(t/2, x))))] ds.$$

We get by using Lemma 5.2, Corollary 5.6 (using the symmetry and the finite moments of Y), again Lemma 5.2 and then Lemma 5.1 (6):

$$\begin{aligned} \|\mathbb{E}[V_0^3 f(X_0(ut/2, X_1(\sqrt{t}Y, X_0(t/2, \cdot)))]\|_{m,L+3} &\leq C\|\mathbb{E}[V_0^3 f(X_0(ut/2, X_1(\sqrt{t}Y, \cdot)))]\|_{m,L+3} \\ &\leq C\|V_0^3 f(X_0(ut/2, \cdot))\|_{2m,L+3} \leq C\|V_0^3 f\|_{2m,L+3} \leq C\|f\|_{2m+3,L} \end{aligned}$$

This gives $\|R_I f(t, x)\|_{m,L+3} \leq Ct^3\|f\|_{2m+3,L}$, for $t \in [0, T]$.

We now expand again and get from (5.8)

$$\begin{aligned} &\mathbb{E}\left[\left(f + \frac{t}{2}V_0f + \frac{t^2}{8}V_0^2f\right)(X_1(\sqrt{t}Y, X_0(t/2, x)))\right] \\ &= f(X_0(t/2, x)) + \frac{t}{2}V_1^2f(X_0(t/2, x)) + \frac{t^2}{2}(V_1^2/2)^2f(X_0(t/2, x)) + \frac{t}{2}V_0f(X_0(t/2, x)) \\ &\quad + \frac{t^2}{2}(V_1^2/2)V_0f(X_0(t/2, x)) + \frac{t^2}{8}V_0^2f(X_0(t/2, x)) + R_{II}f(t, x), \end{aligned}$$

with

$$\begin{aligned} R_{II}f(t, x) &= t^3\mathbb{E}\left[Y^6 \int_0^u \frac{(1-u)^5}{5!} V_1^6 f(X_1(u\sqrt{t}Y, X_0(T/2, x))) du \right. \\ &\quad + \frac{Y^4}{2} \int_0^u \frac{(1-u)^3}{3!} V_1^4 V_0 f(X_1(u\sqrt{t}Y, X_0(T/2, x))) du \\ &\quad \left. + \frac{Y^2}{8} \int_0^u (1-u)V_1^2 V_0^2 f(X_1(u\sqrt{t}Y, X_0(T/2, x))) du \right]. \end{aligned}$$

We use Lemmas 5.7, 5.2 and 5.1 to get, for $t \in [0, T]$,

$$\|R_{II}f(t, \cdot)\|_{m,L+3} \leq Ct^3(\|f\|_{2(m+3),L} + \|V_0f\|_{2(m+2),L+1} + \|V_0^2f\|_{2(m+1),L+2}) \leq Ct^3\|f\|_{2(m+3),L}.$$

Last, we use again (5.7) to get

$$\begin{aligned} &f(X_0(t/2, x)) + \frac{t}{2}[V_1^2 + V_0]f(X_0(t/2, x)) + \frac{t^2}{2}[(V_1^2/2)^2 + (V_1^2/2)V_0 + V_0^2/4]f(X_0(t/2, x)) \\ &= f(x) + \frac{t}{2}V_0f(x) + \frac{t^2}{8}V_0^2f(x) + \frac{t}{2}[V_1^2 + V_0]f(x) + \frac{t^2}{4}[V_0V_1^2 + V_0^2]f(x) \end{aligned}$$

$$+ \frac{t^2}{2} [(V_1^2/2)^2 + (V_1^2/2)V_0 + V_0^2/4]f(x) + R_{III}f(t, x),$$

where again by Lemma 5.7 and 5.1, we have

$$\begin{aligned} \|R_{III}f(t, \cdot)\|_{m, L+3} &\leq Ct^3 (\|f\|_{m+3, L} + \|[V_1^2 + V_0]f\|_{m+2, L+1} \\ &\quad + \|[V_1^2/2]^2 + (V_1^2/2)V_0 + V_0^2/4]f\|_{m+1, L+2}) \\ &\leq Ct^3 \|f\|_{m+5, L}. \end{aligned}$$

Finally, we get $\|\bar{R}f(t, \cdot)\|_{m, L+3} \leq Ct^3 \|f\|_{2(m+3), L}$ with $\bar{R}f := R_I f + R_{II} f + R_{III} f$ and

$$\begin{aligned} \mathbb{E}[f(X_0(t/2, X_1(\sqrt{t}Y, X_0(t/2, x)))] &= f(x) + t[V_0 + V_1^2/2]f(x) \\ &\quad + \frac{t^2}{2} [V_0^2 + V_0V_1^2/2 + (V_1^2/2)V_0 + (V_1^2/2)^2]f(x) + \bar{R}f(t, x) \\ &= f(x) + t\mathcal{L}f(x) + \frac{t^2}{2}\mathcal{L}^2f(x) + \bar{R}f(t, x). \end{aligned} \quad \square$$

5.2. Proof of (\bar{H}_2) . In this section, we mainly prove the following result.

Proposition 5.9. *Let $\sigma^2 \leq 4a$ and $\hat{X}_t^x = \varphi(x, t, \sqrt{t}N)$ be the scheme (3.1) with $N \sim \mathcal{N}(0, 1)$. Let $T > 0$ and $m, L \in \mathbb{N}$ such that $L \geq m$. We define for $n \geq 1$ and $l \in \mathbb{N}$, $Q_l f(x) = \mathbb{E}[f(\hat{X}_{h_l}^x)]$ with $h_l = \frac{T}{n^l}$. Then, there exists a constant $C \in \mathbb{R}_+^*$ such that*

$$\forall f \in \mathcal{C}_{\text{pol}}^{m, L}(\mathbb{R}_+), \forall t \in [0, T], l \geq 0, \|\mathbb{E}[f(X_t)]\|_{m, L} + \max_{0 \leq k \leq n^l} \|Q_l^{[k]} f\|_{m, L} \leq C \|f\|_{m, L}. \quad (5.9)$$

We split the proof in two parts. The first one deals with the semigroup of the CIR process, for which the assumption $\sigma^2 \leq 4a$ is not needed. This is stated in Proposition 5.10, whose proof exploits the particular form of the density of X_t^x . The second part that deals with the approximation scheme is quite technical. We prove in fact in Proposition 5.12 a slightly more general result for the scheme $\hat{X}_t^x = \varphi(x, t, \sqrt{t}Y)$, when Y is a symmetric random variable with a smooth density. However, the conditions needed on the density are quite restrictive. These conditions are satisfied by the standard normal variable by Lemma 5.14. If we want besides to have (5.9) for any m and in addition to match the moments $\mathbb{E}[Y^2] = \mathbb{E}[N^2]$ and $\mathbb{E}[Y^4] = \mathbb{E}[N^4]$ – which is required to have a second-order scheme –, then Theorem 5.16 shows that we necessarily have $Y \sim \mathcal{N}(0, 1)$. This is why we directly state here, for sake of simplicity, Proposition 5.9 with $Y = N \sim \mathcal{N}(0, 1)$.

5.2.1. Upper bound for the semigroup. We first prove the estimate (\bar{H}_2) for the semigroup of the CIR process. To do so, we take back the arguments of [1, Proposition 4.1] that gives polynomial estimates for $(t, x) \mapsto P_t f(x)$. First we remove the polynomial Taylor expansion of the function f at 0, which enables then to do an integration by parts and to get the remarkable formula in Eq. (5.13) below for the iterated derivatives of $P_t f$ that gives then the desired estimate. The polynomial part is analyzed separately in Lemma 5.11 below with standard arguments.

Proposition 5.10. *Let $f \in \mathcal{C}_{\text{pol}}^{m, L}(\mathbb{R}_+)$, $L \geq m$, $T > 0$ and $t \in (0, T]$. Let X^x be the CIR process starting from $x \geq 0$. Then, $\mathbb{E}[f(X_t)] \in \mathcal{C}_{\text{pol}}^{m, L}(\mathbb{R}_+)$ and we have the following estimate for some constant $C_{\text{cir}}(m, L, T) \in \mathbb{R}_+$:*

$$\|\mathbb{E}[f(X_t)]\|_{m, L} \leq C_{\text{cir}}(m, L, T) \|f\|_{m, L}. \quad (5.10)$$

Proof. Let $f \in \mathcal{C}_{\text{pol}}^{m,L}(\mathbb{R}_+)$ and $T_m(f)(x) = \sum_{j=0}^m \frac{f^{(j)}(0)}{j!} x^j$ its Taylor polynomial expansion at 0 of order m . We define $\hat{f}_m = f - T_m(f) \in \mathcal{C}_{\text{pol}}^{m,L}(\mathbb{R}_+)$, so we have $f = \hat{f}_m + T_m(f)$. By Lemma 5.11 below, one gets $\|\mathbb{E}[T_m(f)(X_t)]\|_{m,L} \leq C_{\text{cir}}(m, T) \|T_m(f)\|_{m,L}$ and then

$$\|\mathbb{E}[T_m(f)(X_t)]\|_{m,L} \leq e C_{\text{cir}}(m, T) \|f\|_{m,L}, \quad (5.11)$$

since for all $i \in \{0, \dots, m\}$ and $x \geq 0$

$$\left| \frac{(T_m(f))^{(i)}(x)}{1+x^L} \right| = \left| \sum_{j=0}^{m-i} \frac{f^{(i+j)}(0)}{j!} \frac{x^j}{1+x^L} \right| \leq \sum_{j=0}^{m-i} \frac{1}{j!} \|f\|_{m,L} \leq e \|f\|_{m,L}.$$

We now focus on $\mathbb{E}[\hat{f}_m(X_t)]$. We recall the density of X_t^x (see e.g. [3, Proposition 1.2.11])²

$$p(t, x, z) = \sum_{i=0}^{\infty} \frac{e^{-d_t x/2} (d_t x/2)^i}{i!} \frac{c_t/2}{\Gamma(i+v)} \left(\frac{c_t z}{2}\right)^{i-1+v} e^{-c_t z/2} \quad (5.12)$$

where $c_t = \frac{4k}{\sigma^2(1-e^{-kt})}$, $v = 2a/\sigma^2$ and $d_t = c_t e^{-kt}$. Let us remark that

$$c_t \geq c_{\min} := \begin{cases} \frac{4k}{\sigma^2}, & k > 0 \\ \frac{4}{\sigma^2 T}, & k = 0 \\ \frac{4|k|}{\sigma^2(e^{|k|T}-1)}, & k < 0. \end{cases}$$

We have

$$\mathbb{E}[\hat{f}_m(X_t^x)] = \sum_{i=0}^{\infty} \frac{e^{-d_t x/2} (d_t x/2)^i}{i!} I_i(\hat{f}_m, c_t), \quad t > 0,$$

where

$$I_i(\hat{f}_m, c_t) = \int_0^{\infty} \hat{f}_m(z) \frac{c_t/2}{\Gamma(i+v)} \left(\frac{c_t z}{2}\right)^{i-1+v} e^{-c_t z/2} dz.$$

Differentiating successively, we get that for $j \leq m$, $t \in (0, T]$ and $x \in \mathbb{R}_+$

$$\partial_j^x \mathbb{E}[\hat{f}(X_t^x)] = \sum_{i=0}^{\infty} \frac{e^{-d_t x/2} (d_t x/2)^i}{i!} \Delta_t^j(I_i(\hat{f}_m, c_t)),$$

where $\Delta_t : \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}^{\mathbb{N}}$ is an operator defined on sequences $(I_i)_{i \geq 0} \in \mathbb{R}^{\mathbb{N}}$ by $\Delta_t(I_i) = \frac{d_t}{2}(I_{i+1} - I_i) = \frac{e^{-kt}}{2} c_t (I_{i+1} - I_i)$. An integration by parts gives for $i \geq 1$

$$\begin{aligned} I_i(\hat{f}_m^{(j)}, c_t) &= \int_0^{\infty} \hat{f}_m^{(j-1)}(z) \frac{(c_t/2)^2}{\Gamma(i+v)} \left(\frac{c_t z}{2}\right)^{i-1+v} e^{-c_t z/2} dz \\ &\quad - \int_0^{\infty} \hat{f}_m^{(j-1)}(z) \frac{(c_t/2)^2 (i-1+v)}{\Gamma(i+v)} \left(\frac{c_t z}{2}\right)^{i-2+v} e^{-c_t z/2} dz \\ &= \frac{c_t}{2} (I_i(\hat{f}_m^{(j-1)}, c_t) - I_{i-1}(\hat{f}_m^{(j-1)}, c_t)) = e^{kt} \Delta_t(I_{i-1}(\hat{f}_m^{(j-1)}, c_t)), \end{aligned}$$

²In the case $a = 0$, X_t^x is distributed according to the probability measure $e^{-d_t x/2} \delta_0(dx) + \sum_{i=1}^{\infty} \frac{e^{-d_t x/2} (d_t x/2)^i}{i!} \frac{c_t/2}{\Gamma(i)} \left(\frac{c_t z}{2}\right)^{i-1} e^{-c_t z/2}$. The proof works the same since $\hat{f}_m(0) = 0$, so that $\mathbb{E}[\hat{f}_m(X_t^x)]$ only involves the absolutely continuous part of the distribution.

since $\hat{f}_m^{(j)}(0) = 0$ for all $1 \leq j \leq m$ and $\hat{f}_m^{(j)}$ has a polynomial growth. By iterating, we get for all $t \in (0, T]$ and $x \in \mathbb{R}_+$,

$$\partial_j^x \mathbb{E}[\hat{f}_m(X_t^x)] = \sum_{i=0}^{\infty} \frac{e^{-d_t x/2} (d_t x/2)^i}{i!} I_{i+j}(\hat{f}_m^{(j)}, c_t) e^{-kjt}. \quad (5.13)$$

Note that, since for $j \leq m$, $|\hat{f}_m^{(j)}(z)| \leq \|\hat{f}_m\|_{m,L}(1+z^L)$ and using the well known formula for the L -th raw moment of gamma distribution we have for all $i \in \mathbb{N}$

$$|I_i(\hat{f}_m^{(j)}, c_t)| \leq \|\hat{f}_m\|_{m,L} \left(1 + \left(\frac{2}{c_t} \right)^L \frac{\Gamma(i+L+v)}{\Gamma(i+v)} \right). \quad (5.14)$$

Thus, the derivation of the series (5.13) is valid, and we get that

$$|\partial_j^x \mathbb{E}[\hat{f}_m(X_t^x)]| \leq \|\hat{f}_m\|_{m,L} e^{-kjt} \left(1 + \left(\frac{2}{c_t} \right)^L \sum_{i=0}^{\infty} \frac{e^{-d_t x/2} (d_t x/2)^i}{i!} \frac{\Gamma(i+j+L+v)}{\Gamma(i+j+v)} \right).$$

The quotient $\frac{\Gamma(i+j+L+v)}{\Gamma(i+j+v)}$ is a polynomial function of degree L with respect to i , and we denote $\beta_0^j, \dots, \beta_L^j$ its coefficients in the basis $\{1, i, i(i-1), \dots, i(i-1) \cdots (i-L+1)\}$. Thus, we get that

$$\begin{aligned} |\partial_j^x \mathbb{E}[\hat{f}_m(X_t^x)]| &\leq \|\hat{f}_m\|_{m,L} e^{(-k)^+ jT} \left(1 + \left(\frac{2}{c_t} \right)^L \sum_{i=0}^L |\beta_i^j| \left(\frac{d_t}{2} \right)^i x^i \right) \\ &\leq \|\hat{f}_m\|_{m,L} e^{(-k)^+ mT} \left(1 + \sum_{i=0}^L |\beta_i^j| \left(\frac{2}{c_t} \right)^{L-i} e^{-kit} (1+x^L) \right) \\ &\leq \|\hat{f}_m\|_{m,L} e^{(-k)^+ (m+L)T} \left(1 + \sum_{i=0}^L |\beta_i^j| \left(\frac{2}{c_{\min}} \right)^{L-i} \right) (1+x^L). \end{aligned}$$

By the triangular inequality and (5.11), we get $\|\hat{f}_m\|_{m,L} \leq (1+e)\|f\|_{m,L}$, so one has for all $t \in (0, T]$, $j \leq m$

$$|\partial_j^x \mathbb{E}[\hat{f}_m(X_t^x)]| \leq (1+e) e^{(-k)^+ (m+L)T} \left(1 + \sum_{i=0}^L |\beta_i^j| \left(\frac{2}{c_{\min}} \right)^{L-i} \right) \|f\|_{m,L} (1+x^L), \quad (5.15)$$

and thus for all $t \in (0, T]$:

$$\|\mathbb{E}[\hat{f}_m(X_t)]\|_{m,L} \leq \hat{C} \|f\|_{m,L}, \quad (5.16)$$

where $\hat{C} := (1+e) e^{(-k)^+ (m+L)T} \max_{0 \leq j \leq m} \left(1 + \sum_{i=0}^L |\beta_i^j| \left(\frac{2}{c_{\min}} \right)^{L-i} \right)$. Finally, we get the desired estimate by the triangular inequality, (5.11) and Lemma 5.11:

$$\|\mathbb{E}[f(X_t)]\|_{m,L} \leq \|\mathbb{E}[\hat{f}_m(X_t)]\|_{m,L} + \|\mathbb{E}[T_m(f)(X_t)]\|_{m,L} \leq (\hat{C} + C_{\text{cir}}(m, T)) \|f\|_{m,L}. \quad \square$$

Lemma 5.11. *Let $P \in \mathfrak{P}_m(\mathbb{R}_+)$ be a polynomial function of degree $m \in \mathbb{N}^*$ and $L \in \mathbb{N}^*$ such that $L \geq m$. Then, for $t \in [0, T]$ we have the following estimate*

$$\|\mathbb{E}[P(X_t)]\|_{m,L} \leq C_{\text{cir}}(m, T) \|P\|_{m,L}, \quad (5.17)$$

where $C_{\text{cir}}(m, T) = \max_{t \in [0, T]} \sum_{j=0}^m \sum_{i=0}^j |\tilde{u}_{i,j}(t)|$.

Proof. We consider a polynomial function $P(y) = \sum_{i=0}^m b_i y^i$ of degree m and $L \geq m$. For all $l \in \{0, \dots, m\}$ one has from Lemma 4.3

$$\begin{aligned} \frac{|\partial_x^l \mathbb{E}[P(X_t^x)]|}{1+x^L} &= \left| \sum_{j=0}^m b_j \frac{\partial_x^l \tilde{u}_j(t, x)}{1+x^L} \right| \leq \left| \sum_{j=0}^m b_j \sum_{i=l}^j \tilde{u}_{i,j}(t) \frac{i!}{(i-l)!} \frac{x^{i-l}}{1+x^L} \right| \\ &\leq \sum_{j=0}^m |b_j| j! \sum_{i=l}^j |\tilde{u}_{i,j}(t)| \leq \max_{t \in [0, T]} \sum_{j=0}^m \sum_{i=l}^j |\tilde{u}_{i,j}(t)| \max_{j \in \{0, \dots, L\}} |b_j| j!, \end{aligned}$$

passing to supremum over $x \geq 0$, $l \in \{0, \dots, m\}$ we get (5.17) observing that $|b_j| j! = |P^{(j)}(0)| \leq \|P\|_{m,L}$. \square

5.2.2. *Upper bound for the approximation scheme.* We now prove the estimate (\overline{H}_2) for the approximation of the CIR process. The main result of this paragraph is the following.

Proposition 5.12. *Let $T > 0$, $\sigma^2 \leq 4a$, $m, M \in \mathbb{N}$, Y be a symmetric random variable with density $\eta \in \mathcal{C}^M(\mathbb{R})$ such that for all $i \in \{0, \dots, M\}$, $|\eta^{(i)}(y)| = o(|y|^{2L+i})$ for $|y| \rightarrow \infty$, and $\eta_m^* \geq 0$ for all $1 \leq m \leq M$ (see Lemma 5.13 below for the definition of η_m^*). Let $Q_l f(x) = \mathbb{E}[f(\hat{X}_{h_l}^x)]$ with $\hat{X}_t^x = \varphi(t, x, \sqrt{t}Y)$, $n \geq 1$, $l \in \mathbb{N}$ and $h_l = T/n^l$. Then, we have*

$$\forall L \in \mathbb{N}, \exists C \in \mathbb{R}_+, \forall f \in \mathcal{C}_{\text{pol}}^{m,L}(\mathbb{R}_+), l \in \mathbb{N}, \max_{0 \leq j \leq n^l} \|Q_l^{[j]} f\|_{m,L} \leq C \|f\|_{m,L}.$$

Note that by Lemma 5.14 below, the assumptions of Proposition 5.12 are satisfied by $Y \sim \mathcal{N}(0, 1)$. Therefore, (\overline{H}_2) holds for the scheme of Ninomiya and Victoir (3.1).

Proof. We have $\hat{X}_t^x = \varphi(t, x, \sqrt{t}Y) = X_0(t/2, X_1(\sqrt{t}Y, X_0(t/2, x)))$. Let $f \in \mathcal{C}_{\text{pol}}^{m,L}(\mathbb{R}_+)$. We apply Lemma 5.2 and Lemma 5.13 below to get:

$$\begin{aligned} \|\mathbb{E}[f(X_0(t/2, X_1(\sqrt{t}Y, X_0(t/2, \cdot)))]\|_{m,L} &\leq e^{Kt/2} \|\mathbb{E}[f(X_1(\sqrt{t}Y, X_0(t/2, \cdot)))]\|_{m,L} \\ &\leq e^{Kt/2 + Ct} \|f(X_0(t/2, \cdot))\|_{m,L} \leq e^{(C+K)t} \|f\|_{m,L} \end{aligned}$$

This gives $\max_{0 \leq j \leq n^l} \|Q_l^{[j]} f\|_{m,L} \leq e^{(C+K)T} \|f\|_{m,L}$. \square

Lemma 5.13. *Let $M, L \in \mathbb{N}$. Let Y be a symmetric random variable with density $\eta \in \mathcal{C}^M(\mathbb{R})$ such that for all $i \in \{0, \dots, M\}$, $|\eta^{(i)}(y)| = o(|y|^{2L+i})$ for $|y| \rightarrow \infty$. Then, for all function $f \in \mathcal{C}_{\text{pol}}^{M,L}(\mathbb{R}_+)$, $m \in \{1, \dots, M\}$ and $t \in [0, T]$ one has the following representation*

$$\partial_x^m \mathbb{E}[f(X_1(\sqrt{t}Y, x))] = \int_{-\infty}^{\infty} \int_0^1 (u-u^2)^{m-1} f^{(m)}(w(u, x, y)) \eta_m^*(y) dy du \quad (5.18)$$

where $w(u, x, y) = x + (2u-1)\sigma\sqrt{ty}\sqrt{x} + \sigma^2 ty^2/4$, $\eta_m^*(y) = (-1)^{m-1} \left(\sum_{j=1}^m c_{j,m} y^j \eta^{(j)}(y) \right)$, and the coefficients $c_{j,m}$ are defined by induction, starting from $c_{1,1} = -1$, through the following formula

$$c_{l,m} = \left(\frac{2l}{m-1} - 4 \right) c_{l,m-1} \mathbf{1}_{l < m} + \frac{2}{m-1} c_{l-1,m-1} \mathbf{1}_{l > 2}, \quad l \in \{1, \dots, m\}, m \in \{2, \dots, M\}. \quad (5.19)$$

Furthermore, if the density η is such that $\eta_m^*(y) \geq 0$ for all $y \in \mathbb{R}$, and all $m \in \{1, \dots, M\}$, then there exists $C \in \mathbb{R}_+$ such that

$$\forall m \in \{1, \dots, M\}, \forall t \in [0, T] \quad \|\mathbb{E}[f(X_1(\sqrt{t}Y, \cdot))]\|_{m,L} \leq (1 + Ct)\|f\|_{m,L}. \quad (5.20)$$

Let us stress here two things that are crucial in (5.20): the same norm is used in both sides, and the sharp time dependence of the multiplicative constant $(1 + Ct)$. These properties are used in the proof of Proposition 5.12 to get (\overline{H}_2) .

Proof. We first consider $m = 1$ and $f \in C_{\text{pol}}^{M,L}(\mathbb{R}_+)$. From the symmetry of Y , we have the equality $\mathbb{E}[f(X_1(\sqrt{t}Y, x))] = \mathbb{E}[f(X_1(\sqrt{t}Y, x)) + f(X_1(-\sqrt{t}Y, x))]/2$ and using the notation $\psi_f^\pm(x, y) = f(x + \sigma\sqrt{t}y\sqrt{x} + \sigma^2ty^2/4) \pm f(x - \sigma\sqrt{t}y\sqrt{x} + \sigma^2ty^2/4)$ we can write,

$$\partial_x \mathbb{E}[f(X_1(\sqrt{t}Y, x))] = \frac{1}{2} \int_{-\infty}^{\infty} \partial_x \psi_f^+(x, y) \eta(y) dy.$$

One derivation and a little of algebra show that

$$\begin{aligned} \partial_x \psi_f^+(x, y) &= (1 + \frac{\sigma\sqrt{t}y}{2\sqrt{x}}) f'(x + \sigma\sqrt{t}y\sqrt{x} + \sigma^2ty^2/4) + (1 - \frac{\sigma\sqrt{t}y}{2\sqrt{x}}) f'(x - \sigma\sqrt{t}y\sqrt{x} + \sigma^2ty^2/4) \\ &= \frac{1}{\sigma\sqrt{t}\sqrt{x}} \left((\sigma^2ty/2 + \sigma\sqrt{t}\sqrt{x}) f'(x + \sigma\sqrt{t}y\sqrt{x} + \sigma^2ty^2/4) \right. \\ &\quad \left. - (\sigma^2ty/2 - \sigma\sqrt{t}\sqrt{x}) f'(x - \sigma\sqrt{t}y\sqrt{x} + \sigma^2ty^2/4) \right) \\ &= \frac{1}{\sigma\sqrt{t}\sqrt{x}} \left(\partial_y [f(x + \sigma\sqrt{t}y\sqrt{x} + \sigma^2ty^2/4)] - \partial_y [f(x - \sigma\sqrt{t}y\sqrt{x} + \sigma^2ty^2/4)] \right) \\ &= \frac{\partial_y \psi_f^-(x, y)}{\sigma\sqrt{t}\sqrt{x}}. \end{aligned}$$

Integrating by parts in the variable y , observing that the boundary term vanishes since $|\eta'(y)| =_{|y| \rightarrow \infty} o(|y|^{2L+1})$ and $f'(z) =_{z \rightarrow \infty} O(z^L)$, one has

$$\begin{aligned} \partial_x \mathbb{E}[f(X_1(\sqrt{t}Y, x))] &= -\frac{1}{2} \int_{-\infty}^{\infty} \frac{\psi_f^-(x, y) \eta'(y)}{\sigma\sqrt{t}\sqrt{x}} dy \\ &= -\int_{-\infty}^{\infty} \int_0^1 f'(x + (2u-1)\sigma\sqrt{t}y\sqrt{x} + \sigma^2ty^2/4) \eta'(y) y du dy \\ &= \int_{-\infty}^{\infty} \int_0^1 f'(x + (2u-1)\sigma\sqrt{t}y\sqrt{x} + \sigma^2ty^2/4) (-\eta'(y) y) du dy \end{aligned}$$

since $\partial_u f(x + (2u-1)\sigma\sqrt{t}y\sqrt{x} + \sigma^2ty^2/4) = 2\sigma\sqrt{t}y\sqrt{x} f'(x + (2u-1)\sigma\sqrt{t}y\sqrt{x} + \sigma^2ty^2/4)$. In order to simplify the notation, we define $w(u, x, y) := x + (2u-1)\sigma\sqrt{t}y\sqrt{x} + \sigma^2ty^2/4$ and we write explicitly the partial derivatives of w

$$\begin{cases} \partial_u w(u, x, y) = 2\sigma\sqrt{t}y\sqrt{x}, \\ \partial_x w(u, x, y) = 1 + \frac{(2u-1)\sigma\sqrt{t}y}{2\sqrt{x}}, \\ \partial_y w(u, x, y) = (2u-1)\sigma\sqrt{t}\sqrt{x} + \frac{\sigma^2ty}{2}, \end{cases} \quad (5.21)$$

and we define for $s : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$

$$I_{m,n}^{(l)}(s) = \int_{-\infty}^{\infty} \int_0^1 s(u, y) (u^2 - u)^{m-1} f^{(l)}(w(u, x, y)) \left(\sum_{j=1}^n c_{j,n} y^j \eta^{(j)}(y) \right) dudy, \quad (5.22)$$

so we can rewrite (5.18) as $\partial_x^m \mathbb{E}[f(X_1(\sqrt{t}Y, x))] = I_{m,m}^{(m)}(1)$ where the 1 in the argument has to be intended as the constant map identically equal to 1. So far, we have shown that formula (5.18) is true for $m = 1$, we take now $m \geq 2$ and we prove it by induction over m assuming that the result holds for $m - 1$. We differentiate Eq. (5.18) for $m - 1$ and use the second equality of (5.21) to get

$$\partial_x^m \mathbb{E}[f(X_1(\sqrt{t}Y, x))] = I_{m-1,m-1}^{(m)}(1) + I_{m-1,m-1}^{(m)} \left(\frac{(2u-1)\sigma\sqrt{t}y}{2\sqrt{x}} \right). \quad (5.23)$$

Then, from the third equality of (5.21), one has $\frac{(2u-1)\sigma\sqrt{t}y}{2\sqrt{x}} = \frac{(2u-1)}{\sigma\sqrt{t}\sqrt{x}} \partial_y w - (2u-1)^2$ and so

$$\begin{aligned} \partial_x^m \mathbb{E}[f(X_1(\sqrt{t}Y, x))] &= I_{m-1,m-1}^{(m)}(1 - (2u-1)^2) + I_{m-1,m-1}^{(m)} \left(\frac{(2u-1)}{\sigma\sqrt{t}\sqrt{x}} \partial_y w(u, x, y) \right) \\ &= -4I_{m-1,m-1}^{(m)}(u^2 - u) + I_{m-1,m-1}^{(m)} \left(\frac{(2u-1)}{\sigma\sqrt{t}\sqrt{x}} \partial_y w(u, x, y) \right) \\ &= -4I_{m,m-1}^{(m)}(1) + I_{m-1,m-1}^{(m)} \left(\frac{(2u-1)}{\sigma\sqrt{t}\sqrt{x}} \partial_y w(u, x, y) \right). \end{aligned}$$

We work on the term $I_{m-1,m-1}^{(m)} \left(\frac{(2u-1)}{\sigma\sqrt{t}\sqrt{x}} \partial_y w(u, x, y) \right)$. We use first an integration by parts in the variable y and subsequently one in the variable u . The boundary terms vanishes by using the hypothesis on η since $|f^{(m)}(w(u, x, y))| \leq \|f\|_{m,L}(1 + w(u, x, y)^L) = O(y^{2L})$ and to $|y| \rightarrow \infty$

the fact that the function $u^2 - u$ vanishes in 0 and 1. One gets

$$\begin{aligned} &\int_0^1 \int_{-\infty}^{\infty} \frac{(2u-1)(u^2-u)^{m-2}}{\sigma\sqrt{t}\sqrt{x}} f^{(m)}(w(u, x, y)) \partial_y w(u, x, y) \left(\sum_{j=1}^{m-1} c_{j,m-1} y^j \eta^{(j)}(y) \right) dy du \\ &= - \int_{-\infty}^{\infty} \int_0^1 \frac{(2u-1)(u^2-u)^{m-2}}{\sigma\sqrt{t}\sqrt{x}} f^{(m-1)}(w(u, x, y)) \left(\sum_{j=1}^{m-1} c_{j,m-1} (jy^{j-1} \eta^{(j)}(y) + y^j \eta^{(j+1)}(y)) \right) dudy \\ &= \int_{-\infty}^{\infty} \int_0^1 (u^2-u)^{m-1} f^{(m)}(w(u, x, y)) \left(\sum_{j=1}^{m-1} \frac{2}{m-1} c_{j,m-1} (jy^j \eta^{(j)}(y) + y^{j+1} \eta^{(j+1)}(y)) \right) dudy \\ &= \int_{-\infty}^{\infty} \int_0^1 (u^2-u)^{m-1} f^{(m)}(w(u, x, y)) \frac{2}{m-1} \left(\sum_{j=1}^{m-1} j c_{j,m-1} y^j \eta^{(j)}(y) + \sum_{j=2}^m c_{j-1,m-1} y^j \eta^{(j)}(y) \right) dudy. \end{aligned} \quad (5.24)$$

Rewriting the last equality for $\partial_x^m \mathbb{E}[f(X_1(\sqrt{t}Y, x))]$, one has

$$\begin{aligned} \partial_x^m \mathbb{E}[f(X_1(\sqrt{t}Y, x))] &= \int_{-\infty}^{\infty} \int_0^1 (u^2-u)^{m-1} f^{(m)}(w(u, x, y)) \left(-4 \sum_{j=1}^{m-1} c_{j,m-1} y^j \eta^{(j)}(y) \right) dudy + \\ &+ \int_{-\infty}^{\infty} \int_0^1 (u^2-u)^{m-1} f^{(m)}(w(u, x, y)) \frac{2}{m-1} \left(\sum_{j=1}^{m-1} j c_{j,m-1} y^j \eta^{(j)}(y) + \sum_{j=2}^m c_{j-1,m-1} y^j \eta^{(j)}(y) \right) dudy \end{aligned}$$

$$\begin{aligned}
&= \int_{-\infty}^{\infty} \int_0^1 (u^2 - u)^{m-1} f^{(m)}(w(u, x, y)) \left(\left(\frac{2}{m-1} - 4 \right) c_{1, m-1} y \eta^{(1)}(y) + \right. \\
&\quad \left. \sum_{j=2}^{m-1} \left(\left(\frac{2j}{m-1} - 4 \right) c_{j, m-1} + \frac{2}{m-1} c_{j-1, m-1} \right) y^j \eta^{(j)}(y) + \frac{2}{m-1} c_{m-1, m-1} y^m \eta^{(m)}(y) \right) dy du, \tag{5.25}
\end{aligned}$$

which proves the representation (5.18).

We are now able to prove the estimate using this representation. Defining $\eta_m^*(y) = (-1)^{m-1} \sum_{j=0}^m c_{j, m} y^j \eta^{(j)}(y)$, that is nonnegative for all y by hypothesis, one has

$$\begin{aligned}
|\partial_x^m \mathbb{E}[f(X_1(\sqrt{t}Y, x))]| &\leq \int_0^1 (u - u^2)^{m-1} \int_{-\infty}^{\infty} |f^{(m)}(w(u, x, y))| \eta_m^*(y) dy du \\
&\leq \|f\|_{m, L} \int_0^1 (u - u^2)^{m-1} \int_{-\infty}^{\infty} (1 + w(u, x, y)^L) \eta_m^*(y) dy du \\
&= \|f\|_{m, L} \underbrace{\int_0^1 (u - u^2)^{m-1} \int_{-\infty}^{\infty} \eta_m^*(y) dy du}_A \\
&\quad + \|f\|_{m, L} \underbrace{\int_0^1 (u - u^2)^{m-1} \int_{-\infty}^{\infty} w(u, x, y)^L \eta_m^*(y) dy du}_B.
\end{aligned}$$

The double integral A can be seen by means of representation (5.18) with $f(x) = \frac{x^m}{m!}$ ($f^{(m)} \equiv 1$) as

$$A = \partial_x^m \mathbb{E} \left[\frac{X_1(\sqrt{t}Y, x)^m}{m!} \right] = \frac{1}{m!} \partial_x^m \sum_{j=0}^m \binom{2m}{2j} x^{m-j} \left(\frac{\sigma\sqrt{t}}{2} \right)^j \mathbb{E}[Y^{2j}] = 1,$$

by using the symmetry of the density η . In the same way, B can be seen by means of the representation as

$$\begin{aligned}
B &= \partial_x^m \mathbb{E} \left[\frac{L!}{(L+m)!} X_1(\sqrt{t}Y, x)^{L+m} \right] \\
&= \partial_x^m \sum_{j=0}^{L+m} \binom{2(L+m)}{2j} \frac{L! x^{L+m-j}}{(L+m)!} \left(\frac{\sigma\sqrt{t}}{2} \right)^{2j} \mathbb{E}[Y^{2j}] \\
&= \sum_{j=0}^L \binom{2(L+m)}{2j} \frac{L!(L+m-j)!}{(L-j)!(L+m)!} x^{L-j} \left(\frac{\sigma\sqrt{t}}{2} \right)^{2j} \mathbb{E}[Y^{2j}] \\
&= x^L + t \sum_{j=1}^L \binom{2(L+m)}{2j} \frac{L!(L+m-j)!}{(L-j)!(L+m)!} x^{L-j} \left(\frac{\sigma}{2} \right)^{2j} t^{j-1} \mathbb{E}[Y^{2j}] \\
&\leq x^L + t(1+x^L)(1+\mathbb{E}[Y^{2L}]) \sum_{j=1}^L \binom{2(L+m)}{2j} \left(\frac{\sigma c_T}{2} \right)^{2j} \\
&\leq x^L + Ct(1+x^L)
\end{aligned}$$

where $c_T = \max(1, T)$ and $C = \frac{1}{2} \left((1 + \frac{\sigma c_T}{2})^{2(L+k)} + (1 - \frac{\sigma c_T}{2})^{2(L+k)} \right) (1 + \mathbb{E}[Y^{2L}])$. Putting parts *A* and *B* back together one has

$$\partial_x^m \mathbb{E}[f(X_1(\sqrt{t}Y, x))] \leq \|f\|_{m,L} (1 + x^L + Ct(1 + x^L)) = (1 + x^L)(1 + Ct) \|f\|_{m,L}, \quad (5.26)$$

and this proves the desired norm inequality. \square

Lemma 5.14. *Let $\eta(y) = \frac{1}{\sqrt{2\pi}} e^{-y^2/2}$ be the density of a standard normal variable. Then, we have for $m \geq 1$:*

$$\eta_m^*(y) := (-1)^{m-1} \sum_{j=1}^m c_{j,m} y^j \eta^{(j)}(y) = -c_{m,m} y^{2m} \eta(y), \quad (5.27)$$

so, in particular $\eta_m^*(y) \geq 0$ for all $y \in \mathbb{R}$.

Proof. For $m = 1$, (5.27) is clearly true since $\eta'(y) = -y\eta(y)$. We now take $m \geq 2$, $M \geq m$ and we suppose (5.27) true for $m - 1$: for all $f \in \mathcal{C}_{\text{pol}}^{M,L}(\mathbb{R}_+)$ and $x \in \mathbb{R}_+$, we have

$$\int_{-\infty}^{\infty} \int_0^1 (u - u^2)^{m-2} f^{(m-1)}(w(u, x, y)) (\eta_{m-1}^*(y) + c_{m-1,m-1} y^{2m-2} \eta(y)) \, dudy = 0.$$

Doing one differentiation step with respect to x like in the proof of Lemma 5.13 and using that $\eta'(y) = -y\eta(y)$, we obtain

$$\int_{-\infty}^{\infty} \int_0^1 (u - u^2)^{m-1} f^{(m)}(w(u, x, y)) (\eta_m^*(y) + c_{m,m} y^{2m} \eta(y)) \, dudy = 0.$$

By choosing $f_L(x) := \frac{L!}{(L+m)!} x^{L+m}$ for $L \in \mathbb{N}$, we get

$$\forall L \in \mathbb{N}, x \in \mathbb{R}_+, \int_{-\infty}^{\infty} \int_0^1 (u - u^2)^{m-1} w(u, x, y)^L (\eta_m^*(y) + c_{m,m} y^{2m} \eta(y)) \, dudy = 0.$$

We now take $x = 0$ so that $w(u, 0, y) = \frac{\sigma^2 t}{4} y^2$ and then

$$\forall L \in \mathbb{N}, \int_{-\infty}^{\infty} y^{2L} (\eta_m^*(y) + c_{m,m} y^{2m} \eta(y)) \, dy = 0.$$

We remark also that $\eta_m^*(y) = (\sum_{j=1}^m (-1)^{m+j-1} c_{j,m} y^j H_j(y)) \eta(y)$, where H_j is the j^{th} Hermite polynomial function (defined by $\eta^{(j)}(y) = H_j(y) \eta(y)$). Thus, $\eta_m^*(y) + c_{m,m} y^{2m} \eta(y) = P_{2m}(y) \eta(y)$ where P_{2m} is an even polynomial function of degree $2m$. We therefore obtain $\int_{-\infty}^{\infty} y^l P_{2m}(y) \eta(y) \, dy = 0$ for all $l \in \mathbb{N}$, which gives $P_{2m} = 0$ and thus the claim. \square

Remark 5.15. *Lemma 5.14 gives a remarkable formula of the monomial of order $2m$ $m \in \mathbb{N}^*$ in terms of the first m Hermite polynomials multiplied respectively by the first m monomials*

$$y^{2m} = \sum_{j=1}^m (-1)^{m+j} \frac{c_{j,m}}{c_{m,m}} y^j H_j(y). \quad (5.28)$$

The next result gives a kind of reciprocal result to Lemma 5.14. It explains why we consider a normal random variable for Y in Theorem 3.2, since we use Proposition 5.12 for any $M \in \mathbb{N}$.

Theorem 5.16. *Let Y be a symmetric random variable with a C^∞ probability density function η such that $\mathbb{E}[Y^2] = 1$, $\mathbb{E}[Y^4] = 3$ and $\eta_m^* \geq 0$ for all $m \geq 1$. Then, $Y \sim \mathcal{N}(0, 1)$.*

Proof. By Corollary B.2, there exists a positive Borel measure μ such that $\eta(x) = \int_0^\infty e^{-tx^2} \mu(dt)$. Since $\int_{\mathbb{R}} \eta = 1$, we get $\int_0^\infty \sqrt{\pi/t} \mu(dt) = 1$ and then $\eta(x) = \int_0^\infty \frac{e^{-tx^2}}{\sqrt{\pi/t}} \tilde{\mu}(dt)$ with $\tilde{\mu}(dt) = \sqrt{\pi/t} \mu(dt)$ being a probability measure on \mathbb{R}_+ . We have $\mathbb{E}[Y^2] = \int_0^\infty \int_{\mathbb{R}} x^2 \frac{e^{-tx^2}}{\sqrt{\pi/t}} dx \tilde{\mu}(dt) = \int_0^\infty \frac{1}{2t} \tilde{\mu}(dt)$ and $\mathbb{E}[Y^4] = \int_0^\infty 3 \left(\frac{1}{2t}\right)^2 \tilde{\mu}(dt)$. Therefore, we have

$$\int_0^\infty \frac{1}{2t} \tilde{\mu}(dt) = \int_0^\infty \left(\frac{1}{2t}\right)^2 \tilde{\mu}(dt) = 1.$$

The equality condition in the Cauchy-Schwarz inequality implies that $\tilde{\mu}(dt) = \delta_{1/2}(dt)$, i.e. Y is a standard normal variable. \square

5.3. Proof of Theorem 3.2. We prove the result for $\hat{P}^{2,n}$. By assumption, $f \in \mathcal{C}_{\text{pol}}^{18,L}(\mathbb{R}_+)$, for $L \geq 18$ sufficiently large. From (2.6), we have

$$\begin{aligned} P_T f - \hat{P}^{2,n} f &= \sum_{k=0}^{n-1} Q_1^{[n-(k+1)]} [P_{h_1} - Q_2^{[n]}] Q_1^{[k]} f \\ &\quad + \sum_{k=0}^{n-1} \sum_{k'=0}^{n-(k+2)} P_{(n-(k+k'+2))h_1} [P_{h_1} - Q_1] Q_1^{[k']} [P_{h_1} - Q_1] Q_1^{[k]}, \end{aligned}$$

with $h_l = T/n^l$. Using Proposition 5.9 three times and Proposition 5.3 twice, we get for $k \in \{0, \dots, n-1\}, k' \in \{0, \dots, n-(k+2)\}$:

$$\begin{aligned} \|P_{(n-(k+k'+2))h_1} [P_{h_1} - Q_1] Q_1^{[k']} [P_{h_1} - Q_1] Q_1^{[k]} f\|_{0,L+6} &\leq C \| [P_{h_1} - Q_1] Q_1^{[k']} [P_{h_1} - Q_1] Q_1^{[k]} f \|_{0,L+6} \\ &\leq Ch_1^3 \| Q_1^{[k']} [P_{h_1} - Q_1] Q_1^{[k]} f \|_{6,L+3} \\ &\leq Ch_1^3 \| [P_{h_1} - Q_1] Q_1^{[k]} f \|_{6,L+3} \\ &\leq Ch_1^6 \| Q_1^{[k]} f \|_{18,L} \leq Ch_1^6 \| f \|_{18,L}. \end{aligned}$$

For the other term, we write $P_{h_1} - Q_2^{[n]} = \sum_{k'=0}^{n-1} P_{(n-(k'+1))h_2} [P_{h_2} - Q_2] Q_2^{[k']}$ and get for $k, k' \in \{0, \dots, n-1\}$ by using Proposition 5.9, Proposition 5.3 and Lemma 5.1:

$$\begin{aligned} \|Q_1^{[n-(k+1)]} P_{(n-(k'+1))h_2} [P_{h_2} - Q_2] Q_2^{[k']} Q_1^{[k]} f\|_{0,L+6} &\leq C \| [P_{h_2} - Q_2] Q_2^{[k']} Q_1^{[k]} f \|_{0,L+6} \\ &\leq Ch_2^3 \| Q_2^{[k']} Q_1^{[k]} f \|_{6,L+3} \\ &\leq Ch_2^3 \| f \|_{6,L+3} \leq Ch_2^3 \| f \|_{18,L}. \end{aligned}$$

This gives

$$\|P_T f - \hat{P}^{2,n} f\|_{0,L+6} \leq C \|f\|_{18,L} n^2 (h_1^6 + h_2^3) \leq C \|f\|_{18,L} n^{-4},$$

and in particular $P_T f(x) - \hat{P}^{2,n} f(x) = O(n^{-4})$ for any $x \geq 0$.

We now consider $f \in \mathcal{C}^\infty$ with derivatives of polynomial growth. Therefore, for any $m \in \mathbb{N}$, it exists $L \geq m$ sufficiently large, such that $f \in \mathcal{C}_{\text{pol}}^{m,L}(\mathbb{R}_+)$. We can then apply [4, Theorem 3.10] to get that for some functions $\mathbf{m}, \ell : \mathbb{N}^* \rightarrow \mathbb{N}^*$, we have $\|P_T f - \hat{P}^{\nu,n} f\|_{0,L+\ell(\nu)} \leq C \|f\|_{\mathbf{m}(\nu),L} n^{-2\nu}$, which gives the claim.

6. SIMULATIONS RESULTS

In order to present some numerical test, we first explain how to implement the approximations $\hat{P}^{2,n}$ and $\hat{P}^{3,n}$ defined respectively by (2.7) and (2.8) (let us recall here that $\hat{P}^{1,n}$ is the approximation obtained on the regular time grid $\Pi^0 = \{kT/n, 0 \leq k \leq n\}$). We consider a general case of a scheme that can be written as a function of the starting point, the time step, the Brownian increment and an independent random variable, i.e.

$$Q_l f(x) = \mathbb{E}[\varphi(x, h_l, W_{h_l}, V)].$$

The second order scheme for the CIR (3.1) falls into this framework as well as the second order scheme for the Heston model (6.5) that we introduce below. As illustrated in [4] the approximation $\hat{P}^{2,n}$ is the simplest case for the implementation. It consists in the simulation of two starting schemes on the uniform time grid Π^0 and on the random grid : $\Pi^1 = \Pi^0 \cup \{\kappa T/n + k'T/n^2, 1 \leq k' \leq n-1\}$, where κ is an independent uniform random variable on $\{0, \dots, n-1\}$. We denote by $\hat{X}^{n,0}$ the scheme on Π^0

$$\begin{aligned} \hat{X}_0^{n,0} &= x, \\ \hat{X}_{(k+1)h_1}^{n,0} &= \varphi(\hat{X}_{kh_1}^{n,0}, h_1, W_{(k+1)h_1} - W_{kh_1}, V_k), \quad 0 \leq k \leq n-1, \end{aligned} \quad (6.1)$$

and by $\hat{X}^{n,1}$ the scheme on Π^1 :

$$\begin{aligned} \hat{X}_{kh_1}^{n,1} &= \hat{X}_{kh_1}^{n,0}, & 0 \leq k \leq \kappa, \\ \hat{X}_{\kappa h_1 + (k'+1)h_2}^{n,1} &= \varphi(\hat{X}_{\kappa h_1 + k'h_2}^{n,1}, h_2, W_{\kappa h_1 + (k'+1)h_2} - W_{\kappa h_1 + k'h_2}, V_{n+k'}), & 0 \leq k' \leq n-1, \\ \hat{X}_{(k+1)h_1}^{n,1} &= \varphi(\hat{X}_{kh_1}^{n,1}, h_1, W_{(k+1)h_1} - W_{kh_1}, V_k), & \kappa + 1 \leq k \leq n-1. \end{aligned}$$

Here, $(V_k)_{k \geq 0}$ is an i.i.d. sequence with the same law as V . Finally, we can give the following probabilistic representation

$$\begin{aligned} \hat{P}^{2,n} f &= Q_1^{[n]} f + n \mathbb{E}[Q_1^{[n-(\kappa+1)]} [Q_2^{[n]} - Q_1] Q_1^{[\kappa]} f] \\ &= \mathbb{E}[f(\hat{X}_T^{n,0})] + n \mathbb{E}[f(\hat{X}_T^{n,1}) - f(\hat{X}_T^{n,0})]. \end{aligned} \quad (6.2)$$

The approximation $\hat{P}^{3,n}$ is more involved. Let κ' be an independent uniform random variable on $\{0, \dots, n-1\}$. We define the scheme $\hat{X}^{n,2}$:

$$\begin{aligned} \hat{X}_{kh_1}^{n,2} &= \hat{X}_{kh_1}^{n,1}, \quad \hat{X}_{\kappa h_1 + k'h_2}^{n,2} = \hat{X}_{\kappa h_1 + k'h_2}^{n,1}, \quad 0 \leq k \leq \kappa, \quad 0 \leq k' \leq \kappa', \\ \hat{X}_{\kappa h_1 + \kappa' h_2 + (k''+1)h_3}^{n,2} &= \varphi(\hat{X}_{\kappa h_1 + \kappa' h_2 + k'' h_3}^{n,2}, h_3, W_{\kappa h_1 + \kappa' h_2 + (k''+1)h_3} - W_{\kappa h_1 + \kappa' h_2 + k'' h_3}, V_{2n+k''}), \\ & \quad 0 \leq k'' \leq n-1, \\ \hat{X}_{\kappa h_1 + (k'+1)h_2}^{n,2} &= \varphi(\hat{X}_{\kappa h_1 + k'h_2}^{n,2}, h_2, W_{\kappa h_1 + (k'+1)h_2} - W_{\kappa h_1 + k'h_2}, V_{n+k'}), \quad \kappa + 1 \leq k' \leq n-1, \\ \hat{X}_{(k+1)h_1}^{n,2} &= \varphi(\hat{X}_{kh_1}^{n,2}, h_1, W_{(k+1)h_1} - W_{kh_1}, V_k), \quad \kappa + 1 \leq k \leq n-1. \end{aligned}$$

This is the scheme obtained on the time grid $\Pi^1 \cup \{\kappa T/n + \kappa' T/n^2 + k'' T/n^3, 1 \leq k'' \leq n-1\}$. We have

$$\sum_{k=0}^{n-1} Q_1^{[n-(k+1)]} \left[\sum_{k'=0}^{n-1} Q_2^{[n-(k'+1)]} [Q_3^{[n]} - Q_2] Q_2^{[k']} \right] Q_1^{[k]} f = n^2 \mathbb{E}[f(\hat{X}_T^{n,2}) - f(\hat{X}_T^{n,1})].$$

We now explain how to calculate the second term in (2.8). Let (κ_1, κ_2) be an independent random variable uniformly distributed on the set $\{(k_1, k_2) : 0 \leq k_1 < k_2 < n\}$. We define:

$$\begin{aligned} \hat{X}_{kh_1}^{n,3} &= \hat{X}_{kh_1}^{n,0}, & 0 \leq k \leq \kappa_1, \\ \hat{X}_{\kappa_1 h_1 + (k'+1)h_2}^{n,3} &= \varphi(\hat{X}_{\kappa_1 h_1 + k' h_2}^{n,3}, h_2, W_{\kappa_1 h_1 + (k'+1)h_2} - W_{\kappa_1 h_1 + k' h_2}, V_{3n+k'}), & 0 \leq k' \leq n-1, \\ \hat{X}_{(k+1)h_1}^{n,3} &= \varphi(\hat{X}_{kh_1}^{n,3}, h_1, W_{(k+1)h_1} - W_{kh_1}, V_k), & \kappa_1 + 1 \leq k \leq n-1, \\ \\ \hat{X}_{kh_1}^{n,4} &= \hat{X}_{kh_1}^{n,0}, & 0 \leq k \leq \kappa_2, \\ \hat{X}_{\kappa_2 h_1 + (k'+1)h_2}^{n,4} &= \varphi(\hat{X}_{\kappa_2 h_1 + k' h_2}^{n,4}, h_2, W_{\kappa_2 h_1 + (k'+1)h_2} - W_{\kappa_2 h_1 + k' h_2}, V_{4n+k'}), & 0 \leq k' \leq n-1, \\ \hat{X}_{(k+1)h_1}^{n,4} &= \varphi(\hat{X}_{kh_1}^{n,4}, h_1, W_{(k+1)h_1} - W_{kh_1}, V_k), & \kappa_2 + 1 \leq k \leq n-1, \end{aligned}$$

and

$$\begin{aligned} \hat{X}_{kh_1}^{n,5} &= \hat{X}_{kh_1}^{n,3}, & 0 \leq k \leq \kappa_2, \\ \hat{X}_{\kappa_2 h_1 + (k'+1)h_2}^{n,5} &= \varphi(\hat{X}_{\kappa_2 h_1 + k' h_2}^{n,5}, h_2, W_{\kappa_2 h_1 + (k'+1)h_2} - W_{\kappa_2 h_1 + k' h_2}, V_{4n+k'}), & 0 \leq k' \leq n-1, \\ \hat{X}_{(k+1)h_1}^{n,5} &= \varphi(\hat{X}_{kh_1}^{n,5}, h_1, W_{(k+1)h_1} - W_{kh_1}, V_k), & \kappa_2 + 1 \leq k \leq n-1, \end{aligned}$$

These schemes correspond respectively to the time grids $\Pi^0 \cup \{\kappa_1 T/n + k'T/n^2, 1 \leq k' \leq n-1\}$, $\Pi^0 \cup \{\kappa_2 T/n + k'T/n^2, 1 \leq k' \leq n-1\}$ and $\Pi^0 \cup \{\kappa_1 T/n + k'T/n^2, 1 \leq k' \leq n-1\} \cup \{\kappa_2 T/n + k'T/n^2, 1 \leq k' \leq n-1\}$. We then get

$$\begin{aligned} \hat{P}^{3,n} f &= \mathbb{E}[f(\hat{X}_T^{n,0})] + n \mathbb{E}[f(\hat{X}_T^{n,1}) - f(\hat{X}_T^{n,0})] + n^2 \mathbb{E}[f(\hat{X}_T^{n,2}) - f(\hat{X}_T^{n,1})] \\ &\quad + \frac{n(n-1)}{2} \mathbb{E}[f(\hat{X}_T^{n,5}) - f(\hat{X}_T^{n,4}) - f(\hat{X}_T^{n,3}) + f(\hat{X}_T^{n,0})]. \end{aligned} \quad (6.3)$$

6.1. Simulations result for the CIR process. In this subsection, we want to illustrate the convergence of the approximations $\hat{P}^{2,n}$ and $\hat{P}^{3,n}$, which together with the use of the second order scheme (3.1) guarantee respectively approximations of order four and six by Theorem 3.2. In order to calculate these approximations, we use Monte-Carlo estimators of (6.2) and (6.3), using independent samples for each expectation. The number of samples (up to 10^{10}) is such that we can neglect the statistical error. In Figures 1, 2 and 3 we plot the convergence in function of the time step for different parameters choices, taking advantage of the closed formula for the Laplace transform of the CIR process, see e.g. [3, Proposition 1.2.4]. The three numerical experiments test different levels of the ratio $\sigma^2/4a$ in decreasing order. We observe that the slopes estimated on the log-log plots are close to 2, 4 and 6 respectively, so that they are in accordance with Theorem 3.2. Note however that Theorem 3.2 gives an asymptotic result for $n \rightarrow \infty$, while we are restricted here to rather small values of n since we are using a large number of samples to kill the statistical error. In all the cases shown, the approximations of higher order outperform the one built with the simple second order scheme (3.1). Talking about accuracies, the fourth order approximation for $n = 3$ shows an absolute relative error of about 0.17% in the tests in Figures 1, and 2 and 0.02% in the one in Figure 3; the sixth order approximation already for $n = 3$ exhibits a relative error of 0.002% in each case studied.




FIGURE 1. Parameters: $x = 0.2$, $a = 0.2$, $k = 0.5$, $\sigma = 0.65$, $f(z) = \exp(-10z)$ and $T = 1$. Graphic (A) shows the values of $\hat{P}^{1,n}f$, $\hat{P}^{2,n}f$, $\hat{P}^{3,n}f$ as a function of the time step $1/n$ and the exact value. Graphic (B) draws $\log(|\hat{P}^{i,n}f - P_T f|)$ in function of $\log(1/n)$: the regressed slopes are 1.79, 3.91 and 5.88 for the second, fourth and sixth order respectively.




FIGURE 2. Parameters: $x = 0.3$, $a = 0.4$, $k = 1$, $\sigma = 0.4$, $f(z) = \exp(-8z)$ and $T = 1$. Graphic (A) shows the values of $\hat{P}^{1,n}f$, $\hat{P}^{2,n}f$, $\hat{P}^{3,n}f$ as a function of the time step $1/n$ and the exact value. Graphic (B) draws $\log(|\hat{P}^{i,n}f - P_T f|)$ in function of $\log(1/n)$: the regressed slopes are 1.90, 3.93 and 5.77 for the second, fourth and sixth order respectively.

6.2. Simulations result for the Heston model. In this subsection, we want to test the second order scheme for the Heston model proposed by Alfonsi in [2] along with the approximations of order 4 and 6 obtained with combination of random grids. First, we recall the couple of stochastic differential equations describing this model

$$\begin{cases} dS_t^{(x,s)} = rS_t^{(x,s)}dt + \sqrt{X_t}S_t^{(x,s)}(\rho dW_t + \sqrt{1-\rho^2}dZ_t), & S_0^{(x,s)} = s, \\ dX_t^x = (a - kX_t^x)dt + \sigma\sqrt{X_t^x}dW_t, & X_0^x = x, \end{cases} \quad (6.4)$$




FIGURE 3. Parameters: $x = 10$, $a = 10$, $k = 1$, $\sigma = 0.23$, $f(z) = \exp(-z)$ and $T = 1$. Graphic (A) shows the values of $\hat{P}^{1,n}f$, $\hat{P}^{2,n}f$, $\hat{P}^{3,n}f$ as a function of the time step $1/n$ and the exact value. Graphic (B) draws $\log(|\hat{P}^{i,n}f - P_T f|)$ in function of $\log(1/n)$: the regressed slopes are 1.96, 4.00 and 6.02 for the second, fourth and sixth order respectively.

where W and Z are two independent Brownian motions. We define the two following random variable

$$S_1((x, s), h, Z_h) = \left(x, s \exp\left(\sqrt{x(1-\rho^2)}Z_h\right) \right)$$

$$S_2((x, s), h, W_h) = \left(\varphi(x, h, W_h), \right. \\ \left. s \exp\left(\left(r - \frac{\rho}{\sigma}a\right)h + \left(\frac{\rho}{\sigma}k - \frac{1}{2}\right)\frac{x + \varphi(x, h, W_h)}{2}h + \frac{\rho}{\sigma}(\varphi(x, h, W_h) - x) \right) \right)$$

where φ is defined by (3.2) and corresponds to the second order scheme for the CIR process. We define as in [2] the second order scheme for (6.4) as follows

$$\Phi((x, s), h, (W_h, Z_h), B) = \begin{cases} S_2(S_1((x, s), h, Z_h), h, W_h), & \text{if } B = 1, \\ S_1(S_2((x, s), h, W_h), h, Z_h), & \text{if } B = 0. \end{cases} \quad (6.5)$$

where B is an independent Bernoulli random variable of parameter $1/2$.

To test the order of the approximations $\hat{P}^{2,n}$ and $\hat{P}^{3,n}$ boosting the second order scheme (6.5), we have calculated European put prices taking advantage of the existence of a semi closed formula for this option, see [8]. In Figure 4 we draw the convergence in function of the time step. Again, we noticed that the slopes obtained on the log-log plot are in line with the expected order of convergence. More importantly, we see that the correction terms of the approximations $\hat{P}^{2,n}$ and $\hat{P}^{3,n}$ really improves the precision. They respectively give relative errors of a 0.035% and 0.0023%, already for $n = 3$.

6.3. Optimized implementation of $\hat{P}^{2,n}$. The approximations $\hat{P}^{2,n}$ and $\hat{P}^{3,n}$ defined respectively by (6.2) and (6.3) involve respectively two and four expectations. The larger is ν the more expectations are involved in $\hat{P}^{\nu,n}$. Thus, for simplicity, independent samples were used by Alfonsi and Bally [4] to compute each term. However, it may be interesting to




FIGURE 4. Test function: $f(x, s) = (K - s)^+$. Parameters: $S_0 = 100$, $r = 0$, $x = 0.25$, $a = 0.25$, $k = 1$, $\sigma = 0.65$, $\rho = -0.3$, $T = 1$, $K = 100$. Graphic (A) shows the values of $\hat{P}^{1,n}f$, $\hat{P}^{2,n}f$, $\hat{P}^{3,n}f$ as a function of the time step $1/n$ and the exact value. Graphic (B) draws $\log(|\hat{P}^{i,n}f - P_T f|)$ in function of $\log(1/n)$: the regressed slopes are 1.34, 4.00 and 6.02 for the second, fourth and sixth order respectively.

reuse some samples in order to spare computation time. This is what we investigate in this subsection.

Namely, Equation (6.2) leads naturally to the two following estimators of $P^{2,n}f$:

$$\Theta_i(M_1, M_2, n) = \frac{1}{M_1} \sum_{j=1}^{M_1} f((\hat{X}_T^{n,0})^{(j)}) + \frac{1}{M_2} \sum_{i=M_1+1}^{M_1+M_2} n \left(f((\hat{X}_T^{n,1})^{(i)}) - f((\hat{X}_T^{n,0})^{(i)}) \right), \quad (6.6)$$

$$\Theta_d(M_1, M_2, n) = \frac{1}{M_1} \sum_{j=1}^{M_1} f((\hat{X}_T^{n,0})^{(j)}) + \frac{1}{M_2} \sum_{i=1}^{M_2} n \left(f((\hat{X}_T^{n,1})^{(i)}) - f((\hat{X}_T^{n,0})^{(i)}) \right). \quad (6.7)$$

The first one takes independent samples and we call this estimator Θ_i . This approach is the one used in [4]. In the second case, we reuse the first $M_1 \wedge M_2$ simulations of $f((\hat{X}_T^{n,0})^{(i)})$ in both sums. We call this estimator Θ_d . In terms of variance, we have

$$\text{Var}(\Theta_i(M_1, M_2, n)) = \frac{\text{Var}(f(\hat{X}_T^{n,0}))}{M_1} + \frac{\text{Var}(n(f(\hat{X}_T^{n,1}) - f(\hat{X}_T^{n,0})))}{M_2}, \quad (6.8)$$

$$\text{Var}(\Theta_d(M_1, M_2, n)) = \frac{\text{Var}(f(\hat{X}_T^{n,0}))}{M_1} + 2 \frac{\text{Cov}(f(\hat{X}_T^{n,0}), n(f(\hat{X}_T^{n,1}) - f(\hat{X}_T^{n,0})))}{M_1 \vee M_2} + \frac{\text{Var}(n(f(\hat{X}_T^{n,1}) - f(\hat{X}_T^{n,0})))}{M_2}. \quad (6.9)$$

Let us define t_1 as the time to generate one sample $f(\hat{X}_T^{n,0})$ and t_2 as the one needed for one sample of the correction $n(f(\hat{X}_T^{n,1}) - f(\hat{X}_T^{n,0}))$. The computation time needed to compute Θ_i is given by $g_i(M_1, M_2) = M_1 t_1 + M_2 t_2$, while the one needed to compute Θ_d is $g_d(M_1, M_2) = \mathbf{1}_{M_1 \geq M_2} [(M_1 - M_2)t_1 + M_2 t_2] + \mathbf{1}_{M_1 < M_2} M_2 t_2$. We note $\zeta = \frac{t_2}{t_1}$. From the definition of schemes $\hat{X}^{n,0}$ and $\hat{X}^{n,1}$ in (6.1), we observe that $2 \leq \zeta \leq 3$ and that $\zeta \approx 2.5$ in

average since these schemes are equal up to κh_1 . The advantage of Θ_d is not necessarily in reducing the variance, but in decreasing the number of simulations needed, making it more efficient from a computational time point of view.

We want to find the optimal numbers of simulations M_1 and M_2 for our estimators in order to minimize the execution time for a given variance ε^2 . Let us define $\sigma_2^2(n) = \text{Var}(f(\hat{X}_T^{n,0}))$, $\sigma_4^2(n) = \text{Var}(n(f(\hat{X}_T^{n,1}) - f(\hat{X}_T^{n,0})))$, $\Gamma(n) = \text{Cov}(f(\hat{X}_T^{n,0}), n(f(\hat{X}_T^{n,1}) - f(\hat{X}_T^{n,0})))$. For Θ_i , the minimization of g_i given that $\sigma_2^2(n)/M_1 + \sigma_4^2(n)/M_2 = \varepsilon^2$ leads to $M_1 = \sqrt{\zeta} \frac{\sigma_2(n)}{\sigma_4(n)} M_2$ and then to:

$$M_{1,i} = \left\lceil \frac{1}{\varepsilon^2} \left(\sigma_2^2(n) + \sqrt{\zeta} \sigma_2(n) \sigma_4(n) \right) \right\rceil, \quad M_{2,i} = \left\lceil \frac{1}{\varepsilon^2} \left(\sigma_4^2(n) + \frac{\sigma_2(n) \sigma_4(n)}{\sqrt{\zeta}} \right) \right\rceil. \quad (6.10)$$

To minimize the execution time g_d , one has first to decide whether we take $M_1 \geq M_2$ or $M_1 < M_2$. From (6.9), this amounts to compare $\frac{\sigma_2^2(n)+2\Gamma(n)}{m+\tilde{m}\zeta}$ with $\frac{\sigma_4^2(n)+2\Gamma(n)}{m+\tilde{m}}$ where $m = M_1 \wedge M_2$ and $\tilde{m} \geq 0$ (\tilde{m} simulations of the correction term takes the same time as $\zeta \tilde{m}$ simulations of $f(\hat{X}_T^{n,0})$). Taking the derivative at $\tilde{m} = 0$, we get that $M_1 \geq M_2$ if $\zeta \frac{\sigma_2^2(n)+2\Gamma(n)}{\sigma_4^2(n)+2\Gamma(n)} \geq 1$, and $M_1 < M_2$ otherwise. When $M_1 \geq M_2$, the minimisation of g_d given $\text{Var}(\Theta_d(M_1, M_2, n)) = \varepsilon^2$ leads to

$$\begin{cases} M_{1,d} &= \left\lceil \frac{1}{\varepsilon} \left(\sigma_2^2(n) + 2\Gamma(n) + \sqrt{(\sigma_2^2(n) + 2\Gamma(n)) \sigma_4^2(n) (\zeta - 1)} \right) \right\rceil, \\ M_{2,d} &= \left\lceil \frac{1}{\varepsilon} \left(\sigma_4^2(n) + \sqrt{\frac{(\sigma_2^2(n)+2\Gamma(n)) \sigma_4^2(n)}{\zeta-1}} \right) \right\rceil. \end{cases} \quad (6.11)$$

We have similar formulas when $M_1 < M_2$. In all our numerical experiments below, we are in the case where $\zeta \frac{\sigma_2^2(n)+2\Gamma(n)}{\sigma_4^2(n)+2\Gamma(n)} \geq 1$ and thus taking $M_1 \geq M_2$ is optimal.

Now, we show the performance of the two estimators (6.6) and (6.7). To do this, we calculate the empirical variances $\sigma_2^2(n)$, $\sigma_4^2(n)$ and the empirical covariance $\Gamma(n)$ on a small sampling, fix a desired precision $\varepsilon = 1.96 \sqrt{\text{Var}(\Theta(M_1, M_2, n))}$ for both the estimators, so that all the terms have roughly the same statistical error with a 95% confidence interval half-width equal to ε . We show two tables in which we set the precision ε to 10^{-3} . In Table 1, we have $\sigma_2^2(n) \gg \sigma_4^2(n)$, while in Table 2, $\sigma_2^2(n)$ is still larger than $\sigma_4^2(n)$, but of the same order of magnitude.

	$n = 2$	$n = 3$	$n = 4$	$n = 5$
Θ_i	63.04	96.15	131.84	165.80
Θ_d	51.61	87.24	122.76	152.32

TABLE 1. Computation time (in seconds) needed by the Estimators Θ_i and Θ_d for a precision $\varepsilon = 10^{-3}$. Test function: $f(x, s) = (K - s)^+$. Parameters: $S_0 = 100$, $r = 0$, $x = 0.4$, $a = 0.4$, $k = 1$, $\sigma = 0.2$, $\rho = -0.3$, $T = 1$, $K = 100$.

We observe that we do not have a great gain in using Θ_i when $\sigma_2^2(n) \gg \sigma_4^2(n)$ (Table 1), while we save up to 30% of execution time, using Θ_d instead of Θ_i , when $\sigma_2^2(n)$ is of the same order of magnitude $\sigma_4^2(n)$ (Table 2). Heuristically, this can be understood as follows: when $\sigma_2^2(n)$ is of the same magnitude as $\sigma_4^2(n)$, so are $M_{1,i}$ and $M_{2,i}$, which gives an important gain

	$n = 2$	$n = 3$	$n = 4$	$n = 5$
Θ_i	59.50	102.13	148.45	193.41
Θ_d	37.59	70.43	100.14	136.16

TABLE 2. Computation time (in seconds) needed by the Estimators Θ_i and Θ_d for a precision $\varepsilon = 10^{-3}$. Test function: $f(x, s) = (K - s)^+$. Parameters: $S_0 = 100$, $r = 0$, $x = 0.1$, $a = 0.1$, $k = 1$, $\sigma = 0.63$, $\rho = -0.3$, $T = 1$, $K = 100$.

in reusing the simulation of the correction term. In any case, Θ_d turns out to be faster for each choice of parameters, and therefore we recommend it at the expense of Θ_i .

6.4. Comparison between the second and the fourth order approximation. Subsections 6.1 and 6.2 have confirmed numerically the theoretical results obtained in this paper. However, they do not compare directly the computation time required by the different methods. We now present numerical tests that allow us to prove the real advantage of using the fourth order approximation $\hat{P}^{2,n}$ instead of the simple second order scheme. Namely, we compare the squared L^2 distance of the estimator Θ_d from the true value with the same distance between the estimator of \hat{P}^{1,n^2} with the true value. We plot these quantities in function of the computation time needed. Note that $\hat{P}^{2,n}$ and \hat{P}^{1,n^2} converges at a rate of $O(n^{-4})$ so that their bias have the same order of magnitude.




FIGURE 5. L^2 -square error in function of the execution time in seconds. Test function: $f(x, s) = (K - s)^+$. Parameters in graphic (A) : $S_0 = 100$, $r = 0$, $x = 0.4$, $a = 0.4$, $k = 1$, $\sigma = 0.1$, $\rho = -0.3$, $T = 1$, $K = 100$.

Parameters in graphic (B) : $S_0 = 100$, $r = 0$, $x = 0.1$, $a = 0.1$, $k = 1$, $\sigma = 0.63$, $\rho = -0.3$, $T = 1$, $K = 100$.

Figure 5 shows the results for the calculation of the price of a European put option in the Heston model with two different sets of parameters. In this numerical experience we set a precision ε equal to 10^{-3} . The empirical evidences show that the fourth order estimator Θ_d is the best choice, especially when the ratio $\sigma^2/(4a) \ll 1$ (Figure 5 (A)) where the performance of the fourth order estimator is unparalleled. For example, $\hat{P}^{2,3}$ is twice more accurate and more than twice faster than $\hat{P}^{1,9}$. Even in Figure 5 (B), where the ratio $\sigma^2/(4a)$ is larger, the

fourth order estimator Θ_d is more precise than the second order estimator and is faster from $n = 3$ onward. These experiments illustrate the outperformance of the boosted estimator $\hat{P}^{2,n}$ with respect to $\hat{P}^{1,n}$.

APPENDIX A. PROOFS OF SECTION 4

Proof of Lemma 4.1. (1) Let $f \in \mathfrak{P}_L(\mathbb{R}_+)$. We have $X_0(t, x)^j = \sum_{i=0}^j \binom{j}{i} ((a - \sigma^2/4)\psi_k(t))^{j-i} e^{-kti} x^i$ and thus

$$f(X_0(t, \cdot)) = \sum_{j=0}^L a_j \sum_{i=0}^j \binom{j}{i} ((a - \sigma^2/4)\psi_k(t))^{j-i} e^{-kti} x^i.$$

Therefore, $f(X_0(t, \cdot)) \in \mathfrak{P}_L(\mathbb{R}_+)$ and we have

$$\|f(X_0(t, \cdot))\| \leq \sum_{j=0}^L |a_j| \sum_{i=0}^j \binom{j}{i} (|a - \sigma^2/4|\psi_k(t))^{j-i} e^{-kti} = \sum_{j=0}^L |a_j| \tilde{X}_0(t)^j,$$

with $\tilde{X}_0(t) = e^{-kt} + |a - \sigma^2/4|\psi_k(t)$. For $k \geq 0$, we have $0 \leq \psi_k(t) \leq t$ and thus $\tilde{X}_0(t) \leq (1 + |a - \sigma^2/4|t)$. For $k < 0$, we have $\tilde{X}_0(t) = e^{-kt}(1 + |a - \sigma^2/4|\psi_{-k}(t)) \leq e^{-kt}(1 + |a - \sigma^2/4|t)$. Since $(1 + |a - \sigma^2/4|t)^L \leq 1 + t \sum_{i=1}^j \binom{j}{i} |a - \sigma^2/4|^i (1 \vee T)^i \leq 1 + t(1 + |a - \sigma^2/4|(1 \vee T))^L$, we get $\tilde{X}_0(t)^j \leq (1 \vee e^{-kLt})[1 + t(1 + |a - \sigma^2/4|(1 \vee T))^L]$ for $j \in \{0, \dots, L\}$ and then

$$\|f(X_0(t, \cdot))\| \leq (1 \vee e^{-kLt})(1 + (1 + |a - \sigma^2/4|(1 \vee T))^L t) \|f\|,$$

which gives the claim with $C_{X_0} = 1 + |a - \sigma^2/4|(1 \vee T)$.

(2) Since Y is a symmetric random variable, we have

$$\begin{aligned} \mathbb{E}[f(X_1(\sqrt{t}Y, x))] &= \sum_{j=0}^L a_j \mathbb{E}[X_1(\sqrt{t}Y, x)^j] = \sum_{j=0}^L a_j \sum_{i=0}^{2j} \binom{2j}{i} \left(\frac{\sigma\sqrt{t}}{2}\right)^{2j-i} \mathbb{E}[Y^{2j-i}] x^{i/2} \\ &= \sum_{j=0}^L a_j \sum_{i=0}^j \binom{2j}{2i} \left(\frac{\sigma^2 t}{4}\right)^{j-i} \mathbb{E}[Y^{2(j-i)}] x^i \\ &= \sum_{j=0}^L a_j x^j + t \sum_{j=0}^L a_j \sum_{i=0}^{j-1} \binom{2j}{2i} \left(\frac{\sigma^2}{4}\right)^{j-i} t^{j-i-1} \mathbb{E}[Y^{2(j-i)}] x^i. \end{aligned}$$

This proves that $\mathbb{E}[f(X_1(\sqrt{t}Y, \cdot))] \in \mathfrak{P}_L(\mathbb{R}_+)$. We note that $\mathbb{E}[Y^{2j}] \geq 1$ by Hölder inequality since $\mathbb{E}[Y^2] = 1$, and thus $\mathbb{E}[Y^{2j}] \leq \mathbb{E}[Y^{2L}]$ for $j \in \{0, \dots, L\}$. We get

$$\begin{aligned} \|f(X_1(\sqrt{t}Y, \cdot))\| &\leq \|f\| + t \mathbb{E}[Y^{2L}] \sum_{j=0}^L |a_j| \sum_{i=0}^{j-1} \binom{2j}{2i} \left(\frac{\sigma^2}{4}\right)^{j-i} (1 \vee T)^{j-i} \\ &\leq \|f\| \left(1 + t \mathbb{E}[Y^{2L}] \left(1 + \frac{\sigma}{2} \sqrt{1 \vee T}\right)^{2L}\right), \end{aligned}$$

since $\sum_{i=0}^{j-1} \binom{2j}{2i} \left(\frac{\sigma^2}{4}\right)^{j-i} (1 \vee T)^{j-i} \leq \left(1 + \frac{\sigma}{2} \sqrt{1 \vee T}\right)^{2j} \leq \left(1 + \frac{\sigma}{2} \sqrt{1 \vee T}\right)^{2L}$. This gives the claim with $C_{X_1} = \left(1 + \frac{\sigma}{2} \sqrt{1 \vee T}\right)^2$. \square

Proof of Lemma 4.2. We have $\tilde{u}_0(t, x) = 1$, and in the case $m = 1$, we have $\tilde{u}_1(t, x) = x + \int_0^t (a - k\tilde{u}_1(s, x))ds$ that has the solution:

$$\tilde{u}_1(t, x) = xe^{-kt} + a\psi_k(t)$$

where $\psi_k(t) = \frac{1-e^{-kt}}{k}$ if $k \neq 0$ and $\psi_k(t) = t$ otherwise. This gives the claim for $m = 1$ with $\tilde{u}_{0,1} = a\psi_k(t)$ and $\tilde{u}_{1,1} = e^{-kt}$. We then prove the result by induction and consider $m \geq 2$. Using Itô formula and taking the expected value, one has $\partial_t \tilde{u}_m(t, x) = (am + \sigma^2 m(m-1)/2)\tilde{u}_{m-1}(t, x) - km\tilde{u}_m(t, x)$. Hence, we have

$$\tilde{u}_m(t, x) = (e^{-kt})^m \left(x^m + \int_0^t (am + \sigma^2 m(m-1)/2)(e^{ks})^m \tilde{u}_{m-1}(s, x) ds \right),$$

and we get the following induction relations that give us the representation (4.3)

$$\begin{cases} \tilde{u}_{j,m}(t) = (e^{-kt})^m \int_0^t (am + \sigma^2 m(m-1)/2)(e^{ks})^m \tilde{u}_{j,m-1}(s) ds, & 0 \leq j \leq m-1, \\ \tilde{u}_{m,m}(t) = (e^{-kt})^m. \end{cases}$$

Let $f \in \mathfrak{P}_L(\mathbb{R}_+)$. We clearly get from the preceding result that $\mathbb{E}[f(X_t)] \in \mathfrak{P}_L(\mathbb{R}_+)$ and

$$\|\mathbb{E}[f(X_t)]\| \leq \sum_{m=0}^L |a_m| \sum_{j=0}^m |\tilde{u}_{j,m}(t)| \leq C_{\text{cir}}(L, T) \|f\|. \quad \square$$

APPENDIX B. ASSUMPTION (\overline{H}_1) FOR SYMMETRIC RANDOM VARIABLES

Theorem B.1. *Let $\eta : \mathbb{R} \rightarrow \mathbb{R}_+$ be a C^∞ even function. Then, $\eta_m^* \geq 0$ for all $m \in \mathbb{N}^*$ if and only if $\eta(\sqrt{\cdot})$ is the Laplace transform of a finite positive Borel measure μ on $[0, \infty)$, i.e. $\eta(\sqrt{x}) = \int_0^\infty e^{-tx} \mu(dt)$ for all $x \in \mathbb{R}_+$.*

Proof. We start to prove that $\eta_m^* \geq 0$ for all $m \in \mathbb{N}^*$ implies $\eta(\sqrt{x}) = \int_0^\infty e^{-tx} \mu(dt)$ for all $x \in \mathbb{R}$. To prove this, we use Bernstein's Theorem for completely monotone functions (see e.g. [15, Theorem 12a p. 160]) and show that for all $m \in \mathbb{N}$ and $x \in \mathbb{R}_+$, $(-1)^m \partial_x^m [\eta(\sqrt{x})] \geq 0$. To do so, we prove by induction on m the representation

$$\partial_x^m [\eta(\sqrt{x})] = -\frac{(m-1)!}{2^{2m-1}} x^{-m} \sum_{j=1}^m c_{j,m} x^{\frac{j}{2}} \eta^{(j)}(\sqrt{x}) = (-1)^m \frac{(m-1)!}{2^{2m-1}} x^{-m} \eta_m^*(\sqrt{x}).$$

For $m = 1$, we have $\eta_1^*(\sqrt{x}) = c_{1,1} \sqrt{x} \eta'(\sqrt{x})$ and the representation holds from $\partial_x [\eta(\sqrt{x})] = \frac{1}{2\sqrt{x}} \eta'(\sqrt{x}) = -\frac{1}{2x} \eta_1^*(\sqrt{x})$ using that $c_{1,1} = -1$. Now, let $m \geq 2$ and suppose the representation is true for $m-1$, so

$$\partial_x^m [\eta(\sqrt{x})] = \partial_x (\partial_x^{m-1} [\eta(\sqrt{x})]) = \partial_x \left(-\frac{(m-2)!}{2^{2m-3}} x^{-(m-1)} \sum_{j=1}^{m-1} c_{j,m-1} x^{\frac{j}{2}} \eta^{(j)}(\sqrt{x}) \right).$$

Differentiating and using that $\partial_x(x^{\frac{j}{2}}\eta^{(j)}(\sqrt{x})) = \frac{1}{2x}(jx^{\frac{j}{2}}\eta^{(j)}(\sqrt{x}) + x^{\frac{j+1}{2}}\eta^{(j+1)}(\sqrt{x}))$, we get

$$\begin{aligned}
\partial_x^m[\eta(\sqrt{x})] &= -\frac{(m-2)!}{2^{2m-3}} \left(-\frac{m-1}{x^m} \sum_{j=1}^{m-1} c_{j,m-1} x^{\frac{j}{2}} \eta^{(j)}(\sqrt{x}) \right. \\
&\quad \left. + \frac{1}{2x^m} \sum_{j=1}^{m-1} c_{j,m-1} \left(jx^{\frac{j}{2}} \eta^{(j)}(\sqrt{x}) + x^{\frac{j+1}{2}} \eta^{(j+1)}(\sqrt{x}) \right) \right) \\
&= -\frac{(m-2)!}{2^{2m-3}} x^{-m} \left(\left(\frac{1}{2} - m - 1 \right) c_{1,m-1} x^{\frac{1}{2}} \eta^{(1)}(\sqrt{x}) \right. \\
&\quad \left. + \sum_{j=1}^{m-1} \left(\left(\frac{j}{2} - m + 1 \right) c_{j,m-1} + \frac{1}{2} c_{j-1,m-1} \right) x^{\frac{j}{2}} \eta^{(j)}(\sqrt{x}) \right. \\
&\quad \left. + \frac{1}{2} c_{m-1,m-1} x^{\frac{m}{2}} \eta^{(m)}(\sqrt{x}) \right) \\
&= -\frac{(m-1)!}{2^{2m-1}} x^{-m} \left(\left(\frac{2}{m-1} - 4 \right) c_{1,m-1} x^{\frac{1}{2}} \eta^{(1)}(\sqrt{x}) \right. \\
&\quad \left. + \sum_{j=1}^{m-1} \left(\left(\frac{2j}{m-1} - 4 \right) c_{j,m-1} + \frac{2}{m-1} c_{j-1,m-1} \right) x^{\frac{j}{2}} \eta^{(j)}(\sqrt{x}) \right. \\
&\quad \left. + \frac{2}{m-1} c_{m-1,m-1} x^{\frac{m}{2}} \eta^{(m)}(\sqrt{x}) \right)
\end{aligned}$$

and we conclude using the recursion formula (5.19) for $c_{j,m}$.

We now assume that $\eta(\sqrt{x}) = \int_0^\infty e^{-tx} \mu(dt)$ and show that $\eta_m^* \geq 0$ for all $m \geq 1$. We define $\eta_g(x) = e^{-\frac{x^2}{2}}$ and consider for all $t > 0$ the function $\eta_t(x) = e^{-tx^2}$. We remark that for all $t > 0$, $\eta_t(x) = \eta_g(h_t(x))$ with $h_t(x) = \sqrt{2t}x$ and so we can write by Lemma 5.14

$$(\eta_t)_m^*(x) = (-1)^{m-1} \sum_{j=1}^m c_{j,m} x^j \eta_t^{(j)}(x) = (-1)^{m-1} \sum_{j=1}^m c_{j,m} (\sqrt{2t}x)^j \eta_g^{(j)}(\sqrt{2t}x) = (\eta_g)_m^*(\sqrt{2t}x).$$

Therefore, $(\eta_t)_m^*(x) \geq 0$ for all $t > 0$ and $x \in \mathbb{R}$. We now consider an even function $\eta : \mathbb{R} \rightarrow \mathbb{R}_+$ such that $\eta(\sqrt{x}) = \int_0^\infty e^{-tx} \mu(dt)$ for some Borel measure μ on $[0, \infty)$. We then have for all $x \in \mathbb{R}$, $\eta(x) = \int_0^\infty e^{-tx^2} \mu(dt) = \int_0^\infty \eta_t(x) \mu(dt)$ and thus $\eta^{(j)}(x) = \int_0^\infty \eta_t^{(j)}(x) \mu(dt)$. This gives, for all $m \in \mathbb{N}^*$,

$$\begin{aligned}
\eta_m^*(x) &= (-1)^{m-1} \sum_{j=1}^m c_{j,m} x^j \eta^{(j)}(x) = \int_0^\infty (-1)^{m-1} \sum_{j=1}^m c_{j,m} x^j \eta_t^{(j)}(x) \mu(dt) \\
&= \int_0^\infty (\eta_t)_m^*(x) \mu(dt) \geq 0
\end{aligned}$$

where the last integral is positive for all $x \in \mathbb{R}$ because is an integral of a positive function against a positive measure. \square

Corollary B.2. *All the densities that satisfy the hypothesis of the representation Lemma 5.13 for all $m \in \mathbb{N}^*$ are such that $\eta(\sqrt{\cdot})$ is the Laplace transform of a finite positive Borel measure μ over $[0, \infty)$.*

REFERENCES

- [1] A. Alfonsi. On the discretization schemes for the CIR (and Bessel squared) processes. *Monte Carlo Methods Appl.*, 11(4):355–384, 2005.
- [2] A. Alfonsi. High order discretization schemes for the CIR process: application to affine term structure and Heston models. *Math. Comp.*, 79(269):209–237, 2010.
- [3] A. Alfonsi. *Affine diffusions and related processes: simulation, theory and applications*, volume 6 of *Bocconi & Springer Series*. Springer, Cham; Bocconi University Press, Milan, 2015.
- [4] A. Alfonsi and V. Bally. A generic construction for high order approximation schemes of semigroups using random grids. *Numer. Math.*, 148(4):743–793, 2021.
- [5] M. Bossy and A. Diop. An efficient discretisation scheme for one dimensional SDEs with a diffusion coefficient function of the form $|x|^a$, a in $[1/2, 1)$. (RR-5396):44, 2007. Version 2.
- [6] M. Briani, L. Caramellino, and G. Terenzi. Convergence rate of Markov chains and hybrid numerical schemes to jump-diffusion with application to the Bates model. *SIAM J. Numer. Anal.*, 59(1):477–502, 2021.
- [7] J. C. Cox, J. E. Ingersoll, Jr., and S. A. Ross. A theory of the term structure of interest rates. *Econometrica*, 53(2):385–407, 1985.
- [8] S. L. Heston. A closed-form solution for options with stochastic volatility with applications to bond and currency options. *Rev. Financ. Stud.*, 6(2):327–343, 1993.
- [9] S. Kusuoka. Approximation of expectation of diffusion process and mathematical finance. In *Taniguchi Conference on Mathematics Nara '98*, volume 31 of *Adv. Stud. Pure Math.*, pages 147–165. Math. Soc. Japan, Tokyo, 2001.
- [10] M. Ninomiya and S. Ninomiya. A new higher-order weak approximation scheme for stochastic differential equations and the Runge-Kutta method. *Finance Stoch.*, 13(3):415–443, 2009.
- [11] S. Ninomiya and N. Victoir. Weak approximation of stochastic differential equations and application to derivative pricing. *Appl. Math. Finance*, 15(1-2):107–121, 2008.
- [12] K. Oshima, J. Teichmann, and D. Velušček. A new extrapolation method for weak approximation schemes with applications. *Ann. Appl. Probab.*, 22(3):1008–1045, 2012.
- [13] G. Pagès. Multi-step Richardson-Romberg extrapolation: remarks on variance control and complexity. *Monte Carlo Methods Appl.*, 13(1):37–70, 2007.
- [14] D. Talay and L. Tubaro. Expansion of the global error for numerical schemes solving stochastic differential equations. *Stochastic Anal. Appl.*, 8(4):483–509 (1991), 1990.
- [15] D. V. Widder. *The Laplace Transform*. Princeton Mathematical Series, vol. 6. Princeton University Press, Princeton, N. J., 1941.

AURÉLIEN ALFONSI, CERMICS, ECOLE DES PONTS, MARNE-LA-VALLÉE, FRANCE. MATHRISK, INRIA, PARIS, FRANCE.

Email address: aurelien.alfonsi@enpc.fr

EDOARDO LOMBARDO, CERMICS, ECOLE DES PONTS, MARNE-LA-VALLÉE, FRANCE. MATHRISK, INRIA, PARIS, FRANCE. UNIVERSITÀ DEGLI STUDI DI ROMA TOR VERGATA, ROME, ITALY.

Email address: edoardo.lombardo@enpc.fr