

# THE G-DRAZIN INVERSES OF ANTI-TRIANGULAR BLOCK OPERATOR MATRICES

HUANYIN CHEN AND MARJAN SHEIBANI

ABSTRACT. An element  $a$  in a Banach algebra  $\mathcal{A}$  has g-Drazin inverse if there exists  $b \in \mathcal{A}$  such that  $ab = ba, b = bab$  and  $a - a^2b \in \mathcal{A}^{qnil}$ . In this paper we find new explicit representations of the g-Drazin inverse of the block operator matrix  $\begin{pmatrix} E & I \\ F & 0 \end{pmatrix}$ . We thereby solve a wider kind of singular differential equations posed by Campbell [S.L. Campbell, The Drazin inverse and systems of second order linear differential equations, Linear & Multilinear Algebra, 14(1983), 195–198].

## 1. INTRODUCTION

Let  $\mathcal{A}$  be a complex Banach algebra with an identity. An element  $a$  in a Banach algebra  $\mathcal{A}$  has g-Drazin inverse provided that there exists some  $b \in \mathcal{A}$  such that  $b = bab, ab = ba, a - a^2b \in \mathcal{A}^{qnil}$ . Here,  $\mathcal{A}^{qnil} = \{a \in \mathcal{A} \mid \lim_{n \rightarrow \infty} \|a^n\|^{\frac{1}{n}} = 0\}$ . Such  $b$  is unique, if it exists, and we denote it by  $a^d$ . We always use  $\mathcal{A}^d$  to stand for the set of all g-Drazin invertible elements in  $\mathcal{A}$ . We say  $a$  has Drazin inverse if there exists some  $b \in \mathcal{A}$  such that  $b = bab, ab = ba, a^n = a^{n+1}b$ . Such  $b$  is unique, if exists, and we denote it by  $a^D$ . The smallest integer  $n$  satisfying the preceding equations is called the index of  $a$ . Evidently, every Drazin invertible  $a \in \mathcal{A}$  has g-Drazin inverse and  $a^D = a^d$ .

Let  $E, F$  be block operator matrices and  $I$  be the identity matrix over a Banach space  $X$ . It is attractive to investigate the

---

2010 *Mathematics Subject Classification.* 15A09, 47A11, 47A53.

*Key words and phrases.* g-Drazin inverse; Drazin inverse; operator matrix; Banach algebra.

Drazin inverse of the block matrix  $M = \begin{pmatrix} E & I \\ F & 0 \end{pmatrix}$ . The relationship of computing the Drazin inverse of the block complex matrices  $\begin{pmatrix} E & -F \\ -I & 0 \end{pmatrix}$  to second order differential equations was observed by Campbell (see [2]). The application of Drazin inverse to singular difference equations was also found in [1]. By using Cline's formula (see [7, Theorem 2.1]), we easily see that

$$\begin{pmatrix} E & -F \\ -I & 0 \end{pmatrix}^D = \begin{pmatrix} E & I \\ -I & 0 \end{pmatrix} \left( \begin{pmatrix} E & I \\ F & 0 \end{pmatrix}^D \right)^2 \begin{pmatrix} I & 0 \\ 0 & -F \end{pmatrix}.$$

Thus finding the solution of singular differential equation  $Ax''(t) + Bx(t) + Cx(t) = 0$  is equivalent to finding the explicit representation of the Drazin inverse for the block complex matrix  $M$ .

In 2005, Castro-González and Dopazo considered the representations of the Drazin inverse for a class of block matrix  $\begin{pmatrix} I & I \\ F & 0 \end{pmatrix}$  (see [3]). In 2011, Bu et al. gave the Drazin inverse of the preceding block matrix  $M$  under  $EF = FE$  (see [1]). Patricio and Hartwig investigated the g-Drazin inverse of  $M$  under  $F^\pi E F F^d = 0$ ,  $F^\pi F E = E F F^\pi$  (see [9]). Here,  $F^\pi = I - F F^d$ . In 2016, Zhang studied the g-Drazin inverse of  $M$  under  $F^d E F^\pi = 0$ ,  $F^\pi F E = 0$  and  $F^\pi E F^d = 0$ ,  $E F F^\pi = 0$  (see [11, Theorem 2.6, Theorem 2.8]).

In [5, Theorem 3.1], Deng and Wei gave formula for the g-Drazin inverse of  $M$  under  $EF = 0$ . In [4, Theorem 2.2], Cvetković-Ilić obtained the representation of Drazin inverse  $M^D$  under certain conditions  $P_n = 0$ . The motivation of this paper is to present formulae for the g-Drazin inverse of  $M$  under new conditions. We shall give formula for the g-Drazin inverse  $M^d$  under  $EF^2 = 0$ ,  $EFE = 0$ . Then we give the explicit representation of  $M^d$  under the condition  $F^\pi E F^2 = 0$  and  $F^\pi E F E = 0$ . In particular, we obtain new representations of  $M^D$  for an anti-triangular complex matrix  $M$ . We thereby solve a wider kind of singular differential equations posed by Campbell (see [2]).

Throughout the paper, all matrices are bounded linear operator matrices over a Banach space  $X$ . Let  $a \in \mathcal{A}$  and  $e^2 = e \in \mathcal{A}$ . Then

$a$  has the Pierce decomposition relative to  $e$ , and we denote it by  $\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}_e$ .  $[x]$  stands for the truncates integer of  $x$ .

## 2. MAIN RESULTS

We begin with the following elementary lemma.

**Lemma 2.1.** ([11, Lemma 2.2]) *Let  $A, B, C$  are operators over a Banach space  $X$ . If  $A$  and  $B$  have  $g$ -Drazin inverses, then  $M = \begin{pmatrix} A & 0 \\ C & B \end{pmatrix}$  has  $g$ -Drazin inverse and*

$$M^d = \begin{pmatrix} A^d & 0 \\ X & B^d \end{pmatrix},$$

where

$$X = (B^d)^2 \left( \sum_{i=0}^{\infty} (B^d)^i C A^i \right) A^\pi + B^\pi \left( \sum_{i=0}^{\infty} B^i C (A^d)^i \right) (A^d)^2 - B^d C A^d.$$

**Lemma 2.2.** *Let  $P$  and  $Q$  have  $g$ -Drazin inverses. If  $PQ^2 = 0, PQP = 0$ , then  $P + Q$  has  $g$ -Drazin inverse and*

$$\begin{aligned} & (P + Q)^d \\ &= Q^\pi \sum_{i=0}^{\infty} Q^i (P^d)^{i+1} + \sum_{i=0}^{\infty} (Q^d)^{i+1} P^i P^\pi + Q^\pi \sum_{i=0}^{\infty} Q^i (P^d)^{i+2} Q \\ &+ \sum_{i=0}^{\infty} (Q^d)^{i+3} P^{i+1} P^\pi Q - Q^d P^d Q - (Q^d)^2 P P^d Q. \end{aligned}$$

*Proof.* This was proved in [10, Theorem 2.1] for Drazin inverse of matrices, the proof is similar for  $g$ -Drazin inverse of operators.  $\square$

We are ready to prove the following.

**Theorem 2.3.** *Let  $E$  and  $F$  have  $g$ -Drazin inverses. If  $EF^2 = 0$  and  $EFE = 0$ , then  $M = \begin{pmatrix} E & I \\ F & 0 \end{pmatrix}$  has  $g$ -Drazin inverse. In this case,*

$$M^d = \begin{pmatrix} \Gamma & \Delta \\ \Lambda & \Xi \end{pmatrix},$$

where

$$\begin{aligned}
\Gamma &= E^d + (E^d)^3 F - F^d E^d F - (F^d)^2 E^2 E^d F - F F^d E^d + F^d E E^\pi \\
&+ (F^d)^2 E F - F F^d (E^d)^3 F + (F^d)^3 E^3 E^\pi F + \sum_{i=1}^{\infty} F^i F^\pi E^{2i+1} \\
&+ \sum_{i=1}^{\infty} (F^d)^{i+1} E^{2i+1} E^\pi + \sum_{i=1}^{\infty} F^i F^\pi E^{2i} E^d F + \sum_{i=1}^{\infty} (F^d)^{i+3} E^{2i+3} E^\pi F, \\
\Delta &= (E^d)^2 + (E^d)^4 F - F^d (E^d)^2 F - (F^d)^2 E E^d F - F F^d (E^d)^2 \\
&+ F^d E^\pi - F F^d (E^d)^4 F + (F^d)^3 E^2 E^\pi F + \sum_{i=1}^{\infty} F^i F^\pi E^{2i+1} E^d \\
&+ \sum_{i=1}^{\infty} (F^d)^{i+1} E^{2i} E^\pi + \sum_{i=1}^{\infty} F^i F^\pi E^{2i-1} E^d F + \sum_{i=1}^{\infty} (F^d)^{i+3} E^{2i+2} E^\pi F; \\
\Lambda &= F F^\pi (E^d)^2 + F F^d E^\pi + F F^\pi (E^d)^4 F + (F^d)^2 E^2 E^\pi F \\
&- F F^d (E^d)^2 F - F^d E E^d F + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+2} + \sum_{i=1}^{\infty} (F^d)^i E^{2i} E^\pi \\
&+ \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+4} F + \sum_{i=1}^{\infty} (F^d)^{i+2} E^{2i+2} E^\pi F, \\
\Xi &= -F F^d (E^d)^3 F - F^d E^d F + F F^\pi (E^d)^3 - F F^d E^d \\
&+ F F^\pi (E^d)^5 F - (F^d)^2 E^2 E^d F + (F^d)^2 E F + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+3} \\
&+ \sum_{i=1}^{\infty} (F^d)^i E^{2i-1} E^\pi + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+5} F + \sum_{i=1}^{\infty} (F^d)^{i+2} E^{2i+1} E^\pi F \quad (*)
\end{aligned}$$

*Proof.* Clearly, we have

$$M^2 = \begin{pmatrix} E^2 + F & E \\ FE & F \end{pmatrix} = P + Q,$$

where

$$P = \begin{pmatrix} E^2 & E \\ 0 & 0 \end{pmatrix}, Q = \begin{pmatrix} F & 0 \\ FE & F \end{pmatrix}.$$

In light of Lemma 2.1, we have

$$\begin{aligned}
P^d &= \begin{pmatrix} (E^d)^2 & (E^d)^3 \\ 0 & 0 \end{pmatrix}, P^\pi = \begin{pmatrix} E^\pi & -E^d \\ 0 & I \end{pmatrix}; \\
Q^d &= \begin{pmatrix} F^d & 0 \\ X & F^d \end{pmatrix}; Q^\pi = \begin{pmatrix} F^\pi & 0 \\ -FX & F^\pi \end{pmatrix},
\end{aligned}$$

where

$$\begin{aligned}
X &= F^\pi \sum_{i=0}^{\infty} F^{i+1} E (F^d)^{i+2} + \sum_{i=0}^{\infty} (F^d)^{i+2} F E F^i F^\pi \\
&= (F^d)^2 F E + (F^d)^3 F E F \\
&= F^d E + (F^d)^2 E F.
\end{aligned}$$

For any  $n \geq 2$ , we have

$$\begin{aligned} P^n &= \begin{pmatrix} E^{2n} & E^{2n-1} \\ 0 & 0 \end{pmatrix}; (P^d)^n = \begin{pmatrix} (E^d)^{2n} & (E^d)^{2n+1} \\ 0 & 0 \end{pmatrix}; \\ Q^n &= \begin{pmatrix} F^n & 0 \\ Y_n & F^n \end{pmatrix}, Y_1 = FE, Y_n = Y_{n-1}F + F^nE; \\ (Q^d)^n &= \begin{pmatrix} (F^d)^n & 0 \\ X_n & (F^d)^n \end{pmatrix}, X_1 = X, X_n = X_{n-1}F^d + F^dX_{n-1}. \end{aligned}$$

Since  $EF^2 = 0$  and  $EFE = 0$ , by induction, we verify that

$$X_n = (F^d)^n E + (F^d)^{n+1} EF, Y_n = F^{n-1} EF + F^n E.$$

Moreover, we check that

$$PQ^2 = 0, PQP = 0.$$

In view of Lemma 2.2,  $M^2 = P + Q$  has g-Drazin inverse. Further, we get

$$\begin{aligned} (M^2)^d &= (P + Q)^d \\ &= Q^\pi \sum_{i=0}^{\infty} Q^i (P^d)^{i+1} + \sum_{i=0}^{\infty} (Q^d)^{i+1} P^i P^\pi + Q^\pi \sum_{i=0}^{\infty} Q^i (P^d)^{i+2} Q \\ &+ \sum_{i=0}^{\infty} (Q^d)^{i+3} P^{i+1} P^\pi Q - Q^d P^d Q - (Q^d)^2 P P^d Q. \end{aligned}$$

We easily check that

$$\begin{aligned} Q^d P^d Q &= \begin{pmatrix} F^d (E^d)^2 F & F^d (E^d)^3 F \\ X (E^d)^2 F & X (E^d)^3 F \end{pmatrix}; \\ (Q^d)^2 P P^d Q &= \begin{pmatrix} (F^d)^2 E E^d F & (F^d)^2 E^d F \\ X_2 E E^d F & X_2 E^d F \end{pmatrix}. \end{aligned}$$

If  $i \geq 1$ , we have

$$\begin{aligned} Q^\pi Q^i (P^d)^{i+1} &= \\ &\begin{pmatrix} F^\pi F^i (E^d)^{2i+2} & F^\pi F^i (E^d)^{2i+3} \\ -FXF^i (E^d)^{2i+2} + F^\pi Y_i (E^d)^{2i+2} & -FXF^i (E^d)^{2i+3} + F^\pi Y_i (E^d)^{2i+3} \end{pmatrix}; \\ Q^\pi P^d &= \begin{pmatrix} F^\pi (E^d)^2 & F^\pi (E^d)^3 \\ -FX(E^d)^2 & -FX(E^d)^3 \end{pmatrix}. \end{aligned}$$

If  $i \geq 1$ , we have

$$(Q^d)^{i+1} P^i P^\pi = \begin{pmatrix} (F^d)^{i+1} E^{2i} E^\pi & -(F^d)^{i+1} E^{2i} E^d + (F^d)^{i+1} E^{2i-1} \\ X_{i+1} E^{2i} E^\pi & -X_{i+1} E^{2i} E^d + X_{i+1} E^{2i-1} \end{pmatrix};$$

$$Q^d P^\pi = \begin{pmatrix} F^d E^\pi & -F^d E^d \\ X E^\pi & -X E^d + F^d \end{pmatrix};$$

If  $i \geq 1$ , we have

$$Q^\pi Q^i (P^d)^{i+2} Q = \begin{pmatrix} F^\pi F^i (E^d)^{2i+4} F & F^\pi F^i (E^d)^{2i+5} F \\ -F X F^i (E^d)^{2i+4} F + F^\pi Y_i (E^d)^{2i+4} F & -F X F^i (E^d)^{2i+5} F + F^\pi Y_i (E^d)^{2i+5} F \end{pmatrix};$$

$$Q^\pi (P^d)^2 Q = \begin{pmatrix} F^\pi (E^d)^4 F & F^\pi (E^d)^5 F \\ -F X (E^d)^4 F & -F X (E^d)^5 F \end{pmatrix};$$

$$\begin{aligned} & (Q^d)^{i+3} P^{i+1} P^\pi Q = \\ & \begin{pmatrix} (F^d)^{i+3} E^{2i+2} E^\pi F & -(F^d)^{i+3} E^{2i+2} E^d F + (F^d)^{i+3} E^{2i+1} F \\ X_{i+3} E^{2i+2} E^\pi F & -X_{i+3} E^{2i+2} E^d F + X_{i+3} E^{2i+1} F \end{pmatrix}. \end{aligned}$$

Since  $X E^d = F^d E E^d$  and  $X_2 = (F^d)^2 E + (F^d)^3 E F$ , we have

$$\begin{aligned} Q^d P^d Q &= \begin{pmatrix} F^d (E^d)^2 F & F^d (E^d)^3 F \\ F^d E^d F & F^d (E^d)^2 F \end{pmatrix}; \\ (Q^d)^2 P P^d Q &= \begin{pmatrix} (F^d)^2 E E^d F & (F^d)^2 E^d F \\ (F^d)^2 E^2 E^d F & (F^d)^2 E E^d F \end{pmatrix}. \end{aligned}$$

Moreover, we have

$$\begin{aligned} Q^\pi P^d &= \begin{pmatrix} F^\pi (E^d)^2 & F^\pi (E^d)^3 \\ -F F^d E^d & -F F^d (E^d)^2 \end{pmatrix}; \\ Q^d P^\pi &= \begin{pmatrix} F^d E^\pi & -F^d E^d \\ F^d E E^\pi + (F^d)^2 E F & F^d E^\pi \end{pmatrix}. \end{aligned}$$

Also we check that

$$\begin{aligned} F X F^i (E^d)^{2i+2} &= \begin{cases} F F^d E^d & i = 0; \\ 0 & i \geq 1; \end{cases} \\ F^\pi Y_i (E^d)^{2i+2} &= F^\pi F^i E^{2i+1} \quad (i \geq 1); \\ X_{i+1} E^{2i-1} &= (F^d)^{i+1} E^{2i} \quad (i \geq 1); \\ X_{i+3} E^{2i+1} &= (F^d)^{i+3} E^{2i+2} \quad (i \geq 0). \end{aligned}$$

By virtue of [8, Corollary 2.2],  $M$  has g-Drazin inverse. Therefore

$$M^d = M(M^d)^2 = M(M^2)^d = \begin{pmatrix} E & I \\ F & 0 \end{pmatrix} (M^2)^d = \begin{pmatrix} \Gamma & \Delta \\ \Lambda & \Xi \end{pmatrix},$$

where  $\Gamma, \Delta, \Lambda$  and  $\Xi$  are given as in (\*) by direct computation.  $\square$

For a block complex matrix, the Drazin and g-Drazin inverses coincide with each other, and so we derive

**Corollary 2.4.** *Let  $E$  and  $F$  be  $n \times n$  complex matrices, and let  $M = \begin{pmatrix} E & I \\ F & 0 \end{pmatrix}$ . If  $EF^2 = 0$  and  $EFE = 0$ , then*

$$M^D = \begin{pmatrix} \Gamma & \Delta \\ \Lambda & \Xi \end{pmatrix},$$

where

$$\begin{aligned} \Gamma &= E^D + (E^D)^3 F - F^D E^D F - (F^D)^2 E^2 E^D F - F F^D E^D + F^D E E^\pi \\ &+ (F^D)^2 E F - F F^D (E^D)^3 F + (F^D)^3 E^3 E^\pi F + \sum_{i=1}^t F^i F^\pi E^{2i+1} \\ &+ \sum_{i=1}^{\lfloor \frac{s}{2} \rfloor} (F^D)^{i+1} E^{2i+1} E^\pi + \sum_{i=1}^t F^i F^\pi E^{2i} E^D F + \sum_{i=1}^{\lfloor \frac{s}{2} \rfloor - 1} (F^D)^{i+3} E^{2i+3} E^\pi F, \\ \Delta &= (E^D)^2 + (E^D)^4 F - F^D (E^D)^2 F - (F^D)^2 E E^D F - F F^D (E^D)^2 \\ &+ F^D E^\pi - F F^D (E^D)^4 F + (F^D)^3 E^2 E^\pi F + \sum_{i=1}^t F^i F^\pi E^{2i+1} E^D \\ &+ \sum_{i=1}^{\lfloor \frac{s}{2} \rfloor + 1} (F^D)^{i+1} E^{2i} E^\pi + \sum_{i=1}^t F^i F^\pi E^{2i-1} E^D F + \sum_{i=1}^{\lfloor \frac{s}{2} \rfloor} (F^D)^{i+3} E^{2i+2} E^\pi F; \\ \Lambda &= F F^\pi (E^D)^2 + F F^D E^\pi + F F^\pi (E^D)^4 F + (F^D)^2 E^2 E^\pi F \\ &- F F^D (E^D)^2 F - F^D E E^D F + \sum_{i=1}^{t-1} F^{i+1} F^\pi (E^D)^{2i+2} + \sum_{i=1}^{\lfloor \frac{s}{2} \rfloor + 1} (F^D)^i E^{2i} E^\pi \\ &+ \sum_{i=1}^{t-1} F^{i+1} F^\pi (E^D)^{2i+4} F + \sum_{i=1}^{\lfloor \frac{s}{2} \rfloor} (F^D)^{i+2} E^{2i+2} E^\pi F, \\ \Xi &= -F F^D (E^D)^3 F - F^D E^D F + F F^\pi (E^D)^3 - F F^D E^D \\ &+ F F^\pi (E^D)^5 F - (F^D)^2 E^2 E^D F + (F^D)^2 E F + \sum_{i=1}^{t-1} F^{i+1} F^\pi (E^D)^{2i+3} \\ &+ \sum_{i=1}^{\lfloor \frac{s}{2} \rfloor + 1} (F^D)^i E^{2i-1} E^\pi + \sum_{i=1}^{t-1} F^{i+1} F^\pi (E^D)^{2i+5} F + \sum_{i=1}^{\lfloor \frac{s}{2} \rfloor} (F^D)^{i+2} E^{2i+1} E^\pi F; \end{aligned}$$

$s = i(E)$  and  $t = i(F)$ .

*Proof.* Since  $E^s E^\pi = F^t F^\pi = 0$ , we obtain the result by Theorem 2.3.  $\square$

We give a numerical example to illustrate the representation of Corollary 2.4.

**Example 2.5.** *Let  $M = \begin{pmatrix} E & I \\ F & 0 \end{pmatrix}$ , where*

$$E = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, F = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \in M_2(\mathbb{C}).$$

Then  $EF^2 = 0$  and  $EFE = 0$ , while  $EF \neq 0$ . In this case,

$$M^D = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

*Proof.* Clearly, we have

$$EF^2 = 0, EFE = 0 \text{ and } EF = F \neq 0.$$

To compute the Drazin inverse of  $M$ , we shall apply Corollary 2.4 to  $M$ . One easily verifies that

$$F^2 = F^D = FE = 0, E^2 = E.$$

Using the notations in Corollary 2.4, we directly compute

$$\Gamma = \Delta = E + EF = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} \text{ and } \Lambda = \Xi = 0,$$

as required.  $\square$

We have accumulated all the information necessary to prove the main result.

**Theorem 2.6.** *Let  $E, F, EF^\pi$  have  $g$ -Drazin inverses. If  $F^\pi EF^2 = 0$  and  $F^\pi EFE = 0$ , then  $M = \begin{pmatrix} E & I \\ F & 0 \end{pmatrix}$  has  $g$ -Drazin inverse. In this case,*

$$M^d = \begin{pmatrix} \Theta & \Psi \\ \Phi & \Omega \end{pmatrix} + \sum_{i=0}^{\infty} \begin{pmatrix} 0 & F^d \\ FF^d & -EF^d \end{pmatrix}^{i+1} \begin{pmatrix} F^\pi E + FF^d EF^\pi & F^\pi \\ FF^\pi & 0 \end{pmatrix}^i \begin{pmatrix} I - E\Theta & -E\Psi \\ -F\Theta & I - F\Psi \end{pmatrix},$$



where

$$\begin{aligned}
 \Theta &= [F^\pi E^d + F^\pi (E^d)^3 F F^\pi + F F^\pi (E^d)^3 + F F^\pi (E^d)^5 F F^\pi \\
 &+ \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+3} + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+5} F F^\pi] \\
 &\quad [F^\pi E E^d + F^\pi (E^d)^2 F F^\pi + F F^\pi (E^d)^2 + F F^\pi (E^d)^4 F F^\pi \\
 &+ \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+2} + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+4} F F^\pi] \\
 &+ [F^\pi E E^d + F^\pi (E^d)^2 F F^\pi + F F^\pi (E^d)^2 + F F^\pi (E^d)^4 F F^\pi \\
 &- F^\pi E^d F^\pi E - F^\pi (E^d)^3 F F^\pi E - F F^\pi (E^d)^3 F^\pi E - F F^\pi (E^d)^5 F F^\pi E \\
 &+ \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+2} + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+4} F F^\pi \\
 &- \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+3} F^\pi E - \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+5} F F^\pi E] \\
 &\quad [F^\pi E^d + F^\pi (E^d)^3 F F^\pi + \sum_{i=1}^{\infty} F^i F^\pi E^{2i+1} + \sum_{i=1}^{\infty} F^i F^\pi E^{2i} E^d F F^\pi],
 \end{aligned}$$

$$\begin{aligned}
 \Psi &= [F^\pi E^d F^\pi + F^\pi (E^d)^3 F F^\pi + F F^\pi (E^d)^3 F^\pi + F F^\pi (E^d)^5 F F^\pi \\
 &+ \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+3} F^\pi + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+5} F F^\pi]^2 \\
 &+ [F^\pi E E^d + F^\pi (E^d)^2 F F^\pi + F F^\pi (E^d)^2 + F F^\pi (E^d)^4 F F^\pi \\
 &- F^\pi E^d F^\pi E - F^\pi (E^d)^3 F F^\pi E - F F^\pi (E^d)^3 F^\pi E - F F^\pi (E^d)^5 F F^\pi E \\
 &+ \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+2} + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+4} F F^\pi \\
 &- \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+3} F^\pi E - \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+5} F F^\pi E] \\
 &\quad [F^\pi (E^d)^2 F^\pi + F^\pi (E^d)^4 F F^\pi + \sum_{i=1}^{\infty} F^i F^\pi E^{2i+1} F^\pi E^d F^\pi \\
 &+ \sum_{i=1}^{\infty} F^i F^\pi E^{2i-1} F^\pi E^d F F^\pi],
 \end{aligned}$$

$$\begin{aligned}
 \Phi &= [F F^\pi (E^d)^2 + F F^\pi (E^d)^4 F F^\pi + \sum_{i=1}^{\infty} F^{i+1} F^\pi E^{2i+1} F^\pi E^d \\
 &+ \sum_{i=1}^{\infty} F^{i+1} F^\pi E^{2i-1} F^\pi E^d F F^\pi] \\
 &\quad [F^\pi E E^d + F^\pi (E^d)^2 F F^\pi + F F^\pi (E^d)^2 + F F^\pi (E^d)^4 F F^\pi \\
 &+ \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+2} + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+4} F F^\pi] \\
 &+ [F F^\pi E^d + F F^\pi (E^d)^3 F F^\pi - F F^\pi (E^d)^2 F^\pi E - F F^\pi (E^d)^4 F F^\pi E \\
 &+ \sum_{i=1}^{\infty} F^{i+1} F^\pi E^{2i+1} + \sum_{i=1}^{\infty} F^{i+1} F^\pi E^{2i} E^d F F^\pi \\
 &- \sum_{i=1}^{\infty} F^{i+1} F^\pi E^{2i+1} F^\pi E^d F^\pi E - \sum_{i=1}^{\infty} F^{i+1} F^\pi E^{2i-1} F^\pi E^d F F^\pi E] \\
 &\quad [F^\pi E^d + F^\pi (E^d)^3 F F^\pi + \sum_{i=1}^{\infty} F^i F^\pi E^{2i+1} + \sum_{i=1}^{\infty} F^i F^\pi E^{2i} E^d F F^\pi],
 \end{aligned}$$

$$\begin{aligned}
\Omega &= [FF^\pi(E^d)^2 + FF^\pi(E^d)^4FF^\pi + \sum_{i=1}^{\infty} F^{i+1}F^\pi E^{2i+1}F^\pi E^d \\
&+ \sum_{i=1}^{\infty} F^{i+1}F^\pi E^{2i-1}F^\pi E^d FF^\pi] \\
&\quad [F^\pi E^d F^\pi + F^\pi(E^d)^3 FF^\pi + FF^\pi(E^d)^3 F^\pi + FF^\pi(E^d)^5 FF^\pi \\
&+ \sum_{i=1}^{\infty} F^{i+1}F^\pi(E^d)^{2i+3}F^\pi + \sum_{i=1}^{\infty} F^{i+1}F^\pi(E^d)^{2i+5}FF^\pi] \\
&+ [FF^\pi E^d + F^\pi(E^d)^3 FF^\pi - FF^\pi(E^d)^2 F^\pi E - F^\pi(E^d)^4 FF^\pi E \\
&+ \sum_{i=1}^{\infty} F^{i+1}F^\pi E^{2i+1} + \sum_{i=1}^{\infty} F^{i+1}F^\pi E^{2i} E^d FF^\pi] \\
&\quad [F^\pi(E^d)^2 F^\pi + F^\pi(E^d)^4 FF^\pi + \sum_{i=1}^{\infty} F^i F^\pi E^{2i+1} F^\pi E^d F^\pi \\
&+ \sum_{i=1}^{\infty} F^i F^\pi E^{2i-1} F^\pi E^d FF^\pi] (**).
\end{aligned}$$

*Proof.* Let  $e = \begin{pmatrix} F^\pi & 0 \\ 0 & 0 \end{pmatrix}$ . Then  $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}_e$ , where

$$a = eMe, b = eM(I - e), c = (I - e)Me, d = (I - e)M(I - e).$$

Since  $F^\pi EF^2 = 0$  and  $F^\pi EFE = 0$ , we have

$$\begin{aligned}
a &= \begin{pmatrix} F^\pi E & 0 \\ 0 & 0 \end{pmatrix}, b = \begin{pmatrix} 0 & F^\pi \\ 0 & 0 \end{pmatrix}, \\
c &= \begin{pmatrix} FF^d EF^\pi & 0 \\ FF^\pi & 0 \end{pmatrix}, d = \begin{pmatrix} EFF^d & FF^d \\ F^2 F^d & 0 \end{pmatrix}.
\end{aligned}$$

By direct computation, we get

$$d^d = \begin{pmatrix} 0 & F^d \\ FF^d & -EF^d \end{pmatrix}, d^\pi = \begin{pmatrix} 0 & 0 \\ 0 & F^\pi \end{pmatrix}, dd^\pi = 0.$$

As  $EF^\pi$  has g-Drazin inverse, it follows by Cline's formula,  $F^\pi E$  has g-Drazin inverse. Since  $F^\pi EFF^d = 0$ , it follows by [12, Lemma 2.2] that  $(F^\pi E)^d = F^\pi E^d$ , and so

$$a^d = \begin{pmatrix} F^\pi E^d & 0 \\ 0 & 0 \end{pmatrix}, a^\pi = \begin{pmatrix} F^\pi E^\pi & 0 \\ 0 & 0 \end{pmatrix}.$$

Write  $M = P + Q$ , where

$$P = \begin{pmatrix} a & b \\ c & 0 \end{pmatrix}_e, Q = \begin{pmatrix} 0 & 0 \\ 0 & d \end{pmatrix}_e.$$

Step 1. We prove that  $P$  has g-Drazin inverse. Clearly,  $bd = 0, bc = \begin{pmatrix} FF^\pi & 0 \\ 0 & 0 \end{pmatrix}$  has g-Drazin inverse,  $(bc)^d = 0$  and  $(bc)^\pi = 1$ .

We easily check that  $a(bc)^2 = 0, a(bc)a = 0$ . In light of Theorem 2.3,  $\begin{pmatrix} a & 1 \\ bc & 0 \end{pmatrix}$  has g-Drazin inverse. Moreover, we have

$$\begin{pmatrix} a & 1 \\ bc & 0 \end{pmatrix}^d = \begin{pmatrix} \Gamma & \Delta \\ \Lambda & \Xi \end{pmatrix},$$

where

$$\begin{aligned} (\Gamma)_{11} &= F^\pi E^d + F^\pi (E^d)^3 F F^\pi + \sum_{i=1}^{\infty} F^i F^\pi E^{2i+1} + \sum_{i=1}^{\infty} F^i F^\pi E^{2i} E^d F F^\pi, \\ (\Delta)_{11} &= F^\pi (E^d)^2 + F^\pi (E^d)^4 F F^\pi + \sum_{i=1}^{\infty} F^i F^\pi E^{2i+1} F^\pi E^d + \sum_{i=1}^{\infty} F^i F^\pi E^{2i-1} F^\pi E^d F F^\pi, \\ (\Lambda)_{11} &= F F^\pi (E^d)^2 + F F^\pi (E^d)^4 F F^\pi + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+2} + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+4} F F^\pi, \\ (\Xi)_{11} &= F F^\pi (E^d)^3 + F F^\pi (E^d)^5 F F^\pi + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+3} + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+5} F F^\pi, \\ (\Gamma)_{ij} &= 0, (\Delta)_{ij} = 0, (\Lambda)_{ij} = 0, (\Xi)_{ij} = 0, \text{ otherwise.} \end{aligned}$$

Since

$$\begin{pmatrix} a & bc \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & -a \end{pmatrix}^{-1} \begin{pmatrix} a & 1 \\ bc & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & -a \end{pmatrix},$$

we see that  $\begin{pmatrix} a & bc \\ 1 & 0 \end{pmatrix}$  has g-Drazin inverse and

$$\begin{pmatrix} a & bc \\ 1 & 0 \end{pmatrix}^d = \begin{pmatrix} 0 & 1 \\ 1 & -a \end{pmatrix}^{-1} \begin{pmatrix} a & 1 \\ bc & 0 \end{pmatrix}^d \begin{pmatrix} 0 & 1 \\ 1 & -a \end{pmatrix};$$

hence,

$$\begin{pmatrix} a & bc \\ 1 & 0 \end{pmatrix}^d = \begin{pmatrix} a\Delta + \Xi & a\Gamma + \Lambda - a\Delta a - \Xi a \\ \Delta & \Gamma - \Delta a \end{pmatrix}.$$

Clearly, we have

$$\begin{pmatrix} a & bc \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} a & b \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & c \end{pmatrix}.$$

By using Cline's formula,

$$\begin{aligned}
P^d &= \begin{pmatrix} 1 & 0 \\ 0 & c \end{pmatrix} \left( \begin{pmatrix} a & bc \\ 1 & 0 \end{pmatrix}^d \right)^2 \begin{pmatrix} a & b \\ 1 & 0 \end{pmatrix} \\
&= \begin{pmatrix} a\Delta + \Xi & a\Gamma + \Lambda - a\Delta a - \Xi a \\ c\Delta & c\Gamma - c\Delta a \end{pmatrix} \begin{pmatrix} a\Gamma + \Lambda & a\Delta b + \Xi b \\ \Gamma & \Delta b \end{pmatrix} \\
&= \begin{pmatrix} \gamma & \delta \\ \lambda & \xi \end{pmatrix},
\end{aligned}$$

where

$$\begin{aligned}
\gamma &= (a\Delta + \Xi)(a\Gamma + \Lambda) + (a\Gamma + \Lambda - a\Delta a - \Xi a)\Gamma, \\
\delta &= (a\Delta + \Xi)^2 b + (a\Gamma + \Lambda - a\Delta a - \Xi a)\Delta b, \\
\lambda &= c\Delta(a\Gamma + \Lambda) + c(\Gamma - \Delta a)\Gamma, \\
\xi &= c\Delta(a\Delta + \Xi)b + c(\Gamma - \Delta a)\Delta b.
\end{aligned}$$

We directly compute that

$$\begin{aligned}
a\Delta &= \begin{pmatrix} F^\pi E^d + F^\pi (E^d)^3 F F^\pi & 0 \\ 0 & 0 \end{pmatrix}, \\
a\Gamma &= \begin{pmatrix} F^\pi E E^d + F^\pi (E^d)^2 F F^\pi & 0 \\ 0 & 0 \end{pmatrix}.
\end{aligned}$$

Hence we have

$$\begin{aligned}
(a\Delta + \Xi)_{11} &= F^\pi E^d + F^\pi (E^d)^3 F F^\pi + F F^\pi (E^d)^3 + F F^\pi (E^d)^5 F F^\pi \\
&\quad + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+3} + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+5} F F^\pi, \\
(a\Delta + \Xi)_{ij} &= 0, \text{ otherwise;} \\
(a\Gamma + \Lambda)_{11} &= F^\pi E E^d + F^\pi (E^d)^2 F F^\pi + F F^\pi (E^d)^2 + F F^\pi (E^d)^4 F F^\pi \\
&\quad + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+2} + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+4} F F^\pi, \\
(a\Gamma + \Lambda)_{ij} &= 0, \text{ otherwise.}
\end{aligned}$$

Also we have

$$\begin{aligned}
 & (a\Gamma + \Lambda - a\Delta a - \Xi a)_{11} \\
 = & F^\pi E E^d + F^\pi (E^d)^2 F F^\pi + F F^\pi (E^d)^2 + F F^\pi (E^d)^4 F F^\pi \\
 - & F^\pi E^d F^\pi E - F^\pi (E^d)^3 F F^\pi E - F F^\pi (E^d)^3 F^\pi E - F F^\pi (E^d)^5 F F^\pi E \\
 + & \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+2} + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+4} F F^\pi \\
 - & \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+3} F^\pi E - \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+5} F F^\pi E, \\
 & (a\Gamma + \Lambda - a\Delta a - \Xi a)_{ij} = 0, \text{ otherwise.}
 \end{aligned}$$

Therefore

$$\begin{aligned}
 (\gamma)_{11} &= [F^\pi E^d + F^\pi (E^d)^3 F F^\pi + F F^\pi (E^d)^3 + F F^\pi (E^d)^5 F F^\pi \\
 &+ \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+3} + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+5} F F^\pi] \\
 & [F^\pi E E^d + F^\pi (E^d)^2 F F^\pi + F F^\pi (E^d)^2 + F F^\pi (E^d)^4 F F^\pi \\
 &+ \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+2} + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+4} F F^\pi] \\
 &+ [F^\pi E E^d + F^\pi (E^d)^2 F F^\pi + F F^\pi (E^d)^2 + F F^\pi (E^d)^4 F F^\pi \\
 &- F^\pi E^d F^\pi E - F^\pi (E^d)^3 F F^\pi E - F F^\pi (E^d)^3 F^\pi E - F F^\pi (E^d)^5 F F^\pi E \\
 &+ \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+2} + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+4} F F^\pi \\
 &- \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+3} F^\pi E - \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+5} F F^\pi E] \\
 & [F^\pi E^d + F^\pi (E^d)^3 F F^\pi + \sum_{i=1}^{\infty} F^i F^\pi E^{2i+1} + \sum_{i=1}^{\infty} F^i F^\pi E^{2i} E^d F F^\pi], \\
 (\gamma)_{ij} &= 0, \text{ otherwise.}
 \end{aligned}$$

We see that

$$\begin{aligned}
 (c\Delta)_{11} &= F F^d E F^\pi (E^d)^2 + F F^d E F^\pi (E^d)^4 F F^\pi + \sum_{i=1}^{\infty} F F^d E F^i F^\pi E^{2i+1} F^\pi E^d \\
 &+ \sum_{i=1}^{\infty} F F^d E F^i F^\pi E^{2i-1} F^\pi E^d F F^\pi, \\
 (c\Delta)_{21} &= F F^\pi (E^d)^2 + F F^\pi (E^d)^4 F F^\pi + \sum_{i=1}^{\infty} F^{i+1} F^\pi E^{2i+1} F^\pi E^d \\
 &+ \sum_{i=1}^{\infty} F^{i+1} F^\pi E^{2i-1} F^\pi E^d F F^\pi, \\
 (c\Delta)_{ij} &= 0, \text{ otherwise;} \\
 (\Delta b)_{12} &= F^\pi (E^d)^2 F^\pi + F^\pi (E^d)^4 F F^\pi + \sum_{i=1}^{\infty} F^i F^\pi E^{2i+1} F^\pi E^d F^\pi \\
 &+ \sum_{i=1}^{\infty} F^i F^\pi E^{2i-1} F^\pi E^d F F^\pi, \\
 (\Delta b)_{ij} &= 0, \text{ otherwise;}
 \end{aligned}$$

$$\begin{aligned}
(\Gamma - \Delta a)_{11} &= F^\pi E^d + F^\pi (E^d)^3 F F^\pi - F^\pi (E^d)^2 F^\pi E - F^\pi (E^d)^4 F F^\pi E \\
&+ \sum_{i=1}^{\infty} F^i F^\pi E^{2i+1} + \sum_{i=1}^{\infty} F^i F^\pi E^{2i} E^d F F^\pi \\
&- \sum_{i=1}^{\infty} F^i F^\pi E^{2i+1} F^\pi E^d F^\pi E - \sum_{i=1}^{\infty} F^i F^\pi E^{2i-1} F^\pi E^d F F^\pi E, \\
(\Gamma - \Delta a)_{ij} &= 0, \text{ otherwise};
\end{aligned}$$

Thus,

$$\begin{aligned}
(\delta)_{12} &= [F^\pi E^d F^\pi + F^\pi (E^d)^3 F F^\pi + F F^\pi (E^d)^3 F^\pi + F F^\pi (E^d)^5 F F^\pi \\
&+ \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+3} F^\pi + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+5} F F^\pi]^2 \\
&+ [F^\pi E E^d + F^\pi (E^d)^2 F F^\pi + F F^\pi (E^d)^2 + F F^\pi (E^d)^4 F F^\pi \\
&- F^\pi E^d F^\pi E - F^\pi (E^d)^3 F F^\pi E - F F^\pi (E^d)^3 F^\pi E - F F^\pi (E^d)^5 F F^\pi E \\
&+ \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+2} + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+4} F F^\pi \\
&- \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+3} F^\pi E - \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+5} F F^\pi E] \\
&[F^\pi (E^d)^2 F^\pi + F^\pi (E^d)^4 F F^\pi + \sum_{i=1}^{\infty} F^i F^\pi E^{2i+1} F^\pi E^d F^\pi \\
&+ \sum_{i=1}^{\infty} F^i F^\pi E^{2i-1} F^\pi E^d F F^\pi], \\
(\delta)_{ij} &= 0, \text{ otherwise.}
\end{aligned}$$

Moreover, we have

$$\begin{aligned}
(\lambda)_{11} &= [F F^d E F^\pi (E^d)^2 + F F^d E F^\pi (E^d)^4 F F^\pi + \sum_{i=1}^{\infty} F F^d E F^i F^\pi E^{2i+1} F^\pi E^d \\
&+ \sum_{i=1}^{\infty} F F^d E F^i F^\pi E^{2i-1} F^\pi E^d F F^\pi] \\
&[F^\pi E E^d + F^\pi (E^d)^2 F F^\pi + F F^\pi (E^d)^2 + F F^\pi (E^d)^4 F F^\pi \\
&+ \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+2} + \sum_{i=1}^{\infty} F^{i+1} F^\pi (E^d)^{2i+4} F F^\pi] \\
&+ [F F^d E F^\pi E^d + F F^d E F^\pi (E^d)^3 F F^\pi - F F^d E F^\pi (E^d)^2 F^\pi E \\
&- F F^d E F^\pi (E^d)^4 F F^\pi E + \sum_{i=1}^{\infty} F F^d E F^i F^\pi E^{2i+1} + \sum_{i=1}^{\infty} F F^d E F^i F^\pi E^{2i} E^d F F^\pi \\
&- \sum_{i=1}^{\infty} F F^d E F^i F^\pi E^{2i+1} F^\pi E^d F^\pi E - \sum_{i=1}^{\infty} F F^d E F^i F^\pi E^{2i-1} F^\pi E^d F F^\pi E] \\
&[F^\pi E^d + F^\pi (E^d)^3 F F^\pi + \sum_{i=1}^{\infty} F^i F^\pi E^{2i+1} + \sum_{i=1}^{\infty} F^i F^\pi E^{2i} E^d F F^\pi]
\end{aligned}$$

## THE G-DRAZIN INVERSES OF ANTI-TRIANGULAR BLOCK OPERATOR MATRICES

$$\begin{aligned}
 (\lambda)_{21} &= [FF^\pi(E^d)^2 + FF^\pi(E^d)^4FF^\pi + \sum_{i=1}^{\infty} F^{i+1}F^\pi E^{2i+1}F^\pi E^d \\
 &+ \sum_{i=1}^{\infty} F^{i+1}F^\pi E^{2i-1}F^\pi E^d FF^\pi] \\
 &\quad [F^\pi EE^d + F^\pi(E^d)^2FF^\pi + FF^\pi(E^d)^2 + FF^\pi(E^d)^4FF^\pi \\
 &+ \sum_{i=1}^{\infty} F^{i+1}F^\pi(E^d)^{2i+2} + \sum_{i=1}^{\infty} F^{i+1}F^\pi(E^d)^{2i+4}FF^\pi] \\
 &+ [FF^\pi E^d + FF^\pi(E^d)^3FF^\pi - FF^\pi(E^d)^2F^\pi E - FF^\pi(E^d)^4FF^\pi E \\
 &+ \sum_{i=1}^{\infty} F^{i+1}F^\pi E^{2i+1} + \sum_{i=1}^{\infty} F^{i+1}F^\pi E^{2i} E^d FF^\pi \\
 &- \sum_{i=1}^{\infty} F^{i+1}F^\pi E^{2i+1}F^\pi E^d FF^\pi E - \sum_{i=1}^{\infty} F^{i+1}F^\pi E^{2i-1}F^\pi E^d FF^\pi E] \\
 &\quad [F^\pi E^d + F^\pi(E^d)^3FF^\pi + \sum_{i=1}^{\infty} F^i F^\pi E^{2i+1} + \sum_{i=1}^{\infty} F^i F^\pi E^{2i} E^d FF^\pi], \\
 &(\lambda)_{ij} = 0, \text{ otherwise.}
 \end{aligned}$$

Furthermore, we get

$$\begin{aligned}
 (\xi)_{12} &= [FF^d EF^\pi(E^d)^2 + FF^d EF^\pi(E^d)^4FF^\pi + \sum_{i=1}^{\infty} FF^d EF^i F^\pi E^{2i+1} F^\pi E^d \\
 &+ \sum_{i=1}^{\infty} FF^d EF^i F^\pi E^{2i-1} F^\pi E^d FF^\pi] \\
 &\quad [F^\pi E^d F^\pi + F^\pi(E^d)^3FF^\pi + FF^\pi(E^d)^3F^\pi + FF^\pi(E^d)^5FF^\pi \\
 &+ \sum_{i=1}^{\infty} F^{i+1}F^\pi(E^d)^{2i+3}F^\pi + \sum_{i=1}^{\infty} F^{i+1}F^\pi(E^d)^{2i+5}FF^\pi] \\
 &+ [FF^d EF^\pi E^d + FF^d EF^\pi(E^d)^3FF^\pi - FF^d EF^\pi(E^d)^2F^\pi E \\
 &- FF^d EF^\pi(E^d)^4FF^\pi E + \sum_{i=1}^{\infty} FF^d EF^i F^\pi E^{2i+1} + \sum_{i=1}^{\infty} FF^d EFFF^i F^\pi E^{2i} E^d FF^\pi \\
 &- \sum_{i=1}^{\infty} FF^d EF^i F^\pi E^{2i+1} F^\pi E^d FF^\pi E - \sum_{i=1}^{\infty} FF^d EF^i F^\pi E^{2i-1} F^\pi E^d FF^\pi E] \\
 &\quad [F^\pi(E^d)^2F^\pi + F^\pi(E^d)^4FF^\pi + \sum_{i=1}^{\infty} F^i F^\pi E^{2i+1} F^\pi E^d F^\pi \\
 &+ \sum_{i=1}^{\infty} F^i F^\pi E^{2i-1} F^\pi E^d FF^\pi] \\
 (\xi)_{22} &= [FF^\pi(E^d)^2 + FF^\pi(E^d)^4FF^\pi + \sum_{i=1}^{\infty} F^{i+1}F^\pi E^{2i+1} F^\pi E^d \\
 &+ \sum_{i=1}^{\infty} F^{i+1}F^\pi E^{2i-1} F^\pi E^d FF^\pi] \\
 &\quad [F^\pi E^d F^\pi + F^\pi(E^d)^3FF^\pi + FF^\pi(E^d)^3F^\pi + FF^\pi(E^d)^5FF^\pi \\
 &+ \sum_{i=1}^{\infty} F^{i+1}F^\pi(E^d)^{2i+3}F^\pi + \sum_{i=1}^{\infty} F^{i+1}F^\pi(E^d)^{2i+5}FF^\pi] \\
 &+ [FF^\pi E^d + F^\pi(E^d)^3FF^\pi - FF^\pi(E^d)^2F^\pi E - F^\pi(E^d)^4FF^\pi E \\
 &+ \sum_{i=1}^{\infty} F^{i+1}F^\pi E^{2i+1} + \sum_{i=1}^{\infty} F^{i+1}F^\pi E^{2i} E^d FF^\pi] \\
 &\quad [F^\pi(E^d)^2F^\pi + F^\pi(E^d)^4FF^\pi + \sum_{i=1}^{\infty} F^i F^\pi E^{2i+1} F^\pi E^d F^\pi \\
 &+ \sum_{i=1}^{\infty} F^i F^\pi E^{2i-1} F^\pi E^d FF^\pi], \\
 &(\xi)_{ij} = 0, \text{ otherwise.}
 \end{aligned}$$

Step 2. We claim that  $Q$  has g-Drazin inverse. We have

$$Q^d = \begin{pmatrix} 0 & 0 \\ 0 & d^d \end{pmatrix}_e \quad \text{and} \quad Q^\pi = \begin{pmatrix} e & 0 \\ 0 & d^\pi \end{pmatrix}_e.$$

Step 3. Since  $bd = 0$ , we have  $PQ = 0$ . Clearly, we have

$$\begin{aligned} Q^\pi P^d &= \begin{pmatrix} \gamma & \delta \\ d^\pi \lambda & d^\pi \xi \end{pmatrix}_e \\ &= \gamma + \delta + d^\pi \lambda + d^\pi \xi \\ &= \begin{pmatrix} \gamma_{11} & \delta_{12} \\ \lambda_{21} & \xi_{22} \end{pmatrix} \end{aligned}$$

By direct computation, we have

$$\begin{aligned} (a+c)(\gamma+\delta) &= \begin{pmatrix} E - FF^d E F F^d & 0 \\ FF^\pi & 0 \end{pmatrix} \begin{pmatrix} \gamma_{11} & \delta_{12} \\ 0 & 0 \end{pmatrix} \\ &= \begin{pmatrix} E\gamma_{11} & E\delta_{12} \\ F\gamma_{11} & F\delta_{12} \end{pmatrix} \end{aligned}$$

and

$$\begin{aligned} b(\gamma+\delta) &= \begin{pmatrix} 0 & F^\pi \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \gamma_{11} & \delta_{12} \\ 0 & 0 \end{pmatrix} \\ &= 0. \end{aligned}$$

Hence,

$$\begin{aligned} PP^d &= \begin{pmatrix} a & b \\ c & 0 \end{pmatrix}_e \begin{pmatrix} \gamma & \delta \\ \lambda & \xi \end{pmatrix}_e \\ &= (a+c)(\gamma+\delta) + b(\lambda+\xi) \\ &= \begin{pmatrix} E\gamma_{11} & E\delta_{12} \\ F\gamma_{11} & F\delta_{12} \end{pmatrix}. \end{aligned}$$

Also we see that  $Q^\pi Q^i = 0$  for any  $i \geq 1$ . In view of [6, Theorem 2.3], we have

$$\begin{aligned} M^d &= Q^\pi \sum_{i=0}^{\infty} Q^i (P^d)^{i+1} + \sum_{i=0}^{\infty} (Q^d)^{i+1} P^i P^\pi \\ &= \begin{pmatrix} \Theta & \Psi \\ \Phi & \Omega \end{pmatrix} + \sum_{i=0}^{\infty} \begin{pmatrix} 0 & F^d \\ FF^d & -EF^d \end{pmatrix}^{i+1} \\ &\quad \begin{pmatrix} F^\pi E + FF^d EF^\pi & F^\pi \\ FF^\pi & 0 \end{pmatrix}^i \begin{pmatrix} I - E\Theta & -E\Psi \\ -F\Theta & I - F\Psi \end{pmatrix}, \end{aligned}$$

where  $\Theta = \gamma_{11}$ ,  $\Psi = \delta_{12}$ ,  $\Phi = \lambda_{21}$ ,  $\Omega = \xi_{22}$ , as asserted.  $\square$



**Corollary 2.7.** *Let  $E$  and  $F$  have  $g$ -Drazin inverses. If  $F^\pi EF^2 = 0$  and  $F^\pi EFE = 0$ , then  $M = \begin{pmatrix} E & F \\ I & 0 \end{pmatrix}$  has  $g$ -Drazin inverse.*

*In this case,*

$$M^d = \begin{pmatrix} E\Psi + \Omega & E\Theta - E\Psi E + \Phi - \Omega E \\ \Psi & \Theta - \Psi E \end{pmatrix} + \sum_{i=0}^{\infty} \begin{pmatrix} FF^d & 0 \\ 0 & F^d \end{pmatrix} \begin{pmatrix} 0 & F^d \\ FF^d & -EF^d \end{pmatrix}^i \begin{pmatrix} F^\pi E + FF^d EF^\pi & F^\pi \\ FF^\pi & 0 \end{pmatrix}^i \begin{pmatrix} -E\Psi & I - E\Theta + E\Psi E \\ I - F\Psi & -E - F\Theta + F\Psi E \end{pmatrix},$$

where  $\Theta, \Psi, \Phi, \Omega$  as given in (\*\*).

*Proof.* Obviously, we have

$$\begin{pmatrix} E & F \\ I & 0 \end{pmatrix} = \begin{pmatrix} 0 & I \\ I & -E \end{pmatrix}^{-1} \begin{pmatrix} E & I \\ F & 0 \end{pmatrix} \begin{pmatrix} 0 & I \\ I & -E \end{pmatrix},$$

and so

$$\begin{pmatrix} E & F \\ I & 0 \end{pmatrix}^d = \begin{pmatrix} E & I \\ I & 0 \end{pmatrix} \begin{pmatrix} E & I \\ F & 0 \end{pmatrix}^d \begin{pmatrix} 0 & I \\ I & -E \end{pmatrix}.$$

Applying Theorem 2.6 to the matrix  $\begin{pmatrix} E & I \\ F & 0 \end{pmatrix}$ , we obtain the result. □

#### ACKNOWLEDGEMENT

The authors would like to thank the referee for his/her helpful suggestions for the improvement of this paper.

#### REFERENCES

- [1] C. Bu; K. Zhang and J. Zhao, Representation of the Drazin inverse on solution of a class singular differential equations, *Linear & Multilinear Algebra*, **59**(2011), 863-877.
- [2] S.L. Campbell, The Drazin inverse and systems of second order linear differential equations, *Linear & Multilinear Algebra*, **14**(1983), 195-198.
- [3] N. Castro-González and E. Dopazo, Representations of the Drazin inverse for a class of block matrices, *Linear Algebra Appl.*, **400**(2005), 253-269.

- [4] D.S. Cvetković-Ilić, Some results on the  $(2, 2, 0)$  Drazin inverse problem, *Linear Algebra Appl.*, **438**(2013), 4726-4741.
- [5] C. Deng and Y. Wei, A note on the Drazin inverse of an anti-triangular matrix, *Linear Algebra Appl.*, **431**(2009), 1910-1922.
- [6] D.S. Djordjevic and Y. Wei, Additive results for the generalized Drazin inverse, *J. Austral. Math. Soc.*, **73**(2002), 115-125.
- [7] Y. Liao, J. Chen and J. Cui, Cline's formula for the generalized Drazin inverse, *Bull. Malays. Math. Sci. Soc.*, **37**(2014), 37-42.
- [8] D. Mosić, A note on Cline's formula for the generalized Drazin inverse, *Linear & Multilinear Algebra*, **63**(2014), 1106-1110.
- [9] P. Patricio and R.E. Hartwig, The  $(2,2,0)$  Drazin inverse problem, *Linear Algebra Appl.*, **437**(2012), 2755-2772.
- [10] H. Yang and X. Liu, The Drazin inverse of the sum of two matrices and its applications, *J. Comput. Appl. Math.*, **235**(2011), 1412-1417.
- [11] D. Zhang, Representations for generalized Drazin inverse of operator matrices over a Banach space, *Turk. J. Math.*, **40**(2016), 428-437.
- [12] D. Zhang and D. Mosić, Explicit formulae for the generalized Drazin inverse of block matrices over a Banach algebra, *Filomat*, **32**(2018), 5907-5917.

DEPARTMENT OF MATHEMATICS, HANGZHOU NORMAL UNIVERSITY, HANG  
-ZHOU, CHINA

*Email address:* <huanyinchen@aliyun.com>

WOMEN'S UNIVERSITY OF SEMNAN (FARZANEGAN), SEMNAN, IRAN

*Email address:* <sheibani@fgusem.ac.ir>