

A LEBESGUE–LUSIN PROPERTY FOR GRADIENTS OF FIRST-ORDER LINEAR OPERATORS

ADOLFO ARROYO-RABASA

ABSTRACT. We prove that for a first-order homogeneous linear partial differential operator \mathcal{A} and a map f taking values in the essential range of the operator, there exists a function u of special bounded variation satisfying

$$\mathcal{A}u(x) = f(x) \quad \text{almost everywhere.}$$

This extends a result of G. Alberti for gradients on \mathbf{R}^N . In particular, for $m < N$, it is shown that every integrable m -form field on \mathbf{R}^N is the absolutely continuous part of the boundary of some locally normal current with finite mass.

1. INTRODUCTION

We consider a general constant-coefficient system of linear partial differential equations acting on functions of \mathbf{R}^N -variables. That is a partial differential operator of the form

$$\mathcal{A}u = \sum_{|\alpha|=k}^N A_\alpha \partial^\alpha u, \quad A_\alpha \in \text{Hom}(E; F), \quad (1)$$

where E, F are finite-dimensional \mathbf{R} -spaces. Here $\alpha = (\alpha_1, \dots, \alpha_N) \in (\mathbb{N}_0)^N$ is a multi-index with modulus $|\alpha| = \alpha_1 + \dots + \alpha_N$ and $\partial^\alpha = \partial_1^{\alpha_1} \circ \dots \circ \partial_N^{\alpha_N}$ is the composition of distributional directional partial derivatives. The Fourier transform establishes a one-to-one correspondence between homogeneous operators and their associated principal symbol $\mathbb{A} : \mathbf{R}^N \rightarrow \text{Hom}(V; W)$, which in this context is given by the k -homogeneous tensor-valued polynomial

$$\mathbb{A}(\xi) = \sum_{|\alpha|=k} A_\alpha \xi^\alpha, \quad \xi^\alpha = \xi_1^{\alpha_1} \cdots \xi_N^{\alpha_N}, \quad \xi \in \mathbf{R}^N.$$

Suppose now that $u \in \mathcal{D}'(\Omega; E)$ is a distribution such that $\mathcal{A}u$ is a zero-order distribution, i.e., represented by an F -valued Radon measure. Let $g : \mathbf{R}^N \rightarrow F$ be the polar vector of $\mathcal{A}u$. If additionally \mathcal{A} is a constant-rank operator (see [5, 7, 8]), the De Philippis–Rindler theorem [6] establishes that g is constrained to take values on the *image cone* at *singular points*. More precisely,

$$g(x) \in \mathcal{I}_\mathcal{A} := \bigcup_{|\zeta|=1} \text{Im } \mathbb{A}(\zeta) \quad |\mathcal{A}u| \text{ almost everywhere on } U,$$

for all Borel sets $U \subset \mathbf{R}^N$ with $\mathcal{L}_N(U) = 0$. This property extends a classical result of Alberti [1], which says that if the distributional gradient Du of a map $u : \mathbf{R}^N \rightarrow \mathbf{R}^M$ is represented by an $\mathbf{R}^{M \times N}$ -valued Radon measure, then its polar vector must take values in the cone of rank-one matrices at singular points:

$$\text{rank} \frac{Du}{|Du|}(x) = 1 \quad |Du| \text{ almost everywhere in } U.$$

In counterpart with this restriction over the polar of Du at singular points, Alberti showed in [2] that the absolutely continuous part of a gradient measure is fully unconstrained: if $f \in L^1(\mathbf{R}^N; \mathbf{R}^{M \times N})$, then there exists $u \in BV(\mathbf{R}^N; \mathbf{R}^M)$ such that ⁽¹⁾

$$D^a u = f \mathcal{L}_N.$$

In the context of the De Philippis–Rindler theorem, this begs the answer to the following natural question: *is the absolutely continuous part of an \mathcal{A} -gradient measure fully unconstrained?* Theorem 1 states that, at least when \mathcal{A} is a first-order operator, this is indeed the case.

2. NOTATION AND RESULTS

In all that follows, Ω will denote an open subset of \mathbf{R}^N . We will write \mathcal{L}_N to denote the N dimensional Lebesgue measure and \mathcal{H}_{N-1} to denote the $(N-1)$ dimensional Hausdorff outer measure on \mathbf{R}^N . The space $BV(\Omega; E)$, of E -valued vector fields with bounded variation on Ω , consists of all maps $u : E \rightarrow \Omega$ whose distributional gradient can be represented by a bounded Radon measure. It is well known (see, e.g., [3]) that such maps are Lebesgue continuous outside of a countably \mathcal{H}_{N-1} rectifiable set $J_u \subset \Omega$, called the jump set of u . Moreover, the map u has Lebesgue one-sided limits u^+, u^- across any measure theoretic orientation ν_u of J_u . We define

$$\mathbf{F} := \overline{\{ \mathcal{A}u(x) : u \in C_c^\infty(\mathbf{R}^N; E) \}}^F, \quad (2)$$

the *essential range* of the operator \mathcal{A} . It is easy to see that every distribution $\mathcal{A}u$ takes values in this space, and it is the smallest space with this property. With these considerations in mind, we can state the main result:

Theorem 1. *Let \mathcal{A} be a first-order operator as in (1) and let $f : \Omega \rightarrow \mathbf{F}$ be integrable. Then there exists a map $u \in BV(\Omega; E)$ satisfying*

$$\mathcal{A}u = f \mathcal{L}_N + \mathbb{A}(\nu_u)[u^+ - u^-] \mathcal{H}_{N-1} \quad (3)$$

$$|Du|(\Omega) \leq C \|f\|_{L^1(\Omega)} \quad (4)$$

where C is a constant that only depends on \mathcal{A} .

¹Alberti's result is written originally for functions $f : \Omega \rightarrow \mathbf{R}$. Applying it on each component yields the general case for \mathbf{R}^M -vector fields.

Remark 1 (Second-order operators). A similar statement holds for second-order operators ($k = 2$) with $u \in BV^2$ instead. The proof will be part of a forthcoming paper where we will consider higher-order operators.

Remark 2 (Fourier coefficients). The essential range \mathbf{F} defined in (2) coincides with the space $\text{span} \{ \mathbb{A}(\xi)[e] : \xi \in \mathbf{R}^N, e \in E \} = \text{span} \{ \mathcal{I}_{\mathcal{A}} \}$ of all \mathcal{A} -gradient amplitudes in Fourier space (for a proof see [4, Sec. 2.5]).

Theorem 1 is a corollary of the following equivalence result and Alberti’s original result for gradients. In stating it, it will be convenient to introduce the following short-hand notation

$$P \bullet v = P \underbrace{v \otimes \dots \otimes v}_{(k-1) \text{ times}}, \quad P \in E \otimes \mathbf{R}^{N^k}, v \in \mathbf{R}^N$$

for $k > 1$ and simply

$$P \bullet v = P$$

for $k = 1$.

Proposition 2. *The following are equivalent:*

(a) *Let \mathcal{A} be a k^{th} order linear operator as in (1) and let $f : \Omega \rightarrow \mathbf{F}$ be an integrable map. Then there exists $u \in BV^k(\Omega; E)$ satisfying*

$$\mathcal{A}u = f \mathcal{L}_N + \mathbb{A}(\nu_w)[(w^+ - w^-) \bullet \nu_w] \mathcal{H}_{N-1}$$

$$|D^k f|(\Omega) \leq C \|f\|_{L^1(\Omega)}$$

for some constant C . Here $w = \nabla^{k-1}u$.

(b) *Let $f : \Omega \rightarrow \mathbf{R}_{\text{sym}}^{N^k}$ be an integrable map. Then there exists $u \in BV^k(\Omega)$ satisfying*

$$D^k u = f \mathcal{L}_N + [w^+ - w^-] \otimes \nu_w \mathcal{H}_{N-1}$$

$$\int_{J_u} |w^+ - w^-| d\mathcal{H}_{N-1} \leq C \|f\|_{L^1(\Omega)}$$

for some constant C . Here $w = \nabla^{k-1}u$.

Proof of Proposition 2. The implication (a) \implies (b) is obvious, given that the principal symbol of the k th order Hessian is given by

$$D^k(\xi) = \otimes^k \xi.$$

Therefore the gradient has the form (1) by setting $A_\alpha = M_\alpha$, where the family $\{M_\alpha : |\alpha| = k\}$ is the canonical orthonormal basis of $\mathbf{R}_{\text{sym}}^{N^k}$.

We now show (b) \implies (a). The first step is to use the alternative expression $A[D^k u] = \mathcal{A}u$, where $A : E \otimes \mathbf{R}_{\text{sym}}^{N^k} \rightarrow F$ is the (unique) linear map satisfying

$$A[e \otimes^k \xi] = \mathbb{A}^k(\xi)[e] \quad \text{for all } e \in E. \quad (5)$$

The existence of A is a direct consequence of the universal property of the tensor product and the k -linearity of the principal symbol in the frequency variable ξ . For the sake of simplicity, let us write $\mathbf{E}_k = E \otimes \mathbf{R}_{\text{sym}}^{N^k}$. By construction, we have

$$\text{Im } A = \text{span} \{ \mathbb{A}(\xi)[e] : \xi \in \mathbf{R}^N, e \in E \} = \mathbf{F}.$$

Let us consider the Moore-Penrose quasi-inverse A^\dagger associated with A . This is an element of $\text{Hom}(\mathbf{F}; \mathbf{E}_k)$ satisfying the fundamental identity

$$A \circ A^\dagger = \mathbf{1}_{\text{Im } A} = \mathbf{1}_{\mathbf{F}}. \quad (6)$$

In particular, by our assumption on f , $A^\dagger f(x)$ is well-defined almost everywhere on Ω and $A^\dagger f \in L^1(\Omega; \mathbf{E}_k)$. Now, we make use of the assumption (b) over each E -coordinate to find $u \in BV^k(\Omega; E)$ satisfying

$$D^k u = A^\dagger f \mathcal{L}_N + D^j u, \quad w = \nabla^{k-1} u,$$

where as usual $D^j w = \tilde{g} \mathcal{H}^{N-1} = [w^+ - w^-] \otimes \nu_U \mathcal{H}_{N-1}$ is the jump part of Dw . Pre-composing this expression with A , we arrive at the expression

$$\mathcal{A}u = AD^k u = A \left[A^\dagger f \mathcal{L}_N + \tilde{g} \mathcal{H}^{N-1} \right].$$

Since A is a linear map, we may pull in and distribute A into the densities that belong to the sum of the right-hand side, hence concluding that

$$\mathcal{A}u = A \circ A^\dagger f \mathcal{L}_N + A \tilde{g} \mathcal{H}^{N-1} \stackrel{(5),(6)}{=} f \mathcal{L}_N + g \mathcal{H}^{N-1},$$

where $g = \mathbf{1}_{J_w} \cdot A([w^+ - w^-] \otimes \nu_w)$. In passing to the last equality in the formula above, we have used the almost everywhere pointwise restriction $f(x) \in \mathbf{F}$. Now, we use the fact that at jump points, it holds

$$w^- - w^+ = ([w^- - w^+] \bullet \nu_w) \otimes^k \nu_w.$$

This follows from the curl operator's compatibility conditions on symmetric k^{th} order tensors. From this we obtain that $g = \mathbf{1}_{J_w} \cdot \mathbb{A}(\nu_w)[(w^+ - w^-) \bullet \nu]u$ as desired. Lastly, the bound on the total variation of $D^k u$ follows from the estimates

$$\begin{aligned} \int |A^\dagger f| dx &\leq \|A^\dagger\|_{\mathbf{F} \rightarrow \mathbf{E}_k} \int |f| dx, \\ \int |g| d\mathcal{H}_{N-1} &\leq \|A\|_{\mathbf{E}_k \rightarrow \mathbf{F}} \int |\tilde{g}| d\mathcal{H}_{N-1} \end{aligned}$$

and the fact that $\|A^\dagger\|_{\mathbf{E}_k \rightarrow \mathbf{F}} \leq (\|A\|_{\mathbf{E}_k \rightarrow \mathbf{F}})^{-1}$. This finishes the proof. \square

Proof of Theorem 1. The proof of Theorem 1 follows directly from the validity of the Lusin property for gradients ([2, Thm. 3]), the previous proposition (with $k = 1$), and the equivalence in Remark 3. \square

An application for normal currents. Let m be a non-negative integer. An m -dimensional current T on Ω is an element of the continuous dual $\mathcal{D}_m(\Omega) = \mathcal{D}^m(\Omega)^*$, where

$$\mathcal{D}^m(\Omega) := C_c^\infty(\Omega; \wedge^m \mathbf{R}^N)$$

is the space of smooth and compactly supported m -vector fields on Ω . The (distributional) boundary operator on currents is defined by duality through the rule

$$\partial T(\varphi) = T(d\varphi) \quad \varphi \in \mathcal{D}^m(\Omega)$$

where d is the exterior derivative operator acting on $\mathcal{D}^m(\Omega)$. The boundary operator then maps $\mathcal{D}_m(\Omega)$ into $\mathcal{D}_{m-1}(\Omega)$. The mass of a current $T \in \mathcal{D}_m(\Omega)$ is defined as

$$\mathbf{M}(T) = \sup \left\{ T(\varphi) : \varphi \in \mathcal{D}^m(\Omega), \sup_{x \in \Omega} |\varphi(x)| \leq 1 \right\}.$$

A current T is called *locally normal* if T is representable by a Radon measure and either ∂T is representable by a Radon measure or $m = 0$. The space of locally normal m -dimensional currents on Ω is denoted by $\mathbf{N}_m^{\text{loc}}(\Omega)$.

Corollary 3. *Let $m \in [0, N)$ be an integer and let $S : \Omega \rightarrow \wedge^m \mathbf{R}^N$ be an integrable m -form. Then, there exists a normal current $T \in \mathbf{N}_{m+1}^{\text{loc}}(\Omega)$, with bounded variation and finite mass on Ω , such that*

$$\partial T = S \mathcal{L}_N + (T^+ - T^-) \lrcorner \nu_u \mathcal{H}^{N-1} \llcorner J_T.$$

Moreover,

$$|DT|(\Omega) \leq C_{m,N} \|S\|_{L^1(\Omega)}.$$

Proof. Since the symbol of the exterior derivative is precisely the exterior multiplication (i.e., $d(\xi)a = a \wedge \xi$), the interior product is then precisely the map associated with the symbol of the boundary operator on m -vectors:

$$\partial(\xi)e = \omega \lrcorner \xi, \quad \xi \in \mathbf{R}^N, \omega \in \wedge^m \mathbf{R}^N.$$

In particular, the boundary operator ∂ defines a constant-coefficient first-order operator from $\mathcal{D}^{m+1}(\Omega)$ to $\mathcal{D}^m(\Omega)$. Moreover, since $\text{Im } \partial(\xi) = \wedge^m \mathbf{R}^N$ for all nonzero ξ , it follows that $\wedge^m \mathbf{R}^N = \mathbf{F}_\partial$. We may thus apply Theorem 1 to find $T \in BV_{\text{loc}}(\Omega; \wedge^m \mathbf{R}^N)$ satisfying the desired properties. The fact that $T \in \mathbf{N}_{m+1}^{\text{loc}}(\Omega)$ follows directly from the fact that ∂ is a first-order operator and that T has bounded variation on Ω . \square

Corollary 4. *Let S be as above. There exists a countably \mathcal{H}_{N-1} rectifiable set J with orientation ν , and a Borel $(m+1)$ -form*

$$g : J \rightarrow \wedge^{m+1} \mathbf{R}^N$$

satisfying

$$\partial S = \partial(g \lrcorner \nu_u \mathcal{H}_{N-1} \llcorner J)$$

in the sense of distributions.

Divergence-free fields. Another exciting application of Theorem 1 is the following *rectifiable completion* for systems of vector fields to systems of solenoidal measures.

Corollary 5. *Every vector field $f \in L^1(\Omega; \mathbf{R}^N)$ is the absolutely continuous part of a solenoidal field measure up to a rectifiable measure. More precisely, there exists a countably \mathcal{H}_{N-1} rectifiable set $R \subset \Omega$ and a Borel vector field $g : R \rightarrow \mathbf{R}^N$ satisfying*

$$\operatorname{div}(f \mathcal{L}_N + g \mathcal{H}^{N-1} \llcorner R) = 0 \quad g(x) \in \operatorname{Tan}(R, x) \quad \mathcal{H}_{N-1} \text{ a.e.}$$

and such that

$$\int_R |g| d\mathcal{H}_{N-1} \leq C \|f\|_{L^1(\Omega)}.$$

for a constant C that depends only on N .

Proof. Consider the first-order operator

$$\operatorname{curl}^* V = \partial_j V_{ij} \quad V : \Omega \rightarrow \mathbf{R}_{\text{skew}}^{N^2},$$

whose associated symbol is given by

$$\operatorname{curl}^*(\xi)[P] = P\xi \quad V \in \mathbf{R}_{\text{skew}}^{N^2}.$$

This defines an operator from $\mathbf{R}_{\text{skew}}^{N^2}$ to \mathbf{R}^N . Notice that $\mathbf{F}_{\operatorname{curl}^*} = \mathbf{R}^N$ so that the essential pointwise restriction is trivial in this case. By Theorem 1 we may find $V \in BV(\Omega; \mathbf{R}_{\text{skew}}^{N^2})$ satisfying

$$\mu = \operatorname{curl}^* V = f \mathcal{L}_N + [V^+ - V^-] \nu_V \mathcal{H}_{N-1} \llcorner J_V$$

and

$$|DV|(\Omega) \leq C \|f\|_{L^1(\Omega)}.$$

for some constant C depending only on N (since the symbol of the divergence operator on \mathbf{R}^N depends only on the dimension N). In particular, Since V is a skew-symmetric matrix field, it follows that

$$[V^+ - V^-] \nu_V \cdot \nu_V = 0 \quad \implies \quad \frac{\mu}{|\mu^s|} \in \operatorname{Tan}(J_V, x) \quad \mathcal{H}_{N-1} \text{ almost everywhere.}$$

Moreover,

$$\operatorname{div} \mu = \operatorname{div} \circ \operatorname{curl}^* V = \sum_{i,j=1}^N \partial_{ij} V_{ij} = 0.$$

Setting $R = J_V$ and $g = \mathbf{1}_{J_V} \cdot [V^+ - V^-] \nu_V$ yields the sought assertions. \square

Acknowledgments. I would like to thank Giacomo del Nin and Guido De Philippis for their fascinating insight and comments about the problem discussed in this note.

REFERENCES

- [1] G. Alberti, *Rank one property for derivatives of functions with bounded variation*, Proc. Roy. Soc. Edinburgh Sect. A **123** (1993), no. 2, 239–274, DOI 10.1017/S030821050002566X. MR1215412
 - [2] ———, *A Lusin type theorem for gradients*, J. Funct. Anal. **100** (1991), no. 1, 110–118, DOI 10.1016/0022-1236(91)90104-D. MR1124295
 - [3] L. Ambrosio, N. Fusco, and D. Pallara, *Functions of bounded variation and free discontinuity problems*, Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, New York, 2000. MR1857292
 - [4] A. Arroyo-Rabasa, G. De Philippis, and F. Rindler, *Lower semicontinuity and relaxation of linear-growth integral functionals under PDE constraints*, Adv. Calc. Var. **13** (2020), no. 3, 219–255, DOI 10.1515/acv-2017-0003. MR4116615
 - [5] A. Arroyo-Rabasa and J. Simental, *An elementary approach to the homological properties of constant-rank operators*, arXiv, 2021.
 - [6] G. De Philippis and F. Rindler, *On the structure of \mathcal{A} -free measures and applications*, Ann. of Math. (2) **184** (2016), no. 3, 1017–1039, DOI 10.4007/annals.2016.184.3.10. MR3549629
 - [7] F. Murat, *Compacité par compensation: condition nécessaire et suffisante de continuité faible sous une hypothèse de rang constant*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **8** (1981), no. 1, 69–102 (French). MR616901
 - [8] B. Raiță, *Potentials for \mathcal{A} -quasiconvexity*, Calc. Var. Partial Differential Equations **58** (2019), no. 3, Paper No. 105, 16, DOI 10.1007/s00526-019-1544-x. MR3958799
- Email address:* adolforabasa@gmail.com

UNIVERSITÉ CATHOLIQUE DE LOUVAIN, BELGIUM