

Stochastic integrability of heat-kernel bounds for random walks in a balanced random environment

Xiaoqin Guo ^{*1} and Hung V. Tran ^{†2}

¹ Department of Mathematical Sciences, University of Cincinnati , 2815 Commons Way, Cincinnati, OH 45221, USA

² Department of Mathematics, University of Wisconsin Madison, 480 Lincoln Drive, Madison, WI 53706, USA

September 29, 2022

Abstract

We consider random walks in a balanced i.i.d. random environment in \mathbb{Z}^d for $d \geq 2$ and the corresponding discrete non-divergence form difference operators. We first obtain an exponential integrability of the heat kernel bounds. We then prove the optimal diffusive decay of the semigroup generated by the heat kernel for $d \geq 3$. As a consequence, we deduce a functional central limit theorem for the environment viewed from the particle.

1 Introduction

In this article we consider random walks in a balanced i.i.d. random environment in \mathbb{Z}^d for $d \geq 2$.

1.1 Settings

Let $\mathbb{S}_{d \times d}$ be the set of $d \times d$ positive-definite diagonal matrices. A map

$$\omega : \mathbb{Z}^d \rightarrow \mathbb{S}_{d \times d}$$

2010 *Mathematics subject classification.* 35J15 35J25 35K10 35K20 60G50 60J65 60K37 74Q20 76M50.

Key words and phrases. random walk in a balanced random environment; non-divergence form difference operators; stochastic integrability of heat-kernel bounds; optimal diffusive decay; functional central limit theorem; quantitative stochastic homogenization .

^{*}The work of XG is supported by Simons Foundation through Collaboration Grant for Mathematicians #852943.

[†]HT is supported in part by NSF CAREER grant DMS-1843320 and a Vilas Faculty Early-Career Investigator Award.

is called an *environment*. Denote the set of all environments by Ω and let \mathbb{P} be a probability measure on Ω so that

$$\{\omega(x) = \text{diag}[\omega_1(x), \dots, \omega_d(x)], x \in \mathbb{Z}^d\}$$

are i.i.d. under \mathbb{P} . Expectation with respect to \mathbb{P} is denoted by \mathbb{E} or $E_{\mathbb{P}}$.

Let $\{e_1, \dots, e_d\}$ be the canonical basis for \mathbb{R}^d . For a given function $u : \mathbb{Z}^d \rightarrow \mathbb{R}$ and $\omega \in \Omega$, define the non-divergence form difference operator

$$\text{tr}(\omega(x)\nabla^2 u) = \sum_{i=1}^d \omega_i(x)[u(x + e_i) + u(x - e_i) - 2u(x)], \quad (1)$$

where $\nabla^2 = \text{diag}[\nabla_1^2, \dots, \nabla_d^2]$, and $\nabla_i^2 u(x) = u(x + e_i) + u(x - e_i) - 2u(x)$.

For $r > 0$, $y \in \mathbb{R}^d$ we let

$$\mathbb{B}_r(y) = \{x \in \mathbb{R}^d : |x - y| < r\}, \quad B_r(y) = \mathbb{B}_r(y) \cap \mathbb{Z}^d$$

denote the continuous and discrete balls with center y and radius r , respectively. When $y = 0$, we also write $\mathbb{B}_r = \mathbb{B}_r(0)$ and $B_r = B_r(0)$. For any $B \subset \mathbb{Z}^d$, its *discrete boundary* is the set

$$\partial B := \{z \in \mathbb{Z}^d \setminus B : \text{dist}(z, x) = 1 \text{ for some } x \in B\}.$$

Let $\bar{B} = B \cup \partial B$. Note that with abuse of notation, whenever confusion does not occur, we also use ∂A and \bar{A} to denote the usual continuous boundary and closure of $A \subset \mathbb{R}^d$, respectively.

For $x \in \mathbb{Z}^d$, a *spatial shift* $\theta_x : \Omega \rightarrow \Omega$ is defined by

$$(\theta_x \omega)(\cdot) = \omega(x + \cdot).$$

In a random environment $\omega \in \Omega$, we consider the discrete elliptic Dirichlet problem

$$\begin{cases} \frac{1}{2} \text{tr}(\omega \nabla^2 u(x)) = \frac{1}{R^2} f\left(\frac{x}{R}\right) \psi(\theta_x \omega) & x \in B_R, \\ u(x) = g\left(\frac{x}{|x|}\right) & x \in \partial B_R, \end{cases} \quad (2)$$

where $f \in \mathbb{R}^{\mathbb{B}_1}$, $g \in \mathbb{R}^{\partial \mathbb{B}_1}$ are functions with nice enough regularity properties and $\psi \in \mathbb{R}^{\Omega}$ is bounded and satisfies suitable measurability conditions. Stochastic homogenization studies (for \mathbb{P} -almost all ω) the convergence of u to the solution \bar{u} of a deterministic *effective equation*

$$\begin{cases} \frac{1}{2} \text{tr}(\bar{a} D^2 \bar{u}) = f \bar{\psi} & \text{in } \mathbb{B}_1, \\ \bar{u} = g & \text{on } \partial \mathbb{B}_1, \end{cases} \quad (3)$$

as $R \rightarrow \infty$. Here $D^2 \bar{u}$ denotes the Hessian matrix of \bar{u} and $\bar{a} = \bar{a}(\mathbb{P}) \in \mathbb{S}_{d \times d}$ and $\bar{\psi} = \bar{\psi}(\mathbb{P}, \psi) \in \mathbb{R}$ are *deterministic* and do not depend on the realization of the random environment (see the statement of Proposition C for formulas for \bar{a} and $\bar{\psi}$).

The difference equation (2) is used to describe random walks in a random environment (RWRE) in \mathbb{Z}^d . To be specific, we set

$$\omega(x, x \pm e_i) := \frac{\omega_i(x)}{2\text{tr}\omega(x)} \quad \text{for } i = 1, \dots, d, \quad (4)$$

and $\omega(x, y) = 0$ if $|x - y| \neq 1$. Namely, we normalize ω to get a transition probability. We remark that the configuration of $\{\omega(x, y) : x, y \in \mathbb{Z}^d\}$ is also called a *balanced environment* in the literature [26, 22, 6].

Definition 1. For a fixed $\omega \in \Omega$, the random walk $(X_n)_{n \geq 0}$ in the environment ω is a Markov chain in \mathbb{Z}^d with transition probability P_ω^x specified by

$$P_\omega^x(X_{n+1} = z | X_n = y) = \omega(y, z). \quad (5)$$

The expectation with respect to P_ω^x is written as E_ω^x . When the starting point of the random walk is 0, we sometimes omit the superscript and simply write P_ω^0, E_ω^0 as P_ω and E_ω , respectively. Notice that for random walks (X_n) in an environment ω ,

$$\bar{\omega}^i = \theta_{X_i} \omega \in \Omega, \quad i \geq 0, \quad (6)$$

is also a Markov chain, called the *environment viewed from the particle* process. With abuse of notation, we enlarge our probability space so that P_ω still denotes the joint law of the random walks and $(\bar{\omega}^i)_{i \geq 0}$.

We also consider the *continuous-time* RWRE (Y_t) on \mathbb{Z}^d .

Definition 2. Let $(Y_t)_{t \geq 0}$ be the Markov process on \mathbb{Z}^d with generator

$$L_\omega u(x) = \sum_y \omega(x, y)[u(y) - u(x)] = \frac{1}{2\text{tr}\omega(x)} \text{tr}(\omega(x) \nabla^2 u). \quad (7)$$

With abuse of notation, we also let P_ω^x denote the quenched law of (Y_t) . If there is no ambiguity from the context, we also write, for $x, y \in \mathbb{Z}^d, n \in \mathbb{Z}, t \in \mathbb{R}$, the transition kernels of the discrete and continuous time walks as

$$p_n^\omega(x, y) = P_\omega^x(X_n = y), \quad \text{and} \quad p_t^\omega(x, y) = P_\omega^x(Y_t = y),$$

respectively.

Both the discrete- and continuous-time RWRE share the same trajectory, and their behavior are very much the same. The solutions to the Dirichlet problem can be characterized using the discrete-time RWRE, whereas for the transition kernels it is easier to manipulate the continuous-time case where the derivatives have less cumbersome notation compared to theirs discrete counterparts. Hence we will use both (X_n) and (Y_t) in our paper for convenience.

1.2 Main assumptions

We assume the following points throughout the paper.

- (A1) $\{\omega(x), x \in \mathbb{Z}^d\}$ are i.i.d. under the probability measure \mathbb{P} .
- (A2) $\frac{\omega}{\text{tr}\omega} \geq 2\kappa I$ for \mathbb{P} -almost every ω and some constant $\kappa \in (0, \frac{1}{2d}]$.
- (A3) ψ is a measurable function of the environment with the property that $\{\psi(\theta_x\omega) : x \in \mathbb{Z}^d\}$ are i.i.d. under \mathbb{P} .

In this paper, we use c, C to denote positive constants which may change from line to line but that only depend on the dimension d and the ellipticity constant κ unless otherwise stated. We write $A \lesssim B$ if $A \leq CB$, and $A \asymp B$ if both $A \lesssim B$ and $A \gtrsim B$ hold.

1.3 Earlier results in the literature

The following quenched central limit theorem (QCLT) was proved by Lawler [26], which is a discrete version of Papanicolaou, Varadhan [29].

Theorem A. *Assume (A2) and that law \mathbb{P} of the environment is ergodic under spatial shifts $\{\theta_x : x \in \mathbb{Z}^d\}$. Then*

- (i) *There exists a probability measure $\mathbb{Q} \approx \mathbb{P}$ such that $(\bar{\omega}^i)_{i \geq 0}$ is an ergodic (with respect to time shifts) sequence under law $\mathbb{Q} \times P_\omega$.*
- (ii) *For \mathbb{P} -almost every ω , the rescaled path X_{n^2t}/n converges weakly (under law P_ω) to a Brownian motion with covariance matrix $\bar{a} = E_{\mathbb{Q}}[\omega/\text{tr}\omega] > 0$.*

QCLT for the balanced RWRE in static environments under weaker ellipticity assumptions can be found at [22, 6]. For dynamic balanced random environment, QCLT was established in [14] and finer results concerning the local limit theorem and heat kernel estimates was obtained at [13]. We refer to Zeitouni's lecture notes [31] for a comprehensive account of results and challenges in RWRE.

Denote the Radon-Nikodym derivative of \mathbb{Q} with respect to \mathbb{P} as

$$\rho(\omega) = d\mathbb{Q}/d\mathbb{P}. \quad (8)$$

For any $x \in \mathbb{Z}^d$ and finite set $A \subset \mathbb{Z}^d$, we write

$$\rho_\omega(x) := \rho(\theta_x\omega) \quad \text{and} \quad \rho_\omega(A) = \sum_{x \in A} \rho_\omega(x).$$

It is known that (see, e.g., [13]) for \mathbb{P} -almost all ω , the measure $\rho_\omega(\cdot)$ on \mathbb{Z}^d is the unique (up to a multiplicative constant) invariant measure for the RWRE $(X_n)_{n \geq 0}$. In this sense, \mathbb{Q} is the *steady state* for the environmental process. To investigate the long term behavior of the RWRE and the homogenization of the corresponding diffusion equations, it is crucial to characterize the invariant measure ρ_ω .

Now that \mathbb{Q} is the limiting ergodic measure of the environment, it is expected that, as $t \rightarrow \infty$, $\psi(\bar{\omega}_t) \rightarrow E_{\mathbb{Q}}[\psi]$ almost surely for $\psi \in L^1(\mathbb{P})$. (Recall the process $\bar{\omega}_t$ of the environment as viewed from the particle in (6).) When the balanced environment satisfies a finite range of dependence and ψ is an $L^\infty(\mathbb{P})$ local function, it is shown in [19, Theorem 1.2] that, with overwhelming $\mathbb{P} \otimes P_\omega$ -probability, the average $t^{-1} \int_0^t \psi(\omega_s) ds$ converges to $E_{\mathbb{Q}}[\psi]$ at an algebraic speed $t^{-\alpha}$. For time-reversible random walks in random environment, Kipnis and Varadhan [24] proved that the process of the environment as viewed from the particle has diffusive behavior. Further, in the reversible setting where the walk is generated by divergence form operators $\nabla \cdot a(\omega) \nabla$, algebraic rates for the decay of $E_\omega[\psi(\bar{\omega}_t)]$ are obtained in [28, 18, 11].

One of our goals in this paper is to prove a diffusive behavior for the environmental process, i.e., a CLT for $\frac{1}{\sqrt{t}} \int_0^t (\psi(\omega_s) - E_{\mathbb{Q}}\psi) ds$ and to investigate the decay rate of $E_\omega[\psi(\bar{\omega}_t)] - E_{\mathbb{Q}}\psi$.

As an important feature of the non-divergence form model, ρ_ω does not have deterministic upper and (nonzero) lower bounds. Moreover, the heat kernel $p_t^\omega(\cdot, \cdot)$ is not expected to have deterministic Gaussian bounds. For ergodic balanced random environments, the following stochastic bounds for the invariant measure ρ_ω and the heat kernel were proved in [26, 16, 13].

For $r \geq 0, t > 0$, denote by

$$\mathfrak{h}(r, t) = \frac{r^2}{r \vee t} + r \log\left(\frac{r}{t} \vee 1\right), \quad r \geq 0, t > 0. \quad (9)$$

Theorem B. *Assume (A2) and that the environment is ergodic under spatial shifts $\{\theta_x : x \in \mathbb{Z}^d\}$. There exists a constant $p = p(d, \kappa) > 0$ such that*

$$(i) \quad \mathbb{E}[\rho^{d/(d-1)}] < \infty, \quad \mathbb{E}[\rho^{-p}] < \infty;$$

(ii) \mathbb{P} -almost surely, for any $r > 0$,

$$\rho_\omega(B_r) \geq C \rho_\omega(B_{2r});$$

(iii) \mathbb{P} -almost surely, for all $x \in \mathbb{Z}^d, t > 0$,

$$\frac{c \rho_\omega(0)}{\rho_\omega(B_{\sqrt{t}})} e^{-C|x|^2/t} \leq p_t^\omega(x, 0) \leq \frac{C \rho_\omega(0)}{\rho_\omega(B_{\sqrt{t}})} e^{-c\mathfrak{h}(|x|, t)}. \quad (10)$$

Moreover, for $x \in \mathbb{Z}^d, t > 0$,

$$\begin{aligned} \|p_t^\omega(0, x)\|_{L^{(d+1)/d}(\mathbb{P})} &\leq \frac{C}{(t+1)^{d/2}} e^{-c\mathfrak{h}(|x|, t)}, \\ \|p_t^\omega(0, x)\|_{L^{-p}(\mathbb{P})} &\geq \frac{c}{(t+1)^{d/2}} e^{-C|x|^2/t}. \end{aligned}$$

The positive moment bound in (i) was obtained by Lawler [26], and the negative moment bound in (i) is a special case of the bound proved by Deuschel and the first named author [13, Theorem 11] in the more general time-dependent ergodic environment setting. The volume-doubling property (ii) was proved by Fabes and Stroock [16, Lemma 2.0]. The heat kernel bounds together with its integrability in the form of (iii) were obtained in [13, Theorem 11] for in the more general dynamic ergodic setting.

Roughly speaking, the term $\frac{\rho_\omega(0)}{\rho_\omega(B_{\sqrt{t}})}$ is the long term ratio between the time the RWRE visits the origin and the time it spends in the ball $B_{\sqrt{t}}$. For the special deterministic environment $a \equiv I$, i.e., when the RWRE is a simple random walk, we have $\rho \equiv 1$ and this constant is $Ct^{-d/2}$. However, in a random environment, one should not expect $\frac{\rho_\omega(0)t^{d/2}}{\rho_\omega(B_{\sqrt{t}})}$ to have deterministic bounds. Hence, to understand the bounds of the invariant measure and the heat kernel, it is crucial to obtain their stochastic integrability.

Although for general ergodic environment, positive and negative moment bounds for the invariant measure and the heat kernel were already obtained in Theorem B (i)(iii), it is a natural question whether better mixing properties of the environment would yield better moment bounds for these quantities.

Our another goal in this paper is to show that, in our i.i.d. balanced environment, both ρ_ω and the heat kernel have positive and negative exponential moment bounds.

In the course of our proof of better moment bounds for the invariant measure and the heat kernels, the following quantitative homogenization result for the non-divergence form operator L_ω will be employed.

Proposition C. *Assume (A1), (A2), (A3). Recall the measure \mathbb{Q} in Theorem A. Suppose $g \in C^\alpha(\partial\mathbb{B}_1)$, $f \in C^\alpha(\mathbb{B}_1)$ for some $\alpha \in (0, 1]$, and ψ is a measurable function of $\omega(0)$ with $\|\psi/\text{tr}\omega\|_\infty < \infty$. For any $q \in (0, d)$, there exist a random variable $\mathcal{X}_q = \mathcal{X}_q(\omega, d, \kappa)$ with $\mathbb{E}[\exp(c\mathcal{X}_q^d)] < \infty$, and a constant $\beta = \beta(d, \kappa, q) \in (0, 1)$ such that for any $y \in B_{3R}$, the solution u of*

$$\begin{cases} \frac{1}{2}\text{tr}(\omega\nabla^2 u(x)) = \frac{1}{R^2}f\left(\frac{x-y}{R}\right)\psi(\theta_{x-y}\omega) & x \in B_R(y), \\ u(x) = g\left(\frac{x-y}{|x-y|}\right) & x \in \partial B_R(y) \end{cases}$$

satisfies, with $A_1 = \|f\|_{C^{0,\alpha}(\mathbb{B}_1)}\|\frac{\psi}{\text{tr}(\omega)}\|_\infty + [g]_{C^{0,\alpha}(\partial\mathbb{B}_1)}$,

$$\max_{x \in B_R(y)} \left| u(x) - \bar{u}\left(\frac{x-y}{R}\right) \right| \lesssim A_1(1 + \mathcal{X}_q R^{-q/d})R^{-\alpha\beta}, \quad (11)$$

where \bar{u} solves (3) with $\bar{a} = E_{\mathbb{Q}}[\omega/\text{tr}\omega] > 0$ and $\bar{\psi} = E_{\mathbb{Q}}[\psi/\text{tr}\omega]$.

Remark 3. Our Proposition C in the above is a version of [19, Theorem 1.5], which can be considered as a discrete version of Armstrong, Smart [3, Theorem 1.2]. We remark that, compared to the aforementioned results in [3, 19], a difference in Proposition C is that the ball is allowed to be centered at any point $y \in B_{3R}$ whereas

the random variable \mathcal{X} stays the same. This subtle feature will allow us to define a “homogenization radius” which will be useful later in our proof of the bounds for the Green functions.

The proof of Proposition C, which is a small modification of that of [19, Theorem 1.5], can be found in the Appendix.

In terms of the quantitative homogenization of non-divergence form operators in the PDE setting, Yurinski derived a second moment estimate of the homogenization error in [30] for linear elliptic case, and Caffarelli, Souganidis [10] proved a logarithmic convergence rate for the nonlinear elliptic case. Afterwards, Armstrong, Smart [3], and Lin, Smart [27] achieved an algebraic convergence rate for fully nonlinear elliptic equations, and fully nonlinear parabolic equations, respectively. Armstrong, Lin [2] obtained quantitative estimates for the approximate corrector problems.

1.4 Main results

For RWRE in an i.i.d. balanced, uniformly elliptic environment, we will establish natural bounds for both the invariant measure and the heat kernel (Theorem 4) which possess both positive and negative exponential moments, greatly improving the stochastic integrability in Theorem 4 for the ergodic setting. We then prove the optimal diffusive decay of the semigroup generated by the heat kernel for $d \geq 3$ in Theorem 5. As consequences, we obtain a functional central limit theorem (CLT) for the environment viewed from the particle process, and deduce the existence of a stationary corrector in dimension $d \geq 5$.

Theorem 4. *Assume (A1), (A2), and $d \geq 2$. Let $s = s(d, \kappa) = 2 + \frac{1}{2\kappa} - d \geq 2$. For any $\varepsilon > 0$, there exists a random variable $\mathcal{H}(\omega) = \mathcal{H}(\omega, d, \kappa, \varepsilon) > 0$ with $\mathbb{E}[\exp(c\mathcal{H}^{d-\varepsilon})] < \infty$ such that the following properties hold.*

(a) For \mathbb{P} -almost all ω ,

$$c\mathcal{H}^{-s} \leq \rho(\omega) \leq C\mathcal{H}^{d-1}.$$

In particular, for any $q \in (-\frac{d}{s}, \frac{d}{d-1})$, we have

$$\mathbb{E}[\exp(c\rho^q)] < \infty.$$

(b) Recall the function \mathfrak{h} in (9). For any $r \geq 1$ and \mathbb{P} -almost all ω ,

$$c\mathcal{H}^{-s} \leq \frac{r^d \rho_\omega(0)}{\rho_\omega(B_r)} \leq C\mathcal{H}^{d-1}.$$

(c) For any $x \in \mathbb{Z}^d, t > 0$, and \mathbb{P} -almost all ω ,

$$\begin{aligned} p_t^\omega(x, 0) &\leq C\mathcal{H}^{d-1}(1+t)^{-d/2} e^{-c\mathfrak{h}(|x|, t)}, \\ p_t^\omega(x, 0) &\geq c\mathcal{H}^{-s}(1+t)^{-d/2} e^{-C|x|^2/t}. \end{aligned}$$

Recall the continuous time RWRE $(Y_t)_{t \geq 0}$ in Definition 2. With abuse of notation, we still denote the process of the environment viewed from the particle as

$$\bar{\omega}^t := \theta_{Y_t} \omega.$$

Following Gloria, Neukamm, Otto [18], for any measurable function $\zeta : \Omega \rightarrow \mathbb{R}$, we define its *stationary extension* $\bar{\zeta} : \mathbb{Z}^d \times \Omega \rightarrow \mathbb{R}$ as

$$\bar{\zeta}(x) = \bar{\zeta}(x; \omega) := \zeta(\theta_x \omega).$$

Define the semigroup P_t , $t \geq 0$, on \mathbb{R}^Ω by

$$P_t \zeta(\omega) = E_\omega^0[\zeta(\bar{\omega}^t)] = \sum_z p_t^\omega(0, z) \bar{\zeta}(z; \omega).$$

The following theorem estimates the speed of decorrelation of the environmental process $\bar{\omega}^t$ from the original environment. It gives a rate $t^{-d/4}$ of decay for the semigroup, which is optimal. A function $\zeta : \Omega \rightarrow \mathbb{R}$ is said to be *local* if it depends only on the environment $\{\omega(x) : x \in S\}$ in a finite set $S \subset \mathbb{Z}^d$. Such a set S is called the *support* of ζ and denoted by $\text{Supp}(\zeta)$.

Theorem 5. *Assume (A1), (A2), and $d \geq 3$. For any local measurable function $\zeta : \Omega \rightarrow \mathbb{R}$ with $\|\zeta\|_\infty \leq 1$ and $t \geq 0$, we have, for $C = C(d, \kappa, \#\text{Supp}(\zeta))$,*

$$\text{Var}_{\mathbb{Q}}(P_t \zeta) \leq C(1+t)^{-d/2}; \tag{12}$$

$$\|P_t \zeta - E_{\mathbb{Q}} \zeta\|_{L^1(\mathbb{P})} + \|P_t \zeta - \mathbb{E}[P_t \zeta]\|_{L^p(\mathbb{P})} \leq C_p(1+t)^{-d/4} \quad \text{for all } p \in (0, 2). \tag{13}$$

For divergence form operators, optimal diffusive decay of the semigroup generated by the heat kernel was obtained by Gloria, Neukamm, Otto [18], de Buyer, Mourrat [11]. Our proof of Theorem 5 follows the approach of [18], which uses an Efron-Stein type inequality and the Duhamel representation formula for the vertical derivative. However, unlike the divergence form setting [18], there are no deterministic Gaussian bounds for the heat kernel, and the steady state \mathbb{Q} of the environment process $(\bar{\omega}^t)_{t \geq 0}$ is not the same as the original measure \mathbb{P} . To overcome these difficulties, our heat kernel estimates and the (negative and positive) moment bounds of the Radon-Nikodym derivative $\frac{d\mathbb{P}}{d\mathbb{Q}}$ in Theorem 4 play crucial roles. (Another feature of our non-divergence setting that worth mentioning is that there is no Caccioppoli estimates for $p > 2$. See Lemma 21.)

As a consequence of Theorem 5 and the CLT of [12], we obtain a functional CLT for an additive functional of the environmental process. Recall that $(\bar{\omega}^t)_{t \geq 0}$ is an ergodic sequence under the law $\mathbb{Q} \times P_\omega$ and time shifts. The following CLT says that, when $d \geq 3$, the fluctuation around the ergodic mean is approximately Gaussian under the diffusive rescaling.

Corollary 6. Assume (A1), (A2). Let $d \geq 3$. For \mathbb{P} -almost all ω and any bounded measurable local function ζ of the environment with $E_{\mathbb{Q}}\zeta = 0$, the P_{ω} law of

$$\frac{1}{\sqrt{t}} \int_0^t \zeta(\bar{\omega}^s) ds$$

converges weakly to a Brownian motion with a deterministic diffusivity constant.

Another consequence of Theorem 5 is the existence of a stationary corrector in $d \geq 5$ for the non-divergence form homogenization problem (2).

Corollary 7. Assume (A1), (A2). When $d \geq 5$, for any bounded local measurable function $\zeta : \Omega \rightarrow \mathbb{R}$, there exists $\phi : \Omega \rightarrow \mathbb{R}$ such that $\phi \in L^p(\mathbb{P})$ for all $p \in (0, 2)$, and for \mathbb{P} -almost all ω , its stationary extension $\bar{\phi}(x) = \phi(\theta_x \omega)$ solves

$$L_{\omega} \bar{\phi}(x) = \bar{\zeta}(x) - E_{\mathbb{Q}}[\zeta], \quad \text{for all } x \in \mathbb{Z}^d. \quad (14)$$

Remark 8. The corrector $\bar{\phi}(x)$ plays a crucial role in the quantification of the homogenization error for non-divergence form operators. We remark that our Corollary 7 is a weaker version of [2, Theorem 7.1] where not only the existence of the stationary corrector was proved but a stretched exponential tail was also obtained. Specifically, [2] derived such a result using a large scale $C^{1,1}$ estimate for the homogenization problem. Although our Corollary 7 is an immediate consequence of the optimal diffusive decay rate, it has much weaker stochastic integrability.

In the classical periodic environment setting, it is well-known that the existence of a stationary corrector implies that the optimal homogenization error of problem (2) is generically of scale R^{-1} . Readers may refer to the classical books [4, 23] for the derivation of the rate in the periodic setting, and [21, 20] for discussions on the optimality of the rates.

To obtain the above results, we need to study properties of the Green functions. For $d \geq 2$, $R \geq 1$, denote the exit time from B_R of the RWRE by

$$\tau = \tau_R = \inf\{n \geq 0 : X_n \notin B_R\}. \quad (15)$$

Definition 9. For $R \geq 1$, $\omega \in \Omega$, $x \in \mathbb{Z}^d$, $S \subset \mathbb{Z}^d$, the Green function $G_R(\cdot, \cdot)$ in the ball B_R for the balanced random walk is defined by

$$G_R(x, S) = G_R^{\omega}(x, S) := E_{\omega}^x \left[\sum_{n=0}^{\tau_R-1} \mathbb{1}_{X_n \in S} \right], \quad x \in \bar{B}_R.$$

We also write $G_R(x, y) := G_R^{\omega}(x, \{y\})$ and $G_R(x) := G(x, 0)$.

Note that for $d \geq 3$, by [22, Theorem 1], the RWRE is transient, and so the Green function in the whole space

$$G^{\omega}(x) := \lim_{R \rightarrow \infty} G_R(x) < \infty$$

is well-defined for all $x \in \mathbb{Z}^d$, \mathbb{P} -almost surely. Whereas, when $d = 2$, the RWRE is recurrent, and thus the Green function in the whole \mathbb{Z}^2 is infinity. In this case, the *potential kernel*

$$A(x) = A^\omega(x) = \sum_{n=0}^{\infty} [p_n^\omega(0, 0) - p_n^\omega(x, 0)], \quad x \in \mathbb{Z}^2, \quad (16)$$

is well-defined. Note that G and A are both non-negative functions, and, for $x \in \mathbb{Z}^d$,

$$L_\omega G(x) = -\mathbb{1}_{x=0}, \quad \text{if } d \geq 3,$$

and

$$L_\omega A(x) = \mathbb{1}_{x=0}, \quad \text{if } d = 2.$$

Theorem 10. *Assume (A1), (A2). For $r > 0$, let*

$$U(r) := \begin{cases} -\log r & d = 2, \\ r^{2-d} & d \geq 3. \end{cases} \quad (17)$$

For any $\varepsilon > 0$, there exists a random variable $\mathcal{H} = \mathcal{H}(\omega, d, \kappa, \varepsilon) > 0$ with $\mathbb{E}[\exp(c\mathcal{H}^{d-\varepsilon})] < \infty$ such that, for \mathbb{P} -almost surely, for all $x \in B_R$,

$$\mathcal{H}^{-s}[U(|x| + 1) - U(R + 2)] \lesssim G_R^\omega(x) \lesssim \mathcal{H}^{d-1}[U(|x| + 1) - U(R + 2)],$$

where $s = s(d, \kappa) = 2 + \frac{1}{2\kappa} - d \geq 2$.

We remark that an upper bound for the Green function of the approximate corrector (which is defined in the whole \mathbb{R}^d) was proved by Armstrong, Lin [2, Proposition 4.1].

Our proof of the bounds of G_R^ω follows the idea of Armstrong, Lin [2, Proposition 4.1]. In Theorem 10, we apply their idea to obtain both upper and lower bounds for the Green function G_R in a finite region.

Corollary 11. *Assume (A1), (A2). Let s be as in Theorem 10. For any $\varepsilon > 0$, there exists a random variable $\mathcal{H} = \mathcal{H}(\omega, d, \kappa, \varepsilon) > 0$ with $\mathbb{E}[\exp(c\mathcal{H}^{d-\varepsilon})] < \infty$ such that, \mathbb{P} -almost surely, for all $x \in \mathbb{Z}^d$,*

$$\begin{aligned} \mathcal{H}^{-s} \log(|x| + 1) &\lesssim A^\omega(x) \lesssim \mathcal{H} \log(|x| + 1), \text{ when } d = 2; \\ \mathcal{H}^{-s}(1 + |x|)^{2-d} &\lesssim G^\omega(x) \lesssim \mathcal{H}^{d-1}(1 + |x|)^{2-d}, \text{ when } d \geq 3. \end{aligned}$$

2 Bounds of the Green function in a ball

By the Markov property, $G_R(x, S)$ satisfies $G_R = 0$ on ∂B_R and

$$L_\omega G_R(x, S) = -\mathbb{1}_{x \in S}, \quad x \in B_R. \quad (18)$$

Our proof of the bounds of the Green function G_R (Theorem 10) follows the idea of Armstrong-Lin [2, Proposition 4.1]. The idea, which we learn from [2], is through the comparison of G_R and test functions, and it is explained as follows.

Let us call a function $u : \mathbb{Z}^d \rightarrow \mathbb{R}$ ω -harmonic on $A \subset \mathbb{Z}^d$ if $L_\omega u(y) = 0$ for $y \in A$. Clearly, the Green function $G_R(x, 0)$ is ω -harmonic on $B_R \setminus \{0\}$.

To obtain the upper bound, we construct a function h which is almost ω -harmonic away from the origin and with (almost) zero boundary values, so that $G_R - h$ is subharmonic at places that are either close to the origin or the boundary of B_R . As a result, if $(G_R - h)(x_0) = \max_{B_R} (G_R - h)$ were positive, then the maximum principle forces the maximizer x_0 to be sufficiently far away from both the origin and the boundary. This allows enough space for the homogenization to occur around x_0 , i.e., h is close to its continuous harmonic counterpart up to an algebraic error. On the other hand, by the maximal principle, the ω -harmonic counterpart of $G_R - h$ (which is an algebraic error away from $G_R - h$) cannot achieve its maximum over the ball $\bar{B}_{|x_0|/2}(x_0)$ in the center. This would contradict the assumption that the maximizer is x_0 , if we can exploit the fact that h is strictly ω -superharmonic to give it enough room to absorb the algebraic homogenization error.

The proof of the lower bound follows similar philosophy.

Note that [2, Proposition 4.1] only deals with the Green function of the ‘‘approximate corrector’’ which is defined on the whole \mathbb{R}^d , while our Green function corresponds to the original non-divergence form operator within a finite ball B_R . Hence, in our case, the challenge also lies in the construction of test functions so that they have the desired boundary values and concavity near the discrete boundary.

The following Lemmas contain properties of some deterministic functions that will be useful in our construction of the test functions in the next subsections.

Lemma 12. *Let $\delta = \beta/2$, where $\beta = \beta(d, \kappa, q_\varepsilon)$ is as defined in Proposition C. Define $\bar{\zeta}, \bar{\xi} : (0, \infty) \rightarrow \mathbb{R}$ as*

$$\bar{\zeta}(r) = \begin{cases} -(\log r) \exp(r^{-\delta}/\delta) & d = 2 \\ r^{2-d} \exp(-r^{-\delta}/\delta) & d \geq 3, \end{cases}$$

$$\bar{\xi}(r) = \begin{cases} -(\log r) \exp(-r^{-\delta}/\delta) & d = 2 \\ r^{2-d} \exp(r^{-\delta}/\delta) & d \geq 3. \end{cases}$$

Define two functions $\zeta, \xi : \mathbb{R}^d \setminus \{0\} \rightarrow \mathbb{R}$ as

$$\zeta(y) = \bar{\zeta}(|y|), \quad \text{and} \quad \xi(y) = \bar{\xi}(|y|), \quad y \neq 0.$$

Then, the following statements hold.

(i) $\bar{\zeta}, \bar{\xi}$ are decreasing functions on (C, ∞) . Moreover, for $r \geq C$,

$$-\bar{\zeta}'(r) \asymp r^{1-d}, \quad \text{and} \quad 0.5r^{1-d} \leq -\bar{\xi}'(r) \leq (d - 0.5)r^{1-d}.$$

(ii) For $|y| \geq C$, we have

$$-\Delta \zeta(y) \geq |y|^{-(2+\delta)} |\zeta(y)|, \quad \text{and} \quad -\Delta \xi(y) \leq -|y|^{-(2+\delta)} |\xi(y)|.$$

(iii) For $|y| \geq 2$ and $k \in \mathbb{N}$, there exists $C = C(k, d)$ such that

$$|D^k \zeta(y)| \leq C|y|^{-k} |\zeta(y)|, \quad \text{and} \quad |D^k \xi(y)| \leq C|y|^{-k} |\xi(y)|.$$

Lemma 13. *There exist constants $\alpha_0 \in (0, 1)$ and $A_0 \geq 1$ depending on κ such that, for any $\alpha \in (0, \alpha_0)$, $A \geq A_0$,*

$$L_\omega(e^{-2\alpha|x|/R}) \leq 0 \quad \text{in } B_R \setminus B_{R/2}, \text{ when } R \geq A_0; \quad (19)$$

$$L_\omega(e^{-A|x|^2}) \geq -\mathbb{1}_{x=0}, \quad x \in \mathbb{Z}^d; \quad (20)$$

$$L_\omega(e^{-A|x|^2/R^2}) > 0, \quad x \in B_R \setminus B_{R/2}, \text{ when } R \geq A^2. \quad (21)$$

The proof of Lemma 13 is in Section A.2 of the Appendix.

2.1 Upper bounds of Green's functions

Recall \mathcal{X}_q in Proposition C. For any $\varepsilon \in (0, 1)$, we write

$$R_0 = R_0(\omega, d, \kappa, \varepsilon) := \mathcal{X}_{d-\varepsilon}^{d/(d-\varepsilon)} + K,$$

where K is a sufficiently large constant depending on (d, κ) , and denote the exit time from B_{R_0} as

$$s_0 = \min\{n \geq 0 : X_n \notin B_{R_0}\}. \quad (22)$$

Note that R_0 plays the role of a "homogenization radius" in the sense that for all $R \geq R_0$ and $y \in B_{3R}$, the upper bound in (11) can be replaced by the algebraic term $CA_1 R^{-\alpha\beta}$.

Let $\alpha = \alpha(d, \kappa) > 0$ be a constant to be determined in Lemma 15, and set

$$C_{\alpha, R} := \frac{[\bar{\zeta}(R/2) - \bar{\zeta}(R)]R^{d-2}}{e^{-\alpha+2\alpha/R} - e^{-2\alpha}} \asymp \alpha^{-1}, \quad \text{when } R \geq R_0.$$

Definition 14. Let $\bar{\zeta}, \zeta$ be as in Lemma 12. For any fixed $R \geq 4R_0$, we define a function $h : \bar{B}_R \rightarrow [0, \infty)$ by

$$h(x) = \begin{cases} h_1(x), & x \in B_{R_0}, \\ h_2(x), & x \in B_{R/2} \setminus B_{R_0}, \\ h_3(x), & x \in \bar{B}_R \setminus B_{R/2}, \end{cases}$$

where the functions h_1, h_2, h_3 , are defined as below

$$h_2(y) = R_0^{d-1} [(\alpha^{-1} - 1)(\bar{\zeta}(R/2) - \bar{\zeta}(R)) + \zeta(y) - \bar{\zeta}(R)], \quad y \in \mathbb{R}^d \setminus \{0\},$$

$$h_1(x) = E_\omega^x[h_2(X_{s_0}) + |X_{s_0}| - |x|], \quad x \in \bar{B}_{R_0},$$

$$h_3(y) = R_0^{d-1} \alpha^{-1} C_{\alpha, R} R^{2-d} [e^{-2\alpha(|y|-1)/R} - e^{-2\alpha}], \quad y \in \mathbb{R}^d.$$

Lemma 15. *When $R \geq 4R_0$, there exists a constant $\alpha > 0$ such that the functions h_1, h_2, h_3, h given in Definition 14 have the following properties.*

- (a) $L_\omega h_1(x) = -L_\omega(|x|) \leq -\mathbb{1}_{x=0}$ for $x \in B_{R_0}$;
- (b) $h_1 = h_2$ on ∂B_{R_0} , and $h_2 = h_3$ on $\partial \mathbb{B}_{R/2}$;
- (c) $h_2 \geq h_1$ in $B_{R_0} \setminus B_{R_0/2}$.
- (d) $h_2 \geq h_3$ in $\mathbb{B}_R \setminus \mathbb{B}_{R/2}$, and $h_2 \leq h_3$ in $\mathbb{B}_{R/2} \setminus \mathbb{B}_{R/2-1}$.
- (e) $L_\omega h_3 \leq 0$ in $B_R \setminus B_{R/2}$.

Proof. (a) and (b) are obvious. To see (c), note that $h_2(x) - h_1(x) = E_\omega^x[f(|x|) - f(|X_{s_0}|)]$, where $f(r) = r + R_0^{d-1}\bar{\zeta}(r)$. Since $R_0 \geq K$, by Lemma 12(i), taking K sufficiently large, $f(r)$ is a decreasing function for $r \in [R_0/2, R_0 + 1]$.

Next, we will prove (d). Indeed, we can write, for $y \in \mathbb{R}^d \setminus \{0\}$,

$$h_2(y) - h_3(y) = R_0^{d-1}a(|y|) + A(R_0, R),$$

where $A(R_0, R)$ is a constant, and $a(r) = \bar{\zeta}(r) - \alpha^{-1}C_{\alpha,R}R^{2-d}e^{-2\alpha(r-1)/R}$. For $r \in [R/2 - 1, R]$, by Lemma 12(i), taking $\alpha > 0$ sufficiently small, we have

$$a'(r) \geq -Cr^{1-d} + 2C_{\alpha,R}e^{-2\alpha}R^{1-d} \geq -Cr^{1-d} + c\alpha^{-1}R^{1-d} \geq 0.$$

Hence, $h_2 - h_3$ is radially increasing in $\mathbb{B}_R \setminus \mathbb{B}_{R/2-1}$. Item (d) then follows from the fact that $h_2 - h_3 = 0$ on $\partial \mathbb{B}_{R/2}$.

Item (e) is a consequence of (19) in Lemma 13. \square

Proof of the upper bound in Theorem 10. For $0 < a < b$ and $d \geq 2$, we have an elementary inequality

$$b - a \leq b^{d-1}(U(a) - U(b)). \quad (23)$$

When $R \in (1, 4R_0]$, note that $L_\omega[G_R(x) - (R + 1 - |x|)] \geq 0$, and $G_R = 0 \leq R + 1 - |x|$ on ∂B_R . By the maximum principle, we have, for $x \in B_R$,

$$G_R(x) \leq R + 1 - |x| \stackrel{(23)}{\leq} (R + 2)^{d-1}[U(|x| + 1) - U(R + 2)].$$

Hence the upper bound in Theorem 10 holds when $R \in (1, 4R_0]$. It remains to consider the case $R > 4R_0$.

First, we will prove via contradiction that

$$G_R \leq h \quad \text{in } \bar{B}_R. \quad (24)$$

Assume by contradiction that (24) fails, i.e., $\max_{\bar{B}_R}(G_R - h) > 0$. By Lemma 15(a), $L_\omega(G_R - h) \geq 0$ in B_{R_0} and so $\max_{\bar{B}_R}(G_R - h)$ is achieved outside of B_{R_0} . Further, note that $(G_R - h)|_{\partial B_R} = (-h_3)|_{\partial B_R} \leq 0$. By Lemma 15(e), $L_\omega(G_R - h_3) \geq 0$ in $B_R \setminus B_{R/2}$, and so, by the maximum principle and Lemma 15(d),

$$\max_{\bar{B}_R \setminus B_{R/2}}(G_R - h) \leq \max_{\partial(B_R \setminus B_{R/2})}(G_R - h_3) \leq 0 \vee \max_{B_{R/2} \setminus B_{R_0}}(G_R - h).$$

Hence, if $\max_{\bar{B}_R}(G_R - h) > 0$, then there exists $x_0 \in B_{R/2} \setminus B_{R_0}$ so that

$$(G_R - h)(x_0) = \max_{\bar{B}_R}(G_R - h) > 0.$$

Since $x_0 \in B_{R/2} \setminus B_{R_0}$, by Lemma 15(c)(d),

$$(G_R - h_2)(x_0) \geq \max_{\bar{B}_{|x_0|/2}(x_0)}(G_R - h_2),$$

which is equivalent to

$$(G_R - R_0^{d-1}\zeta)(x_0) \geq \max_{\bar{B}_{|x_0|/2}(x_0)}(G_R - R_0^{d-1}\zeta). \quad (25)$$

Without loss of generality, assume that $\bar{a} = I$, and set

$$\tilde{\zeta}(y) := \zeta(y) + c|x_0|^{-(2+\delta)}|\zeta(x_0)||y - x_0|^2, \quad y \in \mathbb{R}^d \setminus \{0\},$$

where $c > 0$ is chosen so that (by Lemma 12(ii)) $\Delta\tilde{\zeta}(y) \leq 0$ for $y \in \mathbb{B}_{|x_0|/2}(x_0)$. Then (by Lemma 12(iii)) $|D\tilde{\zeta}| \leq C|x_0|^{-1}|\zeta(x_0)|$ in $\mathbb{B}_{1+0.5|x_0|}(x_0)$, and

$$(G_R - R_0^{d-1}\tilde{\zeta})(x_0) \stackrel{(25)}{\geq} \max_{\partial B_{|x_0|/2}(x_0)}(G_R - R_0^{d-1}\tilde{\zeta}) + CR_0^{d-1}|x_0|^{-\delta}|\zeta(x_0)|. \quad (26)$$

Let $\bar{v} : \bar{\mathbb{B}}_1 \rightarrow \mathbb{R}$ and $v : \bar{B}_{|x_0|/2}(x_0) \rightarrow \mathbb{R}$ be the solutions of (Here $\bar{a} = I$.)

$$\begin{cases} \operatorname{tr}(\bar{a}D^2\bar{v}) = \Delta\bar{v} = 0 & x \in \mathbb{B}_1 \\ \bar{v}(x) = R_0^{d-1}\tilde{\zeta}(x_0 + \frac{|x_0|}{2}x) & x \in \partial\mathbb{B}_1, \end{cases}$$

and

$$\begin{cases} L_\omega v(x) = 0 & x \in B_{|x_0|/2}(x_0) \\ v(x) = R_0^{d-1}\bar{v}(\frac{x-x_0}{|x-x_0|}) & x \in \partial B_{|x_0|/2}(x_0). \end{cases}$$

We will show that v can be controlled by $R_0^{d-1}\tilde{\zeta}$ both on the boundary and inside of $B_{|x_0|/2}(x_0)$. Indeed, for $x \in \partial B_{|x_0|/2}(x_0)$,

$$\begin{aligned} |v(x) - R_0^{d-1}\tilde{\zeta}(x)| &= R_0^{d-1}\left|\tilde{\zeta}(x_0 + \frac{|x_0|(x-x_0)}{2|x-x_0|}) - \tilde{\zeta}(x)\right| \\ &\leq R_0^{d-1} \sup_{\bar{\mathbb{B}}_{1+|x_0|/2}(x_0)} |D\tilde{\zeta}| \\ &\leq CR_0^{d-1}|x_0|^{-1}|\zeta(x_0)|. \end{aligned} \quad (27)$$

For $x \in B_{|x_0|/2}(x_0)$, applying Proposition C to the case $\alpha = 1$, there exists $\beta = \beta(d, \kappa, \varepsilon) \in (0, 1)$ such that

$$\begin{aligned} v(x) &\leq \bar{v}(\frac{x-x_0}{|x_0|/2}) + C|x_0|^{-\beta}R_0^{d-1} \sup_{y \in \partial\mathbb{B}_1} \left|D\tilde{\zeta}(x_0 + \frac{|x_0|}{2}y)\right| \\ &\leq \bar{v}(\frac{x-x_0}{|x_0|/2}) + CR_0^{d-1}|x_0|^{-\beta}|\zeta(x_0)|. \end{aligned} \quad (28)$$

Furthermore, using the fact that $\Delta \tilde{\zeta}(x_0 + \frac{|x_0|}{2}x) \leq 0$ for $x \in \mathbb{B}_1$, we get $\bar{v}(x) \leq R_0^{d-1} \tilde{\zeta}(x_0 + \frac{|x_0|}{2}x)$ in \mathbb{B}_1 . This, together with (28), yields, for $x \in B_{|x_0|/2}(x_0)$,

$$v(x) \leq R_0^{d-1} \tilde{\zeta}(x) + CR_0^{d-1} |x_0|^{-\beta} |\zeta(x_0)|. \quad (29)$$

Notice that $(G_R - v)$ is an ω -harmonic function on $B_{|x_0|/2}(x_0)$, and so

$$\max_{B_{|x_0|/2}(x_0)} (G_R - v) \leq \max_{\partial B_{|x_0|/2}(x_0)} (G_R - v).$$

Therefore, combining this inequality and (27), (29), we get

$$\max_{B_{|x_0|/2}(x_0)} (G_R - R_0^{d-1} \tilde{\zeta}) \leq \max_{\partial B_{|x_0|/2}(x_0)} (G_R - R_0^{d-1} \tilde{\zeta}) + CR_0^{d-1} |x_0|^{-\beta} |\zeta(x_0)|$$

which contradicts (26), since $|x_0| \in (R_0, R)$, $\delta = \beta/2$ by definition in Lemma 12, and $R_0 \geq K$ is chosen to be sufficiently large. Inequality (24) is proved.

Finally, when $x \in B_{R/2} \setminus B_{R_0}$, we have $G_R(x) \stackrel{(24)}{\leq} h_2(x)$, and, by Lemma 12(i),

$$\begin{aligned} h_2(x) &\leq R_0^{d-1} \alpha^{-1} [\bar{\zeta}(|x|) - \bar{\zeta}(R)] \\ &\lesssim R_0^{d-1} \int_{|x|}^R (-\bar{\zeta})'(r) dr \\ &\lesssim R_0^{d-1} \int_{|x|}^R r^{1-d} dr \lesssim R_0^{d-1} [U(|x|) - U(R)]. \end{aligned} \quad (30)$$

When $x \in B_{R_0} \setminus \{0\}$,

$$\begin{aligned} G_R(x) &\stackrel{(24)}{\leq} h_1(x) = E_\omega^x [h_2(X_{s_0}) + |X_{s_0}| - |x|] \\ &\stackrel{(30),(23)}{\lesssim} R_0^{d-1} E_\omega^x [U(|X_{s_0}|) - U(R) + U(|x|) - U(|X_{s_0}|)] \\ &= R_0^{d-1} [U(|x|) - U(R)]. \end{aligned}$$

Note that, for $|x| \geq 1$, $U(|x|) - U(R) \lesssim U(|x| + 1) - U(R + 2)$.

When $x \in B_R \setminus B_{R/2}$,

$$\begin{aligned} G_R &\leq h_3 \leq CR_0^{d-1} R^{2-d} (e^{-2\alpha(|x|-1)/R} - e^{-2\alpha}) \\ &\lesssim R_0^{d-1} R^{2-d} (1 - \frac{|x|-1}{R}) = R_0^{d-1} R^{1-d} (R + 1 - |x|) \\ &\stackrel{(23)}{\lesssim} R_0^{d-1} [U(|x| + 1) - U(R + 2)]. \end{aligned}$$

The upper bound in Theorem 10 is proved by putting $\mathcal{H} = R_0$. \square

2.2 Lower bounds of Green's functions

The proof of the lower bound of Theorem 10, which is similar to that of Theorem 10, is via comparing G_R to appropriate test functions. However, unlike the Green function in the whole space, G_R is defined only in bounded region, and so the test functions should be carefully designed to capture the behavior of G_R near the boundary.

Lemma 16. Define $\bar{\eta} : \mathbb{R} \rightarrow (0, \infty)$ as $\bar{\eta}(r) = (1 + r^2)^{-\theta}$, where

$$\theta := 1/(4\kappa) \geq d/2. \quad (31)$$

Define $\eta : \mathbb{R}^d \rightarrow \mathbb{R}$ as

$$\eta(y) = \bar{\eta}(|y|).$$

There exists a constant $C_0 = C_0(d, \kappa) > 0$ such that, for $x \in \mathbb{Z}^d$,

$$L_\omega \eta(x) \geq -\mathbb{1}_{x \in B_{C_0 \theta^2}}.$$

The proof of Lemma 16 is in Section A.2 of the Appendix.

Let $\gamma = \gamma(\kappa) > 0$ be a large constant to be determined, and set

$$C_{\gamma, R} := \frac{[\bar{\xi}(R/2) - \bar{\xi}(R)]R^{d-2}}{e^{-\gamma/4} - e^{-\gamma}} \asymp e^{\gamma/4} \quad \text{when } R \geq R_0.$$

Definition 17. Recall $\xi, \bar{\xi}, \eta, \bar{\eta}, \theta$ be in Lemma 12 and Lemma 16. For any fixed $R \geq 4R_0$, we define three functions ℓ_i , $i = 1, 2, 3$, as

$$\begin{aligned} \ell_2(y) &= R_0^{d-2-2\theta} [(\gamma^{-2} - 1)(\bar{\xi}(R/2) - \bar{\xi}(R)) + \xi(y) - \bar{\xi}(R)], \quad y \in \mathbb{R}^d \setminus \{0\}; \\ \ell_1(x) &= E_\omega^x[\ell_2(X_{s_0}) + \eta(x) - \eta(X_{s_0})], \quad x \in \bar{B}_{R_0}; \\ \ell_3(y) &= R_0^{d-2-2\theta} \gamma^{-2} C_{\gamma, R} R^{2-d} (e^{-\gamma|y|^2/R^2} - e^{-\gamma}), \quad y \in \mathbb{R}^d, \end{aligned}$$

Also, for $R \geq 4R_0$, we define a function $\ell : \bar{B}_R \rightarrow \mathbb{R}$ by

$$\ell(x) = \begin{cases} \ell_1(x), & x \in B_{R_0}, \\ \ell_2(x), & x \in B_{R/2} \setminus B_{R_0}, \\ \ell_3(x), & x \in \bar{B}_R \setminus B_{R/2}. \end{cases}$$

Lemma 18. When $R \geq 4R_0$, there exists a constant $\gamma > 0$ such that the functions $\ell_1, \ell_2, \ell_3, \ell$ given in Definition 17 have the following properties.

- (a) $L_\omega \ell_1 = L_\omega \eta \geq -\mathbb{1}_{x \in B_{C_0 \theta^2}}$ for $x \in B_{R_0}$;
- (b) $\ell_1 = \ell_2$ on ∂B_{R_0} , and $\ell_2 = \ell_3$ on $\partial B_{R/2}$;
- (c) $\ell_2 \leq \ell_1$ in $B_{R_0} \setminus B_{R_0/2}$.

(d) $\ell_2 \leq \ell_3$ in $\mathbb{B}_R \setminus \mathbb{B}_{R/2}$, and $\ell_2 \geq \ell_3$ in $\mathbb{B}_{R/2} \setminus \mathbb{B}_{0.5R-2}$.

(e) $L_\omega \ell_3 \geq 0$ in $B_R \setminus B_{R/2}$.

Proof. (a) follows from Lemma 16, and (b) follows from definition. To see (c), note that $\ell_1 - \ell_2 = E_\omega^x[f(|X_{s_0}|) - f(|x|)]$, where $f(r) = R_0^{d-2-2\theta} \bar{\xi}(r) - \bar{\eta}(r)$. Since $R_0 \geq K$, by taking K sufficiently large and by Lemma 12(i), we have

$$\begin{aligned} f'(r) &\geq -(d-0.5)R_0^{d-2-2\theta}r^{1-d} + 2\theta(1 + \frac{1}{r^2})^{-\theta-1}r^{d-2-2\theta} \\ &\stackrel{(31)}{\geq} (d-0.5)(r^{d-2-2\theta} - R_0^{d-2-2\theta}) \geq 0 \end{aligned}$$

and so $f(r)$ is decreasing for $r \in [R_0/2, R_0]$. Item (c) is proved.

Next, we will show (d). Indeed, we can write, for $y \neq 0$,

$$\ell_2(y) - \ell_3(y) = R_0^{d-2-2\theta} a(|y|) + A(R_0, R),$$

where $A(R_0, R)$ is a constant, and $a(r) = \bar{\xi}(r) - \gamma^{-2} C_{\gamma, R} R^{2-d} e^{-\gamma r^2 / R^2}$. By Lemma 12(i),

$$\begin{aligned} a'(r) &\leq -cr^{1-d} + C\gamma^{-1} C_\gamma R^{-d} e^{-\gamma r^2 / R^2} \\ &\leq cr^{1-d} \left(-1 + C\gamma^{-1} \left(\frac{r}{R}\right)^d \right) < 0 \end{aligned}$$

for $r \in [0.5R-2, R]$ if γ is chosen to be sufficiently large. Hence, $\ell_2 - \ell_3$ is radially decreasing in $\mathbb{B}_R \setminus \mathbb{B}_{0.5R-2}$. Item (d) then follows from the fact that $\ell_2 = \ell_3$ on $\partial\mathbb{B}_{R/2}$. Item (e) is a consequence of (21) in Lemma 13. \square

Proof of the lower bound in Theorem 10: It suffices to show that, for $x \in B_R$,

$$G_R^\omega(x) \gtrsim R_0^{d-2-2\theta} (U(|x| + 1) - U(R + 1)). \quad (32)$$

Recall $U(r)$ in (17). Indeed, (32) is equivalent to the lower bound of Theorem 10 when $x \in B_{R-1}$. When $x \in B_R \setminus B_{R-1}$, $R \geq 2$, taking $y \in B_R$ with $|y| \leq |x| - 1$ and $|x - y|_1 \leq 2d$, by the assumption (A2), inequality (32) yields

$$\begin{aligned} G_R(x) &\geq P_x^\omega(X_{|x-y|_1} = y) G_R(y) \\ &\gtrsim R_0^{d-2-2\theta} (U(|x|) - U(R + 1)) \\ &\gtrsim R_0^{d-2-2\theta} (U(|x| + 1) - U(R + 2)). \end{aligned}$$

Our proof of (32) consists of several steps. Let C_0 be as in Lemma 16 and recall $\tau = \tau_R$ in (15).

When $R \in (1, 2C_0\theta^2)$, by (20), taking $A = A(\kappa) \geq 1$ sufficiently large,

$$L_\omega[G_R - (e^{-A|x|^2} - e^{-AR^2})] \leq 0 \quad \text{in } B_R.$$

Since $G_R = 0 \geq (e^{-A|x|^2} - e^{-AR^2})$ on ∂B_R , by the maximum principle, we have $G_R \geq e^{-A|x|^2} - e^{-AR^2}$ in B_R . Thus, using the inequality $e^a \geq 1 + a$ for $a \geq 0$, we get, for $x \in B_R$, (Note that $R \asymp 1$ in this case.)

$$G_R \geq e^{-AR^2} \left(e^{A(R^2 - |x|^2)} - 1 \right) \geq e^{-AR^2} (R^2 - |x|^2) \gtrsim R - |x|$$

By the fact $a \geq \log(1 + a)$, $a \geq 0$, we have $R - |x| \gtrsim \log \frac{R+1}{|x|+1}$. Moreover, since $R \asymp 1$, for $d \geq 3$, we also have $R - |x| \gtrsim (|x| + 1)^{2-d} - (R + 1)^{2-d}$. Thus (32) holds for this case.

When $R \geq 2C_0\theta^2$, by assumption (A2), for $x \in B_R$ and any $y \in B_{C_0\theta^2}$,

$$\begin{aligned} G_R(x) &\geq \sum_{i=0}^{\infty} P_{\omega}^x(X_i = y, i < \tau_R) P_{\omega}^y(X_{|y|_1} = 0) \\ &\geq G_R(x, y) \kappa^{|y|_1} \gtrsim G_R(x, y), \end{aligned}$$

and so (Recall $G_R(\cdot, \cdot)$ in Definition 9.)

$$G_R(x) \gtrsim G_R(x, B_{C_0\theta^2}) =: H_R(x), \quad x \in B_R. \quad (33)$$

Thus it suffices to obtain the corresponding lower bound for H_R defined above.

When $R \in (2C_0\theta^2, 4R_0]$, since (by Lemma 16) $L_{\omega}(H_R - \eta) \leq 0$ in B_R , by the maximum principle, we have $H_R \geq \eta - \bar{\eta}(R)$ in B_R . Notice that

$$\bar{\eta}(r_1) - \bar{\eta}(r_2) \gtrsim R_0^{d-2-2\theta} (U(r_1 + 1) - U(r_2 + 1)), \quad \forall r_1 < r_2 \leq 4R_0. \quad (34)$$

Indeed, for $d = 2$,

$$\begin{aligned} \bar{\eta}(r_1) - \bar{\eta}(r_2) &\geq \left[\left(\frac{1+r_1^2}{1+r_2^2} \right)^{-\theta} - 1 \right] (1+r_2^2)^{-\theta} \\ &\geq CR_0^{-2\theta} \log \frac{1+r_2^2}{1+r_1^2} \geq CR_0^{-2\theta} \log \frac{1+r_2}{1+r_1}, \end{aligned}$$

where we used the fact $a \geq \log(1 + a)$ for $a \geq 0$ in the second inequality. For $d \geq 3$, recalling that $\theta \geq d/2$ in (31),

$$\begin{aligned} \bar{\eta}(r_1) - \bar{\eta}(r_2) &= (1+r_1^2)^{-\theta} - (1+r_2^2)^{-\theta} \\ &\geq (1+r_2^2)^{0.5d-1-\theta} [(1+r_1^2)^{1-0.5d} - (1+r_2^2)^{1-0.5d}] \\ &\gtrsim R_0^{d-2-2\theta} [(1+r_1)^{2-d} - (1+r_2)^{2-d}] \end{aligned}$$

Hence, we obtain $H_R \gtrsim R_0^{d-2-2\theta} (U(|x| + 1) - U(R + 1))$ for this case.

It remains to consider the case $R \geq 4R_0$. To this end, we will prove

$$(G_R \stackrel{(33)}{\gtrsim}) H_R \geq \ell \quad \text{in } \bar{B}_R. \quad (35)$$

Assume by contradiction that (35) fails, i.e., $\max_{\bar{B}_R}(\ell - H_R) > 0$. By Lemma 18(a), $\max_{\bar{B}_R}(\ell - H_R)$ is achieved outside of B_{R_0} . Further, note that $(\ell - H_R)|_{\partial B_R} \leq 0$. By Lemma 18 (e), (d), and the maximum principle,

$$\max_{\bar{B}_R \setminus B_{R/2}}(\ell - H_R) \leq \max_{\partial(B_R \setminus \bar{B}_{R/2})}(\ell_3 - H_R) \leq 0 \vee \max_{B_{R/2} \setminus B_{R_0}}(\ell - H_R).$$

Hence, if $\max_{\bar{B}_R}(\ell - H_R) > 0$, then there exists $x_0 \in B_{R/2} \setminus B_{R_0}$ such that

$$(\ell - H_R)(x_0) = \max_{\bar{B}_R}(\ell - H_R).$$

Since $x_0 \in B_{R/2} \setminus B_{R_0}$, by Lemma 18(c), (d),

$$(\ell_2 - H_R)(x_0) \geq \max_{\bar{B}_{|x_0|/2}(x_0)}(\ell_2 - H_R),$$

which is equivalent to

$$(R_0^{d-2-2\theta} \xi - H_R)(x_0) \geq \max_{\bar{B}_{|x_0|/2}(x_0)}(R_0^{d-2-2\theta} \xi - H_R).$$

Without loss of generality, assume $\bar{a} = I$, and set, for $y \in \mathbb{R}^d \setminus \{0\}$,

$$\tilde{\xi}(y) := \xi(y) - c|x_0|^{-(2+\delta)}|\xi(x_0)||y - x_0|^2,$$

where $c > 0$ is chosen so that (Lemma 12(ii)) $-\Delta \tilde{\xi} \leq 0$ for $|y| \geq C$. Then

$$(R_0^{d-2-2\theta} \tilde{\xi} - H_R)(x_0) \geq \max_{\partial B_{|x_0|/2}(x_0)}(R_0^{d-2-2\theta} \tilde{\xi} - H_R) + cR_0^{d-2-2\theta}|x_0|^{-\delta}|\xi(x_0)|. \quad (36)$$

Next, let $g(x) = \tilde{\xi}(x_0 + \frac{|x_0|}{2}x)$, and let v be the solution of

$$\begin{cases} L_\omega v(x) = 0 & x \in B_{|x_0|/2}(x_0) \\ v(x) = R_0^{d-2-2\theta} g\left(\frac{x-x_0}{|x-x_0|}\right) & x \in \partial B_{|x_0|/2}(x_0). \end{cases}$$

Note that v is close to $R_0^{d-2-2\theta} \tilde{\xi}$ on $\partial B_{|x_0|/2}(x_0)$ in the sense that $\left|g\left(\frac{x-x_0}{|x-x_0|}\right) - \tilde{\xi}(x)\right| \leq C|x_0|^{-1}|\xi(x_0)|$ for $x \in \partial B_{|x_0|/2}(x_0)$. Comparing the L_ω -harmonic functions v and H_R in $\bar{B}_{|x_0|/2}(x_0)$ via the maximum principle, we have, for $x \in B_{|x_0|/2}(x_0)$,

$$H_R(x) + \max_{\partial B_{|x_0|/2}(x_0)}(R_0^{d-2-2\theta} \tilde{\xi} - H_R) \geq v(x) - cR_0^{d-2-2\theta}|x_0|^{-1}|\xi(x_0)|. \quad (37)$$

By Proposition C, for $x \in B_{|x_0|/2}(x_0)$,

$$v(x) \geq \bar{v}\left(\frac{x-x_0}{|x_0|/2}\right) - CR_0^{d-2-2\theta}|x_0|^{-\beta}[g]_{C^{0,1}(\partial B_1)} \geq \bar{v}\left(\frac{x-x_0}{|x_0|/2}\right) - CR_0^{d-2-2\theta}|x_0|^{-\beta}|\xi(x_0)|, \quad (38)$$

where \bar{v} solves

$$\begin{cases} \operatorname{tr}(\bar{a}D^2\bar{v}) = \Delta\bar{v} = 0 & x \in \mathbb{B}_1 \\ \bar{v}(x) = R_0^{d-2-2\theta} g(x) & x \in \partial\mathbb{B}_1. \end{cases}$$

Furthermore, using the fact that $\Delta g(x) = |x_0|^2 \Delta \tilde{\xi}(x_0 + \frac{|x_0|}{2}x) \geq 0$ for $x \in \mathbb{B}_1$, we get $\bar{v} \geq R_0^{d-2-2\theta} g$ in \mathbb{B}_1 . Therefore, by (37), (38), for $x \in B_{|x_0|/2}(x_0)$,

$$R_0^{d-2-2\theta} g\left(\frac{x-x_0}{|x_0|/2}\right) - H_R(x) \leq \max_{\partial B_{|x_0|/2}(x_0)} (R_0^{d-2-2\theta} \tilde{\xi} - H_R) + C R_0^{d-2-2\theta} |x_0|^{-\beta} |\xi(x_0)|.$$

Noting that $g\left(\frac{x-x_0}{|x_0|/2}\right) = \tilde{\xi}(x)$, the above inequality contradicts (36) since $|x_0| > R_0 \geq K$, where K is chosen to be sufficiently large. Inequality (35) is proved.

When $x \in B_{R/2} \setminus B_{R_0}$, we have $G_R \stackrel{(35)}{\gtrsim} \ell_2$, and, by Lemma 12(i),

$$\begin{aligned} \ell_2 &\geq R_0^{d-2-2\theta} [\bar{\xi}(|x|) - \bar{\xi}(R)] \\ &\gtrsim R_0^{d-2-2\theta} \int_{|x|}^R r^{1-d} dr \\ &\gtrsim R_0^{d-2-2\theta} (U(|x|+1) - U(R+1)). \end{aligned} \quad (39)$$

When $x \in B_{R_0}$,

$$\begin{aligned} G_R &\stackrel{(35)}{\gtrsim} \ell_1 = E_\omega^x[\ell_2(X_{s_0}) + \eta(x) - \eta(X_{s_0})] \\ &\stackrel{(39),(34)}{\gtrsim} R_0^{d-2-2\theta} E_\omega^x \left[U(|X_{s_0}|+1) - U(R+1) + U(|x|+1) - U(|X_{s_0}|+1) \right] \\ &= R_0^{d-2-2\theta} [U(|x|+1) - U(R+1)]. \end{aligned}$$

Finally, when $x \in B_R \setminus B_{R/2}$, using the inequality $e^a \geq 1 + a$ for $a \geq 0$,

$$\begin{aligned} G_R &\stackrel{(35)}{\gtrsim} \ell_3 \gtrsim R_0^{d-2-2\theta} R^{2-d} \left(e^{\gamma(1-|x|^2/R^2)} - 1 \right) \\ &\gtrsim R_0^{d-2-2\theta} R^{2-d} \left(1 - \frac{|x|}{R} \right). \end{aligned}$$

For $d = 2$, note that $1 - \frac{|x|}{R} \asymp \frac{R}{|x|} - 1 \geq \log \frac{R}{|x|}$. For $d = 3$, clearly

$$R^{2-d} \left(1 - \frac{|x|}{R} \right) \gtrsim |x|^{2-d} \left(1 - \left(\frac{|x|}{R} \right)^{d-2} \right).$$

Our proof is complete. □

Lemma 19. *Assume (A1), (A2). When $d = 2$, then for \mathbb{P} -almost all ω ,*

$$c\rho(\omega) \leq \liminf_{|x| \rightarrow \infty} \frac{A^\omega(x)}{\log|x|} \leq \overline{\lim}_{|x| \rightarrow \infty} \frac{A^\omega(x)}{\log|x|} \leq C\rho(\omega).$$

3 Heat kernel bounds and consequences

3.1 Integrability of ρ and the heat kernel bounds

Using bounds of the Green functions, we will obtain the exponential integrability (under \mathbb{P}) of the Radon-Nikodym derivative $\rho(\omega)$ (defined in (8)) and the heat kernel of the RWRE.

The goal of this section is to prove Theorem 4. Recall the continuous-time RWRE in Definition 2 and its transition kernel $p_t(x, y)$. We remark that for the time continuous random walk (Y_t) , setting

$$\tau^Y = \tau_R^Y := \inf\{t \geq 0 : Y_t \notin B_R\}, \quad (40)$$

the corresponding Green functions of (Y_t) can be defined similarly as

$$\int_0^\infty p_t^\omega(x, S) dt, \quad d \geq 3, \quad \text{and} \quad E_\omega^x \left[\int_0^{\tau^Y} \mathbb{1}_{Y_t \in S} dt \right], \quad d \geq 2,$$

and they have the same values as $G(x, S)$ and $G_R(x)$, respectively. Thus we do not need to distinguish notations in discrete and continuous time cases and use $G(x, S)$ and $G_R(x)$ to denote Green's functions in both settings.

Corollary 20. *Assume (A1), (A2) and $d = 2$. For any $\varepsilon > 0$, there exists a random variable $\mathcal{H} = \mathcal{H}(\omega, d, \kappa, \varepsilon) > 0$ with $\mathbb{E}[\exp(c\mathcal{H}^{d-\varepsilon})] < \infty$ such that \mathbb{P} -almost surely, for $R > 0$,*

$$\int_0^{R^2} p_t^\omega(x, 0) dt \lesssim \mathcal{H} \left(1 + \log \frac{R+1}{|x|+1}\right), \quad \forall x \in B_R.$$

Proof. We only consider $R \geq 4$. Let $R_k := 2^{k-1}R$, and define recursively $T_0 = 0$,

$$T_k := \min\{t \geq T_{k-1} : Y_t \notin B_{R_k}\}, \quad k \in \mathbb{N}.$$

Set $N_R = \max\{n \geq 0 : T_n \leq R^2\}$. Then, using the strong Markov property, a.s.,

$$\begin{aligned} E_\omega^x \left[\int_0^{R^2} \mathbb{1}_{Y_t=0} dt \right] &\leq E_\omega^x \left[\sum_{k=1}^\infty \int_{T_{k-1}}^{T_k} \mathbb{1}_{\{Y_t=0, T_{k-1} \leq R^2\}} dt \right] \\ &\leq \sum_{k=1}^\infty E_\omega^x \left[G_{R_k}^\omega(Y_{T_{k-1}}) \mathbb{1}_{\{T_{k-1} \leq R^2\}} \right] \\ &\lesssim \mathcal{H} \left(G_R^\omega(x) + \sum_{k=2}^\infty P_\omega^x(T_{k-1} \leq R^2) \right), \end{aligned}$$

where we used (by Theorem 10) the fact $G_{R_k}^\omega(Y_{T_{k-1}}) \lesssim \mathcal{H}$, $k \geq 2$, in the last inequality. Note that (Y_t) is a martingale. By Hoeffding's inequality, for $k \geq 1$,

$$P_\omega^x(T_k \leq R^2) \leq P_\omega^x \left(\sup_{t \leq R^2} |Y_t| \geq R_k \right) \leq C e^{-cR_k^2/R^2} \leq C \exp(-c4^k).$$

The conclusion follows. \square

Proof of Theorem 4: First, we will show the upper bounds in (a) and (b). To this end, for $r \geq 1$, we take $x_0 \in \partial B_r$. We claim that

$$|x_0|^{2-d} \mathcal{H}^{d-1} \gtrsim \int_{r^2/2}^{r^2} p_t^\omega(x_0, 0) dt \gtrsim r^2 \frac{\rho_\omega(0)}{\rho_\omega(B_r)}. \quad (41)$$

Indeed, the lower bound of (41) follows from integrating the lower bound of (10). For $d = 2$, the upper bound in (41) is a consequence of Corollary 20. When $d \geq 3$, the upper bound of (41) follows from Corollary 11. Note that $|x_0| \asymp r$. The upper bound in (b) is proved. The upper bound in (a) then follows from taking $r \rightarrow \infty$ and the ergodic theorem.

To obtain the lower bound in (b), for $r \geq 5$. Recall τ_r in (15) and $G_r(\cdot, \cdot)$ in Definition 9. For any fixed $y_0 \in \partial B_{r/2}$, the function $v(x) = G_r(y_0, x)/\rho_\omega(x)$ solves the *adjoint equation*

$$L_\omega^* v(x) := \sum_y \omega^*(x, y)[v(y) - v(x)] = 0, \quad x \in B_{r/2}, \quad (42)$$

where

$$\omega^*(x, y) := \rho_\omega(y)\omega(y, x)/\rho_\omega(x). \quad (43)$$

Here, we used the facts that $\sum_y \rho_\omega(y)\omega(y, x) = \rho_\omega(x)$ and $\sum_y G_r(y_0, y)\omega(y, x) = G_r(y_0, x)$. By the Harnack inequality for the adjoint operator [13, Theorem 6], we have $v(0) \asymp v(x)$ for all $x \in B_{r/4}$. Hence

$$G_r(y_0, 0) \frac{\rho_\omega(B_{r/4})}{\rho_\omega(0)} \asymp G_r(y_0, B_{r/4}). \quad (44)$$

Moreover, since $(|X_n|^2 - n)$ is a martingale under P_ω , by the optional stopping lemma we get $E_\omega^{y_0}[|X_{\tau_r}|^2 - \tau_r] = |y_0|^2 \geq 0$, and so

$$G_r(y_0, B_{r/4}) \leq E_\omega^{y_0}[\tau_r] \leq E_\omega^{y_0}[|X_{\tau_r}|^2] \leq Cr^2.$$

The above inequality, together with (44) and Theorem 10, yields

$$\frac{\rho_\omega(0)}{\rho_\omega(B_{r/4})} \gtrsim \frac{G_r(y_0, 0)}{G_r(y_0, B_{r/4})} \gtrsim \frac{\mathcal{H}^{-s} r^{2-d}}{r^2} \gtrsim \mathcal{H}^{-s} r^{-d}.$$

The lower bound in Theorem 4(b) follows. Letting $r \rightarrow \infty$, we also get the lower bound in (a). \square

3.2 Optimal semigroup decay for $d \geq 3$

The Efron-Stein inequality (47) of Boucheron, Bousquet, and Massart [8] will be used in our derivation of the variance decay for the semi-group.

Let $\omega'(x), x \in \mathbb{Z}^d$, be independent copies of $\omega(x), x \in \mathbb{Z}^d$. For any $y \in \mathbb{Z}^d$, let $\omega'_y \in \Omega$ be the environment such that

$$\omega'_y(x) = \begin{cases} \omega(x) & \text{if } x \neq y, \\ \omega'(y) & \text{if } x = y. \end{cases}$$

That is, ω'_y is a modification of ω only at location y . For any measurable function Z of the environment ω , we write, for $y \in \mathbb{Z}^d$,

$$Z'_y = Z(\omega'_y), \quad \partial'_y Z(\omega) = Z'_y - Z, \quad (45)$$

and set

$$V(Z) = \sum_{y \in \mathbb{Z}^d} (\partial'_y Z)^2. \quad (46)$$

By an L_p version of Efron-Stein inequality [8, Theorem 3], for $q \geq 2$,

$$\mathbb{E}[|Z - \mathbb{E}Z|^q] \leq Cq^{q/2} \mathbb{E}[V^{q/2}]. \quad (47)$$

Following the strategy of [18], our proof of the diffusive decay of the semi-group $\{P_t\}$ will make use of the Efron-Stein type inequality (47) and the Duhamel representation formula (50) for the vertical derivative. Let us reemphasize that, in the non-divergence form setting, there is no deterministic Gaussian bounds for the heat kernel, and the steady state \mathbb{Q} of the environment process $(\bar{\omega}')_{t \geq 0}$ is not the same as the original measure \mathbb{P} . To overcome these difficulties, we employ crucially the heat kernel estimates and the (negative and positive) moment bounds of the Radon-Nikodym derivative $\frac{d\mathbb{P}}{d\mathbb{Q}}$ in Theorem 4.

For any $\zeta \in L^1(\Omega)$, we write

$$v(t) := P_t \zeta(\omega).$$

Then, its stationary extension $\bar{v}(t, x)$ solves the parabolic equation

$$\begin{cases} \partial_t \bar{v}(t, x) - L_\omega \bar{v}(t, x) = g(t, x) & t \geq 0, x \in \mathbb{Z}^d, \\ v(0, x) = g_0(x) & x \in \mathbb{Z}^d, \end{cases} \quad (48)$$

with $g(t, x) = 0$ and $g_0(x) = \bar{\zeta}(x; \omega)$. In general, the solution of (48) can be represented by Duhamel's formula

$$\bar{v}(t, x) = \sum_z p_t^\omega(x, z) g_0(z) + \int_0^t p_{t-s}^\omega(x, z) g(s, z) ds. \quad (49)$$

To apply (47), recall notations Z'_y and $\partial'_y Z$ in (45). By enlarging the probability space, we still use \mathbb{P} to denote the joint law of (ω, ω') . For $y \in \mathbb{Z}^d$, applying ∂'_y to (48), we get that $\partial'_y \bar{v}$ satisfies

$$\begin{cases} \partial_t (\partial'_y \bar{v})(t, x) - L_\omega (\partial'_y v)(t, x) = (\partial'_y \omega_i(x)) \nabla_i^2 \bar{v}'_y(t, x) & t \geq 0, x \in \mathbb{Z}^d, \\ (\partial'_y \bar{v})(0, x) = \partial'_y \bar{\zeta}(x), & x \in \mathbb{Z}^d. \end{cases}$$

Here we used the convention of summation over repeated integer indices. Hence, by formula (49), $\partial'_y \bar{v}$ has the representation

$$\begin{aligned} \partial'_y \bar{v}(t, x) &= \sum_z \left[p_t^\omega(x, z) \partial'_y \bar{\zeta}(z) + \sum_{i=1}^d \int_0^t p_{t-s}^\omega(x, z) (\partial'_y \omega_i(z)) \nabla_i^2 \bar{v}'(s, z) ds \right] \\ &= \sum_{z \in y + \text{Supp}(\zeta)} p_t^\omega(x, z) \partial'_y \bar{\zeta}(z) + \sum_{i=1}^d \int_0^t p_{t-s}^\omega(x, y) (\partial'_y \omega_i(y)) \nabla_i^2 \bar{v}'(s, y) ds, \end{aligned} \quad (50)$$

where in the last equality we used the fact $\partial'_y \omega_i(z) = 0$ for $y \neq z$, and that $\partial'_y \bar{\zeta}(z) = 0$ for $z \notin y + \text{Supp}(\zeta)$.

Proof of Theorem 5: Recall the notation $V(\cdot)$ in (46). We write

$$K(y, s) = K(y, s; \omega, \omega') := \sum_{i=1}^d (\partial'_y \omega_i(y)) \nabla_i^2 \bar{v}'(s, y).$$

We also write $\mathcal{H}_z := \mathcal{H}(\theta_z \omega)$ and $S := \text{Supp}(\zeta)$. Without loss of generality, assume $E_{\mathbb{Q}} \zeta = 0$. Using (50), for any $p > 1$,

$$\begin{aligned} \|V(v(t))\|_p &= \left\| \sum_y (\partial'_y \bar{v}(t, 0))^2 \right\|_p \\ &\lesssim \left\| \sum_y \left(\sum_{z \in S} p_t^\omega(0, z + y) \right)^2 \right\|_p + \left\| \sum_y \left(\int_0^t p_{t-s}^\omega(0, y) K(y, s) ds \right)^2 \right\|_p \\ &\lesssim (\#S)^2 \left\| \sum_y p_t^\omega(0, y)^2 \right\|_p + \int_0^t \left\| \sum_y p_{t-s}^\omega(0, y)^2 K(y, s)^2 \right\|_p^{1/2} ds \\ &=: (\#S)^2 \text{I} + \text{II}^2, \end{aligned} \quad (51)$$

where $\#S$ denote the cardinality of S , and in the last inequality we applied the Cauchy-Schwarz inequality and Minkowski's integral inequality to the two norms respectively. Then, by Theorem 4(c),

$$\begin{aligned} \text{I} &= \left\| \sum_y p_t^\omega(0, y)^2 \right\|_p \lesssim (1+t)^{-d} \left\| \sum_z \mathcal{H}_z^{2(d-1)} e^{-c\mathfrak{h}(|z|, t)} \right\|_p \\ &\lesssim (1+t)^{-d} \left\| \sum_z e^{-c\mathfrak{h}(|z|, t)} \mathcal{H}_z^{2(d-1)} \right\|_p \\ &\lesssim (1+t)^{-d/2} \left\| \mathcal{H}^{2(d-1)p} \right\|_1^{1/p}, \end{aligned} \quad (52)$$

where in the last inequality we used the translation invariance of \mathbb{P} .

Further, using Theorem 4(c) again,

$$\begin{aligned} \Pi &\lesssim \int_0^t (1+t-s)^{-d/2} \left(\sum_y e^{-c\mathfrak{b}(|y|,t-s)} \|\mathcal{H}_y^{2(d-1)} K(y,s)^2\|_p \right)^{1/2} ds \\ &\lesssim \int_0^t (1+t-s)^{-d/4} \|\mathcal{H}^{2(d-1)p} K(0,s)^2\|_1^{1/(2p)} ds \end{aligned} \quad (53)$$

where in the last inequality we used the translation-invariance of \mathbb{P} and the fact that $\|\bar{v}\|_\infty \lesssim 1$.

Let $p' > 1$ denote the Hölder conjugate of $p > 1$. Since

$$\begin{aligned} \|\mathcal{H}^{2(d-1)p} K(0,s)^2\|_1 &\leq \|\mathcal{H}^{2(d-1)p}\|_{p'} \|K(0,s)^2\|_p \\ &= C_p \|K(0,s)^2\|_p \\ &\leq C_p \sum_{i=1}^d \|\nabla_i^2 \bar{v}(s,0)\|_{2p}^2 \\ &\leq C_p \sum_{e:|e|=1} \|\bar{v}(s,e) - \bar{v}(s,0)\|_1^{2/p} \\ &\stackrel{\text{Hölder}}{\leq} C_p \|\rho_\omega^{-1/p}\|_{p'}^{1/p} \sum_{e:|e|=1} \|\rho_\omega |\bar{v}(s,e) - \bar{v}(s,0)|^2\|_1^{1/p^2} \end{aligned} \quad (54)$$

where in the third inequality we used the fact $\|\bar{v}\|_\infty \leq 1$. Then, setting

$$u(t) := \text{Var}_{\mathbb{Q}}(v(t)),$$

by (54), Theorem 4(b) and Lemma 21, we obtain

$$\|\mathcal{H}^{2(d-1)p} K(0,s)^2\|_1 \leq C_p \sum_e E_{\mathbb{Q}}[|\bar{v}(s,e) - \bar{v}(s,0)|^2]^{1/p^2} \leq C_p \left(-\frac{d}{dt}u\right)^{1/p^2}. \quad (55)$$

This inequality, together with (51),(52), (53), implies

$$\|V(v(t))\|_p^{1/2} \lesssim_p (1+t)^{-d/4} + \int_0^t (1+t-s)^{-d/4} \left(-\frac{d}{dt}u\right)^{1/(2p^3)} ds,$$

where \lesssim_p means that the multiplicative constant depends on (p, d, κ) .

Furthermore, by Hölder's inequality and Theorem 4(a),

$$u(t) \leq E_{\mathbb{Q}}[(v(t) - \mathbb{E}v(t))^2] \leq \|\rho_\omega\|_{p'} \|v(t) - \mathbb{E}v(t)\|_p \lesssim_p \|V(v(t))\|_p$$

where we applied (47) in the last inequality.

Therefore, we conclude that, for any $p > 1$,

$$u(t)^{1/2} \lesssim_p (1+t)^{-d/4} + \int_0^t (1+t-s)^{-d/4} \left(-\frac{d}{dt}u(s)\right)^{1/(2p^3)} ds.$$

When $d \geq 3$, we can take $p > 1$ sufficiently close to 1 and apply [1, Lemma 3.5]) (Under the notation of [1], we apply it to the case $\gamma = d/4$ and $\delta = 1/p^3$.) to obtain

$$\text{Var}_{\mathbb{Q}}(v(t)) = u(t) \lesssim (1+t)^{-d/2}.$$

Thus, (12) is proved. By Hölder's inequality,

$$\|v(t)\|_1 \leq \|\rho_\omega^{-1}\|_1^{1/2} \|\rho_\omega v(t)^2\|_1^{1/2} \leq C u(t)^{1/2} \leq C(1+t)^{-d/4}.$$

Then, by the triangle inequality, we get $E_{\mathbb{Q}}[(v(t) - \mathbb{E}v(t))^2] \lesssim (1+t)^{-d/2}$. By Hölder's inequality, for any $q > 1$,

$$\| |v(t) - \mathbb{E}v(t)|^{2/q} \|_1 \leq \|\rho_\omega^{-1/q}\|_{q'} \|\rho_\omega^{1/q} |v(t) - \mathbb{E}v(t)|^{2/q}\|_q \lesssim_q E_{\mathbb{Q}}[(v(t) - \mathbb{E}v(t))^2]^{1/q}.$$

Display (13) is proved. \square

Lemma 21. For any bounded measurable function $\zeta \in \mathbb{R}^\Omega$,

$$\frac{d}{dt} \text{Var}_{\mathbb{Q}}(v(t)) \lesssim - \sum_{e:|e|=1} E_{\mathbb{Q}} [(\bar{v}(t, e) - \bar{v}(t, 0))^2],$$

where L_ω only acts on the spatial variable.

Proof. Without loss of generality, assume $\mathbb{E}[v(t)] = 0$. Then

$$\begin{aligned} \frac{d}{dt} E_{\mathbb{Q}}[v(t)^2] &= 2E_{\mathbb{Q}}[v(t)L_\omega \bar{v}(t, 0)] \\ &= E_{\mathbb{Q}} \left[L_\omega(\bar{v}(t, 0)^2) - \sum_{e:|e|=1} \omega(0, e)(\bar{v}(t, e) - \bar{v}(t, 0))^2 \right] \\ &= - \sum_{e:|e|=1} E_{\mathbb{Q}} [\omega(0, e)(\bar{v}(t, e) - \bar{v}(t, 0))^2] \end{aligned}$$

where in the last equality we used the fact that (since $(\bar{\omega}^t)_{t \geq 0}$ is stationary under $\mathbb{Q} \times P_\omega$) for any $f \in L^1(\mathbb{Q})$, $E_{\mathbb{Q}}[L_\omega \bar{f}(0; \omega)] = 0$. The lemma follows by the uniform ellipticity assumption. \square

3.3 Proof of Corollary 7: Existence of a stationary corrector for $d \geq 5$

Proof of Corollary 7: Without loss of generality, assume $E_{\mathbb{Q}}[\zeta] = 0$. By Theorem 5, since $\int_0^\infty (1+t)^{-d/4} dt < \infty$ when $d \geq 5$, the limit

$$\phi(\omega) = \lim_{t \rightarrow \infty} \phi_t(\omega) := - \lim_{t \rightarrow \infty} \int_0^t (P_s \zeta) ds \quad (56)$$

exists in $L^p(\mathbb{P})$ form any $p \in (0, 2)$. By (48), $\bar{\phi}_t(x)$ satisfies

$$L_\omega \bar{\phi}_t(x) = \zeta(\theta_x \omega) - P_t \zeta(\theta_x \omega), \quad x \in \mathbb{Z}^d.$$

Note that by Theorem 5, $\lim_{t \rightarrow \infty} P_t \zeta = 0$ in $L^p(\mathbb{P})$, $\forall p \in (0, 2)$. Therefore, taking $L^p(\mathbb{P})$ limits as $t \rightarrow \infty$, we conclude that $\bar{\phi}$ satisfies $L_\omega \bar{\phi}(x) = \bar{\zeta}(x)$, $x \in \mathbb{Z}^d$. \square

A Appendix

Define the parabolic operator \mathcal{L}_ω as

$$\mathcal{L}_\omega u(x, t) = \sum_{y: y \sim x} \omega(x, y)[u(y, t) - u(x, t)] - \partial_t u(x, t)$$

for every function $u : \mathbb{Z}^d \times \mathbb{R} \rightarrow \mathbb{R}$ which is differentiable in t .

Theorem A.1. ([13, Proposition 5]) Assume $\frac{\omega}{\text{tr}\omega} > 2\kappa I$ for some $\kappa > 0$ and $R > 0$. Any non-negative function u with $\mathcal{L}_\omega u = 0$ in $B_{2R} \times (0, 4R^2)$ satisfies

$$\sup_{B_R \times (R^2, 2R^2)} u \leq C \inf_{B_R \times (3R^2, 4R^2)} u.$$

Corollary A.2. Assume $\frac{\omega}{\text{tr}\omega} > 2\kappa I$ for some $\kappa > 0$ and $(x_0, t_0) \in \mathbb{Z}^d \times \mathbb{R}$. There exists $\gamma = \gamma(d, \kappa) \in (0, 1)$ such that any non-negative function u with $\mathcal{L}_\omega u = 0$ in $B_R(x_0) \times (t_0 - R^2, t_0)$, $R > 0$, satisfies

$$|u(\hat{x}) - u(\hat{y})| \leq C \left(\frac{r}{R}\right)^\gamma \sup_{B_R(x_0) \times (t_0 - R^2, t_0)} u$$

for all $\hat{x}, \hat{y} \in B_r(x_0) \times (t_0 - r^2, t_0)$ and $r \in (0, R)$.

A.1 Proof of Proposition C

Before giving a proof, recall that by [19, Proposition 2.1], for any $p \in (0, d)$, there exists δ_p depending on (d, κ, p) such that for any $R > 0$, the solution $\phi : \bar{B}_R \rightarrow \mathbb{R}$ of

$$\begin{cases} L_\omega \phi = \bar{a} - a & \text{in } B_R, \\ \phi = 0 & \text{on } \partial B_R \end{cases} \quad (57)$$

satisfies

$$\mathbb{P}(\max_{B_R} |\phi| \geq CR^{2-\delta_p}) \leq C \exp(-cR^p). \quad (58)$$

Set $\delta := \delta_1$. For $q \in (0, d)$, let $\gamma = \gamma(d, \kappa, q)$ be the constant

$$\gamma = \min \left\{ \frac{d-q}{d(1+\delta)}, \frac{1}{2} \right\} \quad (59)$$

and set

$$R_0 := R^\gamma, \quad \sigma = \min\{n \geq 0 : X_n - X_0 \notin B_{R_0}\}.$$

Let

$$\omega_0 = \frac{\omega}{\text{tr}(\omega)}, \quad \psi_0 = \psi_0(\omega) = \frac{\psi}{\text{tr}(\omega)}.$$

Following [19, Definition 4.1], we define bad points.

Definition 22. Let $\delta = \delta(d, \kappa)$ be as above. We say that a point is good (and otherwise bad) if for any $\zeta(\omega) \in \{\psi_0, \omega_0\}$,

$$\left| E_\omega \left[\sum_{i=0}^{\sigma-1} (E_{\mathbb{Q}}[\zeta] - \zeta(\bar{\omega}_i)) \right] \right| \leq C \|\zeta\|_\infty R_0^{2-\delta}.$$

Note that by (58), $\mathbb{P}(x \text{ is bad}) \leq C e^{-cR_0}$.

We will give first the proof for the special case $g \in C^{2,\alpha}(\partial\mathbb{B}_1)$. It is a small modification of the proof of [19, Theorem 1.5].

Proof of Proposition C for the case $f \in C^\alpha(\mathbb{B}_1)$, $g \in C^{2,\alpha}(\partial\mathbb{B}_1)$ and $y = 0$: Note that if $g \in C^{2,\alpha}(\partial\mathbb{B}_1)$, it can be extended to be a function $\tilde{g} \in C^{2,\alpha}(\mathbb{B}_2)$ such that

$$|\tilde{g}|_{2,\alpha;\mathbb{B}_2} \leq C |g|_{2,\alpha;\partial\mathbb{B}_1}.$$

By [17, Theorem 6.6],

$$|\bar{u}|_{2,\alpha;\mathbb{B}_1} \lesssim |f|_{0,\alpha;\mathbb{B}_1} \left\| \frac{\psi}{\text{tr}(\omega)} \right\|_\infty + |g|_{2,\alpha;\partial\mathbb{B}_1} =: A \quad (60)$$

Step 1. Set $\bar{u}_R(x) = \bar{u}(x/R)$ for $x \in \bar{\mathbb{B}}_R$. We will show that in B_R , \bar{u}_R is very close to the solution $\hat{u} : \bar{B}_R \rightarrow \mathbb{R}$ of

$$\begin{cases} L_\omega \hat{u} = \frac{1}{2} \text{tr}[\omega_0 D^2 \bar{u}_R] & \text{in } B_R \\ \hat{u} = g\left(\frac{x}{|x|}\right) & \text{on } \partial B_R. \end{cases}$$

To this end, define u_+, u_- by $u_\pm(x) = \tilde{g}\left(\frac{x}{R+1}\right) \pm CA \frac{(R+1)^2 - |x|^2}{R^2}$, $x \in \bar{\mathbb{B}}_{R+1}$. Here A is as defined in (60). Then, for $x \in \mathbb{B}_{R+1}$, taking C large enough,

$$\text{tr}[\bar{a} D^2 (u_+ - \bar{u}_{R+1})] \leq \frac{c}{R^2} (|g|_{2,\alpha;\mathbb{B}_1} + |f|_{0;\mathbb{B}_1} - CA) \leq 0,$$

and similarly $\text{tr}[\bar{a} (u_- - \bar{u}_{R+1})] \geq 0$. The comparison principle then yields

$$u_- \leq \bar{u}_{R+1} \leq u_+ \quad \text{in } \mathbb{B}_{R+1}.$$

In particular, for $x \in \partial B_R$, $|\bar{u}_{R+1}(x) - \tilde{g}\left(\frac{x}{R+1}\right)| \lesssim A \frac{(R+1)^2 - |x|^2}{R^2} \lesssim \frac{A}{R}$ and so

$$\max_{\partial B_R} |\hat{u} - \bar{u}_{R+1}| = \max_{x \in \partial B_R} \left| g\left(\frac{x}{|x|}\right) - \bar{u}_{R+1}(x) \right| \lesssim \frac{A}{R}. \quad (61)$$

Moreover, noting that $D^2 \bar{u}_R(x) = R^{-2} D^2 \bar{u}\left(\frac{x}{R}\right)$, in B_R ,

$$\begin{aligned} |L_\omega(\hat{u} - \bar{u}_{R+1})| &= |\text{tr}[\omega_0 (D^2 \bar{u}_R - \nabla^2 \bar{u}_{R+1})]| \\ &\leq |\text{tr}[\omega_0 (D^2 \bar{u}_R - D^2 \bar{u}_{R+1})]| + |\text{tr}[\omega_0 (D^2 \bar{u}_{R+1} - \nabla^2 \bar{u}_{R+1})]| \\ &\lesssim R^{-2-\alpha} |\bar{u}|_{2,\alpha;\bar{\mathbb{B}}_1} \stackrel{(60)}{\lesssim} AR^{-2-\alpha}. \end{aligned} \quad (62)$$

Hence, by the ABP maximum principle [19, Lemmas 2.3 and 2.4], (61) and (62) imply $\max_{B_R} |\hat{u} - \bar{u}_{R+1}| \lesssim AR^{-\alpha}$, and so, by (60),

$$\max_{B_R} |\hat{u} - \bar{u}_R| \lesssim AR^{-\alpha}.$$

Step 2. Let $v = u - \hat{u}$. Then v solves

$$\begin{cases} L_\omega v = \frac{1}{2} \text{tr}[(\bar{a} - \omega_0) D^2 \bar{u}_R] + \frac{1}{R^2} f\left(\frac{x}{R}\right)(\psi_0 - \bar{\psi}) & \text{in } B_R \\ v = 0 & \text{on } \partial B_R. \end{cases}$$

Note that, for $x \in B_{R-R_0}$ and $y \in B_{R_0}(x)$,

$$\begin{aligned} |D^2 \bar{u}_R(x) - D^2 \bar{u}_R(y)| &\leq R^{-2} \left(\frac{R_0}{R}\right)^\alpha [\bar{u}]_{2,\alpha;\mathbb{B}_1} \stackrel{(60)}{\lesssim} AR^{-2} \left(\frac{R_0}{R}\right)^\alpha, \\ |f\left(\frac{x}{R}\right) - f\left(\frac{y}{R}\right)| &\leq \left(\frac{R_0}{R}\right)^\alpha [f]_{\alpha;\mathbb{B}_1}. \end{aligned}$$

Hence, if $x \in B_{R-R_0}$ is a good point, setting

$$\bar{\omega}_0^i := \omega_0(X_i) \quad \text{and} \quad \psi_0^i := \psi_0(\theta_{X_i} \omega),$$

and noting that $E_\omega^x[\sigma] \leq (R_0 + 1)^2$, we have

$$\begin{aligned} &E_\omega^x[v(X_\sigma) - v(x)] \\ &= E_\omega^x \left[\sum_{i=0}^{\sigma-1} \frac{1}{2} \text{tr}[(\bar{\omega}_0^i - \bar{a}) D^2 \bar{u}_R(X_i)] + \frac{1}{R^2} f\left(\frac{X_i}{R}\right)(\bar{\psi} - \psi_0^i) \right] \\ &\lesssim \text{tr} \left[E_\omega^x \left[\sum_{i=0}^{\sigma-1} (\bar{\omega}_0^i - \bar{a}) D^2 \bar{u}_R(x) \right] + \frac{1}{R^2} f\left(\frac{x}{R}\right) E_\omega^x \left[\sum_{i=0}^{\sigma-1} (\bar{\psi} - \psi_0^i) \right] + A \frac{R_0^\alpha}{R^{2+\alpha}} E_\omega^x[\sigma] \right] \\ &\lesssim \frac{1}{R^2} \left| E_\omega^x \left[\sum_{i=0}^{\sigma-1} (\bar{\omega}_0^i - \bar{a}) \right] \right| [\bar{u}]_{2;\mathbb{B}_1} + \frac{1}{R^2} |f|_{0;\mathbb{B}_1} \left| E_\omega^x \left[\sum_{i=0}^{\sigma-1} (\bar{\psi} - \psi_0^i) \right] \right| + A \left(\frac{R_0}{R}\right)^{2+\alpha} \\ &\lesssim AR^{-2} R_0^{2-\delta} + A \left(\frac{R_0}{R}\right)^{2+\alpha} \lesssim AR^{-2-\alpha\delta\gamma} R_0^2. \end{aligned}$$

Let $\tau_R = \min\{n \geq 0 : X_n \notin B_R\}$ and set

$$w(x) = v(x) + C_1 AR^{-2-\alpha\delta\gamma} E_\omega^x[\tau_R].$$

Then, for any good point $x \in B_{R-R_0}$, by choosing C_1 big enough,

$$E_\omega^x[w(X_\sigma) - w(x)] = E_\omega^x[v(X_\sigma) - v(x)] - C_1 AR^{-2-\alpha\delta\gamma} E_\omega^x[\sigma] < 0,$$

where we used the fact that $E_\omega^x[\sigma] \geq R_0^2$. This implies

$$\partial w(x; B_R) = \emptyset \quad \text{for any good point } x \in B_{R-R_0} \quad (63)$$

where $\partial w(x; B_R)$ denote the sub-differential set of w at x with respect to B_R . For the definition of the sub-differential set and the ABP inequality, we refer to [19, Definition 2.2, Lemmas 2.3 and 2.4]. Next, we will apply the ABP inequality to bound $|v|$ from the above.

By [19, Lemma 2.4], since

$$L_\omega w = L_\omega v - C_1 A R^{-2-\alpha\delta\gamma} \lesssim A R^{-2},$$

we know that $|\partial w(x; B_R)| \lesssim A^d R^{-2d}$ for $x \in B_R$. Let

$$\mathcal{B}_R = \mathcal{B}_R(\omega, \gamma) := \#\text{bad points in } B_{R-R_0},$$

where $\#S$ denotes the cardinality of a set S . Display (63) then yields

$$|\partial w(B_R)| \lesssim [\mathcal{B}_R + \#(B_R \setminus B_{R-R_0})] A^d R^{-2d} \lesssim (\mathcal{B}_R + R^{d-1+\gamma}) A^d R^{-2d}.$$

Hence, by the ABP inequality [19, Lemma 2.4],

$$\min_{B_R} w \geq -C R |\partial w(B_R)|^{1/d} \gtrsim -A(R^{-1} \mathcal{B}_R^{1/d} + R^{-(1-\gamma)/d}).$$

Therefore, noting that $\max_{x \in B_R} E_\omega^x[\tau_R] \leq (R+1)^2$ and choosing $\delta < 1/d$,

$$\min_{B_R} v \geq \min_{B_R} w - C A R^{-\alpha\delta\gamma} \gtrsim -A(R^{-1} \mathcal{B}_R^{1/d} + R^{-\alpha\gamma\delta}).$$

Similar bound for $\min_{B_R} (-v)$ can be obtained by substituting f, g by $-f, -g$ in the problem. Therefore

$$\max_{B_R} |v| \lesssim A(R^{-1} \mathcal{B}_R^{1/d} + R^{-\alpha\gamma\delta}).$$

Step 3. Combining results in Steps 1 and 2, we get

$$\max_{B_R} |u - \bar{u}_R| \lesssim A(R^{-1} \mathcal{B}_R^{1/d} + R^{-\alpha\gamma\delta}).$$

It is shown in Step 6 of [19, Proof of Theorem 1.5] that, with

$$\mathcal{X} = \mathcal{X}(\omega) := \max_{R \geq 1} R^{-\gamma} \mathcal{B}_R^{1/d}, \quad (64)$$

we have $\mathbb{E}[\exp(c\mathcal{X}^d)] < \infty$. Therefore, recalling the values of γ, A in (59), (60), we conclude that

$$\max_{B_R} |u - \bar{u}_R| \lesssim R^{-\alpha\gamma\delta} (1 + R^{-q/d} \mathcal{X}) (|f|_{0,\alpha;\mathbb{B}_1} \|\psi_0\|_\infty + |g|_{2,\alpha;\partial\mathbb{B}_1}). \quad (65)$$

□

In what follows, we will relax the regularity of g to be $C^{0,\alpha}(\partial\mathbb{B}_1)$.

Proof of Proposition C.: First, we consider the case $y = 0$. The function $g \in C^{0,\alpha}(\partial\mathbb{B}_1)$ can be extended into \mathbb{R}^d so that $g \in C^{0,\alpha}(\mathbb{R}^d)$ and

$$|g|_{0,\alpha;\mathbb{B}_2} \leq C|g|_{0,\alpha;\partial\mathbb{B}_1}.$$

We can further obtain a smooth perturbation of it. To this end, let $\rho \in C^\infty(\mathbb{R}^d)$ be a mollifier supported on \mathbb{B}_1 with $\int_{\mathbb{B}_1} \rho dx = 1$. For $h \in (0, 1)$, set $\rho_h(x) := h^{-d} \rho(\frac{x}{h})$, and let $g_h = \rho_h * g$. That is, $g_h(x) = \int_{\mathbb{R}^d} \rho_h(x-z)g(z)dz$. Then g_h satisfies

$$\begin{cases} |g - g_h|_{0,\alpha;\partial\mathbb{B}_1} \leq Ch^\alpha |g|_{0,\alpha;\partial\mathbb{B}_1} \\ |g_h|_{2,\alpha;\partial\mathbb{B}_1} \leq Ch^{-2} |g|_{0,\alpha;\partial\mathbb{B}_1}. \end{cases}$$

Next, for $h \in (0, 1)$, let $v : \bar{B}_R \rightarrow \mathbb{R}$ and $\bar{v} : \bar{\mathbb{B}}_1 \rightarrow \mathbb{R}$ be solutions of

$$\begin{cases} \frac{1}{2}\text{tr}(\omega\nabla^2 v) = \frac{1}{R^2} f(\frac{x}{R})\psi(\theta_x\omega) & \text{in } B_R \\ v(x) = g_h(\frac{x}{R}) & \text{for } x \in \partial B_R \end{cases}$$

and

$$\begin{cases} \text{tr}(\bar{a}D^2\bar{v}) = f\bar{\psi} & \text{in } \mathbb{B}_1 \\ \bar{v} = g_h & \text{on } \partial\mathbb{B}_1. \end{cases}$$

Then, $\max_{\mathbb{B}_1} |\bar{u} - \bar{v}| \leq \max_{\partial\mathbb{B}_1} |g - g_h| \leq h^\alpha |g|_{0,\alpha;\partial\mathbb{B}_1}$, and

$$\max_{B_R} |u - v| \leq \max_{\partial B_R} |g(\frac{x}{R}) - g_h(\frac{x}{R})| \leq Ch^\alpha |g|_{0,\alpha;\partial\mathbb{B}_1}.$$

Moreover, by (65), with $A_1 = \|f\|_{C^{0,\alpha}(\mathbb{B}_1)} \|\frac{\psi}{\text{tr}(\omega)}\|_\infty + [g]_{C^{0,\alpha}(\partial\mathbb{B}_1)}$ as in Proposition C,

$$\begin{aligned} \max_{x \in B_R} |v(x) - \bar{v}(\frac{x}{R})| &\lesssim R^{-\alpha\gamma\delta} (1 + R^{-q/d} \mathcal{X}) (\|f\|_{0,\alpha;\mathbb{B}_1} \|\frac{\psi}{\text{tr}(\omega)}\|_\infty + |g_h|_{2,\alpha;\partial\mathbb{B}_1}) \\ &\lesssim A_1 h^{-2} R^{-\alpha\gamma\delta} (1 + R^{-q/d} \mathcal{X}). \end{aligned}$$

Notice that up to an additive constant, we may assume that $\inf_{\partial B_1} g = 0$, so that $|g|_{0,\alpha;\partial\mathbb{B}_1} \leq C[g]_{0,\alpha;\partial\mathbb{B}_1}$. Therefore, putting $h = R^{-\alpha\gamma\delta/3}$, by the triangle inequality,

$$\max_{x \in B_R} |u - \bar{u}(\frac{x}{R})| \lesssim A_1 R^{-\alpha\gamma\delta/3} (1 + R^{-q/d} \mathcal{X}). \quad (66)$$

We proved Proposition C for the case $y = 0$ with $\beta = \gamma\delta/3$.

Finally, for any $y \in B_{3R}$, it follows from (66) that

$$\max_{x \in B_R(y)} \left| \bar{u}(\frac{x-y}{R}) - u(x) \right| \lesssim A_1 R^{-\alpha\beta} (1 + \mathcal{X}(\theta_y\omega) R^{-q/d}).$$

Let

$$\mathcal{B}_R(y) = \mathcal{B}_R(\theta_y\omega, \gamma) := \#\text{bad points in } B_{R-R_0}(y).$$

Observe that $\mathcal{B}_R(y) \leq \mathcal{B}_{4R}(0)$. Thus, recalling the definition of \mathcal{X} in (64),

$$\mathcal{X}(\theta_y\omega) \leq \max_{R \geq 1} R^{-\gamma} \mathcal{B}_{4R}^{1/d} \leq 4^\gamma \mathcal{X}.$$

Our proof of Proposition C is complete. \square

A.2 Proofs of Lemmas 16 and 13

Proof of Lemma 16: By direct computation, for any $i = 1, \dots, d$, $x \in \mathbb{Z}^d$,

$$\begin{aligned}\partial_i^2 \eta(x) &= \theta(1 + |x|^2)^{-\theta-2} [4(\theta + 1)x_i^2 - 2|x|^2 - 2], \\ |\partial_i^3 \eta(x)| &\leq C\theta^3 |x|(1 + |x|^2)^{-\theta-2}.\end{aligned}\tag{67}$$

Moreover, noting that for any $y \in \mathbb{R}^d$ and $x \notin B_\theta$ with $|y - x| \leq 1$,

$$|\partial_i^3 \eta(y)| \leq C\theta^3 \left(\frac{1+\theta}{\theta}\right)^{\theta+2} |x|(1 + |x|^2)^{-\theta-2} \leq C\theta^3 |x|(1 + |x|^2)^{-\theta-2}$$

and $|\nabla_i^2 \eta(x) - \partial_i^2 \eta(x)| \leq C \sup_{y: |y-x| \leq 1} |\partial_i^3 \eta(y)|$, we have, for a sufficiently large constant $C_0(\kappa) > 1$ and $x \notin B_{C_0\theta^2}$,

$$\begin{aligned}L_\omega \eta(x) &\geq \sum_{i=1}^d \frac{\omega_i(x)}{2\text{tr}\omega(x)} [\partial_i^2 \eta(x) - C\theta^3 |x|(1 + |x|^2)^{-\theta-2}] \\ &\stackrel{(67)}{\geq} \theta(1 + |x|^2)^{-\theta-2} [4\kappa(\theta + 1)|x|^2 - |x|^2 - 1 - C\theta^2 |x|] > 0.\end{aligned}$$

On the other hand, for $x \in B_{C_0\theta^2}$, clearly $L_\omega \eta(x) \geq -\eta(x) \geq -1$.

The lemma is proved. \square

Proof of Lemma 13. Computations show that, for $x \neq 0$, $i = 1, \dots, d$,

$$\begin{aligned}\partial_i^2 e^{-2\alpha|x|/R} &= \left(-\frac{2\alpha}{|x|} + \frac{2\alpha x_i^2}{|x|^3} + \frac{4\alpha^2 x_i^2}{R|x|^2}\right) R^{-1} e^{-2\alpha|x|/R}, \\ \partial_i^3 e^{-2\alpha|x|/R} &= \left(\frac{6\alpha}{R} + \frac{3}{|x|} - \frac{4\alpha x_i^2}{R|x|^2} - \frac{4\alpha^2 x_i^2}{R^2|x|} - \frac{3x_i^2}{|x|^3}\right) \frac{2\alpha x_i}{R|x|^2} e^{-2\alpha|x|/R}.\end{aligned}$$

Note that, for $i = 1, \dots, d$, $x \in B_R \setminus B_{R/2}$,

$$\left|(\partial_i^2 - \frac{1}{2}\nabla_i^2)e^{-2\alpha|x|/R}\right| \leq C \sup_{y: |y-x| \leq 1} \left|\partial_i^3 e^{-2\alpha|y|/R}\right| \leq \frac{C\alpha}{R^3} e^{-2\alpha|x|/R},$$

and so

$$\left|\sum_{i=1}^d \omega(x, x + e_i) (\partial_i^2 - \frac{1}{2}\nabla_i^2) e^{-2\alpha|x|/R}\right| \leq \frac{C\alpha}{R^3} e^{-2\alpha|x|/R}.\tag{68}$$

Further, by taking $K > 0$ sufficiently large (note $R \geq K$), and choosing $\alpha > 0$ to be sufficiently small, we have, for $x \in B_R \setminus B_{R/2}$,

$$\begin{aligned}&\sum_{i=1}^d \omega(x, x + e_i) \partial_i^2 e^{-2\alpha|x|/R} \\ &= \sum_{i=1}^d \omega(x, x + e_i) \left(-\frac{2\alpha}{|x|} + \frac{2\alpha x_i^2}{|x|^3} + \frac{4\alpha^2 x_i^2}{R|x|^2}\right) R^{-1} e^{-2\alpha|x|/R} \\ &\leq \left(-\frac{\alpha}{|x|} + \frac{(1-2\kappa)\alpha|x|^2}{|x|^3} + \frac{2(1-2\kappa)\alpha^2|x|^2}{R|x|^2}\right) R^{-1} e^{-2\alpha|x|/R} \\ &\leq C(-1 + C\alpha) \frac{\alpha}{R^2} e^{-2\alpha|x|/R} \leq -\frac{C\alpha}{R^2} e^{-2\alpha|x|/R}.\end{aligned}$$

This, together with (68), implies,

$$L_\omega e^{-2\alpha|x|/R} \leq -\frac{C\alpha}{R} e^{-2\alpha|x|/R}, \quad \text{for } x \in B_R \setminus B_{R/2}.$$

Display (19) is proved.

To prove (20), note that when $x = 0$, $L_\omega(e^{-A|x|^2}) = e^{-A} - 1 > -1$. When $x \in \mathbb{Z}^2 \setminus \{0\}$, choosing $A > 0$ sufficiently large,

$$\begin{aligned} L_\omega(e^{-A|x|^2}) &= e^{-A|x|^2} \left[\sum_{i=1}^d \omega(x, x + e_i) (e^{2Ax_i - A} + e^{-2Ax_i - A}) - 1 \right] \\ &\geq e^{-A|x|^2} [\kappa e^{2A-A} - 1] > 0. \end{aligned}$$

Display (20) is proved.

It remains to prove (21). Using the inequalities $e^a + e^{-a} \geq 2 + a^2$, $e^a \geq 1 + a$, we get, by taking A sufficiently large, for $x \in B_R \setminus B_{R/2}$,

$$\begin{aligned} L_\omega(e^{-A|x|^2/R^2}) &= e^{-A|x|^2/R^2} \sum_{i=1}^d \omega(x, x + e_i) \left[e^{-A(1+2x_i)/R^2} + e^{-A(1-2x_i)/R^2} - 2 \right] \\ &\geq e^{-A|x|^2/R^2} \sum_{i=1}^d \omega(x, x + e_i) \left[e^{-A/R^2} (2 + 4A^2 x_i^2 / R^4) - 2 \right] \\ &\geq \frac{A}{R^2} e^{-A|x|^2/R^2} \sum_{i=1}^d \omega(x, x + e_i) \left[\frac{4Ax_i^2}{R^2} \left(1 - \frac{A}{R^2}\right) - 1 \right] \\ &\geq \frac{A}{R^2} e^{-A|x|^2/R^2} \left[\frac{2\kappa A|x|^2}{R^2} - 1 \right] > 0. \end{aligned}$$

Our proof is complete. □

References

- [1] S. Andres, S. Neukamm, *Berry-Esseen Theorem and Quantitative homogenization for the Random Conductance Model with degenerate Conductances* Stoch PDE: Anal Comp. 7, 240-296 (2019).
- [2] S. N. Armstrong, J. Lin, *Optimal quantitative estimates in stochastic homogenization for elliptic equations in nondivergence form* Arch. Ration. Mech. Anal. 225 (2017), no. 2, 937-991.
- [3] S. N. Armstrong, C. Smart, *Quantitative Stochastic Homogenization of Elliptic Equations in Nondivergence Form*. Arch. Ration. Mech. Anal. 214 (2014), no. 3, 867-911.
- [4] A. Bensoussan, J.-L. Lions, G. Papanicolaou, *Asymptotic analysis for periodic structures*. AMS Chelsea Publishing, Providence, RI, 2011. xii+398 pp. ISBN: 978-0-8218-5324-5.

- [5] N. Berger, M. Cohen, J.-D. Deuschel, X. Guo, *An elliptic Harnack inequality for random walk in balanced environments*. Ann. Probab. 50 (2022), no. 3, 835-873.
- [6] N. Berger, J.-D. Deuschel, *A quenched invariance principle for non-elliptic random walk in i.i.d. balanced random environment*. Probab. Theory Related Fields 158 (2014), no. 1-2, 91-126.
- [7] M. Biskup, *Recent progress on the random conductance model*. Probab. Surv. 8 (2011), 294-373.
- [8] S. Boucheron, O. Bousquet, G. Lugosi, P. Massart, *Moment inequalities for functions of independent random variables*. Ann. Probab. 33 (2005), no. 2, 514-560.
- [9] S. Boucheron, G. Lugosi, P. Massart, *Concentration Inequalities: A Nonasymptotic Theory of Independence*. Oxford University Press, 2013.
- [10] L. A. Caffarelli, P. E. Souganidis, *Rates of convergence for the homogenization of fully nonlinear uniformly elliptic pde in random media*. Invent. Math. 180(2), 301-360 (2010).
- [11] P. de Buyer, J.-C. Mourrat, *Diffusive decay of the environment viewed by the particle*, Electron. Commun. Probab. 20 (2015), no. 23, 1-12.
- [12] Y. Derriennic, M. Lin, *The central limit theorem for Markov chains started at a point*. Probab. Theory Relat. Fields 125 (2003), 73-76.
- [13] J.-D. Deuschel, X. Guo, *Quenched local central limit theorem for random walks in a time-dependent balanced random*. Probab. Theory Relat. Fields 182 (2022), 111-156.
- [14] J.-D. Deuschel, X. Guo, A. Ramirez, *Quenched invariance principle for random walk in time-dependent balanced random environment*. Ann. Inst. Henri Poincaré Probab. Stat. Vol. 54, No. 1(2018), 363-384.
- [15] L. Escauriaza *Bounds for the fundamental solutions of elliptic and parabolic equations: In memory of Eugene Fabes*. Communications in Partial Differential Equations 25.5-6 (2000): 821-845.
- [16] E. Fabes, D. Stroock, *The L^p -integrability of Green's functions and fundamental solutions for elliptic and parabolic equations*. Duke Math. J. 51 (1984), no. 4, 997-1016.
- [17] D. Gilbarg, N. S. Trudinger, *Elliptic partial differential equations of second order*. Reprint of the 1998 edition. Classics in Mathematics. Springer-Verlag, Berlin, 2001. xiv+517 pp.

- [18] A. Gloria, S. Neukamm, F. Otto, *Quantification of ergodicity in stochastic homogenization: optimal bounds via spectral gap on Glauber dynamics*. Invent. Math. 199 (2015), no. 2, 455-515.
- [19] X. Guo, J. Peterson, H. V. Tran, *Quantitative homogenization in a balanced random environment*. Electron. J. Probab., 2022, to appear.
- [20] X. Guo, T. Sprekeler, H. V. Tran *Characterizations of diffusion matrices in homogenization of elliptic equations in nondivergence-form*. arXiv:2201.01974 [math.AP].
- [21] X. Guo, H. V. Tran, Y. Yu, *Remarks on optimal rates of convergence in periodic homogenization of linear elliptic equations in non-divergence form*. SN Partial Differ. Equ. Appl. 1, 15 (2020).
- [22] X. Guo, O. Zeitouni, *Quenched invariance principle for random walks in balanced random environment*. Probab. Theory Related Fields 152 (2012), 207-230.
- [23] V. V. Jikov, S. M. Kozlov, O. A. Oleinik, *Homogenization of Differential Operators and Integral Functionals*. Translated from the Russian by G.A. Yosifian, Springer-Verlag Berlin Heidelberg 1994, xii+570 pp. ISBN: 3-540-54809-2.
- [24] C. Kipnis, S. R. S. Varadhan, *Central Limit Theorem for Additive Functionals of Reversible Markov Processes and Applications to Simple Exclusions*, Commun.Math.Phys. 104,1-19(1986).
- [25] H. J. Kuo, N. Trudinger, *Positive difference operators on general meshes*. Duke Math. J. 83 (1996), no. 2, 415-433.
- [26] G. Lawler, *Weak convergence of a random walk in a random environment*. Comm. Math. Phys. 87:81–87, (1982).
- [27] J. Lin, C. Smart, *Algebraic error estimates for the stochastic homogenization of uniformly parabolic equations*. Anal. PDE 8 (2015), no. 6, 1497-1539.
- [28] J.-C. Mourrat, *Variance decay for functionals of the environment viewed by the particle*. Ann. Inst. H. Poincaré Probab. Stat. 47(11), 294-327 (2011)
- [29] G. Papanicolaou and S.R.S. Varadhan. *Diffusions with random coefficients*. Statistics and probability: essays in honor of C. R. Rao, pp. 547-552, North-Holland, Amsterdam, (1982).
- [30] V. V. Yurinski, *On the error of averaging of multidimensional diffusions*. Teor. Veroyatnost. i Primenen 33(1), 14-24 (1988) [Eng. transl. in Theory Probab. Appl. 33(1), 11-21 (1988)].

- [31] O. Zeitouni, *Random walks in random environment*. In Lectures on probability theory and statistics, volume 1837 of Lecture Notes in Math., pages 189-312. Springer, Berlin, 2004.

E-mail address, Xiaoqin Guo: `guoxq@ucmail.uc.edu`

E-mail address, Hung Vinh Tran: `hung@math.wisc.edu`