

FREE PRODUCT ON SEMIHYPERGROUPS

CHOITI BANDYOPADHYAY

ABSTRACT. In a previous paper [1], we initiated a systematic study of semihypergroups and had a thorough discussion about some important analytic and algebraic objects associated to this class of objects. In this paper, we investigate free structures on the category of semihypergroups. We show that the natural free product structure along with the natural topology, although fails to give a free-product for topological groups, works well on a vast non-trivial class of ‘pure’ semihypergroups containing most of the well-known examples including non-trivial coset and orbit spaces.

1. INTRODUCTION

The concept of semihypergroups arises naturally in abstract harmonic analysis in terms of left/right coset spaces and orbit spaces of certain actions which appear frequently in different areas of research including Lie groups, coset theory, homogeneous spaces, dynamical systems, ordinary and partial differential equations and orthogonal polynomials, to name a few. These objects arising from locally compact groups, although retain some interesting structures from the parent category, fail to be a topological group, semigroup or even a hypergroup [1, 9]. Hence these kinds of objects that frequently appear while studying the classical theory of topological groups, fall out of the parent category and can not quite be studied or analysed with the existing general theory for hypergroups and topological groups.

The concept of a semihypergroup was first introduced around 1972 by C. Dunkl [4], I. Jewett [9] and R. Spector [19] independently, in pursuit of developing an unified theory involving convolution of measures that includes a wide array of objects arising from locally compact groups. In a sense, they serve as building blocks for hypergroup-axioms.

A semihypergroup, as one would expect, can be perceived as simply a generalization of a locally compact semigroup, where the product of two points is a certain probability measure, rather than a single point. But unlike hypergroups, it does not need to have an identity or involution (analogous to inverses in groups). Hence in a nutshell, a semihypergroup is essentially a locally compact topological space where the measure space is equipped with a certain convolution product, turning it into an associative algebra, which need not have an identity or an involution. It can be seen (details in the following section) that all the orbit

2010 *Mathematics Subject Classification*. Primary: 43A62, 43A60, 43A10, 43A99, 20M12; Secondary: 46G12, 46J20, 46E27.

Key words and phrases. semihypergroups, free products, universal properties, hypergroups, coset spaces, orbit spaces.

Part of this work is included in author’s PhD thesis at the University of Alberta, Canada.

spaces and coset spaces can naturally be given a semihypergroup structure, though not a hypergroup structure.

The fact that semihypergroups contain more generalized objects arising from different fields of research than classical group and hypergroup theory and yet sustains enough structure to allow an independent theory to develop, makes it an intriguing and useful area of study with essentially a broader area of applications. Unlike hypergroups, there is a severe lack of any extensive prior research on semihypergroups since its inception. On the other hand, almost all of the important concepts and results on hypergroups and locally compact groups fail to extend naturally to semihypergroups due to the lack of an involution, an identity, a standardized Haar measure and even an algebraic structure on the underlying space in semihypergroups.

Hence in a previous paper [1] we initiated developing a systematic extensive theory on semihypergroups. Our approach towards developing a systematic theory was to define and explore some basic algebraic and analytic aspects and areas on semihypergroups, which are imperative for the development of any category of analytic objects in general. In the first installment of the study [1] we introduced and developed the concepts and theories of homomorphisms, ideals, kernels and almost periodic and weakly almost periodic functions on a semihypergroup. In this paper, we advance the theory by discussing free structures and universal properties on a class of semihypergroups.

We first discuss how a semihypergroup homomorphism (introduced in [1]) naturally induces a homomorphism (in the classical sense) between measure algebras. Using this property of homomorphisms, we discuss how free products can be defined given a class of semihypergroups, abiding by the general norms of defining a free product given any category of objects in general. We then introduce a sub-class of ‘pure’ non-trivial semihypergroups and investigate in detail the existence of a unique (upto isomorphism) free product given such a family of objects.

The most interesting observation that comes out of this particular path of research is that the natural free product structure with the naturally induced topology, which fails to give a free product structure for topological groups [14], indeed works well here and gives us a natural free product structure for a class of non-trivial semihypergroups. As expected, the existence of a natural free product ensures the abundance of new examples and opens up several new paths of research including compactification of a given semihypergroup and amalgamation of different kinds of semihypergroups in pursuit of specific examples and counter-examples among others. The rest of the paper is designed as below.

In the next, *i.e.*, second section of this article, we recall some preliminary definitions and notations given by Jewett in [9], and introduce some new definitions required for further work. We conclude the section with listing some important useful examples of semihypergroups and hypergroups. In the third section, we initiate the study of a free structure on semihypergroups. We introduce a free product structure and construct a specific topology and convolution for a family of semihypergroups such that the resulting semihypergroup abides by an universal property equivalent to the universal property for free products in the

classical theory of topological groups, thus providing us with a whole new class of examples for semihypergroups.

Finally, this text is the second installment of a systematic study of this unifying category of objects with an intriguing scope for applications in different fields of research discussed above. Hence we conclude the text with some related potential problems and areas which we intend to work on and explore further in near future. Furthermore, introducing and investigating similar structures as in [2, 3, 5, 6, 7, 8, 10, 11, 13, 16, 17, 18] for the case of semihypergroups and thereby realising further where and why this theory deviates from the classical theory of topological semigroups and groups, potentially provides us with deeper insight into different questions in the study of abstract harmonic analysis.

2. PRELIMINARY

Here we first list some of the preliminary set of basic notations that we will use throughout the text, followed by a brief introduction to the tools and concepts needed for the formal definition of a semihypergroup (referred to as ‘semiconvo’ in [9]). Finally, we list some of the well-known natural examples of a semihypergroup [1, 9, 20]. All the topologies throughout this text are assumed to be Hausdorff, unless otherwise specified.

For any locally compact Hausdorff topological space X , we denote by $M(X)$ the space of all regular complex Borel measures on X , where $M^+(X)$ and $P_c(X)$ denote the subsets of $M(X)$ consisting of all finite non-negative regular Borel measures on X and all probability measures with compact support on X respectively. Moreover, $B(X)$, $C(X)$, $C_c(X)$ and $C_0(X)$ denote the function spaces of all bounded functions, bounded continuous functions, compactly supported continuous functions and continuous functions vanishing at infinity on X respectively.

Next, we introduce two very important topologies on the positive measure space and the space of compact subsets for any locally compact topological space X . Unless mentioned otherwise, we will always assume these two topologies on the respective spaces.

The **cone topology** on $M^+(X)$ is defined as the weakest topology on $M^+(X)$ for which the maps $\mu \mapsto \int_X f d\mu$ is continuous for any $f \in C_c^+(X) \cup \{1_X\}$ where 1_X denotes the characteristic function of X . Note that if X is compact then it follows immediately from the Riesz representation theorem that the cone topology coincides with the weak*-topology on $M^+(X)$ in this case.

We denote by $\mathfrak{C}(X)$ the set of all compact subsets of X . The **Michael topology** on $\mathfrak{C}(X)$ is defined to be the topology generated by the sub-basis $\{\mathcal{C}_U(V) : U, V \text{ are open sets in } X\}$, where

$$\mathcal{C}_U(V) = \{C \in \mathfrak{C}(X) : C \cap U \neq \emptyset, C \subset V\}.$$

Note that $\mathfrak{C}(X)$ actually becomes a locally compact Hausdorff space with respect to this natural topology. Moreover if X is compact then $\mathfrak{C}(X)$ is also compact.

For any locally compact Hausdorff space X and any element $x \in X$, we denote by p_x the point-mass measure or the Dirac measure at the point $\{x\}$.

For any three locally compact Hausdorff spaces X, Y, Z , a bilinear map $\Psi : M(X) \times M(Y) \rightarrow M(Z)$ is called **positive continuous** if the following holds :

- (1) $\Psi(\mu, \nu) \in M^+(Z)$ whenever $\mu \in M^+(X), \nu \in M^+(Y)$.
- (2) The map $\Psi|_{M^+(X) \times M^+(Y)}$ is continuous.

Now we are ready to state the formal definition for a semihypergroup. Note that we follow Jewett's notion [9] in terms of the definitions and notations, in most cases.

Definition 2.1. (*Semihypergroup*) A pair $(K, *)$ is called a (topological) semihypergroup if they satisfy the following properties:

- (A1): K is a locally compact Hausdorff space and $*$ defines a binary operation on $M(K)$ such that $(M(K), *)$ becomes an associative algebra.
- (A2): The bilinear mapping $*$: $M(K) \times M(K) \rightarrow M(K)$ is positive continuous.
- (A3): For any $x, y \in K$ the measure $p_x * p_y$ is a probability measure with compact support.
- (A4): The map $(x, y) \mapsto \text{supp}(p_x * p_y)$ from $K \times K$ into $\mathfrak{C}(K)$ is continuous.

Note that for any $A, B \subset K$ the convolution of subsets is defined as the following:

$$A * B := \cup_{x \in A, y \in B} \text{supp}(p_x * p_y).$$

We define the concept of left (or right) topological and semitopological semihypergroups, similar to the concepts of left (or right) topological and semitopological semigroups.

Definition 2.2. A pair $(K, *)$ is called a left topological semihypergroup if it satisfies all the conditions of Definition 2.1 with property (A2) replaced by the following:

- (A2'): The map $(\mu, \nu) \mapsto \mu * \nu$ is positive and for each $\omega \in M^+(K)$ the map $L_\omega : M^+(K) \rightarrow M^+(K)$ given by $L_\omega(\mu) = \omega * \mu$ is continuous.

Definition 2.3. A pair $(K, *)$ is called a right topological semihypergroup if it satisfies all the conditions of Definition 2.1 with property (A2) replaced by the following:

- (A2''): The map $(\mu, \nu) \mapsto \mu * \nu$ is positive and for each $\omega \in M^+(K)$ the map $R_\omega : M^+(K) \rightarrow M^+(K)$ given by $R_\omega(\mu) = \mu * \omega$ is continuous.

Definition 2.4. A pair $(K, *)$ is called a **semitopological semihypergroup** if it is both left and right topological semihypergroup.

Now we list some well known examples [9, 20] of semihypergroups and hypergroups. Please see [1, Section 3] for details on the constructions as well as why most of the structures discussed there, although attain semihypergroup structures, fail to be hypergroups.

Example 2.5. If (S, \cdot) is a locally compact topological semigroup, then $(S, *)$ is a semihypergroup where $p_x * p_y = p_{x \cdot y}$ for any $x, y \in S$. Similarly, if (G, \cdot) is a locally compact topological group, then $(G, *)$ is a hypergroup with the same bilinear operation $*$, identity element e where e is the identity of G and the involution on G defined as $x \mapsto x^{-1}$.

Example 2.6. Take $T = \{e, a, b\}$ and equip it with the discrete topology. Define

$$\begin{aligned} p_e * p_a &= p_a * p_e = p_a \\ p_e * p_b &= p_b * p_e = p_b \\ p_a * p_b &= p_b * p_a = z_1 p_a + z_2 p_b \\ p_a * p_a &= x_1 p_e + x_2 p_a + x_3 p_b \\ p_b * p_b &= y_1 p_e + y_2 p_a + y_3 p_b \end{aligned}$$

where $x_i, y_i, z_i \in \mathbb{R}$ such that $x_1 + x_2 + x_3 = y_1 + y_2 + y_3 = z_1 + z_2 = 1$ and $y_1 x_3 = z_1 x_1$. Then $(T, *)$ is a commutative hypergroup with identity e and the identity function on T taken as involution.

Example 2.7. Let G be a locally compact topological group and H be a compact subgroup of G . Also, μ be the normalized Haar measure of H . Consider the left coset space $S := G/H = \{xH : x \in G\}$ and the double coset space $K := G//H = \{HxH : x \in G\}$ and equip them with the respective quotient topologies. Then $(S, *)$ is a semihypergroup and $(K, *)$ is a hypergroup where the convolutions are given as following for any $x, y \in G$:

$$p_{xH} * p_{yH} = \int_H p_{(xty)H} d\mu(t), \quad p_{HxH} * p_{HyH} = \int_H p_{H(xty)H} d\mu(t).$$

Example 2.8. Let G be a locally compact topological group and H be any compact group. For any continuous affine action [9] π of H on G , consider the orbit space $\mathcal{O} := \{x^H : x \in G\}$, where $x^H = \mathcal{O}(x) = \{\pi(h, x) : h \in H\}$.

Let σ be the normalized Haar measure of H . Consider \mathcal{O} with the quotient topology and the following convolution

$$p_{x^H} * p_{y^H} := \int_H \int_H p_{(\pi(s,x)\pi(t,y))^H} d\sigma(s)d\sigma(t).$$

Then $(\mathcal{O}, *)$ becomes a semihypergroup.

3. FREE STRUCTURE ON SEMIHYPERGROUPS

Free groups and free products of topological groups and semigroups have been an useful tool for a number of reasons. Providing specific examples or counter-examples, studying subgroups and quotients of a topological group, constructing an unified group with specific properties of a number of groups and studying the problem of (topologically) embedding a topological semigroup into a topological group are a few of the areas where the use of a free structure proves to be very helpful.

In this section, we initiate the study of a free structure on semihypergroups. We introduce a free product structure and a specific topology and convolution to a family of semihypergroups such that the resulting semihypergroup abides by an universal property equivalent to the universal property for the free product of a family of topological groups [14]. But before we proceed to the definition and construction of a free product, let us first briefly recall some basic formal structures on an arbitrary set and examine how a homomorphism between two

semihypergroups translates to a specific homomorphism between their measure algebras, in the classical sense.

Recall [1] that given two (semitopological) semihypergroups K and H , a continuous map $\phi : K \rightarrow H$ is called a homomorphism if for any Borel measurable function f on H and for any $x, y \in K$ we have that

$$f \circ \phi(x * y) = f(\phi(x) * \phi(y)).$$

Let A be any nonempty set. Then a **word** or string on A is a finite sequence $a_1 a_2 \dots a_n$ which is either void or $a_i \in A$ for each $1 \leq i \leq n$. Similarly, given a family $\{A_\alpha\}$ of nonempty sets, a word or string on $\{A_\alpha\}$ is a finite sequence $a_1 a_2 \dots a_n$ which is either void or $a_i \in A_\alpha$ for some α , for each $1 \leq i \leq n$. For convenience of notation, here we sometimes write the word $a_1 a_2 \dots a_n$ also as $(a_1 a_2 \dots a_n)$.

Definition 3.1. Let $\{K_\alpha\}$ be a family of semihypergroups and $w = a_1 a_2 \dots a_n$ is a word on $\{K_\alpha\}$ where $a_i \in K_{\alpha_i}$ for $1 \leq i \leq n$. Then w is called a **reduced word** if either w is empty or if w satisfies the following two conditions:

- (1) $a_i \neq 1_{K_\alpha}$ for $1 \leq i \leq n$ where 1_{K_α} is the identity element of K_α (if identity exists in K_α).
- (2) $\alpha_i \neq \alpha_{i+1}$ for $1 \leq i \leq n - 1$.

Given a reduced word $a_1 a_2 \dots a_n$ on a family of semihypergroups $\{K_\alpha\}$, the **length** of the word is defined to be the number of elements in the word, and is written as $l(a_1 a_2 \dots a_n) = n$.

Proposition 3.2. Given a finite set A of cardinality n , we can form the free semihypergroup $(F_A, *)$ generated by A , where F_A is the set of all words (finite sequences) consisting of 0 or more elements from A .

Proof. Given any finite set of cardinality n , we can easily form a free semigroup (F, \cdot) generated by A where F is the set of all words (finite sequences) consisting of 0 or more elements from A and the multiplication of two words $a_1 a_2 \dots a_n, b_1 b_2 \dots b_m$ are simple concatenation given by

$$(a_1 a_2 \dots a_n) \cdot (b_1 b_2 \dots b_m) = a_1 a_2 \dots a_n b_1 b_2 \dots b_m.$$

Note that equipping F with the discrete topology makes it into a topological semigroup. Then as outlined in Example 2.5 we can consider (F, \cdot) as a semihypergroup $(F_A, *)$ where as sets, we have $F = F_A$. □ □

Theorem 3.3. Let K and H be two semihypergroups. Then a homomorphism $\phi : K \rightarrow H$ induces a positive continuous homomorphism $\Gamma_\phi : M(K) \rightarrow M(H)$.

Proof. Define $T_\phi : C_0(H) \rightarrow C_0(K)$ by

$$T_\phi f(x) := f(\phi(x))$$

for each $f \in C_0(H), x \in K$.

Then we know that the adjoint operator $T_\phi^* : C_0(K)^* \rightarrow C_0(H)^*$ is given by

$$T_\phi^*(m)(f) := m(T_\phi f)$$

for each $m \in C_0(K)^*$, $f \in C_0(H)$.

We define $\Gamma_\phi := T_\phi^* : M(K) \rightarrow M(H)$. Hence on point-mass measures p_x for any $x \in K$, we have

$$\Gamma_\phi(p_x)(f) = f(\phi(x)) = p_{\phi(x)}(f)$$

for any $f \in C_0(H)$.

Hence we have Γ_ϕ in the following way. Define

$$\Gamma_\phi(p_x) := p_{\phi(x)}$$

and then extend Γ_ϕ linearly to $M_F^+(K)$. Since $M_F^+(K)$ is dense in $M^+(K)$ we can extend Γ_ϕ to $M^+(K)$ and then finally we extend Γ_ϕ linearly to $M(K)$.

Hence by construction, we get that Γ_ϕ is a positive continuous linear map on $M(K)$. Now to see if it is a homomorphism, it suffices to show that

$$\Gamma_\phi(p_x * p_y) = \Gamma_\phi(p_x) * \Gamma_\phi(p_y)$$

for any $x, y \in K$.

Now pick any $x, y \in K$. Then $p_x * p_y \in M^+(K)$. Hence there exists a net $\{m_\alpha\}$ in $M_F^+(K)$ such that

$$m_\alpha \rightarrow (p_x * p_y).$$

For each α let

$$m_\alpha = \sum_{i=1}^{k_\alpha} c_i^\alpha p_{x_i^\alpha}$$

for some $k_\alpha \in \mathbb{N}$, $c_i^\alpha \in \mathbb{F}$, $x_i^\alpha \in K$. Thus we have

$$\begin{aligned} \Gamma_\phi(p_x * p_y) &= \Gamma_\phi(\lim_\alpha m_\alpha) \\ &= \lim_\alpha \Gamma_\phi\left(\sum_{i=1}^{k_\alpha} c_i^\alpha p_{x_i^\alpha}\right) \\ &= \lim_\alpha \sum_{i=1}^{k_\alpha} c_i^\alpha \Gamma_\phi(p_{x_i^\alpha}) \\ &= \lim_\alpha \sum_{i=1}^{k_\alpha} c_i^\alpha p_{\phi(x_i^\alpha)} = \lim_\alpha \tilde{m}_\alpha \end{aligned}$$

where for each α we set

$$\tilde{m}_\alpha := \sum_{i=1}^{k_\alpha} c_i^\alpha p_{\phi(x_i^\alpha)}.$$

Now pick any $f \in C_0(H)$. We have

$$\begin{aligned}
(p_{\phi(x)} * p_{\phi(y)})(f) &= \int_H f d(p_{\phi(x)} * p_{\phi(y)}) \\
&= f(\phi(x) * \phi(y)) \\
&= (f \circ \phi)(x * y) \\
&= \int_K (f \circ \phi) d(p_x * p_y) \\
&= \int_K (f \circ \phi) d(\lim_{\alpha} m_{\alpha}) = \lim_{\alpha} \int_K (f \circ \phi) d(m_{\alpha})
\end{aligned}$$

where the third equality holds since ϕ is a homomorphism. Thus we have that

$$\begin{aligned}
(p_{\phi(x)} * p_{\phi(y)})(f) &= \lim_{\alpha} \sum_{i=1}^{k_{\alpha}} c_i^{\alpha} \int_K (f \circ \phi) d(p_{x_i^{\alpha}}) \\
&= \lim_{\alpha} \sum_{i=1}^{k_{\alpha}} c_i^{\alpha} f(\phi(x_i^{\alpha})) \\
&= \lim_{\alpha} \sum_{i=1}^{k_{\alpha}} c_i^{\alpha} p_{\phi(x_i^{\alpha})}(f) = \lim_{\alpha} \tilde{m}_{\alpha}(f).
\end{aligned}$$

Since $f \in C_0(H)$ was chosen arbitrarily, it follows that

$$\lim_{\alpha} \tilde{m}_{\alpha} = p_{\phi(x)} * p_{\phi(y)}.$$

Hence for any $x, y \in K$ we have that

$$\begin{aligned}
\Gamma_{\phi}(p_x * p_y) &= p_{\phi(x)} * p_{\phi(y)} \\
&= \Gamma_{\phi}(p_x) * \Gamma_{\phi}(p_y).
\end{aligned}$$

□

□

Now we define the free product of a family of (semitopological) semihypergroups, following the standard norms of defining a free product on a category of objects. Note that the definition follows the definition of free product for topological groups [14, Definition 2.1]. The first two structural conditions essentially remain the same as in the case of topological groups, whereas the universal property is naturally defined in terms of measure spaces, generalising the product to semihypergroups.

Definition 3.4. *Let $\{K_{\alpha}\}$ be a family of semihypergroups. Then a semihypergroup F is called the free (topological) product of $\{K_{\alpha}\}$, if the following conditions are satisfied:*

- (1) *For each α , K_{α} is a sub-semihypergroup of F .*
- (2) *F is algebraically generated by $\cup_{\alpha} K_{\alpha}$ (in terms of finite reduced words).*
- (3) *Given any semihypergroup H , if we have a continuous homomorphism $\phi_{\alpha} : K_{\alpha} \rightarrow H$ for each α , then there exists a unique positive continuous homomorphism $\Gamma : M(F) \rightarrow M(H)$ such that $\Gamma|_{M(K_{\alpha})} = \Gamma_{\phi_{\alpha}}$ for each α .*

We denote F as $\prod_{\alpha}^* K_{\alpha}$.

Now before we prove that such products exists for all the significant examples discussed in the third section, let us first define a specific type of semihypergroups.

Definition 3.5. Let $(K, *)$ be a (semitopological) semihypergroup. We say that K is a **pure semihypergroup** if for any $x, y \in K \setminus \{e\}$ we have that $\text{supp}(p_x * p_y)$ is not singleton whenever $e \in \text{supp}(p_x * p_y)$. Here e is the identity of K .

Equivalently, we can say that K is a **pure semihypergroup** if we have that $p_x * p_y \neq p_e$ for any $x, y \in K \setminus \{e\}$.

Note that if a semihypergroup does not have an identity, then the above conditions are vacuously true and so K is a pure semihypergroup. Also, note that all the non-trivial examples discussed in Examples 2.6, 2.7 and 2.8 are pure semihypergroups.

However, locally compact groups clearly fail to be a pure semihypergroup. In fact, a locally compact semigroup (S, \cdot) with identity e can be seen as a pure semihypergroup (Example 2.5) only if $x \cdot y \neq e$ for any $x, y \in S \setminus \{e\}$, i.e, if none of the elements in S have a left or right inverse.

Theorem 3.6. Let $\{K_{\alpha}\}$ be a family of semihypergroups such that if more than one of the K_{α} 's has an identity element, then all the K_{α} 's are pure semihypergroups. Then there exists a semihypergroup F such that $F = \prod_{\alpha}^* K_{\alpha}$.

Moreover, if more than one of the K_{α} 's has an identity element, then F is also a semihypergroup with identity.

Proof. Set $\Lambda := \{\alpha : K_{\alpha} \text{ has an identity element}\}$. If $|\Lambda| \leq 1$, set

$$K := \{a_1 a_2 \dots a_n : n \in \mathbb{N}, a_1 a_2 \dots a_n \text{ is a reduced word on } \{K_{\alpha}\}\}.$$

Note that if $|\Lambda| = 1$, then we treat the identity as any other arbitrary element in that semihypergroup. For convenience of notations, we rename the identity element as x_e in this case, so that it can be included in finite sequences of reduced words.

If $|\Lambda| > 1$, let e denote the empty word on $\{K_{\alpha}\}$ and set $e = 1_{K_{\alpha}}$ for each $\alpha \in \Lambda$ where $1_{K_{\alpha}}$ denotes the identity element of K_{α} . Now set

$$K := \{e\} \cup \{a_1 a_2 \dots a_n : n \in \mathbb{N}, a_1 a_2 \dots a_n \text{ is a reduced word on } \{K_{\alpha}\}\}.$$

For any $x_1 x_2 \dots x_n, y_1 y_2 \dots y_m \in K$ where $x_i \in K_{\alpha_i}, y_j \in K_{\beta_j}$ for $i = 1, 2, \dots, n; j = 1, 2, \dots, m$ and any subset $A \subset K_{\alpha_0}$ where $\alpha_0 \neq \alpha_n, \beta_1$ we define that:

$$(x_1 \dots x_n)A(y_1 \dots y_m) := \begin{cases} \{x_1 x_2 \dots x_n a y_1 y_2 \dots y_m : a \in A\}, & \text{if } e \notin A. \\ \{(x_1 x_2 \dots x_n y_1 y_2 \dots y_m)\} \cap K \\ \cup \{(x_1 x_2 \dots x_n a y_1 y_2 \dots y_m) : a \in A, a \neq e\}, & \text{if } e \in A. \end{cases}$$

Now for any two sets $A_1 \subset K_{\alpha_1}$, $A_2 \subset K_{\alpha_2}$ where $\alpha_1 \neq \alpha_2$ define

$$\begin{aligned} A_1 A_2 &:= \{a_1 a_2 : a_1 \in A_1, a_2 \in A_2\}, \text{ if } e \notin A_1, A_2. \\ A_1 A_2 &:= \{a_1 a_2 : a_1 \in A_1 \setminus \{e\}, a_2 \in A_2\} \cup \{a_2 : a_2 \in A_2\}, \text{ if } e \in A_1, e \notin A_2. \\ A_1 A_2 &:= \{a_1 a_2 : a_1 \in A_1, a_2 \in A_2 \setminus \{e\}\} \cup \{a_1 : a_1 \in A_1\}, \text{ if } e \notin A_1, e \in A_2. \\ A_1 A_2 &:= \{a_1 a_2 : a_1 \in A_1 \setminus \{e\}, a_2 \in A_2 \setminus \{e\}\} \cup \{a_1 : a_1 \in A_1 \setminus \{e\}\} \\ &\quad \cup \{a_2 : a_2 \in A_2 \setminus \{e\}\} \cup \{e\}, \text{ if } e \in A_1, A_2. \end{aligned}$$

Similarly for $A_1 \subset K_{\alpha_1}$, $A_2 \subset K_{\alpha_2}$, $A_3 \subset K_{\alpha_3}$ where $\alpha_1 \neq \alpha_2 \neq \alpha_3$, we define

$$\begin{aligned} A_1 A_2 A_3 &:= \{a_1 a_2 a_3 : a_1 \in A_1, a_2 \in A_2, a_3 \in A_3\}, \text{ if } e \notin A_1, A_2, A_3. \\ A_1 A_2 A_3 &:= \{a_1 a_2 a_3 : a_1 \in A_1 \setminus \{e\}, a_2 \in A_2, a_3 \in A_3\} \\ &\quad \cup \{a_2 a_3 : a_2 \in A_2, a_3 \in A_3\}, \text{ if } e \in A_1, e \notin A_2, A_3. \\ A_1 A_2 A_3 &:= \{a_1 a_2 a_3 : a_1 \in A_1 \setminus \{e\}, a_2 \in A_2 \setminus \{e\}, a_3 \in A_3\} \\ &\quad \cup \{a_2 a_3 : a_2 \in A_2 \setminus \{e\}, a_3 \in A_3\} \\ &\quad \cup \{a_1 a_3 : a_1 \in A_1 \setminus \{e\}, a_3 \in A_3\} \cup \{a_3 : a_3 \in A_3\}, \\ &\quad \text{if } e \in A_1, A_2, e \notin A_3. \\ A_1 A_2 A_3 &:= \{a_1 a_2 a_3 : a_1 \in A_1 \setminus \{e\}, a_2 \in A_2 \setminus \{e\}, a_3 \in A_3 \setminus \{e\}\} \\ &\quad \cup \{a_1 a_2 : a_1 \in A_1 \setminus \{e\}, a_2 \in A_2 \setminus \{e\}\} \cup \{a_2, a_3 : a_2 \in A_2 \setminus \{e\}, \\ &\quad a_3 \in A_3 \setminus \{e\}\} \cup \{a_1, a_3 : a_1 \in A_1 \setminus \{e\}, a_3 \in A_3 \setminus \{e\}\} \\ &\quad \cup \{a_1 : a_1 \in A_1 \setminus \{e\}\} \cup \{a_2 : a_2 \in A_2 \setminus \{e\}\} \\ &\quad \cup \{a_3 : a_3 \in A_3 \setminus \{e\}\} \cup \{e\}, \text{ if } e \in A_1, A_2, A_3. \end{aligned}$$

We can get the other cases following the same pattern.

In general for any $n \in \mathbb{N}$, consider subsets $A_i \subset K_{\alpha_i}$ for $i = 1, 2, \dots, n$, where $\alpha_i \neq \alpha_{i+1}$ for $i = 1, 2, \dots, (n-1)$. Let $e \in A_{i_k}$ for $k = 1, 2, \dots, m$ where $i_{k_1} < i_{k_2}$ for $k_1 < k_2$ and $e \notin A_i$ whenever $i \neq i_k$ for any $k = 1, 2, \dots, m$.

Then continuing in the same manner as above, we get that

$$A_1 A_2 \dots A_n := \begin{cases} \left\{ \begin{aligned} &\{a_1 a_2 \dots a_n \in K : a_i \in A_i, i = 1, 2, \dots, n\} \\ &\cup (F_n \cap K) \cup \{e\}, \end{aligned} \right. & \text{if } m = n \\ \left\{ \begin{aligned} &\{a_1 a_2 \dots a_n \in K : a_i \in A_i, i = 1, 2, \dots, n\} \\ &\cup (F_n \cap K), \end{aligned} \right. & \text{if } m \neq n \end{cases}$$

where F_n is defined as the following:

$$F_n := \bigcup_{l=1}^m \bigcup_{\substack{1 \leq k_1 < k_2 < \dots \\ \dots < k_l \leq m}} \left\{ (a_1 a_2 \dots a_{i_{k_1}-1} a_{i_{k_1}+1} \dots a_{i_{k_2}-1} a_{i_{k_2}+1} \dots a_{i_{k_l}-1} a_{i_{k_l}+1} \dots \dots a_{n-1} a_n) : a_j \in A_j, j = 1, 2, \dots, n \right\}.$$

Note that from now on unless otherwise mentioned, whenever we write a product $A_1 A_2 \dots A_n$ of subsets $A_i \subset K_{\alpha_i}$, we assume that $\alpha_i \neq \alpha_{i+1}$ for $i = 1, 2, \dots, (n-1)$. Also it follows immediately from the above construction that for any such sets A_1, A_2, \dots, A_n we have

$$A_1 A_2 \dots A_n = A_1 (A_2 A_3 \dots A_n) = A_1 (A_2 A_3 \dots A_{n-1}) A_n = (A_1 A_2 \dots A_{n-1}) A_n.$$

Natural Topology on K :

Now we first introduce a certain topology τ on K . Afterwards we will define a binary operation ‘ $*$ ’ on the measure space $M(K)$ and then investigate how the operation gives rise to an associative algebra on $M(K)$ and how the Borel sets act under the said operation and the cone topology on $M^+(K)$ induced by τ . First, we set

$$\mathcal{B} := \{U_1 U_2, \dots, U_n : U_i \text{ is an open subset of } K_{\alpha_i}, i = 1, 2, \dots, n; n \in \mathbb{N}\}.$$

We need to show that \mathcal{B} serves as a base for a topology on K . Pick any element $x_1 x_2 \dots x_n \in K \setminus \{e\}$ where $x_i \in K_{\alpha_i}$, $i = 1, 2, \dots, n$. Since $x_1 x_2 \dots x_n$ is a reduced word, $x_i \neq 1_{K_{\alpha_i}}$, even if $\alpha_i \in \Lambda$. Thus we can find an open neighbourhood V_{x_i} of x_i in K_{α_i} such that $e \notin V_{x_i}$. Then $x_1 x_2 \dots x_n \in V_{x_1} V_{x_2} \dots V_{x_n} \in \mathcal{B}$.

Also note that if $e \in K$, then for any open neighborhood U of $e = 1_{K_\alpha}$ in K_α where $\alpha \in \Lambda$, we have that $e \in U \in \mathcal{B}$. Now let $U_1 U_2 \dots U_n, V_1 V_2 \dots V_m \in \mathcal{B}$ where $U_i \subset K_{\alpha_i}$ and $V_j \subset K_{\beta_j}$ for $i = 1, 2, \dots, n; j = 1, 2, \dots, m$. Pick any element $x_1 x_2 \dots x_l \in U_1 U_2 \dots U_n \cap V_1 V_2 \dots V_m$. Note that $l \leq \min(n, m)$ and hence for each $k = 1, 2, \dots, l$, there exists some $i_k \in \{1, 2, \dots, n\}$ and $j_k \in \{1, 2, \dots, m\}$ such that $i_1 < i_2 < \dots < i_l$, $j_1 < j_2 < \dots < j_l$ and $x_k \in U_{i_k} \cap V_{j_k}$ for each $k = 1, 2, \dots, l$.

Set $W_k := U_{i_k} \cap V_{j_k}$ for each $k = 1, 2, \dots, l$. Since each W_k is open in $K_{\alpha_{i_k}} = K_{\beta_{j_k}}$ and $x_k \neq e$ for each k , we have that $x_1 x_2 \dots x_l \in W_1 W_2 \dots W_l \in \mathcal{B}$. Now to show that $W_1 W_2 \dots W_l \subset U_1 U_2 \dots U_n \cap V_1 V_2 \dots V_m$, pick any $y_1 y_2 \dots y_t \in W_1 W_2 \dots W_l$. Then for $k = 1, 2, \dots, t$, we have $y_k \in W_{s_k} = U_{i_{s_k}} \cap V_{j_{s_k}}$, for some $s_k \in \{1, 2, \dots, l\}$.

Since $x_1 x_2 \dots x_l \in U_1 U_2 \dots U_n$ we must have that $e \in U_i$ whenever $i \in \{1, 2, \dots, n\} \setminus \{i_k : k = 1, 2, \dots, l\}$. Again since $y_1 y_2 \dots y_t \in W_1 W_2 \dots W_l$, we must have that $e \in W_s = U_{i_s} \cap V_{j_s} \subset U_{i_s}$ whenever $s \in \{1, 2, \dots, l\} \setminus \{s_k : k = 1, 2, \dots, t\}$. Combining these two statements gives us that $e \in U_i$ whenever $i \in \{1, 2, \dots, n\} \setminus \{i_{s_k} : k = 1, 2, \dots, t\}$. Also, we know that $y_k \in U_{i_{s_k}}$ for each $k = 1, 2, \dots, t$. Hence it follows that $y_1 y_2 \dots y_t \in U_1 U_2 \dots U_n$.

Similarly since $x_1 x_2 \dots x_l \in V_1 V_2 \dots V_m$, we must have that $e \in V_j$ whenever $j \in \{1, 2, \dots, m\} \setminus \{j_k : k = 1, 2, \dots, l\}$ and since $y_1 y_2 \dots y_t \in W_1 W_2 \dots W_l$, we must have that $e \in W_s = U_{i_s} \cap V_{j_s} \subset V_{j_s}$ whenever $s \in \{1, 2, \dots, l\} \setminus \{s_k : k = 1, 2, \dots, t\}$, we can proceed in the same manner to get that $y_1 y_2 \dots y_t \in V_1 V_2 \dots V_m$.

Thus finally we see that $x_1 x_2 \dots x_l \in W_1 W_2 \dots W_l \subset (U_1 U_2 \dots U_n \cap V_1 V_2 \dots V_m)$ where $W_1 W_2 \dots W_l \in \mathcal{B}$, implying that \mathcal{B} serves as a base for a topology on K . We denote the topology on K generated by \mathcal{B} as τ .

Natural Binary Operation $*$ on $M(K)$:

As pointed out in the first section of this text, in order to define a binary operation on $M(K)$ it is sufficient to define a binary operation on the point-mass measures on $M(K)$. First of

all, if $\Lambda \neq \emptyset$ then for any $x_1x_2 \dots x_n \in K$ we define

$$p_e * p_e = p_e \quad ; \quad p_e * p_{x_1x_2 \dots x_n} = p_{x_1x_2 \dots x_n} = p_{x_1x_2 \dots x_n} * p_e.$$

Now for any two elements $x = x_1x_2 \dots x_n, y = y_1y_2 \dots y_m \in K \setminus \{e\}$ where $x_i \in K_{\alpha_i}, y_j \in K_{\beta_j}$ for $i = 1, 2, \dots, n; j = 1, 2, \dots, m$ and for any Borel set $E' \subseteq K$ we define:

$$(p_x * p_y)(E') = \begin{cases} p_{(x_1 \dots x_n y_1 \dots y_m)}(E'), & \text{if } \alpha_n \neq \beta_1. \\ (p_{x_n} * p_{y_1})(E), & \text{if } \alpha_n = \beta_1, \alpha_{n-1} \neq \beta_2 \\ & \text{and } E' \cap ((x_1 \dots x_{n-1})K_{\alpha_n}(y_2 \dots y_m)) \\ & = (x_1 \dots x_{n-1})E(y_2 \dots y_m). \\ (p_{x_n} * p_{y_1})(E \cup \{e\}), & \text{if } \alpha_n = \beta_1, \alpha_{n-1} = \beta_2 \\ & \text{and } E' \cap ((x_1 \dots x_{n-1})K_{\alpha_n}(y_2 \dots y_m)) \\ & = (x_1 \dots x_{n-1})E(y_2 \dots y_m) \neq \emptyset. \\ 0, & \text{if } \alpha_n = \beta_1, \alpha_{n-1} = \beta_2 \\ & \text{and } E' \cap ((x_1 \dots x_{n-1})K_{\alpha_n}(y_2 \dots y_m)) \\ & = \emptyset. \end{cases}$$

Note that when $\alpha_n = \beta_1$, then $(p_x * p_y)$ mimics the behaviour of the probability measure $(p_{x_n} * p_{y_1})$ and hence is a probability measure with support $(x_1 \dots x_{n-1}) \text{supp}(p_{x_n} * p_{y_1})(y_2 \dots y_m)$. **$(p_x * p_y)$ has compact support:**

Due to the construction of τ , for any two closed subsets $C_1 \subset K_{\alpha_1}$ and $C_2 \subset K_{\alpha_2}$ where $\alpha_1 \neq \alpha_2$, we have that C_1C_2 is a closed subset of K . This is true since $K_{\alpha_1}K_{\alpha_2}$ is open in K and we have that

$$K_{\alpha_1}K_{\alpha_2} \setminus C_1C_2 = (K_{\alpha_1} \setminus C_1) \cdot K_{\alpha_2} \cup K_{\alpha_1} \cdot (K_{\alpha_2} \setminus C_2),$$

which is open in $K_{\alpha_1}K_{\alpha_2}$. Similarly, we can show that $C_1C_2 \dots C_n$ is a closed set in K for any $n \in \mathbb{N}$ where $C_i \in K_{\alpha_i}$ for $1 \leq i \leq n$ and $\alpha_i \neq \alpha_{i+1}$ for $1 \leq i \leq n-1$. In fact, it is now easy to see that the set $\{C_1C_2 \dots C_n : n \in \mathbb{N}, C_i \text{ is a closed set in some } K_{\alpha_i}\}$ is the family of basic closed sets in K .

Now let both C_1 and C_2 are compact subsets of K_{α_1} and K_{α_2} respectively and let $\{A_\gamma\}_{\gamma \in I}$ be a family of closed subsets of C_1C_2 with finite intersection property. Then we must have that $A_\gamma = B_1^\gamma B_2^\gamma$ for some closed subsets $B_i^\gamma \subset C_i \subset K_{\alpha_i}$ for each $\gamma, i = 1, 2$. Now for any $\{\gamma_1, \gamma_2, \dots, \gamma_n\} \subset I$ we have that $\bigcap_1^n B_1^{\gamma_i} B_2^{\gamma_i} = \bigcap_1^n A_{\gamma_i} \neq \emptyset$.

Hence in particular, $\bigcap_1^n B_1^{\gamma_i} \neq \emptyset$ for any such finite subset $\{\gamma_1, \gamma_2, \dots, \gamma_n\} \subset I$. So $\{B_1^\gamma\}_{\gamma \in I}$ is a family of subsets in C_1 with the finite intersection property. But C_1 is compact in K_{α_1} and hence there exists some $x_1 \in C_1$ such that $x_1 \in \bigcap_{\gamma \in I} B_1^\gamma$. Similarly, since C_2 is also compact in K_{α_2} , there exists some $x_2 \in C_2$ such that $x_2 \in \bigcap_{\gamma \in I} B_2^\gamma$. Thus we see that there exists an

element $x \in C_1C_2$ such that $x \in \cap_{\gamma \in I} A_\gamma$ where x is given as the following:

$$x = \begin{cases} (x_1x_2), & \text{if } x_1, x_2 \neq e. \\ (x_1), & \text{if } x_1 \neq e, x_2 = e. \\ (x_2), & \text{if } x_1 = e, x_2 \neq e. \\ e, & \text{if } x_1 = x_2 = e. \end{cases}$$

Thus we see that C_1C_2 has to be compact too. Now let C_1, C_2, C_3 be compact subsets of $K_{\alpha_1}, K_{\alpha_2}, K_{\alpha_3}$ respectively. Note that if $\alpha_i \neq \alpha_j$ for any $i, j \in \{1, 2, 3\}$ or if $e \notin \cap_{i=1}^3 C_i$ then we can proceed similarly as above to show that $C_1C_2C_3$ gives us a compact set in K . Otherwise, first consider the following case.

Let $C_0 := (a_1 \dots a_s)C(b_1 \dots b_t)$, where C is a compact set in K_{α_0} and $(a_1a_2 \dots a_s), (b_1b_2 \dots b_t) \in K \setminus \{e\}$ where $a_i \in K_{\gamma_i}$ and $b_j \in K_{\delta_j}$ for $1 \leq i \leq s, 1 \leq j \leq t$ such that $\alpha_0 \neq \gamma_s, \delta_1$. We claim that any such C_0 is compact in (K, τ) .

Let $\{U_\alpha\}$ be any basic open cover of C_0 . Hence $U_\alpha = U_1^\alpha U_2^\alpha \dots U_{n_\alpha}^\alpha$ for some $n_\alpha \in \mathbb{N}$ where each U_k^α is an open set in some K_α such that the product exists. Now for each $x \in C \setminus \{e\}$ there exists α_x such that $(a_1 \dots a_s x b_1 \dots b_t) \in U_{\alpha_x}$. Therefore there exists $i_x \in \{1, 2, \dots, n_{\alpha_x}\}$ such that $x \in U_{i_x}^{\alpha_x}$ where $U_{i_x}^{\alpha_x}$ is an open set in K_{α_0} . Thus $\mathcal{U} := \{U_{i_x}^{\alpha_x}\}_{x \in C}$ serves an open cover of the set $C \setminus \{e\}$.

If $e \notin C$, then \mathcal{U} serves as an open cover of C and hence has a finite subcover $\{U_1, U_2, \dots, U_p\}$ where $U_k = U_{i_{x_k}}^{\alpha_{x_k}}$ for some $x_k \in C$ for $1 \leq k \leq p$. Thus $\{U_{\alpha_{x_1}}, U_{\alpha_{x_2}}, \dots, U_{\alpha_{x_p}}\}$ serves as a finite subcover of $\{U_\alpha\}$.

Now let $e \in C$. Pick $\alpha' \in \Lambda$ such that $\alpha' \neq \alpha_0$. Since $K_{\alpha'}$ is locally compact, we can find a compact neighborhood F of e in $K_{\alpha'}$. Let $\mathcal{V} := \{V_\beta\}$ be an open cover of F . Then $C \cup F$ is a compact set in $K_{\alpha_0} \cup K_{\alpha'}$ with open cover $\mathcal{U} \cup \mathcal{V}$. Thus we get a finite subcover of the form $\{U_1, U_2, \dots, U_p, V_{\beta_1}, V_{\beta_2}, \dots, V_{\beta_q}\}$ where $U_k = U_{i_{x_k}}^{\alpha_{x_k}}$ for some $x_k \in C \setminus \{e\}$ for $1 \leq k \leq p$. Note that $e \in V_{\beta_l}$ for some $l \in \{1, 2, \dots, q\}$ here and hence $\{U_{i_{x_k}}^{\alpha_{x_k}}\}_{k=1}^p$ covers $C \setminus \{e\}$.

Thus if $\gamma_s = \delta_1$, then $\{U_{\alpha_{x_1}}, U_{\alpha_{x_2}}, \dots, U_{\alpha_{x_p}}\}$ covers C_0 , as desired. Otherwise if $\gamma_s \neq \delta_1$, then there exists some α' such that $(a_1 \dots a_s b_1 \dots b_t) \in U_{\alpha'}$. Hence in this case, $\{U_{\alpha_{x_1}}, U_{\alpha_{x_2}}, \dots, U_{\alpha_{x_p}}, U_{\alpha'}\}$ serves as a finite subcover of C_0 . Hence we see that any set of the form C_0 is compact in (K, τ) . Using similar argument we can immediately see that sets of the form $C_1C_2C_3$ are also compact where $C_i \in K_{\alpha_i}$ is compact.

In particular, for any two words $x = (x_1x_2 \dots x_n), y = (y_1y_2 \dots y_m) \in K$ where $x_i \in K_{\alpha_i}, y_j \in K_{\beta_j}$ and $\alpha_n = \beta_1$, we have that

$$\text{supp}(p_x * p_y) = (x_1 \dots x_{n-1}) \text{supp}(p_{x_n} * p_{y_1})(y_2 \dots y_m)$$

is compact since $\text{supp}(p_{x_n} * p_{y_1})$ is compact in $K_{\alpha_n} = K_{\beta_1}$. Thus it follows from the construction of convolution products that for any two words $x, y \in K$ we have that $(p_x * p_y)$ is a probability measure with compact support. Also since convolution is associative in each $M(K_\alpha)$ and concatenation of finite words is associative, it follows from the construction that convolution products on $M(K)$ are associative.

τ is Hausdorff:

Now we proceed further to show that $(K, *)$ indeed gives us a semihypergroup. First note that since each K_α is Hausdorff, for any two elements $x_1x_2 \dots x_n, y_1y_2 \dots y_m \in K$ where $x_i \in K_{\alpha_i}, y_j \in K_{\beta_j}$, we can find sets $A_i \subset K_{\alpha_i}, B_j \subset K_{\beta_j}$ for $1 \leq i \leq n, 1 \leq j \leq m$ such that $x_i \in A_i, y_j \in B_j$ and $A_i \cap B_j = \emptyset$ for all i, j . Hence we have that $x_1x_2 \dots x_n \in A_1A_2 \dots A_n, y_1y_2 \dots y_m \in B_1B_2 \dots B_m$ where $A_1A_2 \dots A_n \cap B_1B_2 \dots B_m = \emptyset$. Also if $e \in K$, then for any such $x_1x_2 \dots x_n$ since $x_i \neq e$ for any i , we can choose A_i 's such that $e \notin A_i$ for any i . Then again using the fact that each K_α is Hausdorff, we can choose a neighborhood V around e such that $V \cap A_i = \emptyset$ for all i and hence $V \cap A_1A_2 \dots A_n = \emptyset$. Thus we can easily see that the topology τ on K is Hausdorff.

 τ is locally compact:

Next pick any element $x_1x_2 \dots x_n \in K$ where $x_i \in K_{\alpha_i}$ for $1 \leq i \leq n$. Since each K_{α_i} is locally compact and Hausdorff, we can find a compact neighborhood C_i of x_i in K_{α_i} such that $e \notin C_i$. Let U_i be an open set in K_{α_i} such that $C_i \subset U_i$ and $e \notin U_i$. Now consider the map $\phi : U_1 \times U_2 \times \dots \times U_n \rightarrow U_1U_2 \dots U_n$ given by

$$\phi((a_1, a_2, \dots, a_n)) := (a_1a_2 \dots a_n)$$

for any $(a_1, a_2, \dots, a_n) \in U_1 \times U_2 \times \dots \times U_n$.

Note that since $e \notin U_i$ for any i , the map ϕ is bijective. Moreover, for any basic open set $V_1V_2 \dots V_n \subset U_1U_2 \dots U_n$ we have $\phi^{-1}(V_1V_2 \dots V_n) = V_1 \times V_2 \times \dots \times V_n$ which is open in $U_1 \times U_2 \times \dots \times U_n$ in the product topology. Hence ϕ is continuous, granting us that $C_1C_2 \dots C_n = \phi(C_1 \times C_2 \times \dots \times C_n)$ is a compact neighborhood of $x_1x_2 \dots x_n$ in (K, τ) . Also if $e \in K$ then there exists some α_0 such that $e \in K_{\alpha_0}$. Since K_{α_0} is locally compact we can always find a compact neighborhood for e . Thus we see that (K, τ) is a locally compact Hausdorff space.

Continuity of support with respect to Michael Topology:

Now in order to see if the map $(x, y) \mapsto \text{supp}(p_x * p_y) : K \times K \rightarrow \mathfrak{C}(K)$ is continuous, first recall [9] that the map $\pi : X \rightarrow M(X)$ given by $x \mapsto p_x$ is a homeomorphism onto its image for any locally compact Hausdorff space X . Now let $\{(x_\alpha, y_\alpha)\}$ be a net in $K \times K$ that converges to some (x, y) where $x = x_1x_2 \dots x_n, y = y_1y_2 \dots y_m \in K$ such that $x_i \in K_{\gamma_i}$ and $y_j \in K_{\delta_j}$ for all i, j . Then $\{x_\alpha\}$ and $\{y_\alpha\}$ converges to x and y respectively. Then for each α we have that $x_\alpha = x_1^\alpha x_2^\alpha \dots x_{n_\alpha}^\alpha$ for some $n_\alpha \in \mathbb{N}$ where $x_i^\alpha \in K_{\alpha_i}$ for $1 \leq i \leq n_\alpha$. But x_α eventually lies in any basic open set of the form $U_1U_2 \dots U_n$ around x where U_i is an open neighborhood of x_i in K_{γ_i} that does not contain e . Hence without loss of generality for each α we can assume that $n_\alpha = n$ and $\alpha_i = \gamma_i$ for $1 \leq i \leq n$. Similarly we can assume that $y_\alpha = y_1^\alpha y_2^\alpha \dots y_m^\alpha$ where $y_j^\alpha \in K_{\delta_j}$ for $1 \leq j \leq m$.

Now consider the Michael topology on $\mathfrak{C}(K)$ and recall [16] that the map

$$x \mapsto \{x\} : K \rightarrow \mathfrak{C}(K)$$

is a homeomorphism onto its image which is also closed in $\mathfrak{C}(K)$. If $\gamma_n \neq \delta_1$ then using the same technique as above, we can easily see that the net $\{(x_1^\alpha \dots x_n^\alpha y_1^\alpha \dots y_m^\alpha)\}_\alpha$ converges to $(x_1 \dots x_n y_1 \dots y_m)$ in K . Hence using the homeomorphism mentioned

above we have that $\text{supp}(p_{x_\alpha} * p_{y_\alpha}) = \{(x_1^\alpha \dots x_n^\alpha y_1^\alpha \dots y_m^\alpha)\}$ converges to $\{(x_1 \dots x_n y_1 \dots y_m)\} = \text{supp}(p_x * p_y)$ in $\mathfrak{C}(K)$ in Michael Topology.

Now let $\gamma_n = \delta_1$. For any $i_0 \in \{1, 2, \dots, n\}$, pick and fix open neighborhoods U_i around x_i in K_{γ_i} for all $i \neq i_0$. Then for any open neighborhood V around x_{i_0} in $K_{\gamma_{i_0}}$ we have that $x_1^\alpha x_2^\alpha \dots x_n^\alpha$ lie eventually in $U_1 U_2 \dots U_{i_0-1} V U_{i_0+1} \dots U_n$. Hence we must have that $\{x_{i_0}^\alpha\}$ lie eventually in V , i.e., $\{x_{i_0}^\alpha\}$ converges to x_{i_0} in $K_{\gamma_{i_0}}$. Thus $\{x_i^\alpha\}$ converges to x_i in K_{γ_i} and $\{y_j^\alpha\}$ converges to y_j in K_{δ_j} for each i, j . In particular, we have that the net $\{(x_n^\alpha, y_1^\alpha)\}_\alpha$ converges to (x_n, y_1) in $K_{\gamma_n} \times K_{\gamma_n}$. Since the map $(x, y) \mapsto \text{supp}(p_x * p_y) : K_{\gamma_n} \times K_{\gamma_n} \rightarrow \mathfrak{C}(K_{\gamma_n})$ is continuous, we have that $\text{supp}(p_{x_n^\alpha} * p_{y_1^\alpha})$ converges to $\text{supp}(p_{x_n} * p_{y_1})$ in $\mathfrak{C}(K_{\gamma_n})$.

As discussed before, we also have that the nets $\{\{x_i^\alpha\}\}_\alpha$ and $\{\{y_j^\alpha\}\}_\alpha$ converge to $\{x_i\}$ and $\{y_j\}$ in $\mathfrak{C}(K_{\gamma_i})$ and $\mathfrak{C}(K_{\delta_j})$ respectively for $1 \leq i < n$ and $1 < j \leq m$. Hence it follows from the construction of basic open sets of K that $\text{supp}(p_{x_\alpha} * p_{y_\alpha}) = \{(x_1^\alpha \dots x_{n-1}^\alpha)\} \text{supp}(p_{x_n^\alpha} * p_{y_1^\alpha}) \{(y_2^\alpha \dots y_m^\alpha)\}$ converges to $\{(x_1 \dots x_{n-1})\} \text{supp}(p_{x_n} * p_{y_1}) \{(y_2 \dots y_m)\} = \text{supp}(p_x * p_y)$ in $\mathfrak{C}(K)$.

Finally, it follows immediately from the construction of convolution products that the map $* : M(K) \times M(K) \rightarrow M(K)$ is positive bilinear. Using the same technique as above we can see that the restricted map $*|_{M^+(K) \times M^+(K)} : M^+(K) \times M^+(K) \rightarrow M^+(K)$ is continuous. Thus we have that the pair $(K, *)$ indeed forms a semihypergroup.

K_α as a sub-semihypergroup:

Note that for each α the map $i_\alpha : K_\alpha \rightarrow K$ given by

$$i_\alpha(x) = \begin{cases} (x), & \text{if } x \neq e \\ e, & \text{if } x = e \end{cases}$$

where (x) is a word of length 1 in K , enables us to view K_α as a sub-semihypergroup of K since $i_\alpha(K_\alpha) * i_\alpha(K_\alpha) \subset i_\alpha(K_\alpha)$ and i_α is a homeomorphism onto its image.

Universal property:

Now let H be a semihypergroup and $\phi_\alpha : K_\alpha \rightarrow H$ be a homomorphism for each α . For each $x = x_1 x_2 \dots x_n \in K$ where $x_i \in K_{\alpha_i}$ for $1 \leq i \leq n$ define

$$\Gamma(p_x) := p_{\phi_{\alpha_1}(x_1)} * p_{\phi_{\alpha_2}(x_2)} * \dots * p_{\phi_{\alpha_n}(x_n)}.$$

We can then extend Γ to $M(K)$ as in the proof of Theorem 3.3 to get a positive continuous linear map $\Gamma : M(K) \rightarrow M(H)$.

Since for any $x \in K_\alpha$ we have that $\Gamma(p(x)) = p_{\phi_\alpha(x)}$, it follows immediately from Theorem 3.3 that $\Gamma|_{M(K_\alpha)} = \Gamma \circ \Gamma_{i_\alpha} = \Gamma_{\phi_\alpha}$ for each α . Now to show that Γ is a homomorphism, pick any two words $x = x_1 x_2 \dots x_n, y = y_1 y_2 \dots y_m$ in K where $x_i \in K_{\alpha_i}, y_j \in K_{\beta_j}$ for $1 \leq i \leq n, 1 \leq j \leq m$. Note that if $\alpha_n \neq \beta_1$ then we have that

$$\begin{aligned}
\Gamma(p_x * p_y) &= \Gamma(p_{x_1 \dots x_n y_1 \dots y_m}) \\
&= p_{\phi_{\alpha_1}(x_1)} * \dots * p_{\phi_{\alpha_n}(x_n)} * p_{\phi_{\beta_1}(y_1)} * \dots * p_{\phi_{\beta_m}(y_m)} \\
&= (p_{\phi_{\alpha_1}(x_1)} * \dots * p_{\phi_{\alpha_n}(x_n)}) * (p_{\phi_{\beta_1}(y_1)} * \dots * p_{\phi_{\beta_m}(y_m)}) \\
&= \Gamma(p_{x_1 x_2 \dots x_n}) * \Gamma(p_{y_1 y_2 \dots y_m}) = \Gamma(p_x) * \Gamma(p_y)
\end{aligned}$$

as required.

Now pick any $x \in K_\gamma$ and $y, z \in K_\beta$ where $\gamma \neq \beta$. Then there exists a net $\{m_\alpha\}$ in $M_F^+(K_\beta)$ such that m_α converges to $(p_y * p_z)$ in $M^+(K_\beta)$. Hence $(p_x * m_\alpha)$ converges to $(p_x * (p_y * p_z)) = (p_x * p_y * p_z)$ in $M^+(K)$. Then for each α we have that $m_\alpha = \sum_{i=1}^{n_\alpha} c_i^\alpha p_{x_i^\alpha}$ for some $n_\alpha \in \mathbb{N}$, $c_i^\alpha \in \mathbb{C}$ and $x_i^\alpha \in K_\beta$. Since Γ is continuous on $M^+(K)$, we have that

$$\begin{aligned}
\Gamma(p_x * p_y * p_z) &= \Gamma(\lim_\alpha (p_x * m_\alpha)) \\
&= \lim_\alpha \Gamma(p_x * m_\alpha) \\
&= \lim_\alpha \Gamma\left(p_x * \sum_{i=1}^{n_\alpha} c_i^\alpha p_{x_i^\alpha}\right) \\
&= \lim_\alpha \Gamma\left(\sum_{i=1}^{n_\alpha} c_i^\alpha (p_x * p_{x_i^\alpha})\right) \\
&= \lim_\alpha \sum_{i=1}^{n_\alpha} c_i^\alpha \Gamma(p_x * p_{x_i^\alpha}) \\
&= \lim_\alpha \sum_{i=1}^{n_\alpha} c_i^\alpha \left(\Gamma(p_x) * \Gamma(p_{x_i^\alpha})\right) \\
&= \lim_\alpha \Gamma(p_x) * \left(\sum_{i=1}^{n_\alpha} c_i^\alpha \Gamma(p_{x_i^\alpha})\right) \\
&= \lim_\alpha \Gamma(p_x) * \Gamma\left(\sum_{i=1}^{n_\alpha} c_i^\alpha p_{x_i^\alpha}\right) \\
&= \Gamma(p_x) * \lim_\alpha \Gamma(m_\alpha) = \Gamma(p_x) * \Gamma(p_y * p_z)
\end{aligned}$$

where the fourth equality follows since the convolution $(\mu, \nu) \mapsto \mu * \nu : M(K) \times M(K) \rightarrow M(K)$ is bilinear, the fifth and eighth equality follows from the linearity of Γ on $M(K)$ and finally the seventh equality follows since the map $(\mu, \nu) \mapsto \mu * \nu : M(H) \times M(H) \rightarrow M(H)$ is bilinear.

Similarly for any $x, y \in K_\gamma$ and $z \in K_\beta$ where $\gamma \neq \beta$ we have that

$$\Gamma(p_x * p_y * p_z) = \Gamma(p_x * p_y) * \Gamma(p_z).$$

Using these two equalities simultaneously for any two words $x = x_1x_2 \dots x_n$, $y = y_1y_2 \dots y_m$ in K where $x_i \in K_{\alpha_i}$, $y_j \in K_{\beta_j}$ for $1 \leq i \leq n$, $1 \leq j \leq m$ and $\alpha_n = \beta_1$, we have that

$$\begin{aligned}
\Gamma(p_x * p_y) &= \Gamma(p_{x_1x_2 \dots x_{n-1}} * p_{x_n} * p_{y_1} * p_{y_2y_3 \dots y_m}) \\
&= \Gamma(p_{x_1x_2 \dots x_{n-1}}) * \Gamma(p_{x_n} * p_{y_1}) * \Gamma(p_{y_2y_3 \dots y_m}) \\
&= \Gamma(p_{x_1x_2 \dots x_{n-1}}) * \Gamma_{\phi_{\alpha_n}}(p_{x_n} * p_{y_1}) * \Gamma(p_{y_2y_3 \dots y_m}) \\
&= \Gamma(p_{x_1 \dots x_{n-1}}) * \left(\Gamma_{\phi_{\alpha_n}}(p_{x_n}) * \Gamma_{\phi_{\alpha_n}}(p_{y_1}) \right) * \Gamma(p_{y_2 \dots y_m}) \\
&= \Gamma(p_{x_1 \dots x_{n-1}}) * \left(p_{\phi_{\alpha_n}(x_n)} * p_{\phi_{\alpha_n}(y_1)} \right) * \Gamma(p_{y_2 \dots y_m}) \\
&= \left(p_{\phi_{\alpha_1}(x_1)} * \dots * p_{\phi_{\alpha_{n-1}}(x_{n-1})} \right) * \left(p_{\phi_{\alpha_n}(x_n)} * p_{\phi_{\beta_1}(y_1)} \right) \\
&\quad * \left(p_{\phi_{\beta_2}(y_2)} * \dots * p_{\phi_{\beta_m}(y_m)} \right) \\
&= \left(p_{\phi_{\alpha_1}(x_1)} * \dots * p_{\phi_{\alpha_{n-1}}(x_{n-1})} * p_{\phi_{\alpha_n}(x_n)} \right) \\
&\quad * \left(p_{\phi_{\beta_1}(y_1)} * p_{\phi_{\beta_2}(y_2)} * \dots * p_{\phi_{\beta_m}(y_m)} \right) \\
&= \Gamma(p_{x_1x_2 \dots x_n}) * \Gamma(p_{y_1y_2 \dots y_m}) = \Gamma(p_x) * \Gamma(p_y)
\end{aligned}$$

as required. Note that here the fourth equality follows since $\Gamma_{\phi_{\alpha_n}} : M(K_{\alpha_n}) \rightarrow M(H)$ induced by ϕ_{α_n} is a homomorphism and the fifth equality follows from the construction of $\Gamma_{\phi_{\alpha_n}}$ as shown in Theorem 3.3.

Finally, note that if $\Theta : M(K) \rightarrow M(H)$ is another positive linear continuous homomorphism such that $\Theta|_{M(K_{\alpha})} = \Gamma_{\phi_{\alpha}}$ then for any $x_1x_2 \dots x_n \in K$ where $x_i \in K_{\alpha_i}$ for $1 \leq i \leq n$ we have that

$$\begin{aligned}
\Theta(p_{x_1x_2 \dots x_n}) &= \Theta(p_{x_1} * p_{x_2} * \dots * p_{x_n}) \\
&= \Theta(p_{x_1}) * \Theta(p_{x_2}) * \dots * \Theta(p_{x_n}) \\
&= \Theta|_{M(K_{\alpha_1})}(p_{x_1}) * \Theta|_{M(K_{\alpha_2})}(p_{x_2}) * \dots * \Theta|_{M(K_{\alpha_n})}(p_{x_n}) \\
&= \Gamma_{\phi_{\alpha_1}}(p_{x_1}) * \Gamma_{\phi_{\alpha_2}}(p_{x_2}) * \dots * \Gamma_{\phi_{\alpha_n}}(p_{x_n}) \\
&= \Gamma|_{M(K_{\alpha_1})}(p_{x_1}) * \Gamma|_{M(K_{\alpha_2})}(p_{x_2}) * \dots * \Gamma|_{M(K_{\alpha_n})}(p_{x_n}) \\
&= \Gamma(p_{x_1}) * \Gamma(p_{x_2}) * \dots * \Gamma(p_{x_n}) \\
&= \Gamma(p_{x_1} * p_{x_2} * \dots * p_{x_n}) = \Gamma(p_{x_1x_2 \dots x_n}).
\end{aligned}$$

Thus we see that the map Γ constructed above is unique and hence $(K, *)$ satisfies all the conditions of Definition 3.4 giving us that $K = \prod_{\alpha}^* K_{\alpha}$. \square \square

Remark 3.7. Note that in classical literature, this natural construction fails to provide a free product of topological semigroups or groups. But in the case of pure semihypergroups, the natural construction suffices to provide a free product structure, *i.e.*, in the most trivial case, when applied to a family of semigroups which are pure semihypergroups, the natural construction explained above results in a semihypergroup, rather than a semigroup.

4. OPEN QUESTIONS AND FURTHER WORK

As discussed briefly in the introduction, the lack of prior extensive research since its inception and the significant examples available in coset and orbit spaces of locally compact groups, Lie groups and homogeneous spaces, makes way to a number of exciting new avenues of research on semihypergroups.

We have already discussed [1] spaces of almost periodic and weakly almost periodic functions, amenability, ideals and homomorphisms on a semihypergroup. The following is some immediate potential problems related to this article that we are currently working on and/or intend to work on in near future.

Problem 1: Investigate the behavior of free products on semihypergroups for compact case, thereby exploring amenability properties of the resulting free product. While the compactness of the individual components make some of the deductions easier enabling us to drop some prior restrictions, as in most cases, the lack of an algebra in the underlying space prevents any direct extension of similar results [14, 15] to this setup.

Problem 2: Generalize the construction of the above free product for amalgamated products as well as investigate the properties of the resulting semihypergroup, similar to the works in [15].

Problem 3: Amend and update the construction of the above free product for a family of hypergroups, such that the resulting free product becomes a hypergroup. This needs updating the construction of the natural topology τ so that it takes into account the involution of respective hypergroups, eventually resulting in a natural involution of the form $x_1 x_2 \dots x_{n-1} x_n \mapsto x_n^- x_{n-1}^- \dots x_2^- x_1^-$.

5. ACKNOWLEDGEMENTS

The author would like to sincerely thank her doctoral thesis advisor Dr. Anthony To-Ming Lau, for suggesting the topic of this study and for the helpful discussions during the course of this work. She is also grateful to the referee for the valuable comments and suggestions.

REFERENCES

- [1] Bandyopadhyay, C. : *Analysis on Semihypergroups: Function Spaces, Homomorphisms and Ideals*. Semigroup Forum. 100, 671-697 (2020)
- [2] Berglund, J. F., Junghenn, H. D., Milnes, P. : *Analysis on Semigroups*. Canadian Mathematical Society Series of Monographs and Advanced Texts, Wiley Interscience Publication. 24 (1989)
- [3] Day, M. M. : *Amenable Semigroups*. Illinois J. Math. 1, 509-544 (1957)
- [4] Dunkl, C. F. : *The Measure Algebra of a Locally Compact Hypergroup*. Trans. Amer. Math. Soc. 179, 331-348 (1973)
- [5] Forrest, B. : *Amenability and Ideals in $A(G)$* . J. Austral. Math. Soc. Ser. A 53, no. 2, 143-155 (1992)
- [6] Forrest, B., Spronk, N., Samei, E. : *Convolutions on Compact Groups and Fourier Algebras of Coset Spaces*. Studia Math. 196 no. 3, 223-249 (2010)
- [7] Ghahramani, F., Medgalchi, A. R. : *Compact Multipliers on Weighted Hypergroup Algebras*. Math. Proc. Cambridge Philos. Soc. 98, 493-500 (1985)

- [8] Ghahramani, F., Lau, A. T., Losert, V. : *Isometric Isomorphisms between Banach Algebras Related to Locally Compact Groups*. Trans. Amer. Math. Soc. 321 no. 1, 273-283 (1990)
- [9] Jewett, R. I. : *Spaces with an Abstract Convolution of Measures*. Advances in Mathematics. 18, no. 1, 1-101 (1975)
- [10] Lau, A. T. : *Uniformly Continuous Functions on Banach Algebras*. Colloq. Math. 51, 195-205 (1987)
- [11] Lau, A. T., Dales, H. G., Strauss, D. P. : *Banach Algebras on Semigroups and on their Compactifications*. Mem. Amer. Math. Soc. 205, no. 966, vi+165 pp (2010)
- [12] Michael, E. : *Topologies on Spaces of Subsets*. Trans. Amer. Math. Soc. 71, 152-182 (1951)
- [13] Mitchell, T. : *Topological semigroups and fixed points*. Illinois J. Math. 14, 630-641 (1970)
- [14] Morris, S. A. : *Free Products of Topological Groups*. Bull. Austral. Math. Soc. 4, 17-29 (1971)
- [15] Morris, S. A., Khan, M. S. : *Free products of topological groups with central amalgamation. I*. Trans. Amer. Math. Soc. 273, 405-416 (1982)
- [16] Nachbin, L. : *On the Finite Dimensionality of Every Irreducible Representation of a Compact Group*. Proc. Amer. Math. Soc. 12, 11-12 (1961)
- [17] Pier, J-P. : *Amenable Banach algebras*. Pitman Research Notes in Mathematics Series, 172. Longman Scientific and Technical, Harlow; John Wiley and Sons Inc., New York, (1988)
- [18] Ross, K. A. : *Hypergroups and Centers of Measure Algebras*. Symposia Mathematica, Vol. XXII, 189-203 (1977)
- [19] Spector, R. : *Aperçu de la théorie des hypergroupes, (French) Analyse harmonique sur les groupes de Lie (Sém. Nancy-Strasbourg, 1973-75)*. Lecture Notes in Math. 497, Springer-Verlag, New York, 643-673 (1975)
- [20] Zeuner, H. : *One Dimensional Hypergroups*. Advances in Mathematics 76, no. 1, 1-18 (1989)

DEPARTMENT OF MATHEMATICAL AND STATISTICAL SCIENCES, UNIVERSITY OF ALBERTA, CANADA T6G 2G1

Email address: choiti@ualberta.ca