

THE TWO-PHASE PROBLEM FOR HARMONIC MEASURE IN VMO AND THE CHORD-ARC CONDITION

XAVIER TOLSA AND TATIANA TORO

ABSTRACT. Let $\Omega^+ \subset \mathbb{R}^{n+1}$ be a bounded δ -Reifenberg flat domain, with $\delta > 0$ small enough, possibly with locally infinite surface measure. Assume also that $\Omega^- = \mathbb{R}^{n+1} \setminus \overline{\Omega^+}$ is an NTA domain as well and denote by ω^+ and ω^- the respective harmonic measures of Ω^+ and Ω^- with poles $p^\pm \in \Omega^\pm$. In this paper we show that the condition that $\log \frac{d\omega^-}{d\omega^+} \in \text{VMO}(\omega^+)$ is equivalent to Ω^+ being a chord-arc domain with inner normal belonging to $\text{VMO}(\mathcal{H}^n|_{\partial\Omega^+})$.

1. INTRODUCTION

This paper is devoted to study a two-phase problem for harmonic measure. In this type of problems one considers two disjoint domains $\Omega^+, \Omega^- \subset \mathbb{R}^{n+1}$ whose boundaries have non-empty intersection, and whose respective harmonic measures ω^+, ω^- are usually mutually absolutely continuous in some subset of $\partial\Omega^+ \cap \partial\Omega^-$. Then the objective is to relate the analytic properties of the density $\frac{d\omega^-}{d\omega^+}|_{\partial\Omega^+ \cap \partial\Omega^-}$ to the geometric properties of $\partial\Omega^+ \cap \partial\Omega^-$. For example, the works [AMT1], [AMTV] show that if ω^+ and ω^- are mutually absolutely continuous in some subset $E \subset \partial\Omega^+ \cap \partial\Omega^-$, then the measures $\omega^+|_E$ and $\omega^-|_E$ are n -rectifiable measures, that is, they are concentrated in an n -rectifiable subset of E and they are absolutely continuous with respect to the Hausdorff n -dimensional measure \mathcal{H}^n . Recall that a set $F \subset \mathbb{R}^d$ is called n -rectifiable if there are Lipschitz maps $f_i : \mathbb{R}^n \rightarrow \mathbb{R}^d$, $i = 1, 2, \dots$, such that

$$(1.1) \quad \mathcal{H}^n\left(F \setminus \bigcup_i f_i(\mathbb{R}^n)\right) = 0.$$

See [KPT] for a previous partial related result where Ω^+ and Ω^- are assumed to be complementary NTA domains (see Section 2.3 for the definition of NTA domain), and [AM] for a related work involving elliptic measure.

In other works of more quantitative nature, besides assuming that Ω^+ and Ω^- are complementary NTA domains, one asks also some quantitative condition about the density $\frac{d\omega^-}{d\omega^+}$. For example, in [KT3] the authors prove (among other results) that if $\log \frac{d\omega^-}{d\omega^+} \in \text{VMO}(\omega^+)$ and moreover Ω^+ is δ -Reifenberg flat for some $\delta > 0$ small enough, then Ω^+ is vanishing Reifenberg flat (see again Section 2.3 for the notions of δ -Reifenberg flatness and vanishing Reifenberg flatness). Later on, Engelstein [En] showed that if one strengthens the VMO condition on ω^+ by asking $\log \frac{d\omega^-}{d\omega^+} \in C^\alpha$ for some $\alpha > 0$ (still under the δ -Reifenberg flat assumption) then the inner unit normal N of Ω^+ belongs to C^α and Ω^+ is a $C^{1+\alpha}$ domain. On the other hand, in [AMT2] it is shown that

X.T. was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement 101018680). Also partially supported by MICINN (Spain) under the grant PID2020-114167GB-I00 and the María de Maeztu Program for units of excellence CEX2020-001084-M (Spain). T.T. was partially supported by the Craig McKibben & Sarah Merner Professorship in Mathematics and NSF grant DMS-1954545.

the condition $\omega^- \in A_\infty(\omega^+)$ is equivalent to the fact that Ω^+ and Ω^- have joint big pieces of chord-arc subdomains. See Section 2.3 for the precise definition. See also [BH] for a related result involving the conditions $\log \frac{d\omega^+}{d\mathcal{H}^n|_{\partial\Omega^+}} \in \text{VMO}(\mathcal{H}^n|_{\partial\Omega^+})$ and $\log \frac{d\omega^-}{d\mathcal{H}^n|_{\partial\Omega^+}} \in \text{VMO}(\mathcal{H}^n|_{\partial\Omega^+})$; and see the papers [BET1] and [BET2] for results about the structure of the singular set of the boundary under the assumption that $\log \frac{d\omega^-}{d\omega^+}$ belongs either to $\text{VMO}(\omega^+)$ or to C^α .

In connection with the precise results that we obtain in this work, the following theorem of Prats and the first author of this paper is particularly relevant.

Theorem A ([PT]). *Let $\Omega^+ \subset \mathbb{R}^{n+1}$ be a bounded NTA domain and let $\Omega^- = \mathbb{R}^{n+1} \setminus \overline{\Omega^+}$ be an NTA domain as well. Denote by ω^+ and ω^- the respective harmonic measures with poles $p^+ \in \Omega^+$ and $p^- \in \Omega^-$. Suppose that Ω^+ is a δ -Reifenberg flat domain, with $\delta > 0$ small enough. Then the following conditions are equivalent:*

- (a) ω^+ and ω^- are mutually absolutely continuous and $\log \frac{d\omega^-}{d\omega^+} \in \text{VMO}(\omega^+)$.
- (b) Ω^+ is vanishing Reifenberg flat, Ω^+ and Ω^- have joint big pieces of chord-arc subdomains, and

$$(1.2) \quad \lim_{\rho \rightarrow 0} \sup_{r(B) \leq \rho} \int_B |N - N_B| d\omega^+ = 0,$$

where N_B is the normal to the n -plane L_B pointing to Ω^+ and minimizing

$$(1.3) \quad \max \left\{ \sup_{y \in \partial\Omega^+ \cap B} \text{dist}(y, L_B), \sup_{y \in L_B \cap B} \text{dist}(y, \partial\Omega^+) \right\}.$$

Notice that the condition (b) above provides a geometric characterization of the analytic condition $\log \frac{d\omega^-}{d\omega^+} \in \text{VMO}(\omega^+)$, under the assumption that Ω^+ is a δ -Reifenberg flat domain, with $\delta > 0$ small enough. Remark that the Reifenberg flatness condition on the domain is necessary in the theorem above. This can be easily seen by taking a suitable smooth truncation of the cone $\Omega^+ = \{(x_1, x_2, x_3, x_4) \in \mathbb{R}^4 : x_1^2 + x_2^2 < x_3^2 + x_4^2\}$, for which the harmonic measures ω^+ and ω^- with pole at ∞ coincide.

The main result of the current paper is the following.

Theorem 1.1. *Let $\Omega^+ \subset \mathbb{R}^{n+1}$ be a bounded NTA domain and let $\Omega^- = \mathbb{R}^{n+1} \setminus \overline{\Omega^+}$ be an NTA domain as well. Denote by ω^+ and ω^- the respective harmonic measures with poles $p^+ \in \Omega^+$ and $p^- \in \Omega^-$. Suppose that Ω^+ is a δ -Reifenberg flat domain, with $\delta > 0$ small enough. Then the conditions (a) and (b) in Theorem A are equivalent to the following:*

- (c) Ω^+ is a chord-arc domain and the inner normal N to $\partial\Omega^+$ belongs to $\text{VMO}(\mathcal{H}^n|_{\partial\Omega^+})$.

Recall that a chord-arc domain in \mathbb{R}^{n+1} is an NTA domain with n -AD regular boundary (see (2.1)). The fact that the condition (c) above implies (a) from Theorem A is essentially known. See Section 3.1 for more details. So the main novelty lies in the fact that (a) and/or (b) imply (c). Notice that neither the assumptions in Theorem 1.1 nor the conditions (a) and (b) in Theorem A contain the fact the surface measure $\mathcal{H}^n|_{\partial\Omega^+}$ is locally finite. So the conclusion that Ω^+ is chord-arc, and thus with locally finite surface measure may appear rather surprising at first glance.

It is worth comparing the results from Theorems A and 1.1 with previous results obtained by Kenig and the second author of this paper in connection with the one-phase problem for harmonic

measure in chord-arc domains. By combining some of the results from [KT1], [KT2], [KT4], one gets the following.

Theorem B. *Let $\Omega \subset \mathbb{R}^{n+1}$ be a bounded chord-arc domain which is δ -Reifenberg flat, with $\delta > 0$ small enough. Denote by ω the harmonic measure in Ω with pole $p \in \Omega$ and write $\sigma = \mathcal{H}^n|_{\partial\Omega}$. Then the following conditions are equivalent:*

- (i) $\log \frac{d\omega}{d\sigma} \in \text{VMO}(\sigma)$.
- (ii) Ω is vanishing Reifenberg flat and the inner normal N to $\partial\Omega$ belongs to $\text{VMO}(\sigma)$.
- (iii) The inner normal N to $\partial\Omega$ exists σ -a.e. and it belongs to $\text{VMO}(\sigma)$.

Notice the analogies between the conditions (a), (b), (c) from Theorems A and 1.1 with the preceding conditions (i), (ii), (iii). In fact, in view of these results, it is natural to ask if the conditions (a), (b), (c) from Theorems A and 1.1 are equivalent to the fact that the inner normal N to $\partial\Omega^+$ (which exists ω^+ -a.e.) belongs to $\text{VMO}(\omega^+)$ together with the vanishing Reifenberg flatness of Ω_+ . Indeed, remark that the condition (1.2) implies that $N \in \text{VMO}(\omega^+)$. For the moment, the converse implication is an open problem.

To prove the main implication of Theorem 1.1, namely that Ω^+ is a chord-arc domain with $N \in \text{VMO}(\sigma)$ if either (a) or (b) from Theorem A hold, we will use a geometric corona decomposition which will allow us to estimate the surface measure $\mathcal{H}^n|_{\partial\Omega^+}$. We will split some dyadic lattice of “cubes” from $\partial\Omega^+$ into trees, so that, in each tree, $\partial\Omega^+$ is well approximated by a Lipschitz graph at the location and scales of the cubes from the tree. A key step consists in showing that each tree does not have too many stopping cubes (see (3.4) in Lemma 3.3). The main ingredient for this estimate is the condition (b) in Theorem A.

2. PRELIMINARIES

We denote by C or c some constants that may depend on the dimension and perhaps other fixed parameters. Their value may change at different occurrences. On the contrary, constants with subscripts, like C_0 , retain their values. For $a, b \geq 0$, we write $a \lesssim b$ if there is $C > 0$ such that $a \leq Cb$. We write $a \approx b$ to mean $a \lesssim b \lesssim a$. If we want to emphasize that the implicit constant depends on some parameter $\eta \in \mathbb{R}$, we write $a \approx_\eta b$.

2.1. Measures and rectifiability. All measures in this paper are assumed to be Borel measures. A measure μ in \mathbb{R}^d is called *doubling* if there is some constant $C > 0$ such that

$$\mu(B(x, 2r)) \leq C \mu(B(x, r)) \quad \text{for all } x \in \text{supp } \mu.$$

The measure μ is called *n-AD regular* (or *n-Ahlfors-David regular*) if

$$(2.1) \quad C^{-1}r^n \leq \mu(B(x, r)) \leq Cr^n \quad \text{for all } x \in \text{supp } \mu \text{ and } 0 < r \leq \text{diam}(\text{supp } \mu).$$

Note that *n-AD regular* measures are doubling. A set $E \subset \mathbb{R}^d$ is called *n-AD regular* if $\mathcal{H}^n|_E$ is *n-AD regular*. In case that μ satisfies the second inequality in (2.1), but not necessarily the first one, we say that μ has *n-polynomial growth*. We write “*C-doubling*” or “*C-n-AD regular*” if we wish to mention the constant C involved in the definitions.

A set $E \subset \mathbb{R}^d$ is called *n-rectifiable* if there are Lipschitz maps $f_i : \mathbb{R}^n \rightarrow \mathbb{R}^d$ such that

$$\mathcal{H}^n\left(E \setminus \bigcup_i f_i(\mathbb{R}^n)\right) = 0.$$

Analogously, one says that a measure μ is *n-rectifiable* if there are Lipschitz maps $f_i : \mathbb{R}^n \rightarrow \mathbb{R}^d$, $i = 1, 2, \dots$, such that

$$(2.2) \quad \mu\left(\mathbb{R}^d \setminus \bigcup_i f_i(\mathbb{R}^n)\right) = 0,$$

and moreover μ is absolutely continuous with respect to \mathcal{H}^n . An equivalent definition for rectifiability of sets and measures is obtained if we replace Lipschitz images of \mathbb{R}^n by possibly rotated n -dimensional graphs of C^1 functions.

A measure μ in \mathbb{R}^d is called *uniformly n-rectifiable* (UR) if it is n -AD-regular and there exist constants $\theta, M > 0$ such that for all $x \in \text{supp } \mu$ and all $0 < r \leq \text{diam}(\text{supp } \mu)$ there is a Lipschitz mapping g from the ball $B_n(0, r)$ in \mathbb{R}^n to \mathbb{R}^d with $\text{Lip}(g) \leq M$ such that

$$\mu(B(x, r) \cap g(B_n(0, r))) \geq \theta r^n.$$

A set E is called uniformly n -rectifiable if the measure $\mathcal{H}^n|_E$ is uniformly n -rectifiable. The notion of uniform n -rectifiability is a quantitative version of n -rectifiability introduced by David and Semmes (see [DS]). It is easy to check that uniform n -rectifiability implies n -rectifiability.

Given two doubling measures μ, ν in \mathbb{R}^d with the same support, we write $\nu \in A_\infty(\mu)$ if there are constants $\delta, \varepsilon \in (0, 1)$ such that, for any ball B centered in $\text{supp } \mu$ and any Borel set $E \subset B$, the following holds:

$$\mu(E) \geq \delta \mu(B) \quad \Rightarrow \quad \nu(E) \geq \varepsilon \nu(B).$$

2.2. Harmonic measure. For a bounded domain $\Omega \subset \mathbb{R}^{n+1}$ and $x \in \Omega$, one can construct the harmonic measure ω_Ω^x (see [He, p. 217], for example). In fact, for any continuous function f , the Perron solution for the boundary function f is given by

$$H_f(x) = \int_{\partial\Omega} f(y) d\omega_\Omega^x(y),$$

Remark that constant functions are continuous and since $H_1(x) = 1$, for any $x \in \Omega$, we have that $\omega_\Omega^x(\partial\Omega) = 1$, for any $x \in \Omega$.

Let \mathcal{E} denote the fundamental solution for the Laplace equation in \mathbb{R}^{n+1} , for $n \geq 2$, so that $\mathcal{E}(x) = c_n |x|^{1-n}$ for $n \geq 2$, $c_n > 0$. The Newtonian potential of a measure $\mu \in M_+(\mathbb{R}^{n+1})$ is defined by

$$U\mu(x) = \mathcal{E} * \mu(x),$$

and the Newtonian capacity of a compact set $F \subset \mathbb{R}^{n+1}$ equals

$$\text{Cap}(F) = \sup \left\{ \mu(F) : \mu \in M_+(\mathbb{R}^{n+1}), \text{supp } \mu \subset F, \|U\mu\|_\infty \leq 1 \right\},$$

where $M_+(\mathbb{R}^{n+1})$ is the space of finite Borel measures from \mathbb{R}^{n+1} .

The following result is standard. See [To1, Lemma 2.1] for example.

Lemma 2.1. *Given $n \geq 2$, let $\Omega \subset \mathbb{R}^{n+1}$ be open and let B be a closed ball centered at $\partial\Omega$. Then*

$$\omega_\Omega^x(B) \geq c(n) \frac{\text{Cap}(\frac{1}{4}B \setminus \Omega)}{r(B)^{n-1}} \quad \text{for all } x \in \frac{1}{4}B \cap \Omega,$$

with $c(n) > 0$.

A similar result holds in the case $n = 1$, involving logarithmic capacity instead of Newtonian capacity.

2.3. NTA and Reifenberg flat domains. Given $\Omega \subset \mathbb{R}^{n+1}$ and $C \geq 2$, we say that Ω satisfies the *C-Harnack chain condition* if for every $\rho > 0$, $k \geq 1$, and every pair of points $x, y \in \Omega$ with $\text{dist}(x, \partial\Omega), \text{dist}(y, \partial\Omega) \geq \rho$ and $|x - y| < 2^k \rho$, there is a chain of open balls $B_1, \dots, B_m \subset \Omega$, $m \leq Ck$, with $x \in B_1, y \in B_m, B_k \cap B_{k+1} \neq \emptyset$ and $C^{-1} \text{diam}(B_k) \leq \text{dist}(B_k, \partial\Omega) \leq C \text{diam}(B_k)$. The chain of balls is called a *Harnack chain*. Note that if such a chain exists, then any positive harmonic function $u : \Omega \rightarrow \mathbb{R}$ satisfies

$$u(x) \approx u(y),$$

with the implicit constant depending on m and n . We say that Ω is a *C-corkscrew domain* if for every $\xi \in \partial\Omega$ and $r \in (0, \text{diam}(\partial\Omega))$ there is a ball of radius r/C contained in $B(\xi, r) \cap \Omega$. If $B(x, r/C) \subset B(\xi, r) \cap \Omega$, we call x a *corkscrew point* for the ball $B(\xi, r)$. Finally, we say that Ω is *C-non-tangentially accessible* (or *C-NTA*, or just *NTA*) if it satisfies the *C-Harnack chain condition* and both Ω and $\mathbb{R}^{n+1} \setminus \overline{\Omega}$ are *C-corkscrew domains*. We say that Ω is *C-chord-arc* if, additionally, $\partial\Omega$ is *C-n-AD-regular*. Also, Ω is *two-sided C-NTA* if both Ω and $\Omega_{\text{ext}} := (\overline{\Omega})^c$ are *C-NTA*.

NTA domains were introduced by Jerison and Kenig in [JK]. In that work, the behavior of harmonic measure in this type of domains was studied in detail. Among other results, the authors showed that harmonic measure is doubling in NTA domains, and its support coincides with the whole boundary. They also proved that harmonic measure in these domains satisfies the following important change of pole formula.

Theorem 2.2. *Let $n \geq 1$, Ω be a C-NTA domain in \mathbb{R}^{n+1} and let B be a ball centered in $\partial\Omega$. Let $p_1, p_2 \in \Omega$ be such that $\text{dist}(p_i, B \cap \partial\Omega) \geq c_0^{-1} r(B)$ for $i = 1, 2$. Then, for any Borel set $E \subset B \cap \partial\Omega$,*

$$(2.3) \quad \frac{\omega^{p_1}(E)}{\omega^{p_1}(B)} \approx \frac{\omega^{p_2}(E)}{\omega^{p_2}(B)},$$

with the implicit constant depending only on n, c_0 and C .

For the proof, see Lemma 4.11 from [JK]. See also [MT] for an extension of this result to the so-called uniform domains.

Given two NTA domains $\Omega^+ \subset \mathbb{R}^{n+1}$ and $\Omega^- = \mathbb{R}^{n+1} \setminus \overline{\Omega^+}$, we say that Ω^+ and Ω^- have joint big pieces of chord-arc subdomains if for any ball B centered in $\partial\Omega^+$ with radius at most $\text{diam}(\partial\Omega^+)$ there are two *C-chord-arc domains* $\Omega_B^s \subset \Omega^s$, with $s = +, -$, such that $\mathcal{H}^n(\partial\Omega_B^1 \cap \partial\Omega_B^2 \cap B) \gtrsim r(B)^n$, uniformly on B .

Given a set $E \subset \mathbb{R}^{n+1}$, $x \in \mathbb{R}^{n+1}$, $r > 0$, and P an n -plane, we set

$$(2.4) \quad D_E(x, r, P) = r^{-1} \max \left\{ \sup_{y \in E \cap B(x, r)} \text{dist}(y, P), \sup_{y \in P \cap B(x, r)} \text{dist}(y, E) \right\}.$$

We also define

$$(2.5) \quad D_E(x, r) = \inf_P D_E(x, r, P),$$

where the infimum is over all n -planes P . For a given ball $B = B(x, r)$, we will also write $D_E(B)$ instead of $D_E(x, r)$. Given $\delta, R > 0$, set E is (δ, R) -*Reifenberg flat* (or just δ -*Reifenberg flat*) if $D_E(x, r) < \delta$ for all $x \in E$ and $0 < r \leq R$, and it is *vanishing Reifenberg flat* if

$$\limsup_{r \rightarrow 0} \sup_{x \in E} D_E(x, r) = 0.$$

Let $\Omega \subset \mathbb{R}^{n+1}$ be an open set, and let $0 < \delta < 1/2$. We say that Ω is a (δ, R) -Reifenberg flat domain (or just δ -Reifenberg flat) if it satisfies the following conditions:

- (a) $\partial\Omega$ is (δ, R) -Reifenberg flat.
- (b) For every $x \in \partial\Omega$ and $0 < r \leq R$, denote by $P(x, r)$ an n -plane that minimizes $D_E(x, r)$. Then one of the connected components of

$$B(x, r) \cap \{x \in \mathbb{R}^{n+1} : \text{dist}(x, P(x, r)) \geq 2\delta r\}$$

is contained in Ω and the other is contained in $\mathbb{R}^{n+1} \setminus \Omega$.

If, additionally, $\partial\Omega$ is vanishing Reifenberg flat, then Ω is said to be vanishing Reifenberg flat, too. It is well known that if Ω is a δ -Reifenberg flat domain, with δ small enough, then it is also an NTA domain (see [KT1]).

2.4. The space VMO. Given a Radon measure μ in \mathbb{R}^{n+1} , $f \in L^1_{loc}(\mu)$, and $A \subset \mathbb{R}^{n+1}$, we write

$$m_{\mu, A}(f) = \int_A f d\mu = \frac{1}{\mu(A)} \int_A f d\mu.$$

Assume μ to be doubling. We say that $f \in \text{VMO}(\mu)$ if

$$(2.6) \quad \lim_{r \rightarrow 0} \sup_{x \in \text{supp } \mu} \int_{B(x, r)} |f - m_{\mu, B(x, r)} f| d\mu = 0.$$

It is well known that the space $\text{VMO}(\mu)$ coincides with the closure of the set of bounded uniformly continuous functions on $\text{supp } \mu$ in the BMO norm.

2.5. Dyadic lattices and densities. In [Ch] Michael Christ introduced a dyadic lattice for a doubling metric spaces. We state below a version of this result from Hytönen and Kareima [HK], in the particular setting of $\partial\Omega^+$, which has the advantage that $\partial\Omega^+$ is covered by the family of cubes of any given generation and avoids the problem of the possible existence of uncovered sets of zero measure, as in [Ch]. The construction of David and Mattila in [DM], valid for non-doubling measure in \mathbb{R}^{n+1} , would also work in our context.

Theorem 2.3. *Let Ω^+ be as in Theorem 1.1. There exist a family \mathcal{D} of Borel subsets of $\partial\Omega^+$ and constants $0 < r_0 < 1$, $0 < a_1, \eta, C_1, C_2 < \infty$ such that $\mathcal{D} = \bigcup_{k \in \mathbb{Z}} \mathcal{D}_k$ with $\mathcal{D}_k = \{Q^i\}_{i \in I_k}$, and the following holds:*

- (a) For every $k \in \mathbb{Z}$ we have $\partial\Omega^+ = \bigcup_{Q \in \mathcal{D}_k} Q$.
- (b) For every $k_0 \leq k_1$ and $Q_j \in \mathcal{D}_{k_j}$ for $j \in \{0, 1\}$, then either $Q_1 \subset Q_0$ or $Q_1 \cap Q_0 = \emptyset$.
- (c) For each $Q_1 \in \mathcal{D}_{k_1}$ and each $k_0 < k_1$ there exists a unique cube $Q_0 \in \mathcal{D}_{k_0}$ such that $Q_1 \subset Q_0$.
- (d) For $Q \in \mathcal{D}_k$ there are $z_Q \in Q$ and balls $\tilde{B}_Q = B(z_Q, a_1 r_0^k)$ and $B_Q = B(z_Q, C_1 r_0^k)$ such that $\tilde{B}_Q \cap \partial\Omega^+ \subset Q \subset B_Q$.

We say that $Q \in \mathcal{D}_k$ is a *dyadic cube* of generation k , and write $\ell(Q) := 2C_1 r_0^k$. We call $\ell(Q)$ the side length of Q . The point z_Q in (d) is called center of Q . For $R \in \mathcal{D}$, we denote by $\mathcal{D}(R)$ the family of the cubes $Q \in \mathcal{D}$ which are contained in R .

We say that two cubes $Q, S \in \mathcal{D}_k$ (of the same generation) are *neighbors*, writing $S \in \mathcal{N}(Q)$, if $2B_Q \cap 2B_S \neq \emptyset$. We write $\mathcal{ND}_k := \left\{ \bigcup_{S \in \mathcal{N}(Q)} S \right\}_{Q \in \mathcal{D}_k}$ and $\mathcal{ND} := \bigcup_k \mathcal{ND}_k$. We say that $P \in \mathcal{ND}_k$ is an extended cube of generation k , and write $\ell(P) := 2C_1 r_0^k$.

Many of our arguments on cubes will need to be applied both to the dyadic and the extended cubes. We write $\widehat{\mathcal{D}}_k = \mathcal{D}_k \cup \mathcal{N}\mathcal{D}_k$ and $\widehat{\mathcal{D}} = \mathcal{D} \cup \mathcal{N}\mathcal{D}$. We refer as *cubes* to both the dyadic and the extended cubes.

We need also to introduce “dilations” of cubes. Given $Q \in \widehat{\mathcal{D}}_k$ and $\Lambda > 1$, we write

$$\Lambda Q = \{x \in \partial\Omega^+ : \text{dist}(x, Q) < (\Lambda - 1)\ell(Q)\}.$$

and $\ell(\Lambda Q) := 2C_1(\Lambda r_0)^k$. Obviously, we also define $1Q \equiv Q$.

Let μ be a Radon measure in \mathbb{R}^{n+1} . Given a ball $B \subset \mathbb{R}^{n+1}$, we denote

$$(2.7) \quad \Theta_\mu(B) = \frac{\mu(B)}{r(B)^n}.$$

So $\Theta_\mu(B)$ is the n -dimensional density of μ on B .

3. PROOF OF THEOREM 1.1

3.1. Proof of (c) \Rightarrow (a). The fact that Ω^+ is a two-sided chord-arc domain ensures that $\omega^\pm \in A_\infty(\sigma)$. Also, since $N \in \text{VMO}(\sigma)$, from (iii) \Rightarrow (i) in Theorem B, we infer that $\log \frac{d\omega^\pm}{d\sigma} \in \text{VMO}(\sigma)$. For any ball B centered in $\partial\Omega^+$, writing

$$\log \frac{d\omega^-}{d\omega^+} = \log \frac{d\omega^-}{d\sigma} - \log \frac{d\omega^+}{d\sigma},$$

and denoting $m_B^\pm = \int_B \log \frac{d\omega^\pm}{d\sigma} d\sigma$, we get

$$\int_B \left| \log \frac{d\omega^-}{d\omega^+} - (m_B^- - m_B^+) \right| d\omega^+ \leq \int_B \left| \log \frac{d\omega^-}{d\sigma} - m_B^- \right| d\omega^+ + \int_B \left| \log \frac{d\omega^+}{d\sigma} - m_B^+ \right| d\omega^+.$$

Since $\omega^+ \in A_\infty(\sigma)$, it follows that $\frac{d\omega^+}{d\sigma} \in B_p(\sigma)$ for some $p \in (1, \infty)$ (i.e., it satisfies a reverse Hölder inequality with exponent p). Then we write

$$\begin{aligned} \int_B \left| \log \frac{d\omega^\pm}{d\sigma} - m_B^\pm \right| d\omega^+ &= \frac{\sigma(B)}{\omega^+(B)} \int_B \left| \log \frac{d\omega^\pm}{d\sigma} - m_B^\pm \right| \frac{d\omega^+}{d\sigma} d\sigma \\ &\leq \frac{\sigma(B)}{\omega^+(B)} \left(\int_B \left| \log \frac{d\omega^\pm}{d\sigma} - m_B^\pm \right|^{p'} d\sigma \right)^{1/p'} \left(\int_B \left(\frac{d\omega^+}{d\sigma} \right)^p d\sigma \right)^{1/p} \\ &\lesssim \left(\int_B \left| \log \frac{d\omega^\pm}{d\sigma} - m_B^\pm \right|^{p'} d\sigma \right)^{1/p'}. \end{aligned}$$

By the John-Nirenberg inequality and the fact that $\log \frac{d\omega^\pm}{d\sigma} \in \text{VMO}(\sigma)$, it follows that the right hand side above tends to 0 as $r(B) \rightarrow 0$, uniformly with respect to the center of the ball. Therefore,

$$\int_B \left| \log \frac{d\omega^-}{d\omega^+} - \int_B \log \frac{d\omega^-}{d\omega^+} d\omega^+ \right| d\omega^+ \lesssim \int_B \left| \log \frac{d\omega^-}{d\omega^+} - (m_B^- - m_B^+) \right| d\omega^+ \rightarrow 0 \quad \text{as } r(B) \rightarrow 0,$$

uniformly with respect to the center of the ball. So $\log \frac{d\omega^-}{d\omega^+} \in \text{VMO}(\omega^+)$, as wished.

3.2. Stopping cubes for the proof of (a) \Rightarrow (c). The rest of the paper is devoted to the proof of this implication.

As shown in Corollary 1.2 from [PT], the condition (a) together with the n -AD-regularity of $\partial\Omega^+$ imply that $N \in \text{VMO}(\sigma)$. Hence, it suffices to prove that $\partial\Omega^+$ is n -AD-regular. To this end, we have to show that, for any $x \in \partial\Omega^+$ and any $r > 0$,

$$\mathcal{H}^n(\partial\Omega^+ \cap B(x, r)) \leq C r^n.$$

Remark that the converse estimate (i.e., the lower n -AD-regularity of $\partial\Omega^+$) is an easy consequence of the two-sided corkscrew condition for Ω^+ . We consider the lattice \mathcal{D} from Theorem 2.3. Then, clearly, the preceding condition is equivalent to

$$\mathcal{H}^n(R_0) \leq C \ell(R_0)^n \quad \text{for all } R_0 \in \mathcal{D}.$$

Since (a) \Rightarrow (b), clearly we may assume that Ω^+ is vanishing Reifenberg flat, that Ω^+ and Ω^- have joint big pieces of chord-arc subdomains, and that (1.2) holds.

We consider a fixed cube $R_0 \in \mathcal{D}$. For a small constant $\varepsilon_0 \in (0, 1)$, we assume $\ell(R_0)$ small enough so that

$$(3.1) \quad D_{\partial\Omega^+}(x, r) \leq \varepsilon_0 \quad \text{for all } x \in \partial\Omega^+ \text{ and } 0 < r \leq 100\ell(R_0).$$

Given some parameters $\delta, \alpha \in (0, 1)$ and $M \gg 1$ to be chosen below, we consider some stopping cubes defined as follows: we say that $Q \in \text{Stop}(R_0)$ if $Q \in \mathcal{D}$ is a maximal cube contained in R_0 such that one of the following options holds:

- $\Theta_{\omega^+}(B_Q) > M\Theta_{\omega^+}(B_{R_0})$. We write $Q \in \mathcal{HD}^+(R_0)$.
- $\Theta_{\omega^+}(B_Q) \leq \delta\Theta_{\omega^+}(B_{R_0})$. We write $Q \in \mathcal{LD}^+(R_0)$.
- $|N_{B_{R_0}} - N_{B_Q}| \geq \alpha$. We write $Q \in \mathcal{BS}(R_0)$.

Recall that the vectors $N_{B_{R_0}}$ and N_{B_Q} are defined in (1.3). The notations \mathcal{HD} , \mathcal{LD} and \mathcal{BS} stand for high density, low density, and big slope, respectively.

We denote

$$\text{HD}^+(R_0) = \bigcup_{Q \in \mathcal{HD}^+(R_0)} Q, \quad \text{LD}^+(R_0) = \bigcup_{Q \in \mathcal{LD}^+(R_0)} Q, \quad \text{BS}(R_0) = \bigcup_{Q \in \mathcal{BS}(R_0)} Q$$

and

$$G(R_0) = R_0 \setminus \bigcup_{Q \in \text{Stop}(R_0)} Q.$$

We let $\text{Tree}(R_0)$ be the subfamily of $\mathcal{D}(R_0)$ such that no cube from $\text{Tree}(R_0)$ is properly contained in any cube from $\text{Stop}(R_0)$.

From now on, we also write $N_Q \equiv N_{B_Q}$ and $L_Q = L_{B_Q}$, with L_{B_Q} defined in (1.3).

Lemma 3.1. *For any $\varepsilon > 0$, if $\ell(R_0)$ is small enough, M big enough, and δ small enough, then*

$$\omega^+(\text{HD}^+(R_0) \cup \text{LD}^+(R_0)) \leq \varepsilon \omega^+(R_0).$$

The proof of this lemma is the same as the one of Lemma 4.1 from [PT]. We also have:

Lemma 3.2. *For any $\varepsilon > 0$, if $\ell(R_0)$ is small enough (depending on ε and α), then*

$$\omega^+(\text{BS}(R_0)) \leq \varepsilon \omega^+(R_0).$$

Proof. By (1.2), assuming $\ell(R_0)$ small enough,

$$\int_{R_0} |N - N_{R_0}| d\omega^+ \leq \varepsilon^2,$$

and, analogously,

$$\int_Q |N - N_Q| d\omega^+ \leq \varepsilon^2 \quad \text{for all } Q \in \mathcal{BS}(R_0).$$

Therefore,

$$\begin{aligned} \sum_{Q \in \mathcal{BS}(R_0)} |N_Q - N_{R_0}| \omega^+(Q) &\leq \sum_{Q \in \mathcal{BS}(R_0)} \int_Q |N - N_{R_0}| d\omega^+ + \sum_{Q \in \mathcal{BS}(R_0)} \int_Q |N - N_Q| d\omega^+ \\ &\leq \varepsilon^2 \omega^+(R_0) + \sum_{Q \in \mathcal{BS}(R_0)} \varepsilon^2 \omega^+(Q) \leq 2\varepsilon^2 \omega^+(R_0). \end{aligned}$$

Hence,

$$\omega^+(\mathcal{BS}(R_0)) \lesssim \sum_{Q \in \mathcal{BS}(R_0)} \frac{|N_Q - N_{R_0}|}{\alpha} \omega^+(Q) \lesssim \frac{\varepsilon^2}{\alpha} \omega^+(R_0),$$

and so the lemma follows, since we can assume that $\varepsilon \leq c\alpha$. \square

From the last two lemmas we infer that, for $\ell(R_0)$ small enough,

$$(3.2) \quad \omega^+\left(\bigcup_{Q \in \text{Stop}(R_0)} Q\right) \leq 2\varepsilon \omega^+(R_0).$$

Our next objective is to prove the following related estimates in the lemma below.

Lemma 3.3. *Suppose that $\ell(R_0)$ is small enough and $\varepsilon_0, \varepsilon, \delta, \alpha$, and M are chosen suitably. Then,*

$$(3.3) \quad \mathcal{H}^n(G(R_0)) \lesssim \ell(R_0)^n$$

and

$$(3.4) \quad \sum_{Q \in \text{Stop}(R_0)} \ell(Q)^n \leq \frac{1}{10} \ell(R_0)^n.$$

Before proving this result we will show how this yields Theorem 1.1.

3.3. Proof of Theorem 1.1 using Lemma 3.3. We now construct the family of cubes $\text{Top}(R_0)$ contained in R_0 inductively. First we write $\text{Top}_0(R_0) = \{R_0\}$, and then assuming $\text{Top}_{k-1}(R_0)$ to be defined, we set

$$\text{Top}_k(R_0) = \bigcup_{R \in \text{Top}_{k-1}(R_0)} \text{Stop}(R),$$

and we let $\text{Top}(R_0) = \bigcup_{k \geq 0} \text{Top}_k(R_0)$. We also set

$$G_k(R_0) = \bigcup_{R \in \text{Top}_{k-1}(R_0)} \left(R \setminus \bigcup_{Q \in \text{Stop}(R)} Q \right) = \bigcup_{R \in \text{Top}_{k-1}(R_0)} R \setminus \bigcup_{Q \in \text{Top}_k(R_0)} Q.$$

So we have the partition

$$R_0 = \bigcup_{k=1}^{\infty} G_k(R_0) \cup \bigcap_{k=1}^{\infty} \bigcup_{Q \in \text{Top}_k(R_0)} Q,$$

and thus

$$(3.5) \quad \mathcal{H}^n(R_0) \leq \sum_{k=1}^{\infty} \mathcal{H}^n(G_k(R_0)) + \mathcal{H}^n\left(\bigcap_{k=1}^{\infty} \bigcup_{Q \in \text{Top}_k(R_0)} Q\right).$$

From Lemma 3.3 (applied to the cubes Q below in place of R_0), we deduce that, for every k ,

$$(3.6) \quad \mathcal{H}^n(G_k(R_0)) \leq \sum_{Q \in \text{Top}_{k-1}(R_0)} \mathcal{H}^n(G(Q)) \lesssim \sum_{Q \in \text{Top}_{k-1}(R_0)} \ell(Q)^n$$

and that

$$\sum_{Q \in \text{Top}_k(R_0)} \ell(Q)^n = \sum_{R \in \text{Top}_{k-1}(R)} \sum_{Q \in \text{Stop}(R)} \ell(Q)^n \leq \frac{1}{10} \sum_{R \in \text{Top}_{k-1}(R_0)} \ell(R)^n.$$

Thus,

$$(3.7) \quad \sum_{Q \in \text{Top}_k(R_0)} \ell(Q)^n \leq 10^{-k} \ell(R_0)^n,$$

which, together with (3.6), implies that

$$(3.8) \quad \mathcal{H}^n(G_k(R_0)) \lesssim 10^{-k} \ell(R_0)^n.$$

On the other hand, taking into account that there exists some $r_0 \in (0, 1)$ such that every $Q \in \text{Top}_k(R_0)$ satisfies $\ell(Q) \leq r_0^k \ell(R_0)$, from the definition of \mathcal{H}^n we deduce that

$$(3.9) \quad \mathcal{H}^n\left(\bigcap_{k=1}^{\infty} \bigcup_{Q \in \text{Top}_k(R_0)} Q\right) \lesssim \limsup_{k \rightarrow \infty} \sum_{Q \in \text{Top}_k(R_0)} \ell(Q)^n \leq \limsup_{k \rightarrow \infty} 10^{-k} \ell(R_0)^n = 0.$$

Hence, by (3.5), (3.8), and (3.9), we obtain

$$\mathcal{H}^n(R_0) \lesssim \sum_{k=1}^{\infty} 10^{-k} \ell(R_0)^n \lesssim \ell(R_0)^n.$$

□

4. THE PROOF OF LEMMA 3.3

4.1. Preliminaries for the construction of the approximating Lipschitz graph. For technical reasons we have to work with (extended) cubes from $\mathcal{N}\mathcal{D}$. Given $R_0 \in \mathcal{D}$, we consider the cube $S_0 \in \mathcal{N}\mathcal{D}$ defined by

$$S_0 = \bigcup_{R \in \mathcal{N}(R_0)} R,$$

We denote

$$\text{Stop}(S_0) = \bigcup_{R \in \mathcal{N}(R_0)} \text{Stop}(R), \quad \text{Tree}(S_0) = \bigcup_{R \in \mathcal{N}(R_0)} \text{Tree}(R), \quad G(S_0) = \bigcup_{R \in \mathcal{N}(R_0)} G(R),$$

and we also set

$$z_{S_0} = z_{R_0}, \quad \ell(S_0) = \ell(R_0), \quad L_{S_0} = L_{R_0}, \quad B_{S_0} = 5B_{R_0}.$$

Recall that z_{R_0} stands for the center of R_0 . So z_{S_0} should also be considered as the center of S_0 .

To prove Lemma 3.3 we need first to construct a Lipschitz graph that will approximate well S_0 at the scale of $\text{Stop}(S_0)$ and will contain $G(S_0)$. The arguments involved are very standard and there are many presentations of this. We will follow [To2]. First we need to state some preliminary lemmas.

Lemma 4.1. *Let $P, Q \in \text{Tree}(S_0) \setminus \text{Stop}(S_0)$ be so that $P \subset Q$. If $x \in L_P \cap B_P$, then*

$$\text{dist}(x, L_Q) \lesssim \varepsilon_0 \ell(Q).$$

The proof of this result is standard. See Lemma 7.14 in [To2] for a similar estimate. Recall that the constant ε_0 appears in (3.1), and the parameter δ is the one in the definition of the family \mathcal{LD}^+ . Other relevant parameters are M and α , in the definition of the families \mathcal{HD}^+ and \mathcal{BS} , respectively.

We denote by Π (respectively Π^\perp) the orthogonal projection onto L_{S_0} (respectively, $L_{S_0}^\perp$). Further, we assume that $L_{S_0} = \mathbb{R}^n \times \{0\}$, so that $L_{S_0}^\perp$ is the vertical axis of \mathbb{R}^{n+1} , which we will identify with \mathbb{R} . For any given n -plane L , we denote by Π_L the orthogonal projection on L .

The next lemma is analogous to Lemma 7.15 in [To2].

Lemma 4.2. *Let $Q, P \in \text{Tree}(S_0) \setminus \text{Stop}(S_0)$, $q \in L_Q \cap 2B_Q$ and $p \in L_P \cap 2B_P$. Then*

$$|\Pi^\perp(p) - \Pi^\perp(q)| \leq C\alpha (|\Pi(p) - \Pi(q)| + 2\ell(P) + 2\ell(Q)).$$

Next we need to define the following auxiliary function:

$$(4.1) \quad d(x) := \inf_{Q \in \text{Tree}(S_0) \setminus \text{Stop}(S_0)} (\text{dist}(x, Q) + \ell(Q)) \quad \text{for } x \in \mathbb{R}^{n+1}.$$

The next lemma can be found as Lemma 7.19 of [To2].

Lemma 4.3. *Let $\varepsilon_0 > 0$ and $\alpha > 0$ be sufficiently small. Then for any $x, y \in \mathbb{R}^{n+1}$,*

$$\left| \Pi^\perp(x) - \Pi^\perp(y) \right| \lesssim \alpha |\Pi(x) - \Pi(y)| + d(x) + d(y).$$

We denote

$$Z_0 = \{x \in \mathbb{R}^2 : d(x) = 0\}.$$

It is easy to check that $G(S_0) \subset Z_0$. Also, as in Lemma 7.18 from [To2], we have:

Lemma 4.4. *We have $\mathcal{H}^n(Z_0) < \infty$ and*

$$\omega^+|_{Z_0} = \rho_{Z_0} \mathcal{H}^n|_{Z_0},$$

where ρ_{Z_0} is a function satisfying $C^1\delta \leq \rho_{Z_0} \leq CA$.

Let $\mathcal{D}_{L_{S_0}}$ be the set of dyadic n -dimensional cubes in L_{S_0} . For $p \in L_{S_0}$, we denote

$$D(p) = \inf_{x \in \Pi^{-1}(p)} d(x),$$

and for $I \in \mathcal{D}_{L_{S_0}}$,

$$D(I) := \inf_{p \in I} D(p).$$

Set

$$W := \{I \text{ maximal in } \mathcal{D}_{L_{S_0}} : \ell(I) < 20^{-1} D(I)\}.$$

We summarize the properties of the cubes in W in the following lemma. We index W as $\{R_i\}_{i \in I_W}$.

Lemma 4.5. *The cubes R_i , $i \in I_W$, have disjoint interiors in L_{S_0} and satisfy the following properties:*

(1) *If $x \in 15R_i$, then $5\ell(R_i) \leq D(x) \leq 50\ell(R_i)$.*

(2) *There exists an absolute constant $c > 1$ such that if $15R_i \cap 15R_j \neq \emptyset$, then*

$$(4.2) \quad c^{-1}\ell(R_i) \leq \ell(R_j) \leq c\ell(R_i).$$

(3) *For each $i \in I_W$, there are at most N_0 cubes R_j such that $15R_i \cap 15R_j \neq \emptyset$, where N_0 is some absolute constant depending only on n .*

(4) $L_{S_0} \setminus \Pi(Z_0) = \bigcup_{i \in I_W} R_i$.

The proof of this result is standard. See for example Lemma 7.20 in [To2].

Let

$$(4.3) \quad B_0 = B(\Pi(z_{S_0}), 10r(B_{S_0})).$$

Certainly, $\text{dist}(z_{S_0}, L_{S_0}) < 10^{-1}\ell(S_0)$, and so we have that

$$(4.4) \quad \Pi(S_0) \subset \Pi(B_0) \subset 2B_0 \cap L_{S_0}.$$

Now set

$$I_0 := \{i \in I_W : R_i \cap B_0 \neq \emptyset\}.$$

The next lemma is proven exactly as Lemma 7.21 from [To2].

Lemma 4.6. *The following holds.*

- *If $i \in I_0$, then $\ell(R_i) \lesssim \ell(S_0)$ and $3R_i \subset L_{S_0} \cap 1.2B_0$.*
- *If $i \notin I_0$, then*

$$\ell(R_i) \approx \text{dist}(\Pi(z_{S_0}), R_i) \approx |z_{S_0} - x| \gtrsim \ell(S_0) \text{ for all } x \in R_i.$$

The next lemma is analogous to Lemma 7.22 from [To2].

Lemma 4.7. *Let $i \in I_0$; there exists a cube $Q = Q_i \in \text{Tree}(S_0) \setminus \text{Stop}(S_0)$ such that*

$$(4.5) \quad \ell(R_i) \lesssim \ell(Q_i) \lesssim \ell(R_i);$$

$$(4.6) \quad \text{dist}(R_i, \Pi(Q_i)) \lesssim \ell(R_i).$$

4.2. The approximating Lipschitz graph. Observe that, by Lemma 4.3, for every $x, y \in Z_0$ satisfy

$$\left| \Pi^\perp(x) - \Pi^\perp(y) \right| \lesssim \alpha |\Pi(x) - \Pi(y)|.$$

Therefore, the map $\Pi : Z_0 \rightarrow L_{S_0}$ is injective, and we can define a Lipschitz function A on $\Pi(Z_0)$ by requiring

$$(4.7) \quad A(\Pi(x)) := \Pi^\perp(x) \quad \text{for } x \in Z_0.$$

Next we intend to extend the function A to the whole L_{S_0} . To this end, for $i \in I_0$, first we denote by A_i the affine function $L_{S_0} \rightarrow L_{S_0}^\perp$ whose graph is the n -plane L_{Q_i} . Notice that, for each $i \in I_0$, A_i is Lipschitz with constant $\lesssim \alpha$.

On the other hand, for $i \in I_W \setminus I_0$, we put $A_i \equiv 0$, so that its graph is just L_{S_0} .

The following lemma is proven in the same way as Lemma 7.23 in [To2].

Lemma 4.8. *Assume that $\varepsilon_0 \ll \alpha$. If $10R_i \cap 10R_j \neq \emptyset$ for some $i, j \in I_W$, then*

- (1) $\text{dist}(Q_i, Q_j) \lesssim \ell(R_i)$ if, moreover, $i, j \in I_0$;
- (2) $|A_i(x) - A_j(x)| \lesssim \varepsilon_0 \ell(R_i)$ for $x \in 100R_i$;
- (3) $|\nabla A_i(x) - \nabla A_j(x)| \lesssim \varepsilon_0$.

Remark that [To2, Lemma 7.23] yields estimates like (2) and (3) above with the implicit constants depending on M and δ . Although this dependence would be harmless for our purposes, in our situation the implicit constants above depend neither on M nor on δ due to the Reifenberg flatness of $\partial\Omega^+$.

Next we complete the task of defining A on L_{S_0} . To do so, we recur to a standard construction involving a partition of unity adapted to the Whitney decomposition $\{R_i\}_{i \in I_W}$; this construction goes as follows: for $i \in I_W$, we find a function $\tilde{\phi}_i \in C^\infty(L_{S_0})$ such that

$$(4.8) \quad \chi_{2R_i} \leq \tilde{\phi} \leq \chi_{3R_i},$$

$$(4.9) \quad \|\nabla \tilde{\phi}_i\|_\infty \lesssim \frac{1}{\ell(R_i)},$$

$$(4.10) \quad \|\nabla^2 \tilde{\phi}_i\|_\infty \lesssim \frac{1}{\ell(R_i)^2}.$$

Then for each $i \in I_W$, we put

$$(4.11) \quad \phi_i := \frac{\tilde{\phi}_i}{\sum_{j \in I_W} \tilde{\phi}_j}.$$

Then it is immediate from the construction that $\{\phi_i\}_{i \in I_W}$ is a partition of unity in $L_{S_0} \setminus \Pi(Z_0)$ subordinated to $\{3R_i\}_{i \in I_W}$. Moreover, properties (4.9) and (4.10) together with Lemma 4.5 give that

$$\begin{aligned} \|\nabla \phi_i\|_\infty &\lesssim \ell(R_i)^{-1} \quad \text{and} \\ \|\nabla^2 \phi_i\|_\infty &\lesssim \ell(R_i)^{-2}. \end{aligned}$$

We now define $A : L_{S_0} \setminus \Pi(Z_0) \rightarrow L_{S_0}^\perp$: if $x \in L_{S_0}$, we put

$$A(x) := \sum_{i \in I_W} \phi_i(x) A_i(x) = \sum_{i \in I_0} \phi_i(x) A_i(x).$$

Recall that A is defined in Z_0 by (4.7).

The following lemma, which follows in the same way as Lemmas 7.24 and 7.25 from [To2], will also be useful later on.

Lemma 4.9. *The function $A : L_{S_0} \rightarrow L_{S_0}^\perp$ is supported on $1.2B_0$. It is Lipschitz with slope at most $C\alpha$. Moreover, if $x \in 15R_i$ for $i \in I_W$, then*

$$|\nabla^2 A(x)| \lesssim \frac{\varepsilon_0}{\ell(R_i)}.$$

We will denote the graph of A by Γ , that is

$$(4.12) \quad \Gamma := \{(x, A(x)) \mid x \in L_{S_0}\}.$$

Also, we assume α small enough so that A is Lipschitz with constant at most $1/100$.

Observe now that the estimate (3.3) in Lemma 3.3 is an immediate consequence of the preceding lemma. Indeed,

$$G(R_0) \subset G(S_0) \subset Z_0 \cap B_{S_0} \subset \Gamma \cap B_{S_0},$$

and thus, using that Γ is a Lipschitz graph with slope at most $C\alpha \leq 1$, we get

$$(4.13) \quad \mathcal{H}^n(G(R_0)) \leq \mathcal{H}^n(\Gamma \cap B_{S_0}) \lesssim r(B_{S_0})^n \approx \ell(R_0)^n,$$

which proves (3.3). Further, it is easy to check that $G(R_0)$ (and so $G(S_0)$) is non-empty. Indeed, this is an immediate consequence of (3.2):

$$\omega^+(G(R_0)) = \omega^+(R_0) - \omega^+\left(\bigcup_{Q \in \text{Stop}(R_0)} Q\right) \geq (1 - 2\varepsilon)\omega^+(R_0) > 0.$$

4.3. The Lipschitz graph Γ and $\partial\Omega^+$ are close to each other. The next four lemmas are proven as Lemmas 7.28, 7.29, 7.30, and 7.32 in [To2], although in our case the implicit constants in the estimates below do not depend on M or δ due to the Reifenberg flatness of $\partial\Omega^+$. Anyway, analogously to Lemma 4.8, such dependence would be harmless.

Lemma 4.10. *Every any $x \in 3B_{S_0}$ satisfies*

$$\text{dist}(x, \Gamma) \lesssim d(x).$$

Lemma 4.11. *Let $\varepsilon_0 > 0$ be sufficiently small. If $Q \in \text{Tree}(S_0) \setminus \text{Stop}(S_0)$ and $x \in \Gamma \cap 2B_Q$, then*

$$(4.14) \quad \text{dist}(x, L_Q) \lesssim \varepsilon_0 \ell(Q).$$

Lemma 4.12. *Let $Q \in \text{Tree}(S_0)$. Then every $x \in Q$ satisfies*

$$\text{dist}(x, \Gamma) \lesssim \varepsilon_0 \ell(Q).$$

Lemma 4.13. *We have*

$$\text{dist}(x, L_{S_0}) \lesssim \varepsilon_0 \ell(S_0)$$

for all $x \in \Gamma$.

4.4. The domains \tilde{U}^+ and \tilde{U}^- . Recall that, without loss of generality, we assume that the n -plane L_{S_0} coincides with the horizontal n -plane $\{(x_1, \dots, x_{n+1}) : x_{n+1} = 0\}$. We denote $\Omega_1^+ = \{(x_1, \dots, x_{n+1}) : x_{n+1} > A(x_1, \dots, x_n)\}$. By the Reifenberg flatness property of Ω^+ (recall (3.1)), for every $Q \in \mathcal{D}$ we have

$$B_Q \setminus U_{C\varepsilon_0 r(B_Q)}(L_Q) \subset \Omega^+ \cup \Omega^-$$

(where $U_t(A)$ stands for the t -neighborhood of A) and one component of $B_Q \setminus U_{C\varepsilon_0 r(B_Q)}(L_Q)$ is contained in Ω^+ and the other in Ω^- . In the particular case of B_{S_0} , by rotating the axes if necessary, we assume that

$$(4.15) \quad B_{S_0} \cap \mathbb{R}_+^{n+1} \setminus U_{C\varepsilon_0 r(B_0)}(L_{S_0}) \subset \Omega^+ \quad \text{and} \quad B_{S_0} \cap \mathbb{R}_-^{n+1} \setminus U_{C\varepsilon_0 r(B_0)}(L_{S_0}) \subset \Omega^-.$$

We define two functions $A^\pm : L_{S_0} \rightarrow L_{S_0}^\perp$ as follows, for $x \in L_{S_0}$,

$$A^+(x) = \inf_{y \in G(S_0)} \left(A(\Pi(y)) + \frac{1}{100} |x - \Pi(y)| \right),$$

$$A^-(x) = \sup_{y \in G(S_0)} \left(A(\Pi(y)) - \frac{1}{100} |x - \Pi(y)| \right).$$

Recall that $G(S_0) \neq \emptyset$ and notice that A^\pm are $\frac{1}{100}$ -Lipschitz functions. Remark also that if $x \in \Pi(G(S_0))$, then $A^+(x) = A^-(x) = A(x)$. Indeed, it is clear from the definition of A^+ that $A^+(x) \leq A(x)$. Also, since A is $1/100$ -Lipschitz, for every $y \in G(S_0)$,

$$A(\Pi(y)) + \frac{1}{100} |x - \Pi(y)| \geq A(x),$$

and taking infimum over y we get $A^+(x) \geq A(x)$. Hence $A(x) = A^+(x)$. The proof of $A(x) = A^-(x)$ is analogous.

Next we consider the domains

$$U^+ = \{x \in \mathbb{R}^{n+1} : \Pi^\perp(x) > A^-(\Pi(x))\},$$

$$U^- = \{x \in \mathbb{R}^{n+1} : \Pi^\perp(x) < A^+(\Pi(x))\}.$$

Notice that U^\pm are both Lipschitz domains.

Lemma 4.14. *We have*

$$\Omega^+ \cap B(z_{S_0}, 10r(B_{S_0})) \subset U^+$$

and

$$\Omega^- \cap B(z_{S_0}, 10r(B_{S_0})) \subset U^-.$$

Proof. We will show that $\Omega^+ \cap B(z_{S_0}, 10r(B_{S_0})) \subset U^+$. The other inclusion is proved by similar arguments.

By connectivity, it is enough to show that

$$(4.16) \quad \partial U^+ \cap B(z_{S_0}, 10r(B_{S_0})) \subset \overline{\Omega^-} = \mathbb{R}^{n+1} \setminus \Omega^+,$$

since $U^+ \cap B(z_{S_0}, 10r(B_{S_0})) \cap \Omega^+ \neq \emptyset$, by the first inclusion in (4.15). To prove (4.16), consider $x \in \partial U^+ \cap B(z_{S_0}, 10r(B_{S_0}))$. Our objective is to show that $x \in \overline{\Omega^-}$. In the case $x \in \overline{G(S_0)}$, by construction we have $x \in \partial \Omega^+ = \partial \Omega^-$ and we are done.

In the case $x \notin \overline{G(S_0)}$, denote $x_0 = \Pi(x)$ and let $y \in \overline{G(S_0)}$ be such that

$$(4.17) \quad \Pi^\perp(x) = A^-(x_0) = A(\Pi(y)) - \frac{1}{100} |x_0 - \Pi(y)|.$$

Recall that we are identifying $L_{S_0}^\perp$ with the vertical axis of \mathbb{R}^{n+1} . Denote $\tilde{x} = (x_0, A(x_0))$, so that

$$\tilde{x} - x = (A(x_0) - A^-(x_0)) e_{n+1}.$$

Since A is $C\alpha$ -Lipschitz, we have

$$(4.18) \quad \Pi^\perp(\tilde{x}) = A(x_0) \geq A(\Pi(y)) - C\alpha |x_0 - \Pi(y)|.$$

Thus, combining (4.17) and (4.18), we obtain

$$A^-(x_0) \leq A(x_0) + \left(C\alpha - \frac{1}{100} \right) |x_0 - \Pi(y)| \leq A(x_0) - \frac{1}{200} |x_0 - \Pi(y)|,$$

assuming α small enough. Hence,

$$|\tilde{x} - x| = A(x_0) - A^-(x_0) \geq \frac{1}{200} |x_0 - \Pi(y)|.$$

For $t \in [0, \ell(R_0)/10]$, consider the point

$$x(t) = x - t e_{n+1},$$

so that $x(0) = x$ and

$$|\tilde{x} - x(t)| = |\tilde{x} - x| + t = A(x_0) - A^-(x_0) + t.$$

Let R_i , $i \in I_W$, be such that $x_0 \in R_i$ (in fact, by the definition of I_0 , we have $i \in I_0$). Observe that

$$\ell(R_i) \lesssim \text{dist}(x_0, \Pi(G(S_0))) \leq |x_0 - \Pi(y)| \lesssim |x - \tilde{x}| \leq |x(t) - \tilde{x}|.$$

By Lemma 4.7, there exists $Q_0 \in \text{Tree}(S_0)$ such that $\ell(Q_0) \approx \ell(R_i)$ and $\text{dist}(\Pi(Q_0), R_i) \lesssim \ell(R_i)$. By Lemma 4.12 we have $\text{dist}(Q_0, \Gamma) \lesssim \ell(Q_0)$ and thus it easily follows that

$$(4.19) \quad \text{dist}(\tilde{x}, Q_0) \lesssim \ell(Q_0).$$

Notice also that the estimate in the display above ensures that

$$(4.20) \quad |x(t) - \tilde{x}| \gtrsim \ell(Q_0).$$

Consider the minimal ancestor $Q \in \mathcal{D}$ of Q_0 such that $x(t), \tilde{x} \in 2B_Q$. Suppose first that $\ell(Q) \leq \ell(S_0)$. From the definition of $\text{Tree}(S_0)$, it follows that $Q \in \text{Tree}(S_0)$, and by the minimality of Q , (4.19), and (4.20), we obtain

$$\ell(Q) \approx r(2B_Q) \approx |x(t) - \tilde{x}| + \ell(Q_0) + \text{dist}(\tilde{x}, Q_0) \lesssim |x(t) - \tilde{x}| + \ell(Q_0) \lesssim |x(t) - \tilde{x}|.$$

Let $z \in L_Q$ be such that $\Pi(z) = \Pi(x(t)) = \Pi(\tilde{x})$. Then, by Lemma 4.11,

$$(4.21) \quad |x(t) - \tilde{x}| \leq |x(t) - z| + |\tilde{x} - z| \lesssim \text{dist}(x(t), L_Q) + \text{dist}(\tilde{x}, L_Q) \lesssim \text{dist}(x(t), L_Q) + \varepsilon_0 \ell(Q).$$

Hence,

$$\text{dist}(x(t), L_Q) \geq c|x(t) - \tilde{x}| - C\varepsilon_0 \ell(Q) \gtrsim \ell(Q).$$

This implies that $x(t)$ belongs to one of the two components of $2B_Q \setminus U_{10\varepsilon_0 r(B_Q)} L_Q$, by the Reifenberg flatness of Ω^- . In the case $\ell(Q) > \ell(S_0)$, since $|x(t) - \tilde{x}| \lesssim \ell(S_0)$ by construction, one easily deduces that $\ell(Q) \approx \ell(S_0)$, and the same estimates above are also valid. Indeed, observe for example that the estimate $\text{dist}(\tilde{x}, L_Q) \lesssim \varepsilon_0 \ell(Q)$ used in (4.21) also holds in this case, by Lemma 4.13.

For t large enough, $x(t) \in \Omega^-$, by the assumption (4.15), and then by connectivity we deduce that $x(t) \in \Omega^-$ for all $t \in [0, \ell(R_0)/10]$. In particular, $x = x(0) \in \Omega^-$. \square

Now we define the domains $\tilde{U}^+, \tilde{U}^- \subset \mathbb{R}^{n+1}$ as follows. First we consider the cylinder $K_0 = \Pi(2B_{R_0}) \times [-4r(B_{R_0}), 4r(B_{R_0})]$, and then we set

$$\tilde{U}^+ = U^+ \cap K_0, \quad \tilde{U}^- = U^- \cap K_0.$$

Observe that both \tilde{U}^+ and \tilde{U}^- are bounded Lipschitz domains, and by Lemma 4.14, we have

$$\Omega^+ \cap K_0 \subset \tilde{U}^+, \quad \Omega^- \cap K_0 \subset \tilde{U}^-.$$

Recall that the estimate (3.3) has already been obtained at the end of Section 4.2. So to complete the proof of Lemma 3.3 it only remains to prove (3.4).

4.5. **Proof of (3.4).** Recall that

$$\omega^+(R \setminus G(R)) \leq 2\varepsilon \omega^+(R)$$

for all $R \in \mathcal{N}(R_0)$, by (3.2) applied to these cubes R in place of R_0 . Thus, using also that ω^+ is doubling and that $K_0 \cap \partial\Omega^+ \subset S_0$,

$$(4.22) \quad \omega^+(K_0 \cap \partial\Omega^+ \setminus G(S_0)) \leq \sum_{R \in \mathcal{N}(R_0)} \omega^+(R \setminus G(R)) \leq 2\varepsilon \sum_{R \in \mathcal{N}(R_0)} \omega^+(R) \lesssim \varepsilon \omega^+(K_0 \cap \partial\Omega^+).$$

Consider a point $p_0 \in K_0 \cap \Omega^+$ such that $\text{dist}(p_0, \partial(K_0 \cap \Omega^+)) \approx \ell(R_0)$. From (4.22) and the change of pole formula (2.3) for NTA domains, we derive

$$(4.23) \quad \omega^{+,p_0}(K_0 \cap \partial\Omega^+ \setminus G(S_0)) \lesssim \varepsilon \omega^{+,p_0}(K_0 \cap \partial\Omega^+) \leq \varepsilon.$$

We also have

$$\omega_{\Omega^+ \cap K_0}^{p_0}(K_0 \cap \partial\Omega^+) \gtrsim 1.$$

This follows easily from Lemma 2.1. Indeed, from this lemma we deduce that, for $p_1 = z_{R_0} + \frac{1}{5}r(B_{R_0})e_{n+1}$,

$$\omega_{\Omega^+ \cap K_0}^{p_1}(K_0 \cap \partial\Omega^+) \geq \omega_{\Omega^+ \cap K_0}^{p_1}(B_{R_0} \cap \partial\Omega^+) \gtrsim \frac{\text{Cap}(\frac{1}{4}B_{R_0} \setminus \Omega^+)}{r(B_{R_0})^{n-1}} \gtrsim 1,$$

where the last estimate follows from the fact that, by the exterior corkscrew condition of Ω^+ , there exists some ball $B \subset \frac{1}{4}B_{R_0} \setminus \Omega^+$ with radius comparable to $r(B_{R_0})$.¹ By the Reifenberg flatness of Ω^+ one can find a Harnack chain contained in $\Omega^+ \cap K_0$ joining p_0 and p_1 (to this end notice that the distance from the segment with end-points p_0, p_1 to $(\Omega^+ \cap K_0)^c$ is comparable to $\ell(R_0)$). Thus,

$$\omega_{\Omega^+ \cap K_0}^{p_0}(K_0 \cap \partial\Omega^+) \approx \omega_{\Omega^+ \cap K_0}^{p_1}(K_0 \cap \partial\Omega^+) \gtrsim 1.$$

From (4.23), the maximum principle, and the last estimate, we infer

$$\omega_{\Omega^+ \cap K_0}^{p_0}(K_0 \cap \partial\Omega^+ \setminus G(S_0)) \leq \omega^{+,p_0}(K_0 \cap \partial\Omega^+ \setminus G(S_0)) \lesssim \varepsilon \approx \varepsilon \omega_{\Omega^+ \cap K_0}^{p_0}(K_0 \cap \partial\Omega^+).$$

Hence,

$$(4.24) \quad \begin{aligned} \omega_{\Omega^+ \cap K_0}^{p_0}(K_0 \cap \partial\Omega^+ \cap G(S_0)) &= \omega_{\Omega^+ \cap K_0}^{p_0}(K_0 \cap \partial\Omega^+) - \omega_{\Omega^+ \cap K_0}^{p_0}(K_0 \cap \partial\Omega^+ \setminus G(S_0)) \\ &\geq (1 - C\varepsilon) \omega_{\Omega^+ \cap K_0}^{p_0}(K_0 \cap \partial\Omega^+). \end{aligned}$$

Next we claim that

$$(4.25) \quad \omega_{\Omega^+ \cap K_0}^{p_0}(K_0 \cap \partial\Omega^+) \geq \omega_{\tilde{U}^+}^{p_0}(K_0 \cap \partial U^+).$$

Indeed, for $x \in K_0 \cap \partial\Omega^+$, we have $\omega_{\Omega^+ \cap K_0}^x(K_0 \cap \partial\Omega^+) = 1 \geq \omega_{\tilde{U}^+}^x(K_0 \cap \partial U^+)$. For $x \in \partial K_0 \cap \overline{\Omega^+}$, since $\partial K_0 \cap \overline{\Omega^+} \subset \partial K_0 \cap \overline{\tilde{U}^+}$, it follows also that

$$\omega_{\Omega^+ \cap K_0}^x(K_0 \cap \partial\Omega^+) = 0 = \omega_{\tilde{U}^+}^x(K_0 \cap \partial U^+).$$

So our claim follows from the maximum principle, since both $\omega_{\Omega^+ \cap K_0}^x(K_0 \cap \partial\Omega^+)$ and $\omega_{\tilde{U}^+}^x(K_0 \cap \partial U^+)$ are harmonic functions of $x \in \Omega^+ \cap K_0$.

¹In the planar case, a similar estimate holds works, using an argument involving logarithmic capacity instead of Newtonian capacity.

By the maximum principle again, recalling that $G(S_0) \subset \partial\Omega^+ \cap \partial U^+$, and by (4.24) and (4.25), we get

$$\omega_{\tilde{U}^+}^{p_0}(K_0 \cap G(S_0)) \geq \omega_{\Omega^+ \cap K_0}^{p_0}(K_0 \cap G(S_0)) \geq (1 - C\varepsilon) \omega_{\Omega^+ \cap K_0}^{p_0}(K_0 \cap \partial\Omega^+) \geq (1 - C\varepsilon) \omega_{\tilde{U}^+}^{p_0}(K_0 \cap \partial U^+).$$

Equivalently,

$$\omega_{\tilde{U}^+}^{p_0}(K_0 \cap \partial U^+ \setminus G(S_0)) \leq C\varepsilon \omega_{\tilde{U}^+}^{p_0}(K_0 \cap \partial U^+).$$

Now, since \tilde{U}^+ is a Lipschitz domain, by Dahlberg's theorem $\omega_{\tilde{U}^+}^{p_0}$ is an A_∞ weight with respect to $\mathcal{H}^n|_{\partial\tilde{U}^+}$, and thus we have

$$(4.26) \quad \mathcal{H}^n(K_0 \cap \partial U^+ \setminus G(S_0)) \lesssim \varepsilon^a \mathcal{H}^n(K_0 \cap \partial U^+) \lesssim \varepsilon^a \ell(R_0)^n,$$

for some $a > 0$ (depending only on the A_∞ character of $\omega_{\tilde{U}^+}^{p_0}$ with respect to $\mathcal{H}^n|_{\partial\tilde{U}^+}$).

Our next objective consists of showing that

$$(4.27) \quad \sum_{Q \in \text{Stop}(R_0)} \ell(Q)^n \lesssim \mathcal{H}^n(K_0 \cap \partial U^+ \setminus G(S_0)).$$

Clearly, this estimate, together with (4.26), implies (3.4) for ε small enough. To prove (4.27), notice that, for each $Q \in \text{Stop}(R_0)$, $\frac{1}{4}\tilde{B}_Q \cap \Gamma \neq \emptyset$ by Lemma 4.12 (recall that \tilde{B}_Q is defined in Theorem 2.3(d)). Hence,

$$\ell(Q)^n \lesssim \mathcal{H}^n(\frac{1}{2}\tilde{B}_Q \cap \Gamma).$$

Observe also that the balls $\frac{1}{2}\tilde{B}_Q$, with $Q \in \text{Stop}(R_0)$, are disjoint from $G(S_0)$. Indeed, by definition $G(S_0) = S_0 \setminus \bigcup_{P \in \text{Stop}(S_0)} P$ and so $\tilde{B}_Q \cap \partial\Omega^+ \subset Q \subset \partial\Omega^+ \setminus G(S_0)$. Then we deduce

$$(4.28) \quad \sum_{Q \in \text{Stop}(R_0)} \ell(Q)^n \lesssim \sum_{Q \in \text{Stop}(R_0)} \mathcal{H}^n(\frac{1}{2}\tilde{B}_Q \cap \Gamma) \leq \mathcal{H}^n(B_{R_0} \cap \Gamma \setminus G(S_0)),$$

where in the last inequality we took into account that the balls $\frac{1}{2}\tilde{B}_Q$, with $Q \in \text{Stop}(R_0)$, are pairwise disjoint, contained in B_{R_0} , and disjoint from $G(S_0)$. Using the fact that $G(S_0) \subset \Gamma \cap \partial\tilde{U}^+$ and that Γ and ∂U^+ are Lipschitz graphs, we derive

$$(4.29) \quad \mathcal{H}^n(B_{R_0} \cap \Gamma \setminus G(S_0)) \approx \mathcal{H}^n(\Pi(B_{R_0} \cap \Gamma \setminus G(S_0))) \leq \mathcal{H}^n(K_0 \cap \partial U^+ \setminus G(S_0)).$$

By combining (4.28) and (4.29) we get (4.27). Finally, just note again that (4.26) and (4.27) yield (3.4) and then the proof of Lemma 3.3 is concluded. \square

REFERENCES

- [AM] J. Azzam and M. Mouroglou. *Tangent measures of elliptic measure and applications*. Anal. PDE 12 (2019), no. 8, 1891–1941. [1](#)
- [AMT1] J. Azzam, M. Mouroglou and X. Tolsa. *Mutual absolute continuity of interior and exterior harmonic measure implies rectifiability*. Comm. Pure Appl. Math. 71 (2017), no. 11, 2121–2163. [1](#)
- [AMT2] J. Azzam, M. Mouroglou and X. Tolsa. *A two-phase free boundary problem for harmonic measure and uniform rectifiability*. Trans. Amer. Math. Soc. 373 (2020), no. 6, 4359–4388. [1](#)
- [AMTV] J. Azzam, M. Mouroglou, X. Tolsa, and A. Volberg. *On a two-phase problem for harmonic measure in general domains*. Amer. J. Math. 141 (2019), no. 5, 1259–1279. [1](#)
- [BET1] M. Badger, M. Engelstein, and T. Toro. *Structure of sets which are well approximated by zero sets of harmonic polynomials*. Anal. PDE 10 (2017), no. 6, 1455–1495. [2](#)
- [BET2] M. Badger, M. Engelstein, and T. Toro. *Regularity of the singular set in a two-phase problem for harmonic measure with Hölder data*. Rev. Mat. Iberoam. 36 (2020), no. 5, 1375–1408. [2](#)

- [BH] S. Bortz and S. Hofmann. *A singular integral approach to a two phase free boundary problem*. Proc. Amer. Math. Soc. 144 (2016), no. 9, 3959–3973. [2](#)
- [Ch] M. Christ. *A $T(b)$ theorem with remarks on analytic capacity and the Cauchy integral*. Colloq. Math. 60/61(2) (1990), 601–628. [6](#)
- [DJ] G. David and D. Jerison. *Lipschitz approximation to hypersurfaces, harmonic measure, and singular integrals*. Indiana Univ. Math. J. 39 (1990), no. 3, 831–845.
- [DM] G. David and P. Mattila. *Removable sets for Lipschitz harmonic functions in the plane*. Rev. Mat. Iberoamericana 16(1) (2000), 137–215. [6](#)
- [DS] G. David and S. Semmes. *Singular integrals and rectifiable sets in \mathbb{R}^n : Beyond Lipschitz graphs*, Astérisque, No. 193 (1991). [4](#)
- [En] M. Engelstein. *A two-phase free boundary problem for harmonic measure*. Ann. Sci. Éc. Norm. Supér. (4) 49 (2016), no. 4, 859–905. [1](#)
- [He] Helms, L. L. *Potential theory*, Universitext, Springer, London, 2014, 2nd ed. [4](#)
- [HK] T. Hytönen and A. Kairema. *Systems of dyadic cubes in a doubling metric space*. Colloq. Math. 126 (2012), no. 1, 1–33. [6](#)
- [JK] D. S. Jerison and C. E. Kenig. *Boundary behavior of harmonic functions in nontangentially accessible domains*, Adv. in Math. 46 (1982), no. 1, 80–147. [5](#)
- [KPT] C. Kenig, D. Preiss, and T. Toro. *Boundary structure and size in terms of interior and exterior harmonic measures in higher dimensions*. J. Amer. Math. Soc. 22 (2009), no. 3, 771–796. [1](#)
- [KT1] C.E. Kenig and T. Toro. *Harmonic measure on locally flat domains*. Duke Math. J. 87 (1997), no. 3, 509–551. [3](#), [6](#)
- [KT2] C. Kenig and T. Toro. *Free boundary regularity for harmonic measures and Poisson kernels*, Ann. of Math. 150 (1999) 369–454. [3](#)
- [KT3] C. Kenig and T. Toro. *Free boundary regularity below the continuous threshold: 2 phase problems*. J. Reine Angew. Math. 596 (2006), 1–44. [1](#)
- [KT4] C. Kenig and T. Toro. *Poisson kernel characterization of Reifenberg flat chord arc domains*, Ann. Sci. Ecole Norm. Sup. (4) 36 (2003), no.3, 323–401. [3](#)
- [MT] M. Mourougolou and X. Tolsa. *Harmonic measure and Riesz transform in uniform and general domains*. J. Reine Angew. Math. 758 (2020), 183–221. [5](#)
- [PT] M. Prats and X. Tolsa. *The two-phase problem for harmonic measure in VMO*. Calc. Var. Partial Differential Equations 59 (2020), no. 3, Paper No. 102, 58 pp. [2](#), [8](#)
- [To1] X. Tolsa. *The mutual singularity of harmonic measure and Hausdorff measure of codimension smaller than one*. Int. Math. Res. Not. IMRN 2021, no. 18, 13783–13811. [4](#)
- [To2] X. Tolsa. *Analytic capacity, the Cauchy transform, and non-homogeneous Calderón-Zygmund theory*. Progress in Mathematics, vol. 307, Birkhäuser/Springer, Cham, 2014. [11](#), [12](#), [13](#), [14](#)

ICREA, BARCELONA, DEPT. DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, AND CENTRE DE RECERCA MATEMÀTICA, BARCELONA, CATALONIA.

Email address: xtolsa@mat.uab.cat

DEPARTMENT OF MATHEMATICS, BOX 354350, UNIVERSITY OF WASHINGTON, SEATTLE, WA 98195-4350, USA.

Email address: toro@uw.edu