

CANAL HYPERSURFACES GENERATED BY PSEUDO NULL, PARTIALLY NULL AND NULL CURVES IN LORENTZ-MINKOWSKI 4-SPACE

MUSTAFA ALTIN^{1*}, AHMET KAZAN² AND DAE WON YOON³

ABSTRACT. In this paper, we obtain the parametric expressions of the canal hypersurfaces that are formed as the envelope of a family of pseudo hyperspheres or pseudo hyperbolic hyperspheres whose centers lie on a pseudo null, partially null or null curves in E_1^4 and give their some geometric invariants such as unit normal vector fields, Gaussian curvatures and mean curvatures. Also, we construct some examples for these canal hypersurfaces and finally, we give some characterizations for tubular hypersurfaces in E_1^4 .

1. GENERAL INFORMATION AND BASIC CONCEPTS

The class of surfaces formed by sweeping a sphere is called canal surfaces and they have been investigated by Monge in 1850. So, one can see a canal surface as the envelope of a moving sphere with varying radius, defined by the trajectory $\gamma(s)$ of its centers and a radius function $r(s)$. The canal surface is called a tubular surface or pipe surface if the radius function is constant. After defining canal and tubular surfaces, many mathematicians have studied their different geometric characterizations and these surfaces have been applied to many fields, such as the solid and the surface modeling for CAD/CAM, construction of blending surfaces, shape re-construction. Also, they are useful to represent various objects such as pipe, hose, rope or intestine of a body [22], [36]. For some results about different geometric characterizations of canal and tubular (hyper)surfaces in different three, four or higher dimensional spaces, we refer to [2], [7], [11], [12], [13], [15], [16], [19], [20], [21], [26], [27], [29], [32], [34], [35], [36], [38], [39], and etc.

On the other hand, the extrinsic differential geometry of submanifolds in Lorentz-Minkowski 4-space E_1^4 is of special interest in Relativity Theory. So, many studies about curves and (hyper)surfaces have been done in this space and this motivated us to construct the canal hypersurfaces using pseudo null, partially null and null curves in E_1^4 . Here, we'll recall some basic concepts for curves and hypersurfaces in E_1^4 .

Let $\vec{u} = (u_1, u_2, u_3, u_4)$, $\vec{v} = (v_1, v_2, v_3, v_4)$ and $\vec{w} = (w_1, w_2, w_3, w_4)$ be three vectors in E_1^4 . Then the inner product of two vectors and vector product of three vectors in E_1^4 are given by

$$\langle \vec{u}, \vec{v} \rangle = -u_1v_1 + u_2v_2 + u_3v_3 + u_4v_4 \quad (1.1)$$

and

$$\vec{u} \times \vec{v} \times \vec{w} = \det \begin{bmatrix} -e_1 & e_2 & e_3 & e_4 \\ u_1 & u_2 & u_3 & u_4 \\ v_1 & v_2 & v_3 & v_4 \\ w_1 & w_2 & w_3 & w_4 \end{bmatrix}, \quad (1.2)$$

respectively.

A vector $\vec{u} \in E_1^4 - \{0\}$ is called spacelike if $\langle \vec{u}, \vec{u} \rangle > 0$; timelike if $\langle \vec{u}, \vec{u} \rangle < 0$ and lightlike (null) if $\langle \vec{u}, \vec{u} \rangle = 0$. In particular, the vector $\vec{u} = 0$ is spacelike. Also, the norm of the vector \vec{u} is $\|\vec{u}\| = \sqrt{|\langle \vec{u}, \vec{u} \rangle|}$. A curve $\gamma(s)$ in E_1^4 is spacelike, timelike or lightlike (null), if all its velocity vectors $\gamma'(s)$ are spacelike, timelike or lightlike, respectively and a non-null (i.e. timelike or spacelike) curve has unit speed if $\langle \gamma', \gamma' \rangle = \mp 1$. If $\langle \gamma''(s), \gamma''(s) \rangle = 1$, then the null curve γ is parametrized by

2010 *Mathematics Subject Classification.* 14J70, 53A07, 53A10.

Key words and phrases. Pseudo null curve, Partially null curve, Null curve, Canal hypersurface, Tubular hypersurface.

arclength function s . Also, if the principal normal vector or the binormal vector of a spacelike curve $\gamma(s)$ in E_1^4 is null, then the spacelike curve $\gamma(s)$ is called a pseudo null curve or a partially null curve (for detail, one can see [9], [10], [30], [33], and etc.)

If $\{F_1, F_2, F_3, F_4\}$ is the moving Frenet frame along a curve $\gamma(s)$ in E_1^4 consisting of the unit tangent vector field, principal normal vector field, binormal vector field and trinormal vector field, then we have the following cases according to the causal character of γ (see [9], [10], [17], [37], and etc):

i) If the curve $\gamma(s)$ is pseudo null, then the Frenet formulas are

$$\left. \begin{aligned} F_1' &= k_1 F_2, \\ F_2' &= k_2 F_3, \\ F_3' &= k_3 F_2 - k_2 F_4, \\ F_4' &= -k_1 F_1 - k_3 F_3, \end{aligned} \right\} \quad (1.3)$$

and we have

$$\left. \begin{aligned} \langle F_1, F_1 \rangle &= \langle F_3, F_3 \rangle = 1, \quad \langle F_2, F_2 \rangle = \langle F_4, F_4 \rangle = 0, \\ \langle F_2, F_4 \rangle &= 1, \quad \langle F_1, F_2 \rangle = \langle F_1, F_3 \rangle = \langle F_1, F_4 \rangle = \langle F_2, F_3 \rangle = \langle F_3, F_4 \rangle = 0. \end{aligned} \right\} \quad (1.4)$$

ii) If the curve $\gamma(s)$ is partially null, then the Frenet formulas are

$$\left. \begin{aligned} F_1' &= k_1 F_2, \\ F_2' &= -k_1 F_1 + k_2 F_3, \\ F_3' &= k_3 F_3, \\ F_4' &= -k_2 F_2 - k_3 F_4, \end{aligned} \right\} \quad (1.5)$$

and we have

$$\left. \begin{aligned} \langle F_1, F_1 \rangle &= \langle F_2, F_2 \rangle = 1, \quad \langle F_3, F_3 \rangle = \langle F_4, F_4 \rangle = 0, \\ \langle F_3, F_4 \rangle &= 1, \quad \langle F_1, F_2 \rangle = \langle F_1, F_3 \rangle = \langle F_1, F_4 \rangle = \langle F_2, F_3 \rangle = \langle F_2, F_4 \rangle = 0. \end{aligned} \right\} \quad (1.6)$$

iii) If the curve $\gamma(s)$ is null, then the Frenet formulas are

$$\left. \begin{aligned} F_1' &= k_1 F_2, \\ F_2' &= k_2 F_1 - k_1 F_3, \\ F_3' &= -k_2 F_2 + k_3 F_4, \\ F_4' &= -k_3 F_1, \end{aligned} \right\} \quad (1.7)$$

and we have

$$\left. \begin{aligned} \langle F_2, F_2 \rangle &= \langle F_4, F_4 \rangle = 1, \quad \langle F_1, F_1 \rangle = \langle F_3, F_3 \rangle = 0, \\ \langle F_1, F_3 \rangle &= 1, \quad \langle F_1, F_2 \rangle = \langle F_1, F_4 \rangle = \langle F_2, F_3 \rangle = \langle F_2, F_4 \rangle = \langle F_3, F_4 \rangle = 0. \end{aligned} \right\} \quad (1.8)$$

Here, the first curvature $k_1(s) = 0$, if γ is a straight line and $k_1(s) = 1$ in all other cases for pseudo null and null curves. So, the pseudo null and null curves have two curvatures $k_2(s)$ and $k_3(s)$. Also, the third curvature $k_3(s) = 0$ for each s for a partially null curve and thus, a partially null curve has two curvatures $k_1(s)$ and $k_2(s)$.

Furthermore, if p is a fixed point in E_1^4 and r is a positive constant, then the pseudo-Riemannian hypersphere, pseudo-Riemannian hyperbolic space and pseudo-Riemannian null hypercone are defined by

$$\left. \begin{aligned} S_1^3(p, r) &= \{u \in E_1^4 : \langle u - p, u - p \rangle = r^2\}, \\ H_0^3(p, r) &= \{u \in E_1^4 : \langle u - p, u - p \rangle = -r^2\}, \\ Q_1^3 &= \{u \in E_1^4 : \langle u - p, u - p \rangle = 0\}, \end{aligned} \right\} \quad (1.9)$$

respectively.

In the present study, we construct the canal hypersurfaces in E_1^4 as the envelope of a family of pseudo hyperspheres or pseudo hyperbolic hyperspheres whose centers lie on a pseudo null, partially null or null curve.

Furthermore, the differential geometry of different types of (hyper)surfaces in 4-dimensional spaces has been a popular topic for geometers, recently ([3], [4], [5], [6], [8], [23], [24], [25], [28], and etc). In this context, let Ω be a hypersurface in E_1^4 given by

$$\begin{aligned} \Omega : U \subset E^3 &\longrightarrow E_1^4 \\ (u_1, u_2, u_3) &\longrightarrow \Omega(u_1, u_2, u_3) = (\Omega_1(u_1, u_2, u_3), \Omega_2(u_1, u_2, u_3), \Omega_3(u_1, u_2, u_3), \Omega_4(u_1, u_2, u_3)). \end{aligned} \quad (1.10)$$

Then the Gauss map (i.e., the unit normal vector field), the matrix forms of the first and second fundamental forms are

$$N_\Omega = \frac{\Omega_{u_1} \times \Omega_{u_2} \times \Omega_{u_3}}{\|\Omega_{u_1} \times \Omega_{u_2} \times \Omega_{u_3}\|}, \quad [g_{ij}] = \begin{bmatrix} g_{11} & g_{12} & g_{13} \\ g_{21} & g_{22} & g_{23} \\ g_{31} & g_{32} & g_{33} \end{bmatrix} \quad \text{and} \quad [h_{ij}] = \begin{bmatrix} h_{11} & h_{12} & h_{13} \\ h_{21} & h_{22} & h_{23} \\ h_{31} & h_{32} & h_{33} \end{bmatrix}, \quad (1.11)$$

respectively. Here $g_{ij} = \langle \Omega_{u_i}, \Omega_{u_j} \rangle$, $h_{ij} = \langle \Omega_{u_i u_j}, N_\Omega \rangle$, $\Omega_{u_i} = \frac{\partial \Omega}{\partial u_i}$, $\Omega_{u_i u_j} = \frac{\partial^2 \Omega}{\partial u_i \partial u_j}$, $i, j \in \{1, 2, 3\}$. Also, the matrix of shape operator of the hypersurface (1.10) is

$$S = [a_{ij}] = [g^{ij}] \cdot [h_{ij}], \quad (1.12)$$

where $[g^{ij}]$ is the inverse matrix of $[g_{ij}]$. With the aid of (1.11) and (1.12), the Gaussian curvature and mean curvature of a hypersurface in E_1^4 are given by

$$K = \varepsilon \frac{\det[h_{ij}]}{\det[g_{ij}]} \quad \text{and} \quad 3\varepsilon H = \text{tr}(S), \quad (1.13)$$

respectively. Here, $\varepsilon = \langle N_\Omega, N_\Omega \rangle$. We say that a hypersurface is flat or minimal, if it has zero Gaussian or zero mean curvature, respectively. For more details about hypersurfaces in E_1^4 , we refer to [14], [31] and etc.

2. CANAL HYPERSURFACES GENERATED BY PSEUDO NULL, PARTIALLY NULL AND NULL CURVES IN E_1^4

In this section, firstly we construct the canal hypersurfaces that are formed as the envelope of a family of pseudo hyperspheres or pseudo hyperbolic hyperspheres whose centers lie on a pseudo null curve, partially null curve or null curve in E_1^4 . After that, we obtain some important geometric invariants such as unit normal vector fields, Gaussian curvatures and mean curvatures of these canal hypersurfaces, separately.

Theorem 1. *The canal hypersurfaces that are formed as the envelope of a family of pseudo hyperspheres whose centers lie on a pseudo null curve $\gamma(s)$ with Frenet vector fields F_i , $i \in \{1, 2, 3, 4\}$, in E_1^4 can be parametrized by*

$$\left. \begin{aligned} & \overset{psd-n}{C_1}(s, t, w) = \gamma(s) - r(s)r'(s)F_1(s) \\ & \quad \mp r(s)\sqrt{1-r'(s)^2} \left(g(t, w) \sin(f(t, w))F_2(s) + \cos(f(t, w))F_3(s) + \frac{\sin(f(t, w))}{2g(t, w)}F_4(s) \right), \\ & \overset{psd-n}{C_2}(s, t, w) = \gamma(s) - r(s)r'(s)F_1(s) \\ & \quad \mp r(s)\sqrt{1-r'(s)^2} \left(g(t, w) \cos(f(t, w))F_2(s) + \sin(f(t, w))F_3(s) + \frac{\cos(f(t, w))}{2g(t, w)}F_4(s) \right), \\ & \overset{psd-n}{C_3}(s, t, w) = \gamma(s) - r(s)r'(s)F_1(s) \\ & \quad \mp r(s)\sqrt{1-r'(s)^2} \left(g(t, w) \sinh(f(t, w))F_2(s) + \cosh(f(t, w))F_3(s) - \frac{\sinh(f(t, w))}{2g(t, w)}F_4(s) \right), \\ & \overset{psd-n}{C_4}(s, t, w) = \gamma(s) - r(s)r'(s)F_1(s) \\ & \quad \mp r(s)\sqrt{r'(s)^2-1} \left(g(t, w) \cosh(f(t, w))F_2(s) + \sinh(f(t, w))F_3(s) - \frac{\cosh(f(t, w))}{2g(t, w)}F_4(s) \right), \end{aligned} \right\} \quad (2.1)$$

where we suppose $r'(s)^2 < 1$ for canal hypersurfaces ${}^{psd-n}C_1$, ${}^{psd-n}C_2$, ${}^{psd-n}C_3$ and $r'(s)^2 > 1$ for canal hypersurface ${}^{psd-n}C_4$.

Furthermore, the canal hypersurfaces that are formed as the envelope of a family of pseudo hyperbolic hyperspheres whose centers lie on a pseudo null curve $\gamma(s)$ with Frenet vector fields F_i , $i \in \{1, 2, 3, 4\}$, in E_1^4 can be parametrized by

$$\begin{aligned} {}^{psd-n}C_5(s, t, w) &= \gamma(s) + r(s)r'(s)F_1(s) \\ &\mp r(s)\sqrt{1+r'(s)^2} \left(g(t, w) \cosh(f(t, w))F_2(s) + \sinh(f(t, w))F_3(s) - \frac{\cosh(f(t, w))}{2g(t, w)}F_4(s) \right). \end{aligned} \quad (2.2)$$

Here, the canal hypersurfaces ${}^{psd-n}C_1$, ${}^{psd-n}C_2$, ${}^{psd-n}C_3$, ${}^{psd-n}C_4$ are timelike and the canal hypersurface ${}^{psd-n}C_5$ is spacelike.

Proof. Let the center curve $\gamma : I \subseteq \mathbb{R} \rightarrow E_1^4$ be a pseudo null curve with non-zero curvature with Frenet vector fields $F_1(s)$, $F_2(s)$, $F_3(s)$, $F_4(s)$ called unit tangent, principal normal, binormal and trinormal vectors of $\gamma(s)$, respectively. Then, the parametrization of the envelope of pseudo hyperspheres (or pseudo hyperbolic hyperspheres) defining the canal hypersurfaces ${}^{psd-n}C(s, t, w)$ in E_1^4 can be given by

$${}^{psd-n}C(s, t, w) - \gamma(s) = a_1(s, t, w)F_1(s) + a_2(s, t, w)F_2(s) + a_3(s, t, w)F_3(s) + a_4(s, t, w)F_4(s), \quad (2.3)$$

where $a_i(s, t, w)$ are differentiable functions of s, t, w on the interval I . Furthermore, since ${}^{psd-n}C(s, t, w)$ lies on the pseudo hyperspheres ($\lambda = 1$) (or pseudo hyperbolic hyperspheres ($\lambda = -1$)), we have

$$g({}^{psd-n}C(s, t, w) - \gamma(s), {}^{psd-n}C(s, t, w) - \gamma(s)) = \lambda r^2(s) \quad (2.4)$$

which leads to from (2.3) and (1.4) that

$$a_1^2 + a_3^2 + 2a_2a_4 = \lambda r^2 \quad (2.5)$$

and

$$a_1a_{1_s} + a_3a_{3_s} + a_4a_{2_s} + a_2a_{4_s} = \lambda rr_s, \quad (2.6)$$

where $r(s)$ is the radius function; $r = r(s)$, $r_s = \frac{dr(s)}{ds}$, $a_i = a_i(s, t, w)$, $a_{i_s} = \frac{\partial a_i(s, t, w)}{\partial s}$.

So, differentiating (2.3) with respect to s and using the Frenet formula (1.3), we get

$$\begin{aligned} ({}^{psd-n}C)_s &= (1 - a_4k_1 + a_{1_s})F_1 + (a_1k_1 + a_3k_3 + a_{2_s})F_2 \\ &+ (a_2k_2 - a_4k_3 + a_{3_s})F_3 + (-a_3k_2 + a_{4_s})F_4, \end{aligned} \quad (2.7)$$

where $({}^{psd-n}C)_s = \frac{\partial ({}^{psd-n}C(s, t, w))}{\partial s}$. Furthermore, ${}^{psd-n}C(s, t, w) - \gamma(s)$ is a normal vector to the canal hypersurfaces, which implies that

$$g({}^{psd-n}C(s, t, w) - \gamma(s), ({}^{psd-n}C)_s(s, t, w)) = 0 \quad (2.8)$$

and so, from (2.3), (2.7) and (2.8) we have

$$\left(\begin{array}{l} a_1(1 - a_4k_1 + a_{1_s}) + a_2(-a_3k_2 + a_{4_s}) \\ + a_3(a_2k_2 - a_4k_3 + a_{3_s}) + a_4(a_1k_1 + a_3k_3 + a_{2_s}) \end{array} \right) = 0. \quad (2.9)$$

Using (2.6) in (2.9), we get

$$a_1 = -\lambda rr_s. \quad (2.10)$$

Hence, using (2.10) in (2.5), we reach that

$$a_3^2 + 2a_2a_4 = \lambda r^2(1 - \lambda r_s^2). \quad (2.11)$$

Therefore from (2.10) and (2.11), the canal hypersurfaces $C^{psd-n}(s, t, w)$ that are formed as the envelope of a family of pseudo hyperspheres or pseudo hyperbolic hyperspheres whose centers lie on a pseudo null curve in E_1^4 can be parametrized by (2.1) or (2.2), respectively. \square

Here we must note that, from now on we will state $r' = r_s(s)$, $r'' = \frac{d^2r(s)}{ds^2}$, $f = f(t, w)$, $g = g(t, w)$, $\sin f = \sin(f(t, w))$, $f_t = \frac{\partial f(t, w)}{\partial t}$, $f_{tt} = \frac{\partial^2 f(t, w)}{\partial t^2}$, and so on.

Theorem 2. *The Gaussian and mean curvatures of the canal hypersurfaces C_i^{psd-n} , $i \in \{1, 2, 3, 4, 5\}$, given by (2.1) and (2.2) in E_1^4 are*

$$\left. \begin{aligned}
 & \left. \begin{aligned}
 K_{C_1^{psd-n}} &= \frac{-r(1-r'^2)k_1^2 \sin^2 f + 4r''(1-r'^2 - rr'')g^2 + 2\sqrt{1-r'^2}(1-r'^2 - 2rr'')k_1 g \sin f}{r^2(r\sqrt{1-r'^2}k_1 \sin f - 2(1-r'^2 - rr'')g)^2} \\
 H_{C_1^{psd-n}} &= \frac{2r(1-r'^2)^{3/2}k_1 g \sin f + 3r^2(1-r'^2)k_1^2 \sin^2 f - 4(1-r'^2 - rr'')(2-2r'^2 - 3rr'')g^2}{3(-r^3(1-r'^2)k_1^2 \sin^2 f + 4r(1-r'^2 - rr'')^2 g^2)}
 \end{aligned} \right\}, \\
 & \left. \begin{aligned}
 K_{C_2^{psd-n}} &= \frac{r(1-r'^2)k_1^2 \cos^2 f - 4r''(1-r'^2 - rr'')g^2 - 2\sqrt{1-r'^2}(1-r'^2 - 2rr'')k_1 g \cos f}{r^2(r\sqrt{1-r'^2}k_1 \cos f - 2(1-r'^2 - rr'')g)^2} \\
 H_{C_2^{psd-n}} &= \frac{-2r(1-r'^2)^{3/2}k_1 g \cos f - 3r^2(1-r'^2)k_1^2 \cos^2 f + 4(1-r'^2 - rr'')(2-2r'^2 - 3rr'')g^2}{3(-r^3(1-r'^2)k_1^2 \cos^2 f + 4r(1-r'^2 - rr'')^2 g^2)}
 \end{aligned} \right\}, \\
 & \left. \begin{aligned}
 K_{C_3^{psd-n}} &= \frac{-r(1-r'^2)k_1^2 \sinh^2 f + 4r''(1-r'^2 - rr'')g^2 - 2\sqrt{1-r'^2}(1-r'^2 - 2rr'')k_1 g \sinh f}{r^2(r\sqrt{1-r'^2}k_1 \sinh f + 2(1-r'^2 - rr'')g)^2} \\
 H_{C_3^{psd-n}} &= \frac{-2r(1-r'^2)^{3/2}k_1 g \sinh f + 3r^2(1-r'^2)k_1^2 \sinh^2 f - 4(1-r'^2 - rr'')(2-2r'^2 - 3rr'')g^2}{3(-r^3(1-r'^2)k_1^2 \sinh^2 f + 4r(1-r'^2 - rr'')^2 g^2)}
 \end{aligned} \right\}, \\
 & \left. \begin{aligned}
 K_{C_4^{psd-n}} &= \frac{-r(r'^2 - 1)k_1^2 \cosh^2 f - 4r''(-1+r'^2 + rr'')g^2 + 2\sqrt{-1+r'^2}(-1+r'^2 + 2rr'')k_1 g \cosh f}{r^2(r\sqrt{r'^2 - 1}k_1 \cosh f - 2(-1+r'^2 + rr'')g)^2} \\
 H_{C_4^{psd-n}} &= \frac{-2r(r'^2 - 1)^{3/2}k_1 g \cosh f - 3r^2(r'^2 - 1)k_1^2 \cosh^2 f + 4(-1+r'^2 + rr'')(-2+2r'^2 + 3rr'')g^2}{3(r^3(r'^2 - 1)k_1^2 \cosh^2 f - 4r(-1+r'^2 + rr'')^2 g^2)}
 \end{aligned} \right\}, \\
 & \left. \begin{aligned}
 K_{C_5^{psd-n}} &= \frac{r(1+r'^2)k_1^2 \cosh^2 f + 4r''(1+r'^2 + rr'')g^2 + 2\sqrt{1+r'^2}(1+r'^2 + 2rr'')k_1 g \cosh f}{r^2(r\sqrt{1+r'^2}k_1 \cosh f + 2(1+r'^2 + rr'')g)^2} \\
 H_{C_5^{psd-n}} &= \frac{-2r(1+r'^2)^{3/2}k_1 g \cosh f + 3r^2(1+r'^2)k_1^2 \cosh^2 f - 4(1+r'^2 + rr'')(2+2r'^2 + 3rr'')g^2}{3(r^3(1+r'^2)k_1^2 \cosh^2 f - 4r(1+r'^2 + rr'')^2 g^2)}
 \end{aligned} \right\}.
 \end{aligned} \right\} \quad (2.12)$$

Proof. Here we will obtain the unit normal vector field, Gaussian and mean curvatures of the canal hypersurfaces $C_1^{psd-n}(s, t, w)$ given by

$$\begin{aligned}
 C_1^{psd-n}(s, t, w) &= \gamma(s) - r(s)r'(s)F_1(s) \\
 &+ r(s)\sqrt{1-r'(s)^2} \left(g(t, w) \sin(f(t, w))F_2(s) + \cos(f(t, w))F_3(s) + \frac{\sin(f(t, w))}{2g(t, w)}F_4(s) \right),
 \end{aligned} \quad (2.13)$$

in E_1^4 . Firstly, from (1.3), the first derivatives of the canal hypersurface (2.13) are obtained as

$$\binom{psd-n}{C_1}_s = A_1 F_1 + A_2 F_2 + A_3 F_3 + A_4 F_4, \quad (2.14)$$

$$\binom{psd-n}{C_1}_t = r\sqrt{1-r'^2} (gf_t \cos f + g_t \sin f) F_2 - r\sqrt{1-r'^2} f_t \sin f F_3 \\ + \frac{r\sqrt{1-r'^2}}{2g^2} (gf_t \cos f - g_t \sin f) F_4, \quad (2.15)$$

$$\binom{psd-n}{C_1}_w = r\sqrt{1-r'^2} (gf_w \cos f + g_w \sin f) F_2 - r\sqrt{1-r'^2} f_w \sin f F_3 \\ + \frac{r\sqrt{1-r'^2}}{2g^2} (gf_w \cos f - g_w \sin f) F_4, \quad (2.16)$$

where

$$\left. \begin{aligned} A_1 &= 1 - r'^2 - rr'' - \frac{r\sqrt{1-r'^2}k_1 \sin f}{2g}, \\ A_2 &= \frac{1}{\sqrt{1-r'^2}} \left(r(1-r'^2)k_3 \cos f - r' \left(r\sqrt{1-r'^2}k_1 - (1-r'^2 - rr'')g \sin f \right) \right), \\ A_3 &= \frac{(r(1-r'^2)(2k_2g^2 - k_3) \sin f) + 2r'(1-r'^2 - rr'')g \cos f}{2g\sqrt{1-r'^2}}, \\ A_4 &= \frac{-2r(1-r'^2)k_2g \cos f + r'(1-r'^2 - rr'') \sin f}{2g\sqrt{1-r'^2}}. \end{aligned} \right\}$$

From (1.11) and (2.14)-(2.16), the unit normal vector field of $\binom{psd-n}{C_1}$ in E^4 is

$$N = -r'F_1 + \sqrt{1-r'^2} \left(g \sin f F_2 + \cos f F_3 + \frac{\sin f}{2g} F_4 \right). \quad (2.17)$$

and we get $\langle N, N \rangle = 1$. Also, the coefficients of the first fundamental form are given by

$$\left. \begin{aligned} g_{11} &= \frac{1}{4g^2(1-r'^2)} \left(\begin{aligned} &(1-r'^2) \left(r\sqrt{1-r'^2}k_1 \sin f - 2(1-r'^2 - rr'')g \right)^2 \\ &+ (r(1-r'^2)(2k_2g^2 - k_3) \sin f + 2r'(1-r'^2 - rr'')g \cos f)^2 \\ &+ 4g \begin{pmatrix} 2r(1-r'^2)k_2g \cos f \\ -r'(1-r'^2 - rr'') \sin f \end{pmatrix} \begin{pmatrix} -r(1-r'^2)k_3 \cos f \\ +r' \begin{pmatrix} r\sqrt{1-r'^2}k_1 \\ -(1-r'^2 - rr'')g \sin f \end{pmatrix} \end{pmatrix} \end{aligned} \right), \\ g_{12} = g_{21} &= \frac{r^2}{2g^2} \left(\begin{aligned} &-2(1-r'^2)k_2g^3f_t - \left(r'\sqrt{1-r'^2}k_1 \cos f - (1-r'^2)k_3 \right) gf_t \\ &- (1-r'^2)k_2g^2g_t \sin(2f) + \left(r'\sqrt{1-r'^2}k_1 - (1-r'^2)k_3 \cos f \right) g_t \sin f \end{aligned} \right), \\ g_{13} = g_{31} &= \frac{r^2}{2g^2} \left(\begin{aligned} &-2(1-r'^2)k_2g^3f_w - \left(r'\sqrt{1-r'^2}k_1 \cos f - (1-r'^2)k_3 \right) gf_w \\ &- (1-r'^2)k_2g^2g_w \sin(2f) + \left(r'\sqrt{1-r'^2}k_1 - (1-r'^2)k_3 \cos f \right) g_w \sin f \end{aligned} \right), \\ g_{22} &= \frac{r^2}{g^2} (1-r'^2) (g^2f_t^2 - g_t^2 \sin^2 f), \\ g_{23} = g_{32} &= \frac{r^2}{g^2} (1-r'^2) (g^2f_t f_w - g_t g_w \sin^2 f), \\ g_{33} &= \frac{r^2}{g^2} (1-r'^2) (g^2f_w^2 - g_w^2 \sin^2 f), \end{aligned} \right\} \quad (2.18)$$

and it follows that

$$\det[g_{ij}] = \frac{-r^4(1-r'^2) \left(r\sqrt{1-r'^2}k_1 \sin f - 2(1-r'^2 - rr'')g \right)^2 (g_w f_t - f_w g_t)^2 \sin^2 f}{4g^4}. \quad (2.19)$$

Now, for obtaining the coefficients of the second fundamental form, let us give the second derivatives $\binom{psd-n}{C}_1$ $_{x_i x_j} = \frac{\partial^2 \binom{psd-n}{C}}{\partial x_i \partial x_j}$ of the canal hypersurface (2.13):

$$\binom{psd-n}{C}_1 \Big|_{ss} = B_1 F_1 + B_2 F_2 + B_3 F_3 + B_4 F_4, \quad (2.20)$$

$$\begin{aligned} \binom{psd-n}{C}_1 \Big|_{st} &= \binom{psd-n}{C}_1 \Big|_{ts} = \frac{r\sqrt{1-r'^2}k_1(-gf_t \cos f + g_t \sin f)}{2g^2} F_1 \\ &- \frac{r(1-r'^2)k_3 f_t \sin f - r'(1-r'^2 - rr'')(gf_t \cos f + g_t \sin f)}{\sqrt{1-r'^2}} F_2 \\ &+ \frac{\left(\begin{array}{l} (r(1-r'^2)(2k_2 g^2 - k_3) \cos f - 2r'(1-r'^2 - rr'')g \sin f) g f_t \\ + r(1-r'^2)(2g^2 k_2 + k_3) g_t \sin f \end{array} \right)}{2\sqrt{1-r'^2}g^2} F_3 \\ &+ \frac{\left((2r(1-r'^2)k_2 g \sin f + r'(1-r'^2 - rr'') \cos f) g f_t - r'(1-r'^2 - rr'')g_t \sin f \right)}{2\sqrt{1-r'^2}g^2} F_4 \end{aligned} \quad (2.21)$$

$$\begin{aligned} \binom{psd-n}{C}_1 \Big|_{sw} &= \binom{psd-n}{C}_1 \Big|_{ws} = \frac{r\sqrt{1-r'^2}k_1(-gf_w \cos f + g_w \sin f)}{2g^2} F_1 \\ &+ \frac{1}{\sqrt{1-r'^2}} (-r(1-r'^2)k_3 f_w \sin f + r'(1-r'^2 - rr'')(gf_w \cos f + g_w \sin f)) F_2 \\ &+ \frac{1}{2\sqrt{1-r'^2}g^2} \left(\begin{array}{l} (r(1-r'^2)(2k_2 g^2 - k_3) \cos f - 2r'(1-r'^2 - rr'')g \sin f) g f_w \\ + r(1-r'^2)(2k_2 g^2 + k_3) g_w \sin f \end{array} \right) F_3 \\ &+ \frac{1}{2\sqrt{1-r'^2}g^2} \left(\begin{array}{l} (2r(1-r'^2)k_2 g \sin f + r'(1-r'^2 - rr'') \cos f) g f_w \\ - r'(1-r'^2 - rr'')g_w \sin f \end{array} \right) F_4 \end{aligned} \quad (2.22)$$

$$\begin{aligned} \binom{psd-n}{C}_1 \Big|_{tt} &= r\sqrt{1-r'^2} \left((2f_t g_t + g f_{tt}) \cos f + (-g f_t^2 + g_{tt}) \sin f \right) F_2 \\ &- r\sqrt{1-r'^2} (f_t^2 \cos f + f_{tt} \sin f) F_3 \\ &- \frac{r\sqrt{1-r'^2}}{2g^3} (g(2f_t g_t - g f_{tt}) \cos f + (-2g_t^2 + g(g f_t^2 + g_{tt})) \sin f) F_4 \end{aligned} \quad (2.23)$$

$$\begin{aligned} \binom{psd-n}{C}_1 \Big|_{tw} &= \binom{psd-n}{C}_1 \Big|_{wt} = r\sqrt{1-r'^2} \left((f_t g_w + f_w g_t + g f_{tw}) \cos f + (-g f_t f_w + g_{tw}) \sin f \right) F_2 \\ &- r\sqrt{1-r'^2} (f_t f_w \cos f + f_{tw} \sin f) F_3 \\ &- \frac{r\sqrt{1-r'^2}}{2g^3} (g(f_t g_w + f_w g_t - g f_{tw}) \cos f + (-2g_t g_w + g(g f_t f_w + g_{tw})) \sin f) F_4 \end{aligned} \quad (2.24)$$

and

$$\begin{aligned} \binom{psd-n}{C}_1 \Big|_{ww} &= r\sqrt{1-r'^2} \left((2f_w g_w + g f_{ww}) \cos f + (-g f_w^2 + g_{ww}) \sin f \right) F_2 \\ &- r\sqrt{1-r'^2} (f_w^2 \cos f + f_{ww} \sin f) F_3 \\ &- \frac{r\sqrt{1-r'^2}}{2g^3} (g(2f_w g_w - g f_{ww}) \cos f + (-2g_w^2 + g(g f_w^2 + g_{ww})) \sin f) F_4, \end{aligned} \quad (2.25)$$

where

$$\begin{aligned}
B_1 &= \frac{-r\sqrt{1-r'^2}k'_1 \sin f}{2g} + \frac{k_1(r(1-r'^2)k_2g \cos f - r'(1-r'^2 - rr'') \sin f)}{g\sqrt{1-r'^2}} - 3r'r'' - rr''', \\
B_2 &= \frac{-r\sqrt{1-r'^2}k_1^2 \sin f}{2g} + (1 - 2r'^2 - 2rr'') k_1 \\
&\quad - \frac{1}{2g(1-r'^2)^{3/2}} \left(\begin{aligned} &r(1-r'^2)^2 k_3^2 \sin f + 2(1-r'^2)g \left(\begin{aligned} &r r' \sqrt{1-r'^2} k'_1 \\ &- \left((rk'_3 + 2r'k_3)(1-r'^2) \right) \cos f \end{aligned} \right) \\ &- 2g^2 \left(r(1-r'^2)^2 k_2 k_3 + r''(1-4r'^2 + 3r'^4 - rr'') - rr'r'''(1-r'^2) \right) \sin f \end{aligned} \right), \\
B_3 &= -\frac{1}{2g(1-r'^2)^{3/2}} \left(\begin{aligned} &2(1-r'^2) \left(-(1-r'^2)(rk'_2 + 2r'k_2) + 2rr'r''k_2 \right) g^2 \sin f \\ &- (1-r'^2) \left(-(1-r'^2)(rk'_3 + 2r'k_3) + 2rr'r''k_3 \right) \sin f \\ &- 2g \left(\begin{aligned} &-r(1-r'^2)k_2 \left(r'\sqrt{1-r'^2}k_1 - 2(1-r'^2)k_3 \cos f \right) \\ &+ (r''(1-4r'^2 + 3r'^4 - rr'') - rr'r'''(1-r'^2)) \cos f \end{aligned} \right) \end{aligned} \right), \\
B_4 &= -\frac{1}{2g(1-r'^2)^{3/2}} \left(\begin{aligned} &-2(1-r'^2) \left(-(1-r'^2)(rk'_2 + 2r'k_2) + 2rr'r''k_2 \right) g \cos f \\ &+ \left(\begin{aligned} &2r(1-r'^2)^2 k_2^2 g^2 - r(1-r'^2)^2 k_2 k_3 \\ &- r''(1-4r'^2 + 3r'^4 - rr'') + rr'r'''(1-r'^2) \end{aligned} \right) \sin f \end{aligned} \right).
\end{aligned}$$

Thus, from (1.11), (2.17) and (2.20)-(2.25), the coefficients of the second fundamental form are given by

$$\left. \begin{aligned}
h_{11} &= \frac{-1}{4g^2(1-r'^2)} \left(\begin{aligned} &r(1-r'^2)^2 k_1^2 \sin^2 f + 4r(1-r'^2)^2 k_2^2 g^4 \sin^2 f + r(1-r'^2)^2 k_3^2 \sin^2 f \\ &- 2g^2 \left(r(1-r'^2)^2 k_2 k_3 (3 + \cos(2f)) + 2r''(1-r'^2 - rr'') \right) \\ &+ 2\sqrt{1-r'^2}k_1g(4rr'(1-r'^2)k_2g \cos f - (1-r'^2 - 2rr'') \sin f) \end{aligned} \right), \\
h_{12} = h_{21} &= \frac{-r}{2g^2} \left(\begin{aligned} &-2(1-r'^2)k_2g^3f_t - \left(r'\sqrt{1-r'^2}k_1 \cos f - (1-r'^2)k_3 \right) g f_t \\ &- (1-r'^2)k_2g^2g_t \sin(2f) + \left(r'\sqrt{1-r'^2}k_1 - (1-r'^2)k_3 \cos f \right) g_t \sin f \end{aligned} \right), \\
h_{13} = h_{31} &= \frac{-r}{2g^2} \left(\begin{aligned} &-2(1-r'^2)k_2g^3f_w - \left(r'\sqrt{1-r'^2}k_1 \cos f - (1-r'^2)k_3 \right) g f_w \\ &- (1-r'^2)k_2g^2g_w \sin(2f) + \left(r'\sqrt{1-r'^2}k_1 - (1-r'^2)k_3 \cos f \right) g_w \sin f \end{aligned} \right), \\
h_{22} &= \frac{r(1-r'^2)(g_t^2 \sin^2 f - g^2 f_t^2)}{g^2}, \\
h_{23} = h_{32} &= \frac{r(1-r'^2)(g_t g_w \sin^2 f - g^2 f_t f_w)}{g^2}, \\
h_{33} &= \frac{r(1-r'^2)(g_w^2 \sin^2 f - g^2 f_w^2)}{g^2}
\end{aligned} \right\} \quad (2.26)$$

and it implies

$$\det[h_{ij}] = \frac{r^2(1-r'^2)}{4g^4} \left(\begin{aligned} &r(1-r'^2)k_1^2 \sin^2 f - 4r''(1-r'^2 - rr'')g^2 \\ &- 2\sqrt{1-r'^2}(1-r'^2 - 2rr'')k_1g \sin f \end{aligned} \right) (f_t g_w - f_w g_t)^2 \sin^2 f. \quad (2.27)$$

So, from (1.13), (2.19) and (2.27), we obtain the Gaussian curvature $K_{C_1}^{psd-n}$ as given in (2.12).

Also, using (2.18) and (2.26) in (1.12), the shape operator of the canal hypersurface (2.13) can be obtained and so using the shape operator, we get the mean curvature $H_{C_1}^{psd-n}$ as given in (2.12).

Similarly, the Gaussian and mean curvatures $K_{C_i}^{psd-n}$ and $H_{C_i}^{psd-n}$ of the canal hypersurfaces C_i , $i \in \{2, 3, 4, 5\}$, given by (2.12) can be obtained. \square

Here, from (2.12), we can state the following theorem which gives an important relation between Gaussian and mean curvatures of the canal hypersurfaces:

Proposition 1. *The Gaussian curvatures and the mean curvatures of the canal hypersurfaces C_1 , C_3 and C_4 given by (2.1) in E_1^4 satisfy*

$$3H_{C_i}^{psd-n} - r^2 K_{C_i}^{psd-n} + \frac{2}{r} = 0, \quad i = 1, 3, 4 \quad (2.28)$$

and the Gaussian curvature and the mean curvature of the canal hypersurfaces C_2 and C_5 given by (2.1) and (2.2) in E_1^4 satisfy

$$3H_{C_i}^{psd-n} - r^2 K_{C_i}^{psd-n} - \frac{2}{r} = 0, \quad i = 2, 5. \quad (2.29)$$

Now, we will give some results for the canal hypersurface C_1 given by (2.1) in E_1^4 . Similarly, one can obtain similar results for the canal hypersurfaces C_2 , C_3 , C_4 and C_5 given by (2.1) and (2.2) in E_1^4 , too.

Proposition 2. *Let the pseudo null curve $\gamma(s)$, which generates the canal hypersurface C_1 given by (2.1) in E_1^4 , be a straight line. Then, the canal hypersurface C_1 is flat, if $r(s) = as + b$, $a, b \in \mathbb{R}$, $a \neq \pm 1$.*

Proof. If the pseudo null curve $\gamma(s)$ is a straight line, then we have $k_1(s) = 0$. So from (2.12), we have

$$K_{C_1}^{psd-n} = \frac{r''}{r^2(1 - r'^2 - rr'')}. \quad (2.30)$$

If we use $r(s) = as + b$, $a, b \in \mathbb{R}$, $a \neq \pm 1$ in (2.30), then $K_{C_1}^{psd-n}$ vanishes and this completes the proof. \square

Proposition 3. *Let the pseudo null curve $\gamma(s)$, which generates the canal hypersurface C_1 given by (2.1) in E_1^4 , not be a straight line. If $g(t, w) \neq \sin(f(t, w))$, then the canal hypersurface C_1 cannot be flat.*

Also, if $g(t, w) = \sin(f(t, w))$, then the canal hypersurface C_1 is flat when the equation

$$2(1 - r'^2) \left(\sqrt{1 - r'^2} + 2r'' \right) - r \left(1 - r'^2 + 4r'' \left(\sqrt{1 - r'^2} + r'' \right) \right) = 0 \quad (2.31)$$

holds.

Proof. If the pseudo null curve $\gamma(s)$, which generates the canal hypersurface C_1 given by (2.1) in E_1^4 , isn't a straight line, then we have $k_1(s) = 1$. So, from (2.12) we get

$$K_{C_1}^{psd-n} = \frac{-r(1 - r'^2) \sin^2 f + 4r''(1 - r'^2 - rr'')g^2 + 2\sqrt{1 - r'^2}(1 - r'^2 - 2rr'')g \sin f}{r^2 \left(r\sqrt{1 - r'^2} \sin f - 2(1 - r'^2 - rr'')g \right)^2}. \quad (2.32)$$

From (2.32), if the canal hypersurface C_1 is flat, then we have

$$-r(1 - r'^2) \sin^2 f + 4r''(1 - r'^2 - rr'')g^2 + 2\sqrt{1 - r'^2}(1 - r'^2 - 2rr'')g \sin f = 0. \quad (2.33)$$

Firstly, let us suppose $g(t, w) \neq \sin(f(t, w))$. Since the set $\{\sin^2 f, g^2, g \sin f\}$ is linearly independent, we get from (2.33) that

$$-r(1-r'^2) = 4r''(1-r'^2 - rr'') = 2\sqrt{1-r'^2}(1-r'^2 - 2rr'') = 0 \quad (2.34)$$

and this is a contradiction. Thus, C_1^{psd-n} cannot be flat in this situation and this proves the first part of this Proposition.

Secondly, if $g(t, w) = \sin(f(t, w))$ holds, then from (2.32) we have

$$K_{C_1^{psd-n}} = \frac{2(1-r'^2)\left(\sqrt{1-r'^2} + 2r''\right) - r\left(1-r'^2 + 4r''\left(\sqrt{1-r'^2} + r''\right)\right)}{r^2\left(2 - 2r'^2 - r\left(\sqrt{1-r'^2} + 2r''\right)\right)^2} \quad (2.35)$$

and this completes the proof. \square

Proposition 4. *Let the pseudo null curve $\gamma(s)$, which generates the canal hypersurface C_1^{psd-n} given by (2.1) in E_1^4 , be a straight line. Then, the canal hypersurface C_1^{psd-n} is minimal, if $r(s)$ satisfies*

$$\int \frac{dr}{\sqrt{1-\left(\frac{a}{r}\right)^4}} = \pm s + b, \quad a, b \in \mathbb{R}.$$

Proof. If the pseudo null curve $\gamma(s)$ is a straight line, from (2.12), we have

$$H_{C_1^{psd-n}} = -\frac{2 - 2r'^2 - 3rr''}{3r(1-r'^2 - rr'')}. \quad (2.36)$$

So, if the equation

$$2 - 2r'^2 - 3rr'' = 0 \quad (2.37)$$

holds, then the canal hypersurface C_1^{psd-n} is minimal. The solution of the equation (2.37) can be found in [23] and this completes the proof. \square

Proposition 5. *Let the pseudo null curve $\gamma(s)$, which generates the canal hypersurface C_1^{psd-n} given by (2.1) in E_1^4 , not be a straight line. If $g(t, w) \neq \sin(f(t, w))$, then the canal hypersurface C_1^{psd-n} cannot be minimal.*

Also, if $g(t, w) = \sin(f(t, w))$, then the canal hypersurface C_1^{psd-n} is minimal when the equation

$$8(1-r'^2)^2 - 2r(1-r'^2)\left(\sqrt{1-r'^2} + 10r''\right) - 3r^2(1-r'^2 - 4r''^2) = 0 \quad (2.38)$$

holds.

Proof. If the pseudo null curve $\gamma(s)$, which generates the canal hypersurface C_1^{psd-n} given by (2.1) in E_1^4 , isn't a straight line, then from (2.12) we get

$$H_{C_1^{psd-n}} = \frac{2r(1-r'^2)^{3/2}g \sin f + 3r^2(1-r'^2)\sin^2 f - 4(1-r'^2 - rr'')(2 - 2r'^2 - 3rr'')g^2}{3\left(-r^3(1-r'^2)\sin^2 f + 4r(1-r'^2 - rr'')^2g^2\right)}. \quad (2.39)$$

From (2.39), if the canal hypersurface C_1^{psd-n} is minimal, then we have

$$2r(1-r'^2)^{3/2}g \sin f + 3r^2(1-r'^2)\sin^2 f - 4(1-r'^2 - rr'')(2 - 2r'^2 - 3rr'')g^2 = 0. \quad (2.40)$$

Firstly, let us suppose that $g(t, w) \neq \sin(f(t, w))$. Since the set $\{g \sin f, \sin^2 f, g^2\}$ is linearly independent, we get from (2.40) that

$$2r(1-r'^2)^{3/2} = 3r^2(1-r'^2) = 4(1-r'^2 - rr'')(2 - 2r'^2 - 3rr'') = 0 \quad (2.41)$$

and this is a contradiction. Thus, C_1^{psd-n} cannot be minimal in this situation and this proves the first part of this Proposition.

Secondly, if $g(t, w) = \sin(f(t, w))$ holds, then we have

$$H_{C_1^{psd-n}} = -\frac{8(1-r'^2)^2 - 2r(1-r'^2)(\sqrt{1-r'^2} + 10r'') - 3r^2(1-r'^2 - 4r''^2)}{3r(4(1-r'^2)^2 - 8r(1-r'^2)r'' - r^2(1-r'^2 - 4r''^2))} \quad (2.42)$$

and this completes the proof. \square

Here, let us construct an example for the canal hypersurface $C_1^{psd-n}(s, t, w)$ that is formed as the envelope of a family of pseudo hyperspheres whose center lie on a pseudo null curve in E_1^4 . One can construct the canal hypersurfaces C_2^{psd-n} , C_3^{psd-n} , C_4^{psd-n} and C_5^{psd-n} , similarly.

Example 1. Let us take the pseudo null curve (given in [18])

$$\gamma(s) = \frac{1}{2\sqrt{2}} (\cosh(2s), \sinh(2s), \sin(2s), -\cos(2s)) \quad (2.43)$$

in E_1^4 . The Frenet vectors and curvatures of the curve (2.43) are

$$\left. \begin{aligned} F_1 &= \frac{1}{\sqrt{2}} (\sinh(2s), \cosh(2s), \cos(2s), \sin(2s)), \\ F_2 &= \sqrt{2} (\cosh(2s), \sinh(2s), -\sin(2s), \cos(2s)), \\ F_3 &= \frac{1}{\sqrt{2}} (\sinh(2s), \cosh(2s), -\cos(2s), -\sin(2s)), \\ F_4 &= \frac{1}{2\sqrt{2}} (-\cosh(2s), -\sinh(2s), -\sin(2s), \cos(2s)), \\ k_1 &= 1, \quad k_2 = 4, \quad k_3 = 0. \end{aligned} \right\} \quad (2.44)$$

If we assume that $g(t, w) = t$ and $f(t, w) = w$ in (2.1), the canal hypersurfaces $C_1^{psd-n}(s, t, w)$ can be parametrized by

$$C_1^{psd-n}(s, t, w) = \frac{1}{16\sqrt{2}t} \left(\begin{aligned} &\cosh(2s) (8t + \sqrt{3}s (-1 + 8t^2) \sin w) + 4st (-1 + \sqrt{3} \cos w) \sinh(2s), \\ &\sinh(2s) (8t + \sqrt{3}s (-1 + 8t^2) \sin w) + 4st (-1 + \sqrt{3} \cos w) \cosh(2s), \\ &\sin(2s) (8t - \sqrt{3}s (1 + 8t^2) \sin w) + 4st (-1 - \sqrt{3} \cos w) \cos(2s), \\ &\cos(2s) (-8t + \sqrt{3}s (1 + 8t^2) \sin w) + 4st (-1 - \sqrt{3} \cos w) \sin(2s) \end{aligned} \right) \quad (2.45)$$

where the radius function has been taken as $r(s) = \frac{s}{2}$. From (2.12), the Gaussian and mean curvatures of the canal hypersurfaces C_1^{psd-n} are obtained as

$$\left. \begin{aligned} K_{C_1^{psd-n}} &= -\frac{8}{s^2(s - 2\sqrt{3}t \csc w)}, \\ H_{C_1^{psd-n}} &= \frac{-48t^2 + 2s(2\sqrt{3}t + 3s \sin w) \sin w}{36st^2 - 3s^3 \sin^2 w}. \end{aligned} \right\} \quad (2.46)$$

In Figure 1 (a), one can see the projection of the canal hypersurfaces (2.45) for $w = \frac{\pi}{3}$ into $x_1x_3x_4$ -space.

Theorem 3. The canal hypersurfaces that are formed as the envelope of a family of pseudo hyperspheres whose centers lie on a partially null curve $\gamma(s)$ with Frenet vector fields F_i , $i \in \{1, 2, 3, 4\}$, in

E_1^4 can be parametrized by

$$\left. \begin{aligned} & \overset{\text{part-n}}{C_1}(s, t, w) = \gamma(s) - r(s)r'(s)F_1(s) \\ & \mp r(s)\sqrt{1-r'(s)^2} \left(\sin(f(t, w))F_2(s) + g(t, w)\cos(f(t, w))F_3(s) + \frac{\cos(f(t, w))}{2g(t, w)}F_4(s) \right), \\ & \overset{\text{part-n}}{C_2}(s, t, w) = \gamma(s) - r(s)r'(s)F_1(s) \\ & \mp r(s)\sqrt{1-r'(s)^2} \left(\cos(f(t, w))F_2(s) + g(t, w)\sin(f(t, w))F_3(s) + \frac{\sin(f(t, w))}{2g(t, w)}F_4(s) \right), \\ & \overset{\text{part-n}}{C_3}(s, t, w) = \gamma(s) - r(s)r'(s)F_1(s) \\ & \mp r(s)\sqrt{1-r'(s)^2} \left(\cosh(f(t, w))F_2(s) + g(t, w)\sinh(f(t, w))F_3(s) - \frac{\sinh(f(t, w))}{2g(t, w)}F_4(s) \right), \\ & \overset{\text{part-n}}{C_4}(s, t, w) = \gamma(s) - r(s)r'(s)F_1(s) \\ & \mp r(s)\sqrt{r'(s)^2 - 1} \left(\sinh(f(t, w))F_2(s) + g(t, w)\cosh(f(t, w))F_3(s) - \frac{\cosh(f(t, w))}{2g(t, w)}F_4(s) \right), \end{aligned} \right\} \quad (2.47)$$

where we suppose $r'(s)^2 < 1$ for canal hypersurfaces $\overset{\text{part-n}}{C_1}$, $\overset{\text{part-n}}{C_2}$, $\overset{\text{part-n}}{C_3}$ and $r'(s)^2 > 1$ for canal hypersurface $\overset{\text{part-n}}{C_4}$.

Furthermore, the canal hypersurfaces that are formed as the envelope of a family of pseudo hyperbolic hyperspheres whose centers lie on a partially null curve $\gamma(s)$ with Frenet vector fields F_i , $i \in \{1, 2, 3, 4\}$, in E_1^4 can be parametrized by

$$\begin{aligned} & \overset{\text{part-n}}{C_5}(s, t, w) = \gamma(s) + r(s)r'(s)F_1(s) \\ & \mp r(s)\sqrt{1+r'(s)^2} \left(\sinh(f(t, w))F_2(s) + g(t, w)\cosh(f(t, w))F_3(s) - \frac{\cosh(f(t, w))}{2g(t, w)}F_4(s) \right). \end{aligned} \quad (2.48)$$

Here, the canal hypersurfaces $\overset{\text{part-n}}{C_1}$, $\overset{\text{part-n}}{C_2}$, $\overset{\text{part-n}}{C_3}$, $\overset{\text{part-n}}{C_4}$ are timelike and the canal hypersurface $\overset{\text{part-n}}{C_5}$ is spacelike.

Proof. Let the center curve $\gamma : I \subseteq \mathbb{R} \rightarrow E_1^4$ be a partially null curve with non-zero curvature with Frenet vector fields. Then, the parametrization of the envelope of pseudo hyperspheres (or pseudo hyperbolic hyperspheres) defining the canal hypersurfaces $X(s, t, w)$ in E_1^4 can be given by

$$\overset{\text{part-n}}{C}(s, t, w) - \gamma(s) = a_1(s, t, w)F_1(s) + a_2(s, t, w)F_2(s) + a_3(s, t, w)F_3(s) + a_4(s, t, w)F_4(s). \quad (2.49)$$

Furthermore, since $C(s, t, w)$ lies on the pseudo hyperspheres ($\lambda = 1$) (or pseudo hyperbolic hyperspheres ($\lambda = -1$)), we have

$$g(\overset{\text{part-n}}{C}(s, t, w) - \gamma(s), \overset{\text{part-n}}{C}(s, t, w) - \gamma(s)) = \lambda r^2(s) \quad (2.50)$$

which leads to from (2.49) and (1.6) that

$$a_1^2 + a_2^2 + 2a_3a_4 = \lambda r^2 \quad (2.51)$$

and

$$a_1a_{1s} + a_2a_{2s} + a_4a_{2s} + a_2a_{4s} = \lambda rr_s. \quad (2.52)$$

So, differentiating (2.49) with respect to s and using the Frenet formula (1.5), we get

$$\begin{aligned} (\overset{\text{part-n}}{C})_s &= (1 - a_2k_1 + a_{1s})F_1 + (a_1k_1 - a_4k_2 + a_{2s})F_2 \\ &+ (a_2k_2 + a_3k_3 + a_{3s})F_3 + (-a_4k_3 + a_{4s})F_4. \end{aligned} \quad (2.53)$$

Furthermore, $C(s, t, w) - \gamma(s)$ is a normal vector to the canal hypersurfaces, which implies that

$$g(\overset{\text{part-n}}{C}(s, t, w) - \gamma(s), (\overset{\text{part-n}}{C})_s(s, t, w)) = 0 \quad (2.54)$$

and so, from (2.49), (2.53) and (2.55) we have

$$\begin{pmatrix} a_1(1 - a_2k_1 + a_{1s}) + a_2(a_1k_1 - a_4k_2 + a_{2s}) \\ + a_3(-a_4k_3 + a_{4s}) + a_4(a_2k_2 + a_3k_3 + a_{3s}) \end{pmatrix} = 0. \quad (2.55)$$

Using (2.52) in (2.55), we get

$$a_1 = -\lambda r r_s. \quad (2.56)$$

Hence, using (2.56) in (2.51), we reach that $a_1^2 + a_2^2 + 2a_3a_4 = \lambda r^2$

$$a_2^2 + 2a_3a_4 = \lambda r^2(1 - \lambda r_s^2). \quad (2.57)$$

Therefore from (2.56) and (2.57), the canal hypersurfaces $C^{part-n}(s, t, w)$ that are formed as the envelope of a family of pseudo hyperspheres or pseudo hyperbolic hyperspheres whose centers lie on a partially null curve in E_1^4 can be parametrized by (2.47) or (2.48), respectively. \square

Using the similar procedure with proof of the Theorem 2, we can obtain the following Theorem:

Theorem 4. *The Gaussian and mean curvatures of the canal hypersurfaces C_i^{part-n} , $i \in \{1, 2, 3, 4\}$, given by (2.47) and (2.48) in E_1^4 are*

$$\left. \begin{aligned} & \left. \begin{aligned} K_{C_1^{part-n}} &= \frac{-\tau(1-r'^2)k_1^2 \sin^2 f + r''(1-r'^2 - rr'') + \sqrt{1-r'^2}(1-r'^2 - 2rr'')k_1 \sin f}{r^2(1-r'^2 - rr'' - r\sqrt{1-r'^2}k_1 \sin f)^2} \\ H_{C_1^{part-n}} &= \frac{r(1-r'^2)^{3/2}k_1 \sin f + 3r^2(1-r'^2)k_1^2 \sin^2 f - (1-r'^2 - rr'')(2-2r'^2 - 3rr'')}{3r(-r^2(1-r'^2)k_1^2 \sin^2 f + (1-r'^2 - rr'')^2)} \end{aligned} \right\} \\ & \left. \begin{aligned} K_{C_2^{part-n}} &= \frac{r(1-r'^2)k_1^2 \cos^2 f - r''(1-r'^2 - rr'') - \sqrt{1-r'^2}(1-r'^2 - 2rr'')k_1 \cos f}{r^2(1-r'^2 - rr'' - r\sqrt{1-r'^2}k_1 \cos f)^2} \\ H_{C_2^{part-n}} &= \frac{-\tau(1-r'^2)^{3/2}k_1 \cos f - 3r^2(1-r'^2)k_1^2 \cos^2 f + (1-r'^2 - rr'')(2-2r'^2 - 3rr'')}{3r(-r^2(1-r'^2)k_1^2 \cos^2 f + (1-r'^2 - rr'')^2)} \end{aligned} \right\} \\ & \left. \begin{aligned} K_{C_3^{part-n}} &= \frac{-\tau(1-r'^2)k_1^2 \cosh^2 f + r''(1-r'^2 - rr'') + \sqrt{1-r'^2}(1-r'^2 - 2rr'')k_1 \cosh f}{r^2(1-r'^2 - rr'' - r\sqrt{1-r'^2}k_1 \cosh f)^2} \\ H_{C_3^{part-n}} &= \frac{r(1-r'^2)^{3/2}k_1 \cosh f + 3r^2(1-r'^2)k_1^2 \cosh^2 f - (1-r'^2 - rr'')(2-2r'^2 - 3rr'')}{3r(-r^2(1-r'^2)k_1^2 \cosh^2 f + (1-r'^2 - rr'')^2)} \end{aligned} \right\} \\ & \left. \begin{aligned} K_{C_4^{part-n}} &= -\frac{r(-1+r'^2)k_1^2 \sinh^2 f + r''(-1+r'^2 + rr'') + \sqrt{-1+r'^2}(-1+r'^2 + 2rr'')k_1 \sinh f}{r^2(-1+r'^2 + rr'' + r\sqrt{-1+r'^2}k_1 \sinh f)^2} \\ H_{C_4^{part-n}} &= \frac{r(-1+r'^2)^{3/2}k_1 \sinh f - 3r^2(-1+r'^2)k_1^2 \sinh^2 f + (-1+r'^2 + rr'')(-2+2r'^2 + 3rr'')}{3r(r^2(-1+r'^2)k_1^2 \sinh^2 f - (-1+r'^2 + rr'')^2)} \end{aligned} \right\} \\ & \left. \begin{aligned} K_{C_5^{part-n}} &= \frac{r(1+r'^2)k_1^2 \sinh^2 f + r''(1+r'^2 + rr'') - \sqrt{1+r'^2}(1+r'^2 + 2rr'')k_1 \sinh f}{r^2(1+r'^2 + rr'' - r\sqrt{1+r'^2}k_1 \sinh f)^2} \\ H_{C_5^{part-n}} &= \frac{r(1+r'^2)^{3/2}k_1 \sinh f + 3r^2(1+r'^2)k_1^2 \sinh^2 f - (1+r'^2 + rr'')(2+2r'^2 + 3rr'')}{3r(r^2(1+r'^2)k_1^2 \sinh^2 f - (1+r'^2 + rr'')^2)} \end{aligned} \right\}. \end{aligned} \right\} \quad (2.58)$$

Here, from (2.58), we can state the following theorem which gives an important relation between Gaussian and mean curvatures of the canal hypersurfaces:

Proposition 6. *The Gaussian curvatures and the mean curvatures of canal hypersurfaces C_1^{part-n} , C_3^{part-n} and C_4^{part-n} given by (2.47) in E_1^4 satisfy*

$$3H_{C_i^{part-n}} - r^2K_{C_i^{part-n}} + \frac{2}{r} = 0, \quad i = 1, 3, 4 \quad (2.59)$$

and the Gaussian curvature and the mean curvature of canal hypersurfaces C_2^{part-n} and C_5^{part-n} given by (2.47) and (2.48) in E_1^4 satisfy

$$3H_{C_i^{part-n}} - r^2 K_{C_i^{part-n}} - \frac{2}{r} = 0, \quad i = 2, 5. \quad (2.60)$$

Now, we will give some results for the canal hypersurface C_5^{part-n} given by (2.48) in E_1^4 . Similarly, one can obtain similar results for the canal hypersurfaces C_1^{part-n} , C_2^{part-n} , C_3^{part-n} and C_4^{part-n} given by (2.47) in E_1^4 , too.

Proposition 7. *The canal hypersurface C_5^{part-n} that is formed as the envelope of a family of pseudo hyperbolic hyperspheres whose centers lie on a partially null curve $\gamma(s)$ in E_1^4 is flat, if $k_1(s) = 0$ and the radius function satisfies $r(s) = as + b$, $a, b \in \mathbb{R}$.*

Proof. If we use $k_1(s) = 0$ in (2.58), then we have

$$K_{C_5^{part-n}} = \frac{r''}{r^2(1 + r'^2 + rr'')} \quad (2.61)$$

and this completes the proof. \square

Proposition 8. *The canal hypersurface C_5^{part-n} that is formed as the envelope of a family of pseudo hyperbolic hyperspheres whose centers lie on a partially null curve $\gamma(s)$ in E_1^4 cannot be flat when $k_1(s) \neq 0$.*

Proof. Let the canal hypersurface C_5^{part-n} be flat. Then from (2.58), we get

$$r(1 + r'^2)k_1^2 \sinh^2 f + r''(1 + r'^2 + rr'') - \sqrt{1 + r'^2}(1 + r'^2 + 2rr'')k_1 \sinh f = 0. \quad (2.62)$$

Since the set $\{\sinh f, \sinh^2 f, 1\}$ is linearly independent, we get from (2.62) that

$$r(1 + r'^2)k_1^2 = r''(1 + r'^2 + rr'') = \sqrt{1 + r'^2}(1 + r'^2 + 2rr'')k_1 = 0 \quad (2.63)$$

and this is a contradiction when $k_1(s) \neq 0$. So, this completes the proof. \square

Proposition 9. *The canal hypersurface C_5^{part-n} that is formed as the envelope of a family of pseudo hyperbolic hyperspheres whose centers lie on a partially null curve $\gamma(s)$ in E_1^4 is minimal, if $k_1(s) = 0$ and the radius function $r(s)$ satisfies $\int \frac{dr}{\sqrt{(\frac{a}{r})^{\frac{4}{3}} - 1}} = \pm s + b$, $a, b \in \mathbb{R}$.*

Proof. If we use $k_1(s) = 0$ in (2.58), then we have

$$H_{C_5^{part-n}} = \frac{2 + 2r'^2 + 3rr''}{3r(1 + r'^2 + rr'')}. \quad (2.64)$$

So, if the equation

$$2 + 2r'^2 + 3rr'' = 0 \quad (2.65)$$

holds, then the canal hypersurface C_5^{part-n} is minimal.

Now, let us solve the equation (2.65).

If we take $r'(s) = h(s)$, we get

$$r'' = h' = \frac{dh}{dr} \frac{dr}{ds} = \frac{dh}{dr} h. \quad (2.66)$$

Using (2.66) in (2.65), we have

$$3r \frac{dh}{dr} h + 2h^2 + 2 = 0. \quad (2.67)$$

From (2.65), $r'(s) = h(s) \neq 0$ and so we reach that

$$\frac{-3h}{2(1+h^2)}dh = \frac{dr}{r}. \quad (2.68)$$

By integrating (2.68), we have

$$h = \pm \sqrt{\left(\frac{a}{r}\right)^{\frac{4}{3}} - 1}, \quad (2.69)$$

where a is constant. Since $r' = \frac{dr}{ds} = h$, from (2.69) we get

$$\int \frac{dr}{\sqrt{\left(\frac{a}{r}\right)^{\frac{4}{3}} - 1}} = \pm \int ds \quad (2.70)$$

and this completes the proof. \square

Proposition 10. *The canal hypersurface C_5^{part-n} that is formed as the envelope of a family of pseudo hyperbolic hyperspheres whose centers lie on a partially null curve $\gamma(s)$ in E_1^4 cannot be minimal when $k_1(s) \neq 0$.*

Proof. *The proof can be done with similar method used in the proof of Proposition 8. \square*

Here, let us construct an example for the canal hypersurface $C_5^{part-n}(s, t, w)$ that is formed as the envelope of a family of pseudo hyperspheres whose center lie on a partially null curve in E_1^4 . One can construct the canal hypersurfaces C_1^{part-n} , C_2^{part-n} , C_3^{part-n} and C_4^{part-n} , similarly.

Example 2. *Let us take the partially null curve (given in [18])*

$$\gamma(s) = \frac{1}{2}(2e^s, 2e^s, \cos(2s), \sin(2s)) \quad (2.71)$$

in E_1^4 . The Frenet vectors and curvatures of the curve (2.43) are

$$\left. \begin{aligned} F_1 &= (e^s, e^s, -\sin(2s), \cos(2s)), \\ F_2 &= \frac{1}{2}(e^s, e^s, -2\cos(2s), -2\sin(2s)), \\ F_3 &= \frac{3}{2}(1, 1, 0, 0), \\ F_4 &= \left(-\frac{e^{2s}}{4} - \frac{1}{5}, -\frac{e^{2s}}{4} + \frac{1}{5}, \frac{e^s}{5}(\cos(2s) + 2\sin(2s)), \frac{e^s}{5}(\sin(2s) - 2\cos(2s))\right), \\ k_1 &= 2, \quad k_2 = e^s, \quad k_3 = 0. \end{aligned} \right\} \quad (2.72)$$

If we assume that $g(t, w) = t$ and $f(t, w) = w$ in (2.48), the canal hypersurfaces $C_5^{part-n}(s, t, w)$ can be parametrized by

$$C_5^{part-n}(s, t, w) = \left(\begin{aligned} &\frac{s(4+5e^{2s}+100t^2)\cosh w}{32\sqrt{5}t} + \frac{e^s}{8}(8+2s+\sqrt{5}s\sinh w), \\ &\frac{s(-4+5e^{2s}+100t^2)\cosh w}{32\sqrt{5}t} + \frac{e^s}{8}(8+2s+\sqrt{5}s\sinh w), \\ &\frac{1}{4}\left(2\cos(2s) - s\sin(2s) + \sqrt{5}s\left(-\frac{e^s\cosh w(\cos(2s)+2\sin(2s))}{10t} - \cos(2s)\sinh w\right)\right), \\ &\frac{1}{4}\left(2\sin(2s) + s\cos(2s) + \sqrt{5}s\left(-\frac{e^s\cosh w(\sin(2s)-2\cos(2s))}{10t} - \sin(2s)\sinh w\right)\right) \end{aligned} \right) \quad (2.73)$$

where the radius function has been taken as $r(s) = \frac{s}{2}$. From (2.58), the Gaussian and mean curvatures of the canal hypersurfaces C_5^{part-n} are obtained as

$$\left. \begin{aligned} K_{C_5^{part-n}} &= -\frac{16\sinh w}{s^2(\sqrt{5}-2s\sinh w)}, \\ H_{C_5^{psd-n}} &= -\frac{20-4s(\sqrt{5}+6s\sinh w)\sinh w}{3s(-5+4s^2\sinh^2 w)}. \end{aligned} \right\} \quad (2.74)$$

In Figure 1 (b), one can see the projection of the canal hypersurfaces (2.73) for $w = \frac{\pi}{3}$ into $x_1x_3x_4$ -space.

Theorem 5. *The canal hypersurfaces that are formed as the envelope of a family of pseudo hyperspheres whose centers lie on a null curve $\gamma(s)$ with Frenet vector fields F_i , $i \in \{1, 2, 3, 4\}$, in E_1^4 can be parametrized by*

$$C_1^{\text{null}}(s, t, w) = \gamma(s) + a_1(s, t, w)F_1(s) + a_2(s, t, w)F_2(s) - r(s)r'(s)F_3(s) + a_4(s, t, w)F_4(s), \quad (2.75)$$

where

$$a_2(s, t, w)^2 + a_4(s, t, w)^2 = r(s) (r(s) + 2a_1(s, t, w)r'(s)).$$

Furthermore, the canal hypersurfaces that are formed as the envelope of a family of pseudo hyperbolic hyperspheres whose centers lie on a null curve $\gamma(s)$ with Frenet vector fields F_i , $i \in \{1, 2, 3, 4\}$, in E_1^4 can be parametrized by

$$C_2^{\text{null}}(s, t, w) = \gamma(s) + a_1(s, t, w)F_1(s) + a_2(s, t, w)F_2(s) + r(s)r'(s)F_3(s) + a_4(s, t, w)F_4(s). \quad (2.76)$$

where

$$a_2(s, t, w)^2 + a_4(s, t, w)^2 = -r(s) (r(s) + 2a_1(s, t, w)r'(s)).$$

Proof. Let the center curve $\gamma : I \subseteq \mathbb{R} \rightarrow E_1^4$ be a null curve with non-zero curvature with Frenet vector fields. Then, the parametrization of the envelope of pseudo hyperspheres (or pseudo hyperbolic hyperspheres) defining the canal hypersurfaces $C^{\text{null}}(s, t, w)$ in E_1^4 can be given by

$$C^{\text{null}}(s, t, w) - \gamma(s) = a_1(s, t, w)F_1(s) + a_2(s, t, w)F_2(s) + a_3(s, t, w)F_3(s) + a_4(s, t, w)F_4(s). \quad (2.77)$$

Furthermore, since $X^{\text{null}}(s, t, w)$ lies on the pseudo hyperspheres ($\lambda = 1$) (or pseudo hyperbolic hyperspheres ($\lambda = -1$)), we have

$$g(C^{\text{null}}(s, t, w) - \gamma(s), C^{\text{null}}(s, t, w) - \gamma(s)) = \lambda r^2(s) \quad (2.78)$$

which leads to from (2.77) and (1.8) that

$$a_2^2 + a_4^2 + 2a_1a_3 = \lambda r^2 \quad (2.79)$$

and

$$a_2a_{2_s} + a_4a_{4_s} + a_1a_{3_s} + a_3a_{1_s} = \lambda rr_s. \quad (2.80)$$

So, differentiating (2.77) with respect to s and using the Frenet formula (1.7), we get

$$\begin{aligned} (C^{\text{null}})_s &= (1 + a_2k_2 - a_4k_3 + a_{1_s})F_1 + (a_1k_1 - a_3k_2 + a_{2_s})F_2 \\ &\quad + (-a_2k_1 + a_{3_s})F_3 + (a_3k_3 + a_{4_s})F_4. \end{aligned} \quad (2.81)$$

Furthermore, $C^{\text{null}}(s, t, w) - \gamma(s)$ is a normal vector to the canal hypersurfaces, which implies that

$$g(C^{\text{null}}(s, t, w) - \gamma(s), (C^{\text{null}})_s(s, t, w)) = 0 \quad (2.82)$$

and so, from (2.77), (2.81) and (2.82) we have

$$\begin{pmatrix} a_1(-a_2k_1 + a_{3_s}) + a_2(a_1k_1 - a_3k_2 + a_{2_s}) \\ + a_3(1 + a_2k_2 - a_4k_3 + a_{1_s}) + a_4(a_3k_3 + a_{4_s}) \end{pmatrix} = 0. \quad (2.83)$$

Using (2.80) in (2.83), we get

$$a_3 = -\lambda rr_s. \quad (2.84)$$

Hence, using (2.84) in (2.79), we reach that

$$a_2^2 + a_4^2 = \lambda r(r + 2a_1r_s). \quad (2.85)$$

Therefore from (2.84) and (2.85), the canal hypersurfaces $\overset{null}{C}(s, t, w)$ that are formed as the envelope of a family of pseudo hyperspheres or pseudo hyperbolic hyperspheres whose centers lie on a null curve in E_1^4 can be parametrized by (2.75) or (2.76), respectively. \square

Here, let us construct an example for the canal hypersurface $\overset{null}{C}_1(s, t, w)$ that is formed as the envelope of a family of pseudo hyperspheres whose center lie on a null curve in E_1^4 . One can construct the canal hypersurface $\overset{null}{C}_2$, similarly.

Example 3. Let us take the null curve (given in [1])

$$\gamma(s) = \frac{1}{\sqrt{2}} (\sinh s, \cosh s, \sin s, \cos s) \quad (2.86)$$

in E_1^4 . The Frenet vectors and curvatures of the curve (2.43) are

$$\left. \begin{aligned} F_1 &= \frac{1}{\sqrt{2}} (\cosh s, \sinh s, \cos s, -\sin s), \\ F_2 &= \frac{1}{\sqrt{2}} (\sinh s, \cosh s, -\sin s, -\cos s), \\ F_3 &= \frac{1}{\sqrt{2}} (-\cosh s, -\sinh s, \cos s, -\sin s), \\ F_4 &= \frac{1}{\sqrt{2}} (\sinh s, \cosh s, \sin s, \cos s), \\ k_1 &= 1, \quad k_2 = 0, \quad k_3 = -1. \end{aligned} \right\} \quad (2.87)$$

If we assume that $g(t, w) = t$ and $f(t, w) = w$ in (2.48), the canal hypersurfaces $\overset{null}{C}_1(s, t, w)$ can be parametrized by

$$\overset{null}{C}_1(s, t, w) = \frac{1}{8\sqrt{2}t} \begin{pmatrix} -2st(1 + \sqrt{3}\cos w)\cosh s + (8t + \sqrt{3}s(1 + 2t^2)\sin w)\sinh s, \\ -2st(1 + \sqrt{3}\cos w)\sinh s + (8t + \sqrt{3}s(1 + 2t^2)\sin w)\cosh s, \\ -2st(1 - \sqrt{3}\cos w)\cos s + (8t + \sqrt{3}s(1 - 2t^2)\sin w)\sin s, \\ -2st(-1 + \sqrt{3}\cos w)\sin s + (8t + \sqrt{3}s(1 - 2t^2)\sin w)\cos s \end{pmatrix} \quad (2.88)$$

where the radius function has been taken as $r(s) = \frac{s}{2}$.

In Figure 1 (c), one can see the projection of the canal hypersurfaces (2.88) for $w = \frac{\pi}{3}$ into $x_1x_3x_4$ -space.




FIGURE 1

3. TUBULAR HYPERSURFACES GENERATED BY PSEUDO-NULL, PARTIALLY NULL AND NULL CURVES IN E_1^4

In this section, by taking the radius function $r(s) = r$ is constant in the Section 2, firstly we give the parametric expressions of the tubular hypersurfaces which are formed as the envelope of a family of pseudo hyperspheres or pseudo hyperbolic hyperspheres whose centers lie on a pseudo-null curve, partially null curve or null curve in E_1^4 . After that, we obtain some results for Weingarten tubular hypersurfaces with the aid of the Gaussian curvatures and mean curvatures of these tubular hypersurfaces.

Theorem 6. *The tubular hypersurfaces that are formed as the envelope of a family of pseudo hyperspheres whose centers lie on pseudo null curves with Frenet vector fields F_i , $i \in \{1, 2, 3, 4\}$, in E_1^4 can be parametrized by*

$$\left. \begin{aligned} T_1^{psd-n}(s, t, w) &= \gamma(s) \mp r \left(g(t, w) \sin(f(t, w)) F_2(s) + \cos(f(t, w)) F_3(s) + \frac{\sin(f(t, w))}{2g(t, w)} F_4(s) \right), \\ T_2^{psd-n}(s, t, w) &= \gamma(s) \mp r \left(g(t, w) \cos(f(t, w)) F_2(s) + \sin(f(t, w)) F_3(s) + \frac{\cos(f(t, w))}{2g(t, w)} F_4(s) \right), \\ T_3^{psd-n}(s, t, w) &= \gamma(s) \mp r \left(g(t, w) \sinh(f(t, w)) F_2(s) + \cosh(f(t, w)) F_3(s) - \frac{\sinh(f(t, w))}{2g(t, w)} F_4(s) \right). \end{aligned} \right\} \quad (3.1)$$

Furthermore, the tubular hypersurfaces that are formed as the envelope of a family of pseudo hyperbolic hyperspheres whose centers lie on pseudo null curves with Frenet vector fields F_i , $i \in \{1, 2, 3, 4\}$, in E_1^4 can be parametrized by

$$T_4^{psd-n}(s, t, w) = \gamma(s) \mp r \left(g(t, w) \cosh(f(t, w)) F_2(s) + \sinh(f(t, w)) F_3(s) - \frac{\cosh(f(t, w))}{2g(t, w)} F_4(s) \right). \quad (3.2)$$

Theorem 7. *The Gaussian and mean curvatures of the tubular hypersurfaces T_i^{psd-n} , $i \in \{1, 2, 3, 4\}$, given by (3.1) and (3.2) in E_1^4 are*

$$\left. \begin{aligned} K_{T_1^{psd-n}} &= \frac{k_1}{r^2(2g \csc f - rk_1)}, & H_{T_1^{psd-n}} &= \frac{1}{-r + \frac{2rg}{-4g + 3rk_1 \sin f}}; \\ K_{T_2^{psd-n}} &= \frac{k_1}{r^2(rk_1 - 2g \sec f)}, & H_{T_2^{psd-n}} &= \frac{1}{r + \frac{2rg}{4g - 3rk_1 \cos f}}; \\ K_{T_3^{psd-n}} &= \frac{-k_1}{r^2(2g \csc h f + rk_1)}, & H_{T_3^{psd-n}} &= \frac{1}{-r + \frac{2rg}{-4g - 3rk_1 \sinh f}}; \\ K_{T_4^{psd-n}} &= \frac{k_1}{r^2(2g \operatorname{sech} f + rk_1)}, & H_{T_4^{psd-n}} &= \frac{1}{r + \frac{2rg}{4g + 3rk_1 \cosh f}}. \end{aligned} \right\} \quad (3.3)$$

So from (3.3), we can give the following results:

Proposition 11. i) *If the pseudo null curve $\gamma(s)$, which generates the tubular hypersurface T_1^{psd-n} given by (3.1) in E_1^4 , is a straight line, then the tubular hypersurface is flat.*

ii) *Let the pseudo null curve $\gamma(s)$, which generates the tubular hypersurface T_1^{psd-n} given by (3.1) in E_1^4 , not be a straight line. If $g(t, w) = \sin(f(t, w))$, then the Gaussian curvature is constant with $\frac{1}{r^2(2-r)}$.*

Proposition 12. i) *The tubular hypersurface T_1^{psd-n} given by (3.1) in E_1^4 cannot be minimal.*

ii) *If the pseudo null curve $\gamma(s)$, which generates the tubular hypersurface T_1^{psd-n} given by (3.1) in E_1^4 , is a straight line, then the mean curvature of the tubular hypersurface T_1^{psd-n} given by (3.1) in E_1^4 is constant with $\frac{-2}{3r}$.*

iii) Let the pseudo null curve $\gamma(s)$, which generates the tubular hypersurface T_1^{psd-n} given by (3.1) in E_1^4 , not be a straight line. If $g(t, w) = \sin(f(t, w))$, then the mean curvature of the tubular hypersurface T_1^{psd-n} given by (3.1) in E_1^4 is constant with $\frac{3r-4}{3r(2-r)}$, $2 \neq r \neq \frac{4}{3}$.

Also, if

$$\left. \begin{aligned} \begin{pmatrix} H_{T_i^{psd-n}} \\ H_{T_i^{psd-n}} \\ H_{T_i^{psd-n}} \end{pmatrix}_s \begin{pmatrix} K_{T_i^{psd-n}} \\ K_{T_i^{psd-n}} \\ K_{T_i^{psd-n}} \end{pmatrix}_t - \begin{pmatrix} H_{T_i^{psd-n}} \\ H_{T_i^{psd-n}} \\ H_{T_i^{psd-n}} \end{pmatrix}_t \begin{pmatrix} K_{T_i^{psd-n}} \\ K_{T_i^{psd-n}} \\ K_{T_i^{psd-n}} \end{pmatrix}_s &= 0, \\ \begin{pmatrix} H_{T_i^{psd-n}} \\ H_{T_i^{psd-n}} \\ H_{T_i^{psd-n}} \end{pmatrix}_s \begin{pmatrix} K_{T_i^{psd-n}} \\ K_{T_i^{psd-n}} \\ K_{T_i^{psd-n}} \end{pmatrix}_w - \begin{pmatrix} H_{T_i^{psd-n}} \\ H_{T_i^{psd-n}} \\ H_{T_i^{psd-n}} \end{pmatrix}_w \begin{pmatrix} K_{T_i^{psd-n}} \\ K_{T_i^{psd-n}} \\ K_{T_i^{psd-n}} \end{pmatrix}_s &= 0, \\ \begin{pmatrix} H_{T_i^{psd-n}} \\ H_{T_i^{psd-n}} \\ H_{T_i^{psd-n}} \end{pmatrix}_t \begin{pmatrix} K_{T_i^{psd-n}} \\ K_{T_i^{psd-n}} \\ K_{T_i^{psd-n}} \end{pmatrix}_w - \begin{pmatrix} H_{T_i^{psd-n}} \\ H_{T_i^{psd-n}} \\ H_{T_i^{psd-n}} \end{pmatrix}_w \begin{pmatrix} K_{T_i^{psd-n}} \\ K_{T_i^{psd-n}} \\ K_{T_i^{psd-n}} \end{pmatrix}_t &= 0 \end{aligned} \right\} \quad (3.4)$$

hold on a hypersurface, then we call the hypersurface as $\left(H_{T_i^{psd-n}}, K_{T_i^{psd-n}} \right)_{st}$ -Weingarten, $\left(H_{T_i^{psd-n}}, K_{T_i^{psd-n}} \right)_{sw}$ -

Weingarten, $\left(H_{T_i^{psd-n}}, K_{T_i^{psd-n}} \right)_{tw}$ -Weingarten hypersurface, respectively, where $\left(H_{T_i^{psd-n}} \right)_s = \frac{\partial \left(H_{T_i^{psd-n}} \right)}{\partial s}$ and so on. Thus,

Theorem 8. The tubular hypersurfaces T_i^{psd-n} , $i \in \{1, 2, 3, 4\}$, given by (3.1) and (3.2) in E_1^4 , are $\left(H_{T_i^{psd-n}}, K_{T_i^{psd-n}} \right)_{st}$ -Weingarten, $\left(H_{T_i^{psd-n}}, K_{T_i^{psd-n}} \right)_{sw}$ -Weingarten and $\left(H_{T_i^{psd-n}}, K_{T_i^{psd-n}} \right)_{tw}$ -Weingarten.

Proof. Let us give the proof for $i = 1$. From (3.3), we have

$$\left. \begin{aligned} \begin{pmatrix} H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \end{pmatrix}_s &= \frac{2k'_1 g \sin f}{3(-2g+rk_1 \sin f)^2} & \text{and} & \begin{pmatrix} K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \end{pmatrix}_s &= \frac{2k'_1 g \sin f}{r^2(-2g+rk_1 \sin f)^2} \\ \begin{pmatrix} H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \end{pmatrix}_t &= \frac{2k_1(gf_t \cos f - g_t \sin f)}{3(-2g+rk_1 \sin f)^2} & & \begin{pmatrix} K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \end{pmatrix}_t &= \frac{2k_1(gf_t \cos f - g_t \sin f)}{r^2(-2g+rk_1 \sin f)^2} \\ \begin{pmatrix} H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \end{pmatrix}_w &= \frac{2k_1(gf_w \cos f - g_w \sin f)}{3(-2g+rk_1 \sin f)^2} & & \begin{pmatrix} K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \end{pmatrix}_w &= \frac{2k_1(gf_w \cos f - g_w \sin f)}{r^2(-2g+rk_1 \sin f)^2}, \end{aligned} \right\} \quad (3.5)$$

where $f_t = \frac{\partial f(t, w)}{\partial t}$, $f_w = \frac{\partial f(t, w)}{\partial w}$ and so on.

So, from (3.5) we get

$$\left. \begin{aligned} \begin{pmatrix} H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \end{pmatrix}_s \begin{pmatrix} K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \end{pmatrix}_t - \begin{pmatrix} H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \end{pmatrix}_t \begin{pmatrix} K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \end{pmatrix}_s &= 0, \\ \begin{pmatrix} H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \end{pmatrix}_s \begin{pmatrix} K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \end{pmatrix}_w - \begin{pmatrix} H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \end{pmatrix}_w \begin{pmatrix} K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \end{pmatrix}_s &= 0, \\ \begin{pmatrix} H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \end{pmatrix}_t \begin{pmatrix} K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \end{pmatrix}_w - \begin{pmatrix} H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \\ H_{T_1^{psd-n}} \end{pmatrix}_w \begin{pmatrix} K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \\ K_{T_1^{psd-n}} \end{pmatrix}_t &= 0 \end{aligned} \right\} \quad (3.6)$$

and this completes the proof. Similarly, one can obtain the results for $i = 2, 3, 4$. \square

Theorem 9. The tubular hypersurfaces that are formed as the envelope of a family of pseudo hyperspheres whose centers lie on partially null curves with Frenet vector fields F_i , $i \in \{1, 2, 3, 4\}$, in E_1^4 can be parametrized by

$$\left. \begin{aligned} T_1^{part-n}(s, t, w) &= \gamma(s) \mp r \left(\sin(f(t, w))F_2(s) + g(t, w) \cos(f(t, w))F_3(s) + \frac{\cos(f(t, w))}{2g(t, w)}F_4(s) \right), \\ T_2^{part-n}(s, t, w) &= \gamma(s) \mp r \left(\cos(f(t, w))F_2(s) + g(t, w) \sin(f(t, w))F_3(s) + \frac{\sin(f(t, w))}{2g(t, w)}F_4(s) \right), \\ T_3^{part-n}(s, t, w) &= \gamma(s) \mp r \left(\cosh(f(t, w))F_2(s) + g(t, w) \sinh(f(t, w))F_3(s) - \frac{\sinh(f(t, w))}{2g(t, w)}F_4(s) \right), \end{aligned} \right\} \quad (3.7)$$

Furthermore, the tubular hypersurfaces that are formed as the envelope of a family of pseudo hyperbolic hyperspheres whose centers lie on partially null curves with Frenet vector fields F_i , $i \in \{1, 2, 3, 4\}$, in E_1^4 can be parametrized by

$$T_4^{part-n}(s, t, w) = \gamma(s) \mp r \left(\sinh(f(t, w))F_2(s) + g(t, w) \cosh(f(t, w))F_3(s) - \frac{\cosh(f(t, w))}{2g(t, w)}F_4(s) \right). \quad (3.8)$$

Theorem 10. The Gaussian and mean curvatures of the tubular hypersurfaces T_i^{part-n} , $i \in \{1, 2, 3, 4\}$, given by (3.7) and (3.8) in E_1^4 are

$$\left. \begin{aligned} K_{T_1}^{part-n} &= \frac{k_1}{r^2(\csc f - rk_1)}, & H_{T_1}^{part-n} &= \frac{1}{-r + \frac{r}{-2+3rk_1 \sin f}}; \\ K_{T_2}^{part-n} &= \frac{-k_1}{r^2(\sec f - rk_1)}, & H_{T_2}^{part-n} &= \frac{-1}{-r + \frac{r}{-2+3rk_1 \cos f}}; \\ K_{T_3}^{part-n} &= \frac{k_1}{r^2(\operatorname{sech} f - rk_1)}, & H_{T_3}^{part-n} &= \frac{1}{-r + \frac{r}{-2+3rk_1 \cosh f}}; \\ K_{T_4}^{part-n} &= \frac{-k_1}{r^2(\operatorname{csch} f - rk_1)}, & H_{T_4}^{part-n} &= \frac{-1}{-r + \frac{r}{-2+3rk_1 \sinh f}}. \end{aligned} \right\} \quad (3.9)$$

Thus,

Proposition 13. If $k_1(s) = 0$ for the partially null curve $\gamma(s)$, which generates the tubular hypersurfaces T_i^{part-n} given by (3.7) and (3.8) in E_1^4 , then these tubular hypersurfaces are flat and also, the tubular hypersurfaces T_i^{part-n} have constant mean curvature with $\frac{2(-1)^i}{3r}$.

Theorem 11. The tubular hypersurfaces T_i^{part-n} , $i \in \{1, 2, 3, 4\}$, given by (3.7) and (3.8) in E_1^4 , are $\left(H_{T_i}^{part-n}, K_{T_i}^{part-n} \right)_{st}$ -Weingarten, $\left(H_{T_i}^{part-n}, K_{T_i}^{part-n} \right)_{sw}$ -Weingarten and $\left(H_{T_i}^{part-n}, K_{T_i}^{part-n} \right)_{tw}$ -Weingarten.

Proof. Here we will give the proof for $i = 3$. From (3.9), we have

$$\left. \begin{aligned} \left(H_{T_3}^{part-n} \right)_s &= \frac{k_1' \cosh f}{3(-1+rk_1 \cosh f)^2} & \left(K_{T_3}^{part-n} \right)_s &= \frac{k_1' \cosh f}{r^2(-1+rk_1 \cosh f)^2} \\ \left(H_{T_3}^{part-n} \right)_t &= \frac{k_1 f_t \sinh f}{3(-1+rk_1 \cosh f)^2} & \left(K_{T_3}^{part-n} \right)_t &= \frac{k_1 f_t \sinh f}{r^2(-1+rk_1 \cosh f)^2} \\ \left(H_{T_3}^{part-n} \right)_w &= \frac{k_1 f_w \sinh f}{3(-1+rk_1 \cosh f)^2} & \left(K_{T_3}^{part-n} \right)_w &= \frac{k_1 f_w \sinh f}{r^2(-1+rk_1 \cosh f)^2}. \end{aligned} \right\} \quad (3.10)$$

So, from (3.10) we get

$$\left. \begin{aligned} \left(H_{T_3}^{part-n} \right)_s \left(K_{T_3}^{part-n} \right)_t - \left(H_{T_3}^{part-n} \right)_t \left(K_{T_3}^{part-n} \right)_s &= 0, \\ \left(H_{T_3}^{part-n} \right)_s \left(K_{T_3}^{part-n} \right)_w - \left(H_{T_3}^{part-n} \right)_w \left(K_{T_3}^{part-n} \right)_s &= 0, \\ \left(H_{T_3}^{part-n} \right)_t \left(K_{T_3}^{part-n} \right)_w - \left(H_{T_3}^{part-n} \right)_w \left(K_{T_3}^{part-n} \right)_t &= 0 \end{aligned} \right\} \quad (3.11)$$

and this completes the proof. Similarly, one can obtain the results for $i = 1, 2, 4$. \square

Theorem 12. The tubular hypersurfaces that are formed as the envelope of a family of pseudo hyperspheres whose centers lie on a null curve $\gamma(s)$ with Frenet vector fields F_i , $i \in \{1, 2, 3, 4\}$, in E_1^4 can be parametrized by

$$T_1^{null}(s, t, w) = \gamma(s) + a_1(s, t, w)F_1(s) + a_2(s, t, w)F_2(s) + a_4(s, t, w)F_4(s), \quad (3.12)$$

where

$$a_2(s, t, w)^2 + a_4(s, t, w)^2 = r^2.$$

There is no tubular hypersurfaces that are formed as the envelope of a family of pseudo hyperbolic hyperspheres whose centers lie on a null curve $\gamma(s)$ with Frenet vector fields F_i , $i \in \{1, 2, 3, 4\}$, in E_1^4 .

REFERENCES

- [1] F.K. Aksoyak, İ. Gök and K. İlarşlan; *Generalized Null Bertrand Curves in Minkowski Space-Time*, Ann. Alexandru Ioan Cuza Univ. Math., 60, (2014), 489-502.
- [2] M. Akyığıt, K. Eren and H.H. Kosal; *Tubular Surfaces with Modified Orthogonal Frame in Euclidean 3-Space*, Honam Mathematical J., 43(3), (2021), 453–463.
- [3] M. Altın, *Rotational hypersurfaces generated by weighted mean curvature in E_1^4 with density*, Thermal Science, 26(4A), (2022), 3043-3053.
- [4] M. Altın, A. Kazan and H.B. Karadağ; *Monge Hypersurfaces in Euclidean 4-Space with Density*, Journal of Polytechnic, 23(1), (2020), 207-214.
- [5] M. Altın, A. Kazan, and D.W. Yoon; *2-Ruled hypersurfaces in Euclidean 4-space*, Journal of Geometry and Physics, 166, (2021), 104236.
- [6] M. Altın and A. Kazan; *Rotational Hypersurfaces in Lorentz-Minkowski 4-Space*, Hacettepe Journal of Mathematics and Statistics, 50(5), (2021), 1409-1433.
- [7] S. Aslan and Y. Yaylı; *Canal Surfaces with Quaternions*, Adv. Appl. Clifford Algebr., 26, (2016), 31-38.
- [8] M.E. Aydın and I. Mihai; *On certain surfaces in the isotropic 4-space*, Mathematical Communications, 22(1), (2017), 41-51.
- [9] W.B. Bonnor; *Null curves in a Minkowski space-time*, Tensor, 20, (1969), 229-242.
- [10] W.B. Bonnor; *Curves with null normals in Minkowski space-time*, A random walk in relativity and cosmology, Wiley Easten Limitid, (1985), 33-47.
- [11] F. Doğan and Y. Yaylı; *Tubes with Darboux Frame*, Int. J. Contemp.Math. Sci., 7(16), (2012), 751-758.
- [12] X. Fu, S.D. Jung, J. Qian and M. Su; *Geometric Characterizations of Canal Surfaces in Minkowski 3-space I*, Bull. Korean Math. Soc., 56(4), (2019), 867-883.
- [13] R. Garcia, J. Llibre and J. Sotomayor; *Lines of Principal Curvature on Canal Surfaces in \mathbb{R}^3* , An. Acad. Brasil. Cienc., 78(3), (2006), 405-415.
- [14] E. Güler; *Helical Hypersurfaces in Minkowski Geometry E_1^4* , Symmetry, 12, (2020), 1206.
- [15] E. Hartman; *Geometry and Algorithms for Computer Aided Design*, Dept. of Math. Darmstadt Univ. of Technology, 2003.
- [16] S. Izumiya and M. Takahashi; *On caustics of submanifolds and canal hypersurfaces in Euclidean space*, Topology Appl., 159, (2012), 501-508.
- [17] K. İlarşlan and E. Nesovic; *Some Characterizations of Null, Pseudo Null and Partially Null Rectifying Curves in Minkowski Space-Time*, Taiwanese Journal of Mathematics, 12(5), (2008), 1035-1044.
- [18] K. İlarşlan and E. Nesovic; *Some Characterizations of Pseudo and Partially Null Osculating Curves in Minkowski Space-Time*, International Electronic Journal of Geometry, 4(2), (2011), 1-12.
- [19] M.K. Karacan, H. Es and Y. Yaylı; *Singular Points of Tubular Surfaces in Minkowski 3-Space*, Sarajevo J. Math., 2(14), (2006), 73-82.
- [20] M.K. Karacan and Y. Tuncer; *Tubular Surfaces of Weingarten Types in Galilean and Pseudo-Galilean*, Bull. Math. Anal. Appl., 5(2), (2013), 87-100.
- [21] M.K. Karacan, D.W. Yoon and Y. Tuncer; *Tubular Surfaces of Weingarten Types in Minkowski 3-Space*, Gen. Math. Notes, 22(1), (2014), 44-56.
- [22] M.K. Karacan and B. Bukcu; *An alternative moving frame for a tubular surface around a spacelike curve with a spacelike normal in Minkowski 3-space* Rendiconti del Circolo Matematico di Palermo, 57, (2008), 193-201.
- [23] A.Kazan, M. Altın and D.W. Yoon; *Geometric Characterizations of Canal Hypersurfaces in Euclidean Spaces*, arXiv:2111.04448v1 [math.DG], (2021).
- [24] A.Kazan, M. Altın and D.W. Yoon; *Canal Hypersurfaces generated by Non-Null Curves in Lorentz-Minkowski 4-Space*, arXiv:2206.12572v1 [math.DG], (2022).
- [25] A.Kazan; *Canal Hypersurfaces according to one of the Extended Darboux Frame Field in Euclidean 4-Space*, Thermal Science, 26(4A), (2022), 3029-3041.
- [26] S. Kızıltuğ, M. Dede and C. Ekici; *Tubular Surfaces with Darboux Frame in Galilean 3-Space*, Facta Universitatis Ser. Math. Inform., 34(2), (2019), 253-260.
- [27] Y.H. Kim, H. Liu and J. Qian; *Some Characterizations of Canal Surfaces*, Bull. Korean Math. Soc., 53(2), (2016), 461-477.
- [28] İ. Kişi, G. Öztürk and K. Arslan; *A new type of canal surface in Euclidean 4-space IE^4* , Sakarya University Journal of Science, 23(5), (2019), 801-809.

- [29] S.N. Krivoschapko and C.A.B. Hyeng; *Classification of Cyclic Surfaces and Geometrical Research of Canal Surfaces*, International Journal of Research and Reviews in Applied Sciences, 12(3), (2012), 360-374.
- [30] W. Kuhnel; *Differential geometry: curves-surfaces-manifolds*. American Mathematical Soc., Braunschweig, Wiesbaden, 1999.
- [31] J.M. Lee; *Riemannian Manifolds-An Introduction to Curvature*, Springer-Verlag New York, Inc, 1997.
- [32] T. Maekawa, N.M. Patrikalakis, T. Sakkalis and G. Yu; *Analysis and Applications of Pipe Surfaces*, Comput. Aided Geom. Design, 15, (1998), 437-458.
- [33] B. O'Neill; *Semi-Riemannian geometry with applications to relativity*, Academic Press, New York, 1983.
- [34] M. Peternell and H. Pottmann; *Computing Rational Parametrizations of Canal Surfaces*, J. Symbolic Comput., 23, (1997), 255-266.
- [35] J.S. Ro and D.W. Yoon; *Tubes of Weingarten Type in a Euclidean 3-Space*, Journal of the Chungcheong Mathematical Society, 22(3), (2009), 359-366.
- [36] A. Uçum and K. İlarıslan; *New Types of Canal Surfaces in Minkowski 3-Space*, Adv. Appl. Clifford Algebr., 26, (2016), 449-468.
- [37] J.Walrave; *Curves and surfaces in Minkowski space*, Dissertation, K. U. Leuven, Fac. of Science, Leuven, 1995.
- [38] Z. Xu, R. Feng and J-G. Sun; *Analytic and Algebraic Properties of Canal Surfaces*, J. Comput. Appl. Math., 195, (2006), 220-228.
- [39] D.W. Yoon and Z.K. Yüzbaşı; *Tubular Surfaces with Galilean Darboux Frame in G_3* , Journal of Mathematical Physics, Analysis, Geometry, 15(2), (2019), 278-287.

, ¹TECHNICAL SCIENCES VOCATIONAL SCHOOL, BINGÖL UNIVERSITY, BINGÖL, TURKEY,, ²DEPARTMENT OF COMPUTER TECHNOLOGIES, DOĞANŞEHİR VAHAP KÜÇÜK VOCATIONAL SCHOOL, MALATYA TURGUT ÖZAL UNIVERSITY, MALATYA, TURKEY,, ³DEPARTMENT OF MATHEMATICS EDUCATION AND RINS, GYEONGSANG NATIONAL UNIVERSITY, JINJU 52828, REPUBLIC OF KOREA., *CORRESPONDING AUTHOR: MALTIN@BINGOL.EDU.TR