

Linear Time-Varying Dynamic-Algebraic Equations of Index One on Time Scales

Svetlin G. Georgiev* and Sergey Kryzhevich†

October 3, 2022

Abstract

In this paper, we introduce a class of linear time-varying dynamic-algebraic equations (LTVDAE) of tractability index one on arbitrary time scales. We propose a procedure for the decoupling of the considered class LTVDAE. A projector approach is used to prove the main statement of the paper.

Keywords: decoupling, time scales, linear systems, projectors.

1 Introduction

The time scale dynamics, first introduced by Aulbach and Hilger [1], became an important part of the modern theory of dynamical systems. They appear in biological systems, mechanical systems, describing dynamics with impulse interactions, sociological models, etc. Roughly speaking, those systems combine features of ordinary differential equations and discrete dynamical systems. A brief survey on time scale calculus and time scale dynamics is performed in the book [2], see also [3], [4], [7], references therein, and Section 2 of this paper.

In this paper, we study the problem of decoupling linear systems, investigating the LTVDAE

$$A^\sigma(t)(Bx)^\Delta(t) = C^\sigma(t)x^\sigma(t) + f(t), \quad t \in I, \quad (1.1)$$

where $A : I \rightarrow \mathcal{M}_{n \times m}$, $B : I \rightarrow \mathcal{M}_{m \times n}$, $C : I \rightarrow \mathcal{M}_{n \times n}$, $f : I \rightarrow \mathbb{R}^n$ are given, $x : I \rightarrow \mathbb{R}^n$ is unknown, $I \subseteq \mathbb{T}$, \mathbb{T} is a time scale with forward jump operator and delta differentiation operator σ and Δ , respectively. Here, with $\mathcal{M}_{p \times q}$ we denote the space of C^1 -smooth $p \times q$ matrices with real entries. More precisely, we give conditions for A , B , C , and f under which we construct projectors and matrix chains ensuring decoupling of the LTVDAE (1.1). To our knowledge, no investigations were performed devoted to LTVDAE on arbitrary time scales.

The paper is organized as follows. In the next section, we give some basic facts from time scale calculus. In Section 3, we introduce P -projectors and matrix chains and we deduct some of their properties. In Section 4, we consider a particular case. In Section 5, we investigate standard form index one problems. In Section 6, we give a procedure for decoupling of the equation (1.1) and provide the main result of our paper. In Section 7, we give some examples, illustrating the obtained results. A conclusion is made in Section 8.

*Sorbonne University, Department of Mathematics, Paris, France

†Gdańsk University of Technology, Faculty of Applied Physics and Mathematics, Gdańsk, Poland, serkryzh@pg.edu.pl

2 Time Scales Essentials

In this paper, a time scale denoted by the symbol \mathbb{T} , is any closed non-empty subset of \mathbb{R} .

We suppose that a time scale \mathbb{T} has the topology that inherits from the real numbers with the standard topology.

Definition 2.1. *We say that a function is rd-continuous if it is continuous in that inherited topology.*

Also, in this section only, we assume for simplicity that either $\sup \mathbb{T} = +\infty$ or $\sup \mathbb{T}$ is not an isolated point.

We set $\inf \emptyset = \sup \mathbb{T}$ and $\sup \emptyset = \inf \mathbb{T}$.

Definition 2.2. *For $t \in \mathbb{T}$ we define the forward jump operator $\sigma : \mathbb{T} \mapsto \mathbb{T}$ as follows $\sigma(t) = \inf\{s \in \mathbb{T} : s > t\}$. We note that $\sigma(t) \geq t$ for any $t \in \mathbb{T}$.*

Definition 2.3. *We define the graininess function by the formula $\mu(t) = \sigma(t) - t$. The point is called right-dense if $\mu(t) = 0$ and right-scattered otherwise.*

Definition 2.4. *If $f : \mathbb{T} \mapsto \mathbb{R}$ is a function, then we define $f^\sigma : \mathbb{T} \mapsto \mathbb{R}$ by $f^\sigma(t) = f(\sigma(t))$ for any $t \in \mathbb{T}$, i.e., $f^\sigma = f \circ \sigma$.*

Here is the definition of a segment of a time scale.

Definition 2.5. *Let $a, b \in \mathbb{T}$, $a \leq b$. We define the interval $[a, b]$ in \mathbb{T} by*

$$[a, b] = \{t \in \mathbb{T} : a \leq t \leq b\}.$$

Open intervals, half-open intervals, half-lines and so on are defined accordingly.

Definition 2.6. *Assume that $f : \mathbb{T} \mapsto \mathbb{R}$ is a rd-continuous function and let $t \in \mathbb{T}$. We define $f^\Delta(t)$ to be the number, provided it exists, as follows: for any $\varepsilon > 0$ there is a neighborhood U of t , $U = (t - \delta, t + \delta) \cap \mathbb{T}$ for some $\delta > 0$, such that*

$$|f(\sigma(t)) - f(s) - f^\Delta(t)(\sigma(t) - s)| \leq \varepsilon |\sigma(t) - s| \quad \text{for all } s \in U, s \neq \sigma(t).$$

We say that $f^\Delta(t)$ is the delta or Hilger derivative of f at t .

We say that f is delta or Hilger differentiable, shortly differentiable, in \mathbb{T} if $f^\Delta(t)$ exists for all $t \in \mathbb{T}$. The function $f^\Delta : \mathbb{T} \mapsto \mathbb{R}$ is said to be delta derivative or Hilger derivative, shortly derivative, of f in \mathbb{T} .

If $\mathbb{T} = \mathbb{R}$, then the delta derivative coincides with the classical derivative, if $\mathbb{T} = \mathbb{N}$, this is just the increment $f(n+1) - f(n)$.

We list some basic properties of Hilger derivatives.

Theorem 2.1. *Let $f : \mathbb{T} \mapsto \mathbb{R}$ is a function and let $t \in \mathbb{T}$. Then the following holds.*

1. *If f is differentiable at t , then f is continuous at t .*
2. *If f is continuous at t and t is right-scattered, then f is differentiable at t with*

$$f^\Delta(t) = \frac{f(\sigma(t)) - f(t)}{\mu(t)}.$$

3. If t is right-dense, then f is differentiable if and only if the limit

$$f^\Delta(t) := \lim_{s \rightarrow t} \frac{f(t) - f(s)}{t - s}$$

exists and is finite.

4. If f is differentiable at t , then $f(\sigma(t)) = f(t) + \mu(t)f^\Delta(t)$.

We list some basic properties of the Hilger derivatives.

Theorem 2.2. Assume that $f, g : \mathbb{T} \mapsto \mathbb{R}$ are differentiable at $t \in \mathbb{T}$. Then

1. the sum $f + g$ is differentiable at t with

$$(f + g)^\Delta(t) = f^\Delta(t) + g^\Delta(t).$$

2. for any constant α , the function αf is differentiable at t with

$$(\alpha f)^\Delta(t) = \alpha f^\Delta(t).$$

3. if $g(t), g(\sigma(t)) \neq 0$ then f/g is differentiable at t with

$$(f/g)^\Delta(t) = \frac{f^\Delta(t)g(t) - f(t)g^\Delta(t)}{g(t)g(\sigma(t))}.$$

4. the product fg is differentiable at t with

$$(fg)^\Delta(t) = f^\Delta(t)g(t) + f(\sigma(t))g^\Delta(t) = f(t)g^\Delta(t) + f^\Delta(t)g(\sigma(t)).$$

The integral calculus on time scales is also quite well-developed, see [2]. Now we introduce some basic concepts related to the theory of linear systems of the form

$$x^\Delta = A(t)x \tag{2.1}$$

on time scales.

Properties of solutions of system (2.1) are also studied. In general, they correspond to those for ordinary differential equations with one important exception which can be illustrated by the following example.

Example. Let $\mathbb{T} = \mathbb{N}$. Consider the scalar equation $x^\Delta = -x$. Then any solution of the considered equation with initial conditions $x(n_0) = x_0$ is zero for any $n > n_0$, so there is no backward uniqueness of solutions.

Definition 2.7. We say that the matrix A is regressive with respect to \mathbb{T} provided $E + \mu(t)A(t)$ is invertible for all $t \in \mathbb{T}$.

Theorem 2.3. 1. Any solution of system (2.1) with initial conditions $x(t_0) = x_0$ is correctly defined for any $t \geq t_0$.

2. Solutions of system (2.1) are unique if the matrix is regressive.

3. The matrix-valued function A is regressive if and only if the eigenvalues $\lambda_i(t)$ of $A(t)$ are regressive for all $1 \leq i \leq n$.

An analog of the Lagrange method to solve linear non-homogeneous systems holds true for time scale systems, as well.

3 P -Projectors. Matrix Chains

Below, without loss of generality, we remove the dependence on t for the sake of notational simplicity.

3.1 Properly Stated Matrix Pairs

Suppose that $A : I \rightarrow \mathcal{M}_{n \times m}$ and $B : I \rightarrow \mathcal{M}_{m \times n}$.

Definition 3.1. *We will say that the matrix pair (A, B) is properly stated on I if A and B satisfy*

$$\ker A \oplus \operatorname{im} B = \mathbb{R}^m \quad (3.1)$$

and both $\ker A$ and $\operatorname{im} B$ are \mathcal{C}^1 -spaces. The condition (3.1) is said to be the transversality condition.

Let (A, B) be a properly stated matrix pair on I . Then A , B , and AB have a constant rank on I and

$$\ker AB = \ker B.$$

Indeed, (3.1) implies that if $y \in \operatorname{im} B$ then $Ay \neq 0$.

Note that, by the transversality condition (3.1) together with the \mathcal{C}^1 -smoothness of $\ker A$ and $\operatorname{im} B$, it follows that there is a projector $R \in \mathcal{C}^1(I)$ onto $\operatorname{im} B$ and along $\ker A$ on I .

Theorem 3.1. *Let $N = \ker A$ and $\dim N = m - r$. Then there exists a smooth projector Q onto N on I if and only if N can be spanned by $\{n_1, \dots, n_{m-r}\}$ on I , where $n_j \in \mathcal{C}^1(I)$, $j \in \{1, \dots, m - r\}$.*

Proof. 1. Suppose that there is a smooth projector Q onto N . Take $t_0 \in I$ arbitrarily. Let

$$n_1^0, \dots, n_{m-r}^0 \in \mathbb{R}^m$$

be a basis of $N(t_0)$. We choose n_j , $j \in \{1, \dots, m - r\}$, as the solution of the initial value

$$n_j^\Delta = Q^\Delta n_j \quad \text{on } I,$$

$$n_j(t_0) = n_j^0, \quad j \in \{1, \dots, m - r\}.$$

Let also,

$$P = E - Q.$$

Here E stands for the unit matrix. Then

$$PQ = 0, \quad Pn_j^0 = 0, \quad j \in \{1, \dots, m - r\},$$

and

$$0 = (PQ)^\Delta = P^\Delta Q + P^\sigma Q^\Delta,$$

whereupon

$$P^\Delta Q = -P^\sigma Q^\Delta.$$

Next,

$$P^\sigma n_j^\Delta = P^\sigma Q^\Delta n_j,$$

and

$$\begin{aligned} (Pn_j)^\Delta &= P^\Delta n_j + P^\sigma n_j^\Delta = P^\Delta n_j + P^\sigma Q^\Delta n_j = P^\Delta n_j - P^\Delta Q n_j \\ &= P^\Delta (E - Q) n_j = P^\Delta P n_j \quad \text{on } I, \quad j \in \{1, \dots, m-r\}. \end{aligned}$$

So, we obtain the following IVP

$$(Pn_j)^\Delta = P^\Delta P n_j \quad \text{on } I$$

$$P n_j(t_0) = 0, \quad j \in \{1, \dots, m-r\}.$$

Therefore $P n_j = 0$ on I , $j \in \{1, \dots, m-r\}$, and $n_j \in N$ on I . Since n_1, \dots, n_{m-r} are linearly independent on I (otherwise, the dimension of N is non-constant), we conclude that they span N on I .

2. Let N be spanned by

$$F = (n_1, \dots, n_{m-r}), \quad n_j \in \mathcal{C}^1(I), \quad j \in \{1, \dots, m-r\},$$

on I . Take

$$Q = F (F^T F)^{-1} F^T \quad \text{on } I.$$

Observe that the matrix $F^T F$ is invertible since vectors n_1, \dots, n_{m-r} are linearly independent. Then

$$\begin{aligned} QQ &= F (F^T F)^{-1} F^T F (F^T F)^{-1} F^T \\ &= F (F^T F)^{-1} F^T \\ &= Q. \end{aligned}$$

Note that $Q \in \mathcal{C}^1(I)$. This completes the proof. □

Definition 3.2. *If the null space $N = \ker A$, $\dim N = m - r$, satisfies Theorem 3.1, we will say that N depends on $t \in I$ smoothly.*

3.2 Matrix Chains

Suppose that (A, B) is a properly stated matrix pair on I and $C \in \mathcal{C}(I)$. Set

$$G_0 = AB.$$

Denote with P_0 a projector along $\ker G_0$. Assume that R is a continuous projector onto $\text{im } B$ and along $\ker A$. With B^- we denote the $\{1, 2\}$ -inverse of B , determined by

$$\begin{aligned} BB^-B &= B, \quad B^-BB^- = B^-, \\ B^-B &= P_0, \quad BB^- = R. \end{aligned} \tag{3.2}$$

Note that the matrix B^- is uniquely determined by the conditions (3.2). Set

$$Q_0 = E - P_0.$$

We assume that a matrix $G_1 : I \mapsto \mathcal{M}_{m,m}$ is such that

(A1) G_1 has a constant rank r_1 on I .

(A2) $N_1 = \ker G_1$ satisfies $N_0 \cap N_1 = \{0\}$.

We choose a continuous projector Q_1 onto N_1 such that

(A3) $Q_1 Q_0 = 0$.

Set $P_1 = E - Q_1$ and assume

(A4) $BP_0 P_1 B^- \in C^1(I)$.

Assume that (A1)-(A4) hold and set $C_0 = C$. Define

$$G_1 = G_0 + C_0 Q_0.$$

Remark 3.1. Note that the existence of a projector Q_1 so that $Q_1 Q_0 = 0$ relies on the fact that the condition (A2) makes it possible to choose a projector Q_1 onto N_1 such that $N_0 \subseteq \ker Q_1$. Since Q_0 projects onto N_0 , we have $\text{im} Q_0 = N_0$. Hence, for any $x \in \mathbb{R}^m$, we have

$$Q_0 x \in N_0 \subseteq \ker Q_1$$

and then $Q_1 Q_0 x = 0$, i.e., $Q_1 Q_0 = 0$.

Definition 3.3. The projectors $\{Q_0, Q_1\}$, respectively $\{P_0, P_1\}$, are said to be admissible up to level 1 if (A1), (A2) and (A3) hold.

Definition 3.4. The matrix pair (A, B) is said to be the leading term for the LTVDAE (1.1).

Definition 3.5. Let the leading term of (1.1) is properly stated on I . The equation (1.1) is said to be regular with tractability index 1 if there are projectors Q_0 and Q_1 that are admissible up to level 1, G_0 is singular and G_1 is nonsingular.

Proposition 3.1. Let $\{Q_0, Q_1\}$ be admissible up to level 1. Then $\ker P_0 P_1 = N_0 \oplus N_1$.

Proof. 1. Suppose that $z \in \ker P_0 P_1$. Then

$$P_0 P_1 z = 0. \tag{3.3}$$

Set

$$z_1 = P_1 z = (E - Q_1)z = z - Q_1 z,$$

i.e.,

$$z = z_1 + Q_1 z.$$

Hence, (3.3) holds if and only if $z_1 \in \ker P_0 = N_0$. Note that $Q_1 z \in N_1$. Therefore

$$z \in N_0 \oplus N_1.$$

Because $z \in \ker P_0 P_1$ was arbitrarily chosen and we get that it is an element of $N_0 \oplus N_1$, we arrive at the relation

$$\ker P_0 P_1 \subseteq N_0 \oplus N_1. \tag{3.4}$$

2. Let $z \in N_0 \oplus N_1$ be arbitrarily chosen. Then

$$z = z_0 + z_1,$$

where $z_0 \in N_0$ and $z_1 \in N_1$. Since

$$\ker P_1 = N_1,$$

we have

$$P_1 z_1 = 0.$$

Next, by $z_0 \in N_0$, it follows that there is a $w_0 \in \mathbb{R}^m$ so that

$$z_0 = Q_0 w_0.$$

Therefore

$$\begin{aligned} P_0 P_1 z &= P_0 P_1 (z_0 + z_1) = P_0 P_1 z_0 + P_0 P_1 z_1 = P_0 P_1 Q_0 w_0 \\ &= P_0 (E - Q_1) Q_0 w_0 = P_0 (Q_0 - Q_1 Q_0) w_0 = P_0 Q_0 w_0 = 0, \end{aligned}$$

i.e., $z \in \ker P_0 P_1$. Since $z \in N_0 \oplus N_1$ was arbitrarily chosen and we get that it is an element of $\ker P_0 P_1$, we obtain the relation

$$N_0 \oplus N_1 \subseteq \ker P_0 P_1.$$

By the last inclusion and (3.4), we obtain $\ker P_0 P_1 = N_0 \oplus N_1$. This completes the proof. \square

3.3 Independence of the Matrix Chains

We will start this subsection with the following useful lemmas.

Lemma 3.1. *Let $\{Q_0, Q_1\}$ and $\{\overline{Q}_0, \overline{Q}_1\}$ be two admissible up to level 1 projector sequences and $\{G_0, N_0, G_1, N_1\}$ and $\{\overline{G}_0, \overline{N}_0, \overline{G}_1, \overline{N}_1\}$ be the corresponding sequences. Then*

$$\overline{B}^- = \overline{P}_0 B^-,$$

$$Q_0 \overline{Q}_0 = \overline{Q}_0, \tag{3.5}$$

$$\overline{Q}_0 Q_0 = Q_0, \tag{3.6}$$

$$P_0 \overline{P}_0 = P_0,$$

$$\overline{P}_0 P_0 = \overline{P}_0$$

and

$$G_0 Q_0 = G_0 \overline{Q}_0 = 0.$$

Proof. We have by definition

$$G_0 = \overline{G}_0, \quad N_0 = \overline{N}_0$$

and

$$\ker P_0 = \ker \overline{P}_0.$$

Moreover,

$$BB^{-1}B = B, \quad B^{-1}BB^{-1} = B^{-1}, \quad BB^{-1} = R, \quad B^{-1}B = P_0$$

and

$$B\overline{B}^{-1}B = B, \quad \overline{B}^{-1}B\overline{B}^{-1} = \overline{B}^{-1}, \quad B\overline{B}^{-1} = R, \quad \overline{B}^{-1}B = \overline{P}_0.$$

Hence,

$$\overline{P}_0B^{-1} = \overline{B}^{-1}BB^{-1} = \overline{B}^{-1}R = \overline{B}^{-1}B\overline{B}^{-1} = \overline{B}^{-1}.$$

Since $N_0 = \overline{N}_0$, $\text{im } Q_0 = N_0$, $\text{im } \overline{Q}_0 = \overline{N}_0$, we have

$$\text{im } Q_0 = \text{im } \overline{Q}_0.$$

Take $z \in \mathbb{R}^m$ arbitrarily. If $z \in \ker \overline{Q}_0$, then

$$\overline{Q}_0z = 0$$

and

$$Q_0\overline{Q}_0z = Q_00 = 0.$$

If $z \in \text{im } \overline{Q}_0$, then since \overline{Q}_0 is a projector, we have

$$\overline{Q}_0z = z$$

and

$$z \in \text{im } Q_0 = \text{im } \overline{Q}_0.$$

Therefore

$$Q_0z = z$$

and

$$Q_0\overline{Q}_0z = Q_0z = z = \overline{Q}_0z.$$

Because $z \in \mathbb{R}^m$ was arbitrarily chosen, we conclude that

$$Q_0\overline{Q}_0 = \overline{Q}_0.$$

Let $y \in \mathbb{R}^m$ be arbitrarily chosen. If $y \in \ker Q_0$, then

$$Q_0y = 0$$

and

$$\overline{Q}_0Q_0y = \overline{Q}_00 = 0 = Q_0y.$$

If $y \in \text{im } Q_0$, then $y \in \text{im } \overline{Q}_0$ and

$$\overline{Q}_0y = Q_0y = y.$$

From here,

$$\overline{Q}_0Q_0y = \overline{Q}_0y = y = Q_0y.$$

Because $y \in \mathbb{R}^m$ was arbitrarily chosen, we arrive at

$$\overline{Q}_0 Q_0 = Q_0.$$

By (3.5), we find

$$E - \overline{P}_0 = (E - P_0)(E - \overline{P}_0) = E - P_0 - \overline{P}_0 + P_0 \overline{P}_0,$$

whereupon

$$P_0 \overline{P}_0 = P_0.$$

By (3.6), we find

$$E - P_0 = (E - \overline{P}_0)(E - P_0) = E - P_0 - \overline{P}_0 + \overline{P}_0 P_0,$$

from where

$$\overline{P}_0 P_0 = \overline{P}_0.$$

Since

$$\text{im } Q_0 = N_0 = \ker G_0 = \text{im } \overline{Q}_0,$$

we get

$$G_0 Q_0 = G_0 \overline{Q}_0 = 0.$$

This completes the proof. \square

Lemma 3.2. *Let $\{Q_0, Q_1\}$ and $\{\overline{Q}_0, \overline{Q}_1\}$ be two admissible up to level 1 projector sequences and $\{G_0, N_0, G_1, N_1\}$ and $\{\overline{G}_0, \overline{N}_0, \overline{G}_1, \overline{N}_1\}$ be the corresponding sequences. If*

$$Z_1 = E + Q_0 \overline{Q}_0 P_0,$$

then

$$\overline{G}_1 = G_1 Z_1$$

and

$$\ker P_0 P_1 = \ker \overline{P}_0 \overline{P}_1 = N_0 \oplus N_1 = \overline{N}_0 \oplus \overline{N}_1.$$

Proof. Applying (3.5) and (3.6), we get

$$\begin{aligned} \overline{G}_1 &= G_0 + C \overline{Q}_0 = G_0 + C Q_0 \overline{Q}_0 = G_0 + C Q_0 \overline{Q}_0 (P_0 + Q_0) \\ &= G_0 + C Q_0 \overline{Q}_0 P_0 + C Q_0 \overline{Q}_0 Q_0 = G_0 + C Q_0 \overline{Q}_0 P_0 + C Q_0 Q_0 = G_0 + C Q_0 + C Q_0 \overline{Q}_0 P_0 \\ &= G_1 + C Q_0 \overline{Q}_0 P_0 = G_1 + (G_0 Q_0 + C Q_0^2) \overline{Q}_0 P_0 = G_1 + (G_0 + C Q_0) Q_0 \overline{Q}_0 P_0 \\ &= G_1 + G_1 Q_0 \overline{Q}_0 P_0 = G_1 (E + Q_0 \overline{Q}_0 P_0) = G_1 Z_1. \end{aligned}$$

Hence,

$$N_1 = Z_1^{-1} \overline{N}_1$$

and thus,

$$N_0 \oplus N_1 = \overline{N}_0 \oplus \overline{N}_1.$$

By Proposition 3.1, it follows that

$$\ker P_0 P_1 = \ker \overline{P}_0 \overline{P}_1 = N_0 \oplus N_1 = \overline{N}_0 \oplus \overline{N}_1.$$

This completes the proof. \square

By Lemma 3.1 and Lemma 3.2, it follows the following result.

Theorem 3.2. *The subspaces $\text{im } G_0$, $\text{im } G_1$, N_0 and $N_0 \oplus N_1$ do not all depend on the special choice of the projectors Q_0 and Q_1 .*

4 An Important Particular Case

Consider the equation

$$A^\sigma x^\Delta = C^\sigma x^\sigma + f. \quad (4.1)$$

We will show that the equation (4.1) can be reduced to the equation (1.1). Suppose that P is a \mathcal{C}^1 -projector along $\ker A^\sigma$. Then

$$A^\sigma P = A^\sigma$$

and

$$A^\sigma x^\Delta = A^\sigma P x^\Delta = A^\sigma (P x)^\Delta - A^\sigma P^\Delta x^\sigma.$$

Hence, the equation (4.1) takes the form

$$A^\sigma (P x)^\Delta - A^\sigma P^\Delta x^\sigma = C^\sigma x^\sigma + f$$

or

$$A^\sigma (P x)^\Delta = (A^\sigma P^\Delta + C^\sigma) x^\sigma + f.$$

Set

$$C_1^\sigma = A^\sigma P^\Delta + C^\sigma.$$

Thus, (4.1) takes the form

$$A^\sigma (P x)^\Delta = C_1^\sigma x^\sigma + f, \quad (4.2)$$

i.e., the equation (4.1) is a particular case of the equation (1.1).

5 Standard Form Index One Problems

In this section, we will investigate equation (4.2) where $\ker A$ is a \mathcal{C}^1 -space, $C \in \mathcal{C}(I)$, P is a \mathcal{C}^1 -projector along $\ker A$. Then

$$AP = A.$$

Assume in addition that

$$Q = E - P$$

and

(B1) the matrix

$$A_1 = A + CQ$$

is invertible.

The condition (B1) ensures the equation (4.2) to be regular with tractability index 1. We will start our investigations with the following useful lemma.

Lemma 5.1. *Suppose that (B1) holds. Then*

$$A_1^{-1} A = P$$

and

$$A_1^{-1} CQ = Q.$$

Proof. We have

$$A_1P = (A + CQ)P = AP + CQP = A.$$

Since $Q = E - P$ and $\ker P = \ker A$, we have $\text{im } Q = \ker A$ and

$$AQ = 0.$$

Then

$$A_1Q = (A + CQ)Q = AQ + CQQ = CQ.$$

This completes the proof. \square

Now, we multiply the equation (4.2) with $(A_1^{-1})^\sigma$ and we get

$$(A_1^{-1})^\sigma A^\sigma (Px)^\Delta = (A_1^{-1})^\sigma C^\sigma x^\sigma + (A_1^{-1})^\sigma f.$$

Now, we employ the first equation of Lemma 5.1 and we get

$$P^\sigma (Px)^\Delta = (A_1^{-1})^\sigma C^\sigma x^\sigma + (A_1^{-1})^\sigma f. \quad (5.1)$$

We decompose x in the following way

$$x = Px + Qx.$$

Then the equation (5.1) takes the following form

$$\begin{aligned} P^\sigma (Px)^\Delta &= (A_1^{-1})^\sigma C^\sigma (P^\sigma x^\sigma + Q^\sigma x^\sigma) + (A_1^{-1})^\sigma f \\ &= (A_1^{-1})^\sigma C^\sigma P^\sigma x^\sigma + (A_1^{-1})^\sigma C^\sigma Q^\sigma x^\sigma + (A_1^{-1})^\sigma f. \end{aligned}$$

Using the second equation of Lemma 5.1, the last equation can be rewritten as follows

$$P^\sigma (Px)^\Delta = (A_1^{-1})^\sigma C^\sigma P^\sigma x^\sigma + Q^\sigma x^\sigma + (A_1^{-1})^\sigma f. \quad (5.2)$$

We multiply the equation (5.2) with the projector P^σ and using that

$$PP = P, \quad PQ = 0,$$

we find

$$P^\sigma P^\sigma (Px)^\Delta = P^\sigma (A_1^{-1})^\sigma C^\sigma P^\sigma x^\sigma + P^\sigma Q^\sigma x^\sigma + P^\sigma (A_1^{-1})^\sigma f$$

or

$$P^\sigma (Px)^\Delta = P^\sigma (A_1^{-1})^\sigma C^\sigma P^\sigma x^\sigma + P^\sigma (A_1^{-1})^\sigma f. \quad (5.3)$$

Note that

$$P^\sigma (Px)^\Delta = (PPx)^\Delta - P^\Delta Px = (Px)^\Delta - P^\Delta Px.$$

Hence and using (5.3), we find

$$(Px)^\Delta - P^\Delta Px = P^\sigma (A_1^{-1})^\sigma C^\sigma P^\sigma x^\sigma + P^\sigma (A_1^{-1})^\sigma f$$

or

$$(Px)^\Delta = P^\Delta Px + P^\sigma (A_1^{-1})^\sigma C^\sigma P^\sigma x^\sigma + P^\sigma (A_1^{-1})^\sigma f. \quad (5.4)$$

Now, we multiply the equation (5.2) by Q^σ and we find

$$Q^\sigma P^\sigma (Px)^\Delta = Q^\sigma (A_1^{-1})^\sigma C^\sigma P^\sigma x^\sigma + Q^\sigma Q^\sigma x^\sigma + Q^\sigma (A_1^{-1})^\sigma f$$

or

$$0 = Q^\sigma (A_1^{-1})^\sigma C^\sigma P^\sigma x^\sigma + Q^\sigma x^\sigma + Q^\sigma (A_1^{-1})^\sigma f. \quad (5.5)$$

Set

$$u = Px, \quad v = Qx.$$

Then, by (5.4) and (5.5), we get the system

$$\begin{aligned} u^\Delta &= P^\Delta u + P^\sigma (A_1^{-1})^\sigma C^\sigma u^\sigma + P^\sigma (A_1^{-1})^\sigma f \\ v^\sigma &= -Q^\sigma (A_1^{-1})^\sigma C^\sigma u^\sigma - Q^\sigma (A_1^{-1})^\sigma f. \end{aligned} \quad (5.6)$$

We find $u \in \mathcal{C}^1(I)$ by the first equation of the system (5.6) and then we find $v^\sigma \in \mathcal{C}(I)$ by the second equation of the system (5.6). Then, for the solution x of the equation (4.2) we have the following representation

$$x^\sigma = u^\sigma + v^\sigma = P^\sigma x^\sigma + Q^\sigma x^\sigma.$$

6 Decoupling

Suppose that the equation (1.1) is regular with tractability index 1. In addition, assume that R is a continuous projector onto $\text{im } B$ and along $\ker A$. Set

$$G_0 = AB$$

and take P_0 to be a continuous projector along $\ker G_0$ and denote

$$Q_0 = E - P_0, \quad G_1 = G_0 + CQ_0,$$

$$BB^-B = B, \quad B^-BB^- = B^-, \quad B^-B = P_0, \quad BB^- = R, \quad N_0 = \ker G_0.$$

Since

$$R = BB^-$$

and R is a continuous projector along $\ker A$, we have

$$A = AR = ABB^- = G_0B^-.$$

Then, we can rewrite the equation (1.1) as follows

$$A^\sigma B^\sigma B^{-\sigma} (Bx)^\Delta = C^\sigma x^\sigma + f$$

or

$$G_0^\sigma B^{-\sigma} (Bx)^\Delta = C^\sigma x^\sigma + f.$$

Now, we multiply both sides of the last equation with $(G_1^{-1})^\sigma$ and we find

$$(G_1^{-1})^\sigma G_0^\sigma B^{-\sigma}(Bx)^\Delta = (G_1^{-1})^\sigma C^\sigma x^\sigma + (G_1^{-1})^\sigma f. \quad (6.1)$$

Note that by the definition of G_0 and Q_0 we have $G_0 Q_0 = 0$. Then

$$\begin{aligned} G_1(E - Q_0) &= (G_0 + CQ_0)(E - Q_0) = G_0 + CQ_0 - G_0 Q_0 - CQ_0 Q_0 \\ &= G_0 + CQ_0 - CQ_0 = G_0, \end{aligned}$$

whereupon

$$G_1^{-1}G_0 = E - Q_0.$$

Thus, the equation (6.1) takes the form

$$(E - Q_0^\sigma)B^{-\sigma}(Bx)^\Delta = (G_1^{-1})^\sigma C^\sigma x^\sigma + (G_1^{-1})^\sigma f. \quad (6.2)$$

Note that

$$B^\sigma P_0^\sigma(E - Q_0^\sigma) = B^\sigma(P_0^\sigma - P_0^\sigma Q_0^\sigma) = B^\sigma P_0^\sigma.$$

Then, we multiply the equation (6.2) by $B^\sigma P_0^\sigma$ and we find

$$B^\sigma P_0^\sigma B^{-\sigma}(Bx)^\Delta = B^\sigma P_0^\sigma (G_1^{-1})^\sigma C^\sigma x^\sigma + B^\sigma P_0^\sigma (G_1^{-1})^\sigma f. \quad (6.3)$$

Note that

$$\begin{aligned} B^\sigma P_0^\sigma B^{-\sigma}(Bx)^\Delta &= (BP_0 B^- Bx)^\Delta - (BP_0 B^-)^\Delta Bx \\ &= (BP_0 P_0 x)^\Delta - (BP_0 B^-)^\Delta Bx = (BP_0 x)^\Delta - (BP_0 B^-)^\Delta Bx. \end{aligned}$$

From here, the equation (6.3) can be rewritten as follows

$$(BP_0 x)^\Delta = (BP_0 B^-)^\Delta Bx + B^\sigma P_0^\sigma (G_1^{-1})^\sigma C^\sigma x^\sigma + B^\sigma P_0^\sigma (G_1^{-1})^\sigma f.$$

Now, we use the decomposition

$$x = P_0 x + Q_0 x,$$

to get

$$\begin{aligned} (BP_0 x)^\Delta &= (BP_0 B^-)^\Delta BP_0 x + (BP_0 B^-)^\Delta BQ_0 x \\ &\quad + B^\sigma P_0^\sigma (G_1^{-1})^\sigma C^\sigma P_0^\sigma x^\sigma + B^\sigma P_0^\sigma (G_1^{-1})^\sigma C^\sigma Q_0^\sigma x^\sigma \\ &\quad + B^\sigma P_0^\sigma (G_1^{-1})^\sigma f. \end{aligned}$$

Since $\text{im } Q_0 = \ker B$, we have

$$(BP_0 B^-)^\Delta BQ_0 x = 0.$$

Next,

$$\begin{aligned} E &= G_1^{-1}(G_0 + CQ_0) = G_1^{-1}G_0 + G_1^{-1}CQ_0 \\ &= E - Q_0 + G_1^{-1}CQ_0, \end{aligned}$$

whereupon

$$G_1^{-1}CQ_0 = Q_0. \quad (6.4)$$

Then

$$B^\sigma P_0^\sigma (G_1^{-1})^\sigma C^\sigma Q_0^\sigma = B^\sigma P_0^\sigma Q_0^\sigma = 0$$

and we arrive at the equation

$$(BP_0x)^\Delta = (BP_0B^-)^\Delta BP_0x + B^\sigma P_0^\sigma (G_1^{-1})^\sigma C^\sigma P_0^\sigma x^\sigma + B^\sigma P_0^\sigma (G_1^{-1})^\sigma f.$$

Since

$$B^-BP_0 = P_0P_0 = P_0,$$

the last equation can be rewritten in the form

$$(BP_0x)^\Delta = (BP_0B^-)^\Delta BP_0x + B^\sigma P_0^\sigma (G_1^{-1})^\sigma C^\sigma B^{-\sigma} B^\sigma P_0^\sigma x^\sigma + B^\sigma P_0^\sigma (G_1^{-1})^\sigma f.$$

We set

$$u = BP_0x.$$

Then, we obtain the equation

$$u^\Delta = (BP_0B^-)^\Delta u + B^\sigma P_0^\sigma (G_1^{-1})^\sigma C^\sigma B^{-\sigma} u^\sigma + B^\sigma P_0^\sigma (G_1^{-1})^\sigma f. \quad (6.5)$$

Now, we multiply both sides of (6.2) by Q_0^σ and using that

$$Q_0^\sigma(E - Q_0^\sigma) = Q_0^\sigma - Q_0^\sigma Q_0^\sigma = Q_0^\sigma - Q_0^\sigma = 0,$$

we find

$$0 = Q_0^\sigma (G_1^{-1})^\sigma C^\sigma x^\sigma + Q_0^\sigma (G_1^{-1})^\sigma f,$$

or

$$\begin{aligned} Q_0^\sigma (G_1^{-1})^\sigma f &= -Q_0^\sigma (G_1^{-1})^\sigma C^\sigma x^\sigma \\ &= -Q_0^\sigma (G_1^{-1})^\sigma C^\sigma (P_0^\sigma x^\sigma + Q_0^\sigma x^\sigma) \\ &= -Q_0^\sigma (G_1^{-1})^\sigma C^\sigma P_0^\sigma x^\sigma - Q_0^\sigma (G_1^{-1})^\sigma C^\sigma Q_0^\sigma x^\sigma \\ &= -Q_0^\sigma (G_1^{-1})^\sigma C^\sigma P_0^\sigma x^\sigma - Q_0^\sigma Q_0^\sigma x^\sigma \\ &= -Q_0^\sigma (G_1^{-1})^\sigma C^\sigma P_0^\sigma x^\sigma - Q_0^\sigma x^\sigma, \end{aligned}$$

where we have used (6.4). We set

$$v = Q_0x.$$

Then, we get

$$v^\sigma = -Q_0^\sigma (G_1^{-1})^\sigma C^\sigma u^\sigma - Q_0^\sigma (G_1^{-1})^\sigma f.$$

By the last equation and (6.5), we obtain the system

$$\begin{aligned} u^\Delta &= (BP_0B^-)^\Delta u + B^\sigma P_0^\sigma (G_1^{-1})^\sigma C^\sigma B^{-\sigma} u^\sigma + B^\sigma P_0^\sigma (G_1^{-1})^\sigma f \\ v^\sigma &= -Q_0^\sigma (G_1^{-1})^\sigma C^\sigma u^\sigma - Q_0^\sigma (G_1^{-1})^\sigma f. \end{aligned} \quad (6.6)$$

By the first equation of (6.6) we find u and then by the second equation we find v^σ . Then, for the solution x of (1.1), we have

$$B^{-\sigma}u^\sigma + v^\sigma = B^{-\sigma}B^\sigma P_0^\sigma x^\sigma + Q_0^\sigma x^\sigma = P_0^\sigma x^\sigma + Q_0^\sigma x^\sigma = x^\sigma.$$

Here we write $B^{-\sigma}$ instead of $(B^-)^\sigma$. Now, we will show that the described process above is σ -reversible, i.e., if $u \in \mathcal{C}^1(I)$, $u \in \text{im } BP_0B^-$ and $v^\sigma \in \mathcal{C}(I)$ satisfy (6.6) and

$$x^\sigma = B^{-\sigma}u^\sigma + v^\sigma, \quad (6.7)$$

then x satisfies (1.1). Really, since $u \in \text{im } BP_0B^-$, we have

$$BP_0B^-u = u$$

and

$$BB^-u = BB^-BP_0B^-u = BP_0P_0B^-u = BP_0B^-u = u.$$

By the second equation of (6.6), using that $Q_0^\sigma Q_0^\sigma = Q_0^\sigma$, we find

$$v^\sigma = Q_0^\sigma v^\sigma.$$

Because $\text{im } Q_0 = \ker B$, we have

$$B^\sigma v^\sigma = B^\sigma Q_0^\sigma v^\sigma = 0.$$

Using (6.7), we find

$$\begin{aligned} B^\sigma P_0^\sigma x^\sigma &= B^\sigma P_0^\sigma (B^{-\sigma}u^\sigma + v^\sigma) = B^\sigma P_0^\sigma (B^{-\sigma}u^\sigma + Q_0^\sigma v^\sigma) \\ &= B^\sigma P_0^\sigma B^{-\sigma}u^\sigma = u^\sigma. \end{aligned}$$

Hence,

$$B^{-\sigma}u^\sigma = B^{-\sigma}B^\sigma P_0^\sigma x^\sigma = P_0^\sigma P_0^\sigma x^\sigma = P_0^\sigma x^\sigma = x^\sigma - Q_0^\sigma x^\sigma$$

using (6.7), we find

$$x^\sigma = B^{-\sigma}u^\sigma + v^\sigma = x^\sigma - Q_0^\sigma x^\sigma + v^\sigma,$$

whereupon $v^\sigma = Q_0^\sigma x^\sigma$. Next,

$$\begin{aligned} B^\sigma x^\sigma &= B^\sigma (P_0^\sigma x^\sigma + Q_0^\sigma x^\sigma) = B^\sigma (P_0^\sigma x^\sigma + v^\sigma) \\ &= B^\sigma P_0^\sigma x^\sigma + B^\sigma v^\sigma = u^\sigma. \end{aligned}$$

Note that the equation (6.5) is restated by the equation (6.1). Then, multiplying (6.1) with $B^\sigma P_0^\sigma$, we find

$$B^\sigma P_0^\sigma (G_1^{-1})^\sigma G_0^\sigma B^{-\sigma} (Bx)^\Delta = B^\sigma P_0^\sigma (G_1^{-1})^\sigma C^\sigma x^\sigma + B^\sigma P_0^\sigma (G_1^{-1})^\sigma f,$$

which we premultiply by $B^{-\sigma}$ and using that $B^{-\sigma}B^\sigma P_0^\sigma = P_0^\sigma$, we arrive at

$$P_0^\sigma (G_1^{-1})^\sigma G_0^\sigma B^{-\sigma} (Bx)^\Delta = P_0^\sigma (G_1^{-1})^\sigma C^\sigma x^\sigma + P_0^\sigma (G_1^{-1})^\sigma f,$$

from where

$$P_0^\sigma (G_1^{-1})^\sigma A^\sigma (Bx)^\Delta = P_0^\sigma (G_1^{-1})^\sigma C^\sigma x^\sigma + P_0^\sigma (G_1^{-1})^\sigma f. \quad (6.8)$$

Next, we multiply (6.1) by Q_0^σ and we find

$$Q_0^\sigma (G_1^{-1})^\sigma A^\sigma (Bx)^\Delta = Q_0^\sigma (G_1^{-1})^\sigma C^\sigma x^\sigma + Q_0^\sigma (G_1^{-1})^\sigma f. \quad (6.9)$$

Now, we add (6.8) and (6.9) and we find

$$(G_1^{-1})^\sigma A^\sigma (Bx)^\Delta = (G_1^{-1})^\sigma C^\sigma x^\sigma + (G_1^{-1})^\sigma f,$$

whereupon we get (1.1).

Definition 6.1. *The equation (6.5) is said to be the inherent equation for the equation (1.1).*

Theorem 6.1. *The subspace $\text{im } P_0$ is an invariant subspace for the equation (6.5), i.e.,*

$$u(t_0) \in (\text{im } BP_0)(t_0)$$

for some $t_0 \in I$ if and only if

$$u(t) \in (\text{im } BP_0)(t)$$

for any $t \in I$.

Proof. Let $u \in \mathcal{C}^1(I)$ be a solution to the equation (6.5) so that

$$(BP_0)(t_0)u(t_0) = u(t_0).$$

Hence,

$$\begin{aligned} u(t_0) &= (BP_0)(t_0)u(t_0) = (BP_0P_0P_0)(t_0)u(t_0) = (BP_0B^-BP_0)(t_0)u(t_0) \\ &= (BP_0B^-)(t_0)(BP_0)(t_0)u(t_0) = (BP_0B^-)(t_0)u(t_0). \end{aligned}$$

We multiply the equation (6.5) by $E - B^\sigma P_0^\sigma B^{-\sigma}$ and we get

$$\begin{aligned} (E - B^\sigma P_0^\sigma B^{-\sigma})u^\Delta &= (E - B^\sigma P_0^\sigma B^{-\sigma})(BP_0B^-)^\Delta u \\ &\quad + (E - B^\sigma P_0^\sigma B^{-\sigma})B^\sigma P_0^\sigma G_1^{-1\sigma} C^\sigma B^{-\sigma} u^\sigma \\ &\quad + (E - B^\sigma P_0^\sigma B^{-\sigma})B^\sigma P_0^\sigma (G_1^{-1})^\sigma f \\ &= (E - B^\sigma P_0^\sigma B^{-\sigma})(BP_0B^-)^\Delta u \\ &\quad + (B^\sigma P_0^\sigma - B^\sigma P_0^\sigma B^{-\sigma} B^\sigma P_0^\sigma)G_1^{-1\sigma} C^\sigma B^{-\sigma} u^\sigma \\ &\quad + (B^\sigma P_0^\sigma - B^\sigma P_0^\sigma B^{-\sigma} B^\sigma P_0^\sigma)G_1^{-1\sigma} f \\ &= (E - B^\sigma P_0^\sigma B^{-\sigma})(BP_0B^-)^\Delta u \\ &\quad + (B^\sigma P_0^\sigma - B^\sigma P_0^\sigma)G_1^{-1\sigma} C^\sigma B^{-\sigma} u^\sigma \\ &\quad + (B^\sigma P_0^\sigma - B^\sigma P_0^\sigma)G_1^{-1\sigma} f \\ &= (E - B^\sigma P_0^\sigma B^{-\sigma})(BP_0B^-)^\Delta u, \end{aligned}$$

i.e.,

$$(E - B^\sigma P_0^\sigma B^{-\sigma})u^\Delta = (E - B^\sigma P_0^\sigma B^{-\sigma})(BP_0B^-)^\Delta u. \quad (6.10)$$

Set

$$v = (E - BP_0B^-)u.$$

Then

$$\begin{aligned} v^\Delta &= (E - B^\sigma P_0^\sigma B^{-\sigma})u^\Delta + (E - BP_0B^-)^\Delta u \\ &= (E - B^\sigma P_0^\sigma B^{-\sigma})(BP_0B^-)^\Delta u + (E - BP_0B^-)^\Delta u \\ &= ((E - BP_0B^-)BP_0B^-)^\Delta u \\ &\quad - (E - BP_0B^-)^\Delta BP_0B^- u \\ &\quad + (E - BP_0B^-)^\Delta u \\ &= (BP_0B^- - BP_0B^-BP_0B^-)^\Delta u \\ &\quad + (E - BP_0B^-)^\Delta (E - BP_0B^-)u \\ &= (E - BP_0B^-)^\Delta v, \end{aligned}$$

i.e.,

$$v^\Delta = (E - BP_0B^-)^\Delta v.$$

Note that

$$\begin{aligned} v(t_0) &= u(t_0) - (BP_0B^-)(t_0)u(t_0) \\ &= u(t_0) - u(t_0) \\ &= 0. \end{aligned}$$

Thus, we get the following IVP

$$v^\Delta = (E - BP_0B^-)^\Delta v \quad \text{on } I$$

$$v(t_0) = 0.$$

Therefore $v = 0$ on I and then

$$BP_0B^-u = u \quad \text{on } I.$$

Hence, using that

$$\text{im } BP_0 = \text{im } BP_0B^-,$$

we get

$$BP_0u = u \quad \text{on } I.$$

This completes the proof. \square

7 Examples

Example 7.1. Let $\mathbb{T} = \mathbb{N}_0$ Consider (4.2) for

$$A(t) = \begin{pmatrix} -1 & t+1 & -1 \\ 0 & 0 & 0 \\ 0 & 2t+2 & -1 \end{pmatrix}, \quad C(t) = \begin{pmatrix} 0 & 0 & 1 \\ 0 & -t & 1 \\ 0 & 2 & 1 \end{pmatrix},$$

$$P(t) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & -(t+1) & 1 \end{pmatrix}, \quad t \in \mathbb{T}.$$

We have $\sigma(t) = t+1$, $t \in \mathbb{T}$, and $P(t)P(t) = P(t)$, $A(t)P(t) = A(t)$, $t \in \mathbb{T}$. Then

$$Q(t) = I - P(t) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & t+1 & 0 \end{pmatrix}$$

$$A_1(t) = A(t) + C(t)Q(t) = \begin{pmatrix} -1 & 2(t+1) & -1 \\ 0 & 1 & 0 \\ 0 & 3t+5 & -1 \end{pmatrix},$$

$$A_1^{-1}(t) = \begin{pmatrix} -1 & -(t+3) & 1 \\ 0 & 1 & 0 \\ 0 & 3t+5 & -1 \end{pmatrix}, \quad t \in \mathbb{T}.$$

Hence,

$$(A_1^{-1})^\sigma(t) = \begin{pmatrix} -1 & -(t+4) & 1 \\ 0 & 1 & 0 \\ 0 & 3t+8 & -1 \end{pmatrix},$$

$$C^\sigma(t) = \begin{pmatrix} 0 & 0 & 1 \\ 0 & -t-1 & 1 \\ 0 & 2 & 1 \end{pmatrix}, \quad P^\sigma(t) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & -(t+2) & 1 \end{pmatrix},$$

$$Q^\sigma(t) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & t+2 & 0 \end{pmatrix}, \quad P^\Delta(t) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix}, \quad t \in \mathbb{T},$$

and

$$P^\sigma(t) (A_1^{-1})^\sigma(t) = \begin{pmatrix} -1 & -(t+4) & 1 \\ 0 & 0 & 0 \\ 0 & 2(t+3) & -1 \end{pmatrix},$$

$$P^\sigma(t) (A_1^{-1})^\sigma(t) C^\sigma(t) = \begin{pmatrix} 0 & (t+2)(t+3) & -(t+4) \\ 0 & 0 & 0 \\ 0 & -2(t+2)^2 & 2t+5 \end{pmatrix},$$

$$Q^\sigma(t) (A_1^{-1})^\sigma(t) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & t+2 & 0 \end{pmatrix},$$

$$Q^\sigma(t) (A_1^{-1})^\sigma(t) C^\sigma(t) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & -t-1 & 1 \\ 0 & -(t+1)(t+2) & t+2 \end{pmatrix}, \quad t \in \mathbb{T}.$$

Then the system (5.6) takes the form

$$u^\Delta(t) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix} u(t) + \begin{pmatrix} 0 & (t+2)(t+3) & -(t+4) \\ 0 & 0 & 0 \\ 0 & -2(t+2)^2 & 2t+5 \end{pmatrix} u^\sigma(t) + \begin{pmatrix} -1 & -(t+4) & -1 \\ 0 & 0 & 0 \\ 0 & 2(t+3) & -1 \end{pmatrix} f(t)$$

$$v^\sigma(t) = - \begin{pmatrix} 0 & 0 & 0 \\ 0 & -t-1 & 1 \\ 0 & -(t+1)(t+2) & t+2 \end{pmatrix} u^\sigma(t) - \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & t+2 & 0 \end{pmatrix} f(t), \quad t \in \mathbb{T}.$$

Example 7.2. Let $\mathbb{T} = 2^{\mathbb{N}_0}$. Consider (1.1) for

$$A(t) = \begin{pmatrix} t & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & t^2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad B(t) = \begin{pmatrix} t & 0 & 0 & 0 & 0 \\ 0 & t^2 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{pmatrix},$$

$$C(t) = \begin{pmatrix} 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & t & 1 & 0 \\ 0 & -1 & 0 & 0 & 0 \\ -1 & 1 & 0 & -t^2 & 0 \\ 1 & 0 & 0 & 0 & t^2 \end{pmatrix}, \quad t \in \mathbb{T}.$$

Here $\sigma(t) = 2t$, $t \in \mathbb{T}$, and

$$R(t) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad G_0(t) = A(t)B(t) = \begin{pmatrix} t^2 & 0 & 0 & 0 & 0 \\ 0 & t^2 & 0 & 0 & 0 \\ 0 & 0 & t^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Q_0(t) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad P_0(t) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad t \in \mathbb{T}.$$

Hence,

$$G_1(t) = G_0(t) + C(t)Q_0(t) = \begin{pmatrix} t^2 & 0 & 0 & -1 & 1 \\ 0 & t^2 & 0 & 1 & 0 \\ 0 & 0 & t^2 & 0 & 0 \\ 0 & 0 & 0 & -t^2 & 0 \\ 0 & 0 & 0 & 0 & t^2 \end{pmatrix},$$

$$B^-(t) = \begin{pmatrix} \frac{1}{t} & 0 & 0 \\ 0 & \frac{1}{t^2} & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad t \in \mathbb{T}.$$

Next,

$$\begin{aligned}
B^{-\sigma}(t) &= \begin{pmatrix} \frac{1}{2t} & 0 & 0 \\ 0 & \frac{1}{4t^2} & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & B^{\sigma}(t) &= \begin{pmatrix} 2t & 0 & 0 & 0 & 0 \\ 0 & 4t^2 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{pmatrix}, \\
C^{\sigma}(t) &= \begin{pmatrix} 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & 2t & 1 & 0 \\ 0 & -1 & 0 & 0 & 0 \\ -1 & 1 & 0 & -4t^2 & 0 \\ 1 & 0 & 0 & 0 & 4t^2 \end{pmatrix}, & Q_0^{\sigma}(t) &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \\
P_0^{\sigma}(t) &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, & t &\in \mathbb{T}.
\end{aligned}$$

Note that

$$\det G_1(t) = -t^{10} \neq 0, \quad t \in \mathbb{T},$$

and

$$\begin{aligned}
G_1^{-1}(t) &= \begin{pmatrix} \frac{1}{t^2} & 0 & 0 & -\frac{1}{t^4} & -\frac{1}{t^4} \\ 0 & \frac{1}{t^2} & 0 & \frac{1}{t^4} & 0 \\ 0 & 0 & \frac{1}{t^2} & 0 & 0 \\ 0 & 0 & 0 & -\frac{1}{t^2} & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{t^2} \end{pmatrix}, \\
(G^{-1})^{\sigma}(t) &= \begin{pmatrix} \frac{1}{4t^2} & 0 & 0 & -\frac{1}{16t^4} & -\frac{1}{16t^4} \\ 0 & \frac{1}{4t^2} & 0 & \frac{1}{16t^4} & 0 \\ 0 & 0 & \frac{1}{4t^2} & 0 & 0 \\ 0 & 0 & 0 & -\frac{1}{4t^2} & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{4t^2} \end{pmatrix}, & t &\in \mathbb{T}.
\end{aligned}$$

Therefore

$$\begin{aligned}
B(t)P_0(t)B^{-}(t) &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, & (BP_0B^{-})^{\Delta}(t) &= 0, \\
B^{\sigma}(t)P_0^{\sigma}(t)(G_1^{-1})^{\sigma}(t) &= \begin{pmatrix} \frac{1}{2t} & 0 & 0 & -\frac{1}{8t^3} & -\frac{1}{8t^3} \\ 0 & 1 & 0 & \frac{1}{4t^2} & 0 \\ 0 & 0 & \frac{1}{4t^2} & 0 & 0 \end{pmatrix},
\end{aligned}$$

$$\begin{aligned}
B^\sigma(t)P_0^\sigma(t)(G_1^{-1})^\sigma(t)C^\sigma(t)B^{-\sigma}(t) &= \begin{pmatrix} 0 & -\frac{1}{32t^5} & 0 \\ -\frac{1}{8t^3} & \frac{1}{16t^4} & 2t \\ 0 & -\frac{1}{16t^4} & 0 \end{pmatrix}, \\
Q_0^\sigma(t)(G_1^{-1})^\sigma(t) &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\frac{1}{4t^2} & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{4t^2} \end{pmatrix}, \\
Q_0^\sigma(t)(G_1^{-1})^\sigma(t)C^\sigma(t)B^{-\sigma}(t) &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ \frac{1}{8t^3} & -\frac{1}{16t^4} & 0 \\ \frac{1}{8t^3} & 0 & 0 \end{pmatrix}, \quad t \in \mathbb{T}.
\end{aligned}$$

Then, the decoupling (6.5) takes the form

$$\begin{aligned}
u^\Delta(t) &= \begin{pmatrix} 0 & -\frac{1}{32t^5} & 0 \\ -\frac{1}{8t^3} & \frac{1}{16t^4} & 2t \\ 0 & -\frac{1}{16t^4} & 0 \end{pmatrix} u^\sigma(t) + \begin{pmatrix} \frac{1}{2t} & 0 & 0 & -\frac{1}{8t^3} & -\frac{1}{8t^3} \\ 0 & 1 & 0 & \frac{1}{4t^2} & 0 \\ 0 & 0 & \frac{1}{4t^2} & 0 & 0 \end{pmatrix} f(t) \\
v^\sigma(t) &= - \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ \frac{1}{8t^3} & -\frac{1}{16t^4} & 0 \\ \frac{1}{8t^3} & 0 & 0 \end{pmatrix} u^\sigma(t) - \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\frac{1}{4t^2} & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{4t^2} \end{pmatrix} f(t), \quad t \in \mathbb{T}.
\end{aligned}$$

8 Conclusion

The problem of decoupling is very significant in the theory of dynamical systems of any origin. First of all, it simplifies the process of finding the analytic solution and qualitative studies. For instance, it is widely applied in the Stability Theory (e.g., for hyperbolic, partially hyperbolic, non-uniformly hyperbolic, or regular systems).

The case of an arbitrary time scale is more sophisticated than that of classical ordinary differential equations for the following reasons:

1. time scale calculus involves more sophisticated formulas than the classical ones and the classical theory cannot be immediately translated to the language of time scales;
2. the theory of time scale systems is much less developed than that of ordinary differential equations;
3. autonomous systems are much more difficult to study for generic time scales.

The main result of the paper is Theorem 6.1. Our proofs are constructive.

Although the approach of this paper is quite straightforward, special techniques, based on projector approach were elaborated. The obtained results are illustrated with specific examples.

References

- [1] B. Aulbach, S. Hilger, *Linear Dynamic Processes with Inhomogenous Time Scale*, In Nonlinear Dynamics and Quantum Dynamical Systems (Gaussig, 1990), volume 59 of Math. Res., pages 9–20. Akademie Verlag, Berlin, 1990.
- [2] M. Bohner, A. Peterson. *Dynamic Equations on Time Scales: An Introduction with Applications*, Birkhäuser, Boston, 2001.
- [3] M. Bohner, S. Georgiev. *Multivariable Dynamic Calculus on Time Scales*, Springer, 2016.
- [4] J. J. DaCunha, *Stability for Time Varying Linear Dynamic Systems on Time Scales*, J. Comput. Appl. Math., 176:2 (2005): 381-410.
- [5] Gonçalves Salsa Junior, R., Ma, F., *Linear Systems and Configuration-Space Decoupling Techniques*. In: Advances in the Theory of System Decoupling. Springer, Cham. (2021)
- [6] R. Lamour, R. März and G. Tischendorf, *Differential-Algebraic Equations: A Projector Based Analysis*, Springer, 2013.
- [7] A. A. Martynyuk, *Stability Theory for Dynamic Equations on Time Scales*, Birkhäuser, 2016.