

Dimensionless physics: Planck constant as an element of Minkowski metric

G.E. Volovik^{1,2}

¹*Low Temperature Laboratory, Aalto University,
P.O. Box 15100, FI-00076 Aalto, Finland*

²*Landau Institute for Theoretical Physics,
acad. Semyonov av., 1a, 142432, Chernogolovka, Russia*

(Dated: November 29, 2022)

Several approaches to quantum gravity (including the model of superplastic vacuum; Diakonov tetrads emerging as the bilinear combinations of the fermion fields; and BF -theories of gravity) suggest that in general relativity the metric may have dimension 2, i.e. $[g_{\mu\nu}] = 1/[L]^2$. The important consequence of such metric dimension is that all the diffeomorphism invariant quantities are dimensionless for any dimension of spacetime. These include the action S , interval s , cosmological constant Λ , scalar curvature R , scalar field Φ , etc. Another consequence of the metric dimension $1/[L]^2$ is that the Planck constant \hbar is the element of the Minkowski metric. The Minkowski parameter \hbar is invariant only under Lorentz transformations, and is not diffeomorphism invariant. As a result the Planck constant \hbar has nonzero dimension – the dimension of length $[L]$. Whether this Planck constant length is related to the Planck length scale, is an open question. In principle there can be different Minkowski vacua with their own values of the parameter \hbar . Then in the thermal contact between the two vacua their temperatures obey the analog of the Tolman law: $\hbar_1/T_1 = \hbar_2/T_2$.

PACS numbers:

Contents

I. Introduction.	2
A. Dimensional tetrads	2
B. Dimensional metric	3
II. Dynamics of classical point particle and dimensionless physics	4
III. Minkowski spacetime and Planck constant	4
IV. Schrödinger equation in Minkowski spacetime	6
V. de Sitter spacetime and Planck constant	7
VI. Planck constant and Tolman law	8
VII. Length dimension of Newton constant and Planck length	9
VIII. Conclusion	9
References	10

I. INTRODUCTION.

A. Dimensional tetrads

Several approaches to quantum gravity (including the model of superplastic vacuum; Diakonov tetrads emerging as the bilinear combinations of the fermion fields; BF -theories of gravity; and effective acoustic metric) suggest^{1,2} that in general relativity the metric must have dimension 2, i.e. $[g_{\mu\nu}] = 1/[L]^2$. In particular, the model of the superplastic vacuum³ is described in terms of the so-called elasticity tetrads.⁴⁻⁹

$$E_{\mu}^a = \frac{\partial X^a}{\partial x^{\mu}}, \quad (1)$$

where equations $X^a(x) = 2\pi n_a$ are equations of the (deformed) crystal planes. Since the functions X^a play the role of the geometric $U(1)$ phases and thus are dimensionless, the elasticity tetrads play the role of the gauge fields (translation gauge fields). That is why

tetrads have the same dimension 1 as the dimension of gauge fields:

$$[E_\mu^a] = \frac{1}{[L]}. \quad (2)$$

The dimension n of quantity A means $[A] = [L]^{-n}$, where $[L]$ is dimension of length. The $U(1)$ gauge fields, which form the dimensional tetrads, are not necessarily related to translations and the crystalline structure of the vacuum. The latter is only one of the possible sources of the dimensional tetrads. In particular, the dimension 1 tetrads appear in the Diakonov theory,¹⁰⁻¹³ where the tetrads emerge as bilinear combinations of the quantum fermionic fields:

$$E_\mu^a = \frac{1}{2} \left(\Psi^\dagger \gamma^a \partial_\mu \Psi - \Psi^\dagger \overleftarrow{\partial}_\mu \gamma^a \Psi \right). \quad (3)$$

This means that the metric and thus the space-time distance are both the quantum objects made of the leptons and quarks.¹⁴

In Diakonov theory the fermionic fields Ψ are dimensionless,¹¹ and thus the dimension of the fermionic tetrads is the same as the dimension of the elasticity tetrads in Eq.(2). The fermionic tetrads in Eq.(3) serve as the quantum spinor generalization of the elasticity tetrads.

B. Dimensional metric

Elasticity tetrads in Eq.(1) and fermionic tetrads in Eq.(3) give rise to the metric with lower indices, which is the following bilinear combination of tetrads:

$$g_{\mu\nu} = \eta_{ab} E_\mu^a E_\nu^b. \quad (4)$$

This covariant metric tensor has dimension $n = 2$, while the contravariant metric $g^{\mu\nu}$ has dimension $n = -2$:

$$[g_{\mu\nu}] = \frac{1}{[L]^2}, \quad [g^{\mu\nu}] = [L]^2. \quad (5)$$

The determinant of the tetrads has dimension 4 in the 4-dimensional spacetime:

$$[e] = [\sqrt{-g}] = \frac{1}{[L]^4}. \quad (6)$$

In the N -dimensional spacetime, the dimensions of the metric elements are the same as in Eq.(5), while the tetrad determinant has dimension N :

$$[e] = [\sqrt{-g}] = \frac{1}{[L]^N}. \quad (7)$$

Eq.(4) gives rise to the dimensionless interval:

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu, \quad [s^2] = \frac{1}{[L]^2} \cdot [L]^2 = [1] = 0. \quad (8)$$

What are the consequences of such unusual dimension of metric? Does such metric describe geometry or dynamics? Let us consider this on example of dynamics of classical point particle.

II. DYNAMICS OF CLASSICAL POINT PARTICLE AND DIMENSIONLESS PHYSICS

The interval in Eq.(8) describes in particular the classical dynamics of a point particle with action:

$$S = M \int ds. \quad (9)$$

Here the particle mass M in Eq.(9) is dimensionless, $[M] = [1] = 0$, as well as all other diffeomorphism invariant quantities, such as action S , interval s , Newton constant G , cosmological constant Λ , scalar curvature R , scalar field Φ , etc.^{1,2} This is valid for the arbitrary dimension N of spacetime, and thus universal, which is the most important consequence of the metric dimension $1/[L]^2$.

The variation of action leads to the Hamilton–Jacobi equation expressed in terms of the contravariant metric $g^{\mu\nu}$:

$$g^{\mu\nu} \partial_\mu S \partial_\nu S + M^2 = 0. \quad (10)$$

Both terms in Eq.(10) are dimensionless.

III. MINKOWSKI SPACETIME AND PLANCK CONSTANT

Let us introduce the wave vector:

$$k_\mu = \partial_\mu S, \quad [k_\mu] = \frac{1}{[L]}. \quad (11)$$

It has dimension of inverse length and obeys equation:

$$g^{\mu\nu} k_\mu k_\nu + M^2 = 0. \quad (12)$$

Let us now consider this equation in the flat Minkowski spacetime, where the interval becomes:

$$ds_{\text{Mink}}^2 = |g_{00}^{\text{Mink}}|(-dt^2 + d\mathbf{r}^2) \equiv \frac{1}{\hat{h}^2}(-dt^2 + d\mathbf{r}^2). \quad (13)$$

Here we introduced the parameter $\hat{h} \equiv 1/\sqrt{-g_{00}^{\text{Mink}}}$ with dimension of length, $[\hat{h}] = [L]$. In Minkowski spacetime the wave vector obeys equation:

$$\hat{h}^2(\omega^2 - \mathbf{k}^2) = M^2. \quad (14)$$

This suggests that it is natural to identify \hat{h} with Planck constant:¹⁵

$$\hbar \equiv \hat{h} = \frac{1}{\sqrt{-g_{00}^{\text{Mink}}}}. \quad (15)$$

In general, the dimensional metric leads to the difference between the dimensional frequency, $[\omega_{mn}] = 1/[L]$, and the dimensionless mass and energy:

$$[E] = [M] = [\sqrt{g^{00}}][\omega] = [L] \cdot \frac{1}{[L]} = [1] = 0. \quad (16)$$

The red shift equation, $M_m - M_n = \sqrt{-g^{00}} \omega_{mn}$, is the generalization of the conventional relation between the energy levels and frequency of radiating photon in Minkowski vacuum, $E_m - E_n = \hbar \omega_{mn}$. This supports the suggestion that \hbar is the constant conformal factor in a given Minkowski vacuum, $\hbar \equiv \sqrt{-g_{\text{Mink}}^{00}}$, with dimension of length, $[\hbar] = [L]$. As a results the energy in equation $E = \hbar \omega$ is dimensionless, $[E] = [\hbar][\omega] = [1] = 0$.

Note that as distinct from the dimensionless quantities, which are diffeomorphism invariant, the parameter \hbar is not diffeomorphism invariant. It is determined only in the Minkowski vacuum, and being the element of the Minkowski metric it is invariant only under Lorentz transformations. As a result the Planck constant \hbar is not dimensionless, and has the dimension of length $[L]$. Another parameter of Minkowski spacetime is the speed of light c . It is invariant only under space rotation group $SO(3)$, and has dimension $[L]/[T]$. If the parameter c is taken into account, then the Planck constant has dimension of time $[\hbar] = [t]$. Here we do not consider this parameter and use units with $c = 1$.

Since \hbar enters only the Minkowski metric, in general the commutation relations for position and momentum operators in quantum mechanics do not contains \hbar :

$$[\hat{k}_i, \hat{x}^j] = i\delta_i^j. \quad (17)$$

One can also introduce the momentum p^μ :

$$p^\mu = g^{\mu\nu} k_\nu, \quad g_{\mu\nu} p^\mu p^\nu + M^2 = 0. \quad (18)$$

In Minkowski spacetime one obtains:

$$p^\mu = \hbar^2 k_\nu, \quad [p^\mu] = [L]. \quad (19)$$

As distinct from the momentum in the vacua described by the conventional dimensionless metric, this momentum contains \hbar^2 and has dimension $[L]$ instead of $1/[L]$.

IV. SCHRÖDINGER EQUATION IN MINKOWSKI SPACETIME

Let us consider the quadratic terms in the action for the classical scalar field Φ in the N -dimensional spacetime:

$$S = \int d^N x \sqrt{-g} (g^{\mu\nu} \nabla_\mu \Phi^* \nabla_\nu \Phi + M^2 |\Phi|^2), \quad (20)$$

Taking into account equations (5) and (7), one obtains that the scalar field is dimensionless, $[\Phi] = [1] = 0$, for arbitrary spacetime dimension N . This universal zero dimension differs from the N -dependent dimension of scalar fields in the conventional approach, where the dimension is $n = (N - 2)/2$.

Expanding the Klein-Gordon equation for scalar Φ in Eq.(20) over $1/M$ one obtains the non-relativistic Schrödinger action. In Minkowski spacetime, introducing the Schrödinger wave function ψ :

$$\Phi(\mathbf{r}, t) = \frac{1}{\sqrt{M}} \exp\left(iMt/\sqrt{-g^{00}}\right) \psi(\mathbf{r}, t), \quad (21)$$

one obtains the Schrödinger-type action in the form

$$S_{\text{Schr}} = \int d^3x dt \sqrt{-g} \mathcal{L}, \quad (22)$$

$$2\mathcal{L} = i\sqrt{-g^{00}} (\psi \partial_t \psi^* - \psi^* \partial_t \psi) + \frac{g^{ik}}{M} \nabla_i \psi^* \nabla_k \psi. \quad (23)$$

Since in Minkowski vacuum the metric elements play the role of \hbar :

$$\sqrt{-g_{\text{Mink}}^{00}} \equiv \hbar, \quad g_{\text{Mink}}^{ik} \equiv \hbar^2 \delta^{ik}, \quad (24)$$

one obtains the conventional Schrödinger Lagrangian and the Schrödinger wave equation:

$$i\hbar \partial_t \psi = -\frac{\hbar^2}{2M} \nabla^2 \psi. \quad (25)$$

This is another consequence of the metric with dimension $1/[L]^2$: the quantum mechanical Schrödinger equation for nonrelativistic particle is obtained from the classical relativistic scalar field.

The same equation for nonrelativistic fermions can be obtained from classical action for massive Dirac particles:

$$S = \int d^4x e (ie_a^\mu \bar{\Psi} \gamma^a \nabla_\mu \Psi - M \bar{\Psi} \Psi), \quad (26)$$

where e_a^μ are tetrads that are inverse to E_μ^a , with dimension of length, $[e_a^\mu] = [L]$. In the Minkowski spacetime one has

$$e_a^\mu = \hbar \text{diag}(-1, 1, 1, 1), \quad (27)$$

and in the limit $\hbar|\mathbf{k}| \ll M$ one obtains Eq.(25).

The corresponding Hamiltonian for nonrelativistic particle is dimensionless together with energy E :

$$\mathcal{H} = -\frac{\hbar^2}{2M} \nabla^2, \quad [\mathcal{H}] = [\hbar]^2 \cdot [\nabla]^2 \cdot [M]^{-1} = [L]^2 \cdot \frac{1}{[L]^2} \cdot [1] = [1] = 0. \quad (28)$$

If to follow the elasticity tetrad scenario in Eq. (1), then \hbar with its dimension of length, $[\hbar] = [L]$, plays the role of the "interatomic" distance in the medium called the relativistic quantum vacuum. Then the Schrödinger equation (25) can be valid only in the long wavelength limit, $\hbar|\mathbf{k}| \ll M \ll 1$. Whether this "Planck constant length" is related to the "Planck length scale", is an open question.¹⁶

V. DE SITTER SPACETIME AND PLANCK CONSTANT

The same conformal factor in terms of \hbar is valid for any $D + 1$ Minkowski spacetime including the 4+1 Minkowski spacetime, which determines the de Sitter spacetime:

$$g_{\text{dS}}^{\mu\nu} X^\mu X^\nu = \alpha^2. \quad (29)$$

Here α is dimensionless constant, and $g_{\text{dS}}^{\mu\nu} = 1/\hbar^2$. This equation gives the Hubble parameter:

$$H = \frac{1}{\hbar\alpha}, \quad [H] = \frac{1}{[L]}. \quad (30)$$

The dS metric in the Paineve-Gullstrand form contains two parameters, \hbar and H (actually three parameters if the speed of light is included):

$$ds^2 = \frac{1}{\hbar^2} (-dt^2 + (dr - Hrdt)^2 + r^2 d\Omega^2) . \quad (31)$$

At $r = 0$ the metric is Minkowski, however it is not excluded that in the dS universe the parameter \hbar deviates from its value in Minkowski vacuum and depends on α .

The probability of Hawking radiation of particle with mass M detected by observer at $r = 0$ is

$$w \propto \exp\left(-\frac{2\pi M}{H\sqrt{-g_{\text{Mink}}^{00}}}\right) = \exp\left(-\frac{2\pi M}{\hbar H}\right) . \quad (32)$$

VI. PLANCK CONSTANT AND TOLMAN LAW

Eq.(32) corresponds to the following Hawking temperature:

$$T(r = 0) = \frac{\sqrt{-g_{\text{Mink}}^{00}}H}{2\pi} = \frac{\hbar H}{2\pi} . \quad (33)$$

The Hawking temperature is dimensionless, $[T(r = 0)] = [L] \cdot 1/[L] = [1] = 0$. Note that constant $H/2\pi$ plays the role of Tolman temperature, which enters the Tolman law:

$$T(r) = \frac{T_{\text{Tolman}}}{\sqrt{-g_{00}(r)}} , \quad T_{\text{Tolman}} = \frac{H}{2\pi} . \quad (34)$$

While temperature is dimensionless, $[T] = [M] = 0$, the Tolman temperature has dimension of inverse length, $[T_{\text{Tolman}}] = [H] = 1/[L]$, see also Refs.^{1,2}.

The parameter \hbar determines the ratio between the temperature and Tolman temperature in the Minkowski vacuum. In principle there can be different Minkowski vacua with their own values of \hbar . Then the Tolman law can be applied to the boundary between the two Minkowski vacua with different values of \hbar . In thermal equilibrium they must have the same Tolman temperature, and thus their temperatures obey the rule, $\hbar_1/T_1 = \hbar_2/T_2$. This means that in thermal equilibrium the contacting Minkowski vacua have the same time on imaginary axis, $\tau_1 = \tau_2$.

VII. LENGTH DIMENSION OF NEWTON CONSTANT AND PLANCK LENGTH

In previous works we considered the Newton constant G as dimensionless.^{1,2} This is because we did not take into account the dimension of \hbar . Now, since the Planck constant has dimension of length, the Newton constant also acquires the dimension of length, $[G] = [L]$, and the dimensionless gravitational action has the following form:

$$S = \int d^4x \sqrt{-g} \left(\frac{\hbar}{16\pi G} R + \beta \mathcal{R}^2 + \dots \right) + \gamma \int d^4x E^a \wedge E^b \wedge R_{ab}. \quad (35)$$

Here the $\beta \mathcal{R}^2$ means all the terms which are quadratic in curvature tensor $\mathcal{R}_{\alpha\beta\mu\nu}$, and the last term is Barbero-Immirzi action. The scalar curvature R is dimensionless, $[R] = [1] = 0$, as well as all \mathcal{R}^2 terms, and the parameters β are dimensionless. The parameter γ , which enters the Barbero-Immirzi term, is also dimensionless due to Eq.(2).

As distinct from the $\beta \mathcal{R}^2$ terms and Barbero-Immirzi term, the first term in Eq.(35) explicitly contains the Planck constant \hbar , which demonstrates the quantum mechanical origin of the Einstein–Hilbert action. The Planck length scale has the conventional form $l_{\text{Planck}}^2 = \hbar G$, with $[l_{\text{Planck}}]^2 = [\hbar][G] = [L][L] = [L]^2$. Since the Planck constant \hbar has the same dimension as the Planck length l_{Planck} , this also suggests the close connection between gravity and quantum mechanics.

VIII. CONCLUSION

The important consequence of the metric dimension $1/[L]^2$ is that all the diffeomorphism invariant quantities are dimensionless for any dimension of spacetime. These include the action S , interval s , Newton constant G , cosmological constant Λ , scalar curvature R , scalar field Φ , etc.

Another consequence of the metric dimension $1/[L]^2$, is that the Planck constant \hbar is the property of the Minkowski vacuum. Being the element of the Minkowski metric, \hbar is invariant only under Lorentz transformations, and is not diffeomorphism invariant. As a result the Planck constant \hbar is not dimensionless, and has the dimension of length $[L]$. Whether this Planck constant length \hbar is related to the Planck length scale $\sqrt{\hbar G}$ is an open question. It is possible that the dimensionless quantity \hbar/G can be the fundamental constant, which reflects the symmetry or/and topology of the quantum vacuum.²

In principle it is not excluded that there can be different Minkowski vacua with their own values of the parameter \hbar . Then the thermal contact between the two vacua obeys the Tolman law: in thermal equilibrium their temperatures are connected in the following way, $\hbar_1/T_1 = \hbar_2/T_2$.

All these consequences of the metric with dimension $1/[L]^2$ suggest that such metric describes the dynamics, quantum mechanics and thermodynamics, rather than the geometry.

Acknowledgements. I thank Yuri Obukhov for discussion. This work has been supported by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (Grant Agreement No. 694248).

-
- ¹ G.E. Volovik, Dimensionless physics, *ZhETF* **159**, 815–821 (2021), *JETP* **132**, 727–733 (2021), arXiv:2006.16821.
- ² G.E. Volovik, Dimensionless physics: continuation, accepted in *JETP*, arXiv:2207.05754 [gr-qc].
- ³ F.R. Klinkhamer and G.E. Volovik, Tetrads and q -theory, *Pis’ma ZhETF* **109**, 369–370 (2019), *JETP Lett.* **109**, 364–367 (2019), arXiv:1812.07046.
- ⁴ I.E. Dzyaloshinskii and G.E. Volovick, Poisson brackets in condensed matter physics, *Ann. Phys.* **125**, 67–97 (1980).
- ⁵ J. Nissinen and G.E. Volovik, Elasticity tetrads, mixed axial-gravitational anomalies, and (3+1)-d quantum Hall effect, *Phys. Rev. Research* **1**, 023007 (2019), arXiv:1812.03175.
- ⁶ J. Nissinen, Field theory of higher-order topological crystalline response, generalized global symmetries and elasticity tetrads, arXiv:2009.14184 [cond-mat.str-el].
- ⁷ J. Nissinen, Emergent spacetime and gravitational Nieh-Yan anomaly in chiral $p + ip$ Weyl superfluids and superconductors, *Phys. Rev. Lett.* **124**, 117002 (2020).
- ⁸ S. Laurila and J. Nissinen, Torsional Landau levels and geometric anomalies in condensed matter Weyl systems, *Phys. Rev. B* **102**, 235163 (2020).
- ⁹ L. Gioia, Chong Wang, and A.A. Burkov, Unquantized anomalies in topological semimetals, *Phys. Rev. Research* **3**, 043067 (2021).
- ¹⁰ D. Diakonov, Towards lattice-regularized Quantum Gravity, arXiv:1109.0091.
- ¹¹ A.A. Vladimirov and D. Diakonov, Phase transitions in spinor quantum gravity on a lattice, *Phys. Rev. D* **86**, 104019 (2012).

- ¹² A.A. Vladimirov and D. Diakonov, Diffeomorphism-invariant lattice actions, *Physics of Particles and Nuclei* **45**, 800 (2014).
- ¹³ Y.N. Obukhov and F.W. Hehl, Extended Einstein–Cartan theory a la Diakonov: The field equations, *Phys. Lett. B* **713**, 321–325 (2012).
- ¹⁴ K. Akama, An Attempt at Pregeometry: Gravity with Composite Metric, *Progress of Theoretical Physics* **60**, 1900–1909 (1978).
- ¹⁵ G.E. Volovik, \hbar as parameter of Minkowski metric in effective theory, *Pis'ma ZhETF* **90**, 793–799 (2009); *JETP Lett.* **90**, 697–704 (2009); arXiv:0904.1965.
- ¹⁶ S. Carlip, Spacetime foam: a review, arXiv:2209.14282