

Global hyperbolicity through the eyes of the null distance

Annegret Burtscher* and Leonardo García-Heveling*

October 18, 2022

Abstract

No Hopf–Rinow Theorem is possible in Lorentzian Geometry. Nonetheless, we prove that a spacetime is globally hyperbolic if and only if it is metrically complete with respect to the null distance of a time function. Our approach is based on the observation that null distances behave particularly well for temporal functions, leading also to bi-Lipschitz equivalences and the encoding of the causal relation.

2020 *Mathematics Subject Classification*: 53C50 (primary), 53C23, 53B30, 83C05 (secondary).

Keywords: globally hyperbolic, spacetimes, time functions, temporal functions, null distance, completeness, causality.

1 Introduction

The notion of global hyperbolicity was introduced by Leray [34] in 1952 to prove the uniqueness of solutions for hyperbolic partial differential equations. Shortly thereafter, global hyperbolicity entered the field of General Relativity through the proof of the well-posedness of the Einstein equations of Choquet–Bruhat [21] and the Singularity Theorems of Penrose and Hawking [25, 43]. Via the topological splitting result of Geroch [23] globally hyperbolic spacetimes manifestly settled in Lorentzian Geometry in 1970.

*Department of Mathematics, Institute for Mathematics, Astrophysics and Particle Physics (IMAPP), Radboud University Nijmegen, The Netherlands

Emails: burtscher@math.ru.nl, l.heveling@math.ru.nl

Acknowledgements: This manuscript was completed during an extended research stay at the Fields Institute for Research in Mathematical Sciences in Toronto, in connection with the thematic program "Nonsmooth Riemannian and Lorentzian Geometry" that took place in the Fall of 2022. Both authors gratefully acknowledge funding by the Fields Institute during their stay. AB's research is in part also supported by the Dutch Research Council (NWO), Project number VI.Veni.192.208.

Spacetimes are time-oriented Lorentzian manifolds (M, g) and are the underlying objects in General Relativity. Global hyperbolicity establishes a deep link between the topology of M and the causal structure induced by the metric tensor g . The causal preorder is induced by causal curves, i.e., curves γ with $g(\dot{\gamma}, \dot{\gamma}) \leq 0$. If q can be reached by a future-directed causal curve from p we say that q is in the causal future of p , and write $q \in J^+(p)$ (dually for the causal past $J^-(p)$). Leray's original definition of global hyperbolicity was based on the C^0 -compactness of the set of causal curves between any two points in a spacetime. The modern definition of global hyperbolicity requires compactness of causal diamonds $J^+(p) \cap J^-(q)$ akin to the Heine–Borel property for complete Riemannian manifolds (see [25, Sec. 6.6] and [10]).

Definition 1.1. A spacetime (M, g) is called *globally hyperbolic* if it is *causal* (there is no closed causal curve) and all causal diamonds $J^+(p) \cap J^-(q)$, $p, q \in M$, are compact.

If (M, g) is a noncompact spacetime of dimension $\dim M \geq 3$ then the causal condition can be dropped [27].

A landmark result concerning global hyperbolicity is Geroch's Topological Splitting Theorem [23], later promoted to a smooth orthogonal splitting by Bernal and Sánchez [9]. It states that any globally hyperbolic spacetime (M, g) admits a *Cauchy orthogonal splitting*, i.e., an isometry

$$(M, g) \cong (\mathbb{R} \times \Sigma, -\alpha d\tau^2 + \bar{g}_\tau), \quad (1)$$

for $\alpha: \mathbb{R} \times \Sigma \rightarrow (0, \infty)$ a smooth function and $(\bar{g}_\tau)_\tau$ identifiable with a family of Riemannian metrics on the Cauchy slices $\{\tau\} \times \Sigma$, smoothly varying in τ . The proof of this splitting result is rooted in the construction of a suitable time function τ .

Definition 1.2. Let (M, g) be a spacetime. A continuous function $\tau: M \rightarrow \mathbb{R}$ is said to be a *time function* if

$$q \in J^+(p) \setminus \{p\} \implies \tau(p) < \tau(q).$$

Every *stably causal* spacetime admits a (non-unique) time function [26, 35, 36]. The class of globally hyperbolic spacetimes can conveniently be characterized by the special type of time functions they admit.

Theorem 1.3 (Geroch [23], Bernal–Sánchez [9]). *A spacetime (M, g) is globally hyperbolic if and only if there exists a Cauchy time function τ on M , meaning that each of its level sets $\tau^{-1}(s)$, $s \in \mathbb{R}$, is intersected (exactly once) by every inextendible causal curve.*

From the compactness condition in Definition 1.1 it follows by a result obtained independently by Avez [5] and Seifert [46] that there exists a

length-maximizing geodesic between any two causally related points (length-maximizing with respect to the Lorentzian distance, which is then also finite-valued and continuous [8, Ch. 4]). In that sense globally hyperbolic spacetimes again resemble complete Riemannian manifolds. But here the analogy ends. The Lorentzian distance is far from inducing a metric space structure. Even more troubled is the relationship with geodesic completeness. Neither does global hyperbolicity imply geodesic completeness (the famous Penrose Singularity Theorem [43] actually shows incompleteness under additional curvature bounds) nor the other way round (anti-de Sitter space). In both cases these are actually features rather than bugs of Lorentzian manifolds, and physically highly desired, for instance, for the mathematical existence of black holes. Nonetheless, even the physically undesired assumption of compactness does not guarantee geodesic completeness (Clifton–Pohl torus). Altogether these properties render any Hopf–Rinow type statement for spacetimes virtually a lost case (see the early works of Busemann [16], Beem [6]; and [7, 17, 19, 24] for work on completeness of spacelike submanifolds). We reopen the case and characterize global hyperbolicity in a profoundly new way.

Theorem 1.4. *A spacetime (M, g) is globally hyperbolic if and only if there exists a time function τ such that (M, \hat{d}_τ) is a complete metric space.*

Here \hat{d}_τ is the null distance of Sormani and Vega [48], defined in 2016 with the purpose of studying geometric stability problems in General Relativity by means of a metric (measure) convergence theory, and to develop robust tools for spacetimes of low regularity (partly already realized in [1, 14, 29, 44, 48, 50]). The $\hat{\cdot}$ in \hat{d}_τ indicates the dependence on the causal cone structure and τ the link to the time function.

Definition 1.5. Let (M, g) be a spacetime with time function τ . A *piecewise causal path* $\beta: [a, b] \rightarrow M$ is given by a partition $a = s_0 < s_1 < \dots < s_{k-1} < s_k = b$ on which each restriction $\beta|_{[s_{i-1}, s_i]}$ is either a future- or past-directed causal curve. The *null length* of β is given by

$$\hat{L}_\tau(\beta) := \sum_{i=1}^k |\tau(\beta(s_i)) - \tau(\beta(s_{i-1}))|.$$

The *null distance* between two points $p, q \in M$ is

$$\hat{d}_\tau(p, q) := \inf\{\hat{L}_\tau(\beta) \mid \beta \text{ piecewise causal path between } p \text{ and } q\}.$$

For locally anti-Lipschitz time functions we obtain a conformally invariant length-metric space (M, \hat{d}_τ) that induces the manifold topology (see [48, Thm. 4.6] and [1, Thm. 1.1]). Throughout most of this manuscript we will assume even more, namely that τ is a temporal function.

Definition 1.6. Let (M, g) be a spacetime. A *temporal function* is a smooth function $\tau: M \rightarrow \mathbb{R}$ with past-directed timelike gradient $\nabla\tau$.

This is no restriction since every spacetime that admits a time function also admits a temporal function [9, Thm. 1.2]. The main advantage of temporal functions over other time functions is the orthogonal decomposition $g = -\alpha d\tau^2 + \bar{g}_\tau$ (although no product splitting of the manifold) due to Müller and Sánchez [40, Lem. 3.5]. This property is key in the proof of the following result.

Theorem 1.7. *Let (M, g) be a spacetime, τ be a temporal function, and h be a Riemannian metric on M . Then, for each compact set $K \subseteq M$, there exists a constant $C \geq 1$ such that for all $p, q \in K$,*

$$\frac{1}{C}d_h(p, q) \leq \hat{d}_\tau(p, q) \leq Cd_h(p, q).$$

In fact, we prove Theorem 1.7 for a larger class of time functions, including regular cosmological time functions à la Andersson–Galloway–Howard [4] and Wald–Yip [51]. This result immediately implies that two stably causal spacetimes are metrically equivalent on compact sets in the following sense.

Corollary 1.8. *Let M be a smooth manifold and g and \tilde{g} be spacetime metrics on M with temporal functions τ and $\tilde{\tau}$ (and corresponding null distances \hat{d}_τ and $\hat{d}_{\tilde{\tau}}$), respectively. Then, for each compact set $K \subseteq M$, there exists a constant $C \geq 1$ such that for all $p, q \in K$,*

$$\frac{1}{C}\hat{d}_\tau(p, q) \leq \hat{d}_{\tilde{\tau}}(p, q) \leq C\hat{d}_\tau(p, q).$$

Corollary 1.8 can be viewed as a Lorentzian version of the Riemannian distance theorem obtained by Burtscher [15, Thm. 4.5]. On a global scale the situation is, of course, quite different and it is here where globally hyperbolic spacetimes really shine. We prove the following.

Theorem 1.9. *Let (M, g) be a globally hyperbolic spacetime and τ a Cauchy temporal function. Then the null distance encodes causality, that is, for any $p, q \in M$,*

$$q \in J^+(p) \iff \hat{d}_\tau(p, q) = \tau(q) - \tau(p). \quad (2)$$

The \implies direction in Theorem 1.9 is trivial. Sormani and Vega [48, Thm. 3.25] showed that the converse holds for warped product spacetimes with complete Riemannian fiber. It remained an open problem to determine under which general circumstances causality is encoded. Our Theorem 1.9 provides an answer for Cauchy temporal functions, and is sharp in terms of the causality class of spacetimes allowed (Example 3.8). Independently and simultaneously, Sakovich and Sormani [44, Thm. 4.1] have obtained a

different global causality encoding result where they allow for generalized anti-Lipschitz time functions, but require them to be proper. This properness assumption, in fact, implies global hyperbolicity (see Remarks 3.5 and 3.9). Both approaches yield local encoding of causality on any stably causal spacetime (see Theorem 3.3 and [44, Thm. 1.1]).

The manuscript is structured as follows. In Section 2 we prove a generalization of Theorem 1.7 and Corollary 1.8 for the larger class of admissible time functions (see Definition 2.1). In the proof we make use of the orthogonal decomposition of the spacetime metric with respect to a temporal function due to Müller and Sánchez [40] and techniques developed by Burtscher [15]. We also present counterexamples that show that the admissibility condition cannot be relaxed below locally Lipschitz and locally anti-Lipschitz. In Section 3 we prove Theorem 1.9 and provide counterexamples for spacetimes on lower steps of the causal ladder. Nonetheless, a local result for any temporal function on any stably causal spacetime is also obtained. In Section 4 we prove Theorem 1.4 and show that h -steep Cauchy temporal functions (recently independently introduced by Bernard and Suhr [11] and Minguzzi [37]) guarantee completeness of (M, \hat{d}_τ) . The different sections are independent of each other, except for some technical lemmas of Section 2 appearing again in Section 3.

2 Bi-Lipschitz bounds

In this section we prove Theorem 1.7 and Corollary 1.8 of the introduction. In fact, we prove these results for the larger class of admissible time functions of Definition 2.1.

Throughout this manuscript, we use the sign convention $(-, +, \dots, +)$ for g , that is, a tangent vector $v \in T_p M \setminus \{0\}$ is causal precisely when $g_p(v, v) \leq 0$ and timelike when $g_p(v, v) < 0$. For *causal curves* we use the regularity convention of Chruściel [18, Sec. 2.3]. More precisely, a *future-directed causal path* γ is assumed to be locally Lipschitz with respect to any Riemannian metric with tangent vector $\dot{\gamma}$ being future-directed causal almost everywhere (the existence already follows from Rademacher’s Theorem). A curve is an equivalence class of paths with respect to orientation-preserving reparametrizations. Note that Chruściel defined the local Lipschitz condition only with respect to one complete Riemannian metric, but due to its local nature the statement is independent of the choice of an (even incomplete) Riemannian metric by [15, Thm. 4.5].

We furthermore denote the *causal relation* $J^+ \subseteq M \times M$ by

$$J^+ := \{(p, q) \mid \text{there exists a future-directed causal curve from } p \text{ to } q\},$$

the “+” indicating that we consider future-directed causal curves. This is a

preorder often also denoted by “ \leq ”, in order words,

$$(p, q) \in J^+ \iff q \in J^+(p) \iff p \leq q.$$

Definition 2.1. Let (M, g) be a spacetime with time function $\tau: M \rightarrow \mathbb{R}$. Let h be any Riemannian metric on M and d_h the associated distance function. If for every point x there exists a neighborhood U of x and $C \geq 1$ such that

$$(p, q) \in J^+ \cap (U \times U) \implies \frac{1}{C}d_h(p, q) \leq \tau(q) - \tau(p) \leq Cd_h(p, q), \quad (3)$$

then we say that τ is an *admissible time function*.

Note that the first \leq condition in (3) is called *locally anti-Lipschitz* by Sormani and Vega and always assumed in statements about the null distance to guarantee that it is definite [48, Thm. 4.6] (by Burtscher [15, Thm. 4.5] the choice of h is irrelevant, see also [1, Rem. 3.2]). The second \leq in (3) is a mild additional *locally Lipschitz* assumption for causally related points and satisfied by a large class of time functions. In follow-up sections we will only use temporal functions. Since time functions are, however, also a useful tool in weaker geometric settings (see, for instance, [12, 14, 29]) where differentiability is not available but locally Lipschitz properties are, we decided to present the results here in more generality.

Lemma 2.2. *Temporal functions and regular cosmological time functions (à la Andersson–Galloway–Howard [4] and Wald–Yip [51]) are admissible time functions.*

Proof. For temporal functions, locally Lipschitz follows by smoothness and locally anti-Lipschitz by [48, Cor. 4.16]. For regular cosmological time functions, locally Lipschitz follows by [4, Thm. 1.2(v)] and locally anti-Lipschitz by [48, Thm. 5.4]. \square

Since $\hat{d}_\tau(p, q) = \tau(q) - \tau(p)$ for causally related points, the condition (3) can be viewed as a local metric equivalence. In this section we show that it can be extended to all points in any compact set.

Theorem 2.3. *Let (M, g) be a spacetime, $\tau: M \rightarrow \mathbb{R}$ be an admissible time function, and h be a Riemannian metric on M . Then, for each compact set $K \subseteq M$, there exists a constant $C \geq 1$ such that for all $p, q \in K$,*

$$\frac{1}{C}d_h(p, q) \leq \hat{d}_\tau(p, q) \leq Cd_h(p, q). \quad (4)$$

The proof of Theorem 2.3 makes use of several technical results related to temporal functions obtained in Section 2.1, some of which are also used in Section 3. The proof of Theorem 2.3 is carried out subsequently in Section 2.2.

Theorem 2.3 implies that we can compare the null distances of any two spacetime metrics with entirely different causal cones as follows.

Corollary 2.4. *Let M be a smooth manifold and g and \tilde{g} be spacetime metrics on M with admissible time functions τ and $\tilde{\tau}$ (and corresponding null distances \hat{d}_τ and $\hat{d}_{\tilde{\tau}}$), respectively. Then, for each compact set $K \subseteq M$, there exists a constant $C \geq 1$ such that for all $p, q \in K$,*

$$\frac{1}{C}\hat{d}_\tau(p, q) \leq \hat{d}_{\tilde{\tau}}(p, q) \leq C\hat{d}_\tau(p, q).$$

Remark 2.5. Bi-Lipschitz maps are a crucial tool in Metric Geometry. In particular, they are useful in the study of Gromov–Hausdorff convergence of metric spaces. Moreover, the estimates of Theorem 1.7 can be used to equip suitable spacetimes with a (local) integral current space structure (see [3, 20, 28, 31, 32, 49]) via the approach laid out in Allen and Burtscher [1, Sec. 2.5, 2.6, and 4]. Thanks to our Completeness Theorem 1.4 and [1, Thm. 1.3] we know, for instance, that any globally hyperbolic spacetime viewed as the metric space (M, \hat{d}_τ) with an h -steep temporal function τ is a (local) integral current space (see Section 4). This also makes it possible to use spacetime intrinsic flat convergence to study geometric stability questions in General Relativity, as proposed by Sormani.

In particular, the null distance structures with respect to admissible time functions are equivalent on compact sets of a fixed spacetime.

Corollary 2.6. *Let τ_1, τ_2 be two admissible time functions on a spacetime (M, g) , and $\hat{d}_{\tau_1}, \hat{d}_{\tau_2}$ their associated null distances. Then, for every compact set $K \subseteq M$, there exists a constant $C \geq 1$ such that for all $p, q \in K$,*

$$\frac{1}{C}\hat{d}_{\tau_1}(p, q) \leq \hat{d}_{\tau_2}(p, q) \leq C\hat{d}_{\tau_1}(p, q). \quad (5)$$

Note that Corollary 2.6 was already announced in [1, p. 7739] with an alternative direct proof. The advantage of such a proof is that it does not require the use of temporal functions which, for instance, do not exist in theory of Lorentzian length spaces [14] (while admissible time functions only require local Lipschitz conditions and can still be considered). We therefore provide a short alternative proof of Corollary 2.6 in Section 2.3.

The question remains how optimal all the results just mentioned are with respect to the class of time functions considered. The lower bound in (3) is the standard local anti-Lipschitz assumption on τ and needed to even obtain a sensible metric space (M, \hat{d}_τ) [48, Thm. 4.6]. In Section 2.4 we show that the locally Lipschitz assumption (the upper bound in (3)) cannot be dropped either. Moreover, we show that even in the best of (noncompact) circumstances, a general global version of our weakest result presented in this section, namely Corollary 2.6, cannot be expected. This implies that also Theorem 2.3 and Corollary 2.4 in general do not hold globally.

2.1 Temporal functions and Wick-rotated metrics

The aim of this section is to compare the null distance to the distance obtained with respect to the Wick-rotated metric that exists for a temporal function. Recall that a spacetime admits a smooth temporal function whenever it admits a time function by a well-known result of Bernal and Sánchez [9, Thm. 1.2]. We then make use of a weaker splitting result for temporal functions (compare to (1) in the globally hyperbolic case).

Lemma 2.7 (Müller–Sánchez [40, Lem. 3.5]). *If a spacetime (M, g) admits a temporal function τ , then the metric g admits an orthogonal decomposition*

$$g = -\alpha d\tau^2 + \bar{g}, \quad (6)$$

where $\alpha = |g(\nabla\tau, \nabla\tau)|^{-1} > 0$ and \bar{g} is a symmetric 2-tensor which vanishes on $\nabla\tau$ and is positive definite on the complement.

The temporal function τ is said to be *steep* if $g(\nabla\tau, \nabla\tau) \leq -1$, and hence $\alpha = |g(\nabla\tau, \nabla\tau)|^{-1} \leq 1$. Not every (even causally simple) spacetime admits a steep temporal function [40, Thm. 1.1 and Ex. 3.3], but we can always rewrite (6) as $g = -\alpha(d\tau^2 + \tilde{g})$, for $\tilde{g} = \alpha^{-1}\bar{g}$. Since the null distance is conformally invariant we assume from now on, without loss of generality, that g is of the form

$$g = -d\tau^2 + \bar{g}. \quad (7)$$

Next we perform a standard trick to obtain a Riemannian metric g_W from g and τ . This technique is called Wick-rotation in the physics literature. The *Wick-rotated metric* g_W is given by

$$g_W := d\tau^2 + \bar{g}. \quad (8)$$

We denote the associated norm, length functional, and Riemannian distance by $\|\cdot\|_W$, L_W , and d_W respectively. We proceed to compare d_W and \hat{d}_τ .

Lemma 2.8. *Let (M, g) be a spacetime equipped with a smooth temporal function τ , so that g and g_W are given by (7) and (8), respectively. Let $\beta: [a, b] \rightarrow M$ be a piecewise causal curve. Then*

$$\hat{L}_\tau(\beta) \leq L_W(\beta) \leq \sqrt{2}\hat{L}_\tau(\beta),$$

and thus for any $p, q \in M$,

$$d_W(p, q) \leq \sqrt{2}\hat{d}_\tau(p, q). \quad (9)$$

Proof. By assumption τ is smooth and also the tangent vector $\dot{\beta}$ exists almost everywhere, thus we can write both length functionals in terms of

the integrals

$$\begin{aligned}\hat{L}_\tau(\beta) &= \int_a^b |(\tau \circ \beta)'(s)| ds, \\ L_W(\beta) &= \int_a^b \|\dot{\beta}(s)\|_W ds = \int_a^b \sqrt{|d\tau(\dot{\beta}(s))|^2 + \|\dot{\beta}(s)\|_g^2} ds.\end{aligned}$$

Since $\dot{\beta}^\tau(s) := (\tau \circ \beta)'(s) = d\tau_{\beta(s)}(\dot{\beta}(s))$ and the curve β is piecewise causal, that is, $g(\dot{\beta}, \dot{\beta}) \leq 0$ almost everywhere, we have that

$$|\dot{\beta}^\tau| \geq \|\dot{\beta}\|_{\hat{g}},$$

and the inequalities for the lengths follow immediately.

The second inequality for lengths descends to the level of distances, because the class of piecewise causal curves considered (for \hat{d}_τ) is contained in the class of all locally Lipschitz curves (strictly speaking, d_W is obtained via piecewise smooth curves, but [15, Cor. 3.13] shows that it is the same as the intrinsic metric obtained via the class of absolutely continuous curves, each of which has a locally Lipschitz reparametrization [2, Lem. 1.1.4]). \square

The next lemma proves a reverse inequality to (9), albeit only locally and via a more involved proof (because d_W requires knowledge also about non-causal curves).

Lemma 2.9. *Let (M, g) be a spacetime equipped with a smooth temporal function τ , so that g and g_W are given by (7) and (8), respectively. Then, around every $x \in M$ there is a neighborhood U such that for all $p, q \in U$,*

$$\hat{d}_\tau(p, q) \leq 4d_W(p, q).$$

Proof. Let $x \in M$ and $\dim M = N + 1$. The idea is to use coordinates adapted to the vector field $\nabla\tau$ and the hypersurface $\mathcal{S}_{\tau(x)} := \tau^{-1}(\tau(x))$ to explicitly construct approximating piecewise causal curves β_n in a neighborhood U of x .

Step 1. Construction of U and coordinates. The level set $\mathcal{S}_{\tau(x)}$ is a hypersurface in M (since $d\tau \neq 0$), therefore there exists a neighborhood $V \subseteq \mathcal{S}_{\tau(x)}$ of x and a coordinate map $\varphi: V \rightarrow \varphi(V) \subseteq \mathbb{R}^N$. Let $y \in \varphi(V) \subseteq \mathbb{R}^N$ and let c_y denote the unique integral curve of $-\nabla\tau$ in M with $c_y(0) \in V \subseteq \mathcal{S}_{\tau(x)} \subseteq M$ such that $\varphi(c_y(0)) = y$. Since we have assumed that $d\tau(-\nabla\tau) = -g(\nabla\tau, \nabla\tau) = 1$, we have that

$$\tau(c_y(t)) = \tau(x) + t. \tag{10}$$

The Flow Box Theorem [33, Thm. 2.91] guarantees the existence of a suitable $a > 0$ and a neighborhood of x (which we immediately restrict to our chart

neighborhood on $\mathcal{S}_{\tau(x)}$ and again denote by V) such that the map

$$\begin{aligned}\phi: (-a, a) \times \varphi(V) &\rightarrow M, \\ (t, y) &\mapsto c_y(t),\end{aligned}$$

is well-defined. By the Inverse Function Theorem, ϕ is a local diffeomorphism, hence one obtains a coordinate system (t, y) on a neighborhood U of x in M . At any point $z = (t, y) \in \phi^{-1}(U) \subseteq (-a, a) \times \mathbb{R}^N$, the differential $D_z\phi: \mathbb{R} \times \mathbb{R}^N \rightarrow T_{\phi(z)}M$ maps $(1, 0, \dots, 0)$ to $\dot{c}_y(t) = \nabla\tau|_{\phi(z)}$. Thanks to (10), $D_z\phi$ also maps $\{0\} \times \mathbb{R}^N$ to $T_{\phi(z)}\mathcal{S}_{\tau(\phi(z))} \subseteq T_{\phi(z)}M$. Since $T_{\phi(z)}\mathcal{S}_{\tau(\phi(z))} = (\text{span } \nabla\tau|_{\phi(z)})^\perp$, pulling back the metric tensor g with ϕ , we see that its components in the (t, y) coordinates are

$$g_{00} = -1 \qquad g_{0i} = 0 \quad \text{for all } i \neq 0, \qquad (11)$$

where the subindex 0 corresponds to the t -component and the subindex i to any of the y -components. Finally, by further shrinking U , we may assume that it is g_W -convex.

Step 2. Construction of piecewise causal curves β_n approximating $d_W(p, q)$. Let $p, q \in U$ be given. By assumption on U , there exists a length-minimizing g_W -geodesic $\gamma: [0, L] \rightarrow U$ from p to q , parametrized by arclength. In particular,

$$L = L_W(\gamma) = d_W(p, q).$$

We write $\gamma = (\gamma^\tau, \gamma^1, \dots, \gamma^N)$ to denote the coordinate expression of γ . Consider the sequence $(\beta_n)_n$ of curves given in coordinates by

$$\beta_n(s) := (\gamma^\tau(s) + 3f_n(s), \gamma^1(s), \dots, \gamma^N(s)), \qquad (12)$$

for $|f_n|$ sufficiently small so that β_n is contained in U . Furthermore, we choose $(f_n)_n$ in a way that β_n is piecewise causal, that is,

$$g(\dot{\beta}_n, \dot{\beta}_n) \leq 0 \quad \text{almost everywhere,}$$

and so that $\hat{L}_\tau(\beta_n) \leq 4L_W(\gamma)$ for n sufficiently large.

We show that the functions $f_n: [0, L] \rightarrow \mathbb{R}$, given by

$$\begin{aligned}f_0(s) &:= \begin{cases} s & \text{for } s \in [0, \frac{L}{2}], \\ L - s & \text{for } s \in [\frac{L}{2}, L], \end{cases} \quad \text{and} \\ f_n(s) &:= \frac{1}{2} \begin{cases} f_{n-1}(2s) & \text{for } s \in [0, \frac{L}{2}], \\ f_{n-1}(2s - L) & \text{for } s \in [\frac{L}{2}, L], \end{cases}\end{aligned}$$

as depicted in Figure 1a satisfy all these properties. The curves β_n given by (12) are shown in Figure 1b. By definition, $\dot{\beta}_n$ exists almost everywhere, and by (11) it follows that

$$\begin{aligned} g(\dot{\beta}_n, \dot{\beta}_n) &= -|\dot{\beta}_n^\tau|^2 + \bar{g}(\dot{\beta}_n, \dot{\beta}_n) \\ &= -|\dot{\gamma}_n^\tau + 3f_n'|^2 + (\bar{g}_{ij} \circ \beta_n)\dot{\gamma}^i\dot{\gamma}^j. \end{aligned} \quad (13)$$

Since $|f_n'| = 1$ almost everywhere, and $|\dot{\gamma}_n^\tau| \leq g_W(\dot{\gamma}, \dot{\gamma}) = 1$, it follows that almost everywhere

$$2 \leq 3|f_n'| - |\dot{\gamma}_n^\tau| \leq |\dot{\gamma}_n^\tau + 3f_n'| \leq 3|f_n'| + |\dot{\gamma}_n^\tau| \leq 4. \quad (14)$$

We consider the second term in (13). Since γ is a g_W -geodesic it is smooth and so are all coordinate components $\dot{\gamma}^i$ of tangent vector. Due to compactness of γ the $\dot{\gamma}^i$ must therefore also be bounded, say $|\dot{\gamma}^i| \leq M$ for some $M > 0$. Due to $|f_n(s)| \leq \frac{L}{2^{n+1}} \rightarrow 0$ uniformly as $n \rightarrow \infty$ it follows that $\beta_n^\tau \rightarrow \gamma^\tau$ uniformly, and thus $\beta_n \rightarrow \gamma$ uniformly in coordinates. The coordinate functions $\bar{g}_{ij} \circ \beta_n \rightarrow h_{ij} \circ \gamma$ converge uniformly on $[0, L]$ as $n \rightarrow \infty$ as well. Thus for any $\varepsilon \in (0, \frac{1}{M^2})$ there is an $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$, for all $s \in [0, L]$,

$$\begin{aligned} &\bar{g}_{ij}(\beta_n(s))\dot{\gamma}^i(s)\dot{\gamma}^j(s) \\ &\leq \bar{g}_{ij}(\gamma(s))\dot{\gamma}^i(s)\dot{\gamma}^j(s) + |\bar{g}_{ij}(\beta_n(s)) - \bar{g}_{ij}(\gamma(s))||\dot{\gamma}^i(s)||\dot{\gamma}^j(s)| \\ &\leq g_W(\dot{\gamma}(s), \dot{\gamma}(s)) + \varepsilon M^2 \leq 2. \end{aligned} \quad (15)$$

Together, (13)–(15) imply

$$g(\dot{\beta}_n, \dot{\beta}_n) \leq -2^2 + 2 \leq 0,$$

hence all β_n for $n \geq n_0$ are piecewise causal.

Using again that (14) holds almost everywhere, it follows that for all n


$$\hat{L}_\tau(\beta_n) = \int_0^L |(\tau \circ \beta_n)'(s)| ds = \int_0^L |\dot{\gamma}^\tau(s) + 3f_n'(s)| ds \leq 4L = 4L_W(\gamma),$$

and since γ is a length-minimizing g_W -geodesic we conclude that

$$\hat{d}_\tau(p, q) \leq 4L = 4d_W(p, q). \quad \square$$

2.2 Proof of Theorem 2.3

In Section 2.1 we have already shown that the bi-Lipschitz estimates (4) of Theorem 2.3 hold globally (the first inequality, by Lemma 2.8) and locally (the second inequality, by Lemma 2.9) if we choose τ to be a temporal function and $h = g_W$ to be the corresponding Wick-rotated Riemannian metric. We first extend these local results to admissible time functions and arbitrary Riemannian metrics, and then prove Theorem 2.3 on compact sets.



(a) The function f_2 . (b) The curves γ (dashed) and β_2 (solid).

Figure 1: Illustration of the proof of Lemma 2.9.

Lemma 2.10. *Let (M, g) be a spacetime equipped with an admissible time function τ . Suppose h is a Riemannian metric on M . Then, for every $x \in M$ there exists a neighborhood U of x and a constant $C \geq 1$ such that for all $p, q \in U$*

$$\frac{1}{C}d_h(p, q) \leq \hat{d}_\tau(p, q) \leq Cd_h(p, q). \quad (16)$$

Proof. Let $x \in M$. We consider both inequalities in (16) separately for different Riemannian metrics and choose U to be intersection of the neighborhoods V_1 and V_2 derived in each step. More precisely, the first inequality follows from the local anti-Lipschitz assumption for τ in Definition 2.1, and the second inequality from Lemma 2.9 for an auxiliary temporal functions. Without loss of generality we furthermore assume that U is relatively compact. Then by a result of Burtscher [15, Thm. 4.5] the bi-Lipschitz estimate extends to any Riemannian metric.

Step 1. Lower bound. We show that $\frac{1}{C}d_h(p, q) \leq \hat{d}_\tau(p, q)$ for C and h as in Definition 2.1.

Let U_1 be the neighborhood of x from Definition 2.1. Since \hat{d}_τ induces the manifold topology there is a radius $r > 0$ such that for the open ball $\hat{B}_{3r}^\tau(x) \subseteq U_1$. Consider $p, q \in V_1 := \hat{B}_r^\tau(x)$. There exists a sequence $(\beta_n)_n$ of piecewise causal curves in M between p and q such that

$$\hat{L}_\tau(\beta_n) \leq \hat{d}_\tau(p, q) + \frac{1}{n}.$$

For all $n > \frac{1}{r}$ the curves β_n cannot leave U_1 . Consider one such $\beta_n: [0, 1] \rightarrow U_1$ and its partition $0 = s_0 < s_1 < \dots < s_{k-1} < s_k = 1$ in causal pieces.

Then for C and h as in Definition 2.1

$$\begin{aligned} \frac{1}{C}d_h(p, q) &\leq \frac{1}{C} \sum_{i=1}^k d_h(\beta_n(s_i), \beta_n(s_{i-1})) \\ &\leq \sum_{i=1}^k \hat{d}_\tau(\beta_n(s_i), \beta_n(s_{i-1})) = \hat{L}_\tau(\beta_n) \leq \hat{d}_\tau(p, q) + \frac{1}{n}. \end{aligned}$$

Thus the result follows as $n \rightarrow \infty$.

Step 2. Upper bound. We show that $\hat{d}_\tau(p, q) \leq 5Cd_W(p, q)$ for g_W the Riemannian metric (8) with respect to an auxiliary temporal function and corresponding C of Definition 2.1.

Every spacetime which admits a time function also admits a temporal function $\tilde{\tau}$ by Bernal and Sánchez [9, Thm. 1.2]. We consider the Wick-rotated Riemannian metric g_W defined in (8) with respect to such a fixed $\tilde{\tau}$. Let V_2 be a geodesically convex neighborhood of x with respect to g_W . Without loss of generality we assume that V_2 is contained in the neighborhoods U_2 of Definition 2.1 and Lemma 2.9. Suppose $p, q \in V_2$ and γ is the g_W -length minimizing geodesic γ between p and q in V_2 . In the proof of Lemma 2.9 a piecewise causal curve β in V_2 , sufficiently close to γ , was constructed for which

$$\begin{aligned} L_W(\beta) &= \int_0^L \|\dot{\beta}(s)\|_W ds = \int_0^L \sqrt{|\dot{\beta}^\tau|^2 + h(\dot{\beta}, \dot{\beta})} ds \\ &\leq \int_0^L \sqrt{4^2 + 2} ds < 5L = 5L_W(\gamma). \end{aligned} \tag{17}$$

Since β is piecewise causal, there is a partition $0 = s_0 < s_1 < \dots < s_{k-1} < s_k = L$ such that

$$\hat{L}_\tau(\beta) = \sum_{i=1}^k \hat{d}_\tau(\beta(s_i), \beta(s_{i-1})),$$

and τ being an admissible time function implies that there is a constant $C \geq 1$ (Definition 2.1) such that

$$\hat{L}_\tau(\beta) \leq C \sum_{i=1}^k d_W(\beta(s_i), \beta(s_{i-1})) \leq CL_W(\beta).$$

Together with (17) and the fact that γ is g_W -length minimizing we thus obtain

$$\hat{d}_\tau(p, q) \leq \hat{L}_\tau(\beta) \leq CL_W(\beta) \leq 5CL_W(\gamma) = 5Cd_W(p, q). \quad \square$$

Lemma 2.10 shows that the original local bi-Lipschitz bound (3) for causality related points (as required for admissible time functions in Definition 2.1) extends to the bi-Lipschitz bound (16) on a whole neighborhood

of each point. Finally, we adapt the proof of Burtscher [15, Thm. 4.5] to extend this local estimate to compact sets.

Proof of Theorem 2.3. Let K be a compact subset of M . The argument proceeds by contradiction. Suppose that for any $n \in \mathbb{N}$, there exist points $p_n, q_n \in K$ such that

$$\hat{d}_\tau(p_n, q_n) > nd_h(p_n, q_n). \quad (18)$$

By passing to subsequences, we can assume that $p_n \rightarrow p$ and $q_n \rightarrow q$ (in one and hence both metrics, since they both induce the manifold topology). Since K is compact it is bounded with respect to the null distance \hat{d}_{τ_1} and the inequality (18) furthermore implies that

$$d_h(p_n, q_n) \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

and hence $p = q$. Thus, for n large enough, p_n, q_n are contained in neighborhood U of p that by Lemma 2.10 is small enough so that (16) holds for some $\tilde{C} \geq 1$. But then we conclude that

$$\tilde{C}d_h(p_n, q_n) \geq \hat{d}_\tau(p_n, q_n) > nd_h(p_n, q_n),$$

a contradiction for n large. This proves that there is a constant $C \geq 1$ such that $\hat{d}_\tau(p, q) \leq Cd_h(p, q)$, and the reverse inequality is obtained in the same way. \square

2.3 Direct proof of Corollary 2.6

Corollary 2.6 is a very special case of Theorem 2.3. Here we show that it can also be obtained directly without the use of temporal functions. We first prove a local result.

Lemma 2.11. *Let τ_1, τ_2 be two admissible time functions (see Definition 2.1) on a spacetime (M, g) , and $\hat{d}_{\tau_1}, \hat{d}_{\tau_2}$ their associated null distances. Then, for every point $x \in M$, there exists a neighborhood U of x and a constant $C \geq 1$ such that for all $p, q \in U$,*

$$\frac{1}{C}\hat{d}_{\tau_1}(p, q) \leq \hat{d}_{\tau_2}(p, q) \leq C\hat{d}_{\tau_1}(p, q).$$

Proof. Fix a point $x \in M$ and an arbitrary Riemannian metric h on M . Let $U_x^{\tau_1}$ and $U_x^{\tau_2}$ be the neighborhoods around x where both τ_1 and τ_2 are Lipschitz and anti-Lipschitz with positive constants C_1, C_2 with respect to h as in (3) of Definition 2.1. Since M is locally compact there exists a relatively compact neighborhood $V \subseteq U_x^{\tau_1} \cap U_x^{\tau_2}$ of x . Since both null distances $\hat{d}_{\tau_1}, \hat{d}_{\tau_2}$ induce the manifold topology [48, Thm. 4.6], we find $r > 0$ sufficiently small so that $\hat{B}_{4r}^{\tau_1}(x) \cup \hat{B}_{4r}^{\tau_2}(x) \subseteq V$. Define a neighborhood of x by

$$U := \hat{B}_r^{\tau_1}(x) \cap \hat{B}_r^{\tau_2}(x).$$

We claim that for $i = 1, 2$ and for all $p, q \in U$,

$$\hat{d}_{\tau_i}(p, q) = \inf\{\hat{L}_{\tau_i}(\beta) \mid \beta \text{ a p.w. causal curve in } V \text{ between } p \text{ and } q\}. \quad (19)$$

What we are saying is that the null distances in U can be approximated by sequences of curves that never leave V . To prove this, let $p, q \in U$, consider an arbitrary $\varepsilon \in (0, r)$, and for each $i = 1, 2$ choose a piecewise causal curve $\beta_\varepsilon^i: I \rightarrow M$ from p to q such that

$$\hat{L}_{\tau_i}(\beta_\varepsilon^i) < \hat{d}_{\tau_i}(p, q) + \varepsilon.$$

Then, for all $t \in I$,

$$\begin{aligned} \hat{d}_{\tau_i}(x, \beta_\varepsilon^i(t)) &\leq \hat{d}_{\tau_i}(x, p) + \hat{d}_{\tau_i}(p, \beta_\varepsilon^i(t)) \leq \hat{d}_{\tau_i}(x, p) + \hat{L}_{\tau_i}(\beta_\varepsilon^i) \\ &\leq \hat{d}_{\tau_i}(x, p) + \hat{d}_{\tau_i}(p, q) + \varepsilon < 4r, \end{aligned}$$

hence β_ε^i lies entirely in $\hat{B}_{4r}^{\tau_i}(x) \subseteq V$, proving the claim (19) for $i = 1, 2$.

For any causally related points $(\tilde{p}, \tilde{q}) \in J^+$ in $U_x^{\tau_1} \cap U_x^{\tau_2}$, by (3), we have for $C := C_1 C_2 \geq 1$ that

$$\frac{1}{C}(\tau_1(\tilde{q}) - \tau_1(\tilde{p})) \leq \tau_2(\tilde{q}) - \tau_2(\tilde{p}) \leq C(\tau_1(\tilde{q}) - \tau_1(\tilde{p})). \quad (20)$$

This implies that for all piecewise causal curves β contained in $U_x^{\tau_1} \cap U_x^{\tau_2}$,

$$\frac{1}{C}\hat{L}_{\tau_1}(\beta) \leq \hat{L}_{\tau_2}(\beta) \leq C\hat{L}_{\tau_1}(\beta). \quad (21)$$

It remains to be shown that the bi-Lipschitz bounds on the null lengths extends to the null distances between any two points p and q in the smaller set U . We have already seen that for each $n \in \mathbb{N}$ exists a piecewise causal curve β_n^1 between p and q (by (19) β_n^1 is entirely contained in V !) such that

$$\hat{L}_{\tau_1}(\beta_n^1) \leq \hat{d}_{\tau_1}(p, q) + \frac{1}{n},$$

and by (21) therefore

$$\hat{d}_{\tau_2}(p, q) \leq \hat{L}_{\tau_2}(\beta_n^1) \leq C\hat{L}_{\tau_1}(\beta_n^1) \leq C\hat{d}_{\tau_1}(p, q) + \frac{C}{n}.$$

Since this inequality holds for all n and fixed C , we have that $\hat{d}_{\tau_2}(p, q) \leq C\hat{d}_{\tau_1}(p, q)$ on U . In the same way, for piecewise causal curves β_n^2 approximating $\hat{d}_{\tau_2}(p, q)$, we obtain $\hat{d}_{\tau_1}(p, q) \leq C\hat{d}_{\tau_2}(p, q)$ on U , which proves the Lemma. \square

Proof of Corollary 2.6. Having established the local result in Lemma 2.11, the proof on compact sets is now the same as that of Theorem 2.3. \square

2.4 Counterexamples

Finally, we construct two examples showing that the assumptions and statement of Corollary 2.6, and hence also that of Theorem 2.3 and Corollary 2.4, are sharp.

Example 2.12 (Corollary 2.6 is false globally). Let $M = \mathbb{R}^{1,n}$ be the Minkowski spacetime, and $\tau_1 = t$, $\tau_2 = \exp(t)$ two temporal functions (which even have the same level sets!). Since the exponential function is *not* globally Lipschitz, the time functions τ_1 and τ_2 are not equivalent in the sense that the bi-Lipschitz condition (20) for causally related points does not hold globally. Since $\tau(q) - \tau(p) = \hat{d}_\tau(p, q)$ for all $q \in J^+(p)$, it follows that if (20) fails to hold globally, then the null distances \hat{d}_{τ_1} , \hat{d}_{τ_2} are globally inequivalent.

Example 2.12 shows that although τ_1 and τ_2 are both temporal functions on the *same* spacetime (M, g) , there is a clear distinction *globally* between the metric structures induced by the null distances \hat{d}_{τ_1} and \hat{d}_{τ_2} . In particular, in Section 4 we show that (M, \hat{d}_{τ_1}) is complete while (M, \hat{d}_{τ_2}) is not.

Example 2.13 (Corollary 2.6 is false for time functions that are not locally Lipschitz). Consider the Minkowski spacetime $M = \mathbb{R}^{1,n}$ with respect to the usual global coordinates (t, x) . Let $\tau_1(t, x) := t$ be the standard (smooth and locally anti-Lipschitz) time function and

$$\tau_2(t, x) := \operatorname{sgn}(t)\sqrt{|t|}.$$

Clearly, τ_2 is continuous and a time function. It is, however, *not* locally Lipschitz around $t = 0$ because $\partial_t \tau_2(t, x) \rightarrow \infty$ as $t \rightarrow 0$. Nonetheless, τ_2 is locally anti-Lipschitz (and therefore \hat{d}_{τ_2} induces the manifold topology): Consider a point (t, x) with $t > 0$ and the neighborhood $U = (0, 2t) \times \mathbb{R}^n$. Suppose $q \in J^+(p)$ with $p = (t_p, x_p)$, $q = (t_q, x_q)$. By the Mean Value Theorem for $\sqrt{\cdot}$ on $[0, 2t]$ we have

$$\sqrt{t_q} - \sqrt{t_p} \geq \left(\inf_{r \in (0, 2t)} \frac{d}{dr} \sqrt{r} \right) (t_q - t_p) = \frac{1}{2\sqrt{2t}} |t_q - t_p|.$$

Hence due to the causal relation on Minkowski spacetime we obtain with respect to the Euclidean distance d and with $C := \frac{1}{4\sqrt{2t}} > 0$ that on U

$$q \in J^+(p) \implies \tau_2(q) - \tau_2(p) \geq \frac{1}{4\sqrt{2t}} \sqrt{(t_q - t_p)^2 + |x_q - x_p|^2} \geq Cd(q, p).$$

In the same fashion, τ_2 is anti-Lipschitz in the neighborhood $U = (2t, 0) \times \mathbb{R}$ of a point (t, x) with $t < 0$. If $t = 0$, we can simply use the neighborhood $U = (-1, 1) \times \mathbb{R}^n$ and $C = \frac{1}{4}$ (since the infimum is at $|r| = 1$). Hence τ_2

is locally anti-Lipschitz everywhere (but clearly not globally anti-Lipschitz with respect to the Euclidean distance).

Now assume that $p = (0, 0)$ and let $q_t = (t, 0)$, $t \in (0, 1)$, be arbitrarily close to p . Then

$$\hat{d}_{\tau_1}(p, q_t) = t \leq \sqrt{t} = \hat{d}_{\tau_2}(p, q_t)$$

but the bi-Lipschitz estimate (4) of Theorem 2.6 does not hold because

$$\frac{\hat{d}_{\tau_2}(p, q_t)}{\hat{d}_{\tau_1}(p, q_t)} = \frac{1}{\sqrt{t}} \rightarrow \infty \quad \text{as } t \rightarrow 0.$$

3 Encoding causality

Recall that we defined the causal relation $J^+ \subseteq M \times M$ by

$$J^+ := \{(p, q) \mid \text{there exists a future-directed causal curve from } p \text{ to } q\}.$$

By definition the null distance is achieved for causal curves. We investigate when the converse holds, that is, when

$$(p, q) \in J^+ \iff \hat{d}_\tau(p, q) = \tau(q) - \tau(p), \quad (22)$$

in which case the null distance is said to *encode causality*, an open problem mentioned in [48, Sec. 1]. See Remark 3.5 for a discussion on the recent answers obtained by Sakovich and Sormani [44] and in this paper. We introduce a new notation for the right hand side of (22).

Definition 3.1. Let (M, g) be a spacetime equipped with a time function $\tau: M \rightarrow \mathbb{R}$. We define the *null distance relation* $\hat{R}_\tau^+ \subseteq M \times M$ by

$$(p, q) \in \hat{R}_\tau^+ : \iff \hat{d}_\tau(p, q) = \tau(q) - \tau(p).$$

Clearly, the relation \hat{R}_τ^+ is reflexive and transitive. Antisymmetry requires definiteness of \hat{d}_τ which, for instance, follows if τ is a locally anti-Lipschitz time function [48, Thm. 4.6]. We can thus summarize the basic properties of \hat{R}_τ^+ as follows.

Lemma 3.2. *Suppose (M, g) is a spacetime and τ is a locally anti-Lipschitz time function on M . Then the null distance relation \hat{R}_τ^+ is a partial order on M satisfying $J^+ \subseteq \hat{R}_\tau^+$. \square*

In Section 3.1 we prove that globally hyperbolic spacetimes encode causality when τ is carefully chosen, which can be reformulated in terms of \hat{R}_τ^+ as follows.

Theorem 1.9. *Let (M, g) be a globally hyperbolic spacetime and $\tau: M \rightarrow \mathbb{R}$ be a Cauchy temporal function. Then causality is encoded in \hat{d}_τ , meaning $J^+ = \hat{R}_\tau^+$.*

Subsequently we also prove a local version of this result for any spacetime with a temporal function.

Theorem 3.3. *Let (M, g) be a spacetime and τ be a temporal function. Then, every point $x \in M$ has a neighborhood U such that*

$$J^+ \cap (U \times U) = \hat{R}_\tau^+ \cap (U \times U).$$

In Section 3.2 we show, however, that the assumptions for the global Theorem 1.9 are sharp in the sense that there are (non-globally hyperbolic) spacetimes and temporal functions τ for which $J^+ \subsetneq \hat{R}_\tau^+$.

3.1 Proofs of Theorems 1.9 and 3.3

In this section, the Wick-rotated metric g_W of g introduced in Section 2.1 plays an important role again. Since g_W need not be complete even if the spacetime is globally hyperbolic (see Example 4.5) we furthermore make use of the existence of a conformally equivalent complete Riemannian metric

$$g_R := \Omega^2 g_W, \tag{23}$$

where $\Omega: M \rightarrow [1, \infty)$ is a smooth function [42, Thm. 1]. We denote the corresponding norm, length functional and distance by $\|\cdot\|_R$, L_R and d_R , respectively. The relation of g_R and g_W carries over to the distances as follows, that is, for any curve $\gamma: [a, b] \rightarrow M$,

$$L_W(\gamma) \leq L_R(\gamma),$$

and hence for any $p, q \in M$

$$d_W(p, q) \leq d_R(p, q).$$

In the following lemma, we obtain reverse inequalities on compact sets.

Lemma 3.4. *Let g_R and g_W be two conformally equivalent Riemannian metrics on M as in (23), and suppose K is a compact set in M . Then there exist constants $C_K \geq c_K \geq 1$ such that for any curve γ in K*

$$L_W(\gamma) \leq L_R(\gamma) \leq c_K L_W(\gamma).$$

and for any $p, q \in K$,

$$d_W(p, q) \leq d_R(p, q) \leq C_K d_W(p, q).$$

Proof. The first statement about the lengths follows immediately from the Definition (23), and we may pick $c_K := \max_{x \in K} \Omega(x)$. The first inequality for the distances is trivial and the second inequality follows from [15, Thm. 4.5] (note that, in general, we need to pick $C_K > c_K$ as minimizing curves may leave K , but the proof in [15] guarantees boundedness of C_K). \square

With these tools, we proceed to prove the theorem.

Proof of Theorem 1.9. Trivially, $J^+ \subseteq \hat{R}_\tau^+$, so we only need to prove $\hat{R}_\tau^+ \subseteq J^+$. Suppose that $(p, q) \in \hat{R}_\tau^+$, that is, $\hat{d}_\tau(p, q) = \tau(q) - \tau(p)$. Since $(p, p) \in J^+$ is trivially satisfied, we can assume that $p \neq q$ and thus necessarily $\tau(q) > \tau(p)$. By definition of \hat{d}_τ there exists a sequence of piecewise causal curves $(\beta_n)_n$ such that

$$0 < \tau(q) - \tau(p) \leq \hat{L}_\tau(\beta_n) \leq \tau(q) - \tau(p) + \frac{1}{n}. \quad (24)$$

In what follows we construct a future-directed causal curve β between p and q , which then immediately implies $(p, q) \in J^+$. We proceed as follows: After a preliminary local estimate we construct a candidate limit curve near p in (M, g_R) , and then show that it is both locally Lipschitz and future-directed causal. Finally, we show it naturally extends all the way up to q .

Step 1. A local version of (24). We consider $\beta_n: [0, L_n] \rightarrow M$ on arbitrary subintervals of its domain. Suppose there exists a subinterval $[s, t] \subseteq [0, L_n]$ such that $\hat{L}_\tau(\beta_n|_{[s, t]}) > \tau(\beta_n(t)) - \tau(\beta_n(s)) + \frac{1}{n}$. Then by the additivity of null lengths

$$\begin{aligned} \hat{L}_\tau(\beta_n) &= \hat{L}_\tau(\beta_n|_{[0, s]}) + \hat{L}_\tau(\beta_n|_{[s, t]}) + \hat{L}_\tau(\beta_n|_{[t, L_n]}) \\ &> |\tau(\beta_n(s)) - \tau(p)| + |\tau(\beta_n(t)) - \tau(\beta_n(s)) + \frac{1}{n}| + |\tau(q) - \tau(\beta_n(t))| \\ &\geq \tau(q) - \tau(p) + \frac{1}{n}, \end{aligned}$$

a contradiction to (24). Thus for all subintervals $[s, t]$ of the domain of β_n

$$|\tau(\beta_n(t)) - \tau(\beta_n(s))| \leq \hat{L}_\tau(\beta_n|_{[s, t]}) \leq \tau(\beta_n(t)) - \tau(\beta_n(s)) + \frac{1}{n}. \quad (25)$$

(Note that at this point we cannot yet rule out that $\tau(\beta_n(t)) - \tau(\beta_n(s)) < 0$, which is why we do not want to drop the absolute value on the left hand side.)

Step 2. Construction of a candidate limit curve β near p . Suppose, that each $\beta_n: [0, L_n] \rightarrow M$ is parametrized by g_R -arclength with $\beta_n(0) = p$ and $\beta_n(L_n) = q$. In addition, we attach a future-directed inextendible causal curve $\tilde{\beta}$ at q . Since g_R is complete, $\tilde{\beta}$ must have infinite g_R -length. Thus we can extend all β_n by $\tilde{\beta}$ to obtain future-inextendible piecewise causal curves with g_R -arclength parametrization, again denoted by $\beta_n: [0, \infty) \rightarrow M$ and satisfying (25). In particular, for each n and any $s, t \in [0, \infty)$,

$$d_R(\beta_n(s), \beta_n(t)) \leq L_R(\beta_n|_{[s, t]}) = |t - s|. \quad (26)$$

By construction, $\beta_n(0) = p$ for all n . Fix any other $t_0 \in (0, \infty)$. Then, due to the g_R -arclength parametrization, each curve segment $\beta_n|_{[0, t_0]}$ is contained in the bounded set (which, by the Hopf–Rinow Theorem for g_R is also compact)

$$B_{t_0}(p) := \{x \in M : d_R(p, x) \leq t_0\}.$$

In particular, the family $(\beta_n|_{[0, t_0]})_n$ is uniformly bounded and uniformly equicontinuous. Thus, by the Arzelà–Ascoli Theorem (see, for instance, [41, Thm. 47.1] or [8, Thm. 3.30]), there exists a continuous curve $\beta: [0, \infty) \rightarrow M$, such that a subsequence of $(\beta_n)_n$ converges uniformly to β on all compact subintervals. Since $\frac{1}{n_k} \leq \frac{1}{k}$ for any subsequence, we can denote this subsequence again by $(\beta_n)_n$. Moreover, (26) implies that β is a locally Lipschitz curve with

$$d_R(\beta(s), \beta(t)) \leq |t - s| \quad (27)$$

for all $s, t \in [0, \infty)$.

Step 3. The curve β is future-directed causal. Since β is locally Lipschitz, together with Rademacher’s Theorem, we know that $\dot{\beta}$ exists almost everywhere. To conclude that β is a future-directed causal curve, it remains to be shown that $g(\dot{\beta}, \dot{\beta}) \leq 0$ and $(\tau \circ \beta)' > 0$ almost everywhere.

By [15, Prop. 4.10] the g_R -norm of the analytic derivative and d_R -metric derivative of β exist and coincide almost everywhere. Combined with the Lipschitz estimate (27) for β we thus have for almost all $s \in [0, \infty)$

$$0 \leq \|\dot{\beta}(s)\|_R = \lim_{h \rightarrow 0} \frac{d_R(\beta(s+h), \beta(s))}{|h|} \leq \lim_{h \rightarrow 0} \frac{|h|}{|h|} = 1. \quad (28)$$

In order to show that $\dot{\beta}$ is almost everywhere causal we need to control the τ -component $\dot{\beta}^\tau = (\tau \circ \beta)'$ of the tangent vectors. Suppose $\dot{\beta}(s_0)$ exists for a fixed $s_0 \in (0, \infty)$, and consider the closed (and hence compact) d_R -ball B_ε of radius ε at $\beta(s_0)$ in M . By Lipschitz continuity (27) of β the whole interval $[s_0 - \varepsilon/2, s_0 + \varepsilon/2]$, is mapped into $B_{\varepsilon/2}$. We use the approximating sequence $(\beta_n)_n$ of β next, more precisely, that the β_n are piecewise causal and converge uniformly on compact intervals. Thus for sufficiently large n all $\beta_n([s_0 - \varepsilon/2, s_0 + \varepsilon/2]) \subseteq B_\varepsilon$. By the local estimate (25)

$$\lim_{n \rightarrow \infty} \hat{L}_\tau(\beta_n|_{[s_0 - \varepsilon/2, s_0 + \varepsilon/2]}) = \tau(\beta(s_0 + \varepsilon/2)) - \tau(\beta(s_0 - \varepsilon/2)). \quad (29)$$

On the other hand, due to the g_R -arclength parametrization of β_n and Lemmas 2.8 and 3.4 (with constant $c_\varepsilon = \max_{x \in B_\varepsilon} \Omega(x) \geq 1$)

$$\begin{aligned} \varepsilon &= L_R(\beta_n|_{[s_0 - \varepsilon/2, s_0 + \varepsilon/2]}) \leq c_\varepsilon L_W(\beta_n|_{[s_0 - \varepsilon/2, s_0 + \varepsilon/2]}) \\ &\leq \sqrt{2}c_\varepsilon \hat{L}_\tau(\beta_n|_{[s_0 - \varepsilon/2, s_0 + \varepsilon/2]}). \end{aligned}$$

Hence by (29)

$$\frac{1}{\sqrt{2}c_\varepsilon} \leq \frac{1}{\varepsilon} \lim_{n \rightarrow \infty} \hat{L}_\tau(\beta_n|_{[s_0-\varepsilon/2, s_0+\varepsilon/2]}) = \frac{\tau(\beta(s_0 + \varepsilon/2)) - \tau(\beta(s_0 - \varepsilon/2))}{\varepsilon}.$$

Due to the continuity of Ω , it follows that $c_\varepsilon \rightarrow \Omega(\beta(s_0))$ as $\varepsilon \rightarrow 0$, while the difference quotient of $\tau \circ \beta$ converges to the derivative $\dot{\beta}^\tau(s_0) = (\tau \circ \beta)'(s_0)$. Thus in the limit we obtain

$$\dot{\beta}^\tau(s_0) \geq \frac{1}{\sqrt{2}\Omega(\beta(s_0))} > 0.$$

Due to the Wick-rotation g_W of g as well as the g_R -bound (28)

$$\begin{aligned} g(\dot{\beta}(s_0), \dot{\beta}(s_0)) &= -2|\dot{\beta}^\tau(s_0)|^2 + \|\dot{\beta}(s_0)\|_W^2 \\ &\leq -\Omega^{-2}(\beta(s_0)) + \Omega^{-2}(\beta(s_0))\|\dot{\beta}(s_0)\|_R^2 \leq 0. \end{aligned}$$

This proves that $\dot{\beta}$ is future-directed causal almost everywhere.

Step 4. The point q lies on β . By construction $\beta(0) = p$, and by Step 3 we know that $\tau \circ \beta$ is strictly increasing on $[0, \infty)$. We distinguish two cases:

- (i) If there is an $s_0 \in [0, \infty)$ such that $\tau(\beta(s_0)) = \tau(q)$, then the following argument implies that $\beta(s_0) = q$:

Let $\varepsilon > 0$ be arbitrary. Since both \hat{d}_τ and d_R induce the manifold topology, there exists a $\delta > 0$ such that $d_R(\beta(s_0), x) < \delta$ implies $\hat{d}_\tau(\beta(s_0), x) < \varepsilon$. Due to the convergence $\beta_n(s_0) \rightarrow \beta(s_0)$ (obtained with respect to d_R in Step 2), for any n sufficiently large

$$\hat{d}_\tau(\beta(s_0), \beta_n(s_0)) < \varepsilon. \quad (30)$$

Moreover, due to the continuity of $\tau \circ \beta$, for all $n > \frac{1}{\varepsilon}$ sufficiently large

$$|\tau(\beta_n(s_0)) - \tau(q)| = |\tau(\beta_n(s_0)) - \tau(\beta(s_0))| < \varepsilon.$$

Since $\beta_n(L_n) = q$, the local estimate (25) on $[s_0, L_n]$ (or $[L_n, s_0]$ if $L_n < s_0$) yields

$$\begin{aligned} \hat{d}_\tau(\beta_n(s_0), q) &\leq \hat{L}_\tau(\beta_n|_{[s_0, L_n]}) \\ &\leq \tau(q) - \tau(\beta_n(s_0)) + \frac{1}{n} < 2\varepsilon. \end{aligned} \quad (31)$$

Combining (30) and (31) implies that $\hat{d}_\tau(\beta(s_0), q) < 3\varepsilon$ for any $\varepsilon > 0$. Thus $\beta(s_0) = q$.

- (ii) The only obstruction to the desired conclusion is therefore that for all $s \in [0, \infty)$

$$\tau(\beta(s)) < \tau(q). \quad (32)$$

We show that this case cannot occur. Since τ is a Cauchy time function, (32) implies that the causal curve β is future extendible as a future-directed causal curve. In particular, the future endpoint $x := \lim_{s \rightarrow \infty} \beta(s)$ of β exists (since it is necessarily part of any extension) and by (32), $\tau(x) \leq \tau(q)$. If $x = q$ we are done, otherwise there exists a relatively compact open set W around $\beta \cup \{x\}$ such that $q \notin \overline{W}$.

We will use the approximating curves β_n to show that β must in fact leave W : Since $\beta_n(0) = p \in W$ and $\beta_n(L_n) = q \notin W$ there exists

$$b_n := \sup\{s \mid \beta_n(t) \in W \text{ for all } t \in [0, s]\} \in (0, L_n),$$

and $\beta_n(b_n) \in \partial W$. Since \overline{W} is compact, by Lemmas 2.8 and 3.4, there exists a constant $C > 0$ such that

$$\begin{aligned} b_n &= L_R(\beta_n|_{[0, b_n]}) \leq \sqrt{2}C \hat{L}_\tau(\beta_n|_{[0, b_n]}) \leq \sqrt{2}C \hat{L}_\tau(\beta_n|_{[0, L_n]}) \\ &\leq \sqrt{2}C \left(\tau(q) - \tau(p) + \frac{1}{n} \right). \end{aligned}$$

Hence all b_n are uniformly bounded from above by a constant $a := \sqrt{2}C(\tau(q) - \tau(p) + 1)$. In particular, a subsequence converges to $b := \limsup b_n \in [0, a)$. Let $\varepsilon > 0$. By uniform convergence $\beta_n \rightarrow \beta$ on $[0, a]$ for all sufficiently large n (along the previous subsequence), we have

$$d_R(\beta(b_n), \beta_n(b_n)) < \varepsilon. \quad (33)$$

Since $\beta_n(b_n) \in \partial W$ and ∂W is compact as closed subset of the compact set \overline{W} , there exists a subsequence of points $\beta_{n_k}(b_{n_k})$ that converges to a point $y \in \partial W$, i.e., for k sufficiently large

$$d_R(\beta_{n_k}(b_{n_k}), y) < \varepsilon. \quad (34)$$

Combining (33)–(34) yields

$$d_R(\beta(b), y) \leq d_R(\beta(b), \beta_{n_k}(b_{n_k})) + d_R(\beta_{n_k}(b_{n_k}), y) < 2\varepsilon.$$

In other words, $\beta(b) = y \in \partial W$, a contradiction to the assumption that β is entirely contained in the open set W . Hence the assumption (32) must be false, and thus by case (i) β indeed reaches q .

To sum up, we have constructed a future-directed causal (Step 3) curve β from p (Step 2) to q (Step 4). Therefore, $(p, q) \in J^+$, and thus $J^+ = \hat{R}_\tau^+$. \square

We can adapt the proof of Theorem 1.9 to show its local counterpart Theorem 3.3, which holds for every stably causal Lorentzian manifold (since every such manifold admits a smooth temporal function by [39, Thm. 4.100]).

Proof of Theorem 3.3. Let $x \in M$. Since $J^+ \subseteq \hat{R}_\tau^+$ is always true, the \subseteq inclusion is trivial. In order to show \supseteq we construct a suitable relatively compact open neighborhood U of x and construct causal curves locally similar to the proof of Theorem 1.9. Only Step 2 made use of the complete Riemannian metric g_R and Step 4 used global hyperbolicity of (M, g) and have to be carried out slightly different.

By local compactness of M there is an $r > 0$ sufficiently small such that $\hat{B}_{3r}^\tau(x)$ is relatively compact (the open ball with respect to \hat{d}_τ of radius $3r$). Consider $U := \hat{B}_r^\tau(x)$. Suppose now $p, q \in U$ and $(p, q) \in \hat{R}_\tau^+$, i.e.,

$$\hat{d}_\tau(p, q) = \tau(q) - \tau(p) > 0.$$

By definition of U and the triangle inequality also $\hat{d}_\tau(p, q) < 2r$. Let $(\beta_n)_n$ be a sequence of piecewise causal paths $\beta_n: [0, L_n] \rightarrow M$ that approximates $\hat{d}_\tau(p, q)$. Hence we may fix any $\varepsilon \in (0, \hat{d}_\tau(p, q))$ and assume without loss of generality that for all n ,

$$\hat{L}_\tau(\beta_n) < \hat{d}_\tau(p, q) + \varepsilon < 2r. \quad (35)$$

In particular, contained in $\hat{B}_{3r}^\tau(x)$.

Since τ is assumed to be temporal the Wick-rotated metric g_W of g exists (see Section 2.1). We assume that the β_n are parametrized by g_W -arclength, and therefore $L_n = L_W(\beta_n)$. By Lemma 2.8 and (35) we then obtain the following estimates:

$$\hat{d}_\tau(p, q) \leq \hat{L}_\tau(\beta_n) \leq L_n \leq \sqrt{2}\hat{L}_\tau(\beta_n) \leq \sqrt{2}(\hat{d}_\tau(p, q) + \varepsilon). \quad (36)$$

It is more convenient to have all the β_n defined on the same interval $[0, L]$, which we achieve by extending each β_n as follows: Set $L := \sqrt{2}(\hat{d}_\tau(p, q) + \varepsilon)$. Then attach to each β_n a future-directed causal curve $\tilde{\beta}_n$ starting at q of g_W -length $\tilde{L}_n \leq L - L_n$. Notice that

$$\tilde{L}_n \leq L - L_n < (\sqrt{2} - 1)\hat{d}_\tau(p, q) + \sqrt{2}\varepsilon < 2(\sqrt{2} - 1)r + \sqrt{2}\varepsilon < r$$

for ε small enough. Since $\hat{L}_\tau(\tilde{\beta}_n) \leq \tilde{L}_n$ (by Lemma 2.8) and $\tilde{\beta}_n$ starts at $q \in \hat{B}_r^\tau(x)$, it follows that $\tilde{\beta}_n$ is contained in $\hat{B}_{3r}^\tau(x)$. This also proves that indeed a long enough extension up to $\tilde{L}_n = L - L_n$ exists, given that $\hat{B}_{3r}^\tau(x)$ is relatively compact and the spacetime (M, g) is non-imprisoning (because it admits a time function). We have thus obtained a sequence of piecewise

causal curves, denoted again as $\beta_n: [0, L] \rightarrow \hat{B}_{3r}^\tau(x)$. The curves β_n start at $p = \beta_n(0)$, reach $q = \beta_n(L_n)$, and then continue to their endpoint $\beta_n(L)$. The local estimate (25) of Step 1 holds too.

Let $B := \hat{B}_{3r}^\tau(x)$. By the Arzelà–Ascoli Theorem applied on the compact metric space $(\overline{B}, d_W|_{\overline{B}})$, there exists a subsequence, again denoted by $(\beta_n)_n$, that uniformly converges to a 1-Lipschitz continuous limit curve $\beta: [0, L] \rightarrow \overline{B}$ as in Step 2 of the proof of Theorem 3.3.

Step 3 in the proof of Theorem 3.3 can be used verbatim with $g_R = g_W$ (and $\Omega \equiv 1$) since completeness of g_R was not needed here. Thus β is future-directed causal.

It remains to prove that β reaches the point q , which is now easier thanks to the fact that $(\beta_n)_n$ converges to β uniformly on $[0, L]$. Recall that $\beta_n(L_n) = q$ for all $n \in \mathbb{N}$. By passing to a subsequence, if necessary, we may assume that $L_n \rightarrow L_\infty \leq L$. But then, given $\delta > 0$, for all n large enough we obtain

$$\begin{aligned} d_W(\beta(L_\infty), q) &\leq d_W(\beta(L_\infty), \beta_n(L_\infty)) + d_W(\beta_n(L_\infty), \beta_n(L_n)) + d_W(\beta_n(L_n), q) \\ &\leq \delta + |L_\infty - L_n| + 0 < 2\delta, \end{aligned}$$

and therefore conclude that $\beta(L_\infty) = q$.

Thus β is indeed a future-directed causal curve from p through q and hence $(p, q) \in J^+$. \square

Remark 3.5 (Alternative approaches and results). In earlier work of Sormani and Vega the important class of warped product spacetimes $I \times_f \Sigma$ with interval $I \subseteq \mathbb{R}$ and complete Riemannian fibers Σ was already shown to encode causality [48, Thm. 3.25]. It is also easy to see that \hat{d}_τ encodes causality if all null distances are realized by piecewise causal curves [1, Rem. 3.22]. It remained an open problem to understand causality encoding in the general case. Independently to our approach, Sakovich and Sormani [44] very recently obtained some results that are comparable to Theorem 1.9 and Theorem 3.3. We briefly discuss their setting and how it compares to ours. Their local result [44, Thm. 1.1] is formulated for locally anti-Lipschitz generalized time functions and hence by Lemma 2.2 extends our Theorem 3.3. Their global result [44, Thm. 4.1] is formulated for *proper* generalized time functions, requiring that all time slabs $\tau^{-1}([\tau_1, \tau_2])$ with $[\tau_1, \tau_2] \subseteq \tau(M)$ are compact. By a result of Borde et al [12, Prop. 1] any spacetime admitting a proper generalized time function must be globally hyperbolic. Moreover, using results of Budic et al [13, Thm. 1] and Galloway [22, Cor. 1], one can show that the Cauchy surfaces must be compact. Therefore, our global Theorem 1.3 is applicable to a wider class of spacetimes while their result [44, Thm. 4.1] is applicable to a wider class of time functions (even discontinuous ones). In Section 3.2 we show that global hyperbolicity can, in general, not be relaxed.

3.2 Counterexamples

We conclude this section with a series of examples that show that the local Theorem 3.3 with respect to temporal functions cannot be promoted to a global statement in the spirit of Theorem 1.9, even if the spacetime is causally simple. Causal simplicity is defined by requiring the causal relation J^+ to be antisymmetric and closed, and sits only one step below global hyperbolicity on the causal ladder.

In order to better contextualize our examples, we also consider the K^+ relation. Recall that K^+ is defined as the (unique) smallest closed and transitive relation containing J^+ . The definition of K^+ is due to Sorkin and Woolgar [47]. Furthermore, a spacetime (M, g) is called K^+ -causal if the K^+ relation is antisymmetric, a condition later shown to be equivalent to stable causality by Minguzzi [35], and hence also equivalent to the existence of time function [26, 38].

Example 3.6 ($J^+ \subsetneq K^+ = \hat{R}_\tau^+$). Allen and Burtscher constructed examples [1, Ex. 3.23, 3.24] by removing points and lines from Minkowski space for which causality is not encoded, i.e., $J^+ \neq \hat{R}_\tau^+$. Notably, K -causality is still encoded, meaning that $K^+ = \hat{R}_\tau^+$.

Sormani and Vega gave another example [48, Prop. 3.4] for which one can check that $J^+ \subsetneq K^+ \subsetneq \hat{R}_\tau^+$. Their example is Minkowski space with the time function $\tau = t^3$, which is not locally anti-Lipschitz, and the null distance is not definite. We modify said example to obtain a definite null distance for which neither J^+ nor K^+ are encoded.

Example 3.7 ($J^+ \subsetneq K^+ \subsetneq \hat{R}_\tau^+$). Consider $M := \mathbb{R}^{1,1} \setminus \{(0, x) \mid x \geq 0\}$ equipped with the usual Minkowski metric $g := -dt^2 + dx^2$. We define the function $\tau: M \rightarrow \mathbb{R}$ by

$$\tau(t, x) := \begin{cases} t^3 & \text{if } x > 0, \\ t^3 + tx^2 & \text{if } x \leq 0. \end{cases}$$

First observe that τ is a temporal function on (M, g) because its gradient vector is timelike: For $x > 0$, this is trivial (note that then $t \neq 0$, by definition of M). For $x \leq 0$, since $(0, 0) \notin M$, it follows from

$$\begin{aligned} g(\nabla\tau, \nabla\tau) &= -(\partial_t\tau)^2 + (\partial_x\tau)^2 \\ &= -(3t^2 + x^2)^2 + (2tx)^2 \\ &= -9t^4 - x^4 - 6t^2x^2 + 4t^2x^2 < 0. \end{aligned}$$

Next we show that every pair of points $p = (t_p, x_p)$, $q = (t_q, x_q)$ with $x_p, x_q > 0$ and $t_q < 0 < t_p$ satisfies

$$\hat{d}_\tau(p, q) = \tau(p) - \tau(q), \tag{37}$$




Figure 2: A piecewise causal curve that approximates the distance between the points p and q in Example 3.7, giving the upper bound (40).

despite the fact that clearly not all such p and q are related by J^+ or K^+ : Let $k \in \mathbb{N}$ be large enough so that $t_p > x_p/k$ and $|t_q| > x_q/k$. Consider the points (see Figure 2)

$$q_1 := (x_p/k, x_p), \quad q_2 := (x_p/k, 0), \quad q_3 := (0, -\min\{x_p, x_q\}/k).$$

Since $p \in J^+(q_1)$ and $q_2 \in J^+(q_3)$ trivially

$$\hat{d}_\tau(p, q_1) = t_p^3 - \left(\frac{x_p}{k}\right)^3, \quad \hat{d}_\tau(q_2, q_3) = \left(\frac{x_p}{k}\right)^3. \quad (38)$$

Moreover, taking a piecewise null curve from q_1 to q_2 that consists of $2k$ segments between $t = x_p/k$ and $t = 2x_p/k$ (see Figure 2), we get that

$$\hat{d}_\tau(q_1, q_2) \leq 2k \left(\left(\frac{2x_p}{k}\right)^3 - \left(\frac{x_p}{k}\right)^3 \right) = 14 \frac{x_p^3}{k^2}. \quad (39)$$

Combining (38) and (39) with the triangle inequality proves

$$\hat{d}_\tau(p, q_3) \leq t_p^3 + 14 \frac{x_p^3}{k^2},$$

and by symmetry we obtain the analogous estimate for $\hat{d}_\tau(q, q_3)$, thus

$$\hat{d}_\tau(p, q) \leq \hat{d}_\tau(p, q_3) + \hat{d}_\tau(q_3, q) \leq t_p^3 + 14 \frac{x_p^3 + x_q^3}{k^2} + |t_q|^3, \quad (40)$$

and taking the limit $k \rightarrow \infty$ implies (37), as desired.

In the previous example, we have constructed a minimizing sequence of piecewise causal curves by choosing curves that are close to a “barrier” (the removed positive x axis). This barrier, however, also makes the spacetime causally discontinuous. Since the time function τ is perfectly regular (it is

C^1 and temporal, and can easily be smoothed out), one might suspect that causal discontinuity is the reason that causality is not encoded by \hat{d}_τ in Example 3.7. This motivates the next example, where the time function τ is smooth temporal and *steep*, a notion introduced by Müller and Sánchez [40] that additionally requires $g(\nabla\tau, \nabla\tau) \leq -1$. The underlying spacetime is even causally simple, but causality is still not encoded in \hat{d}_τ . In order to achieve this effect, instead of approaching a barrier, we construct an example with a minimizing sequence of piecewise null curves that runs off to infinity.

Example 3.8 ($J^+ = K^+ \subsetneq \hat{R}_\tau^+$). Let $M := \mathbb{R}^3$ with coordinates (t, x, y) and warped product metric tensor

$$g := \cosh^2(x) (-dt^2 + dy^2) + dx^2. \quad (41)$$

Note that all $\{x = x_0\}$ planes are conformal to Minkowski space, while each $\{y = y_0\}$ plane is isometric to the universal cover of AdS^2 . Therefore, the induced null geodesics $s \mapsto (t_\pm(s), x_\pm(s))$ in the $\{y = y_0\}$ plane going through a point $(t_0, 0)$ are given by

$$\begin{aligned} t_\pm(s) &= 2 \arctan \left(\tanh \left(\frac{s}{2} \right) \right) + t_0, \\ x_\pm(s) &= \pm s, \end{aligned} \quad (42)$$

the subscript $+$ or $-$ indicating the right- or left-going geodesics respectively [30, Sec. 5.10]. Moreover, the function

$$\tau(t, x, y) := \cosh^{-1}(x)t + t^3,$$

is a steep temporal function on (M, g) since

$$\begin{aligned} g(\nabla\tau, \nabla\tau) &= -\cosh^2(x) (\partial_t\tau)^2 + (\partial_x\tau)^2 \\ &= -\cosh^2(x) (\cosh^{-1}(x) + 3t^2)^2 + (-\cosh^{-1}(x) \tanh(x)t)^2 \\ &= -1 - \cosh^2(x)9t^4 - (6 \cosh(x) - \cosh^{-2}(x) \tanh^2(x)) t^2 \leq -1, \end{aligned}$$

where in the last line, we have used that $|\tanh(x)| < 1 \leq \cosh(x)$ for all $x \in \mathbb{R}$.

Consider now two points $p, q \in M$ of the form

$$p := \left(-\frac{\pi}{2}, 0, y_p \right), \quad q := \left(\frac{\pi}{2}, 0, y_q \right),$$

where $|y_p - y_q| > \pi$. We are going to show that $(p, q) \in \hat{R}_\tau^+$, i.e., that

$$\hat{d}_\tau(p, q) = \tau(q) - \tau(p), \quad (43)$$

despite the fact that $(p, q) \notin J^+$ (and since our spacetime is causally simple, $J^+ = K^+$). To see that $(p, q) \notin J^+$, note that by definition of g in (41)




Figure 3: Two piecewise causal curves going through the points p_{s_0} and q_{s_0} (thick) and p_{s_1} and q_{s_1} (thin), respectively, that approximate the null distance between p and q in Example 3.8. The thin curve yields a better approximation.

the $\{x = 0\}$ plane is isometric to 1 + 1-Minkowski space, and that the projection $(t, x, y) \mapsto (t, y)$ of any causal curve from p to q in M to $\{x = 0\}$ remains causal in $\{x = 0\}$. The condition $|y_p - y_q| > \pi$ then implies that $(p, q) \notin J^+_{\{x=0\}}$, and therefore also $(p, q) \notin J^+$.

In order to show (43), we proceed in three steps. First, define

$$p_s = (0, s, y_p), \quad q_s = (0, s, y_q),$$

where $s \in \mathbb{R}$ is arbitrary. By (42), a null geodesic in the $\{y = y_p\}$ plane starting at p eventually reaches the point $(t_+(s), s)$ where $t_+(s) < 0$. It follows that $(p, p_s) \in J^+$, and by the time reversed argument on the $\{y = y_q\}$ plane, that $(q_s, q) \in J^+$, altogether implying that

$$\hat{d}_\tau(p, p_s) = \tau(p_s) - \tau(p) = -\tau(p), \quad \hat{d}_\tau(q_s, q) = \tau(q) - \tau(q_s) = \tau(q). \quad (44)$$

The second step is to estimate the null distance between p_s and q_s . Given that the $\{x = s\}$ plane is conformal to Minkowski space and the null distance is conformally invariant it is easy to construct piecewise causal curves between p_s and q_s . The null distance induced on each plane is different though, and since $\tau|_{\{x=x_s\}} \rightarrow t^3$ as $s \rightarrow \infty$ we have at the “boundary” of our spacetime an indefinite null distance that cannot distinguish any points in the $\{t = 0\}$ slice [48, Prop. 3.4]. We make this intuitive picture precise by constructing in each $\{x = x_s\}$ plane a piecewise null curve $\beta_{s,k}$ that bounces k times between $t = 0$ and $t = |y_p - y_q|/k$ (see Figure 3). Using the curves

$\beta_{s,k}$ to estimate the null distance, we obtain the upper bound

$$\begin{aligned} \hat{d}_\tau(p_s, q_s) &\leq \liminf_{k \rightarrow \infty} \hat{L}_\tau(\beta_{s,k}) \\ &= \liminf_{k \rightarrow \infty} 2k \left(\frac{|y_p - y_q|^3}{k^3} + \cosh^{-1}(s) \frac{|y_p - y_q|}{k} \right) \\ &= 2 \cosh^{-1}(s) |y_p - y_q|. \end{aligned} \tag{45}$$

Finally, the triangle inequality together with (44) and (45) yields

$$\hat{d}_\tau(p, q) \leq \lim_{s \rightarrow \infty} \left(\hat{d}_\tau(p, p_s) + \hat{d}_\tau(p_s, q_s) + \hat{d}_\tau(q_s, q) \right) = \tau(q) - \tau(p).$$

This finishes the proof of (43), since the opposite inequality is always true.

Note that in our proof, $|y_p - y_q|$ can be chosen arbitrarily large while $\hat{d}_\tau(p, q) = \pi + \frac{\pi^3}{4}$ remains the same. Therefore the \hat{d}_τ -ball at p of radius $R > \pi + \frac{\pi^3}{4}$ is unbounded with respect to the usual Euclidean distance.

Remark 3.9 (Global hyperbolicity in Theorem 1.9 is sharp). Given that in the causal ladder of spacetimes, causal simplicity comes just before global hyperbolicity, the previous Example 3.8 shows that the assumptions in Theorem 1.9 are sharp in view of the causal structure required. It is, however, still open whether time functions that are not Cauchy temporal encode causality in globally hyperbolic spacetimes. In view of our results in Section 2, we conjecture that at least for admissible time functions (see Definition 2.1) this should be the case. The next Section, however, shows why instead of dropping the regularity it makes more sense to consider even better temporal functions, namely h -steep ones (see Definition 4.2).

4 Completeness

In this final section we prove our Main Theorem 1.4 which characterizes global hyperbolicity of (M, g) by metric completeness of (M, \hat{d}_τ) . Completeness is a global statement, therefore we cannot expect (M, \hat{d}_τ) to be complete for all choices of τ (even though they are locally equivalent by Section 2). In Section 3 we have observed that Cauchy temporal functions encode causality globally. Therefore, it comes at no surprise that it is precisely this class of time functions that gives rise to our completeness result. We prove the following refined version of Theorem 1.4.

Theorem 4.1. *Let (M, g) be a spacetime.*

- (i) *If τ is a time function such that (M, \hat{d}_τ) is a complete metric space, then τ is a Cauchy time function. In particular, (M, g) is globally hyperbolic.*

(ii) If (M, g) is globally hyperbolic, then there exists an h -steep (with respect to a complete Riemannian metric h) Cauchy temporal function τ such that (M, \hat{d}_τ) is a complete metric space.

The notion of h -steep temporal functions with respect to a (complete) Riemannian metric h was recently defined by Bernard and Suhr [11, p. 473] to study closed cone fields. Independently, f -steep functions with respect to any positive homogenous C^1 function (not just $f = \|\cdot\|_h$) were introduced by Minguzzi [37, p. 2] in the analysis of Lorentz–Finsler spaces.

Definition 4.2. Let (M, g) be a spacetime and let h be a complete Riemannian metric on M . We say that $\tau: M \rightarrow \mathbb{R}$ is an h -steep temporal function if for all causal vectors $v \in TM$

$$d\tau(v) \geq \|v\|_h.$$

It was shown independently by Bernard and Suhr [11, Thm. 3] and Minguzzi [37, Thm. 3.1] that the existence of an h -steep temporal function is equivalent to global hyperbolicity of the spacetime. The earlier notion of a steep temporal function by Müller and Sánchez [40, Thm. 1.1] does not imply global hyperbolicity.

Proof of Theorem 4.1 and Theorem 1.4. (i) Assume that (M, \hat{d}_τ) is complete but τ is not a Cauchy time function. Then there exists, without loss of generality, a future-directed future-inextendible causal curve $\gamma: \mathbb{R} \rightarrow M$ such that $\lim_{s \rightarrow \infty} \tau(\gamma(s)) < \infty$. Consider the sequence $(p_n)_n$ of points given by $p_n = \gamma(n)$. Since the p_n are causally related among each other $\hat{d}_\tau(p_n, p_m) = |\tau(p_n) - \tau(p_m)|$. Then the fact that $\tau \circ \gamma: \mathbb{R} \rightarrow \mathbb{R}$ is strictly increasing and bounded from above implies that $(p_n)_n$ is a Cauchy sequence in (M, \hat{d}_τ) . By completeness there exists a limit point p , and since γ is continuous, $p \in \bar{\gamma}$, a contradiction to the inextendibility of γ . Hence τ must be a Cauchy time function, and (M, g) globally hyperbolic by Theorem 1.3.

(ii) By [11, 37], global hyperbolicity of (M, g) implies that there is a smooth h -steep temporal function τ with respect to a complete Riemannian metric h . We show that τ is anti-Lipschitz with respect to the (complete) distance d_h induced by h , i.e., there is a $C > 0$ such that

$$(p, q) \in J^+ \implies \tau(q) - \tau(p) \geq C d_h(p, q). \quad (46)$$

Pick any $q \in J^+(p)$ and $\gamma: [0, 1] \rightarrow M$ a causal curve with $\gamma(0) = p$, $\gamma(1) = q$. Then by h -steepness of τ

$$\begin{aligned} \tau(q) - \tau(p) &= \int_0^1 d\tau(\dot{\gamma}(s)) ds \\ &\geq \int_0^1 \sqrt{h(\dot{\gamma}(s), \dot{\gamma}(s))} ds = L_h(\gamma) \geq d_h(p, q). \end{aligned}$$

Thus (46) holds globally, and a theorem of Allen and Burtscher [1, Thm. 1.6] implies that (M, \hat{d}_τ) is complete (and definite). \square

Remark 4.3. Recall that (M, \hat{d}_τ) is always a locally compact length-metric space [1, Thm. 1.1]. If (M, \hat{d}_τ) is also complete, the Hopf–Rinow–Cohn–Vossen Theorem implies that any pair of points can be joined by a \hat{d}_τ -length minimizing curve. Beware that the minimizer is, in general, only \hat{d}_τ -rectifiable, but not necessarily piecewise causal [1, Ex. 3.17]. If τ is Cauchy temporal, thanks to Theorem 1.9, we still do know that the null distance between two points is their difference in time precisely when there is a causal curve between them.

In the second part of the proof, we have not used the fact that τ is Cauchy. This observation leads to an alternative proof of the following result, originally due to Bernard and Suhr [11, Thm. 3].

Corollary 4.4. *If a temporal function τ on a spacetime (M, g) is h -steep with respect to a complete Riemannian metric h , then τ is Cauchy.*

Proof. In the proof of Theorem 4.1(ii) we have shown that if a temporal function (not necessarily Cauchy) is steep with respect to a complete Riemannian metric, then (M, \hat{d}_τ) is complete. But if (M, \hat{d}_τ) is complete, then by Theorem 4.1(i), τ must be Cauchy. \square

We conclude our paper with a counterexample that shows that non- h -steep temporal functions on globally hyperbolic spacetimes do, in general, not imply metric completeness.

Example 4.5 (Cauchy temporal function with incomplete null distance). In [45, Sec. 6.4], Sánchez constructs a globally hyperbolic spacetime $(M, g) = (\mathbb{R}^2, -dt^2 + f^2(t, x)dx^2)$ with a certain piecewise defined L^1 -function $f: M \rightarrow (0, \infty)$ and such that t is a Cauchy temporal function, but the Riemannian slice $\{t = 0\} = (\mathbb{R}, f^2(0, x)dx^2)$, that is, the Riemannian manifold $(\mathbb{R}, f^2(0, x)dx^2)$, is geodesically incomplete. Let $(0, x), (0, y)$ be two points on the $\{t = 0\}$ slice. Then we can estimate their null distance by a sequence of piecewise null curves $\gamma_n(s) = (\gamma_n^t(s), s)$ satisfying $0 \leq \gamma_n^t(s) \leq \frac{1}{n}$. We obtain

$$\hat{d}_t((0, x), (0, y)) \leq \hat{L}_t(\gamma_n) = \int_x^y |\dot{\gamma}_n^t(s)| ds = \int_x^y f(\gamma_n^t(s), s) ds.$$

Applying dominated convergence to the right hand side yields

$$\lim_{n \rightarrow \infty} \int_x^y f(\gamma_n^t(s), s) ds = \int_x^y f(0, s) ds \leq \|f(0, \cdot)\|_{L^1(\mathbb{R})} < \infty.$$

This implies that the sequence $(n)_n$ is Cauchy because for any $\varepsilon > 0$, assuming that $m \leq n$ sufficiently large,

$$\hat{d}_t((0, m), (0, n)) \leq \int_m^n f(0, s) ds \leq \int_m^\infty f(0, s) ds < \varepsilon.$$

The hypothetical limit point at ∞ , however, is not in M . Therefore, (M, \hat{d}_t) is incomplete.

References

- [1] B. Allen and A. Burtscher. Properties of the null distance and spacetime convergence. *Int. Math. Res. Not. IMRN* 10:7729–7808, 2022.
- [2] L. Ambrosio, N. Gigli, and G. Savaré. *Gradient flows in metric spaces and in the space of probability measures*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, Second, 2008.
- [3] L. Ambrosio and B. Kirchheim. Currents in metric spaces. *Acta Math.* 185 (1):1–80, 2000.
- [4] L. Andersson, G. J. Galloway, and R. Howard. The cosmological time function. *Classical Quantum Gravity* 15 (2):309–322, 1998.
- [5] A. Avez. Essais de géométrie riemannienne hyperbolique globale. Applications à la relativité générale. *Ann. Inst. Fourier (Grenoble)* 13 (fasc. 2):105–190, 1963.
- [6] J. K. Beem. Globally hyperbolic space-times which are timelike Cauchy complete. *Gen. Relativity Gravitation* 7 (4):339–344, 1976.
- [7] J. K. Beem and P. E. Ehrlich. Geodesic completeness of submanifolds in Minkowski space. *Geom. Dedicata* 18 (2):213–226, 1985.
- [8] J. K. Beem, P. E. Ehrlich, and K. L. Easley. *Global Lorentzian geometry*. Monographs and Textbooks in Pure and Applied Mathematics, vol. 202. Marcel Dekker, Inc., New York, Second, 1996.
- [9] A. N. Bernal and M. Sánchez. Smoothness of time functions and the metric splitting of globally hyperbolic spacetimes. *Comm. Math. Phys.* 257 (1):43–50, 2005.
- [10] ———. Globally hyperbolic spacetimes can be defined as ‘causal’ instead of ‘strongly causal’. *Class. Quant. Grav.* 24:745–750, 2007.
- [11] P. Bernard and S. Suhr. Lyapounov functions of closed cone fields: from Conley theory to time functions. *Comm. Math. Phys.* 359 (2):467–498, 2018.
- [12] A. Borde, H. F. Dowker, R. S. Garcia, R. D. Sorkin, and S. Surya. Causal continuity in degenerate spacetimes. *Classical Quantum Gravity* 16 (11):3457–3481, 1999.
- [13] R. Budic, J. Isenberg, L. Lindblom, and P. B. Yasskin. On determination of Cauchy surfaces from intrinsic properties. *Comm. Math. Phys.* 61 (1):87–95, 1978.
- [14] A. Burtscher and L. García-Heveling. Time functions on Lorentzian length spaces, 2021. arXiv:2108.02693 [gr-qc].
- [15] A. Y. Burtscher. Length structures on manifolds with continuous Riemannian metrics. *New York J. Math.* 21:273–296, 2015.
- [16] H. Busemann. Timelike spaces. *Dissertationes Math. (Rozprawy Mat.)* 53:52, 1967.
- [17] S. Y. Cheng and S. T. Yau. Maximal space-like hypersurfaces in the Lorentz-Minkowski spaces. *Ann. of Math. (2)* 104 (3):407–419, 1976.

- [18] P. T. Chruściel. *Geometry of black holes*. Int. Ser. Monogr. Phys., vol. 169. Oxford: Oxford University Press, 2020.
- [19] F. Erkekoğlu. A survey on sufficient conditions for geodesic completeness of nondegenerate hypersurfaces in Lorentzian geometry. *Int. J. Geom. Methods Mod. Phys.* 13 (3):1630003, 9, 2016.
- [20] H. Federer and W. H. Fleming. Normal and integral currents. *Ann. of Math. (2)* 72:458–520, 1960.
- [21] Y. Fourès-Bruhat. Théorème d’existence pour certains systèmes d’équations aux dérivées partielles non linéaires. *Acta Math.* 88:141–225, 1952.
- [22] G. J. Galloway. Some results on Cauchy surface criteria in Lorentzian geometry. *Illinois J. Math.* 29 (1):1–10, 1985.
- [23] R. Geroch. Domain of dependence. *J. Mathematical Phys.* 11:437–449, 1970.
- [24] S. G. Harris. Closed and complete spacelike hypersurfaces in Minkowski space. *Classical Quantum Gravity* 5 (1):111–119, 1988.
- [25] S. W. Hawking and G. F. R. Ellis. *The large scale structure of space-time*. Cambridge University Press, London-New York, 1973.
- [26] S. Hawking. The existence of cosmic time functions. *Proc. Roy. Soc. Lond. A* A308:433–435, 1968.
- [27] R. A. Hounnonkpe and E. Minguzzi. Globally hyperbolic spacetimes can be defined without the ‘causal’ condition. *Classical Quantum Gravity* 36 (19):197001, 9, 2019.
- [28] J. L. Jauregui and D. A. Lee. Lower semicontinuity of ADM mass under intrinsic flat convergence. *Calc. Var. Partial Differential Equations* 60 (5):Paper No. 193, 42, 2021.
- [29] M. Kunzinger and R. Steinbauer. Null distance and convergence of Lorentzian length spaces. *Ann. Henri Poincaré*, 2022. <https://doi.org/10.1007/s00023-022-01198-6>.
- [30] K. Landsman. *Foundations of General Relativity: From Einstein to Black Holes*. Radboud University Press, Nijmegen, 2021.
- [31] U. Lang. Local currents in metric spaces. *J. Geom. Anal.* 21 (3):683–742, 2011.
- [32] U. Lang and S. Wenger. The pointed flat compactness theorem for locally integral currents. *Comm. Anal. Geom.* 19 (1):159–189, 2011.
- [33] J. M. Lee. *Manifolds and differential geometry*. Graduate Studies in Mathematics, vol. 107. American Mathematical Society, Providence, RI, 2009.
- [34] J. Leray. *Hyperbolic differential equations*. Institute of Advanced Study, 1953, reprinted 1955.
- [35] E. Minguzzi. K -causality coincides with stable causality. *Comm. Math. Phys.* 290 (1):239–248, 2009.
- [36] ———. Time functions as utilities. *Comm. Math. Phys.* 298 (3):855–868, 2010.
- [37] ———. The representation of spacetime through steep time functions. *J. Phys. Conf. Ser.* 968:012009, 2018.
- [38] ———. Causality theory for closed cone structures with applications. *Rev. Math. Phys.* 31 (5):1930001, 139, 2019.
- [39] ———. Lorentzian causality theory. *Living Rev. Rel.* 22 (1):3, 2019.
- [40] O. Müller and M. Sánchez. Lorentzian manifolds isometrically embeddable in \mathbb{L}^N . *Trans. Amer. Math. Soc.* 363 (10):5367–5379, 2011.
- [41] J. R. Munkres. *Topology*. Prentice Hall, Inc., Upper Saddle River, NJ, 2000.

- [42] K. Nomizu and H. Ozeki. The existence of complete Riemannian metrics. *Proc. Amer. Math. Soc.* 12:889–891, 1961.
- [43] R. Penrose. Gravitational collapse and space-time singularities. *Phys. Rev. Lett.* 14:57–59, 1965.
- [44] A. Sakovich and C. Sormani. The null distance encodes causality, 2022. arXiv:2208.01975 [math.DG].
- [45] M. Sánchez. Globally hyperbolic spacetimes: slicings, boundaries and counterexamples, 2021. arXiv:2110.13672 [gr-qc].
- [46] H.-J. Seifert. Global connectivity by timelike geodesics. *Z. Naturforsch 22a*:1356–1360, 1967.
- [47] R. D. Sorkin and E. Woolgar. A causal order for spacetimes with C^0 Lorentzian metrics: proof of compactness of the space of causal curves. *Classical Quantum Gravity* 13 (7):1971–1993, 1996.
- [48] C. Sormani and C. Vega. Null distance on a spacetime. *Classical Quantum Gravity* 33 (8):085001, 2016.
- [49] C. Sormani and S. Wenger. The intrinsic flat distance between Riemannian manifolds and other integral current spaces. *J. Differential Geom.* 87 (1):117–199, 2011.
- [50] C. Vega. Spacetime distances: an exploration, 2021. arXiv:2103.01191 [gr-qc].
- [51] R. M. Wald and P. Yip. On the existence of simultaneous synchronous coordinates in spacetimes with spacelike singularities. *J. Math. Phys.* 22 (11):2659–2665, 1981.