

LINEAR RECURRENCES OF ORDER AT MOST TWO IN NONTRIVIAL SMALL DIVISORS AND LARGE DIVISORS

HÙNG VIỆT CHU, KEVIN HUU LE, STEVEN J. MILLER, YUAN QIU, AND LIYANG SHEN

ABSTRACT. For each positive integer N , define

$$S'_N = \{1 < d < \sqrt{N} : d|N\} \text{ and } L'_N = \{\sqrt{N} < d < N : d|N\}.$$

Recently, Chentouf characterized all positive integers N such that the set of small divisors $\{d \leq \sqrt{N} : d|N\}$ satisfies a linear recurrence of order at most two. We non-trivially extend the result by excluding the trivial divisor 1 from consideration, which dramatically increases the analysis complexity. Our first result characterizes all positive integers N such that S'_N satisfies a linear recurrence of order at most two. Moreover, our second result characterizes all positive N such that L'_N satisfies a linear recurrence of order at most two, thus extending considerably a recent result that characterizes N with L'_N being in an arithmetic progression.

CONTENTS

1. Introduction	1
2. Preliminaries	4
2.1. Regarding S'_N	4
3. Small recurrent numbers	6
3.1. The case $ S'_N \geq 4$	6
3.2. The case $ S'_N \leq 3$	12
4. Large recurrent numbers	12
4.1. The case $ L'_N \geq 4$	12
4.2. The case $ L'_N \leq 3$	17
5. Appendix	17
References	19

1. INTRODUCTION

For a positive integer N , the set of small divisors of N is

$$S_N := \{d : 1 \leq d \leq \sqrt{N}, d \text{ divides } N\}.$$

Since the case $N = 1$ is trivial, we assume throughout the paper that $N > 1$. In 2018, Iannucci characterized all positive integers N whose S_N forms an arithmetic progression (or AP, for short). Iannucci's key idea was to show that if S_N forms an AP, then the size $|S_N|$ cannot exceed 6. Observing that the trivial divisor 1 plays an important role in Iannucci's proofs (see [4, Lemma 3 and Theorem 4]), Chu [3] excluded both 1

and \sqrt{N} from the definition of S_N to obtain a more general theorem that characterizes all N whose

$$S'_N := \{d : 1 < d < \sqrt{N}, d \text{ divides } N\}$$

is in an AP. Interestingly, with the trivial divisor 1 excluded, [2, Theorem 1.1] still gives that $|S'_N| \leq 5$. Recently, Chentouf generalized Iannucci's result from a different perspective by characterizing all N whose S_N satisfies a linear recurrence of order at most two. In particular, for each tuple $(u, v, a, b) \in \mathbb{Z}^4$, there is an integral linear recurrence, denoted by $U(u, v, a, b)$, of order at most two, given by

$$n_i = \begin{cases} u & \text{if } i = 1, \\ v & \text{if } i = 2, \\ an_{i-1} + bn_{i-2} & \text{if } i \geq 3. \end{cases}$$

Noting that the appearance of the trivial divisor 1 contributes nontrivially to the proof of [1, Theorem 3, Lemma 8, Theorem 10], we generalize Chentouf's result in the same manner as [2, Theorem 1.1] generalizes [4, Theorem 4]: we characterize all positive integers N whose S'_N satisfies a linear recurrence of order at most two.

Definition 1.1. A positive integer N is said to be small recurrent if S'_N satisfies a linear recurrence of order at most two. When $|S'_N| \leq 2$, N is vacuously small recurrent.

Theorem 1.2. Let p, q, r denote prime numbers such that $p < q < r$ and k be some positive integer. A positive integer $N > 1$ is small recurrent if and only if N belongs to one of the following forms.

- (1) $N = p^k$ for some $k \geq 1$. In this case, $S'_N = \{p, p^2, \dots, p^{\lfloor (k-1)/2 \rfloor}\}$ satisfies $U(p, p^2, p, 0)$.
- (2) $N = p^k q$ or $N = pq^k$ for some $1 \leq k \leq 3$. A restriction for $N = p^3 q$ is that either $p < q < p^2$ or $p^3 < q$.
- (3) $N = p^k q$ for some $k \geq 4$ and $q > p^k$. In this case, $S'_N = \{p, p^2, p^3, \dots, p^k\}$ satisfies $U(p, p^2, p, 0)$.
- (4) $N = p^k q$ for some $k \geq 4$ and $\sqrt{q} < p < q$. In this case, $S'_N = \{p, q, p^2, pq, \dots\}$ satisfies $U(p, q, 0, p)$.
- (5) $N = pq^k$ for some $k \geq 4$ and $p < q$. In this case, $S'_N = \{p, q, pq, q^2, \dots\}$ satisfies $U(p, q, 0, q)$.
- (6) $N = pq^k r$ for some $k \geq 2$, $p < q$, and $r > pq^k$. In this case, $S'_N = \{p, q, pq, q^2, \dots, pq^{k-1}, q^k, pq^k\}$ satisfies $U(p, q, 0, q)$.
- (7) $N = p^2 q^2$ for some $p < q < p^2$. In this case, $S'_N = \{p, q, p^2\}$.
- (8) $N = pqr$ for some $p < q < r$. If $r < pq$, then $S'_N = \{p, q, r\}$. If $r > pq$, then $S'_N = \{p, q, pq\}$.
- (9) $N = p^3 q^2$ for some $p^{3/2} < q < p^2$. In this case, $S'_N = \{p, q, p^2, pq, p^3\}$ satisfies $U(p, q, 0, p)$.
- (10) $N = p^2 qr$, where $p < q < p^2 < r < pq$, $(q^2 - p^3) | (pq - r)$, $(q^2 - p^3) | (rq - p^4)$, and $r = pq - \sqrt{(q^2 - p^3)(p^2 - q)}$. In this case, $S'_N = \{p, q, p^2, r, pq\}$ satisfies $U\left(p, q, \frac{p(pq-r)}{q^2-p^3}, \frac{rq-p^4}{q^2-p^3}\right)$.

Next, consider the set of large divisors of N

$$L_N := \{d : d \geq \sqrt{N}, d \text{ divides } N\} \text{ and } L'_N := \{d : \sqrt{N} < d < N, d \text{ divides } N\}.$$

The second result of this paper is the characterization of all positive integers N whose L'_N satisfies a linear recurrence of order at most two. This considerably extends [2, Theorem 1].

Definition 1.3. A positive integer N is said to be large recurrent if L'_N satisfies a linear recurrence of order at most two. When $|L'_N| \leq 2$, N is vacuously large recurrent.

Theorem 1.4. Let p, q, r denote prime numbers such that $p < q < r$ and k be some positive integer. A positive integer $N > 1$ is large recurrent if and only if N belongs to one of the following forms.

- (1) $N = p^k$ for some $k \geq 1$. In this case, $L'_N = \{p^{\lceil (k-1)/2 \rceil + 1}, p^{\lceil (k-1)/2 \rceil + 2}, \dots, p^{k-1}\}$ satisfies $U(p^{\lceil (k-1)/2 \rceil + 1}, p^{\lceil (k-1)/2 \rceil + 2}, p, 0)$.
- (2) $N = p^k q$ for some $k \geq 1$ and $q > p^k$. In this case, $L'_N = \{q, pq, p^2 q, \dots, p^{k-1} q\}$ satisfies $U(q, pq, p, 0)$.
- (3) $N = p^k q$ for some $k \geq 2$ and $p^{k-1} < q < p^k$. Then

$$L'_N = \{p^k, pq, p^2 q, \dots, p^{k-1} q\}$$

satisfies $U(p^k, pq, p, 0)$.

- (4) $N = p^k q$ some for $k \geq 3$ and $p < q < p^2$. In this case,

$$L'_N = \begin{cases} \{p^{k/2+1}, p^{k/2} q, p^{k/2+2}, \dots, p^{k-1} q\} & \text{if } 2|k, \\ \{p^{(k-1)/2} q, p^{(k+3)/2}, p^{(k+1)/2} q, \dots, p^{k-1} q\} & \text{if } 2 \nmid k. \end{cases}$$

Observe that L'_N satisfies $U(p^{k/2+1}, p^{k/2} q, 0, p)$ and $U(p^{(k-1)/2} q, p^{(k+3)/2}, 0, p)$ for even and odd k , respectively.

- (5) $N = p^4 q$ with $p^2 < q < p^3$, $(p^5 - q^2)|(p^2 - q)$, and $(p^5 - q^2)|(p^3 - q)$. In this case, $L'_N = \{pq, p^4, p^2 q, p^3 q\}$.
- (6) $N = p^3 q^2$ for $p < q < p^2$. In this case, $L'_N = \{q^2, p^2 q, pq^2, p^3 q, p^2 q^2\}$ satisfies $U(q^2, p^2 q, 0, p)$.
- (7) $N = p^2 q^2$ for some $p < q$. If $p < q < p^2$, then $L'_N = \{q^2, p^2 q, pq^2\}$.
- (8) $N = pq^k$ for some $k \geq 2$ and $p < q$. In this case,

$$L'_N = \begin{cases} \{pq^{\frac{k}{2}}, q^{\frac{k}{2}+1}, \dots, q^k\} & \text{if } 2|k, \\ \{q^{\frac{k+1}{2}}, pq^{\frac{k+1}{2}}, \dots, q^k\} & \text{if } 2 \nmid k. \end{cases}$$

Observe that L'_N satisfies $U(pq^{k/2}, q^{k/2+1}, 0, q)$ and $U(q^{(k+1)/2}, pq^{(k+1)/2}, 0, q)$ for even and odd k , respectively.

- (9) $N = pq^k r$ for some $k \geq 1$ and $p < q < pq^k < r$. In this case, $L'_N = \{r, pr, qr, pqr, q^2 r, \dots, q^k r\}$ satisfies $U(r, pr, 0, q)$.

The paper is structured as follows: Section 2 studies the case when N has a small number of divisors and establish some preliminary results; Section 3 characterizes small recurrent numbers, while Section 4 characterizes large recurrent numbers.

2. PRELIMINARIES

For each $N \in \mathbb{N}$ with the prime factorization $\prod_{i=1}^{\ell} p_i^{a_i}$, the divisor-counting function is

$$\tau(N) := \sum_{d|N} 1 = \prod_{i=1}^{\ell} (a_i + 1). \quad (2.1)$$

It is easy to verify that for $N > 1$,

$$\tau(N) := \begin{cases} 2|S'_N| + 3 = 2|L'_N| + 3 & \text{if } N \text{ is a square,} \\ 2|S'_N| + 2 = 2|L'_N| + 2 & \text{otherwise.} \end{cases} \quad (2.2)$$

Using (2.1), we can characterize all N with $\tau(N) \leq 9$; equivalently, $|S'_N|, |L'_N| \leq 3$.

- (i) If $\tau(N) = 2$ or 3 , (2.1) gives that $N = p$ or p^2 for some prime p ,
- (ii) If $\tau(N) = 4$ or 5 , $N = pq, p^3, p^4$ for some primes $p < q$,
- (iii) If $\tau(N) = 6$ or 7 , $N = p^5, pq^2, p^2q, p^6$ for some primes $p < q$,
- (iv) If $\tau(N) = 8$ or 9 , $N = pqr, pq^3, p^3q, p^7, p^2q^2, p^8$ for some primes $p < q < r$.

2.1. Regarding S'_N .

Proposition 2.1. *If $|S'_N| \geq 2$, then N cannot have two (not necessarily distinct) prime factors p_1 and p_2 at least \sqrt{N}*

Proof. Suppose that N has two prime factors p_1 and p_2 at least \sqrt{N} . Then $p_1p_2 \geq N$ and p_1p_2 divides N ; hence, $N = p_1p_2$, which contradicts that $|S'_N| \geq 2$. \square

Proposition 2.2. *If all elements of S'_N are divisible by some prime p and $|S'_N| \geq 4$, then either $N = p^k$ or $N = p^kq$ for some $k \geq 1$ and some prime $q > p^k$.*

Proof. If all divisors (except 1) of N are divisible by p , then $n = p^k$ for some $k \geq 1$. Assume that N has a prime factor $q \neq p$. Then $q \geq \sqrt{N}$. Proposition 2.1 implies that N cannot have another prime factor at least \sqrt{N} . Hence, $N = p^kq$ for some $k \geq 1$ and $q > p^k$. \square

Let $p < q < r < s$ be distinct prime numbers. Write $S'_N = \{d_2, d_3, d_4, d_5, \dots\}$. (We start with d_2 since the smallest divisor of N is usually denoted by $d_1 = 1$, which is excluded from S'_N .)

Lemma 2.3. *Suppose that the first 4 numbers in S'_N are $p < q < p^2 < r$. If N is small recurrent with $U(p, q, a, b)$, the following hold:*

- i) $\gcd(a, b) = 1$.
- ii) if $d_{2i} \in S'_N$, then $p|d_{2i}$; however, if $d_{2i-1} \in S'_N$, then $p \nmid d_{2i-1}$.
- iii) for $d_i, d_{2i-1} \in S'_N$, we have $\gcd(b, d_i) = \gcd(a, d_{2i-1}) = 1$.
- iv) for $d_i, d_{i+1} \in S'_N$, we have $\gcd(d_i, d_{i+1}) = 1$.
- v) for $d_{2i-1}, d_{2i+1} \in S'_N$, we have $\gcd(d_{2i-1}, d_{2i+1}) = 1$.

Proof. i) Since $r = ap^2 + bq$, we have $\gcd(a, b)|r$. Observe that $\gcd(a, b) = r$ contradicts $p^2 = aq + bp$. Hence, $\gcd(a, b) = 1$.

ii) We prove by induction. The claim is true for $i \leq 2$. Assume that the claim holds for $i = j \geq 2$. We show that it holds for $i = j + 1$. Since $p^2 = aq + bp$, we know that $p|a$. By item i), $p \nmid b$. Write

$$\begin{aligned} d_{2(j+1)} &= ad_{2(j+1)-1} + bd_{2(j+1)-2} = ad_{2j+1} + bd_{2j} \\ &= (a^2 + b)d_{2j} + abd_{2j-1}. \end{aligned}$$

By the inductive hypothesis, $p|d_{2j}$. Since $p|a$, we obtain $p|d_{2(j+1)}$. Furthermore, write

$$d_{2(j+1)-1} = ad_{2(j+1)-2} + bd_{2(j+1)-3} = ad_{2j} + bd_{2j-1}.$$

Since $p \nmid bd_{2j-1}$ by the inductive hypothesis and $p|ad_{2j}$, we obtain $p \nmid d_{2(j+1)-1}$. This completes our proof.

iii) Suppose that $k = \gcd(b, d_i) > 1$ for some i . If $i = 2$, then $p|b$, which contradicts that $p|a$ and $\gcd(a, b) = 1$. If $i = 3$, then $q|b$, which contradicts $p^2 = aq + bp$. If $i \geq 4$, write $d_i = ad_{i-1} + bd_{i-2}$. Since $k|bd_{i-2}$ and $\gcd(a, b) = 1$, we get $k|d_{i-1}$ and so, $k|\gcd(b, d_{i-1})$. By induction, we obtain $k|\gcd(b, d_3)$, which has been shown to be impossible.

Next, suppose that $k = \gcd(a, d_{2i-1}) > 1$ for some i . If $i = 2$, then $q|a$, contradicting $r = ap^2 + bq$. If $i \geq 3$,

$$k \mid d_{2i-1}, d_{2i-1} = ad_{2i-2} + bd_{2i-3}, \text{ and } \gcd(a, b) = 1 \implies k \mid d_{2i-3}.$$

By induction, $k|d_3$; that is, $\gcd(a, d_3) > 1$, which has been shown to be impossible.

iv) The claim holds for $i \leq 4$. For $i \geq 5$, using the recurrence $d_{i+1} = ad_i + bd_{i-1}$, we have

$$\gcd(d_i, d_{i+1}) = \gcd(d_i, bd_{i-1}) \stackrel{\text{iii)}}{=} \gcd(d_i, d_{i-1}).$$

By induction, we obtain $\gcd(d_i, d_{i+1}) = 1$.

v) The claim holds for $i \leq 2$. For $i \geq 3$, we have

$$\gcd(d_{2i-1}, d_{2i+1}) = \gcd(d_{2i-1}, ad_{2i}) \stackrel{\text{iii)}}{=} \gcd(d_{2i-1}, d_{2i}) \stackrel{\text{iv)}}{=} 1.$$

This completes our proof. \square

Lemma 2.4. *Suppose that the first 4 numbers in S'_N are $p < q < r < p^2$. If N is small recurrent with $U(p, q, a, b)$, the following hold:*

- i) $\gcd(a, b) = 1$ and $p \nmid a$.
- ii) For all $d_i \in S'_N$, $\gcd(b, d_i) = 1$.
- iii) For all $d_i, d_{i+1} \in S'_N$, $\gcd(d_i, d_{i+1}) = 1$.
- iv) Let $d_i \in S'_N$. Then $p|d_i$ if and only if $i \equiv 2 \pmod 3$.

Proof. i) We have $\gcd(a, b)$ divides r because $r = aq + bp$. Furthermore, $\gcd(a, b)$ divides p because $p^2 = ar + bq$. Hence, $\gcd(a, b) = 1$.

Since $r = aq + bp$ and r is a prime, $p \nmid a$.

ii) Suppose that $k = \gcd(b, d_i) > 1$ for some i . If $i = 2$, then $k = p$ and $p|b$. It follows from $p^2 = ar + bq$ that $p|a$, which contradicts that $\gcd(a, b) = 1$. If $i = 3$, then $k = q$ and $q|b$. It follows from $r = aq + bp$ that $q|r$, a contradiction. Hence, $i \geq 4$. Write

$$1 < \gcd(b, d_i) = \gcd(b, ad_{i-1} + bd_{i-2}) = \gcd(b, d_{i-1}).$$

By induction, we obtain $\gcd(b, d_3) > 1$, which has been shown to be impossible.

iii) The claim holds for $i \leq 4$. Let $d_i, d_{i+1} \in S'_N$ and $i \geq 5$. Then

$$\gcd(d_i, d_{i+1}) = \gcd(d_i, ad_i + bd_{i-1}) = \gcd(d_i, bd_{i-1}) \stackrel{\text{ii)}}{=} \gcd(d_i, d_{i-1}).$$

By induction, $\gcd(d_i, d_{i+1}) = 1$.

iv) The claim holds for $i \leq 4$. Suppose that the claim holds for all $i \leq j$ for some $j \geq 5$. We show that it also holds for $i = j + 1$ under the assumption that $d_{j+1} \in S'_N$. We have

$$p^2 = ar + bq = a(aq + bp) + bq = (a^2 + b)q + abp.$$

Hence, $p|(a^2 + b)$. Write

$$d_{j+1} = ad_j + bd_{j-1} = a(ad_{j-1} + bd_{j-2}) + bd_{j-1} = (a^2 + b)d_{j-1} + abd_{j-2}.$$

Therefore, $p|d_{j+1}$ if and only if $p|d_{j-2}$. By the inductive hypothesis, we know that $p|d_{j+1}$ if and only if $j + 1 \equiv 2 \pmod{3}$. \square

The proofs of the next two lemmas are similar to those of Lemmas 2.3 and 2.4. Thus, we move their proofs to the Appendix.

Lemma 2.5. *Suppose that the first 4 numbers in S'_N are $p < q < r < pq$. If N is small recurrent with $U(p, q, a, b)$, the following hold:*

- i) $\gcd(a, b) = 1$ and $p \nmid a$.
- ii) For all $d_i \in S'_N$, $\gcd(b, d_i) = 1$.
- iii) For all $d_i, d_{i+1} \in S'_N$, $\gcd(d_i, d_{i+1}) = 1$.
- iv) Let $d_i \in S'_N$. Then $p|d_i$ if and only if $i \equiv 2 \pmod{3}$.
- v) Let $d_i \in S'_N$. Then $q|d_i$ if and only if i is odd.

Lemma 2.6. *Suppose that the first 4 numbers in S'_N are $p < q < r < s$. If N is small recurrent with $U(p, q, a, b)$, the following hold:*

- i) $\gcd(a, b) = 1$.
- ii) For all $d_i \in S'_N$ with $i \geq 3$, $\gcd(b, d_i) = 1$.
- iii) For $d_i, d_{i+1} \in S'_N$, $\gcd(d_i, d_{i+1}) = 1$.
- iv) For $d_i \in S'_N$, $\gcd(a, d_i) = 1$.
- v) For $d_i, d_{i+2} \in S'_N$, $\gcd(d_i, d_{i+2}) = 1$.

3. SMALL RECURRENT NUMBERS

We first find all small recurrent numbers with $|S'_N| \geq 4$. Then we check which N is small recurrent out of all N with $|S'_N| \leq 3$ at the end of this section. As we rely heavily on case analysis, we underline possible forms of N throughout our analysis for the ease of later summary.

3.1. The case $|S'_N| \geq 4$. Let $d_2 = p$ for some prime p . Then d_3 is either p^2 or q for some prime $q > p$. If $d_3 = p^2$, according to Proposition 2.2, we know that $N = p^k$ or $N = p^k q$ for some $k \geq 1$ and $q > p^k$.

Assume, for the rest of this subsection, that $d_3 = q$ for some prime $q > p$. Then $d_4 = pq, p^2, r$ for some prime $r > q$.

3.1.1. When $d_4 = pq$. Since S'_N satisfies $U(p, q, a, b)$, we get $pq = aq + bp$. So, $p|a$ and $q|b$. Write $a = pm$ for some $m \in \mathbb{Z}$ and get $b = (1 - m)q$. Since $p^2 \nmid N$ and q divides $d_5 = apq + bq$, we can write

$$d_5 = p^s q^t r_1^{\ell_1} \cdots r_k^{\ell_k},$$

where $s \leq 1$, $t \geq 1$, and r_i 's are primes strictly greater than pq . If some $\ell_i \geq 1$, then $pq < r_i < d_5$ and $r_i \in S'_N$, a contradiction. Hence, $\ell_i = 0$ for all $i \leq k$ and $d_5 = p^s q^t$. Since $d_5 > pq$ and $s \leq 1$, we know that $t \geq 2$.

- a) If $s = 1$, $d_5 = pq^2 > q^2 > d_4$ and $q^2 | d_5$, so $q^2 \in S'_N$, a contradiction.
- b) If $s = 0$, $d_5 = q^t$ for some $t \geq 2$. Since d_5 is the next number after d_4 in increasing order, $d_5 = q^2$. Using the linear recurrence, we obtain $q^2 = apq + bq$, so $q = ap + b = p^2 m + (1 - m)q$. It follows that $p^2 m = mq$. We arrive at $m = 0$, $a = 0$, and $b = q$. Hence, all elements of S'_N are divisible by either p or q . If N has a prime factor $r \geq \sqrt{N}$, by Proposition 2.1, r is unique. We conclude that $N = pq^k$ or $pq^k r$ for some $k \geq 2$, $p < q$, and $pq^k < r$.

3.1.2. When $d_4 = p^2$. The first few divisors of N are $1 < p < q < p^2$. We have $p^2 = aq + bp$, so $p|a$. Write $a = pm$ and get $b = p - mq$. We argue for possible forms of d_5 . Let r be the largest prime factor of d_5 . If $r > q$, then $r > p^2$ and $d_5 = r$. Otherwise, if $r \leq q$, then $d_5 = p^\ell q^k$ for some $\ell, k \geq 0$. Suppose that $k \geq 2$. We get

$$d_5 \geq q^2 > pq > d_4 \text{ and } pq | N,$$

a contradiction. Hence, $k \leq 1$. If $k = 0$, then

$$d_5 = p^3 > pq > d_4 \text{ and } pq | N,$$

another contradiction. Therefore, $k = 1$ and $d_5 = pq$. We conclude that either $d_5 = r$ for some $r > q$ or $d_5 = pq$.

- a) If $d_5 = pq$, then $bq + ap^2 = pq$. So, $(p - mq)q + mp^3 = pq$, which gives $mq^2 = mp^3$. Hence, $m = 0$, $a = 0$, and $b = p$. We know that elements of S'_N are divisible by either p or q . If N has a prime factor r' at least \sqrt{N} , by Proposition 2.1, r' is unique. Hence, either $N = p^\ell q^k$ or $p^\ell q^k r'$ for some prime $r' > p^\ell q^k$, $\ell \geq 2$, and $k \geq 1$.

Case a.i) $N = p^\ell q^k r'$. We claim that $k = 1$. Indeed, if $k \geq 2$, then

$$q^4 < q^2 r' < N \implies q^2 < \sqrt{N} \implies q^2 \in S'_N.$$

Since $b = p$, we know that $p|d$ for all $d \geq d_4$ and $d \in S'_N$, which contradicts $q^2 \in S'_N$. Hence, $N = p^\ell q r'$ for some $\ell \geq 2$, some prime $r' > p^\ell q$, and $\sqrt{q} < p < q$.

Case a.ii) $N = p^\ell q^k$. As above, $q^2 \notin S'_N$. If $k \geq 2$, then

$$q^2 \geq \sqrt{N} \implies q^4 \geq N = p^\ell q^k = p^{\ell-2} p^2 q^k > q^{k+1}.$$

Hence, $k < 3$, which implies that $k = 2$. In this case, $N = p^\ell q^2$ and

$$q^2 > p^\ell > q^{\ell/2} \implies \ell \leq 3.$$

We conclude that one of the following holds:

- $N = p^\ell q$ for some $\ell \geq 2$ and $\sqrt{q} < p < q$,
- $N = p^2 q^2$ for some $p < q < p^2$,

- $N = p^3q^2$ for some $p^{3/2} < q < p^2$.
- b) $d_5 = r$.

Proposition 3.1. *Suppose that the first 4 numbers in S'_N are $p < q < p^2 < r$. Then $|S_N| \leq 7$. As a result, $|S'_N| \leq 6$.*

Proof. Assume that $|S_N| \geq 2i$ for some $i \geq 4$. We obtain a contradiction by showing that $|S_N| \geq 2i + 2$. By Lemma 2.3 item ii), $p \nmid d_{2i-1}$, $p \mid d_{2i-2}$, and $p \nmid d_{2i-3}$. By Lemma 2.3 item v), $\gcd(d_{2i-1}, d_{2i-3}) = 1$, so $p^2 d_{2i-1} d_{2i-3}$ divides N . Hence, $pd_{2i-3} \in S'_N$.

If $pd_{2i-3} = d_{2i-2}$, then

$$pd_{2i-3} = ad_{2i-3} + bd_{2i-4} \implies d_{2i-3} \mid bd_{2i-4},$$

which contradicts Lemma 2.3 items iii) and iv).

If $pd_{2i-3} = d_{2i}$, then

$$\begin{aligned} pd_{2i-3} &= ad_{2i-1} + bd_{2i-2} = a(ad_{2i-2} + bd_{2i-3}) + bd_{2i-2} \\ &= (a^2 + b)d_{2i-2} + abd_{2i-3}. \end{aligned}$$

Therefore, d_{2i-3} divides $a^2 + b$. It is easy to check that for $d_j \in S'_N$, the sequence $d_j \bmod a^2 + b$ is congruent to

$$1, p, q, p^2, abp, abq, abp^2, (ab)^2p, (ab)^2q, (ab)^2p^2, \dots$$

Hence, we can write

$$d_{2i-3} = (a^2 + b)\ell + a^k b^k s,$$

for some $\ell \in \mathbb{Z}$, some $k \geq 1$, and some $s \in \{p, q, p^2\}$. Since $d_{2i-3} \mid (a^2 + b)$, $d_{2i-3} \mid a^k b^k s$. By Lemma 2.3 item iii), $d_{2i-3} \mid s$; that is, $d_{2i-3} \leq p^2$. However, $d_{2i-3} \geq d_5 > d_4 = p^2$, a contradiction.

We conclude that $pd_{2i-3} \geq d_{2i+2}$. Since $pd_{2i-3} \in S'_N$, we know that $d_{2i+2} \in S'_N$ and $|S_N| \geq 2i + 2$. \square

Proposition 3.2. *Suppose that the first 4 numbers in S'_N are $p < q < p^2 < r$. If N is small recurrent, then $|S'_N| \neq 4, 6$.*

Proof. If $|S'_N| = 4$, then (2.2) gives $\tau(N) = 10$ or 11. Note that N has three distinct prime factors p, q, r and the power of p is at least 2. Since $2^3 \cdot 3 > 11$, N cannot have another prime factor besides p, q, r . Write $N = p^a q^b r^c$, for some $a \geq 2, b \geq 1, c \geq 1$. However, neither $(a+1)(b+1)(c+1) = 10$ nor $(a+1)(b+1)(c+1) = 11$ has a solution. Therefore, $|S'_N| \neq 4$. A similar argument gives $|S'_N| \neq 6$. \square

By Propositions 3.1 and 3.2, we know that $|S'_N| = 5$; that is, $\tau(N) = 12$ or 13. Using the same reasoning as in the proof of Proposition 3.2, we know that $\tau(N) = 12$ and $N = p^2qr$, where $p < q < p^2 < r$.

3.1.3. When $d_4 = r$ for some $r > q$. The possible values for d_5 are p^2, pq, s for some prime $s > r$.

a) If $d_5 = p^2$, we can generalize the method by Chentouf.

Proposition 3.3. *If N is small recurrent and the first four numbers of S'_N are $p < q < r < p^2$, then $|S_N| \leq 7$. As a result, $|S'_N| \leq 6$.*

Proof. Suppose that $|S_N| \geq 8$. We show that $|S_N| \geq 3i + 2$ for all $i \in \mathbb{N}$, which is a contradiction. The claim holds for $i = 2$. Assume that $|S_N| \geq 3j + 2$ for some $j \geq 2$. By Lemma 2.4, $p \nmid d_{3j}d_{3j+1}$ and $\gcd(d_{3j}, d_{3j+1}) = 1$. Hence, $p^2d_{3j}d_{3j+1}$ divides N , which implies that $pd_{3j} \in S'_N$.

If $pd_{3j} = d_{3j+2} = ad_{3j+1} + bd_{3j}$, then d_{3j} divides ad_{3j+1} . By Lemma 2.4, $d_{3j} | a$. Observe that for $d_i \in S'_N$, the sequence $d_i \bmod a$ is

$$1, p, q, bp, bq, b^2p, b^2q, \dots$$

Write $d_{3j} = al + b^k s$, for some $\ell \in \mathbb{Z}$, some $k \geq 0$, and some $s \in \{p, q\}$. We see that $d_{3j} | b^k s$ for some $k \geq 0$ and $s \in \{p, q\}$. By Lemma 2.4, $d_{3j} \leq q$. However,

$$d_{3j} \geq d_6 > d_3 = q,$$

a contradiction.

If $pd_{3j} > d_{3j+2}$, then $pd_{3j} \geq d_{3(j+1)+2}$ by Lemma 2.4. Therefore, $|S_N| \geq 3(j+1) + 2$. \square

Proposition 3.4. *There is no small recurrent N whose the first four numbers of S'_N are $p < q < r < p^2$.*

Proof. By Proposition 3.3, $|S'_N| \in \{4, 5, 6\}$. If $|S'_N| = 4$, then $\tau(N) = 10$ or 11 , none of which can be written as a product of at least three integers, each of which is at least 2. This contradicts (2.2) and the fact that N has three distinct prime factors. We arrive at the same conclusion when $|S'_N| = 6$. For $|S'_N| = 5$, we obtain $N = p^2qr$ for some primes $p < q < r < p^2$. However, this poses another contradiction. Observe that $(pq)^2 < p^2qr$, so the divisors in S'_N are $p < q < r < p^2 < pq$. Since $pq = ap^2 + br$, we get $p|b$, which contradicts Lemma 2.4 item ii). \square

b) Suppose that $d_5 = pq$.

Proposition 3.5. *There is no small recurrent number N such that the first four numbers of S'_N are $p < q < r < pq$.*

Proof. Assume that $|S'_N| \geq 8$. Since $p^2 \notin S'_N$, p divides N exactly. By Lemma 2.5, d_6 is divisible neither by p nor q . Hence, $d_6 = s$ for some prime $s > pq$. The divisor d_7 is divisible by q ; hence, $d_7 = qr$ or q^2 . The divisor d_8 is divisible by p but not by q . So, $d_8 = pr$, which gives that d_7 must be q^2 because $d_7 < d_8$. Now $q|d_9$ and $p \nmid d_9 \implies d_9 = qr$. However, that $\gcd(d_8, d_9) = r$ contradicts Lemma 2.5. Therefore, $|S'_N| \in \{4, 5, 6, 7\}$. Using the same argument as in the proof of Proposition 3.4, we know that $|S'_N| \neq 4, 6$ and so, $|S'_N| \in \{5, 7\}$. By the above argument, if $|S'_N| \geq 5$, then d_6 is a prime greater than pq . Hence, N has at least 4 distinct prime factors, so $\tau(N)$ can be written as a product of

at least 4 integers greater than 1. Clearly, (2.1) rules out the case $|S'_N| = 5$. If $|S'_N| = 7$, the above argument shows that $q^2|N$; hence, $\tau(N)$ can be written as a product of at least 4 integers greater than 1, one of which is greater than 2. This cannot happen as $\tau(N) \in \{16, 17\}$. \square

c) Suppose that $d_5 = s$.

Proposition 3.6. *There is no small recurrent number N such that the first four numbers of S'_N are $p < q < r < s$.*

Proof. Observe that pq and pr are in S'_N . Let $d_j = pv$ be the largest element of S'_N that is divisible by p . Clearly, $v > p$ and $j \geq 7$. By Lemma 2.6, d_j, d_{j-1} , and d_{j-2} are pairwise coprime. Hence, $pvd_{j-1}d_{j-2}$ divides N , so $pd_{j-2} \in S'_N$.

If $pd_{j-2} = d_{j-1}$, then $p|d_{j-1}$ and so, $p|\gcd(d_{j-1}, d_j)$, which contradicts Lemma 2.6 item iii).

If $pd_{j-2} = d_j$, then $d_{j-2} = v$ and $\gcd(d_{j-2}, d_j) = v > 1$, which contradicts Lemma 2.6 item v).

Therefore, we have $pd_{j-2} > d_j$, which, however, contradicts that d_j is the largest element of S'_N that is divisible by p . We conclude that there is no small recurrent number N such that the first four numbers of S'_N are $p < q < r < s$. \square

From the above analysis, we arrive at the following proposition.

Proposition 3.7. *If N is small recurrent and $|S'_N| \geq 4$, then N belongs to one of the following forms.*

- (S1) $N = p^k$ or $N = p^kq$ for some $k \geq 1$ and $q > p^k$.
- (S2) $N = pq^k$ or pq^kr for some $k \geq 2$, $p < q$, and $pq^k < r$.
- (S3) $N = p^kqr$ for some $k \geq 2$, some prime $r > p^kq$, and $\sqrt{q} < p < q$.
- (S4) $N = p^kq$ for some $k \geq 2$ and $\sqrt{q} < p < q$.
- (S5) $N = p^2q^2$ for some $p < q < p^2$.
- (S6) $N = p^3q^2$ for some $p^{3/2} < q < p^2$.
- (S7) $N = p^2qr$, where the first four numbers in S'_N are $p < q < p^2 < r$.

These forms together establish the necessary condition for a small recurrent N to have $|S'_N| \geq 4$. We now refine each form (if necessary) to obtain a necessary and sufficient condition.

- (S1) • If $N = p^k$, then N is small recurrent with $|S'_N| \geq 4$ if $k \geq 9$. In this case, $S'_N = \{p, p^2, p^3, \dots, p^{\lfloor (k-1)/2 \rfloor}\}$ satisfies $U(p, p^2, p, 0)$.
- If $N = p^kq$ for some $k \geq 1$ and $q > p^k$, then N is small recurrent with $|S'_N| \geq 4$ if $k \geq 4$. In this case, $S'_N = \{p, p^2, p^3, \dots, p^k\}$ satisfies $U(p, p^2, p, 0)$.
- (S2) • If $N = pq^k$ for some $k \geq 2$ and $p < q$, then N is small recurrent with $|S'_N| \geq 4$ if $\sqrt{N} = \sqrt{pq^k} > q^2$. Hence, $N = pq^k$ for some $k \geq 4$ and $p < q$. In this case, $S'_N = \{p, q, pq, q^2, \dots\}$ satisfies $U(p, q, 0, q)$.
- If $N = pq^kr$ for some $k \geq 2$, $p < q$, and $r > pq^k$, then N is small recurrent with $|S'_N| \geq 4$. In this case, $S'_N = \{p, q, pq, q^2, \dots, pq^{k-1}, q^k, pq^k\}$ satisfies $U(p, q, 0, q)$.

- (S3) If N belongs to (S3), then $S'_N = \{p, q, p^2, pq, \dots, p^k, p^{k-1}q, p^kq\}$. Since $p^2 = aq + bp$, we know that $p|a$. Write $a = pm$ for some $m \in \mathbb{Z}$ and get $b = p - mq$. Hence,

$$pq = ap^2 + bq = p^3m + (p - mq)q \implies mq^2 = p^3m.$$

Therefore, $(m, a, b) = (0, 0, p)$. However, the largest element in S'_N , p^kq , is not equal to $p \cdot p^k$. We conclude that form (S3) does not give a small recurrent number.

- (S4) If $N = p^kq$ for some $k \geq 2$ and $\sqrt{q} < p < q$, then the nontrivial divisors of N in increasing order is $p < q < p^2 < pq < \dots$. In order that $|S'_N| \geq 4$, we need $(pq)^2 < p^kq$, so $q < p^{k-2}$. Hence, $k \geq 4$. In this case, $S'_N = \{p, q, p^2, pq, \dots\}$ satisfies $U(p, q, 0, p)$.
- (S5) If $N = p^2q^2$ for some $p < q < p^2$, then $\tau(N) = 9$. However, if $|S'_N| \geq 4$, then $\tau(N) \geq 10$ by (2.2). We conclude that form (S5) does not give a small recurrent number.
- (S6) If $N = p^3q^2$ for some $p^{3/2} < q < p^2$, then $S'_N = \{p, q, p^2, pq, p^3\}$ satisfies $U(p, q, 0, p)$.
- (S7) Let N have form (S7). Since the first four numbers of S'_N are $p < q < p^2 < r$ and $\tau(N) = 12$, we know that the fifth number in S'_N must be pq . That $p < q < p^2 < r < pq$ satisfies some $U(p, q, a, b)$ gives $a = \frac{p(pq-r)}{q^2-p^3}$, $b = \frac{rq-p^4}{q^2-p^3}$, and $r = pq - \sqrt{(q^2-p^3)(p^2-q)}$. We conclude that a number of form (S7) is small recurrent if and only if $p < q < p^2 < r < pq$, $(q^2-p^3)|(pq-r)$, $(q^2-p^3)|(rq-p^4)$, and $r = pq - \sqrt{(q^2-p^3)(p^2-q)}$. An example is $(p, q, r) = (2, 3, 5)$. We do not know if $(2, 3, 5)$ is the only set of primes that satisfy all these conditions or not.

From the above analysis, we obtain the proposition, which is a refinement of Proposition 3.7.

Proposition 3.8. *Let p, q, r denote prime numbers and k be some positive integer. A positive integer N is small recurrent with $|S'_N| \geq 4$ if and only if N belongs to one of the following forms.*

- (1) $N = p^k$ for some $k \geq 9$. In this case, $S'_N = \{p, p^2, p^3, \dots, p^{\lfloor (k-1)/2 \rfloor}\}$ satisfies $U(p, p^2, p, 0)$.
- (2) $N = p^kq$ for some $k \geq 4$ and $q > p^k$. In this case, $S'_N = \{p, p^2, p^3, \dots, p^k\}$ satisfies $U(p, p^2, p, 0)$.
- (3) $N = pq^k$ for some $k \geq 4$ and $p < q$. In this case, $S'_N = \{p, q, pq, q^2, \dots\}$ satisfies $U(p, q, 0, q)$.
- (4) $N = pq^k r$ for some $k \geq 2$, $p < q$, and $r > pq^k$. In this case, $S'_N = \{p, q, pq, q^2, \dots, pq^{k-1}, q^k, pq^k\}$ satisfies $U(p, q, 0, q)$.
- (5) $N = p^kq$ for some $k \geq 4$ and $\sqrt{q} < p < q$. In this case, $S'_N = \{p, q, p^2, pq, \dots\}$ satisfies $U(p, q, 0, p)$.
- (6) $N = p^3q^2$ for some $p^{3/2} < q < p^2$. In this case, $S'_N = \{p, q, p^2, pq, p^3\}$ satisfies $U(p, q, 0, p)$.

- (7) $N = p^2qr$, where $p < q < p^2 < r < pq$, $(q^2 - p^3)|(pq - r)$, $(q^2 - p^3)|(rq - p^4)$, and $r = pq - \sqrt{(q^2 - p^3)(p^2 - q)}$. In this case, $S'_N = \{p, q, p^2, r, pq\}$ satisfies $U\left(p, q, \frac{p(pq-r)}{q^2-p^3}, \frac{rq-p^4}{q^2-p^3}\right)$.

3.2. The case $|S'_N| \leq 3$. If $|S'_N| \leq 3$, then $\tau(N) \leq 9$. We use the classifications of those N from the introduction to obtain the following proposition.

Proposition 3.9. *Let p, q, r denote prime numbers and k be some positive integer. A positive integer $N > 1$ is small recurrent with $|S'_N| \leq 3$ if and only if N belongs to one of the following forms.*

- (1) $N = p^k$ for some $k \leq 8$. In this case, $S'_N = \{p, p^2, \dots, p^{\lfloor (k-1)/2 \rfloor}\}$ satisfies $U(p, p^2, p, 0)$.
- (2) $N = pq$ for some $p < q$. In this case, $S'_N = \{p\}$.
- (3) $N = pq^2$ for some $p < q$. In this case, $S'_N = \{p, q\}$.
- (4) $N = p^2q$ for some $p < q$. If $q < p^2$, then $S'_N = \{p, q\}$. If $q > p^2$, then $S'_N = \{p, p^2\}$.
- (5) $N = pq^3$ for some $p < q$. In this case, $S'_N = \{p, q, pq\}$.
- (6) $N = p^3q$ for some $p < q$. If $p < q < p^2$, then $S'_N = \{p, q, p^2\}$. If $p^3 < q$, then $S'_N = \{p, p^2, p^3\}$. (The case $p^2 < q < p^3$ is eliminated because the three elements in S'_N would be $p < p^2 < q$. However, there is no integral solution (a, b) to $q = ap^2 + bp$.)
- (7) $N = p^2q^2$ for some $p < q$. If $p < q < p^2$, then $S'_N = \{p, q, p^2\}$. (The case $p^2 < q$ is eliminated due to the same reason as in item (6).)
- (8) $N = pqr$ for some $p < q < r$. If $r < pq$ and there is an integral solution (a, b) to $r = aq + bp$, then $S'_N = \{p, q, r\}$. If $r > pq$, then $S'_N = \{p, q, pq\}$.

Combining Propositions 3.8 and 3.9, we obtain Theorem 1.2.

4. LARGE RECURRENT NUMBERS

Now we characterize all positive integers N whose L'_N satisfies a linear recurrence of order at most two. By a simple observation, instead of working directly with divisors in L'_N , we work with divisors in S'_N . Again, the set of divisors of a positive integer N is $1 = d_1 < d_2 < \dots < d_{\tau(N)}$ and the set $S'_N = \{d_2, d_3, \dots\}$.

4.1. The case $|L'_N| \geq 4$. Note that $|L'_N| \geq 4$ is equivalent to $|S'_N| \geq 4$.

Lemma 4.1. *For any $d \in L'_N$, we have $N/d \in S'_N$. If N is large recurrent with $|L'_N| \geq 4$, then*

$$ad_{i+2} + bd_{i+1} = \frac{d_{i+1}d_{i+2}}{d_i}, \forall d_i, d_{i+1}, d_{i+2} \in S'_N. \quad (4.1)$$

In particular, we have

$$ad_4 + bd_3 = \frac{d_3d_4}{d_2}. \quad (4.2)$$

Proof. If $d \in L'_N$, then $\sqrt{N} < d < N$. Then $1 < N/d < \sqrt{N}$, so $N/d \in S'_N$. Let

$$d'_i := d_{\tau(n)+1-i} = \frac{N}{d_i} \in L'_N, \forall d_i \in S'_N.$$

If N is large recurrent, then we have

$$d'_i = ad'_{i+1} + bd'_{i+2}, \forall d'_i, d'_{i+1}, d'_{i+2} \in L'_N.$$

Therefore,

$$\frac{N}{d_i} = a \frac{N}{d_{i+1}} + b \frac{N}{d_{i+2}}, \forall d_i, d_{i+1}, d_{i+2} \in S'_N,$$

which gives

$$ad_{i+2} + bd_{i+1} = \frac{d_{i+1}d_{i+2}}{d_i}, \forall d_i, d_{i+1}, d_{i+2} \in S'_N.$$

This completes our proof. \square

Since d_2 is a prime number p and d_3 is either p^2 or a prime number $q > p$, we consider two cases.

4.1.1. When $d_3 = p^2$. Then d_4 is either p^3 or a prime number $q > p^2$.

a) If $d_4 = p^3$, then (4.2) implies that $p^2 = ap + b$.

Claim 4.2. *If $p \neq a$, then $S'_N = \{p, p^2, \dots, p^k\}$ for some $k \geq 4$.*

Proof. We need to show that if $d_i \in S'_N$, then $d_i = p^{i-1}$. Base case: the claim holds for $i \leq 4$. Suppose that there exists a $j \geq 4$ such that $d_i = p^{i-1}$ for all $i \leq j$. Using (4.1), we have

$$ad_{j+1} + bp^{j-1} = ad_{j+1} + bd_j = \frac{d_{j+1}d_j}{d_{j-1}} = \frac{d_{j+1}p^{j-1}}{p^{j-2}} = pd_{j+1},$$

which, combined with $p^2 = ap + b$, gives

$$(p - a)(d_{j+1} - p^j) = 0.$$

Since $p \neq a$, we obtain $d_{j+1} = p^j$, as desired. \square

By Proposition 2.2, we know that when $p \neq a$, either $N = p^k$ or $N = p^k q$ for some $k \geq 1$ and some prime $q > p^k$.

Now suppose that $p = a$. Then $b = 0$. We can write elements in L'_N as $\{g_1, g_2, pg_2, p^2g_2, \dots, p^k g_2\}$ for some $k \geq 2$. Correspondingly, the set S'_N is $\{p, p^2, \dots, p^k, p^{k+1}, p^{k+1}g_2/g_1\}$. If $p^{k+1}g_2/g_1$ is a power of p , then we have the same conclusion about N as when $p \neq a$. If $p^{k+1}g_2/g_1$ is not a power of p , then

$$\frac{p^{k+1}g_2}{g_1} = q, \text{ for some prime } q > p \implies g_1 = p^{k+1} \frac{g_2}{q}.$$

Note that $g_2/q \in \mathbb{N}$. Furthermore, we claim that $g_2/q = p$. Indeed, since $1 < g_2/q < g_1$, we know that $g_2/q \in S'_N$. If $g_2/q = q$, then

$$pq < p^{k+1} \frac{g_2}{q} = g_1,$$

which implies that $pq \in S'_N$, a contradiction. If $g_2/q = p^j$ for some $j > 1$, then

$$p^{k+2} < p^{k+1+j} = p^{k+1} \frac{g_2}{q} = g_1,$$

which implies that $p^{k+2} \in S'_N$, another contradiction. Therefore, $g_2/q = p$, and we obtain $g_1 = p^{k+2}$ and $g_2 = pq$. We conclude that $\underline{N = p^{k+2}q}$ for some $k \geq 2$ and $p^{k+1} < q < p^{k+2}$.

- b) If $d_4 = q$, we claim that $a \neq p$. Suppose otherwise. Applying (4.2) to d_2, d_3 , and d_4 gives $aq + bp^2 = pq$. Hence, $a = p$ implies that $b = 0$. However, applying (4.1) to d_3, d_4 , and d_5 gives $(p^3 - q)d_5 = 0$, a contradiction. Therefore, $a \neq p$. By (4.2), we have

$$d_4 = \frac{bp^2}{p-a} = q \implies q|b \implies b = kq \text{ for some } k \in \mathbb{Z} \setminus \{0\}.$$

Hence, $a = p - kp^2$. By (4.1) applied to d_3, d_4 , and d_5 ,

$$d_5 = \frac{kp^2q^2}{q - p^3 + kp^4}, \quad (4.3)$$

which implies that $p^2|d_5$ since $\gcd(p^2, q - p^3 + kp^4) = 1$. Hence, $d_5 = p^3$, and (4.3) gives $k = \frac{p(p^3-q)}{p^5-q^2}$.

Case b.i) If S'_N has exactly four elements, which are p, p^2, q, p^3 , then $\tau(N) = 10$, which implies that $N = p^4q$. Hence, $L'_N = \{pq, p^4, p^2q, p^3q\}$ with $a = pq \frac{p^2-q}{p^5-q^2}$ and $b = pq \frac{p^3-q}{p^5-q^2}$. We conclude that $\underline{N = p^4q}$ with $p^2 < q < p^3$, $(p^5 - q^2)|(p^2 - q)$, and $(p^5 - q^2)|(p^3 - q)$.

Case b.ii) If $|S'_N| > 4$, then (4.1) gives

$$d_6 = \frac{bp^3q}{p^3 - aq} \implies q|d_6 \implies d_6 = pq.$$

However, since $(a, b) = (p(1 - kp), kq)$, we have

$$pq = d_6 = \frac{bp^3q}{p^3 - aq} = \frac{kp^2q^2}{p^2 - q(1 - kp)},$$

which gives $p^2 = q$, a contradiction.

We summarize our result when $d_3 = p^2$.

Proposition 4.3. *A number N is large recurrent with $|L'_N| \geq 4$ and $(d_2, d_3) = (p, p^2)$ for some prime p if and only if N belongs to one of the following forms.*

- (1) $N = p^k$ for some $k \geq 9$. In this case, $L'_N = \{p^{\lceil (k-1)/2 \rceil + 1}, p^{\lceil (k-1)/2 \rceil + 2}, \dots, p^{k-1}\}$ satisfies $U(p^{\lceil (k-1)/2 \rceil + 1}, p^{\lceil (k-1)/2 \rceil + 2}, p, 0)$.
- (2) $N = p^kq$ for some $k \geq 4$ and $q > p^k$. In this case, $L'_N = \{q, pq, p^2q, \dots, p^{k-1}q\}$ satisfies $U(q, pq, p, 0)$.
- (3) $N = p^kq$ for some $k \geq 4$ and $p^{k-1} < q < p^k$. Then

$$L'_N = \{p^k, pq, p^2q, \dots, p^{k-1}q\}$$

satisfies $U(p^k, pq, p, 0)$.

- (4) $N = p^4q$ with $p^2 < q < p^3$, $(p^5 - q^2)|(p^2 - q)$, and $(p^5 - q^2)|(p^3 - q)$. In this case, $L'_N = \{pq, p^4, p^2q, p^3q\}$.

4.1.2. When $d_3 = q$. By (4.1),

$$p(ad_4 + bq) = d_4q \implies p|d_4.$$

Write $d_4 = kp$ for some integer k . Since $d_2 = p$ and $d_3 = q$, d_4 must be either p^2 or pq .

a) If $d_4 = p^2$, (4.1) gives $ap^2 + bq = pq$. Hence, $q|a$ and $p|b$. Write $a = mq$ and $b = np$ for some integers m, n to get $mp + n = 1$. By (4.1), we see that

$$d_5 = \frac{bp^2q}{p^2 - aq} \implies q|d_5 \implies d_5 = pq.$$

Therefore,

$$\frac{bp^2q}{p^2 - aq} = \frac{np^3q}{p^2 - mq^2} = \frac{(1 - mp)p^3q}{p^2 - mq^2} = pq \implies m(p^3 - q^2) = 0,$$

which gives $m = 0$ and so, $(a, b) = (0, p)$. By (4.1), $d_{i+2} = pd_i$ for all $d_i, d_{i+2} \in S'_N$ and

$$S'_N = \{p, q, p^2, pq, \dots\}.$$

If $|S'_N| = 4$, then $\tau(N) = 10$ and $N = p^4q$ for $p < q < p^2$. Suppose that $|S'_N| \geq 5$, then $p^3 \in S'_N$.

Case a.i) If $q^2|N$, let $k \geq 2$ and $\ell \geq 3$ be the largest power such that $q^k|N$ and $p^\ell|N$, respectively. Since $q^2 \notin S'_N$, we know that

$$q^4 \geq N \geq p^3q^k > q^{k+3/2} \implies k < 5/2.$$

It follows that $k = 2$. That $q^2 < p^2q$ implies that

$$(p^2q)^2 > N \geq p^\ell q^2 \implies 3 \geq \ell \geq 3.$$

Hence, $\ell = 3$. If N does not have any other prime divisors besides p and q , then $N = p^3q^2$ for $p < q < p^2$. If N has a prime divisor $r \neq p, q$, then $r > \sqrt{N}$. So, r must be the unique prime divisor different from p and q . We have $N = p^3q^2r$ for $p < q < p^2$ and $r > p^3q^2$. Then $q^2 \in S'_N$, a contradiction.

Case a.ii) If $q^2 \nmid N$ and N has no prime divisors other than p and q , then $N = p^kq$ some for $k \geq 2$ and $p < q < p^2$.

Case a.iii) If $q^2 \nmid N$ and there exists a prime divisor r other than p or q , then $r > \sqrt{N}$ and r is the unique prime different from p and q . Therefore, $N = p^kqr$ for some $k \geq 2$ and $p < q < p^2 < p^kq < r$. Note that the two largest elements in S'_N are $p^{k-1}q$ and p^kq . Let d be the third largest divisor in S'_N . The relation $d_{i+2} = pd_i$ for all $d_i, d_{i+2} \in S'_N$ gives that $dp = p^kq$ and so, $d = p^{k-1}q$, which contradicts that $p^{k-1}q$ is the second largest in S'_N .

b) If $d_4 = pq$, then $p^2 \nmid N$ since $p^2 < pq$. By (4.2),

$$ap = q - b. \tag{4.4}$$

We see that d_5 is equal to q^2 or r , for some prime $r > pq$.

Case b.i) If $d_5 = q^2$, then (4.1) gives

$$bp = (p - a)q. \tag{4.5}$$

From (4.4) and (4.5), we obtain $a(p^2 - q) = 0$, so $(a, b) = (0, q)$. By (4.1), $d_{i+2} = qd_i$ for all $d_i, d_{i+2} \in S'_N$. Using [1, Proposition 5], we conclude that $N = pq^k$ or $N = pq^k r$ for some $k \geq 2$ and $p < q < pq^k < r$.

Case b.ii) If $d_5 = r$, then we claim that $|S'_N| > 4$. If not, $|S'_N| = 4$ implies that $\tau(N) = 10$, which contradicts that N has three distinct prime divisors. By (4.1), we see that

$$pq(ad_6 + br) = d_6 r,$$

so $pq|d_6$. So, $d_6 \in \{pq^2, pqr\}$. If $d_6 = pq^2$, then $q^2 < d_6$, but q^2 does not appear before d_6 in S'_N , a contradiction. If $d_6 = pqr$, then $pr < d_6$, but pr does not appear before d_6 in S'_N , again a contradiction.

Proposition 4.4. *A number N is large recurrent with $|L'_N| \geq 4$ and $(d_2, d_3) = (p, q)$ for some primes $p < q$ if and only if N belongs to one of the following forms.*

- (1) $N = p^3 q^2$ for $p < q < p^2$. In this case, $L'_N = \{q^2, p^2 q, pq^2, p^3 q, p^2 q^2\}$ satisfies $U(q^2, p^2 q, 0, p)$.
- (2) $N = p^k q$ some for $k \geq 4$ and $p < q < p^2$. In this case,

$$L'_N = \begin{cases} \{p^{k/2+1}, p^{k/2} q, p^{k/2+2}, \dots, p^{k-1} q\} & \text{if } 2|k, \\ \{p^{(k-1)/2} q, p^{(k+3)/2}, p^{(k+1)/2} q, \dots, p^{k-1} q\} & \text{if } 2 \nmid k. \end{cases}$$

Observe that L'_N satisfies $U(p^{k/2+1}, p^{k/2} q, 0, p)$ and $U(p^{(k-1)/2} q, p^{(k+3)/2}, 0, p)$ for even and odd k , respectively.

- (3) $N = pq^k$ for some $k \geq 4$ and $p < q$. In this case,

$$L'_N = \begin{cases} \{pq^{k/2}, q^{k/2+1}, \dots, q^k\} & \text{if } 2|k, \\ \{q^{k/2}, pq^{k/2}, \dots, q^k\} & \text{if } 2 \nmid k. \end{cases}$$

Observe that L'_N satisfies $U(pq^{k/2}, q^{k/2+1}, 0, q)$ and $U(q^{(k+1)/2}, pq^{(k+1)/2}, 0, q)$ for even and odd k , respectively.

- (4) $N = pq^k r$ for some $k \geq 2$ and $p < q < pq^k < r$. In this case, $L'_N = \{r, pr, qr, pqr, q^2 r, \dots, q^k r\}$ satisfies $U(r, pr, 0, q)$.

Combining Propositions 4.3 and 4.4, we obtain the following.

Proposition 4.5. *A number N is large recurrent with $|L'_N| \geq 4$ if and only if N belongs to one of the following forms.*

- (1) $N = p^k$ for some $k \geq 9$. In this case, $L'_N = \{p^{\lceil (k-1)/2 \rceil + 1}, p^{\lceil (k-1)/2 \rceil + 2}, \dots, p^{k-1}\}$ satisfies $U(p^{\lceil (k-1)/2 \rceil + 1}, p^{\lceil (k-1)/2 \rceil + 2}, p, 0)$.
- (2) $N = p^k q$ for some $k \geq 4$ and $q > p^k$. In this case, $L'_N = \{q, pq, p^2 q, \dots, p^{k-1} q\}$ satisfies $U(q, pq, p, 0)$.
- (3) $N = p^k q$ for some $k \geq 4$ and $p^{k-1} < q < p^k$. Then

$$L'_N = \{p^k, pq, p^2 q, \dots, p^{k-1} q\}$$

satisfies $U(p^k, pq, p, 0)$.

- (4) $N = p^k q$ some for $k \geq 4$ and $p < q < p^2$. In this case,

$$L'_N = \begin{cases} \{p^{k/2+1}, p^{k/2} q, p^{k/2+2}, \dots, p^{k-1} q\} & \text{if } 2|k, \\ \{p^{(k-1)/2} q, p^{(k+3)/2}, p^{(k+1)/2} q, \dots, p^{k-1} q\} & \text{if } 2 \nmid k. \end{cases}$$

Observe that L'_N satisfies $U(p^{k/2+1}, p^{k/2}q, 0, p)$ and $U(p^{(k-1)/2}q, p^{(k+3)/2}, 0, p)$ for even and odd k , respectively.

- (5) $N = p^4q$ with $p^2 < q < p^3$, $(p^5 - q^2)|(p^2 - q)$, and $(p^5 - q^2)|(p^3 - q)$. In this case, $L'_N = \{pq, p^4, p^2q, p^3q\}$.
- (6) $N = p^3q^2$ for $p < q < p^2$. In this case, $L'_N = \{q^2, p^2q, pq^2, p^3q, p^2q^2\}$ satisfies $U(q^2, p^2q, 0, p)$.
- (7) $N = pq^k$ for some $k \geq 4$ and $p < q$. In this case,

$$L'_N = \begin{cases} \{pq^{\frac{k}{2}}, q^{\frac{k}{2}+1}, \dots, q^k\} & \text{if } 2|k, \\ \{q^{\frac{k+1}{2}}, pq^{\frac{k+1}{2}}, \dots, q^k\} & \text{if } 2 \nmid k. \end{cases}$$

Observe that L'_N satisfies $U(pq^{k/2}, q^{k/2+1}, 0, q)$ and $U(q^{(k+1)/2}, pq^{(k+1)/2}, 0, q)$ for even and odd k , respectively.

- (8) $N = pq^k r$ for some $k \geq 2$ and $p < q < pq^k < r$. In this case, $L'_N = \{r, pr, qr, pqr, q^2r, \dots, q^k r\}$ satisfies $U(r, pr, 0, q)$.

4.2. The case $|L'_N| \leq 3$. If $|L'_N| \leq 3$, then $\tau(N) \leq 9$. We use the classifications of those N from the introduction to obtain the following proposition

Proposition 4.6. *Let p, q, r denote prime numbers and k be some positive integer. A positive integer $N > 1$ is small recurrent with $|L'_N| \leq 3$ if and only if N belongs to one of the following forms.*

- (1) $N = p^k$ for some $k \leq 8$. In this case, $L'_N = \{p^{\lceil (k-1)/2 \rceil + 1}, \dots, p^{k-1}\}$ satisfies $U(p^{\lceil (k-1)/2 \rceil + 1}, p^{\lceil (k-1)/2 \rceil + 2}, p, 0)$.
- (2) $N = pq$ for some $p < q$. In this case, $L'_N = \{q\}$.
- (3) $N = pq^2$ for some $p < q$. In this case, $L'_N = \{pq, q^2\}$.
- (4) $N = p^2q$ for some $p < q$. If $q < p^2$, then $L'_N = \{p^2, pq\}$. If $q > p^2$, then $L'_N = \{q, pq\}$.
- (5) $N = pq^3$ for some $p < q$. In this case, $L'_N = \{q^2, pq^2, q^3\}$.
- (6) $N = p^3q$ for some $p < q$. If $p < q < p^2$, then $L'_N = \{pq, p^3, p^2q\}$. If $p^3 < q$, then $L'_N = \{q, pq, p^2q\}$. If $p^2 < q < p^3$, then $L'_N = \{p^3, pq, p^2q\}$.
- (7) $N = p^2q^2$ for some $p < q$. If $p < q < p^2$, then $L'_N = \{q^2, p^2q, pq^2\}$. The case $p^2 < q$ is impossible as it gives $L'_N = \{p^2q, pq^2, q^2\}$ and there is no integral solution (a, b) to $apq^2 + bp^2q = q^2$.
- (8) $N = pqr$ for some $p < q < r$. If $r > pq$, then $L'_N = \{r, pr, qr\}$. The case $r < pq$ is impossible as it gives $L'_N = \{pq, pr, qr\}$ and there is no integral solution (a, b) to $apr + bpq = qr$.

Combining Propositions 4.5 and 4.6, we obtain Theorem 1.4.

5. APPENDIX

Proof of Lemma 2.5. i) Since $r = aq + bp$ and $pq = ar + bq$, we know that $\gcd(a, b)|r$ and $\gcd(a, b)|pq$, respectively. Hence, $\gcd(a, b) = 1$.

Since $r = aq + bp$ and r is a prime, $p \nmid a$.

ii) Suppose that $k = \gcd(b, d_i) > 1$ for some $d_i \in S'_N$. If $d_i = d_2$, then $p|b$. Since $pq = ar + bq$, we get $p|a$, which contradicts $\gcd(a, b) = 1$. If $d_i = d_3$, then $q|b$. Since

$r = aq + bp$, we get $q|r$, a contradiction. If $d_i > d_3$, then write

$$\gcd(b, d_i) = \gcd(b, ad_{i-1} + bd_{i-2}) \stackrel{\text{i)}}{=} \gcd(b, d_{i-1}),$$

which, by induction, gives $1 < \gcd(b, d_i) = \gcd(b, d_3)$, which has been shown to be impossible.

iii) The claim holds for $i \leq 4$. Let $d_i, d_{i+1} \in S'_N$ for some $i \geq 5$. We have

$$\gcd(d_i, d_{i+1}) = \gcd(d_i, ad_i + bd_{i-1}) \stackrel{\text{ii)}}{=} \gcd(d_i, d_{i-1}).$$

By induction, we obtain $\gcd(d_i, d_{i+1}) = 1$.

iv) The claim holds for $i \leq 5$. Assume that it holds for all $i \leq j$ for some $j \geq 5$. We show that it holds for $i = j + 1$. We have

$$pq = ar + bq = a(aq + bp) + bq = (a^2 + b)q + abp.$$

Hence, $p|(a^2 + b)$. Write

$$d_{j+1} = ad_j + bd_{j-1} = a(ad_{j-1} + bd_{j-2}) + bd_{j-1} = (a^2 + b)d_{j-1} + abd_{j-2}.$$

Since $p|(a^2 + b)$ and $\gcd(p, ab) = 1$, we know that $p|d_{j+1}$ if and only if $p|d_{j-1}$. By the inductive hypothesis, $p|d_{j-1}$ if and only if $j - 2 \equiv 2 \pmod{3}$, or equivalently, $j + 1 \equiv 2 \pmod{3}$. By induction, we have the desired conclusion.

v) The claim holds for $i \leq 5$. Assume that it holds for all $i \leq j$ for some $j \geq 5$. We show that it holds for $i = j + 1$. That $pq = ar + bq$ implies that $q|a$. Write

$$d_{j+1} = ad_j + bd_{j-1}.$$

By ii), $q|d_{j+1}$ if and only if $q|d_{j-1}$. By the inductive hypothesis, $q|d_{j-1}$ if and only if $j + 1 \equiv 1 \pmod{2}$. This completes our proof. \square

Proof of Lemma 2.6. i) Same as the proof of Lemma 2.5 item i).

ii) Suppose, for a contradiction, that $\gcd(b, d_i) > 1$ for some $i \geq 3$. If $i = 3$, then $q|b$. We have $r = aq + bp$. Since $q|b$, we get $q|r$, a contradiction. If $i \geq 4$, write

$$\gcd(b, d_i) = \gcd(b, ad_{i-1} + bd_{i-2}) = \gcd(b, d_{i-1}).$$

By induction, $1 < \gcd(b, d_i) = \gcd(b, d_3)$, which has been shown to be impossible.

iii) The claim holds for $i \leq 4$. Pick $i \geq 5$. We have

$$\gcd(d_i, d_{i+1}) = \gcd(d_i, ad_i + bd_{i-1}) = \gcd(d_i, bd_{i-1}) \stackrel{\text{ii)}}{=} \gcd(d_i, d_{i-1}).$$

By induction, we obtain $\gcd(d_i, d_{i+1}) = \gcd(d_4, d_5) = 1$.

iv) Assume that $\gcd(a, d_i) > 1$ for some $i \geq 2$. If $i = 2$, then $p|a$, which contradicts the primality of r and the linear recurrence $r = aq + bp$. If $i = 3$, then $q|b$, which contradicts the primality of s and the linear recurrence $s = ar + bq$. Assume that $i \geq 4$. Write

$$\gcd(a, d_i) = \gcd(a, ad_{i-1} + bd_{i-2}) \stackrel{\text{i)}}{=} \gcd(a, d_{i-2}).$$

By induction, either $1 < \gcd(a, d_i) = \gcd(a, d_2)$ or $1 < \gcd(a, d_i) = \gcd(a, d_3)$, neither of which is possible.

v) The claims holds for $i \leq 3$. Pick $i \geq 4$ and suppose that $k = \gcd(d_i, d_{i+2}) > 1$. Since $d_{i+2} = ad_{i+1} + bd_i$, k divides ad_{i+1} . By iii), $\gcd(k, d_{i+1}) = 1$, so $k|a$. However, $\gcd(a, d_i) > 1$ contradicts iv). \square

REFERENCES

- [1] A. A. Chentouf, Linear recurrences of order at most two in small divisors, *J. Integer Seq.* **25** (2022).
- [2] H. V. Chu, When the large divisors of a natural number are in arithmetic progression, *J. Integer Seq.* **23** (2020).
- [3] H. V. Chu, When the nontrivial, small divisors of a natural number are in arithmetic progression, *Quaest. Math.* **45** (2022), 969–977.
- [4] D. E. Iannucci, When the small divisors of a natural number are in arithmetic progression, *Integers* **18** (2018).
- [5] S. Ramanujan, Collected Papers of Srinivasa Ramanujan, *American Mathematical Soc* (2000), 262–275.
- [6] E. Landau, Handbuch der Lehre von der Verteilung der Primzahlen, *B. G. Teubner* (1909).

Email address: hungchu2@illinois.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL 61820, USA

Email address: kevinhle@tamu.edu

MATHEMATICS, TEXAS A&M UNIVERSITY COLLEGE STATION, TX 77843, USA

Email address: sjml@williams.edu, Steven.Miller.MC.96@aya.yale.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, WILLIAMS COLLEGE, WILLIAMSTOWN, MA 01267, USA

Email address: yql@williams.edu

WILLIAMS COLLEGE, WILLIAMSTOWN, MA 01267, USA

Email address: Liyang.Shen@nyu.edu

COURANT INSTITUTE OF MATHEMATICAL SCIENCES, NEW YORK UNIVERSITY, NEW YORK, NY 10012, USA