

TWISTED CONJUGACY IN RESIDUALLY FINITE GROUPS OF FINITE PRÜFER RANK

EVGENIJ TROITSKY

ABSTRACT. Suppose, G is a residually finite group of finite upper rank admitting an automorphism φ with finite Reidemeister number $R(\varphi)$ (the number of φ -twisted conjugacy classes). We prove that such G is soluble-by-finite (in other words, any residually finite group of finite upper rank, which is not soluble-by-finite, has the R_∞ property).

This reduction is the first step in the proof of the second main theorem of the paper: suppose, G is a residually finite group of finite Prüfer rank and φ is its automorphism with $R(\varphi) < \infty$; then $R(\varphi)$ is equal to the number of equivalence classes of finite-dimensional irreducible unitary representations of G , which are fixed points of the dual map $\widehat{\varphi} : [\rho] \mapsto [\rho \circ \varphi]$ (i.e., we prove the TBFT $_f$, the finite version of the conjecture about the twisted Burnside-Frobenius theorem, for such groups).

INTRODUCTION

If $d(H)$ denotes the minimal number of generators of a group H , then the (*Prüfer*) *rank* of a group G is

$$\text{rk}(G) := \sup\{d(H) : H \text{ is a finitely generated subgroup of } G\}$$

and the *upper rank* of G is

$$\text{urk}(G) := \sup\{\text{rk}(\overline{G}) : \overline{G} \text{ is a finite quotient of } G\}.$$

Subgroups of finite rank attracted much interest since the end of 80's, in particular, in connection with the subgroup growth and related questions [17].

The main results of the present paper are as follows.

Theorem A. *Suppose that G is a residually finite group of finite upper rank and $\varphi : G \rightarrow G$ is an automorphism with $R(\varphi) < \infty$. Then G is a soluble-by-finite group.*

A reformulation is: any residually finite group of finite upper rank, which is not soluble-by-finite, has the R_∞ *property*.

Theorem B. *Suppose, G is a residually finite group of finite rank. Then TBFT $_f$ is true for G .*

Theorem A may be considered as a development of [13] and at the same time as a supporting result for the following conjecture [6]: a residually finite finitely generated group with an automorphism with finite Reidemeister number is soluble-by-finite. The proof of Theorem A uses in particular a result of A. Jaikin-Zapirain [14].

2000 *Mathematics Subject Classification.* 20C; 20E45; 22D10; 37C25;

Key words and phrases. Reidemeister number, twisted conjugacy class, Burnside-Frobenius theorem, group of finite rank, residually finite group, soluble group, unitary dual, finite-dimensional representation .

The work is supported by the Russian Science Foundation under grant 21-11-00080.

The initial conjecture about the TBFT was formulated by A. Fel'shtyn and R. Hill as follows: if the Reidemeister number is finite then it coincides with the number of equivalence classes of irreducible unitary representations of G , which are fixed points of the dual map $[\rho] \mapsto [\rho \circ \varphi]$ (see [2]). A counterexample to TBFT from [8] caused a modified version, namely TBFT_f , where only finite-dimensional representations are taken into account. The widest class of groups with TBFT_f , namely polycyclic-by-finite, was detected in [5]. Then we have found several counterexamples, in particular, in [9] an example of a group, which has neither TBFT_f , nor TBFT, was found. Also, TBFT_f was proved for a class of wreath products [21, 22].

We prove in Theorem B the TBFT_f for a wide class which is considered by many authors as a natural extension of the class of polycyclic-by-finite groups.

The interest in Reidemeister theory, R_∞ , and TBFT comes from topological dynamics and other fields (see an overview in [2, 5]). For some overview of results related to R_∞ and recent results one can see [6, 11, 19] (these are several papers from a long publications list).

Finite rank groups (nilpotent torsion-free) in the context of Reidemeister theory were studied in [3, 7]. Here we study a much wider class.

The paper is organized as follows. In Section 1 we recall and prove some necessary statements. Theorem A is proved in Section 2. In Section 3 we prove statements for abelian quotients to be used in the next section. Theorem B is proved in Section 4.

1. PRELIMINARIES AND REMINDING

Suppose, that H is a normal subgroup of G and $p : G \rightarrow G/H$ is the natural projection. Let H be invariant for an endomorphism $\varphi : G \rightarrow G$. Denote by $\varphi' : H \rightarrow H$ and by $\bar{\varphi} : G/H \rightarrow G/H$ the induced homomorphisms. Evidently, p induces a surjection of Reidemeister classes and

$$(1) \quad R(\bar{\varphi}) \leq R(\varphi).$$

Definition 1.1. Denote by $C(\varphi)$ the subgroup of G , formed by elements fixed by $\varphi : G \rightarrow G$: $C(\varphi) := \{g \in G : \varphi(g) = g\}$.

The following theorem appeared in [10], see also [5, 12].

Theorem 1.2. *For the above defined G , H , φ , φ' , and $\bar{\varphi}$, we have the following properties.*

- a) *If $|C(\bar{\varphi})| = n$, then $R(\varphi') \leq R(\varphi) \cdot n$;*
- b) *If $C(\bar{\varphi}) = \{e\}$, then each Reidemeister class of φ' is an intersection of the appropriate Reidemeister class of φ and H ;*

Considering the equality

$$yg\varphi(y^{-1})x = ygx x^{-1}\varphi(y^{-1})x = y(gx)(\tau_{x^{-1}} \circ \varphi)(y^{-1}),$$

where $\tau_x(z) := xzx^{-1}$, one can deduce the following very useful statement (see e.g. [5]).

Lemma 1.3. *The shifts of Reidemeister classes of φ are Reidemeister classes of $\tau_{x^{-1}} \circ \varphi$:*

$$\{g\}_{\varphi}x = \{gx\}_{\tau_{x^{-1}} \circ \varphi}.$$

In particular, $R(\tau_g \circ \varphi) = R(\varphi)$.

Theorem 1.4 ([14]). *Let G be a finite group of rank r admitting an automorphism with m fixed elements. Then G has a characteristic soluble subgroup H , whose index is (m, r) -bounded and whose derived length is r -bounded.*

Definition 1.5. Denote the bounding functions for index and derived length from Theorem 1.4 by $I(m, r)$ and $L(r)$ respectively.

The following statement is well known (see [18, Lemma 3.5]). We sketch here a more simple proof with a more weak estimation.

Theorem 1.6. *The number $a_n(G)$ of subgroups of index n in a finite upper rank group G is finite. More specifically $a_n(G) \leq (n!)^r$, where $r = \text{urk}(G)$.*

Proof. Suppose that G has s distinct subgroups of index n : H_1, \dots, H_s (we do not suppose that the list is exhaustive). Consider $H = \bigcap_i H_i$ and the natural left action of G on the finite set of all left cosets by H . Let H_0 be the kernel of this action, which is a normal subgroup. Since $H_0 \subseteq H$, the natural projection $p : G \rightarrow G/H_0 =: F$ sends H_i to distinct subgroups of index n of the finite group F . Thus, $s \leq a_n(F)$. For a finite (or more generally, finitely generated) group F the following estimate is well known from M. Hall theorem (see e.g. [16, Vol. 1, p. 57]): $a_n(F) \leq (n!)^d$, where d is the number of elements in some generating system of F . Since $d \leq \text{urk}(F) \leq \text{urk}(G)$, we are done. \square

Of course, this is a very general rough estimation. For more sharp ones in specific cases see e.g. [17].

In [13] E. Jabara has proved

Lemma 1.7. *Suppose that $\varphi : F \rightarrow F$ is an automorphism of a finite group G . Then $|C(\varphi)| \leq \alpha(R(\varphi))$, where $\alpha(r) = 2^{2^r}$.*

Using this estimation, it was proved in [13] that the same holds for an automorphism of finite order of a residually finite group, and in [4] for an automorphism of a finitely generated residually finite group. In the same way we prove here the following statement.

Lemma 1.8. *Suppose that $\varphi : G \rightarrow G$ is an automorphism of a residually finite group of finite upper rank and $R := R(\varphi) < \infty$. Then $|C(\varphi)| \leq \alpha(R) < \infty$.*

Proof. Let $\{f_1, \dots, f_n\} \subset C(\varphi)$ be some finite set of φ -fixed elements. Let $H \subset G$ be a normal subgroup of finite index such that, for $p : G \rightarrow G/H$, the elements $p(f_i)$ are pairwise distinct. By Theorem 1.6, we can take the intersection of all subgroups of the same index as H to obtain a characteristic (in particular, normal φ -invariant) subgroup of finite index $H' \subseteq G$. For $p' : G \rightarrow G/H'$, we still have that $p'(f_i)$ are pairwise distinct. Also, $\{p'(f_1), \dots, p'(f_n)\} \subset C(\bar{\varphi})$, where $\bar{\varphi} : G/H' \rightarrow G/H'$ is the induced automorphism of a finite group. Then, by Lemma 1.7 we have

$$n \leq |C(\bar{\varphi})| \leq \alpha(R(\bar{\varphi})) \leq \alpha(R).$$

Hence, $|C(\varphi)| \leq \alpha(R)$. \square

We will work with the following reformulation of the TBFT $_f$. With some additional restriction it was proved in [6].

Lemma 1.9. *Suppose that φ is an automorphism of G with $R(\varphi) < \infty$. Then TBFT $_f$ is true for φ if and only if there is a finite group F with an automorphism φ_F and an equivariant epimorphism $G \rightarrow F$ inducing a bijection of Reidemeister classes of φ and φ_F .*

Proof. Suppose that such an epimorphism p exists. Denote $R = R(\varphi)$. Then there are $\widehat{\varphi_F}$ -fixed irreducible unitary pairwise non-equivalent representations ρ_1, \dots, ρ_R of F . Hence,

$\rho_1 \circ p, \dots, \rho_R \circ p$ are $\widehat{\varphi}$ -fixed irreducible unitary pairwise non-equivalent representations of G . Thus R does not exceed the number of $\widehat{\varphi}$ -fixed representation classes.

Each finite-dimensional $\widehat{\varphi}$ -fixed class of representation ρ has a matrix coefficient $g \mapsto \text{Trace}(J \circ \rho(g))$, where J is an intertwining operator between ρ and $\rho \circ \varphi$. This coefficient is a φ -class function (i.e. it is constant on each Reidemeister class) and only coefficients of this form have this property (see [5] for details of verification). On the other hand, matrix elements of non-equivalent finite-dimensional representations are linear independent (see [1, Corollary (27.13)]). Thus the number of fixed representation classes can not exceed R and the ‘if’ direction is proved.

Conversely, suppose that TBFT_f is true and ρ_1, \dots, ρ_R are $\widehat{\varphi}$ -fixed irreducible unitary pairwise non-equivalent representations of G . Then these representations are finite, i.e. $|\rho_i(G)| < \infty$, $i = 1, \dots, R$, because otherwise the corresponding φ -class function has infinitely many values. Indeed, for $\rho = \rho_1, \dots, \rho_R$, $f_J(g) = \text{Trace}(J \circ \rho(g))$ has to take finitely many values because $R < \infty$. Since f_J is a non-trivial matrix coefficient, its left translations $L_g f_J$ generate a finite-dimensional representation π , which is equivalent to a direct sum of several copies of ρ . This follows from the fact, that each matrix coefficient is of the form $\varphi_A(g) = \text{Trace}(A \circ \rho(g))$ and one obtains an equivariant isomorphism $\varphi_A \mapsto A$, $A \in \text{End}(W_\rho) = W_\rho \otimes W_\rho^*$ between the space of all matrix coefficients and $W_\rho \otimes W_\rho^*$, where W_ρ is the space of the representation ρ and the action on W_ρ^* is trivial. (This can be extracted e.g. from the proof of [15, Theorem 3, p. 124].)

The space S of the representation π has a basis $L_{g_1} f_J, \dots, L_{g_k} f_J$, for some finite collection $\{g_1, \dots, g_k\}$. Then the level sets of any function from S have the form $\cap_i g_i U_j$, where U_j are the level sets of f_J . In particular, it has finitely many values. Taking some unions of these sets (if necessary) we can form a finite partition $G = V_1 \sqcup \dots \sqcup V_m$ such that elements of S are constant on V_i , $j = 1, \dots, m$, and separate them. Thus any left translation permutes V_i and the representation G on S factorizes through (a subgroup of) the permutation group on m elements, i.e. a finite group. Then the subrepresentation ρ is finite as well. It remains to take $H = \cap_{i=1}^R \text{Ker}(\rho_i)$. Then $p : G \rightarrow G/H$ is the desired equivariant epimorphism onto a finite group, where on G/H we consider the induced automorphism $\overline{\varphi}$. Indeed, $G/H \subseteq \rho_1(G) \oplus \dots \oplus \rho_R(G)$. Hence it is finite. By construction, the map is equivariant. It remains to prove that $R(\overline{\varphi}) = R$. For this purpose we will find R distinct $\widehat{\varphi}$ -fixed classes of representations. These classes are $[\rho_i \circ p]$. Indeed, since ρ_i are pairwise non-equivalent and p is an epimorphism, then so are $\rho_i \circ p$. Since

$$\widehat{\varphi}[\rho_i \circ p] = [\rho_i \circ p \circ \varphi] = [\rho_i \circ \overline{\varphi} \circ p] = [\rho_i \circ p],$$

the classes are $\widehat{\varphi}$ -fixed and we are done. □

We will need the following statement.

Lemma 1.10. *Suppose that G is a finite-by-abelian group, where F is a finite normal subgroup and the abelian factor group $A = G/F$ is of finite Prüfer rank $\text{rk}(A) < \infty$. Then the number of distinct inner automorphisms of G does not exceed $|F|! \cdot |F|^{\text{rk}(A)}$.*

Proof. Let $A_0 \subseteq A$ be a subgroup with generators a_1, \dots, a_k , $k \leq \text{rk}(A)$. Then $G_0 = p^{-1}(A_0)$ has a generating set $F \cup \{g_1, \dots, g_k\}$ of cardinality $|F| + k$, where p is the projection $p : G \rightarrow A$ and $g_i \in G_0$ are arbitrary elements such that $p(g_i) = a_i$, $i = 1, \dots, k$. Then any inner automorphism τ_g of G_0 is completely defined by the images of these generators. Since A is abelian, τ_g maps each F -coset to itself. Hence, we obtain not more than $|F|! \cdot |F|^k$ possibilities.

Now suppose that for the entire G there is $s > |F|! \cdot |F|^{\text{rk}(A)}$ distinct inner automorphisms τ_{y_i} , $y_i \in G$, $i = 1, \dots, s$. Then there are elements $x_{ij} \in G$ such that $\tau_{y_i}(x_{ij}) \neq \tau_{y_j}(x_{ij})$. Consider $A_0 \subseteq A$ generated by all $p(y_i)$ and $p(x_{ij})$ and take $G_0 := p^{-1}A_0$. Then τ_{y_i} remain inner automorphisms of G_0 (because $y_i \in G_0$) and remain distinct. This contradicts the first part of the proof. \square

Corollary 1.11. *Suppose, the number of distinct inner automorphisms of G is finite. Then the TBFT $_f$ is true for G . In particular, under suppositions of Lemma 1.10 the TBFT $_f$ is true for G .*

Proof. Let $\varphi : G \rightarrow G$ be an automorphism with a finite Reidemeister number. Consider shifts of Reidemeister classes. By Lemma 1.3, there is only finitely many such sets, namely $n \leq R(\varphi) \cdot t$, where t is the number of distinct inner automorphisms. Thus there is a natural homomorphism $\sigma : G \rightarrow S_n$, where S_n is the permutation group. Denote by R_n its image. The kernel of σ is the intersection of shift stabilizers of Reidemeister classes of φ . Each of these stabilizers is φ -invariant, because, if z belongs to the stabilizer $\text{St}(\{x\}_\varphi)$ of $\{x\}_\varphi$ then

$$\begin{aligned} gx\varphi(g^{-1})\varphi(z) &= gx\varphi(x)\varphi(x^{-1}g^{-1})\varphi(z) = \varphi[\varphi^{-1}(gx)x\varphi(\varphi^{-1}(x^{-1}g^{-1}))z] \\ &= \varphi(hx\varphi(h^{-1})) = (\varphi(h)x^{-1})x\varphi(x\varphi(h^{-1})), \end{aligned}$$

for some h . So $\text{Ker } \sigma$ is φ -invariant and we can consider the induced automorphism $\varphi_R : R_n \rightarrow R_n$. We claim that $\sigma : G \rightarrow R_n$ induces a bijection of Reidemeister classes. To prove this, we need to verify that any Reidemeister class of φ is a union of cosets of $\text{Ker } \sigma$. Consider an arbitrary element $x \in \{x\}_\varphi$ and $z \in \text{St}(\{x\}_\varphi)$. Then $xz = gx\varphi(g^{-1})$, for some g . Hence, $x \in x \text{Ker } \sigma \subseteq x \text{St}(\{x\}_\varphi) \subseteq \{x\}_\varphi$. Thus, each element is included in its Reidemeister class together with some coset of $\text{Ker } \sigma$ and we are done. \square

2. PROOF OF THEOREM A

Proof of Theorem A. Our argument will be close to some parts of [13]. For a residually finite group G consider an exhausting family of subgroups of finite index H_λ , $\lambda \in \Lambda$. By Theorem 1.6, taking the intersection of all subgroups of a fixed index, we may assume H_λ to be characteristic, in particular normal and φ -invariant. Denote $F_\lambda := G/H_\lambda$ and by $\varphi_\lambda : F_\lambda \rightarrow F_\lambda$ the induced automorphism. Then $R(\varphi_\lambda) \leq R(\varphi) =: R < \infty$. Hence by Lemma 1.8 we have $|C(\varphi_\lambda)| \leq \alpha(R)$. Of course, $\text{rk}(F_\lambda) = \text{urk}(F_\lambda) \leq \text{urk}(G) = r$. Applying Theorem 1.4 we can find, for each $\lambda \in \Lambda$, a soluble characteristic subgroup $F_\lambda^0 \subseteq F_\lambda$ of length $\leq L(r) =: L$ and index $\leq I(\alpha(R), r) =: J$. For each λ , denote by $p_\lambda : G \rightarrow F_\lambda$ the natural projection and by H_λ^0 the pre-image $H_\lambda^0 = (p_\lambda)^{-1}(F_\lambda^0)$. Let $H := \bigcap_\lambda H_\lambda^0$. Then each H_λ^0 is a characteristic subgroup of index $\leq J$. Then H is characteristic and of finite index by Theorem 1.6.

It remains to prove that H is soluble. Suppose the opposite: H is not a soluble group of derived length $\leq L$. Thus there exists an expression $\omega(h_1, \dots, h_k) \neq e$ in products and commutators, which has the commutator length $L + 1$. Choose λ such that

$$(2) \quad p_\lambda(\omega(h_1, \dots, h_k)) \neq e \in F_\lambda.$$

Since $p_\lambda(\omega(h_1, \dots, h_k)) = \omega(p_\lambda(h_1), \dots, p_\lambda(h_k))$, $h_i \in H \subseteq H_\lambda$ and $p_\lambda(h_i) \in F_\lambda^0$, the inequality (2) contradicts the property of F_λ^0 to be soluble of length $\leq L$. Thus H is a characteristic soluble subgroup of finite index in G . \square

Remark 2.1. Any *finitely generated* residually finite group of finite upper rank is soluble-by-finite by Mann-Segal [18].

3. ABELIAN FINITE RANK GROUPS AND THEIR AUTOMORPHISMS

We need a separate (from 1.8) study of abelian quotients, which may be not residually finite. We start from the torsion-free case.

Lemma 3.1. (cf. [7]) *An automorphism φ of any torsion free abelian group A of finite rank with finite Reidemeister number has $C(\varphi) = \{e\}$.*

Proof. By Fuchs' lemma (see [20, 15.2.3]), an endomorphism ψ of a torsion-free abelian group A of finite rank is injective if and only if the index of $\text{Im}(\psi)$ is finite. Since $R(\varphi) = |\text{Coker}(1 - \varphi)| < \infty$, we can apply this lemma to $\psi = 1 - \varphi$ and obtain $\{e\} = \text{Ker}(1 - \varphi) = C(\varphi)$. \square

Now suppose that $\varphi : A \rightarrow A$ is an automorphism of a general finite rank abelian group with $R(\varphi) < \infty$. Let T be the torsion subgroup, hence $A_0 := A/T$ is torsion-free. The subgroup T is fully invariant. Denote by $\varphi_T : T \rightarrow T$ and $\varphi_0 : A_0 \rightarrow A_0$ the automorphisms induced by φ .

By Lemma 3.1, $C(\varphi_0) = \{e\}$. Hence, by Theorem 1.2, $R(\varphi_T) < \infty$. Also,

$$(3) \quad C(\varphi) = C(\varphi_T).$$

For p prime, denote by $\mathbb{Z}(p^\infty)$ the *quasicyclic* (or *Prüfer*) *group of type p^∞* . An abelian finite rank torsion group is a direct sum of finitely many quasicyclic and cyclic groups (see e.g. [20, 4.1.1 + 4.3.13]):

$$(4) \quad T = T_1 \oplus \cdots \oplus T_m \oplus F, \quad T_i := \mathbb{Z}(p_i^\infty)^{d_i}, \quad p_i \neq p_j \text{ if } i \neq j, \quad |F| < \infty.$$

We will need the following simple observation.

Lemma 3.2. *Suppose that $G = \mathbb{Z}(p^\infty)^d$ (finite direct sum) and H is a subgroup of G . Then either H is finite or $G/H \cong \mathbb{Z}(p^\infty)^{d'}$ with $d' < d$.*

Proof. The subgroup H is a finite rank p -group, hence $H = H' \oplus F$, where $H' \cong \mathbb{Z}(p^\infty)^{m'}$ and F is a finite sum of cyclic groups. The subgroup H is infinite if and only if $m' \geq 1$. Since H' is a divisible subgroup, it has a complement H'' , so $G = H' \oplus H''$. Then $H'' \cong G/H'$ is a divisible p -group. Hence, $H'' \cong \mathbb{Z}(p^\infty)^{m''}$. Similarly, $G/H \cong \mathbb{Z}(p^\infty)^{d'}$. Since the number of $\mathbb{Z}(p^\infty)$ -summands is an invariant, $d = m' + m''$. Since H'' has a subgroup F' , isomorphic F , and $H''/F' \cong G/H$, then $d' \leq m'' = d - m'$. Hence, if H is infinite, then $d' < d$. \square

Lemma 3.3. *Suppose that $G = \mathbb{Z}(p^\infty)^d$ and $\varphi : G \rightarrow G$ is an endomorphism with $R(\varphi) < \infty$. Then $|C(\varphi)| < \infty$.*

Proof. Consider $C(\varphi) = \text{Ker}(1 - \varphi)$, which is a φ -invariant subgroup. If $|C(\varphi)| = \infty$, consider $\varphi_1 : G_1 \rightarrow G_1$, where $G_1 = G/C(\varphi)$. If $|C(\varphi_1)| = \infty$ then continue the process. By Lemma 3.2 the process will stop in several steps. At that moment the following cases can occur:

1) $C(\varphi_i) = G_i$ and the next factor group will be trivial. Then φ_i is the identity map of an infinite abelian group. Hence, by Theorem 1.2 $\infty = R(\varphi_i) \leq R(\varphi) < \infty$. A contradiction.

2) $|C(\varphi_i)| < \infty$, but $|C(\varphi_{i-1})| = \infty$ (otherwise we stop at the previous step). (Here we denote $G_0 = G$ and $\varphi_0 = \varphi$.) By Theorem 1.2 $R(\varphi_{i-1}) < \infty$. Then by Theorem 1.2 $R(\varphi'_{i-1}) < \infty$, where $\varphi'_{i-1} = \varphi_{i-1}|_{C(\varphi_{i-1})}$. But φ'_{i-1} is the identity map of an infinite abelian group and $R(\varphi'_{i-1}) = \infty$. That is why there were no previous steps, i.e. $i = 0$, and $|C(\varphi)| < \infty$ as desired. \square

Since $T_1 \oplus \cdots \oplus T_m$ is fully invariant as the divisible part, and each its p_i -component T_i is also invariant, we have

$$C(\varphi_T) \leq |F| \cdot \prod_{i=1}^m C(\varphi_{T_i}), \text{ where } \varphi_{T_i} = \varphi_T|_{T_i}.$$

Also, by Theorem 1.2,

$$\prod_{i=1}^m R(\varphi_{T_i}) = R(\varphi_T|_{T_1 \oplus \cdots \oplus T_m}) \leq R(\varphi_T) \cdot |F| < \infty.$$

Applying Lemma 3.3 we obtain the main statement of this section:

Theorem 3.4. *Let φ be an automorphism of an abelian group of finite rank with $R(\varphi) < \infty$. Then $|C(\varphi)| < \infty$.*

4. PROOF OF THEOREM B

Proof of Theorem B. Suppose, $\varphi : G \rightarrow G$ is an automorphism with $R(\varphi) < \infty$. Then Theorem A implies that G is soluble-by-finite.

Let us remark that the soluble subgroup H of finite index constructed in Theorem A is characteristic and one can reduce the proof of $\text{TBF}T_f$ to the case of a soluble G . Indeed, for the projection $p : G \rightarrow G/H$ and the induced automorphism $\bar{\varphi} : G/H \rightarrow G/H$, consider some representatives g_1, \dots, g_r (with $g_1 = e$) of Reidemeister classes of φ . Denote by φ' the restriction of φ to H . For each τ_{g_j} , $j = 1, \dots, r$, one has $R(\tau_{g_j} \circ \varphi') < \infty$ by Theorem 1.2 and since $|G/H| < \infty$. If H has the $\text{TBF}T_f$ property, one can find, for each j , a normal $\tau_{g_j} \circ \varphi'$ -invariant subgroup H_j such that $p_j : H \rightarrow H/H_j$ gives a bijection of Reidemeister classes of $\tau_{g_j} \circ \varphi'$ and of the induced automorphism of H/H_j . Since the rank of H is finite, one can consider the intersection H'_j of all subgroups of H of the same index as H_j . Then by 1.6 H'_j is a characteristic subgroup of H of finite index. The same is true for $H^0 := \bigcap_{j=1}^r H'_j$. So, we can consider an equivariant epimorphism $p^0 : G \rightarrow G/H^0$, where G/H^0 is finite. It remains to verify that p^0 gives a bijective mapping of Reidemeister classes. As in the proof of Corollary 1.11, we need to prove that Reidemeister classes of φ are some unions of cosets of H^0 . For this purpose, consider an arbitrary $g \in G$. Then, for some j , $j = 1, \dots, r$, we have $g \in \{g_j\}_\varphi$. Hence, $g = xg_j\varphi(x^{-1})$, for some $x \in G$. Then $H^0g = H^0xg_j\varphi(x^{-1}) = xH^0g_j\varphi(x^{-1})$. Since $H^0 \subseteq \{e\}_{\tau_{g_j} \circ \varphi'}$, for any $h_0 \in H^0$ and some $h \in H$, we have $h_0 = h(\tau_{g_j} \circ \varphi')(h^{-1})$. Then

$$xh^0g_j\varphi(x^{-1}) = xhg_j(\varphi')(h^{-1})(g_j)^{-1}g_j\varphi(x^{-1}) = xhg_j(\varphi')(h^{-1})\varphi(x^{-1}) \in \{g_j\}_\varphi.$$

Hence,

$$(5) \quad g \in H^0g \subseteq \{g_j\}_\varphi.$$

So any class is a union of H^0 -cosets and the reduction is proved.

Thus we suppose G to be soluble and consider the derived series of G :

$$(6) \quad G = G^{(0)} \supset G' = G^{(1)} \supset G^{(2)} \supset \cdots \supset G^{(k)} = \{e\}, \quad G^{(i+1)} = [G^{(i)}, G^{(i)}],$$

of characteristic subgroups. Denote by $A_i := G^{(i)}/G^{(i+1)}$ the corresponding abelian groups and by $G_i := G/G^{(i)}$ the corresponding factor groups. Denote the induced automorphism by

$$\varphi_i := G^{(i)} \rightarrow G^{(i)}, \quad \alpha_i : A_i \rightarrow A_i, \quad \beta_i : G_i \rightarrow G_i.$$

First we will prove inductively that $R(\varphi_i) < \infty$, and similarly, by Lemma 1.3 applied to the initial φ , that

$$(7) \quad R(\tau_g \circ \varphi_i) < \infty, \quad \text{for any } g \in G, \quad i = 0, \dots, k.$$

Indeed, by Theorem 3.4, $|C(\alpha_0)| < \infty$. Then, by Theorem 1.2.a), $R(\varphi_1) < \infty$. Since the finite rank condition is hereditary for subgroups and factor groups, the next steps of the induction are quite similar to the first one.

Now we will argue by induction to prove the theorem. Since $G^{(k-1)}$ is abelian, TBFT_f is true for it. Suppose that the theorem is proved for $G^{(i+1)}$ and prove it for $G^{(i)}$.

Choose representatives $e = g_1, \dots, g_r$ in $G^{(i)}$ of Reidemeister classes of φ_i . For each g_j ($j = 1, \dots, r$), choose a subgroup H_j realizing the TBFT_f for $\tau_{g_j} \circ \varphi_{i+1}$. In other words, H_j is a normal $\tau_{g_j} \circ \varphi_{i+1}$ -invariant subgroup of $G^{(i+1)}$ and the projection $G^{(i+1)} \rightarrow G^{(i+1)}/H_j$ induces a bijection of Reidemeister classes of $\tau_{g_j} \circ \varphi_{i+1}$ and of those of the induced automorphism of $G^{(i+1)}/H_j$ (equivalently each Reidemeister class of $\tau_{g_j} \circ \varphi_{i+1}$ is a union of some cosets of H_j).

Consider $H_0 := \cap_{j=1}^r H_j$. This is a subgroup of finite index in $G^{(i+1)}$. By Theorem 1.6 the intersection H of all subgroups of $G^{(i+1)}$ of the same index as H_0 is a characteristic subgroup of $G^{(i+1)}$ of finite index. We have $H \subseteq H_0 \subseteq H_j$ ($j = 1, \dots, r$).

Then $G^{(i)} \rightarrow G^{(i)}/H$ induces a bijection of Reidemeister classes. This can be proved literally like for $G \rightarrow G/H^0$ in the beginning of this proof. Finally $G^{(i)}/H$ is finite-by-abelian, namely $(G^{(i+1)}/H)$ -by- A_i . Since A_i has finite rank, we apply Lemma 1.10 and Corollary 1.11 to see that TBFT_f is true for $G^{(i)}/H$ and there is an equivariant projection onto a finite group $G^{(i)}/H \rightarrow F$ inducing a bijection of Reidemeister classes. Then the composition $G^{(i)} \rightarrow G^{(i)}/H \rightarrow F$ gives a bijection of Reidemeister classes, which proves the TBFT_f for $G^{(i)}$. The induction completes the proof. \square

REFERENCES

- [1] CHARLES W. CURTIS AND IRVING REINER. *Representation theory of finite groups and associative algebras*. Pure and Applied Mathematics, Vol. XI. Interscience Publishers, a division of John Wiley & Sons, New York-London, 1962.
- [2] ALEXANDER FEL'SHTYN. Dynamical zeta functions, Nielsen theory and Reidemeister torsion. *Mem. Amer. Math. Soc.* **147**, No. 699, xii+146, 2000.
- [3] ALEXANDER FEL'SHTYN AND BENJAMIN KLOPSCH. Pólya-Carlson dichotomy for coincidence Reidemeister zeta functions via profinite completions. *Indag. Math. (N.S.)* **33**, No. 4, 753–767, 2022.
- [4] ALEXANDER FEL'SHTYN, NIKITA LUCHNIKOV, AND EVGENIJ TROITSKY. Twisted inner representations. *Russian Journal of Mathematical Physics* **22**, No. 3, 301–306, 2015.
- [5] ALEXANDER FEL'SHTYN AND EVGENIJ TROITSKY. Twisted Burnside-Frobenius theory for discrete groups. *J. Reine Angew. Math.* **613**, 193–210, 2007.
- [6] ALEXANDER FEL'SHTYN AND EVGENIJ TROITSKY. Aspects of the property R_∞ . *J. Group Theory* **18**, No. 6, 1021–1034, 2015.
- [7] A. FEL'SHTYN AND E. TROITSKY. Pólya-Carlson dichotomy for dynamical zeta functions and a twisted Burnside-Frobenius theorem. *Russ. J. Math. Phys.* **28**, No. 4, 455–463, 2021.
- [8] ALEXANDER FEL'SHTYN, EVGENIJ TROITSKY, AND ANATOLY VERSHIK. Twisted Burnside theorem for type II_1 groups: an example. *Math. Res. Lett.* **13**, No. 5, 719–728, 2006.
- [9] A. FEL'SHTYN, E. TROITSKY, AND M. ZIETEK. New Zeta Functions of Reidemeister Type and the Twisted Burnside-Frobenius Theory. *Russ. J. Math. Phys.* **27**, No. 2, 199–211, 2020.
- [10] D. GONÇALVES. The coincidence Reidemeister classes on nilmanifolds and nilpotent fibrations. *Topology and Its Appl.* **83**, 169–186, 1998.

- [11] DACIBERG GONÇALVES AND PETER WONG. Twisted conjugacy classes in wreath products. *Internat. J. Algebra Comput.* **16**, No. 5, 875–886, 2006.
- [12] DACIBERG GONÇALVES AND PETER WONG. Twisted conjugacy classes in nilpotent groups. *J. Reine Angew. Math.* **633**, 11–27, 2009.
- [13] ENRICO JABARA. Automorphisms with finite Reidemeister number in residually finite groups. *J. Algebra* **320**, No. 10, 3671–3679, 2008.
- [14] A. JAIKIN-ZAPIRAIN. Finite groups of bounded rank with an almost regular automorphism. *Israel J. Math.* **129**, 209–220, 2002.
- [15] A. A. KIRILLOV. *Elements of the Theory of Representations*. Springer-Verlag, Berlin Heidelberg New York, 1976.
- [16] A. G. KUROSH. *The theory of groups*. Translated from the Russian and edited by K. A. Hirsch. 2nd English ed. 2 volumes. Chelsea Publishing Co., New York, 1960.
- [17] ALEXANDER LUBOTZKY AND DAN SEGAL. *Subgroup growth*, volume 212 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2003.
- [18] AVINOAM MANN AND DAN SEGAL. Uniform finiteness conditions in residually finite groups. *Proc. London Math. Soc. (3)* **61**, No. 3, 529–545, 1990.
- [19] TIMUR R. NASYBULLOV. The R_∞ -property for Chevalley groups of types B_l , C_l , D_l over integral domains. *J. Algebra* **446**, 489–498, 2016.
- [20] DEREK J. S. ROBINSON. *A course in the theory of groups*, volume 80 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1996.
- [21] EVGENIJ TROITSKY. Reidemeister classes in lamplighter-type groups. *Comm. Algebra* **47**, No. 4, 1731–1741, 2019. arXiv:1711.09371.
- [22] EVGENIJ TROITSKY. Twisted conjugacy in some lamplighter-type groups. e-print, 2022. <https://arxiv.org/abs/2208.13018>.

MOSCOW CENTER FOR FUNDAMENTAL AND APPLIED MATHEMATICS, MSU DEPARTMENT,
DEPT. OF MECH. AND MATH., LOMONOSOV MOSCOW STATE UNIVERSITY, 119991 MOSCOW, RUSSIA
Email address: troitsky@mech.math.msu.su