

Chi-Square Goodness-of-Fit Tests for Conditional Distributions*

Miguel A. Delgado[†]

Department of Economics

Universidad Carlos III de Madrid

Julius Vainora

Department of Economics

University of Cambridge

October 4, 2022

Abstract

We propose a cross-classification rule for the dependent and explanatory variables resulting in a contingency table such that the classical trinity of chi-square statistics can be used to check for conditional distribution specification. The resulting Pearson statistic is equal to the Lagrange multiplier statistic. We also provide a Chernoff-Lehmann result for the Pearson statistic using the raw data maximum likelihood estimator, which is applied to show that the corresponding limiting distribution of the Wald statistic does not depend on the number of parameters. The asymptotic distribution of the proposed statistics does not change when the grouping is data dependent. An algorithm allowing to control the number of observations per cell is developed. Monte Carlo experiments provide evidence of the excellent size accuracy of the proposed tests and their good power performance, compared to omnibus tests, in high dimensions.

Keywords: Conditional distribution specification testing; Rosenblatt transform; Pearson statistic; Trinity of chi-square tests.

*Research funded by Ministerio Economía y Competitividad (Spain), grant PID2021-15178NB-100.

[†]Corresponding author. E-mail address: delgado@est-econ.uc3m.es.

1 Introduction

The χ^2 goodness-of-fit tests are widely applied to check for model specification of distribution functions using grouped data. Given a grouping of a random sample into M cells, the χ^2 goodness-of-fit statistics evaluate whether the difference between expected and observed frequencies in each cell arose by chance. The observed frequencies are distributed as a multinomial random vector, and the χ^2 statistics correspond to the classical trinity of tests, based on the grouped data likelihood, for testing that the $M - 1$ linearly independent restrictions imposed by the specified model on the multinomial parameters hold (see, e.g., Rao, 2002, Section 6b and Bickel and Doksum, 2015, Section 6.4). The Lagrange multiplier (LM) statistic is identical to the X^2 statistic introduced by Pearson (1900), both under simple and composite hypotheses, but the Wald statistic is only identical under simple hypotheses. The likelihood ratio statistic (LR), known as G^2 , is usually numerically different from X^2 , but both have the same asymptotic distribution under the null.

Our tests are related to the McFadden (1974) proposal to check for the specification of a discrete choice (multinomial regression) model using choice experimental data, where J randomly chosen individuals with given attributes repeat choice trials among L alternatives. That is, the explanatory variables are fixed. See Read and Cressie (1988) and Osius and Rojek (1992) for later developments. Unlike the classical case, with full information on the data generating process, which includes the fixed explanatory variables setup, the expected frequencies of arbitrarily grouped data are unknown under the conditional distribution model specification, since the marginal distribution of the explanatory variables is unspecified. This makes it impossible to apply the χ^2 statistics using arbitrary groupings of the multivariate data (see Section 2). However, the expected frequencies of any grouped data can be estimated using a preliminary \sqrt{n} -consistent estimator of the model's parameters; for instance, the conditional Maximum Likelihood Estimator (MLE) using raw data. Model checks can then be based on a Wald statistic, i.e., a suitably standardized quadratic form in the difference between observed and estimated expected frequencies. This approach was proposed by Heckman (1984) to test whether the marginal distribution of the dependent variable is compatible with a conditional distribution specification. Horowitz (1985) applied the Wald test to check the specification of discrete choice models, and Andrews (1988a,b) considered specification testing of general conditional distribution models using possibly data-dependent grouping of the observations. This extends the Wald test for the classical marginal distribution specification testing using grouped data introduced by Nikulin (1973) and Rao and Robson (1974). See Moore (1977) for a summary of related results. The limiting distribution of these Wald statistics is a chi-squared random variable with degrees of freedom equal to

the rank of the weighting matrix, which can be estimated from the data.

In this paper, the hypothesis of correct specification is characterized in terms of the [Rosenblatt \(1952\)](#) transform of the dependent variable. Recall that the Rosenblatt transform extends the probability integral transform to the conditional case — the transform of the dependent variable is uniformly distributed on the interval $[0, 1]$, independently of the vector of explanatory variables. The data is cross-classified as follows: the Rosenblatt transform of the dependent variable observations under the assumed model is classified according to L subintervals of $[0, 1]$ on the rows, and the vector of explanatory variable observations according to a partition of its range into J cells on the columns. When the specified model is correct, the joint expected frequency in each cell of the resulting contingency table equals the product of the marginals on the corresponding row and column, and the expected marginal frequency of each row is the length of the $[0, 1]$ subinterval. Therefore, under the null hypothesis, the observed joint frequencies in the contingency table are distributed as a multinomial random vector, whose parameters are the product of the marginal frequencies in columns and rows. Thus, the G^2 statistic is the LR test for checking that the $LJ - 1$ linearly independent multinomial parameters satisfy the restrictions imposed by the specified model, and X^2 is the corresponding LM statistic, which is identical to the Wald statistic under simple hypotheses, as in the classical case. This X^2 is, in fact, the Pearson statistic for testing independence between rows and columns in the contingency table, where the expected marginal frequencies on the rows are known. Under simple hypotheses, there are only $J - 1$ unknown parameters (the marginal frequencies on the columns). Therefore, the statistics are distributed as a $\chi^2_{J(L-1)}$ under simple hypotheses. Under composite hypotheses, the p unknown parameters in the specified model are asymptotically efficiently estimated based only on the information in the contingency table using the true parameter to perform the Rosenblatt transform. Since the resulting estimator is infeasible, we use the equally asymptotically efficient one-step Gauss-Newton estimator starting from an arbitrary \sqrt{n} -consistent estimator. The corresponding trinity of χ^2 statistics is asymptotically distributed as a $\chi^2_{J(L-1)-p}$ under the null — the degrees of freedom are the $LJ - 1$ linear independent restrictions on the multinomial parameters minus the $J - 1 + p$ linearly independent estimated parameters.

We obtain the asymptotic distribution of the X^2 statistic using the raw data, rather than the grouped data, conditional MLE, which extends the [Chernoff and Lehmann \(1954\)](#) result. This result is applied to show that the corresponding Wald statistic is asymptotically distributed as a $\chi^2_{J(L-1)}$ under the null hypothesis, irrespective of the number of parameters estimated in the specified model. The asymptotic properties of all the considered statistics are preserved when the explanatory variables grouping depends on the data.

The rest of the article is organized as follows. [Section 2](#) discusses the difficulties of

using χ^2 goodness-of-fit test statistics to model checks for conditional distributions. Section 3 introduces the proposed grouping and the corresponding trinity of tests under simple hypotheses. Section 4 presents the grouped data MLE and justifies the tests for composite hypotheses. In this section, we also present the asymptotic distribution of the X^2 statistic using the fully efficient raw data MLE, and the corresponding Wald statistic. Section 5 justifies the validity of the tests when the explanatory variables' grouping is data dependent and discusses algorithmic grouping rules. Section 6 provides the power of the tests under contiguous alternatives. Finite sample properties of the tests are studied by means of Monte Carlo experiments in Section 7. In this section we compare the performance of the proposed tests with Andrews (1997) omnibus Kolmogorov-Smirnov type bootstrap test. Proofs are gathered at the end of the article.

2 The Pearson Tests to Check for Conditional Distributions Model Specification

The data set consists of i.i.d. observations $\{Y_i, \mathbf{X}_i\}_{i=1}^n$ of an \mathbb{R}^{1+k} -valued random vector (Y, \mathbf{X}) , where Y is the dependent variable taking values in \mathbb{R} , and \mathbf{X} is the vector of explanatory variables taking values in \mathbb{R}^k . The conditional cumulative distribution function (CDF) of Y given \mathbf{X} , denoted by $F_{Y|\mathbf{X}}(\cdot | \mathbf{X})$, satisfies

Assumption 1. $F_{Y|\mathbf{X}}(\cdot | \mathbf{X})$ is continuous a.s. on \mathbb{R} .

The hypothesis of interest is

$$H_0 : F_{Y|\mathbf{X}} \in \mathcal{F}$$

for a family of parametric continuous conditional CDF's $\mathcal{F} = \{F_{Y|\mathbf{X},\theta} : \theta \in \Theta\}$, given a proper parameter space $\Theta \subset \mathbb{R}^p$. That is, under H_0 , there exists a $\theta_0 \in \Theta$ such that $F_{Y|\mathbf{X}}(y | \mathbf{X}) = F_{Y|\mathbf{X},\theta_0}(y | \mathbf{X})$ a.s. for all $y \in \mathbb{R}$.

Let $\mathbf{C} = \{C_m\}_{m=1}^M$ be a partition of \mathbb{R}^{1+k} into $M > 1$ cells. The Pearson test is designed to detect alternatives to H_0 of the form

$$H_{1\mathbf{C}} : \text{For all } \theta \in \Theta, \Pr((Y, \mathbf{X}) \in C_m) \neq \pi_{C_m}(\theta) \text{ for some } m = 1, \dots, M,$$

where $\pi_{C_m}(\theta) = \int_{(y,\mathbf{x}) \in C_m} F_{Y|\mathbf{X},\theta}(dy | \mathbf{x}) F_{\mathbf{X}}(d\mathbf{x})$, and $F_{\mathbf{X}}$ is the CDF of \mathbf{X} .

The vector of observed frequencies $\hat{\mathbf{O}}_{\mathbf{C}} = (\hat{O}_{C_1}, \dots, \hat{O}_{C_M})'$, with $\hat{O}_{C_m} = \sum_{i=1}^n \mathbf{1}_{\{(Y_i, \mathbf{X}_i) \in C_m\}}$ and $\mathbf{1}_S$ as the indicator function of the event S , is distributed as a multinomial random vector with parameters $\pi_{C_m}^0 = \Pr((Y, \mathbf{X}) \in C_m)$, $m = 1, \dots, M$. Thus, the log-likelihood of

$\boldsymbol{\pi} = (\pi_1, \dots, \pi_M)'$ given $\hat{\boldsymbol{O}}_C$ is given by

$$\mathcal{L}(\boldsymbol{\pi} \mid \hat{\boldsymbol{O}}_C) \propto \sum_{m=1}^M \hat{O}_{C_m} \ln \pi_{C_m}.$$

Hence, the unrestricted MLE of $\pi_{C_m}^0$ is $n^{-1}\hat{O}_{C_m}$ (see, e.g., [Bickel and Doksum, 2015](#), Example 2.2.8). When F_X is known and, hence, the maps $\boldsymbol{\theta} \mapsto \pi_{C_m}(\boldsymbol{\theta})$, $m = 1, \dots, M$, are known, the log-likelihood of $\boldsymbol{\theta}$ given $\hat{\boldsymbol{O}}_C$ is

$$\mathcal{L}(\boldsymbol{\theta} \mid \hat{\boldsymbol{O}}_C) \propto \sum_{m=1}^M \hat{O}_{C_m} \ln \pi_{C_m}(\boldsymbol{\theta}). \quad (1)$$

Let $\bar{\boldsymbol{\theta}}$ be the MLE of $\boldsymbol{\theta}_0$ that maximizes (1). Thus, $\pi_{C_m}(\bar{\boldsymbol{\theta}})$ is the restricted MLE of $\pi_{C_m}^0$. The LR statistic, G^2 , for testing H_0 in the direction H_{1C} is $\bar{G}_C^2(\bar{\boldsymbol{\theta}})$, with

$$\bar{G}_C^2(\boldsymbol{\theta}) = 2 \sum_{m=1}^M \hat{O}_{C_m} \ln \frac{\hat{O}_{C_m}}{n\pi_{C_m}(\boldsymbol{\theta})},$$

and the corresponding X^2 is $\bar{X}_C^2(\bar{\boldsymbol{\theta}})$, with

$$\bar{X}_C^2(\boldsymbol{\theta}) = \sum_{m=1}^M \frac{(\hat{O}_{C_m} - n\pi_{C_m}(\boldsymbol{\theta}))^2}{n\pi_{C_m}(\boldsymbol{\theta})} = \bar{\boldsymbol{\Phi}}_C'(\boldsymbol{\theta})\bar{\boldsymbol{\Phi}}_C(\boldsymbol{\theta}),$$

and $\bar{\boldsymbol{\Phi}}_C(\boldsymbol{\theta}) = \text{diag}^{-1/2}(n\boldsymbol{\pi}_C(\boldsymbol{\theta})) (\hat{\boldsymbol{O}}_C - n\boldsymbol{\pi}_C(\boldsymbol{\theta}))$ with $\boldsymbol{\pi}_C(\boldsymbol{\theta}) = (\pi_{C_1}(\boldsymbol{\theta}), \dots, \pi_{C_M}(\boldsymbol{\theta}))'$. The Pearson X^2 and the LM statistics for testing H_0 in the direction H_{1C} are identical both under simple ($\boldsymbol{\theta}_0$ known) and composite hypotheses ($\boldsymbol{\theta}_0$ unknown). The Wald statistic is also identical to $\hat{X}_D^2(\boldsymbol{\theta}_0)$ under simple hypotheses. Under composite hypotheses, the LM, LR, and Wald tests are typically different. The trinity of test statistics is asymptotically distributed as a χ_{M-1}^2 under simple hypotheses and as a χ_{M-p-1}^2 under composite hypotheses (see, e.g., [Bickel and Doksum, 2015](#), Section 6.4).

The Wald's testing principle using any \sqrt{n} -consistent estimator $\tilde{\boldsymbol{\theta}}$, is

$$\bar{W}_C(\tilde{\boldsymbol{\theta}}) = \bar{\boldsymbol{\Phi}}_C'(\tilde{\boldsymbol{\theta}}) \widehat{\text{Avar}}^{-}(\bar{\boldsymbol{\Phi}}_C(\tilde{\boldsymbol{\theta}})) \bar{\boldsymbol{\Phi}}_C(\tilde{\boldsymbol{\theta}}),$$

where \mathbf{A}^- is a generalized inverse of \mathbf{A} , and, for any vector of statistics $\hat{\boldsymbol{\varphi}}$, $\widehat{\text{Avar}}(\hat{\boldsymbol{\varphi}})$ is a consistent estimator of $\text{Avar}(\hat{\boldsymbol{\varphi}})$, the asymptotic variance of $\hat{\boldsymbol{\varphi}}$. Under H_0 and regularity conditions, $\bar{W}_C(\tilde{\boldsymbol{\theta}})$ is asymptotically distributed as a $\chi_{\text{rank}(\widehat{\text{Avar}}(\bar{\boldsymbol{\Phi}}_C(\tilde{\boldsymbol{\theta}})))}^2$. When $\tilde{\boldsymbol{\theta}}$ is the MLE using raw data, the rank equals $M - 1$ a.s. See [Moore \(1977\)](#) and references therein.

When $F_{\mathbf{X}}$ is unknown, the statistics discussed above are infeasible, since $\boldsymbol{\theta} \mapsto \boldsymbol{\pi}_{\mathbf{C}}(\boldsymbol{\theta})$ is an unknown function. However, $\pi_{C_m}(\boldsymbol{\theta})$ is estimated by

$$\hat{\pi}_{C_m}(\boldsymbol{\theta}) = \frac{1}{n} \sum_{i=1}^n \int_{(y,x) \in C_m} F_{Y|x,\boldsymbol{\theta}}(dy | \mathbf{X}_i), \quad m = 1, \dots, M,$$

and the corresponding Wald statistic, using any \sqrt{n} -consistent estimator $\tilde{\boldsymbol{\theta}}$, is

$$\hat{W}_{\mathbf{C}}(\tilde{\boldsymbol{\theta}}) = \hat{\boldsymbol{\Phi}}_{\mathbf{C}}'(\tilde{\boldsymbol{\theta}}) \widehat{\text{Avar}}^{-1}(\hat{\boldsymbol{\Phi}}_{\mathbf{C}}(\tilde{\boldsymbol{\theta}})) \hat{\boldsymbol{\Phi}}_{\mathbf{C}}(\tilde{\boldsymbol{\theta}}),$$

with $\hat{\boldsymbol{\Phi}}_{\mathbf{C}}(\boldsymbol{\theta}) = \text{diag}^{-1/2}(n\hat{\boldsymbol{\pi}}_{\mathbf{C}}(\boldsymbol{\theta})) (\hat{\mathbf{O}}_{\mathbf{C}} - n\hat{\boldsymbol{\pi}}_{\mathbf{C}}(\boldsymbol{\theta}))$ and $\hat{\boldsymbol{\pi}}_{\mathbf{C}}(\boldsymbol{\theta}) = (\hat{\pi}_{C_1}(\boldsymbol{\theta}), \dots, \hat{\pi}_{C_M}(\boldsymbol{\theta}))'$. This statistics has been proposed by Heckman (1984) for partitions $\mathbf{C} = \{C_m\}_{m=1}^M$ with

$$C_m = \{(y, x) : y \in (a_{m-1}, a_m], x \in \mathbb{R}\}, \quad (2)$$

where $-\infty = a_0 \leq a_1 \leq \dots \leq a_M = \infty$. That is, the data is classified according to Y , and the corresponding $\hat{W}_{\mathbf{C}}$ compares the expected and observed frequencies of the dependent variable. So, the test can only detect the alternatives where the actual marginal CDF of Y is not compatible with the specified conditional CDF. Horowitz (1985) considers the statistic to check for multinomial regression specifications, and Andrews (1988a,b) for general specifications using data-dependent partitions.

Under H_0 and suitable regularity conditions $\hat{W}(\tilde{\boldsymbol{\theta}})$ is asymptotically distributed as a chi-squared random variable with degrees of freedom equal to $\text{rank}(\text{Avar}(\hat{\boldsymbol{\Phi}}_{\mathbf{C}}(\tilde{\boldsymbol{\theta}})))$. When the classes in the partitions are of the form (2) and $\tilde{\boldsymbol{\theta}}$ is the conditional MLE, $\text{rank}(\text{Avar}(\hat{\boldsymbol{\Phi}}_{\mathbf{C}}(\tilde{\boldsymbol{\theta}}))) = M - 1$ (Heckman, 1984). But, in general, $\text{rank}(\text{Avar}(\hat{\boldsymbol{\Phi}}_{\mathbf{C}}(\tilde{\boldsymbol{\theta}})))$ must be estimated.

Under H_0 , $\hat{\boldsymbol{\Phi}}_{\mathbf{C}}(\boldsymbol{\theta}_0) \xrightarrow{d} \mathcal{N}_M(\mathbf{0}, \mathbf{Q}_{\mathbf{C}}(\boldsymbol{\theta}_0))$, where

$$\mathbf{Q}_{\mathbf{C}}(\boldsymbol{\theta}) = \text{diag}^{-1/2}(\boldsymbol{\pi}_{\mathbf{C}}(\boldsymbol{\theta})) \cdot \boldsymbol{\Sigma}_{\mathbf{C}}(\boldsymbol{\theta}) \cdot \text{diag}^{-1/2}(\boldsymbol{\pi}_{\mathbf{C}}(\boldsymbol{\theta})),$$

with $\boldsymbol{\Sigma}_{\mathbf{C}}(\boldsymbol{\theta}) = \mathbb{E}(\text{diag}(\mathbf{p}_{\mathbf{C},\boldsymbol{\theta}}(\mathbf{X})) - \mathbf{p}_{\mathbf{C},\boldsymbol{\theta}}(\mathbf{X})\mathbf{p}_{\mathbf{C},\boldsymbol{\theta}}'(\mathbf{X}))$, $\mathbf{p}_{\mathbf{C},\boldsymbol{\theta}}(\mathbf{X}) = (p_{C_1,\boldsymbol{\theta}}(\mathbf{X}), \dots, p_{C_M,\boldsymbol{\theta}}(\mathbf{X}))'$, and $p_{C_m,\boldsymbol{\theta}}(\mathbf{X}) = \int_{(y,\mathbf{X}) \in C_m} F_{Y|\mathbf{X},\boldsymbol{\theta}}(dy | \mathbf{X})$. The corresponding Pearson X^2 statistic is $\hat{X}_{\mathbf{C}}^2(\boldsymbol{\theta}_0)$ with

$$\hat{X}_{\mathbf{C}}^2(\boldsymbol{\theta}) = \sum_{m=1}^M \frac{(\hat{O}_{C_m} - n\hat{\pi}_{C_m}(\boldsymbol{\theta}))^2}{n\hat{\pi}_{C_m}(\boldsymbol{\theta})} = \hat{\boldsymbol{\Phi}}_{\mathbf{C}}'(\boldsymbol{\theta}) \hat{\boldsymbol{\Phi}}_{\mathbf{C}}(\boldsymbol{\theta}).$$

Therefore, under H_0 ,

$$\hat{X}_{\mathbf{C}}^2(\boldsymbol{\theta}_0) \xrightarrow{d} \sum_{m=1}^M \alpha_m Z_m^2,$$

where the Z_m 's are independent standard normal random variables, and $\{\alpha_m\}_{m=1}^M$ are the eigenvalues of $\mathbf{Q}_C(\boldsymbol{\theta})$. In general, the matrix $\mathbf{Q}_C(\boldsymbol{\theta})$ is not idempotent, and its eigenvalues may depend on the unknown $\boldsymbol{\theta}_0$ and $F_{\mathbf{X}}$. Thus, the asymptotic distribution of $\hat{X}_C^2(\boldsymbol{\theta}_0)$ is not pivotal. However, when $\mathbf{Q}_C(\boldsymbol{\theta})$ is idempotent and the α_m 's take values in $\{0, 1\}$, $\hat{X}_C^2(\boldsymbol{\theta}_0) \xrightarrow{d} \chi_{\sum_{j=1}^M \alpha_m}^2$ under H_0 , but this is rarely the case for general partitions \mathbf{C} , as we discuss in the following remark. Henceforth, for any vector $\mathbf{a} = (a_1, \dots, a_m)'$, $\sqrt{\mathbf{a}} = (\sqrt{a_1}, \dots, \sqrt{a_m})'$.

Remark 1. In the trivial case $\mathbf{p}_{C, \boldsymbol{\theta}_0}(\mathbf{X}) = \boldsymbol{\pi}_C(\boldsymbol{\theta}_0)$ a.s., under H_0 , $\boldsymbol{\Sigma}_C(\boldsymbol{\theta}_0) = \text{diag}(\boldsymbol{\pi}_C(\boldsymbol{\theta}_0)) - \boldsymbol{\pi}_C(\boldsymbol{\theta}_0)\boldsymbol{\pi}_C'(\boldsymbol{\theta}_0)$, $\mathbf{Q}_C(\boldsymbol{\theta}_0) = \mathbf{I}_M - \sqrt{\boldsymbol{\pi}_C(\boldsymbol{\theta}_0)}\sqrt{\boldsymbol{\pi}_C(\boldsymbol{\theta}_0)'}$ is idempotent with rank $\text{tr}(\mathbf{Q}_C(\boldsymbol{\theta}_0)) = M - 1$, and $\bar{X}_C^2(\boldsymbol{\theta}_0) \xrightarrow{d} \chi_{M-1}^2$. This corresponds to the multinomial regression fixed design under product-multinomial sampling. However, for most choices of \mathbf{C} using observational data, $\mathbf{p}_{C, \boldsymbol{\theta}}(\mathbf{X})$ is not constant, which implies that $\mathbf{Q}_C(\boldsymbol{\theta})$ is not idempotent, and the limiting distribution of $\bar{X}_C^2(\boldsymbol{\theta}_0)$ under H_0 is non-pivotal.

3 The Trinity of Tests Under Simple Hypotheses Using Cross-Classified Data

We propose parameter-dependent partitions

$$\mathbf{D}(\boldsymbol{\theta}) = \{D_{\ell j}(\boldsymbol{\theta}) : \ell = 1, \dots, L, j = 1, \dots, J\}$$

with

$$D_{\ell j}(\boldsymbol{\theta}) = \{(y, \mathbf{x}) \in \mathbb{R} \times \mathbb{R}^k : F_{Y|\mathbf{X}, \boldsymbol{\theta}}(y | \mathbf{x}) \in U_\ell, \mathbf{x} \in A_j\},$$

where $\mathbf{U} = \{U_\ell\}_{\ell=1}^L$ is a partition of the interval $[0, 1]$, $U_\ell = (\vartheta_{\ell-1}, \vartheta_\ell]$ with $0 = \vartheta_0 < \vartheta_1 < \dots < \vartheta_L = 1$, and $\mathbf{A} = \{A_j\}_{j=1}^J$ is a partition of \mathbb{R}^k .

The data set $\{Y_i, \mathbf{X}_i\}_{i=1}^n$ is grouped according to the partition $\mathbf{D}(\boldsymbol{\theta})$, which cross-classifies $\{V_i(\boldsymbol{\theta}), \mathbf{X}_i\}_{i=1}^n$ into $M = LJ$ classes, where $V_i(\boldsymbol{\theta}) = F_{Y|\mathbf{X}, \boldsymbol{\theta}}(Y_i | \mathbf{X}_i)$ is the Rosenblatt transform of Y_i according to the specified model. This transform was proposed by [Rosenblatt \(1952\)](#) for omnibus specification testing of marginal multivariate continuous CDF's, and applied by [Delgado and Stute \(2008\)](#) to omnibus distribution-free specification testing of conditional CDF index models. Each observation (Y_i, \mathbf{X}_i) is simultaneously cross-classified according to the Rosenblatt transform, $V_i(\boldsymbol{\theta})$, into one of the L classes of the interval $[0, 1]$ partition, \mathbf{U} , on the rows, and according to the explanatory variables vector, \mathbf{X}_i , into one of the J classes of the \mathbb{R}^k partition, \mathbf{A} , on the columns. The vector of observed frequencies in each of the JL cells, for a given $\boldsymbol{\theta} \in \boldsymbol{\Theta}$, is

$$\hat{\mathbf{O}}_{\mathbf{D}(\boldsymbol{\theta})} = \left(\hat{O}_{D_{11}(\boldsymbol{\theta})}, \dots, \hat{O}_{D_{1J}(\boldsymbol{\theta})}, \dots, \hat{O}_{D_{L1}(\boldsymbol{\theta})}, \dots, \hat{O}_{D_{LJ}(\boldsymbol{\theta})} \right)'$$

and $\hat{O}_{D_{\ell j}}(\boldsymbol{\theta}) = \sum_{i=1}^n \mathbb{1}_{\{V_i(\boldsymbol{\theta}) \in U_\ell, \mathbf{X}_i \in A_j\}} = \sum_{i=1}^n \eta_{\ell i}(\boldsymbol{\theta}) \cdot \zeta_{ji}$, where $\eta_{\ell i} = \mathbb{1}_{\{V_i(\boldsymbol{\theta}) \in U_\ell\}}$ and $\zeta_{ji} = \mathbb{1}_{\{\mathbf{X}_i \in A_j\}}$. The joint and marginal frequencies are summarized as a contingency table (Table 1), where $\hat{q}_j = n^{-1} \sum_{\ell=1}^L \hat{O}_{D_{\ell j}}(\boldsymbol{\theta}) = n^{-1} \sum_{i=1}^n \mathbb{1}_{\{\mathbf{X}_i \in A_j\}}$ and $\hat{v}_\ell = n^{-1} \sum_{j=1}^J \hat{O}_{D_{\ell j}}(\boldsymbol{\theta}) = n^{-1} \sum_{i=1}^n \mathbb{1}_{\{V_i(\boldsymbol{\theta}) \in U_\ell\}}$ are the observed marginal frequencies on the columns and rows, with corresponding expected frequencies $q_j = \mathbb{E}(\hat{q}_j) = \Pr(\mathbf{X} \in A_j)$ and $v_\ell(\boldsymbol{\theta}) = \Pr(V_1(\boldsymbol{\theta}) \in U_\ell)$, respectively.

	$\mathbf{X} \in A_1$	\cdots	$\mathbf{X} \in A_J$	Sum
$V(\boldsymbol{\theta}) \in U_1$	$\hat{O}_{D_{11}}(\boldsymbol{\theta})$	\cdots	$\hat{O}_{D_{1J}}(\boldsymbol{\theta})$	$n\hat{v}_1(\boldsymbol{\theta})$
\vdots	\vdots	\ddots	\vdots	\vdots
$V(\boldsymbol{\theta}) \in U_L$	$\hat{O}_{D_{L1}}(\boldsymbol{\theta})$	\cdots	$\hat{O}_{D_{LJ}}(\boldsymbol{\theta})$	$n\hat{v}_L(\boldsymbol{\theta})$
Sum	$n\hat{q}_1$	\cdots	$n\hat{q}_J$	1

Table 1: Contingency table corresponding to $\mathbf{D}(\boldsymbol{\theta})$.

For a given $\boldsymbol{\theta} \in \Theta$, $\hat{\mathbf{O}}_{\mathbf{D}(\boldsymbol{\theta})}$ is distributed as a multinomial random vector with mean $\mathbb{E}(\hat{\mathbf{O}}_{\mathbf{D}(\boldsymbol{\theta})}) = n\boldsymbol{\pi}_{\mathbf{D}(\boldsymbol{\theta})}^0$, with

$$\boldsymbol{\pi}_{\mathbf{D}(\boldsymbol{\theta})}^0 = \left(\pi_{D_{11}}^0(\boldsymbol{\theta}), \dots, \pi_{D_{1J}}^0(\boldsymbol{\theta}), \dots, \pi_{D_{L1}}^0(\boldsymbol{\theta}), \dots, \pi_{D_{LJ}}^0(\boldsymbol{\theta}) \right)',$$

and

$$\pi_{D_{\ell j}}^0(\boldsymbol{\theta}) = \mathbb{E}(\zeta_{j1} \cdot \eta_{\ell 1}(\boldsymbol{\theta})) = \int_{\mathbf{x} \in A_j} \int_{F_{Y|\mathbf{X},\boldsymbol{\theta}}^{-1}(\vartheta_{\ell-1} | \mathbf{x})}^{F_{Y|\mathbf{X},\boldsymbol{\theta}}^{-1}(\vartheta_\ell | \mathbf{x})} F_{Y|\mathbf{X}}(\mathrm{d}y | \mathbf{x}) F_{\mathbf{X}}(\mathrm{d}\mathbf{x}).$$

Under H_0 , $\{V_i(\boldsymbol{\theta}_0)\}_{i=1}^n$ are independently distributed as uniform random variables on $[0, 1]$ independently of $\{\mathbf{X}_i\}_{i=1}^n$. Thus, under H_0 , $v_\ell(\boldsymbol{\theta}_0) = \vartheta_\ell - \vartheta_{\ell-1} = v_\ell$ for $\ell = 1, \dots, L$, and

$$\mathbb{E}(\hat{\mathbf{O}}_{\mathbf{D}^0}) = n \cdot \mathbb{E}(\boldsymbol{\eta}_1(\boldsymbol{\theta}_0) \otimes \boldsymbol{\zeta}_1) = n \cdot \mathbb{E}(\boldsymbol{\eta}_1(\boldsymbol{\theta}_0)) \otimes \mathbb{E}(\boldsymbol{\zeta}_1) = n \cdot \mathbf{v} \otimes \mathbf{q},$$

where, henceforth, $\mathbf{D}^0 = \mathbf{D}(\boldsymbol{\theta}_0)$, $\mathbf{v} = (v_1, \dots, v_L)'$, $\mathbf{q} = (q_1, \dots, q_J)'$, $\boldsymbol{\zeta}_i = (\zeta_{i1}, \dots, \zeta_{iJ})'$, and $\boldsymbol{\eta}_i(\boldsymbol{\theta}) = (\eta_{1i}(\boldsymbol{\theta}), \dots, \eta_{Li}(\boldsymbol{\theta}))'$. The alternative of interest with the proposed partition is

$$H_{1D} : \boldsymbol{\pi}_{\mathbf{D}(\boldsymbol{\theta})}^0 \neq \mathbf{v} \otimes \mathbf{q} \text{ for all } \boldsymbol{\theta} \in \Theta.$$

This is, in fact, the alternative hypothesis for the independence between rows and columns in the contingency table, where the expected marginal frequencies on the rows, \mathbf{v} , are known. Therefore, there are $LJ - 1$ linear independence restrictions and, since \mathbf{v} is known, $J - 1$ free parameters (the $J - 1$ linearly independent components of \mathbf{q}) to be estimated.

The log-likelihood of $\boldsymbol{\pi} = (\pi_{11}, \dots, \pi_{1J}, \dots, \pi_{L1}, \dots, \pi_{LJ})'$ for the observed joint frequen-

cies $\hat{\mathbf{O}}_{D^0}$ is

$$\mathcal{L}(\boldsymbol{\pi} \mid \hat{\mathbf{O}}_{D^0}) \propto \sum_{\ell=1}^L \sum_{j=1}^J \hat{O}_{D_{\ell j}(\boldsymbol{\theta}_0)} \ln \pi_{\ell j}. \quad (3)$$

Thus, the unrestricted MLE of $\boldsymbol{\pi}_{D^0}^0$ is $n^{-1}\hat{\mathbf{O}}_{D^0}$, and the corresponding restricted estimator is $\mathbf{v} \otimes \hat{\mathbf{q}}$, with $\hat{\mathbf{q}} = (\hat{q}_1, \dots, \hat{q}_J)'$. Therefore, under simple hypotheses, the X^2 statistic is, in this case, $\hat{X}^2(\boldsymbol{\theta}_0)$, where

$$\hat{X}_D^2(\boldsymbol{\theta}) = \sum_{\ell=1}^L \sum_{j=1}^J \frac{(\hat{O}_{D_{\ell j}(\boldsymbol{\theta})} - n \cdot v_{\ell} \cdot \hat{q}_j)^2}{n \cdot v_{\ell} \cdot \hat{q}_j} = \hat{\boldsymbol{\Phi}}_D'(\boldsymbol{\theta}) \hat{\boldsymbol{\Phi}}_D(\boldsymbol{\theta}),$$

and

$$\hat{\boldsymbol{\Phi}}_D(\boldsymbol{\theta}) = \text{diag}^{-1/2}(n(\mathbf{v} \otimes \hat{\mathbf{q}})) \left(\hat{\mathbf{O}}_{D(\boldsymbol{\theta})} - n(\mathbf{v} \otimes \hat{\mathbf{q}}) \right). \quad (4)$$

Under H_0 , the information matrix of $\hat{\mathbf{O}}_{D^0}$ is

$$\mathcal{I}_D = \frac{1}{n} \text{Var} \left(\hat{\mathbf{O}}_{D^0} - n(\mathbf{v} \otimes \mathbf{q}) \right) = (\text{diag}(\mathbf{v}) - \mathbf{v}\mathbf{v}') \otimes \text{diag}(\mathbf{q}).$$

Therefore, $\text{Avar} \left(\hat{\boldsymbol{\Phi}}_D(\boldsymbol{\theta}_0) \right) = \mathbf{Q}_U$ for any grouping of the explanatory variables \mathbf{A} , and

$$\mathbf{Q}_U = \text{diag}^{-1/2}(\mathbf{v} \otimes \mathbf{q}) \cdot \mathcal{I}_D \cdot \text{diag}^{-1/2}(\mathbf{v} \otimes \mathbf{q}) = \left(\mathbf{I}_L - \sqrt{\mathbf{v}}\sqrt{\mathbf{v}'} \right) \otimes \mathbf{I}_J \quad (5)$$

is idempotent with $\text{rank}(\mathbf{Q}_U) = \text{tr}(\mathbf{Q}_U) = J(L-1)$.

The Wald statistic based on (3), with known $\boldsymbol{\theta}_0$, is

$$\hat{W}_D(\boldsymbol{\theta}_0) = \hat{\boldsymbol{\Phi}}_D'(\boldsymbol{\theta}_0) \cdot \text{Avar}^{-1} \left(\hat{\boldsymbol{\Phi}}_D(\boldsymbol{\theta}_0) \right) \cdot \hat{\boldsymbol{\Phi}}_D(\boldsymbol{\theta}_0).$$

In order to derive the LM statistic, we need to obtain the information matrix inverse of the grouped data $\hat{\mathbf{O}}_{D^0}$. It suffices to consider $LJ-1$ free parameters $\boldsymbol{\pi}^* = (\pi_1^*, \dots, \pi_{LJ-1}^*)'$, where we remove π_{LJ} from $\boldsymbol{\pi}$, i.e., $\pi_s^* = \pi_{\ell j}$ for $s = (\ell-1)J+j$ with $(\ell, j) \in (\{1, \dots, L\} \times \{1, \dots, J\}) \setminus \{(L, J)\}$. Notice that the score vector is $\hat{\boldsymbol{\Psi}}_D(\boldsymbol{\theta}_0)$, where $\hat{\boldsymbol{\Psi}}_D(\boldsymbol{\theta}) = n^{-1} \left(\hat{\psi}_1(\boldsymbol{\theta}), \dots, \hat{\psi}_{LJ-1}(\boldsymbol{\theta}) \right)'$ and

$$\hat{\psi}_s(\boldsymbol{\theta}) = \frac{\partial}{\partial \pi_s^*} \mathcal{L}(\boldsymbol{\pi} \mid \hat{\mathbf{O}}_{D(\boldsymbol{\theta})}) \Big|_{\boldsymbol{\pi}=\mathbf{v} \otimes \mathbf{q}} = \frac{\hat{O}_{D_{\ell j}(\boldsymbol{\theta})}}{v_{\ell} \hat{q}_j} - \frac{\hat{O}_{D_{LJ}(\boldsymbol{\theta})}}{v_L \hat{q}_J}$$

for $s = (\ell-1)J+j$ with $(\ell, j) \in (\{1, \dots, L\} \times \{1, \dots, J\}) \setminus \{(L, J)\}$. The information matrix inverse of $\hat{\mathbf{O}}_{D^0}$ is

$$\mathcal{I}_D^{-1} = \text{Var} \left(\frac{\hat{\mathbf{O}}_{D^0}^*}{\sqrt{n}} \right) = \text{diag}(\boldsymbol{\pi}^*) - \boldsymbol{\pi}^* \boldsymbol{\pi}^{*'},$$

where $\hat{\mathbf{O}}_{\mathbf{D}(\boldsymbol{\theta})}^* = (\hat{O}_1^*(\boldsymbol{\theta}), \dots, \hat{O}_{LJ-1}^*(\boldsymbol{\theta}))'$, $\hat{O}_s^*(\boldsymbol{\theta}) = \hat{O}_{D_{\ell_j}(\boldsymbol{\theta})}$, $\boldsymbol{\pi}^* = (\pi_1^*, \dots, \pi_{LJ-1}^*)'$, and $\pi_s^* = v_\ell q_j$ under H_0 . Thus, $\mathcal{I}_{\mathbf{D}}^{-1}$ is estimated by $\hat{\mathcal{I}}_{\mathbf{D}}^{-1}(\boldsymbol{\theta}) = \text{diag}(\hat{\boldsymbol{\pi}}^*(\boldsymbol{\theta})) - \hat{\boldsymbol{\pi}}^*(\boldsymbol{\theta})\hat{\boldsymbol{\pi}}^{*\prime}(\boldsymbol{\theta})$, where $\hat{\boldsymbol{\pi}}^* = (\hat{\pi}_1^*, \dots, \hat{\pi}_{LJ-1}^*)'$ and $\hat{\pi}_s^* = v_\ell \hat{q}_j$. The LM statistic is $\widehat{LM}_{\mathbf{D}}(\boldsymbol{\theta}_0)$, with

$$\widehat{LM}_{\mathbf{D}}(\boldsymbol{\theta}) = n \cdot \hat{\boldsymbol{\Psi}}_{\mathbf{D}}'(\boldsymbol{\theta}) \cdot \hat{\mathcal{I}}_{\mathbf{D}}^{-1} \cdot \hat{\boldsymbol{\Psi}}_{\mathbf{D}}(\boldsymbol{\theta}).$$

The LR statistic based on (3), with known $\boldsymbol{\theta}_0$, is $\hat{G}_{\mathbf{D}}^2(\boldsymbol{\theta}_0)$ with

$$\hat{G}_{\mathbf{D}}^2(\boldsymbol{\theta}) = 2 \sum_{\ell=1}^L \sum_{j=1}^J \hat{O}_{D_{\ell_j}(\boldsymbol{\theta})} \ln \frac{\hat{O}_{D_{\ell_j}(\boldsymbol{\theta})}}{nv_\ell \hat{q}_j}.$$

There are $LJ - 1$ linearly independent restrictions on the multinomial parameters under H_0 , and $J - 1$ estimated free parameters q_1, \dots, q_{J-1} . Therefore, the trinity of test statistics are asymptotically distributed as a $\chi_{J(L-1)}^2$ under H_0 (see, e.g., [Bickel and Doksum, 2015](#), Sections 6.3.2 and 6.4.1).

The next theorem, proved in the Appendix as the rest of results, establishes that $\hat{W}_{\mathbf{D}}(\boldsymbol{\theta})$, $\widehat{LM}_{\mathbf{D}}(\boldsymbol{\theta})$, and $\hat{X}_{\mathbf{D}}^2(\boldsymbol{\theta})$ are identical, and provides the asymptotic distribution of all the statistics.

Theorem 1. *Let Assumption 1 be satisfied. Then*

- (i) $\hat{X}_{\mathbf{D}}^2(\boldsymbol{\theta}_0) \xrightarrow{d} \chi_{J(L-1)}^2$ under H_0 ,
- (ii) $\hat{X}_{\mathbf{D}}^2(\boldsymbol{\theta}) = \widehat{LM}_{\mathbf{D}}(\boldsymbol{\theta})$ a.s. for all $\boldsymbol{\theta} \in \Theta$,
- (iii) $\hat{X}_{\mathbf{D}}^2(\boldsymbol{\theta}_0) = \hat{W}_{\mathbf{D}}(\boldsymbol{\theta}_0)$ a.s.,
- (iv) $\hat{G}_{\mathbf{D}}^2(\boldsymbol{\theta}_0) = \hat{X}_{\mathbf{D}}^2(\boldsymbol{\theta}_0) + o_p(1)$ under H_0 .

4 Composite Hypotheses

We first derive the asymptotic distribution of $\hat{\boldsymbol{\Phi}}_{\mathbf{D}}(\tilde{\boldsymbol{\theta}})$ for any \sqrt{n} -consistent estimator $\tilde{\boldsymbol{\theta}}$, which is essential to justify the χ^2 tests for composite hypotheses.

Assumption 2. $\boldsymbol{\theta}_0$ is an interior point of Θ , which is a compact subset of \mathbb{R}^p . $F_{Y|\mathbf{X},\boldsymbol{\theta}}(y | \mathbf{X})$ is continuously differentiable on Θ for all $y \in \mathbb{R}$ a.s.

Define

$$\mathbf{B}_{\mathbf{D}}(\boldsymbol{\theta}) = \text{diag}^{-1/2}(\mathbf{v} \otimes \mathbf{q}) \cdot \mathbb{E}(\boldsymbol{\tau}_{\boldsymbol{\theta}}(\mathbf{X}_1) \otimes \boldsymbol{\zeta}_1),$$

where $\boldsymbol{\tau}_\theta(\mathbf{X}) = (\tau_{\theta_1}(\mathbf{X})', \dots, \tau_{\theta_L}(\mathbf{X})')'$ and

$$\tau_{\theta_\ell}(\mathbf{X}) = \frac{\partial}{\partial \boldsymbol{\theta}} F_{Y|\mathbf{X},\theta}(F_{Y|\mathbf{X},\theta}^{-1}(\vartheta_\ell | \mathbf{X}) | \mathbf{X}) - \frac{\partial}{\partial \boldsymbol{\theta}} F_{Y|\mathbf{X},\theta}(F_{Y|\mathbf{X},\theta}^{-1}(\vartheta_{\ell-1} | \mathbf{X}) | \mathbf{X}).$$

Theorem 2. *Let Assumptions 1 and 2 be satisfied, and $\tilde{\boldsymbol{\theta}} = \boldsymbol{\theta}_0 + O_p(n^{-1/2})$. Then, under H_0 ,*

$$\hat{\boldsymbol{\Phi}}_D(\tilde{\boldsymbol{\theta}}) = \hat{\boldsymbol{\Phi}}_D(\boldsymbol{\theta}_0) - \mathbf{B}_D(\boldsymbol{\theta}_0)\sqrt{n}(\tilde{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) + o_p(1).$$

Most \sqrt{n} -consistent estimators of $\boldsymbol{\theta}_0$ satisfy the following asymptotic representation.

Assumption 3. $\tilde{\boldsymbol{\theta}}$ satisfies $\tilde{\boldsymbol{\theta}} = \boldsymbol{\theta}_0 + n^{-1} \sum_{i=1}^n \ell_{\boldsymbol{\theta}_0}(Y_i, \mathbf{X}_i) + o_p(n^{-1/2})$, where $\ell_{\boldsymbol{\theta}}: \mathbb{R}^{1+k} \rightarrow \mathbb{R}^p$ is such that, for every $\boldsymbol{\theta} \in \Theta$,

$$\int_{\mathbb{R}} \ell_{\boldsymbol{\theta}}(y, \mathbf{X}) F_{Y|\mathbf{X},\theta}(dy | \mathbf{X}) = \mathbf{0} \text{ a.s.} \quad \text{and} \quad \left\| \int_{\mathbb{R}} \ell_{\boldsymbol{\theta}}(y, \mathbf{X}) \ell'_{\boldsymbol{\theta}}(y, \mathbf{X}) F_{Y|\mathbf{X},\theta}(dy | \mathbf{X}) \right\| < \infty \text{ a.s.}$$

The asymptotic distribution of $\hat{\boldsymbol{\Phi}}_D(\tilde{\boldsymbol{\theta}})$, with $\tilde{\boldsymbol{\theta}}$ satisfying Assumption 3, follows from Theorem 2. Define

$$\boldsymbol{\Omega}_{D,\tilde{\boldsymbol{\theta}}} = \mathbf{Q}_U + \mathbf{B}_D(\boldsymbol{\theta}_0)\mathbf{L}(\boldsymbol{\theta}_0)\mathbf{B}'_D(\boldsymbol{\theta}_0) - \mathbf{B}_D(\boldsymbol{\theta})\boldsymbol{\Upsilon}_D(\boldsymbol{\theta}_0) - \boldsymbol{\Upsilon}'_D(\boldsymbol{\theta}_0)\mathbf{B}'_D(\boldsymbol{\theta}),$$

with

$$\mathbf{L}(\boldsymbol{\theta}) = \mathbb{E}(\ell_{\boldsymbol{\theta}}(Y, \mathbf{X})\ell'_{\boldsymbol{\theta}}(Y, \mathbf{X})), \quad \boldsymbol{\Upsilon}_D(\boldsymbol{\theta}) = \text{diag}^{-1/2}(\mathbf{v} \otimes \mathbf{q}) \mathbb{E}(((\boldsymbol{\eta}_1(\boldsymbol{\theta}) - \mathbf{v}) \otimes \boldsymbol{\zeta}_1)\ell'_{\boldsymbol{\theta}}(Y, \mathbf{X})).$$

Theorem 3. *Let Assumptions 1 and 2 be satisfied and $\tilde{\boldsymbol{\theta}}$ be as in Assumption 3. Then, under H_0 , $\hat{\boldsymbol{\Phi}}_D(\tilde{\boldsymbol{\theta}}) \xrightarrow{d} \mathcal{N}_{LJ}(\mathbf{0}, \boldsymbol{\Omega}_{D,\tilde{\boldsymbol{\theta}}})$.*

In order to obtain the trinity of tests under composite hypotheses, we must estimate $\boldsymbol{\theta}_0$ efficiently using the grouped data. For a given $\boldsymbol{\theta}_* \in \Theta$, the vector of observed frequencies $\hat{\mathbf{O}}_{D(\boldsymbol{\theta}_*)}$ is distributed, conditional on $\{\mathbf{X}_i\}_{i=1}^n$, as a multinomial random vector with mean $\mathbb{E}(\hat{\mathbf{O}}_{D(\boldsymbol{\theta}_*)} | \{\mathbf{X}_i\}_{i=1}^n) = \hat{\boldsymbol{\pi}}_{D(\boldsymbol{\theta}_*)}(\boldsymbol{\theta}_0)$ under H_0 , with

$$\hat{\boldsymbol{\pi}}_{D(\boldsymbol{\theta}_*)}(\boldsymbol{\theta}) = \left(\hat{\boldsymbol{\pi}}_{D_{11}(\boldsymbol{\theta}_*)}(\boldsymbol{\theta}), \dots, \hat{\boldsymbol{\pi}}_{D_{1J}(\boldsymbol{\theta}_*)}(\boldsymbol{\theta}), \dots, \hat{\boldsymbol{\pi}}_{D_{L1}(\boldsymbol{\theta}_*)}(\boldsymbol{\theta}), \dots, \hat{\boldsymbol{\pi}}_{D_{LJ}(\boldsymbol{\theta}_*)}(\boldsymbol{\theta}) \right)'$$

and

$$\hat{\boldsymbol{\pi}}_{D_{\ell j}(\boldsymbol{\theta}_*)}(\boldsymbol{\theta}) = \frac{1}{n} \sum_{i=1}^n \zeta_{ji} \int_{F_{Y|\mathbf{X},\theta_*}^{-1}(\vartheta_{\ell-1} | \mathbf{X}_i)}^{F_{Y|\mathbf{X},\theta_*}^{-1}(\vartheta_\ell | \mathbf{X}_i)} F_{Y|\mathbf{X},\theta}(d\vartheta | \mathbf{X}_i).$$

Therefore, the conditional log-likelihood of $\boldsymbol{\theta}$ for given observed data $\hat{\mathbf{O}}_{D(\boldsymbol{\theta}_*)}$ is

$$\mathcal{L}^c(\boldsymbol{\theta} \mid \hat{\mathbf{O}}_{D(\boldsymbol{\theta}_*)}) \propto \sum_{\ell=1}^L \sum_{j=1}^J \hat{O}_{D_{\ell j}(\boldsymbol{\theta}_*)} \ln \hat{\pi}_{D_{\ell j}(\boldsymbol{\theta}_*)}(\boldsymbol{\theta}),$$

and the conditional MLE, $\hat{\boldsymbol{\theta}}_{D(\boldsymbol{\theta}_*)}$, maximizes it over $\boldsymbol{\Theta}$. The next theorem provides the asymptotic distribution of the infeasible MLE, $\hat{\boldsymbol{\theta}}_{D^0}$, assuming the following identifiability condition.

Assumption 4. $\text{rank}(\mathbf{B}_D(\boldsymbol{\theta}_0)) = p$ and, for any $\delta > 0$, there exists an $\varepsilon > 0$ such that

$$\inf_{\|\boldsymbol{\theta} - \boldsymbol{\theta}_0\| \geq \delta} \sum_{\ell=1}^L \sum_{j=1}^J v_{\ell} q_j \log \frac{v_{\ell} q_j}{\pi_{D_{\ell j}^0}(\boldsymbol{\theta})} \geq \varepsilon.$$

These assumptions are standard for the consistency of the MLE in multinomial models (see, e.g., Rao, 2002, Section 5.e). However, Assumption 4 does not hold for certain partitions.

Remark 2. Consider $Y = \mu(\mathbf{X}, \boldsymbol{\beta}_0) + \sigma_0 \varepsilon$, where $\boldsymbol{\theta}_0 = (\boldsymbol{\beta}_0', \sigma_0)' \in \mathbb{R}^{p-1} \times \mathbb{R}^+$, $\mu(\mathbf{X}, \cdot)$ is differentiable on \mathbb{R}^{p-1} a.s., $\mathbb{E}[\partial \mu(\mathbf{X}, \boldsymbol{\beta}_0) / \partial \boldsymbol{\beta}]$ exists, $\mathbb{E}[\varepsilon] = 0$, $\text{Var}[\varepsilon] = 1$, ε is independent of \mathbf{X} and admits a Lebesgue density. If $L = 2$ and μ has an intercept, then $\text{rank}(\mathbf{B}_D(\boldsymbol{\theta})) < p$ for all $\boldsymbol{\theta} \in \mathbb{R}^p$. In general, necessary rank and order conditions for $\text{rank}(\mathbf{B}_D(\boldsymbol{\theta}_0)) = p$ are $\mathbb{E}[\partial \mu(\mathbf{X}, \boldsymbol{\beta}_0) / \partial \boldsymbol{\beta} \otimes \boldsymbol{\zeta}_1] = p - 1$ and $J \geq p - 1$, respectively.

For the next theorem define $\boldsymbol{\Sigma}_D(\boldsymbol{\theta}) = \mathbf{B}'_D(\boldsymbol{\theta}) \mathbf{B}_D(\boldsymbol{\theta})$.

Theorem 4. *Let Assumptions 1, 2, and 4 be satisfied. Then, under H_0 ,*

$$\sqrt{n} (\hat{\boldsymbol{\theta}}_{D^0} - \boldsymbol{\theta}_0) \xrightarrow{d} \mathcal{N}_p(\mathbf{0}, \boldsymbol{\Sigma}_D^{-1}(\boldsymbol{\theta}_0)).$$

We can use a feasible asymptotically equally efficient one-step ahead Gauss-Newton estimator starting from any preliminary \sqrt{n} -consistent estimator $\tilde{\boldsymbol{\theta}}$, i.e.,

$$\hat{\boldsymbol{\theta}}_{D(\tilde{\boldsymbol{\theta}})}^{(1)} = \tilde{\boldsymbol{\theta}} + n^{-1/2} \hat{\boldsymbol{\Sigma}}_D^{-1}(\tilde{\boldsymbol{\theta}}) \hat{\mathbf{B}}'_D(\tilde{\boldsymbol{\theta}}) \hat{\boldsymbol{\Phi}}_D(\tilde{\boldsymbol{\theta}}), \quad (6)$$

where $\hat{\boldsymbol{\Sigma}}_D(\boldsymbol{\theta}) = \hat{\mathbf{B}}'_D(\boldsymbol{\theta}) \hat{\mathbf{B}}_D(\boldsymbol{\theta})$ and $\hat{\mathbf{B}}_D(\boldsymbol{\theta}) = \text{diag}^{-1/2}(\mathbf{v} \otimes \hat{\mathbf{q}}) n^{-1} \sum_{i=1}^n \boldsymbol{\tau}_{\boldsymbol{\theta}}(\mathbf{X}_i) \otimes \boldsymbol{\zeta}_i$. Higher order efficiency gains can be achieved by iterating $\hat{\boldsymbol{\theta}}_{D(\tilde{\boldsymbol{\theta}})}^{(1)}$ (Robinson, 1988).

Corollary 1. *Under the conditions in Theorem 4, for any $\tilde{\boldsymbol{\theta}} = \boldsymbol{\theta}_0 + O_p(n^{-1/2})$,*

$$\sqrt{n} (\hat{\boldsymbol{\theta}}_{D(\tilde{\boldsymbol{\theta}})}^{(1)} - \boldsymbol{\theta}_0) \xrightarrow{d} \mathcal{N}_p(\mathbf{0}, \boldsymbol{\Sigma}_D^{-1}(\boldsymbol{\theta}_0)).$$

In order to obtain the asymptotic variance of $\hat{\Phi}_D(\hat{\theta}_{D^0})$ and $\hat{\Phi}_D(\hat{\theta}_{D(\tilde{\theta})}^{(1)})$, note that in this case $\mathbf{L}(\boldsymbol{\theta}) = \boldsymbol{\Sigma}_D^{-1}(\boldsymbol{\theta})$ for all $\boldsymbol{\theta} \in \Theta$, and $\Upsilon_D(\boldsymbol{\theta}_0)\mathbf{B}'_D(\boldsymbol{\theta}) = \mathbf{B}_D(\boldsymbol{\theta})\boldsymbol{\Sigma}_D^{-1}(\boldsymbol{\theta})\mathbf{B}'_D(\boldsymbol{\theta})$. Hence,

$$\hat{\Phi}_D(\hat{\theta}_{D(\tilde{\theta})}^{(1)}) \xrightarrow{d} \mathcal{N}_{JL}(\mathbf{0}, \boldsymbol{\Omega}_{D, \hat{\theta}_{D^0}}),$$

where $\boldsymbol{\Omega}_{D, \hat{\theta}_{D^0}} = \mathbf{Q}_U - \mathbf{B}_D(\boldsymbol{\theta}_0)\boldsymbol{\Sigma}_D^{-1}(\boldsymbol{\theta}_0)\mathbf{B}'_D(\boldsymbol{\theta}_0)$, which is idempotent with rank $J(L-1) - p$.

The trinity of tests in the composite case is derived from the results in the previous section. The LM statistic is identical to the X^2 by (ii) of Theorem 1, i.e., $\widehat{LM}_D(\hat{\theta}_{D^0}) = \hat{X}_D^2(\hat{\theta}_{D^0})$ a.s. The Wald statistic, $\hat{W}_D(\hat{\theta}_{D^0})$, and the LR statistic, $\hat{G}_D^2(\hat{\theta}_{D^0})$, have the same asymptotic distribution as $\hat{X}_D^2(\hat{\theta}_{D^0})$.

Theorem 5. *Let Assumptions 1, 2, and 4 be satisfied. Then, under H_0 , $\hat{X}_D^2(\hat{\theta}_{D^0})$, $\hat{G}_D^2(\hat{\theta}_{D^0})$, and $\hat{W}_D(\hat{\theta}_{D^0})$ are asymptotically distributed as a $\chi_{J(L-1)-p}^2$. The asymptotic distribution does not change when $\hat{\theta}_{D^0}$ is replaced by $\hat{\theta}_{D(\tilde{\theta})}^{(1)}$ for any $\tilde{\boldsymbol{\theta}} = \boldsymbol{\theta}_0 + O_p(n^{-1/2})$.*

A sensible starting estimator in (6) is the MLE based on the raw data $\{Y_i, \mathbf{X}_i\}_{i=1}^n$,

$$\hat{\boldsymbol{\theta}}^{(0)} = \arg \max_{\boldsymbol{\theta} \in \Theta} \sum_{i=1}^n \log f_{Y|\mathbf{X}, \boldsymbol{\theta}}(Y_i | \mathbf{X}_i),$$

where $f_{Y|\mathbf{X}, \boldsymbol{\theta}}(y | \mathbf{X}) dy = F_{Y|\mathbf{X}, \boldsymbol{\theta}}(dy | \mathbf{X})$. In this case, assuming that $f_{Y|\mathbf{X}, \boldsymbol{\theta}}(y | \mathbf{X})$ is twice continuously differentiable on Θ a.s. for all $y \in \mathbb{R}$ and other regularity conditions,

$$\ell_{\boldsymbol{\theta}}(Y, \mathbf{X}) = \mathbb{I}^{-1}(\boldsymbol{\theta}) \frac{\partial}{\partial \boldsymbol{\theta}} \ln f_{Y|\mathbf{X}, \boldsymbol{\theta}}(Y | \mathbf{X}), \quad (7)$$

where

$$\mathbb{I}(\boldsymbol{\theta}) = -\mathbb{E} \left(\frac{\partial^2}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}'} \ln f_{Y|\mathbf{X}, \boldsymbol{\theta}}(Y | \mathbf{X}) \right)$$

is taken by granted to be positive definite uniformly in Θ , and $\mathbb{I}(\boldsymbol{\theta}_0)$ is the conditional information matrix of $\{Y_i\}_{i=1}^n$ given $\{\mathbf{X}_i\}_{i=1}^n$ under H_0 . Therefore, $\mathbf{L}(\boldsymbol{\theta}) = \mathbb{I}^{-1}(\boldsymbol{\theta})$, and, under H_0 ,

$$\mathbb{E} \left(\frac{\partial}{\partial \boldsymbol{\theta}} \ln f_{Y|\mathbf{X}, \boldsymbol{\theta}}(Y | \mathbf{X}) \Big|_{\boldsymbol{\theta}=\boldsymbol{\theta}_0} \right) = \mathbf{0}.$$

Notice that, in this case, we can write

$$\mathbf{B}_D(\boldsymbol{\theta}_0) = \text{diag}^{-1/2}(\mathbf{v} \otimes \mathbf{q}) \cdot \mathbb{E} \left((\boldsymbol{\eta}_1(\boldsymbol{\theta}_0) \otimes \boldsymbol{\zeta}_1) \frac{\partial}{\partial \boldsymbol{\theta}'} \ln f_{Y|\mathbf{X}, \boldsymbol{\theta}}(Y | \mathbf{X}) \Big|_{\boldsymbol{\theta}=\boldsymbol{\theta}_0} \right).$$

Therefore, $\Upsilon_D \mathbf{B}'_D(\boldsymbol{\theta}_0) = \mathbf{B}_D(\boldsymbol{\theta}_0) \mathbb{I}^{-1}(\boldsymbol{\theta}_0) \mathbf{B}'_D(\boldsymbol{\theta}_0)$, and

$$\text{Avar} \left(\hat{\boldsymbol{\Phi}}_D \left(\hat{\boldsymbol{\theta}}^{(0)} \right) \right) = \boldsymbol{\Omega}_{D, \hat{\boldsymbol{\theta}}^{(0)}} = \mathbf{Q}_U - \mathbf{B}_D(\boldsymbol{\theta}_0) \mathbb{I}^{-1}(\boldsymbol{\theta}_0) \mathbf{B}'_D(\boldsymbol{\theta}_0).$$

The next theorem provides the asymptotic distribution of X^2 based on the raw data MLE $\hat{\boldsymbol{\theta}}^{(0)}$, which depends on the unknown $\boldsymbol{\theta}_0$ and $F_{\mathbf{X}}$. The result is applied to obtain the asymptotic distribution of the corresponding Wald statistic,

$$\hat{W}_D \left(\hat{\boldsymbol{\theta}}^{(0)} \right) = \hat{\boldsymbol{\Phi}}'_D \left(\hat{\boldsymbol{\theta}}^{(0)} \right) \widehat{\text{Avar}}^{-1} \left(\hat{\boldsymbol{\Phi}}_D \left(\hat{\boldsymbol{\theta}}^{(0)} \right) \right) \hat{\boldsymbol{\Phi}}_D \left(\hat{\boldsymbol{\theta}}^{(0)} \right)$$

with

$$\widehat{\text{Avar}} \left(\hat{\boldsymbol{\Phi}}_D \left(\hat{\boldsymbol{\theta}}^{(0)} \right) \right) = \mathbf{Q}_U - \hat{\mathbf{B}}_D \left(\hat{\boldsymbol{\theta}}^{(0)} \right) \hat{\mathbb{I}}^{-1} \left(\hat{\boldsymbol{\theta}}^{(0)} \right) \hat{\mathbf{B}}'_D \left(\hat{\boldsymbol{\theta}}^{(0)} \right),$$

where

$$\hat{\mathbb{I}}^{-1}(\boldsymbol{\theta}) = -\frac{1}{n} \sum_{i=1}^n \frac{\partial^2}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}'} \ln f_{Y|\mathbf{X}, \boldsymbol{\theta}}(Y_i | \mathbf{X}_i).$$

The theorem extends the results in [Chernoff and Lehmann \(1954\)](#) to the conditional case. Let $\{\lambda_j\}_{j=J(L-1)-p+1}^{J(L-1)}$ be the p roots of the determinantal equation

$$|\mathbf{B}'_D(\boldsymbol{\theta}_0) \mathbf{B}_D(\boldsymbol{\theta}_0) - (1 - \lambda) \mathbb{I}(\boldsymbol{\theta}_0)| = 0,$$

which always satisfy $0 \leq \lambda_i < 1$, and $0 < \lambda_i < 1$ when $\mathbb{I}(\boldsymbol{\theta}_0) - \mathbf{B}'(\boldsymbol{\theta}_0) \mathbf{B}(\boldsymbol{\theta}_0)$ is positive definite.

Assumption 5. $\mathbb{I}(\boldsymbol{\theta}) - \mathbf{B}'(\boldsymbol{\theta}) \mathbf{B}(\boldsymbol{\theta})$ is positive definite for all $\boldsymbol{\theta} \in \Theta$.

Assumption 3'. $\hat{\boldsymbol{\theta}}^{(0)}$ satisfies Assumption 3 with $\ell_{\boldsymbol{\theta}}$ defined in (7).

Theorem 6. *Let Assumptions 1, 2, and 3' be satisfied. Then, under H_0 ,*

$$\hat{X}_D^2 \left(\hat{\boldsymbol{\theta}}^{(0)} \right) \xrightarrow{d} \sum_{j=1}^{J(L-1)-p} Z_j^2 + \sum_{j=J(L-1)-p+1}^{J(L-1)} \lambda_j Z_j^2,$$

where Z_j 's are i.i.d. standard normal random variables with $0 \leq \lambda_j < 1$.

Corollary 2. *Let Assumptions 1, 2, 3', and 5 be satisfied. Then, under H_0 , $\hat{W}_D \left(\hat{\boldsymbol{\theta}}^{(0)} \right) \xrightarrow{d} \chi_{J(L-1)}^2$.*

That is, the Wald test based on $\hat{\mathbf{O}}_D \left(\hat{\boldsymbol{\theta}}^{(0)} \right)$ is asymptotically pivotal and invariant to the number of estimated parameters.

5 Sample-Dependent Grouping and Classification Algorithms

5.1 Asymptotics with sample-dependent grouping

The Chernoff and Lehmann (1954) result in the classical case, which provides the asymptotic distribution of the χ^2 evaluated at the raw data MLE, was extended to partitions with random cells by Roy (1956) and Watson (1957). They showed that the asymptotic behavior of χ^2 statistics is unaffected when class boundaries depend on data. See Chibisov (1971), Moore (1971), and Pollard (1979) for further developments. Andrews (1988b) extends the result to the Wald's statistic for conditional CDF's specification testing.

In view of (4), the asymptotic distribution of the χ^2 statistics is non-pivotal when we use a sample-dependent \mathbf{U} . Since $V = F_{Y|\mathbf{X}}(Y | \mathbf{X})$ is uniformly distributed on $[0, 1]$ independently of \mathbf{X} , it is sensible to use partitions $\mathbf{U} = \{(\vartheta_{\ell-1}, \vartheta_{\ell}]\}_{\ell=1}^L$ with $\vartheta_{\ell} = \ell/L$. These partitions, called balanced, are easy to use and, in the classical case, lead to unbiased tests (Cohen and Sackrowitz, 1975). In this section, we show that the asymptotic distribution of the χ^2 statistics does not change when the explanatory variable observations' grouping is sample dependent.

Using Pollard (1979) notation, the cells A_j in the partitions \mathbf{A} are chosen from a class \mathcal{A} of measurable cells. The partitions of \mathbb{R}^k into cells in a class \mathcal{A} correspond to elements of the class

$$\mathcal{H} = \left\{ \mathbf{A} \in \mathcal{A}^J : A_1, \dots, A_J \text{ disjoint and } \cup_{j=1}^J A_j = \mathbb{R}^k \right\}.$$

Equip \mathcal{H} with its product topology and Borel structure. A partition of \mathbb{R}^k into data-dependent cells $\hat{A}_1, \dots, \hat{A}_J$ determines a map $\hat{\mathbf{A}}$ from the underlying probability space into \mathcal{H} . Call $\hat{\mathbf{A}} = \{\hat{A}_j\}_{j=1}^J$ a random element of \mathcal{H} if it is a measurable map. We assume that the random cells $\hat{\mathbf{A}}$ converge in probability (in the sense of the topology on \mathcal{H}) to a set of fixed cells $\mathbf{A} \in \mathcal{H}$; that is, $F_{\mathbf{X}} \left\{ \hat{A}_j \cup A_j \setminus \hat{A}_j \cap A_j \right\} \xrightarrow{P} 0$ for $j = 1, \dots, J$, where $F_{\mathbf{X}} \{S\} = \int_{x \in S} dF_{\mathbf{X}}$.

Assumption 6. \mathcal{A} is a Vapnik-Chervonenkis (VC) class. $\hat{\mathbf{A}} = \{\hat{A}_j\}_{j=1}^J$ is a sequence of random elements of \mathcal{H} converging in probability to a fixed $\mathbf{A} = \{A_j\}_{j=1}^J \in \mathcal{H}$ such that for each component A_j , $q_j > 0$, $j = 1, \dots, J$.

Andrews (1988a,b) points out that the set of VC classes includes most reasonable partitions, such as those based on rectangles, regular cylinders, clusters obtained using k -means or any other algorithm that yields a finite number of straight edges. See (Andrews, 1988a,b) for further discussion. The next theorem states that using sample-dependent partitions

$$\hat{\mathbf{D}}(\boldsymbol{\theta}) = \left\{ \hat{D}_{\ell_j}(\boldsymbol{\theta}) : \ell = 1, \dots, L, j = 1, \dots, J \right\}$$

with

$$\hat{D}_{\ell_j}(\boldsymbol{\theta}) = \left\{ (y, \mathbf{x}) \in \mathbb{R}^{1+k} : F_{Y|\mathbf{X},\boldsymbol{\theta}}(y | \mathbf{x}) \in U_{\ell}, \mathbf{x} \in \hat{A}_j \right\},$$

rather than $\mathbf{D}(\boldsymbol{\theta})$, does not have any effect on the asymptotic distribution of the different statistics under the null.

Theorem 7. *Let Assumptions 1, 2, 3, and 6 hold. Then, under H_0 , $\hat{X}_{\hat{\mathbf{D}}}^2(\hat{\boldsymbol{\theta}}^{(1)})$, $\hat{W}_{\hat{\mathbf{D}}}(\hat{\boldsymbol{\theta}}^{(1)})$, and $\hat{G}_{\hat{\mathbf{D}}}^2(\hat{\boldsymbol{\theta}}^{(1)})$ share the same limiting distribution with $\hat{X}_{\mathbf{D}}^2(\hat{\boldsymbol{\theta}}^{(1)})$. If, in addition, Assumption 3' holds, then $\hat{W}_{\hat{\mathbf{D}}}(\hat{\boldsymbol{\theta}}^{(0)})$ share the same limiting distribution with $\hat{W}_{\mathbf{D}}(\hat{\boldsymbol{\theta}}^{(0)})$.*

5.2 Algorithms

In this section we propose a partitioning algorithm allowing to control the number of points in its cells and, hence, amenable to use with large k .

Gessaman (1970) has proposed a very simple, deterministic rule to obtain cells \hat{A}_j containing approximately the same number of points $\{\mathbf{X}_i\}_{i=1}^n$. Let an integer $T \geq 2$ such that $n \geq T^k$ be given, and let $\mathbf{X} = (X_1, \dots, X_k)'$. Start by splitting $\{\mathbf{X}_i\}_{i=1}^n$ into T sets having the same number of points, except for perhaps the boundary sets, using hyperplanes perpendicular to the axis of X_1 . If $k = 1$, the process terminates here. If $k > 1$, one can proceed recursively by next partitioning each of the obtained T cylindrical sets into T additional subsets by cutting along the axis of X_2 and making sure that the resulting subsets again have roughly the same number of points. The procedure yields $\hat{\mathbf{A}} = \{\hat{A}_j\}_{j=1}^J$ with $J = T^k$ so that $n \geq T^k$ is required to make sure that the cells are non-empty. It can be shown that

$$\max_{j=1, \dots, J} \hat{N}_j - \min_{j=1, \dots, J} \hat{N}_j \leq 1, \quad (8)$$

where $\hat{N}_j = \sum_{i=1}^n \mathbb{1}_{\{\mathbf{X}_i \in \hat{A}_j\}}$ is the number of points in \hat{A}_j , $j = 1, \dots, J$. One can expect cells with approximately the same number of points to improve finite sample behavior. However, even with $T = L = 2$, the total number of cells, $LJ = 2^{k+1}$, is rapidly increasing in k and demands an alternative approach for relatively large values of k .

We propose a Random Tree Partition (RTP) resulting into significantly fewer than T^k cells while preserving the possibility of controlling the number of points per cell. Let $T \geq 2$ and $r \geq 1$ be integers. Let $\mathcal{G} \leftarrow (1, \dots, 1, 2, \dots, 2, \dots, k, \dots, k)'$ be a $kr \times 1$ vector repeating each of $1, \dots, k$ for r times.

0. Start with a tree containing a single node (the root) associated with $A \leftarrow \mathbb{R}^k$.
1. Uniformly at random select $j \in \mathcal{G}$.

2. Using $T-1$ hyperplanes perpendicular to the X_j axis, partition A into T cells $\hat{A}_1, \dots, \hat{A}_T$ such that each cell contains approximately the same number of points $\{\mathbf{X}_i\}_{i=1}^n$. Split the current node associated with A by adding T child nodes associated with $\hat{A}_1, \dots, \hat{A}_T$.
3. Remove one instance of j from \mathcal{G} .
4. If \mathcal{G} has no elements left, the procedure terminates here. Otherwise, select a terminal node associated with \hat{A} that contains the highest number of points and repeat Step 1 with $A \leftarrow \hat{A}$.

Each $j \in \mathcal{G}$ induces a single split, resulting in kr splits in total. As we start with a single-node tree and each split increases the number of terminal nodes by $T - 1$, the final tree has $J = 1 + kr(T - 1)$ terminal nodes. To be able to perform all kr splits and avoid empty terminal nodes, we require $n \geq 1 + kr(T - 1)$ hereafter. It can be shown that

$$\max_{j=1, \dots, J} \hat{N}_j - \min_{j=1, \dots, J} \hat{N}_j \leq 1 + (T - 1) \min_{j=1, \dots, J} \hat{N}_j \iff \frac{\max_{j=1, \dots, J} \hat{N}_j}{\min_{j=1, \dots, J} \hat{N}_j} \leq T + \frac{1}{\min_{j=1, \dots, J} \hat{N}_j}.$$

That is, the number of points in any two cells, including those with the largest and smallest number of points, differs by at most a little more than T times. For example, if $T = 2$ and the smallest cell contains 5 points, the largest cell contains up to 11 points. The inequality becomes equality if the terminal nodes are not at the same depth. However, the self-balancing Step 4 guarantees that if $J = 1 + kr(T - 1)$ is a power of T , then all the terminal nodes are at the same depth and (8) holds. It is equivalent to requiring the number of splits, or the length of \mathcal{G} , to be equal to $kr = (T^q - 1)/(T - 1)$ for some $q \in \mathbb{N}$. If the equality does not hold for chosen k and r , it can be forced by instead defining \mathcal{G} as a $(T^q - 1)/(T - 1) \times 1$ vector, where each of $1, \dots, k$ no longer appears the same number of times. For simplicity and as to avoid high values of J with large k , in Section 7 we consider binary trees with median splits so that $T = 2$, and the number of cells equals $J = 1 + kr$.

We emphasize that, for $n \geq J = 1 + kr(T - 1)$, the algorithm guarantees that each \hat{A}_j , $j = 1, \dots, J$, will be nonempty, which is crucial as then $\hat{q}_j > 0$, $j = 1, \dots, J$. As a result, the X^2 and Wald statistics are always computable when using an RTP. On the other hand, G^2 additionally requires that $\hat{O}_{D_{\ell_j}(\boldsymbol{\theta})} > 0$ for all $j = 1, \dots, J$ and $\ell = 1, \dots, L$. The latter, however, cannot be guaranteed by *any* procedure without making \mathbf{U} data dependent, unless $L = J = JL = 1$.

6 Power

Consider contiguous alternatives, as in [Delgado and Stute \(2008\)](#),

$$H_{1n} : \frac{F_{Y|X}(dy | \mathbf{X})}{F_{Y|X, \theta_0}(dy | \mathbf{X})} = 1 + \frac{t_{n\theta_0}(y, \mathbf{X})}{\sqrt{n}} \text{ a.s. for some } \theta_0 \in \Theta \text{ and all } y \in \mathbb{R},$$

where $t_{n\theta}$ and t_θ are such that

$$\int_{\mathbb{R}} t_{n\theta}(y, \mathbf{X}) F_{Y|X, \theta}(dy | \mathbf{X}) = 0, \quad \int_{\mathbb{R}} (t_{n\theta} - t_\theta)^2(y, \mathbf{X}) F_{Y|X, \theta}(dy | \mathbf{X}) = o(1) \text{ as } n \rightarrow \infty \text{ a.s.}$$

This allows modeling departures from H_0 that are proper conditional CDF's. See [Delgado and Stute \(2008\)](#) for examples and discussion. Let us consider first the limiting distribution of $\hat{\Phi}_{\hat{D}}(\tilde{\theta})$ under H_{1n} for any \sqrt{n} -consistent estimator $\tilde{\theta}$. The next theorem is essential to derive the limiting distribution of the statistics under H_{1n} . Define $\mathbf{T}_D^{(1)} = [\mathbf{T}_D^{(1)}]_{\ell_j}$, $\ell = 1, \dots, L$, $j = 1, \dots, J$, with

$$[\mathbf{T}_D^{(1)}]_{\ell_j} = \frac{1}{v_{\ell} q_j} \int_{x \in A_j} \int_{F_{Y|X, \theta_0}^{-1}(\vartheta_{\ell-1} | \mathbf{x})}^{F_{Y|X, \theta_0}^{-1}(\vartheta_{\ell} | \mathbf{x})} t_{\theta_0}(y, \mathbf{x}) F_{Y|X, \theta_0}(dy | \mathbf{x}) F_X(d\mathbf{x}).$$

Theorem 8. *Let Assumptions 1, 2, 3, and 6. Then, under H_{1n} ,*

$$\hat{\Phi}_{\hat{D}}(\tilde{\theta}) = \hat{\Phi}_D(\theta_0) - \mathbf{B}(\theta_0) \sqrt{n} (\tilde{\theta} - \theta_0) + \mathbf{T}_D^{(1)} + o_p(1).$$

Under H_{1n} , the ℓ_{θ_0} term in Assumption 3 is not centered anymore (see [Behnen and Neuhaus, 1975](#)). Now $\sqrt{n}(\tilde{\theta} - \theta_0)$ has asymptotic mean

$$\boldsymbol{\mu} = \int_{\mathbb{R}^{1+k}} \ell_{\theta_0}(y, \mathbf{x}) t_{\theta_0}(y, \mathbf{x}) F_{Y|X, \theta_0}(dy | \mathbf{x}) F_X(d\mathbf{x}).$$

This results in the additional shift $\mathbf{T}_D^{(2)} = \mathbf{B}_D(\theta_0) \cdot \boldsymbol{\mu}$. Define $\mathbf{T}_D = \mathbf{T}_D^{(1)} - \mathbf{T}_D^{(2)}$. Then $\sqrt{n} \hat{\Phi}_{\hat{D}}(\tilde{\theta}) - \mathbf{T}_D$ under H_{1n} has the same asymptotic distribution as $\sqrt{n} \hat{\Phi}_{\hat{D}}(\tilde{\theta})$ under H_0 .

The next result provides the limiting distribution of the different statistics under H_{1n} . Henceforth, $\chi_N^2(\Lambda) \stackrel{d}{=} \sum_{j=1}^N (Z_j + \omega_j)^2$ is a non-central chi-squared random variable with N degrees of freedom and non-centrality parameter $\Lambda = \sum_{j=1}^N \omega_j^2$.

Corollary 3. *Let H_{1n} hold. If Assumptions 1, 2, 4, and 6 are satisfied, then $\hat{X}_D^2(\hat{\theta}_D^{(1)})$, $\hat{W}_D(\hat{\theta}_D^{(1)})$, and $\hat{G}_D^2(\hat{\theta}_D^{(1)})$ are asymptotically distributed as a $\chi_{J(L-1)-p}^2(\mathbf{T}_D' \mathbf{T}_D)$. Under Assumptions 1, 3', and 6, $\hat{W}_D(\hat{\theta}_D^{(0)}) \xrightarrow{d} \chi_{J(L-1)}^2(\mathbf{T}_D' \mathbf{T}_D)$.*

7 Monte Carlo

To evaluate the finite sample performance of the Pearson test we consider the null hypothesis

$$H_0 : Y \mid \mathbf{X} \sim \mathcal{N}(\mu(\mathbf{X}), \sigma^2),$$

where the regression model is multivariate linear, i.e., $\mu(\mathbf{X}) = \beta_{00} + \mathbf{X}'\boldsymbol{\beta}_0$, with $\boldsymbol{\beta}_0 = (\beta_{10}, \dots, \beta_{k0})'$. We set $\beta_{00} = \dots = \beta_{k0} = \sigma^2 = 1$. Data consists of $\{Y_i, \mathbf{X}_i\}_{i=1}^n$, where $\{\mathbf{X}_i\}_{i=1}^n$ are i.i.d. observations of \mathbf{X} , a $k \times 1$ vector of independent random variables uniformly distributed on $[0, 1]$, and

$$Y_i = \mu(\mathbf{X}_i) + \delta(\mathbf{X}_i) + \sigma(\mathbf{X}_i) \cdot \varepsilon_i,$$

where $\{\varepsilon_i\}_{i=1}^n$ are i.i.d. with mean zero and variance one,

$$\delta(\mathbf{x}) = a \cdot \sum_{i=1}^k \ln(x_i), \quad \text{and} \quad \sigma(\mathbf{x}) = \exp\left(-b \cdot \sum_{i=1}^k x_i\right) \frac{\sqrt{(2b)^k e^{2bk}}}{\sqrt{(e^{2b} - 1)^k}},$$

where $\sigma(\cdot)$ is such that $\text{Var}[\sigma(\mathbf{X})\varepsilon_i] = 1$ and $a, b \geq 0$. Thus, under H_0 , $\{\varepsilon_i\}_{i=1}^n$ are i.i.d. standard normal and $a = b = 0$. Let $\mu_k = \mathbb{E}[\varepsilon_i^k]$. We consider the following types of departures from H_0 .

1. From linear regression specification: $a = 1, b = 0$, and $\{\varepsilon_i\}_{i=1}^n$ are i.i.d. $\mathcal{N}(0, 1)$.
2. From conditional homoskedasticity: $a = 0, b = \frac{2}{3}$, and $\{\varepsilon_i\}_{i=1}^n$ are i.i.d. $\mathcal{N}(0, 1)$.
3. From conditional symmetry: $a = b = 0$, and $\{\varepsilon_i\}_{i=1}^n$ are i.i.d. and follow a skewed generalized t (SGT) distribution proposed by [Theodossiou \(1998\)](#), with parameters (λ, p, q) . This allows to generate asymmetric distributions (with $\mu_3 > 0$) without excess of kurtosis ($\mu_4 = 0$). We report results of mild (SGT_{1/3}) and strong (SGT_{1/3}) asymmetry using the following parameter values.

(i) $(\lambda, p, q) = (0.2537088, 2.976928, 3.840017)$ with $\mu_1 = 0, \mu_2 = 1, \mu_3 = 1/3, \mu_4 = 3$.

(ii) $(\lambda, p, q) = (0.9988789, 3.273291, 8.60737)$ with $\mu_1 = 0, \mu_2 = 1, \mu_3 = 2/3, \mu_4 = 3$.

4. From conditional mesokurtosis: $a = b = 0$, and standardized $\{\varepsilon_i\}_{i=1}^n$ are i.i.d. and follow a t distribution. We provide two cases, t_5 and $t_{2,1}$, with the latter one having heavier tails and, unlike t_5 , infinite variance.

(i) $\sqrt{5/3} \cdot \varepsilon_i \sim t_5, i = 1, \dots, n$, with $\mu_1 = 0, \mu_2 = 1, \mu_3 = 0, \mu_4 = 9$.

(ii) $\sqrt{21} \cdot \varepsilon \sim t_{2,1}, i = 1, \dots, n$, with $\mu_1 = 0, \mu_2 = 1, \mu_3$ undefined, $\mu_4 = \infty$.

We have run simulations for tests based on $\hat{X}_D^2(\hat{\boldsymbol{\theta}}^{(1)})$, $\hat{G}_D^2(\hat{\boldsymbol{\theta}}^{(1)})$, $\hat{W}_D(\hat{\boldsymbol{\theta}}^{(1)})$, and $\hat{W}_D(\hat{\boldsymbol{\theta}}^{(0)})$. We use $\hat{\boldsymbol{\theta}}^{(1)}$ iterated until convergence as it significantly improves the tests' size accuracy. We also compare these tests with the omnibus conditional Kolmogorov-Smirnov (KS) test proposed by Andrews (1997), which is based on the difference between the sample joint distribution and its restricted version imposing the conditional CDF specification under H_0 . This test is not pivotal and, hence, is implemented with the assistance of a parametric bootstrap. Results are based on 4000 Monte Carlo iterations, and the KS bootstrap test is based on 2000 resamples. We report results for $n = 50, 100$, and 500.

We present two sets of results. We first consider balanced partitions \mathbf{U} , i.e., partitions with $\vartheta_\ell = \ell/L$, $\ell = 1, \dots, L$, which do not favor, in principle, any alternative. Next we consider unbalanced partitions, with small cells on the tails. This has been proposed in the classical literature, to check model specification of marginal distribution, in order to favor alternatives with heavy tails (see, e.g., Kallenberg et al., 1985). The power of χ^2 tests, particularly X^2 , improves for any type of alternatives, not only leptokurtic ones. but also the other alternatives considered. In all the tables we consider $k = 1, 5, 10$ to assess the curse of dimensionality effect on the different tests. We employ the RTP algorithm with $T = 2$ to partition \mathbb{R}^k .

We first consider balanced partitions \mathbf{U} with $\vartheta_\ell = \ell/L$, $\ell = 1, \dots, L$, for $L = 3, 6, 12, 24$, where $L = 3, 6$ are combined with $r = 1, 2, 5, 10$, and $L = 12, 24$ are used only with $r = 1$. Additionally, we consider $J = 1$, i.e., without partitioning \mathbb{R}^k at all, with $L = 2, 3$, which cannot be used with the X^2 statistic, because Assumption 4 fails to hold in this case $\text{rank}(\mathbf{B}_D(\boldsymbol{\theta}_0)) < p$ when $L = 2$ or $J = 1 < k + 1$ (see Remark 2). The rejection rates are not provided when $J = 1 + kr > n$.

Recall that G^2 can be computed only when $\hat{O}_{D_{e_j}}(\boldsymbol{\theta}) > 0$ for all $j = 1, \dots, J$ and $\ell = 1, \dots, L$. For a given n , after a certain point, the probability of no empty cells rapidly decreases as JL increases. For example, with equiprobable cells and $n = 100$, it is approximately 98.5% under $LJ = 3 \cdot 5$ and 33.4% under $LJ = 3 \cdot 10$. As a result, G^2 cannot be computed in many cases. When available, the results for G^2 are almost identical to those of X^2 and, hence, they will not be reported. The results for $\hat{W}_D(\hat{\boldsymbol{\theta}}^{(1)})$ are also very similar. Thus, we only report results for $\hat{X}_D^2(\hat{\boldsymbol{\theta}}^{(1)})$ and $\hat{W}_D(\hat{\boldsymbol{\theta}}^{(0)})$, which will be referred to as X^2 and W , respectively.

Table 2 reports the proportion of rejections under H_0 . All the tests exhibit excellent size accuracy. It is worth noticing that the χ^2 tests are almost as accurate as the KS bootstrap test.

The rejection rates under H_1 are reported in Table 3. The performance of all the tests is

L	r	J	k	$n = 50$			$n = 100$			$n = 500$		
				0.01	0.05	0.10	0.01	0.05	0.10	0.01	0.05	0.10
W test using $\hat{\theta}^{(0)}$												
2	–	1	1	0.01	0.03	0.10	0.01	0.06	0.13	0.01	0.04	0.08
3	–	1	1	0.01	0.05	0.11	0.01	0.05	0.11	0.01	0.06	0.10
3	1	2	1	0.01	0.05	0.10	0.01	0.05	0.10	0.01	0.05	0.09
3	2	3	1	0.01	0.04	0.10	0.01	0.05	0.10	0.01	0.05	0.10
3	5	6	1	0.01	0.05	0.09	0.01	0.05	0.10	0.01	0.04	0.10
3	10	11	1	0.01	0.04	0.09	0.01	0.05	0.10	0.01	0.05	0.10
6	1	2	1	0.01	0.05	0.09	0.01	0.05	0.09	0.01	0.04	0.09
6	2	3	1	0.01	0.05	0.09	0.01	0.05	0.10	0.01	0.05	0.09
6	5	6	1	0.00	0.04	0.09	0.01	0.05	0.10	0.01	0.06	0.11
6	10	11	1	0.00	0.04	0.08	0.01	0.04	0.09	0.01	0.05	0.10
12	1	2	1	0.01	0.05	0.10	0.01	0.05	0.10	0.01	0.05	0.10
24	1	2	1	0.01	0.05	0.09	0.01	0.04	0.10	0.01	0.05	0.10
2	–	1	10	0.01	0.03	0.10	0.01	0.07	0.14	0.01	0.04	0.09
3	–	1	10	0.01	0.05	0.11	0.01	0.05	0.10	0.01	0.05	0.09
3	1	11	10	0.01	0.06	0.14	0.01	0.06	0.12	0.01	0.05	0.10
3	2	21	10	0.00	0.05	0.12	0.01	0.05	0.11	0.01	0.06	0.11
3	5	51	10	–	–	–	0.00	0.02	0.07	0.01	0.05	0.10
3	10	101	10	–	–	–	–	–	–	0.01	0.04	0.09
6	1	11	10	0.01	0.04	0.10	0.01	0.06	0.12	0.01	0.05	0.10
6	2	21	10	0.00	0.03	0.09	0.01	0.04	0.10	0.01	0.04	0.09
6	5	51	10	–	–	–	0.00	0.02	0.06	0.01	0.05	0.10
6	10	101	10	–	–	–	–	–	–	0.01	0.04	0.08
12	1	11	10	0.01	0.05	0.11	0.01	0.05	0.10	0.01	0.05	0.10
24	1	11	10	0.01	0.05	0.10	0.01	0.04	0.09	0.01	0.05	0.10
X^2 test using Iterated $\hat{\theta}^{(1)}$												
3	1	2	1	0.01	0.05	0.15	0.01	0.06	0.11	0.01	0.06	0.11
3	2	3	1	0.01	0.05	0.11	0.01	0.05	0.11	0.01	0.05	0.10
3	5	6	1	0.01	0.05	0.10	0.01	0.05	0.11	0.01	0.04	0.09
3	10	11	1	0.00	0.04	0.10	0.01	0.04	0.09	0.01	0.05	0.10
6	1	2	1	0.01	0.05	0.09	0.01	0.04	0.10	0.01	0.05	0.09
6	2	3	1	0.01	0.05	0.09	0.01	0.05	0.10	0.01	0.05	0.10
6	5	6	1	0.01	0.04	0.09	0.01	0.05	0.09	0.01	0.05	0.10
6	10	11	1	0.01	0.04	0.08	0.01	0.05	0.09	0.01	0.05	0.10
12	1	2	1	0.01	0.05	0.09	0.01	0.04	0.09	0.01	0.05	0.10
24	1	2	1	0.01	0.06	0.10	0.01	0.06	0.10	0.01	0.05	0.09
3	1	11	10	0.23	0.30	0.36	0.06	0.13	0.21	0.01	0.06	0.13
3	2	21	10	0.00	0.04	0.11	0.01	0.04	0.09	0.01	0.05	0.10
3	5	51	10	–	–	–	0.00	0.02	0.07	0.01	0.03	0.08
3	10	101	10	–	–	–	–	–	–	0.00	0.04	0.08
6	1	11	10	0.08	0.15	0.23	0.04	0.08	0.15	0.01	0.06	0.11
6	2	21	10	0.00	0.01	0.06	0.01	0.03	0.07	0.01	0.04	0.09
6	5	51	10	–	–	–	0.00	0.01	0.05	0.01	0.04	0.08
6	10	101	10	–	–	–	–	–	–	0.00	0.03	0.07
12	1	11	10	0.06	0.11	0.16	0.03	0.07	0.12	0.01	0.04	0.09
24	1	11	10	0.05	0.08	0.12	0.02	0.06	0.11	0.01	0.05	0.10
Kolmogorov-Smirnov test using $\hat{\theta}^{(0)}$												
–	–	–	1	0.01	0.05	0.09	0.01	0.05	0.10	0.01	0.05	0.10
			5	0.01	0.05	0.10	0.01	0.05	0.11	0.01	0.05	0.10
			10	0.01	0.05	0.10	0.01	0.05	0.10	0.01	0.05	0.11

Table 2: Size accuracy using balanced partitions ($k = 1, 10$).

similar under the nonlinear regression ($a = 1, b = 0$) for $k = 1$, although the X^2 test appears to be more sensitive to the partition choice than W . Both the X^2 and KS tests suffer from the curse of dimensionality going from simple ($k = 1$) to multiple ($k = 5, 10$) regression, while W is more robust to it. The χ^2 tests perform very well under conditional heteroskedasticity for any k , while the KS test exhibits very poor power under this alternative, but the proportion of

L	r	J	k	$a = 1$			$b = 2/3$			SGT $_{1/3}$			SGT $_{2/3}$			t_5			$t_{2.1}$		
				50	100	500	50	100	500	50	100	500	50	100	500	50	100	500	50	100	500
W test using $\hat{\theta}^{(0)}$																					
2	–	1	1	0.04	0.09	0.13	0.04	0.07	0.05	0.06	0.13	0.33	0.15	0.39	0.95	0.07	0.13	0.10	0.27	0.41	0.45
3	–	1	1	0.06	0.08	0.21	0.06	0.07	0.12	0.09	0.16	0.66	0.28	0.54	1.00	0.16	0.29	0.81	0.66	0.91	1.00
3	1	2	1	0.11	0.15	0.59	0.09	0.14	0.57	0.08	0.12	0.50	0.21	0.44	1.00	0.15	0.22	0.75	0.64	0.89	1.00
3	2	3	1	0.13	0.26	0.91	0.07	0.12	0.51	0.07	0.10	0.45	0.17	0.36	0.99	0.13	0.21	0.70	0.60	0.87	1.00
3	5	6	1	0.20	0.44	1.00	0.07	0.11	0.47	0.06	0.09	0.31	0.13	0.26	0.98	0.11	0.17	0.60	0.53	0.82	1.00
3	10	11	1	0.14	0.35	0.99	0.06	0.09	0.36	0.05	0.07	0.24	0.09	0.18	0.92	0.08	0.13	0.48	0.42	0.74	1.00
6	1	2	1	0.10	0.16	0.62	0.10	0.18	0.81	0.07	0.11	0.51	0.22	0.51	1.00	0.18	0.31	0.90	0.69	0.92	1.00
6	2	3	1	0.13	0.23	0.89	0.09	0.16	0.78	0.07	0.10	0.42	0.16	0.41	1.00	0.16	0.27	0.86	0.64	0.91	1.00
6	5	6	1	0.16	0.37	0.99	0.07	0.14	0.69	0.05	0.08	0.29	0.12	0.28	1.00	0.12	0.21	0.77	0.55	0.85	1.00
6	10	11	1	0.12	0.29	0.99	0.05	0.10	0.54	0.04	0.07	0.21	0.08	0.19	0.98	0.10	0.16	0.65	0.48	0.79	1.00
12	1	2	1	0.10	0.15	0.59	0.09	0.15	0.79	0.07	0.10	0.46	0.21	0.53	1.00	0.19	0.29	0.90	0.67	0.92	1.00
24	1	2	1	0.09	0.14	0.53	0.09	0.12	0.70	0.06	0.09	0.35	0.18	0.42	1.00	0.19	0.30	0.89	0.65	0.91	1.00
2	–	1	10	0.06	0.24	0.86	0.03	0.08	0.08	0.04	0.12	0.35	0.09	0.33	0.94	0.05	0.11	0.09	0.13	0.30	0.40
3	–	1	10	0.08	0.22	0.93	0.20	0.59	1.00	0.07	0.13	0.61	0.14	0.39	1.00	0.10	0.21	0.78	0.40	0.81	1.00
3	1	11	10	0.09	0.14	0.74	0.17	0.44	1.00	0.08	0.07	0.22	0.10	0.16	0.90	0.09	0.11	0.45	0.27	0.60	1.00
3	2	21	10	0.07	0.12	0.70	0.10	0.32	1.00	0.05	0.06	0.16	0.07	0.12	0.78	0.06	0.09	0.33	0.21	0.52	1.00
3	5	51	10	–	0.06	0.59	–	0.13	0.99	–	0.02	0.11	–	0.05	0.53	–	0.03	0.21	–	0.34	1.00
3	10	101	10	–	–	0.46	–	–	0.97	–	–	0.08	–	–	0.35	–	–	0.14	–	–	0.99
6	1	11	10	0.08	0.13	0.76	0.15	0.47	1.00	0.07	0.06	0.20	0.08	0.15	0.96	0.09	0.14	0.61	0.30	0.68	1.00
6	2	21	10	0.06	0.11	0.69	0.10	0.35	1.00	0.04	0.06	0.14	0.05	0.11	0.86	0.06	0.10	0.48	0.21	0.58	1.00
6	5	51	10	–	0.05	0.56	–	0.14	1.00	–	0.02	0.10	–	0.04	0.59	–	0.04	0.31	–	0.39	1.00
6	10	101	10	–	–	0.38	–	–	0.98	–	–	0.07	–	–	0.35	–	–	0.20	–	–	1.00
12	1	11	10	0.06	0.11	0.71	0.14	0.43	1.00	0.05	0.06	0.17	0.07	0.11	0.95	0.08	0.13	0.64	0.30	0.67	1.00
24	1	11	10	0.06	0.09	0.62	0.13	0.37	1.00	0.05	0.06	0.13	0.05	0.09	0.87	0.08	0.13	0.60	0.29	0.66	1.00
X^2 test using Iterated $\hat{\theta}^{(1)}$																					
3	1	2	1	0.05	0.07	0.12	0.10	0.20	0.68	0.04	0.06	0.06	0.04	0.06	0.05	0.08	0.09	0.08	0.20	0.33	0.47
3	2	3	1	0.17	0.30	0.93	0.10	0.13	0.55	0.05	0.05	0.05	0.06	0.06	0.05	0.08	0.09	0.07	0.14	0.25	0.43
3	5	6	1	0.20	0.43	1.00	0.08	0.11	0.44	0.06	0.05	0.05	0.05	0.05	0.06	0.06	0.07	0.06	0.09	0.19	0.39
3	10	11	1	0.13	0.30	0.99	0.05	0.08	0.32	0.04	0.05	0.04	0.04	0.05	0.06	0.04	0.07	0.05	0.07	0.16	0.35
6	1	2	1	0.05	0.07	0.13	0.10	0.17	0.81	0.06	0.07	0.20	0.12	0.18	0.67	0.06	0.05	0.08	0.11	0.20	0.61
6	2	3	1	0.09	0.18	0.87	0.08	0.13	0.77	0.06	0.06	0.16	0.10	0.14	0.57	0.05	0.05	0.06	0.10	0.15	0.56
6	5	6	1	0.14	0.33	1.00	0.08	0.12	0.66	0.05	0.06	0.11	0.08	0.11	0.40	0.05	0.04	0.06	0.08	0.13	0.47
6	10	11	1	0.09	0.23	0.98	0.06	0.08	0.50	0.04	0.05	0.09	0.06	0.08	0.28	0.04	0.05	0.05	0.06	0.10	0.40
12	1	2	1	0.05	0.06	0.13	0.08	0.14	0.78	0.07	0.08	0.31	0.21	0.47	1.00	0.05	0.06	0.12	0.11	0.16	0.72
24	1	2	1	0.06	0.07	0.14	0.08	0.13	0.66	0.07	0.08	0.30	0.19	0.42	1.00	0.06	0.07	0.17	0.13	0.22	0.93
3	1	11	10	0.33	0.14	0.09	0.32	0.35	0.89	0.28	0.13	0.07	0.29	0.13	0.08	0.29	0.17	0.08	0.34	0.36	0.39
3	2	21	10	0.04	0.04	0.14	0.05	0.12	0.79	0.04	0.04	0.05	0.04	0.04	0.06	0.04	0.05	0.05	0.05	0.10	0.31
3	5	51	10	–	0.03	0.15	–	0.05	0.60	–	0.02	0.04	–	0.02	0.04	–	0.02	0.04	–	0.04	0.23
3	10	101	10	–	–	0.14	–	–	0.41	–	–	0.03	–	–	0.04	–	–	0.03	–	–	0.17
6	1	11	10	0.17	0.08	0.17	0.16	0.14	0.78	0.15	0.08	0.08	0.15	0.08	0.29	0.16	0.09	0.06	0.21	0.24	0.41
6	2	21	10	0.02	0.04	0.17	0.02	0.06	0.72	0.02	0.03	0.06	0.02	0.04	0.20	0.02	0.03	0.05	0.02	0.06	0.32
6	5	51	10	–	0.01	0.15	–	0.02	0.54	–	0.01	0.05	–	0.01	0.12	–	0.01	0.04	–	0.02	0.24
6	10	101	10	–	–	0.13	–	–	0.34	–	–	0.04	–	–	0.07	–	–	0.03	–	–	0.16
12	1	11	10	0.11	0.07	0.23	0.11	0.10	0.79	0.10	0.07	0.10	0.10	0.07	0.52	0.10	0.07	0.06	0.14	0.13	0.23
24	1	11	10	0.10	0.07	0.21	0.09	0.08	0.77	0.09	0.06	0.09	0.08	0.07	0.46	0.08	0.06	0.07	0.10	0.09	0.29
Kolmogorov-Smirnov test using $\hat{\theta}^{(0)}$																					
–	–	–	1	0.09	0.20	0.90	0.07	0.07	0.13	0.13	0.17	0.63	0.30	0.57	1.00	0.15	0.23	0.83	0.58	0.89	1.00
–	–	–	5	0.07	0.20	0.99	0.08	0.10	0.17	0.06	0.09	0.19	0.10	0.14	0.59	0.05	0.07	0.25	0.11	0.29	0.98
–	–	–	10	0.09	0.05	0.21	0.10	0.06	0.09	0.05	0.05	0.07	0.05	0.07	0.11	0.08	0.04	0.05	0.12	0.04	0.22

Table 3: Power under deviations in mean, variance, skewness, and kurtosis using balanced partitions ($k = 1, 10$ and $\alpha = 5\%$).

rejections clearly increases with the sample size when $k = 1$. The results under asymmetry (SGT_{1/3} and SGT_{2/3}) are mixed. Obviously, all the tests detect the stronger asymmetry ($\mu_3 = 2/3$) more easily than the mild one ($\mu_3 = 1/3$), particularly the X^2 test, which exhibits trivial power with many partitions in the latter case. Even though the W test performs much better under $\mu_3 = 2/3$, there are still some partitions that lead to trivial power. It seems preferable to choose large L for these asymmetric alternatives, though the X^2 test's power is still very poor with $\mu_3 = 1/3$ even with this choice. Similar comments can be made for the leptokurtic alternatives. In this case, the W test exhibits excellent power, much better than the X^2 and KS tests. The X^2 test also exhibits almost trivial power for most partitions under the milder leptokurtic alternative (t_5), but performs much better with stronger leptokurtosis ($t_{2.1}$). The W test outperforms the KS test under all partitions and leptokurtosis alternatives. The KS test is much more sensitive to the curse of dimensionality than any of the χ^2 tests.

We have observed a poor performance of the X^2 test for the mildest alternatives, particularly in the direction of asymmetry and kurtosis. Next we present results for unbalanced partitions \mathbf{U} with $\vartheta_\ell \neq \ell/L$ and small cells on the tails in the hope of closing the gap between the two χ^2 tests. We report results for the following partitions ($\vartheta_0, \vartheta_1, \dots, \vartheta_L$):

- (i) 6* with (0, 0.01, 0.25, 0.5, 0.75, 0.99, 1),
- (ii) 6** with (0, 0.05, 0.1, 0.5, 0.9, 0.95, 1),
- (iii) 8* with (0, 0.01, 0.06, 0.16, 0.5, 0.84, 0.94, 0.99, 1).
- (iv) 8** with (0, 0.01, 0.10, 0.33, 0.5, 0.66, 0.90, 0.99, 1).

These unbalanced partitions are intended, in principle, to improve the power in the direction of heavy-tailed alternatives. They have also proven to improve other alternatives in the classical case (Kallenberg et al., 1985).

In Table 4 we observe that for $k = 1$ the χ^2 tests exhibit a similar size accuracy to that of the omnibus bootstrap test but suffer size distortions for the smaller sample sizes when $k = 10$, except when $n = 500$.

Table 5 reports the proportion of rejections under the alternative for the chosen unbalanced partitions. The power of the X^2 test improves when the partitions are unbalanced and is similar to that of the W test. Both χ^2 tests are much more robust to the curse of dimensionality than the omnibus KS test with most partitions.

L	r	J	k	$n = 50$			$n = 100$			$n = 500$		
				0.01	0.05	0.10	0.01	0.05	0.10	0.01	0.05	0.10
W test using $\hat{\theta}^{(0)}$												
6*	1	2	1	0.02	0.07	0.11	0.01	0.05	0.09	0.01	0.05	0.10
6**	1	2	1	0.01	0.05	0.10	0.01	0.05	0.10	0.01	0.04	0.10
8*	1	2	1	0.01	0.04	0.09	0.01	0.05	0.10	0.01	0.05	0.10
8**	1	2	1	0.01	0.05	0.09	0.01	0.05	0.10	0.01	0.05	0.10
6*	1	11	10	0.09	0.17	0.23	0.04	0.11	0.16	0.02	0.07	0.11
6**	1	11	10	0.04	0.11	0.17	0.02	0.07	0.13	0.01	0.06	0.10
8*	1	11	10	0.05	0.13	0.19	0.03	0.09	0.14	0.01	0.06	0.10
8**	1	11	10	0.06	0.14	0.20	0.03	0.09	0.14	0.02	0.06	0.11
X^2 test using Iterated $\hat{\theta}^{(1)}$												
6*	1	2	1	0.02	0.06	0.10	0.01	0.05	0.09	0.01	0.05	0.10
6**	1	2	1	0.02	0.07	0.13	0.02	0.06	0.11	0.01	0.06	0.11
8*	1	2	1	0.01	0.05	0.10	0.01	0.05	0.09	0.01	0.05	0.11
8**	1	2	1	0.01	0.05	0.09	0.01	0.05	0.09	0.01	0.05	0.10
6*	1	11	10	0.32	0.40	0.45	0.17	0.29	0.36	0.03	0.09	0.15
6**	1	11	10	0.28	0.38	0.46	0.12	0.23	0.32	0.02	0.08	0.14
8*	1	11	10	0.29	0.37	0.42	0.16	0.28	0.37	0.02	0.08	0.14
8**	1	11	10	0.26	0.34	0.39	0.14	0.26	0.34	0.03	0.08	0.14
Kolmogorov-Smirnov test using $\hat{\theta}^{(0)}$												
—	—	—	1	0.01	0.05	0.09	0.01	0.05	0.10	0.01	0.05	0.10
—	—	—	5	0.01	0.05	0.10	0.01	0.05	0.11	0.01	0.05	0.10
—	—	—	10	0.01	0.05	0.10	0.01	0.05	0.10	0.01	0.05	0.11

Table 4: Size accuracy using unbalanced partitions ($k = 1, 10$).

L	r	J	k	$a = 1$			$b = 2/3$			SGT _{1/3}			SGT _{2/3}			t_5			$t_{2.1}$		
				50	100	500	50	100	500	50	100	500	50	100	500	50	100	500	50	100	500
W test using $\hat{\theta}^{(0)}$																					
6*	1	2	1	0.10	0.16	0.65	0.14	0.21	0.84	0.07	0.09	0.45	0.17	0.38	1.00	0.21	0.30	0.84	0.67	0.91	1.00
6**	1	2	1	0.07	0.13	0.54	0.08	0.17	0.89	0.08	0.10	0.43	0.27	0.70	1.00	0.08	0.15	0.65	0.39	0.68	1.00
8*	1	2	1	0.11	0.16	0.67	0.12	0.20	0.91	0.06	0.09	0.42	0.20	0.63	1.00	0.20	0.30	0.86	0.62	0.87	1.00
8**	1	2	1	0.11	0.17	0.65	0.13	0.21	0.90	0.06	0.09	0.38	0.20	0.39	0.99	0.21	0.31	0.87	0.70	0.92	1.00
Best balanced			1	0.20	0.44	1.00	0.10	0.18	0.81	0.09	0.16	0.66	0.28	0.54	1.00	0.19	0.31	0.90	0.69	0.92	1.00
6*	1	11	10	0.24	0.26	0.77	0.49	0.71	1.00	0.16	0.10	0.14	0.17	0.18	0.87	0.33	0.35	0.69	0.57	0.80	1.00
6**	1	11	10	0.09	0.07	0.52	0.05	0.06	0.96	0.11	0.09	0.15	0.12	0.14	1.00	0.06	0.04	0.22	0.03	0.20	0.99
8*	1	11	10	0.15	0.16	0.72	0.29	0.49	1.00	0.10	0.07	0.14	0.13	0.13	0.99	0.21	0.21	0.63	0.39	0.69	1.00
8**	1	11	10	0.19	0.21	0.76	0.42	0.67	1.00	0.12	0.08	0.12	0.14	0.13	0.75	0.27	0.30	0.70	0.53	0.81	1.00
Best balanced			10	0.09	0.24	0.93	0.20	0.59	1.00	0.08	0.13	0.61	0.14	0.39	1.00	0.10	0.21	0.78	0.40	0.81	1.00
X^2 test using Iterated $\hat{\theta}^{(1)}$																					
6*	1	2	1	0.12	0.16	0.58	0.15	0.23	0.89	0.04	0.05	0.22	0.09	0.08	0.61	0.23	0.31	0.81	0.63	0.85	1.00
6**	1	2	1	0.07	0.08	0.28	0.10	0.19	0.92	0.10	0.12	0.42	0.37	0.79	1.00	0.05	0.04	0.11	0.06	0.12	0.78
8*	1	2	1	0.10	0.14	0.53	0.13	0.22	0.93	0.06	0.09	0.47	0.23	0.68	1.00	0.14	0.19	0.69	0.39	0.66	1.00
8**	1	2	1	0.11	0.15	0.54	0.15	0.23	0.91	0.06	0.08	0.40	0.18	0.38	0.99	0.19	0.25	0.74	0.55	0.78	1.00
Best balanced			1	0.20	0.43	1.00	0.10	0.20	0.81	0.07	0.08	0.31	0.21	0.47	1.00	0.08	0.09	0.17	0.20	0.33	0.93
6*	1	11	10	0.38	0.27	0.77	0.44	0.55	1.00	0.40	0.27	0.09	0.40	0.27	0.27	0.41	0.38	0.64	0.59	0.68	1.00
6**	1	11	10	0.35	0.16	0.51	0.41	0.26	0.87	0.41	0.24	0.20	0.40	0.27	0.85	0.40	0.24	0.06	0.53	0.27	0.15
8*	1	11	10	0.30	0.19	0.76	0.42	0.51	1.00	0.36	0.27	0.16	0.36	0.29	0.78	0.40	0.36	0.48	0.61	0.57	0.98
8**	1	11	10	0.30	0.22	0.78	0.41	0.54	1.00	0.34	0.25	0.13	0.34	0.24	0.59	0.37	0.38	0.55	0.59	0.62	0.99
Best balanced			10	0.33	0.14	0.23	0.32	0.35	0.89	0.28	0.13	0.10	0.29	0.13	0.52	0.29	0.17	0.08	0.34	0.36	0.41
Kolmogorov-Smirnov test using $\hat{\theta}^{(0)}$																					
–	–	–	1	0.09	0.20	0.90	0.07	0.07	0.13	0.13	0.17	0.63	0.30	0.57	1.00	0.15	0.23	0.83	0.58	0.89	1.00
			5	0.07	0.20	0.99	0.08	0.10	0.17	0.06	0.09	0.19	0.10	0.14	0.59	0.05	0.07	0.25	0.11	0.29	0.98
			10	0.09	0.05	0.21	0.10	0.06	0.09	0.05	0.05	0.07	0.05	0.07	0.11	0.08	0.04	0.05	0.12	0.04	0.22

Table 5: Power under deviations in mean, variance, skewness, and kurtosis using unbalanced partitions ($k = 1, 10$ and $\alpha = 5\%$).

Appendix

Proof of Theorem 1. By the law of large numbers (LLN) and the central limit theorem (CLT), under H_0 ,

$$\hat{\Phi}_D(\boldsymbol{\theta}_0) = \left[\text{diag}^{-1/2}(\mathbf{v} \otimes \mathbf{q}) + o_p(1) \right] \frac{1}{\sqrt{n}} \sum_{i=1}^n ((\boldsymbol{\eta}_i(\boldsymbol{\theta}_0) - \mathbf{v}) \otimes \boldsymbol{\zeta}_i) \xrightarrow{d} \mathcal{N}_{LJ}(\mathbf{0}, \mathbf{Q}_U), \quad (9)$$

where \mathbf{Q}_U was defined in (5). Since \mathbf{Q}_U is idempotent and

$$\text{rank}(\mathbf{Q}_U) = \text{tr}(\mathbf{Q}_U) = \text{tr} \left((\mathbf{I}_L - \sqrt{\mathbf{v}}\sqrt{\mathbf{v}'}) \otimes \mathbf{I}_J \right) = J(L-1),$$

using the circular property of the trace, after applying the continuous mapping theorem (CMT),

$$\hat{X}_D^2(\boldsymbol{\theta}_0) = \hat{\Phi}_D'(\boldsymbol{\theta}_0) \hat{\Phi}_D(\boldsymbol{\theta}_0) \xrightarrow{d} \sum_{j=1}^{J(L-1)} Z_j^2 \stackrel{d}{=} \chi_{J(L-1)}^2,$$

where, henceforth, Z_j , $j = 1, 2, \dots, J(L-1)$ are i.i.d. standard normal random variables, which proves (i).

Applying a Taylor expansion $\ln(x+1) = x - x^2/2 + o(x^2)$ we get that

$$\ln \left(\frac{\hat{O}_{D_{\ell j}^0}}{nv_{\ell} \hat{q}_j} \right) = \frac{\hat{O}_{D_{\ell j}^0} - nv_{\ell} \hat{q}_j}{nv_{\ell} \hat{q}_j} - \frac{1}{2} \left(\frac{\hat{O}_{D_{\ell j}^0} - nv_{\ell} \hat{q}_j}{nv_{\ell} \hat{q}_j} \right)^2 + o \left(\left(\frac{\hat{O}_{D_{\ell j}^0} - nv_{\ell} \hat{q}_j}{nv_{\ell} \hat{q}_j} \right)^2 \right) \quad \text{a.s.}$$

so that

$$\begin{aligned} \hat{G}_D^2(\boldsymbol{\theta}_0) &= 2 \sum_{\ell=1}^L \sum_{j=1}^J \hat{O}_{D_{\ell j}^0} \ln \left(\frac{\hat{O}_{D_{\ell j}^0}}{nv_{\ell} \hat{q}_j} \right) \\ &= 2 \sum_{\ell=1}^L \sum_{j=1}^J \left[(\hat{O}_{D_{\ell j}^0} - nv_{\ell} \hat{q}_j) \ln \left(\frac{\hat{O}_{D_{\ell j}^0}}{nv_{\ell} \hat{q}_j} \right) + nv_{\ell} \hat{q}_j \ln \left(\frac{\hat{O}_{D_{\ell j}^0}}{nv_{\ell} \hat{q}_j} \right) \right] \\ &= 2 \sum_{\ell=1}^L \sum_{j=1}^J \left\{ \left[\frac{(\hat{O}_{D_{\ell j}^0} - nv_{\ell} \hat{q}_j)^2}{nv_{\ell} \hat{q}_j} + o \left(\frac{(\hat{O}_{D_{\ell j}^0} - nv_{\ell} \hat{q}_j)^2}{nv_{\ell} \hat{q}_j} \right) \right] \right. \\ &\quad \left. + \left[(\hat{O}_{D_{\ell j}^0} - nv_{\ell} \hat{q}_j) - \frac{1}{2} \frac{(\hat{O}_{D_{\ell j}^0} - nv_{\ell} \hat{q}_j)^2}{nv_{\ell} \hat{q}_j} + o \left(\frac{(\hat{O}_{D_{\ell j}^0} - nv_{\ell} \hat{q}_j)^2}{nv_{\ell} \hat{q}_j} \right) \right] \right\} \\ &= \hat{X}_D^2(\boldsymbol{\theta}_0) + o_p(1) \end{aligned}$$

applying (9) and noticing that $\sum_{\ell=1}^L \sum_{j=1}^J \hat{O}_{D_{\ell j}^0} = n \sum_{\ell=1}^L v_{\ell} \sum_{j=1}^J \hat{q}_j = n$, which shows (iv).

Define $\mathbf{v}^\dagger = (v_1, \dots, v_{L-1})'$, and let $\mathbf{1}_R$ be an $R \times 1$ vector of 1's. Applying [Sherman and Morrison \(1950\)](#) formula, $(\text{diag}(\mathbf{v}^\dagger) - \mathbf{v}^\dagger \mathbf{v}^\dagger)^{-1} = \text{diag}^{-1}(\mathbf{v}^\dagger) + \mathbf{1}_{L-1} \mathbf{1}'_{L-1} / v_L$. Hence,

$$[(\text{diag}(\mathbf{v}) - \mathbf{v} \mathbf{v}') \otimes \text{diag}(\hat{\mathbf{q}})]^{-} = \begin{pmatrix} \text{diag}^{-1}(\mathbf{v}^\dagger) + \mathbf{1}_{L-1} \mathbf{1}'_{L-1} / v_L & \mathbf{0}_{LJ-1} \\ \mathbf{0}_{LJ-1} & 0 \end{pmatrix} \otimes \text{diag}^{-1}(\hat{\mathbf{q}}),$$

and,

$$\begin{aligned} \hat{W}_D(\boldsymbol{\theta}_0) &= \hat{\boldsymbol{\Phi}}_D'(\boldsymbol{\theta}_0) \mathbf{Q}_U^- \hat{\boldsymbol{\Phi}}_D(\boldsymbol{\theta}_0) \\ &= \frac{1}{n} (\hat{\mathbf{O}}_{D(\boldsymbol{\theta}_0)} - n \mathbf{v} \otimes \hat{\mathbf{q}})' \left[\begin{pmatrix} \text{diag}^{-1}(\mathbf{v}^\dagger) + \mathbf{1}_{L-1} \mathbf{1}'_{L-1} / v_L & \mathbf{0}_{LJ-1} \\ \mathbf{0}'_{LJ-1} & 0 \end{pmatrix} \otimes \text{diag}^{-1}(\hat{\mathbf{q}}) \right] \\ &\quad \times (\hat{\mathbf{O}}_{D(\boldsymbol{\theta}_0)} - n \mathbf{v} \otimes \hat{\mathbf{q}}) \\ &= \frac{1}{n} \sum_{\ell=1}^{L-1} \sum_{j=1}^J \frac{(\hat{O}_{D_{\ell j}}(\boldsymbol{\theta}_0) - n v_\ell \hat{q}_j)^2}{v_\ell \hat{q}_j} + \frac{1}{n} \sum_{j=1}^J \frac{(\sum_{\ell=1}^{L-1} (\hat{O}_{D_{\ell j}}(\boldsymbol{\theta}_0) - n v_\ell \hat{q}_j))^2}{v_L \hat{q}_j} \\ &= \frac{1}{n} \sum_{\ell=1}^{L-1} \sum_{j=1}^J \frac{(\hat{O}_{D_{\ell j}}(\boldsymbol{\theta}_0) - n v_\ell \hat{q}_j)^2}{v_\ell \hat{q}_j} + \frac{1}{n} \sum_{j=1}^J \frac{(\hat{O}_{D_{Lj}}(\boldsymbol{\theta}_0) - n v_L \hat{q}_j)^2}{v_L \hat{q}_j} \\ &= \hat{X}_D^2(\boldsymbol{\theta}_0), \end{aligned}$$

which shows (iii). Notice that $\hat{\boldsymbol{\Phi}}_D'(\boldsymbol{\theta}) \mathbf{Q}_U^- \hat{\boldsymbol{\Phi}}_D(\boldsymbol{\theta}) = \hat{X}_D^2(\boldsymbol{\theta})$ for all $\boldsymbol{\theta} \in \Theta$.

To reduce the notational burden, let $\hat{O}_{\ell j} = \hat{O}_{D_{\ell j}}(\boldsymbol{\theta})$. Then, for all $\boldsymbol{\theta} \in \Theta$,

$$\begin{aligned} \widehat{LM}_D(\boldsymbol{\theta}) &= n \hat{\boldsymbol{\Psi}}_D'(\boldsymbol{\theta}) \hat{\mathcal{I}}_D^{-1} \hat{\boldsymbol{\Psi}}_D(\boldsymbol{\theta}) \\ &= n \sum_{r=1}^{LJ-1} \left(\frac{\hat{O}_r^*}{n \hat{\pi}_r^*} - \frac{\hat{O}_{LJ}}{n v_L \hat{q}_J} \right)^2 \hat{\pi}_r^* - n \sum_{r=1}^{LJ-1} \sum_{s=1}^{LJ-1} \left(\frac{\hat{O}_r^*}{n \hat{\pi}_r^*} - \frac{\hat{O}_{LJ}}{n v_L \hat{q}_J} \right) \left(\frac{\hat{O}_s^*}{n \hat{\pi}_s^*} - \frac{\hat{O}_{LJ}}{n v_L \hat{q}_J} \right) \hat{\pi}_r^* \hat{\pi}_s^*. \end{aligned} \tag{10}$$

The second term in (10) equals

$$-n \left[\sum_{r=1}^{LJ-1} \left(\frac{\hat{O}_r^*}{n \hat{\pi}_r^*} - \frac{\hat{O}_{LJ}}{n v_L \hat{q}_J} \right) \hat{\pi}_r^* \right]^2 = -n \left(\frac{\hat{O}_{LJ}}{n v_L \hat{q}_J} - 1 \right)^2$$

using the fact that $\sum_{r=1}^{LJ-1} \hat{O}_r^* = n - \hat{O}_{LJ}$ and $\sum_{r=1}^{LJ-1} \hat{\pi}_r^* = 1 - v_L \hat{q}_J$. To simplify the first sum in (10) we write

$$\frac{\hat{O}_r^*}{n \hat{\pi}_r^*} - \frac{\hat{O}_{LJ}}{n v_L \hat{q}_J} = \left\{ v_L \hat{q}_J \left(\frac{\hat{O}_r^*}{n} - \hat{\pi}_r^* \right) - \hat{\pi}_r^* \left(\frac{\hat{O}_{LJ}}{n} - v_L \hat{q}_J \right) \right\} \frac{1}{\hat{\pi}_r^* v_L \hat{q}_J}.$$

Then, because

$$\sum_{r=1}^{LJ-1} \left(\frac{\hat{O}_r^*}{n} - \hat{\pi}_r^* \right) = - \left(\frac{\hat{O}_{LJ}}{n} - v_L \hat{q}_J \right),$$

the first term in (10) becomes

$$\begin{aligned} & n \left\{ \sum_{r=1}^{LJ-1} \frac{1}{\hat{\pi}_r^*} \left(\frac{\hat{O}_r^*}{n} - \hat{\pi}_r^* \right)^2 + \frac{2}{v_L \hat{q}_J} \left(\frac{\hat{O}_{LJ}}{n} - v_L \hat{q}_J \right)^2 + \frac{1}{v_L^2 \hat{q}_J^2} (1 - v_L \hat{q}_J) \left(\frac{\hat{O}_{LJ}}{n} - v_L \hat{q}_J \right)^2 \right\} \\ &= n \left\{ \sum_{\ell=1}^L \sum_{j=1}^J \frac{1}{v_L^2 \hat{q}_J^2} \left(\frac{\hat{O}_{\ell j}}{n} - v_L \hat{q}_j \right)^2 + \frac{1}{v_L^2 \hat{q}_J^2} \left(\frac{\hat{O}_{LJ}}{n} - v_L \hat{q}_J \right)^2 \right\}, \end{aligned}$$

which proves (ii). \square

The next lemma, which is an application of Lemma 1 by [Delgado and Stute \(2008\)](#), is used in several places below. Define, for $V_i = F_{Y|\mathbf{X}}(Y_i | \mathbf{X}_i)$, real $\kappa_1, \dots, \kappa_n$ that range in a compact interval, and

$$\hat{\beta}_{\ell j}(\kappa_1, \dots, \kappa_n) = \frac{1}{\sqrt{n}} \sum_{i=1}^n \left[\mathbb{1}_{\{V_i \leq \vartheta_\ell + \kappa_i n^{-1/2}\}} - \mathbb{1}_{\{V_i \leq \vartheta_\ell\}} - \kappa_i n^{-1/2} \right] \mathbb{1}_{\{\mathbf{X}_i \in A_j\}}$$

for $\ell = 1, \dots, L$ and $j = 1, \dots, J$.

Lemma 1. *For each finite K ,*

$$\sup_{\{|\kappa_i| \leq K, i=1, \dots, n\}} \left| \hat{\beta}_{\ell j}(\kappa_1, \dots, \kappa_n) \right| = o_p(1), \quad \ell = 1, \dots, L \text{ and } j = 1, \dots, J.$$

Define

$$\alpha_{\ell i}(\boldsymbol{\theta}_1, \boldsymbol{\theta}_2) = \frac{\partial}{\partial \boldsymbol{\theta}} F_{Y|\mathbf{X}, \boldsymbol{\theta}} \left(F_{Y|\mathbf{X}, \boldsymbol{\theta}_2}^{-1}(\vartheta_\ell | \mathbf{X}_i) \right) \Big|_{\boldsymbol{\theta} = \boldsymbol{\theta}_1}.$$

Proof of Theorem 2. Notice that under H_0 , for all $j = 1, \dots, J$ and $\ell = 1, \dots, L$,

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n \left(\mathbb{1}_{\{V_i(\tilde{\boldsymbol{\theta}}) \leq \vartheta_\ell\}} - \vartheta_\ell \right) \zeta_{ji} = \frac{1}{\sqrt{n}} \sum_{i=1}^n \left(\mathbb{1}_{\{V_i \leq F_{Y|\mathbf{X}} \left(F_{Y|\mathbf{X}, \tilde{\boldsymbol{\theta}}}^{-1}(\vartheta_\ell | \mathbf{X}_i) \mid \mathbf{X}_i \right)\}} - \vartheta_\ell \right) \zeta_{ji}. \quad (11)$$

Now, applying the mean value theorem (MVT), for all $i = 1, \dots, n$, under H_0 ,

$$F_{Y|\mathbf{X}, \boldsymbol{\theta}_0} \left(F_{Y|\mathbf{X}, \tilde{\boldsymbol{\theta}}}^{-1}(\vartheta_\ell | \mathbf{X}_i) \mid \mathbf{X}_i \right) = \vartheta_\ell + \alpha_{\ell i}(\tilde{\boldsymbol{\theta}}_i^*, \tilde{\boldsymbol{\theta}}) (\boldsymbol{\theta}_0 - \tilde{\boldsymbol{\theta}}) \quad (12)$$

for some $\tilde{\boldsymbol{\theta}}_i^* \in \mathbb{R}^p$ such that $\|\tilde{\boldsymbol{\theta}}_i^* - \boldsymbol{\theta}_0\| \leq \|\tilde{\boldsymbol{\theta}} - \boldsymbol{\theta}_0\|$. Since $\alpha_{\ell i}(\tilde{\boldsymbol{\theta}}_i^*, \tilde{\boldsymbol{\theta}})$ is bounded in a neighborhood of $\boldsymbol{\theta}_0$ for $i = 1, 2, \dots, n$, $\kappa_{i\ell j} = \alpha_{\ell i}(\tilde{\boldsymbol{\theta}}_i^*, \tilde{\boldsymbol{\theta}}) \zeta_{ji} \sqrt{n} (\boldsymbol{\theta}_0 - \tilde{\boldsymbol{\theta}})$ range, with high probability,

in a possibly large but compact set. Then (11) equals to

$$\begin{aligned} & \frac{1}{\sqrt{n}} \sum_{i=1}^n \left(\mathbb{1}_{\{V_i \leq \vartheta_\ell\}} - \vartheta_\ell \right) + \frac{1}{n} \sum_{i=1}^n \alpha_{\ell i} \left(\tilde{\boldsymbol{\theta}}_i^*, \tilde{\boldsymbol{\theta}} \right) \zeta_{ji} \left(\boldsymbol{\theta}_0 - \tilde{\boldsymbol{\theta}} \right) \\ & + \frac{1}{\sqrt{n}} \sum_{i=1}^n \left[\mathbb{1}_{\{V_i \leq \vartheta_\ell + \kappa_{i\ell j} n^{-1/2}\}} - \mathbb{1}_{\{V_i \leq \vartheta_\ell\}} - \kappa_{i\ell j} n^{-1/2} \right] \zeta_{ji}, \end{aligned}$$

and the last term is $o_p(1)$ by Lemma 1. Then, applying Lemma 1 and noticing that $\tau_{\boldsymbol{\theta}\ell}(\mathbf{X}_i) = \alpha_{\ell i}(\boldsymbol{\theta}, \boldsymbol{\theta}) - \alpha_{(\ell-1)i}(\boldsymbol{\theta}, \boldsymbol{\theta})$ under H_0 ,

$$\begin{aligned} \hat{\boldsymbol{\Phi}}_D(\tilde{\boldsymbol{\theta}}) &= \text{diag}^{-1/2} \left(\mathbf{v} \otimes \frac{1}{n} \sum_{i=1}^n \boldsymbol{\zeta}_i \right) \frac{1}{\sqrt{n}} \sum_{i=1}^n \left(\boldsymbol{\eta}_i(\tilde{\boldsymbol{\theta}}) - \mathbf{v} \right) \otimes \boldsymbol{\zeta}_i \\ &= \text{diag}^{-1/2}(\mathbf{v} \otimes \hat{\mathbf{q}}) \frac{1}{\sqrt{n}} \sum_{i=1}^n \left[(\boldsymbol{\eta}_i(\boldsymbol{\theta}_0) - \mathbf{v}) \otimes \boldsymbol{\zeta}_i - (\boldsymbol{\tau}_{\boldsymbol{\theta}}(\mathbf{X}_i) \otimes \boldsymbol{\zeta}_i) (\tilde{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) \right] + o_p(1) \\ &= \hat{\boldsymbol{\Phi}}_D(\boldsymbol{\theta}_0) - \hat{\mathbf{B}}_D(\boldsymbol{\theta}_0) \sqrt{n} (\tilde{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) + o_p(1) \end{aligned} \quad (13)$$

after applying the LLN. \square

Proof of Theorem 3. Define $\ell_{\boldsymbol{\theta}_0 i} = \ell_{\boldsymbol{\theta}_0}(Y_i, \mathbf{X}_i)$. Notice that by the CLT,

$$\begin{aligned} & \begin{pmatrix} \hat{\boldsymbol{\Phi}}_{D^0}(\boldsymbol{\theta}_0) \\ \mathbf{B}_D(\boldsymbol{\theta}_0) \sqrt{n} (\tilde{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) \end{pmatrix} = \frac{1}{\sqrt{n}} \sum_{i=1}^n \begin{pmatrix} \text{diag}^{-1/2}(\mathbf{v} \otimes \mathbf{q})(\boldsymbol{\eta}_i(\boldsymbol{\theta}_0) - \mathbf{v}) \otimes \boldsymbol{\zeta}_i \\ \mathbf{B}_D(\boldsymbol{\theta}_0) \ell_{\boldsymbol{\theta}_0 i} \end{pmatrix} + o_p(1) \\ & \xrightarrow{d} \mathcal{N}_{LJ+p} \left(\mathbf{0}, \begin{pmatrix} \mathbf{Q}_U & \mathbf{B}_D(\boldsymbol{\theta}_0) \boldsymbol{\Upsilon}_D(\boldsymbol{\theta}_0) \\ \boldsymbol{\Upsilon}'_D(\boldsymbol{\theta}_0) \mathbf{B}'_D(\boldsymbol{\theta}_0) & \mathbf{L}(\boldsymbol{\theta}_0) \end{pmatrix} \right). \end{aligned}$$

Then, applying Theorem 2,

$$\hat{\boldsymbol{\Phi}}_D(\tilde{\boldsymbol{\theta}}) = \begin{pmatrix} \mathbf{I}_{LJ} & -\mathbf{I}_{LJ} \end{pmatrix} \begin{pmatrix} \hat{\boldsymbol{\Phi}}_D(\boldsymbol{\theta}_0) \\ \mathbf{B}_D(\boldsymbol{\theta}_0) \sqrt{n} (\tilde{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) \end{pmatrix} + o_p(1) \xrightarrow{d} \mathcal{N}_{LJ}(\mathbf{0}, \boldsymbol{\Omega}_{D,\tilde{\boldsymbol{\theta}}}).$$

\square

Proof of Theorem 4. First, $\hat{\boldsymbol{\theta}}_{D^0} = \boldsymbol{\theta}_0 + o_p(1)$ by Assumption 4 and (Rao, 2002, Section 5.e.2 (i)). Recall that $\hat{\boldsymbol{\theta}}_{D^0}$ solves the equations

$$\mathbf{0} = \sum_{\ell=1}^L \sum_{j=1}^J \hat{O}_{D_0 \ell j} \frac{\partial \hat{\pi}_{D_0 \ell j}(\boldsymbol{\theta}) / \partial \boldsymbol{\theta} \Big|_{\boldsymbol{\theta} = \hat{\boldsymbol{\theta}}_{D^0}}}{\hat{\pi}_{D_0 \ell j}(\hat{\boldsymbol{\theta}}_{D^0})} \quad (14)$$

and $\hat{\pi}_{D(\boldsymbol{\theta})}(\boldsymbol{\theta}) = \mathbf{v} \otimes \hat{\mathbf{q}}$ for all $\boldsymbol{\theta} \in \Theta$. Taking into account that

$$\sum_{\ell=1}^L \frac{\partial}{\partial \boldsymbol{\theta}} \hat{\pi}_{D_{\ell_j}^0}(\boldsymbol{\theta}) = \frac{1}{n} \sum_{i=1}^n \frac{\partial}{\partial \boldsymbol{\theta}} \sum_{\ell=1}^L \int_{F_{Y|\mathbf{X},\boldsymbol{\theta}_0}^{-1}(\vartheta_{\ell-1} | \mathbf{X}_i)}^{F_{Y|\mathbf{X},\boldsymbol{\theta}_0}^{-1}(\vartheta_{\ell} | \mathbf{X}_i)} F_{Y|\mathbf{X},\boldsymbol{\theta}}(d\vartheta | \mathbf{X}_i) = \mathbf{0}, \quad (15)$$

we can express (14) as

$$\begin{aligned} \mathbf{0} &= \sum_{\ell=1}^L \sum_{j=1}^J \left(\frac{\hat{O}_{D_{\ell_j}^0}}{n \cdot \hat{\pi}_{D_{\ell_j}^0}(\hat{\boldsymbol{\theta}}_{D^0})} - 1 \right) \frac{\partial}{\partial \boldsymbol{\theta}} \hat{\pi}_{D_{\ell_j}^0}(\boldsymbol{\theta}) \Big|_{\boldsymbol{\theta}=\hat{\boldsymbol{\theta}}_{D^0}} \\ &= n^{-1/2} \cdot \text{diag}^{-1}(\hat{\boldsymbol{\pi}}_{D^0}(\hat{\boldsymbol{\theta}}_{D^0})) \cdot \text{diag}(\mathbf{v} \otimes \hat{\mathbf{q}}) \cdot \hat{\mathbf{B}}'_D(\hat{\boldsymbol{\theta}}_{D^0}) \\ &\quad \times \left[\text{diag}^{-1/2}(n \cdot \mathbf{v} \otimes \hat{\mathbf{q}}) \cdot (\hat{\mathbf{O}}_{D^0} - n \cdot \mathbf{v} \otimes \hat{\mathbf{q}}) - \text{diag}^{-1/2}(\mathbf{v} \otimes \hat{\mathbf{q}}) \cdot \sqrt{n} (\hat{\boldsymbol{\pi}}_{D^0}(\hat{\boldsymbol{\theta}}_{D^0}) - \mathbf{v} \otimes \hat{\mathbf{q}}) \right]. \end{aligned}$$

Hence, under H_0 , applying the LLN, the CMT, and a MVT argument,

$$\mathbf{0} = \mathbf{B}'_D(\boldsymbol{\theta}_0) \left[\hat{\boldsymbol{\Phi}}_D(\boldsymbol{\theta}_0) - \mathbf{B}_D(\boldsymbol{\theta}_0) \sqrt{n} (\hat{\boldsymbol{\theta}}_{D^0} - \boldsymbol{\theta}_0) \right] (1 + o_p(1))$$

so that

$$\sqrt{n} (\hat{\boldsymbol{\theta}}_{D^0} - \boldsymbol{\theta}_0) = [\mathbf{B}'_D(\boldsymbol{\theta}_0) \mathbf{B}_D(\boldsymbol{\theta}_0)]^{-1} \mathbf{B}'_D(\boldsymbol{\theta}_0) \hat{\boldsymbol{\Phi}}_D(\boldsymbol{\theta}_0) (1 + o_p(1)).$$

To prove the result apply arguments from the proof of Theorem 1, taking into account that $\mathbf{B}'_D(\boldsymbol{\theta}_0) \mathbf{Q}_U = \mathbf{B}'_D(\boldsymbol{\theta}_0)$ by (15). \square

Proof of Corollary 1. By the LLN and CMT, $\hat{\mathbf{B}}_D(\tilde{\boldsymbol{\theta}}) = \mathbf{B}_D(\boldsymbol{\theta}_0) + o_p(1)$ and, hence,

$$\begin{aligned} \sqrt{n} (\hat{\boldsymbol{\theta}}_{D(\tilde{\boldsymbol{\theta}})}^{(1)} - \boldsymbol{\theta}_0) &= \sqrt{n} (\tilde{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) + \hat{\boldsymbol{\Sigma}}_D^{-1}(\tilde{\boldsymbol{\theta}}) \hat{\mathbf{B}}'_D(\tilde{\boldsymbol{\theta}}) \hat{\boldsymbol{\Phi}}_D(\tilde{\boldsymbol{\theta}}) \\ &= \sqrt{n} (\tilde{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) + [\boldsymbol{\Sigma}_D^{-1}(\boldsymbol{\theta}_0) \mathbf{B}'_D(\boldsymbol{\theta}_0) + o_p(1)] \hat{\boldsymbol{\Phi}}_D(\tilde{\boldsymbol{\theta}}). \end{aligned}$$

Applying Theorem 3,

$$\sqrt{n} (\hat{\boldsymbol{\theta}}_{D(\tilde{\boldsymbol{\theta}})}^{(1)} - \boldsymbol{\theta}_0) = \boldsymbol{\Sigma}_D^{-1}(\boldsymbol{\theta}_0) \mathbf{B}'_D(\boldsymbol{\theta}_0) \hat{\boldsymbol{\Phi}}_D(\boldsymbol{\theta}_0) + o_p(1) \xrightarrow{d} \mathcal{N}_p(\mathbf{0}, \boldsymbol{\Sigma}_D^{-1}(\boldsymbol{\theta}_0)).$$

\square

Proof of Theorem 6. Since $\mathbb{I}(\boldsymbol{\theta}_0)$ is positive definite, and all the components of $\sqrt{\mathbf{v}} \sqrt{\mathbf{v}'} \otimes \mathbf{I}_J$ are nonnegative, there exists an orthonormal matrix \mathbf{C} that simultaneously diagonalizes

$\sqrt{\mathbf{v}}\sqrt{\mathbf{v}'} \otimes \mathbf{I}_J$, $\mathbf{G}_D(\boldsymbol{\theta}_0)$, and $\mathbf{S}_D(\boldsymbol{\theta}_0) = \mathbf{B}_D(\boldsymbol{\theta}_0)\mathbb{I}^{-1}(\boldsymbol{\theta}_0)\mathbf{B}'_D(\boldsymbol{\theta}_0)$, and which satisfies

$$[\mathbf{C}(\sqrt{\mathbf{v}}\sqrt{\mathbf{v}'} \otimes \mathbf{I}_J)\mathbf{C}'\mathbf{1}_{LJ}]_j = \begin{cases} 0 & \text{if } j = 1, \dots, J(L-1), \\ 1 & \text{if } j = J(L-1) + 1, \dots, JL, \end{cases} \quad (16)$$

$$[\mathbf{C}\mathbf{G}(\boldsymbol{\theta}_0)\mathbf{C}'\mathbf{1}_{LJ}]_j = \begin{cases} 0 & \text{if } j = 1, \dots, J(L-1) - p, J(L-1) + 1, JL, \\ 1 & \text{if } j = J(L-1) - p + 1, \dots, J(L-1), \end{cases} \quad (17)$$

$$[\mathbf{C}\mathbf{S}_D(\boldsymbol{\theta}_0)\mathbf{C}'\mathbf{1}_{LJ}]_j = \begin{cases} 0 & \text{if } j = 1, \dots, J(L-1) - p, J(L-1) + 1, JL, \\ 1 - \lambda_j & \text{if } j = J(L-1) - p + 1, \dots, J(L-1). \end{cases}$$

It follows reasoning as in the proof of [Moore and Spruill \(1975, Lemma 5.1\)](#). Since $\sqrt{\mathbf{v}}\sqrt{\mathbf{v}'} \otimes \mathbf{I}_J$ and $\mathbf{S}_D(\boldsymbol{\theta}_0)$ are commuting matrices, there exists a common orthonormal matrix \mathbf{C} that diagonalizes them. Moreover, $\sqrt{\mathbf{v}}\sqrt{\mathbf{v}'} \otimes \mathbf{I}_J$ and $\mathbf{S}_D(\boldsymbol{\theta}_0)$ are orthogonal projections having ranks J and p , respectively, so that by proper choice of basis we can take \mathbf{C} to satisfy (16) and (17). $\mathbf{S}_D(\boldsymbol{\theta}_0)$ has rank p and range contained in the range of $\mathbf{G}_D(\boldsymbol{\theta}_0)$. It follows that the eigenvalues of $\mathbf{S}_D(\boldsymbol{\theta}_0)$ are 0 except for those associated with the eigenvectors in the range of $\mathbf{G}_D(\boldsymbol{\theta}_0)$. That these are $1 - \lambda_j$ follows from the fact that they are roots of the determinantal equation $|\mathbf{S}_D(\boldsymbol{\theta}_0) - \beta\mathbf{I}_{LJ}| = 0$. Hence, we have

$$\begin{aligned} [\mathbf{C}\boldsymbol{\Omega}_{D,\hat{\boldsymbol{\theta}}^{(0)}}\mathbf{C}'\mathbf{1}_{LJ}]_j &= [\mathbf{C}(\mathbf{I}_{LJ} - (\sqrt{\mathbf{v}}\sqrt{\mathbf{v}'} \otimes \mathbf{I}_J) - \mathbf{S}_D(\boldsymbol{\theta}_0))\mathbf{C}']_j \\ &= \begin{cases} 1 & \text{if } j = 1, \dots, J(L-1) - p, \\ \lambda_j & \text{if } j = J(L-1) - p + 1, \dots, J(L-1), \\ 0 & \text{if } j = J(L-1) + 1, \dots, LJ. \end{cases} \end{aligned}$$

□

Proof of Corollary 2. The result follows from the proof of [Theorem 6](#) and [Assumption 5](#) since $0 < \lambda_j < 1$, $j = 1, \dots, LJ$. □

Proof of Theorem 7. First we show that, for any $\tilde{\boldsymbol{\theta}}$ satisfying [Assumption 3](#),

$$\hat{\mathbf{q}}_{\hat{\mathbf{A}}} = \mathbf{q} + o_p(1), \quad (18)$$

$$\hat{\mathbf{B}}_{\hat{\mathbf{D}}}(\tilde{\boldsymbol{\theta}}) = \mathbf{B}_D(\boldsymbol{\theta}_0) + o_p(1) \quad \text{under } H_0, \quad (19)$$

$$\hat{\boldsymbol{\Phi}}_{\hat{\mathbf{D}}}(\tilde{\boldsymbol{\theta}}) \xrightarrow{d} \mathcal{N}_{LJ}(\mathbf{0}, \boldsymbol{\Omega}_{D,\tilde{\boldsymbol{\theta}}}) \quad \text{under } H_0, \quad (20)$$

with $\hat{\mathbf{D}} = (\mathbf{U}, \hat{\mathbf{A}})$. The same arguments as those in [Andrews \(1988a, Lemmas 5 and 6\)](#) and

the LLN prove (18) and (19). In order to prove (20), take into account that, because \mathcal{A} is a VC class and, hence, Glivenko-Cantelli, Lemma 1 holds uniformly in \mathcal{A} , i.e.,

$$\sup_{\substack{A \in \mathcal{A} \\ \{\|\kappa_i| \leq K, i=1, \dots, n\}}} \left| \frac{1}{\sqrt{n}} \sum_{i=1}^n \left[\mathbb{1}_{\{V_i \leq \vartheta_\ell + \kappa_i n^{-1/2}\}} - \mathbb{1}_{\{V_i \leq \vartheta_\ell\}} - \kappa_i n^{-1/2} \right] \mathbb{1}_{\{\mathbf{X}_i \in \hat{A}\}} \right| = o(1) \text{ a.s.} \quad (21)$$

for real $\kappa_1, \dots, \kappa_n$ that range in a compact interval. Thus, mimicking the arguments in (13), (21) implies that, under H_0 ,

$$\hat{\Phi}_D(\tilde{\boldsymbol{\theta}}) = \text{diag}^{-1/2}(\mathbf{v} \otimes \mathbf{q}) \hat{\Delta}_{(\mathbf{U}, \hat{\mathbf{A}})}(\boldsymbol{\theta}_0) - \mathbf{B}_D(\boldsymbol{\theta}_0) \sqrt{n} (\tilde{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) + o_p(1), \quad (22)$$

with

$$\hat{\Delta}_{(\mathbf{U}, \hat{\mathbf{A}})}(\boldsymbol{\theta}) = \frac{1}{\sqrt{n}} \sum_{i=1}^n (\boldsymbol{\eta}_i(\boldsymbol{\theta}) - \mathbf{v}) \otimes \boldsymbol{\zeta}_i,$$

where \mathbf{U} is fixed and \mathbf{A} takes values in \mathcal{H} . Thus, under H_0 , $\{\hat{\Delta}_{(\mathbf{U}, \hat{\mathbf{A}})}(\boldsymbol{\theta}_0) : \mathbf{A} \in \mathcal{H}\}$ is an $LJ \times 1$ vector of marked empirical processes indexed by sets $\mathbf{A} \in \mathcal{H}$ under \mathbf{U} is a fixed partition of $[0, 1]$. Applying Andrews (1988a, Lemma 3), under H_0 ,

$$\hat{\Delta}_{(\mathbf{U}, \cdot)}(\boldsymbol{\theta}_0) \xrightarrow{d} \Delta_{(\mathbf{U}, \cdot)}^0 \text{ as a process on } \mathcal{H},$$

where $\Delta_{(\mathbf{U}, \cdot)}^0$ is an \mathbb{R}^{LJ} -valued tied down Gaussian process with bounded sample paths and covariance structure

$$\mathbb{E}(\Delta_{(\mathbf{U}, \mathbf{A}_1)}^0 \Delta_{(\mathbf{U}, \mathbf{A}_2)}^0) \stackrel{d}{=} (\text{diag}(\mathbf{v}) - \mathbf{v}\mathbf{v}') \otimes \mathbf{K}_{\mathbf{A}_1, \mathbf{A}_2},$$

where $\mathbf{A}_r = (A_{r1}, \dots, A_{rJ})'$, and $\mathbf{K}_{\mathbf{A}_1, \mathbf{A}_2}$ is a $J \times J$ matrix with jk -th element $q_{A_{1j} \cap A_{2k}}$, i.e., $\mathbf{K}_{\mathbf{A}, \mathbf{A}} = \text{diag}(\mathbf{q})$. Under H_0 ,

$$\sqrt{n} (\tilde{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) = \hat{\varphi} + o_p(1),$$

where $\hat{\varphi} = n^{-1/2} \sum_{i=1}^n \ell_{\boldsymbol{\theta}_i} \xrightarrow{d} \varphi$, φ follows $\mathcal{N}_p(\mathbf{0}, \mathbf{L}(\boldsymbol{\theta}_0))$, and

$$\mathbb{E}(\Delta_{(\mathbf{U}, \mathbf{A})}^0 \cdot \varphi) = \text{diag}(\mathbf{v} \otimes \mathbf{q}) \Upsilon_D(\boldsymbol{\theta}_0).$$

The above shows that under H_0 ,

$$\left(\hat{\Delta}_{(\mathbf{U}, \cdot)}(\boldsymbol{\theta}_0), \hat{\mathbf{A}}, \varphi \right) \xrightarrow{d} \left(\Delta_{(\mathbf{U}, \cdot)}^0, \mathbf{A}, \varphi \right) \text{ as a process on } \mathcal{H}. \quad (23)$$

Therefore, (20) follows by (22), (23), and the CMT. Reasoning as in the proof of Corollary (1),

using (19) and (20),

$$\sqrt{n} \left(\hat{\boldsymbol{\theta}}_{\hat{\boldsymbol{\theta}}}^{(1)} - \boldsymbol{\theta}_0 \right) \xrightarrow{d} \mathcal{N}_p \left(\mathbf{0}, \boldsymbol{\Sigma}_D^{-1}(\boldsymbol{\theta}_0) \right).$$

The rest of the proof follows combining (18)–(20) with the arguments in the proofs of Theorems 5 and 6. \square

Proof of Theorem 8. Recall that

$$F_{Y|\mathbf{X}}(\mathrm{d}y \mid \mathbf{x}) = \left[1 + n^{-1/2} t_{n\boldsymbol{\theta}_0}(y, \mathbf{x}) \right] F_{Y|\mathbf{X}, \boldsymbol{\theta}_0}(\mathrm{d}y \mid \mathbf{x}).$$

Then, compared to the proof of Theorem 2, we have to add another term, namely

$$\frac{1}{\sqrt{n}} \int_{F_{Y|\mathbf{X}, \hat{\boldsymbol{\theta}}}^{-1}(\vartheta_{\ell-1} \mid \mathbf{X}_i)}^{F_{Y|\mathbf{X}, \hat{\boldsymbol{\theta}}}^{-1}(\vartheta_{\ell} \mid \mathbf{X}_i)} t_{n\boldsymbol{\theta}_0}(y, \mathbf{X}_i) F_{Y|\mathbf{X}, \boldsymbol{\theta}_0}(\mathrm{d}y \mid \mathbf{X}_i),$$

to the right hand side of (12). Then, the theorem follows mimicking the arguments in Theorem 2 and using the properties of $t_{n\boldsymbol{\theta}_0}$. \square

Proof of Corollary 3. Apply arguments similar to those in the proofs of Theorems 5–7. \square

References

- ANDREWS, D.W.K. (1988a): “Chi-Square Diagnostic Tests for Econometric Models: Introduction and Applications,” *Journal of Econometrics*, 37, 135–156.
- (1988b): “Chi-Square Diagnostic Tests for Econometric Models: Theory,” *Econometrica*, 56, 1419–1453.
- (1997): “A Conditional Kolmogorov test,” *Econometrica*, 65, 1097–1128.
- BEHNEN, K. AND G. NEUHAUS (1975): “A Central Limit Theorem under Contiguous Alternatives,” *The Annals of Statistics*, 3, 1349–1353.
- BICKEL, P.J. AND K.A. DOKSUM (2015): *Mathematical Statistics: Basic Ideas and Selected Topics*, vol. 1, Chapman and Hall/CRC.
- CHERNOFF, H. AND E.L. LEHMANN (1954): “The Use of Maximum Likelihood Estimates in χ^2 Tests for Goodness of Fit,” *The Annals of Mathematical Statistics*, 25, 579–586.
- CHIBISOV, D.M. (1971): “Certain Chi-Square Type Tests for Continuous Distributions,” *Theory of Probability & Its Applications*, 16, 1–22.
- COHEN, A. AND H.B. SACKROWITZ (1975): “Unbiasedness of the Chi-Square, Likelihood Ratio, and Other Goodness of Fit Tests for the Equal Cell Case,” *The Annals of Statistics*, 959–964.
- DELGADO, M.A. AND W. STUTE (2008): “Distribution-free specification tests of conditional models,” *Journal of Econometrics*, 143, 37–55.
- GESSAMAN, M.P. (1970): “A Consistent Nonparametric Multivariate Density Estimator Based on Statistically Equivalent Blocks,” *The Annals of Mathematical Statistics*, 41, 1344–1346.
- HECKMAN, J.J. (1984): “The χ^2 Goodness of Fit Statistic for Models with Parameters Estimated from Microdata,” *Econometrica*, 52, 1543–1547.

- HOROWITZ, J.L. (1985): "Testing Probabilistic Discrete Choice Models of Travel Demand by Comparing Predicted and Observed Aggregate Choice Shares," *Transportation Research Part B: Methodological*, 19, 17–38.
- KALLENBERG, W.C.M., J. OOSTERHOFF, AND B.F. SCHRIEVER (1985): "The Number of Classes in Chi-Squared Goodness-of-Fit Tests," *Journal of the American Statistical Association*, 80, 959–968.
- MCFADDEN, D. (1974): "Conditional logit analysis of qualitative choice behavior," in *Frontiers in Econometrics*, ed. by P. Zarembka, Academic Press, 105–142.
- MOORE, D.S. (1971): "A Chi-Square Statistic with Random Cell Boundaries," *The Annals of Mathematical Statistics*, 42, 147–156.
- (1977): "Generalized Inverses, Wald's Method, and the Construction of Chi-Squared Tests of Fit," *Journal of the American Statistical Association*, 72, 131–137.
- MOORE, D.S. AND M.C. SPRUILL (1975): "Unified Large-Sample Theory of General Chi-Squared Statistics for Tests of Fit," *The Annals of Statistics*, 599–616.
- NIKULIN, M.S. (1973): "Chi-Square Test for Continuous Distributions with Shift and Scale Parameters," *Theory of Probability & Its Applications*, 18, 559–568.
- OSIUS, G. AND D. ROJEK (1992): "Normal Goodness-of-Fit Tests for Multinomial Models With Large Degrees of Freedom," *Journal of the American Statistical Association*, 87, 1145–1152.
- PEARSON, K. (1900): "On the Criterion that a given System of Deviations from the Probable in the Case of a Correlated System of Variables is such that it can be reasonably supposed to have arisen from Random Sampling," *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 50, 157–175.
- POLLARD, D. (1979): "General Chi-square Goodness-of-fit Tests with Data-dependent Cells," *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 50, 317–331.
- RAO, C.R. (2002): *Linear Statistical Inference and Its Applications*, Wiley.
- RAO, K.C. AND B.S. ROBSON (1974): "A Chi-Square Statistic for Goodness-of-Fit Tests Within the Exponential Family," *Communications in Statistics*, 3, 1139–1153.
- READ, T.R.C. AND N.A.C. CRESSIE (1988): *Goodness-of-Fit Statistics for Discrete Multivariate Data*, Springer.
- ROBINSON, P.M. (1988): "The Stochastic Difference Between Econometric Statistics," *Econometrica*, 56, 531–548.
- ROSENBLATT, M. (1952): "Remarks on a Multivariate Transformation," *The Annals of Mathematical Statistics*, 23, 470–472.
- ROY, A.R. (1956): "On χ^2 -Statistics with Variable Intervals," Tech. rep., Department of Statistics, Stanford University, California, technical Report No. 1.
- SHERMAN, J. AND W.J. MORRISON (1950): "Adjustment of an Inverse Matrix Corresponding to a Change in One Element of a Given Matrix," *The Annals of Mathematical Statistics*, 21, 124–127.
- THEODOSSIOU, P. (1998): "Financial Data and the Skewed Generalized T Distribution," *Management Science*, 44, 1650–1661.
- WATSON, G.S. (1957): "The χ^2 Goodness-of-Fit Test for Normal Distributions," *Biometrika*, 336–348.