

Weights characterizing balls via averaging

Nikolay Kuznetsov

Laboratory for Mathematical Modelling of Wave Phenomena,
Institute for Problems in Mechanical Engineering, Russian Academy of Sciences,
V.O., Bol'shoy pr. 61, St Petersburg 199178, Russian Federation
E-mail: nikolay.g.kuznetsov@gmail.com

Abstract

A new theorem characterizing analytically balls in the Euclidean space \mathbb{R}^m is proved. The proof involves just a single weight function, whose volume mean over the domain under consideration is compared with that over a ball having the same volume mean. A class of appropriate test weights is described.

In 1962, Epstein published the one-page long note [1], in which he proved the following theorem.

Let D be a simply connected plane domain of finite area and t a point of D such that, for every function u harmonic in D and integrable over D , the mean value of u over the area of D equals $u(t)$. Then D is a disc and t is its center.

Thereafter, further studies in this area dealt solely with various inverse mean value properties of *harmonic functions* for almost six decades. (The term “inverse mean value property” coined by Hansen and Netuka [2] for assertions, characterizing analytically various domains, became widely accepted.) These studies are reviewed in the extensive survey article [7], Sections 7 and 8; see also [4], where results not covered in [7] are described.

Recently, such properties were established for real-valued solutions of other partial differential equations; namely, the modified Helmholtz equation

$$\nabla^2 v - \mu^2 v = 0, \quad \mu \in \mathbb{R} \setminus \{0\}; \quad (1) \text{ ?MHH?}$$

and the Helmholtz equation $\nabla^2 u + \lambda^2 u = 0$, $\lambda \in \mathbb{R} \setminus \{0\}$; see the notes [3] and [5], respectively. Here $\nabla = (\partial_1, \dots, \partial_m)$ is the gradient operator, $\partial_i = \partial/\partial x_i$. Before formulating one of these results we introduce some notation.

Let $x = (x_1, \dots, x_m)$ be a point in \mathbb{R}^m , $m \geq 2$, by $B_r(x) = \{y \in \mathbb{R}^2 : |y - x| < r\}$ we denote the open ball (disc when $m = 2$) of radius r centred at x . The ball is called admissible with respect to a domain $D \subset \mathbb{R}^m$ provided $\overline{B_r(x)} \subset D$. If D has a finite Lebesgue measure and a function f is integrable over D , then

$$M(f, D) = \frac{1}{|D|} \int_D f(x) dx$$

is its area mean value over D ; here $|D|$ is the is the volume (area) of D .

Now we are in a position to formulate the following.

Theorem 1 (Kuznetsov [3]). *Let $D \subset \mathbb{R}^m$, $m \geq 2$, be a bounded domain, and let $r > 0$ be such that $|B_r| = |D|$. Suppose that there exists a point $x_0 \in D$ such that for some $\mu > 0$ the identity*

$$v(x_0) \Gamma\left(\frac{m}{2} + 1\right) \frac{I_{m/2}(\mu r)}{(\mu r/2)^{m/2}} = M(v, D)$$

holds for every positive function v satisfying equation (1) in the union of D and $\cup_{x \in \partial D} B_r(x)$, then $D = B_r(x_0)$. Here, I_ν denotes the modified Bessel function of order ν , whereas Γ stands for the Gamma function.

The following two-dimensional analogue of this theorem is based on weighted means.

Theorem 2 (Kuznetsov [6]). *Let $D \subset \mathbb{R}^2$ be a bounded domain, and let $r > 0$ be such that $|D| \geq \pi r^2$. If for a point $x_0 \in D$ and some $\mu > 0$ the weighted mean value identity*

$$a(\mu r) v(x_0) = \frac{1}{|D|} \int_D v(y) \log \frac{r}{|x_0 - y|} dy, \quad a(t) = \frac{2[I_0(t) - 1]}{t^2}, \quad (2) \text{ ?MW' ?}$$

holds for every positive function v satisfying equation (1) in the union of D and $\cup_{x \in \partial D} B_r(x)$, then $D = B_r(x_0)$.

Rethinking the proof of this theorem, it becomes clear that balls can be characterized without exploiting solutions of (1) or of any other equation, but just averaging some weight over the domain under investigation is sufficient. The first example of this kind is based on the weight used in Theorem 2.

Theorem 3. *Let $D \subset \mathbb{R}^m$ be a bounded domain. If the identity*

$$\frac{1}{m} = \frac{1}{|D|} \int_D \log \frac{r}{|x - y|} dy \quad (3) \text{ ?MW' ?}$$

holds with $x \in D$ and $r > 0$ such that $|D| \geq |B_r|$, then $D = B_r(x)$.

Proof. One readily obtains by a direct calculation that

$$\frac{1}{m} = \frac{1}{|B_r|} \int_{B_r(x)} \log \frac{r}{|x - y|} dy \quad \text{for every } r > 0 \text{ and } x \in \mathbb{R}^m. \quad (4) \text{ ?MW' ?}$$

Without loss of generality, we suppose that D is located so that x coincides with the origin. Let us show that a contradiction follows from the assumption $D \neq B_r(0)$. For this purpose we consider two open sets $G_i = D \setminus \overline{B_r(0)}$ (nonempty in view of conditions imposed on D and r) and $G_e = B_r(0) \setminus \overline{D}$ (possibly empty); both are bounded.

Let us write (3) as follows:

$$\frac{|D|}{m} = \int_D \log \frac{r}{|y|} dy, \quad (5) \text{ ?1?}$$

Since identity (4) holds for $x = 0$ and r , we write it in the same way:

$$\frac{|B_r|}{m} = \int_{B_r(0)} \log \frac{r}{|y|} dy. \quad (6) \text{ ?2?}$$

Subtracting (6) from (5), we obtain

$$\frac{|D| - |B_r|}{m} = \int_{G_i} \log \frac{r}{|y|} dy - \int_{G_e} \log \frac{r}{|y|} dy. \quad (7) \text{ ?3?}$$

Here the difference on the right-hand side is negative. Indeed, $\log(r/|y|) < 0$ on $G_i \neq \emptyset$, because $|y| > r$ there. Hence, the first term is negative. If $G_e \neq \emptyset$, then the second integral is positive because $\log(r/|y|) > 0$ on G_e , where $|y| < r$, and it vanishes otherwise. On the other hand, the expression on the left-hand side is nonnegative. The obtained contradiction proves the theorem. \square

Another weight for which the same result is true, is as follows:

$$|x - y|^{\alpha-m} - r^{\alpha-m}, \quad \text{where } r > 0 \text{ and } \alpha \in (0, m).$$

By a direct calculation one obtains that

$$\frac{1}{|B_r|} \int_{B_r(x)} [|x - y|^{\alpha-m} - r^{\alpha-m}] \, dy = (m\alpha^{-1} - 1) r^{\alpha-m} \quad \text{for any } x \in \mathbb{R}^m.$$

This value is positive for $r > 0$ in view of the assumption about α , thus allowing us to use it in the proof instead of (4).

In the present note, we describe a class of weights (say W) each of which, under an appropriate condition (the latter concerns the weight's average over a domain and is similar to assumption (4) in Theorem 3), yields that the domain is a ball. Of course, W includes both weights mentioned above.

Definition 1. A continuous function of the form

$$(0, \infty) \times (0, \infty) \ni (t, r) \mapsto w \in \mathbb{R}$$

is called a *weight characterizing balls* ($w \in W$ for short) if

- (i) $w(t, r) > 0$ when $t < r$, $w(t, r) < 0$ when $t > r$ and $w(r, r) = 0$;
- (ii) for every $x \in \mathbb{R}^m$ and every $r > 0$, the superposition $w(|x - y|, r)$ is a locally integrable function of $y \in \mathbb{R}^m$.

If $w \in W$, then it immediately follows that the mean value over a ball, namely,

$$\mathbf{m}(w, r) := M(w(|x - \cdot|, r), B_r(x))$$

is finite, positive and independent of $x \in \mathbb{R}^m$ for every $r > 0$. This allows us to generalise Theorem 3 as follows.

Theorem 4. Let $D \subset \mathbb{R}^m$ be a bounded domain and let $w \in W$. If the identity

$$\mathbf{m}(w, r) = \frac{1}{|D|} \int_D w(|x - y|, r) \, dy$$

holds with $x \in D$ and $r > 0$ such that $|D| \geq |B_r|$, then $D = B_r(x)$.

The proof by contradiction literally repeats that of Theorem 3, but equality (7) must be changed to

$$(|D| - |B_r|) \mathbf{m}(w, r) = \left[\int_{G_i} - \int_{G_e} \right] w(|y|, r) \, dy.$$

Here the difference on the right-hand side is negative according to condition (i). Indeed, the integrand is negative on $G_i \neq \emptyset$, whereas the second integral is positive if $G_e \neq \emptyset$ and vanishes otherwise. On the other hand, the expression on the left-hand side is nonnegative.

To provide a nonsingular illustration of this theorem we consider the weight function

$$w_\alpha(t, r) = r^\alpha - t^\alpha \quad \text{with } \alpha > 0.$$

It is clear that it satisfies conditions of Definition 1, whereas

$$\mathbf{m}(w_\alpha, r) = \left[1 - \frac{m}{m + \alpha}\right] r^\alpha > 0 \quad \text{when } r > 0,$$

is independent of $x \in \mathbb{R}^m$. Hence, Theorem 4 with $w = w_\alpha$ yields another characterization of balls via averaging this weight over D with $|D| \geq |B_r|$.

It is easy to construct more examples of weights (singular and nonsingular) that satisfy conditions of Definition 1, thus characterizing balls via averaging.

References

- [E][1] B. Epstein, “On the mean-value property of harmonic functions”, *Proc. Amer. Math. Soc.* **13** (1962), 830.
- [HN1][2] W. Hansen, I. Netuka, “Inverse mean value property of harmonic functions”, *Math. Ann.* **297** (1993), 147–156. “Corrigendum”, *Math. Ann.* **303** (1995), 373–375.
- [Ku1][3] N. Kuznetsov, “Characterization of balls via solutions of the modified Helmholtz equation”, *Comptes Rendus Math.* **359** (2021), 945–948.
- [Ku3][4] N. Kuznetsov, “Inverse mean value properties (a survey)”, *J. Math. Sci.* **262** (2022), 275–290.
- [Ku2][5] N. Kuznetsov, “On characterization of balls via solutions to the Helmholtz equation”, *J. Math. Sci.* **264** (2022), 603–608.
- [Ku4][6] N. Kuznetsov, “Characterizations of discs via weighted means”, Preprint arXiv:2209.10281v1 (2022).
- [NV][7] I. Netuka, J. Veselý, “Mean value property and harmonic functions”, *Classical and Modern Potential Theory and Applications*, Kluwer, Dordrecht, 1994, pp. 359–398.