

SCALING LIMIT OF A LONG-RANGE RANDOM WALK IN TIME-CORRELATED RANDOM ENVIRONMENT

GUANGLIN RANG, JIAN SONG, AND MENG WANG

ABSTRACT. This paper concerns a long-range random walk in random environment in dimension $1 + 1$, where the environmental disorder is independent in space but correlated in time. We prove that the rescaled partition function converges weakly to the Stratonovich solution of a fractional stochastic heat equation with multiplicative Gaussian noise which is white in space and colored in time.

CONTENTS

1. Introduction	1
1.1. Notations and known results	2
1.2. Model description	3
1.3. Main result, strategy and discussions	4
2. On the SPDE	7
2.1. Preliminaries on Gaussian spaces	7
2.2. Mild Stratonovich solution	12
3. On weak convergences	16
3.1. Weak convergence of U-statistics	17
3.2. Weak convergence of rescaled partition functions	19
3.3. L^1 -bound of rescaled partition functions	23
Appendix A. Physical Wick product	25
Appendix B. Some preliminaries on convergence of probability measures	27
Appendix C. Miscellaneous results	28
References	30

1. INTRODUCTION

The model of directed polymer in random environment was first introduced by Huse and Henley [20] in the study of the Ising model and became a canonical model for disordered systems (see e.g. the lecture notes by Comets [8]). In recent years, it has attracted much attention in particular due to its intimate connections to the stochastic heat equation (parabolic Anderson model), the stochastic Burgers equation, and the Kardar-Parisi-Zhang (KPZ) equation and its universality class (see [8] and [9] for a review).

For a simple symmetric random walk in i.i.d. random environment on $\mathbb{Z}_+ \times \mathbb{Z}$, Alberts et al. [1] proved that the rescaled partition function converges weakly to the *Itô-Skorohod* solution of the stochastic heat equation (SHE) with multiplicative space-time white Gaussian noise. This result was extended by Caravenna et al. [6] to a long-range random walk in i.i.d. random environment and to other disordered models (disordered pinning model and random field Ising model). Recently, Rang [32] proved that, for a simple random walk in random environment which is white in time but correlated in space, the rescaled partition function converges weakly to the *Itô-Skorohod* solution of SHE with Gaussian noise white in time and colored in space; if the random environment is given by the occupation field of a Poisson system of independent random walks on \mathbb{Z} which is now correlated in both time and space, Shen et al. [36] showed that the scaling limit is the *Stratonovich* solution of SHE with space-time colored Gaussian noise whose covariance coincides with the heat kernel. It is suggested by the results and methodologies in the above-mentioned papers that the *temporal independence* of the random environment plays a critical rule when identifying the scaling limits for partition functions, which will be further discussed in Section 1.3.

In this paper, we aim to study the scaling limit of the partition function for a long-range random walk in random environment which is independent in space but correlated in time on $\mathbb{Z}_+ \times \mathbb{Z}$.

We remark that the model of directed polymer in space-time correlated random environment whose covariance has a power-law decay was first considered in physics literature by Medina et al. [28] where the Burgers equation with colored noise was studied and then applied to analyse directed polymer and interface growth. We also would like to mention that Rovira and Tindel [34] introduced Brownian polymer in a centered Gaussian field that is white in time and correlated in space on $\mathbb{R}_+ \times \mathbb{R}$ and studied the asymptotic behavior of the partition function; in the two subsequent papers, Bezerra et al. [4] obtained the superdiffusivity and Lacoïn [26] investigated the effect of strong spatial correlation. Finally, we remind the reader that the model of long-range directed polymer has been studied by Comets [7], and more recently by Wei [39].

1.1. Notations and known results. For the reader's convenience, we collect the mathematical notations that will be used throughout this article.

Notations 1.1. Let \mathbb{N} denote the set of natural numbers without 0, i.e., $\mathbb{N} \stackrel{\text{def}}{=} \{1, 2, \dots\}$; for $N \in \mathbb{N}$, $[[N]] \stackrel{\text{def}}{=} \{1, 2, \dots, N\}$; for $a \in \mathbb{R}$, $[a]$ means the greatest integer that is not greater than a ; $\|\cdot\|$ is used for the Euclidean norm; let $\mathbf{k} \stackrel{\text{def}}{=} (k_1, \dots, k_d)$, $\mathbf{l} \stackrel{\text{def}}{=} (l_1, \dots, l_d)$, $\mathbf{x} \stackrel{\text{def}}{=} (x_1, \dots, x_d)$, $\mathbf{y} \stackrel{\text{def}}{=} (y_1, \dots, y_d)$ etc stand for vectors in \mathbb{Z}^d or \mathbb{R}^d depending on the context; we use C to denote a generic positive constant that may change from line to line; we say $f(x) \lesssim g(x)$ if $f(x) \leq Cg(x)$ for all x ; we write $f(x) \sim g(x)$ (as $x \rightarrow \infty$), if $\lim_{x \rightarrow \infty} f(x)/g(x) = 1$. We use \xrightarrow{d} to denote the convergence in distribution (also called weak convergence) for random variables/vectors. For a random variable X , $\|X\|_{L^p} = (\mathbb{E}[|X|^p])^{1/p}$ for $p \geq 1$.

Let $S = \{S_i, i \in \mathbb{N}\}$ be a random walk in \mathbb{Z} and $\omega = \{\omega(i, k), (i, k) \in \mathbb{N} \times \mathbb{Z}\}$ be a family of random variables independent of S serving as the random environment (disorder). We shall use \mathbb{P}_S and \mathbb{E}_S (resp. \mathbb{P}_ω and \mathbb{E}_ω) to denote the probability and expectation in the probability space of S (resp. ω), respectively. The probability and expectation in the product probability space of (S, ω) is denoted by \mathbb{P} and \mathbb{E} , respectively.

The *partition function* is defined by

$$(1.1) \quad Z_\omega^{(N)}(\beta) = \sum_S e^{\beta \sum_{i=1}^N \omega(i, S_i)} \mathbb{P}(S) = \mathbb{E}_S \left[e^{\beta \sum_{i=1}^N \omega(i, S_i)} \right],$$

where $\beta > 0$ is the inverse temperature, and the *point-to-line partition function* (or *partition function* for simplicity) is given by

$$(1.2) \quad Z_\omega^{(N)}(\beta, k) \stackrel{\text{def}}{=} \mathbb{E}_S \left[e^{\beta \sum_{i=1}^N \omega(i, S_i)} \middle| S_0 = k \right], \quad \text{for } k \in \mathbb{Z}.$$

When S is a simple symmetric random walk with i.i.d. increments and $\omega(i, k)$ are i.i.d. random variables with $\lambda(\beta) \stackrel{\text{def}}{=} \log \mathbb{E}[e^{\beta \omega(i, k)}] < \infty$ for some β sufficiently small, Alberts et al. [1] introduced the so-called intermediate disorder regime. More precisely, if β is scaled in the way

$$\beta \rightarrow \hat{\beta}_N \stackrel{\text{def}}{=} \beta N^{-\frac{1}{4}},$$

one has the following weak convergence

$$e^{-N\lambda(\hat{\beta}_N)} Z_\omega^{(N)}(\hat{\beta}_N, N^{1/2}x) \xrightarrow{d} u(1, x) \text{ as } N \rightarrow \infty,$$

where $u(t, x)$ is the Itô-Skorohod solution of the equation

$$(1.3) \quad \begin{cases} \frac{\partial u(t, x)}{\partial t} = \frac{1}{2} \Delta u(t, x) + \sqrt{2} \beta u(t, x) \dot{W}(t, x), & t > 0, x \in \mathbb{R}, \\ u(0, x) = 1. \end{cases}$$

Here, \dot{W} is space-time Gaussian white noise and Itô-Skorohod solution means that the product in $u(t, x) \dot{W}(t, x)$ is a Wick product, or equivalently the associated stochastic integral is an Itô-Skorohod integral.

Under the same setting except that the random walk S may be long-range (i.e., the increments of S now lie in the domain of attraction of a ρ -stable law with $\rho \in (1, 2]$), the result of [1] was extended in Caravenna et al. [6] where a unified framework based on the Lindeberg principle for polynomial chaos expansion was developed to study the scaling limits for pinning model, directed polymer model, and Ising model. Still using the setting of [1] but assuming that the disorder $\{\omega(i, k), (i, k) \in \mathbb{N} \times \mathbb{Z}\}$ is correlated in space (but still independent in time), Rang [32] obtained the weak convergence of the rescaled partition function to the Itô-Skorohod solution of (1.3) with the Gaussian noise $\dot{W}(t, x)$ being colored in space (still white in time).

1.2. Model description. Motivated by the above-mentioned works, we aim to study the weak convergence of the rescaled partition function for a long-range random walk $\{S_n, n \in \mathbb{N}\}$ in random environment on $\mathbb{N} \times \mathbb{Z}$, where the disorder $\{\omega(i, k), i \in \mathbb{N}, k \in \mathbb{Z}\}$ is correlated in time and independent in space.

In our model, the directed polymer (random walk) is given by $S_n = S_0 + \sum_{i=1}^n Y_i$, where $S_0 \in \mathbb{Z}$ and $\{Y_i\}_{i=1}^\infty$ are independent and identically distributed random variables which have a 1-lattice distribution belonging to the domain of attraction of a ρ -stable distribution with density function denoted by $\mathfrak{g}_\rho(\cdot)$. We assume $1 < \rho \leq 2$ and

$$(1.4) \quad \begin{cases} \mathbb{P}(Y_i = k) \lesssim |k|^{-1-\rho} \text{ for } k \in \mathbb{Z} \setminus \{0\}, & \text{if } \rho \in (1, 2), \\ \mathbb{E}[Y_i] = 0 \text{ and } \mathbb{E}[Y_i^2] = 1, & \text{if } \rho = 2. \end{cases}$$

Let $\psi(u) \stackrel{\text{def}}{=} \mathbb{E}[e^{uY_i}]$ be the characteristic function of Y_i . Then the 1-lattice distribution of Y_i implies that $\psi(u)$ is periodic with period 2π , and furthermore, $|\psi(u)| < 1$ for all $u \in [-\pi, \pi] \setminus \{0\}$ (see, e.g., [21, Theorem 1.4.2]); this property of $\psi(u)$ usually makes analysis easier. The assumption of 1-lattice distribution on Y_i also yields that the random walk $S_n = \sum_{i=1}^n Y_i$ has period 1 and hence is *aperiodic*. We remind the reader that a simple symmetric random walk has period 2.

Denote by $P_n(k) = \mathbb{P}(S_n = k)$ for $n \in \mathbb{N}, k \in \mathbb{Z}$ the probability of S being at k at time n . Then by the local limit theorem for ρ -stable distribution (see [35, Theorem 6.1]), the convergence

$$(1.5) \quad n^{1/\rho} P_n(k) - \mathfrak{g}_\rho(k/n^{1/\rho}) \rightarrow 0$$

holds uniformly in k . Noting that \mathfrak{g}_ρ is bounded, (1.5) yields

$$(1.6) \quad P_n(k) \lesssim n^{-1/\rho}.$$

It is well known that

$$\frac{S_n}{n^{1/\rho}} \xrightarrow{d} \xi, \text{ as } n \rightarrow \infty,$$

where ξ has the symmetric ρ -stable distribution with characteristic function $\exp\{-c_\rho|\eta|^\rho\}$ for some $c_\rho > 0$. Letting $\mathfrak{g}_\rho(t, x)$ be the density function of the corresponding ρ -stable process, we have

$$\int_{\mathbb{R}} e^{\eta x} \mathfrak{g}_\rho(t, x) dx = e^{-c_\rho t |\eta|^\rho}.$$

Throughout the rest of the paper we will omit the subscript ρ and use $\mathfrak{g}(t, x)$ exclusively to denote the density function of the ρ -stable process X . Note that \mathfrak{g} has the following scaling property and the upper bound

$$(1.7) \quad \mathfrak{g}(t, x) = t^{-\frac{1}{\rho}} \mathfrak{g}(1, t^{-\frac{1}{\rho}} x) \lesssim t^{-\frac{1}{\rho}}.$$

We assume that the disorder in the environment is given by a family of Gaussian random variables $\{\omega(n, k), n \in \mathbb{N}, k \in \mathbb{Z}\}$ with mean zero and covariance

$$(1.8) \quad \mathbb{E}[\omega(n, k)\omega(n', k')] = \gamma(n - n')\delta_{kk'},$$

where $\delta_{kk'}$ is the Kronecker delta function, i.e., $\delta_{kk'} = 1$ if $k = k'$ and $\delta_{kk'} = 0$ otherwise, and $\gamma(n)$ has a power law decay:

$$(1.9) \quad \gamma(n) \lesssim |n|^{2H-2} \wedge 1, \text{ for } n \in \mathbb{Z},$$

where $H \in (1/2, 1]$ denotes a fixed constant throughout this paper. We further assume that for all $t \in \mathbb{R} \setminus \{0\}$,

$$(1.10) \quad \lim_{N \rightarrow \infty} N^{2-2H} \gamma([Nt]) = |t|^{2H-2}.$$

1.3. Main result, strategy and discussions. Consider the stochastic fractional heat equation on \mathbb{R}

$$(1.11) \quad \begin{cases} \frac{\partial u(t, x)}{\partial t} = -(-\Delta)^{\frac{\rho}{2}} u(t, x) + \beta u(t, x) \dot{W}(t, x), \\ u(0, x) = 1, \end{cases}$$

where $\rho \in (1, 2]$ and $\dot{W}(t, x)$ is Gaussian noise with covariance function given by

$$(1.12) \quad \mathbb{E}[\dot{W}(t, x)\dot{W}(s, y)] = K(t - s, x - y) \stackrel{\text{def}}{=} |t - s|^{2H-2} \delta(x - y),$$

with $H \in (1/2, 1]$ and $\delta(\cdot)$ being the Dirac delta function.

The following is the main result of this paper:

Theorem 1.2. *Let the random walk S and the disorder ω be given as in Section 1.2. Let H and ρ be parameters satisfying*

$$(1.13) \quad H \in (1/2, 1], \rho \in (1, 2], \text{ and } \theta \stackrel{\text{def}}{=} H - \frac{1}{2\rho} > \frac{1}{2}.$$

Consider the rescaled partition function $Z_\omega^{(N)}(\hat{\beta}_N, N^{1/\rho}x_0)$ given in (1.2) under the scaling

$$\beta \rightarrow \hat{\beta}_N \stackrel{\text{def}}{=} \beta N^{-\theta}.$$

Let $u(t, x)$ be the Stratonovich solution of (1.11). Then we have

$$Z_\omega^{(N)}(\hat{\beta}_N, N^{1/\rho}x_0) \xrightarrow{d} u(1, x_0), \text{ as } N \rightarrow \infty.$$

In the Theorem above, Stratonovich solution means that the product $u(t, x)\dot{W}(t, x)$ in (1.11) is an ordinary product, or equivalently the associated stochastic integral is a Stratonovich integral. See Section 2.1 for details.

The proof of Theorem 1.2 will be presented in Section 3. For the reader's convenience, here we explain the strategy briefly. By Taylor's expansion, the rescaled partition function can be written as

$$Z_\omega^{(N)}(\hat{\beta}_N, N^{1/\rho}x_0) = \sum_{m=0}^{\infty} \frac{1}{m!} \mathbb{S}_m^{(N)},$$

where

$$(1.14) \quad \mathbb{S}_m^{(N)} = \hat{\beta}_N^m \sum_{n_1, \dots, n_m \in \llbracket N \rrbracket} \sum_{k_1, \dots, k_m \in \mathbb{Z}} \omega(n_1, k_1) \cdots \omega(n_m, k_m) P_{\mathbf{n}^*}(N^{1/\rho}x_0; k_1, \dots, k_m),$$

with $P_{\mathbf{n}^*}$ (see (3.4)) being the product of the transition densities of the random walk S . Meanwhile, the solution of the continuum equation (1.11) has the series representation (see equation (2.27)):

$$u(1, x_0) = \sum_{m=0}^{\infty} \beta^m \mathbb{I}_m(\mathbf{g}_m(x_0; \cdot)),$$

where $\mathbb{I}_m(\cdot)$ is a multiple Stratonovich integral (see Section 2.1.2) and \mathbf{g}_m (see (2.28)) is the product of the transition densities of the limiting stable process X . We remind the reader that the continuum multiple Stratonovich integral $\mathbb{I}_m(\mathbf{g}_m(x_0; \cdot))$ resembles the discrete sum $\mathbb{S}_m^{(N)}$ up to a time reversal.

Under condition (1.13), one can obtain the L^1 -convergence of the series of $u(t, x)$ (see Proposition 2.10) and the uniform (in N) L^1 -convergence of the series of $Z_\omega^{(N)}(\hat{\beta}_N, N^{1/\rho}x_0)$ (see (3.37)). Thus, in order to prove the weak convergence $Z_\omega^{(N)}(\hat{\beta}_N, N^{1/\rho}x_0) \xrightarrow{d} u(1, x_0)$, in light of Lemma B.1, it suffices to obtain joint weak convergences for $\mathbb{S}_m^{(N)}$'s, that is, for $k \in \mathbb{N}$ and $l_1, \dots, l_k \in \mathbb{N}$,

$$\left(\frac{1}{l_1!} \mathbb{S}_{l_1}^{(N)}, \dots, \frac{1}{l_k!} \mathbb{S}_{l_k}^{(N)} \right) \xrightarrow{d} \left(\beta^{l_1} \mathbb{I}_{l_1}(\mathbf{g}_{l_1}), \dots, \beta^{l_k} \mathbb{I}_{l_k}(\mathbf{g}_{l_k}) \right), \text{ as } N \rightarrow \infty,$$

which is proved in Proposition 3.4.

Multiple Wiener integrals $\mathbf{I}_m(f)$ and multiple Stratonovich integrals $\mathbb{I}_m(f)$ are linked via the celebrated Hu-Meyer's formula (2.19). Usually it is more convenient to deal with multiple Wiener integrals $\mathbf{I}_m(f)$ whose second moment is easier to calculate. For this reason, in Section 3.1 we first consider U -statistics given in (3.6) which are multi-linear Wick polynomials of $\omega(n_i, k_i)$'s, and prove that they converge weakly to multiple Wiener integrals (see Proposition 3.3). This together with Hu-Meyer's formula (2.19) then yields Proposition 3.4.

As can be seen, our approach is very much inspired by [1] where the random variables of disorder are i.i.d. However, in our setting the disorder is correlated in time, and it turns out that this difference is critical in the sense that it prevents us from using the trick of "modified partition function" (see in [1, equation (4)], which was also employed in [32]), as this method relies on the temporal independence of the disorder. For the same reason, the Mayer expansion (see [6, equations (1.5) and (1.7)]), which was employed in [6] to linearize the exponential while introducing an error

term that comes from the Itô correction, cannot be applied here either, since it works well only if the disorder is independent in both time and space.

We conclude the introduction by making a few remarks on Theorem 1.2 and related results.

(i) In the polymer model, the temporal independence of the disorder plays a critical role. Assuming that the disorder is correlated in time and white in space in our setting, the rescaled partition function converges weakly to a *Stratonovich* solution. Recently, directed polymer in time-space correlated random environment was studied in [36] and it was shown that the rescaled partition function converges weakly to a Stratonovich solution. In contrast, if the disorder possesses temporal independence, after a proper scaling, the limit of the rescaled partition function is an *Itô-Skorohod* solution, as has been shown in [1, 6, 32]. Among others, a technical explanation for the critical role of the temporal independence of the disorder is the following: when the temporal independence of disorder appears, by using the “modified partition function” trick, the term corresponding to $\mathbb{S}_m^{(N)}$ given by (1.14) becomes a multi-linear polynomial of $\omega(n_i, k_i)$ for $i = 1, \dots, m$ with $n_1 < n_2 < \dots < n_m$, which converges weakly to a multiple Wiener integral due to the independence (see [29, 6]).

(ii) In contrast to the case that the disorder is independent in time where one can employ the “modified partition function” trick (or Mayer’s expansion for the case that the disorder is independent both in time and space) to make time indices in the summation distinct from each other, we expand the partition function directly via Taylor’s expansion, and as a consequence the time indices n_1, \dots, n_m in the summation (1.14) can be repeated. It turns out that this in fact gives negligible contribution due to the condition (1.13) which is more restrictive than the conditions proposed in [1, 6, 32]. For instance, when $m = 2$, the expectation of the sum of the diagonal terms in (1.14) is $\beta\gamma(0)N^{1-2\theta}$ which converges to zero as N tends to infinity, due to the condition $\theta > 1/2$.

(iii) As indicated in [28], it is also of interest to study directed polymer in space-time correlated random environment. In light the result of [36] and the discussion in (i), we expect that our result and methodology can be extended to the model in dimension $1 + d$ with disorder induced by space-time fractional Brownian motion whose Hurst parameters are bigger than $1/2$. Note that the corresponding continuum SPDEs have been investigated in [19, 37].

(iv) In our model, we assume the Gaussianity of the disorder ω for technical reasons. The loss of Gaussian property in ω shall cause a lot more complexity in computations (see e.g. (A.7) and (A.8) for a comparison of the computations of moments for Gaussian and non-Gaussian random variables). The functional analytic approach developed in [36] might be a choice to circumvent this difficulty. Nevertheless, we conjecture that our result still holds for sub-Gaussian disorder.

(v) By Skorohod’s representation theorem, if the disorder ω and the limiting noise \dot{W} live in the same probability space, we can get strong convergence (a.s. convergence) in Theorem 1.2. In this sense, Theorem 1.2 can be also considered as an approximation result for the equation (1.11). See e.g. Foondun et al. [12] and Joseph et al. [24] for related results.

(vi) As a byproduct, we obtain a Jensen type inequality for the integrals induced by fractional Brownian motion (see Lemmas 2.2 and C.1) which is new in the literature to our best knowledge. We provide a new approach to prove the exponential integrability of the weighted self-intersection local time of the ρ -stable process in Proposition 2.6 which plays a critical rule in obtaining the Feynman-Kac formula for the solution of (1.11); this approach also simplifies the proof in [19] which works exclusively for the weighted self-intersection local time of Brownian motion.

The rest of the paper is organized as follows. In Section 2, we provide some preliminaries on Gaussian spaces and then study the limiting SPDE (1.11). The main result Theorem 1.2 is proved in Section 3. In Appendices A, B, and C, we collect some preliminaries on Wick products, convergence of probability measures, and some other miscellaneous results that are used in the analysis.

2. ON THE SPDE

In this section, we first recall some preliminaries on Gaussian spaces and then we study the limiting continuum SPDE (1.11).

2.1. Preliminaries on Gaussian spaces. In this subsection, we provide some preliminaries on Gaussian spaces. We refer to [18, 22, 30] for more details.

2.1.1. *Banach spaces associated with \dot{W} .* On probability space (Ω, \mathcal{F}, P) satisfying usual conditions, let $\dot{W} = \{\dot{W}(t, x) : t \in [0, 1], x \in \mathbb{R}\}$ be real-valued centered Gaussian noise with covariance

$$(2.1) \quad \mathbb{E}[\dot{W}(t, x)\dot{W}(s, y)] = K(t - s, x - y) = |t - s|^{2H-2}\delta(x - y),$$

where δ is the Dirac delta function. The Hilbert space \mathcal{H} associated with \dot{W} is the completion of smooth functions with compact support under the inner product

$$(2.2) \quad \begin{aligned} \langle f, g \rangle_{\mathcal{H}} &= \int_0^1 \int_0^1 \int_{\mathbb{R}} \int_{\mathbb{R}} f(s, x)K(s - t, x - y)g(t, y)dx dy ds dt \\ &= \int_0^1 \int_0^1 \int_{\mathbb{R}} |s - t|^{2H-2} f(s, x)g(t, x)dx ds dt. \end{aligned}$$

We assume $H \in (1/2, 1]$. For the case $H = 1$, the noise $\dot{W}(x)$ is indeed a spatial white noise which does not depend on time.

We remark that the Hilbert space \mathcal{H} contains distributions (see [31]). For our purpose, it suffices to just consider classical measurable functions. We introduce the following Banach space \mathcal{B} which is a subset of the set $\mathcal{B}([0, 1] \times \mathbb{R})$ of measurable functions on $[0, 1] \times \mathbb{R}$.

Definition 2.1. *We define*

$$(2.3) \quad \mathcal{B} \stackrel{\text{def}}{=} \left\{ f \in \mathcal{B}([0, 1] \times \mathbb{R}) : \|f\|_{\mathcal{B}} \stackrel{\text{def}}{=} \left(\int_0^1 \int_0^1 \int_{\mathbb{R}} |s - t|^{2H-2} |f(s, x)| |f(t, x)| dx ds dt \right)^{\frac{1}{2}} < \infty \right\}.$$

Clearly \mathcal{B} is a dense subset of \mathcal{H} , and if $f \in \mathcal{H}$ is a nonnegative function, f also belongs to \mathcal{B} .

For $N \in \mathbb{N}$ and $f \in \mathcal{B}^{\otimes m}$, we denote by $\mathcal{A}_N(f)$ the conditional expectation of f with respect to the σ -algebra $\mathcal{B}_N^m = \sigma(\mathfrak{R}_N^m)$ generated by the set \mathfrak{R}_N^m of rectangles

$$(2.4) \quad \mathfrak{R}_N^m := \left\{ \left(\frac{\mathbf{i} - \vec{1}_m}{N}, \frac{\mathbf{i}}{N} \right] \times \left(\frac{\mathbf{k} - \vec{1}_m}{N^{1/\rho}}, \frac{\mathbf{k}}{N^{1/\rho}} \right] : \mathbf{i} \in \llbracket N \rrbracket^m, \mathbf{k} \in \mathbb{Z}^m \right\},$$

where $\vec{1}_m$ is the m -dimensional vector of all ones. That is, $\mathcal{A}_N(f)$ is defined by the average values of f on the blocks $B \in \mathfrak{R}_N^m$:

$$(2.5) \quad \mathcal{A}_N(f)(\mathbf{t}, \mathbf{x}) = \frac{1}{|B|} \int_B f(\mathbf{s}, \mathbf{y}) ds d\mathbf{y} \cdot \mathbf{1}_B(\mathbf{t}, \mathbf{x}),$$

where $\mathbf{1}_B$ is an indicator function and $|B|$ is the Lebesgue measure of B .

The following Jensen type of inequality will be used to prove the weak convergence of U-statistics in Section 3.1.

Lemma 2.2. *For $m, N \in \mathbb{N}$, consider $f \in \mathcal{B}^{\otimes m}$ and let $\mathcal{A}_N(f)$ be given by (2.5). Then, we have*

$$(2.6) \quad \|\mathcal{A}_N(f)\|_{\mathcal{B}^{\otimes m}} \leq C^m \|f\|_{\mathcal{B}^{\otimes m}},$$

where C is a constant depending only on H .

Proof We prove (2.6) for $m = 1$ and the general case can be proved in a similar way. Note that by Fubini's theorem, $\mathcal{A}_N(f) = \mathcal{A}_N^t(\mathcal{A}_N^x(f)) = \mathcal{A}_N^x(\mathcal{A}_N^t(f))$, where \mathcal{A}_N^t (resp. \mathcal{A}_N^x) means taking average in time (resp. space) only. Therefore, we have

$$\|\mathcal{A}_N(f)\|_{\mathcal{B}} = \|\mathcal{A}_N^t(\mathcal{A}_N^x(f))\|_{\mathcal{B}} \leq C\|\mathcal{A}_N^x(f)\|_{\mathcal{B}},$$

for some C depending on H only, where the inequality follows from Lemma C.1.

Now, it suffices to prove $\|\mathcal{A}_N^x(f)\|_{\mathcal{B}} \leq \|f\|_{\mathcal{B}}$. Using the identity, for $H \in (\frac{1}{2}, 1)$,

$$|s - t|^{2H-2} = c_H \int_{\mathbb{R}} |s - \tau|^{H-\frac{3}{2}} |t - \tau|^{H-\frac{3}{2}} d\tau$$

where c_H is a finite number only depending on H , we have

$$\begin{aligned} \|\mathcal{A}_N^x(f)\|_{\mathcal{B}}^2 &= c_H \int_{\mathbb{R}} dx \int_0^1 ds \int_0^1 dt \int_{\mathbb{R}} d\tau |s - \tau|^{H-\frac{3}{2}} |t - \tau|^{H-\frac{3}{2}} |\mathcal{A}_N^x(f)|(s, x) |\mathcal{A}_N^x(f)|(t, x) \\ &= c_H \int_{\mathbb{R}} dx \int_{\mathbb{R}} d\tau \left(\int_0^1 ds |s - \tau|^{H-\frac{3}{2}} |\mathcal{A}_N^x(f)|(s, x) \right)^2 \\ &\leq c_H \int_{\mathbb{R}} dx \int_{\mathbb{R}} d\tau \left(\mathcal{A}_N^x \left(\int_0^1 ds |s - \tau|^{H-\frac{3}{2}} |f|(s, \cdot) \right) (\tau, x) \right)^2 \\ &\leq c_H \int_{\mathbb{R}} dx \int_{\mathbb{R}} d\tau \left(\int_0^1 ds |s - \tau|^{H-\frac{3}{2}} |f|(s, x) \right)^2 \\ &= \|f\|_{\mathcal{B}}^2, \end{aligned}$$

where the last inequality follows from the classical Jensen's inequality. \square

2.1.2. *Chaos expansion, Wick product, multiple integrals, etc.* Recall that \mathcal{H} is the Hilbert space associated with Gaussian noise \dot{W} with covariance (2.1). Let $\{W(f), f \in \mathcal{H}\}$ be an isonormal Gaussian process with covariance

$$(2.7) \quad \mathbb{E}[W(f)W(g)] = \langle f, g \rangle_{\mathcal{H}}.$$

In particular, if $f = \mathbf{1}_{[0,t] \times [0,x]}$, we denote

$$W(t, x) \stackrel{\text{def}}{=} W(\mathbf{1}_{[0,t] \times [0,x]}).$$

Then the fractional noise $\dot{W}(t, x)$ can be viewed as the partial derivative $\frac{\partial^2}{\partial t \partial x} W(t, x)$ in the sense of distribution. For $f \in \mathcal{H}$, we also use the integral form to denote the Wiener integral $W(f)$:

$$\int_0^1 \int_{\mathbb{R}} f(s, y) W(ds, dy) \stackrel{\text{def}}{=} W(f).$$

For $m \in \mathbb{N} \cup \{0\}$, let

$$(2.8) \quad H_m(x) \stackrel{\text{def}}{=} (-1)^m e^{x^2/2} \frac{d^m}{dx^m} e^{-x^2/2}, \quad x \in \mathbb{R}$$

be the m th Hermite polynomial. For $g \in \mathcal{H}$ with $\|g\|_{\mathcal{H}} = 1$, the multiple Wiener integral of $g^{\otimes m} \in \mathcal{H}^{\otimes m}$ can be defined via (see e.g. [17, 30])

$$(2.9) \quad \mathbf{I}_m(g^{\otimes m}) \stackrel{\text{def}}{=} H_m(W(g)).$$

In particular, we have $W(g) = \mathbf{I}_1(g)$ and $\mathbf{I}_m(g^{\otimes m}) =: W(g)^m$, where $:W(g)^m$ is a physical Wick product (see Section A).

For $f \in \mathcal{H}^{\otimes m}$, let \hat{f} be its symmetrization, i.e.,

$$\hat{f}(t_1, x_1, \dots, t_m, x_m) = \frac{1}{m!} \sum_{\sigma \in \mathcal{P}_m} f(t_{\sigma(1)}, x_{\sigma(1)}, \dots, t_{\sigma(m)}, x_{\sigma(m)}),$$

where \mathcal{P}_m is the set of all permutations of $\llbracket m \rrbracket = \{1, 2, \dots, m\}$. Let $\mathcal{H}^{\hat{\otimes} m}$ be the symmetrization of $\mathcal{H}^{\otimes m}$. Then for $f \in \mathcal{H}^{\hat{\otimes} m}$, one can define the m th multiple Wiener integral $\mathbf{I}_m(f)$ via (2.9) by a limiting argument. Moreover, for $f \in \mathcal{H}^{\hat{\otimes} m}$ and $g \in \mathcal{H}^{\hat{\otimes} n}$, we have

$$(2.10) \quad \mathbb{E}[\mathbf{I}_m(f)\mathbf{I}_n(g)] = m! \langle f, g \rangle_{\mathcal{H}^{\otimes m}} \delta_{mn},$$

where we recall that δ_{mn} is the Kronecker delta function. For $f \in \mathcal{H}^{\otimes m}$ which is not necessarily symmetric, we simply let $\mathbf{I}_m(f) \stackrel{\text{def}}{=} \mathbf{I}_m(\hat{f})$. We also take the following notation for multiple Wiener integrals:

$$(2.11) \quad \int_{([0,1] \times \mathbb{R})^m} f(\mathbf{t}, \mathbf{x}) W(dt_1, dx_1) \diamond \dots \diamond W(dt_m, dx_m) \stackrel{\text{def}}{=} \mathbf{I}_m(f), \text{ for } f \in \mathcal{H}^{\otimes m}.$$

For $f \in \mathcal{H}^{\hat{\otimes} m}$ and $g \in \mathcal{H}^{\hat{\otimes} n}$, their contraction of r indices for $1 \leq r \leq m \wedge n$ is defined by

$$\begin{aligned} & (f \otimes_r g)(t_1, x_1, \dots, t_{m+n-2r}, x_{m+n-2r}) \\ & \stackrel{\text{def}}{=} \int_{[0,1]^{2r}} \int_{\mathbb{R}^{2r}} f(t_1, x_1, \dots, t_{n-r}, x_{n-r}, u_1, y_1, \dots, u_r, y_r) \prod_{i=1}^r K(u_i - v_i, y_i - z_i) \\ & \quad \times g(t_{n-r+1}, x_{n-r+1}, \dots, t_{n+m-2r}, x_{n+m-2r}, v_1, z_1, \dots, v_r, z_r) dy dz du dv, \end{aligned}$$

and we have the following recursive formulas:

$$(2.12) \quad \mathbf{I}_m(f)\mathbf{I}_n(g) = \sum_{r=0}^{m \wedge n} r! \binom{n}{r} \binom{m}{r} \mathbf{I}_{m+n-2r}(f \otimes_r g).$$

In particular, when $n = 1$ we have,

$$(2.13) \quad \mathbf{I}_m(f)\mathbf{I}_1(g) = \mathbf{I}_{m+1}(f \otimes g) + m\mathbf{I}_{m-1}(f \otimes_1 g).$$

Square integrable random variables have a unique chaos expansion as stated below.

Proposition 2.3. *Let \mathcal{G} be the σ -field generated by W . Then for any $F \in L^2(\Omega, \mathcal{G}, P)$, it admits a unique chaos expansion*

$$F = \sum_{m=0}^{\infty} \mathbf{I}_m(f_m) \quad \text{with } f_m \in \mathcal{H}^{\hat{\otimes} m},$$

where the series converges in L^2 . Moreover,

$$\mathbb{E}[F^2] = \sum_{m=0}^{\infty} m! \|f_m\|_{\mathcal{H}^{\otimes m}}^2.$$

For $f \in \mathcal{H}^{\otimes m}$ and $g \in \mathcal{H}^{\otimes n}$, the probabilistic Wick product (or Wick product) of $\mathbf{I}_m(f)$ and $\mathbf{I}_n(g)$ is defined by

$$(2.14) \quad \mathbf{I}_m(f) \diamond \mathbf{I}_n(g) \stackrel{\text{def}}{=} \mathbf{I}_{m+n}(f \otimes g).$$

Unlike the *physical* Wick product, the Wick product defined by (2.14) has the same properties as the ordinary product. As an example, by (2.14) and (2.13), we can show that for $f \in \mathcal{H}^{\otimes m}$ and $g \in \mathcal{H}$,

$$\mathbf{I}_m(f) \diamond \mathbf{I}_1(g) = \mathbf{I}_m(f)\mathbf{I}_1(g) - m\mathbf{I}_{m-1}(f \otimes_1 g).$$

For two square integrable random variables $F = \sum_{m=0}^{\infty} \mathbf{I}_m(f_m)$ and $G = \sum_{n=0}^{\infty} \mathbf{I}_n(g_n)$, their Wick product is given by

$$F \diamond G = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \mathbf{I}_{m+n}(f_m \otimes g_n),$$

as long as the right-hand side is well defined. For instance, let $\varepsilon(u)$ be the *exponential vector* of u for $u \in \mathcal{H}$:

$$\varepsilon(u) \stackrel{\text{def}}{=} \exp\left(W(u) - \frac{\|u\|_{\mathcal{H}}^2}{2}\right).$$

Then $\varepsilon(u) \diamond \varepsilon(v) = \varepsilon(u+v)$ for $u, v \in \mathcal{H}$. We also recall that for a centered Gaussian random variable F , its *Wick exponential* is given by

$$(2.15) \quad \exp^{\diamond}(F) \stackrel{\text{def}}{=} \exp\left(F - \frac{1}{2}\mathbb{E}[F^2]\right) = \sum_{m=0}^{\infty} \frac{1}{m!} F^{\diamond m}.$$

As mentioned in Section A, the physical and probabilistic Wick products of a Gaussian vector coincide. In particular, for $f_i \in \mathcal{H}, i = 1, \dots, n$, we have

$$(2.16) \quad \cdot \mathbf{I}_1(f_1) \cdots \mathbf{I}_1(f_n) := \mathbf{I}_1(f_1) \diamond \cdots \diamond \mathbf{I}_1(f_n).$$

In contrast to the *multiple Wiener integral* given in (2.9), the *multiple Stratonovich integral* of $g^{\otimes m}$ for $g \in \mathcal{H}$ is defined by ([16, 18]):

$$(2.17) \quad \mathbb{I}_m(g^{\otimes m}) \stackrel{\text{def}}{=} W(g)^m.$$

By a limiting argument, one can define $\mathbb{I}_m(g)$ for g in some appropriate space. We also take the following notation for $\mathbb{I}_m(f)$ (in comparison with $\mathbf{I}_m(f)$ in (2.11)):

$$\begin{aligned} \mathbb{I}_m(f) &= \int_{([0,1] \times \mathbb{R})^m} f(\mathbf{t}, \mathbf{x}) W(dt_1, dx_1) \cdots W(dt_m, dx_m) \\ &= \int_{([0,1] \times \mathbb{R})^m} f(\mathbf{t}, \mathbf{x}) \prod_{i=1}^m W(dt_i, dx_i). \end{aligned}$$

For $f \in \mathcal{H}^{\otimes m}$, define the k th order trace $\text{Tr}^k f$ of f by

$$(2.18) \quad \text{Tr}^k f(t_1, x_1, \dots, t_{m-2k}, x_{m-2k}) \stackrel{\text{def}}{=} \int_{([0,1] \times \mathbb{R})^{2k}} f(s_1, y_1, \dots, s_{2k}, y_{2k}, t_1, x_1, \dots, t_{m-2k}, x_{m-2k}) \\ K(s_1 - s_2, y_1 - y_2) \cdots K(s_{2k-1} - s_{2k}, y_{2k-1} - y_{2k}) ds dy.$$

The following Hu-Meyer's formula (see [16], [11], [23]) connects *multiple Stratonovich integrals* $\mathbb{I}_m(f)$ with *multiple Wiener integrals* $\mathbf{I}_m(f)$,

$$(2.19) \quad \mathbb{I}_m(f) = \sum_{k=0}^{\lfloor \frac{m}{2} \rfloor} \frac{m!}{k!(m-2k)!2^k} \mathbf{I}_{m-2k}(\text{Tr}^k \hat{f}), \quad f \in \mathcal{H}^{\otimes m},$$

as long as the right-hand side is well-defined, i.e., $\text{Tr}^k \hat{f} \in \mathcal{H}^{\otimes(m-2k)}$ for $k = 0, 1, \dots, \lfloor m/2 \rfloor$. In this case, we have

$$(2.20) \quad \|f\|_{\mathfrak{S}_m}^2 \stackrel{\text{def}}{=} \mathbb{E}[|\mathbb{I}_m(f)|^2] = \sum_{k=0}^{\lfloor \frac{m}{2} \rfloor} \frac{1}{(m-2k)!} \left(\frac{m!}{k!2^k}\right)^2 \|\text{Tr}^k \hat{f}\|_{\mathcal{H}^{\otimes(m-2k)}}^2.$$

To end this section, we introduce the stochastic Fubini theorem which will be used in the proof of Proposition 2.8. Note that stochastic Fubini theorem has been proved in different contexts (see e.g. [10, 25]), and here we provide a version working for multiple Wiener and Stratonovich integrals.

Let $(\mathbf{X}, \mathcal{X}, \mu)$ be a measurable space with $\mu(\mathbf{X}) < \infty$. For a measurable function $f : \mathbf{X} \rightarrow \mathbb{R}$, we take the notation

$$\mu(f) \stackrel{\text{def}}{=} \int_{\mathbf{X}} f(x) \mu(dx).$$

The following result holds for general Gaussian spaces associated with a *separable* Hilbert space.

Proposition 2.4. *For $m \in \mathbb{N}$, let $h : \mathbf{X} \rightarrow \mathcal{H}^{\otimes m}$ be a measurable mapping such that*

$$\mu(\mathbb{E}[|\mathbf{I}_m(h)|^2]) = \mu(\|h\|_{\mathcal{H}^{\otimes m}}^2) < \infty.$$

Then the following stochastic Fubini theorem holds,

$$(2.21) \quad \mu(\mathbf{I}_m(h)) = \mathbf{I}_m(\mu(h)).$$

Similarly, if we assume

$$\mu\left(\mathbb{E}[|\mathbb{I}_m(h)|^2]\right) = \mu(\|h\|_{\mathbb{S}_m}^2) = \sum_{k=0}^{\lfloor \frac{m}{2} \rfloor} \frac{1}{(m-2k)!} \left(\frac{m!}{k!2^k}\right)^2 \mu\left(\|\text{Tr}^k \hat{h}\|_{\mathcal{H}^{\otimes(m-2k)}}^2\right) < \infty,$$

we also have

$$(2.22) \quad \mu(\mathbb{I}_m(h)) = \mathbb{I}_m(\mu(h)).$$

Proof Let $\{e_i, i \in \mathbb{N}\}$ be an orthonormal basis of \mathcal{H} . Suppose

$$h(x) = \sum_{i_1, \dots, i_m=1}^{\infty} \alpha_{i_1, \dots, i_m}(x) e_{i_1} \otimes \cdots \otimes e_{i_m}.$$

Then the condition $\mu(\|h\|_{\mathcal{H}^{\otimes m}}^2) < \infty$ implies $\sum_{i_1, \dots, i_m} \mu(\alpha_{i_1, \dots, i_m}^2) < \infty$. This further implies

$$(2.23) \quad \mu(h) = \sum_{i_1, \dots, i_m} \mu(\alpha_{i_1, \dots, i_m}) e_{i_1} \otimes \cdots \otimes e_{i_m},$$

where the series on the right-hand side converges in the Hilbert space $\mathcal{H}^{\otimes m}$ due to Jensen's inequality and the assumption $\mu(\mathbf{X}) < \infty$.

Consider $g \in \mathcal{H}^{\otimes m}$ and suppose

$$g = \sum_{i_1, \dots, i_m=1}^{\infty} \beta_{i_1, \dots, i_m} e_{i_1} \otimes \cdots \otimes e_{i_m}$$

with $\|g\|_{\mathcal{H}^{\otimes m}}^2 = \sum_{i_1, \dots, i_m} \beta_{i_1, \dots, i_m}^2 < \infty$. We have

$$(2.24) \quad \mu(\langle h, g \rangle_{\mathcal{H}^{\otimes m}}) = \mu\left(\sum_{i_1, \dots, i_m} \alpha_{i_1, \dots, i_m} \beta_{i_1, \dots, i_m}\right) = \sum_{i_1, \dots, i_m} \mu(\alpha_{i_1, \dots, i_m}) \beta_{i_1, \dots, i_m} = \langle \mu(h), g \rangle_{\mathcal{H}^{\otimes m}},$$

where the second equality follows from classical Fubini's theorem, noting that

$$\mu\left(\sum_{i_1, \dots, i_m} |\alpha_{i_1, \dots, i_m} \beta_{i_1, \dots, i_m}|\right) \leq \mu\left(\left(\sum_{i_1, \dots, i_m} \alpha_{i_1, \dots, i_m}^2\right)^{1/2}\right) \left(\sum_{i_1, \dots, i_m} \beta_{i_1, \dots, i_m}^2\right)^{1/2} < \infty,$$

and the last equality follows from (2.23).

By (2.24) we have $\mathbb{E}[\mu(\mathbf{I}_m(h))\mathbf{I}_m(g)] = \mathbb{E}[\mathbf{I}_m(\mu(h))\mathbf{I}_m(g)]$ for any $g \in \mathcal{H}^{\otimes m}$, and thus the desired equality (2.21) holds. Finally, equation (2.22) follows from (2.21) and (2.19). \square

2.2. Mild Stratonovich solution. In this subsection, we consider the fractional SHE (1.11) with initial value $u(0, x) = u_0(x)$, $0 \leq t \leq 1, x \in \mathbb{R}$. In light of Duhamel's principle, we call $u(t, x)$ a *mild Stratonovich solution* of (1.11) if $u(t, x)$ is an adapted process (with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$ where $\mathcal{F}_t = \sigma(W(s, x), 0 \leq s \leq t, x \in \mathbb{R})$) satisfying the following integral equation

$$(2.25) \quad u(t, x) = \int_{\mathbb{R}} \mathbf{g}(t, x - y) u_0(y) dy + \beta \int_0^t \int_{\mathbb{R}} \mathbf{g}(t - r, x - y) u(r, y) W(dr, dy),$$

where $\mathbf{g}(t, x)$ is the density function of the ρ -stable process $\{X_t, t \geq 0\}$ with $X_0 = 0$ and the integral on the right-hand side is understood in the sense of Stratonovich which will be specified below.

Assuming $u_0(x) \equiv 1$, we have

$$(2.26) \quad u(t, x) = 1 + \beta \int_0^t \int_{\mathbb{R}} \mathbf{g}(t - r, x - y) u(r, y) W(dr, dy).$$

Iterate the equation and we have a series expansion for a mild Stratonovich solution:

$$(2.27) \quad \begin{aligned} u(t, x) &= 1 + \sum_{m=1}^{\infty} \beta^m \int_{[0, t]_{<}^m} \int_{\mathbb{R}^m} \prod_{i=1}^m \mathbf{g}(t_{i+1} - t_i, x_{i+1} - x_i) W(dt_i, dx_i) \\ &= 1 + \sum_{m=1}^{\infty} \beta^m \int_{[0, t]_{<}^m} \int_{\mathbb{R}^m} \mathbf{g}_m(x; \mathbf{t}, \mathbf{x}) \prod_{i=1}^m W(dt_i, dx_i) \end{aligned}$$

where

$$(2.28) \quad \mathbf{g}_m(x; \mathbf{t}, \mathbf{x}) \stackrel{\text{def}}{=} \prod_{i=1}^m \mathbf{g}(t_{i+1} - t_i, x_{i+1} - x_i) \mathbf{1}_{[0, 1]_{<}^m}(t_1, \dots, t_m)$$

with

$$[0, t]_{<}^m \stackrel{\text{def}}{=} \{0 < t_1 < \dots < t_m < t\}$$

and $t_{m+1} \stackrel{\text{def}}{=} t, x_{m+1} \stackrel{\text{def}}{=} x$, and on the right-hand side are *multiple Stratonovich integrals*. We also refer to [19, 37] for an equivalent description.

Knowing that $u(t, x)$ solves a stochastic heat equation, we now aim to derive a Feynman-Kac type representation for $u(t, x)$. Given a path of X , the following Wiener integral

$$(2.29) \quad \mathcal{I}_{t,x}^\varepsilon \stackrel{\text{def}}{=} \int_0^t \int_{\mathbb{R}} p_\varepsilon(X_{t-r}^x - y) W(dr, dy),$$

is well defined, where $X_s^x \stackrel{\text{def}}{=} X_s + x$ and $p_\varepsilon(x) \stackrel{\text{def}}{=} \frac{1}{\sqrt{2\pi\varepsilon}} e^{-\frac{x^2}{2\varepsilon}}$ is the heat kernel.

As for the discrete model, we use \mathbb{E}_X and \mathbb{E}_W to denote the expectations in the probability space of X and W , respectively, and we abuse the notation $\mathbb{E} = \mathbb{E}_W \times \mathbb{E}_X$ in this section.

Proposition 2.5. *Assume the condition (1.13). Then, for each $\varepsilon > 0$, $p_\varepsilon(X_{t-}^x - \cdot)$ belongs to \mathcal{H} a.s. and the family of random variables $\mathcal{I}_{t,x}^\varepsilon$ defined by (2.29) converges in L^2 to a limit denoted by*

$$(2.30) \quad \mathcal{I}_{t,x} \stackrel{\text{def}}{=} \int_0^t \int_{\mathbb{R}} \delta(X_{t-r}^x - y) W(dr, dy),$$

where $\delta(X_{t-}^x - \cdot)$ is an \mathcal{H} -valued random variable given by the L^2 -limit of $p_\varepsilon(X_{t-}^x - \cdot)$. Moreover, $\delta(X_{t-}^x - \cdot) \in L^2(\Omega_X, \mathcal{G}, P; \mathcal{H})$, where $(\Omega_X, \mathcal{G}, P)$ is the probability space of X .

Conditional on X , $\mathcal{I}_{t,x}$ is a Gaussian random variable with mean 0 and variance

$$(2.31) \quad \text{Var}[\mathcal{I}_{t,x}|X] = \int_0^t \int_0^t |r - s|^{2H-2} \delta(X_r - X_s) dr ds.$$

Proof For $\varepsilon, \sigma > 0$, by (2.2) we have for $t \leq 1$,

$$\begin{aligned}
(2.32) \quad & \left\langle p_\varepsilon(X_{t-}^x - \cdot) \mathbf{I}_{[0,t]}, p_\sigma(X_{t-}^x - \cdot) \mathbf{I}_{[0,t]} \right\rangle_{\mathcal{H}} \\
&= \int_0^t \int_0^t \int_{\mathbb{R}} |s-r|^{2H-2} p_\varepsilon(X_s^x - y) p_\sigma(X_r^x - y) dy ds dr \\
&= \int_0^t \int_0^t |s-r|^{2H-2} p_{\varepsilon+\sigma}(X_s - X_r) ds dr.
\end{aligned}$$

By (1.7), we have

$$\mathbb{E}[p_{\varepsilon+\sigma}(X_s - X_r)] = \int_{\mathbb{R}} p_{\varepsilon+\sigma}(y) \mathfrak{g}(|r-s|, y) dy \lesssim |r-s|^{-1/\rho}.$$

Thus, the condition (1.13) yields

$$\mathbb{E} \left\langle p_\varepsilon(X_{t-}^x - \cdot), p_\sigma(X_{t-}^x - \cdot) \right\rangle_{\mathcal{H}} \lesssim \int_0^t \int_0^t |s-r|^{2H-2-\frac{1}{\rho}} ds dr < \infty,$$

hence $p_\varepsilon(X_{t-}^x - \cdot)$ belongs to \mathcal{H} for all $\varepsilon > 0$ almost surely and

$$(2.33) \quad \mathbb{E}[\mathcal{I}_{t,x}^\varepsilon \mathcal{I}_{t,x}^\sigma] = \mathbb{E} \left\langle p_\varepsilon(X_{t-}^x - \cdot), p_\sigma(X_{t-}^x - \cdot) \right\rangle_{\mathcal{H}} = \mathbb{E} \int_0^t \int_0^t |s-r|^{2H-2} p_{\varepsilon+\sigma}(X_s - X_r) ds dr.$$

As $(\varepsilon, \sigma) \rightarrow 0$, by the dominated convergence theorem we have

$$\lim_{(\varepsilon, \sigma) \rightarrow 0} \mathbb{E}[\mathcal{I}_{t,x}^\varepsilon \mathcal{I}_{t,x}^\sigma] = \mathbb{E} \int_0^t \int_0^t |s-r|^{2H-2} \delta(X_s - X_r) ds dr = C \int_0^t \int_0^t |s-r|^{2H-2-\frac{1}{\rho}} ds dr < \infty$$

for some proper constant C . As a consequence, we have $\lim_{(\varepsilon, \sigma) \rightarrow 0} \mathbb{E} \left[\left(\mathcal{I}_{t,x}^\varepsilon - \mathcal{I}_{t,x}^\sigma \right)^2 \right] = 0$.

This implies that, as ε converges to zero, $p_\varepsilon(X_{t-}^x - \cdot)$ is a Cauchy sequence in $L^2(\Omega_X, \mathcal{G}, P; \mathcal{H})$, and the limit is denoted by $\delta(X_{t-}^x - \cdot)$. Therefore, $\mathcal{I}_{t,x}^\varepsilon$ converges to $\mathcal{I}_{t,x}$ given in (2.30) in L^2 . Finally, (2.31) can be proven by a similar argument. \square

Proposition 2.6. *Under the condition (1.13), we have*

$$(2.34) \quad \mathbb{E} \left[\exp \left(\beta \int_0^1 \int_0^1 |r-s|^{2H-2} \delta(X_r - X_s) dr ds \right) \right] < \infty \text{ for all } \beta > 0.$$

Proof The proof essentially follows from that of [37, Theorem 3.3] with some minor modifications. Taylor expansion and Fubini's theorem yield

$$\begin{aligned}
(2.35) \quad & \mathbb{E} \left[\exp \left(\beta \int_0^1 \int_0^1 |r-s|^{2H-2} \delta(X_r - X_s) dr ds \right) \right] \\
&= \sum_{m=0}^{\infty} \frac{1}{m!} \beta^m \int_{[0,1]^{2m}} \int_{\mathbb{R}^m} \prod_{i=1}^m |s_{2i} - s_{2i-1}|^{2H-2} \mathbb{E} \left[e^{i \sum_{i=1}^m \xi_i(X_{s_{2i}} - X_{s_{2i-1}})} \right] d\xi ds \\
&= \sum_{m=0}^{\infty} \frac{1}{m!} \beta^m \sum_{\sigma \in \mathcal{P}_{2m}} \int_{[0,1]^{2m}} \int_{\mathbb{R}^m} \prod_{i=1}^m |s_{\sigma(2i)} - s_{\sigma(2i-1)}|^{2H-2} \mathbb{E} \left[e^{i \sum_{i=1}^m \xi_i(X_{s_{\sigma(2i)}} - X_{s_{\sigma(2i-1)}})} \right] d\xi ds,
\end{aligned}$$

where we recall that $[0, 1]_{<}^{2m} = \{0 < s_1 < s_2 < \dots < s_{2m} < 1\}$ and \mathcal{P}_{2m} is the set of all permutations on $[2m]$.

Let $\sigma \in \mathcal{P}_{2m}$ be arbitrarily chosen and fixed. Consider $(s_1, \dots, s_{2m}) \in [0, 1]_{<}^{2m}$, and for each pair $(s_{\sigma(2i-1)}, s_{\sigma(2i)})$, we let $t_{2i}^* \stackrel{\text{def}}{=} s_{\sigma(2i-1)} \vee s_{\sigma(2i)}$ and t_{2i-1}^* be the unique s_j which is the closest point

to t_{2i}^* from the left. Then clearly, we have

$$\prod_{i=1}^m |s_{\sigma(2i)} - s_{\sigma(2i-1)}|^{2H-2} \leq \prod_{i=1}^m |t_{2i}^* - t_{2i-1}^*|^{2H-2}.$$

Noting that

$$\sum_{i=1}^m \xi_i(X_{s_{\sigma(2i)}} - X_{s_{\sigma(2i-1)}}) = \sum_{j=2}^{2m} \eta_j(X_{s_j} - X_{s_{j-1}})$$

where each η_j is a linear combination of ξ_i 's for $i = 1, \dots, m$. Then we have

$$\mathbb{E} \left[e^{i \sum_{i=1}^m \xi_i(X_{s_{\sigma(2i)}} - X_{s_{\sigma(2i-1)}})} \right] = \prod_{j=2}^{2m} e^{-c_\rho |\eta_j|^\rho (s_j - s_{j-1})} \leq \prod_{i=1}^m e^{-c_\rho |\tilde{\eta}_i|^\rho (t_{2i}^* - t_{2i-1}^*)},$$

where the inequality holds because we only keep the factors resulting from the characteristic function of $X_{t_{2i}^*} - X_{t_{2i-1}^*}$ for $i \in \llbracket m \rrbracket$ and drop all the others. Here, for each $i \in \llbracket m \rrbracket$, we define $\tilde{\eta}_i = \eta_j$ where j is the unique index such that $s_j = t_{2i}^* = s_{\sigma(2i)} \vee s_{\sigma(2i-1)}$.

Thus, we have

$$\begin{aligned} & \int_{[0,1]_{<}^{2m}} \int_{\mathbb{R}^m} \prod_{i=1}^m |s_{\sigma(2i)} - s_{\sigma(2i-1)}|^{2H-2} \mathbb{E} \left[e^{i \sum_{i=1}^m \xi_i(X_{s_{\sigma(2i)}} - X_{s_{\sigma(2i-1)}})} \right] d\xi ds \\ (2.36) \quad & \leq \int_{[0,1]_{<}^{2m}} \int_{\mathbb{R}^m} \prod_{i=1}^m |t_{2i}^* - t_{2i-1}^*|^{2H-2} \prod_{i=1}^m e^{-c_\rho |\tilde{\eta}_i|^\rho (t_{2i}^* - t_{2i-1}^*)} d\xi ds \\ & = C^m \int_{[0,1]_{<}^{2m}} \prod_{i=1}^m |t_{2i}^* - t_{2i-1}^*|^{2H-2-\frac{1}{\rho}} ds \\ & \leq \frac{C^m}{\Gamma\left(m(2H - \frac{1}{\rho}) + 1\right)}. \end{aligned}$$

where the equality follows from the fact that the Jacobian determinant $|\partial \xi_i / \partial \tilde{\eta}_j|_{m \times m}$ is one and that $\int_{\mathbb{R}} e^{-a|\xi|^\rho} d\xi = Ca^{-1/\rho}$ for $a > 0$, and the last step follows from Lemma C.2.

Therefore, by (2.35) and (2.36), and noting that $|\mathcal{P}_{2m}| = (2m)!$, we have

$$\mathbb{E} \left[\exp \left(\beta \int_0^1 \int_0^1 |r-s|^{2H-2} \delta(X_r - X_s) dr ds \right) \right] \leq \sum_{m=0}^{\infty} C^m \beta^m \frac{(2m)!}{m! \Gamma\left(m(2H - \frac{1}{\rho}) + 1\right)} < \infty,$$

where the last step follows from Sterling's formula and $2H - \frac{1}{\rho} > 1$. \square

Remark 2.7. In the proof of [37, Theorem 3.3], the Markov property instead of the independent increment property of X was invoked, and as a consequence, [37, Lemma 2.2] was needed therein. As shown in the proof of Proposition 2.6, if we utilise the independent increment property of X , [37, Lemma 2.2] is no longer needed, and this is important for the proof of uniform L^1 -bound for rescaled partition functions in Section 3.3.

Proposition 2.8. The mild Stratonovich solution given in (2.27) has the following Feynman-Kac representation:

$$(2.37) \quad u(t, x) = \mathbb{E}_X \left[\exp \left(\beta \int_0^t \int_{\mathbb{R}} \delta(X_{t-r}^x - y) W(dr, dy) \right) \right].$$

Proof First, we prove the integrability of the right-hand side of (2.37). For all $\beta \in \mathbb{R}$,

$$\begin{aligned}
(2.38) \quad & \mathbb{E}_W \left[\mathbb{E}_X \left[\exp \left(\beta \int_0^t \int_{\mathbb{R}} \delta(X_{t-r}^x - y) W(dr, dy) \right) \right] \right] \\
&= \mathbb{E}_X \left[\mathbb{E}_W \left[\exp \left(\beta \int_0^t \int_{\mathbb{R}} \delta(X_{t-r}^x - y) W(dr, dy) \right) \right] \right] \\
&= \mathbb{E}_X \left[\exp \left(\frac{1}{2} \beta^2 t^{2H-\frac{1}{\rho}} \int_0^1 \int_0^1 |r-s|^{2H-2} \delta(X_r - X_s) dr ds \right) \right] \\
&< \infty,
\end{aligned}$$

where the second equality follows from (2.31) and the self-similarity of X , and the last step follows from Proposition 2.6. Thus $u(t, x)$ given by (2.37) is well-defined and is L^p -integrable for all $p > 0$.

The Feynman-Kac formula (2.37) now follows from (2.27) and the following equation:

$$\begin{aligned}
(2.39) \quad & \mathbb{E}_X \left[\left(\int_0^t \int_{\mathbb{R}} \delta(X_{t-r}^x - y) W(dr, dy) \right)^m \right] \\
&= \int_{[0,t]^m} \int_{\mathbb{R}^m} \mathbb{E}_X \left[\prod_{i=1}^m \delta(X_{t-r_i}^x - y_i) \right] W(dr_1, dy_1) \dots W(dr_m, dy_m) \\
&= m! \int_{[0,t]^m} \int_{\mathbb{R}^m} \mathfrak{g}_m(x; \mathbf{r}, \mathbf{y}) \prod_{i=1}^m W(dr_i, dy_i),
\end{aligned}$$

where the first equality follows from (2.17) and the stochastic Fubini theorem (Proposition 2.4). \square

Remark 2.9. A direct corollary of Propositions of 2.6 and 2.8 is

$$(2.40) \quad \|\mathfrak{g}_m\|_{\mathbf{S}_m} < \infty,$$

where \mathfrak{g}_m is given in (2.28) and $\|\cdot\|_{\mathbf{S}_m}$ in (2.20). Indeed,

$$\|\mathfrak{g}_m\|_{\mathbf{S}_m}^2 = \mathbb{E}[|\mathbb{I}_m(\mathfrak{g}_m)|^2] = \left(\frac{1}{m!}\right)^2 \mathbb{E}_W \left[\left(\mathbb{E}_X[\mathcal{I}_{t,x}^m] \right)^2 \right] \leq \left(\frac{1}{m!}\right)^2 \mathbb{E}[\mathcal{I}_{t,x}^{2m}] \leq \left(\frac{1}{m!}\right)^2 \mathbb{E}[e^{\mathcal{I}_{t,x}}] < \infty,$$

where $\mathcal{I}_{t,x}$ is given in (2.30), the second equality follows from (2.39), and the last inequality follows from (2.38).

To close this subsection, we prove the L^1 -convergence of the series representation for $u(1, x)$, which will be used in the proof of Theorem 1.2.

Proposition 2.10. The mild Stratonovich solution of (2.26) given in (2.27) converges in L^1 .

Proof By (2.39) we have

$$\begin{aligned}
& \sum_{m=0}^{\infty} \beta^m \mathbb{E} \left[\left| \int_{[0,t]^m} \int_{\mathbb{R}^m} \mathfrak{g}_m(x; \mathbf{t}, \mathbf{y}) \prod_{i=1}^m W(dt_i, dy_i) \right| \right] \\
&\leq \sum_{m=0}^{\infty} \frac{\beta^m}{m!} \mathbb{E} \left[\left| \int_0^t \int_{\mathbb{R}} \delta(X_{t-r}^x - y) W(dr, dy) \right|^m \right] \\
&= \mathbb{E} \left[\exp \left(\beta \left| \int_0^t \int_{\mathbb{R}} \delta(X_{t-r}^x - y) W(dr, dy) \right| \right) \right] \\
&\leq \mathbb{E} \left[\exp \left(\beta \int_0^t \int_{\mathbb{R}} \delta(X_{t-r}^x - y) W(dr, dy) \right) \right] + \mathbb{E} \left[\exp \left(-\beta \int_0^t \int_{\mathbb{R}} \delta(X_{t-r}^x - y) W(dr, dy) \right) \right],
\end{aligned}$$

and the right-hand side is finite due to (2.38). \square

3. ON WEAK CONVERGENCES

In this section, we aim to prove the main result Theorem 1.2.. Recall that $\theta = H - \frac{1}{2\rho} \in (\frac{1}{2}, 1)$. For $N \in \mathbb{N}$, under the scaling $\beta \rightarrow \hat{\beta}_N = \beta N^{-\theta}$, the rescaled point-to-line partition function is given by

$$(3.1) \quad Z_\omega^{(N)}(\hat{\beta}_N, N^{1/\rho}x_0) = \mathbb{E}_S \left[\exp \left(\hat{\beta}_N \sum_{n=1}^N \omega(n, S_n) \right) \middle| S_0 = N^{1/\rho}x_0 \right],$$

where $x_0 \in \mathbb{R}$ is fixed such that $N^{1/\rho}x_0 \in \mathbb{Z}$. Taylor expansion yields

$$(3.2) \quad Z_\omega^{(N)}(\hat{\beta}_N, N^{1/\rho}x_0) = \sum_{m=0}^{\infty} \frac{1}{m!} \mathbb{S}_m^{(N)},$$

where $\mathbb{S}_m^{(N)}$ is the m th moment of the Hamiltonian (with a factor $\hat{\beta}_N$) given by

$$(3.3) \quad \begin{aligned} \mathbb{S}_m^{(N)} &= \mathbb{E}_S \left[\left(\hat{\beta}_N \sum_{n=1}^N \omega(n, S_n) \right)^m \middle| S_0 = N^{1/\rho}x_0 \right] \\ &= \hat{\beta}_N^m \mathbb{E}_S \left[\sum_{\mathbf{n} \in \llbracket N \rrbracket^m} \omega(n_1, S_{n_1}) \cdots \omega(n_m, S_{n_m}) \middle| S_0 = N^{1/\rho}x_0 \right]. \end{aligned}$$

For every $\mathbf{n} = (n_1, \dots, n_m) \in \llbracket N \rrbracket^m$, the components can be arranged in an increasing order and the resulted sequence is denoted by $\mathbf{n}^* = (n_1^*, \dots, n_m^*)$ with $n_1^* \leq n_2^* \leq \dots \leq n_m^*$. For each $\mathbf{n} \in \llbracket N \rrbracket^m$, there is a permutation σ of $\llbracket N \rrbracket$ such that $n_i^* = n_{\sigma_i}$ for $i = 1, \dots, m$. Denote

$$(3.4) \quad \begin{aligned} P_{\mathbf{n}^*} &= P_{\mathbf{n}^*}(N^{1/\rho}x_0; k_1, \dots, k_m) \\ &= P_{n_1^*}(k_{\sigma_1} - N^{1/\rho}x_0) P_{n_2^* - n_1^*}(k_{\sigma_2} - k_{\sigma_1}) \cdots P_{n_m^* - n_{m-1}^*}(k_{\sigma_m} - k_{\sigma_{m-1}}), \end{aligned}$$

where we use the convention $P_0(0) = 1$ and $P_0(k) = 0$ for $k \neq 0$. We remark that $P_{\mathbf{n}^*}$ is *symmetric* in its n arguments. The m th moment in (3.3) can be written as

$$(3.5) \quad \mathbb{S}_m^{(N)} = \hat{\beta}_N^m \sum_{n_1, \dots, n_m \in \llbracket N \rrbracket} \sum_{k_1, \dots, k_m \in \mathbb{Z}} \omega(n_1, k_1) \cdots \omega(n_m, k_m) P_{\mathbf{n}^*}(N^{1/\rho}x_0; k_1, \dots, k_m).$$

We introduce the following general U-statistics $\mathbf{I}_m^{(N)}(f)$ which will be proven to converge weakly to a multiple Wiener integral. For $f \in \mathcal{B}^{\otimes m}$ where we recall that \mathcal{B} is given in (2.3), we denote

$$(3.6) \quad \begin{aligned} \mathbf{I}_m^{(N)}(f) &\stackrel{\text{def}}{=} N^{-m(\theta+1/\rho)} \sum_{n_1, \dots, n_m \in \llbracket N \rrbracket} \sum_{k_1, \dots, k_m \in \mathbb{Z}} \left[: \omega(n_1, k_1) \cdots \omega(n_m, k_m) : \right. \\ &\quad \left. \times \mathcal{A}_N(f) \left(\frac{n_1}{N}, \frac{k_1}{N^{1/\rho}}, \dots, \frac{n_m}{N}, \frac{k_m}{N^{1/\rho}} \right) \right] \\ &\stackrel{\text{def}}{=} N^{-m(\theta+1/\rho)} \sum_{\mathbf{n} \in \llbracket N \rrbracket^m} \sum_{\mathbf{k} \in \mathbb{Z}^m} : \omega_{\llbracket m \rrbracket} : \mathcal{A}_N(f)(\mathbf{t}, \mathbf{x}), \end{aligned}$$

where $: \omega_{\llbracket m \rrbracket} : \stackrel{\text{def}}{=} : \omega_1 \cdots \omega_m :$ is the physical Wick product of $\omega_i \stackrel{\text{def}}{=} \omega(n_i, k_i)$, $\mathcal{A}_N(f)$ is given in (2.5), and

$$(3.7) \quad \begin{aligned} \mathbf{t} &= (t_1, \dots, t_m) \stackrel{\text{def}}{=} \mathbf{n}/N = (n_1/N, \dots, n_m/N), \\ \mathbf{x} &= (x_1, \dots, x_m) \stackrel{\text{def}}{=} \mathbf{x}/N^{1/\rho} = (k_1/N^{1/\rho}, \dots, k_m/N^{1/\rho}). \end{aligned}$$

3.1. Weak convergence of U-statistics. In this subsection, we shall prove the joint weak convergence for U -statistics $\mathbf{I}_m^{(N)}(f)$ defined by (3.6).

We first introduce the following lemma which provides a uniform (in N) bound for the second moment of $\mathbf{I}_m^{(N)}(f)$ with $f \in \mathcal{B}^{\otimes m}$, which will be used to prove Propositions 3.2 and 3.3.

Lemma 3.1. *Assume the disorder ω is given as in Section 1.2. Then, there exists a constant C such that for all $f \in \mathcal{B}^{\otimes m}$,*

$$(3.8) \quad \mathbb{E} \left[\left(\mathbf{I}_m^{(N)}(f) \right)^2 \right] \leq C^m m! \|\hat{f}\|_{\mathcal{B}^{\otimes m}}^2,$$

where \hat{f} is the symmetrization of f .

Proof For $m = 1$, we have

$$(3.9) \quad \begin{aligned} & \mathbb{E} \left[\left(\mathbf{I}_1^{(N)}(f) \right)^2 \right] \\ &= N^{-2(\theta+1/\rho)} \sum_{n, n' \in \llbracket N \rrbracket} \sum_{k, k' \in \mathbb{Z}} \mathbb{E}[\omega(n, k)\omega(n', k')] \mathcal{A}_N(f) \left(\frac{n}{N}, \frac{k}{N^{1/\rho}} \right) \mathcal{A}_N(f) \left(\frac{n'}{N}, \frac{k'}{N^{1/\rho}} \right) \\ &= N^{-2(\theta+1/\rho)} \sum_{n, n' \in \llbracket N \rrbracket} \sum_{k \in \mathbb{Z}} \gamma(n - n') \mathcal{A}_N(f) \left(\frac{n}{N}, \frac{k}{N^{1/\rho}} \right) \mathcal{A}_N(f) \left(\frac{n'}{N}, \frac{k}{N^{1/\rho}} \right). \end{aligned}$$

For $t \in \mathbb{R}$, let

$$(3.10) \quad \gamma_N(t) \stackrel{\text{def}}{=} N^{2-2H} \gamma(\lfloor tN \rfloor).$$

Then, by (1.9) and (1.10), we have for all $t \in \mathbb{R}$

$$(3.11) \quad \gamma_N(t) \lesssim |t|^{2H-2} \text{ and } \lim_{N \rightarrow \infty} \gamma_N(t) = |t|^{2H-2}.$$

Then, using the notations $t = n/N, t' = n'/N, x = k/N^{1/\rho}, x' = k'/N^{1/\rho}$, we have

$$(3.12) \quad \begin{aligned} \mathbb{E} \left[\left(\mathbf{I}_1^{(N)}(f) \right)^2 \right] &= N^{-2-1/\rho} \sum_{n, n' \in \llbracket N \rrbracket} \sum_{k \in \mathbb{Z}} N^{2-2H} \gamma(n - n') \mathcal{A}_N(f)(t, x) \mathcal{A}_N(f)(t', x) \\ &\lesssim \int_0^1 \int_0^1 \int_{\mathbb{R}} |t - t'|^{2H-2} |\mathcal{A}_N(f)(t, x) \mathcal{A}_N(f)(t', x)| dx dt dt' \\ &= \|\mathcal{A}_N(|f|)\|_{\mathcal{B}}^2 \lesssim \|f\|_{\mathcal{B}}^2, \end{aligned}$$

where the last inequality follows from Lemma 2.2.

For general m ,

$$(3.13) \quad \mathbb{E} \left[\left(\mathbf{I}_m^{(N)}(f) \right)^2 \right] = N^{-2m(\theta+1/\rho)} \sum_{\mathbf{n}, \mathbf{n}'} \sum_{\mathbf{k}, \mathbf{k}'} \mathbb{E} \left[: \omega_{\llbracket m \rrbracket} :: \omega'_{\llbracket m \rrbracket} : \right] \mathcal{A}_N(f)(\mathbf{t}, \mathbf{x}) \mathcal{A}_N(f)(\mathbf{t}', \mathbf{x}'),$$

where $: \omega_{\llbracket m \rrbracket} := \prod_{i=1}^m \omega(n_i, k_i) :$ and similarly for $: \omega'_{\llbracket m \rrbracket} :=$

Recalling that we have assumed that ω is Gaussian, then by (A.8) we have

$$\mathbb{E} \left[: \omega_{\llbracket m \rrbracket} :: \omega'_{\llbracket m \rrbracket} : \right] = \mathbb{E} \left[: \prod_{i=1}^m \omega(n_i, k_i) :: \prod_{i=1}^m \omega(n'_i, k'_i) : \right] = \sum_{\sigma \in \mathcal{P}_m} \prod_{i=1}^m \gamma(n_i - n'_{\sigma(i)}) \delta_{k_i k'_{\sigma(i)}},$$

where the summation $\sum_{\sigma \in \mathcal{P}_m}$ is taken over the set \mathcal{P}_m of all the permutations of $[[m]]$. Note that by the symmetry of the summation $\sum_{\mathbf{n}, \mathbf{n}'} \sum_{\mathbf{k}, \mathbf{k}'}$, we have $\mathbf{I}_m^{(N)}(f) = \mathbf{I}_m^{(N)}(\hat{f})$. Thus, by the symmetry of $\mathcal{A}_N(\hat{f})$, we have

$$\mathbb{E} \left[\left(\mathbf{I}_m^{(N)}(\hat{f}) \right)^2 \right] = m! N^{-2m(\theta+1/\rho)} \sum_{\mathbf{n}, \mathbf{n}'} \sum_{\mathbf{k}} \left(\prod_{i=1}^m \gamma(n_i - n'_i) \right) \mathcal{A}_N(\hat{f})(\mathbf{t}, \mathbf{x}) \mathcal{A}_N(\hat{f})(\mathbf{t}', \mathbf{x}).$$

Then, similarly to the case $m = 1$, we can get $\mathbb{E} \left[\left(\mathbf{I}_m^{(N)}(f) \right)^2 \right] \lesssim \|\hat{f}\|_{\mathcal{B}^{\otimes m}}^2$. \square

Now, we are ready to prove the weak convergence for $\mathbf{I}_m^{(N)}(f)$ defined in (3.6). We first prove the weak convergence for $m = 1$.

Proposition 3.2. *Let $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ be a function in \mathcal{B} . Then, $\mathbf{I}_1^{(N)}(f)$ converges weakly to the Wiener integral $\mathbf{I}_1(f)$ as $N \rightarrow \infty$. Moreover, for any $k \in \mathbb{N}$ and $f_1, \dots, f_k \in \mathcal{B}$, we have the joint convergence in distribution:*

$$(3.14) \quad \left(\mathbf{I}_1^{(N)}(f_1), \dots, \mathbf{I}_1^{(N)}(f_k) \right) \xrightarrow{d} \left(\mathbf{I}_1(f_1), \dots, \mathbf{I}_1(f_k) \right), \text{ as } N \rightarrow \infty.$$

Proof First we consider an indicator function $f = \mathbf{1}_{[s,t] \times [y,z]}$ with $0 \leq s \leq t \leq 1$ and $y \leq z$. In this case,

$$\mathbf{I}_1^{(N)}(f) = N^{-\theta-1/\rho} \sum_{n \in [Ns, Nt] \cap \mathbb{N}} \sum_{k \in [N^{1/\rho}y, N^{1/\rho}z] \cap \mathbb{Z}} \omega(n, k).$$

Clearly it has mean zero and the covariance is

$$\mathbb{E} \left[\left(\mathbf{I}_1^{(N)}(f) \right)^2 \right] = N^{-2-1/\rho} \sum_{n, n' \in [Ns, Nt] \cap \mathbb{N}} \sum_{k \in [N^{1/\rho}y, N^{1/\rho}z] \cap \mathbb{Z}} \gamma_N(r - r'),$$

where γ_N is given by (3.10) and $r = n/N, x = k/N^{1/\rho}$. By (3.11) and the dominated convergence theorem, we get

$$(3.15) \quad \lim_{N \rightarrow \infty} \mathbb{E} \left[\left(\mathbf{I}_1^{(N)}(f) \right)^2 \right] = \int_s^t \int_s^t \int_y^z |r - r'|^{2H-2} dx dr dr' = \|f\|_{\mathcal{H}}^2$$

where we recall that the inner product $\langle \cdot, \cdot \rangle_{\mathcal{H}}$ is given in (2.2). Thus, we have $\mathbf{I}_1^{(N)}(f) \xrightarrow{d} \mathbf{I}_1(f)$ as $N \rightarrow \infty$, noting that $\mathbf{I}_1^{(N)}(f), N \in \mathbb{N}$ and $\mathbf{I}_1(f)$ are Gaussian random variables and that $\mathbf{I}_1(f)$ has zero mean and variance $\|f\|_{\mathcal{H}}^2$.

Similarly, one can show that (3.15) holds for each simple function f , i.e., f is a linear combination of indicator functions. For a general function $f \in \mathcal{B}$, there exists an approximation sequence of simple functions $\{f^{(n)}\}_{n \in \mathbb{N}}$ in \mathcal{B} . On the one hand, $\mathbf{I}_1^{(N)}(f^{(n)})$ converges weakly to $\mathbf{I}_1(f^{(n)})$ as $N \rightarrow \infty$ for each fixed n . On the other hand, by Lemma 3.1, we have $\mathbf{I}_1^{(N)}(f^{(n)}) \rightarrow \mathbf{I}_1^{(N)}(f)$ in L^2 uniformly in N as $n \rightarrow \infty$. Combining the above convergences together with the obvious L^2 -convergence of $\mathbf{I}_1(f^{(n)}) \rightarrow \mathbf{I}_1(f)$ as $n \rightarrow \infty$, by Lemma B.1 we obtain the weak convergence $\mathbf{I}_1^{(N)}(f) \xrightarrow{d} \mathbf{I}_1(f)$ as $N \rightarrow \infty$ for $f \in \mathcal{B}$. The argument can be presented in a diagram:

$$\begin{array}{ccc} \mathbf{I}_1^{(N)}(f^{(n)}) & \xrightarrow[\substack{\text{in } L^2, \text{ uniformly in } N \\ n \rightarrow \infty}]{} & \mathbf{I}_1^{(N)}(f) \\ \downarrow \substack{d \\ N \rightarrow \infty} & & \downarrow \substack{d \\ N \rightarrow \infty} \\ \mathbf{I}_1(f^{(n)}) & \xrightarrow[\substack{\text{in } L^2 \\ n \rightarrow \infty}]{} & \mathbf{I}_1(f) \end{array}$$

Finally, the joint weak convergence (3.14) follows from the linearity of $\mathbf{I}_1^{(N)}$ and the Cramér-Wold theorem (see Theorem B.4). \square

The main result in this section is presented below.

Proposition 3.3. *For each $m \in \mathbb{N}$ and $f \in \mathcal{B}^{\otimes m}$, we have*

$$(3.16) \quad \mathbf{I}_m^{(N)}(f) \xrightarrow{d} \mathbf{I}_m(f), \text{ as } N \rightarrow \infty,$$

where $\mathbf{I}_m^{(N)}(f)$ is given in (3.6) and $\mathbf{I}_m(f)$ is an m th multiple Wiener integral. Moreover, for any $k \in \mathbb{N}$ and f_1, \dots, f_k with $f_i \in \mathcal{B}^{\otimes l_i}$ with $l_i \in \mathbb{N}$, we have the joint convergence in distribution:

$$(3.17) \quad \left(\mathbf{I}_{l_1}^{(N)}(f_1), \dots, \mathbf{I}_{l_k}^{(N)}(f_k) \right) \xrightarrow{d} \left(\mathbf{I}_{l_1}(f_1), \dots, \mathbf{I}_{l_k}(f_k) \right), \text{ as } N \rightarrow \infty.$$

Proof We only prove (3.16) and (3.17) can be proved in a similar way in the light of the Cramér-Wold theorem (Theorem B.4). We only need to consider $m \geq 2$. First, we consider $\mathbf{I}_m^{(N)}(f)$ for $f = h_1^{\otimes p_1} \otimes \dots \otimes h_l^{\otimes p_l}$ with $h_i \in \mathcal{B}, k_i \in \mathbb{N}$ for $i = 1, 2, \dots, l$ satisfying $p_1 + \dots + p_l = m$. For such f , we have

$$\begin{aligned} \mathbf{I}_m^{(N)}(f) &= N^{-m(\theta+1/\rho)} \sum_{n_1, \dots, n_m \in [N]} \sum_{k_1, \dots, k_m \in \mathbb{Z}} : \omega_{[m]} : h_1^{\otimes p_1} \otimes \dots \otimes h_l^{\otimes p_l} \\ &= : \left(\mathbf{I}_1^{(N)}(h_1) \right)^{p_1} \dots \left(\mathbf{I}_1^{(N)}(h_l) \right)^{p_l} :, \end{aligned}$$

where the term on the right-hand side is a physical Wick product (see Section A). By Proposition 3.2, we have $(\mathbf{I}_1^{(N)}(h_1), \dots, \mathbf{I}_1^{(N)}(h_l)) \xrightarrow{d} (\mathbf{I}_1(h_1), \dots, \mathbf{I}_1(h_l))$ as $N \rightarrow \infty$.

For each $i = 1, \dots, l$, $\mathbf{I}_1^{(N)}(h_i)$ is a centered Gaussian random variable with variance bounded by $\|h_i\|_{\mathcal{B}}^2$ with some constant multiplicative factor by Lemma 3.1. Therefore, for any $q \in \mathbb{N}$, by the Gaussianity of ω and (3.12), we have $\mathbb{E} \left[\left(\mathbf{I}_1^{(N)}(h_i) \right)^{2q} \right] \lesssim (2q-1)!! \|h_i\|_{\mathcal{B}}^{2q} < \infty$. Then we can apply Lemma B.3 and get that, as $N \rightarrow \infty$,

$$\begin{aligned} \mathbf{I}_m^{(N)}(f) &\xrightarrow{d} \left(\mathbf{I}_1(h_1) \right)^{\diamond p_1} \diamond \dots \diamond \left(\mathbf{I}_1(h_l) \right)^{\diamond p_l} = \mathbf{I}_{p_1}(h_1^{\otimes p_1}) \diamond \dots \diamond \mathbf{I}_{p_l}(h_l^{\otimes p_l}) \\ &= \mathbf{I}_m(h_1^{\otimes p_1} \otimes \dots \otimes h_l^{\otimes p_l}) = \mathbf{I}_m(f). \end{aligned}$$

Then, for any $f \in \mathcal{B}^{\otimes m}$, one can find a sequence $\{f^{(n)}\}_{n \in \mathbb{N}}$ which are the linear combinations of the functions of the form $h_1^{\otimes p_1} \otimes \dots \otimes h_l^{\otimes p_l}$ such that $f^{(n)} \rightarrow f$ in $\mathcal{B}^{\otimes m}$. By the preceding argument, for each $f^{(n)}$, we have $\mathbf{I}_m^{(N)}(f^{(n)}) \xrightarrow{d} \mathbf{I}_m(f^{(n)})$ as $N \rightarrow \infty$. Then similarly as in the proof of Proposition 3.2, we have, recalling Lemma 3.1,

$$\begin{array}{ccc} \mathbf{I}_m^{(N)}(f^{(n)}) & \xrightarrow[n \rightarrow \infty]{\text{in } L^2, \text{ uniformly in } N} & \mathbf{I}_m^{(N)}(f) \\ \downarrow d \Big|_{N \rightarrow \infty} & & \\ \mathbf{I}_m(f^{(n)}) & \xrightarrow[n \rightarrow \infty]{\text{in } L^2} & \mathbf{I}_m(f) \end{array}$$

This together with Lemma B.1 yields $\mathbf{I}_m^{(N)}(f) \xrightarrow{d} \mathbf{I}_m(f)$ as $N \rightarrow \infty$ for $f \in \mathcal{B}^{\otimes m}$. \square

3.2. Weak convergence of rescaled partition functions. Recall that the so-called m th moment $\mathbb{S}_m^{(N)}$ appearing in the rescaled partition function (3.2) has the expression (3.5). We first prove the joint weak convergence of moments.

Proposition 3.4. *Assume condition 1.13. For each $m \in \mathbb{N}$, we have*

$$(3.18) \quad \frac{1}{m!} \mathbb{S}_m^{(N)} \xrightarrow{d} \beta^m \mathbb{I}_m(\mathfrak{g}_m), \text{ as } N \rightarrow \infty,$$

where $\mathfrak{g}_m = \mathfrak{g}_m(x_0, \mathbf{t}, \mathbf{x})$ is given in (2.28) and $\mathbb{I}_m(\mathfrak{g}_m)$ is an m th multiple Stratonovich integral. Moreover, for any $k \in \mathbb{N}$ and $l_1, \dots, l_k \in \mathbb{N}$, we have the joint convergence in distribution:

$$(3.19) \quad \left(\frac{1}{l_1!} \mathbb{S}_{l_1}^{(N)}, \dots, \frac{1}{l_k!} \mathbb{S}_{l_k}^{(N)} \right) \xrightarrow{d} \left(\beta^{l_1} \mathbb{I}_{l_1}(\mathfrak{g}_{l_1}), \dots, \beta^{l_k} \mathbb{I}_{l_k}(\mathfrak{g}_{l_k}) \right), \text{ as } N \rightarrow \infty.$$

Proof We first prove (3.18). Note that $\text{Tr}^k \hat{\mathfrak{g}}_m \in \mathcal{B}^{\otimes(m-2k)}$ for $k = 0, 1, \dots, [\frac{m}{2}]$ (see Remark 2.9). Thus, by Proposition 3.3, we have (without loss of generality, assuming m is an odd integer),

$$(3.20) \quad \begin{aligned} & \left(\mathbf{I}_1^{(N)} \left(\text{Tr}^{[\frac{m}{2}]} \hat{\mathfrak{g}}_m \right), \mathbf{I}_3^{(N)} \left(\text{Tr}^{[\frac{m}{2}]-1} \hat{\mathfrak{g}}_m \right), \dots, \mathbf{I}_m^{(N)} \left(\hat{\mathfrak{g}}_m \right) \right) \\ & \xrightarrow{d} \left(\mathbf{I}_1 \left(\text{Tr}^{[\frac{m}{2}]} \hat{\mathfrak{g}}_m \right), \mathbf{I}_3 \left(\text{Tr}^{[\frac{m}{2}]-1} \hat{\mathfrak{g}}_m \right), \dots, \mathbf{I}_m \left(\hat{\mathfrak{g}}_m \right) \right), \text{ as } N \rightarrow \infty. \end{aligned}$$

Recalling (3.5), by the symmetry of the summation, we can write $\mathbb{S}_m^{(N)}$ as

$$\mathbb{S}_m^{(N)} = \hat{\beta}_N^m \sum_{n_1, \dots, n_m \in \llbracket N \rrbracket} \sum_{k_1, \dots, k_m \in \mathbb{Z}} \omega_{\llbracket m \rrbracket} P_{\mathbf{n}^*},$$

where $P_{\mathbf{n}^*}$ is given in (3.4) and $\omega_{\llbracket m \rrbracket} = \prod_{i=1}^m \omega_i$ with $\omega_i = \omega(n_i, k_i)$. It follows from (A.5) and (A.4) that

$$\omega_{\llbracket m \rrbracket} = \sum_{j=0}^{[\frac{m}{2}]} \sum_{\substack{B \subset \llbracket m \rrbracket \\ |B|=m-2j}} : \omega_B : \mathbb{E} \left[\omega_{\llbracket m \rrbracket \setminus B} \right].$$

Therefore,

$$(3.21) \quad \begin{aligned} \mathbb{S}_m^{(N)} &= \hat{\beta}_N^m \sum_{n_1, \dots, n_m} \sum_{k_1, \dots, k_m} \left(\sum_{j=0}^{[\frac{m}{2}]} \sum_{\substack{B \subset \llbracket m \rrbracket \\ |B|=m-2j}} : \omega_B : \mathbb{E} \left[\omega_{\llbracket m \rrbracket \setminus B} \right] \right) P_{\mathbf{n}^*} \\ &= \sum_{j=0}^{[\frac{m}{2}]} \sum_{\substack{B \subset \llbracket m \rrbracket \\ |B|=m-2j}} \left(\hat{\beta}_N^m \sum_{n_1, \dots, n_m} \sum_{k_1, \dots, k_m} : \omega_B : \mathbb{E} \left[\omega_{\llbracket m \rrbracket \setminus B} \right] P_{\mathbf{n}^*} \right). \end{aligned}$$

For each $j = 0, 1, \dots, [\frac{m}{2}]$, all the terms in the summation $\sum_{\substack{B \subset \llbracket m \rrbracket \\ |B|=m-2j}}$ are equal by the symmetry. We fix $j \in \{0, 1, \dots, [\frac{m}{2}]\}$. Note that there are in total $\binom{m}{m-2j}$ subsets in $\llbracket m \rrbracket$ with cardinality $m - 2j$. Without loss of generality, we assume $B = \{2j + 1, \dots, m\}$. Then by (A.4), we have

$$(3.22) \quad \mathbb{E}[\omega_{\llbracket m \rrbracket \setminus B}] = \mathbb{E}[\omega_{\llbracket 2j \rrbracket}] = \sum_V \prod_{\{\ell_1, \ell_2\} \in V} \mathbb{E}[\omega_{\ell_1} \omega_{\ell_2}],$$

where the sum \sum_V is taken over all pair partitions of $\llbracket 2j \rrbracket$. By the symmetry again, the summations

$$\sum_{n_1, \dots, n_m} \sum_{k_1, \dots, k_m} : \omega_B : \prod_{\{\ell_1, \ell_2\} \in V} \mathbb{E}[\omega_{\ell_1} \omega_{\ell_2}] P_{\mathbf{n}^*}$$

coincide with each other for all the partitions V , and note that there are in total $(2j - 1)!!$ pair partitions of $\llbracket 2j \rrbracket$. Thus, we have the following equation:

$$(3.23) \quad \begin{aligned} \mathbb{S}_m^{(N)} &= \sum_{j=0}^{\lfloor \frac{m}{2} \rfloor} \binom{m}{m-2j} (2j-1)!! \left[\hat{\beta}_N^m \sum_{n_1, \dots, n_m} \sum_{k_1, \dots, k_m} : \omega_B : \prod_{\ell=1}^j \mathbb{E}[\omega_{2\ell-1} \omega_{2\ell}] P_{\mathbf{n}^*} \right] \\ &= \sum_{j=0}^{\lfloor \frac{m}{2} \rfloor} \frac{m!}{j!(m-2j)!2^j} \left[\hat{\beta}_N^m \sum_{n_1, \dots, n_m} \sum_{k_1, \dots, k_m} : \omega_B : \prod_{\ell=1}^j \mathbb{E}[\omega_{2\ell-1} \omega_{2\ell}] P_{\mathbf{n}^*} \right], \end{aligned}$$

where we recall $B = \{2j + 1, \dots, m\}$.

Compare (3.23) with Hu-Meyer's formula (2.19) and also note (3.20). It then follows easily that in order to prove (3.18), it suffices to show, setting $\beta = 1$ without loss of generality,

$$(3.24) \quad \begin{aligned} Y_N &\stackrel{\text{def}}{=} \frac{1}{m!} N^{-m\theta} \sum_{n_1, \dots, n_m} \sum_{k_1, \dots, k_m} : \omega_B : \prod_{\ell=1}^j \mathbb{E}[\omega_{2\ell-1} \omega_{2\ell}] P_{\mathbf{n}^*} - \mathbf{I}_{m-2j}^{(N)}(\text{Tr}^j \hat{\mathbf{g}}_m) \\ &= \mathbf{I}_{m-2j}^{(N)} \left(N^{-2j\theta} N^{(m-2j)\frac{1}{\rho}} \sum_{n_i: i \in \llbracket 2j \rrbracket} \sum_{k_i: i \in \llbracket 2j \rrbracket} \prod_{\ell=1}^j \mathbb{E}[\omega_{2\ell-1} \omega_{2\ell}](P_{\mathbf{n}^*}/m!) \right) - \mathbf{I}_{m-2j}^{(N)}(\text{Tr}^j \hat{\mathbf{g}}_m) \\ &\rightarrow 0 \text{ in } L^2 \text{ as } N \rightarrow \infty. \end{aligned}$$

for all $j = 0, 1, \dots, \lfloor \frac{m}{2} \rfloor$, where we recall that $\mathbf{I}_{m-2j}^{(N)}(f)$ is given in (3.6).

We prove (3.24) for $j = 0$, and the general case can be proved in a similar spirit. When $j = 0$, we have

$$Y_N = \mathbf{I}_m^{(N)} \left(N^{m/\rho} P_{\mathbf{n}^*}/m! - \hat{\mathbf{g}}_m \right).$$

Denoting $\mathfrak{D}_N = \mathfrak{D}_N(\mathbf{n}, \mathbf{k}) = N^{m/\rho} P_{\mathbf{n}^*}/m! - \mathcal{A}_N(\hat{\mathbf{g}}_m)(\mathbf{n}/N, \mathbf{k}/N^{1/\rho})$, $\mathfrak{D}'_N = \mathfrak{D}_N(\mathbf{n}', \mathbf{k}')$ and $\tilde{\mathfrak{D}}'_N = \mathfrak{D}_N(\mathbf{n}', \mathbf{k})$, we have

$$(3.25) \quad \begin{aligned} \mathbb{E}[Y_N^2] &= N^{-2m(\theta + \frac{1}{\rho})} \sum_{\mathbf{n}, \mathbf{n}' \in \llbracket N \rrbracket^m} \sum_{\mathbf{k}, \mathbf{k}' \in \mathbb{Z}^m} \mathbb{E} \left[: \omega_{\llbracket m \rrbracket} : : \omega'_{\llbracket m \rrbracket} : \right] \mathfrak{D}_N \mathfrak{D}'_N \\ &= m! N^{-2m(\theta + \frac{1}{\rho})} \sum_{\mathbf{n}, \mathbf{n}' \in \llbracket N \rrbracket^m} \sum_{\mathbf{k} \in \mathbb{Z}^m} \prod_{i=1}^m \gamma(n_i - n'_i) \mathfrak{D}_N \tilde{\mathfrak{D}}'_N \\ &\lesssim m! \int_{[0,1]^{2m} \times \mathbb{R}^m} \prod_{i=1}^m |t_i - t'_i|^{2H-2} \mathfrak{D}_N \tilde{\mathfrak{D}}'_N dt dt' d\mathbf{x} \end{aligned}$$

where the second equality follows from (A.8) and we use the notations that link the discrete summations and continuum integrals: $t_i = n_i/N$, $t'_i = n'_i/N$ and $x_i = k_i/N^{1/\rho}$. In light of Lemma 2.2, we have

$$(3.26) \quad \mathbb{E}[Y_N^2] \lesssim m! \int_{[0,1]^{2m} \times \mathbb{R}^m} \prod_{i=1}^m |t_i - t'_i|^{2H-2} \left| \mathbf{D}_N \tilde{\mathbf{D}}'_N \right| dt dt' d\mathbf{x},$$

where $\mathbf{D}_N = \mathbf{D}_N(\mathbf{n}, \mathbf{k}) = N^{m/\rho} P_{\mathbf{n}^*}/m! - \hat{\mathbf{g}}_m(\mathbf{t}, \mathbf{x})$ and $\tilde{\mathbf{D}}'_N = \mathbf{D}_N(\mathbf{n}', \mathbf{k})$.

Inspired by [6], we decompose $I \stackrel{\text{def}}{=} [0, 1]^{2m} \times \mathbb{R}^m$ as $I \cap (I_1 \cup I_2 \cup I_3)$, where

$$I_1 \stackrel{\text{def}}{=} \bigcap_{\substack{i, i' \in \llbracket 2m \rrbracket \\ i \neq i'}} \left\{ |t_i - t_{i'}| > \varepsilon \right\} \cap \bigcap_{i=1}^m \left\{ |x_i| < M \right\},$$

$$I_2 \stackrel{\text{def}}{=} \bigcup_{\substack{i, i' \in \llbracket 2m \rrbracket \\ i \neq i'}} \{|t_i - t_{i'}| \leq \varepsilon\}, \text{ and } I_3 \stackrel{\text{def}}{=} \bigcup_{i=1}^m \{|x_i| \geq M\},$$

for some fixed $\varepsilon, M > 0$. On $I \cap I_1$, $\prod_{i=1}^m |t_i - t'_i|^{2H-2} |\mathbf{D}_N \tilde{\mathbf{D}}'_N|$ is uniformly bounded and hence the integral converges to 0 by the local limit theorem. Then, we can get $\lim_{N \rightarrow \infty} \mathbb{E}[Y_N^2] = 0$, if we can show the integral on $I \cap I_2$ (resp. $I \cap I_3$) can be arbitrarily small if we choose ε sufficiently small (resp. M sufficiently large).

By (1.6) and (1.7), we have

$$(3.27) \quad \begin{aligned} (N^{m/\rho} P_{\mathbf{n}^*}) (N^{m/\rho} P_{(\mathbf{n}')^*}) &\lesssim (t_1^*(t_2^* - t_1^*) \cdots (t_m^* - t_{m-1}^*))^{-1/\rho} N^{m/\rho} P_{(\mathbf{n}')^*}, \\ \hat{\mathbf{g}}_m(x_0; \mathbf{t}, \mathbf{x}) \hat{\mathbf{g}}_m(x_0; \mathbf{t}', \mathbf{x}) &\lesssim (t_1^*(t_2^* - t_1^*) \cdots (t_m^* - t_{m-1}^*))^{-1/\rho} \hat{\mathbf{g}}_m(x_0; \mathbf{t}', \mathbf{x}), \end{aligned}$$

and similarly for the cross terms $N^{m/\rho} P_{\mathbf{n}^*} \hat{\mathbf{g}}_m(\mathbf{t}', \mathbf{x})$ and $N^{m/\rho} P_{(\mathbf{n}')^*} \hat{\mathbf{g}}_m(\mathbf{t}, \mathbf{x})$, where $t_i^* = t_{\sigma(i)}$ for some permutation σ on $\llbracket m \rrbracket$ such that $t_1^* \leq t_2^* \leq \cdots \leq t_m^*$. These inequalities and the fact for all $\mathbf{t} \in [0, 1]^m$,

$$\int_{\mathbb{R}^m} N^{m/\rho} P_{\mathbf{n}^*} d\mathbf{x} = \int_{\mathbb{R}^m} (m!) \hat{\mathbf{g}}_m d\mathbf{x} = 1,$$

yield that $\int_I \prod_{i=1}^m |t_i - t'_i|^{2H-2} |\mathbf{D}_N \tilde{\mathbf{D}}'_N| dt dt' d\mathbf{x}$ is bounded uniformly in N by (up to a multiplicative constant)

$$(3.28) \quad C_m \stackrel{\text{def}}{=} \frac{1}{m!} \int_{[0,1]^{2m}} \prod_{i=1}^m |t_i - t'_i|^{2H-2} (t_1^*(t_2^* - t_1^*) \cdots (t_m^* - t_{m-1}^*))^{-1/\rho} dt dt',$$

which is finite due to $2H - 2 > -1$ and $-1/\rho > -1$. This implies that the integral on $I \cap I_2$ can be arbitrarily small by choosing ε sufficiently small. Similarly, for the integral on $I \cap I_3$,

$$\begin{aligned} \int_{I \cap I_3} \prod_{i=1}^m |t_i - t'_i|^{2H-2} |\mathbf{D}_N \tilde{\mathbf{D}}'_N| dt dt' d\mathbf{x} &\lesssim C_m \left(P \left\{ \sup_{0 \leq t \leq 1} |X_t| \geq M \mid X_1 = x_0 \right\} \right. \\ &\quad \left. + P \left\{ \max_{0 \leq m \leq N} |S_m| \geq N^{1/\rho} M \mid S_0 = N^{1/\rho} x_0 \right\} \right). \end{aligned}$$

Noting that $P \left\{ \max_{0 \leq m \leq N} |S_m| \geq N^{1/\rho} M \mid S_0 = N^{1/\rho} x_0 \right\}$ converges to $P \left\{ \sup_{0 \leq t \leq 1} |X_t| \geq M \mid X_1 = x_0 \right\}$ as N goes to infinity, we can make the integral on $I \cap I_3$ as small uniformly in N as we want by choosing M sufficiently large. This proves (3.24) for $j = 0$.

For general $j = 0, 1, \dots, \lfloor \frac{m}{2} \rfloor$, using the same argument leading to (3.26), we have

$$(3.29) \quad \mathbb{E}[Y_N^2] \lesssim (m - 2j)! \int_{[0,1]^{2(m-2j)} \times \mathbb{R}^{m-2j}} \prod_{i=1}^{m-2j} |t_i - t'_i|^{2H-2} |\mathbf{D}_N \tilde{\mathbf{D}}'_N| dt dt' d\mathbf{x},$$

where now

$$(3.30) \quad \begin{aligned} \mathbf{D}_N &= \mathbf{D}_N(\mathbf{n}, \mathbf{k}) = N^{-2j\theta} N^{(m-2j)\frac{1}{\rho}} \sum_{n_i: i \in \llbracket 2j \rrbracket} \sum_{k_i: i \in \llbracket 2j \rrbracket} \prod_{\ell=1}^j \mathbb{E}[\omega_{2\ell-1} \omega_{2\ell}] (P_{\mathbf{n}^*} / m!) - \text{Tr}^j \hat{\mathbf{g}}_m \\ &\stackrel{\text{def}}{=} \mathcal{Q}_N - \text{Tr}^j \hat{\mathbf{g}}_m \end{aligned}$$

and $\tilde{\mathbf{D}}'_N = \mathbf{D}_N(\mathbf{n}', \mathbf{k})$. Then, we estimate the right-hand side of (3.29) in a similar way as for the case $j = 0$, i.e., we split $[0, 1]^{2(m-2j)} \times \mathbb{R}^{m-2j}$ as the union of its restrictions on I_1, I_2 and I_3 and then analyse the integral restricted on I_i 's separately for $i = 1, 2, 3$. The analysis for the integral

on I_1 is the same as for $j = 0$. To argue that the integral on I_2 and I_3 can be arbitrarily small if we choose ε sufficiently small and M sufficiently large, it suffices to find a uniform (in N) upper bound for

$$(3.31) \quad A \stackrel{\text{def}}{=} \int_{[0,1]^{2(m-2j)} \times \mathbb{R}^{m-2j}} \prod_{i=1}^{m-2j} |t_i - t'_i|^{2H-2} \text{Tr}^j \hat{\mathbf{g}}_m(x_0; \mathbf{t}, \mathbf{x}) \text{Tr}^j \hat{\mathbf{g}}_m(x_0; \mathbf{t}', \mathbf{x}) dt dt' d\mathbf{x}$$

and

$$(3.32) \quad B \stackrel{\text{def}}{=} \int_{[0,1]^{2(m-2j)} \times \mathbb{R}^{m-2j}} \prod_{i=1}^{m-2j} |t_i - t'_i|^{2H-2} \mathcal{Q}_N \tilde{\mathcal{Q}}'_N dt dt' d\mathbf{x}$$

where \mathcal{Q}_N is given in (3.30) and $\tilde{\mathcal{Q}}'_N = \mathcal{Q}_N(\mathbf{n}', \mathbf{k})$. This is true, noting that Remark 2.9 yields

$$A = \mathbb{E} \left[\left| \mathbf{I}_{m-2j}(\text{Tr}^j \hat{\mathbf{g}}_m) \right|^2 \right] \leq \mathbb{E} \left[|\mathbb{I}_m(\mathbf{g}_m)|^2 \right] < \infty,$$

and similarly, recalling that $\mathbb{S}_m^{(N)}$ is given in (3.5), we have $B \lesssim \mathbb{E}[|\mathbb{S}_m^{(N)}|^2]$ which has a uniform upper bound by (3.36).

In the above, we have assumed m is an odd integer. If m is even, the analysis is the same except for the case $j = \frac{m}{2}$. In this case, we have

$$Y_N = N^{-m\theta} \sum_{n_i: i \in \llbracket m \rrbracket} \sum_{k_i: i \in \llbracket m \rrbracket} \prod_{\ell=1}^{m/2} \mathbb{E}[\omega_{2\ell-1} \omega_{2\ell}] (P_{\mathbf{n}^*} / m!) - \text{Tr}^{m/2} \hat{\mathbf{g}}_m,$$

which is deterministic and converges to 0 by the local limit theorem and a similar argument proving $\lim_{N \rightarrow \infty} \mathbb{E}[Y_N^2] = 0$ for $j = 0$.

Finally, one can prove the weak convergence for linear combinations of $(\mathbb{S}_{l_1}^{(N)}, \dots, \mathbb{S}_{l_k}^{(N)})$ in a similar way, and hence (3.19) holds due to Theorem B.4. \square

Now we are ready to prove our main result.

Proof of Theorem 1.2. By Proposition 3.4, we have for all $M \in \mathbb{N}$,

$$\mathcal{Z}_\omega^{(N,M)} \stackrel{\text{def}}{=} \sum_{m=0}^M \frac{1}{m!} \mathbb{S}_m^{(N)} \xrightarrow{d} \mathcal{Z}^{(M)} \stackrel{\text{def}}{=} \sum_{m=0}^M \beta^m \mathbb{I}_m(\mathbf{g}_m(\mathbf{x}_0; \cdot)), \text{ as } N \rightarrow \infty.$$

Recalling Proposition 2.10 which yields the L^1 -convergence of $\mathcal{Z}^{(M)}$ to \mathcal{Z} as $M \rightarrow \infty$, we only need to show that $\mathcal{Z}_\omega^{(N,M)}$ converges to $\mathcal{Z}_\omega^{(N)} = \mathcal{Z}_\omega^{(N)}(\hat{\beta}_N, N^{1/\rho} x_0)$ given in (3.2) in probability uniformly in N as $M \rightarrow \infty$ by Lemma B.1. This follows from (3.37) obtained in Section 3.3.

3.3. L^1 -bound of rescaled partition functions. Recalling that ω is Gaussian, we have

$$\begin{aligned} \mathbb{E} \left[\exp \left(N^{-\theta} \sum_{n=1}^N \omega(n, S_n) \right) \right] &= \mathbb{E} \left[\exp \left(N^{-\theta} \sum_{n=1}^N \sum_{k \in \mathbb{Z}} \omega(n, k) \mathbf{1}_{[S_n=k]} \right) \right] \\ &= \mathbb{E} \left[\exp \left(\frac{1}{2} N^{-2\theta} \sum_{n, n'=1}^N \sum_{k, k' \in \mathbb{Z}} \gamma(n - n') \delta_{kk'} \mathbf{1}_{[S_n=k]} \mathbf{1}_{[S_{n'}=k']} \right) \right] \\ &= \mathbb{E} \left[\exp \left(\frac{1}{2} N^{-2\theta} \sum_{n, n'=1}^N \gamma(n - n') \mathbf{1}_{[S_n=S_{n'}]} \right) \right]. \end{aligned}$$

In this section, we shall prove a discretised version of Proposition 2.6, i.e., to show that the above exponential moment is uniformly bounded.

Noting that $S_n \in \mathbb{Z}$ and using the identity

$$\mathbf{1}_{[S_n=S_{n'}]} = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{i(S_n-S_{n'})\lambda} d\lambda,$$

we have

$$\begin{aligned} & \mathbb{E} \left[\exp \left(\frac{1}{2} N^{-2\theta} \sum_{n,n'=1}^N \gamma(n-n') \mathbf{1}_{[S_n=S_{n'}]} \right) \right] \\ &= \sum_{m=0}^{\infty} \frac{1}{m!} (4\pi)^{-m} N^{-2m\theta} \sum_{n_1, \dots, n_m=1}^N \sum_{n'_1, \dots, n'_m=1}^N \int_{[-\pi, \pi]^m} \prod_{i=1}^m \gamma(n_i - n'_i) \mathbb{E} \left[e^{i \sum_{i=1}^m (S_{n_i} - S_{n'_i}) \lambda_i} \right] d\boldsymbol{\lambda}. \end{aligned}$$

Using changes of variables $n_i = Nt_i$, $n'_i = Nt'_i$ and $\lambda_i = u_i/N^{1/\rho}$ for $i = 1, \dots, m$, we have for each term on the right-hand side of the above equation,

$$(3.33) \quad \begin{aligned} A_m &\stackrel{\text{def}}{=} \frac{1}{m!} (4\pi)^{-m} N^{-2m} \sum_{\mathbf{n}, \mathbf{n}'} \prod_{i=1}^m \left[N^{2-2H} \gamma(N(t_i - t'_i)) \right] \\ &\quad \times \int_{[-\pi N^{1/\rho}, \pi N^{1/\rho}]^m} \mathbb{E} \left[e^{i \sum_{i=1}^m (S_{n_i} - S_{n'_i}) u_i / N^{1/\rho}} \right] d\mathbf{u}. \end{aligned}$$

Recall that $S_n = S_0 + \sum_{j=1}^n Y_j$ and that $\psi(u) = \mathbb{E}[e^{2uY_j}]$ is the characteristic function of Y_j . The 1-lattice distribution of Y_j implies that, for any $\varepsilon > 0$, one can find a positive constant c such that (see [33, eq. (5.14)])

$$(3.34) \quad |\psi(u/N^{1/\rho})| \leq e^{-c|u|^{\rho-\varepsilon}/N}, \text{ for } 1 \leq |u| \leq \pi N^{1/\rho}.$$

Thus, for $n \in \llbracket N \rrbracket$, we have

$$(3.35) \quad \begin{aligned} \int_{-\pi N^{1/\rho}}^{\pi N^{1/\rho}} |\psi(u/N^{1/\rho})|^n du &\leq 2 + \int_{\{1 \leq |u| \leq \pi N^{1/\rho}\}} |\psi(u/N^{1/\rho})|^n du \\ &\leq 2 + \int_{\mathbb{R}} e^{-c|u|^{\rho-\varepsilon}n/N} du \\ &\leq C(n/N)^{-\frac{1}{\rho-\varepsilon}}. \end{aligned}$$

Under the condition (1.13), we may choose $\varepsilon > 0$ sufficiently small such that $2H - \frac{1}{\rho-\varepsilon} > 1$. To estimate A_m given in (3.33), we combine the estimate (3.35) with the argument used in the proof of Proposition 2.6 which leads to (2.36), and we can get for all $N \in \mathbb{N}$,

$$\begin{aligned} A_m &\leq \frac{(2m)!}{m!} C^m \int_{[0,1]_{<}^{2m}} \prod_{i=1}^m |t_{2i}^* - t_{2i-1}^*|^{2H-2-\frac{1}{\rho-\varepsilon}} ds \\ &\leq \frac{(2m)! C^m}{m! \Gamma\left(m(2H - \frac{1}{\rho-\varepsilon}) + 1\right)} \end{aligned}$$

Therefor, we have uniformly in N ,

$$\sum_{m=0}^{\infty} A_m \leq \sum_{m=0}^{\infty} \frac{(2m)! C^m}{m! \Gamma\left(m(2H - \frac{1}{\rho-\varepsilon}) + 1\right)} < \infty,$$

where the finiteness follows from Stirling's formula and $2H - \frac{1}{\rho-\varepsilon} > 1$. This implies

$$(3.36) \quad \sup_{N \in \mathbb{N}} \mathbb{E} \left[\exp \left(N^{-\theta} \sum_{n=1}^N \omega(n, S_n) \right) \right] < \infty.$$

Finally, the above analysis also yields

$$(3.37) \quad \lim_{M \rightarrow \infty} \sup_{N \in \mathbb{N}} \sum_{m=M+1}^{\infty} \mathbb{E} \left[\left| N^{-\theta} \sum_{n=1}^N \omega(n, S_n) \right|^m \right] = 0,$$

which shall be used to prove the weak convergence of rescaled partition functions.

APPENDIX A. PHYSICAL WICK PRODUCT

In order to expand the partition function (1.1) in a proper way to obtain its weak convergence, we shall invoke the notion of *physical* Wick product for general random variables (in contrast to the *probabilistic* Wick product defined via Wiener chaos in Section 2.1). The physical Wick product (also known as Wick power, Wick polynomial or Wick renormalization) was introduced by Wick [40] in the study of quantum field theory. We collect some facts on physical Wick products in this subsection. The reader is referred to [2, 3, 13, 14, 27, 38] for more readable account.

Let $\{X_i\}_{i \in \mathbb{N}}$ be a family of real random variables with finite moments of all orders. The physical Wick product $:X_1 \cdots X_n:$ is defined recursively as follows. For $n = 0$, we set $: := 1$, and for $n \geq 1$,

$$\frac{\partial}{\partial X_i} :X_1 \cdots X_n := :X_1 \cdots \hat{X}_i \cdots X_n:, \quad \mathbb{E}[:X_1 \cdots X_n:] = 0,$$

where \hat{X}_i means the absence of X_i in the product. For example,

$$\begin{aligned} :X_1: &= X_1 - \mathbb{E}[X_1]; \\ :X_1 X_2: &= X_1 X_2 - X_1 \mathbb{E}[X_2] - X_2 \mathbb{E}[X_1] + 2\mathbb{E}[X_1] \mathbb{E}[X_2] - \mathbb{E}[X_1 X_2]. \end{aligned}$$

We remark that different indices may refer to the same random variable. In this situation, as an example, we also write $:X^n Y^m := \stackrel{\text{def}}{=} :X_1 \cdots X_n Y_1 \cdots Y_m:$, if $X_1 = \cdots = X_n = X$ and $Y_1 = \cdots = Y_m = Y$.

If we assume $\mathbb{E} \left[e^{\beta \sum_{i=1}^n |X_i|} \right] < \infty$ for some $\beta > 0$, the physical Wick product can be equivalently defined by

$$(A.1) \quad :X_1 \cdots X_n := \stackrel{\text{def}}{=} \frac{\partial^n}{\partial z_1 \cdots \partial z_n} G(z_1, \dots, z_n; X_1, \dots, X_n) \Big|_{z=0},$$

where

$$G(z_1, \dots, z_n; X_1, \dots, X_n) \stackrel{\text{def}}{=} \frac{e^{\sum_{i=1}^n z_i X_i}}{\mathbb{E} \left[e^{\sum_{i=1}^n z_i X_i} \right]}$$

is called the generating function (or Wick exponential). If $\{X_i\}_{i \in \mathbb{N}}$ is a centred Gaussian family, the generating function is simply $G(z, \mathbf{X}) = e^{z \cdot \mathbf{X} - z \cdot Q z / 2}$ where $Q = (\mathbb{E}[X_i X_j])_{n \times n}$ is the covariance matrix of $\mathbf{X} = (X_1, \dots, X_n)$, and the resulting Wick products are related to Hermit polynomials (see (2.8) in Section 2.1). In particular, for a Gaussian random vector (X_1, \dots, X_n) , the *physical* Wick product $:X_1 \cdots X_n:$ coincides with *probabilistic* Wick product $X_1 \diamond X_2 \diamond \cdots \diamond X_n$ defined in Section 2.1.

We collect some basic properties of physical Wick products. Clearly $:X_1 X_2 \cdots X_n:$ only involves the random variables X_1, \dots, X_n and their joint moments up to order n , and it is *symmetric* and *multilinear* in (X_1, \dots, X_n) (multilinear means linear in terms of each $X_i, i = 1, \dots, n$). If

two groups of random variables $\{X_1, X_2, \dots, X_m\}$ and $\{X_{m+1}, \dots, X_n\}$ are independent of each other, then $:X_1 \cdots X_n := X_1 \cdots X_m :: X_{m+1} \cdots X_n :$ (see [2, eq. (2.4)]). We remind that the physical Wick product is no longer associative, which is different from the ordinary product. For instance, assuming $\mathbb{E}[X] = 0$, we have $:XXX := X^3 - 3X\mathbb{E}[X^2] - \mathbb{E}[X^3]$ which is different from $:XY :|_{Y=:X^2:} = X^3 - X\mathbb{E}[X^2] - \mathbb{E}[X^3]$. Thus a physical Wick product is a single term whose value is determined by the definition, and cannot be viewed as a composition of two (or several) physical Wick products. For instance, all the physical Wick products $:X^3 :$, $:X^2X :$ and $:XX^2 :$ mean the same $:XXX :$, and in particular $:XX^2 : \neq :XY :|_{Y=:X^2:}$.

For any finite index set $A = \{i_1, \dots, i_n\} \subset \mathbb{N}$, we denote $:X_A \stackrel{\text{def}}{=} X_{i_1} \cdots X_{i_n} :$ and similarly, we take the notation $X_A \stackrel{\text{def}}{=} \prod_{i \in A} X_i$ for the ordinary product. We use X^A to denote the set $\{X_i, i \in A\}$ of random variables.

We recall some facts about cumulants. Let $\kappa(X^A)$ denote the joint cumulant of $X^A = \{X_i, i \in A\}$. Then

$$(A.2) \quad \kappa(X^A) = \sum_V (|V| - 1)! (-1)^{|V|-1} \prod_{i=1}^{|V|} \mathbb{E}[X_{V_i}]$$

and

$$(A.3) \quad \mathbb{E}[X_A] = \sum_V \prod_{i=1}^{|V|} \kappa(X^{V_i}),$$

where the sum \sum_V is taken over all partitions $V = \{V_1, \dots, V_k\}, k \geq 1$ of A , and $|V| = k$ is the partition number. For instance,

$$\kappa(X_1) = \mathbb{E}[X_1], \quad \kappa(X_1, X_2) = \mathbb{E}[X_1 X_2] - \mathbb{E}[X_1] \mathbb{E}[X_2],$$

and

$$\kappa(X_1, X_2, X_3) = \mathbb{E}[X_1 X_2 X_3] - \mathbb{E}[X_1 X_2] \mathbb{E}[X_3] - \mathbb{E}[X_1 X_3] \mathbb{E}[X_2] - \mathbb{E}[X_2 X_3] \mathbb{E}[X_1] + 2\mathbb{E}[X_1] \mathbb{E}[X_2] \mathbb{E}[X_3].$$

If $\{X_i\}_{i \in \mathbb{N}}$ is *Gaussian*, we have $\kappa(X^A) = 0$ if $|A| \geq 3$. If we assume further $\mathbb{E}[X_i] = 0$, the formula (A.3) reduces to the Wick's theorem:

$$(A.4) \quad \mathbb{E}[X_A] = \begin{cases} 0, & \text{if } |A| \text{ is odd,} \\ \sum_V \prod_{\{i,j\} \in V} \mathbb{E}[X_i X_j], & \text{if } |A| \text{ is even,} \end{cases}$$

where the summation \sum_V is taken over all *pair partitions* $V = \{V_1, \dots, V_{|A|/2}\}$ of A .

Ordinary products and physical Wick products are connected by the following formula (see [38, Prop. 1] or [3, Appendix B]),

$$(A.5) \quad X_A = \sum_{B \subset A} :X_B : \sum_V \prod_{i=1}^{|V|} \kappa(X^{V_i}) = \sum_{B \subset A} :X_B : \mathbb{E}[X_{A \setminus B}],$$

and

$$(A.6) \quad :X_A := \sum_{B \subset A} X_B \sum_V (-1)^{|V|} \prod_{i=1}^{|V|} \kappa(X^{V_i}),$$

where the sum $\sum_{B \subset A}$ is taken over all subsets $B \subset A$ including $B = \emptyset$, and the sum \sum_V is over all partitions $V = \{V_1, \dots, V_k\}, k \geq 1$ of the set $A \setminus B$. We use the convention $X_\emptyset = :X_\emptyset := \kappa(X_\emptyset) = 1$.

The following formula (see [3, Appendix B] or [15, Lemma 4.5]) will be used

$$(A.7) \quad \mathbb{E}[:X_A :: X_B :] = \sum_V \prod_{i=1}^{|V|} \kappa(X^{V_i}),$$

where the summation \sum_V is taken over all partitions $V = \{V_1, \dots, V_k\}, k \geq 1$ of $A \cup B$ satisfying $V_i \cap A \neq \emptyset \neq V_i \cap B$ for each V_i . In particular, if we assume $\{X_i\}_{i \in \mathbb{N}}$ is a centered *Gaussian* family, equations (A.7) and (A.4) yield

$$(A.8) \quad \mathbb{E}[:X_A :: X_B :] = \begin{cases} 0, & \text{if } |A| \neq |B| \\ \sum_V \prod_{\{i,j\} \in V} \mathbb{E}[X_i X_j], & \text{if } |A| = |B|, \end{cases}$$

where the summation \sum_V is taken over all *pair partitions* $V = \{V_1, \dots, V_{|A|}\}$ of $A \cup B$ such that $V_k = \{i, j\}$ with $i \in A, j \in B$ for $k = 1, \dots, |A|$.

APPENDIX B. SOME PRELIMINARIES ON CONVERGENCE OF PROBABILITY MEASURES

The following result can be found in [5, Theorem 3.2, Chapter 1].

Lemma B.1. *Consider random variables $Y_n^{(N)}, Y^{(N)}, Y_n$ and Y , such that $Y_n^{(N)} \xrightarrow{d} Y_n$ as $N \rightarrow \infty$, $Y_n \xrightarrow{d} Y$ as $n \rightarrow \infty$, and $Y_n^{(N)}$ converges in probability to $Y^{(N)}$ uniformly in N as $n \rightarrow \infty$, then we have $Y^{(N)} \xrightarrow{d} Y$ as $N \rightarrow \infty$. That is, assuming*

$$\begin{array}{ccc} Y_n^{(N)} & \xrightarrow[n \rightarrow \infty]{\text{in probability, uniformly in } N} & Y^{(N)} \\ \downarrow d \text{ } N \rightarrow \infty & & \\ Y_n & \xrightarrow[n \rightarrow \infty]{d} & Y, \end{array}$$

we have $Y^{(N)} \xrightarrow{d} Y$ as $N \rightarrow \infty$.

Lemma B.2. *Let $k \geq 1$ be an integer. If $(X_1^{(n)}, \dots, X_k^{(n)}) \xrightarrow{d} (X_1, \dots, X_k)$ as $n \rightarrow \infty$, we have*

$$f(X_1^{(n)}, \dots, X_k^{(n)}) \xrightarrow{d} f(X_1, \dots, X_k)$$

for any continuous function f .

Lemma B.3. *Let $k \geq 1$ be an integer. Suppose $(X_1^{(n)}, \dots, X_k^{(n)}) \xrightarrow{d} (X_1, \dots, X_k)$ as $n \rightarrow \infty$ and assume that for any subset A of $\{1, \dots, k\}$, $\left\{ \prod_{i \in A} X_i^{(n)} \right\}_{n \in \mathbb{N}}$ is uniformly integrable. Then $:X_1^{(n)} \dots X_k^{(n)} : \xrightarrow{d} :X_1 \dots X_k :$ as $n \rightarrow \infty$.*

Proof Lemma B.2 yields the weak convergence of the ordinary product $\prod_{i \in A} X_i^{(n)} \xrightarrow{d} \prod_{i \in A} X_i$ for any $A \subset \{1, \dots, k\}$ as $n \rightarrow \infty$. By (A.6), it suffices to prove the convergence of the cumulants

$$\lim_{n \rightarrow \infty} \kappa(X_i^{(n)}, i \in A) = \kappa(X_i, i \in A).$$

This follows from the Skorohod representation theorem, equation (A.2), and the assumption of uniform integrability for the products of $X_1^{(n)}, \dots, X_k^{(n)}$. \square

Theorem B.4. [Cramér-Wold Theorem] As $n \rightarrow \infty$, $(X_1^{(n)}, \dots, X_k^{(n)}) \xrightarrow{d} (X_1, \dots, X_k)$ if and only if

$$\sum_{i=1}^k a_i X_i^{(n)} \xrightarrow{d} \sum_{i=1}^k a_i X_i, \quad \text{for all } (a_1, \dots, a_k) \in \mathbb{R}^k.$$

APPENDIX C. MISCELLANEOUS RESULTS

Let $f : [0, 1]^m \rightarrow \mathbb{R}$ be an integrable function. For a fixed $N \in \mathbb{N}$, let $t_i = i/N$ for $i = 0, 1, \dots, N$ and denote $I_i = (t_{i-1}, t_i]$. Define

$$(B.1) \quad f_N(t_1, \dots, t_m) \stackrel{\text{def}}{=} N^m \int_{I_{\ell_1} \times \dots \times I_{\ell_m}} f(s_1, \dots, s_m) \mathbf{d}\mathbf{s},$$

if $(t_1, \dots, t_m) \in I_{\ell_1} \times \dots \times I_{\ell_m}$ for some $\ell_i \in \llbracket N \rrbracket$. That is, f_N is the conditional expectation of f with respect to the σ -field generated by the rectangles of the form $I_{\ell_1} \times \dots \times I_{\ell_m}$. Then we have the following inequality of Jensen type.

Lemma C.1. Assume $H \in (1/2, 1]$ and suppose $f, g : [0, 1]^m \rightarrow \mathbb{R}$ is an integrable function. Then, for each $N \in \mathbb{N}$,

$$(B.2) \quad \begin{aligned} & \int_{[0, 1]^{2m}} \prod_{i=1}^m |s_i - t_i|^{2H-2} |f_N(s_1, \dots, s_m)| |g_N(t_1, \dots, t_m)| \mathbf{d}\mathbf{s} \mathbf{d}\mathbf{t} \\ & \leq C^m \int_{[0, 1]^{2m}} \prod_{i=1}^m |s_i - t_i|^{2H-2} |f(s_1, \dots, s_m)| |g(t_1, \dots, t_m)| \mathbf{d}\mathbf{s} \mathbf{d}\mathbf{t}, \end{aligned}$$

for some constant C depending on H only.

Proof We only prove the case $m = 1$, and the other cases $m > 1$ follows from an induction argument together with Fubini's theorem. It suffices to prove (B.2) with $m = 1$ for nonnegative functions, i.e., for $f, g \geq 0$,

$$(B.3) \quad \int_{I_i \times I_j} |s - t|^{2H-2} f_N(s) g_N(t) \mathbf{d}s \mathbf{d}t \leq C \int_{I_i \times I_j} |s - t|^{2H-2} f(s) g(t) \mathbf{d}s \mathbf{d}t, \quad 1 \leq i, j \leq N,$$

where recalling $I_i = (t_{i-1}, t_i]$. We shall prove (B.3) for simple nonnegative functions. The general case can be proved by a limiting argument and thus is omitted.

For $i = j$, we assume that on I_i ,

$$f(s) = \sum_{\ell=1}^k a_\ell \mathbf{1}_{A_\ell}(s), \quad g(t) = \sum_{\ell=1}^k b_\ell \mathbf{1}_{A_\ell}(t),$$

where $A_\ell = (t_{i-1} + (\ell - 1)/Nk, t_{i-1} + \ell/Nk]$ with $\ell = 1, \dots, k$ form a uniform partition of the interval I_i , and a_ℓ, b_ℓ are nonnegative numbers. Then, denoting $\bar{a} = \frac{1}{k} \sum_{\ell=1}^k a_\ell$, $\bar{b} = \frac{1}{k} \sum_{\ell=1}^k b_\ell$,

$$f_N(s) = \bar{a}, \quad g_N(t) = \bar{b}, \quad \text{on } I_i.$$

The left-hand side of (B.3) is

$$(B.4) \quad \bar{a} \bar{b} \int_{I_i \times I_i} |s - t|^{2H-2} \mathbf{d}s \mathbf{d}t = \sum_{\ell=1}^k \sum_{m=1}^k a_\ell b_m \left(\frac{1}{k^2} \int_{I_i \times I_i} |s - t|^{2H-2} \mathbf{d}s \mathbf{d}t \right),$$

and the right-hand side without the constant C is

$$(B.5) \quad \sum_{\ell=1}^k \sum_{m=1}^k a_{\ell} b_m \left(\int_{A_{\ell} \times A_m} |s-t|^{2H-2} ds dt \right).$$

Then, (B.3) follows from (B.4), (B.5) and the following inequality

$$\frac{1}{k^2} \int_{I_i \times I_i} |s-t|^{2H-2} ds dt \leq C \min_{\ell, m \in \llbracket k \rrbracket} \int_{A_{\ell} \times A_m} |s-t|^{2H-2} ds dt,$$

where C is a constant depending on H only. Indeed, by change of variables, the above inequality is equivalent to

$$\frac{1}{k^2} \int_0^1 \int_0^1 |s-t|^{2H-2} ds dt \leq C \min_{\ell, m \in \llbracket k \rrbracket} \int_{(m-1)/k}^{m/k} \int_{(\ell-1)/k}^{\ell/k} |s-t|^{2H-2} ds dt,$$

which holds for $C = \int_0^1 \int_0^1 |s-t|^{2H-2} ds dt$, noting that for all $\ell, m \in \llbracket k \rrbracket$, we have $|s-t| \leq 1$ on $[(\ell-1)/k, \ell/k] \times [(m-1)/k, m/k]$, and hence $\int_{(m-1)/k}^{m/k} \int_{(\ell-1)/k}^{\ell/k} |s-t|^{2H-2} ds dt \geq 1/k^2$. This proves (B.3) for $i = j$.

Now we prove (B.3) for $i \neq j$. By symmetry, we only need to consider $j > i$. Let

$$f(s) = \sum_{\ell=1}^k a_{\ell} \mathbf{1}_{A_{\ell}}(s) \text{ on } I_i = (t_{i-1}, t_i],$$

and

$$g(t) = \sum_{\ell=1}^k b_{\ell} \mathbf{1}_{B_{\ell}}(t) \text{ on } I_j = (t_{j-1}, t_j],$$

where a_{ℓ}, b_{ℓ} are nonnegative numbers, and $\{A_{\ell}, \ell = 1, \dots, k\}$ and $\{B_{\ell}, \ell = 1, \dots, k\}$ are uniform partitions of I_i and I_j , respectively. Then,

$$f_N(s) = \bar{a} \text{ on } I_i \text{ and } g_N(t) = \bar{b} \text{ on } I_j,$$

where $\bar{a} = \frac{1}{k} \sum_{\ell=1}^k a_{\ell}$ and $\bar{b} = \frac{1}{k} \sum_{\ell=1}^k b_{\ell}$. The left-hand side of (B.3) is

$$(B.6) \quad \bar{a} \bar{b} \int_{I_i \times I_j} |s-t|^{2H-2} ds dt = \sum_{\ell=1}^k \sum_{m=1}^k a_{\ell} b_m \left(\frac{1}{k^2} \int_{I_i \times I_j} |s-t|^{2H-2} ds dt \right),$$

and the right-hand side of (B.3) without C is

$$(B.7) \quad \sum_{\ell=1}^k \sum_{m=1}^k a_{\ell} b_m \left(\int_{A_{\ell} \times B_m} |s-t|^{2H-2} ds dt \right).$$

To get (B.3), it suffices to show

$$(B.8) \quad \frac{1}{k^2} \int_{I_i \times I_j} |s-t|^{2H-2} ds dt \leq C \min_{\ell, m \in \llbracket k \rrbracket} \int_{A_{\ell} \times B_m} |s-t|^{2H-2} ds dt$$

for some constant C depending on H only. Note that by change of variables, (B.8) is equivalent to the following equality

$$(B.9) \quad \frac{1}{k^2} \int_{j-i}^{j-i+1} \int_0^1 |s-t|^{2H-2} ds dt \leq C \min_{\ell, m \in \llbracket k \rrbracket} \int_{j-i+(m-1)/k}^{j-i+m/k} \int_{(\ell-1)/k}^{\ell/k} |s-t|^{2H-2} ds dt$$

We shall prove (B.9) for $j = i + 1$ and $j \geq i + 2$ separately. If $j = i + 1$, (B.9) follows directly by choosing

$$C = 2^{2-2H} \int_1^2 \int_0^1 |s-t|^{2H-2} ds dt,$$

noting that

$$\int_{1+(m-1)/k}^{1+m/k} \int_{(\ell-1)/k}^{\ell/k} |s-t|^{2H-2} ds dt \geq 2^{2H-2}/k^2, \text{ for all } \ell, m \in \llbracket k \rrbracket.$$

When $n \stackrel{\text{def}}{=} j - i \geq 2$, to prove (B.9), it suffices to prove that the biggest integral among the integrals over the regions $\{(n + (m - 1)/k, n + m/k) \times ((\ell - 1)/k, \ell/k), \ell \in \llbracket k \rrbracket\}$ can be dominated by the smallest one, i.e.,

$$\int_n^{n+1/k} \int_{1-1/k}^1 |s-t|^{2H-2} ds dt \leq C \int_{n+1-1/k}^{n+1} \int_0^{1/k} |s-t|^{2H-2} ds dt$$

for some constant C only depending on H . This is true because there exists a finite constant depending on H only such that

$$|n-1|^{2H-2} \leq 3^{2-2H} |n+1|^{2H-2} \text{ for all } n \geq 2.$$

This proves (B.9), and hence completes the proof of (B.3) for $i \neq j$. \square

The following result can be calculated by a direct calculation.

Lemma C.2. *Suppose $\alpha_i < 1$ for $i = 1, \dots, n$ and let $\alpha = \sum_{i=1}^n \alpha_i$. Then*

$$\int_{[0 < r_1 < \dots < r_n < r_{n+1} = t]} \prod_{i=1}^n (r_{i+1} - r_i)^{-\alpha_i} d\mathbf{r} = \frac{\prod_{i=1}^n \Gamma(1 - \alpha_i)}{\Gamma(n - \alpha + 1)} t^{n-\alpha},$$

where $\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt$ is the Gamma function.

Acknowledgement The authors would like to thank Rongfeng Sun for his helpful comments. G. Rang is partially supported by the National Natural Science Foundation of China (Grants no. 11971361, 11731012 and 12131019). J. Song is partially supported by Shandong University (Grant No. 11140089963041) and the National Natural Science Foundation of China (Grant No. 12071256).

REFERENCES

- [1] Tom Alberts, Konstantin Khanin, and Jeremy Quastel. The intermediate disorder regime for directed polymers in dimension $1 + 1$. *The Annals of Probability*, 42(3):1212–1256, 2014.
- [2] Florin Avram and Murad S Taqqu. Noncentral limit theorems and appell polynomials. *The Annals of Probability*, pages 767–775, 1987.
- [3] Florin Avram and Murad S Taqqu. On a szegő type limit theorem and the asymptotic theory of random sums, integrals and quadratic forms. In *Dependence in probability and statistics*, pages 259–286. Springer, 2006.
- [4] Sérgio Bezerra, Samy Tindel, and Frederi Viens. Superdiffusivity for a brownian polymer in a continuous gaussian environment. *The Annals of Probability*, 36(5):1642–1675, 2008.
- [5] Patrick Billingsley. *Convergence of probability measures*. John Wiley & Sons, 1999.
- [6] Francesco Caravenna, Rongfeng Sun, and Nikos Zygouras. Polynomial chaos and scaling limits of disordered systems. *Journal of the European Mathematical Society*, 19(1):1–65, 2016.
- [7] Francis Comets. Weak disorder for low dimensional polymers: The model of stable laws. *arXiv preprint math/0603390*, 2006.
- [8] Francis Comets. *Directed polymers in random environments*. Springer, 2017.
- [9] Ivan Corwin. The Kardar–Parisi–Zhang equation and universality class. *Random matrices: Theory and applications*, 1(01):1130001, 2012.
- [10] Giuseppe Da Prato and Jerzy Zabczyk. *Stochastic equations in infinite dimensions*. Cambridge university press, 2014.
- [11] A Dasgupta and G Kallianpur. Chaos decomposition of multiple fractional integrals and applications. *Probability theory and related fields*, 115(4):527–548, 1999.
- [12] Mohammud Foondun, Mathew Joseph, and Shiu-Tang Li. An approximation result for a class of stochastic heat equations with colored noise. *The Annals of Applied Probability*, 28(5):2855–2895, 2018.

- [13] L Giraitis and D Surgailis. Multivariate appell polynomials and the central limit theorem. In *Dependence in probability and statistics*, pages 21–71. Springer, 1986.
- [14] Håkon Gjessing, Helge Holden, Tom Lindstrøm, J Ubøe, and T Zhang. The wick product. *Frontiers in Pure and Applied Probability*, 1:29–67, 1993.
- [15] Martin Hairer and Hao Shen. A central limit theorem for the kpz equation. *The Annals of Probability*, 45(6B):4167–4221, 2017.
- [16] Yao-Zhong Hu and Paul-André Meyer. Sur les intégrales multiples de stratonovitch. In *Séminaire de Probabilités XXII*, pages 72–81. Springer, 1988.
- [17] Yao-zhong Hu and Jia-an Yan. Wick calculus for nonlinear gaussian functionals. *Acta Mathematicae Applicatae Sinica, English Series*, 25(3):399–414, 2009.
- [18] Yaozhong Hu. *Analysis on Gaussian spaces*. World Scientific, 2016.
- [19] Yaozhong Hu, David Nualart, and Jian Song. Feynman–kac formula for heat equation driven by fractional white noise. *The Annals of Probability*, 39(1):291–326, 2011.
- [20] David A Huse and Christopher L Henley. Pinning and roughening of domain walls in Ising systems due to random impurities. *Physical review letters*, 54(25):2708, 1985.
- [21] I. A. Ibragimov and Yu. V. Linnik. *Independent and stationary sequences of random variables*. Wolters-Noordhoff Publishing, Groningen, 1971. With a supplementary chapter by I. A. Ibragimov and V. V. Petrov, Translation from the Russian edited by J. F. C. Kingman.
- [22] Svante Janson et al. *Gaussian hilbert spaces*. Number 129. Cambridge university press, 1997.
- [23] Maria Jolis. On a multiple stratonovich-type integral for some gaussian processes. *Journal of Theoretical Probability*, 19(1):121–133, 2006.
- [24] Mathew Joseph, Davar Khoshnevisan, and Carl Mueller. Strong invariance and noise-comparison principles for some parabolic stochastic pdes. *The Annals of Probability*, 45(1):377–403, 2017.
- [25] Ida Kruk, Francesco Russo, and Ciprian A Tudor. Wiener integrals, malliavin calculus and covariance measure structure. *Journal of Functional Analysis*, 249(1):92–142, 2007.
- [26] Hubert Lacoin. Influence of spatial correlation for directed polymers. *The Annals of Probability*, 39(1):139–175, 2011.
- [27] Jani Lukkarinen and Matteo Marozzi. Wick polynomials and time-evolution of cumulants. *Journal of Mathematical Physics*, 57(8):083301, 2016.
- [28] Ernesto Medina, Terence Hwa, Mehran Kardar, and Yi-Cheng Zhang. Burgers equation with correlated noise: Renormalization-group analysis and applications to directed polymers and interface growth. *Physical Review A*, 39(6):3053, 1989.
- [29] Elchanan Mossel, Ryan O’Donnell, and Krzysztof Oleszkiewicz. Noise stability of functions with low influences: invariance and optimality. *Ann. of Math. (2)*, 171(1):295–341, 2010.
- [30] David Nualart. *The Malliavin calculus and related topics*, volume 1995. Springer, 2006.
- [31] Vladas Pipiras and Murad S Taqqu. Integration questions related to fractional Brownian motion. *Probability theory and related fields*, 118(2):251–291, 2000.
- [32] Guanglin Rang. From directed polymers in spatial-correlated environment to stochastic heat equations driven by fractional noise in $1+1$ dimensions. *Stochastic Processes and their Applications*, 130(6):3408–3444, 2020.
- [33] Jay Rosen. Random walks and intersection local time. *The Annals of Probability*, pages 959–977, 1990.
- [34] Carles Rovira and Samy Tindel. On the Brownian-directed polymer in a gaussian random environment. *Journal of Functional Analysis*, 222(1):178–201, 2005.
- [35] EL Rvaceva. On domains of attraction of multi-dimensional distributions. *Select. Transl. Math. Statist. and Probability*, 2:183–205, 1962.
- [36] Hao Shen, Jian Song, Rongfeng Sun, and Lihu Xu. Scaling limit of a directed polymer among a poisson field of independent walks. *Journal of Functional Analysis*, 281(5):109066, 2021.
- [37] Jian Song. On a class of stochastic partial differential equations. *Stochastic Processes and their Applications*, 127(1):37–79, 2017.
- [38] D Surgailis. On poisson multiple stochastic integrals and associated equilibrium markov processes. In *Theory and application of random fields*, pages 233–248. Springer, 1983.
- [39] Ran Wei. On the long-range directed polymer model. *Journal of Statistical Physics*, 165(2):320–350, 2016.
- [40] Gian-Carlo Wick. The evaluation of the collision matrix. *Physical review*, 80(2):268, 1950.

SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY, WUHAN 430072, CHINA; COMPUTATIONAL SCIENCE HUBEI KEY LABORATORY, WUHAN UNIVERSITY, WUHAN, 430072, CHINA

Email address: glrang.math@whu.edu.cn

RESEARCH CENTER FOR MATHEMATICS AND INTERDISCIPLINARY SCIENCES, SHANDONG UNIVERSITY, QINGDAO 266237, CHINA; SCHOOL OF MATHEMATICS, SHANDONG UNIVERSITY, JINAN 250100, SHANDONG, CHINA

Email address: txjsong@sdu.edu.cn

SCHOOL OF MATHEMATICS, SHANDONG UNIVERSITY, JINAN, SHANDONG, 250100, CHINA

Email address: wangmeng22@mail.sdu.edu.cn