

Bernstein-Sato Polynomials of Semi-weighted-homogeneous Polynomials of Brieskorn-Pham Type

Morihiro Saito

Abstract. Let f be a semi-weighted-homogeneous polynomial having an isolated singularity at the origin. Let $\alpha_{f,k}$ ($k \in [1, \mu_f]$) be the spectral numbers of f with μ_f the Milnor number. As a consequence of theorems of Malgrange and Varchenko, there are non-negative integers r_k such that the $\alpha_{f,k} - r_k$ are the roots of the local Bernstein-Sato polynomial divided by $s+1$ up to sign, that is, the roots are given up to sign by spectral numbers shifted by non-negative integers. However, it is not easy in general to determine explicitly these shifts on the parameter space of miniversal μ -constant deformation of a weighted homogeneous polynomial. Assuming that the latter is a Brieskorn-Pham type polynomial, we can obtain a surprisingly simple algorithm to determine these shifts, which can be realized by using C (without employing Gröbner bases). This implies a refinement of classical calculations by Kato and Cassou-Noguès in two variable cases, showing that the stratification of the parameter space of the μ -constant deformation can be controlled by using the (partial) additive semigroup structure of the weights of parameters. As a corollary we can produce examples where one root of the Bernstein-Sato polynomial is quite distant from the others, and also an example of a semi-homogeneous polynomial with roots of its local Bernstein-Sato polynomial nonconsecutive.

Introduction

Let $f \in \mathbb{C}\{x\}$ be a convergent power series of n variables having an *isolated* singularity at 0, where $f(0) = 0$ and $n \geq 2$. Set $(X, 0) := (\mathbb{C}^n, 0)$. Let $\alpha_{f,k}$ ($k \in [1, \mu_f]$) be the *spectral numbers* of f counted with multiplicities, where μ_f is the Milnor number, see [St 77b] (and also [DiSa 14]). The local Bernstein-Sato polynomial $b_f(s)$ (see [Be 72], [SaSh 72], [Bj 73], [Sat 75], [Kas 76]) is called the *BS polynomial* for short in this paper. This is a factor of the *global* BS polynomial in the f polynomial case (using finite determinacy of holomorphic functions with isolated singularities), since f may have other singularities. As a consequence of theorems of Malgrange [Ma 75] and Varchenko [Va 81], it is quite well known that there are *non-negative integers* r_k ($k \in [1, \mu_f]$) such that

$$(1) \quad \tilde{R}_f = \{\alpha_k - r_k\}_k \quad \text{with} \quad \min\{\alpha_{f,k}\}_k = \min\{\alpha_{f,k} - r_k\}_k,$$

forgetting the multiplicities, where $\tilde{R}_f \subset \mathbb{R}_{>0}$ is the set of roots of the *reduced* BS polynomial $\tilde{b}_f(s) := b_f(s)/(s+1)$ up to sign. We say that $\alpha_{f,k} - r_k$ is a *shifted* root up to sign of the reduced BS polynomial $\tilde{b}_f(s)$ if $r_k \geq 1$. It is not easy to determine these shifts r_k on the base space of the miniversal μ -constant deformation of a weighted homogeneous polynomial. This is a quite interesting problem, see for instance [Kat 81], [Kat 82], [Ca 87], [NaTa 21]. (Recall that the spectral numbers are invariant by μ -constant deformations, and coincide with the roots of $\tilde{b}_f(s)$ up to sign in the weighted homogeneous case.)

The above assertion (1) can be verified by defining the *saturated Hodge filtration* \tilde{F} on λ -eigenspaces of the *vanishing cohomology* $H^{n-1}(F_f, \mathbb{C})_\lambda$ (with F_f the Milnor fiber), where the *Brieskorn lattice* $\mathcal{H}_f'' := \Omega_{X,0}^n / df \wedge d\Omega_{X,0}^{n-2}$ (see [Br 70]) is replaced by its *saturation* $\tilde{\mathcal{H}}_f'' := \sum_{i \geq 0} (\partial_t)^i \mathcal{H}_f''$ (see [Ma 75]) in the formula for the Hodge filtration F in [ScSt 85, (4.5)] (or [Sa 89, (2.6.3)] or (2.1.8) below) if we consider the meaning of the bigraded pieces $\text{Gr}_{\tilde{F}}^{p+r} \text{Gr}_F^p H^{n-1}(F_f, \mathbb{C})_\lambda$, since r corresponds to the *shift*. Here $H^{n-1}(F_f, \mathbb{C})_\lambda$ is identified with $\text{Gr}_V^\alpha \mathcal{G}_f$ for $\lambda = e^{-2\pi i \alpha}$ with $\mathcal{G}_f := \tilde{\mathcal{H}}_f''[\partial_t]$ the localization of $\tilde{\mathcal{H}}_f''$ by the action of ∂_t^{-1} , which is called the *Gauss-Manin system* (see for instance [Sa 89]), and V denotes the filtration of

Kashiwara and Malgrange on the regular holonomic $\mathcal{D}_{\mathbb{C},0}$ -module \mathcal{G}_f indexed by \mathbb{Q} . (This filtration was originally indexed by \mathbb{Z} , see also [Sa 83], [Sa 84, §3.4] about the reason for which V must be indexed by \mathbb{Q} .)

We assume in the introduction the monodromy T is *semi-simple*, that is, $N := \log T_u = 0$ with $T = T_s T_u$ the Jordan decomposition. Let ω_k ($k \in [1, \mu_f]$) be free generators of the Brieskorn lattice \mathcal{H}_f'' over $\mathbb{C}\{\{\partial_t^{-1}\}\}$ (see for instance [Sa 89] for $\mathbb{C}\{\{\partial_t^{-1}\}\}$). We assume that the ω_k give a \mathbb{C} -basis of $\Omega_f^n := \Omega_{X,0}^n / df \wedge \Omega_{X,0}^{n-1} = \mathcal{H}_f'' / \partial_t^{-1} \mathcal{H}_f''$ in a compatible way with the V -filtration (inducing a \mathbb{C} -basis of $\text{Gr}_V^\bullet \Omega_f^n$); for instance, the ω_k are associated with an *opposite filtration* in the sense of [Sa 89]. Using the *semi-simplicity* of T , we can define

$$\mathcal{G}_f^{(\alpha)} := \text{Ker}(\partial_t t - \alpha) \subset \mathcal{G}_f \quad (\alpha \in \mathbb{Q}).$$

so that we can get the *asymptotic expansions*

$$(2) \quad \omega_k = \sum_{\alpha \geq \alpha_{f,k}} \omega_k^{(\alpha)} \quad \text{with} \quad \omega_k^{(\alpha)} \in \mathcal{G}_f^{(\alpha)} \quad (k \in [1, \mu_f]),$$

and $v_k := \omega_k^{(\alpha_{f,k})} \neq 0$ (renumbering the ω_k if necessary). The v_k form a free basis of \mathcal{G}_f over $\mathbb{C}\{\{\partial_t^{-1}\}\}[\partial_t]$, and we have the *power series expansions*

$$(3) \quad \omega_k = \sum_{k'=1}^{\mu_f} g_{k,k'} v_{k'} \quad \text{with} \quad g_{k,k'} \in \mathbb{C}\{\{\partial_t^{-1}\}\}[\partial_t].$$

Assume furthermore the ω_k give a \mathbb{C} -basis of $\text{Gr}_{\tilde{F}}^\bullet \text{Gr}_F^\bullet H^{n-1}(F_f, \mathbb{C})_\lambda$ (inducing a *bisplitting* of F, \tilde{F}) using the above identification of $H^{n-1}(F_f, \mathbb{C})_\lambda$ with $\text{Gr}_V^\alpha \mathcal{G}_f$. This condition is not necessarily satisfied in the semi-weighted-homogeneous case if we consider only *monomial bases*. It is, however, trivially satisfied if the following condition holds:

$$(4) \quad \dim H^{n-1}(F_f, \mathbb{C})_\lambda \leq 1 \quad (\forall \lambda),$$

for instance, if the exponents of Brieskorn-Pham type polynomials are *mutually prime*. (In general, it is not easy to find free generators ω_k satisfying the above condition related to $\text{Gr}_{\tilde{F}}^\bullet \text{Gr}_F^\bullet$ before determining the saturation $\widetilde{\mathcal{H}}_f''$. So this approach is not good for explicit calculations of $\widetilde{\mathcal{H}}_f''$ unless condition (4) is satisfied at least partially.)

We can easily verify the following.

Proposition 1. *Under the above assumptions, the saturation $\widetilde{\mathcal{H}}_f''$ is generated over $\mathbb{C}\{\{\partial_t^{-1}\}\}$ by the $\omega_k^{(\alpha)}$ for $k \in [1, \mu_f]$, $\alpha \in [\alpha_{f,k}, n - \tilde{\alpha}_f]$, where $\tilde{\alpha}_f$ is the minimal spectral number.*

Combining this with the expansions in (3), we get the following.

Corollary 1. *With the above notation and assumptions, let $m_{k,k'}$ be the pole order of $g_{k,k'}$ as a power series of ∂_t^{-1} having a pole. Set $r_{k'} := \max\{m_{k,k'}\}_{k \in [1, \mu_f]}$. Then the $\alpha_{f,k} - r_k$ ($k \in [1, \mu_f]$) are the roots of $b_f(s)/(s+1)$ up to sign forgetting the multiplicities.*

Assume f is a *semi-weighted-homogeneous* polynomial $\sum_{\beta \geq 1} f_\beta$, where the f_β are weighted homogeneous polynomials of weighted degree β , which vanish except for a finite number of β . (Recall that convergent power series with isolated singularities have *finite determinacy*, see for instance [GLS 07]). We assume that the lowest weighted degree part f_1 is a *Brieskorn-Pham type* polynomial $\sum_{i=1}^n x_i^{e_i}$ with $e_i \geq 3$; in particular, f has an isolated singularity at 0, and the weight w_i of x_i is $1/e_i$.

It is well known (see [St 77a], [Va 82a]) that the *spectral numbers* of f are given by

$$(5) \quad \Sigma_f = \{ \alpha_{\mathbf{w}}(\nu) \mid \nu = (\nu_i) \in E_f \} \quad \text{with} \\ \alpha_{\mathbf{w}}(\nu) := \sum_{i=1}^n \nu_i / e_i, \quad E_f := \prod_{i=1}^n \mathbb{Z} \cap [1, e_i - 1].$$

Set $\omega^\nu := [x^{\nu-1}dx] \in \mathcal{H}_f''$ for $\nu \in \mathbb{Z}_{>0}^n$. Here $dx := dx_1 \wedge \cdots \wedge dx_n$ and $\mathbf{1} := (1, \dots, 1)$ (with $\mathbf{w} = (w_1, \dots, w_n)$). It is easy to see that the ω^ν for $\nu \in E_f$ form a free basis of \mathcal{H}_f'' and $\alpha_{\mathbf{w}}(\nu) = \alpha_V(\omega^\nu)$ ($\nu \in E_f$) using an argument similar to [Sa 88], where

$$\alpha_V(\omega) := \max\{\alpha \in \mathbb{Q} \mid \omega \in V^\alpha \mathcal{G}_f\} \quad (\omega \in \mathcal{G}_f).$$

We now assume that the exponents e_i are *mutually prime* so that condition (4) is satisfied. Let h_j ($j \in J$) be the monomials such that h_j is not contained in the Jacobian ideal (∂f_1) and moreover $\alpha_{\mathbf{w}}(h_j) > 1$. Here $\alpha_{\mathbf{w}}(h_j) := \alpha_{\mathbf{w}}(\nu)$ if $h_j = x^\nu$. So $\alpha_{\mathbf{w}}(h_j) = \alpha_V([h_j dx]) - \tilde{\alpha}_f$ with $\tilde{\alpha}_f := \alpha_V([dx]) = \alpha_{\mathbf{w}}(\mathbf{1})$. Note that the first condition is equivalent to that $\nu + \mathbf{1} \in E_f$, and the second may be replaced by $\alpha_{\mathbf{w}}(h_j) \geq 1$ assuming the first (using the condition on the e_i). By [Va 82b] we have $|J| = m_{f_1}$, the *modality* of f_1 . We may assume that f is written as

$$(6) \quad f = f_1 + \sum_{j \in J} u_j h_j \quad (u_j \in \mathbb{C}).$$

Here the u_j are identified with the coordinates of the parameter space of the miniversal μ -constant deformation of f_1 , and have weights $\gamma_j := \alpha_{\mathbf{w}}(h_j) - 1$ for $j \in J$ (up to sign). We can then calculate the Gauss-Manin connection rather explicitly, and get the following.

Theorem 1. *For semi-weighted-homogeneous polynomials of Brieskorn-Pham type, there is an efficient algorithm (without using Gröbner bases) for calculating the theoretically lowest coefficients of the $g_{k,k'} \in \mathbb{C}\{\{\partial_t^{-1}\}\}[\partial_t]$ in (3).*

In the two variable case (or more precisely, if $\tilde{\alpha}_f \geq (n-2)/2$), we have either $r_\nu = 1$ or 0 ($\forall \nu \in E_f$). So it is enough to calculate the coefficients $g_{k,k'}^{(1)}$ of ∂_t^1 in the $g_{k,k'} \in \mathbb{C}\{\{\partial_t^{-1}\}\}[\partial_t]$. We see that the $g_{k,k'}^{(1)}$ are *weighted homogenous* polynomials in the parameters u_j of the μ -constant deformation, and their *weighted degrees* are given by

$$(7) \quad \deg_{\mathbf{w}} g_{k,k'}^{(1)} = \deg_{\mathbf{w}} \omega_{k'} - \deg_{\mathbf{w}} \omega_k - 1,$$

with $\deg_{\mathbf{w}} u_j = \gamma_j$. Note that $\deg_{\mathbf{w}} g_{k,k'}^{(1)} = \gamma_j (= \alpha_{\mathbf{w}}(h_j) - 1)$ if $\omega_k = [dx]$, $\omega_{k'} = [h_j dx]$. It not difficult to realize this algorithm using Singular [DGPS 20] (or even C). The computation of *necessary terms* takes a few seconds if the (usual) polynomial degree of f_1 is at most 8 in the two variable case. We have, however, a problem of *integer overflow* even for $f_1 = x^9 + y^7$.

It is well known that there is a stratification of the parameter space of the miniversal μ -constant deformation of f_1 such that the BS polynomial is constant on each stratum. This stratification can be describes as follows: We identify E_f with $\mathbb{Z} \cap [1, \mu_k]$ in such a way that the spectral numbers $\alpha_{f,k}$ ($k \in [1, \mu_k]$) are *strictly increasing*; in particular $\mathbf{1} \in \mathbb{N}^n$ corresponds to $1 \in [1, \mu_k]$. We note $\nu' \succ \nu$ when $\nu'_i \geq \nu_i$ ($\forall i$), and similarly for \prec . For $j \in J$, there is a unique $\nu^{(j)} \in E_f$ with $x^{\nu^{(j)}-1} = h_j$. For $j, k \in J$, we note $j \succ k$ when $\nu^{(j)} \succ \nu^{(k)}$ (similarly for \prec). If $K \subset J$, set

$$K^{\succ j} := \{k \in K \mid k \succ j\} \quad (\text{similarly for } K^{\prec j}).$$

We say that $j \in K \subset J$ is *minimal* if $K^{\prec j} = \{j\}$. Let $\text{SG}(K) \subset \mathbb{Q}_{>0}$ be the *semigroup generated additively* by the $\gamma_k \in \mathbb{Q}_{>0}$ ($k \in K$). Set

$$V_K := \{u = (u_j) \in \mathbb{C}^J \mid u_j = 0 \ (j \notin K)\}.$$

Put $r_{(j)} := r_{\nu^{(j)}}$ ($j \in J$). Using (7) together with *non-vanishing* of certain coefficients, the following is easily verified:

Proposition 2. *Let $j \in K \subset J$. Then*

- (i) *We have $r_{(j')} = 1$ for any $j' \in J^{\succ j}$ if $\gamma_j \notin \text{SG}(K \setminus \{j\})$ and $u \in V_K$ with $u_j \neq 0$.*
- (ii) *We have $r_{(j')} = 1$ for any $j' \in J^{\succ j}$ if $\gamma_j \in \text{SG}(K \setminus \{j\})$ and u_k ($k \in K \setminus \{j\}$) are sufficiently general with $u_{k'}$ ($k' \notin K \setminus \{j\}$) fixed (even if $j' \notin K$ and $u_{j'} = 0$).*
- (iii) *We have $r_{(j)} = 0$ for some $u \in V_K$ with $u_j \neq 0$ if $\gamma_j \in \text{SG}(K \setminus \{j\})$ and $j \in K$ is minimal.*

Here “sufficiently general” means that it is contained in a non-empty Zariski-open subset. Proposition 2 shows the importance of the (partial) additive *semigroup structure* of the weights γ_j ($j \in J$) for the determination of the $r_{(j)}$. It gives the first *affine stratification* of the parameter space of the μ -constant deformation of f_1 with coordinates u_j ($j \in J$). Its strata correspond to *bistable* subsets $K \subset J$ (where K may be J or \emptyset). Here a subset $K \subset J$ is called *bistable* if the following two conditions are satisfied:

- (a) If $j \in J$, $k \in K$, and $j \succ k$, then $j \in K$.
- (b) If $j \in J$ and $\gamma_j \in \text{SG}(K)$, then $j \in K$.

These are called respectively the *upper* and *semigroup stability* conditions. The closure of the stratum corresponding to K is given by V_K , and we delete the closed subspaces corresponding to bistable proper subsets of K , that is, the stratum corresponding K is given by

$$V_K^\circ := V_K \setminus \bigcup_{K'} V_{K'},$$

where K' runs over bistable proper subsets of K . Note that bistable subsets are stable by intersections, and the closure of the corresponding stratum is compatible with intersections.

Remark 1. We can determine the bistable subsets K of J by *decreasing* induction on $|K|$. We first determine those with $|K| = |J| - 1$ by deleting each element from J and verifying the two conditions of bistability, where the obtained bistable subsets are ordered increasingly using the identification $J = \{1, \dots, m_{f_1}\}$ such that the γ_j are increasing. We say that an element is *removable* if its complement is bistable. There is at least one removable element, since the two conditions are satisfied by deleting j with γ_j minimal. We apply the same to the obtained bistable subsets. If we get an already obtained bistable subset, it is of course neglected. We can then proceed by decreasing induction on $|K|$. Note that for any proper bistable subset K' of a bistable subset K , there is a removable element of K not contained in K' . (Take an element $j \in K \setminus K'$ with γ_j minimal.) This implies that the strata of the first stratification are affine varieties.

If $K \subset J$ is a *bistable* subset, we have at a sufficiently general point of V_K

$$(8) \quad r_{(j)} = 1 \iff j \in K.$$

On some locally closed subspace of V_K , however, the equivalence (8) can hold only after replacing K with a suitable subset of K . We thus have to consider a further stratification of each stratum of the first affine stratification, depending on the (partial) additive *semigroup structure* of the corresponding bistable subset $K \subset J$. In simple cases as in [Kat 81], [Kat 82], [Ca 87], where the semigroup structure is not quite complicated, the stratification of the parameter space of the μ -constant deformation can be described completely by using the bistable subsets of J , see 1.1–2 below. (The relation to the stratification by Tjurina numbers does not seem very clear.)

We say that a root of a BS polynomial of a weighted homogeneous polynomial f_1 with an isolated singularity is *shiftable* if it is shifted by a μ -constant deformation, which is given by a semi-weighted-homogeneous polynomial f , see [Va 82b]. It is well known (and easy to show) that this condition is equivalent to that the root up to sign is strictly greater than $\tilde{\alpha}_f + 1$. We denote by $R_{f_1}^{\text{sh}} (\cong J)$ the set of *shiftable roots* up to sign of $b_{f_1}(s)$. (Note that $R_{f_1}^{\text{sh}} \subset \tilde{R}_{f_1}$, since 1 is unshiftable.) We say that $\alpha \in R_{f_1}^{\text{sh}}$ is a *unique unshifted shiftable root* up to sign of $b_f(s)$ if $\alpha \in \tilde{R}_f$ (that is, unshifted) and any $\alpha' \in R_{f_1}^{\text{sh}} \setminus \{\alpha\}$ does not belong to \tilde{R}_f (that is, shifted). This phenomenon is often seen if an unshifted root up to sign is close to $\tilde{\alpha}_f + 1$, for instance if the root is associated with a removable element of J . If α is a unique unshifted shiftable root up to sign of $b_f(s)$, we set $\text{SI}(f, \alpha) := |R_{f_1}^{\text{sh}, < \alpha}| / |R_{f_1}^{\text{sh}}|$, called the *solitude index*, where $R_{f_1}^{\text{sh}, < \alpha} = R_{f_1}^{\text{sh}} \cap (0, \alpha)$. As this index becomes large, we have more complexity of the

defining equations of the subspace on which α is the unique unshifted shiftable root up to sign of the BS polynomial. We can show an example with $\text{SI}(f, \alpha) = \frac{9}{16} > \frac{1}{2}$ after a computer calculation using C, where $n = 3$, see 1.5 below.

Note finally that a similar argument can be applied to the homogeneous case. We can indeed construct an example of a semi-homogeneous polynomial such that the roots of its local BS polynomial are *nonconsecutive* (that is, not given by the intersection of a connected interval in \mathbb{R} and $\frac{1}{d}\mathbb{Z}$ with d the degree of the lowest homogeneous part of f), for instance, $f = \frac{1}{10}x^{10} + \frac{1}{10}y^{10} + x^3y^8 + x^8y^3 + 8x^6y^6 - 128x^7y^7$, where $\tilde{R}_f = \{\frac{2}{10}, \dots, \frac{14}{10}\} \cup \{\frac{16}{10}\}$ according to Singular [DGPS 20], see 1.6 below. (Note that $\tilde{R}_f \subset \frac{1}{d}\mathbb{Z} \cap (0, n)$ if f is semi-homogeneous and d is the degree of the lowest homogeneous part which has an isolated singularity.) It does not seem easy to find an example as above with $d \leq 9$, $n = 2$. This situation is quite different from the case of homogeneous polynomials with non-isolated singularities, where the roots supported at the origin are consecutive as far as calculated, see for instance [Sa 16], [Sa 20].

In Section 1 we explain some examples. In Section 2 we describe the algorithm after reviewing some basics of Brieskorn lattices.

This work was partially supported by JSPS Kakenhi 15K04816.

1. Examples

In this section we explain some examples.

1.1. Example I. Let $f_1 = x^7 + y^5$ or $x^9 + y^4$, see [Kat 81], [Kat 82]. The weights γ_j ($j \in J$) multiplied by 35 or 36 and the exponents of the corresponding monomials are as follows:

$$(1.1.1) \quad \begin{array}{cccccc} 1 & 6 & 11 & & 3, 3 & 4, 3 & 5, 3 \\ & & 4 & & & & 5, 2 \end{array} \quad \text{or} \quad \begin{array}{cccccc} 2 & 6 & 10 & & 5, 2 & 6, 2 & 7, 2 \\ & & 1 & & & & 7, 1 \end{array}$$

We see that the number of strata of the first affine stratification is 6 or 5, where the nonempty bistable subsets are as below:

$$(1.1.2) \quad \begin{array}{cccccc} 1 & 6 & 11 & & 6 & 11 & & 6 & 11 & & 11 & & 11 \\ & & 4 & & & 4 & & & & & 4 & & & \end{array}$$

or

$$(1.1.3) \quad \begin{array}{cccccc} 2 & 6 & 10 & & 2 & 6 & 10 & & 6 & 10 & & 10 \\ & & 1 & & & & & & & & & & \end{array}$$

The open stratum of the first affine stratification corresponding to $K = J$ contains one or two subspaces on which the assertion (8) holds with J replaced by $J \setminus \{2\}$ or $J \setminus \{2\}$ and $J \setminus \{3\}$, that is,

$$(1.1.4) \quad \begin{array}{cccccc} 1 & 6 & 11 & & 6 & 10 & & 2 & 10 \\ & & & & & 1 & & & 1 \end{array} \quad \text{or} \quad \begin{array}{cccccc} 6 & 10 & & 2 & 10 \\ & 1 & & & 1 \end{array} \quad \text{and} \quad \begin{array}{cccccc} 2 & 10 \\ & 1 \end{array}$$

using Proposition 2. The corresponding subspaces are as follows:

$$(1.1.5) \quad \{u_2 = cu_1^4\} \quad \text{or} \quad \{u_2 = c'u_1^2\} \quad \text{and} \quad \{u_2 = c''u_1^2, u_3 = c'''u_1^6\}.$$

These follow from (7) looking at the weights γ_j and (1.1.1). Note that $\gamma_2 = 4\gamma_1$ or $\gamma_2 = 2\gamma_1$ and $\gamma_3 = 6\gamma_1$. Here we identify J with $\{1, \dots, m_{f_1}\}$ so that the γ_j are *strictly increasing*, where m_{f_1} denotes the *modality* of f_1 , which coincides with the *inner modality*, see [NaTa 21], [Va 82b]. (We have to add the minimal spectral number $\frac{12}{35}$ or $\frac{13}{36}$ plus one to the weight γ_j in order to get the spectral number corresponding to γ_j , see a remark just after (7).) It is not necessarily easy to determine the constants $c, c', c'', c''' \in \mathbb{C}^*$. Using a computer, we can get that

$$(1.1.6) \quad c = \frac{6}{175}, \quad c' = \frac{1}{3}, \quad c'' = \frac{7}{18}, \quad c''' = \frac{2429}{1259712}.$$

To determine c''' , we use the vanishing of $g_{2,23}^{(1)} = -u_2 + \frac{7}{18}u_1^2$ (which gives c'') and that of

$$(1.1.7) \quad g_{1,23}^{(1)} = -u_3 - \frac{7}{72}u_2^3 + \frac{7}{27}u_1^2u_2^2 - \frac{175}{1458}u_1^4u_2 + \frac{595}{39366}u_1^6,$$

where $23 = \mu_f - 1$. (Notice that the *sign* depends only on the degrees of monomials. This holds in general applying repeatedly (2.2.1) below.) For c' , we need the vanishing of $g_{1,22}^{(1)} = -u_2 + \frac{1}{3}u_1^2$. (For c with $f_1 = x^7 + y^5$, we use the vanishing of $g_{1,22}^{(1)} = -u_2 + \frac{6}{175}u_1^4$ with $22 = \mu_f - 2$.)

For a μ -constant deformation f of $f_1 = x^9 + y^4$ having a unique unshifted shiftable root $\frac{55}{36}$ up to sign of $\tilde{b}_f(s)$, the distribution of roots up to sign is as below:

$$\bullet \bullet \bullet \bullet \quad \bullet \bullet \bullet \bullet \quad \bullet \quad \bullet \bullet \quad \circ \circ \quad \bullet \quad \circ$$

$$\frac{50}{36} \quad \frac{55}{36} \quad \frac{59}{36}$$

Here the black and white vertices represent respectively the roots of $b_f(s)$ up to sign and those of $b_{f_1}(s)$ which are not roots of $b_f(s)$ up to sign (that is, shifted). We have $\text{SI}(f, \frac{55}{36}) = \frac{1}{2}$.

1.2. Example II. Let $f_1 = x^7 + y^6$. The weights of parameters multiplied by 42 and the exponents of the corresponding monomials are as follows:

$$(1.2.1) \quad \begin{array}{ccccccc} 4 & 10 & 16 & & 3, 4 & 4, 4 & 5, 4 \\ & & 3 & 9 & & 4, 3 & 5, 3 \\ & & & 2 & & & 5, 2 \end{array}$$

Following Remark 1 in the introduction, we get ten nonempty bistable subsets of J as below:

$$(1.2.2) \quad \begin{array}{ccccc} 4 & 10 & 16 & 4 & 10 & 16 & 4 & 10 & 16 & 4 & 10 & 16 & 10 & 16 \\ & 3 & 9 & & 3 & 9 & & 9 & & 9 & & 9 & 3 & 9 \\ & & 2 & & & & & 2 & & & & & & \\ 10 & 16 & & 4 & 10 & 16 & & 10 & 16 & & 16 & & & 16 \\ & 9 & & & & & & & & & 9 & & & 9 \end{array}$$

The first and third subsets have respectively one and two subsets as below, which are not bistable, but correspond to the sets of *shifted* roots up to sign of BS polynomial $b_f(s)$ by adding 13 and dividing it by 42:

$$(1.2.3) \quad \begin{array}{ccc} 10 & 16 & \\ 3 & 9 & \\ 2 & & \end{array} \quad \text{and} \quad \begin{array}{ccc} 10 & 16 & 4 & 16 \\ 9 & & 9 & 9 \\ 2 & & 2 & 2 \end{array}$$

These are determined by using the (partial) *semigroup structure* of K (more precisely, $4 = 2 \cdot 2$ and $10 = 5 \cdot 2 = 4 + 3 \cdot 2 = 2 \cdot 4 + 2$). The corresponding subspaces are respectively as follows:

$$(1.2.4) \quad \{u_3 = cu_1^2\} \quad \text{and} \quad \{u_2 = 0, u_3 = c'u_1^2\}, \quad \{u_2 = 0, u_3 = c''u_1^2, u_5 = c'''u_1^5\}.$$

The first subset of (1.2.2) (that is, J) has only one subset, since $5 \in J$ is *over* $2 \in J$ with $\gamma_5 = \frac{10}{42}$ and $\gamma_2 = \frac{3}{42} \notin \text{SG}(J \setminus \{2\})$. We have

$$(1.2.5) \quad c = c' = \frac{2}{7}, \quad c'' = \frac{5}{14}, \quad c''' = -\frac{5}{16464}.$$

It is not necessarily easy to determine c''' . (There is another method used in [Sa 22b].) Here we calculate it by combining the vanishing of $g_{1,38}^{(1)} = -u_2$, $g_{2,41}^{(1)} = -u_3 + \frac{5}{14}u_1^2$ (which gives c''), and that of

$$(1.2.6) \quad g_{1,41}^{(1)} = -u_5 - \frac{25}{84}u_2^2u_3 - \frac{25}{84}u_1u_3^2 + \frac{25}{98}u_1^2u_2^2 + \frac{25}{147}u_1^3u_3 - \frac{95}{4116}u_1^5,$$

looking at [1;7,7], [5;7,7] in the output of the code in 2.6 below and using a variation of Proposition 2, where we assume $u_1 u_6 \neq 0$. For $u_1 = u_6 = 1$, we then obtain that $u_9 = 8$, $u_{17} = -128$ as is written at the end of the introduction. By a similar argument, we can see, assuming $u_1 u_6 \neq 0$, that the root $\frac{15}{10}$ up to sign is shifted if

$$u_9 \neq 4u_1^2 + \frac{7}{2}u_6^2, \quad u_9 \neq \frac{7}{2}u_1^2 + 4u_6^2,$$

looking at [2;6,7], [3;7,6] in the output of the code in 2.6 below. It is easy to see that the roots $\frac{13}{10}$, $\frac{14}{10}$ up to sign are unshifted. Note that the argument is rather different from the weighted homogeneous case, since condition (4) is unsatisfied.

2. Explanation of the algorithm

In this section we describe the algorithm after reviewing some basics of Brieskorn lattices.

2.1. Brieskorn lattices. Let $f \in \mathbb{C}\{x\}$ be a convergent power series of n variables having an isolated singularity at 0 with $f(0) = 0$ and $n \geq 2$. Set $(X, 0) := (\mathbb{C}^n, 0)$. The Brieskorn lattice \mathcal{H}_f'' (see [Br 70]) is defined by

$$(2.1.1) \quad \mathcal{H}_f'' := \Omega_{X,0}^n / df \wedge d\Omega_{X,0}^{n-2}.$$

This is a free module of rank μ_f (with μ_f the Milnor number of f) over $\mathbb{C}\{t\}$ and also over $\mathbb{C}\{\{\partial_t^{-1}\}\}$, see for instance [Sa 89]. We can define the actions of t and ∂_t^{-1} respectively by multiplication by f and

$$(2.1.2) \quad \partial_t^{-1}[\omega] = [df \wedge \eta] \quad \text{if} \quad d\eta = \omega \quad \text{for} \quad \omega \in \Omega_{X,0}^n, \eta \in \Omega_{X,0}^{n-1}.$$

The *Gauss-Manin system* can be defined by

$$\mathcal{G}_f := \mathcal{H}_f''[\partial_t],$$

which is the localization of \mathcal{H}_f'' by the action of ∂_t^{-1} . This is a regular holonomic $\mathcal{D}_{X,0}$ -module with quasi-unipotent monodromy, and has the V -filtration of Kashiwara and Malgrange indexed by \mathbb{Q} so that $\partial_t t - \alpha$ is nilpotent on the graded quotients $\text{Gr}_V^\alpha \mathcal{G}_f$ ($\mathbb{Q} \in \mathbb{Q}$). The latter is identified with

$$(2.1.3) \quad \mathcal{G}_f^{(\alpha)} := \text{Ker}((\partial_t t - \alpha)^i : \mathcal{G}_f \rightarrow \mathcal{G}_f) \quad (i \gg 0),$$

and we have the inclusion (see for instance [Sa 89]):

$$(2.1.4) \quad \mathcal{G}_f \hookrightarrow \prod_{\alpha \in \mathbb{Q}} \mathcal{G}_f^{(\alpha)}.$$

This implies for $\omega \in \mathcal{G}_f$ the *asymptotic expansion*

$$(2.1.5) \quad \omega = \sum_{\alpha \in \mathbb{Q}} \omega^{(\alpha)} \quad \text{with} \quad \omega^{(\alpha)} \in \mathcal{G}_f^{(\alpha)} \quad (\alpha \in \mathbb{Q}).$$

Set

$$(2.1.6) \quad \alpha_V(\omega) := \min\{\alpha \in \mathbb{Q} \mid \omega^{(\alpha)} \neq 0\}, \quad \text{Gr}^V \omega := \omega^{(\alpha_V(\omega))}.$$

We have the canonical isomorphism

$$(2.1.7) \quad \text{Gr}_V^\alpha \mathcal{G}_f = H^{n-1}(F_f, \mathbb{C})_\lambda \quad (\lambda = e^{-2\pi i \alpha}),$$

with F_f the Milnor fiber, and moreover

$$(2.1.8) \quad \begin{aligned} \text{Gr}_V^\alpha \mathcal{H}_f'' &= F^{n-1-p} H^{n-1}(F_f, \mathbb{C})_\lambda \\ \text{for } \alpha &= \beta + p, \beta \in (0, 1], p \in \mathbb{Z}, \end{aligned}$$

see [ScSt 85, (4.5)], [Va 81] (and also [Sa 89, (2.6.3)]).

Remark 2.1a. In the *weighted homogeneous* case, the variable x_i has weight ω_i so that the weighted degree of f is 1, and the filtration V is induced by the filtration on $\Omega_{X,0}^n$

by the *weighted degree*, where the weight of dx_i is w_i . Indeed, we have the Euler field $\xi = \sum_{i=1}^n w_i x_i \partial_{x_i}$ such that $\xi(f) = f$, and

$$(2.1.9) \quad d(\iota_\xi \omega) = L_\xi \omega, \quad df \wedge \iota_\xi \omega = f\omega \quad (\omega \in \Omega_{X,0}^n),$$

where L_ξ, ι_ξ denote respectively the *interior product* and the *Lie derivation* respectively, see also [Sa 22b, 1.1.7].

Remark 2.1b. Assume $f = f_1 + f_{>1}$ is a *semi-weighted-homogeneous* deformation of a weighted homogeneous polynomial f_1 having an isolated singularity at 0. Let V be the decreasing filtration on $\Omega_{X,0}^n$ defined by the condition that the weighted degree is at least α . This induces the V -filtration on the Gauss-Manin system \mathcal{G}_f , see for instance [Sa 22b]. Moreover we have the canonical isomorphism

$$(2.1.10) \quad \mathrm{Gr}_V^\bullet \mathcal{G}_f = \mathrm{Gr}_V^\bullet \mathcal{G}_{f_1}.$$

This can be shown for instance considering $\mathrm{Gr}_V^\bullet \Omega_{X,0}^n / \mathrm{Gr}_V^1 df \wedge d \mathrm{Gr}_V^\bullet \Omega_{X,0}^{n-2}$.

Remark 2.1c. Let $\alpha_{f,1}, \dots, \alpha_{f,\mu_f}$ be the *spectral numbers* of f , see [St 77b] (and also [DiSa 14], [JKSY 22]). It is well known (see for instance [ScSt 85], [Sa 89], [Va 81] and also (2.1.8) that we have the equality

$$(2.1.11) \quad \dim_{\mathbb{C}} \mathrm{Gr}_V^\alpha \Omega_f^n = \#\{k \in [1, \mu_f] \mid \alpha_{f,k} = \alpha\} \quad (\forall \alpha \in \mathbb{Q}),$$

with

$$(2.1.12) \quad \Omega_f^n := \Omega_{X,0}^n / df \wedge \Omega_{X,0}^{n-1} = \mathcal{H}_f'' / \partial_t^{-1} \mathcal{H}_f''.$$

Remark 2.1d. By [Ma 75], the *reduced* BS polynomial $b_f(s)/(s+1)$ is equal to the minimal polynomial of the action of $-\partial_t t$ on

$$\widetilde{\mathcal{H}}_f'' / t \widetilde{\mathcal{H}}_f''.$$

Here $t \widetilde{\mathcal{H}}_f''$ may be replaced by $\partial_t^{-1} \widetilde{\mathcal{H}}_f''$, see for instance [Sa 22b].

2.2. Proof of Theorem 1. We have $f = f_1 + \sum_{j \in J} u_j h_j$ as in the introduction, where the u_j are viewed as constants. For $\nu \in \mathbb{Z}_{>0}^n$, set $\omega^\nu := [x^{\nu-1} dx] \in \mathcal{H}_f''$. We can easily verify that

$$(2.2.1) \quad (\partial_t t - \alpha_w(\nu)) \omega^\nu = -\sum_{j \in J} \gamma_j u_j \partial_t \omega^{\nu + \nu^{(j)}},$$

using (2.1.9), where $h_j = x^{\nu^{(j)}}$, see also [Sa 22b]. Comparing the *asymptotic expansions* of both sides (using (2.1.10)), we can determine the theoretically lowest term of the asymptotic expansion of ω^ν by decreasing induction on $\alpha_w(\nu)$. Here what is important is the ratio of the weight of the variable in which one is interested and the minimal weight of the variables,

since this gives the maximal number of procedures to which one applies (2.2.1), see also the sample codes in 2.5-6 below for more details. This finishes the proof of Theorem 1.

2.3. Proof of Proposition 1. Since the maximal spectral number is $n - \tilde{\alpha}_f$, the assertion follows from the generalized Jordan decomposition, see for instance [Sa 22a, Rem. A.7c]. This finishes the proof of Proposition 1.

2.4. Proof of Proposition 2. This follows by applying (2.2.1) inductively. Note that the sign depends only on the total degree of the u_j . Here we use (2.1.10) in an essential way. This finishes the proof of Proposition 2.

Remark 2.4. The minimal root of $b_{f_1}(s)$ is not a root of $b_f(s)$ if $u_j \neq 0$ for some $j \in J$. Indeed, let $j_0 \in J$ with $u_{j_0} \neq 0$ and γ_{j_0} minimal. Let h_{j_0} be the corresponding monomial. Set $g := \prod_{i=1}^n x_i^{e_i-2}/h_{j_0}$. Consider the asymptotic expansion of $[gdx]$, and apply (2.2.1).

2.5. Sample code for Example III. For the convenience of the reader we note here a sample code for Singular to calculate Example III. Since it is written in a condensed way, it may be better to add line breaks appropriately after copying and pasting it in a text file. One may modify a,b as long as $a+b \leq 15$ and $(a,b)=1$. (If ‘‘Division Error’’ appears, one has to increase the size of the vector iv.) This code cannot be applied to $f_1 = x^9 + y^7$, although it works at least for $(a,b) = (7,6), (9,4), (7,5)$. (Please verify whether the list of weights in the last line is correct.)

```
ring R = 0, (u_1,u_2,u_3,u_4,u_5,u_6,u_7), ds;
int a,b,rs,num,i,j,wd,n,p,iq,ir,jq,jr,kk,e,li,lj,rs,maxp,MA,MB,NuM;a=8;b=7;
int k,q,od,di,ip,jp,wp,ie,je,we,pp,rp,MM,mxn,mxk,maxdiv,am,bm,m,mm;poly Sub;m=a*b;
vector iv=[1,1/2,1/3,1/4,1/5,1/6,1/7,1/8,1/9,1/10,1/11,1/12];intvec wt=0,0,0,0,0,0,0;
am=a-1;bm=b-1;mm=am*bm;if(a+b>14){rs=5;}else{rs=4;}mxn=7;MA=2000;MB=am*bm;NuM=1000;
intmat Co[mxn][2];intmat Nu[NuM][NuM];matrix M[MA][MB];intmat O[MA][MB];
vector va=[u_1,u_2,u_3,u_4,u_5,u_6,u_7];mxk=2*(m-a-b);p=1;for(k=m+1;k<mxk&&p<=mxn;k++)
{for(i=1;i<a-1&&k>b*i;i++){j=(k-b*i)div a;if((k-b*i)%a==0&&j<b-1){Co[p,1]=i;Co[p,2]=j;
wt[p]=k-m;p++;}}num=p-1;rp=wt[num]div wt[1];MM=wt[num]+m*rp;p=0;maxdiv=0;for(wd=MM;
wd>0;wd--){n=wd div b;for(i=0;i<n;i++){j=(wd-b*i)div a;if((wd-b*i)%a==0){p++;
Nu[i+1,j+1]=p;ir=i*a;iq=i div a;jr=j*b;jq=j div b;li=1;for(e=1;e<=iq;e++){li=li*(i-e*
a+1);}lj=1;for(e=1;e<=jq;e++){lj=lj*(j-e*b+1);}if(ir!=a-1&&jr!=b-1){M[p,ir+am*jr+1]=
li*lj*iv[a]^iq*iv[b]^jq;O[p,ir+am*jr+1]=iq+jq;}for(q=1;q<=num;q++){ip=i+Co[q,1];jp=j+
Co[q,2];wp=b*ip+a*jp;if(wp<=MM){pp=Nu[ip+1,jp+1];for(e=1;e<=mm;e++){ie=(e-1)%am;je=
(e-1)div am;we=b*ie+a*je;if(M[pp,e]!=0&&ie+je<a+b-rs&&we>m){od=0[pp,e];di=we+m*(od-1)
-wd;if(di<=wt[num]){Sub=M[pp,e]*wt[q]*iv[di]*va[q];if(O[p,e]==od-1){if(di>maxdiv)
{maxdiv=di;}M[p,e]=M[p,e]-Sub;}if(O[p,e]>od-1||O[p,e]<od-1&&M[p,e]==0){if(di>maxdiv)
{maxdiv=di;}M[p,e]=0-Sub;O[p,e]=od-1;}}}}}} maxp=p;if(size(iv)<maxdiv){sprintf(
"Division Error %s",maxdiv);} for(i=1;i<=maxp;i++){for(e=1;e<=mm;e++){ie=(e-1)%am;je=
(e-1)div am;we=b*ie+a*je;if(O[i,e]==-1&&ie+je<a+b-rs&&we>m){sprintf("[%s;%s,%s]:",
maxp+1-i,(e-1)%am,(e-1)div am);M[i,e];}} sprintf("wt=%s",wt);
```

In case one needs a computation for $\frac{1}{a}x^a + \frac{1}{b}y^b$ (instead of $x^a + y^b$) as in Example IV, one can do it by removing $*iv[a]^iq*iv[b]^jq$.

2.6. Sample code for Example IV. We note also a sample code for Example IV using C. (This is possible, since the algorithm is extremely simple.) One can copy and past it in a text file and compile it using gcc in Unix (or clang in Mac, etc.). In the first several lines, *line breaks* must be inserted before $\#$. Do not forget to remove page numbers, etc. (It may be necessary to replace $'$ with a *character from keyboard* if ‘‘Preview’’ in Mac is used.) When one runs $./a.out$, one is asked to type three numbers (and press the return key), which are 9 7 7 and 10 10 18 in the case of Examples IV and VI. The last number is the number of variables of the parameter space in which one is interested. (It is not assumed that the exponents are mutually prime. Note however that the code can calculate only the negative grading part of the parameter space of the miniversal μ -constant deformation if the exponents are not mutually prime.) One can type C before these numbers if one needs a calculation for $f_1 = x^a + y^b$ instead of $f_1 = \frac{1}{a}x^a + \frac{1}{b}y^b$. So one may enter C7 5 3, C9 4 3, C7 6 5, C8 7 7 for examples in 1.1–3.

```

#include<stdio.h> #include<stdlib.h> #define ME 11 #define MJ 18 #define MN 3000 #define MNM 30
#define PN 20 #define Num 5000 long long L0,L1,L2,L3;int P1[PN],P2[PN],P3[PN],PP[PN],AA[PN];
int BB[PN],DI[PN],Dia[MJ][MJ],Cn[ME][ME],J[MJ][3],JW[MJ][PN],Ad[MN][2],num,maxn,maxp,Ca,e;
int M[MN][MJ][MNM][PN+MJ],Ord[MN][MJ],Mprf[PN+MJ],N[MN][MJ],Nu[Num][NuM],Prm[PN],PRMS[10*PN];
long long L[MN][MJ][MNM];void prf(long long LLL);long long summ(void);long long pfac(long long
mmm);int main(void){int c,cc,i,j,k,l,n,nn,nnn,p,q,wd,rp,a,b,m,fl,mw,am,bm,mm,MM,gcd,ad,bd,iq;
ir,jr,jk,ok,od,ip,jp,wp,ie,je,we,pp,ee;long long li,lj,ll,lli;
for(i=2;i<10*PN;i++){PRMS[i]=1;}for(i=2;i<10*PN;i++){for(j=2;j<10*PN;j++){PRMS[i*j]=0;}}
for(i=j=0;i<10*PN&&j<PN;i++){if(PRMS[i]==1){Prm[j]=i;j++;}}if(j<PN){printf("Prime Error\n"),
exit(1);}printf("Exponents and the number of interesting monomials:");cc=getchar();if(cc=='C')
{Ca=1;}else{Ca=0;}while(cc<'0' || '9'<cc){cc=getchar();}for(e=0;'0'<=cc&&cc<='9';cc=getchar())
{e=e*10+cc-'0';}a=i=e;while(cc<'0' || '9'<cc){cc=getchar();}for(e=0;'0'<=cc&&cc<='9';cc=
getchar()){e=e*10+cc-'0';}b=j=e;while(cc<'0' || '9'<cc){cc=getchar();}for(e=0;'0'<=cc&&cc<='9';
cc=getchar()){e=e*10+cc-'0';}num=e;if(a>ME||b>ME){printf("Too big exponents\n");}if(Ca==1){
printf("C-mod. ");}printf("Exponents:%d,%d. Numbers of monomials:%d.\n",a,b,num);pfac(a);for
(i=0;i<PN;i++){AA[i]=PP[i];}pfac(b);for(i=0;i<PN;i++){BB[i]=PP[i];}for(i=0;i<PN;i++){if(AA[i]
<BB[i]){PP[i]=AA[i];}else{PP[i]=BB[i];}}for(i=0,gcd=1;i<PN;i++){for(j=0;j<PP[i];j++){gcd=gcd*
Prm[i];}ad=a/gcd;bd=b/gcd;m=ad*bd*gcd;for(i=0;i<MJ;i++){for(j=0;j<MJ;j++){Dia[i][j]=0;}}for(i=
0;i<Mj;i++){Dia[i][i]=1;}am=a-1;bm=b-1,mm=am*bm;mw=m-2*(ad+bd);for(i=0;i<ME;i++){for(j=0;j<ME;
j++){Cn[i][j]=0;}}for(p=0,k=m+1;k<2*m&&p<num;k++){for(i=1;i<a-1&&k<bd*i;i++){if((k-bd*i)%ad==0
&&(j=(k-bd*i)/ad)<b-1){J[p][0]=k-m;J[p][1]=i;J[p][2]=j;Cn[i][j]=p+1;lll=pfac(J[p][0]);if(lll!=1
){printf("Too large weight%d\n",J[p][0]),exit(1);}for(l=0;l<PN;l++){JW[p][l]=PP[l];}p++;}}if(
num>p||num>MJ){printf("num=%d is replaced by %d!\n\n",num,p);num=p;rp=J[num-1][0]/J[0][0];MM
=J[num-1][0]+m*rp;printf("Weights and exponents:\n");for(i=0;i<num;i++){printf("%d %d %d",
J[i][0],J[i][1],J[i][2]);}printf("\nMM=%d, rp=%d\n",MM,rp);Nu[0][0]=Nu[1][0]=Nu[0][2][0]
=Nu[1][1]=Nu[0][2]=-1;for(p=wd=0;wd<=MM;wd++){nnn=wd/bd;for(i=0;i<=nnn;i++){j=(wd-bd*i)/ad;if
((wd-bd*i)%ad==0&&Nu[i][j]<0){if(p>=MN){printf("Too many monomials to calculate\n"),exit(1);}
Nu[i][j]=p+1;Ad[p][0]=i;Ad[p][1]=j;for(k=0;k<num;k++){if(i+J[k][1]<Num&&j+J[k][2]<Num&&
Nu[i+J[k][1]][j+J[k][2]]<=0){Nu[i+J[k][1]][j+J[k][2]]=-1;}}p++;}}printf("%d monomials\n",p);
for(p--;p>=0;p--){i=Ad[p][0];j=Ad[p][1];wd=ad*j+bd*i;ir=i/a;iq=i/a;jr=j/b;jq=j/b;for(li=e=1;
e<=iq;e++){li=li*(1-e*a+1);}for(lj=e=1;e<=jq;e++){lj=1*j*(j-e*b+1);}ee=Cn[ir][jr]-1;num=PP[ee];
if(ir!=am&&jr!=bm&&ee>=0){L[p][ee][n]=pfac(li*lj);for(l=0;l<PN;l++){M[p][ee][n][l]=0-PP[l];}
if(Ca==1){for(l=0;l<PN;l++){M[p][ee][n][l]=M[p][ee][n][l]+AA[l]*iq+BB[l]*jq;}Ord[p][ee]=iq+jq;
N[p][ee]++;if(N[p][ee]>maxn)maxn=N[p][ee];if(N[p][ee]>=MNM){printf("Too many terms%d\n",N[p][ee]),
exit(1);}for(q=0;q<num;q++){ip=i+J[q][1];jp=j+J[q][2];wp=bd*ip+ad*jp;if(wp<=MM){pp=Nu[ip][jp]-
1;for(e=0;e<num;e++){nn=N[pp][e];ie=J[e][1];je=J[e][2];we=bd*ie+ad*je;od=Ord[pp][e];di=we+m*(od
-1)-wd;if(nn!=0&&di<=J[num-1][0]){ll=pfac(di);if(ll!=1){printf("Too large di=%d\n\n",di);}for(l
=0;l<PN;l++){DI[l]=PP[l];}if(Ord[p][e]==od-1&&N[p][e]!=0){for(cc=0;cc<nn;cc++){n=N[p][e];for
(fl=c=0;c<=n&&fl==0;c++){for(l=0;l<num&&M[p][e][c][PN+1]==M[pp][e][cc][PN+1]+Dia[q][l];l++){}
if(l==num){for(l=0;l<PN;l++){P1[l]=M[p][e][c][l];P2[l]=M[pp][e][cc][l]-JW[q][l]+DI[l];}L1=L0=
L[p][e][c];L2=0-L[pp][e][cc];L[p][e][c]=summ(c);for(l=0;l<PN;l++){M[p][e][c][l]=P3[l];}fl++;}}
if(fl==0){L[p][e][n]=0-L[pp][e][cc];for(l=0;l<PN;l++){M[p][e][n][l]=M[pp][e][cc][l]-JW[q][l]+
DI[l];}for(l=0;l<num;l++){M[p][e][n][l+PN]=M[pp][e][cc][l+PN]+Dia[q][l];}N[p][e]++;if(N[p][e]
>maxn)maxn=N[p][e];if(N[p][e]>=MNM){printf("Too many terms%d\n",N[p][e]),exit(1);}}if(Ord[p][e]
>od-1||N[p][e]==0){for(cc=0;cc<N[pp][e];cc++){L[p][e][cc]=0-L[pp][e][cc];for(l=0;l<PN;l++){
M[p][e][cc][l]=M[pp][e][cc][l]-JW[q][l]+DI[l];}for(l=0;l<num;l++){M[p][e][cc][l+PN]=
M[pp][e][cc][l+PN]+Dia[q][l];}N[p][e]=N[pp][e];Ord[p][e]=od-1;}}}}for(i=5;i>=0;i--){for(e=
0;e<num;e++){if(Ord[i][e]==-1){printf("%d;%d,%d):\n",i+1,J[e][1],J[e][2]);for(j=N[i][e]-1;j
>=0;j--){for(l=0;l<PN+num;l++){Mprf[l]=M[i][e][j][l];}prf(L[i][e][j]);}}}}
void prf(long long LLL){int j,k,l,fl;long long De;fl=0;for(De=1,l=0;l<PN;l++){if(Mprf[l]>0){for
(j=0;j<Mprf[l];j++){De=De*Prm[l];}}for(l=0;l<PN;l++){if(Mprf[l]<0){for(j=0;j<0-Mprf[l];
j++){LLL=LLL*Prm[l];}}if(LLL>0&&De>1){printf("+%lld/%lld",LLL,De);fl=1;}else if(LLL>1&&De==1)
{printf("+%lld",LLL);fl=1;}else if(LLL<0&&De>1){printf("%lld/%lld",LLL,De);fl=1;}else if(LLL<
-1&&De==1){printf("%lld",LLL);fl=1;}else if(LLL==1&&De==1){printf("-");}else if(LLL==1&&De==1)
{printf("+");}else if(LLL==0){printf("0!!!!");}for(l=0;l<num;l++){if(fl==0&&Mprf[l+PN]!=0){fl=
1;}else if(fl!=0&&Mprf[l+PN]!=0){printf("*");}if(Mprf[l+PN]>1){printf("u_%d^%d",l+1,Mprf[l+PN])
;}else if(Mprf[l+PN]==1){printf("u_%d",l+1);}}printf("\n");}long long summ(void){int i,j,k;for(
i=0;i<PN;i++){if(P1[i]>P2[i]){for(k=1,j=0;j<P1[i]-P2[i];j++){k=k*Prm[i];}L2=L2*k;P3[i]=P1[i];}
else if(P1[i]<P2[i]){for(k=1,j=0;j<P2[i]-P1[i];j++){k=k*Prm[i];}L1=L1*k;P3[i]=P2[i];}else{P3[i]
=P2[i];}L3=L1+L2;if((L1>0&&L2>0&&L3<0)||(L1<0&&L2<0&&L3>0){printf("Overflow at (%d,%d)\n\n",
J[e][1],J[e][2]);}L3=pfac(L3);for(i=0;i<PN;i++){P3[i]=P3[i]-PP[i];}return(L3);}
long long pfac(long long mmm){lldiv_t dvn;int i,flg,fl2,sgn=1;if(mmm<0){mmm=0-mmm;sgn=-1;}for(i
=0;i<PN;i++){PP[i]=0;}for(flg=0;mmm>1&&flg==0;){for(i=fl2=0;i<PN&&fl2==0;i++){dvn=lldiv(mmm,
Prm[i]);if(dvn.rem==0){fl2++;if(i>maxp){maxp=i;}}if(fl2==0){flg++;}else{mmm=dvn.quot;PP[i-1]++
;}}if(sgn<0){mmm=-mmm;}return(mmm);}

```

Remark 2.6. This algorithm was at first suspected to be too simple to be true. However, many explicit computations using the codes in 2.5–6 combined with “bernstein” in Singular seem to show that it is correct. Please inform us in case one gets different results using the

above two codes, where the code in 2.5 can be modified as is noted at the end of it. (Since Gröbner bases are not used in the second code, the orderings of monomials are different in general, although they may coincide in simple cases as a consequence of the algorithm.) These are still experimental, and might contain a bug that does not appear for relatively simple examples.

REFERENCES

- [Be 72] Bernstein, I.N., The analytic continuation of generalized functions with respect to a parameter, *Functional Analysis and its Applications* 6 (1972), 273–285.
- [Bj 73] Björk, J.E., Dimensions over algebras of differential operators (preprint, 1973).
- [Br 70] Brieskorn, E., Die Monodromie der isolierten Singularitäten von Hyperflächen, *Manuscripta Math.*, 2 (1970), 103–161.
- [Ca 87] Cassou-Noguès, P., Etude du comportement du polynôme de Bernstein lors d’une déformation à μ constant de $X^a + Y^b$ avec $(a, b) = 1$, *Compos. Math.* 63 (1987), 291–313.
- [DGPS 20] Decker, W., Greuel, G.-M., Pfister, G., Schönemann, H., SINGULAR 4.2.0 — A computer algebra system for polynomial computations, available at <http://www.singular.uni-kl.de> (2020).
- [DiSa 14] Dimca, A., Saito, M., Some remarks on limit mixed Hodge structures and spectrum, *An. Şt. Univ. Ovidius Constanţa Ser. Mat.* 22 (2014), 69–78.
- [GLS 07] Greuel, G.-M., Lossen, C., Shustin, E., *Introduction to Singularities and Deformations*, Springer, Berlin, 2007.
- [JKSY 22] Jung, S.-J., Kim, I.-K., Saito, M., Yoon, Y., Hodge ideals and spectrum of isolated hypersurface singularities, *Ann. Inst. Fourier* 72 (2022), 465–510 (arXiv:1904.02453).
- [Kas 76] Kashiwara, M., B -functions and holonomic systems, *Inv. Math.* 38 (1976/77), 33–53.
- [Kat 81] Kato, M., The b -function of μ -constant deformation of $x^7 + y^5$, *Bull. College Sci., Univ. Ryukyus* 32 (1981), 5–10.
- [Kat 82] Kato, M., The b -function of μ -constant deformation of $x^9 + y^4$, *Bull. College Sci., Univ. Ryukyus* 33 (1982), 5–8.
- [Ma 75] Malgrange, B., Le polynôme de Bernstein d’une singularité isolée, *Lect. Notes in Math.* 459, Springer, Berlin, 1975, pp. 98–119.
- [NaTa 21] Nabeshima, K., Tajima, S., Methods for computing b -functions associated with μ -constant deformations: case of inner modality two, *Kyushu J. Math.* 75 (2021), 55–76.
- [Sa 83] Supplement to: “Gauss-Manin system and mixed Hodge structure”, *Astérisque* 101–102, Soc. Math, France, 1983, 320–331.
- [Sa 84] Saito, M., Hodge filtrations on Gauss-Manin systems I, *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* 30 (1984), 489–498.
- [Sa 88] Saito, M., Exponents and Newton polyhedra of isolated hypersurface singularities, *Math. Ann.* 281 (1988), 411–417.
- [Sa 89] Saito, M., On the structure of Brieskorn lattice, *Ann. Inst. Fourier* 39 (1989), 27–72.
- [Sa 16] Saito, M., Bernstein-Sato polynomials of hyperplane arrangements, *Selecta Math. (N.S.)* 22 (2016), 2017–2057.
- [Sa 20] Roots of Bernstein-Sato polynomials of certain homogeneous polynomials with two-dimensional singular loci, *Pure Appl. Math. Q.* 16 (2020), 1219–1280.
- [Sa 22a] Saito, M., Notes on regular holonomic \mathcal{D} -modules for algebraic geometers (arXiv:2201.01507).
- [Sa 22b] Saito, M., Length of $\mathcal{D}_X f^{-\alpha}$ in the isolated singularity case (arXiv:2208.08977).
- [Sat 75] Sato, M. (ed.), *Singularities of Hypersurfaces and b -Function* (Proceedings of workshop in 1973), RIMS Kokyuroku 225 (Japanese), 1975.
- [SaSh 72] Sato M., Shintani, T., On zeta functions associated with prehomogeneous vector spaces, *Proc. Nat. Acad. Sci. USA* 69 (1972), 1081–1082.
- [ScSt 85] Scherk, J., Steenbrink, J. H. M., On the mixed Hodge structure on the cohomology of the Milnor fibre, *Math. Ann.* 271 (1985), 641–665.
- [St 77a] Steenbrink, J. H. M., Intersection form for quasi-homogeneous singularities, *Compos. Math.* 34 (1977), 211–223.
- [St 77b] Steenbrink, J. H. M., Mixed Hodge structure on the vanishing cohomology, in *Real and complex singularities*, Sijthoff and Noordhoff, Alphen aan den Rijn, 1977, pp. 525–563.
- [Va 81] Varchenko, A. N., Asymptotic mixed Hodge structure in vanishing cohomologies, *Izv. Akad. Nauk SSSR Ser. Mat.* 45 (1981), 540–591.

- [Va 82a] Varchenko, A. N., The complex singularity index does not change along the stratum $\mu = \text{const}$, Funk. Anal. Pri. 16 (1982), 1–12.
- [Va 82b] Varchenko, A. N., A lower bound for the codimension of the stratum $\mu = \text{constant}$ in terms of the mixed Hodge structure, Vestnik Moskov Univ. Ser. I Mat. Mekh. 37 (1982), 28–31.

RIMS Kyoto University, Kyoto 606-8502 Japan