

The limit in the $(k + 2, k)$ -Problem of Brown, Erdős and Sós exists for all $k \geq 2$

Michelle Delcourt * Luke Postle †

October 4, 2022

Abstract

Let $f^{(r)}(n; s, k)$ be the maximum number of edges of an r -uniform hypergraph on n vertices not containing a subgraph with k edges and at most s vertices. In 1973, Brown, Erdős and Sós conjectured that the limit

$$\lim_{n \rightarrow \infty} n^{-2} f^{(3)}(n; k + 2, k)$$

exists for all positive integers $k \geq 2$. They proved this for $k = 2$. In 2019, Glock proved this for $k = 3$ and determined the limit. Quite recently, Glock, Joos, Kim, Kühn, Lichev and Pikhurko proved this for $k = 4$ and determined the limit. We combine their work with a new reduction to fully resolve the conjecture by proving that indeed the limit exists for all positive integers $k \geq 2$.

1 Introduction

In an r -uniform hypergraph, an (s, k) -*configuration* is a collection of k edges which span at most s vertices. An r -uniform hypergraph is said to be (s, k) -*free* if it contains no (s, k) -configuration. In 1973, Brown, Erdős and Sós [2] initiated the study of $f^{(r)}(n; s, k)$, the maximum number of edges in an (s, k) -free r -uniform hypergraph on n vertices, and they made the following conjecture:

Conjecture 1.1 (Brown, Erdős and Sós [2]). *The limit $\lim_{n \rightarrow \infty} n^{-2} f^{(3)}(n; k + 2, k)$ exists for all $k \geq 2$.*

The main result of this paper is that we prove Conjecture 1.1. In [2], Brown, Erdős and Sós proved Conjecture 1.1 for $k = 2$ (the limit in this case is $1/6$, with extremal examples given by Steiner triple systems). In 2019, Glock [5] proved this for $k = 3$ and determined the limit (namely $1/5$). Quite recently, Glock, Joos, Kim, Kühn, Lichev and Pikhurko [7] proved this for $k = 4$ and determined the limit (namely $7/36$). According to [7], the best known general upper bound to date for $\limsup_{n \rightarrow \infty} n^{-2} f^{(3)}(n; k + 2, k)$ is $\frac{k-1}{3k}$ by Brown, Erdős and Sós [2] in 1973.

For lower bounds, this problem is related to a famous conjecture of Erdős [4] from 1973 on the existence of high girth Steiner triple systems. Namely, Erdős conjectured that: for every k and large enough n satisfying the necessary divisibility conditions, there exists a Steiner triple system on n vertices with no $(\ell + 2, \ell)$ -configuration for every $\ell \in [2, k]$. Erdős' conjecture implies a lower bound of $1/6$ on the $\liminf_{n \rightarrow \infty} n^{-2} f^{(3)}(n; k + 2, k)$. In 1993, Lefmann, Phelps, and Rödl [13] proved a weak form of Erdős' conjecture which implies a lower bound here of c_k for some small positive constant c_k depending on k . In the last few years, Erdős' conjecture was proved asymptotically by Glock, Kühn, Lo, and Osthus [9], and independently by Bohman and Warnke [1], which yields the following corollary:

*Department of Mathematics, Toronto Metropolitan University (formerly named Ryerson University), Toronto, Ontario M5B 2K3, Canada mdelcourt@ryerson.ca. Research supported by NSERC under Discovery Grant No. 2019-04269.

†Combinatorics and Optimization Department, University of Waterloo, Waterloo, Ontario N2L 3G1, Canada lpotle@uwaterloo.ca. Partially supported by NSERC under Discovery Grant No. 2019-04304.

Theorem 1.2 ([1, 9]). *Let $k \geq 2$ be an integer. Then*

$$\liminf_{n \rightarrow \infty} n^{-2} f^{(3)}(n; k+2, k) \geq \frac{1}{6}.$$

We remark that Erdős' conjecture was proven in full just this year by Kwan, Sah, Sawhney and Simkin [12]. Of course by not requiring the 3-uniform hypergraph to be linear (i.e. $(4, 2)$ -free), more efficient lower bound constructions are in theory possible (such as were done above for $k = 3$ and $k = 4$).

Glock, Kühn, Lo, and Osthus [9] and Keevash and Long [11] conjectured a generalization of Erdős' conjecture: for every k and large enough n satisfying the necessary divisibility conditions, there exists an (n, r, t) -Steiner system with no $((r-t)\ell + t, \ell)$ -configuration for every $\ell \in [2, k]$. Note for $k = 2$ this is the famous Existence of Designs Conjecture, only recently solved by Keevash [10] (and more recently using purely combinatorial methods by Glock, Kühn, Lo, and Osthus [8]). So this conjecture is a common generalization of Erdős' conjecture and the Existence Conjecture. In light of this conjecture, it is natural to study $n^{-t} f^{(r)}(n; (r-t)k + t, k)$ and indeed in 2020, Shangquan and Tamo [14] conjectured the limit of this exists for all integers $r > t \geq 2$ and $k \geq 2$.

This conjecture was proved asymptotically by the authors [3] and independently by Glock, Joos, Kim, Kühn, and Lichev [6] which provides a lower bound on the limit infimum of $n^{-t} f^{(r)}(n; (r-t)k + t, k)$, generalizing Theorem 1.2, as follows:

Theorem 1.3 ([3, 6]). *Let $k \geq 2$ be an integer. Then*

$$\liminf_{n \rightarrow \infty} n^{-t} f^{(r)}(n; (r-t)k + t, k) \geq \frac{1}{t! \binom{r}{t}}.$$

In fact, both groups actually developed a general theory of *conflict-free hypergraph matchings* (or equivalently hypergraph matchings avoiding forbidden submatchings) and provided theorems where conflict-free almost perfect matchings are found provided various necessary degree and codegree conditions hold. The proofs use different approaches, a random greedy process versus the nibble method, but the general theory is similar (except the nibble approach also works for coloring and for very sparse hypergraphs). This general theory then yields Theorem 1.3 as a simple application.

In the recent paper of Glock, Joos, Kim, Kühn, Lichev and Pikhurko [7] proving the limit exists for $k = 4$, they in fact prove a more general version of Theorem 1.3 by invoking the general theory of conflict-free hypergraph matchings. Their most general version is useful for proving better lower bounds, but we only need the following result:

Theorem 1.4 (Corollary 3.2 in [7]). *Fix integers $k \geq 2$, $r \geq 3$ and $t \in [2, r-1]$. Also, fix a $((r-t)k + t, k)$ -free r -uniform hypergraph F on m vertices which is also $((r-t)\ell + t - 1, \ell)$ -free for all $\ell \in [2, k-1]$. Then,*

$$\liminf_{n \rightarrow \infty} n^{-t} f^{(r)}(n; (r-t)k + t, k) \geq \frac{|F|}{m^t}.$$

Specifically, our proof relies on the above result for $r = 3$ and $t = 2$, namely:

Theorem 1.5 (Theorem 1.4 with $r = 3$, $t = 2$). *Fix integer $k \geq 2$. Also, fix a $(k+2, k)$ -free 3-uniform hypergraph F on m vertices which is also $(\ell+1, \ell)$ -free for all $\ell \in [2, k-1]$. Then,*

$$\liminf_{n \rightarrow \infty} n^{-2} f^{(3)}(n; k+2, k) \geq \frac{|F|}{m^2}.$$

We introduce the function $g^{(3)}(n; k+2, k)$ denoting the maximum number of edges in a $(k+2, k)$ -free 3-uniform hypergraph on n vertices that is also $(\ell+1, \ell)$ -free for all $\ell \in [2, k-1]$. We note that f is at least g .

In this language, the following is a corollary of Theorem 1.5:

Corollary 1.6. *For all positive integers $k \geq 2$,*

$$\liminf_{n \rightarrow \infty} n^{-2} f^{(3)}(n; k+2, k) \geq \limsup_{n \rightarrow \infty} n^{-2} g^{(3)}(n; k+2, k).$$

Our main technical result is the following then:

Theorem 1.7. *For all positive integers $k \geq 2$,*

$$\limsup_{n \rightarrow \infty} n^{-2} f^{(3)}(n; k+2, k) = \limsup_{n \rightarrow \infty} n^{-2} g^{(3)}(n; k+2, k).$$

As an immediate corollary of Corollary 1.6 and Theorem 1.7, we have:

Theorem 1.8. *For all positive integers $k \geq 2$, $\lim_{n \rightarrow \infty} n^{-2} f^{(3)}(n; k+2, k)$ exists and*

$$\lim_{n \rightarrow \infty} n^{-2} f^{(3)}(n; k+2, k) = \limsup_{n \rightarrow \infty} n^{-2} g^{(3)}(n; k+2, k).$$

We also prove the following lemma:

Lemma 1.9. *Let F be a $(k+2, k)$ -free 3-uniform hypergraph that is also $(\ell+1, \ell)$ -free for all $\ell \in [2, k-1]$. Then*

$$e(F) \leq \frac{k-1}{2k-1} \cdot \binom{v(F)}{2} \leq \frac{k-1}{4k-2} \cdot v(F)^2.$$

As a corollary of Lemma 1.9 and Theorem 1.7 we improve the general upper bound of $\limsup_{n \rightarrow \infty} n^{-2} f^{(3)}(n; k+2, k)$ from $\frac{k-1}{3k}$ to $\frac{k-1}{4k-2}$ as follows:

Theorem 1.10.

$$\lim_{n \rightarrow \infty} n^{-2} f^{(3)}(n; k+2, k) = \limsup_{n \rightarrow \infty} n^{-2} g^{(3)}(n; k+2, k) \leq \frac{k-1}{4k-2}.$$

We also note that the proof of the main technical result (Theorem 3.1) of [7] actually constructs graphs which are not only $((r-t)k+t, k)$ -free but are also $(r-t)\ell+t-1, \ell)$ -free for all $\ell \in [2, k-1]$. Hence the f in the liminf in the statement of Theorem 3.1 can actually be replaced by the function g we defined. Hence as a corollary one obtains the following slightly stronger statement of Theorem 1.4.

Theorem 1.11. *Fix integers $k \geq 2$, $r \geq 3$ and $t \in [2, r-1]$. Also, if F is $((r-t)k+t, k)$ -free r -uniform hypergraph on m vertices which is also $((r-t)\ell+t-1, \ell)$ -free for all $\ell \in [2, k-1]$, then,*

$$\liminf_{n \rightarrow \infty} n^{-t} g^{(r)}(n; (r-t)k+t, k) \geq \frac{|F|}{m^t}.$$

It follows that

$$\liminf_{n \rightarrow \infty} n^{-t} g^{(r)}(n; (r-t)k+t, k) = \limsup_{n \rightarrow \infty} n^{-t} g^{(r)}(n; (r-t)k+t, k),$$

that is $\lim_{n \rightarrow \infty} n^{-t} g^{(r)}(n; (r-t)k+t, k)$ exists.

Given Theorem 1.11, Theorem 1.7 actually yields the following:

Theorem 1.12. *For all positive integers $k \geq 2$, $\lim_{n \rightarrow \infty} n^{-2} f^{(3)}(n; k+2, k)$ exists and*

$$\lim_{n \rightarrow \infty} n^{-2} f^{(3)}(n; k+2, k) = \lim_{n \rightarrow \infty} n^{-2} g^{(3)}(n; k+2, k).$$

2 Proof

In [7], an approach to proving Conjecture 1.1 in full was suggested, namely can one prove that one can always delete $o(m^2)$ edges from a $(k+2, k)$ -free 3-uniform hypergraph F to make it $(\ell+1, \ell)$ -free for all $\ell \in [2, k-1]$? While this strategy is intriguing, it seems difficult in general.

Here we pursue a different approach. Instead we prove (Lemma 2.3 below) that provided that $|F| \geq \frac{1}{6} \cdot (1 - \frac{1}{2k}) \cdot v(F)^2$, there is a subgraph F' with $v(F') \geq \frac{v(F)}{\sqrt{2k}}$ and $\frac{|F'|}{v(F')^2} \geq \frac{|F|}{v(F)^2}$ that is $(\ell+1, \ell)$ -free for all $\ell \in [2, k-1]$. That is, we prove that having $(\ell+1, \ell)$ -configurations is ‘inefficient’ for the extremal $(k+2, k)$ -free graphs. Given Lemma 2.3, Theorem 1.7 follows straightforwardly (but we include a proof for completeness).

To pursue this approach, we first prove a structural lemma. Let $k \geq 2$ be a positive integer and let $\ell \in [2, k-1]$. We say an $(\ell+1, \ell)$ -configuration S in a $(k+2, k)$ -free 3-uniform hypergraph F is k -maximal if there does not exist an $(\ell'+1, \ell')$ -configuration S' with $S \subseteq S'$ and $\ell < \ell' \leq k-1$.

Lemma 2.1. *Let $k \geq 2$ be integer. Let F be a $(k+2, k)$ -free 3-uniform hypergraph. Let S be a k -maximal $(\ell+1, \ell)$ -configuration of F with $\ell \in [2, k-1]$. Then all of the following hold:*

1. $e(F[V(S)]) \leq k-1$,
2. there does not exist an edge $e \in E(F)$ with $|e \cap V(S)| = 2$,
3. there do not exist edges $e, f \in E(F)$ with $|e \cap V(S)| = |f \cap V(S)| = 1$ and $e \setminus V(S) = f \setminus V(S)$, and
4. the graph H with $V(H) = V(F) \setminus V(S)$ and $E(H) = \{T \in \binom{V(H)}{2} : \exists e \in E(F), |e \cap S| = 1, e \setminus V(S) = T\}$ is a forest whose components have size at most $k-\ell$,

and hence

$$e(F) - e(F \setminus V(S)) \leq \left(1 - \frac{1}{k}\right) \cdot (v(F) - v(S)) + (k-1).$$

Proof. First we prove (1). Suppose not. Then $e(F[V(S)]) \geq k$ and hence $E(F[V(S)])$ is a (k, k) -configuration, contradicting that F is $(k+2, k)$ -free. This proves (1).

Next we prove (2). Suppose not. That is, there exists an edge $e \in E(F)$ with $|e \cap V(S)| = 2$. If $\ell = k-1$, then $S \cup e$ is a $(k+1, k)$ -configuration contradicting that F is $(k+2, k)$ -free. So we may assume that $\ell < k-1$. But then $S \cup e$ is an $(\ell+2, \ell+1)$ -configuration contradicting that S is k -maximal. This proves (2).

Next we prove (3). Suppose not. That is, there exist edges $e, f \in E(F)$ with $|e \cap V(S)| = |f \cap V(S)| = 1$ and $e \setminus V(S) = f \setminus V(S)$. If $\ell = k-1$, then $S \cup e$ is a $(k+2, k)$ -configuration contradicting that F is $(k+2, k)$ -free. So we may assume that $\ell < k-1$. If $\ell = k-2$, then $S \cup \{e, f\}$ is a $(k+1, k)$ -configuration contradicting that F is $(k+2, k)$ -free. So we may assume that $\ell < k-2$. But then $S \cup \{e, f\}$ is an $(\ell+3, \ell+2)$ -configuration contradicting that S is k -maximal. This proves (3).

Next we prove (4). Suppose not. That is, H has a component of size greater than $k-\ell$ or H has a cycle of length at most $k-\ell$. First suppose that H has a component of size greater than $k-\ell$. Thus H contains a tree T on exactly $k-\ell+1$ vertices. Let $E(T) = \{e_1, e_2, \dots, e_{k-\ell}\}$. For each $i \in [k-\ell]$, let $f_i \in E(F)$ such that $f_i \setminus V(S) = e_i$. But then $S \cup \{f_i : i \in [k-\ell]\}$ is a set of $|S| + (k-\ell) = k$ edges on a set of $|V(S)| + (k-\ell+1) = k+2$ vertices, contradicting that F is $(k+2, k)$ -free.

So we may assume that H has a cycle C of length j where $j \leq k-\ell$. Let $E(C) = \{e_1, \dots, e_j\}$. For each $i \in [j]$, let $f_i \in E(F)$ such that $f_i \setminus V(S) = e_i$. Let $S' = S \cup \{f_i : i \in [j]\}$. Hence S' is a set of $\ell+j$ edges on a set of $\ell+j+1$ vertices. If $j = k-\ell$, then S' is a $(k+1, k)$ -configuration, contradicting that F is $(k+2, k)$ -free. So we assume that $j < k-\ell$. But then S' is an $(\ell+j+1, \ell+j)$ -configuration where $\ell+j \leq k-1$ and $\ell+j > \ell$, contradicting that S is k -maximal. This proves (4).

Finally we prove the formula. For each $i \in [3]$, let $E_i := \{e \in E(F) : |e \cap V(S)| = i\}$. Since F is a 3-uniform hypergraph, we have that

$$e(F) - e(F \setminus V(S)) = \sum_{i=1}^3 |E_i|.$$

By (1), we have that $|E_3| \leq k - 1$. By (2), we have that $|E_2| = 0$. By (3), we have that $|E_1| = e(H)$. Combining, we find that

$$e(F) - e(F \setminus V(S)) \leq k - 1 + e(H).$$

Let a be the number of components of H . By (4), each component of H has size at most $k - \ell \leq k$ and hence

$$a \geq \frac{1}{k} \cdot v(F \setminus V(S)).$$

By (4), H is a forest and hence

$$\begin{aligned} e(H) &= v(F \setminus V(S)) - a \leq \left(1 - \frac{1}{k}\right) \cdot v(F \setminus V(S)) \\ &= \left(1 - \frac{1}{k}\right) \cdot (v(F) - v(S)). \end{aligned}$$

Combining, we find that

$$e(F) - e(F \setminus V(S)) \leq e(H) + k - 1 \leq \left(1 - \frac{1}{k}\right) \cdot (v(F) - v(S)) + (k - 1),$$

as desired. \square

We now apply Lemma 2.1 to find an induced subgraph that is $(\ell + 1, \ell)$ -free for all $\ell \in [2, k - 1]$ where not too many edges are lost in proportion to the number of vertices deleted.

Lemma 2.2. *Let $k \geq 2$ be integer. Let F be a $(k + 2, k)$ -free 3-uniform hypergraph. There exists a $(k + 2, k)$ -free subgraph F' of F that is also $(\ell + 1, \ell)$ -free for all $\ell \in [2, k - 1]$ such that*

$$e(F) - e(F') \leq \frac{1}{6} \cdot \left(1 - \frac{1}{k}\right) \cdot (v(F) - v(F')) \cdot (v(F) + v(F') + 2k).$$

Proof. We proceed by induction on $v(F)$. First suppose F is also $(\ell + 1, \ell)$ -free for all $\ell \in [2, k - 1]$. Then $F' = F$ is as desired. So we may assume that there exists an $(\ell + 1, \ell)$ -configuration in F .

Let S be a k -maximal $(\ell + 1, \ell)$ -configuration in F with $\ell \in [2, k - 1]$. By Lemma 2.1, we have that

$$e(F) - e(F \setminus V(S)) \leq \left(1 - \frac{1}{k}\right) \cdot (v(F) - v(S)) + (k - 1).$$

Note that $v(S) \geq 3$ since F is 3-uniform hypergraph and hence $v(F \setminus V(S)) < v(F)$. Thus by induction, we find that there exists a $(k + 2, k)$ -free subgraph F' of $F \setminus V(S)$ that is also $(\ell + 1, \ell)$ -free for all $\ell \in [2, k - 1]$ such that

$$\begin{aligned} e(F \setminus V(S)) - e(F') &\leq \frac{1}{6} \cdot \left(1 - \frac{1}{k}\right) \cdot (v(F \setminus V(S)) - v(F')) \cdot (v(F \setminus V(S)) + v(F') + 2k) \\ &= \frac{1}{6} \cdot \left(1 - \frac{1}{k}\right) \cdot (v(F) - v(S) - v(F')) \cdot (v(F) - v(S) + v(F') + 2k) \\ &= \frac{1}{6} \cdot \left(1 - \frac{1}{k}\right) \cdot (v(F) - v(F')) \cdot (v(F) + v(F') + 2k) \\ &\quad - \frac{1}{6} \cdot \left(1 - \frac{1}{k}\right) \cdot v(S) \cdot (2 \cdot v(F) + 2k - v(S)). \end{aligned}$$

Now using that $2 \cdot v(F) - v(S) \geq 2(v(F) - v(S))$, we have that

$$\frac{1}{6} \cdot \left(1 - \frac{1}{k}\right) \cdot v(S) \cdot (2 \cdot v(F) + 2k - v(S)) \geq \frac{2 \cdot v(S)}{6} \cdot \left(\left(1 - \frac{1}{k}\right) \cdot (v(F) - v(S)) + (k - 1) \right)$$

Using that $v(S) \geq 3$, we find that

$$\frac{1}{6} \cdot \left(1 - \frac{1}{k}\right) \cdot v(S) \cdot (2 \cdot v(F) + 2k - v(S)) \geq \left(1 - \frac{1}{k}\right) \cdot (v(F) - v(S)) + (k - 1).$$

Altogether then, we find that

$$e(F) - e(F') \leq \frac{1}{6} \cdot \left(1 - \frac{1}{k}\right) \cdot (v(F) - v(F')) \cdot (v(F) + v(F') + 2k),$$

as desired. \square

We need an upper bound on the number of edges in a $(k + 2, k)$ -free 3-uniform hypergraph that is also $(\ell + 1, \ell)$ -free for all $\ell \in [2, k - 1]$. While a trivial upper bound would suffice for our purposes, finding a better upper bound is also of interest in light of Theorem 1.7. Thus we found Lemma 1.9 which we now prove, but first a definition.

The *t-shadow* of an r -uniform hypergraph F is the t -uniform hypergraph H with $V(H) = V(F)$ and $E(H) = \{S \in \binom{V(H)}{t} : \exists e \in E(F) \text{ with } S \subseteq e\}$. If Y is a subset of $E(H)$, then we define the *t-shadow of Y in F* as the t -shadow of the subgraph F' that is the union of the edges in Y (so $V(F') = \bigcup_{e \in Y} V(e)$ and $E(F') = Y$).

Proof of Lemma 1.9. Let H be the graph with $V(H) = E(F)$ and

$$E(H) = \left\{ \{S, T\} \in \binom{V(H)}{2} : |S \cap T| \geq 2 \right\}.$$

Claim 2.2.1. *If T is a connected subgraph of H , then $v(T) \leq k - 1$ and $V(T)$ (as a set of edges of F) spans exactly $v(T) + 2$ vertices of F .*

Proof. Let T' be a connected subgraph of H . It follows from the definition of H that $V(T')$ (as a set of edges of F) spans at most $v(T') + 2$ vertices of F and hence $V(T')$ is a $(v(T') + 2, v(T'))$ -configuration of F .

Since F is $(k + 2, k)$ -free, it follows that there does not exist a connected subgraph T' of H with $v(T') \geq k$. Thus $v(T) \leq k - 1$. Moreover, if $V(T)$ spans at most $v(T) + 1$ vertices of F , then $V(T)$ is a $(v(T) + 1, v(T))$ -configuration of F , contradicting that F is $(\ell + 1, \ell)$ -free for all $\ell \in [2, k - 1]$. \square

Claim 2.2.2. *If T is a connected subgraph of H , then the 2-shadow of $V(T)$ in F has exactly $2 \cdot v(T) + 1$ edges.*

Proof. We proceed by induction on $v(T)$. Let a be the number of edges in the 2-shadow of $V(T)$ in F . If $v(T) = 1$, then $a = 3$ as desired.

So we may assume that $v(T) \geq 2$. Let X_T be the set of vertices of F that $V(T)$ spans. By Claim 2.2.1, $|X_T| = v(T) + 2$. Since T is connected, there exists a connected subgraph T' of T such that $v(T') = v(T) - 1$. Let $V(T) \setminus V(T') = \{e\}$. Let $X_{T'}$ be the set of vertices of F that $V(T')$ spans. By Claim 2.2.1, $|X_{T'}| = v(T') + 2$.

Note that $X_{T'} \subseteq X_T$. Let $X_T \setminus X_{T'} = \{v\}$. It follows that $v \in e$ and yet $v \notin f$ for all $f \in V(T')$. Let $e = vv_1v_2$. Hence vv_1, vv_2 are in the 2-shadow of $V(T)$ in F but are not in the 2-shadow of $V(T')$ in F . By definition of H , it then follows that v_1v_2 is in the 2-shadow of $V(T')$.

By induction, the number of edges in the 2-shadow of $V(T')$ is exactly $2 \cdot v(T') + 1$. Thus the number of edges in the 2-shadow of $V(T)$ is exactly $(2 \cdot v(T') + 1) + 2 = 2 \cdot v(T) + 1$, as desired. \square

Let T_1, \dots, T_r be the components of H . For each $i \in [r]$, let F_i be the 2-shadow of $V(T_i)$ in F . By Claim 2.2.2, we have for each $i \in [r]$ that $e(F_i) = 2 \cdot v(T_i) + 1$. By the definition of H , it follows that $E(F_i) \cap E(F_j) = \emptyset$ for each $i \neq j \in [r]$. By Claim 2.2.1, $v(T_i) \leq k - 1$ for all $i \in [r]$. It follows that for each $i \in [r]$, we have that

$$v(T_i) = \frac{v(T_i)}{2 \cdot v(T_i) + 1} \cdot e(F_i) \leq \frac{k-1}{2k-1} \cdot e(F_i).$$

Since the F_i are edge-disjoint, we find that

$$\binom{v(F)}{2} \geq \sum_{i=1}^r e(F_i) \geq \frac{2k-1}{k-1} \sum_{i=1}^r v(T_i) = \frac{2k-1}{k-1} \cdot e(F).$$

Rearranging, we find that

$$e(F) \leq \frac{k-1}{2k-1} \cdot \binom{v(F)}{2} \leq \frac{k-1}{4k-2} \cdot v(F)^2,$$

as desired. \square

We are now prepared to state and prove Lemma 2.3 about finding a large $(\ell + 1, \ell)$ -free subgraph for all $\ell \in [2, \ell - 1]$ that is as dense as the original $(k + 2, k)$ -free 3-uniform hypergraph.

Lemma 2.3. *Let F be a $(k+2, k)$ -free 3-uniform hypergraph with $v(F) \geq 8k^2$ and $e(F) \geq \frac{1}{6} \left(1 - \frac{1}{2k}\right) \cdot v(F)^2$. Then there exists a $(k+2, k)$ -free subgraph F' of F that is also $(\ell + 1, \ell)$ -free for all $\ell \in [2, k - 1]$ such that both of the following hold:*

1. $v(F') \geq \frac{v(F)}{\sqrt{2k}}$, and
2. $\frac{e(F')}{v(F')^2} \geq \frac{e(F)}{v(F)^2}$.

Proof. By Lemma 2.2, there exists a $(k + 2, k)$ -free subgraph F' of F that is also $(\ell + 1, \ell)$ -free for all $\ell \in [2, k - 1]$ such that

$$e(F) - e(F') \leq \frac{1}{6} \cdot \left(1 - \frac{1}{k}\right) \cdot (v(F) - v(F')) \cdot (v(F) + v(F') + 2k).$$

By Lemma 1.9, we have that

$$e(F') \leq \frac{k-1}{4k-2} \cdot v(F')^2 \leq \frac{1}{4} \cdot v(F')^2.$$

Combining, we find that

$$e(F) \leq \frac{1}{4} \cdot v(F')^2 + \frac{1}{6} \cdot \left(1 - \frac{1}{k}\right) \cdot (v(F) - v(F')) \cdot (v(F) + v(F') + 2k).$$

Since $e(F) \geq \frac{1}{6} \left(1 - \frac{1}{2k}\right) \cdot v(F)^2$, we have that as $k \geq 2$,

$$\frac{1}{12k} \cdot v(F)^2 \leq \frac{1}{12} \cdot v(F')^2 + \frac{k-1}{3} \cdot (v(F) - v(F'))$$

Multiplying by $12k$ and upper bounding $(k - 1)(v(F) - v(F'))$ by $k \cdot v(F)$ yields

$$v(F)^2 \leq k \cdot v(F')^2 + 4k^2 \cdot v(F) \leq v(F')^2 + \frac{1}{2} \cdot v(F)^2,$$

where for the last inequality we used that $v(F) \geq 8k^2$. Rearranging gives

$$\frac{1}{2}v(F)^2 \leq k \cdot v(F')^2,$$

Taking square roots, it now follows that

$$\frac{v(F)}{\sqrt{2k}} \leq v(F').$$

This proves (1).

Now we prove (2). Let $a = \frac{e(F)}{v(F)^2}$. Suppose towards a contradiction that $\frac{e(F')}{v(F')^2} < a$. Note this implies that $v(F') < v(F)$. Moreover, it implies that

$$e(F) - e(F') \geq a \cdot (v(F)^2 - v(F')^2) = a \cdot (v(F) - v(F')) \cdot (v(F) + v(F')).$$

But then using that $v(F) - v(F') > 0$, it follows that

$$a \cdot (v(F) + v(F')) \leq \frac{e(F) - e(F')}{v(F) - v(F')} \leq \frac{1}{6} \cdot \left(1 - \frac{1}{k}\right) \cdot (v(F) + v(F') + 2k).$$

Using that $a \geq \frac{1}{6} \left(1 - \frac{1}{2k}\right)$, we find that

$$\frac{1}{12k} \cdot (v(F) + v(F')) \leq \frac{k-1}{3}.$$

But this implies that

$$v(F) \leq 4k^2,$$

contradicting that $v(F) \geq 8k^2$. This proves (2). \square

We now prove Theorem 1.7 which stated that the $\limsup n^{-2}f^{(3)}(n; k+2, k)$ and $\limsup n^{-2}g^{(3)}(n; k+2, k)$ were equal.

Proof of Theorem 1.7. Suppose not. Let $a = \limsup_{n \rightarrow \infty} n^{-2}f^{(3)}(n; k+2, k)$ and $b = \limsup_{n \rightarrow \infty} n^{-2}g^{(3)}(n; k+2, k)$. Let $\varepsilon = a - b$. Since $f^{(3)}(n; k+2, k) \geq g^{(3)}(n; k+2, k)$ for all positive integers n , we find that $a \geq b$. Since $a \neq b$, we have that $\varepsilon > 0$.

Let $c = \max\{a - \frac{\varepsilon}{2}, \frac{1}{6} \cdot (1 - \frac{1}{2k})\}$. Since $a \geq \frac{1}{6}$ by Theorem 1.2, we find that $c < a$. Yet by definition, $c \geq a - \frac{\varepsilon}{2} > b$. Let

$$I_f = \{n \in \mathbb{N} : n^{-2}f^{(3)}(n; k+2, k) \geq c\}.$$

Let

$$I_g = \{n \in \mathbb{N} : n^{-2}g^{(3)}(n; k+2, k) \geq c\}.$$

Since $c < a$, $|I_f|$ is infinite. Since $c > b$, I_g is finite. Let $n_0 = \max\{n : n \in I_g\}$. Let

$$n'_0 = \max\left\{\sqrt{2k} \cdot n_0 + 1, 8k^2\right\}.$$

Since $|I_f|$ is infinite, there exists $n_1 \in I_f$ such that $n_1 \geq n'_0$. That is, there exists a $(k+2, k)$ -free 3-uniform hypergraph F on n_1 vertices such that $e(F) \geq c \cdot v(F)^2$.

Note that $v(F) \geq 8k^2$ by definition of n'_0 . Since $c \geq \frac{1}{6} \cdot (1 - \frac{1}{2k})$, we have that $e(F) \geq \frac{1}{6} \cdot (1 - \frac{1}{2k}) \cdot v(F)^2$. Hence by Lemma 2.3, there exists a $(k+2, k)$ -free subgraph F' of F that is also $(\ell+1, \ell)$ -free for all $\ell \in [2, k-1]$ such that both of the following hold:

1. $v(F') \geq \frac{v(F)}{\sqrt{2k}}$, and
2. $\frac{e(F')}{v(F')^2} \geq \frac{e(F)}{v(F)^2}$.

By (1) and the definition of n'_0 , we find that $v(F') > n_0$. By (2), we find that $e(F') \geq c \cdot v(F')^2$. Hence $v(F') \in I_g$, contradicting that n_0 is the maximum value in I_g . \square

References

- [1] Tom Bohman and Lutz Warnke. Large girth approximate Steiner triple systems. *Journal of the London Mathematical Society*, 100(3):895–913, 2019.
- [2] William Brown, Paul Erdős, and Vera Sós. Some extremal problems on r -graphs, *New directions in the theory of graphs*. pages 53–63, 1973.
- [3] Michelle Delcourt and Luke Postle. Finding an almost perfect matching in a hypergraph avoiding forbidden submatchings. *arXiv:2204.08981*, 2022.
- [4] Paul Erdős. Problems and results in combinatorial analysis. In *Colloq. Internat. Theor. Combin. Rome*, pages 3–17, 1973.
- [5] Stefan Glock. Triple systems with no three triples spanning at most five points. *Bulletin of the London Mathematical Society*, 51(2):230–236, 2019.
- [6] Stefan Glock, Felix Joos, Jaehoon Kim, Marcus Kühn, and Lyuben Lichev. Conflict-free hypergraph matchings. *arXiv:2205.05564*, 2022.
- [7] Stefan Glock, Felix Joos, Jaehoon Kim, Marcus Kühn, Lyuben Lichev, and Oleg Pikhurko. On the $(6,4)$ -problem of Brown, Erdős, and Sós. *arXiv:2209.14177*, 2022.
- [8] Stefan Glock, Daniela Kühn, Allan Lo, and Deryk Osthus. The existence of designs via iterative absorption: hypergraph F -designs for arbitrary F . *To appear in Memoirs of the American Mathematical Society*, *arXiv preprint arXiv:1611.06827*, 2016.
- [9] Stefan Glock, Daniela Kühn, Allan Lo, and Deryk Osthus. On a conjecture of Erdős on locally sparse Steiner triple systems. *Combinatorica*, 40(3):363–403, 2020.
- [10] Peter Keevash. The existence of designs. *arXiv preprint arXiv:1401.3665*, 2014.
- [11] Peter Keevash and Jason Long. The Brown-Erdős-Sós conjecture for hypergraphs of large uniformity. *arXiv:2007.14824*, 2020.
- [12] Matthew Kwan, Ashwin Sah, Mehtaab Sawhney, and Michael Simkin. High-girth Steiner triple systems. *arXiv preprint arXiv:2201.04554*, 2022.
- [13] Hanno Lefmann, Kevin T Phelps, and Vojtěch Rödl. Extremal problems for triple systems. *Journal of Combinatorial Designs*, 1(5):379–394, 1993.
- [14] Chong Shangquan and Itzhak Tamo. Degenerate turán densities of sparse hypergraphs. *Journal of Combinatorial Theory, Series A*, 173:105228, 2020.