

QUANTUM SUPERSYMMETRIC PAIRS AND \imath SCHUR DUALITY OF TYPE AIII

YAOLONG SHEN

ABSTRACT. Expanding the classic work of Letzter and Kolb, we construct quantum supersymmetric pairs $(\mathbf{U}, \mathbf{U}^s)$ of type AIII and formulate their basic properties. We establish an \imath Schur duality between the \imath quantum supergroup \mathbf{U}^s and a Hecke algebra of type B acting on a tensor space, providing a super generalization of the \imath Schur duality of type AIII. Furthermore, a (quasi) K -matrix for general parameters is constructed and used to realize the Hecke algebra action on the tensor space.

CONTENTS

1. Introduction	2
1.1. Background	2
1.2. Goal	3
1.3. Main results	3
1.4. Organization	5
2. Quantum supergroup of type A	5
2.1. The general linear Lie superalgebra	5
2.2. Root system of $\mathfrak{gl}(m n)$	6
2.3. Quantum supergroup of type A	7
3. Braid group operators	11
3.1. Odd reflections	11
3.2. Braid group operators	12
4. Quantum supersymmetric pair of type AIII	17
4.1. \imath Quantum supergroup of type AIII	18
4.2. Coideal subalgebra property	20
4.3. Quantum \imath Serre relations	22
4.4. Natural filtration	25
5. \imath Schur duality	26
5.1. Bimodule structure	26
5.2. \imath Schur duality of type B	28
6. Quasi K -matrix	30
6.1. Preparation	31
6.2. A recursive formula	32
6.3. Technical Lemmas	33
6.4. Construction of Υ	35
7. K -matrix and the H_0 -action	38
7.1. K -matrix	38
7.2. Realizing H_0 via K -matrix	40

1. INTRODUCTION

1.1. Background. The Schur duality between the general linear group and the symmetric group is one of the most fundamental topics in representation theory. Jimbo [Jim86] provided the Schur-Jimbo duality between the type A quantum group and the Hecke algebra of type A.

Let \mathfrak{g} be a Lie algebra of any finite type and θ be an involution on \mathfrak{g} , the theory of quantum symmetric pairs $(\mathbf{U}_q(\mathfrak{g}), \mathbf{U}^\iota)$ as the quantization of $(\mathfrak{g}, \mathfrak{g}^\theta)$ has been systematically developed by Letzter [Let99], [Let02], where $\mathbf{U}_q(\mathfrak{g})$ is the Drinfeld-Jimbo quantum group associated to \mathfrak{g} . The algebra \mathbf{U}^ι is referred to as an ι quantum group. Kolb [Ko12] has further studied and generalized this theory to the Kac-Moody case. For example, $(\mathbf{U}_q(\mathfrak{g}) \otimes \mathbf{U}_q(\mathfrak{g}), \mathbf{U}_q(\mathfrak{g}))$ is a quantum symmetric pair of diagonal type. Thus ι quantum groups can always be viewed as generalizations of quantum groups. Bao and Wang have developed a general theory of the canonical basis for ι quantum groups arising from quantum symmetric pairs of arbitrary finite type in [BW18b] and of Kac-Moody type in [BW21].

In recent years, the Schur-Jimbo duality has been naturally generalized along the development of the ι quantum groups. In [BW18a], the authors show that the Hecke algebra of type B and the ι quantum group of type AIII satisfy a double centralizer property (see also [Bao17]). Moreover, the Kazhdan-Lusztig basis of type B was shown to coincide with the ι canonical bases [BW18b] arising from tensor product module of ι quantum group. Later on it was generalized to a multi-parameter setting [BWW18]. More recently, a common generalization of both type A and type B Schur dualities has been constructed in [SW21]. Cui and Shen [CS22] also established Schur dualities between ι quantum groups of type AI and AII with Wenzl's q -Brauer algebra [We12].

On the other hand, Chung [Ch19] has studied quantum symmetric pairs $(\mathbf{U}_\pi, \mathbf{U}_\pi^\iota)$ for quantum covering algebras \mathbf{U}_π which is introduced in [CHW13] and specializes to the Lusztig quantum group when $\pi = 1$ and quantum supergroups of anisotropic type when $\pi = -1$; see also [Ch21].

A fundamental property of a quantum symmetric pair is the existence of the quasi K -matrix. The quasi K -matrix for a quantum symmetric pair was first introduced in [BW18a, §2.3] as the intertwiner between the embedding of the ι quantum group into the underlying quantum group and its bar-conjugated embedding with some conditions imposed on the parameters; also cf. [BK19]. In [AV20], the authors reformulated quasi K -matrix without reference to the bar involution, allowing for more general parameters. More recently, Wang and Zhang have given a conceptual reformulation of quasi K -matrix in general parameters as an intertwiner of anti-involutions in [WZ22].

In the Schur-Jimbo duality, it is well known that the action of the generators H_i ($1 \leq i \leq d - 1$) of the type A Hecke algebra can be realized via the R -matrix, cf. [Jim86]. In [BW18a, Theorem 2.18], the authors constructed a K -matrix \mathcal{T} which is an \mathbf{U}^ι -module isomorphism for any finite-dimensional \mathbf{U} -module (see also [BK19] for generalizations). Furthermore, Bao and Wang showed in [BW18a, Lemma 5.3] that the H_0 -action in the type B ι Schur duality can be realized via the K -matrix \mathcal{T} . This result was further generalized in [SW21, §5.4].

From now on we fix $\mathfrak{g} = \mathfrak{gl}(m|n)$ to be a Lie superalgebra of type A; a well-known fact is that the Dynkin diagrams of \mathfrak{g} are not unique (cf. [CW12]). Sergeev [S85] has extended the Schur duality in the setting of $\mathfrak{gl}(m|n)$.

The *quantum supergroup* \mathbf{U} , as a Drinfeld-Jimbo quantization of \mathfrak{g} , has been defined in [Ya94] associated to the standard Dynkin diagram of \mathfrak{g} ; also cf. [KT91, Mi06]. Moreover, it was shown in [Mi06] that the type A quantum supergroup associated to the standard Dynkin diagram and the Hecke algebra of type A satisfy a double centralizer property.

Lusztig's braid group action [Lus93] is one of the fundamental constructions for usual quantum groups. The braid group operators can be viewed as quantization of the reflections on the weight data. For Lie superalgebras, the fundamental systems of the root system associated to \mathfrak{g} are not conjugated under the Weyl group actions because of the existence of odd roots; cf. [CW12]. The reflections associated to the odd (resp. even) roots are called odd (resp. even) reflections.

Note that different presentations of \mathbf{U} correspond to different choices of simple roots, despite the common labeling of generators. In [C16], the author has developed such a super analogue of Lusztig's braid group operators in the setting of $\mathfrak{gl}(m|1)$. While the braid group operators associated to the even simple roots remain to be algebra isomorphisms of \mathbf{U} , the ones associated to the odd simple roots must be treated as isomorphisms of \mathbf{U} with two different presentations.

1.2. Goal. Let $(\mathfrak{g}, \mathfrak{g}^\theta)$ be a *supersymmetric pair* of type AIII; see (4.1). Our first goal is to construct a suitable quantization $(\mathbf{U}, \mathbf{U}^\imath)$ of $(\mathfrak{g}, \mathfrak{g}^\theta)$ following [Let99, Let02], which is called a *quantum supersymmetric pair*. \mathbf{U}^\imath is referred to as an *\imath quantum supergroup*.

A type AIII Satake diagram (4.1) consists of $(I = I_\bullet \cup I_\circ, \tau)$ where τ is the diagram involution. For example, $(\mathfrak{gl}(m_1 + m_2|n_1 + n_2), \mathfrak{gl}(m_1|n_1) \oplus \mathfrak{gl}(m_2|n_2))$ is a supersymmetric pair of type AIII. Note that the Satake diagrams we consider in this paper allow odd simple roots to appear in both I_\circ and I_\bullet .

In order to achieve the first goal, we are led to consider \mathbf{U} associated to different Dynkin diagrams, since the definition of \mathbf{U}^\imath involves braid group operator associated to all simple roots in I_\bullet . The construction will be greatly simplified if we assume that I_\bullet consists of even simple roots only, which is still new.

Our next goal is to establish a Schur duality between \mathbf{U}^\imath and the Hecke algebra of type B acting on the tensor module of the natural representation of \mathbf{U} , which generalizes the \imath Schur dualities in [BW18a, BWW18, SW21].

Our third goal is to construct the quasi K -matrix Υ associated to $(\mathbf{U}, \mathbf{U}^\imath)$ generalizing the conceptual reformulation in [WZ22]; also cf. [BW18a, BK19]. The quasi K -matrix plays a fundamental role for a quantum supersymmetric pair as the quasi R -matrix for a quantum supergroup; cf. [KT91].

Our last goal is to construct a K -matrix \mathcal{T} following [BW18a]. Moreover, after imposing special parameters so that the \imath Schur duality holds, we can show that \mathcal{T} realizes the H_0 -action on the tensor module of the fundamental representations of \mathbf{U} . This gives us a conceptual explanation on the commutativity between the H_0 -actions and the actions of \mathbf{U}^\imath .

1.3. Main results. As explained in §1.2, we first generalize Clark's result on $\mathfrak{gl}(m|1)$ to $\mathfrak{gl}(m|n)$ setting (see Theorem 3.5) by checking the braid group operators are indeed algebra isomorphisms between the same algebra with different presentations. The braid group operators are usually denoted by T_j in this paper and are shown to satisfy the braid group relations.

Then we use the braid group operators to define a *quantum supersymmetric pair* $(\mathbf{U}, \mathbf{U}^\iota)$ of type AIII. More specifically, starting with a diagram $\tilde{\Gamma}$ as in (4.1) satisfying (4.2) with an index set $I = I_\circ \cup I_\bullet$. Ignoring the parity, we can imitate the non-super case to define a longest element w_\bullet associated to the Weyl group of the Levi subalgebra associated to I_\bullet . Applying w_\bullet to $\tilde{\Gamma}$ we obtain another diagram Γ which also satisfies (4.2) and is indexed by I as well. The algebra $\mathbf{U}(\Gamma)$ (resp. $\mathbf{U}(\tilde{\Gamma})$) is generated by $E_j, F_j, K_j^{\pm 1}$ (resp. $\tilde{E}_j, \tilde{F}_j, \tilde{K}_j^{\pm 1}$). In particular, the ι quantum group \mathbf{U}^ι in the pair $(\mathbf{U}(\Gamma), \mathbf{U}^\iota)$ is defined in terms of generators E_j, F_j ($j \in I_\bullet$),

$$B_j = F_j + \varsigma_j T_{w_\bullet}(\tilde{E}_{\tau_j})K_j^{-1}, \quad \text{for } j \in I_\circ$$

and certain Cartan elements.

Note again that a reader can consider the set I_\bullet to consist of even simple roots only in order to avoid the complexity as in this case we already obtain sufficiently new results. In particular, when I_\bullet consists of even simple roots only, Γ and $\tilde{\Gamma}$ coincide and we only need to use Lusztig's braid group actions in the definition of B_j .

We also establish the coideal subalgebra property of \mathbf{U}^ι in \mathbf{U} (see Proposition 4.6) generalizing the non-super constructions (cf. [Let99, Let02, Ko12]). The original methods in [Let99, Let02, Ko12] does not directly apply to the super case and we give a new proof for the super type AIII case.

We then establish new quantum ι Serre relations (Proposition 4.12) by applying the projection technique [Let99, Let02, Ko12] and obtain a natural filtration on \mathbf{U}^ι (Proposition 4.14), whose associated graded algebra $gr\mathbf{U}^\iota$ is essentially isomorphic to the negative half of the quantum supergroup \mathbf{U} (modulo the Cartan elements).

After establishing the quantum supersymmetric pair $(\mathbf{U}, \mathbf{U}^\iota)$, we proceed to establish a multi-parameter ι Schur duality of type AIII between \mathbf{U}^ι and the Hecke algebra of type B, cf. [BW18a, BWW18, SW21].

Let \mathbb{W} be the natural representation of \mathbf{U} . We show that $\mathbb{W}^{\otimes d}$ possesses a right \mathcal{H}_{B_d} -module structure (Proposition 5.1) and a left \mathbf{U}^ι -module structure via the comultiplication in the same time. Under the assumption on the parameters (5.8), we show that the actions of \mathbf{U}^ι and \mathcal{H}_{B_d} commute with each other and form a double centralizer property; see Theorem 5.8.

In [CL22], a duality between type AIII ι Schur superalgebras for a very special Sataki diagram where $I_\bullet = \emptyset$ and the Hecke algebra of type B was studied. Our \mathbf{U}^ι -actions indeed factor through the corresponding ι Schur superalgebra when the corresponding diagrams are the same.

For the construction of the quasi K -matrix Υ and the K -matrix \mathcal{T} we impose one more condition (6.1); i.e. I_\bullet consists of even simple roots only. Following intertwining relations from [WZ22] and strategies from [BW18a, AV20, Ko21], we construct the quasi K -matrix $\Upsilon = \sum \Upsilon_\mu$ with $\Upsilon_0 = 1$ and $\Upsilon_\mu \in \mathbf{U}_\mu^+$.

With the quasi K -matrix Υ being established, we impose one more constraint on the parameters and construct a unique involutive bar involution ψ_ι on \mathbf{U}^ι (Corollary 6.19), which is a super analogue of the bar involution established in [BW18b, Ko21]. The bar involution ψ on the quantum supergroup \mathbf{U} and ψ_ι on \mathbf{U}^ι is intertwined by Υ such that $\psi_\iota(x)\Upsilon = \Upsilon\psi(x)$, $\forall x \in \mathbf{U}^\iota$.

At last, following the construction in [BW18b], we formulate the K -matrix \mathcal{T} . We show that \mathcal{T} induces an \mathbf{U}^ι -isomorphism on \mathbb{W} and we compute that its action on \mathbb{W} coincides with the H_0 -action. Thus the H_0 -action on $\mathbb{W}^{\otimes d}$ is realized by $\mathcal{T} \otimes 1^{\otimes d-1}$.

1.4. Organization. This paper is organized as follows. In §2 we establish a basic construction of the quantum supergroup \mathbf{U} of type A associated to any Dynkin diagram. In §3 we study the odd reflections of the underlying Lie superalgebra and extend Clark's braid group operators to the general case. Then we define the first family of quantum supersymmetric pair of type AIII in §4 and study new ι Serre relations and obtain a natural filtration. In §5 we construct the bimodule structure on the tensor product of the natural representation of \mathbf{U} and prove the double centralizer property. In §6 we construct the quasi K -matrix of $(\mathbf{U}, \mathbf{U}^2)$ and the bar involution on \mathbf{U} and give an example of Υ in closed formulas. In §7 we further establish the K -matrix \mathcal{T} and use it to realize the H_0 -action on the natural representation of \mathbf{U} .

Acknowledgement. We would like to express our great gratitude to Weiqiang Wang for his many insightful advice and extremely helpful discussions. We also thank Weinan Zhang for his useful suggestions. YS is supported by Graduate Research Assistantship from Wang's NSF grant (DMS-2001351) and a semester fellowship from University of Virginia.

2. QUANTUM SUPERGROUP OF TYPE A

In this section we recall the basic definition of a type A Lie superalgebra and introduce the notation of a Quantum supergroup associated to any Dynkin diagram of type A.

2.1. The general linear Lie superalgebra. We adopt basic notations from [CW12]. Let $V = V_{\bar{0}} \oplus V_{\bar{1}}$ be a vector superspace such that $End(V)$ is an associative superalgebra. Then $End(V)$, equipped with the supercommutator, forms a Lie superalgebra, called the general linear Lie superalgebra and is denoted by $\mathfrak{gl}(V)$ or $\mathfrak{gl}(\mathfrak{m}|\mathfrak{n})$ where $dim V_{\bar{0}} = \mathfrak{m}$, $dim V_{\bar{1}} = \mathfrak{n}$.

Choose bases for $V_{\bar{0}}$ and $V_{\bar{1}}$ such that they combine to a homogeneous basis for V . We will make it a convention to parameterize such a basis by the set

$$I(\mathfrak{m}|\mathfrak{n}) = \{\bar{1}, \dots, \bar{\mathfrak{m}}, \underline{1}, \dots, \underline{\mathfrak{n}}\}$$

with total order

$$\bar{1} < \dots < \bar{\mathfrak{m}} < 0 < \underline{1} < \dots < \underline{\mathfrak{n}}.$$

Here 0 is inserted for convention. With such an ordered basis, $\mathfrak{gl}(\mathfrak{m}|\mathfrak{n})$ can be realized as $(\mathfrak{m} + \mathfrak{n}) \times (\mathfrak{m} + \mathfrak{n})$ complex matrices of the block form

$$(2.1) \quad g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

where a, b, c and d are respectively $\mathfrak{m} \times \mathfrak{m}$, $\mathfrak{m} \times \mathfrak{n}$, $\mathfrak{n} \times \mathfrak{m}$ and $\mathfrak{n} \times \mathfrak{n}$ matrices.

The even subalgebra $\mathfrak{g}_{\bar{0}}$ consists of matrices of the form 2.1 with $b = c = 0$, while the odd subspace $\mathfrak{g}_{\bar{1}}$ consists of those with $a = d = 0$. For each element g , we define the supertrace to be

$$str(g) = tr(a) - tr(d).$$

The supertrace str on the general linear Lie superalgebra gives rise to a non-degenerate supersymmetric bilinear form

$$(\cdot, \cdot) : \mathfrak{gl}(\mathfrak{m}|\mathfrak{n}) \times \mathfrak{gl}(\mathfrak{m}|\mathfrak{n}) \rightarrow \mathbb{C}, \quad (x, y) = str(xy).$$

2.2. Root system of $\mathfrak{gl}(m|n)$. Let $\mathfrak{g} = \mathfrak{gl}(m|n)$ and \mathfrak{h} be the Cartan subalgebra of diagonal matrices.

Restricting the supertrace to the Cartan subalgebra \mathfrak{h} , we obtain a non-degenerate symmetric bilinear form on it. Denote by $\{\epsilon_{\bar{i}}, \epsilon_{\underline{j}}\}_{i,j}$ the basis of \mathfrak{h}^* dual to $\{E_{\bar{ii}}, E_{\underline{j}j}\}_{i,j}$.

Its root system $\Phi = \Phi_{\bar{0}} \oplus \Phi_{\bar{1}}$ is given by

$$(2.2) \quad \begin{aligned} \Phi_{\bar{0}} &= \{\epsilon_a - \epsilon_b \mid a \neq b \in I(m|n), a, b > 0 \text{ or } a, b < 0\}, \\ \Phi_{\bar{1}} &= \{\pm(\epsilon_a - \epsilon_b) \mid a < 0 < b\}. \end{aligned}$$

A fundamental system of $\mathfrak{gl}(m|n)$ consists of $m+n-1$ roots

$$\epsilon_{i_1} - \epsilon_{i_2}, \dots, \epsilon_{i_{m+n-1}} - \epsilon_{i_{m+n}},$$

where $\{i_1, \dots, i_{m+n}\} = I(m|n)$. We denote even simple roots by \bullet and odd simple roots by \otimes . Then the corresponding Dynkin diagram is of the form

$$(2.3) \quad \begin{array}{ccccccc} \circ & \text{---} & \circ & \text{---} & \dots & \text{---} & \circ & \text{---} & \circ \\ \epsilon_{i_1} - \epsilon_{i_2} & & \epsilon_{i_2} - \epsilon_{i_3} & & & & \epsilon_{i_{m+n-1}} - \epsilon_{i_{m+n}} & & \end{array}$$

where \circ is either \bullet or \otimes .

Example 2.1. The standard Dynkin diagram is given by

$$\begin{array}{ccccccc} \bullet & \text{---} & \bullet & \text{---} & \dots & \text{---} & \otimes & \text{---} & \dots & \text{---} & \bullet & \text{---} & \bullet \\ \epsilon_{\bar{1}} - \epsilon_{\bar{2}} & & \epsilon_{\bar{2}} - \epsilon_{\bar{3}} & & & & \epsilon_{\bar{m}} - \epsilon_{\underline{1}} & & & & \epsilon_{\underline{n-1}} - \epsilon_{\underline{n}} & & \end{array}$$

Given a Dynkin diagram of the form (2.3). Let

$$\Pi = \{\alpha_j = \epsilon_{i_j} - \epsilon_{i_{j+1}} \mid j = 1, \dots, m+n-1\}$$

denote the set of simple roots with the index set $I = \{1, \dots, m+n-1\}$. We see that I is a disjoint union of two subsets $I = I_{\bar{0}} \cup I_{\bar{1}}$. Let p be the parity function on I such that

$$p(k) = \begin{cases} 0 & \text{if } k \in I_{\bar{0}}, \\ 1 & \text{if } k \in I_{\bar{1}}. \end{cases}$$

We define the weight lattice $P = \mathbb{Z}\Pi = \bigoplus_{b \in I(m|n)} \mathbb{Z}\epsilon_b$ while the symmetric bilinear form on P is given by

$$(2.4) \quad (\epsilon_a, \epsilon_{a'}) = \begin{cases} 1 & \text{if } \bar{1} \leq a = a' \leq \bar{m}, \\ -1 & \text{if } \underline{1} \leq a = a' \leq \underline{n}, \\ 0 & \text{else.} \end{cases}$$

We define the coweight lattice $P^\vee = \bigoplus_{b \in I(m|n)} \mathbb{Z}\epsilon_b^\vee$ and we have the pairing $\langle \cdot, \cdot \rangle : P \times P \rightarrow \mathbb{Z}$ with $\langle \epsilon_a^\vee, \epsilon_b \rangle = \delta_{a,b}$. Then $\Pi^\vee = \{h_j \mid j \in I\}$, the set of simple coroots, is given by the following data:

$$\begin{aligned} \langle h_j, \alpha_j \rangle &= 2, & j \in I_{\bar{0}}, \\ \langle h_j, \alpha_j \rangle &= 0, & j \in I_{\bar{1}}. \end{aligned}$$

More specifically, we have

$$(2.5) \quad h_j = \epsilon_{i_j}^\vee - (-1)^{p(j)} \epsilon_{i_{j+1}}^\vee.$$

We suppose that there are non-zero integers ℓ_j such that

$$(2.6) \quad \ell_j \langle h_j, \lambda \rangle = (\alpha_j, \lambda) \text{ for any } \lambda \in P.$$

2.3. Quantum supergroup of type A. Quantum supergroups have been defined in several articles such as [KT91] and [Ya94]. Following [Mi06], we define a quantum supergroup associated to any fixed Dynkin diagram of the form (2.3).

Denote the quantum integers and quantum binomial coefficients by, for $a \in \mathbb{Z}, k \in \mathbb{N}$,

$$[a] = \frac{q^a - q^{-a}}{q - q^{-1}}, \quad \begin{bmatrix} a \\ k \end{bmatrix} = \frac{[a][a-1] \cdots [a-k+1]}{[k]}.$$

It will be convenient for us to introduce the following notation. We will say $i, j \in I$ are *connected* if $i = j \pm 1$ and write $i \sim j$. Likewise, we say *not connected* if $i \neq j, j \pm 1$ and write $i \not\sim j$.

Let $K_i = q^{\ell_i h_i}$, we recall the definition of $\mathbf{U}_q(\mathfrak{gl}(m|n))$ to be the unital associative algebra over $\mathbb{Q}(q)$ with generators q^h ($h \in P^\vee$), E_i, F_i ($i \in I$) which satisfy the following defining relations:

$$(2.7) \quad \begin{aligned} (R1) \quad & q^h = 1, \quad \text{for } h = 0, \\ (R2) \quad & q^{h_1} q^{h_2} = q^{h_1+h_2}, \\ (R3) \quad & q^h E_j = q^{\langle h, \alpha_j \rangle} E_j q^h \quad \text{for } j \in I, \\ (R4) \quad & q^h F_j = q^{-\langle h, \alpha_j \rangle} F_j q^h, \quad \text{for } j \in I, \\ (R5) \quad & [E_j, F_k] = E_j F_k - (-1)^{p(j)p(k)} F_k E_j = \delta_{j,k} \frac{K_j - K_j^{-1}}{q^{\ell_i} - q^{-\ell_i}}, \quad \text{for } j, k \in I, \\ (R6) \quad & E_j^2 = F_j^2 = 0, \quad \text{for } j \in I_{\overline{1}}, \\ (R7) \quad & E_j E_k = (-1)^{p(j)p(k)} E_k E_j, \quad F_j F_k = (-1)^{p(j)p(k)} F_k F_j, \quad \text{for } j \not\sim k, \\ (R8) \quad & E_j^2 E_k - [2] E_j E_k E_j + E_k E_j^2 = 0, \quad \text{for } j \sim k, p(j) = 0, \\ (R9) \quad & F_j^2 F_k - [2] F_j F_k F_j + F_k F_j^2 = 0, \quad \text{for } j \sim k, p(j) = 0, \\ (R10) \quad & S_{p(k), p(\ell)}(E_k, E_j, E_\ell) = 0, \quad \text{for } k \sim j \sim \ell, k < \ell, p(j) = 1, \\ (R11) \quad & S_{p(k), p(\ell)}(F_k, F_j, F_\ell) = 0, \quad \text{for } k \sim j \sim \ell, k < \ell, p(j) = 1. \end{aligned}$$

where $S_{t_1, t_2}(x_1, x_2, x_3) \in \mathbb{Q}(q)\langle x_1, x_2, x_3 \rangle$ is the polynomial in three non-commuting variables for $t_1, t_2 \in \{\overline{0}, \overline{1}\}$ given by

$$(2.8) \quad \begin{aligned} S_{t_1, t_2}(x_1, x_2, x_3) = & [2]x_2x_3x_1x_2 - [((-1)^{t_1}x_2x_3x_2x_1 + (-1)^{t_1+t_1t_2}x_1x_2x_3x_2) \\ & + ((-1)^{t_1t_2+t_2}x_2x_1x_2x_3 + (-1)^{t_2}x_3x_2x_1x_2)]. \end{aligned}$$

The super q -Serre relations (R8)–(R11) can be found in [CHW16, Proposition 2.7] (cf. [C16, §4.3]).

Moreover, let $q_j := q^{\frac{(\alpha_j, \alpha_j)}{2}}$, we define maps $\sigma, \wp, \text{op}, \bar{\cdot}$ on $\mathbf{U}_q(\mathfrak{gl}(\mathfrak{m}|\mathfrak{n}))$ satisfying:

$$(2.9) \quad \begin{aligned} \sigma(E_j) &= E_j, & \sigma(F_j) &= F_j, & \sigma(K_j) &= (-1)^{p(j)} K_j^{-1}, & \sigma(xy) &= \sigma(y)\sigma(x), \\ \wp(E_j) &= q_j K_j F_j, & \wp(F_j) &= q_j^{-1} E_j K_j^{-1}, & \wp(K_j) &= K_j, & \wp(xy) &= \wp(y)\wp(x), \\ \text{op}(E_j) &= E_j, & \text{op}(F_j) &= F_j, & \text{op}(K_j) &= K_j, & \text{op}(xy) &= \text{op}(y)\text{op}(x), \\ \overline{E_j} &= E_j, & \overline{F_j} &= F_j, & \overline{K_j} &= K_j^{-1}, & \overline{q} &= q^{-1}, & \overline{xy} &= \overline{x} \cdot \overline{y}. \end{aligned}$$

In general, $\mathbf{U}_q(\mathfrak{gl}(\mathfrak{m}|\mathfrak{n}))$ is a Hopf superalgebra (cf. [C16, Lemma 2.1]) but not a Hopf algebra. In order to give $\mathbf{U}_q(\mathfrak{gl}(\mathfrak{m}|\mathfrak{n}))$ a Hopf algebra structure, we define an involutive operator ϱ of parity 0 on it by

$$(2.10) \quad \varrho(q^h) = q^h, \quad \varrho(E_j) = (-1)^{p(j)} E_j \text{ and } \varrho(F_j) = (-1)^{p(j)} F_j, \quad \forall j \in I.$$

Let

$$\mathbf{U} = \mathbf{U}_q(\mathfrak{gl}(\mathfrak{m}|\mathfrak{n})) \oplus \mathbf{U}_q(\mathfrak{gl}(\mathfrak{m}|\mathfrak{n}))\varrho.$$

Then \mathbf{U} is the algebra with the additional multiplication law given by

$$(2.11) \quad \varrho^2 = 1, \quad \varrho^{-1}x\varrho = \varrho(x) \text{ for any } x \in \mathbf{U}_q(\mathfrak{gl}(\mathfrak{m}|\mathfrak{n})).$$

We naturally extend the maps in (2.9) from $\mathbf{U}_q(\mathfrak{gl}(\mathfrak{m}|\mathfrak{n}))$ to \mathbf{U} by setting

$$\sigma(\varrho) = \wp(\varrho) = \text{op}(\varrho) = \overline{\varrho} = \varrho.$$

Next we want to establish the Hopf algebra structure of \mathbf{U} . Let U^ϱ denote the algebra generated by the same set of generators as \mathbf{U} and with the relations (R1)–(R7) in (2.7) and (2.11). Then we have

Proposition 2.2. *\mathbf{U} is a Hopf algebra comultiplication Δ , counit ϵ , antipode S are as follows,*

$$(2.12) \quad \begin{aligned} \Delta(\varrho) &= \varrho \otimes \varrho, \\ \Delta(q^h) &= q^h \otimes q^h \quad \text{for } h \in P^\vee, \\ \Delta(E_j) &= E_j \otimes 1 + \varrho^{p(j)} K_j \otimes E_j \quad \text{for } j \in I, \\ \Delta(F_j) &= F_j \otimes K_j^{-1} + \varrho^{p(j)} \otimes F_j \quad \text{for } j \in I, \\ \epsilon(\varrho) &= \epsilon(q^h) = 1, \text{ for } h \in P^\vee, \quad \epsilon(E_j) = \epsilon(F_j) = 0, \quad \text{for } j \in I, \\ S(\varrho) &= \varrho, \quad S(q^h) = q^{-h} \quad \text{for } h \in P^\vee, \\ S(F_j) &= -\varrho^{p(j)} F_j K_j, \quad S(E_j) = -\varrho^{p(j)} K_j^{-1} E_j \quad \text{for } j \in I. \end{aligned}$$

Proof. We only check for Δ to be an algebra homomorphism as it's straightforward to check for ϵ and S .

The relations (R1)–(R4) are easy to check. To verify the relation (R5), first suppose $j = k \in I$. Then

$$\begin{aligned} & [\Delta(E_j), \Delta(F_j)] \\ &= E_j F_j \otimes K_j^{-1} + E_j \varrho^{p(j)} \otimes F_j + \varrho^{p(j)} K_j F_j \otimes E_j K_j^{-1} + K_j \otimes E_j F_j \\ &\quad - (-1)^{p(j)} [F_j E_j \otimes K_j^{-1} + F_j \varrho^{p(j)} K_j \otimes K_j^{-1} E_j + \varrho^{p(j)} E_j \otimes F_j + K_j \otimes F_j E_j] \\ &= \frac{(K_j - K_j^{-1}) \otimes K_j^{-1} + K_j \otimes (K_j - K_j^{-1})}{q^{\ell_j} - q^{-\ell_j}} = \frac{K_j \otimes K_j - K_j^{-1} \otimes K_j^{-1}}{q^{\ell_j} - q^{-\ell_j}}. \end{aligned}$$

Now if $j \neq k$, we have

$$\begin{aligned} & [\Delta(E_j), \Delta(F_k)] \\ &= E_j F_k \otimes K_k^{-1} + E_j \varrho^{p(k)} \otimes F_k + \varrho^{p(j)} K_j F_k \otimes E_j K_k^{-1} + \varrho^{p(k)} \varrho^{p(j)} K_j \otimes E_j F_k \\ &\quad - (-1)^{p(j)p(k)} [F_k E_j \otimes K_k^{-1} + F_k \varrho^{p(j)} K_j \otimes K_k^{-1} E_j \\ &\quad\quad + \varrho^{p(k)} E_j \otimes F_k + \varrho^{p(k)} \varrho^{p(j)} K_j \otimes F_k E_j] \\ &= (E_j F_k - (-1)^{p(j)p(k)} F_k E_j) \otimes K_k^{-1} \\ &\quad + (\varrho^{p(j)} K_j F_k \otimes E_j K_k^{-1} - (-1)^{p(j)p(k)} F_k \varrho^{p(j)} K_j \otimes K_k^{-1} E_j) \\ &\quad + (E_j \varrho^{p(k)} \otimes F_k - (-1)^{p(j)p(k)} \varrho^{p(k)} E_j \otimes F_k) \\ &\quad + \varrho^{p(k)} \varrho^{p(j)} K_j \otimes (E_j F_k - (-1)^{p(j)p(k)} F_k E_j) \\ &= 0. \end{aligned}$$

To verify the relation (R6), suppose $j \in I_{\bar{1}}$, then we have

$$\Delta(E_j)^2 = E_j^2 \otimes 1 + E_j \varrho^{-1} K_j \otimes E_j + \varrho^{-1} K_j E_j \otimes E_j + K_j^2 \otimes E_j^2 = 0.$$

To verify the relation (R7), suppose $j \approx k$, then we have

$$\begin{aligned} & \Delta(E_j) \Delta(E_k) \\ &= E_j E_k \otimes 1 + E_j \varrho^{p(k)} K_k \otimes E_k + \varrho^{p(j)} K_j E_k \otimes E_j + \varrho^{p(k)} \varrho^{p(j)} K_j K_k \otimes E_j E_k \\ &= (-1)^{p(j)p(k)} \Delta(E_k) \Delta(E_j). \end{aligned}$$

Similarly we have

$$\Delta(F_j) \Delta(F_k) = (-1)^{p(j)p(k)} \Delta(F_k) \Delta(F_j).$$

□

Define

$$(2.13) \quad \begin{aligned} x_{j,k}^+ &= E_j^2 E_k - (q + q^{-1}) E_j E_k E_j + E_k E_j^2, \\ x_{j,k}^- &= F_j^2 F_k - (q + q^{-1}) F_j F_k F_j + F_k F_j^2. \end{aligned}$$

Moreover, let \mathcal{G} denote the ideal of U^e generated by $\{x_{j,k}^\pm \mid p(j) = 0, j \sim k\}$ and

$$\{S_{p(k),p(\ell)}(E_k, E_j, E_\ell), S_{p(k),p(\ell)}(F_k, F_j, F_\ell) \mid p(j) = 1, k \sim j \sim \ell\}.$$

Then we see that $\mathbf{U} \cong U^e / \mathcal{G}$. The next two lemmas are essential to prove the Hopf algebra structure on \mathbf{U} .

Lemma 2.3. *Let $j \sim k$ and $p(j) = 0$, then we have*

$$\Delta(x_{j,k}^{\pm}) \in \mathfrak{G} \otimes U^e + U^e \otimes \mathfrak{G}, \quad S(x_{j,k}^{\pm}) \in \mathfrak{G}.$$

Proof. The proof follows similarly from the proof of [Jan95, Lemma 4.10]. \square

Lemma 2.4. *Let $p(j) = 1$ and $k \sim j \sim \ell$, then we have*

$$\begin{aligned} \Delta(S_{p(k),p(\ell)}(E_k, E_j, E_\ell)) &\in \mathfrak{G} \otimes U^e + U^e \otimes \mathfrak{G}, & S(S_{p(k),p(\ell)}(E_k, E_j, E_\ell)) &\in \mathfrak{G}, \\ \Delta(S_{p(k),p(\ell)}(F_k, F_j, F_\ell)) &\in \mathfrak{G} \otimes U^e + U^e \otimes \mathfrak{G}, & S(S_{p(k),p(\ell)}(F_k, F_j, F_\ell)) &\in \mathfrak{G}. \end{aligned}$$

Proof. We only prove for E 's. We see that

$$(2.14) \quad \begin{aligned} &\Delta(S_{p(k),p(\ell)}(E_k, E_j, E_\ell)) \\ &= S_{p(k),p(\ell)}(E_k, E_j, E_\ell) \otimes 1 + \varrho^{p(k)+p(\ell)} K_j^2 K_\ell K_k \otimes S_{p(k),p(\ell)}(E_k, E_j, E_\ell) + (\clubsuit). \end{aligned}$$

It suffices to prove that the coefficients of terms in (\clubsuit) vanish in (2.14). We take the term

$$K_j E_j E_\ell E_k \varrho \otimes E_j$$

with coefficient A for example. We see that the term $K_j E_j E_\ell E_k \otimes E_j$ appears in

$$\Delta(E_j E_\ell E_k E_j), \quad \Delta(E_j E_\ell E_j E_k) \quad \text{and} \quad \Delta(E_j E_k E_j E_\ell)$$

as A_1, A_2 and A_3 respectively. Note that since $p(j) = 1$, we can assume without loss of generality that $(\alpha_j, \alpha_k) = 1$ while $(\alpha_j, \alpha_\ell) = -1$. Now we compute that

$$\begin{aligned} A_1 &= E_j E_\ell E_k \varrho^{p(j)} K_j \otimes E_j = K_j E_j E_\ell E_k \varrho^{p(j)} \otimes E_j, \\ A_2 &= E_j E_\ell \varrho^{p(j)} K_j E_k \otimes E_j = q K_j E_j E_\ell \varrho^{p(j)} E_k \otimes E_j, \\ A_3 &= E_j E_k \varrho^{p(j)} K_j E_\ell \otimes E_j = q^{-1} K_j E_j E_k \varrho^{p(j)} E_\ell \otimes E_j. \end{aligned}$$

Recall (2.8), then we have

$$A = (q + q^{-1}) - (-1)^{p(k)} (-1)^{p(k)} q - (-1)^{p(\ell)} (-1)^{p(\ell)} q^{-1} = 0.$$

It is similar to show the coefficients of other terms in (2.14) vanish. Thus we conclude that

$$\Delta(S_{p(k),p(\ell)}(E_k, E_j, E_\ell)) \in \mathfrak{G} \otimes U^e + U^e \otimes \mathfrak{G}.$$

On the other hand, one can compute directly that $S(S_{p(k),p(\ell)}(E_k, E_j, E_\ell)) \in \mathfrak{G}$. \square

Proposition 2.5. *\mathbf{U} is a Hopf superalgebra whose comultiplication Δ , counit ϵ , antipode S are defined as in (2.12).*

Proof. Combining Lemma 2.3 and Lemma 2.4, we see that Δ induces a homomorphism

$$U^e/\mathfrak{G} \rightarrow (U^e \otimes U^e)/(U^e \otimes \mathfrak{G} + \mathfrak{G} \otimes U^e) \cong (U^e/\mathfrak{G}) \otimes (U^e/\mathfrak{G}),$$

and S induces an anti-homomorphism

$$\mathbf{U} \cong U^e/\mathfrak{G} \rightarrow U^e/\mathfrak{G} \cong \mathbf{U}.$$

Moreover, it is straightforward to check these maps satisfy the defining relation for \mathbf{U} to be a Hopf algebra. \square

Remark 2.6. The choice of the comultiplication we made here is different from [Mi06]. When \mathfrak{g} is a Lie algebra, this Δ is compatible with the comultiplication in [Lus93].

As usual, we have a triangular decomposition $\mathbf{U} = \mathbf{U}^+ \mathbf{U}^0 \mathbf{U}^-$ where \mathbf{U}^+ (resp. \mathbf{U}^-) denotes the subalgebra of \mathbf{U} generated by E_j (resp. F_j), $j \in I$ and \mathbf{U}^0 denotes the subalgebra of \mathbf{U} generated by $\{K_j^{\pm 1}, \varrho \mid j \in I\}$.

3. BRAID GROUP OPERATORS

In order to define quantum supersymmetric pairs, we need to study the braid group operators on \mathbf{U} . In this section we study the odd reflections and braid group operators associated to them, which is an analogue of Lusztig's braid group operator as in [Lus93].

3.1. Odd reflections. As observed in [C16, §4], fix a simple root α , the braid operator associate to it lifts the action of s_α on the weight data. The main difference between odd and even reflections is that odd reflections change the generalized Cartan matrix while even ones do not.

Note that the fundamental systems of the root system Φ associated to $\mathfrak{gl}(\mathfrak{m}|\mathfrak{n})$ are not conjugated under the Weyl group actions because of the existence of odd roots (cf. [CW12, §1.3.6]). In fact, we have the following lemma,

Lemma 3.1. [CW12, Lemma 1.26] *Let α be an odd simple root of $\mathfrak{gl}(\mathfrak{m}|\mathfrak{n})$ in a positive system Φ^+ . Then,*

$$\Phi_\alpha^+ := \{-\alpha\} \cup \Phi^+ \setminus \{\alpha\}$$

is a new positive system, whose corresponding fundamental system Π_α is given by

$$(3.1) \quad \Pi_\alpha = \{\beta \in \Pi \mid (\beta, \alpha) = 0, \beta \neq \alpha\} \cup \{\beta + \alpha \mid \beta \in \Pi, (\beta, \alpha) \neq 0\} \cup \{-\alpha\}.$$

The process of obtaining Π_α from Π is denoted by s_α and called an odd reflection. When $\beta \in \Pi$ is an even simple root, we abuse the notation s_β to denote the usual reflection associated to β .

For a diagram as in (2.3), we let $s_j := s_{\alpha_j}$, $j \in I$ and we denote by $\mathfrak{S}_{\mathfrak{m}+\mathfrak{n}}$ the Weyl group of $\mathfrak{gl}(\mathfrak{m} + \mathfrak{n})$ ignoring the parity. Essentially, when $p(j) = 1$, the s_j -action can be translated to a transposition between $\epsilon_{i_{j-\frac{1}{2}}}$ and $\epsilon_{i_{j+\frac{1}{2}}}$.

Remark 3.2. By ignoring the parity and focusing on the action on the weight lattice, we observe that the braid group relations still hold for reflections of any parity.

The following lemma tells us how this action changes parities, which in turn allows us to determine the matrix units.

Lemma 3.3. *If $j, k, \ell \in I$ with $j \sim k$ and $j \not\sim \ell$. Then for any $X \in \mathcal{D}_{\mathfrak{m}, \mathfrak{n}}$ we have*

$$(3.2) \quad p_{s_j(X)}(j) = p_X(j), \quad p_{s_j(X)}(k) = p_X(k) + p_X(j), \quad p_{s_j(X)}(\ell) = p_X(\ell).$$

From Lemma (3.3) we see that even reflections will not change the parity of any simple root while odd reflections change the parities of the ones adjacent to it.

Example 3.4.

$$\begin{array}{ccc} \otimes \text{---} \otimes & \xrightarrow{s_1} & \otimes \text{---} \bullet \\ \epsilon_{\overline{1}} - \epsilon_{\underline{1}} & & \epsilon_{\underline{1}} - \epsilon_{\overline{1}} \end{array} \quad \begin{array}{ccc} \epsilon_{\underline{1}} - \epsilon_{\overline{2}} & & \epsilon_{\overline{1}} - \epsilon_{\underline{2}} \end{array}$$

More precisely, for any two fundamental systems Π and Π' of a basic Lie superalgebra of any classical type, there exists a sequence consisting of even and odd reflections s_1, \dots, s_k such that $s_1 \cdots s_k(\Pi) = \Pi'$. (cf. [CW12])

3.2. Braid group operators. Let $\mathcal{D}_{m,n}$ denote the set of all the possible Dynkin diagram for $\mathfrak{gl}(m|n)$. Now for each $X \in \mathcal{D}_{m,n}$, we can associate a quantum enveloping algebra $\mathbf{U}(X)$ with generators $E_{X,i}, F_{X,i}, K_{X,i}^{-1}$ and ϱ_X as in §2.1. Equipped with this family of algebras, we now introduce the braid group operator.

Theorem 3.5. (Compare [C16, Theorem 4.5]) *Let $i \in I$, $X \in \mathcal{D}_{m,n}$, $e = \pm 1$ and set $Y = s_i(X)$. There exists a $\mathbb{Q}(q)$ -linear algebra isomorphisms $T'_{i,e}, T''_{i,e} : \mathbf{U}(X) \rightarrow \mathbf{U}(Y)$ satisfying*

$$(3.3) \quad T'_{i,-e}(E_{X,j}) = \begin{cases} -(-1)^{p_Y(i)} K_{Y,i}^{-e} F_{Y,i}, & \text{if } j = i, \\ E_{Y,j} E_{Y,i} - (-1)^{p_Y(i)p_Y(j)} q^{e(\alpha_{Y,i}, \alpha_{Y,j})} E_{Y,i} E_{Y,j} & \text{if } j \sim i, \\ E_{Y,j} & \text{if } j \not\sim i. \end{cases}$$

$$(3.4) \quad T'_{i,-e}(F_{X,j}) = \begin{cases} -(-1)^{p_Y(i)} E_{Y,i} K_{Y,i}^e, & \text{if } j = i, \\ F_{Y,i} F_{Y,j} - (-1)^{p_Y(i)p_Y(j)} q^{-e(\alpha_{Y,i}, \alpha_{Y,j})} F_{Y,j} F_{Y,i} & \text{if } j \sim i, \\ F_{Y,j} & \text{if } j \not\sim i. \end{cases}$$

$$(3.5) \quad T'_{i,-e}(K_{X,j}) = \begin{cases} (-1)^{p_Y(i)} K_{Y,i}^{-1}, & \text{if } j = i, \\ (-1)^{p_Y(i)p_Y(j)} K_{Y,i} K_{Y,j} & \text{if } j \sim i, \\ K_{Y,j} & \text{if } j \not\sim i. \end{cases}$$

$$(3.6) \quad T'_{i,-e}(\varrho_X) = \varrho_Y.$$

and

$$(3.7) \quad T''_{i,e}(E_{X,j}) = \begin{cases} -F_{Y,i} K_{Y,i}^e, & \text{if } j = i, \\ E_{Y,i} E_{Y,j} - (-1)^{p_Y(i)p_Y(j)} q^{e(\alpha_{Y,i}, \alpha_{Y,j})} E_{Y,j} E_{Y,i} & \text{if } j \sim i, \\ E_{Y,j} & \text{if } j \not\sim i. \end{cases}$$

$$(3.8) \quad T''_{i,e}(F_{X,j}) = \begin{cases} -K_{Y,i}^{-e} E_{Y,i}, & \text{if } j = i, \\ F_{Y,j} F_{Y,i} - (-1)^{p_Y(i)p_Y(j)} q^{-e(\alpha_{Y,i}, \alpha_{Y,j})} F_{Y,i} F_{Y,j} & \text{if } j \sim i, \\ F_{Y,j} & \text{if } j \not\sim i. \end{cases}$$

$$(3.9) \quad T''_{i,e}(K_{Y,j}) = \begin{cases} (-1)^{p_Y(i)} K_{Y,i}^{-1}, & \text{if } j = i, \\ (-1)^{p_Y(i)p_Y(j)} K_{Y,i} K_{Y,j} & \text{if } j \sim i, \\ K_{Y,j} & \text{if } j \not\sim i. \end{cases}$$

$$(3.10) \quad T''_{i,e}(\varrho_X) = \varrho_Y.$$

In the remainder of this subsection we will prove Theorem 3.5 and then verify the braid relations. To do so succinctly, recall σ , $\circ p$ and $\bar{\cdot}$ from (2.9), we observe that

$$(3.11) \quad T'_{i,-e} = \sigma T''_{i,e} \sigma, \quad T'_{i,-e} = \bar{\cdot} T'_{i,e} \bar{\cdot} = \circ p T'_{i,e} \circ p, \quad T''_{i,e} = (T'_{i,-e})^{-1}.$$

One can check these identities on the generators. Thus to prove Theorem 3.5, it suffices to prove in the case of $T'_{i,1}$. More specifically, We need to verify that the images of the generators of $\mathbf{U}(X)$ under $T'_{i,1}$ satisfy the relations in (2.7). In order to keep the proof readable, we will break up the verification into lemmas and recall a few results from [C16]. Since the

calculations get quite involved, we will drop the subscripts on the generators of $\mathbf{U}(Y)$ for the sake of readability. Moreover, from now on we always omit the subscript of ϱ since it's clear from the context where it acts on.

First we take a look at (2.10), we have

Lemma 3.6. *If $j, k \in I$, then*

$$\begin{aligned}\varrho T'_{j,-1}(E_{X,k})\varrho^{-1} &= T'_{j,-1}(\varrho(E_{X,k})) = (-1)^{p_X(k)}T'_{j,-1}(E_{X,k}), \\ \varrho T'_{j,-1}(F_{X,k})\varrho^{-1} &= T'_{j,-1}(\varrho(F_{X,k})) = (-1)^{p_X(k)}T'_{j,-1}(F_{X,k}), \\ \varrho T'_{j,-1}(K_{X,k})\varrho^{-1} &= T'_{j,-1}(\varrho(K_{X,k})) = (-1)^{p_X(k)}T'_{j,-1}(K_{X,k}).\end{aligned}$$

Proof. We prove for E and the other two are similar.

When $j = k$ or $j \approx k$, we have $p(k) = p_X(k)$ and hence

$$\varrho T_j(E_{X,k})\varrho^{-1} = (-1)^{p(k)}T_j(E_{X,k}) = (-1)^{p_X(k)}T_j(E_{X,k}).$$

When $j \sim k$, we have $p(k) = p_X(j) + p_X(k)$ and thus $p(k) + p(j) = p_X(k)$. Hence

$$\varrho T_j(E_{X,k})\varrho^{-1} = (-1)^{p(k)+p(j)}T_j(E_{X,k}) = (-1)^{p_X(k)}T_j(E_{X,k}).$$

This proves the lemma. \square

Recall the defining relations of \mathbf{U} from (2.7). The relations (R1)–(R4) can be verified directly. For the relation (R6), we have

Lemma 3.7. [C16, Lemma 4.7] *If $j, k \in I$ such that $p_X(k) = 1$, then*

$$T'_{j,-1}(E_{X,k})^2 = T'_{j,-1}(F_{X,k})^2 = 0.$$

To verify the relation (R5), we split into two cases.

Lemma 3.8. [C16, Lemma 4.9] *If $j = k \in I$, then*

$$\begin{aligned}& T'_{j,-1}(E_{X,k})T'_{j,-1}(F_{X,k}) - (-1)^{p_X(k)}T'_{j,-1}(F_{X,k})T'_{j,-1}(E_{X,k}) \\ &= \frac{T'_{j,-1}(K_{X,k}) - T'_{j,-1}(K_{X,k})^{-1}}{q^{\ell_k} - q^{-\ell_k}}\end{aligned}$$

To verify the relation (R5) when $k \neq l$, note that we need the following lemma.

Lemma 3.9. *If $k = j - 1$ and $\ell = j + 1$, then we have*

$$(3.12) \quad p_X(k)p_X(\ell) + p(k)p(\ell) + p(j)p(\ell) + p(j)p(k) \equiv p(j) \pmod{2}.$$

Proof. This lemma follows from (3.3) directly. \square

Lemma 3.10. (Compare [C16, Lemma 4.8]) *If $j, k, \ell \in I$ with $k \neq \ell$, then*

$$T'_{j,-1}(E_{X,k})T'_{j,-1}(F_{X,\ell}) = (-1)^{p_X(k)p_X(\ell)}T'_{j,-1}(F_{X,\ell})T'_{j,-1}(E_{X,k})$$

Proof. Let $c_{k,\ell} = T'_{j,-1}(E_{X,k})T'_{j,-1}(F_{X,\ell}) - (-1)^{p_X(k)p_X(\ell)}T'_{j,-1}(F_{X,\ell})T'_{j,-1}(E_{X,k})$. We want to prove $c_{k,\ell} = 0$ for all $k \neq \ell$.

If one of them is not connected to j , let's say $j \approx k$, then $T'_{j,-1}(E_{X,k}) = E_k$ and $p(j) = p_X(j)$. On the other hand, $T'_{j,-1}(E_{X,\ell})$ is a polynomial in the elements K_j, F_j, F_ℓ and F_j with

$p(T'_{j,-1}(F_{X,\ell})) = p_X(\ell)$. Since E_k super-commutes with all of those elements, the statement follows.

The remaining cases are when k and ℓ are both either equal or connected to j .

The case when one of them connects with j and the other equals to j has been verified in [C16, Lemma 4.8].

When k and ℓ are both connected to j , without loss of generality, we assume that $k = j - 1$ and $\ell = j + 1$. Using (3.12) and the relation (R5) in (2.7) repeatedly we get

$$\begin{aligned}
c_{k,\ell} &= (E_k E_j - (-1)^{p(j)p(k)} q^{(\alpha_j, \alpha_k)} E_j E_k) (F_j F_\ell - (-1)^{p(j)p(\ell)} q^{-(\alpha_j, \alpha_\ell)} F_\ell F_j) \\
&\quad - (-1)^{p_X(k)p_X(\ell)} (F_j F_\ell - (-1)^{p(j)p(\ell)} q^{-(\alpha_j, \alpha_\ell)} F_\ell F_j) \\
&\quad \quad (E_k E_j - (-1)^{p(j)p(k)} q^{(\alpha_j, \alpha_k)} E_j E_k) \\
&= E_k [E_j, F_j] F_\ell - q^{(\alpha_j, \alpha_k)} [E_j, F_j] E_k F_\ell - q^{-(\alpha_j, \alpha_\ell)} E_k F_\ell [E_j, F_j] \\
&\quad + (-1)^{p(k)p(\ell)} q^{(\alpha_j, \alpha_k) - (\alpha_j, \alpha_\ell)} F_\ell [E_j, F_j] E_k \\
&= \frac{1}{q^{\ell_j} - q^{-\ell_j}} E_k ((1 - q^{2(\alpha_j, \alpha_k)} - 1 + q^{2(\alpha_j, \alpha_k)}) K_j \\
&\quad - (1 - 1 - q^{-2(\alpha_j, \alpha_\ell)} + q^{-2(\alpha_j, \alpha_\ell)}) K_j^{-1}) F_\ell = 0.
\end{aligned}$$

This proves the lemma. \square

We remark that [C16, Lemma 4.8] is insufficient to prove Lemma 3.10 as it only deals with $\mathfrak{gl}(m|1)$ cases. Now we need to verify the various Serre relations.

The next lemma checks the relation (R7) for E , the proof for F is similar.

Lemma 3.11. *If $j, k, \ell \in I$ such that $k \approx \ell$, then*

$$T'_{j,-1}(E_{X,k}) T'_{j,-1}(E_{X,\ell}) = (-1)^{p_X(k)p_X(\ell)} T'_{j,-1}(E_{X,\ell}) T'_{j,-1}(E_{X,k}).$$

Proof. If either k or ℓ is not connected to j , we are done. So we suppose that $k = j - 1$ and $\ell = j + 1$. We break the proof into two cases:

(Case-1) Assume $p_X(j) = 0$. Observe that in this case we must have $(\alpha_j, \alpha_k) = (\alpha_j, \alpha_\ell)$. Without loss of generality we assume $(\alpha_j, \alpha_k) = (\alpha_j, \alpha_\ell) = 1$. Then we have $T'_{j,-1}(E_{X,k}) = E_k E_j - q E_j E_k$ and $T'_{j,-1}(E_{X,\ell}) = E_\ell E_j - q E_j E_\ell$ and thus

$$\begin{aligned}
T'_{j,-1}(E_{X,k}) T'_{j,-1}(E_{X,\ell}) &= E_k E_j E_\ell E_j - q E_j E_k E_\ell E_j - q E_k E_j^2 E_\ell + q^2 E_j E_k E_j E_\ell, \\
T'_{j,-1}(E_{X,\ell}) T'_{j,-1}(E_{X,k}) &= E_\ell E_j E_k E_j - q E_j E_\ell E_k E_j - q E_\ell E_j^2 E_k + q^2 E_j E_\ell E_j E_k.
\end{aligned}$$

First we see that $E_j E_k E_\ell E_j = (-1)^{p(k)p(\ell)} E_j E_\ell E_k E_j = (-1)^{p_X(k)p_X(\ell)} E_j E_\ell E_k E_j$. Thus, by applying (R8) repeatedly we get

$$\begin{aligned}
 & T'_{j,-1}(E_{X,k})T'_{j,-1}(E_{X,\ell}) - (-1)^{p_X(k)p_X(\ell)} T'_{j,-1}(E_{X,\ell})T'_{j,-1}(E_{X,k}) \\
 &= E_k E_j E_\ell E_j - q(E_k E_j^2 - q E_j E_k E_j) E_\ell \\
 &\quad - (-1)^{p_X(k)p_X(\ell)} [E_\ell E_j E_k E_j - q(E_\ell E_j^2 - q E_j E_\ell E_j) E_k] \\
 &= E_k E_j E_\ell E_j - q(q^{-1} E_j E_k E_j - E_j^2 E_k) E_\ell \\
 &\quad - (-1)^{p_X(k)p_X(\ell)} [E_\ell E_j E_k E_j - q(q^{-1} E_j E_\ell E_j - E_j^2 E_\ell) E_k] \\
 &= \frac{1}{q + q^{-1}} [E_k (E_j^2 E_\ell + E_\ell E_j^2) - (E_k E_j^2 + E_j^2 E_k) E_\ell \\
 &\quad - (-1)^{p_X(k)p_X(\ell)} E_\ell (E_k E_j^2 + E_j^2 E_k) + (-1)^{p_X(k)p_X(\ell)} (E_j^2 E_\ell + E_\ell E_j^2) E_k] \\
 &= 0.
 \end{aligned}$$

(Case-2) Assume $p_X(j) = 1$. In this case we always have $(\alpha_j, \alpha_k) = -(\alpha_j, \alpha_\ell)$. Again we may assume that $(\alpha_j, \alpha_k) = 1$. Then $(\alpha_j, \alpha_\ell) = -1$. Thus we have $T'_{j,-1}(E_{X,k}) = E_k E_j - (-1)^{p(k)} q E_j E_k$ and $T'_{j,-1}(E_{X,\ell}) = E_\ell E_j - (-1)^{p(\ell)} q^{-1} E_j E_\ell$ and thus

$$\begin{aligned}
 T'_{j,-1}(E_{X,k})T'_{j,-1}(E_{X,\ell}) &= E_k E_j E_\ell E_j - (-1)^{p(k)} q E_j E_k E_\ell E_j + (-1)^{p(k)+p(\ell)} E_j E_k E_j E_\ell, \\
 T'_{j,-1}(E_{X,\ell})T'_{j,-1}(E_{X,k}) &= E_\ell E_j E_k E_j - (-1)^{p(\ell)} q^{-1} E_j E_\ell E_k E_j + (-1)^{p(k)+p(\ell)} E_j E_\ell E_j E_k.
 \end{aligned}$$

By taking a difference of the above two equations and unravelling the relation (R10) we can conclude that

$$T'_{j,-1}(E_{X,k})T'_{j,-1}(E_{X,\ell}) = (-1)^{p_X(k)p_X(\ell)} T'_{j,-1}(E_{X,\ell})T'_{j,-1}(E_{X,k}).$$

This proves the lemma. \square

The verifying of the relations (R8) and (R9) is no different as in the $\mathfrak{gl}(m|1)$ case. Hence we have

Lemma 3.12. [C16, Lemma 4.11] *If $j, k, \ell \in I$ such that $p_X(k) = 0$ and $k \sim \ell$, then*

$$\begin{aligned}
 & T'_{j,-1}(E_{X,k})^2 T'_{j,-1}(E_{X,\ell}) - (q + q^{-1}) T'_{j,-1}(E_{X,k}) T'_{j,-1}(E_{X,\ell}) T'_{j,-1}(E_{X,k}) \\
 &\quad + T'_{j,-1}(E_{X,\ell}) T'_{j,-1}(E_{X,k})^2 = 0, \\
 & T'_{j,-1}(F_{X,k})^2 T'_{j,-1}(F_{X,\ell}) - (q + q^{-1}) T'_{j,-1}(F_{X,k}) T'_{j,-1}(F_{X,\ell}) T'_{j,-1}(F_{X,k}) \\
 &\quad + T'_{j,-1}(F_{X,\ell}) T'_{j,-1}(F_{X,k})^2 = 0.
 \end{aligned}$$

Finally we need to verify the relations (R10) and (R11).

Lemma 3.13. *Let $z, k, j, \ell \in I$ with $k \sim j \sim \ell$, $k < \ell$ and $p_X(j) = 1$, then*

$$\begin{aligned}
 & S_{p_X(k), p_X(\ell)}(T'_{z,-1}(E_{X,k}), T'_{z,-1}(E_{X,j}), T'_{z,-1}(E_{X,\ell})) = 0, \\
 & S_{p_X(k), p_X(\ell)}(T'_{z,-1}(F_{X,k}), T'_{z,-1}(F_{X,j}), T'_{z,-1}(F_{X,\ell})) = 0.
 \end{aligned}$$

Proof. We only prove the first equality as the second one can be proved similarly. If none of k, j, ℓ is connected or equal to z , there is nothing to prove.

Now we first suppose that $j \approx z$ and $k \sim z$. In this case we have $T'_{z,-1}(E_{X,k}) = E_k E_z - (-1)^{p(z)p(k)} q^{(\alpha_k, \alpha_z)} E_z E_k$, $T'_{z,-1}(E_{X,j}) = E_j$, $T'_{z,-1}(E_{X,\ell}) = E_\ell$. Thus

$$\begin{aligned} & S_{p_X(k), p_X(\ell)}(T'_{z,-1}(E_{X,k}), T'_{z,-1}(E_{X,j}), T'_{z,-1}(E_{X,\ell})) \\ &= S_{p_X(k), p_X(\ell)}(E_k E_z, E_j, E_\ell) - (-1)^{p(z)p(k)} q^{(\alpha_z, \alpha_k)} S_{p_X(k), p_X(\ell)}(E_z E_k, E_j, E_\ell) \end{aligned}$$

When $p(z) = 0$, E_z commutes with E_j and E_ℓ . Also we have $p_X(k) = p(k)$, $p_X(\ell) = p(\ell)$. Hence

$$\begin{aligned} & S_{p_X(k), p_X(\ell)}(E_k E_z, E_j, E_\ell) = S_{p(k), p(\ell)}(E_k, E_j, E_\ell) E_z = 0, \\ & S_{p_X(k), p_X(\ell)}(E_z E_k, E_j, E_\ell) = E_z S_{p_X(k), p_X(\ell)}(E_k, E_j, E_\ell) = 0. \end{aligned}$$

When $p(z) = 1$, since $E_z E_j = (-1)^{p(z)} E_j E_z$, $E_z E_\ell = (-1)^{p(z)p(\ell)} E_\ell E_z$, $p(k) = p_X(k) + 1$ and $p(\ell) = p_X(\ell)$, again we have

$$\begin{aligned} & S_{p_X(k), p_X(\ell)}(E_k E_z, E_j, E_\ell) = S_{p(k), p(\ell)}(E_k, E_j, E_\ell) E_z = 0, \\ & S_{p_X(k), p_X(\ell)}(E_z E_k, E_j, E_\ell) = E_z S_{p_X(k), p_X(\ell)}(E_k, E_j, E_\ell) = 0. \end{aligned}$$

Note that the case when $j \approx h$, $\ell \sim h$ is similar.

Next, suppose $z \sim j$ and wlog that $z = k$. We further assume that $(\alpha_j, \alpha_k) = -1$, thus $(\alpha_j, \alpha_\ell) = -1$. Note that when $p(k)p(\ell) = 0$, the proof is already given in [C16, Lemma 4.12]. So we only need to consider the case when $p(z) = p(k) = p(\ell) = 1$. In this case we have $p_X(k) = p_X(\ell) = 1$, $p(j) = 0$ and $T'_{z,-1}(E_{X,j}) = E_j E_z - q^{-1} E_z E_j$, $T'_{z,-1}(E_{X,\ell}) = E_{X,\ell}$ and $T'_{z,-1}(E_{X,k}) = K_k^{-1} F_k$. Let $\check{\partial}_{abcd} = T'_{z,-1}(E_{X,a}) T'_{z,-1}(E_{X,b}) T'_{z,-1}(E_{X,c}) T'_{z,-1}(E_{X,d})$. The goal is to prove that

$$(q + q^{-1}) \check{\partial}_{j\ell k j} = -\check{\partial}_{j\ell j k} + \check{\partial}_{k j \ell j} + \check{\partial}_{j k j \ell} - \check{\partial}_{\ell j k j}.$$

Note the identities

$$\begin{aligned} & T'_{z,-1}(E_{X,k}) T'_{z,-1}(E_{X,\ell}) = -T'_{z,-1}(E_{X,\ell}) T'_{z,-1}(E_{X,k}), \\ & T'_{z,-1}(E_{X,j}) T'_{z,-1}(E_{X,k}) = -q^{-1} T'_{z,-1}(E_{X,k}) T'_{z,-1}(E_{X,j}) + q^{-1} E_j. \end{aligned}$$

With the above identities and Lemma 3.7 we see that

$$\begin{aligned} \check{\partial}_{j\ell j k} &= -q^{-1} \check{\partial}_{j\ell k j} + q^{-1} E_j E_k E_\ell E_j - q^{-2} E_k E_j E_\ell E_j, \\ \check{\partial}_{k j \ell j} &= -q \check{\partial}_{j k \ell j} + E_j E_\ell E_j E_k - q^{-1} E_j E_k E_\ell E_j, \\ \check{\partial}_{j k j \ell} &= E_j E_k E_j E_\ell, \\ \check{\partial}_{\ell j k j} &= E_\ell E_j E_k E_j \end{aligned}$$

Thus using Serre relation (R8) repeatedly we have

$$(q + q^{-1}) \check{\partial}_{j\ell k j} = -\check{\partial}_{j\ell j k} + \check{\partial}_{k j \ell j} + \check{\partial}_{j k j \ell} - \check{\partial}_{\ell j k j}.$$

Finally we suppose that $z = j$. Again when $p(k)p(\ell) = 0$ the proof is given in [C16, Lemma 4.12]. So we only need to consider the case when $p(k) = p(\ell) = 1$. Thus we have $p_X(k) = p_X(\ell) = 0$. Wlog we assume that $(\alpha_k, \alpha_j) = -(\alpha_j, \alpha_\ell) = -1$. Then $T'_{z,-1}(E_{X,k}) =$

$E_k E_j + q^{-1} E_j E_k$, $T'_{z,-1}(E_{X,\ell}) = E_\ell E_j + q E_j E_\ell$ and $T'_{z,-1}(E_{X,j}) = K_j^{-1} F_j$. Note that we have the identities

$$\begin{aligned} T'_{j,-1}(E_{X,k}) T'_{j,-1}(E_{X,j}) &= q^{-1} T'_{j,-1}(E_{X,j}) T'_{j,-1}(E_{X,k}) + q^{-1} E_k, \\ T'_{j,-1}(E_{X,\ell}) T'_{j,-1}(E_{X,j}) &= q T'_{j,-1}(E_{X,j}) T'_{j,-1}(E_{X,\ell}) - q E_\ell. \end{aligned}$$

Thus we have

$$\begin{aligned} \check{\partial}_{j\ell jk} &= -K_j^{-1} F_j (q E_\ell E_k E_j + E_\ell E_j E_k), \\ \check{\partial}_{kjlj} &= -E_k E_\ell - K_j^{-1} F_j (E_k E_j E_\ell + q^{-1} E_j E_k E_\ell), \\ \check{\partial}_{jkj\ell} &= K_j^{-1} F_j (q^{-1} E_k E_\ell E_j + E_k E_j E_\ell), \\ \check{\partial}_{\ell jk j} &= -E_\ell E_k + K_j^{-1} F_j (E_\ell E_j E_k + q E_j E_\ell E_k), \\ \check{\partial}_{j k \ell j} &= \check{\partial}_{j \ell k j} = K_j F_j^{-1} (E_k E_\ell E_j - E_j E_k E_\ell). \end{aligned}$$

Then we conclude that

$$(q + q^{-1}) \check{\partial}_{j \ell k j} = \check{\partial}_{j \ell j k} + \check{\partial}_{k j \ell j} + \check{\partial}_{j k j \ell} + \check{\partial}_{\ell j k j}.$$

This proves the lemma. \square

We have now proved that $T'_{j,e}$ and $T''_{j,e}$ are algebra isomorphisms for all $j \in I$ and $e = \pm 1$. Next we need to show they satisfy the braid group relations of type A . The next proposition follows from [C16, Lemma 4.13-4.15].

Proposition 3.14. *Let $j, k, \ell \in I$ and $X \in \mathcal{D}_{m,n}$.*

- (1) *If $j \approx k$, then $T'_{j,e} T'_{k,e} = T'_{k,e} T'_{j,e}$ and $T''_{j,e} T''_{k,e} = T''_{k,e} T''_{j,e}$.*
- (2) *If $j \sim k$ and $Y = r_j r_k(X)$, then*

$$\begin{aligned} T'_{j,-e} T'_{k,-e}(E_{X,j}) &= T''_{j,e} T''_{k,e}(E_{X,j}) = E_{Y,k}, \\ T'_{j,-e} T'_{k,-e}(F_{X,j}) &= T''_{j,e} T''_{k,e}(F_{X,j}) = F_{Y,k}, \\ T'_{j,-e} T'_{k,-e}(K_{X,j}) &= T''_{j,e} T''_{k,e}(K_{X,j}) = K_{Y,k}. \end{aligned}$$

- (3) *If $j \sim k$, then $T'_{j,e} T'_{k,e} T'_{j,e} = T'_{k,e} T'_{j,e} T'_{k,e}$ and $T''_{j,e} T''_{k,e} T''_{j,e} = T''_{k,e} T''_{j,e} T''_{k,e}$.*

Define $T_j := T''_{j,1}$, the next two lemmas can be proved similarly as in [Jan95]

Lemma 3.15. *Let $X \in \mathcal{D}_{m,n}$ and $\alpha, \beta \in \Pi_X$ with $\alpha \neq \beta$. Let w be an element generated by s_α and s_β . Moreover, suppose $Y = w(X)$ and $w(\alpha) > 0$ in the root system associated to X , then $T_w(E_{X,\alpha})$ is contained in the subalgebra generated by $E_{Y,\alpha}$ and $E_{Y,\beta}$. If $w(\alpha) \in \Pi_X$, then $T_w(E_{X,\alpha}) = E_{Y,w(\alpha)}$.*

Lemma 3.16. *Let $w \in W$, $X \in \mathcal{D}_{m,n}$, $Y = w(X)$ and $\alpha \in \Pi_X$. If $w(\alpha) > 0$ in the root system associated to X , then $T_w(E_{X,\alpha}) \in \mathbf{U}(Y)^+$. If $w(\alpha) \in \Pi_X$, then $T_w(E_{X,\alpha}) = E_{Y,w(\alpha)}$.*

Similar results hold for F and K as well.

4. QUANTUM SUPERSYMMETRIC PAIR OF TYPE AIII

In this section we define the quantum supersymmetric pairs and the corresponding ι quantum supergroups of type AIII. From now on we denote by T_i the braid operator $T''_{i,1}$ defined in Theorem 3.5.

4.1. **ι Quantum supergroup of type AIII.** For a real number $x \in \mathbb{R}$ and $m \in \mathbb{N}$, we denote $[x, x+m] = \{x, x+1, \dots, x+m\}$. For $a \in \mathbb{Z}_{\geq 1}$, we denote by

$$\mathbb{I}_a = \left[\frac{1-a}{2}, \frac{a-1}{2} \right].$$

Fix

$$n = \frac{m}{2} \in \frac{1}{2}\mathbb{N}.$$

We consider the Satake diagram of type AIII with $m-1 = 2n-1$ black nodes and r pairs of white nodes, together with a diagram involution τ :

$$(4.1) \quad \begin{array}{c} \begin{array}{ccc} -n-r+1 & & -n \\ \circ & \cdots & \circ \\ \circ & \cdots & \circ \\ n+r-1 & & n \end{array} \end{array}$$

where \circ stands for white dots and \blacksquare stands for black dots. Both white and black dots allow different parities satisfying the following assumption:

$$(4.2) \quad \begin{aligned} \#\{p(j) = 1 \mid j \in I_\bullet\} &\equiv 0 \pmod{2}, \\ p(j) &= p(\tau(j)), \quad \forall i \in I_\circ. \end{aligned}$$

where $I_\bullet = [1-n, n-1]$, $I_\circ = I \setminus I_\bullet$. (In case $n = 0$, the black nodes are dropped; the nodes n and $-n$ are identified and fixed by τ .)

We denote by $\tilde{\Gamma} \in \mathcal{D}_{m,n}$ the underlying Dynkin diagram and we let $I = \mathbb{I}_{m+2r-1} = I_\circ \cup I_\bullet$ to be the index set. We will denote the white even roots, black even roots, black odd roots and white odd roots respectively by \circ , \bullet , \ominus and \otimes .

Ignoring the parity, we denote by w_\bullet the longest element in the Weyl group of the Levi subalgebra associated to I_\bullet . Following [BW18b], we define

$$(4.3) \quad \begin{aligned} X^\iota &= X / \{\mu + w_\bullet \tau(\mu) \mid \mu \in X\}, \\ Y^\iota &= \{\nu - w_\bullet \tau(\nu) \mid \nu \in Y\}. \end{aligned}$$

We call an element in X^ι an ι -weight and X^ι the ι -weight lattice.

Without the diagram involution τ , the diagram (4.1) corresponds to a Lie superalgebra $\mathfrak{gl}(\mathfrak{m}|\mathfrak{n})$ for some $c, d \in \mathbb{Z}_{\geq 0}$ with $\mathfrak{m} + \mathfrak{n} = 2r + m$. The simple roots are given by

$$\tilde{\Pi} = \{\tilde{\alpha}_k = \epsilon_{i_{k-\frac{1}{2}}} - \epsilon_{i_{k+\frac{1}{2}}} \mid k \in I\}$$

where $\{\tilde{i}_{k \pm \frac{1}{2}} \mid k \in I\} = I(\mathfrak{m}|\mathfrak{n})$.

Recall ℓ_j from (2.6). For future use, we give two useful lemmas under the setting that $\tilde{\Gamma} \in \mathcal{D}_{m,n}$.

Lemma 4.1. For $\tilde{\Gamma} \in \mathcal{D}_{m,n}$, we have $\ell_j = (-1)^{p(j)}\ell_{-j}$, $\forall j \in I$.

Proof. By observation we clearly have $\ell_j = (-1)^{p(i_{j-\frac{1}{2}})}$. Moreover, since $p(j) = p(-j)$ for all $j \in I$, we have $\ell_j = (-1)^{p(j)}\ell_{-j}$, $\forall j \in I$. \square

Lemma 4.2. Suppose that $\Gamma = w_\bullet(\tilde{\Gamma})$, then $\Gamma \in \mathcal{D}_{m,n}$ and satisfies (4.2).

Proof. Recall that

$$\tilde{\Pi} = \{\tilde{\alpha}_k = \epsilon_{\tilde{i}_{k-\frac{1}{2}}} - \epsilon_{\tilde{i}_{k+\frac{1}{2}}} \mid k \in I\}$$

is the simple system of $\tilde{\Gamma}$ where $\{i_{k \pm \frac{1}{2}} \mid k \in I\} = I(m|n)$. Then for each $k \in I$, we have

$$(4.4) \quad \alpha_k := w_\bullet(\tilde{\alpha}_k) = \begin{cases} \epsilon_{i_{k-\frac{1}{2}}} - \epsilon_{i_{k+\frac{1}{2}}} & \text{if } |k| > n, \\ \epsilon_{i_{-n+\frac{1}{2}}} - \epsilon_{i_{n+\frac{1}{2}}} & \text{if } k = n, \\ \epsilon_{i_{-k+\frac{1}{2}}} - \epsilon_{i_{-k-\frac{1}{2}}} & \text{if } -n < k < n, \\ \epsilon_{i_{-n-\frac{1}{2}}} - \epsilon_{i_{n-\frac{1}{2}}} & \text{if } k = -n. \end{cases}$$

From (4.4) we can see that $\tilde{\alpha}_k = \alpha_k$ if $|k| > n$ and so is the parity. For $|k| = n$, suppose $p(\tilde{\alpha}_{-n}) = p(\tilde{\alpha}_n) = 0$, then $\epsilon_{\tilde{i}_{-n-\frac{1}{2}}}$ and $\epsilon_{\tilde{i}_{-n+\frac{1}{2}}}$ have the same parity while $\epsilon_{\tilde{i}_{n-\frac{1}{2}}}$ and $\epsilon_{\tilde{i}_{n+\frac{1}{2}}}$ have the same parity. Thus $p(\alpha_n) = p(\alpha_{-n})$. It can be checked similarly that when $p(\tilde{\alpha}_{-n}) = p(\tilde{\alpha}_n) = 1$, we still have $p(\alpha_n) = p(\alpha_{-n})$.

For $-n < k < n$, we see that $\alpha_k = -\alpha_{\Gamma, -k}$. Thus the number of black odd roots stay unchanged. \square

For convention, we define according to (4.4) that

$$(4.5) \quad i_j = \begin{cases} \tilde{i}_j, & \text{if } j > n - \frac{1}{2} \text{ or } j \leq -n - \frac{1}{2}, \\ \tilde{i}_{-j}, & \text{if } -n - \frac{1}{2} < j \leq n - \frac{1}{2} \end{cases}$$

such that $\Pi = \{\alpha_k = \epsilon_{i_{k-\frac{1}{2}}} - \epsilon_{i_{k+\frac{1}{2}}} \mid k \in I\}$.

Let $\mathbf{U}(\Gamma)$ denote the quantum supergroup with generators $\varrho, E_j, F_j, K_j, j \in I$ associated to Γ and $\mathbf{U}(\tilde{\Gamma})$ denote the same algebra with different generators $\varrho, \tilde{E}_j, \tilde{F}_j, \tilde{K}_j, j \in I$ and a different presentation corresponding to $\tilde{\Gamma}$. We note that the comultiplication Δ depends on the presentation we choose to work with, see (2.12). For simplicity we abuse the notation Δ and the parity function p for different presentations as it's clear from the context.

The ι quantum supergroup of type AIII, denoted by $\mathbf{U}^\iota = \mathbf{U}^\iota(\Gamma)$, depends on the parameters $\varsigma_j \in \mathbb{Q}(q)$, for $j \in I_\circ$, which satisfy the conditions $\varsigma_j = \varsigma_{-j}$, for $j \in I_\circ \setminus \{\pm n\}$ [Let02] (also cf. [BK15, BW21]). More precisely, $\mathbf{U}^\iota(\Gamma)$ is the $\mathbb{Q}(q)$ -subalgebra of $\mathbf{U}(\Gamma)$ generated by q^μ ($\mu \in Y^\iota$), E_j, F_j ($j \in I_\bullet$), ϱ and

$$(4.6) \quad B_j = F_j + \varsigma_j T_{w_\bullet}(\tilde{E}_{\tau(j)})K_j^{-1}, \text{ for } j \in I_\circ.$$

(In case $n = 0$, B_0 will be allowed to take a more general form $B_0 = F_0 + \varsigma_0 E_0 K_0^{-1} + \kappa_0 K_0^{-1}$, for an additional parameter $\kappa_0 \in \mathbb{Q}(q)$.)

We note that w_\bullet can be written into reduced expressions $w_\bullet = s_{j_1} \cdots s_{j_l}$. Thus we can write $T_{w_\bullet} = T_{j_1} \cdots T_{j_l}$. By Proposition 3.14, T_{w_\bullet} is a well-defined operator as a product of braid operators associated to both odd and even simple roots in I_\bullet .

Then $(\mathbf{U}(\Gamma), \mathbf{U}^\iota(\Gamma))$ forms a quantum supersymmetric pair of type AIII [Let99, Let02] (cf. [BW18a, BK19]). The algebra \mathbf{U}^ι satisfies the relations

$$\begin{aligned} q^\mu B_j &= q^{-(\mu, \alpha_j)} B_j q^\mu, \quad \forall j \in I_\circ, \\ q^\mu F_j &= q^{-(\mu, \alpha_j)} F_j q^\mu, \quad q^\mu E_j = q^{(\mu, \alpha_j)} E_j q^\mu, \quad \forall j \in I_\bullet, \mu \in Y^\iota, \\ \varrho(B_j) &= (-1)^{p(j)} B_j, \quad \forall j \in I. \end{aligned}$$

and additional Serre type relations. By definition we see the following relation holds in \mathbf{U}^ι .

Lemma 4.3. *For any $j \in I_\bullet$, $k \in I$ we have*

$$(4.7) \quad E_j B_k - (-1)^{p(j)p(k)} B_k E_j = \delta_{jk} \frac{K_j - K_j^{-1}}{q^{\ell_j} - q^{-\ell_j}}.$$

For future use, we let \mathbf{U}_\bullet denote the subalgebra of \mathbf{U} generated by $\{E_j, F_j, K_j^{\pm 1}, \varrho \mid j \in I_\bullet\}$. Let $\mathbf{U}^{\iota 0}$ denote the subalgebra of \mathbf{U}^ι generated by $\{q^\mu, \varrho \mid \mu \in Y^\iota\}$.

The next lemma will help us pin down one of the conditions on the parameters.

Lemma 4.4. *If $\varsigma_j \neq \varsigma_{-j}$ for $j \in I_\circ \setminus \{\pm n\}$, then $(K_j^{-1} K_{-j}^{-1}) \in \mathbf{U}^{\iota 0}$.*

Proof. This claim follows from the relation

$$B_j B_{-j} - (-1)^{p(j)} B_{-j} B_j = \varsigma_{-j} (-1)^{p(j)} \frac{K_j - K_j^{-1}}{q^{\ell_j} - q^{-\ell_j}} K_{-j}^{-1} + \varsigma_j \frac{K_{-j} - K_{-j}^{-1}}{q^{\ell_{-j}} - q^{-\ell_{-j}}} K_j^{-1}$$

According to Lemma 4.1 we have $\ell_j = (-1)^{p(j)} \ell_{-j}$. Thus $q^{\ell_j} - q^{-\ell_j} = (-1)^{p(j)} (q^{\ell_{-j}} - q^{-\ell_{-j}})$. Hence the lemma follows. \square

By the above lemma we see that $\mathbf{U}^\iota \cap \mathbf{U}^0 = \mathbf{U}^{\iota 0}$ can only be satisfied if the parameters satisfying $\varsigma_j = \varsigma_{-j}$ for $j \in I_\circ \setminus \{\pm n\}$. From now on we assume the parameters $\{\varsigma_j\}$ always satisfy this condition.

Furthermore, we determine the action of T_{w_\bullet} on \mathbf{U}_\bullet .

Lemma 4.5. *For all $j \in I_\bullet$, we have*

$$(4.8) \quad \begin{aligned} T_{w_\bullet}(\tilde{E}_j) &= -F_{-j} K_{-j}, & T_{w_\bullet}(\tilde{F}_j) &= -K_{-j}^{-1} E_{-j}, & T_{w_\bullet}(\tilde{K}_j) &= K_{-j}^{-1}, \\ T_{w_\bullet}^{-1}(\tilde{E}_j) &= -K_{-j}^{-1} F_{-j}, & T_{w_\bullet}^{-1}(\tilde{F}_j) &= -E_{-j} K_{-j}, & T_{w_\bullet}^{-1}(\tilde{K}_j) &= K_{-j}^{-1}. \end{aligned}$$

Proof. The proof of this lemma follows from the same argument in [Ko12, Lemma 3.4] and Lemma 3.16. \square

4.2. Coideal subalgebra property. One of the key properties of the quantum group is that it is a coideal subalgebra of the underlying Hopf algebra rather than a Hopf subalgebra. Here we develop such a structure for \mathbf{U}^ι as well.

Proposition 4.6. *\mathbf{U}^ι is a right coideal subalgebra of \mathbf{U} .*

Proof. Clearly, \mathbf{U}_\bullet and \mathbf{U}^{j0} are Hopf subalgebras of \mathbf{U} . Thus it suffices to show that

$$(4.9) \quad \Delta(B_j) \in \mathbf{U}^j \otimes \mathbf{U}, \quad \forall j \in I_\circ.$$

Recall Δ from (2.12). It's straightforward to compute for $j \in I_\circ \setminus \{\pm n\}$ that

$$(4.10) \quad \Delta(B_j) - B_j \otimes K_j^{-1} \in \mathbf{U}_\bullet^+ \mathbf{U}^{j0} \otimes \mathbf{U}.$$

Now suppose $j = -n$. For any reduced expression $\underline{w}_\bullet^{(y)} = s_{y_1} \cdots s_{y_\ell}$ of w_\bullet , we define

$$\Gamma_t^{(y)} = s_{y_t} \cdots s_{y_\ell}(\tilde{\Gamma}), \quad 1 \leq t \leq \ell.$$

More specifically, in this case we choose

$$\underline{w}_\bullet^{(y)} = (s_{n-1}s_{n-2} \cdots s_{-n+1}) \cdots (s_{n-1}s_{n-2})(s_{n-1}).$$

For convention we drop (y) in the following proof. Then we first observe that

$$(4.11) \quad \begin{aligned} T_{w_\bullet}(\tilde{E}_{-n}) &= T_{s_{n-1}} \cdots T_{s_{-n+1}}(E_{-n}^{\Gamma_{2n}}) \\ &= T_{s_{n-1}} \cdots T_{s_{-n+2}}(E_{-n+1}^{\Gamma_{2n-1}} E_{-n}^{\Gamma_{2n-1}} \\ &\quad - (-1)^{p(\alpha_{-n+1}^{\Gamma_{2n-1}})p(\alpha_{-n}^{\Gamma_{2n-1}})} q^{(\alpha_{-n+1}^{\Gamma_{2n-1}}, \alpha_{-n}^{\Gamma_{2n-1}})} E_{-n}^{\Gamma_{2n-1}} E_{-n+1}^{\Gamma_{2n-1}}) \\ &= T_{s_{n-1}} \cdots T_{s_{-n+2}}(E_{-n+1}^{\Gamma_{2n-1}}) E_{-n} - z E_{-n} T_{s_{n-1}} \cdots T_{s_{-n+2}}(E_{-n+1}^{\Gamma_{2n-1}}) \end{aligned}$$

where $z = (-1)^{p(\alpha_{-n+1}^{\Gamma_{2n-1}})p(\alpha_{-n}^{\Gamma_{2n-1}})} q^{(\alpha_{-n+1}^{\Gamma_{2n-1}}, \alpha_{-n}^{\Gamma_{2n-1}})}$.

By (4.11) we see that in order to prove (4.10) for $j = -n$, it suffices to prove that

$$(4.12) \quad \Delta(T_{s_{n-1}} \cdots T_{s_{-n+1}}(E_{-n}^{\Gamma_{2n}})) \in T_{s_{n-1}} \cdots T_{s_{-n+1}}(E_{-n}^{\Gamma_{2n}}) \otimes 1 + \mathbf{U}_\bullet^+ \mathbf{U}^0 K_n \otimes \mathbf{U}.$$

We prove (4.12) by proving the following claim.

Claim: For any $1 \leq k \leq 2n-1$, we have

$$(4.13) \quad \begin{aligned} \Delta(T_{s_{n-1}} \cdots T_{s_{-n+k}}(E_{-n+k-1}^{\Gamma_{2n+1-k}})) \\ \in T_{s_{n-1}} \cdots T_{s_{-n+k}}(E_{-n+k-1}^{\Gamma_{2n+1-k}}) \otimes 1 + \mathbf{U}_\bullet^+ \mathbf{U}^0 \varrho^{p(-n+k-1)} K_{-n+k-1} \otimes \mathbf{U}. \end{aligned}$$

We prove (4.12) through induction on $2n-1-k$. When $k = 2n-1$, we have

$$\begin{aligned} \Delta(T_{s_{n-1}}(E_{n-2}^{\Gamma_2})) &= \Delta(E_{n-1}E_{n-2} - (-1)^{p(\alpha_{n-1})p(\alpha_{n-2})} q^{(\alpha_{n-1}, \alpha_{n-2})} E_{n-2}E_{n-1}) \\ &\in T_{s_{n-1}}(E_{n-2}^{\Gamma_2}) \otimes 1 + \mathbf{U}_\bullet^+ \mathbf{U}^0 \varrho^{p(n-2)} K_{n-2} \otimes \mathbf{U}. \end{aligned}$$

Now suppose the claim is true for $k = 2$, that is

$$(4.14) \quad \Delta(T_{s_{n-1}} \cdots T_{s_{-n+2}}(E_{-n+1}^{\Gamma_{2n-1}})) = T_{s_{n-1}} \cdots T_{s_{-n+2}}(E_{-n+1}^{\Gamma_{2n-1}}) \otimes 1 + \sum_{\ell} x_{\ell} \varrho^{p(-n+1)} K_{-n+1} \otimes y_{\ell},$$

for some $x_{\ell} \in \mathbf{U}_\bullet^+ \mathbf{U}^0$, $y_{\ell} \in \mathbf{U}$.

In view of (2.12), (4.11) and (4.14), we see that

$$\begin{aligned}
& \Delta(T_{s_{n-1}} \cdots T_{s_{-n+2}}(E_{-n+1}^{\Gamma_{2n-1}}))\Delta(E_{-n}) \\
& \in T_{s_{n-1}} \cdots T_{s_{-n+2}}(E_{-n+1}^{\Gamma_{2n-1}})E_{-n} \otimes 1 \\
& \quad + \sum_{\ell} x_{\ell} \varrho^{p(-n+1)} K_{-n+1} E_{-n} \otimes y_{\ell} + \mathbf{U}_{\bullet}^{+} \mathbf{U}^0 \varrho^{p(-n)} K_{-n} \otimes \mathbf{U}, \\
(4.15) \quad & \Delta(E_{-n})\Delta(T_{s_{n-1}} \cdots T_{s_{-n+2}}(E_{-n+1}^{\Gamma_{2n-1}})) \\
& \in E_{-n} T_{s_{n-1}} \cdots T_{s_{-n+2}}(E_{-n+1}^{\Gamma_{2n-1}}) \otimes 1 \\
& \quad + \sum_{\ell} E_{-n} x_{\ell} \varrho^{p(-n+1)} K_{-n+1} \otimes y_{\ell} + \mathbf{U}_{\bullet}^{+} \mathbf{U}^0 \varrho^{p(-n)} K_{-n} \otimes \mathbf{U}.
\end{aligned}$$

By comparing both sides of (4.14) we see that each x_{ℓ} is a monomial of the form $a_{(i)} Z_{i_1} \cdots Z_{i_{2n-2}}$ where

$$Z_{i_t} \in \{E_{i_t}, \varrho^{p(i_t)} K_{i_t}\}, \quad a_{(i)} \in \mathbb{Q}(q).$$

and $\{i_1, \dots, i_{2n-2}\} = \{n-1, \dots, -n+2\}$. Hence each $x_{\ell} \varrho^{p(-n+1)}$ supercommutes with E_{-n} as follows:

$$E_{-n} x_{\ell} \varrho^{p(-n+1)} = (-1)^{p(-n)} (-1)^{p(n-1)+\cdots+p(-n+2)} x_{\ell} E_{-n} \varrho^{p(-n+1)} = x_{\ell} \varrho^{p(-n+1)} E_{-n}.$$

The last equality comes from (4.2). Moreover, we compute directly that

$$(4.16) \quad p(\alpha_{-n+1}^{\Gamma_{2n-1}}) = p(s_{-n+1}(s_{n-1} \cdots s_{-n+2})\tilde{\alpha}_{-n+1}) = 0, \quad (\alpha_{-n+1}^{\Gamma_{2n-1}}, \alpha_{-n}^{\Gamma_{2n-1}}) = (\alpha_{-n+1}, \alpha_{-n}).$$

Now consider $\Delta(T_{w_{\bullet}}(\tilde{E}_{-n}))$, from (4.11), (4.15) and (4.16) we have

$$\begin{aligned}
& \Delta(T_{w_{\bullet}}(\tilde{E}_{-n})) = \Delta(T_{s_{n-1}} \cdots T_{s_{-n+1}}(E_{-n}^{\Gamma_{2n}})) \\
(4.17) \quad & = \Delta(T_{s_{n-1}} \cdots T_{s_{-n+2}}(E_{-n+1}^{\Gamma_{2n-1}}))\Delta(E_{-n}) - z\Delta(E_{-n})\Delta(T_{s_{n-1}} \cdots T_{s_{-n+2}}(E_{-n+1}^{\Gamma_{2n-1}})) \\
& \in T_{w_{\bullet}}(\tilde{E}_{-n}) \otimes 1 + \mathbf{U}_{\bullet}^{+} \mathbf{U}^0 \varrho^{p(-n)} K_{-n} \otimes \mathbf{U}.
\end{aligned}$$

Thus the claim is proved. Similarly for $j = n$ we also have

$$(4.18) \quad \Delta(T_{w_{\bullet}}(\tilde{E}_n)) \in T_{w_{\bullet}}(\tilde{E}_n) \otimes 1 + \mathbf{U}_{\bullet}^{+} \mathbf{U}^0 K_n \otimes \mathbf{U}$$

Thus we conclude from (4.10), (4.17) and (4.18) that

$$(4.19) \quad \Delta(B_j) - B_j \otimes K_j^{-1} \in \mathbf{U}_{\bullet}^{+} \mathbf{U}^0 \otimes \mathbf{U} \quad \forall j \in I_{\circ}.$$

This proves the proposition. \square

4.3. Quantum ι Serre relations. Recall the Serre relations (R5) – (R11) from (2.7). In this subsection we explore the Serre relations of \mathbf{U}^{ι} . For convention, we extend the definition of B_j by setting $B_j = F_j$ for $j \in I_{\bullet}$. Moreover, we define $\mathbf{U}^{\iota,+}$ (resp. $\mathbf{U}^{\iota,-}$) to be the subalgebra of \mathbf{U} generated by \mathbf{U}^{+} (resp. \mathbf{U}^{-}) and ϱ .

The triangular decomposition of \mathbf{U} implies a direct sum decomposition

$$(4.20) \quad \mathbf{U} = \bigoplus_{\mu \in P} \mathbf{U}^{\iota,+} K_{\mu} S(\mathbf{U}^{\iota,-})$$

Let P_μ denote the projection $\mathbf{U} \rightarrow \mathbf{U}^{\varrho,+} K_\mu S(\mathbf{U}^{\varrho,-})$ with respect to this decomposition. It follows from (2.12) that

$$(4.21) \quad \Delta \circ P_\mu(x) = (id \otimes P_\mu) \circ \Delta(x), \quad \forall x \in \mathbf{U}.$$

This relation implies the following grading:

Lemma 4.7. *We have $\mathbf{U}^\varrho = \bigoplus_{\mu \in P} P_\mu(\mathbf{U}^\varrho)$.*

Proof. For any $b \in \mathbf{U}^\varrho$ and $\mu \in P$, one see that

$$\Delta \circ P_\mu(b) = (id \otimes P_\mu) \circ \Delta(b) \in \mathbf{U}^\varrho \otimes \mathbf{U}.$$

Apply $id \otimes \epsilon$ to the above equation and we see that $P_\mu(b) \in \mathbf{U}^\varrho$. □

We may also consider the direct sum decomposition

$$\mathbf{U} = \bigoplus_{\alpha, \beta \in P^+} \mathbf{U}_\alpha^+ \mathbf{U}^0 \mathbf{U}_{-\beta}^-$$

and let $\pi_{\alpha, \beta}$ denote the corresponding projection.

Lemma 4.8. *For $j \in I_{\overline{1}}$, we have $B_j^2 = 0$.*

Proof. It suffices to check for $j \in I_0 \cap I_{\overline{1}}$. We have

$$\begin{aligned} B_j^2 &= (F_j + \varsigma_j T_{w_\bullet}(\tilde{E}_{-j}) K_j^{-1})^2 \\ &= F_j^2 + \varsigma_j [F_j, T_{w_\bullet}(\tilde{E}_{-j})] K_j^{-1} + \varsigma_j^2 T_{w_\bullet}(\tilde{E}_{-j})^2 K_j^{-2} \\ &= 0. \end{aligned}$$

This proves the lemma. □

We define two special weights $\lambda_{j,k} = (1 + |(\alpha_j, \alpha_k)|)\alpha_j + \alpha_k$ and $\lambda_j = 2\alpha_j + \alpha_{j-1} + \alpha_{j+1}$ in order to apply the projection technique in [Ko12]. Furthermore, we define

$$S(x_1, x_2) = x_1^2 x_2 - [2]x_1 x_2 x_1 + x_2 x_1^2, \quad \forall x_1, x_2 \in \mathbf{U}$$

and recall $S_{t_1, t_2}(x_1, x_2, x_3)$ from (2.8).

Lemma 4.9. (1) *Assume $j \in I_\bullet$, $k \in I$ and $j \approx k \in I$, then we have $[B_j, B_k] = B_j B_k - (-1)^{p(j)p(k)} B_k B_j = 0$.*

(2) *For $j \in I_\bullet \cap I_{\overline{0}}$ and $k \sim j$, we have $S(B_j, B_k) = 0$.*

(3) *For $j \in I_\bullet \cap I_{\overline{1}}$ and $k \sim j \sim \ell$, we have $S_{p(k), p(\ell)}(B_k, B_j, B_\ell) = 0$.*

Proof. (1) In general we have

$$[B_j, B_k] = [F_j, F_k] + \varsigma_k [F_j, T_{w_\bullet}(\tilde{E}_{-k}) K_k^{-1}].$$

Now if $j, k \in I_\bullet$, there is nothing to prove. If k is in I_0 then in this case we can rewrite F_j as $-T_{w_\bullet}(\tilde{E}_{-j}) K_j^{-1}$ according to Lemma 4.5 and one computes that

$$[T_{w_\bullet}(\tilde{E}_{-j}) K_j^{-1}, T_{w_\bullet}(\tilde{E}_{-k}) K_k^{-1}] = T_{w_\bullet}([\tilde{E}_{-j} \tilde{K}_{-j}, \tilde{E}_{-k} \tilde{K}_{-k}]) = 0.$$

(2) If $k \in I_\bullet$, there is nothing to prove. Thus we can assume that $k \in I_\circ$. In this case we have

$$\begin{aligned} F_{j,k}(B_j, B_k) &= F_{j,k}(F_j, \varsigma_k T_{w_\bullet}(\tilde{E}_{-k})K_k^{-1}) \\ &= \varsigma_k F_{j,k}(T_{w_\bullet}(\tilde{E}_{-j})K_j^{-1}, T_{w_\bullet}(\tilde{E}_{-k})K_k^{-1}) \\ &= \varsigma_k z F_{j,k}(T_{w_\bullet}(\tilde{E}_{-j}), T_{w_\bullet}(\tilde{E}_{-k}))K_j^{-2}K_k^{-1} \end{aligned}$$

for some $z \in \mathbb{Z}[q, q^{-1}]$. Since T_{w_\bullet} is an algebra homomorphism, we see that $F_{j,k}(B_j, B_k) = 0$.

(3) Without loss of generality, we assume $k = j - 1$ and $\ell = j + 1$. Note that only one of k and ℓ can belong to I_\circ . Suppose that $k \in I_\bullet$, $\ell \in I_\circ$, then we have

$$\begin{aligned} S_{p(k),p(\ell)}(B_k, B_j, B_\ell) &= \varsigma_\ell S_{p(k),p(\ell)}(F_k, F_j, T_{w_\bullet}(\tilde{E}_{-\ell})K_\ell^{-1}) \\ &= -\varsigma_\ell S_{p(k),p(\ell)}(T_{w_\bullet}(\tilde{E}_{-k})K_k^{-1}, T_{w_\bullet}(\tilde{E}_{-j})K_j^{-1}, T_{w_\bullet}(\tilde{E}_{-\ell})K_\ell^{-1}) \\ &= -\varsigma_\ell S_{p(k),p(\ell)}(T_{w_\bullet}(\tilde{E}_{-k}), T_{w_\bullet}(\tilde{E}_{-j}), T_{w_\bullet}(\tilde{E}_{-\ell}))K_k^{-1}K_j^{-2}K_\ell^{-1} = 0. \end{aligned}$$

The case $k \in I_\circ$, $\ell \in I_\bullet$ can be proved similarly. \square

Lemma 4.10. (1) Assume $j \approx k \in I$, then we have $P_{-\lambda_{j,k}}([B_j, B_k]) = 0$.

(2) Assume $j \sim k$ and $j \in I_{\overline{0}}$, then we have $P_{-\lambda_{j,k}}(S(B_j, B_k)) = 0$.

(3) Assume $k \sim j \sim \ell$ and $j \in I_{\overline{1}}$, then we have $P_{-\lambda_j}(S_{p(k),p(\ell)}(B_k, B_j, B_\ell)) = 0$.

Proof. (1) According to Lemma 4.9, it suffices to consider when $j \approx k \in I_\circ$. One sees that in this case $[B_j, B_k] = 0$ unless $j = -k$. Now if $j = -k \neq \pm n$, it follows from Lemma 4.4 that $P_{-\lambda_{j,k}}([B_j, B_k]) = 0$. If $j = -k = n$, then there exists a scalar $a_1 \in \mathbb{Q}(q)$ such that

$$P_{-\lambda_{-n,n}}([B_n, B_{-n}]) = a_1 [T_{w_\bullet}(\tilde{E}_{-n}), T_{w_\bullet}(\tilde{E}_n)]K_n^{-1}K_{-n}^{-1} = 0.$$

(2) According to Lemma 4.9, it suffices to consider when $j \in I_\circ \cap I_{\overline{0}}$. In this case there exists a scalar $a_2 \in \mathbb{Q}(q)$ such that

$$\begin{aligned} P_{-\lambda_{j,k}}(S(B_j, B_k)) &= S(\varsigma_j T_{w_\bullet}(\tilde{E}_{-j})K_j^{-1}, \varsigma_k T_{w_\bullet}(\tilde{E}_{-k})K_k^{-1}) \\ &= a_2 S(T_{w_\bullet}(\tilde{E}_{-j}), T_{w_\bullet}(\tilde{E}_{-k}))K_j^{-2}K_k^{-1} = 0. \end{aligned}$$

(3) According to Lemma 4.9, it suffices to consider when $j \in I_\circ \cap I_{\overline{1}}$. In this case there exists a scalar $a_3 \in \mathbb{Q}(q)$ such that

$$\begin{aligned} P_{-\lambda_j}(S_{p(k),p(\ell)}(B_k, B_j, B_\ell)) &= S_{p(k),p(\ell)}(\varsigma_k T_{w_\bullet}(\tilde{E}_{-k})K_k^{-1}, \varsigma_j T_{w_\bullet}(\tilde{E}_{-j})K_j^{-1}, \varsigma_\ell T_{w_\bullet}(\tilde{E}_{-\ell})K_\ell^{-1}) \\ &= a_3 S_{p(k),p(\ell)}(T_{w_\bullet}(\tilde{E}_{-k}), T_{w_\bullet}(\tilde{E}_{-j}), T_{w_\bullet}(\tilde{E}_{-\ell}))K_j^{-2}K_k^{-1}K_\ell^{-1} = 0. \end{aligned}$$

This proves the lemma. \square

Remark 4.11. The assumption (4.2) is required for Lemma 4.9 and Lemma 4.10.

For any $J = (j_1, \dots, j_r) \in I^r$ define $wt(J) = \sum_{i=1}^r \alpha_{j_i}$ and

$$(4.22) \quad E_J = E_{j_1} \cdots E_{j_r}, \quad F_J = F_{j_1} \cdots F_{j_r}, \quad B_J = B_{j_1} \cdots B_{j_r}.$$

In this case we also define $|J| = r$. Let \mathcal{J} denote a fixed subset of $\cup_{s \in \mathbb{Z}_{\geq 0}} I^s$ such that $\{F_J \mid J \in \mathcal{J}\}$ is a basis of \mathbf{U}^- . Moreover, applying the projection technique one can conclude that

Proposition 4.12. *In \mathbf{U}^ι one has the relation*

$$(4.23) \quad \begin{aligned} (1) [B_j, B_k] &\in \sum_{\{J | wt(J) < \lambda_{jk}\}} \mathbf{U}_\bullet^+ \mathbf{U}^{\iota_0} B_J, && \text{for all } j \approx k \in I, \\ (2) S(B_j, B_k) &\in \sum_{\{J | wt(J) < \lambda_{j,k}\}} \mathbf{U}_\bullet^+ \mathbf{U}^{\iota_0} B_J, && \text{for all } j \sim k \in I, j \in I_{\overline{0}}, \\ (3) S_{p(k), p(\ell)}(B_k, B_j, B_\ell) &\in \sum_{\{J | wt(J) < \lambda_j\}} \mathbf{U}_\bullet^+ \mathbf{U}^{\iota_0} B_J, && \text{for all } k \sim j \sim \ell \in I, j \in I_{\overline{1}}. \end{aligned}$$

Proof. We only proof for (3) and the proof for (1) and (2) is similar.

Let $\Xi = S_{p(k), p(\ell)}(B_k, B_j, B_\ell)$, by relation (4.7) and (4.19) we have

$$(4.24) \quad \Delta(\Xi) = \Xi \otimes K_{-\lambda_j} + \sum_{\{J | wt(J) < \lambda_j\}} \mathbf{U}_\bullet^+ \mathbf{U}^{\iota_0} B_J \otimes \mathbf{U}.$$

Moreover, relation (4.21) and (4.24) implies

$$(4.25) \quad \Delta(P_{-\lambda_j}(\Xi)) \in \Xi \otimes K_{-\lambda_j} + \sum_{\{J | wt(J) < \lambda_j\}} \mathbf{U}_\bullet^+ \mathbf{U}^{\iota_0} B_J \otimes \mathbf{U}^+ K_{-\lambda_j} S(\mathbf{U}^-).$$

Since $P_{-\lambda_j}(\Xi) = 0$ according to Lemma 4.10, by applying the counit to the second tensor factor in (4.25) we get $S_{p(k), p(\ell)}(B_k, B_j, B_\ell) \in \sum_{\{J | wt(J) < \lambda_j\}} \mathbf{U}_\bullet^+ \mathbf{U}^{\iota_0} B_J$. \square

4.4. Natural filtration. Define a filtration \mathcal{F}^* of \mathbf{U}^- by

$$\mathcal{F}^t(\mathbf{U}^-) = \text{span}\{F_J \mid J \in I^s, s \leq t\}, \quad t \in \mathbb{Z}_{\geq 0}.$$

As the quantum Serre relations for \mathbf{U} are homogeneous, the set $\{F_J \mid J \in \mathcal{J}, |J| \leq t\}$ forms a basis for $\mathcal{F}^t(\mathbf{U}^-)$.

Proposition 4.13. *The set $\{B_J \mid J \in \mathcal{J}\}$ is a basis of the left $\mathbf{U}^+ \mathbf{U}^0$ -module \mathbf{U}^ι .*

Proof. First for any $J \in \mathcal{J}$ such that $|J| = t$ we have $F_J - B_J \in \mathbf{U}^+ \mathbf{U}^0 \mathcal{F}^{t-1}(\mathbf{U}^-)$. Thus by induction on t we can conclude that each F_J is contained in the left $\mathbf{U}^+ \mathbf{U}^0$ -module generated by $\{B_J \mid J \in \mathcal{J}\}$.

It remains to show $\{B_J \mid J \in \mathcal{J}\}$ is linearly independent. Assume there exists a non-empty finite subset $\mathcal{J}' \subset \mathcal{J}$ such that $\sum_{J \in \mathcal{J}'} a_J B_J = 0$. Let $t_0 = \max\{|J| \mid J \in \mathcal{J}'\}$. In view of the definition of B_j , we have

$$\sum_{J \in \mathcal{J}', |J|=t_0} a_J F_J = 0.$$

The linear independence of $\{F_J \mid J \in \mathcal{J}\}$ implies $a_J = 0$ for all $J \in \mathcal{J}', |J| = t_0$. Then through induction we conclude the desired result. \square

By Proposition 4.13 any element in \mathbf{U}^ι can be written as a linear combination of elements in $\{B_J \mid J \in \mathcal{J}\}$ with coefficients in $\mathbf{U}^+ \mathbf{U}^0$. We want to further show that the coefficients are from $\mathbf{U}_\bullet^+ \mathbf{U}^{\iota_0}$.

Proposition 4.14. *The set $\{B_J \mid J \in \mathcal{J}\}$ is a basis of the left $\mathbf{U}_\bullet^+ \mathbf{U}^{\iota_0}$ -module \mathbf{U}^ι .*

Proof. First of all, since $\{B_J \mid J \in \mathcal{J}\}$ is linearly independent over $\mathbf{U}^+\mathbf{U}^0$, it's also independent over $\mathbf{U}_\bullet^+\mathbf{U}^0$.

Secondly, let $L \in I^t$, according to Proposition 4.12 one sees that

$$B_L - \sum_{J \in \mathcal{J} \mid |J|=t} a_J B_J \in \sum_{s < t} \sum_{J \in I^s} \mathbf{U}_\bullet^+\mathbf{U}^0 B_J.$$

Thus through induction we see that $\{B_J \mid J \in \mathcal{J}\}$ spans the left $\mathbf{U}_\bullet^+\mathbf{U}^0$ -module \mathbf{U}^t . \square

5. \imath SCHUR DUALITY

In this section, we explore the fundamental representation \mathbb{W} of \mathbf{U} and establish a commuting action between the \imath quantum supergroup \mathbf{U}^t and the Hecke algebra of type B on $\mathbb{W}^{\otimes d}$.

5.1. Bimodule structure. Recall the basic setting from (4.4) and (4.5) that

$$\Pi = w_\bullet(\tilde{\Pi}) = \{\alpha_j = \epsilon_{i_{j-\frac{1}{2}}} - \epsilon_{i_{j+\frac{1}{2}}} \mid j \in I\}$$

is the set of simple roots of Γ . Let \mathbb{W} be the vector superspace with an ordered basis $\{w_j := v_{i_j} \mid j \in \mathbb{I}_{r|m|r}\}$ parametrized by $I(m|n)$ as in §2.1. We introduce a new notation for \mathbb{I}_{2r+m} to indicate a fixed set partition and to parametrize \mathbb{W} properly:

$$(5.1) \quad \mathbb{I}_{r|m|r} := \mathbb{I}_{2r+m}, \quad \mathbb{I}_{r|m|r} = \mathbb{I}_\circ^- \cup \mathbb{I}_\bullet \cup \mathbb{I}_\circ^+$$

where the subsets

$$(5.2) \quad \mathbb{I}_\circ^+ = \left[\frac{m+1}{2}, r + \frac{m-1}{2} \right], \quad \mathbb{I}_\bullet = \left[\frac{1-m}{2}, \frac{m-1}{2} \right], \quad \mathbb{I}_\circ^- = -\mathbb{I}_\circ^+,$$

The Weyl group $W = W_d$ of type B_d is generated by s_i , for $0 \leq i \leq d-1$, subject to the Coxeter relations: $s_i^2 = 1$, $(s_i s_{i+1})^3 = 1$, $(s_0 s_1)^4 = 1$, and $(s_i s_j)^2 = 1$ ($|i-j| > 1$).

We view $f \in \mathbb{I}_{r|m|r}^d$ as a map $f : \{1, \dots, d\} \rightarrow \mathbb{I}_{r|m|r}$, and identify $f = (f(1), \dots, f(d))$, with $f(i) \in \mathbb{I}_{r|m|r}$. We define a right action of the Weyl group W_d on $\mathbb{I}_{r|m|r}^d$ such that, for $f \in \mathbb{I}_{r|m|r}^d$ and $0 \leq j \leq d-1$,

$$(5.3) \quad f^{s_j} = f \cdot s_j = \begin{cases} (\dots, f(j+1), f(j), \dots), & \text{if } j > 0; \\ (-f(1), f(2), \dots, f(d)), & \text{if } j = 0, f(1) \in \mathbb{I}_\circ^- \cup \mathbb{I}_\circ^+; \\ (f(1), f(2), \dots, f(d)), & \text{if } j = 0, f(1) \in \mathbb{I}_\bullet. \end{cases}$$

The only nontrivial relation $(s_0 s_1)^4 = 1$ can be verified by case-by-case inspection depending on whether or not $f(1), f(2) \in \mathbb{I}_\bullet$. We sometimes write

$$f^\omega = f \cdot \omega = (f(\omega(1)), \dots, f(\omega(d))),$$

where it is understood that

$$f(\omega(i)) = \begin{cases} f(\omega(i)), & \text{if } \omega(i) > 0; \\ f(-\omega(i)), & \text{if } \omega(i) < 0, f(-\omega(i)) \in \mathbb{I}_\bullet; \\ -f(-\omega(i)), & \text{if } \omega(i) < 0, f(-\omega(i)) \in \mathbb{I}_\circ^- \cup \mathbb{I}_\circ^+. \end{cases}$$

Consider the $\mathbb{Q}(Q, q)$ -vector superspace

$$(5.4) \quad \mathbb{W} = \bigoplus_{a \in \mathbb{I}_{r|m|r}} \mathbb{Q}(Q, q)w_a.$$

where each w_a has weight ϵ_{i_a} and parity $p(w_a) = p(\epsilon_{i_a})$.

Given $f = (f(1), \dots, f(d)) \in \mathbb{I}_{r|m|r}^d$, we denote

$$M_f = w_{f(1)} \otimes w_{f(2)} \otimes \dots \otimes w_{f(d)}.$$

We shall call f a weight and $\{M_f \mid f \in \mathbb{I}_{r|m|r}^d\}$ the standard basis for $\mathbb{W}^{\otimes d}$.

The $\mathbb{Q}(Q, q)$ -vector superspace \mathbb{W} can be identified with the natural representation of $\mathbf{U}(\Gamma)$, where

$$(5.5) \quad \begin{aligned} E_j w_k &= \delta_{j+\frac{1}{2}, k} w_{k-1}, & F_j w_k &= \delta_{j-\frac{1}{2}, k} w_{k+1}, \\ K_j w_k &= q^{(\alpha_j, \epsilon_{i_k})} w_k, & \varrho(w_k) &= (-1)^{p(w_k)} w_k, \forall k \in \mathbb{I}_{r|m|r}. \end{aligned}$$

Since \mathbf{U}^ι is a subalgebra of \mathbf{U} , \mathbb{W} is naturally a left \mathbf{U}^ι -module. The tensor product $\mathbb{W}^{\otimes d}$ is naturally a \mathbf{U}^ι -module via the comultiplication Δ . For the rest of this subsection we define a right \mathcal{H}_{B_d} -module structure on \mathbb{W} .

The Iwahori-Hecke algebra of type A, denoted by $\mathcal{H}_{\mathfrak{S}_d}$, is a $\mathbb{Q}(q)$ -algebra generated by H_1, \dots, H_{d-1} , subject to the following relations:

$$\begin{aligned} (H_i - q)(H_i + q^{-1}) &= 0, \\ H_i H_{i+1} H_i &= H_{i+1} H_i H_{i+1}, \quad \text{for } i \geq 1; \\ H_i H_j &= H_j H_i, \quad \text{for } |i - j| > 1, \end{aligned}$$

while the Iwahori-Hecke algebra of type B, denoted by \mathcal{H}_{B_d} , is the $\mathbb{Q}(Q, q)$ -algebra generated by $\mathcal{H}_{\mathfrak{S}_d}$ and H_0 where

$$(5.6) \quad (H_0 - Q)(H_0 + Q^{-1}) = 0, \quad H_0 H_1 H_0 H_1 = H_1 H_0 H_1 H_0.$$

If $\omega \in W_d$ has a reduced expression $\omega = s_{i_1} \cdots s_{i_k}$, we denote $H_\omega = H_{i_1} \cdots H_{i_k}$. It is well known that $\{H_\omega \mid \omega \in \mathfrak{S}_d\}$ form a basis for \mathcal{H}_{B_d} .

The following proposition is a multi-parameter version of [CL22, Proposition 2.10].

Proposition 5.1. *There is an right action of \mathcal{H}_{B_d} on $\mathbb{W}^{\otimes d}$ as follows:*

$$M_f \cdot H_i = \begin{cases} (-1)^{p(w_{f(i)})p(w_{f(i+1)})} M_{f \cdot s_i} + (q - q^{-1})M_f, & \text{if } f(i) < f(i+1), i > 0; \\ (-1)^{p(w_{f(i)})p(w_{f(i+1)})} M_{f \cdot s_i}, & \text{if } f(i) > f(i+1), i > 0; \\ \frac{(-1)^{p(w_{f(i)})}(q + q^{-1}) + q - q^{-1}}{2} M_f, & \text{if } f(i) = f(i+1), i > 0; \\ (-1)^{p(w_{f(1)})} M_{f \cdot s_i} + (Q - Q^{-1})M_f, & \text{if } f(1) \in \mathbb{I}_\circ^+, i = 0; \\ (-1)^{p(w_{f(1)})} M_{f \cdot s_i}, & \text{if } f(1) \in \mathbb{I}_\circ^-, i = 0; \\ QM_f, & \text{if } f(1) \in \mathbb{I}_\bullet, i = 0. \end{cases}$$

Proof. We already know from [Mi06] that this defines an action of $\mathcal{H}_{\mathfrak{S}_d}$ on $\mathbb{W}^{\otimes d}$. The only nontrivial relation left to verify is $H_0H_1H_0H_1 = H_1H_0H_1H_0$, which follows from a case-by-case checking. \square

5.2. ι Schur duality of type B. We first recall results from [Mi06] which establish a type A Schur duality between the quantum supergroup and the Hecke algebra of type A.

We let ${}^{st}\mathbf{U}_q^{\mathfrak{g}}(\mathfrak{gl}(m|n))$ denote the quantum supergroup corresponding to the standard Dynkin diagram as in Example 2.1. The actions we define in Proposition (5.1) coincides with [Mi06, (3.1)(3.2)]. We denote by Φ^{st} (resp. Φ) the homomorphism from ${}^{st}\mathbf{U}_q^{\mathfrak{g}}(\mathfrak{gl}(m|n))$ (resp. \mathbf{U}) to $End(\mathbb{W}^{\otimes d})$. It is not hard to see from the definitions that $\Phi(\mathbf{U}) = \Phi^{st}({}^{st}\mathbf{U}_q^{\mathfrak{g}}(\mathfrak{gl}(m|n)))$. Therefore we have the following theorem.

Theorem 5.2. *The actions of \mathbf{U} and $\mathcal{H}_{\mathfrak{S}_d}$ on $\mathbb{W}^{\otimes d}$ commutes with each other:*

$$\mathbf{U} \overset{\Phi}{\curvearrowright} \mathbb{W}^{\otimes d} \overset{\Psi}{\curvearrowleft} \mathcal{H}_{\mathfrak{S}_d}.$$

Moreover, $\Phi(\mathbf{U})$ and $\Psi(\mathcal{H}_{\mathfrak{S}_d})$ form double centralizers in $End(\mathbb{W}^{\otimes d})$.

Proof. This follows from [Mi06] and the fact that $\Phi(\mathbf{U}) = \Phi^{st}({}^{st}\mathbf{U}_q^{\mathfrak{g}}(\mathfrak{gl}(m|n)))$. \square

Following the strategy of [SW21], we develop a type B ι Schur duality between \mathbf{U}^{ι} and \mathcal{H}_{B_d} . For any reduced expression $\underline{w}_{\bullet}^{(y)} = s_{y_1} \cdots s_{y_{\ell}}$ of w_{\bullet} , we define

$$\Gamma_t^{(y)} = s_{y_t} \cdots s_{y_{\ell}}(\tilde{\Gamma}), \quad 1 \leq t \leq \ell.$$

Note that $\Gamma = \Gamma_1^{(y)}$ for any $\underline{w}_{\bullet}^{(y)}$.

In the next two lemmas we compute explicitly the action of B_j ($j \in I_{\circ}$) on \mathbb{W} .

Lemma 5.3. *For $a \in \mathbb{I}_{r|m|r}$ and $i \in I_{\circ} = [1 - n - r, -n] \cup [n, n + r - 1]$, we have*

$$T_{w_{\bullet}}(\tilde{E}_{\tau(i)})(w_a) = \begin{cases} E_{-i}(w_a), & |i| > n; \\ E_{-n+1}E_{-n+2} \cdots E_{n-1}E_n(w_a), & i = -n; \\ \hbar_m E_{-n}E_{-n+1} \cdots E_{n-2}E_{n-1}(w_a), & i = n. \end{cases}$$

where

$$(5.7) \quad \hbar_m = (-1)^{m-1+p(\alpha_{-n+1}^{\Gamma_{2n-1}})p(\alpha_{-n}^{\Gamma_{2n-1}})+\cdots+p(\alpha_{n-1}^{\Gamma_1})p(\alpha_{n-2}^{\Gamma_1})} q^{(\tilde{\alpha}_{-n+1}, \tilde{\alpha}_{-n})+\cdots+(\tilde{\alpha}_{n-1}, \tilde{\alpha}_{n-2})}.$$

Proof. The computation follows similarly as in [SW21, Lemma 4.2].

Take $i = n$ for example. We choose

$$\underline{w}_{\bullet}^{(y)} = (s_{n-1}s_{n-2} \cdots s_{-n+1}) \cdots (s_{n-1}s_{n-2})(s_{n-1}).$$

For convention we drop (y) in the following proof. Then we compute $T_{w_\bullet}(\tilde{E}_{\tau(n)})(w_a)$ as follows:

$$\begin{aligned}
 & T_{w_\bullet}(\tilde{E}_{\tau(n)})(w_a) \\
 &= T_{s_{n-1}} \cdots T_{s_{-n+1}}(E_{-n}^{\Gamma_{2n}})(w_a) \\
 &= T_{s_{n-1}} \cdots T_{s_{-n+2}}(E_{-n+1}^{\Gamma_{2n-1}} E_{-n}^{\Gamma_{2n-1}} \\
 &\quad - (-1)^{p(\alpha_{-n+1}^{\Gamma_{2n-1}})p(\alpha_{-n}^{\Gamma_{2n-1}})} q^{(\alpha_{-n+1}^{\Gamma_{2n-1}}, \alpha_{-n}^{\Gamma_{2n-1}})} E_{-n}^{\Gamma_{2n-1}} E_{-n+1}^{\Gamma_{2n-1}}) w_a \\
 &= - (-1)^{p(\alpha_{-n+1}^{\Gamma_{2n-1}})p(\alpha_{-n}^{\Gamma_{2n-1}})} q^{(\tilde{\alpha}_{-n+1}, \tilde{\alpha}_{-n})} E_{-n} T_{s_{n-1}} \cdots T_{s_{-n+2}}(E_{-n+1}^{\Gamma_{2n-1}})(w_a)
 \end{aligned}$$

By induction on n , recalling $m = 2n$ we have

$$T_{w_\bullet}(\tilde{E}_{\tau(n)})(w_a) = \hbar_m E_{-n} E_{-n+1} \cdots E_{n-2} E_{n-1}(w_a).$$

This proves the lemma. \square

Lemma 5.3 together with the formula for B_j immediately imply the following.

Lemma 5.4. *Let $a \in \mathbb{I}_{r|m|r}$ and $j \in I_\circ$. The action of B_j on \mathbb{W} is given by:*

$$B_{-n}(w_a) = \begin{cases} w_{-n+\frac{1}{2}}, & \text{if } a = -n - \frac{1}{2}; \\ \varsigma_{-n} w_{-n+\frac{1}{2}}, & \text{if } a = n + \frac{1}{2}; \\ 0, & \text{else,} \end{cases}$$

$$B_i(w_a) = \begin{cases} w_{i+\frac{1}{2}}, & \text{if } a = i - \frac{1}{2}; \\ \varsigma_i w_{-i-\frac{1}{2}}, & \text{if } a = -i + \frac{1}{2}; \\ 0, & \text{else,} \end{cases} \quad \text{for } |i| > n,$$

and (recall $m = 2n$)

$$B_n(w_a) = \begin{cases} w_{n+\frac{1}{2}} + q^{(-1)^{p(w_{n-\frac{1}{2}})}} \hbar_m \varsigma_n w_{-n-\frac{1}{2}}, & \text{if } a = n - \frac{1}{2}; \\ 0, & \text{else.} \end{cases}$$

Remark 5.5. When $p(j) = 0$ for all $j \in I_\bullet$, the computations can be greatly simplified and we have

$$\hbar_m = (-1)^{m-1} q^{(-1)^{p(w_{n-\frac{1}{2}})}(m-1)}.$$

For the rest of this section we fix the parameters to be

$$(5.8) \quad \begin{cases} \varsigma_j = (-1)^{p(j)}, & \text{if } j \neq \pm n, \\ \varsigma_{-n} = (-1)^{p(w_{n+\frac{1}{2}})} Q, \\ \varsigma_n = (-1)^{p(w_{n+\frac{1}{2}})} q^{(-1)^{p(w_{n-\frac{1}{2}})}} Q^{-1} \hbar_m^{-1}. \end{cases} \quad \text{where } m = 2n \in \mathbb{Z}_{\geq 1},$$

Introduce the $\mathbb{Q}(Q, q)$ -subspaces of \mathbb{W} :

$$\begin{aligned}\mathbb{W}_- &= \bigoplus_{a \in \mathbb{I}_\circ^+} \mathbb{Q}(Q, q)(w_a - (-1)^{p(w-a)} Q w_{-a}), & \mathbb{W}_\bullet &= \bigoplus_{a \in \mathbb{I}_\bullet} \mathbb{Q}(Q, q)w_a, \\ \mathbb{W}_+ &= \bigoplus_{a \in \mathbb{I}_\circ^+} \mathbb{Q}(Q, q)(w_a + (-1)^{p(w-a)} Q^{-1} w_{-a}).\end{aligned}$$

Lemma 5.6. *Assume (5.8). Then \mathbb{W}_- and $\mathbb{W}_\bullet \oplus \mathbb{W}_+$ are \mathbf{U}^\natural -submodules of \mathbb{W} . Hence, we have a \mathbf{U}^\natural -module decomposition $\mathbb{W} = (\mathbb{W}_\bullet \oplus \mathbb{W}_+) \oplus \mathbb{W}_-$.*

Proof. Follows by a direct computation using the formulas (5.5) and Lemma 5.4. \square

The decomposition of \mathbb{W} above is also compatible with the H_0 -action.

Lemma 5.7. *The Hecke generator H_0 acts on \mathbb{W}_- as $(-Q^{-1})\text{Id}$ and acts on $\mathbb{W}_\bullet \oplus \mathbb{W}_+$ as $Q \cdot \text{Id}$.*

Proof. Follows by Lemma 5.4. \square

Theorem 5.8. *Suppose the parameters satisfy (5.8). Then the actions of \mathbf{U}^\natural and \mathcal{H}_{B_d} on $\mathbb{W}^{\otimes d}$ commutes with each other:*

$$\mathbf{U}^\natural \overset{\Psi}{\curvearrowright} \mathbb{W}^{\otimes d} \overset{\Phi}{\curvearrowleft} \mathcal{H}_{B_d}.$$

Moreover, $\Psi(\mathbf{U}^\natural)$ and $\Phi(\mathcal{H}_{B_d})$ form double centralizers in $\text{End}(\mathbb{V}^{\otimes d})$.

Proof. By Theorem 5.2, we know that the action of \mathbf{U} commutes with the action of H_i , for $1 \leq i \leq d-1$. Thus, to show the commuting actions of \mathbf{U}^\natural and \mathcal{H}_{B_d} , it remains to check the commutativity of the actions of H_0 and the generators of \mathbf{U}^\natural .

To that end, it suffices to consider $d = 1$ (thanks to the coideal property of \mathbf{U}^\natural and the fact that the action of H_0 depends solely on the first tensor factor). In this case, the commutativity between \mathbf{U}^\natural -action and H_0 -action on \mathbb{W} follows directly from Lemmas 5.6 and 5.7.

The double centralizer property is equivalent to a multiplicity-free decomposition of $\mathbb{W}^{\otimes d}$ as an $\mathbf{U}^\natural \otimes \mathcal{H}_{B_d}$ -module, which reduces by a deformation argument to the $q = 1$ setting. At the specialization $q \mapsto 1$, \mathbf{U}^\natural becomes the enveloping algebra of a direct sum of two type A Lie superalgebras, $\mathbb{W} = (\mathbb{W}_\bullet \oplus \mathbb{W}_+) \oplus \mathbb{W}_-$ becomes the natural representation of it, on which $s_0 \in W_d$ acts as $(\text{Id}_{\mathbb{W}_\bullet \oplus \mathbb{W}_+}, -\text{Id}_{\mathbb{W}_-})$. The multiplicity-free decomposition of $\mathbb{W}^{\otimes d}$ at $q = 1$ can be established by a standard approach as in [CW12, Theorem 3.9]. \square

6. QUASI K -MATRIX

Quasi K -matrices were first formulated in [BW18a] as an intertwiner between the embedding $\mathbf{U}^\natural \rightarrow \mathbf{U}$ and a bar-involution conjugated embedding (with constraints on the parameters); a proof of greater generality was given in [BK19]. Appel and Vlaar [AV20] reformulated the intertwining relation and removed the constraints as a consequence. Moreover, Wang and Zhang introduced new intertwining relations of the quasi K -matrix and proved the factorisation of the quasi K -matrix into products of rank 1 quasi K -matrix in [WZ22].

From this section on we impose one extra condition on the Satake diagrams we are working with:

$$(6.1) \quad p(j) = 0, \quad \forall j \in I_\bullet.$$

6.1. Preparation. Under the assumption (6.1), the braid group operators T_i for $i \in I_\bullet$ reduce to the ones of Lusztig's and we do not need to work with different presentations of \mathbf{U} anymore.

Suppose $\Gamma \in \mathcal{D}_{m,n}$ satisfy (4.2) and (6.1). Again we let $\mathbf{U}(\Gamma)$ denote the quantum supergroup with generators $\varrho, E_j, F_j, K_j, j \in I$ associated to Γ . Recall that $\mathbf{U}^\iota(\Gamma)$ is the $\mathbb{Q}(q)$ -subalgebra of $\mathbf{U}(\Gamma)$ generated by q^μ ($\mu \in Y^\iota$), E_j, F_j ($j \in I_\bullet$), ϱ and

$$(6.2) \quad B_j = F_j + \varsigma_j T_{w_\bullet}(E_{\tau_j}) K_j^{-1}, \quad \text{for } j \in I_\circ.$$

Abusing the notation τ , the diagram involution τ gives rise to the following involution on \mathbf{U} :

Proposition 6.1. *Under the assumption (6.1), there is an involution τ on \mathbf{U} such that*

$$(6.3) \quad \tau(E_j) = E_{\tau_j}, \quad \tau(F_j) = F_{\tau_j}, \quad \tau(K_j) = (-1)^{p(j)} K_{\tau_j}, \quad \tau(\varrho) = \varrho$$

for all $j \in I$.

Proof. The proof follows from checking on the generators. \square

For future use we let $\mathbf{U}^{\varrho, \geq}$ (resp. $\mathbf{U}^{\varrho, \leq}$) denote the subalgebra generated by \mathbf{U}^+ (resp. \mathbf{U}^-), and \mathbf{U}^0 .

We first recall some basic constructions from [CHW13]. The super skew derivations (cf. [CHW13, §1.5]) ${}_i r$ and on \mathbf{U}^+ satisfy $r_i(E_j) = \delta_{i,j}$, ${}_i r(E_j) = \delta_{i,j}$, and

$$(6.4) \quad \begin{aligned} {}_i r(xy) &= (-1)^{p(y)p(i)} {}_i r(x)y + q^{(\alpha_i, \mu)} x {}_i r(y), \\ r_i(xy) &= (-1)^{p(y)p(i)} q^{(\alpha_i, \nu)} r_i(x)y + x r_i(y) \end{aligned}$$

for all $x \in \mathbf{U}_\mu^+$, $y \in \mathbf{U}_\nu^+$.

Moreover, there is a bilinear pairing $\langle \cdot, \cdot \rangle$ on $\mathbf{U}^{\varrho, \leq} \times \mathbf{U}^{\varrho, \geq}$ such that for all $x, x' \in \mathbf{U}^{\varrho, \geq}$, $y, y' \in \mathbf{U}^{\varrho, \leq}$ and $\mu, \nu \in X$, we have

$$(6.5) \quad \begin{aligned} \langle y, x x' \rangle &= \langle \Delta(y), x' \otimes x \rangle, & \langle y y', x \rangle &= \langle y \otimes y', \Delta(x) \rangle, \\ \langle q^\mu, q^\nu \rangle &= q^{-(\mu, \nu)}, & \langle F_j, E_k \rangle &= \delta_{j,k}, \\ \langle K_\mu, E_j \rangle &= 0, & \langle F_j, K_\mu \rangle &= 0, \\ \langle K_\mu, \varrho \rangle &= \langle \varrho, \varrho \rangle = 1, & \langle E_j, \varrho \rangle &= \langle F_j, \varrho \rangle = 0. \end{aligned}$$

The next lemma is a super analogue of [Jan95, §6.14].

Lemma 6.2. *For all $x \in \mathbf{U}^+$, $y \in \mathbf{U}^-$ and $j \in I$ one has*

$$(6.6) \quad \langle F_i y, x \rangle = (-1)^{p(x)p(j)} \langle F_i, E_i \rangle \langle y, {}_i r(x) \rangle, \quad \langle y F_i, x \rangle = \langle F_i, E_i \rangle \langle y, r_i(x) \rangle.$$

Proof. Suppose $x \in \mathbf{U}_\mu^+$, then we have that

$$\begin{aligned} \Delta(x) &= x \otimes 1 + \sum_{i \in I} r_i(x) q^{p(i)} K_i \otimes E_i + (\text{rest})_1, \\ \Delta(x) &= q^{p(x)} K_\mu \otimes x + \sum_{i \in I} (-1)^{p(x)p(i)} E_i q^{p(\mu - \alpha_i)} K_{\mu - \alpha_i} \otimes {}_i r(x) + (\text{rest})_2 \end{aligned}$$

where $(\text{rest})_1, (\text{rest})_2 \in \varrho \mathbf{U}_{\mu-\nu}^+ K_\nu \otimes \mathbf{U}_\nu^+$ or $\mathbf{U}_{\mu-\nu}^+ K_\nu \otimes \mathbf{U}_\nu^+$ with $\nu > 0$, $\nu \notin \Pi$. Hence the lemma follows from (6.5). \square

The next lemma is a crucial ingredient to construct the quasi K -matrix.

Lemma 6.3. *For all $x \in \mathbf{U}^+$, we have*

$$(6.7) \quad [x, F_j] = xF_j - (-1)^{p(x)p(j)} F_j x = \frac{1}{q^{\ell_j} - q^{-\ell_j}} (r_j(x)K_j - K_j^{-1}{}_j r(x)).$$

Proof. We induct on $ht(x)$. When $x = E_j$ for some $j \in I$, (6.7) follows from the definition. Now if $x = uv$ where $ht(u) < ht(x)$ and $ht(v) < ht(x)$. Then we have

$$\begin{aligned} uvF_j &= (-1)^{p(v)p(i)} uF_j v + \frac{ur_j(v)K_j - uK_j^{-1}{}_j r(v)}{q^{\ell_j} - q^{-\ell_j}} \\ &= (-1)^{p(uv)p(j)} F_j uv + \frac{ur_j(v)K_j - q^{(\alpha_j, |u|)} K_j^{-1} u_j r(v)}{q^{\ell_j} - q^{-\ell_j}} \\ &\quad + (-1)^{p(v)p(j)} \frac{r_j J(u)K_j v - K_j^{-1}{}_j r(u)v}{q^{\ell_j} - q^{-\ell_j}} \\ &= (-1)^{p(uv)p(j)} F_j uv + \frac{r_j(uv)K_j - K_j^{-1}{}_i r(uv)}{q^{\ell_j} - q^{-\ell_j}}. \end{aligned}$$

□

6.2. A recursive formula. Let X_0^\pm denote the subset of X^+ consists of weights of even parity. Extending (4.6), we write $B_i = F_i$ for $i \in I_\bullet$. Following the intrinsic intertwining relation [WZ22, (3.20)] (also cf. [AV20, Ko21]), we establish the next lemma.

Lemma 6.4. *Let $\Upsilon = \sum_{\mu \in X_0^+} \Upsilon_\mu$ with $\Upsilon_\mu \in \mathbf{U}_\mu^+$, the following are equivalent.*

(1) *For all $i \in I$, we have*

$$(6.8) \quad B_i \Upsilon = \Upsilon \tau \circ \sigma(B_{\tau i}).$$

(2) *For all $i \in I$, we have*

$$(6.9) \quad B_i \Upsilon = \Upsilon(F_i + (-1)^{(2\rho_\bullet, \alpha_i)} q^{(\alpha_i, 2\rho_\bullet + w_\bullet \alpha_{\tau i})} \varsigma_{\tau i} \overline{T_{w_\bullet}(E_{\tau i})} K_i).$$

(3) *The element Υ satisfy the following relations:*

$$(6.10) \quad \begin{aligned} r_i(\Upsilon_\mu) &= -(q^{\ell_i} - q^{-\ell_i}) \Upsilon_{\mu - \alpha_i - w_\bullet(\alpha_{\tau i})} (-1)^{(2\rho_\bullet, \alpha_i)} q^{(\alpha_i, 2\rho_\bullet + w_\bullet \alpha_{\tau i})} \varsigma_{\tau i} \overline{T_{w_\bullet}(E_{\tau i})}, \\ {}_i r(\Upsilon_\mu) &= -(q^{\ell_i} - q^{-\ell_i}) q^{(\alpha_i, w_\bullet(\alpha_{\tau i}))} \varsigma_i T_{w_\bullet}(E_{\tau i}) \Upsilon_{\mu - \alpha_i - w_\bullet(\alpha_{\tau i})}. \end{aligned}$$

Moreover, if these relations hold then additionally we have

$$(6.11) \quad x\Upsilon = \Upsilon x \text{ for all } x \in \mathbf{U}^{u0} \mathbf{U}_\bullet.$$

and

$$(6.12) \quad \Upsilon_\mu = 0 \text{ unless } w_\bullet \tau(\mu) = \mu.$$

Proof. Note that

$$\tau \circ \sigma(B_{\tau i}) = F_i + \varsigma_{\tau i} K_i T_{w_\bullet}^{-1}(E_{\tau i}).$$

Thus the equivalence of (1) and (2) follows from [BW18b, Lemma 4.17]. The equivalence of (2) and (3) follows from Lemma 6.3. Moreover, (6.12) follows from an induction argument on $ht(\mu)$; cf. [BK19, Proposition 6.1]. □

The next lemma tells us that every non-vanishing term of the quasi K -matrix is supposed to have parity 0.

Lemma 6.5. *For $\mu \in X^+$, if $w_\bullet \tau(\mu) = \mu$, then $p(\mu) = 0$.*

Proof. Let's use induction on $ht(\mu)$. Now we can write $\mu = \sum_{t=1}^{\ell} a_t \alpha_{j_t}$ where $a_t > 0$ for $1 \leq t \leq \ell$. Now if all α_{j_t} are even roots, then we have $p(\mu) = 0$. On the other hand, suppose $p(\alpha_{j_1}) = 1$. Since $w_\bullet \tau(\mu) = \mu$, we have $\mu' = \mu - (\alpha_{j_1} + w_\bullet \tau(\alpha_{j_1})) \in X^+$ and $w_\bullet \tau(\mu') = \mu'$ and $ht(\mu') < ht(\mu)$. Thus by the inductive hypothesis we have $p(\mu') = 0$. Also, according to (4.2), we have $p(\alpha_{j_1}) = p(w_\bullet \tau(\alpha_{j_1}))$. Thus we have $p(\mu) = 0$ as well. \square

Define

$$(6.13) \quad \zeta'_i = (-1)^{(2\rho_\bullet, \alpha_i)} q^{(\alpha_i, 2\rho_\bullet + w_\bullet \alpha_{\tau i})} \zeta_{\tau i}, \text{ for all } i \in I.$$

The system of equations (6.10) for all $i \in I$ gives an equivalent condition for the Υ to exist and we want to solve it recursively by the following Proposition.

Proposition 6.6. (cf. [BK19, Proposition 6.3]) *Let $\mu \in X_0^+$ with $ht(\mu) \geq 2$ and fix $A_i, {}_i A \in \mathbf{U}_{\mu - \alpha_i}^+$ for all $i \in I$. The following are equivalent.*

(1) *There exists an element $\Xi \in \mathbf{U}_\mu^+$ such that*

$$r_i(\Xi) = A_i, \quad {}_i r(\Xi) = {}_i A, \quad \forall i \in I.$$

(2) *The elements A_i and ${}_i A$ satisfy the following properties.*

(2a) *For all $i, j \in I$, we have*

$$(6.14) \quad r_i({}_j A) = (-1)^{p(i)p(j)} {}_i r(A_j).$$

(2b) *For all $i \in I_{\overline{1}}$, we have*

$$(6.15) \quad \langle F_i, A_i \rangle = 0.$$

(2c) *For all $i \approx j \in I$, we have*

$$(6.16) \quad \langle F_i, A_j \rangle = (-1)^{p(i)p(j)} \langle F_j, A_i \rangle.$$

(2d) *For all $i \in I_{\overline{0}}$ and $j \sim i$, we have*

$$(6.17) \quad \langle F_i^2, A_j \rangle - [2] \langle F_i F_j, A_i \rangle + \langle F_j F_i, A_i \rangle = 0.$$

(2e) *For all $i \in I_{\overline{1}}$ and $j \sim i \sim k$, we have*

$$(6.18) \quad \begin{aligned} [2] \langle F_i F_k F_j, A_i \rangle &= (-1)^{p(j)} \langle F_i F_k F_i, A_j \rangle + (-1)^{p(j)+p(j)p(k)} \langle F_j F_i F_k, A_i \rangle \\ &+ (-1)^{p(k)} \langle F_i F_j F_i, A_k \rangle + (-1)^{p(k)+p(j)p(k)} \langle F_k F_i F_j, A_i \rangle. \end{aligned}$$

Proof. Follows by a rerun of proof of [BK19, Proposition 6.3]. \square

6.3. Technical Lemmas. Now in order to construct the quasi K -matrix Υ recursively, it suffices to show that

$$(6.19) \quad \begin{aligned} A_i &= -(q^{\ell_i} - q^{-\ell_i}) \Upsilon_{\mu - \alpha_i - w_\bullet \alpha_{\tau i}} \zeta'_i \overline{T_{w_\bullet}(E_{\tau i})}, \\ {}_i A &= -(q^{\ell_i} - q^{-\ell_i}) q^{(\alpha_i, w_\bullet \alpha_{\tau i})} \zeta_i T_{w_\bullet}(E_{\tau i}) \Upsilon_{\mu - \alpha_i - w_\bullet \alpha_{\tau i}} \end{aligned}$$

for all $i \in I$ satisfy relations (6.14)–(6.18). Following the strategy from [BK19] we develop several lemmas as follows.

Lemma 6.7. For any $i, j \in I$, we have

$$r_i \circ_j r = (-1)^{p(i)p(j)} {}_j r \circ r_i.$$

Proof. If $u = E_k$ or 1, then we certainly have $r_i \circ_j r(u) = (-1)^{p(i)p(j)} {}_j r \circ r_i(u)$. Thus it's enough to show that $r_i \circ_j r(xy) = {}_j r \circ r_i(xy)$ for any $x \in \mathbf{U}_\mu^+$, $y \in \mathbf{U}_\nu^+$.

We have

$$\begin{aligned} & r_i \circ_j r(xy) \\ &= r_i((-1)^{p(y)p(j)} {}_j r(x)y + q^{(\alpha_j, \mu)} x {}_j r(y)) \\ &= (-1)^{p(y)p(j)} [(-1)^{p(y)p(i)} q^{(\alpha_i, \nu)} r_i \circ_j r(x)y + {}_j r(x)r_i(y)] \\ &\quad + q^{(\alpha_j, \mu)} [(-1)^{p(jr(y))p(i)} q^{(\alpha_i, \nu - \alpha_j)} r_i(x) {}_j r(y) + x r_i \circ_j r(y)], \\ & {}_j r \circ r_i(xy) \\ &= {}_j r((-1)^{p(y)p(i)} q^{(\alpha_i, \nu)} r_i(x)y + x r_i(y)) \\ &= (-1)^{p(y)p(i)} q^{(\alpha_i, \nu)} [(-1)^{p(y)p(j)} {}_j r \circ r_i(x)y + q^{(\alpha_j, \mu - \alpha_i)} r_i(x) {}_j r(y)] \\ &\quad + [(-1)^{p(ir(y))p(j)} {}_j r(x)r_i(y) + q^{(\alpha_j, \mu)} x {}_j r \circ r_i(y)]. \end{aligned}$$

Now since $p({}_k r(y)) = p(y) \pm p(k)$ for any $k \in I$, we have $r_i \circ_j r = (-1)^{p(i)p(j)} {}_j r \circ r_i$. \square

Lemma 6.8. For all $u \in \mathbf{U}_\mu^+$, we have

$$(6.20) \quad \sigma \circ_i r(u) = (-1)^{p(i)(p(u)+1)} r_i \circ \sigma(u).$$

Proof. We prove by induction on $ht(\mu)$. When $u = E_j$ or $u = 1$ the equality holds by definition. Now suppose $x \in U_{\mu_1}^{\rho, +}$, $y \in \mathbf{U}_{\mu_2}^+$ where $u = xy$, $\mu = \mu_1 + \mu_2$ and $\mu_1, \mu_2 > 0$. Then we have

$$\begin{aligned} \sigma \circ_i r \circ \sigma(xy) &= \sigma((-1)^{p(x)p(i)} {}_i r \circ \sigma(y)\sigma(x) + q^{(\alpha_i, \mu_2)} \sigma(y) {}_i r \circ \sigma(x)) \\ &= (-1)^{p(x)p(i)} x \sigma \circ_i r \circ \sigma(y) + q^{(\alpha_i, \mu_2)} \sigma \circ_i r \circ \sigma(x)y \\ &= (-1)^{p(xy)p(i)+p(i)} r_i(xy). \end{aligned}$$

This proves the lemma. \square

Lemma 6.9. For all $x \in \mathbf{U}_\mu^+$, we have

$$(6.21) \quad \overline{r}_i(x) = q^{(\alpha_i, \alpha_i - \mu)} {}_i r(x).$$

Proof. We prove by induction on $ht(\mu)$. When $u = E_j$ or $u = 1$ the equality holds by definition. Now suppose $x \in U_{\mu_1}^+$, $y \in \mathbf{U}_{\mu_2}^+$ where $u = xy$, $\mu = \mu_1 + \mu_2$ and $\mu_1, \mu_2 > 0$. Then we have

$$\begin{aligned} \overline{r}_i(xy) &= x \overline{r}_i(y) + (-1)^{p(y)p(i)} q^{-(\alpha_i, \mu_2)} \overline{r}_i(x)y \\ &= q^{(\alpha_i, \alpha_i - \mu_2)} x {}_i r(y) + (-1)^{p(y)p(i)} q^{(\alpha_i, \alpha_i - \mu_1 - \mu_2)} {}_i r(x)y \\ &= q^{(\alpha_i, \alpha_i - \mu)} [q^{(\alpha_i, \mu_1)} {}_i r(y) + (-1)^{p(y)p(i)} {}_i r(x)y] \\ &= q^{(\alpha_i, \alpha_i - \mu)} {}_i r(xy). \end{aligned}$$

This proves the lemma. \square

Lemma 6.10. *For all $i \in I_o$, we have*

$$(6.22) \quad \overline{r_i(T_{w_\bullet}(E_i))} = (-1)^{(\alpha_i, 2\rho_\bullet)} q^{(\alpha_i, \alpha_i - w_\bullet \alpha_i - 2\rho_\bullet)} \sigma \circ \tau(r_{\tau i}(T_{w_\bullet}(E_{\tau i}))).$$

Proof. Follow from a rerun of the proof of [BK15, Lemma 2.9]. \square

Lemma 6.11. *For all $i \in I_o$, we have*

$$(6.23) \quad \sigma \circ \tau(r_i(T_{w_\bullet}(E_i))) = r_i(T_{w_\bullet}(E_i)).$$

Proof. Follow from a rerun of the proof of [BK15, Proposition 2.3]. \square

Combining Lemma 6.10 and Lemma 6.11 we get

Corollary 6.12. *For all $i \in I_o$, we have*

$$(6.24) \quad \overline{r_i(T_{w_\bullet}(E_i))} = (-1)^{(\alpha_i, 2\rho_\bullet)} q^{(\alpha_i, \alpha_i - w_\bullet \alpha_i - 2\rho_\bullet)} r_{\tau i}(T_{w_\bullet}(E_{\tau i})).$$

6.4. Construction of Υ . Now we are ready to check that

$$(6.25) \quad \begin{aligned} A_i &= -(q^{\ell_i} - q^{-\ell_i}) \Upsilon_{\mu - \alpha_i - w_\bullet(\alpha_{\tau i})} \zeta_i' \overline{T_{w_\bullet}(E_{\tau i})}, \\ {}_i A &= -(q^{\ell_i} - q^{-\ell_i}) q^{(\alpha_i, w_\bullet(\alpha_{\tau i}))} \zeta_i T_{w_\bullet}(E_{\tau i}) \Upsilon_{\mu - \alpha_i - w_\bullet(\alpha_{\tau i})} \end{aligned}$$

for all $i \in I$ indeed satisfy relations (6.14)–(6.18).

Lemma 6.13. *The relation $r_i({}_j A) = (-1)^{p(i)p(j)} {}_j r(A_i)$ holds for all $i, j \in \mathbb{I}$.*

Proof. We calculate that

$$\begin{aligned} & \frac{1}{-(q^{\ell_j} - q^{-\ell_j})} r_i({}_j A) \\ &= q^{(\alpha_j, w_\bullet \alpha_{\tau j})} \zeta_j [(-1)^{p(i)p(\mu)} q^{(\alpha_i, \mu - \alpha_j - w_\bullet \alpha_{\tau j})} r_i(T_{w_\bullet}(E_{\tau j})) \Upsilon_{\mu - \alpha_j - w_\bullet \alpha_{\tau j}} \\ & \quad + T_{w_\bullet}(E_{\tau j}) r_i(\Upsilon_{\mu - \alpha_j - w_\bullet \alpha_{\tau j}})] \\ &= q^{(\alpha_j, w_\bullet \alpha_{\tau j})} \zeta_j [(-1)^{p(i)p(\mu)} q^{(\alpha_i, \mu - \alpha_j - w_\bullet \alpha_{\tau j})} r_i(T_{w_\bullet}(E_{\tau j})) \Upsilon_{\mu - \alpha_j - w_\bullet \alpha_{\tau j}} \\ & \quad - (q^{\ell_i} - q^{-\ell_i}) T_{w_\bullet}(E_{\tau j}) \Upsilon_{\mu - \alpha_j - w_\bullet \alpha_{\tau j} - \alpha_i - w_\bullet(\alpha_{\tau i})} \zeta_i' \overline{T_{w_\bullet}(E_{\tau i})}], \\ & \frac{1}{-(q^{\ell_i} - q^{-\ell_i})} {}_j r(A_i) \\ &= (-1)^{p(w_\bullet(\alpha_{\tau i}))p(j)} {}_j r(\Upsilon_{\mu - \alpha_i - w_\bullet(\alpha_{\tau i})} \zeta_i' \overline{T_{w_\bullet}(E_{\tau i})} \\ & \quad + q^{(\alpha_j, \mu - \alpha_i - w_\bullet(\alpha_{\tau i}))} \zeta_i' \Upsilon_{\mu - \alpha_i - w_\bullet \alpha_{\tau i}} {}_j r(\overline{T_{w_\bullet}(E_{\tau i})})) \\ &= -(-1)^{p(i)p(j)} (q^{\ell_j} - q^{-\ell_j}) q^{(\alpha_j, w_\bullet \alpha_{\tau j})} \zeta_j T_{w_\bullet}(E_{\tau j}) \cdot \\ & \quad \Upsilon_{\mu - \alpha_j - w_\bullet \alpha_{\tau j} - \alpha_i - w_\bullet \alpha_{\tau i}} \zeta_i' \overline{T_{w_\bullet}(E_{\tau i})} \\ & \quad + q^{(\alpha_j, \mu - \alpha_i - w_\bullet \alpha_{\tau i})} \zeta_i' \Upsilon_{\mu - \alpha_i - w_\bullet(\alpha_{\tau i})} {}_j r(\overline{T_{w_\bullet}(E_{\tau i})}) \end{aligned}$$

Recall that Υ_μ vanishes whenever $p(\mu) = 1$. By comparing the two equations we see that the relation $r_i({}_j A) = (-1)^{p(i)p(j)} {}_j r(A_i)$ holds if and only if

$$(6.26) \quad \begin{aligned} & q^{(\alpha_j, w_\bullet \alpha_{\tau j})} \zeta_j q^{(\alpha_i, \mu - \alpha_j - w_\bullet \alpha_{\tau j})} r_i(T_{w_\bullet}(E_{\tau j})) \Upsilon_{\mu - \alpha_j - w_\bullet \alpha_{\tau j}} \\ &= (-1)^{p(i)p(j)} \frac{q^{\ell_j} - q^{-\ell_j}}{q^{\ell_i} - q^{-\ell_i}} q^{(\alpha_j, \mu - \alpha_i - w_\bullet \alpha_{\tau i})} \zeta_i' \Upsilon_{\mu - \alpha_i - w_\bullet \alpha_{\tau i}} {}_j r(\overline{T_{w_\bullet}(E_{\tau i})}) \end{aligned}$$

We may assume $i = \tau j$, otherwise both sides of (6.26) vanish. According to (6.21) we have

$$\tau_i r(\overline{T_{w_\bullet}(E_{\tau i})}) = q^{(\alpha_{\tau i}, w_\bullet \alpha_{\tau i} - \alpha_{\tau i})} \overline{r_{\tau i}(T_{w_\bullet}(E_{\tau i}))}.$$

Substituting this together with Lemma 4.1 we see that (6.26) is equivalent to

$$(6.27) \quad \begin{aligned} & \varsigma_{\tau i} q^{(\alpha_i, \mu - \alpha_{\tau i} + w_\bullet \alpha_{\tau i} - w_\bullet \alpha_i)} r_i(T_{w_\bullet}(E_i)) \Upsilon_{\mu - \alpha_{\tau i} - w_\bullet \alpha_i} \\ &= q^{(\alpha_{\tau i}, \mu - \alpha_i - \alpha_{\tau i})} \varsigma_i' \Upsilon_{\mu - \alpha_i - w_\bullet \alpha_{\tau i}} \overline{r_{\tau i}(T_{w_\bullet}(E_{\tau i}))} \end{aligned}$$

Observe that $\mu - \alpha_{\tau i} - w_\bullet \alpha_i = \mu - \alpha_i - w_\bullet \alpha_{\tau i}$ and $w_\bullet \alpha_i - w_\bullet \alpha_{\tau i} = \alpha_i - \alpha_{\tau i}$. We may further assume that $\Upsilon_{\mu - \alpha_{\tau i} - w_\bullet(\tilde{\alpha}_i)} \neq 0$, thus $w_\bullet \tau(\mu) = \mu$ and hence $(\alpha_i - \alpha_{\tau i}, \mu) = 0$. Thus we see (6.27) is equivalent to

$$(6.28) \quad \begin{aligned} & q^{(\alpha_{\tau i}, \alpha_i)} \varsigma_{\tau i} r_i(T_{w_\bullet}(E_i)) = \varsigma_i' \overline{r_{\tau i}(T_{w_\bullet}(E_{\tau i}))} \\ & \stackrel{(6.24)}{=} \varsigma_i' (-1)^{(\alpha_{\tau i}, 2\rho_\bullet)} q^{(\alpha_{\tau i}, \alpha_{\tau i} - w_\bullet \alpha_{\tau i} - 2\rho_\bullet)} r_i(T_{w_\bullet}(E_i)), \end{aligned}$$

which follows from the definition of ς_i' (6.13). \square

The next lemma verifies the relation (6.15).

Lemma 6.14. *For all $i \in I$, we have*

$$\langle F_i, A_i \rangle = 0.$$

Proof. Since $wt(A_i) = \mu - \alpha_i$. We see that $\langle F_i, A_i \rangle$ is zero unless $\mu = 2\alpha_i$. But in this case we always have $\mu - \alpha_i - w_\bullet \alpha_{\tau i} \notin X^+$. Hence $A_i = 0$. \square

To verify the relation (6.16), we have

Lemma 6.15. *For all $i \approx j \in I$, we have*

$$\langle F_i, A_j \rangle = (-1)^{p(i)p(j)} \langle F_j, A_i \rangle.$$

Proof. According to [BK19, Lemma 6.4], we can assume that $j = \tau i \in I_\circ \setminus \{\pm n\}$ and $\mu = \alpha_i + \alpha_j$, otherwise all terms vanish. In this case we have $\varsigma_i = \varsigma_j$, $A_i = -(q^{\ell_i} - q^{-\ell_i}) \varsigma_i' E_{\tau i}$, and $A_j = -(q^{\ell_j} - q^{-\ell_j}) \varsigma_j' E_{-j}$. Thus we have

$$\langle F_i, A_{\tau i} \rangle = -(q^{\ell_{\tau i}} - q^{-\ell_{\tau i}}) \varsigma_{\tau i}' = -(-1)^{p(i)} (q^{\ell_i} - q^{-\ell_i}) \varsigma_i' = (-1)^{p(i)} \langle F_{\tau i}, A_i \rangle.$$

This proves the lemma. \square

To verify the relation (6.17), we have

Lemma 6.16. *For all $i \in I_\overline{0}$ and $j \sim i$, we have*

$$\langle F_i^2, A_j \rangle - [2] \langle F_i F_j, A_i \rangle + \langle F_j F_i, A_i \rangle = 0.$$

Proof. First observe that we can assume $\mu = 2\alpha_i + \alpha_j$, otherwise all terms in the above sum vanish. But in this case we see that $w_\bullet \circ \tau(\mu) \neq \mu$ for all $j \sim i \in I$. Hence all terms still vanish. \square

To verify the relation (6.18), we have

Lemma 6.17. *For all $i \in I_{\overline{1}}$ and $j \sim i \sim k$, we have*

$$\begin{aligned} [2]\langle F_i F_k F_j, A_i \rangle &= (-1)^{p(j)} \langle F_i F_k F_i, A_j \rangle + (-1)^{p(j)+p(j)p(k)} \langle F_j F_i F_k, A_i \rangle \\ &\quad + (-1)^{p(k)} \langle F_i F_j F_i, A_k \rangle + (-1)^{p(k)+p(j)p(k)} \langle F_k F_i F_j, A_i \rangle. \end{aligned}$$

Proof. Again we may assume that $\mu = 2\alpha_i + \alpha_k + \alpha_j$. But in this case we see that $w_{\bullet} \circ \tau(\mu) \neq \mu$ for all $j \sim i \sim k$. Hence all terms still vanish. \square

Therefore, we conclude that

$$\begin{aligned} A_i &= -(q - q^{-1}) \Upsilon_{\mu - \alpha_i - w_{\bullet} \alpha_{\tau i}} \overline{\zeta'_i T_{w_{\bullet}}(E_{\tau i})}, \\ iA &= -(q - q^{-1}) q^{(\alpha_i, w_{\bullet} \alpha_{\tau i})} \zeta'_i T_{w_{\bullet}}(E_{\tau i}) \Upsilon_{\mu - \alpha_i - w_{\bullet} \alpha_{\tau i}} \end{aligned}$$

for all $i \in I$ satisfy relations (6.14)–(6.18).

Thus we can conclude the main result of this section.

Theorem 6.18. *There exists a uniquely determined element $\Upsilon = \sum_{\mu \in X_0^+} \Upsilon_{\mu}$ with $\Upsilon_0 = 1$ and $\Upsilon_{\mu} \in \mathbf{U}_{\mu}^+$, such that the equality*

$$B_i \Upsilon = \Upsilon(\tau \circ \sigma(B_{\tau i}))$$

holds for all $i \in I$.

Moreover, $\Upsilon_{\mu} = 0$ unless $w_{\bullet} \tau(\mu) = \mu$.

Once Υ is constructed, we can define a unique bar involution on \mathbf{U}^t with certain assumption on the parameters as follows.

Corollary 6.19. *Under the assumption that $\overline{\zeta}_j = \zeta'_j$ for all $j \in I$, there is a unique bar involution ψ_i on \mathbf{U}^t , defined by*

$$\psi_i(x) = \Upsilon \overline{x} \Upsilon^{-1}, \text{ for all } x \in \mathbf{U}^t$$

and such that

$$\psi_i(q^h) = q^{-h}, \psi_i(B_j) = B_j, \psi_i(E_k) = E_k, \psi_i(F_k) = F_k,$$

for $j \in I_{\circ}, k \in I_{\bullet}$.

Proof. For all $i \in I$, it follows from Lemma 6.4 that $B_i \Upsilon = \Upsilon(\tau \circ \sigma(B_{\tau i}))$ is equivalent to (6.9). Under the assumption $\zeta'_i = \overline{\zeta}_i$ we see that (6.9) is equivalent to

$$B_i \Upsilon = \Upsilon \overline{B}_i.$$

\square

Remark 6.20. One can construct Υ associated to more general Satake diagrams. For example, one can replace (6.1) by a weaker condition:

$$p(j) = p(\tau j), \quad \forall j \in I_{\bullet}.$$

Under this assumption formally we still have $w_{\bullet}(\Gamma) = \Gamma$. Thus $T_{w_{\bullet}}$ can still be treated as an automorphism on $\mathbf{U}(\Gamma)$ although it is a composition of even and odd braid group operators.

In the last of this subsection we give an example of Υ .

Example 6.21. Consider the following Satake diagram

$$\begin{array}{c} \otimes \xrightarrow{k-\vartheta} \otimes \\ -\frac{1}{2} \quad \frac{1}{2} \end{array}$$

We have $(\alpha_{\frac{1}{2}}, \alpha_{-\frac{1}{2}}) = 1$, $\ell_{-\frac{1}{2}} = 1 = -\ell_{\frac{1}{2}}$ and

$$\begin{aligned} B_{\frac{1}{2}} &= F_{\frac{1}{2}} + \varsigma_{\frac{1}{2}} E_{-\frac{1}{2}} K_{\frac{1}{2}}^{-1}, \\ B_{-\frac{1}{2}} &= F_{-\frac{1}{2}} + \varsigma_{-\frac{1}{2}} E_{\frac{1}{2}} K_{-\frac{1}{2}}^{-1}. \end{aligned}$$

In this case, following the constructions in this section we get

$$\Upsilon = \left(\sum_{k \geq 0} \frac{(\varsigma_{\frac{1}{2}})^k}{\{k\}!} \right) (E_{\frac{1}{2}} E_{-\frac{1}{2}} + q E_{-\frac{1}{2}} E_{\frac{1}{2}})^k \left(\sum_{k \geq 0} \frac{(\varsigma_{-\frac{1}{2}})^k}{\{k\}!} \right) (E_{-\frac{1}{2}} E_{\frac{1}{2}} + q E_{\frac{1}{2}} E_{-\frac{1}{2}})^k.$$

7. K -MATRIX AND THE H_0 -ACTION

In this subsection we follow [BW18b] (also cf. [BK19]) to construct a \mathbf{U}^ℓ -module intertwiner (or K -matrix).

7.1. K -matrix. Recall that we assume that I_\bullet consists of simple roots of even parity only. We review several basic lemmas from [BW18b] below. Recall σ and \wp from (2.9).

Lemma 7.1. For all $i, j \in I$ and $e = \pm 1$, we have

$$(7.1) \quad \begin{aligned} \wp(T''_{i,e}(E_j)) &= (-q_i)^{e(\alpha_i, \alpha_j)} T'_{i,-e}(\wp(E_j)), \\ \wp(T'_{i,e}(E_j)) &= (-q_i)^{-e(\alpha_i, \alpha_j)} T''_{i,-e}(\wp(E_j)). \end{aligned}$$

Proof. It follows from a rerun of the proof of [BW18b, Lemma 4.4] in the super case. \square

Lemma 7.2. ([BW18b, Corollary 4.5]) For $i \in I_\circ$ and $e = \pm 1$, we have

$$(7.2) \quad \begin{aligned} \wp(T''_{w_\bullet, e}(E_i)) &= (-1)^{(2\rho_\bullet, \alpha_i)} q^{e(2\rho_\bullet, \alpha_i)} T'_{w_\bullet, -e}(\wp(E_i)), \\ \wp(T'_{w_\bullet, e}(E_i)) &= (-1)^{(2\rho_\bullet, \alpha_i)} q^{-e(2\rho_\bullet, \alpha_i)} T''_{w_\bullet, -e}(\wp(E_i)). \end{aligned}$$

Under the assumption (6.1), the anti-involution \wp on \mathbf{U} restricts to an anti-involution \wp on \mathbf{U}^ℓ ; see [BW18b, Proposition 4.6].

Following [BW18b, §4.5], under the assumption (6.1), we define the following automorphism of \mathbf{U} obtained by composition $\vartheta = \sigma \circ \wp \circ \tau$ such that

$$(7.3) \quad \begin{aligned} \vartheta(E_j) &= (-1)^{p(j)} q_{\tau j} F_{\tau j} K_{\tau j}^{-1}, & \vartheta(F_j) &= (-1)^{p(j)} q_{\tau j}^{-1} K_{\tau j} E_{\tau j}, \\ \vartheta(K_j) &= K_{\tau j}^{-1}, & \vartheta(\varrho) &= \varrho, \quad \text{for all } j \in I. \end{aligned}$$

For any finite-dimensional \mathbf{U} -module M , we define a new \mathbf{U} -module ${}^\vartheta M$ as follows:

- ${}^\vartheta M = M$ as an $\mathbb{Q}(q)$ -vector space,
- We denote a vector in ${}^\vartheta M$ by ${}^\vartheta m$ for $m \in M$,
- the action of $u \in \mathbf{U}$ on ${}^\vartheta M$ is given by $\vartheta(u) {}^\vartheta m = {}^\vartheta (um)$.

Let

$$(7.4) \quad g : X \rightarrow \mathbb{Q}(q)$$

be a function such that for all $\mu \in X$, we have the following two recursive relations of g :

$$(7.5) \quad g(\mu) = -q_{\tau_j} q_j^{2(\alpha_j, \mu)} g(\mu + \alpha_j), \quad \forall j \in I_{\bullet}.$$

$$(7.6) \quad g(\mu) = g(\mu - \alpha_j) (-1)^{p(j)} \varsigma_j (-1)^{(2\rho_{\bullet}, \alpha_j)} q^{(2\rho_{\bullet}, \alpha_j)} q_j q^{(\alpha_j, w_{\bullet} \mu)} q^{-(\alpha_j, \mu)}, \quad \forall j \in I_{\circ}.$$

Such a function g exists; cf. [BW18b, (4.15)]. Note that under our assumption (6.1), we always have $q_j = q_{\tau_j}$ for all $j \in I$.

Lemma 7.3. *For any $\mu \in X$, we have*

$$(7.7) \quad g(\mu) = g(\mu - w_{\bullet} \alpha_j) (-1)^{p(j)} \varsigma_j q^{(\alpha_j, \mu)} q_j q^{-(\alpha_j, w_{\bullet} \mu)}, \quad \forall j \in I_{\circ}.$$

Proof. Recall the following identity [BW18b, (4.18)]:

$$(7.8) \quad g(\mu - \alpha_j) = g(\mu - w_{\bullet} \alpha_j) (-1)^{(2\rho_{\bullet}, \alpha_j)} q^{-(2\rho_{\bullet}, \alpha_j)} q^{2(\alpha_j - w_{\bullet} \alpha_j, \mu)}, \quad \forall j \in I_{\circ}.$$

Then applying (7.6) to (7.8) we get (7.7). \square

The function g induces a $\mathbb{Q}(q)$ -linear map on any finite dimensional \mathbf{U} -module M :

$$\tilde{g} : M \rightarrow M, \quad \tilde{g}(m) = g(\mu)m, \quad m \in M_{\mu}.$$

The next theorem gives the construction of the K -matrix.

Theorem 7.4. (cf. [BW18b, Theorem 4.18]) *For any finite-dimensional \mathbf{U} -module M , we have the following isomorphism of \mathbf{U}^{ι} -modules*

$$\mathcal{T} := \Upsilon \circ \tilde{g} \circ T_{w_{\bullet}}^{-1} : M \rightarrow {}^{\vartheta}M.$$

Proof. It suffices to verify that \mathcal{T} defines a homomorphism of \mathbf{U}^{ι} -modules. We shall prove the following identity

$$(7.9) \quad \mathcal{T}(\vartheta(u) \cdot m) = u \cdot \mathcal{T}(m), \quad \text{for } u \in \mathbf{U}^{\iota}, m \in M_{\mu}.$$

It is straightforward to check (7.9) for $u = K_{\mu}, \varrho$. Also, for $u = F_j, E_j$ ($j \in I_{\bullet}$), the proof is essentially the same as that of [BW18b, Case(2)-(3), Theorem 4.18]. Thus we only verify for $u = B_j$ ($j \in I_{\circ}$) as below.

For $u = B_j$ ($j \in I_{\circ}$), first we see that

$$\begin{aligned} \vartheta(B_i) &= (-1)^{p(i)} q_{\tau_i}^{-1} K_{\tau_i} E_{\tau_i} \varsigma_i \sigma \circ \varrho \circ \tau \circ T_{w_{\bullet}}(E_{\tau_i}) K_{\tau_i} \\ &\stackrel{(7.2)}{=} (-1)^{p(i)} q_{\tau_i}^{-1} K_{\tau_i} E_{\tau_i} + \varsigma_i (-1)^{(2\rho_{\bullet}, \alpha_i)} q^{(2\rho_{\bullet}, \alpha_i)} T_{w_{\bullet}}(\sigma \circ \varrho(E_i)) K_{\tau_i} \\ &= (-1)^{p(i)} q_{\tau_i}^{-1} K_{\tau_i} E_{\tau_i} + (-1)^{p(i)} \varsigma_i (-1)^{(2\rho_{\bullet}, \alpha_i)} q^{(2\rho_{\bullet}, \alpha_i)} q_i T_{w_{\bullet}}(F_i) T_{w_{\bullet}}(K_i^{-1}) K_{\tau_i}. \end{aligned}$$

Therefore we have

$$\begin{aligned} T_{w_{\bullet}}^{-1} \circ \vartheta(B_i) &= (-1)^{p(i)} q_{\tau_i}^{-1} T_{w_{\bullet}}^{-1}(K_{\tau_i}) T_{w_{\bullet}}^{-1}(E_{\tau_i}) \\ &\quad + \varsigma_i (-1)^{p(i)} (-1)^{(2\rho_{\bullet}, \alpha_i)} q^{(2\rho_{\bullet}, \alpha_i)} q_i F_i K_i^{-1} T_{w_{\bullet}}^{-1}(K_{\tau_i}). \end{aligned}$$

Now we have

$$\begin{aligned}
\mathcal{J}(\vartheta(B_i)(m)) &= \Upsilon \circ \tilde{g} \circ T_{w_\bullet}^{-1}(\vartheta(B_i)m) = \Upsilon \circ \tilde{g} (T_{w_\bullet}^{-1} \circ \vartheta(B_i)(T_{w_\bullet}^{-1}(m))) \\
&= \Upsilon \circ \tilde{g}((-1)^{p(i)} q_{\tau_i}^{-1} T_{w_\bullet}^{-1}(K_{\tau_i}) T_{w_\bullet}^{-1}(E_{\tau_i}) T_{w_\bullet}^{-1}(m) \\
&\quad + (-1)^{p(i)} \varsigma_i (-1)^{(2\rho_\bullet, \alpha_i)} q^{(2\rho_\bullet, \alpha_i)} q_i F_i K_i^{-1} T_{w_\bullet}^{-1}(K_{\tau_i}) T_{w_\bullet}^{-1}(m)) \\
&= \Upsilon(g(w_\bullet \mu + w_\bullet \alpha_{\tau_i}) (-1)^{p(i)} q_{\tau_i}^{-1} T_{w_\bullet}^{-1}(K_{\tau_i}) T_{w_\bullet}^{-1}(E_{\tau_i}) T_{w_\bullet}^{-1}(m) \\
&\quad + (-1)^{p(i)} g(w_\bullet \mu - \alpha_i) \varsigma_i (-1)^{(2\rho_\bullet, \alpha_i)} q^{(2\rho_\bullet, \alpha_i)} q_i F_i K_i^{-1} T_{w_\bullet}^{-1}(K_{\tau_i}) T_{w_\bullet}^{-1}(m)) \\
&= \Upsilon(g(w_\bullet \mu + w_\bullet \alpha_{\tau_i}) (-1)^{p(i)} q_{\tau_i}^{-1} q^{(\alpha_{\tau_i}, \mu + \alpha_{\tau_i})} T_{w_\bullet}^{-1}(E_{\tau_i}) T_{w_\bullet}^{-1}(m) \\
&\quad + (-1)^{p(i)} g(w_\bullet \mu - \alpha_i) \varsigma_i (-1)^{(2\rho_\bullet, \alpha_i)} q^{(2\rho_\bullet, \alpha_i)} q_i q^{(\alpha_{\tau_i}, \mu)} q^{-(\alpha_i, w_\bullet \mu)} F_i T_{w_\bullet}^{-1}(m)).
\end{aligned}$$

On the other hand, we have

$$B_i \Upsilon = \Upsilon B_{\tau_i}^{\sigma\tau} = \Upsilon (F_i + \varsigma_{\tau_i} K_i T_{w_\bullet}^{-1}(E_{\tau_i})).$$

Thus

$$\begin{aligned}
B_i \cdot \mathcal{J}(m) &= B_i (\Upsilon \circ \tilde{g} \circ T_{w_\bullet}^{-1}(m)) \stackrel{(6.8)}{=} \Upsilon (F_i (\tilde{g} \circ T_{w_\bullet}^{-1}(m)) + \varsigma_{\tau_i} K_i T_{w_\bullet}^{-1}(E_{\tau_i}) (\tilde{g} \circ T_{w_\bullet}^{-1}(m))) \\
&= \Upsilon (g(w_\bullet \mu) F_i T_{w_\bullet}^{-1}(m) \\
&\quad + g(w_\bullet \mu) \varsigma_{\tau_i} q^{(\alpha_i, w_\bullet \mu + w_\bullet \alpha_{\tau_i})} T_{w_\bullet}^{-1}(E_{\tau_i}) T_{w_\bullet}^{-1}(m)).
\end{aligned}$$

Now the identity (7.9) for $u = B_j$ follows from comparing the coefficients using (7.6) and (7.7). \square

7.2. Realizing H_0 via K -matrix. In this subsection the goal is to realize H_0 -action on \mathbb{W} as in Proposition 5.1 via the K -matrix \mathcal{J} .

We assume the parameters to satisfy (5.8) so that the ι Schur duality holds between \mathbf{U}^q and \mathcal{H}_{B_d} , and moreover, Υ and ψ_i in Corollary 6.19 uniquely exist. We also assume that $Q \in q^{\mathbb{Z}}$ thus $\psi_i(Q) = Q^{-1}$.

Recall ϑ from (7.3). A \mathbf{U} -module ${}^\vartheta M$ is simple if and only if M is simple itself. Let λ be a dominant integral weight and $L(\lambda)$ denote the unique irreducible highest weight module with highest weight vector η_λ . Moreover, we define a lowest weight \mathbf{U} -module ${}^\omega L(\lambda)$ of weight $-\lambda$ which has the same underlying vector space as $L(\lambda)$ but with the action twisted by the automorphism ω where

$$(7.10) \quad \omega(E_j) = F_j, \quad \omega(F_j) = (-1)^{p(j)} E_j, \quad \omega(K_\mu) = K_{-\mu}.$$

When we consider η_λ as a vector in ${}^\omega L(\lambda)$, we shall denote it by $\xi_{-\lambda}$. We check by definition that

$${}^\vartheta L(\lambda) \cong {}^\omega L(\tau\lambda).$$

A basic example of $L(\lambda)$ is our fundamental representation $\mathbb{W} = L(\epsilon_{i_{-n-r+\frac{1}{2}}})$. We check by definition that

$$(7.11) \quad {}^\vartheta \mathbb{W} = {}^\vartheta L(\epsilon_{i_{-n-r+\frac{1}{2}}}) \cong {}^\omega L(-\epsilon_{i_{n+r-\frac{1}{2}}}) = L(\epsilon_{i_{-n-r+\frac{1}{2}}}).$$

Recall $\mathcal{J} = \Upsilon \circ \tilde{g} \circ T_{w_\bullet}^{-1}$ and Theorem 7.4. Together with (7.11) we see that \mathcal{J} induces an \mathbf{U}^q -automorphism on \mathbb{W} and send $T_{w_\bullet}^{-1}(\eta_{\epsilon_{i_{-n-r+\frac{1}{2}}}})$ to $\xi_{\epsilon_{i_{n+r-\frac{1}{2}}}}$, cf. [BW18b, Theorem 4.18]. Thus we have the following corollary:

Corollary 7.5. *The K -matrix \mathcal{T} is an \mathbf{U}^ι -module automorphism of \mathbb{W} :*

$$\mathcal{T} : \mathbb{W} \rightarrow \mathbb{W}, \quad w_{-n-r+\frac{1}{2}} \mapsto (-1)^{p(w_{-n-r+\frac{1}{2}})} w_{n+r-\frac{1}{2}}.$$

Proposition 7.6. *The action of H_0 on $\mathbb{W}^{\otimes d}$ in Proposition 5.1 is realized via the K -matrix as $\mathcal{T} \otimes Id^{\otimes d-1}$.*

Proof. According to Corollary 7.5, we have $\mathcal{T}(w_{-n-r+\frac{1}{2}}) = (-1)^{p(w_{-n-r+\frac{1}{2}})} w_{n+r-\frac{1}{2}}$. Recall the parameters satisfy (5.8).

Suppose $a \in \mathbb{I}_\circ^-$, a simple induction on a shows that

$$\begin{aligned} \mathcal{T}(w_a) &= \mathcal{T}(B_{a-\frac{1}{2}} B_{a-\frac{3}{2}} \cdots B_{-n-r+1} w_{-n-r+\frac{1}{2}}) \\ &= B_{a-\frac{1}{2}} B_{a-\frac{3}{2}} \cdots B_{-n-r+1} \mathcal{T}(w_{-n-r+\frac{1}{2}}) = (-1)^{p(w_a)} w_{-a} = w_a \cdot H_0. \end{aligned}$$

Now suppose $a = -n - \frac{1}{2}$, we have

$$\begin{aligned} \mathcal{T}(w_{-n+\frac{1}{2}}) &= \mathcal{T}(B_{-n} w_{-n-\frac{1}{2}}) = B_{-n} \mathcal{T}(w_{-n-\frac{1}{2}}) = (-1)^{p(w_{-n+\frac{1}{2}})} B_{-n} w_{n+\frac{1}{2}} \\ &= Q w_{-n+\frac{1}{2}} = w_{-n+\frac{1}{2}} \cdot H_0. \end{aligned}$$

Thus for any $a \in \mathbb{I}_\bullet$, we have

$$\begin{aligned} \mathcal{T}(w_a) &= \mathcal{T}(F_{a-\frac{1}{2}} F_{a-\frac{3}{2}} \cdots F_{-n+1} w_{-n+\frac{1}{2}}) = F_{a-\frac{1}{2}} F_{a-\frac{3}{2}} \cdots F_{-n+1} \mathcal{T}(w_{-n+\frac{1}{2}}) \\ &= Q w_a = w_a \cdot H_0. \end{aligned}$$

Next we suppose $a = n + \frac{1}{2}$, we have

$$\begin{aligned} \mathcal{T}(w_{n+\frac{1}{2}}) &= \mathcal{T}(B_n(w_{n-\frac{1}{2}}) - (-1)^{p(w_{n+\frac{1}{2}})} Q^{-1} w_{-n-\frac{1}{2}}) \\ &= Q B_n(w_{n-\frac{1}{2}}) - Q^{-1} w_{n+\frac{1}{2}} \\ &= (-1)^{p(w_{n+\frac{1}{2}})} w_{-n-\frac{1}{2}} + (Q - Q^{-1}) w_{n+\frac{1}{2}} = w_{n+\frac{1}{2}} \cdot H_0. \end{aligned}$$

Thus for any $a \in \mathbb{I}_\circ^+$, another simple induction on a shows that

$$\begin{aligned} \mathcal{T}(w_a) &= \mathcal{T}(B_{a-\frac{1}{2}} B_{a-\frac{3}{2}} \cdots B_{n+1} w_{n+\frac{1}{2}}) = B_{a-\frac{1}{2}} B_{a-\frac{3}{2}} \cdots B_{n+1} \mathcal{T}(w_{n+\frac{1}{2}}) \\ &= (-1)^{p(w_a)} w_{-a} + (Q - Q^{-1}) w_a = w_a \cdot H_0. \end{aligned}$$

In case $m = 0$ or 1 , the non-super specialization of Proposition 7.6 is established in [BW18a, BWW18]. The property of a K -matrix \mathcal{T} in Corollary 7.5 also provides a conceptual explanation for the commutativity of H_0 and \mathbf{U}^ι acting on $\mathbb{W}^{\otimes d}$. □

REFERENCES

- [AV20] A. Appel and B. Vlaar, *Universal K -matrix for quantum Kac-Moody algebras*, arXiv:2007.09218
- [Bao17] H. Bao, *Kazhdan-Lusztig theory of super type D and quantum symmetric pairs*, Represent. Theory **21** (2017), 247–276.
- [BK15] M. Balagovic and S. Kolb, *The bar involution for quantum symmetric pairs*, Represent. Theory **19** (2015), 186–210.
- [BK19] M. Balagovic and S. Kolb, *Universal K -matrix for quantum symmetric pairs*, J. Reine Angew. Math. **747** (2019), 299–353.

- [BW18a] H. Bao and W. Wang, *A new approach to Kazhdan-Lusztig theory of type B via quantum symmetric pairs*, *Astérisque* **402** (2018), vii+134 pp, arXiv:1310.0103
- [BW18b] H. Bao and W. Wang, *Canonical bases arising from quantum symmetric pairs*, *Invent. Math.* **213** (2018), 1099–1177.
- [BW21] H. Bao and W. Wang, *Canonical bases arising from quantum symmetric pairs of Kac-Moody type*, *Compositio Math.* **157** (2021), 1507–1537.
- [BWW18] H. Bao, W. Wang and H. Watanabe, *Multiparameter quantum Schur duality of type B*, *Proc. Amer. Math. Soc.* **146** (2018), 3203–3216.
- [C16] S. Clark, *Canonical bases for the quantum enveloping algebra of $\mathfrak{gl}(m|1)$ and its modules*, arXiv:1605.04266
- [Ch19] C. Chung, *A Serre presentation for the \imath quantum covering groups*, arXiv:1912.09281
- [Ch21] C. Chung, *Canonical bases arising from \imath quantum covering groups*, arXiv:2107.06322
- [CS22] W. Cui and Y. Shen, *Canonical basis of q -Brauer algebras and \imath Schur dualities*, arXiv:2203.02082
- [CHW13] S. Clark, D. Hill and W. Wang, *Quantum Supergroups I. Foundations*, *Transform. Groups* **18** (2013), 1019–1053.
- [CHW16] S. Clark, D. Hill and W. Wang, *Quantum shuffles and quantum supergroups of basic type.*, *Quantum Topol.* **7** (2016), 553–638.
- [CL22] J. Chen and L. Luo, *Multiplication formulas and isomorphism theorem of \imath Schur superalgebras* arXiv:2202.02564
- [CW12] S-J. Cheng, W. Wang, *Dualities and representations of Lie superalgebras*, *Graduate Studies in Mathematics*, **144**. Amer. Math. Soc., Providence, RI, 2012.
- [De87] V. Deodhar, *On Some Geometric Aspects of Bruhat Orderings II. The Parabolic Analogue of Kazhdan-Lusztig Polynomials*, *J. Algebra* **111** (1987), 483–506.
- [Jim86] M. Jimbo, *A q -analogue of $U(\mathfrak{gl}(N + 1))$, Hecke algebra, and the Yang-Baxter equation*, *Lett. Math. Phys.* **11** (1986), 247–252.
- [Jan95] J. Jantzen, *Lectures on quantum groups*, *Graduate Studies in Math.* **6** (1995).
- [Ko12] S. Kolb, *Quantum symmetric Kac-Moody pairs*, *Adv. Math.* **267** (2014), 395–469.
- [Ko21] S. Kolb, *The bar involution for quantum symmetric pairs—hidden in plain sight*, arXiv:2104.06120
- [KT91] S. Khoroshkin and V. Tolstoy, *Universal R -matrix for quantized (super)algebras*, *Commun. Math. Phys.* **141** (1991), 599–617.
- [Let99] G. Letzter, *Symmetric pairs for quantized enveloping algebras*, *J. Algebra* **220** (1999), 729–767.
- [Let02] G. Letzter, *Coideal subalgebras and quantum symmetric pairs*, *New directions in Hopf algebras* (Cambridge), MSRI publications, **43**, Cambridge Univ. Press, (2002), 117–166.
- [Lus93] G. Lusztig, *Introduction to quantum groups*, *Modern Birkhäuser Classics*, Reprint of the 1993 Edition, Birkhäuser, Boston, 2010.
- [Lus03] G. Lusztig, *Hecke algebras with unequal parameters*, *CRM Monograph Series* **18**, Amer. Math. Soc., Providence, RI, 2003; for an updated and enlarged version see arXiv:0208154v2
- [LW20] L. Luo and W. Wang, *Lectures on dualities ABC in representation theory*, "Forty Years of Algebraic Groups, Algebraic Geometry, and Representation Theory in China", (eds. J. Du, J. Wang, L. Lin), World scientific, 2022. arXiv:2012.07203
- [Mi06] H. Mitsuhashi, *Schur-Weyl reciprocity between the quantum superalgebra and the Iwahori-Hecke algebra*, *Algebr. Represent. Theory* **9** (2006), 309–322.
- [S85] A. Sergeev, *The tensor algebra of the identity representation as a module over the Lie superalgebras $Gl(n, m)$ and $Q(n)$* , *Math. USSR Sbornik.* **51** (1985), 419–427.
- [SW21] Y. Shen and W. Wang, *\imath Schur duality and Kazhdan-Lusztig basis expanded*, arXiv:2108.00630
- [We12] H. Wenzl, *A q -Brauer algebra*, *J. Algebra* **358** (2012), 102–127.
- [WZ22] W. Wang and W. Zhang, *An intrinsic approach to relative braid group symmetries on \imath quantum groups*, arXiv:2201.01803
- [Ya94] H. Yamane, *Quantized Enveloping algebras associated with simple Lie superalgebras and their universal R -matrices*, *Publ. RIMS.* **30** (1994), 15–87.