

BOUNDARY CROSSING PROBLEMS AND FUNCTIONAL TRANSFORMATIONS FOR ORNSTEIN-UHLENBECK PROCESSES

Aria Ahari^{1*}, Larbi Alili^{1†}, Massimiliano Tamborrino^{1‡}

¹ Department of Statistics, University of Warwick, Coventry, CV4 7AL, United Kingdom.

Abstract

We are interested in the law of the first passage time of driftless Ornstein-Uhlenbeck processes to time varying thresholds. We show that this problem is connected to the law of the first passage time of the process to some two-parameter family of functional transformations which, for specific values of the parameters, appears in a realisation of a standard Ornstein-Uhlenbeck bridge. We provide three different proofs of this connection. The first proof is based on a similar result to the case of the Brownian motion, the second uses a generalisation of the so-called Gauss-Markov processes and the third relies on the Lie group symmetry method applied to the Fokker-Planck equation of the Ornstein-Uhlenbeck process. We investigate the properties of this transformation and study the algebraic and analytical properties of an involution operator which is used in constructing it. We also show that this transformation maps the space of solutions of Sturm-Liouville equations into the space of solutions of the associated nonlinear ordinary differential equations. Lastly, we discuss the interpretation of such transformations through the method of images and give new examples of curves with explicit first passage time densities.

Keywords: First passage times; Lie algebras; Sturm-Liouville equations; Ornstein-Uhlenbeck; Fokker Planck equation; Ornstein-Uhlenbeck bridge; Brownian motion.

2020 Mathematics Subject Classification: Primary 35K05, 60J50, 60J60.

1 Introduction

Let $U := (U_t)_{t \geq 0}$ be a one-dimensional driftless Ornstein-Uhlenbeck (OU for short) process defined on a filtered probability space $(\Omega, (\mathcal{F})_{t \geq 0}, \mathcal{F}, \mathbb{P})$ as the unique solution to the following stochastic differential equation (SDE for short)

$$dU_t = -kU_t dt + dB_t, \quad U_0 = x \in \mathbb{R}, \quad (1)$$

where $(B_t)_{t \geq 0}$ is a standard Brownian motion (BM) starting at 0 and $k \in \mathbb{R}$ is a constant. The OU process is a Gauss-Markov process with transition density function given by

$$p_t(x, y) := \frac{\partial}{\partial y} \mathbb{P}(U_t \leq y | U_0 = x) = \frac{e^{kt}}{\sqrt{r(t)}} \phi\left(\frac{ye^{kt} - x}{\sqrt{r(t)}}\right), \quad (2)$$

where $\phi(x) = e^{-\frac{x^2}{2}}/\sqrt{2\pi}$ is the probability density function of the standard normal distribution and $r(t) = (e^{2kt} - 1)/2k$ with its inverse given by $s(t) = \ln(2kt + 1)/2k$. It is well known by the

*aria.ahari@warwick.ac.uk

†l.alili@warwick.ac.uk

‡massimiliano.tamborrino@warwick.ac.uk

Dambis, Dubins-Schwartz theorem (see, e.g., Theorem V.1.6 in [25]), that the OU process can be written in terms of a time changed BM $(W_t)_{t \geq 0}$ as

$$U_t = e^{-kt}(x + W_{r(t)}), \quad t \geq 0. \quad (3)$$

We are interested in the distribution of the first passage time (FPT) or the random stopping time

$$\hat{T}^f = \inf\{t : U_t = f(t)\},$$

where $x = 0$, the time-varying threshold $f \in \mathcal{C}([0, \infty), \mathbb{R})$ satisfies $f(0) \neq 0$ and the usual convention $\inf\{\emptyset\} = \infty$ applies. $\mathcal{C}(I, K)$, for some subintervals I and $K \subseteq \mathbb{R}$, denotes the space of continuous functions from I into K . Such a boundary crossing problem for the case when the process is a BM, i.e. $k = 0$, has been of extensive study for over a century. We refer to [2] for some recent review on this topic. For the OU process with drift, well known explicit results only exist for constant [1] or hyperbolic type boundaries [9]. Such a hitting time problem is of great interest, as the OU process has been used in many applications to model objects such as interest rates in finance or the evolution of the neuronal membrane voltages in neuroscience, see e.g. [1] and citations therein. In this paper, we focus on the OU without drift, as the results for the OU process with drift can be directly obtained from our results after some transformations.

The goal of this paper is to derive, through different methods, an explicit analytical expression between the distribution of the FPT of the OU process to a continuously differentiable boundary f and some two-parameter family of curves, $\hat{S}^{\alpha, \beta} f$ (equation (13)), further developing the results for the BM obtained in [3]. Note that the continuously differentiable assumption of f is linked to Starsen's [29] result for the BM and is crucial in ensuring that the distribution of \hat{T}^f has a continuous density with respect to the Lebesgue measure on $(0, \infty)$. After introducing some notations and recalling some results for the FPT of the Brownian motion, in Section 2 we recall the OU bridge and its properties, which will be essential in understanding the properties of the two-parameter transformation $\hat{S}^{\alpha, \beta}$ studied in Section 3, together with the properties of other transformations of interest.

The main result of this paper, Theorem 4.1, is presented in Section 4, and it is proved in three alternative ways. In the first proof, we directly use the relationship between the FPT of an OU and that of a BM and use Theorem 1.2 in [3] to obtain Theorem 4.1. The second proof consists of using a generalisation of the Gauss-Markov processes which were introduced in Section 3.2 of [3] and finding an analogue version of that proof in our case. We develop a third different approach using the Lie algebra to find the symmetries of the OU Fokker-Planck equation

$$\frac{\partial h}{\partial t}(x, t) = \frac{1}{2} \frac{\partial^2 h}{\partial x^2}(x, t) + kx \frac{\partial h}{\partial x}(x, t) + kh(x, t), \quad (4)$$

and then use these symmetries (Lemma 4.7) to construct a certain function, $\hat{h}^{\alpha, \beta}(x, t)$ (equation (23)), derive our transformation $\hat{S}^{\alpha, \beta} f$ and relate the FPT distribution of \hat{T}^f to $\hat{T}^{\hat{S}^{\alpha, \beta} f}$. Within the same section, we also discuss the asymptotics of the distribution of $\hat{T}^{\hat{S}^{\alpha, \beta} f}$. We provide the analogue of the Kolmogorov–Erdős–Petrovski transience test [11] in the OU case and show its connection to the asymptotic behaviour of the FPT. Lastly, in Section 5, we use the method of images to obtain new classes of boundaries such that explicit FPT distributions can be obtained and use our $\hat{S}^{\alpha, \beta} f$ transformation (13) to produce new examples. A limitation of this method is that it only works for boundaries with certain properties given in Lemma 5.3.

As we were finalising the paper, we discovered that the $\hat{S}^{\alpha, \beta}$ transformation, a variant of the boundary crossing identity (15) in Theorem 4.1 and the Lie symmetries (24) had previously appeared in [20] (our (13) can be obtained by setting $A = k^2, B = 0$ in equation (39) therein), using the Lie approach. When comparing our results, we found misprints in one of their Lie symmetries and boundary crossing identity, as discussed in Section 4.3.

1.1 Notations and previous results

Here, we introduce the notations that will be used throughout the paper. Some of the discussed operators and notations can also be found in Alili and Patie [3], but we recall them here for convenience.

The nonlinear operator τ is defined on the space of functions whose reciprocals are square integrable in some (possibly infinite) interval of \mathbb{R}^+ by $\tau f(t) = \int_0^t [f(s)]^{-2} ds$ and we use it to define

$$\hat{A}(a, b) = \left\{ \pm f \in \mathcal{C}([0, a], \mathbb{R}^+) : \tau f(\operatorname{sgn}(k)s(\operatorname{sgn}(k)a)) = \frac{b}{1 - \mathbb{1}_{\{k < 0\}} 2kb} \right\},$$

where $a, b \in \mathbb{R}^+$ and $\operatorname{sgn}(k)$ is the sign of k . We also denote by \hat{A}_∞ the set of continuous functions which are of constant sign on some nonempty interval $[0, l]$, $l > 0$, yielding the following decomposition: $\hat{A}_\infty = \bigcup_{a>0} \bigcup_{b>0} \hat{A}(a, b)$. Now, for $\alpha \in \mathbb{R}^*, \beta \in \mathbb{R}$, where $\mathbb{R}^* = \mathbb{R} \setminus \{0\}$, we introduce

the following family of nonlinear operators acting on \hat{A}_∞

$$\Pi^{\alpha, \beta} f(t) = f(t)(\alpha + \beta \tau f(t)). \quad (5)$$

As explained in the beginning of Section 2 of [3] and [25](Appendix 8), these operators are closely related to the Sturm-Liouville equation

$$\phi'' = \mu \phi, \quad (6)$$

where μ denotes a positive Radon measure on \mathbb{R}^+ and ϕ'' is the second derivative in the sense of distributions. In fact, if ϕ solves (6), then the vectorial space $\{\Pi^{\alpha, \beta} \phi; \alpha, \beta \in \mathbb{R}\}$ is the set of solutions to the same equation. Moreover, all positive solutions are convex and described by the set $\{\Pi^{\alpha, \beta} \varphi; \alpha > 0, \beta \geq 0\}$, where φ is the unique, positive, decreasing solution such that $\varphi(0) = 1$. Next, we let ρ to be the inversion operator acting on the space of continuous monotone functions i.e., $\rho f \circ f(t) = t$, where \circ denotes the composition of functions. In the paper by Alili and Patie [3], the problem of the first hitting time of a BM to a certain transformation of the boundary f is discussed. The latter transformation with its decomposition is the following

$$S^{\alpha, \beta} f(t) = \Sigma \circ \Pi^{\alpha, -\beta} \circ \Sigma f = \left(\frac{1 + \alpha \beta t}{\alpha} \right) f \left(\frac{\alpha^2 t}{1 + \alpha \beta t} \right), \quad (7)$$

where $f \in \mathcal{C}([0, \infty), \mathbb{R})$ satisfies $f(0) \neq 0$ and $\alpha \in \mathbb{R}^*, \beta \in \mathbb{R}$. The involution $\Sigma : A_\infty \rightarrow A_\infty$, i.e., $\Sigma \circ \Sigma = id$, is specified by

$$\Sigma f(t) = \frac{1}{f(\rho \circ \tau f(t))},$$

where

$$A_\infty = \bigcup_{a>0} \bigcup_{b>0} A(a, b) \text{ and } A(a, b) = \{ \pm f \in \mathcal{C}([0, a], \mathbb{R}^+) : \tau f(a) = b \}.$$

Let $T^f = \inf\{t : B_t = f(t)\}$ be the FPT of the standard BM $(B_t)_{t \geq 0}$. For $t < -1/\alpha\beta$ if $\alpha\beta < 0$ or $t < \infty$ otherwise, Theorem 1.2 in [3] connects the FPT distribution of B_t to $S^{\alpha, \beta} f$ to that to f as

$$\mathbb{P}(T^{S^{\alpha, \beta} f} \in dt) = \alpha^3 (1 + \alpha \beta t)^{-\frac{5}{2}} e^{-\frac{\alpha \beta}{2(1 + \alpha \beta t)} (S^{\alpha, \beta} f(t))^2} S^{\alpha, \beta}(\mathbb{P}(T^f \in dt)). \quad (8)$$

This relationship is crucial and will be used later in the direct proof of Theorem 4.1. We now introduce a new nonlinear operator $\hat{\Sigma}$ defined on \hat{A}_∞ by

$$\hat{\Sigma} = \Lambda^{-1} \circ \Sigma \circ \Lambda, \quad (9)$$

where $\Lambda f(t) = e^{ks(t)} f(s(t)) = \sqrt{2kt+1} f(s(t))$, where Λ is defined on $[0, \infty)$ for $k > 0$ and on $[0, -1/2k)$ for $k < 0$, with its inverse given by $\Lambda^{-1} f(t) = e^{-kt} f(r(t))$. Note that, $\hat{\Sigma}$ can also be represented in the following way

$$\hat{\Sigma} f(t) = \frac{e^{-k\rho\tau f(r(t))-kt}}{f(\rho\circ\tau f(r(t)))}. \quad (10)$$

We then define the family $(\hat{S}^{\alpha,\beta})_{\alpha\in\mathbb{R}^*,\beta\in\mathbb{R}}$ of operators, $\hat{S}^{\alpha,\beta} : \hat{A}_\infty \rightarrow \hat{A}_\infty$, as follows

$$\hat{S}^{\alpha,\beta} = \hat{\Sigma} \circ \Pi^{\alpha,-\beta} \circ \hat{\Sigma}. \quad (11)$$

Finally, for all $a > 0$, we set

$$a_{\alpha,\beta} = \begin{cases} \frac{a}{\alpha^2 - \alpha\beta a} & \text{if } \alpha^2 - \alpha\beta a > 0, k \geq 0; \\ \frac{a}{\alpha^2 - (\alpha\beta + 2k\alpha^2)a} & \text{if } 0 < \frac{2ka}{(\alpha\beta + 2k\alpha^2)a - \alpha^2} < 1, k < 0; \\ +\infty & \text{otherwise,} \end{cases}$$

and for $f \in A_\infty$, we write

$$a_{\alpha,\beta}^f = \begin{cases} a & \text{if } \alpha^2 - \alpha\beta a > 0, k \geq 0; \\ \frac{a}{1-2ka} & \text{if } 0 < \frac{2ka}{(\alpha\beta + 2k\alpha^2)a - \alpha^2} < 1, k < 0; \\ \rho \circ \tau f \left(s \left(\frac{\alpha}{\beta} \right) \right) & \text{otherwise.} \end{cases}$$

Notice that, as $a \rightarrow \infty$, we have $a_{\alpha,\beta} \rightarrow r(\zeta_{\alpha,\beta})$ where

$$\zeta_{\alpha,\beta} = \begin{cases} s \left(-\frac{1}{\alpha\beta} \right) & \text{if } \alpha\beta < 0, k \geq 0; \\ s \left(-\frac{1}{\alpha\beta + 2k\alpha^2} \right) & \text{if } 0 < \frac{2k}{\alpha\beta + 2k\alpha^2} < 1, k < 0; \\ +\infty & \text{otherwise.} \end{cases}$$

For convenience, all key transformations that are used throughout this paper are listed in Table 1 in Appendix A.

2 Ornstein-Uhlenbeck bridges

We start by defining the OU bridge and then recall its different representations, using the detailed analysis of both Wiener and OU bridges provided in [5].

Definition 2.1. *An OU bridge process $U^{br} = \{U_t^{br} : t \in [0, T]\}$ from a to b of length T is characterised by the following properties:*

- (i) $U_0^{br} = a$ and $U_T^{br} = b$ (each with probability 1).
- (ii) U^{br} is a Gaussian process.
- (iii) $\mathbb{E}[U_t^{br}] = a \frac{\sinh(k(T-t))}{\sinh(kt)} + b \frac{\sinh(kt)}{\sinh(kT)}$.
- (iv) $\text{Cov}(U_s^{br}, U_t^{br}) = \frac{\sinh(ks) \sinh(k(T-t))}{k \sinh(kT)}$, $0 \leq s \leq t < T$.
- (v) The process U_t^{br} is almost surely continuous on $[0, T]$.

In what follows, we present three different representations of OU bridge processes. First, consider the following linear SDE

$$dU_t^{\text{br}} = \left(-k \coth(k(T-t))U_t^{\text{br}} + k \frac{b}{\sinh(k(T-t))} \right) dt + dB_t, \quad 0 \leq t < T, \quad (12)$$

with initial condition $U_0^{\text{br}} = a$. This SDE has unique strong solution given by

$$U_t^{\text{ir}} := U_t^{\text{br}} = \begin{cases} a \frac{\sinh(k(T-t))}{\sinh(kT)} + b \frac{\sinh(kt)}{\sinh(kT)} + \int_0^t \frac{\sinh(k(T-s))}{\sinh(k(T-s))} dB_s & \text{if } 0 \leq t < T, \\ b & \text{if } t = T. \end{cases}$$

This is referred to as the integral representation (ir) of the OU bridge. The anticipative (av) and the space-time (st) transform versions can also be found by transforming the different representations of the Wiener bridge

$$U_t^{\text{av}} = a \frac{\sinh(k(T-t))}{\sinh(kT)} + b \frac{\sinh(kt)}{\sinh(kT)} + \left(U_t - \frac{\sinh(kt)}{\sinh(kT)} U_T \right), \quad 0 \leq t < T,$$

$$U_t^{\text{st}} = a \frac{\sinh(k(T-t))}{\sinh(kT)} + b \frac{\sinh(kt)}{\sinh(kT)} + e^{-kt} \left(\frac{r(T) - r(t)}{r(T)} \right) W_{\left(\frac{r(T)r(t)}{r(T)-r(t)} \right)}, \quad 0 \leq t < T.$$

The space-time transformation is of our main interest and we will later show its connection to our transformation $\hat{S}^{\alpha,\beta}$. One thing to notice is that the space-time and anticipative versions are only weak solutions to the SDE (12). If this was not the case, then by the definition of strong solutions we would get $\mathbb{P}(U_t^{\text{av}} = U_t^{\text{ir}} \quad \forall t \in [0, T]) = 1$, which is not true. This tells us that the sample path behaviours differ for each version. Section 3.2 of [5] provides a detailed analysis of such behaviours. Of course, U^{br} can be thought of as $(U_t, t \leq T)$ conditioned on $U_T = b$. For conditioned processes and Markovian bridges, see [12], [25].

3 Properties of transformations

We now state a result that gives an insight into our $\hat{S}^{\alpha,\beta}$ transformation.

Theorem 3.1. (1) Let (α, β) and $(\alpha', \beta') \in \mathbb{R}^* \times \mathbb{R}$. The mapping $\hat{S}^{\alpha,\beta} : \hat{A}_\infty \rightarrow \hat{A}_\infty$ defined by (11) admits the following representation

$$\hat{S}^{\alpha,\beta} f(t) = \left(\frac{1 + \alpha\beta r(t)}{\alpha} \right) \sqrt{2k \frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)} + 1} e^{-kt} f \left(s \left(\frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)} \right) \right), \quad (13)$$

with the property $\hat{S}^{\alpha,\beta} \circ \hat{S}^{\alpha',\beta'} = \hat{S}^{\alpha\alpha', \alpha\beta' + \frac{\beta}{\alpha}}$. In particular, $\hat{S}^{1,\beta}$ is a semigroup. Furthermore, if $f \in \hat{A}(a, b)$, then $\hat{S}^{\alpha,\beta} f \in \hat{A}(a_{\alpha,\beta}, b_{\alpha,\beta}^f)$.

(2) Let μ be a positive Radon measure on \mathbb{R}^+ . Then, there exists a unique positive and differentiable function \mathfrak{f} with $\mathfrak{f}(0) = 1$ and $\mathfrak{f}' + k\mathfrak{f} + \frac{k}{(2k\tau\mathfrak{f}+1)\mathfrak{f}} > 0$, which satisfies the following nonlinear differential equation

$$f^3 f'' - k^2 f^4 = \frac{-\mu(s(\tau f)) + k^2}{(2k(\tau f) + 1)^2}. \quad (14)$$

Moreover, $\{\hat{S}^{\alpha,\beta}\mathfrak{f}; \alpha > 0, \beta \in \mathbb{R}\}$ spans the set of positive solutions of (14).

Before proving Theorem 3.1, we start by discussing some of the properties of the operators $\hat{\Sigma}$ and $\Pi^{\alpha,\beta}$.

Proposition 3.2. For any $a, b > 0, \lambda \in \mathbb{R}$ and $f \in \hat{A}_\infty$, we have the following assertions.

(1) $\hat{\Sigma}$ is an involution operator, that is $\hat{\Sigma} \circ \hat{\Sigma} = I_d$ on \hat{A}_∞ .

(2) $\tau f(t) = r(\rho \circ \tau \circ \hat{\Sigma} f(r(t)))$.

(3) $\hat{\Sigma}(\hat{A}(a, b)) = \hat{A}(b, a)$.

(4) $\Pi^{\alpha, \beta} \Lambda = \Lambda \Pi^{\alpha, \beta}$.

Proof. (1) Use the decomposition (9) of $\hat{\Sigma}$ and the fact that Σ is an involution.

(2) By using the functional representation of $\hat{\Sigma}$ (10),

$$\tau \circ \hat{\Sigma} f(t) = \int_0^t f^2(\rho \circ \tau f(r(y))) e^{2ky + 2k\rho \circ \tau f(r(y))} dy = \int_0^{\rho \circ \tau f(r(t))} e^{2kz} dz = r(\rho \circ \tau f(r(t))),$$

which follows by a change of variable. Now use the fact that $\hat{\Sigma}$ is an involution.

(3) First, consider $k < 0$ and take $f \in \hat{A}(a, b)$. Observe that $\rho \circ \tau f(b/1 - 2kb) = -s(-a)$. Now, by part (2), $\tau \circ \hat{\Sigma} f(-s(-b)) = r(\rho \circ \tau f(b/1 - 2kb)) = a/1 - 2ka$ and so $\hat{\Sigma} f \in \hat{A}(b, a)$. Similarly, the result holds for $k > 0$.

(4) $\tau \Lambda f(t) = \int_0^t [e^{2ks(y)} f^2(s(y))]^{-1} dy = \tau f(s(t))$, which follows by a change of variable. The identity can easily be derived now. \square

proof of Theorem 3.1. (1) By using identity (4) from Proposition 3.2 and the decomposition of $S^{\alpha, \beta}$ given in (7) respectively, we can write

$$\hat{S}^{\alpha, \beta} = \hat{\Sigma} \circ \Pi^{\alpha, -\beta} \circ \hat{\Sigma} = \Lambda^{-1} \circ (\Sigma \circ \Pi^{\alpha, -\beta} \circ \Sigma) \circ \Lambda = \Lambda^{-1} \circ S^{\alpha, \beta} \circ \Lambda.$$

Now, if we use the functional definitions of the operators, we get the desired result, as illustrated in Figure 1 in Appendix A. By construction, the $S^{\alpha, \beta}$ transformation plays a key role in obtaining our $\hat{S}^{\alpha, \beta}$ transformation. This procedure can be interpreted as we first take the boundary f to its corresponding first hitting time boundary for the standard BM using the Λ transformation, then use $S^{\alpha, \beta}$ on this curve and then revert it back to the OU problem.

Next, the property of \hat{S} stated in the theorem follows by the similar semigroup property of the operator S (Proposition 2.3 of [3]). Lastly, using the identity $\tau \hat{S}^{\alpha, \beta} f(t) = \tau f\left(s\left(\frac{\alpha^2 r(t)}{1 + \alpha \beta r(t)}\right)\right)$ which follows readily by a change of variables, we get that if $f \in \hat{A}(a, b)$, then $\hat{S}^{\alpha, \beta} f \in \hat{A}(a_{\alpha, \beta}, b_{\alpha, \beta}^f)$.

(2) As discussed in Section 1.1, if ϕ solves the Sturm-Liouville equation (6), then the vectorial space $\{\Pi^{\alpha, \beta} \phi; \alpha, \beta \in \mathbb{R}\}$ is the set of solutions to the same equation. Moreover, all positive solutions are convex and described by the set $\{\Pi^{\alpha, \beta} \varphi; \alpha > 0, \beta \geq 0\}$, where φ is the unique, positive, decreasing solution such that $\varphi(0) = 1$. Now, we need to show that the image of (14) by $\hat{\Sigma}$ is (6) and vice-versa. We start by investigating the operator $\hat{\Sigma}$. Let $\phi(t) = \hat{\Sigma} f(t)$. Then, we obtain $\phi(s(\tau f(t))) f(t) = e^{-ks(\tau f(t)) - kt}$. Differentiating both side once, we get

$$\phi'(s(\tau f(t))) = -e^{-ks(\tau f(t)) - kt} (2k\tau f(t) + 1) \left(kf(t) + f'(t) + \frac{k}{(2k\tau f(t) + 1)f(t)} \right).$$

Differentiating again, we obtain

$$\phi''(s(\tau f(t))) = -(2k\tau f(t) + 1)^2 e^{-ks(\tau f(t)) - kt} \left(f^2(t) f''(t) - k^2 f^3(t) - \frac{k^2}{(2k\tau f(t) + 1)^2 f(t)} \right).$$

So if ϕ satisfies (6), we get (14) and vice versa. Now, by letting $f = \hat{\Sigma}\varphi$, we see that it satisfies the required properties. \square

Remark 3.3. Notice that the law of the process $(\hat{S}^{1,-\beta}U_t, 0 \leq t < \zeta_{1,-\beta})$ for a given β corresponds to the law of a bridge associated to the OU. Indeed, set $\beta = -1/r(T)$ for some $T > 0$. Then, for any $0 \leq t < T$, by using (13), we see that $(\hat{S}^{1,-1/r(T)}U_t, t \leq T)$ is a representation of the OU bridge of length T from 0 to 0.

4 Main result and proofs

We are now ready to state the main result of this paper which relates the distributions of the family of stopping times $(\hat{T}^{\hat{S}^{\alpha,\beta}f})_{\alpha \in \mathbb{R}^*, \beta \in \mathbb{R}}$ to that of \hat{T}^f . Recall that $\mathcal{C}^1(I, K)$, for some subintervals I and $K \subseteq \mathbb{R}$, denotes the space of continuously differentiable functions from I into K .

Theorem 4.1. *Let $f \in C^1([0, \infty), \mathbb{R})$ be such that $f(0) \neq 0$. Let $\alpha, \beta \in \mathbb{R}, \alpha \neq 0$. Then, for $t < \zeta_{\alpha,\beta}$, we have the following relationship*

$$\mathbb{P}\left(\hat{T}^{\hat{S}^{\alpha,\beta}f} \in dt\right) = e^{3kt} \alpha^3 (1 + \alpha\beta r(t))^{-5/2} \left(2k \frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)} + 1\right)^{-3/2} e^{-\frac{\alpha\beta}{2(1+\alpha\beta r(t))} (\hat{S}^{\alpha,\beta}f)^2 e^{2kt}} \hat{S}^{\alpha,\beta} \left(\mathbb{P}(\hat{T}^f \in dt)\right). \quad (15)$$

We provide three different proofs of Theorem 4.1. The first method is a direct approach that uses results from [3]. The second approach is more interesting, as it uses a generalisation of the so-called Gauss-Markov processes and gives an insight into some of the results given in Theorem 3.1. The third proof relies on the Lie group techniques applied to the Fokker-Planck equation of the OU. Before proving the theorem, we see an example of its applications.

Example 4.2. *Theorem 3.1 in [1] gives an expression for the FPT density of an OU process starting at 0 hitting a constant threshold $f(t) = a > 0$. Writing $\hat{p}^a(t)dt = \mathbb{P}(\hat{T}^a \in dt)$, the latter expression is*

$$\hat{p}^a(t) = -ke^{-ka^2/2} \sum_{j=1}^{\infty} \frac{D_{\nu_{j,-a\sqrt{t}}}(0)}{D'_{\nu_{j,-a\sqrt{t}}}(-a\sqrt{2k})} \exp(-k\nu_{j,-a\sqrt{2k}}t),$$

where $D_\nu(\cdot)$ is the parabolic cylinder function with index $\nu \in \mathbb{R}$ [16], $\nu_{j,b}$ the ordered sequence of positive zeros of $D_\nu(b)$, $D_{\nu_{j,-a\sqrt{t}}}(0) = 2^{\frac{\nu_{j,-a\sqrt{t}}}{2}} \frac{\Gamma(1/2)}{\Gamma((1-\nu_{j,-a\sqrt{t}})/2)}$ and $D'_{\nu_{j,b}}(b) = \frac{\partial D_\nu(b)}{\partial \nu} |_{\nu=\nu_{j,b}}$. Now, we use the $\hat{S}^{\alpha,\beta}$ transformation with $f(t) = a$, $\alpha, \beta \in \mathbb{R}, \alpha \neq 0$ to get the following family of curves

$$\hat{S}^{\alpha,\beta}f(t) = \frac{ae^{-kt}}{2k\alpha} \left\{ (\alpha\beta e^{2kt} - \alpha\beta + 2k) \left[(2k\alpha^2 + \alpha\beta)e^{2kt} - 2k\alpha^2 - \alpha\beta + 2k \right] \right\}^{1/2}.$$

By Theorem 4.1,

$$\begin{aligned} \hat{p}^{\hat{S}^{\alpha,\beta}f}(t) &= e^{2kt} \alpha^2 \frac{(1 + \alpha\beta r(t))^{-3/2}}{\left(2k \frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)} + 1\right)} e^{-\frac{\alpha\beta}{2(1+\alpha\beta r(t))} (\hat{S}^{\alpha,\beta}f(t))^2 e^{2kt}} \hat{p}^a \left(s \left(\frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)} \right) \right) \\ &= \frac{-ke^{2kt} \alpha^2 (1 + \alpha\beta r(t))^{-5/2}}{(2\alpha^2 r(t) + \alpha\beta r(t) + 1)} \exp \left(-\frac{\alpha\beta}{2(1 + \alpha\beta r(t))} (\hat{S}^{\alpha,\beta}f(t))^2 e^{2kt} - ka^2/2 \right) \\ &\quad \times \sum_{j=1}^{\infty} \frac{D_{\nu_{j,-a\sqrt{t}}}(0)}{D'_{\nu_{j,-a\sqrt{t}}}(-a\sqrt{2k})} \exp \left(-k\nu_{j,-a\sqrt{2k}} s \left(\frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)} \right) \right). \end{aligned}$$

Such curves for $\beta = 1$, $k = 1/2$ and different values of α are plotted in figure 2 in Appendix A. In the special case of $\alpha\beta = 2k$, we get the family of curves $\hat{S}^{\alpha,\beta} f(t) = \frac{\alpha}{\alpha} \sqrt{(\alpha^2 + 1)e^{2kt} - \alpha^2}$, with FPT density given by

$$\hat{p}^{\hat{S}^{\alpha,\beta} f}(t) = \frac{-k\alpha^2 e^{-ka^2 \left(\frac{\alpha^2+1}{\alpha^2}\right) e^{2kt} + \frac{ka^2}{2} + kt}}{(\alpha^2 + 1)e^{2kt} - \alpha^2} \sum_{j=1}^{\infty} \frac{D_{\nu_{j,-a\sqrt{t}}}(0)}{D'_{\nu_{j,-a\sqrt{t}}}(-a\sqrt{2k})} e^{-k\nu_{j,-a\sqrt{2k}} s\left(\frac{\alpha^2 r(t)}{1+\alpha\beta r(t)}\right)}.$$

Before providing the proofs of Theorem 4.1, we mention that, similarly as in the BM case, it is sufficient to only consider the case where $\alpha > 0$. This is because, by the symmetry of the BM and the OU process starting at 0 without drift, hitting negative or positive valued thresholds for the first time have the same probability. Hence for any $(\alpha, \beta) \in \mathbb{R}^* \times \mathbb{R}$, we have $\hat{T}^{\hat{S}^{\alpha,\beta} f} \stackrel{d}{=} \hat{T}^{\hat{S}^{|\alpha|, \text{sgn}(\alpha)\beta} f}$, where $\stackrel{d}{=}$ denotes equality in distribution.

4.1 First proof of Theorem 4.1 via a direct approach

Theorem 1.2 in [3] connects the FPT distribution of $S^{\alpha,\beta} f$ to that of f . We can connect the FPTs of OU and BM in the following manner,

$$\hat{T}^f = \inf\{t : U_t = f(t)\} = \inf\{s(t) : W_t = e^{ks(t)} f(s(t))\} = s(T^{\Lambda f}). \quad (16)$$

Now, we proceed with the first proof.

First proof of Theorem 4.1. Note that $\hat{T}^{\hat{S}^{\alpha,\beta} f} = s(T^{S^{\alpha,\beta} \Lambda f})$. Writing $\hat{p}^f(t)dt = \mathbb{P}(\hat{T}^f \in dt)$ and $p^f(t)dt = \mathbb{P}(T^f \in dt)$, we get

$$\hat{p}^{\hat{S}^{\alpha,\beta} f}(t) = e^{2kt} p^{S^{\alpha,\beta} \Lambda f}(r(t)). \quad (17)$$

Now, using (7) and (8), we can rewrite the right hand side as follows:

$$\begin{aligned} p^{S^{\alpha,\beta} \Lambda f}(r(t)) &= \alpha^3 (1 + \alpha\beta r(t))^{-5/2} e^{-\frac{\alpha\beta}{2(1+\alpha\beta r(t))} (S^{\alpha,\beta} \Lambda f(r(t)))^2} S^{\alpha,\beta} \left(p^{\Lambda f}(r(t)) \right) \\ &= \alpha^2 (1 + \alpha\beta r(t))^{-3/2} e^{-\frac{\alpha\beta}{2(1+\alpha\beta r(t))} (S^{\alpha,\beta} \Lambda f(r(t)))^2} p^{\Lambda f} \left(\frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)} \right). \end{aligned} \quad (18)$$

By using (16), we get

$$p^{\Lambda f} \left(\frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)} \right) = \frac{1}{2k \frac{\alpha^2 r(t)}{1+\alpha\beta r(t)} + 1} \hat{p}^f \left(s \left(\frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)} \right) \right). \quad (19)$$

Notice that $S^{\alpha,\beta} \Lambda f(r(t)) = e^{kt} \hat{S}^{\alpha,\beta} f(t)$. Now, plugging (19) into (18) and then into (17), we get

$$\begin{aligned} \hat{p}^{\hat{S}^{\alpha,\beta} f}(t) &= e^{2kt} \alpha^2 \frac{(1 + \alpha\beta r(t))^{-3/2}}{\left(2k \frac{\alpha^2 r(t)}{1+\alpha\beta r(t)} + 1\right)} e^{-\frac{\alpha\beta}{2(1+\alpha\beta r(t))} (\hat{S}^{\alpha,\beta} f(t))^2} e^{2kt} \hat{p}^f \left(s \left(\frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)} \right) \right) \\ &= e^{3kt} \alpha^3 (1 + \alpha\beta r(t))^{-5/2} \left(2k \frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)} + 1\right)^{-3/2} e^{-\frac{\alpha\beta}{2(1+\alpha\beta r(t))} (\hat{S}^{\alpha,\beta} f(t))^2} e^{2kt} \hat{S}^{\alpha,\beta} \left(\hat{p}^f(t) \right), \end{aligned}$$

where we used (13) to obtain the last equation. \square

4.2 Second proof of Theorem 4.1 via generalised Gauss-Markov processes

We take $\phi \in \hat{A}(a, b) \cap AC([0, b])$, where $AC([0, b])$ is the space of absolutely continuous functions on $[0, b]$. We introduce the process $X := (X_t)_{0 \leq t < b}$ to be the generalised Gauss-Markov process of OU type with parameters (ϕ, k) , which is defined as the unique strong solution to the following SDE

$$dX_t = \left(\frac{\phi'(t)}{\phi(t)} + k \right) X_t dt + e^{kt} dB_t, \quad 0 \leq t < b,$$

with $X_0 = x \in \mathbb{R}$, i.e.,

$$X_t = \phi(t)e^{kt} \left(x + \int_0^t \frac{1}{\phi(s)} dB_s \right), \quad 0 \leq t < b.$$

We also denote by $\mathbb{P}^{(\phi, k)} = \left(\mathbb{P}_x^{(\phi, k)} \right)_{x \in \mathbb{R}}$ the family of probability measures corresponding to the process X . We assume throughout that $\phi(0) = 1$. Notice that when $\phi(t) = e^{-kt}$, we get $X_t = W_{r(t)}$, where $(W_t, t \geq 0)$ is another BM. The motivation for introducing the generalised Gauss-Markov process is the connection between its boundary crossing problem and that of the OU.

Lemma 4.3. *For any $y \in \mathbb{R}$, set $T_y = \inf\{0 < t < b : \phi(t)e^{kt} \int_0^t \frac{1}{\phi(s)} dB_s = y\}$. Then, for any $f \in \hat{A}(a, b)$, setting $\phi = \hat{\Sigma}f$, the identity*

$$\hat{T}f = s(\tau\phi(T_1))$$

holds almost surely.

Proof. Using (3), we have a.s.

$$T_1 = \inf\{t > 0 : \phi(t)e^{kt}W_{\tau\phi(t)} = 1\} = \inf\left\{t : U_{s(\tau\phi(t))} = \frac{e^{-kt - ks(\tau\phi(t))}}{\phi(t)}\right\} = \rho \circ \tau\phi(r(\hat{T}f))$$

and the result follows. \square

Next, we introduce the following notation

$$H_t^k(x) = \sqrt{\frac{\alpha\phi(t)}{\Pi^{\alpha, \beta}\phi(t)}} \exp\left(\frac{\beta}{2} \frac{x^2 e^{-2kt}}{\phi(t)\Pi^{\alpha, \beta}\phi(t)}\right).$$

Our aim now is to show that the parametric families of distributions $\left(\mathbb{P}^{(\Pi^{\alpha, \beta}\phi, k)} \right)_{(\alpha, \beta) \in \mathbb{R}^* \times \mathbb{R}}$ of generalised Gauss-Markov processes are related by some space-time harmonic transformations.

Proposition 4.4. *For $(\alpha, \beta) \in \mathbb{R}^* \times \mathbb{R}$ and ϕ as above, the process $(H_t^k(X_t))_{0 \leq t < a_{\alpha, -\beta}^\phi}$ is a $\mathbb{P}^{(\phi, k)}$ -martingale. Furthermore, the absolute-continuity relationship*

$$d\mathbb{P}_{x|\mathcal{F}_t}^{(\Pi^{\alpha, \beta}\phi, k)} = \frac{H_t^k(X_t)}{H_0^k(x)} d\mathbb{P}_{x|\mathcal{F}_t}^{(\phi, k)}$$

holds for all $x \in \mathbb{R}$ and $t < a_{\alpha, -\beta}^\phi$. Consequently, for any reals x and y , we have

$$\mathbb{P}_x^{(\Pi^{\alpha, \beta}\phi, k)}(T_y \in dt) = \frac{H_t^k(y)}{H_0^k(x)} \mathbb{P}_x^{(\phi, k)}(T_y \in dt), \quad t < a_{\alpha, -\beta}^\phi.$$

Proof. The proof is similar to that of Lemma 3.2 in [3]. As that proof has a few typos, we will give the full proof here. From the Itô formula, we have

$$\frac{\beta}{2} \frac{X_t^2 e^{-2kt}}{\phi(t)\Pi^{\alpha,\beta}\phi(t)} = \beta \int_0^t \frac{X_s e^{-ks}}{\phi(s)\Pi^{\alpha,\beta}\phi(s)} dB_s - \frac{\beta^2}{2} \int_0^t \left(\frac{X_s e^{-ks}}{\phi(s)\Pi^{\alpha,\beta}\phi(s)} \right)^2 ds + \frac{1}{2} \ln \left(\frac{\alpha + \beta\tau\phi(t)}{\alpha} \right).$$

Now, as $\mathbb{E}[e^{-\frac{\lambda}{2}B_t^2}] = (1 + \lambda t)^{-1/2}$, $\lambda > -1/t$, we deduce that, for all $t < a_{\alpha,-\beta}^\phi$, we have

$$\mathbb{E}[H_t^k(X_t)] = \mathbb{E} \left[\sqrt{\frac{\alpha}{\alpha + \beta\tau\phi}} e^{-\frac{\beta}{2} \frac{W_{\tau\phi}^2}{\alpha + \beta\tau\phi}} \right] = 1$$

Hence, it is a true martingale. Next, notice that

$$d \left\langle \beta \int_0^\cdot \frac{X_s e^{-ks}}{\phi(s)\Pi^{\alpha,\beta}\phi(s)} dB_s, X \right\rangle_t = \beta \frac{X_t}{\phi(t)\Pi^{\alpha,\beta}\phi(t)} dt,$$

where $\langle \cdot, \cdot \rangle$ is the quadratic variation process. Also,

$$\frac{(\Pi^{\alpha,\beta}\phi(t))'}{\Pi^{\alpha,\beta}\phi(t)} = \frac{\phi'(t)}{\phi(t)} + \frac{\beta}{\phi(t)\Pi^{\alpha,\beta}\phi(t)},$$

and hence the absolute continuity relationship follows by an application of Girsanov's theorem. Now, Doob's optional stopping theorem implies that

$$\begin{aligned} \mathbb{P}_x^{(\Pi^{\alpha,\beta}\phi,k)}(T_y \leq t) &= \mathbb{E}_x^{(\phi,k)} \left[\mathbb{1}_{\{T_y \leq t\}} \frac{H_t^k(X_t)}{H_0^k(x)} \right] = \mathbb{E}_x^{(\phi,k)} \left[\mathbb{1}_{\{T_y \leq t\}} \mathbb{E}_x^{(\phi,k)} \left[\frac{H_t^k(X_t)}{H_0^k(x)} \middle| \mathcal{F}_{t \wedge T_y} \right] \right] \\ &= \mathbb{E}_x^{(\phi,k)} \left[\mathbb{1}_{\{T_y \leq t\}} \frac{H_{T_y}^k(y)}{H_0^k(x)} \right], \end{aligned}$$

where $\mathbb{1}_A$ is the indicator function of the set A . The result follows by differentiation. \square

Now, we are ready to give the second proof of Theorem 4.1.

Second proof of Theorem 4.1. We assume that our boundary $f \in \mathcal{C}^1((0, \infty), \mathbb{R}^+)$. This assumption is crucial, as it implies that the curves Λf are also continuously differentiable and, according to Strassen [29], that $T^{\Lambda f}$ has a continuous density with respect to the Lebesgue measure on $(0, \infty)$. By the relation between the two stopping times (16), \hat{T}^f also has a continuous density. Let $\phi = \hat{\Sigma}f$, $\hat{f}^{\alpha,\beta} = \hat{S}^{\alpha,\beta}f$ and thus, by definition, $\hat{f}^{\alpha,\beta} = \hat{\Sigma} \circ \Pi^{\alpha,-\beta}\phi$. Since $\hat{\Sigma}$ is an involution, from Lemma 4.3 we get $\hat{T}^{\hat{f}^{\alpha,\beta}} = s(\tau \circ \Pi^{\alpha,-\beta}\phi(T_1))$ a.s Using identity (2) from Proposition 3.2, we get

$$r(\rho \circ \tau \circ \Pi^{\alpha,-\beta}\phi(r(t))) = r(\rho \circ \tau \circ \hat{\Sigma}\hat{f}^{\alpha,\beta}(r(t))) = \tau\hat{f}^{\alpha,\beta}(t).$$

Now, for $t < \zeta_{\alpha,\beta}$, writing $\hat{p}^{\hat{f}^{\alpha,\beta}}(t)dt = \mathbb{P}(\hat{T}^{\hat{f}^{\alpha,\beta}} \in dt)$ and $\tilde{p}^{(\phi,k)}(t)dt = \mathbb{P}_0^{(\phi,k)}(T_1 \in dt)$, we get

$$\begin{aligned} \hat{p}^{\hat{f}^{\alpha,\beta}}(t) &= \frac{1}{(2k(\tau\hat{f}^{\alpha,\beta}) + 1)(\hat{f}^{\alpha,\beta})^2} \tilde{p}^{(\Pi^{\alpha,-\beta}\phi,k)}(s(\tau\hat{f}^{\alpha,\beta}(t))) \\ &= \frac{1}{(2k(\tau\hat{f}^{\alpha,\beta}) + 1)(\hat{f}^{\alpha,\beta})^2} \tilde{p}^{(\Pi^{\alpha,-\beta}\phi,k)} \left(s \left(\tau f \left(s \left(\frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)} \right) \right) \right) \right), \end{aligned}$$

where we used the identity $\tau \hat{f}^{\alpha, \beta}(t) = \tau f \left(s \left(\frac{\alpha^2 r(t)}{1 + \alpha \beta r(t)} \right) \right)$ which follows readily by a change of variable. Now, using Proposition 4.4 with the identities

$$\begin{aligned} \left(\phi \left(s \circ \tau f \left(s \left(\frac{\alpha^2 r(t)}{1 + \alpha \beta r(t)} \right) \right) \right) \right)^{-1} &= \left(\phi \left(\rho \circ \tau \phi \left(r \circ s \left(\frac{\alpha^2 r(t)}{1 + \alpha \beta r(t)} \right) \right) \right) \right)^{-1} \\ &= e^{k\rho \circ \tau \hat{\Sigma} f \left(\frac{\alpha^2 r(t)}{1 + \alpha \beta r(t)} \right) + ks \left(\frac{\alpha^2 r(t)}{1 + \alpha \beta r(t)} \right)} f \left(s \left(\frac{\alpha^2 r(t)}{1 + \alpha \beta r(t)} \right) \right) \\ &= e^{ks(\tau \hat{f}^{\alpha, \beta}(t)) + ks \left(\frac{\alpha^2 r(t)}{1 + \alpha \beta r(t)} \right)} f \left(s \left(\frac{\alpha^2 r(t)}{1 + \alpha \beta r(t)} \right) \right), \end{aligned}$$

and

$$\begin{aligned} (\Pi^{\alpha, -\beta} \phi(s(\tau \hat{f}^{\alpha, \beta}(t))))^{-1} &= (\Pi^{\alpha, -\beta} \phi(\rho \circ \tau \circ \Pi^{\alpha, -\beta} \phi(r(t))))^{-1} = e^{k\rho \circ \tau \Pi^{\alpha, -\beta} \phi(r(t)) + kt} \hat{f}^{\alpha, \beta}(t) \\ &= e^{ks(\tau \hat{f}^{\alpha, \beta}(t)) + kt} \hat{f}^{\alpha, \beta}(t), \end{aligned}$$

yields, for any $t < \zeta_{\alpha, \beta}$,

$$\begin{aligned} \tilde{p}^{(\Pi^{\alpha, -\beta} \phi, k)} \left(s \left(\tau f \left(s \left(\frac{\alpha^2 r(t)}{1 + \alpha \beta r(t)} \right) \right) \right) \right) &= \\ \sqrt{1 + \alpha \beta r(t)} e^{-\frac{\alpha \beta}{2(1 + \alpha \beta r(t))} (\hat{f}^{\alpha, \beta}(t))^2} e^{2kt} \tilde{p}^{(\phi, k)} \left(s \left(\tau f \left(s \left(\frac{\alpha^2 r(t)}{1 + \alpha \beta r(t)} \right) \right) \right) \right). \end{aligned}$$

Hence,

$$\hat{p}^{\hat{f}^{\alpha, \beta}}(t) = \frac{e^{-\frac{\alpha \beta}{2(1 + \alpha \beta r(t))} (\hat{f}^{\alpha, \beta}(t))^2} e^{2kt} \sqrt{1 + \alpha \beta r(t)}}{(2k(\tau \hat{f}^{\alpha, \beta}) + 1)(\hat{f}^{\alpha, \beta})^2} \tilde{p}^{(\phi, k)} \left(s \left(\tau f \left(s \left(\frac{\alpha^2 r(t)}{1 + \alpha \beta r(t)} \right) \right) \right) \right).$$

Using Lemma 4.3 again, we finally get (15). □

4.3 Third proof of Theorem 4.1 via the Lie group symmetries

We now provide the last proof for Theorem 4.1 using the Lie symmetries approach [23]. In general, this technique can be used to find solutions to differential equations with new boundary conditions from known ones. For example, the Lie point symmetries of the heat equation

$$\frac{\partial h}{\partial t} = \frac{1}{2} \frac{\partial^2 h}{\partial x^2}$$

can be found in Section 3.3 of [3] and an excellent book on this technique is [23].

Before going through the proof, we first discuss the connections between the two boundary value problems (20) corresponding to the heat equation and the OU Fokker-Planck equation. As in the second proof via generalised Gauss-Markov processes, we let f to be a continuously differentiable boundary on \mathbb{R}^+ . Set

$$D^f = \{(x, t) \in \mathbb{R} \times \mathbb{R}^+ : x \leq f(t)\}.$$

We introduce the following boundary value problems $\mathcal{H}(f)$ and $\hat{\mathcal{H}}(f)$ for the BM and OU, respectively:

$$\mathcal{H}(f) = \begin{cases} \frac{\partial}{\partial t} h(x, t) = \frac{1}{2} \frac{\partial^2}{\partial x^2} h(x, t) & \text{on } D^f; \\ h(f(t), t) = 0 & \text{for all } t > 0; \\ h(., 0) = \delta_0(.) & \text{on } (-\infty, f(0)), \end{cases} \quad \hat{\mathcal{H}}(f) = \begin{cases} \frac{\partial}{\partial t} \hat{h}(x, t) = k \frac{\partial}{\partial x} (x \hat{h}(x, t)) + \frac{1}{2} \frac{\partial^2}{\partial x^2} \hat{h}(x, t) & \text{on } D^f; \\ \hat{h}(f(t), t) = 0 & \text{for all } t > 0; \\ \hat{h}(., 0) = \delta_0(.) & \text{on } (-\infty, f(0)). \end{cases} \quad (20)$$

Note that the first equations in $\mathcal{H}(f)$ and $\hat{\mathcal{H}}(f)$ are the heat equation and the OU Fokker-Planck equation, respectively. By Proposition 5.4.3.1 of [6], solutions to both problems admit the following probabilistic representations:

$$h(x, t)dx = \mathbb{P}(W_t \in dx, t < T^f) \quad \text{and} \quad \hat{h}(x, t)dx = \mathbb{P}(U_t \in dx, t < \hat{T}^f). \quad (21)$$

Using (3) and (16), we connect the two solutions directly in the following way:

$$\mathbb{P}(W_t < x, t < T^{\Lambda f}) = \mathbb{P}(e^{-ks(t)}W_t < e^{-ks(t)}x, s(t) < s(T^{\Lambda f})) = \mathbb{P}(U_{s(t)} < e^{-ks(t)}x, s(t) < \hat{T}^f)$$

and so

$$h(x, t) = e^{-ks(t)}\hat{h}(e^{-ks(t)}x, s(t)) \quad \text{and} \quad \hat{h}(x, t) = e^{kt}h(e^{kt}x, r(t)), \quad (22)$$

where h now denotes a solution to $\mathcal{H}(\Lambda f)$. This shows that the solutions to the Fokker-Planck equation of the OU on D^f are directly connected to the solutions to the heat equation on $D^{\Lambda f}$. As will become clear in the third proof of Theorem 4.1 at the end of this section, the aim is to find a solution to the Fokker-Planck equation of the OU such that it vanishes on our desired transformed boundary $\hat{S}^{\alpha, \beta}f$. To do this, we first present a proposition for the OU which resembles Proposition 3.5 in [3] for the Brownian motion.

Proposition 4.5. *Let \hat{h} be the solution to the boundary value problem $\hat{\mathcal{H}}(f)$ in (20). Then, for any $\alpha > 0$, $\beta \in \mathbb{R}$, the following mapping*

$$\hat{h}^{\alpha, \beta}(x, t) = \frac{\alpha e^{kt}}{\sqrt{1 + \alpha\beta r(t)}} e^{-\frac{\alpha\beta e^{2kt}x^2}{2(1 + \alpha\beta r(t))} - ks\left(\frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)}\right)} \hat{h}\left(\frac{\alpha e^{-ks\left(\frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)}\right) + kt}}{1 + \alpha\beta r(t)}x, s\left(\frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)}\right)\right) \quad (23)$$

is the solution to the boundary value problem $\hat{\mathcal{H}}(\hat{S}^{\alpha, \beta}f)$.

Proof. Firstly, if f is an infinitely continuously differentiable function, then so is its transformation $\hat{S}^{\alpha, \beta}f$. From Proposition 3.5 in [3], the following function

$$h^{\alpha, \beta}(x, t) = \frac{\alpha}{\sqrt{1 + \alpha\beta t}} e^{-\frac{\alpha\beta x^2}{2(1 + \alpha\beta t)}} h\left(\frac{\alpha x}{1 + \alpha\beta t}, \frac{\alpha^2 t}{1 + \alpha\beta t}\right),$$

is a solution to $\mathcal{H}(S^{\alpha, \beta}\Lambda f)$ because h solves $\mathcal{H}(\Lambda f)$. Now using relation (22), we define $\tilde{h}^{\alpha, \beta}$ as $\tilde{h}^{\alpha, \beta} = e^{kt}h^{\alpha, \beta}(e^{kt}x, r(t))$. Then, this expression is a solution to the boundary value problem $\hat{\mathcal{H}}(\hat{S}^{\alpha, \beta}f)$. Using relation (22) again, we can write

$$h\left(\frac{\alpha e^{kt}x}{1 + \alpha\beta r(t)}, \frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)}\right) = e^{-ks\left(\frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)}\right)} \hat{h}\left(\frac{\alpha e^{-ks\left(\frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)}\right) + kt}}{1 + \alpha\beta r(t)}x, s\left(\frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)}\right)\right),$$

and so we get that $\tilde{h}^{\alpha, \beta}$ and $\hat{h}^{\alpha, \beta}$ coincide, which implies that $\hat{h}^{\alpha, \beta}$ is indeed a solution to $\hat{\mathcal{H}}(\hat{S}^{\alpha, \beta}f)$. Moreover, we have that $\hat{h}^{\alpha, \beta}(x, t) = 0 \iff x = \hat{S}^{\alpha, \beta}f(t)$, because by assumption \hat{h} is a solution to $\hat{\mathcal{H}}(f)$, hence $\hat{h}(f(\cdot), \cdot) = 0$. Also, $\hat{h}^{\alpha, \beta}(x, 0) = e^{-\frac{\alpha\beta x^2}{2}} \hat{h}^{\alpha, 0}(x, 0)$. Now, let us investigate $\hat{h}^{\alpha, 0}$. Using relation (22) again,

$$\begin{aligned} \hat{h}^{\alpha, 0}(x, t) &= \alpha e^{kt - ks(\alpha^2 r(t))} \hat{h}(\alpha e^{-ks(\alpha^2 r(t)) + kt}x, s(\alpha^2 r(t))) \\ &= \alpha e^{kt - ks(e^{-2(-\ln(\alpha))}r(t))} \hat{h}(e^{-ks(e^{-2(-\ln(\alpha))}r(t)) + kt - (-\ln(\alpha))}x, s(e^{-2(-\ln(\alpha))}r(t))) \\ &= e^{kt}h(e^{-\epsilon}e^{kt}x, e^{-2\epsilon}r(t)) = e^{kt}h_{\epsilon}^{(4)}(e^{kt}x, r(t)), \end{aligned}$$

where $\epsilon = -\ln(\alpha)$. Now, the latter $h_{\epsilon}^{(4)}$ is the fourth symmetry of the heat equation listed in Section 3.3 of [3], meaning that it satisfies the heat equation. By using relation (22), we see that $e^{kt}h_{\epsilon}^{(4)}(e^{kt}x, r(t))$ indeed satisfies the Fokker-Planck equation of the OU and so $\hat{h}^{\alpha, 0}$ is a solution to $\hat{\mathcal{H}}(\hat{S}^{\alpha, 0}f)$ and thus in particular $\hat{h}^{\alpha, 0}(\cdot, 0) = \delta_0(\cdot)$ on $(-\infty, \frac{f(0)}{\alpha})$. Hence, we get

$$\hat{h}^{\alpha, \beta}(x, 0) = e^{-\frac{\alpha\beta x^2}{2}} \hat{h}^{\alpha, 0}(x, 0) = \delta_0(x),$$

which concludes the proof. \square

4.3.1 Lie symmetries of OU Fokker-Planck equation

The interesting result given in this section is constructing the $\hat{h}^{\alpha,\beta}$ function (23) directly from the symmetries of the OU Fokker Planck equation (4). Using similar techniques discussed in Section 2.4 in [23], after some lengthy and tedious calculations, one finds that the Lie algebra of infinitesimal symmetries of the OU Fokker-Planck equation is spanned by six vector fields, where x, t are the two independent variables and h is the dependent variable,

$$\begin{aligned}\mathbf{v}_1 &= e^{-kt} \frac{\partial}{\partial x}, & \mathbf{v}_2 &= \frac{\partial}{\partial t}, & \mathbf{v}_3 &= h \frac{\partial}{\partial h}, & \mathbf{v}_4 &= e^{kt} \frac{\partial}{\partial x} - 2kxhe^{kt} \frac{\partial}{\partial h} \\ \mathbf{v}_5 &= -kxe^{-2kt} \frac{\partial}{\partial x} + e^{-2kt} \frac{\partial}{\partial t} + khe^{-2kt} \frac{\partial}{\partial h}, & \mathbf{v}_6 &= kxe^{2kt} \frac{\partial}{\partial x} + e^{2kt} \frac{\partial}{\partial t} - 2k^2x^2he^{2kt} \frac{\partial}{\partial h}\end{aligned}$$

and by the infinite-dimensional subalgebra $\mathbf{v}_\alpha = u(x, t) \frac{\partial}{\partial h}$, where $u(x, t)$ is an arbitrary solution of the OU Fokker-Planck equation. Now, by exponentiating the basis (for more details on this procedure, see Section 2.1, page 89 of [23]), we can produce the one-parameter group of transformations leaving invariant $\mathcal{H}(f)$. Doing this procedure for each of the vectors fields, we obtain the one-parameter groups G_i generated by the \mathbf{v}_i . Since each group G_i is a symmetry group, if $h = \hat{h}(x, t)$ is a solution to (4), then the following are also solutions to (4) (examples of this procedure are given in Section 2.2 [23]):

$$\begin{aligned}\hat{h}_\epsilon^{(1)}(x, t) &= \hat{h}(x - \epsilon e^{-kt}, t) \\ \hat{h}_\epsilon^{(2)}(x, t) &= \hat{h}(x, t - \epsilon) \\ \hat{h}_\epsilon^{(3)}(x, t) &= e^\epsilon \hat{h}(x, t) \\ \hat{h}_\epsilon^{(4)}(x, t) &= e^{-2kx\epsilon e^{kt} + k\epsilon^2 e^{2kt}} \hat{h}(x - \epsilon e^{kt}, t) \\ \hat{h}_\epsilon^{(5)}(x, t) &= \frac{e^{kt}}{\sqrt{e^{2kt} - 2k\epsilon}} \hat{h}\left(\frac{xe^{kt}}{\sqrt{e^{2kt} - 2k\epsilon}}, \frac{\ln(e^{2kt} - 2k\epsilon)}{2k}\right) \\ \hat{h}_\epsilon^{(6)}(x, t) &= e^{\frac{-2k^2x^2\epsilon}{e^{-2kt} + 2k\epsilon}} \hat{h}\left(\frac{xe^{-kt}}{\sqrt{e^{-2kt} + 2k\epsilon}}, \frac{\ln(e^{-2kt} + 2k\epsilon)}{-2k}\right).\end{aligned}\tag{24}$$

In Proposition 4.5, we proved that the function $\hat{h}^{\alpha,\beta}$ is a solution to the boundary value problem $\hat{\mathcal{H}}(\hat{S}^{\alpha,\beta} f)$ by directly using the relation between the two boundary value problems in (20). We can also construct this function directly using the symmetries above.

Remark 4.6. Proposition 5 of [20] gives a general class of symmetries for SDEs with a drift term specified in Proposition 4 therein. The expression therein contains misprints in $P_{1,2}^\epsilon$ and $\mathcal{T}_{p_{1,2}}$. Using the notation in the original article, the correct expressions are

$$\begin{aligned}P_{1,2}^\epsilon &: \frac{u(x, t)}{\mathcal{U}(\mathcal{X}, \mathcal{T})} = \frac{\theta_{\mathcal{F}_2}(x)}{\theta_{\mathcal{F}_2}(\mathcal{X})} \frac{e^{\pm 2\sqrt{A}(\nu - \frac{1}{4} \pm \frac{1}{4})t \mp \frac{\sqrt{A}}{2}(x+2B/A)^2}}{e^{\pm 2\sqrt{A}(\nu - \frac{1}{4} \pm \frac{1}{4})\mathcal{T} \mp \frac{\sqrt{A}}{2}(\mathcal{X}+2B/A)^2}} \\ \mathcal{T}_{p_{1,2}} &= \mp \ln(e^{\mp 2\sqrt{A}t} + \epsilon) / (2\sqrt{A}).\end{aligned}$$

If we set $A = k^2, B = 0, \nu = 0$ and $\theta_{\mathcal{F}_2}(x) = e^{-kx^2/2}$, we recover the above symmetries (24) for the OU process.

Lemma 4.7.

$$\hat{h}^{\alpha,\beta} = \left(\hat{h}^{(2)}\left(\frac{-\ln(\alpha)}{k}\right) \circ \hat{h}^{(5)}\left(\frac{\alpha^2 - 1}{2k}\right) \right) \circ \left(\hat{h}^{(6)}\left(\frac{\beta}{2k(2k\alpha - \beta)}\right) \circ \hat{h}^{(2)}\left(\frac{1}{k} \ln\left(\frac{2k\alpha - \beta}{2k\alpha}\right)\right) \circ \hat{h}^{(5)}\left(\frac{\beta}{2k(2k\alpha - \beta)}\right) \right).$$

Although the calculation is tedious, it can easily be shown that the expression given in Lemma 4.7 holds. One thing to notice is that $h^{\alpha,\beta}(x,t)$ from Proposition 3.5 of [3] is a composition of three symmetries of the heat equation and so it satisfies the heat equation. Now, because the function $\hat{h}^{\alpha,\beta}$ is a composition of solutions to the OU Fokker-Planck equation, then is itself a solution to the OU Fokker-Planck equation. The Lemma 4.7 can then be used to shorten the first part of the proof given in Proposition 4.5. Now, we give the third proof of Theorem 4.1.

Third proof of Theorem 4.1. By proposition 4.5 and Lemma 4.7, we deduce that $\hat{h}^{\alpha,\beta}$ in (23) is a solution to the boundary value problem $\hat{\mathcal{H}}(\hat{S}^{\alpha,\beta}f)$. Now, for a general boundary f , $\mathbb{P}(\hat{T}^f \leq t) = 1 - \int_{-\infty}^{f(t)} \hat{h}(x,t)dx$. Setting $\hat{p}^f(t)dt = \mathbb{P}(\hat{T}^f \in dt)$, we get

$$\hat{p}^f(t) = -\frac{d}{dt} \left(\int_{-\infty}^{f(t)} \hat{h}(x,t)dx \right) = -\hat{h}(f(t),t)f'(t) - \int_{-\infty}^{f(t)} \frac{\partial}{\partial t} \hat{h}(x,t)dx = - \int_{-\infty}^{f(t)} \frac{\partial}{\partial t} \hat{h}(x,t)dx,$$

because \hat{h} vanishes on the boundary. Using the OU Fokker-Planck equation, we get

$$\hat{p}^f(t) = - \int_{-\infty}^{f(t)} -\frac{\partial}{\partial x} [-k(x)\hat{h}(x,t)] + \frac{1}{2} \frac{\partial^2}{\partial x^2} \hat{h}(x,t) dx = -\frac{1}{2} \frac{\partial}{\partial x} \hat{h}(x,t)|_{x=f(t)}. \quad (25)$$

Using the $\hat{h}^{\alpha,\beta}$ function with its corresponding boundary $\hat{S}^{\alpha,\beta}f$ gives us expression (15) from Theorem 4.1. □

Remark 4.8. When we take the limit as k goes to 0 of the symmetries in (24), we only recover the first three symmetries of the heat equation (Section 3.3 of [3]). The rest of the symmetries of the heat equation can be recovered by using a combination of the symmetries of the OU Fokker-Planck equation.

Remark 4.9. This Lie approach led to a variant of our boundary crossing identity (15) in [20] (their equation 40) up to some misprints therein. In particular, by setting $\beta' = \beta/2k\alpha$, $\alpha' = 1/\alpha^2 - \beta/2k\alpha - 1$, $A = k^2$, $B = 0$, $\nu = 0$, $x_0 = 0$ and $\theta_{\mathcal{F}_2}(x) = e^{-kx^2/2}$, we noted that the first term in (40) should be $\mathcal{T}_-^{-1}\mathcal{T}_+^{-1/2}\sqrt{\alpha' + \beta' + 1}$ instead of $\mathcal{T}_+^{-1/2}/\sqrt{\alpha' + \beta' + 1}$.

4.4 Asymptotic behaviour of the FPT densities

Here, we discuss the asymptotic behaviour of the FPT distribution of $\hat{T}^{\hat{S}^{\alpha,\beta}f}$. As before, we assume that our boundary $f \in \mathcal{C}^1((0,\infty),\mathbb{R}^+)$ and, as discussed before Section 4.1, we only consider $\alpha > 0$. We write $\mathbb{P}(\hat{T}^f \in dt) = \hat{p}^f(t)dt$ and from Theorem 4.1, for $\beta > 0$, observe that we get the following asymptotic identity

$$\hat{p}^{\hat{S}^{\alpha,\beta}f}(t) \sim e^{2kt} \alpha^2 \frac{(\alpha\beta r(t))^{-1/2}}{(2k\alpha^2 r(t) + \alpha\beta r(t))} e^{-\frac{\alpha\beta}{2(1+\alpha\beta r(t))}(\hat{S}^{\alpha,\beta}f(t))^2} e^{2kt} \hat{p}^f \left(s \left(\frac{\alpha}{\beta} \right) \right) \quad \text{as } t \rightarrow \infty, \quad (26)$$

where $h(t) \sim I(t)$ as $t \rightarrow \zeta$ means that $h(t)/I(t) \rightarrow 1$ as $t \rightarrow \zeta$, for some $\zeta \in [0,\infty]$. Now, for $\beta < 0$, we need a different approach. Set $r = \mathbb{P}(\hat{T}^f < \infty)$. We say that f is transient if $r < 1$. We now give a necessary and sufficient condition for when a given curve is transient.

Theorem 4.10. *Let $f \in \mathcal{C}^1((0,\infty),\mathbb{R}^+)$. If $f(t)e^{kt}/\sqrt{r(t)}$ is increasing for sufficiently large t , then f is transient if and only if*

$$\int_{s(1)}^{\infty} \frac{e^{3kt} f(t)}{r(t)^{3/2}} e^{-e^{2kt} f(t)^2 / 2r(t)} dt < \infty.$$

Proof. By relation (16), we deduce that

$$r = \mathbb{P}(\hat{T}^f < \infty) = \mathbb{P}(s(T^{\Lambda f}) < \infty) = \mathbb{P}(T^{\Lambda f} < \infty).$$

Now, by the classical Kolmogorov-Eröds-Petrovski theorem [11], if $t^{-1/2}\Lambda f(t)$ (i.e. $f(t)e^{kt}/\sqrt{r(t)}$) is increasing for sufficiently large t , then Λf is a transient curve for the Brownian motion if and only if

$$\int_1^\infty t^{-3/2}\Lambda f(t)e^{-(\Lambda f)^2/2t}dt < \infty,$$

and the result follows by a change of variable. \square

For $\beta < 0$, the lifetime of our boundaries is $\zeta_{\alpha,\beta} = s(-1/(\alpha\beta))$, so $s\left(\frac{\alpha^2 r(t)}{1+\alpha\beta r(t)}\right) \rightarrow \infty$ as $t \rightarrow \zeta_{\alpha,\beta}$. In this case, whenever f is transient and satisfies suitable conditions, we can derive the asymptotic expression of the FPT density of the OU by making use of the corresponding result for the BM by Anderson and Pitt [4], as proved in the following Proposition.

Proposition 4.11. *Assume that f is transient and satisfies the following conditions*

- (i) Λf is increasing, concave, twice differentiable on $(0, \infty)$ and of regular variation at ∞ with index $\alpha \in [1/2, 1)$,
- (ii) $\Lambda f(t)/\sqrt{t}$ is increasing at ∞ , and $\Lambda f(t)/t$ is convex and decreases to 0 for sufficiently large t ,
- (iii) There exist positive constants $c < 1$ and c' such that the inequalities $tf'(s(t)) \leq (c(2kt + 1) - kt)f(s(t))$ and $|t^2/(2kt + 1)^2(f''(s(t)) - k^2f(s(t)))| \leq c'f(s(t))$ are met for sufficiently large t .

Then, for any $\beta < 0$, we have

$$\hat{p}^{\hat{S}^{\alpha,\beta}f}(t) \sim \sqrt{\frac{|\alpha\beta|^3}{2\pi}} \left(\frac{2k}{|\alpha\beta|} + 1 \right) (1-r)\tilde{f}(t) \quad \text{as } t \rightarrow \zeta_{\alpha,\beta},$$

where

$$\tilde{f}(t) = \frac{\left(\frac{k\alpha^2 r(t)}{1+\alpha\beta r(t)} + 1 \right) f\left(s\left(\frac{\alpha^2 r(t)}{1+\alpha\beta r(t)}\right)\right) - \left(\frac{\alpha^2 r(t)}{1+\alpha\beta r(t)}\right) f'\left(s\left(\frac{\alpha^2 r(t)}{1+\alpha\beta r(t)}\right)\right)}{\sqrt{2k\frac{\alpha^2 r(t)}{1+\alpha\beta r(t)} + 1}}.$$

Proof. By (17), we have $\hat{p}^f(t) = e^{2kt}p^{\Lambda f}(r(t))$. By Anderson and Pitt [4] result for the BM, if conditions (i)-(iii) are satisfied, we can write

$$\hat{p}^f(t) \sim e^{2kt}(1-r)\frac{\Lambda f(r(t)) - r(t)(\Lambda f(r(t)))'}{\sqrt{2\pi r(t)^{3/2}}}e^{-(\Lambda f(r(t)))^2/2r(t)} \quad \text{as } t \rightarrow \infty.$$

The result follows by simplifying this, combining it with (15) and using the assumption that $\Lambda f(t)/t \downarrow 0$ as $t \rightarrow \infty$. \square

For $\beta > 0$, regardless of whether f is transient or not, if $\hat{S}^{\alpha,\beta}f(t)e^{kt}/\sqrt{r(t)}$ is increasing for sufficiently large t , $0 < \beta\sqrt{2k(\alpha/\beta) + 1}f(s(\alpha/\beta)) < \infty$ and $f(0) > 0$, we have

$$\begin{aligned} \int_{s(1)}^\infty \frac{e^{3kt}\hat{S}^{\alpha,\beta}f(t)}{r(t)^{3/2}}e^{-e^{2kt}(\hat{S}^{\alpha,\beta}f(t))^2/2r(t)}dt &= \int_1^\infty \frac{1}{\sqrt{z}}g(s(z))e^{-g(s(z))^2z/2}dz \\ &< \int_1^\infty g(s(z))e^{-g(s(z))^2z/2}dz < \infty, \end{aligned}$$

where we performed a change of variable and noted that $g(t) = e^{kt}\hat{S}^{\alpha,\beta}f(t)/r(t) \sim \beta\sqrt{2k(\alpha/\beta) + 1}f(s(\alpha/\beta))$ as $t \rightarrow \infty$. The finiteness of the integral follows by the fact that g

eventually stabilises and the assumptions ensure that g never blows up or vanishes, so its minimum and the maximum are always attained and by Theorem 4.10, $\hat{S}^{\alpha,\beta} f$ is always transient under these assumptions. However, we cannot use Proposition 4.11 directly, because $\Lambda \hat{S}^{\alpha,\beta} f(t)/t \sim \beta \sqrt{2k(\alpha/\beta) + 1} f(s(\alpha/\beta)) > 0$ as $t \rightarrow \infty$ and so it does not decrease to 0, violating assumption (ii). However, the general formula (26) still holds.

Remark 4.12. We noted a misprint in Section 4.3 of [2]. For the Brownian motion, when $\beta > 0$, the one parameter family of transformed functions $S^\beta f$ is always transient whenever $t^{-1/2} S^\beta f$ is increasing for sufficiently large t , $0 < \beta f(\alpha/\beta) < \infty$ and $f(0) > 0$.

5 Interpretation via the method of images

In his lecture notes [17], Lerche did a thorough investigation of the method of images for the standard BM. Here, we apply the method of images to the OU process and use it to produce new examples of curves with explicit FPT densities. As in the BM case, we would like to construct a function \hat{h} satisfying the OU Fokker-Planck equation. Then, by the uniqueness of such solutions, as seen in Section 4.3, \hat{h} would also satisfy (21) and be the solution to the boundary value problem $\hat{\mathcal{H}}(f)$. We proceed as follows. First, we assume to have a positive σ -finite measure F with $\int_0^\infty \phi(\sqrt{\epsilon\theta}) F(d\theta) < \infty$ for all $\epsilon > 0$. Then, for $a > 0$, we define the \hat{h} function by

$$\hat{h}(x, t) := p_t(0, x) - \frac{1}{a} \int_0^\infty p_t(\theta, x) F(d\theta) = \frac{e^{kt}}{\sqrt{r(t)}} \phi\left(\frac{xe^{kt}}{\sqrt{r(t)}}\right) - \frac{1}{a} \int_0^\infty \frac{e^{kt}}{\sqrt{r(t)}} \phi\left(\frac{xe^{kt} - \theta}{\sqrt{r(t)}}\right) F(d\theta).$$

We know that \hat{h} vanishes on the boundary f , and so by simplifying, we get that $\hat{h}(x, t) = 0$ is equivalent to

$$l\left(\frac{xe^{kt}}{r(t)}, \frac{1}{r(t)}\right) = a, \quad (27)$$

where $l(y, s) = \int_0^\infty e^{\theta y - \frac{1}{2}\theta^2 s} F(d\theta)$. The following theorem gives us a way of calculating the density of \hat{T}^f explicitly.

Theorem 5.1. *The FPT density of \hat{T}^f is given by*

$$\mathbb{P}(\hat{T}^f \in dt) = \frac{e^{2kt}}{2r(t)^{3/2}} \phi\left(\frac{f(t)e^{kt}}{\sqrt{r(t)}}\right) E(\theta | (f(t), t)) dt,$$

where

$$E(\theta | (f(t), t)) = \frac{\int_0^\infty \theta \phi\left(\frac{f(t)e^{kt} - \theta}{\sqrt{r(t)}}\right) F(d\theta)}{\int_0^\infty \phi\left(\frac{f(t)e^{kt} - \theta}{\sqrt{r(t)}}\right) F(d\theta)}.$$

Proof. We just use relation (25) to get the result. □

Remark 5.2. If the support of the measure F is on \mathbb{R} and $F(\{0\}) = 0$, then we define

$$\hat{h}(x, t) := p_t(0, x) - \frac{1}{a} \int_{-\infty}^\infty p_t(\theta, x) F(d\theta).$$

Then, there exist positive and negative valued functions f_+ and f_- , with $f_- < f_+$ and the properties $\hat{h}(f_+(t), t) = 0$ and $\hat{h}(f_-(t), t) = 0$ for all $t < t_a$ for a certain $t_a \leq \infty$. Then, the stopping time can also be defined as the first time the process leaves the region $(f_-(t), f_+(t))$, i.e.

$$\hat{T}^{f_\pm} = \inf\{0 < t < t_a \mid U_t \notin (f_-(t), f_+(t))\}. \quad (28)$$

The following lemma gives us a characterisation of the boundaries obtained through the method of images.

Lemma 5.3 (Characterisation of boundaries). *The boundaries f obtained through the method of images have the following properties:*

1. f is infinitely often continuously differentiable;
2. $\Lambda f(t)/t$ (i.e. $f(t)e^{kt}/r(t)$) is monotone decreasing;
3. $f''(t) - k^2 f(t) \leq 0$.

Proof. In Lemma 1.1 of [17], it is shown that the curves $\eta(t)$ that satisfy the equation $l(\frac{\eta(t)}{t}, \frac{1}{t}) = a$ are infinitely continuously differentiable and concave, with $\eta(t)/t$ monotone decreasing. As in our case we have $a = l\left(\frac{f(t)e^{kt}}{r(t)}, \frac{1}{r(t)}\right)$, by changing $t \rightarrow s(t)$, we get $a = l\left(\frac{\Lambda f(t)}{t}, \frac{1}{t}\right)$. So $\eta(t) := \Lambda f(t)$ must satisfy the three properties given in Lemma 1.1 of [17], so (1) and (2) follow directly. For (3),

$$f''(t) = k^2 e^{-kt} \eta(r(t)) + e^{3kt} \eta''(r(t)) = k^2 f(t) + e^{3kt} \eta''(r(t))$$

and the result follows as $\eta(t)$ is concave. \square

Remark 5.4. From Theorem 3.1, if $\mu(s(\tau f)) > k^2$, then we get that the solutions to nonlinear ordinary differential equations (14) satisfy $f'' - k^2 f \leq 0$.

Let us now consider some examples of the boundaries that arise for specific measures F via Theorem 5.1.

Example 5.5. Consider $F(d\theta) = \delta_{2z}$ for some $z > 0$. Using $\hat{h}(f(t), t) = 0$ and simplifying, we get

$$f(t) = \frac{\ln(a)}{2zk} \sinh(kt) + ze^{-kt}.$$

By Theorem 5.1,

$$\mathbb{P}(\hat{T}^f \in dt) = \frac{ze^{2kt}}{r(t)^{3/2}} \phi\left(\frac{\frac{\ln(a)}{2z} r(t) + z}{\sqrt{r(t)}}\right) dt.$$

Remark 5.6. In [9], the FPT of a mean-reverting OU with parameter μ hitting a hyperbolic type boundary of the form $\mu + Ae^{kt} + Be^{-kt}$ is studied. This reduces to the FPT of a standard OU hitting a curve of the form $Ae^{kt} + Be^{-kt}$ for arbitrary constants A and B which is the curve given in this example. Also, applying the \hat{S} transformation to these curves gives the same family of curves with different constant coefficients.

Example 5.7. Consider $F(d\theta) = \frac{d\theta}{\sqrt{2\pi}}$ on \mathbb{R} . Then,

$$f_{\pm}(t) = \pm e^{-kt} \sqrt{r(t) \ln\left(\frac{a^2 e^{2kt}}{r(t)}\right)}, \quad t \leq t_a = \begin{cases} -s(-a^2) & \text{if } a^2 < 1/(2k), \\ +\infty & \text{otherwise.} \end{cases}$$

This is a two sided boundary, with exit time $\hat{T}^{f_{\pm}}$ defined in (28) and FPT density given by

$$\mathbb{P}(\hat{T}^{f_{\pm}} \in dt) = \frac{e^{2kt}}{2r(t)} \phi\left(\sqrt{\ln\left(\frac{a^2 e^{2kt}}{r(t)}\right)}\right) \sqrt{\ln\left(\frac{a^2 e^{2kt}}{r(t)}\right)} dt.$$

Applying the \hat{S} transformation (13) to this curve, we get

$$\hat{S}^{\alpha, \beta} f_{\pm}(t) = \pm \sqrt{r(t)} \sqrt{1 + \alpha \beta r(t)} e^{-kt} \sqrt{\ln\left[\frac{a^2(2k\alpha^2 r(t) + \alpha \beta r(t) + 1)}{\alpha^2 r(t)}\right]}. \quad (29)$$

The FPT density is given by

$$\begin{aligned}\hat{p}^{\hat{S}^{\alpha,\beta} f_{\pm}}(t) &= e^{2kt} \alpha^2 \frac{(1 + \alpha\beta r(t))^{-3/2}}{(2k \frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)} + 1)} e^{-\frac{\alpha\beta}{2(1 + \alpha\beta r(t))} (\hat{S}^{\alpha,\beta} f_{\pm}(t))^2} e^{2kt} \hat{p}^{f_{\pm}} \left(s \left(\frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)} \right) \right) \\ &= \frac{e^{2kt}}{\sqrt{8\pi}} \sqrt{\frac{\ln \left[\frac{a^2(2k\alpha^2 r(t) + \alpha\beta r(t) + 1)}{\alpha^2 r(t)} \right]}{1 + \alpha\beta r(t)}} e^{-\frac{1}{2} \ln \left[\frac{a^2(2k\alpha^2 r(t) + \alpha\beta r(t) + 1)}{\alpha^2 r(t)} \right] (\alpha\beta r(t) + 1)}.\end{aligned}$$

Now, for example, by setting $a = 1, \alpha = 2, \beta = k$, we get

$$\hat{S}^{2,k} f_{\pm}(t) = \pm \sqrt{\frac{e^{2kt} - 1}{2k}} \sqrt{\ln \left[\frac{k}{4} (\coth(kt) + 9) \right]},$$

with FPT density given by

$$\hat{p}^{\hat{S}^{2,k} f_{\pm}}(t) = \frac{e^{kt}}{\sqrt{8\pi}} \sqrt{\ln \left[\frac{k}{4} (\coth(kt) + 9) \right]} e^{-\frac{e^{2kt}}{2} \ln \left[\frac{k}{4} (\coth(kt) + 9) \right]}.$$

In Appendix A, Figures 3 and 4 illustrate such curves for different values of k, α and β . When $\beta = 0$, $\hat{S}^{\alpha,0} f_{\pm}$ converges to a constant level as $t \rightarrow \infty$, see Figure 3. Moreover, whenever α and β have opposite signs, we get a closed shaped graph defined on $t \in [0, \zeta_{0.1, -0.5}]$, see Figure 4.

Remark 5.8. Back to our \hat{S} transformation, using (27) with the time-transformation $t \rightarrow s \left(\frac{\alpha^2 r(t)}{1 + \alpha\beta r(t)} \right)$, we get

$$a = \int_0^{\infty} e^{\frac{\theta}{\alpha} \left[\frac{\hat{f}^{\alpha,\beta}(t) e^{kt}}{r(t)} \right] - \frac{1}{2} \left[\frac{(\theta/\alpha)^2}{r(t)} \right]} F^{\alpha,\beta}(d\theta),$$

where $F^{\alpha,\beta}(d\theta) = e^{-\frac{\beta}{\alpha} \frac{\theta^2}{2}} F(d\theta)$ is the measure corresponding to the curve $\hat{S}^{\alpha,\beta} f(t)$.

Acknowledgements

We are grateful to Dmitry Muravey for a discussion on the Lie symmetries while finalising the paper. AA was funded by EPSRC grant [EP/V520226/1] as part of the Warwick CDT in Mathematics and Statistics. For the purpose of open access, the authors have applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

References

- [1] Alili, L. Patie, P. Pedersen, J. (2005) Representations of the first hitting time density of an Ornstein-Uhlenbeck process. *Stoch. Mod.*, 21 (4), 967-980.
- [2] Alili, L. Patie, P. (2010) Boundary crossing identities for diffusions having the time-inversion property. *J. Theoret. Probab.*, 23 (1), 65-84.
- [3] Alili, L. Patie, P. (2014) Boundary crossing identities for Brownian motion and some non linear ODE's. *Proc. Am. Math. Soc.*, 142 (11), 3811-3824.
- [4] Anderson, J.M. Pitt, L.D. (1997) Large time asymptotics for Brownian hitting densities of transient concave curves. *J. Theoret. Probab.*, 10 (4), 921-934.
- [5] Barczy, M., Kern, P. (2010) Sample path deviations of the Wiener and the Ornstein-Uhlenbeck process from its bridges. *Braz. J. Probab. Stat.*, 27 (4), 437-466.

- [6] Blanc, J. Yor, M. Chesney M. (2009) *Mathematical Methods for Financial Markets*. Springer London.
- [7] Billingsley, P. (1995) *Probability and measure*. Wiley, New York.
- [8] Breiman, L. (1967) First exit times from a square root boundary. *Berke. Symp. Math. Stat. Probab.*, 5.2B, 9-16.
- [9] Buonocore, A., Nobile, A. G., Ricciardi, L. M. (1987) A new integral equation for the evaluation of first-passage-time probability densities. *Adv. in Appl. Probab.*, 19 (4), 784–800.
- [10] Ditlevsen, S., Lansky, P. (2005) Estimation of the input parameters in the Ornstein-Uhlenbeck neuronal model. *Phys. Rev. E*, 71. 011907.
- [11] Erdős, P. (1942) On the law of the iterated logarithm. *Ann. of Math.*, 43 (3), 419–436.
- [12] Fitzsimmons, P.J., Pitman, J., Yor, M. (1993) Markovian bridges, construction, Palm interpretation, and splicing. *Seminar on Stochastic processes*, Birkhäuser, Boston, E. Çinlar and K.L. Chung and M.J. Sharpe, 101-134.
- [13] Groeneboom, P. (1989) Brownian motion with a parabolic drift and Airy functions. *Probab. Theory Relat. Fields.*, 81 (1), 79–109.
- [14] Jennen, C, Lerche, H.R. (1981) First exit densities of Brownian motion through one-sided moving boundaries, *Z. Wahrscheinlichkeitstheorie verw. Gebiete.*, 55, 133-148.
- [15] Karatzas, I, Shreve, S. E. (1988) *Brownian motion and stochastic calculus*. New York: Springer-Verlag.
- [16] Lebedev, N.N. (1972) *Special Functions and their Applications*. Dover Publications, New York.
- [17] Lerche, H.R. (1986) *Boundary Crossing of Brownian Motion: Its Relation to the Law of the Iterated Logarithm and to Sequential Analysis*. *Lec. Not. Stat.*, 40.
- [18] Lescot, P, Zambrini, J. (2008). Probabilistic deformation of contact geometry, diffusion processes and their quadratures, *Semi. Stoch. Anal, Rand. Field. App. V.*, 59, 203–226.
- [19] Levy, P. (1965) *Processus stochastiques et mouvement brownien*. Paris: Gauthier-Villars. Second edition.
- [20] Muravey, D. (2020) Lie Symmetries Methods in Boundary Crossing Problems for Diffusion Processes. *Acta Appl Math* 170, 347–372.
- [21] Mörters, P., Peres, Y. (2010) *Brownian motion*. *Camb. Ser. Stat. Probab. Math.*
- [22] Novikov, A.A. (1971) The stopping times of a Wiener process. *Theor. Probab. Appl.*, 16 (3), 449-456.
- [23] Olver, P. (1993) *Applications of Lie groups to differential equations*, *Grad. Text. Math.* Springer., 107.
- [24] Pötzelberger, K. Wang, L. (2001) Boundary crossing probability for Brownian motion. *J. Appl. Prob.*, 38:152–164.
- [25] Revuz, D, Yor, M. (1999) *Continuous martingales and Brownian motion*, *Fundamental Principles of Mathematical Sciences*, 293, Springer-Verlag, Berlin.
- [26] Robbins, H, Siegmund, D. (1970). Boundary crossing probabilities for the Wiener process and partials sums, *Ann. Math. Statist.* 41, 1410-1429.

- [27] Rosencrans, S. (1976). Perturbation algebra of an elliptic operator. *J. Math. Anal. App.* 56. 317-329.
- [28] Salminen, P. (1988) On the first hitting time and the last exit time for a Brownian motion to/from a moving boundary. *Adv. Appl. Probab.* 20 (2), 411–426.
- [29] Strassen, V. (1967) Almost sure behaviour of sums of independent random variables and martingales, *Proc. Fifth Berkeley Symp. Math. Statist. Probab.*, Univ. of Cal. Press, Vol. 111, Part I., 315-343.

A Figures and charts

Notations	Transformations
$r(t)$	$(e^{2kt} - 1)/2k$
$s(t)$	$\ln(2kt + 1)/2k$
Λf	$\sqrt{2kt + 1}f(s(t))$
$\Lambda^{-1}f$	$e^{-kt}f(r(t))$
τf	$\int_0^t [f(s)]^{-2} ds$
$\Pi^{\alpha,\beta} f$	$f(t)(\alpha + \beta\tau f(t))$
Σf	$\frac{1}{f(\rho\circ\tau f(t))}$
$\hat{\Sigma} f$	$\Lambda^{-1} \circ \Sigma \circ \Lambda f(t) = \frac{e^{-k\rho\circ\tau f(r(t)) - kt}}{f(\rho\circ\tau f(r(t)))}$
$S^{\alpha,\beta} f$	$\Sigma \circ \Pi^{\alpha,-\beta} \circ \Sigma f(t) = \left(\frac{1+\alpha\beta t}{\alpha}\right) f\left(\frac{\alpha^2 t}{1+\alpha\beta t}\right)$
$\hat{S}^{\alpha,\beta} f$	$\hat{\Sigma} \circ \Pi^{\alpha,-\beta} \circ \hat{\Sigma} f(t) = \left(\frac{1+\alpha\beta r(t)}{\alpha}\right) \sqrt{2k \frac{\alpha^2 r(t)}{1+\alpha\beta r(t)} + 1} e^{-kt} f\left(s\left(\frac{\alpha^2 r(t)}{1+\alpha\beta r(t)}\right)\right)$

Table 1: Key transformations introduced in the paper.

$$\begin{array}{ccc}
 \text{OU} & f(t) & \xrightarrow{\hat{S}^{\alpha,\beta}} \left(\frac{1+\alpha\beta r(t)}{\alpha}\right) \sqrt{2k \frac{\alpha^2 r(t)}{1+\alpha\beta r(t)} + 1} e^{-kt} f\left(s\left(\frac{\alpha^2 r(t)}{1+\alpha\beta r(t)}\right)\right) \\
 & \downarrow \Lambda & \uparrow \Lambda^{-1} \\
 \text{BM} & \sqrt{2kt + 1}f(s(t)) & \xrightarrow{S^{\alpha,\beta}} \left(\frac{1+\alpha\beta t}{\alpha}\right) \sqrt{2k \frac{\alpha^2 t}{1+\alpha\beta t} + 1} f\left(s\left(\frac{\alpha^2 t}{1+\alpha\beta t}\right)\right)
 \end{array}$$

Figure 1: Flow chart of \hat{S} .




Figure 2: $\hat{S}^{\alpha,1}a$ for fixed $\beta = 1$, $k = 1/2$, $a = 1$ and $\alpha = 1$ (red), 2 (green), 3 (orange), 5 (purple), 20 (black).



Figure 3: Transformed curves (29) for $\beta = 0$, $k = 1$, $a = 1$ and $\alpha = -1$ (orange), 0.5 (green), 0.1 (blue).




Figure 4: Transformed curves (29) for fixed $k = 1, a = 1$. Orange: $\beta = -0.5, \alpha = 0.1$. Green: $\beta = 0.1, \alpha = 0.1$. Blue: $\beta = 1, \alpha = 0.1$.