

L_p -Sampling recovery for non-compact subclasses of L_∞

Glenn Byrenheid[†], Serhii Stasyuk^{‡,a}, Tino Ullrich^{‡*}

[†]Friedrich-Schiller-University Jena, Institute of Mathematics

[‡]Chemnitz University of Technology, Faculty of Mathematics

^aInstitute of Mathematics of NAS of Ukraine

October 5, 2022

Abstract

In this paper we study the sampling recovery problem for certain relevant multivariate function classes which are not compactly embedded into L_∞ . Recent tools relating the sampling numbers to the Kolmogorov numbers in the uniform norm are therefore not applicable. In a sense, we continue the research on the small smoothness problem by considering “very” small smoothness in the context of Besov and Triebel-Lizorkin spaces with dominating mixed regularity. There is not much known on the recovery of such functions except of an old result by Oswald in the univariate situation. As a first step we prove the uniform boundedness of the ℓ_p -norm of the Faber-Schauder coefficients in a fixed level. Using this we are able to control the error made by a (Smolyak) truncated Faber-Schauder series in L_q with $q < \infty$. It turns out that the main rate of convergence is sharp. As a consequence we obtain results also for $S_{1,\infty}^1 F([0, 1]^d)$, a space which is “close” to the space $S_1^1 W([0, 1]^d)$ which is important in numerical analysis, especially numerical integration, but has rather bad Fourier analytical properties.

1 Introduction

In this paper we continue studying the approximation power of Smolyak sparse grid sampling recovery for multivariate classes with mixed smoothness $S_{p,q}^r B([0, 1]^d)$ and $S_{p,q}^r F([0, 1]^d)$ based on the Faber-Schauder representation [3, 9] of f

$$f = \sum_{j \in \mathbb{N}_{-1}^d} \sum_{k \in \mathbb{D}_j} d_{j,k}^2(f) v_{j,k}.$$

Here $v_{j,k}$ represents the tensorized dilated and translated hat function (L_∞ -normalized) and

$$d_{j,k}^2(f) = \prod_{i=1}^d \left\{ \begin{array}{ll} \Delta_{2^{-j_i-1}}^{2,i} & : j_i \geq 0 \\ \text{Id} & : j_i = -1 \end{array} \right\} f(k_1 2^{-(j_1)_+}, \dots, k_d 2^{-(j_d)_+})$$

*Corresponding author, Email: tino.ullrich@math.tu-chemnitz.de

represent the Faber-Schauder coefficients of f . The idea to approximate a function f by a truncated Faber-Schauder series

$$I_n f := \sum_{|j|_1 \leq n} \sum_k d_{j,k}^2(f) v_{j,k} \tag{1.1}$$

has been proposed by Triebel [28] and, independently, Dinh Dũng [4]. It provides a powerful tool for studying various situations of errors measured in L_q . The latter reference is even more general and uses so-called B-spline representations of functions, where the Faber system is a special case. A systematic discretization of multivariate functions with mixed smoothness in terms of Faber coefficients has been given [9] and [3], see also [5] for further history. Very recently, the authors in [10] used the Faber system to analyze the path regularity Brownian motion. Surprisingly, the proposed method turned out to be sharp in several regimes. In this paper we study an endpoint situation, where $r = 1/p$ and $q \leq 1$. This still allows for an embedding into the continuous functions (and makes therefore function evaluations possible). However, the embedding is not compact.

Recent observations regarding the problem of optimal sampling recovery of function classes in L_2 bring classes with mixed smoothness to the focus again. Since several newly developed techniques only work for Hilbert-Schmidt operators [11], [16], [1] or, more generally, in situations where certain asymptotic characteristics (approximation numbers) are square summable [12], [6] we need new techniques in situations where this is not the case. In [26, 25] the range of small smoothness has been considered where one is far away from square summability of corresponding widths. Still we have the compact embedding into L_∞ for those examples. This embedding seems to be of crucial importance since the sampling numbers in L_2 can be related to the Kolmogorov numbers in L_∞ . It has been shown in [24, 1]

$$\varrho_{bm}(\mathbf{F}, L_2) \leq \frac{C}{(b-1)^{3/2}} d_m(\mathbf{F}, L_\infty). \tag{1.2}$$

with some absolute constant $C > 0$ and an oversampling constant $b > 1$. In other words, a compact embedding into L_∞ allows us to use the decaying Kolmogorov numbers for controlling the sampling numbers. As discussed in [26] sparse grid techniques [2] perform asymptotically worse by a log-factor for classes with mixed smoothness compactly embedded into L_∞ .

In this paper we continue research in this direction. Note, that there are several relevant (multivariate) function classes \mathbf{F} which are continuously embedded in L_∞ but not compactly. Thus, (1.2) is not applicable. Almost nothing is known how to reconstruct functions from samples, which only satisfy a Besov regularity with smoothness $r = 1/p$ or Sobolev type regularity with $r = 1$ and $p = 1$. As a first step we prove the following relation for the Faber-Schauder coefficients

$$\sup_{j \in \mathbb{N}_{-1}^d} \left(\sum_{k \in \mathbb{Z}^d} |d_{j,k}^2(f)|^p \right)^{1/p} \lesssim \|f\|_{S_{p,1}^{1/p} B(\mathbb{R}^d)}.$$

This relation directly implies that the truncated Faber-Schauder representation (1.1) still works well when we consider errors in L_q with $q < \infty$. We make progress towards the solution of an open problem mentioned in [31]. The univariate class $B_{p,1}^{1/p}$ and its approximation by equidistant samples in $[0, 1]$ has been considered decades ago by Oswald [17]. Our main results read as follows. In case $1 \leq p < \infty$ there is a constant C depending on p and d such that

$$\|f - I_n f\|_p \leq C 2^{-n/p} n^{d-1} \|f\|_{S_{p,1}^{1/p} B([0, 1]^d)}$$

holds for all $n \in \mathbb{N}$. In the case $1 \leq p < q < \infty$ we are able to prove the following bound.

$$\|f - I_n f\|_q \leq C 2^{-n/q} n^{(d-1)/q} \|f\|_{S_{p,1}^{1/p} B([0,1]^d)}.$$

Due to embeddings the latter bound also holds true for $S_{1,\infty}^1 F([0,1]^d)$. All these bounds can be rephrased in terms of linear sampling numbers. In case $1 \leq p < q < \infty$ we obtain, for instance,

$$m^{-1/q} \lesssim \varrho_m^{\text{lin}}(S_{p,1}^{1/p} B([0,1]^d), L_q([0,1]^d)) \lesssim (m^{-1} \log^2(d-1) m)^{1/q}.$$

Notation. As usual \mathbb{N} denotes the natural numbers, $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, $\mathbb{N}_{-1} = \mathbb{N}_0 \cup \{-1\}$, \mathbb{Z} denotes the integers, \mathbb{R} the real numbers, and \mathbb{C} the complex numbers. The letter d is always reserved for the underlying dimension in $\mathbb{R}^d, \mathbb{Z}^d$ etc. For $a \in \mathbb{R}$ we denote $a_+ := \max\{a, 0\}$. For $0 < p \leq \infty$ and $x \in \mathbb{R}^d$ we denote $|x|_p = (\sum_{i=1}^d |x_i|^p)^{1/p}$ with the usual modification in the case $p = \infty$. We further denote $x_+ := ((x_1)_+, \dots, (x_d)_+)$ and $|x|_+ := |x_+|_1$. By $(x_1, \dots, x_d) > 0$ we mean that each coordinate is positive. If X and Y are two (quasi-)normed spaces, the (quasi-)norm of an element x in X will be denoted by $\|x\|_X$. The symbol $X \hookrightarrow Y$ indicates that the identity operator is continuous. For two sequences a_n and b_n we will write $a_n \lesssim b_n$ if there exists a constant $c > 0$ such that $a_n \leq c b_n$ for all n . We will write $a_n \asymp b_n$ if $a_n \lesssim b_n$ and $b_n \lesssim a_n$.

2 The tensor Faber bases

As a main tool we will use decompositions of functions in terms of a Faber series expansion.




Figure 1: Univariate hierarchical Faber basis on $[0, 1]$ for levels $j \in \{0, 1\}$ and their union.

2.1 The univariate Faber basis

Let us briefly recall the basic facts about the Faber basis taken from [28, 3.2.1, 3.2.2]. For $j \in \mathbb{N}_0$ and $k \in \mathbb{D}_j := \{0, 1, \dots, 2^j - 1\}$ we denote by $I_{j,k}$ the dyadic interval

$$I_{j,k} = [2^{-j}k, 2^{-j}(k+1)].$$

Definition 2.1. [The univariate Faber system] Let

$$h(t) = \begin{cases} 1 & : t \in [0, 1/2], \\ -1 & : t \in [1/2, 1], \\ 0 & : \text{otherwise.} \end{cases}$$

the Haar function and $v(x)$ be the integrated Haar function, i.e.,

$$v(x) := 2 \int_0^x h(t) dt \quad , \quad x \in \mathbb{R},$$

and for $j \in \mathbb{N}_0, k \in \mathbb{D}_j$

$$v_{j,k}(\cdot) = v(2^j \cdot -k).$$

For notational reasons we let $v_{-1,0} := x$ and $v_{-1,1} := 1 - x$ for $j = -1$ and obtain the Faber system

$$F := \{v_{j,k} : j \in \mathbb{N}_{-1}, k \in \mathbb{D}_j\},$$

where $\mathbb{D}_{-1} := \mathbb{D}_1 = \{0, 1\}$.

Faber [7] observed that every continuous (non-periodic) function f on $[0, 1]$ can be represented as

$$f(x) = f(0) \cdot (1 - x) + f(1) \cdot x - \frac{1}{2} \sum_{j=0}^{\infty} \sum_{k=0}^{2^j-1} \Delta_{2^{-j-1}}^2(f, 2^{-j}k) v_{j,k}(x)$$

with uniform convergence, see e.g. [28, Theorem 2.1, Step 4]. The analysis of Triebel-Lizorkin and Besov spaces on \mathbb{R} as defined in section 4 requires a version of the Faber-Schauder representation acting on \mathbb{R} . For this purpose we extend the number of translations to the whole integers and obtain

$$f(x) = \sum_{k \in \mathbb{Z}} f(k) v_{-1,k}(x) - \frac{1}{2} \sum_{j=0}^{\infty} \sum_{k \in \mathbb{Z}} \Delta_{2^{-j-1}}^2(f, 2^{-j}k) v_{j,k}(x), \quad (2.1)$$

where $v_{-1,k}(\cdot) := v_{0,0}((\cdot + 1 + k)/2)$.

2.2 The tensor Faber basis

Let now $f(x_1, \dots, x_d)$ be a d -variate function $f \in C(\mathbb{R}^d)$. By fixing all variables except x_i we obtain by $g(\cdot) = f(x_1, \dots, x_{i-1}, \cdot, x_{i+1}, \dots, x_d)$ a univariate continuous function. By applying (2.1) in every such component we obtain the representation

$$f(x) = \sum_{j \in \mathbb{N}_{-1}^d} \sum_{k \in \mathbb{Z}^d} d_{j,k}^2(f) v_{j,k}(x) \quad (2.2)$$

in $C(K)$, $K \subset \mathbb{R}^d$ compact, where

$$v_{j,k}(x_1, \dots, x_d) := v_{j_1, k_1}(x_1) \cdots v_{j_d, k_d}(x_d) \quad , \quad j \in \mathbb{N}_{-1}^d, k \in \mathbb{Z}^d,$$

and

$$d_{j,k}^2(f) = (-2)^{-|e(j)|} \Delta_{2^{-(j+1)}}^{2, e(j)}(f, x_{j,k}) \quad , \quad j \in \mathbb{N}_{-1}^d, k \in \mathbb{Z}^d.$$

Here we put $e(j) = \{i : j_i \neq -1\}$ and $x_{j,k} = (2^{-(j_1)+k_1}, \dots, 2^{-(j_d)+k_d})$.

In Section 6 we apply the Faber-Schauder series expansion for functions on the d -variate unit cube $[0, 1]^d$. For this purpose we simply truncate the series expansion to all translations whose support has a non-empty intersection with $[0, 1]^d$. That is

$$\mathbb{D}_j := \mathbb{D}_{j_1} \times \cdots \times \mathbb{D}_{j_d} \quad , \quad j = (j_1, \dots, j_d) \in \mathbb{N}_{-1}^d.$$

With similar arguments as above we obtain for $f \in C([0, 1]^d)$ the representation

$$f(x) = \sum_{j \in \mathbb{N}_{-1}^d} \sum_{k \in \mathbb{D}_j} d_{j,k}^2(f) v_{j,k}(x). \quad (2.3)$$

3 Faber-Schauder coefficients and bandlimited functions

In the sequel we denote with \mathbb{I}^d one of the tensor domains $[0, 1]^d$ or \mathbb{R}^d . The space $L_p(\mathbb{I}^d)$, $0 < p \leq \infty$, contains all measurable functions $f : \mathbb{I}^d \rightarrow \mathbb{C}$ satisfying

$$\|f\|_p = \left(\int_{\mathbb{I}^d} |f(x)|^p dx \right)^{1/p} < \infty$$

with the usual modification in case $p = \infty$. The space $C(\mathbb{I}^d)$ is often used as a replacement for $L_\infty(\mathbb{I}^d)$. It denotes the collection of all continuous and bounded d -variate functions equipped with the uniform norm. The computation of the Fourier transform (and its inverse) of an L_1 -integrable d -variate function is performed by the integrals ($\xi \in \mathbb{R}^d$)

$$\mathcal{F}f(\xi) = \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} f(x) e^{-i\xi \cdot x} dx \quad , \quad \mathcal{F}^{-1}f(\xi) = \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} f(x) e^{i\xi \cdot x} dx .$$

To begin with, we recall the concept of a dyadic decomposition of unity. The space $C_0^\infty(\mathbb{R}^d)$ consists of all infinitely many times differentiable compactly supported functions.

Definition 3.1. *Let $\Phi(\mathbb{R})$ be the collection of all systems $\varphi = \{\varphi_n(x)\}_{n=0}^\infty \subset C_0^\infty(\mathbb{R})$ satisfying*

- (i) $\text{supp } \varphi_0 \subset \{x : |x| \leq 2\}$,
- (ii) $\text{supp } \varphi_n \subset \{x : 2^{n-1} \leq |x| \leq 2^{n+1}\}$, $n = 1, 2, \dots$,
- (iii) For all $\ell \in \mathbb{N}_0$ it holds $\sup_{x,n} 2^{n\ell} |D^\ell \varphi_n(x)| \leq c_\ell < \infty$,
- (iv) $\sum_{n=0}^\infty \varphi_n(x) = 1$ for all $x \in \mathbb{R}$.

Now we fix a system $\varphi = \{\varphi_n\}_{n \in \mathbb{Z}} \in \Phi(\mathbb{R})$, where we put $\varphi_n \equiv 0$ if $n < 0$. For $\ell = (\ell_1, \dots, \ell_d) \in \mathbb{Z}^d$ let the building blocks f_ℓ be given by

$$f_\ell(x) = \mathcal{F}^{-1}[\varphi_{\ell_1}(\xi_1) \cdots \varphi_{\ell_d}(\xi_d) \mathcal{F}f(\xi)](x) \quad , \quad x \in \mathbb{R}^d . \quad (3.1)$$

Because of the Paley-Wiener theorem the functions f_ℓ are entire analytic functions and therefore continuous. The goal of this section is to derive bounds for fixed ‘‘levels’’ j of the Faber-Schauder expansion (2.2) of such a bandlimited function f_ℓ . To be more precise, we aim at bounds for $\left\| \sum_{k \in \mathbb{Z}^d} d_{j,k}^2(f_\ell) v_{j,k}(\cdot) \right\|_p$. Clearly, due to the compact support of $v_{j,k}$ we may replace $v_{j,k}$ by the characteristic function $\chi_{j,k}$ of the parallelepiped $[2^{-j_1} k_1, 2^{-j_1} (k_1 + 1)] \times \cdots \times [2^{-j_d} k_d, 2^{-j_d} (k_d + 1)]$. Note, that for any continuous function $f \in C(\mathbb{R}^d)$

$$\left\| \sum_{k \in \mathbb{Z}^d} d_{j,k}^2(f) v_{j,k}(\cdot) \right\|_p \asymp \left\| \sum_{k \in \mathbb{Z}^d} d_{j,k}^2(f) \chi_{j,k}(\cdot) \right\|_p \asymp \left(2^{-|j|_1} \sum_{k \in \mathbb{Z}^d} |d_{j,k}^2(f)|^p \right)^{1/p} .$$

To do so we need some tools from harmonic analysis. We state a mixed version of the Peetre maximal inequality, proved in [22, 1.6.4].

Lemma 3.2. *[Peetre maximal inequality] Let $0 < p < \infty$ and $a > 1/p$. Let further $f \in L_1(\mathbb{R}^d)$ such that $\text{supp } \mathcal{F}f \subset [-b_1, b_1] \times \cdots \times [-b_d, b_d]$. Then there is a constant $c > 0$, only depending on a and p but not on f and b , such that*

$$\left\| \sup_{y \in \mathbb{R}^d} \frac{|f(x+y)|}{(1+b_1|y_1|)^a \cdots (1+b_d|y_d|)^a} \right\|_p \leq c \|f\|_p .$$

The following univariate pointwise estimate connecting differences of bandlimited functions to Peetre maximal operator is taken from [30, Lem. 3.3.1].

Lemma 3.3. *Let $a, b > 0$ and $f \in L_1(\mathbb{R})$ with $\text{supp } \mathcal{F}f \subset [-b, b]$. Then there exists a constant $C > 0$ such that*

$$|\Delta_h^m f(x)| \leq C \min\{1, |bh|^m\} \max\{1, |bh|^a\} \sup_{y \in \mathbb{R}} \frac{|f(x+y)|}{(1+|2^{j+\ell}y|)^a}.$$

The bound may be slightly improved when replacing pointwise estimates by estimates involving L_p norms. We have the following.

Lemma 3.4. *Let $j, \ell \in \mathbb{N}_0$ and $f \in C(\mathbb{R})$. Then*

$$\left\| \sup_{|h| \lesssim 2^{-j}} |f(\cdot + h)| \right\|_p \lesssim 2^{\ell/p} \left\| \sup_{|h| \lesssim 2^{-(j+\ell)}} |f(x+h)| \right\|_p \lesssim 2^{\ell/p} \left\| \sup_{y \in \mathbb{R}} \frac{f(x+y)}{(1+|2^{j+\ell}y|)^a} \right\|_p.$$

Proof. We start with a pointwise estimate

$$\begin{aligned} \sup_{|h| \leq 2^{-j}} |f(x+h)| &\leq \sup_k \sup_{|h| \leq 2^{-(j+\ell)}} |f(x+k2^{-(j+\ell)}+h)| \\ &\leq \left(\sum_k \sup_{|h| \lesssim 2^{-(j+\ell)}} |f(x+k2^{-(j+\ell)}+h)|^p \right)^{1/p}. \end{aligned}$$

Taking L_p -norms on both sides gives

$$\begin{aligned} \left\| \sup_{|h| \leq 2^{-j}} |f(x+h)| \right\|_p &\leq \left(\sum_k \int_{\mathbb{R}} \sup_{|h| \leq 2^{-(j+\ell)}} |f_{j+\ell}(x+k2^{-(j+\ell)}+h)|^p dx \right)^{1/p} \\ &\lesssim 2^{\ell/p} \left\| \sup_{|h| \lesssim 2^{-(j+\ell)}} |f(x+h)| \right\|_p. \end{aligned}$$

■

In the next lemma we combine both univariate bounds and derive a multivariate estimate via iteration with respect to coordinate directions. Let $f \in L_1(\mathbb{R}^d)$ and $f_{j+\ell}$ denotes the bandlimited function from (3.1).

Lemma 3.5. *Let $j \in \mathbb{N}_{-1}^d$ and $\ell \in \mathbb{Z}^d$. Then we have*

$$\left\| \sum_{k \in \mathbb{Z}^d} d_{j,k}^2(f_{j+\ell}) \chi_{j,k}(\cdot) \right\|_p \lesssim \|f_{j+\ell}\|_p \prod_{i=1}^d \min\{2^{2\ell_i}, 1\} \max\{2^{\ell_i/p}, 1\}.$$

Proof. *Step 1.* To provide a technically transparent proof of this Lemma we start with the univariate case ($d = 1$). In the second part of this proof we deal with the multivariate case which requires more involved notation. We put for $x \in \mathbb{R}$

$$F_{j,\ell}(x) := \sum_{k \in \mathbb{Z}} d_{j,k}^2(f_{j+\ell}) \chi_{j,k}(x).$$

Let $x \in [2^{-j}k, 2^{-j}(k+1)]$. For this x we have

$$|d_{j,k}^2(f_{j+\ell}) \chi_{j,k}(x)| \lesssim |f_{j+\ell}(k \cdot 2^{-j} + 2 \cdot 2^{-(j+1)})| + 2|f_{j+\ell}(k \cdot 2^{-j} + 2^{-(j+1)})|$$

$$+|f_{j+l}(k \cdot 2^{-j})| \lesssim \sup_{|h| \lesssim 2^{-j}} |f_{j+l}(x+h)|.$$

Since $\chi_{j,k}(x)$ do not have overlap we receive

$$|F_{j,l}(x)| = \left| \sum_{k \in \mathbb{Z}} d_{j,k}^2(f_{j+l}) \chi_{j,k}(x) \right| \lesssim \sup_{|h| \lesssim 2^{-j}} |f_{j+l}(x+h)|.$$

By Lemma 3.4 we find

$$\|F_{j,l}\|_p \lesssim \max\{2^{\ell/p}, 1\} \left\| \sup_{y \in \mathbb{R}} \frac{f_{j+l}(x+y)}{(1+|2^{j+\ell}y|)^a} \right\|_p \quad (3.2)$$

for some $a > 0$ which is at our disposal.

In case $l \leq 0$ we may continue arguing pointwise. First of all, we have

$$|F_{j,l}(x)| \leq \sup_{|y| \lesssim 2^{-j}} |\Delta_{2^{-j-1}}^2 f_{j+l}(x+y)|.$$

Using Lemma 3.3 and the fact that $l \leq 0$ we obtain

$$|F_{j,l}(x)| \leq 2^{2l} \sup_{y \in \mathbb{R}} \frac{|f(x+y)|}{(1+|2^{j+\ell}y|)^a}. \quad (3.3)$$

Combining (3.2) and (3.3) gives

$$\|F_{j,l}(x)\|_p \lesssim \left\| \sup_{y \in \mathbb{R}} \frac{|f(x+y)|}{(1+|2^{j+\ell}y|)^a} \right\|_p \max\{1, 2^{l/p}\} \min\{1, 2^{2l}\}.$$

Choosing $a > 1/p$ and applying the Peetre maximal inequality in Lemma 3.2 gives the result for $d = 1$.

Step 2. We deal with the multivariate case and start with a pointwise estimate of

$$F_{j,\ell}(x) := \sum_{k \in \mathbb{Z}^d} d_{j,k}^2(f_{j+\ell}) \chi_{j,k}(x),$$

where we apply the above procedure in every direction. In order not to drown in notation we introduce the following direction-wise maximal operator

$$\mathfrak{M}_j^i f(x) := \sup_{|h| \lesssim 2^{-j}} |f(x + e_i h)|,$$

where $e_i = (\underbrace{0, \dots, 0}_{i-1}, 1, \underbrace{0, \dots, 0}_{d-i})$. Clearly, we have for $x \in \mathbb{R}^d$

$$|F_{j,\ell}(x)| \lesssim \left(\prod_{i:\ell_i > 0} \mathfrak{M}_j^i \right) \circ \left(\prod_{i:\ell_i \leq 0} \mathfrak{M}_j^i \right) \circ \left(\prod_{i:\ell_i \leq 0} \Delta_{2^{-j-1}}^{2,i} \right) f(x).$$

Here we use that we have in every direction

$$|\Delta_{2^{-j-1}}^{2,i} f_{j+\ell}| \leq \mathfrak{M}_j^i f_{j+\ell}(x)$$

including the case $j = -1$ where the difference is replaced by the function value at the respective point. This case is included in $\left(\prod_{i:\ell_i > 0} \mathfrak{M}_j^i \right)$ since for $\ell < 0$ there is nothing to prove in this case. We use the triangle inequality in order to estimate the difference by point evaluations.

Taking the sup leads to the direction-wise maximal operator. In case $\ell \leq 0$ we keep the direction-wise difference. Nevertheless, in order to get rid of the characteristic function we have to additionally apply a direction-wise sup also in this case. Since it holds that

$$\mathfrak{M}_{j_i}^i f(x) \leq \mathfrak{M}_{j_i}^i g(x)$$

with $|f| \leq |g|$ we first estimate

$$\left| \left(\prod_{i:\ell_i \leq 0} \Delta_{2^{-j-1}}^{2,i} \right) f_{j+\ell}(x) \right|$$

pointwise from above using Lemma 3.3 iteratively. This gives

$$\left(\prod_{i:\ell_i \leq 0} \Delta_{2^{-j-1}}^{2,i} \right) f_{j+\ell}(x) \lesssim \sup_{\substack{y_i \in \mathbb{R} \\ i:\ell_i \leq 0}} \frac{|f_{j+\ell}(x + \sum_{i:\ell_i \leq 0} y_i e_i)|}{\prod_{i:\ell_i \leq 0} (1 + 2^{\ell_i + j_i} |y_i|)^a} \prod_{i:\ell_i \leq 0} 2^{2\ell_i}.$$

for any $a > 0$. The maximal operators $(\prod_{i:\ell_i \leq 0} \mathfrak{M}_{j_i}^i)$ go on the same coordinates. Since $\ell_i \leq 0$ we clearly have also

$$\left(\prod_{i:\ell_i \leq 0} \mathfrak{M}_{j_i}^i \right) \left(\prod_{i:\ell_i \leq 0} \Delta_{2^{-j-1}}^{2,i} \right) f_{j+\ell}(x) \lesssim \sup_{\substack{y_i \in \mathbb{R} \\ i:\ell_i \leq 0}} \frac{|f(x + \sum_{i:\ell_i \leq 0} y_i e_i)|}{\prod_{i:\ell_i \leq 0} (1 + 2^{\ell_i + j_i} |y_i|)^a} \prod_{i:\ell_i \leq 0} 2^{2\ell_i}.$$

It remains to apply $(\prod_{i:\ell_i > 0} \mathfrak{M}_{j_i}^i)$ and take the L_p -norm. Here we use Lemma 3.4 iteratively.

$$\begin{aligned} & \int_{\mathbb{R}^d} |F_{j,\ell}(x)|^p dx \\ & \lesssim \int \cdots \int_{\{(x_i)_{i:\ell_i \leq 0}\}} \int \cdots \int_{\{(x_i)_{i:\ell_i > 0}\}} \left| \left(\prod_{i:\ell_i > 0} \mathfrak{M}_{j_i}^i \right) \circ \left(\prod_{i:\ell_i \leq 0} \mathfrak{M}_{j_i}^i \right) \circ \left(\prod_{i:\ell_i \leq 0} \Delta_{2^{-j-1}}^{2,i} \right) f_{j+\ell}(x) \right|^p \prod_{i:\ell_i > 0} dx_i \prod_{i:\ell_i \leq 0} dx_i \\ & \lesssim \prod_{i=1}^d \min\{2^{2p\ell_i}, 1\} \max\{2^{\ell_i}, 1\} \times \\ & \quad \times \int \cdots \int_{\{(x_i)_{i:\ell_i \leq 0}\}} \int \cdots \int_{\{(x_i)_{i:\ell_i > 0}\}} \left| \left(\prod_{i:\ell_i > 0} \mathfrak{M}_{j_i + \ell_i}^i \right) \sup_{\substack{y_i \in \mathbb{R} \\ i:\ell_i \leq 0}} \frac{|f_{j+\ell}(x + \sum_{i:\ell_i \leq 0} y_i e_i)|}{\prod_{i:\ell_i \leq 0} (1 + 2^{\ell_i + j_i} |y_i|)^a} \right|^p \prod_{i:\ell_i > 0} dx_i \prod_{i:\ell_i \leq 0} dx_i. \end{aligned}$$

Finally, we use the trivial estimate (already used in Lemma 3.4) to replace the operators $\mathfrak{M}_{j_i + \ell_i}^i$ by the Peetre maximal function. This gives

$$\begin{aligned} \int_{\mathbb{R}^d} |F_{j,\ell}(x)|^p dx & \lesssim \prod_{i=1}^d \min\{2^{2p\ell_i}, 1\} \max\{2^{\ell_i}, 1\} \int_{\mathbb{R}^d} \left| \sup_{y \in \mathbb{R}^d} \frac{|f_{j+\ell}(x+y)|}{\prod_{i=1}^d (1 + |2^{\ell_i + j_i} y_i|)^a} \right|^p dx \\ & \lesssim \prod_{i=1}^d \min\{2^{2p\ell_i}, 1\} \max\{2^{\ell_i}, 1\} \|f_{j+\ell}\|_p^p, \end{aligned}$$

if we choose $a > 1/p$. ■

4 Besov-Triebel-Lizorkin spaces with mixed smoothness

For the definition of the corresponding function spaces on \mathbb{R}^d we refer to [22], [32], and [3]. The corresponding function spaces on $[0, 1]^d$ are defined via restrictions of functions on \mathbb{R}^d , see [3, Sect. 3.4]. In this section mainly focus on the definition of Besov and Triebel-Lizorkin spaces with dominating mixed smoothness on \mathbb{R}^d since they are crucial for our subsequent analysis. We closely follow [22, Chapter 2] and use the building blocks $f_j(\cdot)$ defined in (3.1).

Definition 4.1. (*Mixed Besov and Triebel-Lizorkin spaces*)

(i) Let $0 < p, q \leq \infty$ and $r > \sigma_p := (1/p - 1)_+$. Then $S_{p,q}^r B(\mathbb{R}^d)$ is defined as the collection of all $f \in L_1(\mathbb{R}^d)$ such that

$$\|f|S_{p,q}^r B(\mathbb{R}^d)\|^\varphi := \left(\sum_{j \in \mathbb{N}_0^d} 2^{|j|1rq} \|f_j\|_p^q \right)^{1/q}$$

is finite (usual modification in case $q = \infty$).

(ii) Let $0 < p < \infty$ and $r > \sigma_p := (1/p - 1)_+$. Then $S_{p,q}^r F(\mathbb{R}^d)$ is defined as the collection of all $f \in L_1(\mathbb{R}^d)$ such that

$$\|f|S_{p,q}^r F(\mathbb{R}^d)\|^\varphi := \left\| \left(\sum_{j \in \mathbb{N}_0^d} 2^{|j|1rq} |f_j(x)|^q \right)^{1/q} \right\|_p$$

is finite.

Recall, that this definition is independent of the chosen system φ in the sense of equivalent (quasi-)norms. Moreover, in case $\min\{p, q\} \geq 1$ the defined spaces are Banach spaces, whereas they are quasi-Banach spaces in case $\min\{p, q\} < 1$. For details confer [22, 2.2.4]. In the next lemma there appears the condition $r > (1/p - 1)_+$ which is caused by the parameter range in our Definition 4.1. All the subsequent embeddings of course also hold true for the general situation, where $r \in \mathbb{R}$. We have the following elementary embeddings, see [22, 2.2.3].

Lemma 4.2. Let $0 < p < \infty$, $r > (1/p - 1)_+$, and $0 < q \leq \infty$. Let further $A \in \{F, B\}$.

(i) If $\varepsilon > 0$ and $0 < v \leq \infty$ then

$$S_{p,q}^{r+\varepsilon} A(\mathbb{R}^d) \hookrightarrow S_{p,v}^r A(\mathbb{R}^d).$$

(ii) If $p < u < \infty$, $0 < q, q_1, q_2 \leq \infty$ and $r - 1/p = t - 1/u$ then

$$\begin{aligned} S_{p,q}^r B(\mathbb{R}^d) &\hookrightarrow S_{u,q}^t B(\mathbb{R}^d), \\ S_{p,q_1}^r F(\mathbb{R}^d) &\hookrightarrow S_{u,q_2}^t F(\mathbb{R}^d). \end{aligned}$$

(iib) (Jawerth-Franke embedding I) If $0 < p < u < \infty$, $0 < w \leq \infty$ and $r - 1/p = t - 1/u$ then

$$S_{p,u}^r B(\mathbb{R}^d) \hookrightarrow S_{u,w}^t F(\mathbb{R}^d).$$

(iic) (Jawerth-Franke embedding II) If $0 < p < u < \infty$, $0 < q \leq \infty$ and $r - 1/p = t - 1/u$ then

$$S_{p,q}^r F(\mathbb{R}^d) \hookrightarrow S_{u,p}^t B(\mathbb{R}^d).$$

(iiia) If $r > 1/p$ then

$$S_{p,q}^r B(\mathbb{R}^d) \hookrightarrow C(\mathbb{R}^d).$$

(iiib) if $r = 1/p$ and $q \leq 1$ then we still have

$$S_{p,q}^{1/p} B(\mathbb{R}^d) \hookrightarrow C(\mathbb{R}^d).$$

(iiic) It holds

$$S_{1,\infty}^1 F(\mathbb{R}^d) \hookrightarrow C(\mathbb{R}^d).$$

(iv) If $1 < p < \infty$ and $r > 0$ then

$$S_{p,\min\{p,q\}}^r B(\mathbb{R}^d) \hookrightarrow S_{p,q}^r F(\mathbb{R}^d) \hookrightarrow S_{p,\max\{p,q\}}^r B(\mathbb{R}^d).$$

Proof. The embeddings (i), (ia), (iia), (iiib), (iv) are standard and can be found in [22, Chapt. 2]. As for the Jawerth-Franke type embeddings we refer to [5, Lem. 3.4.2, 3.4.3]. See also [5, Rem. 3.4.4] for further references, especially for the mixed smoothness case. Finally, the embedding in (iiic) is a consequence of (iic) and (iiib). \blacksquare

4.1 Spaces on domains

We aim for approximating functions defined on the unit cube $[0, 1]^d$ with the above regularity assumptions. This requires the definition of function spaces on domains. The domain $\Omega \subset \mathbb{R}^d$ represents an open connected set. Later, when dealing with continuous bounded functions we may use as well compact sets like $[0, 1]^d$.

Definition 4.3. Let Ω be a domain in \mathbb{R}^d .

(i) Let additionally $0 < p, q \leq \infty$ and $r > (1/p - 1)_+$. Then we define

$$S_{p,q}^r B(\Omega) := \{f \in L_1(\Omega) : \exists g \in S_{p,q}^r B(\mathbb{R}^d) \text{ with } g|_{\Omega} = f\},$$

where

$$\|f|_{S_{p,q}^r B(\Omega)}\| := \inf\{\|g|_{S_{p,q}^r B(\mathbb{R}^d)}\| : g \in S_{p,q}^r B(\mathbb{R}^d), g|_{\Omega} = f\}.$$

(ii) Let $0 < p < \infty$ and $0 < q \leq \infty$ then we define

$$S_{p,q}^r F(\Omega) := \{f \in L_1(\Omega) : \exists g \in S_{p,q}^r F(\mathbb{R}^d) \text{ with } g|_{\Omega} = f\},$$

where

$$\|f|_{S_{p,q}^r F(\Omega)}\| := \inf\{\|g|_{S_{p,q}^r F(\mathbb{R}^d)}\| : g \in S_{p,q}^r F(\mathbb{R}^d), g|_{\Omega} = f\}.$$

On bounded domains Ω all the embeddings in Lemma 4.2 keep valid. In addition we have the following embeddings. If $0 < p_2 < p_1 \leq \infty$ (F-case: $p_i < \infty$), $0 < q \leq \infty$ and $|\Omega| < \infty$, then

$$S_{p_1,q}^r F(\Omega) \hookrightarrow S_{p_2,q}^r F(\Omega)$$

and

$$S_{p_1,q}^r B(\Omega) \hookrightarrow S_{p_2,q}^r B(\Omega).$$

Clearly, this is a trivial consequence of the embedding

$$L_{p_1}(\Omega) \hookrightarrow L_{p_2}(\Omega).$$

It is well-known that spaces $S_{p,q}^r B([0, 1]^d)$ with sufficiently large smoothness, namely $r > 1/p$, are compactly embedded into L_∞ . This is a direct consequence of results on entropy numbers of the classes $S_{\infty,q}^r B([0, 1]^d)$ in L_∞ , see [15, Cor. 23,(iii)] or [26, Thm. 6.2] and the embeddings stated in Lemma 4.2 above.

However, in case $r = 1/p$ we do not have a compact embedding. For the convenience of the reader we give a direct proof.

Lemma 4.4. *[Non-compactness of limiting embeddings] (i) Let $0 < p < \infty$ and $q \leq 1$. Then the embedding*

$$S_{p,q}^{1/p} B([0, 1]^d) \hookrightarrow L_\infty([0, 1]^d) \quad (4.1)$$

is not compact.

(ii) *If $p \leq 1$ and $0 < q \leq \infty$ then the embedding*

$$S_{p,q}^{1/p} F([0, 1]^d) \hookrightarrow L_\infty([0, 1]^d) \quad (4.2)$$

is not compact.

Proof. We show the non-compactness of the embedding (4.1) first in case $d = 1$. Clearly, by standard (tensorization) arguments this would imply the non-compactness also in higher dimensions. Note further, that the non-compactness of (4.2) is implied by the Jawerth-Franke embedding (see Lemma 4.2, (iib))

$$S_{p_1,p_2}^{1/p_1} B([0, 1]^d) \hookrightarrow S_{p_2,q}^{1/p_2} F([0, 1]^d)$$

together with the non-compactness of (4.1). So, it remains to prove the non-compactness of $B_{p,q}^{1/p}([0, 1]) \hookrightarrow L_\infty([0, 1])$ for any $q \leq 1$ and $0 < p < \infty$. This fact is certainly known and can be found in the literature. However, we would like to give a direct argument here since it fits to the scope of the paper. The idea is to find a sequence $(g_j)_j$ of functions with $\|g_j|_{B_{p,q}^{1/p}}\| \leq 1$ for all j which is not convergent in $L_\infty([0, 1])$. A straight-forward choice for the g_j are the Faber-Schauder basis functions (L_∞ -normalized hat functions) for different levels j . To be more precise we let $j = 0, 1, 2, \dots$ and consider the sequence $(g_j)_j := (v_{j,0})_{j \in \mathbb{N}}$. Clearly, we always have

$$\sup_{|h| \leq 2^{-k}} \|\Delta_h^m v_{j,0}\|_p \lesssim 2^{-j/p}.$$

In case $k > j$ we even obtain (due to cancellation)

$$\sup_{|h| \leq 2^{-k}} \|\Delta_h^m v_{j,0}\|_p \lesssim 2^{-k/p} 2^{j-k},$$

see for instance [9, (3.19),(3.20)]. Note, that here we have to use differences of sufficiently high order $m > s$ since $1/p$ may get large. This implies

$$\begin{aligned} \|v_{j,0}|_{B_{p,q}^s}\| &\asymp \|v_{j,0}\|_1 + \left(\sum_{k=0}^{\infty} [2^{ks} \sup_{|h| \leq 2^{-k}} \|\Delta_h v_{j,0}\|_p]^q \right)^{1/q} \\ &\asymp 2^{-j} + \left(\sum_{k \leq j} [2^{ks} 2^{-j/p}]^q + \sum_{k > j} [2^{ks} 2^{-k/p} 2^{j-k}]^q \right)^{1/q} \\ &\asymp 1 \end{aligned}$$

in case $s \leq 1/p$. Hence, the elements $v_{j,0}$ have uniformly (in j) bounded quasi-norm in $B_{p,q}^s$. However, it holds

$$\|g_j - g_\ell\|_\infty \geq 1$$

if $j \neq \ell$, see for instance Figure 1. This directly disproves the compactness of the unit ball of $B_{p,q}^{1/p}$ in L_∞ . \blacksquare

Remark 4.5. *We need to formulate the definition of the Kolmogorov widths to demonstrate some relations with approximative characteristics considering here. For a compact set $\mathbf{F} \hookrightarrow X$ of a Banach space X define the Kolmogorov widths:*

$$d_m(\mathbf{F}, X) := \inf_{\{u_i\}_{i=1}^m \subset X} \sup_{f \in \mathbf{F}} \inf_{c_i} \left\| f - \sum_{i=1}^m c_i u_i \right\|_X, \quad m = 1, 2, \dots,$$

and

$$d_0(\mathbf{F}, X) := \sup_{f \in \mathbf{F}} \|f\|_X.$$

Clearly, considering \mathbf{F} as the unit ball in $S_{p,1}^{1/p} B([0, 1]^d)$ and $X = L_\infty([0, 1]^d)$ then

$$d_m(S_{p,q}^{1/p} B([0, 1]^d), L_\infty([0, 1]^d)) \rightarrow 0$$

by Lemma 4.4.

5 The decay of the Faber-Schauder coefficients

Now we are ready for proving our main tool: an assertion about the decay of the Faber-Schauder coefficients of functions from $S_{p,1}^{1/p} B(\mathbb{R}^d)$. In order to do so we need to define doubly indexed sequence spaces first.

Definition 5.1. *Let $0 < p, q \leq \infty$ and $r \geq 1/p$.*

(i) *The doubly-index sequence space $s_{p,q}^r b$ is the collection of all sequences $\{\lambda_{j,k}\}_{j \in \mathbb{N}_{-1}^d, k \in \mathbb{Z}^d}$ such that*

$$\|\lambda_{j,k} |s_{p,q}^r b\| := \left[\sum_{j \in \mathbb{N}_{-1}^d} 2^{|j|_1(r-1/p)q} \left(\sum_{k \in \mathbb{Z}^d} |\lambda_{j,k}|^p \right)^{q/p} \right]^{1/q}$$

is finite.

(ii) *Let $\Omega \subset \mathbb{R}^d$ be a compact domain. We define the index set $\mathbb{D}_j(\Omega)$ to be the set of all $k \in \mathbb{Z}^d$ such that $x_{j,k} \in \Omega$. The space $s_{p,q}^r b(\Omega)$ is defined as the space of all doubly-indexed sequences $\{\lambda_{j,k}\}_{j \in \mathbb{N}_{-1}^d, k \in \mathbb{D}_j(\Omega)}$ such that*

$$\|\lambda_{j,k} |s_{p,q}^r b(\Omega)\| := \left[\sum_{j \in \mathbb{N}_{-1}^d} 2^{|j|_1(r-1/p)q} \left(\sum_{k \in \mathbb{D}_j(\Omega)} |\lambda_{j,k}|^p \right)^{q/p} \right]^{1/q}.$$

Remark 5.2. (i) *In case $\Omega = [0, 1]^d$ we have $\mathbb{D}_j(\Omega) = \mathbb{D}_j$, which was defined right before Definition 2.1. In a certain sense the elements of $s_{p,q}^r b(\Omega)$ are restrictions of elements in $s_{p,q}^r b$ to indices related to Ω .*

(ii) *These sequence spaces already appeared in [32, Def. 2.1, 3.2]. Let $0 < p, q \leq \infty$ and $r \in \mathbb{R}$. The spaces $s_{p,q}^r b$ and $s_{p,q}^r b(\Omega)$ are Banach space if $\min\{p, q\} \geq 1$. In case $\min\{p, q\} < 1$ the space $s_{p,q}^r b$ is a quasi-Banach space. Moreover, if $u := \min\{p, q, 1\}$ it is a u -Banach space, i.e.,*

$$\|\lambda + \mu |s_{p,q}^r b\|^u \leq \|\lambda |s_{p,q}^r b\|^u + \|\mu |s_{p,q}^r b\|^u, \quad \lambda, \mu \in s_{p,q}^r b.$$

Here is the first main result.

Proposition 5.3. *Let $1 \leq p < \infty$. Then there exists a constant $c > 0$ (independent of f) such that*

$$\|d_{j,k}^2(f)|s_{p,\infty}^{1/p}b\| \leq c \|f|S_{p,1}^{1/p}B(\mathbb{R}^d)\| \quad (5.1)$$

for all $f \in S_{p,1}^{1/p}B(\mathbb{R}^d)$.

Proof. We make use of the decomposition (3.1) in a slightly modified way. For fixed $j \in \mathbb{N}_{-1}^d$ we write $f = \sum_{\ell \in \mathbb{Z}^d} f_{j+\ell}$. Putting this into (5.1) and using the triangle inequality yields

$$\begin{aligned} \|d_{j,k}^2(f)|s_{p,\infty}^{1/p}b\| &\asymp \sup_{j \in \mathbb{N}_{-1}^d} 2^{|j|_1/p} \left\| \sum_{k \in \mathbb{Z}^d} d_{j,k}^2(f) \chi_{j,k}(x) \right\|_p \\ &\leq \sup_{j \in \mathbb{N}_{-1}^d} 2^{|j|_1/p} \sum_{\ell \in \mathbb{Z}^d} \left\| \sum_{k \in \mathbb{Z}^d} d_{j,k}^2(f_{j+\ell}) \chi_{j,k}(x) \right\|_p. \end{aligned}$$

Applying Lemma 3.5 we obtain

$$\begin{aligned} \|d_{j,k}^2(f)|s_{p,\infty}^{1/p}b\| &\lesssim \sup_{j \in \mathbb{N}_{-1}^d} \sum_{\ell \in \mathbb{Z}^d} 2^{j+\ell|_1/p} \|f_{j+\ell}\|_p \prod_{i=1}^d 2^{-\ell_i/p} \min\{2^{2\ell_i}, 1\} \max\{2^{\ell_i/p}, 1\} \\ &\lesssim \left(\sum_{\ell \in \mathbb{Z}^d} 2^{j+\ell|_1/p} \|f_{j+\ell}\|_p \right) \sup_{\ell \in \mathbb{Z}^d} \prod_{i=1}^d 2^{-\ell_i/p} \min\{2^{2\ell_i}, 1\} \max\{2^{\ell_i/p}, 1\} \\ &\lesssim \|f|S_{p,1}^{1/p}B(\mathbb{R}^d)\|, \end{aligned}$$

since the sup stays finite since $1 \leq p \leq \infty$. ■

As a direct consequence we have the following restricted version

Corollary 5.4. *Let $1 \leq p < \infty$. Then there exists a constant $c > 0$ such that*

$$\|d_{j,k}^2(f)|s_{p,\infty}^{1/p}b(\Omega)\| \leq c \|f|S_{p,1}^{1/p}B(\Omega)\|$$

for all $f \in S_{p,1}^{1/p}B(\Omega)$.

Proof. Assume $f \in S_{p,1}^{1/p}B(\Omega)$. Then, by Definition 4.3 there is a $g \in S_{p,1}^{1/p}B(\mathbb{R}^d)$ with $g|_{\Omega} = f$. By Proposition 5.3 we obtain

$$\|d_{j,k}^2(g)|s_{p,\infty}^{1/p}b\| \leq c \|g|S_{p,1}^{1/p}B\|.$$

Since $d_{j,k}^2(g) = d_{j,k}^2(f)$ for $j \in \mathbb{N}_{-1}^d$ and $k \in \mathbb{D}_j(\Omega)$ we have

$$\|d_{j,k}^2(f)|s_{p,\infty}^{1/p}b(\Omega)\| \leq \|d_{j,k}^2(g)|s_{p,\infty}^{1/p}b\|$$

we obtain

$$\|d_{j,k}^2(f)|s_{p,\infty}^{1/p}b(\Omega)\| \leq c \|g|S_{p,1}^{1/p}B\|.$$

The last inequality holds for every extension g of f . Taking the infimum yields the result. ■

6 Application for sampling recovery in L_q with $q < \infty$

In this section we would like to apply the Faber-Schauder embedding in Corollary 5.4 for sampling recovery on the unit cube $\Omega = [0, 1]^d$. As we have seen in Lemma 4.4 we can not expect an error decay of a sampling recovery operator in the worst-case when we measure the error in $L_\infty([0, 1]^d)$. Hence we focus on the recovery in $L_q([0, 1]^d)$ with $q < \infty$. Based on the Faber-Schauder representation we will use a sparse grid truncation in order to obtain a recovery operator. Let

$$I_n f = \sum_{|j|_1 \leq n} \sum_{k \in \mathbb{D}_j} d_{j,k}^2(f) v_{j,k}. \quad (6.1)$$

with the notation from Section 2.

Lemma 6.1. *The following estimates hold true for $\alpha > 0$*

$$(i) \quad \sum_{|j|_1 > n} 2^{-\alpha|j|_1} \asymp n^{d-1} 2^{-\alpha n}$$

$$(ii) \text{ and } \quad \sum_{|j|_1 \leq n} 2^{|j|_1} \asymp n^{d-1} 2^n.$$

Proof. We refer to [23, p.10, Lemma D] ■

We first consider the case where $p = q$.

Theorem 6.1. Let $1 \leq p < \infty$. Then there is a constant $C > 0$ (independent of n and f) such that

$$\|f - I_n f\|_p \leq C 2^{-n/p} n^{d-1} \|f\|_{S_{p,1}^{1/p} B([0, 1]^d)}$$

holds for all $n \in \mathbb{N}$.

Proof. The representation in (2.3) allows us to express and estimate the error by

$$\begin{aligned} \|f - I_n f\|_p &= \left\| \sum_{|j|_1 > n} \sum_{k \in \mathbb{D}_j} d_{j,k}^2(f) v_{j,k} \right\|_p \\ &\leq \sum_{|j|_1 > n} \left\| \sum_{k \in \mathbb{D}_j} d_{j,k}^2(f) v_{j,k} \right\|_p 2^{|j|_1/p} 2^{-|j|_1/p} \\ &\leq \sup_j 2^{|j|_1/p} \left\| \sum_{k \in \mathbb{D}_j} d_{j,k}^2(f) v_{j,k} \right\|_p \sum_{|j|_1 > n} 2^{-|j|_1/p}. \end{aligned}$$

Finally applying Proposition 5.3 or Corollary 5.4, respectively, yields

$$\begin{aligned} \|f - I_n f\|_p &\lesssim \left\| d_{j,k}^2(f) \Big|_{S_{p,\infty}^{1/p} b([0, 1]^d)} \right\| \sum_{|j|_1 > n} 2^{-|j|_1/p} \\ &\lesssim 2^{-n/p} n^{d-1} \|f\|_{S_{p,1}^{1/p} B([0, 1]^d)}, \end{aligned}$$

where the sum is estimated by Lemma 6.1. ■

Remark. An one-dimensional version of the above result has been proved by Oswald [17] almost forty years ago.

In the situation $p < q$ we loose in the main rate. This phenomenon has been observed earlier in the literature, see [5]. We will use Jawerth-Franke type embeddings to improve the order of the logarithmic term.

Theorem 6.2. Let $1 \leq p < q < \infty$. Then

$$\|f - I_n f\|_q \leq C 2^{-n/q} n^{(d-1)/q} \|f\|_{S_{p,1}^{1/p} B([0,1]^d)}.$$

Proof. Using Jawerth-Franke I (see Lemma 4.2) we obtain

$$\begin{aligned} \|f - I_n f\|_q &\lesssim \|f - I_n f\|_{S_{p,q}^{1/p-1/q} B(\Omega)} \\ &\lesssim \left(\sum_{|j|_1 > n} 2^{q|j|_1[1/p-1/q]} \left\| \sum_{k \in \mathbb{D}_j} d_{j,k}^2(f) \chi_{j,k} \right\|_p^q \right)^{1/q} \\ &\lesssim \left[\sup_{j \in \mathbb{N}_{-1}^d} 2^{j|1|(1/p)} \left\| \sum_{k \in \mathbb{D}_j} d_{j,k}^2(f) \chi_{j,k} \right\|_p \right] \left(\sum_{|j|_1 > n} 2^{-|j|_1} \right)^{\frac{1}{q}}, \end{aligned}$$

where we used the inverse Faber-Schauder characterization for spaces with positive smoothness, cf. [3, Theorem 4.18]. This reference deals with the \mathbb{R}^d -case but can be easily extended by standard arguments as shown for instance in the proof of Corollary 5.4 or following the arguments in [3, Theorem 4.25] to the unit cube setting. Finally applying Proposition 5.3 yields

$$\|f - I_n f\|_q \lesssim 2^{-n/q} n^{(d-1)/q} \|f\|_{S_{p,1}^{1/p} B([0,1]^d)}.$$

■

Let us now deal with the spaces $S_{1,\infty}^1 F([0,1]^d)$, which are embedded into $C([0,1]^d)$ as Lemma 4.2, (iiic) shows. By Jawerth-Franke embedding II (Lemma 4.2, (iic)) we even know that for every $1 < p < \infty$ we have $S_{1,\infty}^1 F([0,1]^d) \hookrightarrow S_{p,1}^{1/p} B([0,1]^d)$. As a direct corollary of Theorems 6.1, 6.2 we obtain the following

Theorem 6.3. (i) It holds for any small $\varepsilon > 0$

$$\|f - I_n f\|_1 \lesssim_\varepsilon 2^{-n(1-\varepsilon)} \|f\|_{S_{1,\infty}^1 F([0,1]^d)}.$$

(ii) For any $1 < q < \infty$ we have

$$\|f - I_n f\|_q \lesssim 2^{-n/q} n^{(d-1)/q} \|f\|_{S_{1,\infty}^1 F([0,1]^d)}.$$

Proof. By the embedding $S_{1,\infty}^1 F([0,1]^d) \hookrightarrow S_{p,1}^{1/p} B([0,1]^d)$ for $p > 1$ we may apply Theorem 6.2 to obtain the result. For (i) we simply choose $q > 1$ small and use the fact that the L_q -norm dominates the L_1 -norm. Clearly, the log-term can be dropped in this regime. ■

6.1 Sampling numbers

Let us sharpen our considerations to the problem of optimal sampling recovery. We compare the number m of samples an algorithm is using to the resulting error and call the quantity

$$\varrho_m(S_{p,\theta}^r A([0,1]^d), L_q([0,1]^d)) := \inf_{\substack{X_m \subset [0,1]^d, |X_m|=m \\ \varphi: \mathbb{C}^m \rightarrow L_q([0,1]^d)}} \sup_{\|f\|_{S_{p,\theta}^r A([0,1]^d)} \leq 1} \|f - \varphi(f(X_m))\|_{L_q([0,1]^d)},$$

m -th sampling width. If, in addition, the mapping $\varphi : \mathbb{C}^n \rightarrow L_q$ is linear then we obtain the linear sampling widths

$$\varrho_m^{\text{lin}}(S_{p,\theta}^r A([0, 1]^d), L_q([0, 1]^d)) := \inf_{\substack{X_m \subset [0, 1]^d, |X_m|=m \\ \varphi: \mathbb{C}^m \rightarrow L_q([0, 1]^d) \\ \text{linear}}} \sup_{\|f\|_{S_{p,\theta}^r A([0, 1]^d)} \leq 1} \|f - \varphi(f(X_m))\|_{L_q([0, 1]^d)}.$$

Theorem 6.4. Let $1 \leq p < \infty$. Then

$$\varrho_m(S_{p,1}^{1/p} B([0, 1]^d), L_p([0, 1]^d)) \lesssim m^{-1/p} (\log^{d-1} m)^{\frac{1}{p}+1}$$

for all $m \in \mathbb{N}$.

Proof. The upper bound is due to Theorem 6.1 recognizing that the algorithm I_n in (6.1) samples f in $m \asymp 2^n n^{d-1}$ nodes. This can be trivially checked applying Lemma 6.1.(ii). For sake of completeness we refer to [3, Section 5.1] where further properties of this operator were studied. \blacksquare

Theorem 6.5. Let $1 \leq p < q < \infty$. Then

(i)

$$\varrho_m(S_{p,1}^{1/p} B([0, 1]^d), L_q([0, 1]^d)) \lesssim m^{-1/q} (\log^{d-1} m)^{2/q}$$

and

(ii)

$$\varrho_m(S_{1,\infty}^1 F([0, 1]^d), L_q([0, 1]^d)) \lesssim m^{-1/q} (\log^{d-1} m)^{2/q}$$

for all $m \in \mathbb{N}$.

Proof. The upper bound in (i) is due to Theorem 6.2 taking the number of sampling nodes into account, see Theorem 6.4. The upper bound in (ii) is due to Theorem 6.3. \blacksquare

6.2 Lower bounds

The linear width of a class \mathbf{F} in a normed space X has been introduced by Tikhomirov [27] more than 60 years ago. It is defined by

$$\lambda_m(\mathbf{F}, X) := \inf_{\substack{A: X \rightarrow X \\ \text{rank } A \leq m}} \sup_{f \in \mathbf{F}} \|f - A(f)\|_X.$$

A.S. Romanyuk [19, 20] proved for \mathbf{F} , the unit ball in $S_{p,1}^{1/p} B([0, 1]^d)$, that in case $1 \leq p \leq q \leq 2$ and $q > 1$

$$\begin{aligned} \lambda_m(S_{p,1}^{1/p} B([0, 1]^d), L_q([0, 1]^d)) &\geq d_m(S_{p,1}^{1/p} B([0, 1]^d), L_q([0, 1]^d)) \\ &\gtrsim (m^{-1} \log^{d-1} m)^{1/q}. \end{aligned}$$

From these bounds we obtain the following lower bounds for the (linear) sampling numbers.

Theorem 6.6. Let $1 \leq p \leq q < \infty$. Then we have

(i)

$$\varrho_n(S_{p,1}^{1/p} B([0, 1]^d), L_q([0, 1]^d)) \gtrsim n^{-1/q}.$$

(ii) If additionally $1 \leq p \leq q \leq 2$ then

$$\varrho_n^{\text{lin}}(S_{p,1}^{1/p} B([0, 1]^d), L_q([0, 1]^d)) \gtrsim (n^{-1} \log^{d-1} n)^{1/q}.$$

7 Outlook and discussion

In this paper we have shown that for the sampling recovery problem the compact embedding into L_∞ is not necessary. There are several relevant multivariate function spaces which fall under this scope, like limiting Besov and Triebel-Lizorkin spaces with smoothness $r = 1/p$ and further parameter conditions to ensure the embedding into the space of continuous functions. We were able to give upper bounds for the so-called sampling numbers in L_q with $q < \infty$ which are sharp in the polynomial main rate. What concerns the right order of the logarithm the situation is completely open. The results presented here are, as far as the authors know, the first results in this direction apart from a univariate result of Oswald [17].

Let us comment on the particular case of L_1 -smoothness spaces. Smoothness spaces built upon L_1 with smoothness $r = 1$ play an important role for numerical integration. This includes for instance the space $S_1^1 W([0, 1]^d)$ defined via weak derivatives. This space can not be described via Fourier analytical means and is therefore difficult to handle. See for instance [14]. However, these spaces fall into the scope of this paper since we also have $r = 1/p$ and a non-compact embedding into L_∞ . In [14] a Faber-Schauder characterization is shown, similar as above, namely

$$\sup_{j \in \mathbb{N}_{-1}^d} \sum_{k \in \mathbb{Z}^d} |d_{j,k}^2(f)| \lesssim \|f|S_1^1 W(\mathbb{R}^d)\|. \quad (7.1)$$

In particular, this would imply the following so-called sampling inequality

$$\sum_{k \in \mathbb{Z}^d} |f(k)| \lesssim \|f|S_1^1 W(\mathbb{R}^d)\|.$$

This extends the result in [21, Prop. 2] in several directions. On the one hand we consider the multivariate case and on the other hand the space $S_1^1 W(\mathbb{R}^d)$ is larger than $S_{1,1}^1 B(\mathbb{R}^d)$.

Having (7.1) at hand the following slightly sharper version of Theorem 6.3 is immediate.

Theorem 7.1. (i) It holds

$$\|f - I_n f\|_1 \lesssim 2^{-n} n^{d-1} \|f|S_1^1 W([0, 1]^d)\|.$$

(ii) For any $1 < q < \infty$ we have

$$\|f - I_n f\|_q \lesssim 2^{-n/q} n^{(d-1)/q} \|f|S_1^1 W([0, 1]^d)\|.$$

A proper Fourier analytical replacement of the spaces $S_1^1 W([0, 1]^d)$ are the spaces $S_{1,q}^1 F([0, 1]^d)$. However these spaces are not really comparable, especially when $q = \infty$. Using the methods in [8, Thm. 1.9] there is strong evidence for proving a version of (7.1) also for the spaces $S_{1,\infty}^1 F(\mathbb{R}^d)$. This would imply a version of Theorem 7.1. By well-known arguments a cubature formula with performance

$$\left| f - \sum_{i=1}^N \lambda_i f(x^i) \right| \lesssim N^{-1} \log N^{2(d-1)} \|f|S_{1,\infty}^1 F([0, 1]^d)\|$$

can be constructed by integrating the approximand. Note, that there were efforts in the literature to treat such limiting cases, see for instance [29, Cor. 6.5]. Suboptimal bounds were proven there. Note, that results for such limiting cases are related to the Kokhmsa-Hlawka inequality, saying that QMC-cubature in $S_1^1 W$ is related to the star discrepancy of the cubature nodes.

Acknowledgments Special thanks go to Winfried Sickel for stimulating discussions. The second named author is supported by the German Academic Exchange Service (DAAD, Grant 57588362) and by the Philipp Schwartz initiative of the Alexander von Humboldt Foundation.

References

- [1] F. Bartel, M. Schäfer, and T. Ullrich. Constructive subsampling of finite frames with applications in optimal function recovery. *arXiv e-prints*, pages 1–37, 2022.
- [2] H.-J. Bungartz and M. Griebel. Sparse grids. *Acta Numer.*, 13:147–269, 2004.
- [3] G. Byrenheid. *Sparse representation of multivariate functions based on discrete point evaluations*. Dissertation, Institut fuer Numerische Simulation, Universitaet Bonn, 2018.
- [4] D. Dũng. B-spline quasi-interpolant representations and sampling recovery of functions with mixed smoothness. *J. Complexity*, 27(6):541–567, 2011.
- [5] D. Dũng, V. Temlyakov, and T. Ullrich. *Hyperbolic cross approximation*. Advanced Courses in Mathematics. CRM Barcelona. Birkhäuser/Springer, Cham, 2018. Edited and with a foreword by Sergey Tikhonov.
- [6] M. Dolbeault, D. Krieg, and M. Ullrich. A sharp upper bound for sampling numbers in L_2 . *arXiv:2204.12621*, 2022.
- [7] G. Faber. Über stetige Funktionen. *Math. Ann.*, 66:81–94, 1909.
- [8] G. Garrigós, A. Seeger, and T. Ullrich. Haar frame characterizations of besov-sobolev spaces and optimal embeddings into their dyadic counterparts. *arXiv e-prints*, pages 1–44, 2022.
- [9] A. Hinrichs, L. Markhasin, J. Oettershagen, and T. Ullrich. Optimal quasi-Monte Carlo rules on order 2 digital nets for the numerical integration of multivariate periodic functions. *Numer. Math.*, 134(1):163–196, 2016.
- [10] H. Kempka, C. Schneider, and J. Vybiral. Path regularity of Brownian motion and Brownian sheet. *arXiv:2202.10114*, 2022.
- [11] D. Krieg and M. Ullrich. Function values are enough for L_2 -approximation. *Found. Comp. Math.*, 21(4):1141–1151, 2021.
- [12] D. Krieg and M. Ullrich. Function values are enough for L_2 -approximation: Part II. *J. Complexity*, to appear.
- [13] I. Limonova and V. Temlyakov. On sampling discretization in L_2 . *J. Math. Anal. Appl.*, 515(2):Paper No. 126457, 14, 2022.
- [14] K. Lüttgen and T. Ullrich. Sobolev spaces with dominating mixed smoothness for $p = 1$. *in preparation*, 2022.
- [15] S. Mayer and T. Ullrich. Entropy numbers of finite dimensional mixed-norm balls and function space embeddings with small mixed smoothness. *Constr. Approx.*, 53(2):249–279, 2021.

- [16] N. Nagel, M. Schäfer, and T. Ullrich. A new upper bound for sampling numbers. *Found. Comp. Math.*, 22:445–468, 2022.
- [17] P. Oswald. L^p -Approximation durch Reihen nach dem Haar-Orthogonalsystem und dem Faber-Schauder-System. *J. Approx. Theory*, 33(1):1–27, 1981.
- [18] K. Pozharska and T. Ullrich. A note on sampling recovery of multivariate functions in the uniform norm. *SIAM J. Numer. Anal.*, 60(3):1363–1384, 2022.
- [19] A. S. Romanyuk. Approximation of the Besov classes of periodic functions of several variables in the space L_q . *Ukr. Math. J.*, 43(10):1297–1306, 1991.
- [20] A. S. Romanyuk. Kolmogorov and trigonometric widths of the Besov classes $B_{p,\theta}^r$ of multivariate periodic functions. *Sb. Math.*, 197(1):69–93, 2006.
- [21] H.-J. Schmeisser and W. Sickel. Sampling theory and function spaces. In *Applied mathematics reviews, Vol. 1*, pages 205–284. World Sci. Publ., River Edge, NJ, 2000.
- [22] H.-J. Schmeisser and H. Triebel. *Topics in Fourier analysis and function spaces*. A Wiley-Interscience Publication. John Wiley & Sons, Ltd., Chichester, 1987.
- [23] V. Temlyakov. Approximation of functions with bounded mixed derivative. *Proc. Steklov Inst. Math.*, (1(178)):vi+121, 1989. A translation of Trudy Mat. Inst. Steklov 178 (1986), Translated by H. H. McFaden.
- [24] V. Temlyakov. On optimal recovery in L_2 . *J. Complexity*, 65:Paper No. 101545, 11, 2021.
- [25] V. Temlyakov and T. Ullrich. Bounds on Kolmogorov widths and sampling recovery for classes with small mixed smoothness. *J. Complexity*, 67:Paper No. 101575, 19, 2021.
- [26] V. N. Temlyakov and T. Ullrich. Approximation of functions with small mixed smoothness in the uniform norm. *J. Approx. Theory*, 277:Paper No. 105718, 23, 2022.
- [27] V. M. Tikhomirov. Diameters of sets in functional spaces and the theory of best approximations. *Russ. Math. Surv.*, 15(3):75–111, 1960.
- [28] H. Triebel. *Bases in function spaces, sampling, discrepancy, numerical integration*, volume 11 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2010.
- [29] M. Ullrich and T. Ullrich. The role of Frolov’s cubature formula for functions with bounded mixed derivative. *SIAM J. Numer. Anal.*, 54(2):969–993, 2016.
- [30] T. Ullrich. Function spaces with dominating mixed smoothness, characterization by differences. Technical report, Jenaer Schriften zur Math. und Inform., Math/Inf/05/06, 2006.
- [31] T. Ullrich. *Smolyak’s algorithm, sparse grid approximation and periodic function spaces with dominating mixed smoothness*. Friedrich-Schiller-Universität Jena, 2007. PhD thesis.
- [32] J. Vybiral. Function spaces with dominating mixed smoothness. *Dissertationes Math.*, 436:73, 2006.