

# Algebraic logic for the negation fragment of classical logic

Luciano J. González

*CONICET (Argentina) – Universidad Nacional de La Pampa. Facultad de Ciencias Exactas y Naturales.  
Santa Rosa, Argentina.*

---

## Abstract

The general aim of this article is to study the negation fragment of classical logic within the framework of contemporary (Abstract) Algebraic Logic. More precisely, we shall find the three classes of algebras that are canonically associated with a logic in Algebraic Logic, that is, we find the classes  $\text{Alg}^*$ ,  $\text{Alg}$  and the intrinsic variety of the negation fragment of classical logic. In order to achieve this, firstly we propose a Hilbert-style axiomatization for this fragment. Then, we characterize the reduced matrix models and the full generalized matrix models of this logic. Also, we classify the negation fragment in the Leibniz and Frege hierarchies.

*Keywords:* Classical logic; Classical negation; Algebraic logic; Reduced matrix models; Full models

---

## 1. Introduction

It is clear that the negation fragment of classical propositional logic (from now on CPL) is a very inexpressive logic from the syntactic point of view, we cannot express much with it. It only can be considered a proposition  $p$  and its negation  $\neg p$ , and no other compound sentential can be built up. However, we think that this fragment has several algebra-based semantics (in the different senses of this term) that are not so trivial. We intend to describe these algebra-base semantics of the negation fragment of CPL from the point of view of algebraic logic.

In the framework of (abstract) algebraic logic, there are essentially three classes of algebras associated with a propositional logic. These classes are obtained from different procedures, which intend to be a kind of generalization or abstraction of the Tarski-Lindenbaum method applied to classical (intuitionistic) propositional logic. The class  $\text{Alg}^*(\mathcal{S})$  is the class of algebras that is canonically associated with a logic  $\mathcal{S}$  according to the theory of logical

---

*Email address:* [lucianogonzalez@exactas.unlpam.edu.ar](mailto:lucianogonzalez@exactas.unlpam.edu.ar) (Luciano J. González)

matrices. More precisely,  $\text{Alg}^*(\mathcal{S})$  is the class of the algebraic reducts of the reduced matrix models of  $\mathcal{S}$ . The intrinsic variety of a logic  $\mathcal{S}$  (denoted by  $\mathbb{V}(\mathcal{S})$ ) is defined as the variety generated by a quotient algebra on the formula algebra. And the class  $\text{Alg}(\mathcal{S})$  is determined from the reduced generalized matrix models, that is,  $\text{Alg}(\mathcal{S})$  is the class of algebraic reducts of the reduced g-models of  $\mathcal{S}$ . We refer the reader to [1, 3, 4] for the specific definitions. In general, for any logic  $\mathcal{S}$ , we always have that  $\text{Alg}^*(\mathcal{S}) \subseteq \text{Alg}(\mathcal{S}) \subseteq \mathbb{V}(\mathcal{S})$ . In many cases, the three classes coincide, and in many others cases the inclusions are proper. From the point of view of algebraic logic, the class of algebras what is more representative for a logic  $\mathcal{S}$  or which can be considered as the *algebraic counterpart* of  $\mathcal{S}$  is the class  $\text{Alg}(\mathcal{S})$ . We refer the reader to [1, 3, 4] for a deeper explanation of the aims and goals of algebraic logic and the corresponding algebra-based semantics. We purpose to obtain the classes  $\text{Alg}^*$ , the intrinsic variety and the class  $\text{Alg}$  for the negation fragment of CPL.

The article is organized as follows. In Section 2, we introduce the very basics to start with the work. Throughout the paper, we will introduce the needed concepts and results of algebraic logic. We assume that the reader is familiar with the very basics of algebraic logic, for instance, with the notions of propositional logic (or sentential logic), logical filter, a theory of a logic, logical matrices, generalized matrices, etc. We shall give all those notions that might be less usual for the reader. We refer the reader to [1, 3] for further information on algebraic logic. Section 3 presents a Hilbert-style system for the negation fragment of CPL and shows a completeness theorem. In Section 4, we characterize the reduced matrix model and we describe the class  $\text{Alg}^*$  of the negation fragment. Section 5 is devoted to obtain the intrinsic variety of the negation fragment. In Section 6, we characterize the full g-models of the negation fragment. In order to obtain this, we describe the logical filters generated by a set on an arbitrary algebra and present several properties of these logical filters. Then, we find the class  $\text{Alg}$  of the negation fragment. Finally, in Section 7, we classify the negation fragment in the Leibniz and Frege hierarchies.

## 2. The $\{\neg\}$ -fragment of classical propositional logic (CPL)

Let  $\mathcal{L} = \{\neg\}$  be an algebraic language of type (1) and let  $\text{Fm}$  be the algebra of formulas over the language  $\mathcal{L}$  and generated by a countably infinite set  $\text{Var}$ . Unless otherwise stated, all the algebras considered in the paper are defined over the algebraic language  $\mathcal{L}$ . Let us denote by  $\mathcal{S}_N = \langle \text{Fm}, \vdash_N \rangle$  the  $\{\neg\}$ -fragment of CPL, where  $\vdash_N$  is the corresponding consequence relation. Let  $\mathbf{2}_\neg = \langle \{0, 1\}, \neg \rangle$  be the  $\{\neg\}$ -reduct of the two-element Boolean algebra  $\mathbf{2}_B$ . Then, it is clear that for all  $\Gamma \cup \{\varphi\} \subseteq \text{Fm}$ ,

$$\Gamma \vdash_N \varphi \iff \forall h \in \text{Hom}(\text{Fm}, \mathbf{2}_\neg)(h[\Gamma] \subseteq \{1\} \implies h\varphi = 1).$$

Let  $\mathbb{N}$  be the set of all positive integers and  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ . Let  $x \in \text{Var}$ . For each  $n \in \mathbb{N}_0$ , we define recursively  $\neg^n x$  as usual:

$$\begin{cases} \neg^0 x & = x \\ \neg^n x & = \neg \neg^{n-1} x \quad \forall n \geq 1. \end{cases}$$

**Lemma 2.1.** *For every  $\alpha \in \text{Fm}$ , there is  $x \in \text{Var}$  and  $n \in \mathbb{N}_0$  such that  $\alpha = \neg^n x$ .*

*Proof.* It is straightforward because  $\text{Fm}$  is the absolutely free algebra of type  $\{\neg\}$  over  $\text{Var}$ .  $\square$

### 3. Hilbert-style axiomatization for the $\{\neg\}$ -fragment of CPL

In [8, Theorem 1], it is claimed that every two-valued logic (a logic defined as usual by a two-valued matrix  $M = \langle A, \{1\} \rangle$ , that is,  $A = \{0, 1\}$  is a two-element algebra) has a finite Hilbert-style axiomatization. Hence, it follows that  $\mathcal{S}_N$  has a finite Hilbert-style axiomatization. In [8, pp. 322], it is mentioned that a Hilbert-style axiomatization for  $\mathcal{S}_N$  is easily established, but it isn't explicitly presented. Here let us present a Hilbert-style system for  $\mathcal{S}_N$  and prove a completeness theorem.

**Definition 3.1.** Let  $\mathcal{S}_\neg = \langle \text{Fm}, \vdash_\neg \rangle$  be the propositional logic defined, as usual, by the following Hilbert-style system:

$$(R1) \quad x, \neg x \vdash y$$

$$(R2) \quad x \vdash \neg \neg x$$

$$(R3) \quad \neg \neg x \vdash x.$$

Then, our goal is to show that the logics  $\mathcal{S}_N$  and  $\mathcal{S}_\neg$  coincide. To this end, we need some auxiliary results.

**Proposition 3.2.** *Let  $\Gamma \subseteq \text{Fm}$  be consistent. If  $\Gamma \vdash_\neg \alpha$ , then there is  $\gamma \in \Gamma$  such that  $\gamma \vdash_\neg \alpha$ .*

*Proof.* Suppose that  $\Gamma \vdash_\neg \alpha$ . We proceed by induction on the length of the proofs from  $\Gamma$ . That is, we prove that for all  $n \in \mathbb{N}$ , if  $\alpha_1, \dots, \alpha_n$  is a proof from  $\Gamma$ , then there is  $\gamma \in \Gamma$  such that  $\gamma \vdash_\neg \alpha_n$ .

If  $n = 1$ , then  $\alpha_1$  is a proof from  $\Gamma$ . Then  $\alpha_1 \in \Gamma$ . Hence, there is  $\gamma := \alpha_1 \in \Gamma$  such that  $\gamma \vdash_\neg \alpha_1$ . Now suppose that for each proof  $\alpha_1, \dots, \alpha_k$  from  $\Gamma$  of length  $k < n$ , there

is  $\gamma \in \Gamma$  such that  $\gamma \vdash_{\neg} \alpha_k$ . Let  $\alpha_1, \dots, \alpha_n$  be a proof from  $\Gamma$ . Then,  $\alpha_n \in \Gamma$  or there is  $i < n$  such that  $\alpha_n$  is derivable from the rules (R2) or (R3) and  $\alpha_i$  (notice that  $\alpha_n$  cannot be derivable from the rule (R1) because  $\Gamma$  is consistent). Thus, in any case,  $\alpha_i \vdash_{\neg} \alpha_n$ . By inductive hypothesis, there is  $\gamma \in \Gamma$  such that  $\gamma \vdash_{\neg} \alpha_i$ . Then  $\gamma \vdash_{\neg} \alpha_n$ .  $\square$

**Proposition 3.3.** *Let  $\alpha, \beta \in \text{Fm}$ . Then,  $\alpha \vdash_{\neg} \beta$  if and only if there is  $x \in \text{Var}$  and  $n, k \in \mathbb{N}_0$  such that  $\alpha = \neg^n x$ ,  $\beta = \neg^k x$  and  $n \equiv_2 k$ .*

*Proof.* ( $\Rightarrow$ ) Assume that  $\alpha \vdash_{\neg} \beta$ . We proceed by induction on the length of the proofs from  $\alpha$ . That is, we prove that for all  $m \in \mathbb{N}$ , if  $\alpha_1, \dots, \alpha_m$  is a proof from  $\alpha$ , then there is  $x \in \text{Var}$  and  $n, k \in \mathbb{N}_0$  such that  $\alpha = \neg^n x$ ,  $\alpha_m = \neg^k x$  and  $n \equiv_2 k$ . If  $m = 1$ , then  $\alpha = \alpha_1$ . From Lemma 2.1, we have  $\alpha = \neg^n x = \alpha_1$  for some  $x \in \text{Var}$  and  $n \in \mathbb{N}_0$ . Now suppose that it holds for all  $i < m$ . Let  $\alpha_1, \dots, \alpha_m$  be a proof from  $\alpha$ . Then,  $\alpha_m = \alpha$  or  $\alpha_m$  is derivable from  $\alpha_i$  with  $i < m$  by an application of (R2) or (R3) (it cannot be derivable by (R1) because  $\{\alpha\}$  is consistent). It is straightforward when  $\alpha_m = \alpha$ . Suppose that  $\alpha_m$  is derivable from  $\alpha_i$  with  $i < m$  by an application of (R2) or (R3). By inductive hypothesis, there is  $x \in \text{Var}$  and  $n, k \in \mathbb{N}_0$  such that  $\alpha = \neg^n x$ ,  $\alpha_i = \neg^k x$  and  $n \equiv_2 k$ . On the one hand, if  $\alpha_m$  is derivable from (R2), then  $\alpha_m = \neg\neg\alpha_i$ . Hence  $\alpha_m = \neg\neg\alpha_i = \neg^{k+2}x$  and  $n \equiv_2 k + 2$ . On the other hand, if  $\alpha_m$  is derivable from (R3), then  $\alpha_i = \neg\neg\alpha_m$ . Since  $\neg\neg\alpha_m = \alpha_i = \neg^k x$ , we have that  $k \geq 2$ . Then  $\alpha_m = \neg^{k-2}x$  and  $k - 2 \geq 0$ . Hence,  $\alpha = \neg^n x$ ,  $\alpha_m = \neg^{k-2}x$  and  $n \equiv_2 k - 2$ .

( $\Leftarrow$ ) Suppose that there are  $x \in \text{Var}$  and  $n, k \in \mathbb{N}_0$  such that  $\alpha = \neg^n x$ ,  $\beta = \neg^k x$  and  $n \equiv_2 k$ . Suppose that  $n \leq k$ . Thus  $0 \leq k - n = 2q$ , for some  $q \in \mathbb{N}_0$ . Then, we have  $\alpha = \neg^n x \vdash_{\neg} \neg^{n+2}x \vdash_{\neg} \dots \vdash_{\neg} \neg^{n+2q}x = \neg^k x = \beta$  (this can be proved by induction). If  $n > k$ , then  $n - 2q = k$ . Hence  $\alpha = \neg^n x \vdash_{\neg} \neg^{n-2}x \vdash_{\neg} \dots \vdash_{\neg} \neg^{n-2q}x = \beta$ .  $\square$

**Remark 3.4.** Notice that for all  $\alpha, \beta \in \text{Fm}$ ,  $\alpha \vdash_{\neg} \beta$  if and only if  $\beta \vdash_{\neg} \alpha$ . In other words,  $\alpha \vdash_{\neg} \beta$  if and only if  $\alpha \dashv\vdash_{\neg} \beta$ .

**Corollary 3.5.** *Let  $\Gamma \cup \{\alpha, \beta\} \subseteq \text{Fm}$ .*

1.  $\alpha \vdash_{\neg} \beta \implies \neg\alpha \not\vdash_{\neg} \beta$ .
2.  $\alpha \vdash_{\neg} \beta \iff \neg\beta \vdash_{\neg} \neg\alpha$ .
3.  $\Gamma, \alpha \vdash_{\neg} \beta$  and  $\Gamma, \neg\alpha \vdash_{\neg} \beta \implies \Gamma \vdash_{\neg} \beta$ .

*Proof.* (1) Suppose  $\alpha \vdash_{\neg} \beta$ . Thus, there are  $x \in \text{Var}$  and  $n, k \in \mathbb{N}_0$  such that  $\alpha = \neg^n x$ ,  $\beta = \neg^k x$  and  $n \equiv_2 k$ . Then  $\neg\alpha = \neg^{n+1}x$  and  $n + 1 \not\equiv_2 k$ . Hence  $\neg\alpha \not\vdash_{\neg} \beta$ .

(2) ( $\Rightarrow$ ) Suppose that  $\alpha \vdash_{\neg} \beta$ . Then, there are  $x \in \text{Var}$  and  $n, k \in \mathbb{N}_0$  such that  $\alpha = \neg^n x$ ,  $\beta = \neg^k x$  and  $n \equiv_2 k$ . Thus  $\neg\alpha = \neg^{n+1}x$ ,  $\neg\beta = \neg^{k+1}x$  and  $n+1 \equiv_2 k+1$ . Hence  $\neg\beta \vdash_{\neg} \neg\alpha$ .  
( $\Leftarrow$ ) It is straightforward by the above and by rules (R2) and (R3).

(3) Assume that  $\Gamma, \alpha \vdash_{\neg} \beta$  and  $\Gamma, \neg\alpha \vdash_{\neg} \beta$ . Suppose that  $\Gamma \not\vdash_{\neg} \beta$ . Then, by Proposition 3.2, it follows that  $\alpha \vdash_{\neg} \beta$  and  $\neg\alpha \vdash_{\neg} \beta$ . This is a contradiction by property (1). Hence  $\Gamma \vdash_{\neg} \beta$ .  $\square$

**Proposition 3.6.** *Let  $\alpha \in \text{Fm}$  and let  $\Delta$  be a maximal theory relative to  $\alpha$  of the logic  $\mathcal{S}_{\neg}$  (that is,  $\Delta$  is a maximal theory among all the consistent theories not containing  $\alpha$ ). Then, for all  $\beta \in \text{Fm}$ ,  $\beta \in \Delta$  iff  $\neg\beta \notin \Delta$ .*

*Proof.* Let  $\beta \in \text{Fm}$ .

( $\Rightarrow$ ) Assume  $\beta \in \Delta$ . Since  $\Delta$  is consistent ( $\alpha \notin \Delta$ ), it follows by rule (R1) that  $\neg\beta \notin \Delta$ .

( $\Leftarrow$ ) Suppose that  $\neg\beta \notin \Delta$ . We suppose, towards a contradiction, that  $\beta \notin \Delta$ . Let  $\Gamma_{\beta}$  and  $\Gamma_{\neg\beta}$  be the theories generated by  $\Delta \cup \{\beta\}$  and  $\Delta \cup \{\neg\beta\}$ , respectively. Since  $\Delta \subset \Gamma_{\beta}$  and  $\Delta \subset \Gamma_{\neg\beta}$ , it follows by the maximality of  $\Delta$  that  $\alpha \in \Gamma_{\beta} \cap \Gamma_{\neg\beta}$ . Then

$$\Delta, \beta \vdash \alpha \quad \text{and} \quad \Delta, \neg\beta \vdash \alpha.$$

Hence, by (3) of Corollary 3.5, we obtain that  $\Delta \vdash \alpha$ , which is a contradiction. Therefore,  $\beta \in \Delta$ .  $\square$

Now we are ready to show that the rules (R1)–(R3) are an axiomatization for  $\mathcal{S}_N$ .

**Theorem 3.7.** *The Hilbert calculus formed by the rules (R1), (R2) and (R3) is an axiomatization for the  $\{\neg\}$ -fragment of CPL.*

*Proof.* We need to show that  $\vdash_N = \vdash_{\neg}$ . Recall that  $\vdash_N$  is defined by the matrix  $\langle \mathbf{2}_{\neg}, \{1\} \rangle$ .

( $\vdash_{\neg} \subseteq \vdash_N$ ) (Soundness) It is a routine argument.

( $\vdash_N \subseteq \vdash_{\neg}$ ) Suppose that  $\Gamma \not\vdash_{\neg} \alpha$ . By [1, Lem. 1.43], there is a theory  $\Delta$  such that  $\Gamma \subseteq \Delta$ ,  $\alpha \notin \Delta$ , and  $\Delta$  is maximal relative to  $\alpha$ . We define  $v: \text{Fm} \rightarrow \mathbf{2}_{\neg}$  as follows: for all  $\varphi \in \text{Fm}$ ,  $v(\varphi) = 1 \iff \varphi \in \Delta$ . By Proposition 3.6, we obtain that  $v$  is a homomorphism such that  $v[\Gamma] \subseteq \{1\}$  and  $v(\alpha) = 0$ . Hence  $\Gamma \not\vdash_N \alpha$ .  $\square$

#### 4. Reduced models and the class $\text{Alg}^*(\mathcal{S}_N)$

In this section, we characterize the reduced matrix models of the logic  $\mathcal{S}_N$  and we describe the class  $\text{Alg}^*(\mathcal{S}_N)$ . We recall some needed notions.

Recall that a *logical matrix* (*matrix* for short) is a pair  $\langle A, F \rangle$  where  $A$  is an algebra and  $F \subseteq A$ . The *Leibniz congruence*, denoted by  $\Omega^A F$ , of a matrix  $\langle A, F \rangle$  can be defined as follows (see [1, Theo. 4.23]): for all  $a, b \in A$ ,

$$(a, b) \in \Omega^A F \iff \text{for all } \delta(x, \vec{z}) \in \text{Fm} \text{ and all } \vec{c} \in \vec{A}, \quad (4.1)$$

$$[\delta^A(a, \vec{c}) \in F \iff \delta^A(b, \vec{c}) \in F].$$

A matrix  $\langle A, F \rangle$  is said to be *reduced* when  $\Omega^A F = \text{Id}_A$ . Recall that a matrix  $\langle A, F \rangle$  is a *model* of a logic  $\mathcal{S}$  when for all  $\Gamma \cup \{\alpha\} \subseteq \text{Fm}$ ,

$$\Gamma \vdash_{\mathcal{S}} \alpha \implies \forall h \in \text{Hom}(\text{Fm}, A)(h[\Gamma] \subseteq F \implies h(\alpha) \in F). \quad (4.2)$$

Then, the class  $\text{Alg}^*(\mathcal{S})$  is defined as follows;

$$\text{Alg}^*(\mathcal{S}) = \{A : \text{there is some } F \subseteq A \text{ such that } \langle A, F \rangle \text{ is a reduced model of } \mathcal{S}\}.$$

Recall also that a subset  $F$  of an algebra  $A$  is called an  $\mathcal{S}$ -*filter* of  $A$  if condition (4.2) is satisfied. That is, a subset  $F \subseteq A$  is an  $\mathcal{S}$ -filter of  $A$  if and only if the matrix  $\langle A, F \rangle$  is a model of  $\mathcal{S}$ .

By (4.1) and taking into account that every formula  $\alpha \in \text{Fm}$  is of the form  $\alpha = \neg^n x$  for some  $x \in \text{Var}$  and  $n \in \mathbb{N}_0$ , we obtain the following characterization of the Leibniz congruences.

**Proposition 4.1.** *For every algebra  $A$  and every  $F \subseteq A$ ,*

$$(a, b) \in \Omega^A F \iff \forall n \in \mathbb{N}_0 (\neg^n a \in F \iff \neg^n b \in F)$$

for all  $a, b \in A$ .

**Proposition 4.2.** *Let  $A$  be an algebra and  $F \subseteq A$ . Then,  $F$  is an  $\mathcal{S}_N$ -filter if and only if the following conditions hold:*

- (1)  $a, \neg a \in F \implies F = A$ ;
- (2)  $a \in F \iff \neg\neg a \in F$ .

*Proof.* It is straightforward by rules (R1)–(R3). □

We are ready to characterize the reduced models of  $\mathcal{S}_N$ .

**Theorem 4.3.** *Let  $\langle A, F \rangle$  be a matrix. Then,  $\langle A, F \rangle$  is a reduced model of  $\mathcal{S}_N$  if and only if the following conditions hold:*

- (1)  $A \models x \approx \neg\neg x$ ,
- (2)  $F = \{a_0\}$  for some  $a_0 \in A$  such that  $a_0 \neq \neg a_0$ , and
- (3)  $2 \leq |A| \leq 3$ .

*Proof.* ( $\Rightarrow$ ) Assume that  $\langle A, F \rangle$  is a reduced model of  $\mathcal{S}_N$ . Thus  $F$  is an  $\mathcal{S}_N$ -filter and  $\Omega^A F = \text{Id}_A$ .

(1) Let  $a \in A$ . From Proposition 4.2, we obtain

$$\forall n \in \mathbb{N}_0 (\neg^n a \in F \iff \neg^n(\neg\neg a) \in F).$$

Then,  $(a, \neg\neg a) \in \Omega^A F$ . Hence  $a = \neg\neg a$ .

(2) We are assumed that the algebra  $A$  is not trivial, that is,  $|A| \geq 2$ . Then,  $F \neq \emptyset$  (otherwise  $\Omega^A F = A \times A \neq \Delta_A$ ). Let  $a, b \in F$ . By (1), we have that  $a = \neg^{2k} a$  and  $b = \neg^{2k} b$  for all  $k \in \mathbb{N}_0$ . Thus,  $\neg^{2k} a, \neg^{2k} b \in F$ , for all  $k \in \mathbb{N}_0$ . On the other hand, since  $a, b \in F$  and  $F$  is proper (otherwise  $\langle A, F \rangle$  is not reduced), it follows by Proposition 4.2 that  $\neg a, \neg b \notin F$ . Then,  $\neg^{2k+1} a, \neg^{2k+1} b \notin F$ , for all  $k \in \mathbb{N}_0$ . Hence, we have that  $\forall n \in \mathbb{N}_0 (\neg^n a \in F \iff \neg^n b \in F)$ . Thus,  $(a, b) \in \Omega^A F$ . Then,  $a = b$ . Therefore,  $F = \{a_0\}$  for some  $a_0 \in A$ . Moreover  $a_0 \neq \neg a_0$ . Otherwise,  $a_0 = \neg a_0 \in F$ , and by Proposition 4.2 we have  $F = A$ , which is a contradiction because  $A$  is not trivial.

(3) Since  $A$  is not trivial, we have  $|A| \geq 2$ . By (2), we have that  $F = \{a_0\}$  and  $a_0 \neq \neg a_0$ . Let  $a, b \in A$  be such that  $a, b \notin \{a_0, \neg a_0\}$ . Then,  $a, b, \neg a, \neg b \notin F = \{a_0\}$ . Thus, it holds that  $a \in F \iff b \in F$  and  $\neg a \in F \iff \neg b \in F$ . Now by (1), it follows that  $\forall n \in \mathbb{N}_0 (\neg^n a \in F \iff \neg^n b \in F)$ . Then  $(a, b) \in \Omega^A F$ . Hence,  $a = b$ . Therefore,  $|A| \leq 3$ .

( $\Leftarrow$ ) Let  $\langle A, F \rangle$  be a matrix such that satisfies (1)–(3). By (1) and (2), it follows that  $F = \{a_0\}$  is an  $\mathcal{S}_N$ -filter, that is,  $\langle A, F \rangle$  is a model of  $\mathcal{S}_N$ . Let us see that  $\langle A, F \rangle$  is reduced. Let  $a, b \in A$  and assume that  $(a, b) \in \Omega_A F$ . Thus, we have  $a \in F \iff b \in F$  and  $\neg a \in F \iff \neg b \in F$ . If  $a \in F$  or  $\neg a \in F$ , then  $a = b$ . Similarly, if  $b \in F$  or  $\neg b \in F$ , then  $a = b$ . Now suppose that  $a, b, \neg a, \neg b \notin F = \{a_0\}$ . Thus  $a, b \notin \{a_0, \neg a_0\}$ . Since  $|A| \leq 3$ , we obtain that  $a = b$ . Hence  $\Omega^A F = \Delta_A$ . Therefore,  $\langle A, F \rangle$  is a reduced model of  $\mathcal{S}_N$ .  $\square$

Therefore, we have that all reduced matrix models  $\langle A, F \rangle$  are of the form, up to isomorphism:

1.  $A_1 = \{a\}$  and  $F = \emptyset$ ;
2.  $A_2 = \{a, b\}$  such that  $\neg a = b$  and  $\neg b = a$ , with  $F = \{a\}$ ; or

3.  $A_3 = \{a, b, c\}$  such that  $\neg a = b$ ,  $\neg b = a$  and  $\neg c = c$ , with  $F = \{a\}$ .

**Corollary 4.4.**  $\text{Alg}^*(\mathcal{S}_N) = \mathbb{I}(\{A_1, A_2, A_3\})$ .

## 5. The intrinsic variety of $\mathcal{S}_N$

If one insists on having a variety associated with a logic  $\mathcal{S}$ , the *intrinsic variety* of  $\mathcal{S}$  might be the adequate choice. The significance of the intrinsic variety as an algebra-based semantics for a logic is in general weak, because no general theory asserts that the algebraic counterpart of a logic should be a variety. In order to define the intrinsic variety and for what follows in the article, we recall some needed concepts.

Recall that a *generalized matrix* (*g-matrix*) is a pair  $\langle A, \mathcal{C} \rangle$  where  $A$  is an algebra and  $\mathcal{C}$  is a closure system on  $A$ . By the correspondence between closure systems and closure operators, we can also consider a g-matrix as a pair  $\langle A, C \rangle$  where  $C$  is a closure operator on  $A$ .

The *Tarski congruence* of a g-matrix  $\langle A, C \rangle$ , denoted by  $\tilde{\Omega}^A C$ , is defined as the largest congruence  $\theta$  on  $A$  satisfying the following

$$(a, b) \in \theta \implies C(a) = C(b).$$

Given a logic  $\mathcal{S}$ , let us denote by  $\mathcal{T}h(\mathcal{S})$  the closure system of all theories of  $\mathcal{S}$ . Consider the Tarski congruence  $\tilde{\Omega}^{\text{Fm}} \mathcal{T}h(\mathcal{S})$  of the g-matrix  $\langle \text{Fm}, \mathcal{T}h(\mathcal{S}) \rangle$ . The *intrinsic variety* of  $\mathcal{S}$ , denoted by  $\mathbb{V}(\mathcal{S})$ , is defined as the variety generated by the quotient algebra  $\text{Fm}/\tilde{\Omega}^{\text{Fm}} \mathcal{T}h(\mathcal{S})$ . That is,  $\mathbb{V}(\mathcal{S}) = \mathbb{V}(\text{Fm}/\tilde{\Omega}^{\text{Fm}} \mathcal{T}h(\mathcal{S}))$ . Now, since the interderivability relation  $\dashv\vdash_N$  is a congruence on  $\text{Fm}$  (that is, the logic  $\mathcal{S}_N$  is selfextensional [1]), we have that

$$\mathbb{V}(\mathcal{S}_N) \models \alpha \approx \beta \iff \text{Fm}/\tilde{\Omega}^{\text{Fm}} \mathcal{T}h(\mathcal{S}) \models \alpha \approx \beta \iff (\alpha, \beta) \in \tilde{\Omega}^{\text{Fm}} \mathcal{T}h(\mathcal{S}) \iff \alpha \dashv\vdash_N \beta.$$

**Theorem 5.1.**  $\mathbb{V}(\mathcal{S}_N) = \{A : A \models x \approx \neg\neg x\}$ .

*Proof.* Let  $V := \{A : A \models x \approx \neg\neg x\}$ . Notice that if  $\alpha = \neg^n x$ , then for all  $h \in \text{Hom}(\text{Fm}, A)$  with  $A \in V$ , it follows that

$$\begin{cases} h\alpha = h(\neg^n x) = hx & \text{if } n \text{ is even} \\ h\alpha = h(\neg^n x) = \neg hx & \text{if } n \text{ is odd} \end{cases} \quad (5.1)$$

Let  $\alpha, \beta \in \text{Fm}$ . Let us prove that

$$V \models \alpha \approx \beta \iff \exists x \in \text{Var}, \exists n, k \in \mathbb{N}_0 \text{ such that } \alpha = \neg^n x, \beta = \neg^k x \text{ and } n \equiv_2 k.$$



Suppose that  $V \models \alpha \approx \beta$ . By Lemma 2.1, there are  $x, y \in \text{Var}$  and  $n, k \in \mathbb{N}_0$  such that  $\alpha = \neg^n x$  and  $\beta = \neg^k y$ . If  $x \neq y$ , then taking  $A = \{a, b\}$  with  $\neg a = a$  and  $\neg b = b$ , we obtain that  $A \in V$  and  $A \not\models \alpha \approx \beta$ . A contradiction. Hence  $x = y$ . Thus  $\alpha = \neg^n x$  and  $\beta = \neg^k x$ . Suppose that  $n \not\equiv_2 k$ . Thus  $n$  is even and  $k$  is odd (or vice versa). Taking  $A_2 = \{a, b\}$  as in page 7 and  $hx = a$ , we obtain by (5.1) that

$$h\alpha = h(\neg^n x) = hx = a \quad \text{and} \quad h\beta = h(\neg^k x) = \neg hx = \neg a = b.$$

Hence  $A \not\models \alpha \approx \beta$ , which is a contradiction. Then  $n \equiv_2 k$ .

Now assume that  $\alpha = \neg^n x$ ,  $\beta = \neg^k x$  and  $n \equiv_2 k$ , for some  $x \in \text{Var}$  and  $n, k \in \mathbb{N}_0$ . Let  $A \in V$  and  $h \in \text{Hom}(\text{Fm}, A)$ . We have that  $n$  and  $k$  are both even or odd. In any case, by (5.1) we obtain that  $h\alpha = h\beta$ . Hence  $V \models \alpha \approx \beta$ .

Hence, for all  $\alpha, \beta \in \text{Fm}$ , we have

$$\begin{aligned} \mathbb{V}(\mathcal{S}_N) \models \alpha \approx \beta &\iff \alpha \dashv\vdash_N \beta \\ &\iff \exists x \in \text{Var}, \exists n, k \in \mathbb{N}_0 \text{ s.t. } \alpha = \neg^n x, \beta = \neg^k x \text{ and } n \equiv_2 k \\ &\iff V \models \alpha \approx \beta. \end{aligned}$$

Therefore,  $\mathbb{V}(\mathcal{S}_N) = \{A : A \models x \approx \neg\neg x\}$ . □

## 6. Full g-models for $\mathcal{S}_N$ and the class $\text{Alg}(\mathcal{S}_N)$

In this section, we obtain the class  $\text{Alg}(\mathcal{S}_N)$ , which is considered in the framework of algebraic logic as *the* algebraic counterpart of  $\mathcal{S}_N$ . To this end, we need first to obtain a characterization of the full g-models of  $\mathcal{S}_N$ . The class of full g-models of a logic  $\mathcal{S}$  is a particular class of g-models of  $\mathcal{S}$  that behave particularly well, in the sense that some interesting metalogical properties of a propositional logic are precisely those shared by its full g-models. Then, the class  $\text{Alg}(\mathcal{S})$  is obtained as the algebraic reducts of the reduced full g-models of  $\mathcal{S}$ . It is claimed (see [3, pp. 3]) that the notion of full g-model is the “right” notion of model of a propositional logic and that the class  $\text{Alg}(\mathcal{S})$  is the “right” class of algebras to be canonically associated with a propositional logic. We refer the reader to [1, 3, 4] for a more detailed argumentation about the full g-models of a logic  $\mathcal{S}$  and the class  $\text{Alg}(\mathcal{S})$ .

We start presenting some needed notions. A g-matrix  $\langle A, C \rangle$  is said to be a *g-model* of a logic  $\mathcal{S}$  when for all  $\Gamma \cup \{\alpha\} \subseteq \text{Fm}$ ,

$$\Gamma \vdash_{\mathcal{S}} \alpha \implies \text{for all } h \in \text{Hom}(\text{Fm}, A), h\alpha \in C(h\Gamma).$$

Recall from the previous section the notion of Tarski congruence of a g-matrix. A g-matrix  $\langle A, \mathcal{C} \rangle$  is said to be *reduced* if  $\tilde{\Omega}^A \mathcal{C} = \text{Id}_A$ . Then, the class  $\text{Alg}(\mathcal{S})$  is defined as follows:

$\text{Alg}(\mathcal{S}) = \{A : \text{there is a closure system } \mathcal{C} \text{ on } A \text{ such that } \langle A, \mathcal{C} \rangle \text{ is a reduced g-model of } \mathcal{S}\}.$

An  $h \in \text{Hom}(A_1, A_2)$  is called a *strict homomorphism* from the g-matrix  $\langle A_1, C_1 \rangle$  to the g-matrix  $\langle A_2, C_2 \rangle$  when

$$a \in C_1(X) \iff h(a) \in C_2(h(X)) \quad \text{for all } X \cup \{a\} \subseteq A_1.$$

Let  $A$  be an algebra. We denote by  $\mathcal{F}i_{\mathcal{S}}(A)$  the closure system of all  $\mathcal{S}$ -filters of the algebra  $A$  of a logic  $\mathcal{S}$ . Given a subset  $X \subseteq A$ , let us denote by  $\text{Fig}_{\mathcal{S}}^A(X)$  the  $\mathcal{S}$ -filter of  $A$  generated by  $X$ , that is,  $\text{Fig}_{\mathcal{S}}^A(X)$  is the least  $\mathcal{S}$ -filter of  $A$  containing  $X$ .

A g-matrix  $\langle A, \mathcal{C} \rangle$  is said to be a *full g-model* of a logic  $\mathcal{S}$  if there is an algebra  $A_1$  and a surjective strict homomorphism  $h: \langle A, \mathcal{C} \rangle \rightarrow \langle A_1, \mathcal{F}i_{\mathcal{S}}(A_1) \rangle$ . The above g-matrix  $\langle A_1, \mathcal{F}i_{\mathcal{S}}(A_1) \rangle$  can be chosen in a particular, significant way:

**Proposition 6.1** ([1, Prop. 5.85]). *A g-matrix  $\langle A, \mathcal{C} \rangle$  is a full g-model of a logic  $\mathcal{S}$  if and only if*

$$\{F/\tilde{\Omega}^A \mathcal{C} : F \in \mathcal{C}\} = \mathcal{F}i_{\mathcal{S}}\left(A/\tilde{\Omega}^A \mathcal{C}\right).$$

**Proposition 6.2** ([1, Corollary 5.88(3)]). *Given a logic  $\mathcal{S}$ ,*

$$\text{Alg}(\mathcal{S}) = \{A : \text{there is a closure system } \mathcal{C} \text{ on } A \text{ such that} \\ \langle A, \mathcal{C} \rangle \text{ is a reduced full g-model of } \mathcal{S}\}.$$

In order to characterize the full g-models of  $\mathcal{S}_N$ , it will be useful to obtain a characterization of the  $\mathcal{S}_N$ -filters generated. We start with some basic properties.

**Proposition 6.3.** *Let  $A$  be an algebra. Then:*

$$(B1) \text{ Fig}_{\mathcal{S}_N}^A(\emptyset) = \emptyset.$$

$$(B2) \text{ Fig}_{\mathcal{S}_N}^A(a, \neg a) = A, \text{ for all } a \in A.$$

$$(B3) \text{ Fig}_{\mathcal{S}_N}^A(a) = \text{Fig}_{\mathcal{S}_N}^A(\neg\neg a).$$

*Proof.* (1) is a consequence from the fact that the logic  $\mathcal{S}_N$  have no theorems, and (2) and (3) are straightforward by rules (R1)–(R3).  $\square$

Now let us to obtain a characterization of  $\text{Fig}_{\mathcal{S}_N}^A(X)$ . Let  $\mathcal{E} = \{n \in \mathbb{N}_0 : n \text{ is even}\}$  and  $\mathcal{O} = \{n \in \mathbb{N}_0 : n \text{ is odd}\}$ . The next is a key proposition.

**Proposition 6.4.** *Let  $A$  be an algebra. Let  $B \subseteq A$  be such that the following condition holds:*

$$\forall b, b' \in B, \forall (s, t) \in \mathcal{E} \times \mathcal{O} (\neg^s b \neq \neg^t b'). \quad (6.1)$$

Then,

$$\text{Fig}_{\mathcal{S}_N}^A(B) = \{a \in A : \exists b \in B, \exists s, t \in \mathcal{E} (\neg^s a = \neg^t b)\} \quad \text{and} \quad \text{Fig}_{\mathcal{S}_N}^A(B) \neq A.$$

*Proof.* Let

$$F = \{a \in A : \exists b \in B, \exists s, t \in \mathcal{E} (\neg^s a = \neg^t b)\}.$$

We prove in several steps that  $\text{Fig}_{\mathcal{S}_N}^A(B) = F$ .

- $F \neq A$ . If  $B = \emptyset$ , then  $F = \emptyset \neq A$ . Suppose that  $B \neq \emptyset$ , and let  $b \in B$ . We show that  $\neg b \notin F$ . Suppose that  $\neg b \in F$ . Thus, there is  $b' \in B$  and  $s, t \in \mathcal{E}$  such that  $\neg^s(\neg b) = \neg^t b'$ . Then  $\neg^{s+1} b = \neg^t b'$  with  $(t, s+1) \in \mathcal{E} \times \mathcal{O}$ , which is a contradiction by (6.1). Hence  $\neg b \notin F$ . Therefore,  $F \neq A$ .

- We show that  $F$  is an  $\mathcal{S}_N$ -filter. We need to verify conditions (1) and (2) of Proposition 4.2. (1) Since  $F \neq A$ , we need to show that for all  $a \in A$ ,  $a \notin F$  or  $\neg a \notin F$ . Let  $a \in A$ . Suppose that  $a, \neg a \in F$ . Thus, there are  $b, b' \in B$  and  $s, t, s', t' \in \mathcal{E}$  such that  $\neg^s a = \neg^t b$  and  $\neg^{s'}(\neg a) = \neg^{t'} b'$ . Then  $\neg^{s+s'+1} a = \neg^{t+t'+1} b$  and  $\neg^{s+s'+1} a = \neg^{s+t'} b'$ . Hence  $\neg^{t+t'+1} b = \neg^{s+t'} b'$  with  $b, b' \in B$  and  $(s+t', t+t'+1) \in \mathcal{E} \times \mathcal{O}$ , which is a contradiction by (6.1). Thus  $a \notin F$  or  $\neg a \notin F$ . (2) Let  $a \in A$ . If  $a \in F$ , then there is  $b \in B$  and  $s, t \in \mathcal{E}$  such that  $\neg^s a = \neg^t b$ . Hence  $\neg^s(\neg a) = \neg^{t+2} b$ . By definition of  $F$ , we obtain that  $\neg a \in F$ . Conversely, suppose that  $\neg a \in F$ . Thus, there is  $b \in B$  and  $s, t \in \mathcal{E}$  such that  $\neg^s(\neg a) = \neg^t b$ . Hence  $\neg^{s+2} a = \neg^t b$  and  $s+2, t \in \mathcal{E}$ . Then  $a \in F$ . Therefore, it follows by Proposition 4.2 that  $F$  is an  $\mathcal{S}_N$ -filter.

- $B \subseteq F$ . It is obvious by definition of  $F$ .

- $F$  is the least  $\mathcal{S}_N$ -filter of  $A$  containing  $B$ . Let  $G$  be an  $\mathcal{S}_N$ -filter of  $A$  such that  $B \subseteq G$ . Let  $a \in F$ . Thus, there is  $b \in B$  and  $s, t \in \mathcal{E}$  such that  $\neg^s a = \neg^t b$ . By Proposition 3.3, we have that  $x \vdash_N \neg^t x$ . Since  $b \in B \subseteq G$  and  $G$  is an  $\mathcal{S}_N$ -filter, it follows that  $\neg^t b \in G$ . Thus  $\neg^s a \in G$ . Since  $\neg^s x \vdash_N x$ , it follows that  $a \in G$ . Hence  $F \subseteq G$ . Therefore, we have proved that  $\text{Fig}_{\mathcal{S}_N}^A(B) = F$ .  $\square$

**Proposition 6.5.** *Let  $A$  be an algebra and  $B \subseteq A$ . The following conditions are equivalent.*

(1) *There exist  $b, b' \in B$  and  $(s, t) \in \mathcal{E} \times \mathcal{O}$  such that  $\neg^s b = \neg^t b'$ .*

(2)  $\text{Fig}_{\mathcal{S}_N}^A(B) = A$ .

*Proof.* (1)  $\Rightarrow$  (2) Let  $b, b' \in B$  and  $(s, t) \in \mathcal{E} \times \mathcal{O}$  such that  $\neg^s b = \neg^t b'$ . We have that  $b, b' \in \text{Fig}_{\mathcal{S}_N}^A(B)$ . Since  $s$  is even, it follows that  $x \vdash_N \neg^s x$ . Then  $\neg^s b \in \text{Fig}_{\mathcal{S}_N}^A(B)$ . Thus  $\neg^t b' \in \text{Fig}_{\mathcal{S}_N}^A(B)$ . Since  $t$  is odd, it follows by Proposition 3.3 that  $\neg^t x \vdash_N \neg x$ . Then  $\neg b' \in \text{Fig}_{\mathcal{S}_N}^A(B)$ . That is,  $b', \neg b' \in \text{Fig}_{\mathcal{S}_N}^A(B)$ . Hence  $\text{Fig}_{\mathcal{S}_N}^A(B) = A$ .

(2)  $\Rightarrow$  (1) It follows by Proposition 6.4.  $\square$

From Propositions 6.4 and 6.5, we have that for every algebra  $A$  and every  $B \subseteq A$ ,

$$\text{Fig}_{\mathcal{S}_N}^A(B) = A \quad \text{or} \quad \text{Fig}_{\mathcal{S}_N}^A(B) = \{a \in A : \exists b \in B, \exists s, t \in \mathcal{E} (\neg^s a = \neg^t b)\}.$$

**Proposition 6.6.** *Let  $A$  be an algebra and  $b \in A$  such that*

$$\forall (s, t) \in \mathcal{E} \times \mathcal{O} (\neg^s b \neq \neg^t b). \quad (6.2)$$

*Then,*

$$(1) \text{Fig}_{\mathcal{S}_N}^A(b) = \{a \in A : \exists s, t \in \mathcal{E} (\neg^s a = \neg^t b)\};$$

$$(B4) a \in \text{Fig}_{\mathcal{S}_N}^A(b) \implies b \in \text{Fig}_{\mathcal{S}_N}^A(a);$$

$$(B5) a \in \text{Fig}_{\mathcal{S}_N}^A(b) \implies \neg b \in \text{Fig}_{\mathcal{S}_N}^A(\neg a).$$

*Proof.* (1) is an immediate consequence of Proposition 6.4. (2) and (3) follow by (1) and from Proposition 3.3.  $\square$

Let  $A$  be an algebra. Notice from Proposition 6.5 that an element  $b \in A$  satisfies condition (6.2) if and only if  $\text{Fig}_{\mathcal{S}_N}^A(b) \neq A$ . Moreover if  $B \subseteq A$  satisfying condition (6.1), then every element  $b \in B$  satisfies condition (6.2). Thus, the next proposition is a consequence of Proposition 6.4 and by (1) of Proposition 6.6.

**Proposition 6.7.** *Let  $A$  be an algebra and  $B \subseteq A$ . Then,*

$$(B6) \text{ if } a \in \text{Fig}_{\mathcal{S}_N}^A(B) \text{ and } \text{Fig}_{\mathcal{S}_N}^A(B) \neq A, \text{ then there is } b \in B \text{ such that } a \in \text{Fig}_{\mathcal{S}_N}^A(b).$$

**Proposition 6.8.** *Let  $\langle A, C \rangle$  be a  $g$ -matrix. Then*

$$(a, b) \in \tilde{\Omega}^A C \iff \forall n \in \mathbb{N}_0 (C(\neg^n a) = C(\neg^n b)).$$

*Proof.* It follows from the fact that  $\tilde{\Omega}^A C = \bigcap_{F \in \mathcal{C}} \Omega^A F$  (see [1, Lem. 5.31]) and by Proposition 4.1.  $\square$

**Proposition 6.9.** *Let  $\langle A, C \rangle$  be a g-matrix such that for each  $a \in A$ ,  $C(a) = C(\neg a)$ . Then, for all  $a, b \in A$ ,*

$$(a, b) \in \tilde{\Omega}^A C \iff C(a) = C(b) \text{ and } C(\neg a) = C(\neg b).$$

*Proof.* The implication  $\Rightarrow$  follows by the previous proposition.

( $\Leftarrow$ ) Let  $a, b \in A$  be such that  $C(a) = C(b)$  and  $C(\neg a) = C(\neg b)$ . We prove that  $\forall n \in \mathbb{N}_0 (C(\neg^n a) = C(\neg^n b))$  by induction on  $n$ . It is clear that holds for  $n = 0$  and  $n = 1$ . Let  $n \geq 2$  and suppose that for all  $k < n$ ,  $C(\neg^k a) = C(\neg^k b)$ . Since  $n \geq 2$ , it follows that

$$C(\neg^n a) = C(\neg^{n-2} a) \stackrel{I.H.}{=} C(\neg^{n-2} b) = C(\neg^n b).$$

Hence,  $\forall n \in \mathbb{N}_0 (C(\neg^n a) = C(\neg^n b))$ . Therefore  $(a, b) \in \tilde{\Omega}^A C$ .  $\square$

Now we are ready to present a characterization for the full g-models of the negation fragment  $\mathcal{S}_N$ .

**Theorem 6.10.** *Let  $\langle A, C \rangle$  be a g-matrix. Then,  $\langle A, C \rangle$  is a full g-model of the logic  $\mathcal{S}_N$  if and only if it satisfies:*

(F1)  $C(\emptyset) = \emptyset$ .

(F2)  $C(a, \neg a) = A$ , for all  $a \in A$ .

(F3)  $C(a) = C(\neg \neg a)$ , for all  $a \in A$ .

(F4) For all  $a, b \in A$ , if  $a \in C(b)$  and  $C(b) \neq A$ , then  $b \in C(a)$ .

(F5) For all  $a, b \in A$ , if  $a \in C(b)$  and  $C(b) \neq A$ ,  $\neg b \in C(\neg a)$ .

(F6) For all  $B \cup \{a\} \subseteq A$ , if  $a \in C(B)$  and  $C(B) \neq A$ , then there is  $b \in B$  such that  $a \in C(b)$ .

(F7) For every  $B \subseteq A$ , if  $C(B) = A$ , then there are  $b, b' \in B$  such that  $C(b) = C(\neg b')$ .

*Proof.* Let  $\langle A, C \rangle$  be a g-matrix and let  $\mathcal{C}$  be the closure system associated with  $C$ .

( $\Rightarrow$ ) Assume that  $\langle A, C \rangle$  is a full g-model of the logic  $\mathcal{S}_N$ . Thus, there is an algebra  $A_1$  and a surjective strict homomorphism  $h: \langle A, \mathcal{C} \rangle \rightarrow \langle A_1, \mathcal{F}i_{\mathcal{S}_N}(A_1) \rangle$ . Conditions (F1)–(F3)

follow by Proposition 6.3 and from the fact that these conditions are preserved by surjective strict homomorphisms (see [1, Proposition 5.90]).

(F4) Let  $a \in C(b)$  and suppose that  $C(b) \neq A$ . Since  $h$  is a strict homomorphism, it follows that  $ha \in \text{Fig}_{\mathcal{S}_N}^{A_1}(hb)$  and  $\text{Fig}_{\mathcal{S}_N}^{A_1}(hb) \neq A$ . Thus, by Proposition 6.6, we have that  $hb \in \text{Fig}_{\mathcal{S}_N}^{A_1}(ha)$ . Then  $b \in C(a)$ .

(F5) It is similar to the proof of (F4).

(F6) Let  $a \in C(B)$  and suppose that  $C(B) \neq A$ . Then, we have that  $ha \in \text{Fig}_{\mathcal{S}_N}^{A_1}(hB)$  and  $\text{Fig}_{\mathcal{S}_N}^{A_1}(hB) \neq A$ . By Proposition 6.7, there is  $b \in B$  such that  $ha \in \text{Fig}_{\mathcal{S}_N}^{A_1}(hb)$ . Then  $a \in C(b)$  with  $b \in B$ .

(F7) Let  $B \subseteq A$  be such that  $C(B) = A$ . Since  $h$  is a surjective strict homomorphism, it follows that  $\text{Fig}_{\mathcal{S}_N}^{A_1}(hB) = A_1$ . By Proposition 6.5, there are  $b, b' \in B$  and  $(s, t) \in \mathcal{E} \times \mathcal{O}$  such that  $\neg^s hb = \neg^t hb'$ . Then,  $\text{Fig}_{\mathcal{S}_N}^{A_1}(h(\neg^s b)) = \text{Fig}_{\mathcal{S}_N}^{A_1}(h(\neg^t b'))$ . Thus  $C(\neg^s b) = C(\neg^t b')$ . By (F3), it follows that  $C(b) = C(\neg^s b) = C(\neg^t b') = C(\neg b')$ .

( $\Leftarrow$ ) Assume that  $\langle A, C \rangle$  is a g-matrix satisfying conditions (F1)–(F7). Let  $\langle A^*, C^* \rangle$  be the reduction of  $\langle A, C \rangle$ . That is,  $A^* = A/\tilde{\Omega}^A C$  and  $C^* = \{F/\tilde{\Omega}^A C : F \in \mathcal{C}\}$ . By Proposition 6.1, it is enough to prove that  $C^* = \mathcal{F}i_{\mathcal{S}_N}(A^*)$ . Notice that the natural homomorphism  $\pi: A \rightarrow A^*$  is a surjective strict homomorphism from the g-matrix  $\langle A, C \rangle$  onto its reduction  $\langle A^*, C^* \rangle$ . For each  $a \in A$ , let  $\bar{a} = a/\tilde{\Omega}^A C$ .

( $\subseteq$ ) By (F2) and (F3), we have that the g-matrix  $\langle A, C \rangle$  is a g-model of  $\mathcal{S}_N$ . Then,  $\langle A^*, C^* \rangle$  is a g-model of  $\mathcal{S}_N$ . Hence  $C^* \subseteq \mathcal{F}i_{\mathcal{S}_N}(A^*)$ .

( $\supseteq$ ) In order to prove the inverse inclusion, let us show that  $C^*(B^*) \subseteq \text{Fig}_{\mathcal{S}_N}^{A^*}(B^*)$ , for all  $B \subseteq A$  where  $B^* = \{\bar{b} : b \in B\}$ . Let  $B \subseteq A$ . If  $\text{Fig}_{\mathcal{S}_N}^{A^*}(B^*) = A^*$ , then  $C^*(B^*) \subseteq \text{Fig}_{\mathcal{S}_N}^{A^*}(B^*)$ . Assume that  $\text{Fig}_{\mathcal{S}_N}^{A^*}(B^*) \neq A^*$ . Let us show that  $C^*(B^*) \neq A^*$ . Suppose by contradiction that  $C^*(B^*) = A^*$ . Thus  $C(B) = A$ . By (F7), there are  $a, a' \in B$  such that  $C(a) = C(\neg a')$ . Now we show that  $C(a) \neq A$ . Suppose that  $C(a) = A$ . Thus, by (F7), we obtain that  $C(a) = C(\neg a)$ . By (F3), we have that  $C(\neg a) = C(\neg(\neg a))$ . Then, by Proposition 6.9, it follows that  $\bar{a} = \overline{\neg a} = \neg \bar{a}$ . Since  $\bar{a} \in B^*$ , we have that  $\bar{a}, \neg \bar{a} \in \text{Fig}_{\mathcal{S}_N}^{A^*}(B^*)$ . Then  $\text{Fig}_{\mathcal{S}_N}^{A^*}(B^*) = A^*$ , which is a contradiction. Hence  $C(\neg a') = C(a) \neq A$ . It follows by (F5) and (F3) that  $C(\neg a) = C(a')$ . Then  $\bar{a} = \overline{\neg a'}$ . Given that  $\bar{a}, \overline{\neg a'} \in B^*$ , we obtain that  $\text{Fig}_{\mathcal{S}_N}^{A^*}(B^*) = A^*$ . A contradiction. Hence  $C^*(B^*) \neq A^*$ . And thus  $C(B) \neq A$ . Now, let  $\bar{a} \in C^*(B^*)$ . So  $a \in C(B)$ . By (F6), there is  $a_0 \in B$  such that  $a \in C(a_0)$ . Since  $C(B) \neq A$ , it follows that  $C(a_0) \neq A$ . Thus, since  $a \in C(a_0) \neq A$ , it follows by (F4) and (F5) that  $a_0 \in C(a)$  and  $\neg a_0 \in C(\neg a)$ . Thus  $C(a) = C(a_0)$ . Since  $C(a) = C(a_0) \neq A$  and  $a_0 \in C(a)$ , we have by (F5) that  $\neg a \in C(\neg a_0)$ . Then, we obtain that  $C(a) = C(a_0)$  and

$C(\neg a) = C(\neg a_0)$ . Thus  $\bar{a} = \bar{a}_0 \in B^*$ . Hence  $\bar{a} \in \text{Fig}_{\mathcal{S}_N}^{A^*}(B^*)$ . Hence,  $C^*(B^*) \subseteq \text{Fig}_{\mathcal{S}_N}^{A^*}(B^*)$ .

We have proved that  $C^* = \mathcal{F}i_{\mathcal{S}_N}(A^*)$ . Therefore,  $\langle A, C \rangle$  is a full g-model of  $\mathcal{S}_N$ .  $\square$

Notice that conditions (F1)–(F6) coincide with conditions (B1)–(B6) when they are considered on the g-matrices  $\langle A, \mathcal{F}i_{\mathcal{S}_N}(A) \rangle$ . And condition (F7) coincides to the implication (2) $\Rightarrow$ (1) of Proposition 6.5.

Let  $\langle A, C \rangle$  be a g-matrix. The *Frege relation*, denoted by  $\Lambda_A C$ , of  $C$  on  $A$  is defined as follows:

$$(a, b) \in \Lambda_A C \iff C(a) = C(b)$$

for all  $a, b \in A$ . Notice that the Tarski congruence is the largest congruence below  $\Lambda_A C$ .

**Proposition 6.11.** *Let  $\langle A, C \rangle$  be a g-matrix satisfying conditions (F5) and (F7). Then, the Frege relation  $\Lambda_A C$  is a congruence on  $A$ .*

*Proof.* Let  $a, b \in A$  be such that  $(a, b) \in \Lambda_A C$ . Thus  $C(a) = C(b)$ . If  $C(a) = C(b) \neq A$ , then we have  $a \in C(b) \neq A$ . Then, we obtain by (F5) that  $\neg b \in C(\neg a)$ . Analogously,  $\neg a \in C(\neg b)$ . Hence  $C(\neg a) = C(\neg b)$ . On the other hand, if  $C(a) = C(b) = A$ , it follows by (F7) that  $C(a) = C(\neg a)$  and  $C(b) = C(\neg b)$ . Then  $C(\neg a) = C(\neg b)$ . Hence  $(\neg a, \neg b) \in \Lambda_A C$ .  $\square$

The following proposition tells us that for an algebra  $A$  in the intrinsic variety of  $\mathcal{S}_N$ , the proper generated  $\mathcal{S}_N$ -filters on  $A$  are quite simple.

**Proposition 6.12.** *Let  $A \in \mathbb{V}(\mathcal{S}_N)$ . For all  $B \subseteq A$ , if  $\text{Fig}_{\mathcal{S}_N}^A(B) \neq A$ , then  $\text{Fig}_{\mathcal{S}_N}^A(B) = B$ .*

*Proof.* Let  $A \in \mathbb{V}(\mathcal{S}_N)$ . Suppose that  $\text{Fig}_{\mathcal{S}_N}^A(B) \neq A$  and let  $a \in \text{Fig}_{\mathcal{S}_N}^A(B)$ . By Proposition 6.4, there is  $b \in B$  and  $s, t \in \mathcal{E}$  such that  $\neg^s a = \neg^t b$ . Since  $A \models x \approx \neg \neg x$ , it follows that  $a = b \in B$ . Hence  $\text{Fig}_{\mathcal{S}_N}^A(B) = B$ .  $\square$

**Theorem 6.13.**

$$\text{Alg}(\mathcal{S}_N) = \{A : A \models x \approx \neg \neg x \text{ and } A \models (x \approx \neg x \ \& \ y \approx \neg y \implies x \approx y)\}.$$

*Proof.* ( $\subseteq$ ) Let  $A \in \text{Alg}(\mathcal{S}_N)$ . Thus  $\langle A, \mathcal{F}i_{\mathcal{S}_N}(A) \rangle$  is a reduced full g-model of  $\mathcal{S}_N$ . Then  $\langle A, \mathcal{F}i_{\mathcal{S}_N}(A) \rangle$  satisfies conditions (F1)–(F7). By the previous proposition, it follows that  $\Lambda_A \mathcal{F}i_{\mathcal{S}_N}(A) = \tilde{\Omega}^A \mathcal{F}i_{\mathcal{S}_N}(A) = \text{Id}_A$ . Then, by (B3) (or (F3)), we obtain that  $a = \neg \neg a$ , for all  $a \in A$ . Hence  $A \models x \approx \neg \neg x$ . Now let  $a, b \in A$  and assume that  $a = \neg a$  and  $b = \neg b$ . By (B2), we have  $\text{Fig}_{\mathcal{S}_N}^A(a) = A = \text{Fig}_{\mathcal{S}_N}^A(b)$ . Thus  $(a, b) \in \Lambda_A \mathcal{F}i_{\mathcal{S}_N}(A)$ . Then  $a = b$ . Hence  $A \models (x \approx \neg x \ \& \ y \approx \neg y \implies x \approx y)$ .

Classes Fragments	$\text{Alg}^*(\mathcal{S}_N)$	$\text{Alg}(\mathcal{S}_N)$	$\mathbb{V}(\mathcal{S}_N)$	For instances, see
$\{\neg\}$	$\mathbb{I}(A_1, A_2, A_3)$	$A \models x \approx \neg\neg x$ and $A \models (x \approx \neg x \ \& \ y \approx \neg y \implies x \approx y)$	$A \models x \approx \neg\neg x$	
$\{\wedge\}$	$\mathbb{I}(1_\wedge, 2_\wedge)$	(meet)-Semilattices	(meet)-Semilattices	[5] (also [1, pp.208–209])
$\{\vee\}$	(Join)-Semilattices with 1 + $a < b \implies \exists c(a \vee c \neq 1 \ \& \ b \vee c = 1)$	(join)-Semilattices	(join)-Semilattices	[5]
$\{\rightarrow\}$ or $\{\vee, \rightarrow\}$	Implication algebras (or Tarski algebras)	Implication algebras (or Tarski algebras)	Implication algebras (or Tarski algebras)	[7] (also [1, pp. 85])
$\{\wedge, \vee\}$	Distributive lattices with 1 + $a < b \implies \exists c(a \vee c \neq 1 \ \& \ b \vee c = 1)$	Distributive lattices	Distributive lattices	[2, 6]
$\{\wedge, \rightarrow\}$ or $\{\wedge, \vee, \rightarrow\}$	Relatively complemented distributive lattices with upper bound 1	Relatively complemented distributive lattices with upper bound 1	Relatively complemented distributive lattices with upper bound 1	
$\{\wedge, \neg\}$ , $\{\vee, \neg\}$ or $\{\rightarrow, \neg\}$	Boolean algebras	Boolean algebras	Boolean algebras	[1]

Table 1: The fragments of CPL and their classes of algebras.

( $\supseteq$ ) Let  $A$  be an algebra such that  $A \models x \approx \neg\neg x$  and  $A \models (x \approx \neg x \ \& \ y \approx \neg y \implies x \approx y)$ . It is straightforward that the g-matrix  $\langle A, \mathcal{F}i_{\mathcal{S}_N}(A) \rangle$  is a full g-model of  $\mathcal{S}_N^1$ . Let us show that the Frege relation of  $\langle A, \mathcal{F}i_{\mathcal{S}_N}(A) \rangle$  is the identity relation. Let  $(a, b) \in \Lambda_A \mathcal{F}i_{\mathcal{S}_N}(A)$ . Thus  $\text{Fig}_{\mathcal{S}_N}^A(a) = \text{Fig}_{\mathcal{S}_N}^A(b)$ . If  $\text{Fig}_{\mathcal{S}_N}^A(a) = \text{Fig}_{\mathcal{S}_N}^A(b) \neq A$ , it follows by Proposition 6.12 that  $a = b$ . Assume that  $\text{Fig}_{\mathcal{S}_N}^A(a) = \text{Fig}_{\mathcal{S}_N}^A(b) = A$ . From  $\text{Fig}_{\mathcal{S}_N}^A(a) = A$ , it follows by Proposition 6.5 that there is  $(s, t) \in \mathcal{E} \times \mathcal{O}$  such that  $\neg^s a = \neg^t a$ . Then  $a = \neg a$ . Similarly,  $b = \neg b$ . Since  $A \models (x \approx \neg x \ \& \ y \approx \neg y \implies x \approx y)$ , we obtain that  $a = b$ . Hence,  $\Lambda_A \mathcal{F}i_{\mathcal{S}_N}(A) = \text{Id}_A$ . Therefore, the full g-model  $\langle A, \mathcal{F}i_{\mathcal{S}_N}(A) \rangle$  is reduced, and thus  $A \in \text{Alg}(\mathcal{S}_N)$ .  $\square$

## 7. Concluding remarks

Throughout this article we have obtained the three classes of algebras that are canonically associated with the logic  $\mathcal{S}_N$  in the context of algebraic logic:

$$\text{Alg}^*(\mathcal{S}_N) = \mathbb{I}(A_1, A_2, A_3) \quad \mathbb{V}(\mathcal{S}) = \{A : A \models x \approx \neg\neg x\}$$

$$\text{Alg}(\mathcal{S}_N) = \{A : A \models (x \approx \neg\neg x) \ \& \ (x \approx \neg x \ \& \ y \approx \neg y \implies x \approx y)\}.$$

With this, we can complete the table with all the fragments of CPL and their corresponding classes of algebras canonically associated, see Table 1.

Notice that it is clear that the inclusions  $\text{Alg}^*(\mathcal{S}_N) \subset \text{Alg}(\mathcal{S}_N) \subset \mathbb{V}(\mathcal{S}_N)$  are proper. For instance, the algebra  $A_2 = \{a, b\}$ , with  $\neg a = a$  and  $\neg b = b$ , belongs to  $\mathbb{V}(\mathcal{S}_N)$  but not to

<sup>1</sup>Actually, for every algebra  $A$ , the g-matrices of the form  $\langle A, \mathcal{F}i_{\mathcal{S}}(A) \rangle$  are called *basic full g-models* of a logic  $\mathcal{S}$ , see [1].



$\text{Alg}(\mathcal{S}_N)$ . The  $\{\neg\}$ -reduct of the four-element Boolean algebra belongs to  $\text{Alg}(\mathcal{S}_N)$  but not to  $\text{Alg}^*(\mathcal{S}_N)$ . Also notice that the class  $\text{Alg}(\mathcal{S}_N)$  is a quasi-variety but not a variety.

From the results that we have obtained, we can classify the  $\{\neg\}$ -fragment of CPL in the Leibniz hierarchy and in the Frege hierarchy (the two hierarchies of algebraic logic, see [1, Chap. 6 and 7]) and respond to an open problem proposed in [1, pp. 418]. The  $\{\neg\}$ -fragment  $\mathcal{S}_N$  of CPL is out of the Leibniz hierarchy because it is neither protoalgebraic nor truth-equational: Notice that the logic  $\mathcal{S}_N$  is non-trivial, and thus  $\mathcal{S}_N$  is not almost inconsistent. Since the unique protoalgebraic logic without theorems is the almost inconsistent one, it follows that  $\mathcal{S}_N$  is not protoalgebraic. Moreover since the logic  $\mathcal{S}_N$  has no theorems, it follows that  $\mathcal{S}_N$  is not truth-equational. Now we turn our attention to the Frege hierarchy (see [1, pp. 414]). It is straightforward that the  $\{\neg\}$ -fragment of CPL is Fregean because  $\mathcal{S}_N$  is a two-valued logic. Moreover, the logic  $\mathcal{S}_N$  is also fully selfextensional: Let  $A \in \text{Alg}(\mathcal{S}_N)$ . In the proof of Theorem 6.13 we have proved that the Frege relation  $\Lambda_A \mathcal{F}i_{\mathcal{S}_N}(A)$  is the identity relation. Hence  $\mathcal{S}_N$  is fully selfextensional. However, we will show that  $\mathcal{S}_N$  is not fully Fregean. In order to show this, consider the following relation: let  $\mathcal{S}$  be a logic,  $A$  an algebra and  $F \subseteq A$ ,

$$\Lambda_{\mathcal{S}}^A F = \{(a, b) \in A \times A : \text{Fig}_{\mathcal{S}}^A(F, a) = \text{Fig}_{\mathcal{S}}^A(F, b)\}.$$

**Proposition 7.1** ([1, Prop. 7.56]). *A logic  $\mathcal{S}$  is fully Fregean if and only if  $\Lambda_{\mathcal{S}}^A F \subseteq \Omega^A F$  for every  $F \in \mathcal{F}i_{\mathcal{S}}(A)$  and every algebra  $A$ .*

Let  $A_3 = \{a, b, c\}$  be the algebra given on page 8. Let  $F = \{a\}$ . We know that  $F \in \mathcal{F}i_{\mathcal{S}_N}(A_3)$ . We have that  $\text{Fig}_{\mathcal{S}_N}^{A_3}(F, b) = A$  (because  $b, \neg b \in \text{Fig}_{\mathcal{S}_N}^{A_3}(F, b)$ ) and  $\text{Fig}_{\mathcal{S}_N}^{A_3}(F, c) = A$  (because  $c, \neg c \in \text{Fig}_{\mathcal{S}_N}^{A_3}(F, c)$ ). Hence  $(b, c) \in \Lambda_{\mathcal{S}_N}^A F$ . But  $(b, c) \notin \Omega^A F$  because  $\neg b \in F$  and  $\neg c \notin F$  (see Proposition 4.1). Thus  $\Lambda_{\mathcal{S}_N}^A F \not\subseteq \Omega^A F$ . Therefore, the  $\{\neg\}$ -fragment of CPL is not fully Fregean. This answers negatively to the question: Is the class of fully Fregean logics the intersection of the classes of the Fregean and the fully selfextensional ones? That is,  $\mathcal{S}_N$  is a Fregean and fully selfextensional logic but is not fully Fregean one.

## Funding

This work was supported by the Fondo para la Investigación Científica y Tecnológica (Argentina) [grant number PICT-2019-0674, PICT-2019-0882]; and the Universidad Nacional de La Pampa [grant number P.I. 78M].

## References

- [1] J. M. Font. *Abstract Algebraic Logic. An Introductory Textbook*, volume 60 of *Studies in Logic*. College Publications, London, 2016.
- [2] J. M. Font, F. Guzmán, and V. Verdú. Characterization of the reduced matrices for the  $\{\wedge, \vee\}$ -fragment of classical logic. *Bulletin of the Section of Logic*, 20(3/4):124–128, 1991.
- [3] J. M. Font and R. Jansana. *A general algebraic semantics for sentential logics*, volume 7 of *Lecture Notes in Logic*. A.S.L., second edition, 2009.
- [4] J. M. Font, R. Jansana, and D. Pigozzi. A survey of abstract algebraic logic. *Studia Logica*, 74(1-2):13–97, 2003.
- [5] J. M. Font and T. Moraschini. A note on congruences of semilattices with sectionally finite height. *Algebra universalis*, 72(3):287–293, 2014.
- [6] J. M. Font and V. Verdú. Algebraic logic for classical conjunction and disjunction. *Studia Logica*, 50(3-4):391–419, 1991.
- [7] H. Rasiowa. *An Algebraic Approach to Non-Classical Logics*. North-Holland, 1974.
- [8] W. Rautenberg. 2-element matrices. *Studia Logica*, 40(4):315–353, 1981.