

# A categorical proof of the Carathéodory extension theorem

*Ruben Van Belle* \*

## Abstract

The Carathéodory extension theorem is a fundamental result in measure theory. Often we do not know what a general measurable subset looks like. The Carathéodory extension theorem states that to define a measure we only need to assign values to subsets in a generating Boolean algebra.

To prove this result categorically, we represent (pre)measures and outer measures by certain (co)lax and strict transformations. The Carathéodory extension then corresponds to a Kan extension of strict transformations. We develop a general framework for extensions of transformations between poset-valued functors and give several results on the existence and construction of extensions of these transformations. We proceed by showing that transformations and functors corresponding to measures satisfy these results, which proves the Carathéodory extension theorem.

## Contents

<b>1</b>	<b>Introduction</b>	2
<b>2</b>	<b>Categories of (co)lax transformations</b>	3
<b>3</b>	<b>Extensions of transformations</b>	6
<b>4</b>	<b>The (co)monad formula</b>	11
<b>5</b>	<b>The extension formula</b>	18
<b>6</b>	<b>Premeasures as transformations</b>	19
	6.1 Premeasures	20
	6.2 Outer/inner premeasures	23
<b>7</b>	<b>The Carathéodory extension theorem</b>	28
	7.1 The right Carathéodory extension	29
	7.2 The left Carathéodory extension	31
<b>A</b>	<b>Lax coends</b>	32
<b>B</b>	<b>Weakly final functors</b>	33
<b>C</b>	<b>Finite outer premeasures</b>	34

---

\*School of Mathematics, University of Edinburgh; ruben.van.belle@ed.ac.uk

## 1 Introduction

The Carathéodory extension theorem is an important result in measure theory. It guarantees the existence of the Lebesgue measure (or more generally the Lebesgue-Stieltjes measure) and of product measures. But it also is a key result in the proof of the Kolmogorov extensions theorem, which guarantees the existence of Brownian motion and which is closely related to martingale convergence results.

If we want to define a measure on a measurable space  $(X, \Sigma)$ , we need to assign to every subset  $A \in \Sigma$  a real number. However, we often work with  $\sigma$ -algebras of the form  $\sigma(\mathcal{B})$ , where  $\mathcal{B}$  is an algebra of subsets of  $X$ , i.e. closed under finite unions and complements. In this case it can be very difficult to know what a general measurable subset looks like and to assign real numbers to them in a  $\sigma$ -additive way. This problem is solved by the Carathéodory extension theorem. It states that a  $\sigma$ -additive map  $\mathcal{B} \rightarrow [0, \infty]$  can be extended to a  $\sigma$ -additive map  $\sigma(\mathcal{B}) \rightarrow [0, \infty]$ . Here,  $\sigma$ -additive should be interpreted as ‘ $\sigma$ -additive whenever unions of countable pairwise disjoint collections exist’. Moreover, this extension is very often, but not always, unique.

$$\begin{array}{ccc}
 \mathcal{B} & \xrightarrow{\quad} & [0, \infty] \\
 \downarrow & \nearrow \text{---} & \\
 \sigma(\mathcal{B}) & & 
 \end{array}$$

The classical proof for this result is relatively long and technical (see for example Theorem 1.41 in [3]). It consists of different steps of extending and restricting back and requires several smart ‘tricks’ and constructions. In this paper we will give a categorical proof for the Carathéodory extension theorem using results on extensions of lax/colax/strict transformations between functors. Several parts of this proof look similar to steps in the classical proof. However, in our proof all constructions follow from the Kan extension formulas and gives an approach that is a lot more structured. Moreover, viewing the Carathéodory extension theorem in this categorical framework, allows us to compare it to extension results in other areas of mathematics. Furthermore, this technique allows us to easily generalize Carathéodory’s result to measures taking values in other posets or spaces.

To do this we start by studying categories of certain transformations between functors and extensions of transformations along transformations. In section 2 and 3 we do this on a fairly abstract level. We discuss (co)laxification (co)monads and are in particular interested in the case that their (co)algebras are strict transformations. We give several abstract conditions for extensions of certain transformations to exist.

In section 4 and 5, we give concrete constructions for the operations and extensions discussed in 2 and 3. In this part (co)lax coends play an important role.

In the last two sections of the paper we apply the previous sections to measure theory. We start by representing inner and outer premeasures by certain transformations between certain functors. By applying the extension results on these functors, we obtain

our categorical proof for Carathédory's extension result. We furthermore, make a distinction between *left* and *right* Carathédory extensions. The right one always exists and is the same as the extension in the classical proof, the left one however does not always exist. This also explains why inner measures and outer measures are, surprisingly, not as symmetric as one might expect. Moreover, we immediately obtain a way to characterize these extensions by a universal property, which is interesting when uniqueness is not guaranteed.

	Categories of transformations	Extensions of transformations
Abstract theory	2	3
Concrete constructions	4	5
Applications to measures	6	7

## 2 Categories of (co)lax transformations

In this section we will introduce and discuss categories of (co)lax  $\Sigma$ -transformations and strict transformations. In particular we will be interested in the embeddings between these and when these are (co)reflective. Furthermore, we will look at the (co)monads these adjunctions induce and their co/algebras.

Let  $\mathcal{C}$  be a small strict 2-category. Let  $\mathbf{Cat}$  be the strict 2-category of small categories, functors and natural transformations. Let  $F$  and  $G$  be strict 2-functors  $\mathcal{C} \rightarrow \mathbf{Cat}$  such that  $GA$  is a poset for all  $A \in \text{ob } \mathcal{C}$ . Let  $\Sigma$  be a collection of morphisms in  $\mathcal{C}$ . A collection of maps  $\tau := (\tau_A : FA \rightarrow GA)_{A \in \text{ob } \mathcal{C}}$  is called a  $\Sigma$ -**transformation**, if for all  $(f : A \rightarrow B) \in \Sigma$

$$Gf \circ \tau_A = \tau_B \circ Ff.$$

A  $\Sigma$ -transformations  $\tau$  is called **lax** if for all  $f : A \rightarrow B$  in  $\mathcal{C}$

$$Gf \circ \tau_A \geq \tau_B \circ Ff,$$

which is an equality whenever  $f \in \Sigma$ . It is called **colax** if for all  $f : A \rightarrow B$  in  $\mathcal{C}$ ,

$$Gf \circ \tau_A \leq \tau_B \circ Ff,$$

where this is an equality when  $f \in \Sigma$ . We say that a  $\Sigma$ -transformation is a **strict transformation** or a **strict  $\Sigma$ -transformation** if  $Gf \circ \tau_A = \tau_B \circ Ff$  for *all* morphisms  $f$  in  $\mathcal{C}$ . If we want to emphasize that a  $\Sigma$ -transformation is not necessarily lax or colax, we will talk about **general  $\Sigma$ -transformations**. In the case that  $\Sigma = \emptyset$ , we will omit ' $\Sigma$ ' in the terminology and notation.

The set of  $\Sigma$ -transformations is partially ordered. The order is defined by

$$\tau^1 \leq \tau^2 :\Leftrightarrow \tau_A^1(x) \leq \tau_A^2(x)$$

for all  $A \in \text{ob } \mathcal{C}$  and  $x \in FA$  and for  $\Sigma$ -transformations  $\tau^1$  and  $\tau^2$ .

The poset of  $\Sigma$ -transformations is denoted by  $[F, G]^\Sigma$ ; the subposets of lax, colax and strict  $\Sigma$ -transformations are denoted by  $[F, G]_l^\Sigma$ ,  $[F, G]_c^\Sigma$  and  $[F, G]_s$  respectively. We clearly have the following pullback square of inclusions.

$$\begin{array}{ccc}
 & [F, G]_s & \\
 \swarrow & \downarrow \vee & \searrow \\
 [F, G]_l^\Sigma & & [F, G]_c^\Sigma \\
 \searrow U_l & & \swarrow U_c \\
 & [F, G]^\Sigma &
 \end{array}$$

**Proposition 2.1.** *Suppose  $GA$  is cocomplete<sup>1</sup> for all  $A \in \text{ob}\mathcal{C}$  and suppose  $Gf$  preserves all joins for all  $f \in \Sigma$ . Then  $[F, G]_l^\Sigma$  and  $[F, G]^\Sigma$  are cocomplete and the inclusion  $U_l : [F, G]_l^\Sigma \rightarrow [F, G]^\Sigma$  preserves all joins.*

*Proof.* Let  $(\lambda^i)_{i \in I}$  be a collection of  $\Sigma$ -transformations. Define for all  $A \in \text{ob}\mathcal{C}$  and  $x \in FA$ ,

$$\lambda_A(x) := \bigvee_{i \in I} \lambda_A^i(x).$$

For  $f : A \rightarrow B \in \Sigma$  and  $x \in FA$ ,

$$Gf(\lambda_A(x)) = Gf\left(\bigvee_{i \in I} \lambda_A^i(x)\right) = \bigvee_{i \in I} Gf(\lambda_A^i(x)) = \bigvee_{i \in I} \lambda_B^i(Ff(x)) = \lambda_B(Ff(x)).$$

Therefore  $\lambda$  is a  $\Sigma$ -transformation and it is easy to see that it is the join of  $(\lambda^i)_{i \in I}$  in  $[F, G]$ . If  $\lambda^i$  is lax for every  $i \in I$ , then for a map  $f : A \rightarrow B$  in  $\mathcal{C}$  and  $x \in FA$ ,

$$Gf(\lambda_A(x)) = Gf\left(\bigvee_{i \in I} \lambda_A^i(x)\right) \leq \bigvee_{i \in I} Gf(\lambda_A^i(x)) \leq \bigvee_{i \in I} \lambda_B^i(Ff(x)) = \lambda_B(Ff(x)).$$

Here we used that  $Gf$  is order-preserving and that  $\lambda^i$  is lax for all  $i \in I$ . It follows that  $\lambda$  is a lax  $\Sigma$ -transformation. It is easy to see that this is the join of  $(\lambda^i)_{i \in I}$  and clearly  $U_l$  preserves this join.  $\square$

Using Proposition 2.1, we immediately obtain the following corollary.

**Corollary 2.2.** *If  $GA$  is complete for all  $A \in \text{ob}\mathcal{C}$  and if  $Gf$  preserves joins for all  $f \in \Sigma$ , then  $U_l : [F, G]_l^\Sigma \rightarrow [F, G]^\Sigma$  has a right adjoint  $R_l : [F, G]^\Sigma \rightarrow [F, G]_l^\Sigma$ .*

<sup>1</sup> Since  $GA$  is a poset for every object  $A$  in  $\mathcal{C}$ ,  $GA$  is also complete.

Since  $U_l$  is full and faithful, the unit is an equality, i.e.  $\lambda = R_l U_l \lambda$  for a lax  $\Sigma$ -transformation  $\lambda$ . Clearly, we also have dual results for Proposition 2.1 and Corollary 2.2. In particular, if  $GA$  is cocomplete for all  $A \in \text{ob } \mathcal{C}$  and  $Gf$  preserves meets for all  $f \in \Sigma$ , then  $U_c : [F, G]_c^\Sigma \rightarrow [F, G]^\Sigma$  has a left adjoint  $L_c : [F, G]^\Sigma \rightarrow [F, G]_c^\Sigma$  and  $\sigma = L_c U_c \sigma$  for a colax  $\Sigma$ -transformation  $\sigma$ .

Assume from now on that  $GA$  is (co)complete and that  $Gf$  preserves all joins and meets for  $f \in \Sigma$ . For a lax  $\Sigma$ -transformation  $\lambda$ , we denote

$$\bar{\lambda} := L_c U_l(\lambda).$$

and for a colax  $\Sigma$ -transformation  $\sigma$ , we write

$$\underline{\sigma} := R_l U_c(\sigma).$$

Note that the map  $\overline{(-)} : [F, G]_l^\Sigma \rightarrow [F, G]_c^\Sigma$  is left adjoint to  $\underline{(-)} : [F, G]_c^\Sigma \rightarrow [F, G]_l^\Sigma$ . The following diagram summarizes this;

$$\begin{array}{ccc}
 & \overline{(-)} & \\
 & \curvearrowright & \\
 [F, G]_l^\Sigma & \xrightarrow{U_l} & [F, G]^\Sigma & \xrightarrow{L_c} & [F, G]_c^\Sigma \\
 & \xleftarrow{R_l} & & \xleftarrow{U_c} & \\
 & \curvearrowleft & & \curvearrowright & \\
 & \underline{(-)} & & & 
 \end{array}$$

The monad on  $[F, G]_l^\Sigma$  induced by this adjunction is denoted by  $T$  and the induced comonad on  $[F, G]_c^\Sigma$  is denoted by  $S$ .

**Proposition 2.3.** *The following are equivalent:*

1.  $\bar{\lambda}$  is strict for all  $\lambda \in [F, G]_l^\Sigma$ ,
2.  $([F, G]_l^\Sigma)^T = [F, G]_s$ ,
3.  $\underline{\sigma}$  is strict for all  $\sigma \in [F, G]_c^\Sigma$
4.  $([F, G]_c^\Sigma)^S = [F, G]_s$

*Proof.* 1  $\Rightarrow$  2: For a strict transformation  $\tau$ , we clearly have that  $T\tau = \tau$ , making it an algebra. For an algebra  $\lambda$ , we find that  $\bar{\lambda} = T\lambda = \lambda$ . This means that  $\lambda$  is colax and therefore strict.

2  $\Rightarrow$  3: Because  $\underline{\sigma} = T\underline{\sigma}$ , the claim follows.

3  $\Rightarrow$  4: Every strict transformation  $\tau$  is a coalgebra since  $S\tau = \tau$ . For a coalgebra  $\sigma$ , we have  $\underline{\sigma} = S\underline{\sigma} = \sigma$ . Therefore  $\sigma$  is lax and thus strict.

4  $\Rightarrow$  1: This follows from the fact that  $\bar{\lambda} = S\bar{\lambda}$ . □

We say that the triple  $(F, G, \Sigma)$  satisfies the **strictness condition** if it satisfies the conditions in Proposition 2.3. In this case it is clear that the inclusion  $[F, G]_s \rightarrow [F, G]_l^\Sigma$  has a right adjoint  $R : [F, G]_l^\Sigma \rightarrow [F, G]_s$  and that the inclusion  $[F, G]_s \rightarrow [F, G]_c^\Sigma$  has a left adjoint  $L : [F, G]_c^\Sigma \rightarrow [F, G]_s$ .

Using the properties of left adjoints it is easy to prove the following corollary.

**Corollary 2.4.** *Suppose that  $GA$  is complete for all  $A \in \text{ob } \mathcal{C}$  and that  $Gf$  preserves meets for all  $f \in \Sigma$ . Suppose that  $(F, G, \Sigma)$  satisfies the strictness condition. Let  $(\tau^i)_{i \in I}$  be a collection in  $[F, G]_s$ . Then their join exists and is given by*

$$\bigvee_{i \in I} \tau^i = \bar{\lambda},$$

where  $\lambda_A(x) := \bigvee_{i \in I} \tau_A^i(x)$  for all  $A \in \text{ob } \mathcal{C}$  and  $x \in FA$ .

### 3 Extensions of transformations

In this section, we will focus on extensions of (co)lax and strict  $\Sigma$ -transformations. We will give results on their existence and their properties.

Let  $\mathcal{C}$  be a small strict 2-category and fix a collection  $\Sigma$  of morphisms in  $\mathcal{C}$ . Let  $F, G, H$  be strict 2-functors  $\mathcal{C} \rightarrow \mathbf{Cat}$  and assume that  $GA$  is complete for all  $A \in \text{ob } \mathcal{C}$  and that  $Gf$  preserves joins and meets for all  $f \in \Sigma$ . Furthermore, let  $\iota : F \rightarrow H$  be a strict transformation.

Let  $\lambda : F \rightarrow G$  be a lax (resp. colax, strict)  $\Sigma$ -transformation. The **left extension of  $\lambda$  along  $\iota$**  is a lax (resp. colax, strict) transformation  $\text{Lan}_\iota \lambda : H \rightarrow G$  such that  $\text{Lan}_\iota \lambda \circ \iota \geq \lambda$  and such that for every other  $\lambda' : H \rightarrow G$  with  $\lambda' \circ \iota \geq \lambda$ , we have  $\lambda' \leq \text{Lan}_\iota \lambda$ ;<sup>2</sup>

$$\begin{array}{ccc} F & \xrightarrow{\lambda} & G \\ \downarrow \iota & \nearrow \text{Lan}_\iota \lambda & \uparrow \\ H & \xrightarrow{\lambda'} & G \end{array}$$

We say that the extension is **proper** if  $\text{Lan}_\iota \lambda \circ \iota = \lambda$  and we call the extension **objectwise** if for all  $A \in \text{ob } \mathcal{C}$ ,

$$(\text{Lan}_\iota \lambda)_A = \text{Lan}_{\iota_A} \lambda_A.$$

Dually, we can define the **right extension of  $\lambda$  along  $\iota$** .

If an extension is objectwise, then we can reduce everything to Kan extensions of functors. This gives us for example the following lemma.

<sup>2</sup> The universal property determines the transformation, therefore we can talk about *the* left extensions

**Lemma 3.1.** *If the left extension of  $\lambda$  along  $\iota$  is objectwise and if  $\iota_A$  is full and faithful for every  $A \in \text{ob } \mathcal{C}$ , then the extension is proper.*

*Proof.* Because  $\iota_A$  is full and faithful for all  $A \in \text{ob } \mathcal{C}$ , we know that  $(\text{Lan}_{\iota_A} \lambda_A) \circ \iota_A = \lambda_A$  for all  $A \in \text{ob } \mathcal{C}$ . Because the extensions of  $\lambda$  along  $\iota$  is objectwise, we conclude that

$$(\text{Lan}_{\iota} \lambda) \circ \iota = \lambda.$$

Therefore the extension is proper.  $\square$

Just as for Kan extensions of functors, extending transformations is adjoint to restricting transformations. The proof is essentially the same as the result for Kan extensions of functors (see for example Proposition 6.1.5 in [5]).

**Lemma 3.2.** *If the restriction map  $-\circ \iota : [H, G]_{\bullet} \rightarrow [F, G]_{\bullet}$  has a right (resp. left) adjoint, then the right (resp. left) extension of  $\lambda$  along  $\iota$  exists for all  $\lambda \in [F, G]_{\bullet}$ . Moreover, the right (resp. left) adjoint is given by  $\text{Ran}_{\iota} -$  (resp.  $\text{Lan}_{\iota} -$ ).*

In the rest of this section we will look at conditions for extensions to exist and for them to be proper or objectwise. Extensions of general transformations always exist and are the best behaved.

**Proposition 3.3.** *The right and left extension of  $\tau$  along  $\iota$  exists for every transformation  $\tau \in [F, G]$ .*

*Proof.* It is clear by the proof of Proposition 2.1 and its dual, that  $[F, G]^{\Sigma}$  and  $[H, G]^{\Sigma}$  are complete and that  $-\circ \iota$  preserves all joins and meets. It follows now that the restriction map  $-\circ \iota$  has a left and a right adjoint. The claim now follows from Lemma 3.2.  $\square$

For lax transformation things become more difficult. However, we still have that right extensions exist.

**Proposition 3.4.** *The right extension of  $\lambda$  along  $\iota$  exists for all  $\lambda \in [F, G]_{\iota}^{\Sigma}$ .*

*Proof.* It is easy to see that  $[F, G]_{\iota}^{\Sigma}$  is cocomplete and that  $-\circ \iota$  preserves all joins. Therefore the restriction map has a right adjoint.  $\square$

**Remark 3.5.** We have the following inequality

$$\begin{array}{ccc} [H, G]_{\iota}^{\Sigma} \leftarrow \text{Ran}_{\iota} - & \text{---} & [F, G]_{\iota}^{\Sigma} \\ \downarrow U_{\iota} & \leq & \downarrow U_{\iota} \\ [H, G]^{\Sigma} \leftarrow \text{Ran}_{\iota} - & \text{---} & [F, G]^{\Sigma} \end{array}$$

If this is an equality, then right extensions of lax  $\Sigma$ -transformations along  $\iota$  inherit objectwiseness and properness from right extensions of (general)  $\Sigma$ -transformations.

We have the following useful property about the existence of extensions of lax  $\Sigma$ -transformations and when they inherit properties from (general)  $\Sigma$ -transformations.

**Proposition 3.6.** *We have the following inequality*

$$\begin{array}{ccc} [H, G]_l^\Sigma & \xrightarrow{-\circ\iota} & [F, G]_l^\Sigma \\ \uparrow R_l & \leq & \uparrow R_l \\ [H, G]^\Sigma & \xrightarrow{-\circ\iota} & [F, G]^\Sigma \end{array}$$

*If this is an equality, then left extensions of lax  $\Sigma$ -transformations along  $\iota$  exist and inherit objectwiseness and properness from left extensions of (general)  $\Sigma$ -transformations. Moreover, right extensions of lax  $\Sigma$ -transformations inherit properness from right extensions of (general)  $\Sigma$ -transformations.*

*Proof.* The map  $[H, G]^\Sigma \xrightarrow{-\circ\iota} [F, G]^\Sigma$  has a left adjoint by Proposition 3.3 and the following diagram commutes

$$\begin{array}{ccc} [H, G]_l^\Sigma & \xrightarrow{-\circ\iota} & [F, G]_l^\Sigma \\ \downarrow U_l & & \downarrow U_l \\ [H, G]^\Sigma & \xrightarrow{-\circ\iota} & [F, G]^\Sigma \end{array}$$

Since  $U_l$  is full and faithful, it follows from the adjoint lifting theorem in [2] that  $[H, G]_l^\Sigma \xrightarrow{-\circ\iota} [F, G]_l^\Sigma$  has a left adjoint. By the hypothesis we have that  $R_l(-\circ\iota) = (R_l-) \circ \iota$ . These are all right adjoint, so therefore their left adjoints commute as well, this means that the following diagram commutes

$$\begin{array}{ccc} [H, G]_l^\Sigma & \leftarrow \text{Lan}_\iota - & [F, G]_l^\Sigma \\ \downarrow U_l & & \downarrow U_l \\ [H, G]^\Sigma & \leftarrow \text{Lan}_\iota - & [F, G]^\Sigma \end{array}$$

It follows now easily that properness and objectwiseness are inherited from left extensions of (general)  $\Sigma$ -transformations.

Suppose now that right extensions of (general)  $\Sigma$ -transformations along  $\iota$  are proper. Because  $U_l(-\circ\iota) = (U_l-) \circ \iota$ , their right adjoint commute as well. Together with the



hypothesis, this leads to the following commutative square.

$$\begin{array}{ccccc}
 [F, G]_l^\Sigma & \xrightarrow{\text{Ran}_\iota} & [H, G]_l^\Sigma & \xrightarrow{-\circ\iota} & [F, G]_l^\Sigma \\
 \uparrow R_l & & \uparrow R_l & & \uparrow R_l \\
 [F, G]^\Sigma & \xrightarrow{\text{Ran}_\iota} & [H, G]^\Sigma & \xrightarrow{-\circ\iota} & [F, G]^\Sigma \\
 & \searrow & \xrightarrow{1_{[F, G]}} & \nearrow & \\
 & & & & 
 \end{array}$$

We now have for a lax  $\Sigma$ -transformation  $\lambda : F \rightarrow G$ ,

$$\text{Ran}_\iota \lambda \circ \iota = \text{Ran}_\iota R_l (U_l) \lambda \circ \iota = R_l U_l \lambda = \lambda.$$

□

We have dual result for Proposition 3.4 and Proposition 3.6 for colax  $\Sigma$ -transformation.

Under even more conditions, we can guarantee the existence of extensions of strict transformations. Let  $T_1$  be the monad on  $[H, G]_l^\Sigma$  as described above and let  $T_2$  be the monad on  $[F, G]_l^\Sigma$ .

**Proposition 3.7.** *Suppose  $-\circ\iota$  is a strict morphism of monads  $([H, G]_l^\Sigma, T_1) \rightarrow ([F, G]_l^\Sigma, T_2)$ . Then  $-\circ\iota$  induces an order-preserving map  $([H, G]_l^\Sigma)^{T_1} \rightarrow ([F, G]_l^\Sigma)^{T_2}$  and this map has a right adjoint. Moreover, if right extensions of lax  $\Sigma$ -transformations along  $\iota$  are proper (resp. objectwise), then so are right extensions of algebras along  $\iota$ .*

*Proof.* Since  $-\circ\iota$  is a morphism of monads, it induces an order-preserving map between the algebras. Furthermore, we have the following commutative square

$$\begin{array}{ccc}
 ([H, G]_l^\Sigma)^{T_1} & \xrightarrow{-\circ\iota} & ([F, G]_l^\Sigma)^{T_2} \\
 \uparrow & & \uparrow \\
 [H, G]_l & \xrightarrow{-\circ\iota} & [F, G]_l
 \end{array}$$

By the adjoint lifting theorem in [2], the right adjoint of  $-\circ\iota : [H, G]_l^\Sigma \rightarrow [F, G]_l^\Sigma$  can be lifted to a right adjoint of the algebras. Because the left adjoints commute, so do the right adjoints. It follows that properness and objectwiseness are inherited from the extensions of lax  $\Sigma$ -transformations. □

Applying this to the case that the strictness condition holds, immediately gives us the following corollary.

**Corollary 3.8.** *Suppose that  $(F, G, \Sigma)$  and  $(H, G, \Sigma)$  satisfy the strictness condition and that  $\bar{\lambda} \circ \iota = \overline{\lambda \circ \iota}$  for all  $\lambda \in [H, G]_\iota$ . Then,  $[H, G]_s \xrightarrow{-\circ\iota} [F, G]_s$  has a right adjoint. Moreover, if right extensions of lax transformations along  $\iota$  are proper (resp. objectwise), then so are right extensions of strict transformation along  $\iota$ .*

Again, there are dual results of Proposition 3.7 and Corollary 3.8 for left extensions of strict transformations.

In general, the existence and properties of left and right extensions are not connected. We can for example have that right extensions don't exist, but left extensions do and are well-behaved. However, we do have the following connection between left and right extensions of strict transformations.

**Proposition 3.9.** *Suppose that  $[H, G]_s \xrightarrow{-\circ\iota} [F, G]_s$  has a left and a right adjoint. Then, left extensions along  $\iota$  are proper if and only if right extensions along  $\iota$  are proper.*

*Proof.* Let  $\tau \in [F, G]_s$ . By the universal property of extensions, we have  $(\text{Ran}_\iota \tau)_A \leq \text{Ran}_{\iota_A} \tau_A$  and  $\text{Lan}_{\iota_A} \tau_A \leq (\text{Lan}_\iota \tau)_A$  for all  $A \in \text{ob } \mathcal{C}$ . We have the following inequalities for all  $A \in \text{ob } \mathcal{C}$ :

$$(\text{Ran}_\iota \tau \circ \iota)_A \leq \text{Ran}_{\iota_A} \tau_A \circ \iota_A \leq \tau_A \leq \text{Lan}_{\iota_A} \tau_A \circ \iota_A \leq (\text{Lan}_\iota \tau \circ \iota)_A$$


Suppose that left extensions along  $\iota$  are proper, i.e.  $\text{Lan}_\iota \tau \circ \iota = \tau$ . By the universal property of right extensions, it follows that

$$\text{Lan}_\iota \tau \leq \text{Ran}_\iota \tau.$$

Therefore  $\tau = \text{Lan}_\iota \tau \circ \iota = \text{Ran}_\iota \tau \circ \iota$ .

Similarly, if right extensions along  $\iota$  are proper, we find by the universal property of left extensions that  $\text{Lan}_\iota \tau \leq \text{Ran}_\iota \tau$  and we can conclude that  $\text{Lan}_\iota \tau \circ \iota = \text{Ran}_\iota \tau \circ \iota = \tau$ .  $\square$

An overview of the maps we *always* have is given by the following diagram. We assume that  $(F, G, \Sigma)$  and  $(H, G, \Sigma)$  satisfy the strictness condition. In the diagram, squares of the same colour commute.



## 4 The (co)monad formula

In this section we will give concrete constructions for the operation  $\overline{(-)}$  and  $\underline{(-)}$  discussed in the previous section, in the case that the functors are well-behaved.

Let  $\mathcal{C}$  be a small strict 2-category and let  $\Sigma$  be a collection of morphisms in  $\mathcal{C}$ . Let  $F$  and  $G$  be strict 2-functors  $\mathcal{C} \rightarrow \mathbf{Cat}$  such that  $GA$  is a complete poset for all  $A \in \text{ob } \mathcal{C}$  and such that  $Gf$  preserves meets and joins for all  $f \in \Sigma$ . Assume also that  $FA$  is a poset for every  $A \in \text{ob } \mathcal{C}$ .

Intuitively it is clear that  $(F, G, \emptyset)$  satisfying the strictness property is stronger than  $(F, G, \Sigma)$  satisfying the strictness property. This is the content of the following results. This proposition motivates that it is enough to give a construction of these operations in the case that  $\Sigma = \emptyset$ .

**Proposition 4.1.** *If  $(F, G, \emptyset)$  satisfies the strictness condition, then  $\overline{\lambda}^\Sigma = \overline{\lambda}^\emptyset$  for all  $\lambda \in [F, G]_l^\Sigma$ . In particular,  $(F, G, \Sigma)$  satisfies the strictness condition.<sup>3</sup>*

*Proof.* The inclusions  $i_c : [F, G]_c^\Sigma \rightarrow [F, G]_c^\emptyset$  and  $i : [F, G]^\Sigma \rightarrow [F, G]^\emptyset$  are meet-preserving maps between complete posets. Therefore, they have left adjoints  $F_c^\Sigma : [F, G]_c^\emptyset \rightarrow [F, G]_c^\Sigma$  and  $F^\Sigma : [F, G]^\emptyset \rightarrow [F, G]^\Sigma$ . Moreover, the following square commutes, since their right adjoints do

$$\begin{array}{ccc} [F, G]^\Sigma & \xleftarrow{F^\emptyset} & [F, G]^\emptyset \\ \downarrow L_c^\Sigma & & \downarrow L_c^\emptyset \\ [F, G]_c^\Sigma & \xleftarrow{F_c^\Sigma} & [F, G]_c^\emptyset \end{array}$$

Using this it can be shown that we have a commutative diagram

$$\begin{array}{ccc} [F, G]_l^\Sigma & \longrightarrow & [F, G]_l^\emptyset \\ \downarrow \overline{(-)}^\Sigma & & \downarrow \overline{(-)}^\emptyset \\ [F, G]_c^\Sigma & \xleftarrow{F_c^\Sigma} & [F, G]_c^\emptyset \end{array}$$

For a lax  $\Sigma$ -transformation  $\lambda : F \rightarrow G$ , we know by the hypothesis that  $\overline{\lambda}^\emptyset$  is strict and therefore it is a  $\Sigma$ -transformation. Therefore  $\overline{\lambda}^\Sigma = F_c^\Sigma \overline{\lambda}^\emptyset = \overline{\lambda}^\emptyset$ .  $\square$

<sup>3</sup> Here  $\overline{(-)}^\Sigma : [F, G]_l^\Sigma \rightarrow [F, G]_c^\Sigma$  and  $\overline{(-)}^\emptyset : [F, G]_l^\emptyset \rightarrow [F, G]_c^\emptyset$  refer to the operations described in section 2.

For the rest of this section, let  $\Sigma := \emptyset$ . We will write  $[F, G]_{\bullet}$  to mean  $[F, G]_{\bullet}^{\emptyset}$  and we will refer to  $\emptyset$ -transformation as just *transformations*.

To obtain the constructions for  $\overline{(-)}$  and  $\underline{(-)}$ , we will define new functors  $F_{\#}, F_{\#}^{\#}$  and  $F^{\#}$  and strict transformations

$$\begin{array}{ccc}
 & F & \\
 c' \nearrow & & \nwarrow l' \\
 F_{\#} & & F^{\#} \\
 & c \searrow & \nearrow l \\
 & F_{\#}^{\#} &
 \end{array}$$

such that  $[F_{\#}, G]_s \cong [F, G]_c$ ;  $[F_{\#}^{\#}, G]_s \cong [F, G]$  and  $[F^{\#}, G]_s \cong [F, G]_l$ .

Using these isomorphisms, we can rewrite  $\overline{(-)} : [F, G]_l \rightarrow [F, G]_c$  as

$$[F, G]_l \cong [F^{\#}, G]_s \xrightarrow{-\circ l} [F_{\#}^{\#}, G]_s \xrightarrow{\text{Lan}_{c^-}} [F_{\#}, G]_s \cong [F, G]_c$$

and similarly, we can write  $\underline{(-)} : [F, G]_c \rightarrow [F, G]_l$  as

$$[F, G]_c \cong [F_{\#}, G]_s \xrightarrow{-\circ c} [F_{\#}^{\#}, G]_s \xrightarrow{\text{Ran}_{l^-}} [F^{\#}, G]_s \cong [F, G]_l.$$

If the extensions in these compositions are objectwise, we can give an explicit expression for  $\overline{(-)}$  and  $\underline{(-)}$ . In this section we will discuss conditions for when this is the case.

The strict 2-functor  $F^{\#} : \mathcal{C} \rightarrow \mathbf{Cat}$  is defined by sending every object  $A$  in  $\mathcal{C}$  to the lax coend<sup>4</sup>

$$F^{\#} A := \oint^{B \in \text{ob } \mathcal{C}} \mathcal{C}(B, A) \times FB.$$

The following proposition gives a concrete description of  $F^{\#} A$  for  $A \in \text{ob } \mathcal{C}$ .

**Proposition 4.2.** *Let  $A \in \text{ob } \mathcal{C}$ . Objects of the category  $F^{\#} A$  are pairs  $(B \xrightarrow{g} A, y)$ , where  $g$  is a morphism in  $\mathcal{C}$  and  $y$  is an element of  $FB$ . A morphism from objects  $(B_1 \xrightarrow{g_1} A, y_1)$  to  $(B_2 \xrightarrow{g_2} A, y_2)$ , is a triple  $(s : B_2 \rightarrow B_1, \alpha : g_1 s \Rightarrow g_2, y_1 \rightarrow F s(y_2))$ .*

*Proof.* Note first that this indeed defines a category and let us denote it by  $\mathcal{D}$ . We will now show that  $\mathcal{D}$  satisfies the universal property of lax coends.

<sup>4</sup> Lax coends are explained in the Appendix A

For  $B \in \text{ob } \mathcal{C}$ , there clearly is a functor  $e_B : \mathcal{C}(B, A) \times FB \rightarrow \mathcal{D}$ . For  $s : B_2 \rightarrow B_1$ , we have a 2-cell

$$\begin{array}{ccc} \mathcal{C}(B_1, A) \times FB_2 & \xrightarrow{\mathcal{C}(s, A) \times \text{Id}} & \mathcal{C}(B_2, A) \times FB_2 \\ \downarrow \text{Id} \times Fs & \nearrow e_s & \downarrow e_{B_2} \\ \mathcal{C}(B_1, A) \times FB_1 & \xrightarrow{e_{B_1}} & \mathcal{D} \end{array} .$$

Let  $\tilde{\mathcal{D}}$  be a category and let  $(\tilde{e}_B : \mathcal{C}(B, A) \times FB)_{B \in \text{ob } \mathcal{C}}$  be a collection of functors with for every morphisms  $s : B_2 \rightarrow B_1$  a 2-cell

$$\begin{array}{ccc} \mathcal{C}(B_1, A) \times FB_2 & \xrightarrow{\mathcal{C}(s, A) \times \text{Id}} & \mathcal{C}(B_2, A) \times FB_2 \\ \downarrow \text{Id} \times Fs & \nearrow \tilde{e}_s & \downarrow \tilde{e}_{B_2} \\ \mathcal{C}(B_1, A) \times FB_1 & \xrightarrow{\tilde{e}_{B_1}} & \tilde{\mathcal{P}} \end{array}$$

There is a functor  $e : \mathcal{D} \rightarrow \tilde{\mathcal{D}}$  that sends  $(B \xrightarrow{g} A, y)$  to  $\tilde{e}_B(g, y)$ . This is the unique functor such that  $e \circ e_B = \tilde{e}_B$  for all  $B \in \text{ob } \mathcal{C}$ . This shows the univernal property  $\square$

Let  $[\mathcal{C}, \mathbf{Cat}]_l$  be the strict 2-category of strict 2-functors  $\mathcal{C} \rightarrow \mathbf{Cat}$  and lax transformations and modifications and let  $[\mathcal{C}, \mathbf{Cat}]_s$  be the 2-subcategory of strict 2-functors and strict transformations and modifications. The following result is Theorem 3.16 in [1] together with section 7.1.2 in [4]. In [1] the result follows from a more general theorem. Here we will also give a direct proof.

**Proposition 4.3.** *The inclusion strict 2-functor  $[\mathcal{C}, \mathbf{Cat}]_s \rightarrow [\mathcal{C}, \mathbf{Cat}]_l$  has a left adjoint, which is given by the assignment  $F \mapsto F^\#$ .*

*Proof.* It is enough to show that  $[F^\#, G]_s \cong [F, G]_l$  for all functors  $F$  and  $G$ .

Given a strict transformation  $\tau : F^\# \rightarrow G$ , we can define for every  $A \in \text{ob } \mathcal{C}$  an order-preserving  $\lambda_A : FA \rightarrow GA$  by sending  $x$  to  $\tau_A(1_A, x)$ . These form a lax transformation.

Given a lax transformations  $\lambda : F \rightarrow G$ , we can define a strict transformations  $\tau : F^\# \rightarrow G$ , by the assignment

$$\tau_A(g : B \rightarrow A, y) := \lambda_B(Fg(y)).$$

These define an isomorphism of categories.  $\square$

The counit of the adjunction in Proposition 4.3 is a strict transformation  $l' : F^\# \rightarrow F$  and  $l'_A(g : B \rightarrow A, y) = Fg(y)$  for all  $A \in \text{ob } \mathcal{C}$  and  $(g, y) \in F^\# A$ .

Dually, we define the functor  $F_{\#} : \mathcal{C} \rightarrow \mathbf{Cat}$  by sending every object  $A$  in  $\mathcal{C}$  to the colax coend

$$F_{\#}A := \oint^{B:\mathcal{C}} \mathcal{C}(B, A) \times FB.$$

The following result is the dual of Proposition 4.2.

**Proposition 4.4.** *For  $A \in \text{ob}\mathcal{C}$ . Objects of the category  $F_{\#}A$  are pairs  $(B \xrightarrow{g} A, y)$ , where  $g$  is a morphism in  $\mathcal{C}$  and  $y$  is an element of  $FB$ . A morphism from objects  $(B_1 \xrightarrow{g_1} A, y_1)$  to  $(B_2 \xrightarrow{g_2} A, y_2)$ , is a triple  $(s : B_1 \rightarrow B_2, \alpha : g_1 \Rightarrow g_2 s, Fs(y_1) \rightarrow y_2)$ .*

Let  $[\mathcal{C}, \mathbf{Cat}]_c$  be the strict 2-category of strict 2-functors  $\mathcal{C} \rightarrow \mathbf{Cat}$ , colax transformations and modifications. The dual of Proposition 4.3 is given by the following the result.

**Proposition 4.5.** *The inclusion strict 2-functor  $[\mathcal{C}, \mathbf{Cat}]_s \rightarrow [\mathcal{C}, \mathbf{Cat}]_c$  has a left adjoint, which is given by the assignment  $F \mapsto F_{\#}$ .*

The counit is a strict transformation  $c' : F_{\#} \rightarrow F$ , such that  $c'_A(g, y) = Fg(y)$  for all  $A \in \text{ob}\mathcal{C}$  and  $(g, y) \in F_{\#}A$ .

Finally, consider the functor  $F_{\#}^{\#} : \mathcal{C} \rightarrow \mathbf{Cat}$  that is defined by sending every object  $A$  in  $\mathcal{C}$  to

$$F_{\#}^{\#}A := \sum_{B \in \text{ob}\mathcal{C}} \mathcal{C}(B, A) \times FB.$$

Let  $[\mathcal{C}, \mathbf{Cat}]$  be the 2-category of functors  $\mathcal{C} \rightarrow \mathbf{Cat}$  and transformations and modifications. The following result can be proven in a similar way as the proof of Proposition 4.3 and Proposition 4.5.

**Proposition 4.6.** *The inclusion strict 2-functor  $[\mathcal{C}, \mathbf{Cat}]_s \rightarrow [\mathcal{C}, \mathbf{Cat}]$  has a left adjoint, which is given by the assignment  $F \mapsto F_{\#}^{\#}$ .*

The unit  $F \rightarrow F_{\#}^{\#}$  is an element of  $[F, F_{\#}^{\#}]_l \subseteq [F, F_{\#}^{\#}]$  and therefore it corresponds to a strict transformation  $l : F_{\#}^{\#} \rightarrow F_{\#}$ . We have that  $l_A(g, y) = (g, y)$  for all  $A \in \text{ob}\mathcal{C}$  and  $(g, y) \in F_{\#}^{\#}A$ . Similarly, the unit  $F \rightarrow F_{\#}$  is an element of  $[F, F_{\#}]_c \subseteq [F, F_{\#}]$  and also corresponds to a strict transformations  $c : F_{\#}^{\#} \rightarrow F_{\#}$  such that  $c_A(g : B \rightarrow A, y) = (g, y)$  for all  $A$  in  $\mathcal{C}$  and  $(g, y)$  in  $F_{\#}^{\#}A$ .

Recall that we assumed that  $FA$  is a poset for all objects  $A$  in  $\text{ob}\mathcal{C}$ . As explained above, writing the transformation in terms of  $F_{\#}$  and  $F_{\#}^{\#}$  allows us to describe the operation  $\overline{(-)}$  as a left extension. For a lax extension  $\lambda : F \rightarrow G$ , let  $\tilde{\lambda} : F_{\#}^{\#} \rightarrow G$  be the corresponding strict transformation. For  $A \in \text{ob}\mathcal{C}$  and  $x \in FA$ , we can write

$$\overline{\lambda}_A(x) = \text{Lan}_c(\tilde{\lambda} \circ l)_A(1_A, x).$$

Suppose now that this left extension is objectwise. We then would have that  $\bar{\lambda}_A(x)$  is equal to

$$\begin{aligned} & \text{colim}(c \downarrow (1_A, x) \rightarrow F_{\#}^{\#} A \xrightarrow{l} F_{\#}^{\#} A \xrightarrow{\tilde{\lambda}} GA) \\ &= \bigvee \left\{ \tilde{\lambda}_A(g, y) \mid g : C \rightarrow A; y \in FC \text{ such that } Fg(y) \leq x \right\} \\ &= \bigvee \left\{ Gg(\lambda_C(y)) \mid g : C \rightarrow A; y \in FC \text{ such that } Fg(y) \leq x \right\} \end{aligned}$$

In what follows, we will give conditions for when this is indeed the case.

**Proposition 4.7.** *For  $\lambda \in [F, G]^{\Sigma}$ . For  $A \in \text{ob } \mathcal{C}$ , define  $\tau_A : FA \rightarrow GA$  by*

$$\tau_A(x) := \bigvee \left\{ Gg(\lambda_C(y)) \mid g : C \rightarrow A; y \in FC \text{ such that } Fg(y) \leq x \right\},$$

for  $x \in FA$ . If  $\tau$  is colax, then  $\tau = L_c \lambda$ . If moreover,  $\lambda$  is lax, then  $\tau = \bar{\lambda}$ .

*Proof.* Let  $\tilde{\lambda}$  be the strict transformation  $F_{\#}^{\#} \rightarrow G$ , corresponding to  $\lambda$  and let  $\tilde{\tau}$  be the strict transformation  $F_{\#} \rightarrow G$  corresponding to  $\tau$ . We want to show that  $\tilde{\tau}$  is the left extension of  $\tilde{\lambda}$  along  $c$ . For  $(f : C \rightarrow A, y) \in F_{\#}^{\#} A$ ,

$$\begin{aligned} (\tilde{\tau} \circ c)_A(f, y) &= Gf(\tau_C(y)) = Gf \bigvee \left\{ Gg(\lambda_{C'}(y')) \mid g : C' \rightarrow C; y' \in FC' \text{ such that } Fg(y') \leq y \right\} \\ &\geq \bigvee \left\{ Gf(Gg(\lambda_{C'}(y'))) \mid g : C' \rightarrow C; y' \in FC' \text{ such that } Fg(y') \leq y \right\} \\ &\geq Gf(\lambda_C(y)) = \tilde{\lambda}_A(f, y), \end{aligned}$$

which means that  $\tilde{\tau} \circ c \geq \tilde{\lambda}$ .

Suppose now that  $\nu : F_{\#} \rightarrow G$  is a strict transformation such that  $\nu \circ c \geq \tilde{\lambda}$ . Let  $(f : C \rightarrow A, y) \in F_{\#}^{\#} A$ . Consider  $g : C' \rightarrow C$  and  $y' \in FC'$  such that  $Fg(y') \leq y$ , then  $(g, y') \in F_{\#}^{\#} C$  and  $c_C(g, y') \leq (1_C, y)$  in  $F_{\#} C$  and therefore

$$\nu_C(1_C, y) \geq (\nu \circ c)_C(g, y') \geq \tilde{\lambda}_C(g, y') = Gg\lambda_{C'}(y').$$

Taking the supremum over all such  $(g, y')$ , we have

$$\nu_C(1_C, y) \geq \tau_C(y)$$

and by applying  $Gf$  to both sides,

$$\nu_A(f, y) = Gf\nu_A(1_C, y) \geq Gf\tau_C(y) = \tilde{\tau}_A(f, y).$$

Here we used that  $\nu$  is strict and the definition  $\tilde{\tau}$ . This shows that  $\tilde{\tau}$  is the left extension of  $\tilde{\lambda} \circ l$  along  $c$ .  $\square$

Let  $A \in \text{ob } \mathcal{C}$  and  $x \in FA$ . The objects of the category  $l'_A \downarrow x$  are pairs  $(g : C \rightarrow A, y)$  where  $g$  is a morphism in  $\mathcal{C}$  and  $y \in FC$  such that  $Fg(y) \leq x$ . A morphism from  $(g_1 : C_1 \rightarrow A, y_1)$  to  $(g_2 : C_2 \rightarrow A, y_2)$ , is a morphism  $s : C_2 \rightarrow C_1$  together with a 2-cell  $\alpha : g_1 s \Rightarrow g_2$  such that  $y_1 \leq Fs(y_2)$ . Let  $(l'_A \downarrow x)_=$  be the subcategory of objects  $(g, y)$  such that  $Fg(y) = x$ .

**Theorem 4.8.** *Let  $\kappa$  be a regular cardinal. Suppose that*

(G<sub>1</sub>)  *$Gf$  preserves  $\kappa$ -directed joins for all morphisms  $f$  in  $\mathcal{C}$ ,*

(F<sub>1</sub>)  *$(l'_A \downarrow x)_= \subseteq l'_A \downarrow x$  is weakly final<sup>5</sup>,*

(F<sub>2</sub>)  *$\mathcal{C}$  has and  $F$  preserves  $\kappa$ -wide pullbacks.*

*Then  $\bar{\lambda}$  is a strict transformation and for  $A \in \text{ob } \mathcal{C}$  and  $x \in FA$ ,*

$$\bar{\lambda}_A(x) = \bigvee \{Gg(\lambda_C(y)) \mid g : C \rightarrow A; y \in FC \text{ such that } Fg(y) \leq x\}.$$

*Proof.* For  $A \in \text{ob } \mathcal{C}$  and  $x \in FA$  denote

$$S_{A,x} := \{Gg\lambda_C(y) \mid g : C \rightarrow A; y \in FC \text{ such that } Fg(y) \leq x\}$$

and write  $\tau_A(x) := \bigvee S_{A,x}$ . Let  $I$  be a set such that  $|I| < \kappa$ . And consider a collection  $(g_i : C_i \rightarrow A, y_i)_{i \in I}$ , where  $y_i \in FC_i$  and  $Fg_i(y_i) \leq x$  for all  $i \in I$ . From (F<sub>1</sub>) it follows that for every  $i \in I$ , there exists a  $(\tilde{g}_i : \tilde{C}_i \rightarrow A, \tilde{y}_i)$ , where  $\tilde{y}_i \in F\tilde{C}_i$  such that  $F\tilde{g}_i(\tilde{y}_i) = x$  and a map  $s_i : \tilde{C}_i \rightarrow C_i$  and a 2-cell  $\alpha_i : g_i s_i \Rightarrow \tilde{g}_i$  such that  $y_i \leq Fs_i(\tilde{y}_i)$ .<sup>6</sup> The maps  $(\tilde{g}_i)_{i \in I}$  form a  $\kappa$ -wide pullback diagram. Let  $C$  be the pullback of this diagram and let  $(p_i : C \rightarrow \tilde{C}_i)_{i \in I}$  be the collection of projections.

$$\begin{array}{ccc}
 & \tilde{C}_i & \\
 p_i \nearrow & & \searrow \tilde{g}_i \\
 C & & A \\
 p_j \searrow & & \nearrow \tilde{g}_j \\
 & \tilde{C}_j & \\
 & \vdots & \\
 & \tilde{C}_i & 
 \end{array}$$

<sup>5</sup> Weakly final functors are explained in Appendix B.

<sup>6</sup> Note that we might need the axiom of choice here.



Because  $F\tilde{g}_i(\tilde{y}_i) = x$  for all  $i \in I$ , we have that  $(\tilde{y}_i)_{i \in I} \in \times_{FA} FC_i$ . Since  $F$  preserves this limit, there exists a unique  $y \in FC$  such that  $Fp_i(y) = \tilde{y}_i$ . Let  $g : C \rightarrow A$  be the morphism that is equal to  $\tilde{g}_i p_i$  for all  $i \in I$ . We find for all  $i \in I$  that

$$\begin{aligned} Gg_i(\lambda_{C_i}(y_i)) &\leq Gg_i(\lambda_{C_i}(Fs_i(\tilde{y}_i))) \\ &= Gg_i(\lambda_{C_i}(Fs_i Fp_i(y))) \\ &\leq (Gg_i s_i p_i)(\lambda_C(y)) \\ &\leq (G\tilde{g}_i s_i)(\lambda_C(y)) = Gg\lambda_C(y) \end{aligned}$$

We conclude that  $S$  is  $\kappa$ -directed. Using  $(G_1)$ , for a morphism  $f : A \rightarrow B$ , we see that

$$\begin{aligned} Gf\tau_A(x) &= \bigvee \{Gf(Gg(\lambda_C(y))) \mid g : C \rightarrow A; y \in FC \text{ such that } Fg(y) \leq x\} \\ &\leq \bigvee S_{B, Ff(x)} = \tau_B(Ff(x)). \end{aligned}$$

This shows that  $\tau$  is a colax transformation. It is now not difficult to see that  $\bar{\lambda} = \tau$ . This shows the second claim.

To prove the first claim we need to show that

$$L := \bigvee \{Gf(Gg(\lambda_C(y))) \mid g : C \rightarrow A; y \in FC \text{ such that } Fg(y) \leq x\} \geq \bigvee S_{B, Ff(x)}$$

Consider a morphism  $g : C \rightarrow B$  and an element  $y \in FC$  such that  $Fg(y) \leq Ff(x)$ . By  $(F_1)$  there exists maps  $s : \tilde{C} \rightarrow C$  and  $\tilde{g} : \tilde{C} \rightarrow B$ , an element  $\tilde{y} \in F\tilde{C}$  and a 2-cell  $\alpha : gs \Rightarrow \tilde{g}$  such that  $y \leq Fs(\tilde{y})$  and  $F\tilde{g}(\tilde{y}) = x$ .

Let  $P$  be the pullback of  $f$  and  $\tilde{g}$  and let  $p_1 : P \rightarrow A$  and  $p_2 : P \rightarrow \tilde{C}$  be the projection maps.

$$\begin{array}{ccc} P & \xrightarrow{p_1} & A \\ \downarrow & \lrcorner & \downarrow \\ p_2 & & f \\ \downarrow & & \downarrow \\ \tilde{C} & \xrightarrow{\tilde{g}} & B \end{array}$$

Clearly  $(x, \tilde{y})$  is an element of  $FA \times_{FB} F\tilde{C}$ . Because  $F$  preserves pullbacks, there exists a unique  $z \in F(A \times_B \tilde{C})$  such that  $Fp_1(z) = x$  and  $Fp_2(z) = \tilde{y}$ . We know have

$$\begin{aligned} Gg(\lambda_C(y)) &\leq Gg(\lambda_C(Fs(\tilde{y}))) \\ &= Gg(\lambda_C(Fs Fp_2(z))) \\ &\leq (Gg s p_2)(\lambda_P(z)) \\ &\leq (G\tilde{g} p_2)(\lambda_P(z)) \\ &= Gf(Gp_1(\lambda_P(z))) \leq L. \end{aligned}$$

Since  $(g, y)$  was arbitrary, this shows that  $\bigvee S_{B, Ff(x)} \leq L$ . This proves that  $\bar{\lambda}$  is a strict transformation.  $\square$

**Remark 4.9.** In the case of Theorem 4.8,  $(F, G, \emptyset)$  satisfies the strictness conditions. Therefore, Proposition 4.1 can be applied.

## 5 The extension formula

Let  $\mathcal{C}$  be a small strict 2-category and let  $\Sigma$  be a collection of morphisms in  $\mathcal{C}$ . Let  $F, G, H$  be strict 2-functors  $\mathcal{C} \rightarrow \mathbf{Cat}$ . Suppose that  $GA$  is a complete poset for all  $A \in \text{ob } \mathcal{C}$  and that  $Gf$  preserves meets and joins for all  $f \in \Sigma$ .

In this section we will give sufficient conditions for extensions to be objectwise. This will allow us to give concrete formulas for the extensions of (co)lax  $\Sigma$ -transformations. For the rest of this section, let  $F, G, H$  be strict 2-functors  $\mathcal{C} \rightarrow \mathbf{Cat}$  and let  $\iota : F \rightarrow H$  be a strict transformation.

We will start with a result on the extensions of (general)  $\Sigma$ -transformations.

**Theorem 5.1.** *Suppose that  $Ff$  and  $Hf$  have left adjoints  $(Ff)_*$  and  $(Hf)_*$  for all  $f : A \rightarrow B$  in  $\Sigma$ , such that the following square commutes*

$$\begin{array}{ccc} FA & \longleftarrow (Ff)_* & \text{---} & FB \\ \downarrow \iota_A & & & \downarrow \iota_B \\ HA & \longleftarrow (Hf)_* & \text{---} & HB \end{array}$$

*Then left extensions of  $\Sigma$ -transformations along  $\iota$  are objectwise.*

*Proof.* Let  $\tau : F \rightarrow G$  be a  $\Sigma$ -transformation. It is enough to show that the transformation  $(\text{Lan}_{\iota_A} \tau_A)_{A \in \text{ob } \mathcal{C}}$  is a  $\Sigma$ -transformation. Let  $f : A \rightarrow B$  be in  $\Sigma$  and let  $x \in HA$ ,

$$\begin{aligned} Gf \text{Lan}_{\iota_A} \tau_A(x) &= Gf \text{colim}(\iota_A \downarrow x \rightarrow FA \xrightarrow{\tau_A} GA) \\ &= \text{colim}(\iota_A \downarrow x \rightarrow FA \xrightarrow{\tau_A} GA \xrightarrow{Gf} GB) \\ &= \text{colim}(\iota_A \downarrow x \rightarrow \iota_B \downarrow Hf(x) \rightarrow FB \xrightarrow{\tau_B} GB) \\ &= \text{colim}(\iota_B \downarrow Hf(x) \rightarrow FB \xrightarrow{\tau_B} GB) = \text{Lan}_{\iota_B} \tau_B(Hf(x)) \end{aligned}$$

The second equality follows from the fact that  $Gf$  preserves joins, since  $f \in \Sigma$ . Because  $Gf\tau_A = \tau_B Hf$ , we have the third equality. Using the hypothesis, it can be shown that

$$\iota_A \downarrow x \rightarrow \iota_B \downarrow Hf(x) \cong \iota_A (Ff)_* \downarrow x$$

is right adjoint and therefore final. The fourth equality now follows. □

**Theorem 5.2.** *Let  $\kappa$  be a regular cardinal.*

(G1) Suppose that  $Gf$  preserves  $\kappa$ -directed joins for all morphisms  $f$  in  $\mathcal{C}$ .

( $\iota$ 1) Suppose that  $\iota_A$  is  $\kappa$ -flat for all  $A \in \text{ob } \mathcal{C}$ , i.e.  $\iota_A \downarrow x$  is  $\kappa$ -directed for all  $x \in FA$ .

( $\iota$ 2)  $Ff$  and  $Hf$  have left adjoints  $(Ff)_*$  and  $(Hf)_*$  for all  $f : A \rightarrow B$  in  $\Sigma$ , such that the following square commutes

$$\begin{array}{ccc} FA & \leftarrow (Ff)_* & FB \\ \downarrow \iota_A & & \downarrow \iota_B \\ HA & \leftarrow (Hf)_* & HB \end{array}$$

Then left extensions of colax  $\Sigma$ -transformations along  $\iota$  are objectwise.

*Proof.* Let  $\sigma : F \rightarrow G$  be a colax  $\Sigma$ -transformation. It is enough to show that the transformation  $(\text{Lan}_{\iota_A} \sigma_A)_{A \in \text{ob } \mathcal{C}}$  is a colax  $\Sigma$ -transformation. Let  $A \rightarrow B$  be a morphism in  $\text{ob } \mathcal{C}$  and let  $x \in HA$ ,

$$\begin{aligned} Gf \text{Lan}_{\iota_A} \sigma_A(x) &= Gf \text{colim}(\iota_A \downarrow x \rightarrow FA \xrightarrow{\sigma_A} GA) \\ &= \text{colim}(\iota_A \downarrow x \rightarrow FA \xrightarrow{\sigma_A} GA \xrightarrow{Gf} GB) \\ &\leq \text{colim}(\iota_A \downarrow x \rightarrow \iota_B \downarrow Hf(x) \rightarrow FB \xrightarrow{\sigma_B} GB) \\ &\leq \text{colim}(\iota_B \downarrow Hf(x) \rightarrow FB \xrightarrow{\sigma_B} GB) = \text{Lan}_{\iota_B} \sigma_B(Hf(x)) \end{aligned}$$

In the second equality we use that  $\iota_A \downarrow x$  is  $\kappa$ -directed and that  $Gf$  preserves  $\kappa$ -directed joins. The inequality in the third line follows from the fact that  $\sigma$  is colax and the last inequality is induced by the inclusion  $\iota_A \downarrow x \rightarrow \iota_B \downarrow Hf(x)$ . This shows that the transformation is colax.

To show that the transformation is a  $\Sigma$ -transformation suppose that  $f \in \Sigma$ . In this case the first inequality becomes an equality because  $\sigma$  is a  $\Sigma$ -transformation and therefore  $Gf \sigma_A = \Sigma_B Ff$ . From ( $\iota$ 2) it is straightforward to check that

$$\iota_A \downarrow x \rightarrow \iota_B \downarrow Hf(x) \cong \iota_A (Ff)_* \downarrow x$$

is right adjoint and therefore final. It follows that the second inequality is an equality in this case.  $\square$

## 6 Premeasures as transformations

In this section we will represent certain premeasures by certain transformations between functors. Roughly speaking, colax and lax transformation correspond to inner and outer

premeasures and strict transformations correspond to measures. Using these representations, we can apply the results from the previous sections to obtain results and formulas for joins and meets of certain premeasures and for operations to turn a certain kind of premeasure in a different kind in a universal way.

We will study two approaches. In the first approach we will look at functors whose domain is  $\mathbf{Set}_c^f$ , a subcategory of countable sets and functions. In the second approach we use the category  $\mathbf{Part}_c^f$ , a subcategory of countable sets and *partial* functions. The advantage of the first way of representing premeasures is that  $\mathbf{Set}_c^f$  is slightly easier to work with than  $\mathbf{Part}_c^f$ . The second approach is more flexible to obtain variations of the representation result. For example, if we restrict to *probability* premeasures, we need to use the second point of view to represent inner and outer probability premeasures.

In this section we will use the notation  $\uplus$  and  $\cup$  to emphasize that we are working with *disjoint* unions of subsets.

## 6.1 Premeasures

A **premeasurable space** is a set  $X$  set together with an algebra  $\mathcal{B}$  of subsets, i.e. a collection of subsets closed under finite unions and complements. Let  $(X_1, \mathcal{B}_1)$  and  $(X_2, \mathcal{B}_2)$  be premeasurable space. A map  $f : X_1 \rightarrow X_2$  is called **premeasurable** if  $f^{-1}(\mathcal{B}_2) \subseteq \mathcal{B}_1$ . We denote the category of premeasurable spaces and premeasurables maps by  $\mathbf{PreMble}$  and interpret it as a strict 2-category where the hom-categories are discrete.

For a countable set  $A$ , let  $\mathcal{F}_A$  be the algebra of finite and cofinite sets. A map  $f : A \rightarrow B$  is called **finite** if  $f^{-1}(\mathcal{F}_B) \subseteq \mathcal{F}_A$ . Let  $\mathbf{Set}_c^f$  be the category of countable sets and finite maps, which we interpret as a strict 2-category. Endowing a countable set  $A$  with the algebra  $\mathcal{F}_A$  of finite and cofinite sets, yields a premeasurable space. This gives a functor  $i : \mathbf{Set}_c^f \rightarrow \mathbf{PreMble}$ . Now define the strict 2-functor  $F_{\mathcal{B}}$  as

$$\mathbf{Set}_c^s \xrightarrow{\mathbf{PreMble}((X, \mathcal{B}), i-)} \mathbf{Cat}.$$

For a countable set  $A$ ,  $F_{\mathcal{B}}A$  can be identified with the set of  $A$ -indexed premeasurable partitions of  $X$ .

Let  $A$  be a countable set. The set  $[0, \infty]^A$  together with the pointwise order, is a poset which we will denote by  $GA$ . For a finite map  $f : A \rightarrow B$  of countable sets, the assignment

$$(p_a)_{a \in A} \mapsto \left( \sum_{f(a)=b} p_a \right)_{b \in B}$$

defines an order-preserving map  $Gf : GA \rightarrow GB$ . We obtain a strict 2-functor  $G : \mathbf{Set}_c^s \rightarrow \mathbf{Cat}$ .

A **premeasure** on a premeasurable space  $(X, \mathcal{B})$  is a map  $\mu : \mathcal{B} \rightarrow [0, \infty]$  such that  $\mu(\uplus_{n=1}^{\infty} B_n) = \sum_{n=1}^{\infty} \mu(B_n)$  for pairwise disjoint  $(B_n)_{n=1}^{\infty}$  in  $\mathcal{B}$  such that  $\uplus_{n=1}^{\infty} B_n$  is also

an element of  $\mathcal{B}$ . For premeasures  $\mu_1$  and  $\mu_2$ , we write  $\mu_1 \leq \mu_2$  if  $\mu_1(B) \leq \mu_2(B)$  for all  $B \in \mathcal{B}$ . This turns the set of premeasures on  $(X, \mathcal{B})$  into a poset, which we will denote by  $M(X, \mathcal{B})$ .

For a premeasure  $\mu$  and a countable set  $A$ , define  $\tau_A^\mu : F_{\mathcal{B}}A \rightarrow GA$  by the assignment

$$(B_a)_{a \in A} \mapsto (\mu(B_a))_{a \in A}.$$

**Proposition 6.1.** *Let  $\mu$  be a premeasure, then  $(\tau_A)_A$  is a strict transformation  $F_{\mathcal{B}} \rightarrow G$ .*

*Proof.* Let  $f : A \rightarrow B$  be a finite map. For  $(B_a)_{a \in A} \in F_{\mathcal{B}}A$  and  $b \in B$

$$Gf(\tau_A((B_a)_a))_b = \sum_{f(a)=b} \mu(B_a) = \mu \left( \bigsqcup_{f(a)=b} B_a \right).$$

Here we used the  $\sigma$ -additivity of  $\mu$ . □

If  $\mu_1 \leq \mu_2$ , then clearly  $\tau^{\mu_1} \leq \tau^{\mu_2}$ . It follows that the assignment  $\mu \mapsto \tau^\mu$  defines an order-preserving map  $\tau^{(-)} : M(X, \mathcal{B}) \rightarrow [F_{\mathcal{B}}, G]_s$ .

**Theorem 6.2.** *The map  $\tau^{(-)}$  is an isomorphism of posets.*

*Proof.* Let  $\tau : F_{\mathcal{B}} \rightarrow G$  be a strict transformation. For  $n \geq 1$ , write  $\mathbf{n} := \{0, 1, \dots, n-1\}$ . For  $B \in \mathcal{B}$ , define a map  $\mu_\tau : \mathcal{B} \rightarrow [0, \infty]$  by

$$\mu_\tau(B) := \tau_2(B^C, B)_1.$$

We will now show that this map is a premeasure. Let  $t : \mathbf{1} \rightarrow \mathbf{2}$  be the map that sends 0 to 0. We find

$$\mu_\tau(\emptyset) = \tau_2(X, \emptyset)_1 = \tau_2(F_{\mathcal{B}}t(X))_1 = Gt(\tau_1(X))_1 = 0.$$

Now consider  $(B_n)_{n=1}^\infty$  pairwise disjoint elements in  $\mathcal{B}$  such that  $B := \bigsqcup_{n=1}^\infty B_n$  is also an element in  $\mathcal{B}$ . Writing  $B_0 := B^c$ , yields an element  $(B_n)_{n \in \mathbb{N}} \in F_{\mathcal{B}}\mathbb{N}$ . For an element  $n \geq 1$ , let  $s_n : \mathbb{N} \rightarrow \mathbf{2}$  be the finite map that sends  $n$  to 1 and every other element to 0. Furthermore, define a map  $s_0 : \mathbb{N} \rightarrow \mathbf{2}$  that sends 0 to 0 and every other element to 1. This map is also finite.

For  $n \in \mathbb{N}$ , we have a commutative diagram

$$\begin{array}{ccc} F_{\mathcal{B}}\mathbb{N} & \xrightarrow{\tau_{\mathbb{N}}} & G\mathbb{N} \\ \downarrow F s_n & & \downarrow G s_n \\ F_{\mathcal{B}}\mathbf{2} & \xrightarrow{\tau_2} & G\mathbf{2} \end{array}$$

It follows that

$$\begin{aligned}\mu_\tau(B) &= \tau_{\mathbf{2}}(F_{\mathcal{B}}s_0((B_m)_{m \in \mathbb{N}}))_1 \\ &= Gs_0(\tau_{\mathbb{N}}((B_m)_{m \in \mathbb{N}}))_1 \\ &= \sum_{n=1}^{\infty} \tau_{\mathbb{N}}((B_m)_{m \in \mathbb{N}})_n.\end{aligned}$$

For  $n \geq 1$ , we find

$$\begin{aligned}\mu_\tau(B_n) &= \tau_{\mathbf{2}}(F_{\mathcal{B}}s_n((B_m)_{m \in \mathbb{N}}))_1 \\ &= Gs_n(\tau_{\mathbb{N}}((B_m)_{m \in \mathbb{N}}))_1 \\ &= \tau_{\mathbb{N}}((B_m)_{m \in \mathbb{N}})_n.\end{aligned}$$

Combining the above gives  $\sum_{n=1}^{\infty} \mu_\tau(B_n) = \mu_\tau(B)$ , which shows that  $\mu$  is a premeasure. Sending  $\tau$  to  $\mu_\tau$  defines an order-preserving map  $\mu_{(-)} : [F_{\mathcal{B}}, G]_s \rightarrow M(X, \mathcal{B})$ .

Let  $A$  be a countable set and let  $a_0 \in A$ . Consider the finite map  $s_{a_0} : A \rightarrow \mathbf{2}$  that sends  $a_0$  to 1 and every other element to 0. For a strict transformation  $\tau : F_{\mathcal{B}} \rightarrow G$  and for  $(B_a)_{a \in A} \in F_{\mathcal{B}}A$ ,

$$\tau_A^{\mu_\tau}((B_a)_a)_{a_0} = \mu_\tau(B_{a_0}) = \tau_{\mathbf{2}}(F_{\mathcal{B}}s_{a_0}((B_a)_a))_1 = Gs_{a_0}(\tau_A((B_a)_a))_1 = \tau_A((B_a)_a)_{a_0}$$

This shows that  $\tau^{\mu_\tau} = \tau$ . For a premeasure  $\mu$  and  $B \in \mathcal{B}$ ,

$$\mu_{\tau^\mu}(B) = \tau_{\mathbf{2}}^\mu(B^C, B)_1 = \mu(B).$$

We conclude that  $\mu_{\tau^\mu} = \mu$ . □

**Lemma 6.3.** *The category  $\mathbf{Set}_c^f$  has pullbacks and  $i : \mathbf{Set}_c^f \rightarrow \mathbf{PreMble}$  preserves those.*

*Proof.* Let  $s : A \rightarrow C$  and  $t : B \rightarrow C$  be finite map of countable sets. Define  $P := \{(a, b) \mid s(a) = t(b)\}$  and let  $p_1 : P \rightarrow A$  and  $p_2 : P \rightarrow B$  be the projection maps. Because  $p_1^{-1}(\{a\}) = t^{-1}(s(a))$  for all  $a \in A$ ,  $p_1$  is a finite map. Similarly,  $p_2$  is a finite map. It is now easy to see that  $(P, p_1, p_2)$  forms the pullback of  $s$  and  $t$  in  $\mathbf{Set}_c^f$ .

To show that  $i$  preserves this pullback, we need to show that the algebra generated by  $p_1$  and  $p_2$  contains all singletons. But this follows immediately from the fact that  $p_1^{-1}(\{a\}) \cap p_2^{-1}(\{b\}) = \{(a, b)\}$  for all  $(a, b) \in P$ . □

**Proposition 6.4.** *For  $f : A \rightarrow B$  in  $\mathbf{Set}_c^f$ ,  $Gf$  preserves directed joins.*

*Proof.* Let  $(p^i)_{i \in I}$  be a directed collection of elements in  $GA$ . Define  $p_a := \sup_{i \in I} p_a^i$ . For  $b \in B$ , we always have that

$$Gf((p_a)_a)_b = \sum_{f(a)=b} p_a \geq \sup_{i \in I} \sum_{f(a)=b} p_a^i = \sup_{i \in I} Gf(p^i).$$

To prove the other inequality, consider an element  $b \in B$ . Suppose first that there exists  $a \in f^{-1}(\{b\})$  such that  $p_a = \infty$ . In that case, for every  $N \geq 1$ , there exists  $i_N \in I$  such that  $N \leq p_a^{i_N}$ . In that case

$$N \leq p_a^{i_N} \leq \sum_{f(a)=b} p_a^{i_N} \leq \sup_{i \in I} \sum_{f(a)=b} p_a^i.$$

By taking  $N \rightarrow \infty$ , we conclude that  $\sup_{i \in I} \sum_{f(a)=b} p_a^i = \infty$ , which shows the equality.

Suppose now that  $p_a < \infty$  for all  $a \in f^{-1}(\{b\})$  and consider a finite set  $A' \subseteq f^{-1}(\{b\})$  and  $\epsilon > 0$ . For every  $a \in A'$  there exists  $i_a \in I$  such that

$$p_a - \frac{\epsilon}{|A'|} \leq p_a^{i_a}.$$

Because  $(p^i)_{i \in I}$  is directed and  $A'$  is finite, there exists an  $i_0 \in I$  such that  $p^{i_a} \leq p^{i_0}$  for all  $a \in A'$ . Summing over  $A'$  leads to

$$\sum_{a \in A'} p_a - \epsilon \leq \sum_{a \in A'} p_a^{i_a} \leq \sum_{a \in A'} p_a^{i_0} \leq \sum_{f(a)=b} p_a^{i_0} \leq \sup_{i \in I} \sum_{f(a)=b} p_a^i.$$

The claim now follows by first taking  $\epsilon \rightarrow 0$  and then taking the supremum over all finite subsets of  $f^{-1}(\{b\})$ .  $\square$

**Proposition 6.5.** *The triple  $(F_{\mathcal{B}}, G, \emptyset)$  satisfies the strictness property. Moreover, for  $\lambda \in [F_{\mathcal{B}}, G]_l$ ,*

$$\mu^{\bar{\lambda}}(B) = \sup \left\{ \sum_{a \in A} \lambda_A(C_a) \mid \bigcup_{a \in A} B_a = B \right\},$$

for all  $B \in \mathcal{B}$ .

*Proof.* By Proposition 6.4,  $Gf$  preserves directed joins for all  $f \in \mathbf{Set}_c^f$ . For a countable set  $A$  and  $x \in F_{\mathcal{B}}(A)$ ,  $(l'_A \downarrow x)_= = l'_A \downarrow x$ , since  $F_{\mathcal{B}}(A)$  is a discrete poset. Finally, Lemma 6.3 and the fact that hom-functors preserve pullbacks imply that  $F_{\mathcal{B}}$  preserves pullbacks. The result now follows by Theorem 4.8.  $\square$

## 6.2 Outer/inner premeasures

Let  $(X_1, \mathcal{B}_1)$  and  $(X_2, \mathcal{B}_2)$  be premeasurable spaces. A partial map  $f : X_1 \rightarrow X_2$  is **premeasurable** if  $f^{-1}(B) \in \mathcal{B}_1$  for all  $B \in \mathcal{B}_2$ . For premeasurable partial maps  $f_1, f_2 : A \rightarrow B$ , we write  $f_1 \leq f_2$  if  $\text{dom} f_1 \subseteq \text{dom} f_2$  and  $f_1(x) = f_2(x)$  for all  $x \in \text{dom} f_1$ . The strict 2-category of premeasurable spaces and premeasurable partial maps, which are ordered, is denoted as **PartPreMble**.

A partial map  $f : A \rightarrow B$  of countable sets is called **finite** if  $f^{-1}(B') \in \mathcal{F}_A$  for all  $B' \in \mathcal{F}_B$ . We write **Part** $_c^s$  for the strict 2-category of countable sets and partial finite maps,

which are ordered. The collection of injective partial finite maps is denoted by  $\Sigma$ . Sending a countable set  $A$  to  $(A, \mathcal{F}_A)$  again yields a strict 2-functor  $j : \mathbf{Part}_c^f \rightarrow \mathbf{PartPreMble}$ . For a premeasurable space  $(X, \mathcal{B})$ , define the strict 2-functor  $F_{\mathcal{B}}$  as

$$\mathbf{Part}_c^f \xrightarrow{\mathbf{PartPreMble}(X, \mathcal{B}, j^-)} \mathbf{Cat}.$$

For a countable set  $A$ , the poset  $F_{\mathcal{B}}A$  is the set of  $A$ -indexed collections of pairwise disjoint premeasurable subsets, whose union is again premeasurable, endowed with the pointwise order.

Let  $A$  be a countable set. Let  $GA$  be the poset  $[0, \infty]^A$ , where the order is defined pointwise. For a finite partial map  $f : A \rightarrow B$ , sending  $(p_a)_{a \in A}$  to

$$\left( \sum_{f(a)=b} p_a \right)_{b \in B}$$

gives an order-preserving map  $Gf : GA \rightarrow GB$ . This yields a strict 2-functor  $G : \mathbf{Part}_c^s \rightarrow \mathbf{Cat}$ .

An order-preserving map  $\mu : \mathcal{B} \rightarrow [0, \infty]$  such that  $\mu(\emptyset) = 0$  is called an **outer premeasure** if  $\mu(B) \leq \sum_{n=1}^{\infty} \mu(B_n)$  for pairwise disjoint subsets  $(B_n)_{n=1}^{\infty}$  in  $\mathcal{B}$  such that  $B := \bigcup_{n=1}^{\infty} B_n$  is also an element in  $\mathcal{B}$ . It is called an **inner premeasure** if  $\mu(B) \geq \sum_{n=1}^{\infty} \mu(B_n)$  for pairwise disjoint subsets  $(B_n)_{n=1}^{\infty}$  in  $\mathcal{B}$  such that  $B := \bigcup_{n=1}^{\infty} B_n$  is also an element in  $\mathcal{B}$ . The set of outer (inner) premeasures together with objectwise order is denoted by  $M_{\leq}(X, \mathcal{B})$  ( $M_{\geq}(X, \mathcal{B})$ ).

For an order-preserving map  $\mu : \mathcal{B} \rightarrow [0, \infty]$  such that  $\mu(\emptyset) = 0$ , we define maps  $(\tau_A^\mu : F_{\mathcal{B}}A \rightarrow GA)_A$  by the assignment

$$(E_a)_{a \in A} \mapsto \mu(E_a)_{a \in A}.$$

**Proposition 6.6.** *If  $\mu$  is an outer (inner) premeasure, then  $\tau^\mu$  is a lax (colax)  $\Sigma$ -transformation.*

*Proof.* For a partial finite map  $f : A \rightarrow B$  and  $(E_a)_{a \in A}$  and  $b \in B$ , we have

$$Gf(\tau_A^\mu((E_a)_a))_b = \sum_{f(a)=b} \mu(E_a) \geq \mu \left( \bigcup_{f(a)=b} E_a \right) = \tau^\mu(F_{\mathcal{B}}((E_a)_a))_b.$$

If  $f$  is injective, then  $f^{-1}(\{b\})$  is either a singleton or the empty set; in both cases the inequality becomes an equality.  $\square$

Clearly, if  $\mu$  is a premeasure,  $\tau^\mu$  is a strict transformation. This leads to following



order-preserving maps.

$$\begin{array}{ccccc}
[F_{\mathcal{B}}, G]_l^\Sigma & \longleftarrow & [F_{\mathcal{B}}, G]_s & \longrightarrow & [F_{\mathcal{B}}, G]_c^\Sigma \\
\uparrow \lambda^{(-)} & & \uparrow \tau^{(-)} & & \uparrow \sigma^{(-)} \\
M_{\leq}(X, \mathcal{B}) & \longleftarrow & M(X, \mathcal{B}) & \longrightarrow & M_{\geq}(X, \mathcal{B})
\end{array}$$

**Theorem 6.7.** *The maps  $\lambda^{(-)}$ ,  $\tau^{(-)}$  and  $\sigma^{(-)}$  are isomorphisms of posets.*

*Proof.* Let  $\lambda : F_{\mathcal{B}} \rightarrow G$  be a lax  $\Sigma$ -transformation. For  $n \geq 1$ , define  $\mathbf{n} := \{0, 1, \dots, n-1\}$ . For  $B \in \mathcal{B}$ , define

$$\mu_\lambda(B) := \lambda_{\mathbf{1}}(B).$$

We will now show that this is an outer premeasure. Let  $s$  be the unique injective map  $\emptyset \rightarrow \mathbf{1}$  and let  $*$  be the unique element in  $F_{\mathcal{B}}\emptyset$ . We find

$$\mu_\lambda(\emptyset) = \lambda_{\mathbf{1}}(\emptyset) = \lambda_{\mathbf{1}}(F_{\mathcal{B}}(*)) = Gs(\lambda_\emptyset(*)) = 0.$$

Let  $x := (B_n)_{n \in \mathbb{N}}$  be a collection of pairwise disjoint premeasurable subsets such that  $B := \bigsqcup_{n \in \mathbb{N}} B_n$  is also an element of  $\mathcal{B}$ . Let  $s : \mathbb{N} \rightarrow \mathbf{1}$  be the map that sends every element to 0 and for  $m \in \mathbb{N}$ , let  $s_m : \mathbb{N} \rightarrow \mathbf{1}$  be the partial map that is only defined on  $\{m\}$ . Note that these maps are finite partial maps and that  $s_m$  is injective for all  $m \in \mathbb{N}$ . We have

$$\begin{aligned}
\mu_\lambda(B) &= \lambda_{\mathbf{1}}(F_{\mathcal{B}}s(x)) \\
&\leq Gs(\lambda_{\mathbb{N}}(x)) = \sum_{n \in \mathbb{N}} \lambda_{\mathbb{N}}(x)_n
\end{aligned}$$

and for every  $n \in \mathbb{N}$ ,

$$\lambda_{\mathbb{N}}(x)_n = Gs_n(\lambda_{\mathbb{N}}(x)) = \lambda_{\mathbf{1}}(F_{\mathcal{B}}s_n(x)) = \lambda_{\mathbf{1}}(B_n) = \mu_\lambda(B_n).$$

This shows that  $\mu_\lambda$  is an outer premeasure and this gives an order-preserving map  $\mu_{(-)} : [F_{\mathcal{B}}, G]_l^\Sigma \rightarrow M_{\leq}(X, \mathcal{B})$ .

Let  $\lambda \in [F_{\mathcal{B}}, G]_l^\Sigma$  and let  $A$  be a countable set. Let  $a_0 \in A$  and let  $x := (B_a)_{a \in A} \in F_{\mathcal{B}}A$ . Let  $s_{a_0} : A \rightarrow \mathbf{1}$  be the finite partial map that is only defined on  $\{a_0\}$ . This map is injective. From the equalities,

$$\lambda^{\mu_\lambda}(x)_{a_0} = \mu_\lambda(B_{a_0}) = \lambda_{\mathbf{1}}(F_{\mathcal{B}}s_{a_0}(x)) = \lambda_A(x)_{a_0},$$

we conclude that  $\lambda^{\mu_\lambda} = \lambda$ .

For an outer premeasure  $\mu$ , we have  $\mu_{\lambda^\mu}(B) = \lambda_{\mathbf{1}}^\mu(B) = \mu(B)$  for all  $B \in \mathcal{B}$ .

We conclude that  $\mu_{(-)}$  is the inverse of  $\lambda^{(-)}$ . The claim for strict and colax  $\Sigma$ -transformations follows from a similar argument.  $\square$

**Lemma 6.8.** *The category  $\mathbf{Part}_c^f$  has pullbacks and  $j : \mathbf{Part}_c^f \rightarrow \mathbf{PartPreMble}$  preserves these.*

*Proof.* Let  $\mathbf{1}$  be the set with one element. It can be shown that  $\mathbf{1}/\mathbf{Set}_c^f \cong \mathbf{Part}_c^f$  and that  $\mathbf{1}/\mathbf{PreMble} \cong \mathbf{PartPreMble}$ . Under these equivalences,  $j \cong \mathbf{1}/i$ , where  $i : \mathbf{Set}_c^f \rightarrow \mathbf{PreMble}$ . The claim now follows from Lemma 6.3.  $\square$

**Proposition 6.9.** *For  $f : A \rightarrow B$  in  $\mathbf{Part}_c^f$ ,  $Gf$  preserves directed joins.*

*Proof.* The proof is the same as the proof of Proposition 6.4.  $\square$

Using the previous results we can now obtain a formula to turn an outer premeasure into a premeasure in a universal way.

**Proposition 6.10.** *The triple  $(F_{\mathcal{B}}, G, \Sigma)$  satisfies the strictness property. Moreover, the map  $M_{\leq}(X, \mathcal{B}) \cong [F_{\mathcal{B}}, G]_l^{\Sigma} \rightarrow [F_{\mathcal{B}}, G]_s \cong M(X, \mathcal{B})$ , sends an outer premeasure  $\mu$  to  $\bar{\mu}$ , which is defined as*

$$\bar{\mu}(B) := \sup \left\{ \sum_{n=1}^{\infty} \mu(B_n) \mid \bigcup_{n=1}^{\infty} B_n \subseteq B \right\},$$

for all  $B \in \mathcal{B}$ .

*Proof.* By Proposition 6.9, we know that  $Gf$  preserves directed joins. From Lemma 6.8 and the fact that hom-functors preserve pullbacks, we know that  $F_{\mathcal{B}}$  preserves pullbacks.

Let  $A$  be a countable set and let  $(X_a)_{a \in A} \in F_{\mathcal{B}}A$ . Consider  $g : C \rightarrow A$  and  $(Y_c)_{c \in C} \in F_{\mathcal{B}}C$  such that  $\bigcup_{g(c)=a} Y_c \subseteq X_a$  for all  $a \in A$ . Because  $\bigcup_{c \in C} Y_c \in \mathcal{B}$  and  $g$  is finite, we know that  $Z_a := \bigcup_{g(c)=a} Y_c \in \mathcal{B}$  for all  $a \in A$ . For  $d \in C \amalg A =: D$ , define

$$\tilde{Y}_d := \begin{cases} Y_c & \text{if } d = c \\ X_a \setminus Z_a & \text{if } d = a \end{cases}.$$

It is clear that  $(\tilde{Y}_d)_{d \in D}$  is an element of  $F_{\mathcal{B}}D$ . Consider the finite partial map  $g+1 : D \rightarrow A$ , which is the same as  $g$  on  $C$  and the identity on  $A$ . Let  $s : D \rightarrow C$  be the finite partial map that is the identity on  $C$  and is not defined on  $A$ .

We have the following inequality

$$\begin{array}{ccc} C & \xrightarrow{g} & A \\ \uparrow s & \leq & \nearrow g+1 \\ D & & \end{array}$$

and we have that  $\bigcup_{g(d)=c} \tilde{Y}_d = Y_c$  for all  $c \in C$  and

$$\bigcup_{g+1(d)=a} \tilde{Y}_d = Z_a \cup (X_a \setminus Z_a) = X_a.$$

Therefore it follows that  $(l'_A \downarrow x)_= \rightarrow (l'_A \downarrow x)$  is weakly final. Theorem 4.8 now says that  $(F_{\mathcal{B}}, G, \Sigma)$  satisfies the strictness condition and that for  $\lambda \in [F_{\mathcal{B}}, G]_l$  and  $x \in F_{\mathcal{B}}A$

$$\bar{\lambda}_A(x) = \bigvee \{Gg\lambda_C(y) \mid g : C \rightarrow A; y \in F_{\mathcal{B}}C; F_{\mathcal{B}}g \leq x\}.$$

Let  $\mu$  be an outer premeasure, then  $\bar{\mu} = \mu_{\bar{\lambda}\mu}$ . For  $B \in \mathcal{B}$  this means,

$$\begin{aligned} \bar{\mu}(B) &= \bar{\lambda}\mu_{\mathbf{1}}(B) \\ &= \sup \left\{ Gg\lambda^{\mu}((B_c)_c) \mid g : C \rightarrow \mathbf{1}; \bigcup_{c \in \text{dom}g} B_c \subseteq B \right\} \\ &= \sup \left\{ \sum_{n=1}^{\infty} \mu(B_n) \mid \bigcup_{n=1}^{\infty} B_n \subseteq B \right\}. \end{aligned}$$

□

The following proposition now immediately follows from Corollary 2.4.

**Proposition 6.11.** *The poset  $M(X, \mathcal{B})$  has all joins and for  $(\mu_i)_i$  in  $M(X, \mathcal{B})$ ,*

$$\left( \bigvee_{i \in I} \mu_i \right) (B) = \sup \left\{ \sum_{n=1}^{\infty} \bigvee_{i \in I} \mu_i(B_n) \mid \bigcup_{n=1}^{\infty} B_n \subseteq B \right\}$$

for  $B \in \mathcal{B}$ .

The next result gives us a way to turn an order-preserving map into an outer measure.

**Proposition 6.12.** *Let  $\sigma \in [F_{\mathcal{B}}, G]^{\Sigma}$ . For a countable set  $A$  and  $x := (X_a)_{a \in A} \in F_{\mathcal{B}}A$  and  $a \in A$ ,*

$$(R_l\sigma)_A(x)_a = \inf \left\{ \sum_{n=1}^{\infty} \sigma_{\mathbf{1}}(B_n) \mid X_a \subseteq \bigcup_{n=1}^{\infty} B_n \right\}.$$

*Proof.* Let  $s_c : C \rightarrow \mathbf{1}$  be the injective finite partial map that is only defined on  $\{c\}$ . For  $y := (B_c)_{c \in C} \in F_{\mathcal{B}}C$ , we have

$$\sigma_C(y)_c = Gs_c\sigma_C(y) = \sigma_{\mathbf{1}}(B_c).$$

Define for a countable set  $A$  and  $x := (X_a)_{a \in A} \in F_{\mathcal{B}}A$ ,

$$\begin{aligned} \tau_A(x)_a &:= \bigwedge \{Gg\sigma_C(y) \mid g : C \rightarrow A; y \in F_{\mathcal{B}}C; Fg(y) \geq x\}_a \\ &= \inf \left\{ \sum_{g(c')=a} \sigma_C((B_c)_c)_{c'} \mid g : C \rightarrow A; X_a \subseteq \bigcup_{g(c')=a} B_c \right\} \\ &= \inf \left\{ \sum_{n=1}^{\infty} \sigma_{\mathbf{1}}(B_n) \mid X_a \subseteq \bigcup_{n=1}^{\infty} B_n \right\} \end{aligned}$$

Consider a finite partial map  $f : A \rightarrow B$  and write  $f^{-1}(\{b\}) =: \{a_1, a_2, \dots\}$ . For all  $n \in \mathbb{N}$ , there exists  $X_{a_n} \subseteq (B_m^n)_{m=1}^\infty$  such that

$$\tau_A(x)_{a_n} + \frac{\epsilon}{2^n} \geq \sum_{m=1}^{\infty} \sigma_1(B_m^n) \geq \sum_{m=1}^{\infty} \sigma_1(B_m^n \cap X_{a_n})$$

Summing over  $n$  gives

$$Gf\tau_A(x)_b + \epsilon \geq \sum_{n,m} \sigma_1(B_m^n \cap X_{a_n}) \geq \tau_B(F_{\mathcal{B}}(x))_b$$

for all  $b \in B$ . Here we used that  $(B_m^n \cap X_{a_n})_{n,m}$  are pairwise disjoint premeasurable subsets, whose union equals  $Ff(x)_b$ . Taking  $\epsilon \rightarrow 0$ , shows that  $\tau$  is lax. By the dual of Proposition 4.7 it follows that  $R_l\sigma = \tau$ . □

This leads to a way of turning inner premeasures into premeasures in a universal way.

**Corollary 6.13.** *The map  $M_{\geq}(X, \mathcal{B}) \cong [F_{\mathcal{B}}, G]_c^{\Sigma} \rightarrow [F_{\mathcal{B}}, G]_s \cong M(X, \mathcal{B})$ , sends an inner premeasure  $\mu$  to  $\underline{\mu}$ , which is defined as*

$$\underline{\mu}(B) := \inf \left\{ \sum_{n=1}^{\infty} \mu(B_n) \mid B \subseteq \bigcup_{n=1}^{\infty} B_n \right\},$$

for all  $B \in \mathcal{B}$ .

*Proof.* If we apply Proposition 6.12 on an inner premeasure  $\mu$ , we obtain for  $B \in \mathcal{B}$ ,

$$\underline{\mu}(B) := \mu_{\underline{\sigma}^{\mu}}(B) = \underline{\sigma}^{\mu}_1(B) = (R_l\sigma^{\mu})_1(B) = \inf \left\{ \sum_{n=1}^{\infty} \mu(B_n) \mid B \subseteq \bigcup_{n=1}^{\infty} B_n \right\}.$$

□

## 7 The Carathéodory extension theorem

In this last section, we will apply the results of extensions of transformations from sections 3 and 5 to the transformations that represent measures and premeasures. This will lead to a proof of the Carathéodory extension theorem.

Let  $(X, \mathcal{B})$  be a premeasurable space. Let  $\sigma(\mathcal{B})$  be the sigma-algebra generated by  $\mathcal{B}$ . Let  $\Sigma, F_{\mathcal{B}}, F_{\sigma(\mathcal{B})}$  and  $G$  as in subsection 6.2. For a countable set  $A$ , we have an inclusion  $\iota_A : F_{\mathcal{B}}A \rightarrow F_{\sigma(\mathcal{B})}A$ . These define a strict transformation  $\iota : F_{\mathcal{B}} \rightarrow F_{\sigma(\mathcal{B})}$ . Our goal is to prove that we can extend strict transformation  $F_{\mathcal{B}} \rightarrow G$  along  $\iota$  and that these extensions are proper.

We will show that right extensions of strict transformations always exist, which will imply the Carathéodory extension theorem. We will refer to this extension as the *right Carathéodory extension*. Moreover, this characterizes the right Carathéodory extension by a universal property. We will also briefly discuss left extensions. However, we will show that the *left Carathéodory extension* does not exist in general.

## 7.1 The right Carathéodory extension

We first look at a useful result for right extensions of (general)  $\Sigma$ -transformations.

**Proposition 7.1.** *For  $\tau \in [F_{\mathcal{B}}, G]^{\Sigma}$ , the right extension of  $\tau$  along  $\iota$  exists and*

$$(\text{Ran}_\iota \lambda)_I = \text{Ran}_{\iota_I} \lambda_I.$$

*Proof.* We will first show that  $[F_{\mathcal{B}}, G]^{\Sigma} \cong [F_{\mathcal{B}\mathbf{1}}, G\mathbf{1}]$ . There is an order-preserving map  $[F_{\mathcal{B}}, G]^{\Sigma} \rightarrow [F_{\mathcal{B}\mathbf{1}}, G\mathbf{1}]$  that sends  $\tau \rightarrow \tau_{\mathbf{1}}$ .

For a functor  $f : F_{\mathcal{B}\mathbf{1}} \rightarrow G\mathbf{1}$ , define

$$\tau_A^f((B_a)_{a \in A}) := (f(B_a))_{a \in A}.$$

Let  $\tau \in [F_{\mathcal{B}\mathbf{1}}, G\mathbf{1}]^{\Sigma}$ . For a countable set  $A$  and  $a_0 \in A$ , let  $s_{a_0} : A \rightarrow \mathbf{1}$  be the finite partial map that is only defined on  $\{a_0\}$ . Since this map is injective we find that

$$\tau_A((B_a)_{a \in A})_{a_0} = Gs_{a_0}(\tau_A((B_a)_{a \in A}))_{\mathbf{1}} = \tau_{\mathbf{1}}(B_{a_0}).$$

This shows that  $\tau^{\tau_{\mathbf{1}}} = \tau$ . For a functor  $f : F_{\mathcal{B}\mathbf{1}} \rightarrow G\mathbf{1}$ , we find that  $\tau_{\mathbf{1}}^f = f$ . This shows that  $\tau \mapsto \tau_{\mathbf{1}}$  is an isomorphism.

Clearly,  $[F_{\sigma(\mathcal{B})}\mathbf{1}, G\mathbf{1}] \xrightarrow{-\circ \iota_{\mathbf{1}}} [F_{\mathcal{B}\mathbf{1}}, G\mathbf{1}]$  has a right adjoint. It is now easy to see that under the identification, this gives a right extension of  $\lambda$  along  $\iota$  and the claimed equality.  $\square$

We will now describe what right extensions of lax  $\Sigma$ -transformations look like.

**Proposition 7.2.** *Let  $\lambda \in [F_{\mathcal{B}}, G]_I^{\Sigma}$ . The right extensions of  $\lambda$  along  $\iota$  exist and is proper. Moreover,*

$$[\text{Ran}_\iota \lambda]_A(x)_a = \inf \left\{ \sum_{n=1}^{\infty} \lambda_I(B_n) \mid X_a \subseteq \bigcup_{n=1}^{\infty} B_n \right\}$$

for all countable sets  $A$  and  $x := (X_a)_{a \in A} \in F_{\sigma(\mathcal{B})}A$ .

*Proof.* The existence follows from Proposition 3.4. The left adjoints in the following diagram commute

$$\begin{array}{ccc} [F_{\sigma(\mathcal{B})}, G]_I^{\Sigma} & \xrightarrow{-\circ \iota} & [F_{\mathcal{B}}, G]_I^{\Sigma} \\ \downarrow U_I & & \downarrow U_I \\ [F_{\sigma(\mathcal{B})}, G]^{\Sigma} & \xrightarrow{-\circ \iota} & [F_{\mathcal{B}}, G]^{\Sigma} \end{array}$$

Therefore their right adjoints also commute. Let  $\tau$  be the (general)  $\Sigma$ -transformation that is the right extension of the (general)  $\Sigma$ -transformation  $\lambda$  along  $\iota$ . We have,

$$\text{Ran}_\iota \lambda = R_l \tau.$$

Let  $A$  be a countable set and let  $x := (X_a)_{a \in A} \in F_{\sigma(\mathcal{B})}A$ . By Proposition 6.12 and Proposition 7.1 together with the fact that  $\text{Ran}_{\iota_1} \lambda_1 \circ \iota_1 = \lambda_1$ , since  $\iota_1$  is full and faithful, we find that

$$\begin{aligned} [\text{Ran}_\iota \lambda]_A(x)_a &= \inf \left\{ \sum_{n=1}^{\infty} \tau_1(A_n) \mid X_a \subseteq \bigcup_{n=1}^{\infty} A_n \right\} \\ &= \inf \left\{ \sum_{n=1}^{\infty} \text{Ran}_{\iota_1} \lambda_1(A_n) \mid X_a \subseteq \bigcup_{n=1}^{\infty} A_n \right\} \\ &\leq \inf \left\{ \sum_{n=1}^{\infty} \text{Ran}_{\iota_1} \lambda_1(B_n) \mid X_a \subseteq \bigcup_{n=1}^{\infty} B_n \right\} \\ &= \inf \left\{ \sum_{n=1}^{\infty} \lambda_1(B_n) \mid X_a \subseteq \bigcup_{n=1}^{\infty} B_n \right\}. \end{aligned}$$

It is easy to see that  $\text{Ran}_{\iota_1} \lambda_1(A) = \inf \{ \lambda_1(A) \mid A \subseteq B \}$ . Let  $(A_n)_{n=1}^{\infty}$  be pairwise disjoint in  $\sigma(\mathcal{B})$  such that  $X_a \subseteq \bigcup_{n=1}^{\infty} A_n$ . Let  $\epsilon > 0$ . For every  $n \geq 1$ , there exists  $A_n \subseteq B_n$  such that

$$\text{Ran}_{\iota_1} \lambda_1(A_n) + \frac{\epsilon}{2^n} \geq \lambda_1(B_n) \geq \lambda_1(\tilde{B}_n),$$

where  $\tilde{B}_1 := B_1$  and  $\tilde{B}_n := B_n \setminus \tilde{B}_{n-1}$  for  $n > 1$ . Summing over  $n \geq 1$  gives

$$\sum_{n=1}^{\infty} \text{Ran}_{\iota_1} \lambda_1(A_n) + \epsilon \geq \sum_{n=1}^{\infty} \lambda_1(\tilde{B}_n) \geq \inf \left\{ \sum_{n=1}^{\infty} \lambda_1(B_n) \mid X_a \subseteq \bigcup_{n=1}^{\infty} B_n \right\}.$$

Taking  $\epsilon \rightarrow 0$  and taking the infimum over all such  $(A_n)_{n=1}^{\infty}$ , yields the claimed equality.

Using Proposition 6.12, it is now easy to see that

$$\text{Ran}_\iota \lambda \circ \iota = R_l(\lambda) = \lambda.$$

This shows that the extension is proper.  $\square$

**Theorem 7.3.** *Right extensions along  $\iota$  of strict transformations exist and are proper.*

*Proof.* Corollary C.3 states that  $\overline{\mu} \upharpoonright_{\mathcal{B}} = \overline{\mu} \upharpoonright_{\mathcal{B}}$  for all outer measures  $\mu$ . This shows that  $\overline{\lambda} \circ \iota = \overline{\lambda} \circ \iota$  for all  $\lambda \in [F_{\sigma(\mathcal{B})}, G]_l^{\Sigma}$ . Because  $(F_{\mathcal{B}}, G, \Sigma)$  and  $(F_{\sigma(\mathcal{B})}, G, \Sigma)$  satisfy the strictness condition, the results now follows from Corollary 3.8.  $\square$

**Theorem 7.4** (Carathéodory). *Let  $\rho$  be a premeasure on a premeasurable space  $(X, \mathcal{B})$ . There exists a measure  $\mu$  on  $(X, \sigma(\mathcal{B}))$ , such that  $\mu(B) = \rho(B)$  for all  $B \in \mathcal{B}$ .*

*Proof.* Let  $\mu$  be the measure corresponding with the right extension along  $\iota$  of the strict transformation  $\tau^\rho : F_{\mathcal{B}} \rightarrow G$ . Since the extension is proper, it follows that  $\mu(B) = \rho(B)$  for all  $B \in \mathcal{B}$ .  $\square$

## 7.2 The left Carathéodory extension

We start by showing that extensions of colax  $\Sigma$ -transformations along  $\iota$  are objectwise and proper.

**Proposition 7.5.** *Left extensions along  $\iota$  of colax  $\Sigma$ -transformations exist and are objectwise and proper.*

*Proof.* We know from Proposition 6.9 that  $Gf$  preserves directed joins. Consider a countable set  $A$  and  $x := (X_a)_{a \in A} \in F_{\sigma(\mathcal{B})}A$  and  $b^1 := (B_a^1)_{a \in A}$  and  $b^2 := (B_a^2)_{a \in A}$  in  $F_{\mathcal{B}}A$  such that  $b^1 \leq x$  and  $b^2 \leq x$ . Then  $(B_a^1 \cup B_a^2)_{a \in A}$  are pairwise disjoint premeasurable subsets and clearly  $B_a^1 \cup B_a^2 \subseteq X_a$  for all  $a \in A$ . Therefore  $\iota_A \downarrow x$  is directed. For an injective finite partial map  $f : A \rightarrow B$ , define  $(F_{\mathcal{B}}f)_* : FB \rightarrow FA$  by sending  $(B_b)_{b \in B}$  to  $(B_f(a))_{a \in A}$ . This map is well-defined because  $f$  is injective. It is furthermore, easy to check that  $(F_{\mathcal{B}}f)_*$  is left adjoint to  $F_{\mathcal{B}}f$  and these left adjoints satisfy condition  $(\iota_2)$  in Theorem 5.2. It follows now from that same theorem that left extensions along  $\iota$  exist and are objectwise. Since  $\iota_A$  is full and faithful for all countable sets  $A$ , we know by Lemma 3.1 that the extension is proper.  $\square$

However, left extensions along  $\iota$  of *strict* transformations might not exist in general. In other words, the left Carathéodory extension of a premeasure does not always exist, as can be seen from the following counterexample, based on Example 4.20 in [6].

**Example 7.6.** Let  $\mathcal{B}$  be the algebra of subsets of  $\mathbb{Q}$ , generated by  $\{(a, b] \cap \mathbb{Q} \mid a, b \in \mathbb{Q}\}$ . Let  $\rho : \mathcal{B} \rightarrow [0, \infty]$  be the premeasure that takes the value  $\infty$  for all non-empty subsets in  $\mathcal{B}$ . Suppose that this premeasure has a left Carathéodory extension  $\mu$ . Note that since right Carathéodory extensions exist and are proper, we know by Proposition 3.9 that the left Carathéodory extension has to be proper. For  $r > 0$ , there is a measure  $\mu_r$  on  $\mathbb{Q}$  that is defined by  $\mu_r(\{q\})$  for all  $q \in \mathbb{Q}$ . We clearly have that  $\mu_r(B) = \rho(B)$  for all  $B \in \mathcal{B}$ . By the universal property of left extensions it follows that  $\mu \leq \mu_r$ , i.e. for all  $r > 0$  and  $q \in \mathbb{Q}$ , we have

$$\mu(\{q\}) \leq r.$$

This implies that  $\mu = 0$ , which is clearly a contradiction.

**Remark 7.7.** It follows immediately from the  $\pi$ - $\lambda$  theorem (Theorem 1.19 in [3]), that a finite measure is determined by its values on a generating algebra of subsets. Using the above results, we can conclude that for finite premeasures, there exist a unique extension to a measure. In particular, the left and right Carathéodory extension exist and are equal.

## A Lax coends

In this section we will give the definition of lax coends. More details about lax coends can be found in chapter 7 of [4].

Let  $\mathcal{A}$  and  $\mathcal{B}$  be strict 2-categories and let  $S : \mathcal{A}^{\text{op}} \times \mathcal{A} \rightarrow \mathcal{B}$  be a strict 2-functor.

**Definition A.1.** A lax cowedge  $\omega$  consists of

- an object  $B$  in  $\mathcal{B}$ ,
- a morphism  $\omega_A : S(A, A) \rightarrow B$  for every object  $A$  in  $\mathcal{A}$  and
- for every morphism  $f : A_1 \rightarrow A_2$ , a 2-cell  $\omega_f$

$$\begin{array}{ccc}
 S(A_1, A_2) & \xrightarrow{S(f \times 1_{A_2})} & S(A_2, A_2) \\
 \downarrow S(1_{A_1} \times f) & \nearrow \omega_f & \downarrow \omega_{A_2} \\
 S(A_1, A_1) & \xrightarrow{\omega_{A_1}} & B
 \end{array} \cdot$$

such that these are consistent with the 2-category structure of  $\mathcal{A}$ .

**Definition A.2.** Let  $\omega^1 := (B^1, (\omega_A^1)_A, (\omega_f^1)_f)$  and  $\omega^2 := (B^2, (\omega_A^2)_A, (\omega_f^2)_f)$  be lax cowedges of  $S$ . A **morphism of cowedges** from  $\omega^1$  to  $\omega^2$  is a morphism  $s : B_1 \rightarrow B_2$  such that

$$\begin{array}{ccc}
 S(A_1, A_2) & \xrightarrow{S(f \times 1_{A_2})} & S(A_2, A_2) \\
 \downarrow S(1_{A_1} \times f) & \nearrow \omega_f^1 & \downarrow \omega_{A_2} \\
 S(A_1, A_1) & \xrightarrow{\omega_{A_1}} & B_1 \\
 & \searrow \omega_{A_1}^2 & \downarrow \omega_{A_2}^2 \\
 & & B_2
 \end{array}
 =
 \begin{array}{ccc}
 S(A_1, A_2) & \xrightarrow{S(f \times 1_{A_2})} & S(A_2, A_2) \\
 \downarrow S(1_{A_1} \times f) & \nearrow \omega_f^2 & \downarrow \omega_{A_2}^2 \\
 S(A_1, A_1) & \xrightarrow{\omega_{A_1}^2} & B_2
 \end{array}$$

$\omega_{A_1}^2$  is a 2-cell from  $\omega_{A_1}$  to  $\omega_{A_1}^2$ , and  $s$  is a morphism from  $B_1$  to  $B_2$ .

for all morphism  $f : A_1 \rightarrow A_2$ .

The category of lax cowedges of  $S$  and their morphisms is denoted by  $\mathbf{Cowedge}_l(S)$ .

**Definition A.3.** The lax coend of  $S$  is an initial object in  $\mathbf{Cowedge}_l(S)$ .



If  $(B, (\omega_A)_A, (\omega_f)_f)$  is the lax coend of  $S$ , i.e. the initial object in  $\mathbf{Cowedge}_l(S)$ , then we also write

$$B =: \oint^{A \in \text{ob } \mathcal{A}} S(A, A).$$

Dually, we can define a category of colax cowedges of  $S$ ,  $\mathbf{Cowedge}_c(S)$  and define colax coends as initial objects in this category. The notation we will use for the colax coend of  $S$  is  $\oint^{A \in \text{ob } \mathcal{A}} S(A, A)$ .

## B Weakly final functors

In this section we will look at final functors and their properties. We will generalize this to *weakly* final functors.

**Definition B.1.** A functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  is called **final** if  $d/F$  is non-empty and connected for all objects  $d$  in  $\mathcal{D}$ .

A functor  $F : \mathbf{1} \rightarrow \mathcal{D}$  is final if and only if  $F(0)$  is final in  $\mathcal{D}$ . This motivates the name. Final functors have the very useful following characterization

**Proposition B.2.** *A functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  is final if and only if the map*

$$\text{colim}DF \rightarrow \text{colim}D$$

*is an isomorphism for all cocomplete categories  $\mathcal{E}$  and functors  $D : \mathcal{D} \rightarrow \mathcal{E}$ .*

We will now introduce weakly final functors.

**Definition B.3.** A functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  is **weakly final** if  $d/F$  is non-empty for all objects  $d$  in  $\mathcal{D}$ .

A functor  $F : \mathbf{1} \rightarrow \mathcal{D}$  is weakly final if and only if  $F(0)$  is weakly final. Clearly, every final functor is weakly final. Furthermore, we also have a characterization for weakly final functors.

**Proposition B.4.** *A functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  is weakly final if and only if*

$$\text{colim}DF = \text{colim}D$$

*for all complete posets  $P$  and all functors  $D : \mathcal{D} \rightarrow P$ .*

*Proof.* Suppose that  $F$  is weakly final. We always have that  $\text{colim}DF \leq \text{colim}D$ . For  $d \in \text{ob } \mathcal{D}$ , there is an object  $c$  in  $\mathcal{C}$  and a morphism  $d \rightarrow Fc$ . Therefore,  $Dd \leq DFc \leq \text{colim}DF$ . Since,  $d$  was arbitrary, we conclude that  $\text{colim}D \leq \text{colim}DF$ .

Define a functor  $D : \mathcal{D} \rightarrow \mathbf{2}$ , that sends  $d$  to 0 if  $d/F$  is non-empty and 1 otherwise. Clearly  $DF$  is the constant 0 functor. It follows that  $\text{colim}D = 0$ , which means that  $D(d) = 0$  for all objects  $d$  in  $\mathcal{D}$ . In other words,  $d/F$  is non-empty for all objects  $d$  in  $\mathcal{D}$ .  $\square$

## C Finite outer premeasures

An outer premeasure  $\mu$  on  $(X, \mathcal{B})$  is called **finite** if  $\overline{\mu}(X) < \infty$ .<sup>7</sup>

**Theorem C.1.** *Let  $(X, \mathcal{B})$  be a premeasurable space and let  $\mu$  be an outer premeasure on  $(X, \sigma(\mathcal{B}))$ . Then  $\mu$  is finite if and only if  $\mu|_{\mathcal{B}}$  is finite. In this case for every  $\epsilon > 0$  and  $A \in \sigma(\mathcal{B})$ , there exists a  $B \in \mathcal{B}$  such that  $\mu(A \Delta B) < \epsilon$ . Also in this case,*

$$\overline{\mu}(X) = \overline{\mu|_{\mathcal{B}}}(X).$$

*Proof.* If  $\mu$  is finite then,  $\overline{\mu|_{\mathcal{B}}}(X) \leq \overline{\mu}(X) < \infty$  and therefore  $\mu|_{\mathcal{B}}$  is finite.

Suppose now that  $\mu|_{\mathcal{B}}$  is finite. Define the set

$$S := \{A \in \sigma(\mathcal{B}) \mid \forall \epsilon > 0 \exists B \in \mathcal{B} : \mu(A \Delta B) < \epsilon\}$$

If  $A \in S$  and  $\epsilon > 0$ , then there exists  $B$  such that  $\mu(A \Delta B) < \epsilon$ . Therefore  $\mu(A^C \Delta B) = \mu(A \Delta B^C) < \epsilon$ . We conclude that  $A^C \in S$  and therefore  $S$  is closed under complements.

Let  $\epsilon > 0$ . For  $A_1, A_2 \in S$ , there exists  $B_1, B_2 \in \mathcal{B}$  such that  $\mu(A_1 \Delta B_1) < \frac{\epsilon}{2}$  and  $\mu(A_2 \Delta B_2) < \frac{\epsilon}{2}$ . Since  $(A_1 \cup A_2) \Delta (B_1 \cup B_2) \subseteq (A_1 \Delta B_1) \cup (A_2 \Delta B_2)$ , it follows that  $\mu((A_1 \cup A_2) \Delta (B_1 \cup B_2)) \leq \epsilon$ . Therefore  $S$  is closed under finite unions.

Let  $\epsilon > 0$ . For  $(A_n)_{n=1}^{\infty}$  pairwise disjoint in  $S$ . For  $N \geq 1$ , there exist  $(B_n)_{n=1}^N$  such that  $\mu(A_n \Delta B_n) \leq \frac{\epsilon}{N}$  for all  $n \in \{1, \dots, N\}$ . We can choose  $(B_n)_{n=1}^N$  such that they are pairwise disjoint.<sup>8</sup> It follows now that

$$\sum_{n=1}^N \mu(A_n) \leq \sum_{n=1}^N (\mu(A_n \cap B_n) + \mu(A_n \Delta B_n)) \leq \sum_{n=1}^N \left( \mu(B_n) + \frac{\epsilon}{N} \right) = \sum_{n=1}^N \mu(B_n) + \epsilon \leq \overline{\mu|_{\mathcal{B}}}(X) + \epsilon$$

By letting  $N \rightarrow \infty$  and then letting  $\epsilon \rightarrow 0$ , we find for pairwise disjoint  $(A_n)_{n=1}^{\infty}$  in  $S$ , that

$$\sum_{n=1}^{\infty} \mu(A_n) \leq \overline{\mu|_{\mathcal{B}}}(X) < \infty \quad (1)$$

<sup>7</sup> Here we use the notation of Proposition 6.10

<sup>8</sup> Because  $A_n \in S$ , there exists  $\tilde{B}_n \in \mathcal{B}$  such that  $\mu(A_n \Delta \tilde{B}_n) < \frac{\epsilon}{N^2}$ . Now define  $B_1 := \tilde{B}_1$  and  $B_n := \tilde{B}_n \setminus \bigcup_{k=1}^{n-1} B_k$  for  $n \geq 2$ . Clearly  $(B_n)_{n=1}^N$  is a pairwise disjoint collection of elements in  $\mathcal{B}$ . Moreover, since  $B_n \subseteq \tilde{B}_n$  and  $A_n \subseteq A_{n-1}^C$  for all  $1 \leq n \leq N$ , we have

$$\begin{aligned} A_n \Delta B_n &= (A_n \cap B_n) \cup (A_n \cap B_n^c) \\ &= (A_n \cap B_n) \cup (A_n \cap \tilde{B}_n^c) \cup (A_n \cap B_{n-1}) \cup \dots \cup (A_n \cap B_1) \\ &\subseteq (A_n \cap \tilde{B}_n) \cup (A_n \cap \tilde{B}_n^c) \cup (A_{n-1}^c \cap \tilde{B}_{n-1}) \cup \dots \cup (A_1^c \cap \tilde{B}_1) \\ &\subseteq (A_n \Delta \tilde{B}_n) \cup (A_{n-1} \Delta \tilde{B}_{n-1}) \cup \dots \cup (A_1 \Delta \tilde{B}_1) \end{aligned}$$

Therefore  $\mu(A_n \Delta B_n) \leq \frac{n\epsilon}{N^2} \leq \frac{\epsilon}{N}$ .

It follows that there exists  $N \geq 1$  such that  $\sum_{n=N+1}^{\infty} \mu(A_n) < \frac{\epsilon}{2}$ . Because

$$\left( \bigcup_{n=1}^{\infty} A_n \right) \Delta \left( \bigcup_{n=1}^N B_n \right) \subseteq \left( \bigcup_{n=1}^N A_n \Delta B_n \right) \cup \bigcup_{n=N+1}^{\infty} A_n,$$

we have that

$$\mu \left( \bigcup_{n=1}^{\infty} A_n \Delta \bigcup_{n=1}^N B_n \right) \leq \sum_{n=1}^N \mu(A_n \Delta B_n) + \sum_{n=N+1}^{\infty} \mu(A_n) \leq \epsilon.$$

We conclude that  $S$  is closed under countable disjoint unions. This means that  $S$  is a  $\sigma$ -algebra that contains  $\mathcal{B}$  and therefore  $S = \sigma(\mathcal{B})$ . This already proves the second claim in the statement. By (1) it also follows that  $\bar{\mu}(X) \leq \overline{\mu}_{\mathcal{B}}(X)$ . This shows that  $\mu$  is finite and the last claim in the statement.  $\square$

**Corollary C.2.** *Let  $\mu$  be an outer measure on  $(X, \sigma(\mathcal{B}))$ , then  $\bar{\mu}(X) = \overline{\mu}_{\mathcal{B}}(X)$ .*

*Proof.* If  $\mu$  is finite, then this follows from Theorem C.1. If  $\mu$  is not finite then,  $\mu|_{\mathcal{B}}$  is also not finite by Theorem C.1 and the result also follows.  $\square$

**Corollary C.3.** *Let  $\mu$  be an outer measure on  $(X, \sigma(\mathcal{B}))$ , then  $\overline{\mu}_{\mathcal{B}}(B) = \bar{\mu}(B)$  for all  $B \in \mathcal{B}$ .*

*Proof.* Let  $B \in \mathcal{B}$ . Consider the algebra on  $B$ ,  $\mathcal{B}' := \{C \in \mathcal{B} \mid C \subseteq B\}$ . It can be checked that  $\sigma(\mathcal{B}') = \sigma(\mathcal{B})'$ .<sup>9</sup> Let  $\mu'$  be the outer measure on  $(B, \sigma(\mathcal{B}'))$  defined by  $\mu'(C) := \mu(C)$  for all  $C \in \sigma(\mathcal{B}')$ . Using the formula from Proposition 6.10, it is now easy to see that  $\bar{\mu}'(B) = \bar{\mu}(B)$  and  $\overline{\mu'}_{\mathcal{B}'}(B) = \overline{\mu}_{\mathcal{B}}(B)$ . Using Corollary C.2, we conclude that  $\overline{\mu}_{\mathcal{B}}(B) = \bar{\mu}(B)$  for all  $B \in \mathcal{B}$ .  $\square$

## References

- [1] R. Blackwell, G. M. Kelly, and A. J. Power. Two-dimensional monad theory. *J. Pure Appl. Algebra*, 59(1):1–41, 1989.
- [2] P. T. Johnstone. Adjoint lifting theorems for categories of algebras. *Bull. London Math. Soc.*, 7(3):294–297, 1975.
- [3] A. Klenke. *Probability Theory: A Comprehensive Course*. Universitext. Springer International Publishing, 2020.

<sup>9</sup> Clearly  $\mathcal{B}' \subseteq \sigma(\mathcal{B})'$  and since  $\sigma(\mathcal{B})'$  is a  $\sigma$ -algebra, we have  $\sigma(\mathcal{B}') \subseteq \sigma(\mathcal{B})'$ . For the other inclusion, consider

$$S := \{A \subseteq X \mid A \cap B \in \sigma(\mathcal{B}')\}.$$

It is easy to see that  $S$  is a  $\sigma$ -algebra containing  $\mathcal{B}$ . Therefore  $\sigma(\mathcal{B}) \subseteq S$ . Thus for  $A \in \sigma(\mathcal{B})'$ , we have that  $A \in S$ , which means that  $A = A \cap B \in \sigma(\mathcal{B}')$ .

- 
- [4] F. Loregian. *(Co)end Calculus*, volume 468 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2021.
  - [5] E. Riehl. *Category Theory in Context*. Aurora: Dover Modern Math Originals. Dover Publications, 2017.
  - [6] G. L. Wise and E. B. Hall. *Counterexamples in Probability and Real Analysis*. The Clarendon Press, Oxford University Press, New York, 1993.