

HIGHER INTERNAL COVERS

MOSHE KAMENSKY

ABSTRACT. We define and study a higher-dimensional version of model theoretic internality, and relate it to higher-dimensional definable groupoids in the base theory.

1. INTRODUCTION

The model theoretic notions of *internality* and the binding group came up originally in work of Zil'ber on categorical theories ([13]), and shortly after by Poizat ([11]) in the ω -stable context, where it was also noticed that differential Galois theory occurs as a special case. The stability hypothesis was completely removed in [3, App. B], where it was shown that the crucial hypothesis is *stable embeddedness* of the base theory.

Internality is a condition on a definable set \mathbf{Q} in an expansion \mathcal{T}^* of a theory \mathcal{T} to “almost” be interpretable in \mathcal{T} : It is interpretable, after adding a set of parameters to \mathcal{T}^* . In this situation, the theory provides a *definable* group \mathbf{G} in \mathcal{T}^* , acting definably on \mathbf{Q} as its group of automorphisms fixing all elements in the reduct \mathcal{T} . It is important here that the binding group \mathbf{G} is defined in \mathcal{T}^* rather than in \mathcal{T} : In applications, one often understands groups in \mathcal{T} better than in \mathcal{T}^* . The group \mathbf{G} itself is also internal to \mathcal{T} , and as a result can be identified with a definable group \mathbf{H} in \mathcal{T} , but only non-canonically (and in general, only after adding parameters). In the context of differential Galois theory, this is related to the fact that the group of points of the (algebraic) Galois group of a differential equation does not act on the set of solutions, and its identification with the group of automorphisms is not canonical.

The non-canonicity was explained by Hrushovski in [4], where it is shown that the natural object that appears in this context is a definable *groupoid* in \mathcal{T} , with the different groups \mathbf{H} occurring as the groups of automorphisms of each object. In fact, it is shown there that there is a correspondence between groupoids definable in the base theory \mathcal{T} and internal sorts in expansions of \mathcal{T} . This correspondence is reviewed in §2. It is also suggested in [4] that sorts of \mathcal{T}^* internal to \mathcal{T} should be viewed as generalised sorts of \mathcal{T} , obtained as a quotient by the corresponding definable groupoid, just like an imaginary sort is obtained from a definable equivalence relation (which is a special case). In the current paper we try to follow this suggestion, by

Date: October 7, 2022.

2020 *Mathematics Subject Classification.* 03C40.

considering what should be the correct notion of internality, after viewing these new sorts as “legitimate” definable sorts.

Our approach is motivated by topology. There, a typical example of a groupoid arises as the fundamental groupoid $\pi_1(X)$ of a space X , i.e., the groupoid whose objects are the points of X , and whose morphisms are homotopy classes of paths. For sufficiently nice X , this groupoid can be described in terms of the category of *local systems* (locally constant sheaves) on X : each point of X determines a functor to the category of sets, satisfying suitable properties (for example, it commutes with products), and each path determines a map between such functors (which depends only on the homotopy class since the system is locally constant). We propose to view internality as analogous to this picture: definable sets in the theory corresponding to a definable groupoid \mathbf{G} in \mathcal{T} can be viewed as local systems (of definable sets) on \mathbf{G} , and conversely. This point of view is explained in §2.2.3 (the base theory \mathcal{T} corresponds to a contractible space in this approach, so definable sets in it correspond to constant systems).

By definition, the local systems on X do not tell us anything about the homotopy type of X above homotopical dimension 1. To encode higher homotopical information, we may try looking at families of spaces rather than of sets. A space X is called *n-truncated* if $\pi_k(X, x)$ is trivial for all $k > n$ and base points $x \in X$. Such spaces are represented in homotopy theory by what we call in this paper *n-groupoids* (Def. 3.2.3 in the definable setting; these are equivalent to *n-categories* in the sense of [8, §2.3.4] which are groupoids). In the case $n = 1$, these are usual groupoids, and the previous paragraph suggests studying them by systems of 0-truncated spaces, i.e., sets. Going one dimension higher, one expects to recover 2-groupoids from systems of 1-truncated spaces. In the definable context, we decided to identify such spaces with internal sorts, we consider “local systems” of internal sorts, i.e., internal sorts of an expanded theory.

Our main result, Theorem 3.3.9, shows one direction of the expected correspondence: we associate to a 2-groupoid \mathbf{G} in the theory \mathcal{T} a theory $\mathcal{T}_{\mathbf{G}}$ expanding it, and a collection of internal sorts of $\mathcal{T}_{\mathbf{G}}$, which we view as “higher local systems”. The statement is that the canonical 2-groupoid associated to this datum recovers (up to weak equivalence) the original one (part of the other direction is indicated briefly, but is mostly left for future work).

We mention that this result is one possible generalisation of the results of [4] to higher dimensions. Other such generalisations include the papers [1], [2] and [12], but they all appear to go in different directions. We also mention that in the context of usual (rather than definable) homotopy theory, analogous results are well known (for example, the main part of Theorem 3.3.9 is really a version of the higher dimensional Yoneda lemma), but the methods in the proof of these results do not translate easily to the definable setting. In fact, the situation here is more similar to the one described

in [8, §6.5], though made simpler by the existence of models (i.e., we have “enough points”).

1.1. Structure of the paper. It is very simple: in Section 2, we review the situation in the one dimensional case. This serves both as a motivating analogy and to complete some background used later. Most of the material there appears in some form in [4] (sometimes implicitly), but we include a few easy remarks regarding morphisms and equivalence, interpretation in terms of “local systems”, and a different description of the groupoid associated to an internal cover (which already appeared slightly differently in [7]).

In Section 3, we expose the higher dimensional picture, concentrate on dimension 2. We first define our higher internal covers, then review the theory of (truncated) Kan complexes and n -categories, with a few remarks special to the definable setting, and then prove the main result mentioned above (Theorem 3.3.9).

1.2. Conventions and terminology. For simplicity, we assume our theories \mathcal{T} to admit elimination of quantifiers. By a \mathcal{T} -structure we mean a substructure of some model of \mathcal{T} . If A is such a \mathcal{T} -structure, by \mathcal{T}_A we mean the expansion of \mathcal{T} by constants for the elements of A , along with the usual axioms describing A . If A was not mentioned, we mean “for some A ”.

We will also assume \mathcal{T} eliminates imaginaries (this could be included in the general treatment, but would complicate the exposition). Our usage of elimination of imaginaries will often be in the (equivalent) form of the existence of internal Homs: For every two definable sets \mathbf{X} and \mathbf{Y} , there are ind-definable set $\underline{\mathbf{Hom}}(\mathbf{X}, \mathbf{Y})$ and map $\mathbf{ev} : \mathbf{X} \times \underline{\mathbf{Hom}}(\mathbf{X}, \mathbf{Y}) \rightarrow \mathbf{Y}$, identifying the A -points of $\underline{\mathbf{Hom}}(\mathbf{X}, \mathbf{Y})$, for each \mathcal{T} -structure A , with the set of A -definable maps from \mathbf{X} to \mathbf{Y} . It follows that the subset $\underline{\mathbf{Iso}}(\mathbf{X}, \mathbf{Y})$ of definable isomorphisms is also ind-definable.

Finally, we assume that each theory is generated by one sort, and finitely many relations. Similar to the case in [4], it can be seen that this assumption is not restrictive, since all our constructions commute with adding structure.

1.3. Acknowledgement and dedication. I am delighted to thank Tomer Schlank’s gang at the Hebrew University (aka “Ha’arakya”), and particularly Asaf Horev, Shaul Barkan, Shai Keidar and Tomer himself for answering what must have been rather bizarre and elementary questions in homotopy theory. They form a very helpful and fun community.

This paper is a late expansion on some vague ideas that I presented in the postponed online conference that Honoured Ehud Hrushovski for his 60th birthday. It is a pleasure to thank Udi again for his guidance and dedicate the paper (hopefully clearer than the talk!) to him.

2. A REVIEW OF THE CLASSICAL THEORY

2.1. Internal covers. We recall the following classical definition of internal covers:

Definition 2.1.1. An expansion \mathcal{T}^* of a theory \mathcal{T} is an *internal cover* if \mathcal{T} is stably embedded in \mathcal{T}^* , and for some expansion \mathcal{T}_A^* of \mathcal{T}^* by a set of constants A , each definable set in \mathcal{T}_A^* is definably isomorphic to a definable set in \mathcal{T}_{A_0} .

We recall that *stably embedded* here means that for every definable set \mathbf{X} in \mathcal{T} , every subset of \mathbf{X} definable in \mathcal{T}^* with parameters from \mathcal{T}^* is definable in \mathcal{T} , with parameters from \mathcal{T} .

It was noted in [6] that this condition can be reformulated as follows: If $i : \mathcal{T} \rightarrow \mathcal{T}^*$ is an expansion and \mathbf{X}, \mathbf{Y} are definable in \mathcal{T} , there is a natural (\mathcal{T}^* -definable) map $i(\underline{\mathbf{Hom}}_{\mathcal{T}}(\mathbf{X}, \mathbf{Y})) \rightarrow \underline{\mathbf{Hom}}_{\mathcal{T}^*}(i(\mathbf{X}), i(\mathbf{Y}))$, and i is stably embedded precisely if this map is a bijection (for all \mathbf{X}, \mathbf{Y} definable in \mathcal{T}). We note that in this case, the restriction of this map to the subset $\underline{\mathbf{Iso}}(\mathbf{X}, \mathbf{Y})$ of isomorphisms is also a bijection, and that taking into account parameters, no new structure is induced on \mathcal{T}^* . In particular, for any \mathcal{T} -structure A , the expansion \mathcal{T}_A^* is well defined.

As already done above, the expansion i can be viewed as a functor from the category of definable sets in \mathcal{T} to those in \mathcal{T}^* , which preserves finite limits. In these terms, i is a stable embedding precisely if it is Cartesian, in the terminology of [10]. We may apply the same definition to a more general interpretation, so we say that an interpretation $i : \mathcal{T}_1 \rightarrow \mathcal{T}_2$ is *stable* if it is Cartesian as a functor. In these terms, the definition of internal covers can be reformulated as follows:

Proposition 2.1.2. *An expansion \mathcal{T}^* of a theory \mathcal{T} is an internal cover if it is stable, and \mathcal{T}^* admits a stable interpretation p in \mathcal{T}_A over \mathcal{T} .*

By “over \mathcal{T} ” we mean that the restriction of p to \mathcal{T} coincides with the expansion by constants.

Proof. let \mathbf{Q} be a definable set in \mathcal{T}^* , generating it over \mathcal{T} . Assume first that \mathcal{T}^* is an internal cover of \mathcal{T} , so there is a sort \mathbf{X} of \mathcal{T}^* , an expansion by a constant symbol $a \in \mathbf{X}$, and a definable bijection $g_a : \mathbf{Q} \rightarrow \mathbf{Q}_a$, with \mathbf{Q}_a definable in \mathcal{T} . By stable-embeddedness, \mathbf{Q}_a is definable by a parameter a_0 in \mathcal{T} . The assignment $\mathbf{Q} \mapsto \mathbf{Q}_a$ extends uniquely to an interpretation x_{a_0} of \mathcal{T}^* in \mathcal{T}_{a_0} , over \mathcal{T} . Since g_a determines a definable isomorphism between \mathbf{Q} and \mathbf{Q}_a (and similarly for any definable set it generates), this interpretation is stable.

Conversely, assume we have a stable interpretation $p : \mathcal{T}^* \rightarrow \mathcal{T}_{A_0}$ over i . We still denote by i the extension of i to the expansion $\mathcal{T}_{A_0} \rightarrow \mathcal{T}_{A_0}^*$ which is the identity on A_0 (it is still stable). Denoting $\mathbf{C} = p(\mathbf{Q})$, the set $\underline{\mathbf{Iso}}(p(\mathbf{Q}), p(i(\mathbf{C}))) = \underline{\mathbf{Iso}}(\mathbf{C}, \mathbf{C})$ is non-empty, since it contains the identity on \mathbf{C} . Since p is stable, the left hand side admits a definable bijection with $\underline{\mathbf{Iso}}(\mathbf{Q}, i(\mathbf{C}))$, so is non-empty as well. Any point a of this set shows that \mathcal{T}^* is an internal cover. \square

2.2. Definable groupoids. A definable groupoid will be denoted as $\mathbf{G} = \langle \mathbf{G}_0, \mathbf{G}_1 \rangle$, with definable set \mathbf{G}_0 of objects and a definable set \mathbf{G}_1 of isomorphisms, where the domain and codomain maps are denoted $\mathbf{d}, \mathbf{c} : \mathbf{G}_1 \rightarrow \mathbf{G}_0$, respectively, and composition denoted by \circ . For objects $a, b \in \mathbf{G}_0$, we write $\mathbf{G}(a, b)$ for the a, b -definable set $\langle \mathbf{d}, \mathbf{c} \rangle^{-1}(\langle a, b \rangle)$ of morphisms from a to b . A map $f : \mathbf{G} \rightarrow \mathbf{H}$ of definable groupoids is a definable functor: a pair of maps $f_0 : \mathbf{G}_0 \rightarrow \mathbf{H}_0$ and $f_1 : \mathbf{G}_1 \rightarrow \mathbf{H}_1$ commuting with the domain, codomain and composition maps. We will say that f is a *weak equivalence*, denoted $f : \mathbf{G} \xrightarrow{\sim} \mathbf{H}$, if it induces an equivalence of categories on all models (this terminology will be generalised in Def 3.2.7).

In [4, §2], a definable groupoid is attached to each internal cover. This groupoid also admits two descriptions. The first as a definable groupoid \mathbf{G}^* in \mathcal{T}^* : its objects are, in the notation of the proof of Prop. 2.1.2, complete types of elements $a \in \mathbf{X}$ over \mathcal{T} , along with an additional object $*$. Since \mathcal{T} is stably embedded, this set of types is definable in \mathcal{T} . The set of isomorphisms from $*$ to such a type p to $*$ is given by the realisations of p . Given another object q , a morphism from p to q is given by a 2-type extending p and q (over \mathcal{T}). We denote by \mathbf{G} the full subgroupoid of \mathbf{G}^* on the same objects excluding $*$. Then \mathbf{G} is defined entirely in \mathcal{T} .

2.2.1. To give a second description, consider, for each \mathcal{T} -structure A , the groupoid $I(A) = I_{\mathcal{T}^*/\mathcal{T}}(A)$ whose objects are stable interpretations of \mathcal{T}^* in \mathcal{T}_A , that are the identity on \mathcal{T}_A , and whose morphisms are isomorphisms of such interpretations, which are the identity when restricted to \mathcal{T} . Here again we may enlarge I to obtain I^* , by adding an additional object $*$, which is described explicitly as the identity interpretation of \mathcal{T}^* , and again morphisms are given by A -definable isomorphisms of interpretations over \mathcal{T} (where A is now a \mathcal{T}^* -structure). The following statement appeared in a slightly different form in [7]:

Proposition 2.2.2. *With notation as in 2.2, for each \mathcal{T} -structure A , there is fully-faithful embedding $i_A : \mathbf{G}^*(A) \rightarrow I^*(A)$, preserving the vertex, and commuting with automorphism action on A . If A is a model, i_A is an equivalence of categories.*

Proof. This is essentially [4, Thm 3.2]. The functor i_A was described in the proof of Prop. 2.1.2: to an object p of $\mathbf{G}(A)$, viewed as a type over \mathcal{T} (definable over A_0), we attach the interpretation $x_b = x_p$ described there, with b any realization of p (as explained there, x_b depends only on $p \in A_0$ and not on b). Each such realization determines an isomorphism g_b from x_b to $*$, again as above, which describes the functor on morphisms from p to $*$. If q is another object, with realization c , i_A assigns $g_c^{-1} \circ g_b : x_p \rightarrow x_q$ to the type r of the pair (b, c) . This depends only on r , since the code for this composition lies in \mathcal{T} , by stable embeddedness. This code also determines r completely, so the functor is fully faithful.

To prove the final statement, let $i : \mathcal{T}^* \rightarrow \mathcal{T}$ be any interpretation over a model M_0 . The internality assumption implies that for some $p \in \mathbf{G}(M_0)$,

the set Y of isomorphisms between $*$ and p is non-empty. Since M_0 is a model, there is a point b in $i(Y)(M_0)$. Then g_b is an isomorphism from x_b to i . \square

To summarise, to each stable embedding of \mathcal{T} in \mathcal{T}^* , we had attached a groupoid $I_{\mathcal{T}^*/\mathcal{T}}$ classifying stable interpretations of \mathcal{T}^* back in \mathcal{T} . The embedding is an internal cover precisely if the groupoid is non-empty, and in this case, the groupoid I is equivalent to a definable one (and to the classical binding groupoid). Conversely, starting with a definable groupoid \mathbf{G} in \mathcal{T} , there is an internal cover $\mathcal{T}^* = \mathcal{T}_{\mathbf{G}}$ and an equivalence $\mathbf{G} \rightarrow I_{\mathcal{T}^*/\mathcal{T}}$ (this construction is recalled below).

2.2.3. Definable \mathbf{G} -sets. If $\mathbf{G} = \langle \mathbf{G}_0, \mathbf{G}_1 \rangle$ is a definable groupoid in \mathcal{T} , by a (left) \mathbf{G} -set we mean a definable set \mathbf{X} , a definable map $\pi : \mathbf{X} \rightarrow \mathbf{G}_0$ to the set of objects \mathbf{G}_0 of \mathbf{G} , and an ‘‘action’’ map $a : \mathbf{G}_1 \times_{\mathbf{G}_0} \mathbf{X} \rightarrow \mathbf{X}$, over \mathbf{G}_0 , satisfying the usual action axioms. Thus, a morphism $g : a \rightarrow b$ in \mathbf{G} determines a bijection $a_g : \mathbf{X}_a \rightarrow \mathbf{X}_b$, where $\mathbf{X}_t = \pi^{-1}(t)$, and we will sometimes write gx in places of $a_g(x)$ (a pair $\langle \mathbf{G}, \mathbf{X} \rangle$ as above is called a *concrete groupoid* in [4, §3]). We think of \mathbf{G} -sets as analogs of local systems over \mathbf{G} .

Let \mathbf{X} be a \mathbf{G} -set. If $\mathbf{H} = \langle \mathbf{H}_0, \mathbf{H}_1 \rangle$ is another groupoid, and $i : \mathbf{G} \rightarrow \mathbf{H}$ is a definable map of groupoids, we set

$$i_!(\mathbf{X}) = \{ \langle h, x \rangle \in \mathbf{H}_1 \times \mathbf{X} \mid i(\pi(x)) = \mathbf{d}(h) \} / \sim,$$

where $\langle h, gx \rangle \sim \langle h \circ i(g), x \rangle$ for $g \in \mathbf{G}_1$ satisfying $\mathbf{d}(g) = \pi(x)$ and $i(\mathbf{c}(g)) = \mathbf{d}(h)$. This is an \mathbf{H} -set, with structure map induced by $\langle h, x \rangle \mapsto \mathbf{c}(h)$ and action induced by $\langle h', \langle h, x \rangle \rangle \mapsto \langle h'h, x \rangle$. On the other hand, if \mathbf{Y} is an \mathbf{H} -set, we set $i^*(\mathbf{Y}) = \mathbf{G}_0 \times_{\mathbf{H}_0} \mathbf{Y}$, with the projection to \mathbf{G}_0 as the structure map, and action given by $\langle g, y \rangle \mapsto i(g)y$ for $y \in \mathbf{Y}$ with $\pi(y) = i(\mathbf{d}(g))$. As the notation suggests, $i_!$ is left adjoint to i^* .

With this notion, we have the following description of definable sets in \mathcal{T}^* as local systems over \mathbf{G} :

Corollary 2.2.4. *If \mathcal{T}^* is an internal cover of \mathcal{T} , corresponding to the definable groupoid \mathbf{G} in \mathcal{T} , then the category of definable sets in \mathcal{T}^* is equivalent to the category of \mathbf{G} -sets in \mathcal{T} . Definable sets from \mathcal{T} correspond to themselves, with trivial action.*

Proof. To a definable set \mathbf{X}^* in \mathcal{T}^* we assign the definable set $\mathbf{X} = \coprod_{p \in \mathbf{G}_0} p(\mathbf{X}^*)$. It follows from the uniformity of p that \mathbf{X} is definable in \mathcal{T} . By definition, \mathbf{X} admits a definable map to \mathbf{G}_0 . The action is given tautologically by the identification of the morphisms in \mathbf{G}_0 with maps of interpretations.

In the other direction, let \mathbf{G}^* be the canonical extension of \mathbf{G} in \mathcal{T}^* (we identify \mathbf{G} with its image in \mathcal{T}^*), let $i : \mathbf{G} \rightarrow \mathbf{G}^*$ be the inclusion, and let j be the inclusion of the canonical object $*$ of \mathbf{G}^* , along with its automorphism group \mathbf{H} , into \mathbf{G}^* . A definable \mathbf{G} -set \mathbf{X} in \mathcal{T} , viewed again as embedded in

\mathcal{T}^* , corresponds then to $\mathbf{X}^* = j^*(i_!(\mathbf{X}))$ (and the resulting action by \mathbf{H} is the natural action by automorphisms.) \square

We note that each definable set in \mathcal{T}^* comes equipped with an action of the binding group $\mathbf{Aut}(\ast)$, and with it, the first direction could likewise be described as $\mathbf{X} = i^*(j_!(\mathbf{X}^*))$.

2.2.5. Pushouts. Let $g : \mathbf{K} \rightarrow \mathbf{G}$ and $h : \mathbf{K} \rightarrow \mathbf{H}$ be maps of definable groupoids, and assume that g is fully faithful. We construct another definable groupoid $\mathbf{G} \otimes_{\mathbf{K}} \mathbf{H}$ that can be viewed as the pushout of \mathbf{G} and \mathbf{H} over \mathbf{K} , as follows: For objects, we let $(\mathbf{G} \otimes_{\mathbf{K}} \mathbf{H})_0 = \mathbf{G}_0 \amalg \mathbf{H}_0$. If a, b are two such objects, we define the morphisms as follows:

- (1) If $a, b \in \mathbf{H}_0$, then $(\mathbf{G} \otimes_{\mathbf{K}} \mathbf{H})(a, b) = \mathbf{H}(a, b)$
- (2) If $a \in \mathbf{G}_0$ and $b \in \mathbf{H}_0$, morphisms from a to b are equivalence classes $v \otimes u$ of pairs $\langle v, u \rangle$, where $u \in \mathbf{G}(a, g(c))$, $v \in \mathbf{H}(h(c), b)$ for some $c \in \mathbf{K}_0$, and $\langle v, g(w) \circ u \rangle$ is equivalent to $\langle v \circ h(w), u \rangle$ for all $w \in \mathbf{K}_1$ for which the composition is defined. Morphisms from b to a are defined analogously.
- (3) If $a, b \in \mathbf{G}_0$ are both in the essential image of g , a morphism from a to b is similarly defined as an equivalence class $u' \otimes v \otimes u$, with $u, u' \in \mathbf{G}_1$ and $v \in \mathbf{H}_1$.
- (4) If either of $a, b \in \mathbf{G}_0$ is not in the essential image of g , then morphisms are the same as in \mathbf{G} .

The composition $(u' \otimes v \otimes u) \circ (u'_1 \otimes v_1 \otimes u_1)$ is defined as follows: There are $a, b \in \mathbf{K}_0$ such that $u \circ u'_1$ is a morphism from $g(a)$ to $g(b)$. Since g is fully faithful, it has the form $g(w)$ for a unique morphism w from a to b in \mathbf{K} . We define the composition to be $u' \otimes (v \circ h(w) \circ v_1) \otimes u_1$. It is clear that this is independent of the choices of representatives. The composition in the other cases is defined similarly.

There is an obvious map $h' : \mathbf{H} \rightarrow \mathbf{G} \otimes_{\mathbf{K}} \mathbf{H}$, and we define $g' : \mathbf{G} \rightarrow \mathbf{G} \otimes_{\mathbf{K}} \mathbf{H}$ by sending each object to itself, each morphism between objects not in the essential image of g to itself as well, and for $a, b \in \mathbf{G}_0$ in the essential image of g , and u a morphism from a to b , we set $g'(u) = (u \circ u'^{-1}) \otimes \mathbf{1}_{h(c)} \otimes u'$, where $u' : a \rightarrow g(c)$ is any morphism and $c \in \mathbf{K}_0$. We have an isomorphism α from $h' \circ h$ to $g' \circ g$, given on an object $c \in \mathbf{K}_0$ by $\mathbf{1}_{g(c)} \otimes \mathbf{1}_{h(c)}$. It is routine to check that everything is well defined, and also that the following statement holds.

Proposition 2.2.6. *Let $g : \mathbf{K} \rightarrow \mathbf{G}$, $h : \mathbf{K} \rightarrow \mathbf{H}$ and the rest of the notation be as above.*

- (1) *Given definable maps of groupoids $g_1 : \mathbf{G} \rightarrow \mathbf{F}$ and $h_1 : \mathbf{H} \rightarrow \mathbf{F}$, and an isomorphism $\beta : h_1 \circ h \rightarrow g_1 \circ g$, there is a unique map of groupoids $f : \mathbf{G} \otimes_{\mathbf{K}} \mathbf{H} \rightarrow \mathbf{F}$ that coincides with g_1 and h_1 on the objects, $f \circ g' = g_1$, $f \circ h' = h_1$ and such that $f \cdot \alpha = \beta$.*
- (2) *$h' : \mathbf{H} \rightarrow \mathbf{G} \otimes_{\mathbf{K}} \mathbf{H}$ is fully faithful. If g is a weak equivalence, then so is h' .*

Remark 2.2.7. We could make a similar construction where the set of objects is $\mathbf{G}_0 \coprod_{\mathbf{K}_0} \mathbf{H}_0$ in place of the disjoint union, and with α the identity. The last proposition provides a map from $\mathbf{G} \otimes_{\mathbf{K}} \mathbf{H}$ to this variant, which is easily seen to be a weak equivalence. We will use the two constructions interchangeably. \square

Remark 2.2.8. Without the assumption that one of the maps is fully-faithful, the pushout need not be definable. For example, when all groupoids are groups, this is the usual free product with amalgamation. \square

2.2.9. Maps of groupoids and of interpretations. With stable interpretations over \mathcal{T} , the assignment $\mathcal{T}^* \mapsto I_{\mathcal{T}^*/\mathcal{T}}$ are contravariantly functorial in \mathcal{T}^* , and fully-faithful: a stable interpretation $i : \mathcal{T}_1 \rightarrow \mathcal{T}_2$ over \mathcal{T} induces a functor $i^* : I_{\mathcal{T}_2/\mathcal{T}} \rightarrow I_{\mathcal{T}_1/\mathcal{T}}$ by composition.

In the other direction, if $f : \mathbf{G} \rightarrow \mathbf{H}$ is a map of definable groupoids, corresponding to internal covers $\mathcal{T}_{\mathbf{G}}$ and $\mathcal{T}_{\mathbf{H}}$, f determines a stable interpretation $i^f : \mathcal{T}_{\mathbf{H}} \rightarrow \mathcal{T}_{\mathbf{G}}$ over \mathcal{T} , that can be described in at least two ways:

- (1) An interpretation of $\mathcal{T}_{\mathbf{H}}$ over \mathcal{T} is determined by its value on the extended groupoid \mathbf{H}^* defined in $\mathcal{T}_{\mathbf{H}}$. We set $i^f(\mathbf{H}^*) = \mathbf{G}^* \otimes_{\mathbf{G}} \mathbf{H}$ (with respect to the given map f). This makes sense since the inclusion of \mathbf{G} in \mathbf{G}^* is a weak equivalence, and is an interpretation since the embedding of \mathbf{H} in $\mathbf{G}^* \otimes_{\mathbf{G}} \mathbf{H}$ is a weak equivalence that misses precisely one object $*$, and this completely determines its theory. To see that it is stable, we may first choose a parameter in \mathbf{G}^* . But then i^f is identified with one of the standard interpretations into \mathcal{T} .
- (2) Alternatively, we may use Cor 2.2.4 to identify definable sets in $\mathcal{T}_{\mathbf{G}}$ and in $\mathcal{T}_{\mathbf{H}}$ with \mathbf{G} - and \mathbf{H} -sets in \mathcal{T} . Then i^f is identified with f^* (in this approach, it is less direct to see that one gets a stable interpretation).

It is easy to verify that $(i^f)^* = f$ (after identifying \mathbf{G} with its image in $I_{\mathcal{T}_{\mathbf{G}}/\mathcal{T}}$ via 2.2.2, and similarly for \mathbf{H}). However, not every stable interpretation $i : \mathcal{T}_{\mathbf{H}} \rightarrow \mathcal{T}_{\mathbf{G}}$ (over \mathcal{T}) is of the form i^f for some $f : \mathbf{G} \rightarrow \mathbf{H}$. The other source of interpretations comes from the other operation described in §2.2.3: when f is a weak equivalence, the composition of f with the inclusion of \mathbf{H} in \mathbf{H}^* is a weak equivalence, so restricting to the image of f (on the objects), we obtain an interpretation i_f of \mathbf{G}^* (hence of $\mathcal{T}_{\mathbf{G}}$).

Proposition 2.2.10. *Let \mathbf{G} and \mathbf{H} be two definable groupoids, with associated covers $\mathcal{T}_{\mathbf{G}}$ and $\mathcal{T}_{\mathbf{H}}$. Then every stable interpretation $i : \mathcal{T}_{\mathbf{G}} \rightarrow \mathcal{T}_{\mathbf{H}}$ over \mathcal{T} is obtained as a composition $i = i^f \circ i_g$, for some definable groupoid \mathbf{K} , definable map $f : \mathbf{H} \rightarrow \mathbf{K}$ and weak equivalence $g : \mathbf{G} \xrightarrow{\sim} \mathbf{K}$.*

Proof. \mathbf{H} embeds in $I_{\mathcal{T}_{\mathbf{H}}/\mathcal{T}}$ via Prop 2.2.2, which maps via i^* to $I_{\mathcal{T}_{\mathbf{G}}/\mathcal{T}}$. We set $f : \mathbf{H} \rightarrow \mathbf{K}$ to be the restriction of i^* to \mathbf{H} , where \mathbf{K} denotes any definable weakly equivalent subgroupoid of $I_{\mathcal{T}_{\mathbf{G}}/\mathcal{T}}$, which also contains \mathbf{G} . Then g is the inclusion of \mathbf{G} in \mathbf{K} . \square

As in the construction of the pushout, we may choose \mathbf{K} so that its objects are the disjoint union of the objects of \mathbf{G} and \mathbf{H} , and we will always assume that this is the case. In the case when i is a bi-interpretation, we recover the notion of equivalence from [4, §3].

2.2.11. Composition and isomorphisms. Assume that for groupoids \mathbf{F} , \mathbf{G} and \mathbf{H} in \mathcal{T} , we are given interpretations $i : \mathcal{T}_{\mathbf{F}} \rightarrow \mathcal{T}_{\mathbf{G}}$ and $j : \mathcal{T}_{\mathbf{G}} \rightarrow \mathcal{T}_{\mathbf{H}}$, represented by $g_1 : \mathbf{F} \xrightarrow{\sim} \mathbf{K}_1$, $f_1 : \mathbf{G} \rightarrow \mathbf{K}_1$, $g_2 : \mathbf{G} \xrightarrow{\sim} \mathbf{K}_2$ and $f_2 : \mathbf{H} \rightarrow \mathbf{K}_2$ as in 2.2.10. Since g_2 is a weak equivalence, we may form the pushout $\mathbf{K} = \mathbf{K}_1 \otimes_{\mathbf{G}} \mathbf{K}_2$. By Prop 2.2.6, the map from \mathbf{K}_1 to \mathbf{K} is a weak equivalence, and therefore so is the composed map g . Hence, g along with the composed map $f : \mathbf{H} \rightarrow \mathbf{K}$ represent a stable interpretation of $\mathcal{T}_{\mathbf{F}}$ in $\mathcal{T}_{\mathbf{H}}$. To conform with the decision about the objects of the representing groupoid \mathbf{K} , we remove the intermediate two copies of \mathbf{G}_0 , and denote the resulting groupoid by $\mathbf{K}_2 \circ_{\mathbf{G}} \mathbf{K}_1 = \mathbf{K}_2 \circ_{\mathbf{G}} \mathbf{K}_1$ (though it does depend on the additional data). The following is a direct calculation.

Corollary 2.2.12. *In the above situation, the maps $g : \mathbf{F} \xrightarrow{\sim} \mathbf{K}_2 \circ_{\mathbf{G}} \mathbf{K}_1$ and $f : \mathbf{H} \rightarrow \mathbf{K}_2 \circ_{\mathbf{G}} \mathbf{K}_1$ represent the composed interpretation $j \circ i$.*

Finally, we consider isomorphisms of interpretations between (stable) interpretations of internal covers over \mathcal{T} .

Corollary 2.2.13. *Let $i, j : \mathcal{T}_{\mathbf{G}_1} \rightarrow \mathcal{T}_{\mathbf{G}_2}$ be two stable interpretations of internal covers over \mathcal{T} . Assume i is represented by groupoid maps $i_1 : \mathbf{G}_1 \xrightarrow{\sim} \mathbf{H}_1$ and $i_2 : \mathbf{G}_2 \rightarrow \mathbf{H}_1$, and j by $j_1 : \mathbf{G}_1 \xrightarrow{\sim} \mathbf{H}_2$, $j_2 : \mathbf{G}_2 \rightarrow \mathbf{H}_2$, where each set of objects of \mathbf{H}_n the disjoint union of the objects of \mathbf{G}_1 and \mathbf{G}_2 (realized by the object parts of i_k and j_k).*

Then there is a natural bijection between isomorphisms $\alpha : i \rightarrow j$ (over \mathcal{T}) and isomorphisms $\tilde{\alpha} : \mathbf{H}_1 \rightarrow \mathbf{H}_2$ which are the identity on the images of $\mathbf{G}_1, \mathbf{G}_2$.

As an example, if \mathbf{G}_1 and \mathbf{G}_2 are groups, then each \mathbf{H}_i corresponds to a $\mathbf{G}_1 - \mathbf{G}_2$ bi-torsor, and an isomorphism of the corresponding interpretations corresponds to an isomorphism of such bi-torsors.

Proof. Let \mathbf{P}_l be the set of arrows in \mathbf{H}_l with domain in \mathbf{G}_1 and codomain in \mathbf{G}_2 . This is a \mathbf{G}_1 -set, with structure given by the domain map and composition. The interpretation i takes \mathbf{P}_1 to the \mathbf{G}_2 -set given by viewing the arrows in \mathbf{P}_1 in the other direction, and likewise with j and \mathbf{P}_2 . So the map α maps \mathbf{P}_1 to \mathbf{P}_2 , compatibly with the composition. This is the same as giving an isomorphism $\tilde{\alpha}$ as in the statement. The rest of the structure is induced by the \mathbf{P}_i , so α is determined by $\tilde{\alpha}$. Conversely, each $\tilde{\alpha}$ as in the statement extends to an interpretation. \square

The description above exhibits the groupoid associated to an expansion as interpretations of \mathcal{T}^* in \mathcal{T} . In [4], it was suggested that definable sets of an internal cover of \mathcal{T} can be viewed a generalised imaginary sorts of

\mathcal{T} . With this point of view, it is natural to ask for the structure classifying interpretations of such sorts as well. However, such generalised sorts have more structure: in addition to the sorts themselves and maps between them (interpretations), we also have maps between maps. The notion of equivalence should be modified as well: it is no longer reasonable to expect a bijection on the level of morphisms. In fact, as the 1-dimensional case already shows, it not reasonable to expect even a map.

3. GENERALISED IMAGINARIES

We now suggest how internal covers can play the role of definable sets in the above description, by going one dimension higher.

3.1. Higher internal covers.

Definition 3.1.1. Let \mathcal{T} be a theory, \mathcal{T}_1 and \mathcal{T}_2 internal covers of \mathcal{T} . For every set of parameters A for \mathcal{T} , we denote by $\text{Hom}_{\mathcal{T}}(\mathcal{T}_1, \mathcal{T}_2)(A)$ the groupoid whose objects are stable interpretations of \mathcal{T}_1 in \mathcal{T}_{2A} , over \mathcal{T}_A , and whose morphisms are isomorphisms of interpretations over \mathcal{T} .

Thus, what we denoted above I is $\text{Hom}_{\mathcal{T}}(\mathcal{T}^*, \mathcal{T})$. Similar to that case, $\text{Hom}_{\mathcal{T}}(\mathcal{T}_1, \mathcal{T}_2)$ is definable in \mathcal{T} : if $\mathcal{T}_i = \mathcal{T}_{\mathbf{G}_i}$ for \mathcal{T} -definable groupoids $\mathbf{G}_1, \mathbf{G}_2$, each interpretation above can be described like in §2.2.9 as given by certain definable maps $\mathbf{G}_i \rightarrow \mathbf{H}$, a definable condition. Similarly, isomorphisms between interpretations are given by the definable families of maps as in Cor 2.2.13, uniform in the \mathbf{H}_i (cf. §3.3.6 for a more detailed description.)

An interpretation between theories extends to internal sorts: If $i : \mathcal{T} \rightarrow \mathcal{S}$ is an interpretation, and $\tilde{\mathcal{T}}$ is an internal cover of \mathcal{T} , associated to the groupoid \mathbf{G} in \mathcal{T} , we denote by $i(\tilde{\mathcal{T}})$ the internal cover of \mathcal{S} associated to $i(\mathbf{G})$.

We now wish to define (slightly) higher analogs of stable embeddings and internal covers. One discrepancy with the 1-dimensional case occurs as follows: If \mathcal{T} is an internal cover of \mathcal{T}_0 , we might be interested in only part of the structure on \mathcal{T} when considering, for example, the Galois group. As long as this partial structure includes the definable sets witnessing the internality, this can be done by replacing \mathcal{T} with a reduct including only those definable sets. In the higher version, definable sets are replaced by definable groupoids in \mathcal{T} (equivalently, internal covers), and again we may wish to restrict to a partial collection. However, there is no reason to expect that this partial collection is the full collection of definable groupoids in some reduct of \mathcal{T} . Furthermore, the internality condition for 0-definable sets automatically implies it for definable sets over parameters. Again, there is no reason to expect a similar statement for groupoids. For this reason, our definition depends on the auxiliary data Γ consisting of families of definable groupoids, which are the groupoids we wish to preserve. More precisely, we have the following.

Definition 3.1.2. Let \mathcal{T} be a theory. The data of *distinguished covers* for \mathcal{T} consists of the following:

- (1) A definable family Γ_0 of internal covers of \mathcal{T} (equivalently, of definable groupoids in \mathcal{T}).
- (2) A definable family of interpretations over \mathcal{T} between any two covers $\mathcal{T}_1, \mathcal{T}_2 \in \Gamma_0$, depending definably on $\mathcal{T}_1, \mathcal{T}_2$ and closed under composition (the full definable family is denoted by Γ_1).
- (3) For every two interpretations $f, g : \mathcal{T}_1 \rightarrow \mathcal{T}_2$ in Γ_1 , a definable family of isomorphisms from f to g , closed under composition and restricting to the identity on \mathcal{T} . Again we assume that the family of all such isomorphisms is uniformly definable in f, g , and denote it by Γ_2 .

If \mathcal{T}_0 is a reduct of \mathcal{T} , we will say that $\Gamma = \langle \Gamma_0, \Gamma_1, \Gamma_2 \rangle$ is over \mathcal{T}_0 if the parameters for the definable families above range over definable sets in \mathcal{T}_0 .

We note that in terms of definable groupoids, the closure under composition translates to closure under the composition operation from §2.2.11.

If a theory \mathcal{T} is given with a collection Γ of distinguished covers, we will often omit further explicit reference to Γ , and call them *admissible covers*. We modify notions like bi-interpretation etc., to be with respect to Γ . In particular, the notation $\text{Hom}_{\mathcal{T}}$ will refer to admissible covers and admissible maps.

Definition 3.1.3. Let $i : \mathcal{T} \rightarrow \mathcal{S}$ be an interpretation, and let Γ be a collection of distinguished covers of \mathcal{T} . We say that the interpretation i is *2-stable* (with respect to Γ) if for every two internal covers $\mathcal{T}_1, \mathcal{T}_2$ in Γ over each \mathcal{T} -structure A , the natural map $i(\text{Hom}_{\mathcal{T}_A}(\mathcal{T}_1, \mathcal{T}_2)) \rightarrow \text{Hom}_{\mathcal{S}_A}(i(\mathcal{T}_1), i(\mathcal{T}_2))$ is an equivalence.

If Γ is omitted, we take it to be all definable groupoids in \mathcal{T} , and all definable morphisms among them.

The expression $i(\text{Hom}_{\mathcal{T}}(\mathcal{T}_1, \mathcal{T}_2))$ makes sense, since, as we had noted above, $\text{Hom}_{\mathcal{T}}(\mathcal{T}_1, \mathcal{T}_2)$ is definable in \mathcal{T} .

We note:

Proposition 3.1.4. *A stable interpretation $i : \mathcal{T}_1 \rightarrow \mathcal{T}_2$ is 2-stable.*

Proof. We may replace \mathcal{T} by \mathcal{T}_A , and thus assume that $A = \emptyset$. Let $\mathcal{T}_1, \mathcal{T}_2$ be internal covers of \mathcal{T} . The statement is invariant when replacing each cover with a bi-interpretable one (over \mathcal{T}). Hence, we may assume that $\mathcal{T}_1 = \mathcal{T}_{\mathbf{G}_1}$ and $\mathcal{T}_2 = \mathcal{T}_{\mathbf{G}_2}$, the covers associated to definable connected groupoids $\mathbf{G}_1, \mathbf{G}_2$ in \mathcal{T} .

According to §2.2.9, we may choose \mathbf{G}_1 and \mathbf{G}_2 so that a stable interpretation of $i(\mathcal{T}_1)$ in $i(\mathcal{T}_2)$ corresponds to a definable map of groupoids from $i(\mathbf{G}_2)$ to $i(\mathbf{G}_1)$. Since i is stable, this map comes from a map in \mathcal{T} (and similarly for morphisms). \square

The definition of a 2-cover is analogous to that of an internal cover, as formulated in 2.1.2:

Definition 3.1.5. A *2-internal cover* of a theory \mathcal{T} consists of a theory \mathcal{T}^* , a collection Γ of families of internal covers of \mathcal{T}^* over \mathcal{T} , and a stable embedding $\mathcal{T} \rightarrow \mathcal{T}^*$ such that $\langle \mathcal{T}^*, \Gamma \rangle$ admits a 2-stable interpretation p in \mathcal{T}_A , over \mathcal{T} .

More explicitly, we require that each internal cover in Γ is bi-interpretable, over parameters in \mathcal{T}^* , with one coming from \mathcal{T} , in a manner coherent with interpretations over \mathcal{T}^* . Or, via the equivalence with groupoids, that for each family of definable groupoids in Γ there is a set of parameters B in \mathcal{T}^* such that each groupoid in the family is equivalent, over B , to one coming from \mathcal{T} (again, in a coherent manner).

As in the 1-dimensional case, the typical examples will come from higher dimensional groupoids, which we review next.

3.2. Higher categories and higher groupoids. We recall a few definitions from homotopy theory and higher category theory, adapted to our language and setup. Though our main references are [8] and [9], the ideas seem to originate in [5] (and in the main case of groupoids, much more classically). We will be interested in the notions of n -category and n -groupoid discussed in [8, §2.3.4] (through most other parts of the paper we are interested in the case $n = 2$, but here it is convenient and harmless to work in general.) There, they are defined as special cases of quasi-categories and Kan simplicial sets, respectively, but for us it will be more convenient to use terminology that makes explicit the finite nature of these structures. The following definitions are a variant of the description in [8, §2.3.4.9], which gives an equivalent condition (in the case of simplicial sets).

For each $i \in \mathbb{N}$, we denote by $[i]$ the ordered set $\{0, \dots, i\}$. For $k \in [i]$, we identify k with the map $[0] \rightarrow [i]$ taking 0 to k (writing k_i if needed), and we let $\widehat{k} = \widehat{k}_i : [i-1] \rightarrow [i]$ be the unique increasing map with k not in the image. We fix a natural number n (one could also allow $n = \infty$ to obtain the usual definitions of quasi-categories and spaces, but we will not use them).

Definition 3.2.1. The signature Σ_n of n -simplicial sets consists of:

- (1) A sort \mathbf{G}_i , for $0 \leq i \leq n+1$
- (2) For each weakly increasing map $t : [i] \rightarrow [j]$, where $i, j \leq n+1$, a function symbol $d_t : \mathbf{G}_j \rightarrow \mathbf{G}_i$.

3.2.2. Notation. We define the following auxiliary notation. We let \mathbf{G}_{-1} be the one element set. For each $0 \leq m \leq n+1$, and $i \leq m$, the map $d_{\widehat{i}} : \mathbf{G}_m \rightarrow \mathbf{G}_{m-1}$ is called the *i -th face map*. We denote by $\partial = \partial_m : \mathbf{G}_m \rightarrow \mathbf{G}_{m-1}^{m+1}$ the Cartesian product of these maps, and by $\widehat{\partial}^i : \mathbf{G}_m \rightarrow \mathbf{G}_{m-1}^m$ the Cartesian product with i omitted. If $g \in \mathbf{G}_m$ and $t : [k] \rightarrow [m]$, we sometimes write g_t in place of $d_t(g)$ (in particular, for $t = \widehat{i}$ or $t = i$).

For $m \geq -1$, the set \mathbf{G}_{m+1}° of $m+1$ -cycles is the definable subset of \mathbf{G}_m^{m+2} given by the conjunction of the equations $d_i(x_j) = d_k(x_l)$ for all $0 \leq j, l \leq m+1$, $0 \leq i, k \leq m$ satisfying $\widehat{j} \circ \widehat{i} = \widehat{l} \circ \widehat{k} : [m-1] \rightarrow [m+1]$ (So no conditions when $m = 0$. Note that $m+1$ -cycles are potential boundaries of $m+1$ -dimensional elements, but are themselves m -dimensional. This is compatible with the notation in [8].)

For each $0 \leq i \leq m+1$, the set $\Lambda_i^{m+1}(\mathbf{G})$ of i -th $m+1$ -horns is the subset of \mathbf{G}_m^{m+1} defined by the same conditions, with x_i omitted. Hence, the projection $\pi_{\widehat{i}} : \mathbf{G}_m^\circ \rightarrow \mathbf{G}_{m-1}^m$ omitting the i -th coordinate takes values in $\Lambda_i^m(\mathbf{G})$.

We extend the notation by inductively setting $\mathbf{G}_m = \mathbf{G}_m^\circ$ for $m > n+1$, and $d_i : \mathbf{G}_m \rightarrow \mathbf{G}_{m-1}$ the i -th projection ($0 \leq i \leq m$). Consequently, all the above notation makes sense for arbitrary natural number m .

Definition 3.2.3. Let $n \geq 1$. The theory \mathcal{C}_n of n -categories in this signature says:

- (1) $d_{tos} = d_s \circ d_t$ for s, t composable, d_t is the identity whenever t is. It follows that ∂_m takes values in \mathbf{G}_m° and $\widehat{\partial}_m^i$ in $\Lambda_i^m(\mathbf{G})$.
- (2) For each $0 < m \leq n+1$ and each $0 < i < m$, the map $\widehat{\partial}_m^i : \mathbf{G}_m \rightarrow \Lambda_i^m(\mathbf{G})$ is surjective.
- (3) For $m = n+1, n+2$, $\widehat{\partial}_m^i$ is bijective for each $0 < i < m$.

The theory \mathcal{G}_n of n -groupoids is the extension of \mathcal{C}_n where the conditions above are required to also hold for $i = 0, m$.

Note that by the third condition, the set \mathbf{G}_{n+1} is completely determined by the rest of the data. However, it is still convenient to have it for the statement of the axiom. It follows from the axioms that the unique map $d_f : \mathbf{G}_0 \rightarrow \mathbf{G}_m$ is injective, and we will use it to identify \mathbf{G}_0 with its image in each \mathbf{G}_m , writing a or a^m for $d_f(a)$ (this map assigns to each object a the m -dimensional identity morphisms at a).

The first condition (when $n = \infty$) is the usual definition of a simplicial set, the second is the definition of quasi-category (or space, in the case of a groupoid, where it is called the Kan condition), and the third specifies that the object is an n -category, rather than a quasi-category. By a *definable n -category* or a *definable n -groupoid* in \mathcal{T} we mean an interpretation in \mathcal{T} of the respective theory.

The intuition is, roughly, that the horns represent configurations of (higher) composable arrows, but the composition (represented by the element g) need not be uniquely determined, except on the highest dimension. We refer to the first chapters of [8] for further explanations, but explain how the case $n = 1$ of the formalism recovers usual categories and groupoids:

Example 3.2.4. A category can be viewed as a 1-category in the above sense by taking \mathbf{G}_0 the set of objects, \mathbf{G}_1 the set of morphisms, and \mathbf{G}_2 the set of pairs of composable morphisms (as we are forced by the axioms). The maps

$d_{\widehat{0}}, d_{\widehat{1}} : \mathbf{G}_1 \rightarrow \mathbf{G}_0$ are the codomain and domain maps, the unique map $\mathbf{G}_0 \rightarrow \mathbf{G}_1$ assigns to each object its identity, and the maps $d_{\widehat{0}}, d_{\widehat{2}}, d_{\widehat{1}} : \mathbf{G}_2 \rightarrow \mathbf{G}_1$ are the two projections and the composition. The only non-trivial instances of the third conditions are when $m = 2$ and $i = 1$, which asserts that any two composable arrows have a unique composition, and when $m = 3$, which corresponds to associativity of the composition.

Conversely, each 1-category determines a category by reversing this process (and likewise for groupoids). \square

As in the 1-dimensional case, the axioms imply that for $0 < i < n + 1$, the relation \mathbf{G}_{n+1}° is the graph of a ‘‘composition’’ function $c_i : \Lambda_i^n(\mathbf{G}) \rightarrow \mathbf{G}_n$, by projecting to the i -coordinate. For n -groupoids, we also have such maps for $i = 0, n + 1$.

Remark 3.2.5. If \mathbf{G} is an n -category, and $m > n$, our extension of the notation determines a canonical way of viewing \mathbf{G} as a Σ_m structure, and as such it is an m -category. Consequently, we will view \mathbf{G} as an m -category for each $m > n$. If \mathbf{G} was an n -groupoid, it will similarly be an m -groupoid for $m > n$. \square

3.2.6. Homotopy sets. The definition of homotopy sets admit a definable version. Let \mathbf{G} be an n -groupoid, and let $b \in \mathbf{G}_m^\circ$ ($m \geq 0$). We let $\mathbf{S}(\mathbf{G}, b) = \partial_m^{-1}(b)$ be the set of elements of \mathbf{G}_m with boundary b . For $\alpha, \beta \in \mathbf{S}(\mathbf{G}, b)$, we write $\alpha \sim \beta$ (or $\alpha \sim_b \beta$) if some $h \in \mathbf{G}_{m+1}$ satisfies $h_{\widehat{0}} = \alpha$, $h_{\widehat{1}} = \beta$ and $h_{\widehat{i}} = d_t(\alpha)_{\widehat{i}}$, where $t : [m + 1] \rightarrow [m]$ is the surjective map with $t(1) = 0$ (so h is a homotopy from α to β , relative to the boundary b). This is an equivalence relation by the Kan condition. Note that when $m \geq n$, this relation coincides with equality.

For $a \in \mathbf{G}_0$ and $k \geq 0$, we write $\mathbf{S}_k(\mathbf{G}, a)$ for $\mathbf{S}(\mathbf{G}, b)$, where b is the constant boundary with value a in \mathbf{G}_k° . These are a -definable sets, whose elements correspond to the set of pointed maps from the k -sphere to \mathbf{G} with base point a (note that $\mathbf{K}_0(\mathbf{G}, a) = \mathbf{G}_0$ does not actually depend on a). The k -th *homotopy set* of \mathbf{G} at a is the quotient $\pi_k(\mathbf{G}, a) = \mathbf{S}_k(\mathbf{G}, a) / \sim$ (in the case of usual simplicial sets, this is one of the equivalent definitions by [9, 00W1]).

If $f : \mathbf{G} \rightarrow \mathbf{H}$ is groupoid map, it commutes with all the structure above, and therefore induces definable maps of sets $\pi_k(f, a) : \pi_k(\mathbf{G}, a) \rightarrow \pi_k(\mathbf{H}, f(a))$.

Definition 3.2.7. A definable map $f : \mathbf{G} \rightarrow \mathbf{H}$ of n -groupoids is a *weak equivalence* if $\pi_k(f, a) : \pi_k(\mathbf{G}, a) \rightarrow \pi_k(\mathbf{H}, f(a))$ is a bijection for all $0 \leq k \leq n$ and $a \in \mathbf{G}_0$.

Remark 3.2.8. More explicitly, for non-empty \mathbf{G} , the map $f : \mathbf{G} \rightarrow \mathbf{H}$ is a weak equivalence if and only if the following conditions are satisfied for each $n \geq k \geq 0$ and each $a \in \mathbf{G}_0$:

- (1) For every $g_0, g_1 \in \mathbf{S}_k(\mathbf{G}, a)$, if $f(g_1) \sim f(g_2)$ then $g_1 \sim g_2$.

(2) For every $h \in \mathbf{S}_k(\mathbf{H}, f(a))$, there is $g \in \mathbf{S}_k(\mathbf{G}, a)$ with $f(g) \sim h$.

Alternatively, f is a weak equivalence if and only if it induces a surjective map on \mathbf{S} -classes, i.e., for each \mathbf{G} -cycle b , and each $v \in \mathbf{S}(\mathbf{H}, f(b))$, there is $u \in \mathbf{S}(\mathbf{G}, b)$ with $f(u) \sim v$. (To prove these equivalences, it suffices to show that they hold in each model, where each of these conditions is equivalent to homotopy equivalence, [9, 00WV].) \square

Definition 3.2.9. The n -groupoids \mathbf{G}_1 and \mathbf{G}_2 are *equivalent* if there are weak equivalences $f_1 : \mathbf{G}_1 \rightarrow \mathbf{H}$ and $f_2 : \mathbf{G}_2 \rightarrow \mathbf{H}$ for some \mathbf{H} .

Example 3.2.10. Let \mathbf{G}, \mathbf{H} be definable groupoids, identified with a 1-groupoid as in Example 3.2.4. A map $f : \mathbf{G} \rightarrow \mathbf{H}$ is a functor. For $k = 0$, the first condition for says that if a, b are objects of \mathbf{G} , and there is a morphism between $f(a)$ and $f(b)$ in \mathbf{H} , then there is a morphism from a to b in \mathbf{G} . The second condition says that every object h of \mathbf{H} has a morphism to an object in the image of f . Together, this part implies that f induces a bijection on isomorphism classes.

For $k = 1$, the first condition says that if g_1, g_2 are automorphisms of a such that $f(g_1) = f(g_2)$, then $g_1 = g_2$, i.e., that f is faithful. The second condition says that f is full.

Hence, f is a weak equivalence if and only if it is a weak equivalence in the sense of §2.2. In particular, our notion of equivalence coincides with [4, §3]. \square

As in §2.2.11, equivalence of n -groupoids is an equivalence relation: If \mathbf{H} and \mathbf{H}' witness that \mathbf{G}_2 is equivalent to \mathbf{G}_1 and \mathbf{G}_3 , respectively, the (suitably defined) pushout $\mathbf{H} \otimes_{\mathbf{G}_2} \mathbf{H}'$ witnesses the equivalence of $\mathbf{G}_1, \mathbf{G}_3$.

Remark 3.2.11. The group operation on $\pi_k(\mathbf{G}, a)$ (for $k > 0$) is also definable, but we will not use this. \square

Remark 3.2.12. The equivalence of our definitions of homotopy groups and weak equivalence with other formulations that appear, for example, in [9, 00V2] does not hold in the definable setting, in general. For example, the analogue of Whitehead's Theorem ([9, 00WV]) is usually false (as seen already in the one-dimensional setting). \square

3.2.13. *Morphism groupoids.* Our next goal is to define the space of morphisms between two objects a, b of an n -category \mathbf{G} , and obtain a (weak) version of the Yoneda embedding that will make sense in the definable setting.

Let \mathbf{G} be an n -category, and let $a, b \in \mathbf{G}_0$ be two objects. As in [8, §1.2.2], we define the Σ_{n-1} -structure $\underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, b)$ by

$$\underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, b)_k = \{g \in \mathbf{G}_{k+1} \mid g_0 = a, g_{\bar{0}} = b^k\}, \quad (1)$$

For $k \leq n$. The structure maps are given by $t \mapsto d_{t^+}^{\mathbf{G}}$, where $t^+ : [u+1] \rightarrow [k+1]$ is given by $t^+(i+1) = t(i) + 1$ for $i \in [u]$ and $t^+(0) = 0$. It is clear that $\underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, b)$ is uniformly definable over a, b when \mathbf{G} is definable. It

follows from [8, §§4.2.1.8, 2.3.4.18, 2.3.4.19] that this structure is equivalent to an $n - 1$ -groupoid, but since we are not working up to equivalence, we need to prove that it is already an $n - 1$ by itself (which we do below).

If we fix a “generic” object $v \in \mathbf{G}_0$, the assignment $b \mapsto \underline{\mathbf{Hom}}^L(v, b)$ looks like the object part of the Yoneda embedding for usual categories. One could hope that this is part of a higher Yoneda embedding in our situation as well. However, since there is no composition function for morphisms in \mathbf{G} , such an embedding does not exist as a functor (it exists non-canonically for set-theoretic quasi-categories, but not definably). Instead, we have the following situation (explained in [8, §2.1]): There is an n -category $\mathbf{G}_{v/}$ ([8, §2.3.4.10]), defined by $(\mathbf{G}_{v/})_k = \{g \in \mathbf{G}_{k+1} | g_0 = v\}$, and a map of n -categories $\pi : \mathbf{G}_{v/} \rightarrow \mathbf{G}$, given by $\pi(g) = g_{\widehat{0}}$. By definition, the fibre of this map over $b \in \mathbf{G}_0$ is $\underline{\mathbf{Hom}}_{\mathbf{G}}^L(v, b)$. Moreover, this map is a *left fibration* ([8, §2.1.22]): Given $g \in \Lambda_i^k(\mathbf{G}_{v/})$, for $i < k$, any “filling” $h \in \mathbf{G}_k$ of $\pi(g)$ (so that $\widehat{\partial}^i(h) = \pi(g)$) can be lifted to a filling $\tilde{h} \in \mathbf{G}_{v/}$ with $\widehat{\partial}^i(\tilde{h}) = g$ and $\pi(\tilde{h}) = h$. It follows from this that the association $b \mapsto \pi^{-1}(b)$ behaves like a functor of b , but this is only precisely true in the homotopy category.

We will show that in the case that h above is invertible, the statement above holds for our definable version of equivalence. To do this, we show that the map π behaves like a local system: the fibres can be continued along (suitable) contractible pieces. The pieces we have in mind, which we denote by D^l , for $l \geq 0$, are the simplicial sets defined by $D_k^l = \{0, \dots, l\}^{\{0, \dots, k\}}$ (all maps, no necessarily increasing, from $[k]$ to $[l]$), and structure maps given by composition. We will often write elements of D_k^l as words of length $k + 1$ in the “digits” $0, \dots, l$. By the usual Yoneda lemma, maps $D^l \rightarrow D^m$ correspond (via composition) to functions $\{0, \dots, l\} \rightarrow \{0, \dots, m\}$ (via composition). Note that homotopically, all these maps are weak equivalences, and in particular the map to the point D^0 , so that all D^l are contractible.

We now extend the definition of morphisms as follows: for \mathbf{G} a definable n -category, let $a \in \mathbf{G}_0$ be an object, and let $f : D^l \rightarrow \mathbf{G}$ be a map of simplicial sets (perhaps over parameters). We define a Σ_{n-1} -structure $\underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, f)$ as follows: For each $k \leq n$,

$$\underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, f)_k = \{\langle g, e \rangle \in \mathbf{G}_{k+1} \times D_k^l | g_0 = a, g_{\widehat{0}} = f(e)\} \quad (2)$$

with structure maps given as before by $\langle g, e \rangle \mapsto \langle d_{t^+}(g), e \circ t \rangle$ for each weakly increasing function $t : [u] \rightarrow [k]$. In other words, $\underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, f)$ is the pullback under f of the map $\pi : \mathbf{G}_{v/} \rightarrow \mathbf{G}$ above. For $l = 0$ and f mapping the point D^0 to b , we recover the previous definition. In general, the projection determines a map $\underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, f) \rightarrow D^l$ of simplicial sets, which can be viewed as the “restriction” of π to D^l . If $h : D^r \rightarrow D^l$ is a map of simplicial sets, there is an induced map $\hat{h} : \underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, f \circ h) \rightarrow \underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, f)$, given by $\hat{h}(\langle g, e \rangle) = \langle g, h(e) \rangle$.

Proposition 3.2.14. *Let \mathbf{G} , $a \in \mathbf{G}_0$ and $f : D^l \rightarrow \mathbf{G}$ be as above.*

- (1) The structure $\underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, f)$ is an $n - 1$ -groupoid.
(2) For any map $h : \{0, \dots, r\} \rightarrow \{0, \dots, l\}$ (identified with the corresponding map $D^r \rightarrow D^l$), the induced map $\hat{h} : \underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, f \circ h) \rightarrow \underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, f)$ is a weak equivalence.

Proof. (1) Let $\mathbf{H} = \underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, f)$. It is clear that \mathbf{H} is a simplicial definable set. To check the Kan condition, we prove a stronger claim, namely, that the projection $\pi : \mathbf{H} \rightarrow D^l$ is a Kan fibration: given a horn element $h \in \Lambda_i^m(\mathbf{H})$ and a pre-image $d \in D_m^l$ with $\widehat{\partial}_m^i(d) = \pi(h)$, there is $\tilde{d} \in \mathbf{H}_m$ with $\pi(\tilde{d}) = d$ and $\widehat{\partial}_m^i(\tilde{d}) = h$.

Let $h \in \Lambda_i^m(\mathbf{H})$ be a horn element as above, with $0 \leq i \leq m \leq n + 2$. Such an element is given by a matching sequence of element $h^j = \langle g^j, e^j \rangle$, for $j \in [m]$, $j \neq i$, with $g^j \in \mathbf{G}_m$ and with $\pi(h) = \langle e^0, \dots, e^m \rangle$ an element of $\Lambda_i^m(D^l)$. In D^l , each such horn element comes from a *unique* element of D_m^l . Let $e \in D_m^l$ be this element, and let $g^{-1} = f(e) \in \mathbf{G}_m$.

We claim that $\tilde{g} = \langle g^{-1}, g^0, \dots, g^m \rangle \in \mathbf{G}_m^{m+1}$ is in $\Lambda_{i+1}^{m+1}(\mathbf{G})$. To show that, we need to show that if $\widehat{b} \circ \widehat{a} = \widehat{d} \circ \widehat{c} : [m-1] \rightarrow [m+1]$ for some $b, d \in [m+1]$, $b, d \neq i+1$ and $a, c \in [m]$, then $g^{b-1}_{\widehat{a}} = g^{d-1}_{\widehat{c}}$.

Assume first that $a, b, c, d \geq 1$. The assumption on h implies that

$$h^{b-1}_{\widehat{a-1}_{m-1}} = h^{d-1}_{\widehat{c-1}_{m-1}} \quad (3)$$

whenever a, b, c, d satisfy

$$\widehat{b-1}_m \circ \widehat{a-1}_{m-1} = \widehat{d-1}_m \circ \widehat{c-1}_{m-1} \quad (4)$$

Equation (3) implies that $g^{b-1}_{\widehat{a-1}_{m-1}^+} = g^{d-1}_{\widehat{c-1}_{m-1}^+}$ under this condition. But $\widehat{j}_{m-1}^+ = \widehat{j+1}_m$ for all $j \in [m-1]$, so we find that $g^{b-1}_{\widehat{a}_m} = g^{d-1}_{\widehat{c}_m}$ whenever eq. (4) holds. But eq. (4) is equivalent to

$$\widehat{b}_{m+1} \circ \widehat{a}_m = \widehat{d}_{m+1} \circ \widehat{c}_m \quad (5)$$

so we obtain the required condition when $a, b, c, d \geq 1$.

If $b = 0$ or $d = 0$, the corresponding element of \mathbf{G}_m is g^{-1} . In this case, the condition follows from the definition of $\underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, f)$: for example, if $b = 0$ we must have $c = 0$ and $d = a + 1$, so we need to show that $g^a_{\widehat{0}} = f(e)_{\widehat{a}} = f(e_{\widehat{a}}) = f(e^a)$, and we are done. If $a = 0$ or $c = 0$, the condition forces $b = 0$ or $d = 0$, so we are back to the same case.

This concludes the proof that $\tilde{g} \in \Lambda_{i+1}^{m+1}(\mathbf{G})$. If $i < m$, the Kan condition on \mathbf{G} implies that that we may find $g \in \mathbf{G}_{m+1}$ restricting to \tilde{g} . It follows that $g_0 = a$ and $g_{\widehat{0}} = f(e)$, so that $\langle g, e \rangle$ solves the lifting problem. It follows from [8, §1.2.5.1] that the case $i < m$ is sufficient.

When $m = n$ or $m = n + 1$, the injectivity follows similarly from injectivity for \mathbf{G} (and for D^l).

- (2) We use Remark 3.2.8. An element in $\underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, f \circ h)_0$ is given by $g \in \mathbf{G}_1$ with $g_0 = a$ and $g_1 = f(h(e))$, where $e \in [u]$. Assume that $\langle s, c \rangle, \langle t, d \rangle \in \underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, f \circ h)_k$ satisfy $s_{\hat{0}} = t_{\hat{0}} = f(h(c)) = f(h(d)) = h(e)$ and $s_{\hat{i}} = t_{\hat{i}} = g$ for $k \geq i > 0$, so that they are elements of $\mathbf{S}_k(\underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, f \circ h))$. Assume also that we are given some $w \in \mathbf{G}_{k+2}$ satisfying $w_{\hat{1}} = s$, $w_{\hat{2}} = t$ and $w_{\hat{i}} = g$ for $i > 2$, and some $v \in D^u_{k+1}$ with $v_{\hat{0}} = c$, $v_{\hat{1}} = d$ and $v_{\hat{i}} = e$ for $i > 1$, and with $f(h(v)) = w_{\hat{0}}$ (this is a homotopy from $\langle s, c \rangle$ to $\langle t, d \rangle$). Then $\langle w, h(v) \rangle$ is a homotopy from $\langle s, h(c) \rangle$ to $\langle t, h(d) \rangle$. The argument for $k = 0$ is similar (using that D^u is connected).

For the second condition of Remark 3.2.8, let $g \in \mathbf{G}_1$ be such that $g_0 = a, g_1 = f(e)$ for some $e \in [l]$, so that $b = \langle g, e \rangle$ represents a basepoint of $\underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, f)$, and let $\langle s, c \rangle \in \mathbf{S}_k(\underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, f), b)$. Then $c \in \mathbf{S}_k(D^l, e)$ is the constant function e . Let $e' \in [u]$, and let $\gamma \in D^l$ be some path from $h(e')$ to e . By the Kan condition above, there is an element s' of $\underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, f)$ above γ , restricting to s . This s' serves as a homotopy from s to an element over $h(e')$, which is thus in the image of \hat{h} . \square

Corollary 3.2.15. *Let \mathbf{G} be an n -category, $v, a, b \in \mathbf{G}_0$ objects. Each isomorphism $t \in \mathbf{G}_1$ from a to b determines an equivalence $e_t : \underline{\mathbf{Hom}}_{\mathbf{G}}^L(v, a) \rightarrow \underline{\mathbf{Hom}}_{\mathbf{G}}^L(v, b)$. Each isomorphism $m : t \rightarrow s$ determines an isomorphism $e_m : e_t \rightarrow e_s$.*

We describe the equivalence explicitly in the 2-dimensional case, which will be most relevant for us:

Example 3.2.16. Let \mathbf{G} be a 2-groupoid, $v, a, b \in \mathbf{G}_0$ and $f \in \mathbf{G}_1$ with $f_0 = a$ and $f_1 = b$. Since \mathbf{G} is a groupoid, there is $h \in \mathbf{G}_2$ (not necessarily unique) with $h_{\hat{0}} = f$ and $h_{\hat{1}} = b$. We denote $f^{-1} = h_{\hat{2}}$. By the 2-groupoid axioms, h has a uniquely determined inverse h^{-1} . Let $\gamma : D^1 \rightarrow \mathbf{G}$ be the unique map with $\gamma(101) = h$, so that $\gamma(01) = g$ and $\gamma(10) = g^{-1}$. Then $\mathbf{H} = \underline{\mathbf{Hom}}_{\mathbf{G}}^L(v, \gamma)$ can be described as follows:

- (1) $\mathbf{H}_0 = \underline{\mathbf{Hom}}_{\mathbf{G}}^L(v, a)_0 \amalg \underline{\mathbf{Hom}}_{\mathbf{G}}^L(v, b)_0$ (this is just the union if $a \neq b$, but if $a = b$ we take disjoint copies).
- (2) Let $\mathbf{X} = \{g \in \mathbf{G}_2 | g_0 = v, g_{\hat{0}} = f\}$, and let $\mathbf{X}^{-1} = \{g \in \mathbf{G}_2 | g_0 = v, g_{\hat{0}} = f^{-1}\}$ (again taking disjoint copies if $f = f^{-1}$). Then

$$\mathbf{H}_1 = \underline{\mathbf{Hom}}_{\mathbf{G}}^L(v, a)_1 \amalg \underline{\mathbf{Hom}}_{\mathbf{G}}^L(v, b)_1 \cup \mathbf{X} \cup \mathbf{X}^{-1}$$

with $d_0^{\mathbf{X}} = d_1^{\mathbf{G}}$, $d_1^{\mathbf{X}} = d_2^{\mathbf{G}}$ and vice versa for \mathbf{X}^{-1} (and the structure coming from $\underline{\mathbf{Hom}}^L$ on the other parts).

- (3) Composition is defined again as in $\underline{\mathbf{Hom}}^L$ on the corresponding parts. The composition of $h \circ g$ for $g \in \underline{\mathbf{Hom}}_{\mathbf{G}}^L(v, a)$ and $h \in \mathbf{X}$ is the composition in \mathbf{G} of the three elements $g, h, i \in \mathbf{G}_2$, where

i is the identity morphism of the object f of $\underline{\mathbf{Hom}}_{\mathbf{G}}^R(a, b)$. Similarly for the compositions $g' \circ h$, $h' \circ g'$, $g \circ h'$, $h \circ h'$ and $h' \circ h$, for $g' \in \underline{\mathbf{Hom}}_{\mathbf{G}}^L(v, b)$ and $h' \in \mathbf{X}^{-1}$ (in each case, the two elements of \mathbf{G}_2 along with i form three faces of a 2-horn, with vertices a, a, b, v or a, b, b, v , and the result is the uniquely determined fourth face)

It is clear, by construction, that the inclusions of $\underline{\mathbf{Hom}}^L(v, a)$ and $\underline{\mathbf{Hom}}^L(v, b)$ into \mathbf{H} determine fully faithful functors. As in the general proof, they are also essentially surjective by the Kan property. \square

Corollary 3.2.17. *Let \mathbf{G} be a 2-groupoid, and let $\gamma : D^2 \rightarrow \mathbf{G}$ be a fixed map, and $a \in \mathbf{G}_0$ a fixed vertex. Then $\underline{\mathbf{Hom}}^L(a, \gamma) = \underline{\mathbf{Hom}}^L(a, \gamma \circ \widehat{2}) \otimes_{\underline{\mathbf{Hom}}^L(a, \gamma \circ \widehat{1})} \underline{\mathbf{Hom}}^L(a, \gamma \circ \widehat{0})$ (canonical isomorphism), and $\underline{\mathbf{Hom}}^L(a, \gamma \circ \widehat{1}) = \underline{\mathbf{Hom}}^L(a, \gamma \circ \widehat{0}) \circ_{\underline{\mathbf{Hom}}^L(a, \gamma \circ \widehat{1})} \underline{\mathbf{Hom}}^L(a, \gamma \circ \widehat{2})$*

In other words, the composition of two morphism groupoids (in the sense of §2.2.11) is given by composition in the homotopy category.

Proof. By definition, both sides have the same sets of objects. Prop. 2.2.6 provides the required map, and since on both sides we also have a weak equivalence (by the second part of Prop. 2.2.6 and by Prop. 3.2.14), this map is an isomorphism. The second part again follows directly from the definition, as both sides are the restriction to the same set of objects. \square

3.3. The theory associated with a groupoid. We continue to fix $n \in \mathbb{N}$. To each definable n -groupoid in the theory \mathcal{T} we define an associated expansion $\mathcal{T}_{\mathbf{G}}$ of \mathcal{T} , directly generalizing (a variant of) the one-dimensional case.

Definition 3.3.1. Let \mathbf{G} be a definable n -groupoid in a theory \mathcal{T} . The expansion $\mathcal{T}_{\mathbf{G}}$ of \mathcal{T} is obtained by adding additional sorts \mathbf{G}_i^* for $0 \leq i \leq n+1$, function symbols $e_i : \mathbf{G}_i \rightarrow \mathbf{G}_i^*$, and a constant symbol $*$ in \mathbf{G}_0^* , and the axioms expressing:

- (1) \mathbf{G}^* is an n -groupoid, and e_* is a map of simplicial sets (i.e., commutes with the structure maps). We identify \mathbf{G} with its image.
- (2) $\mathbf{G}_0^* = \mathbf{G}_0 \cup \{*\}$.
- (3) The inclusion of \mathbf{G} in \mathbf{G}^* is a weak homotopy equivalence (Remark 3.2.8), and an isomorphism onto the sub-groupoid of \mathbf{G}^* spanned by \mathbf{G}_0 .

For each natural number r , there is a definable family $\Gamma_r = \underline{\mathbf{Hom}}^L(*, f)$ of groupoids, parametrised by the definable set of maps $f : D^r \rightarrow \mathbf{G}^*$. This is our collection Γ of admissible groupoids, in the sense of Def 3.1.3.

We note that our choice of Γ does satisfy the assumption on composition, by Cor. 3.2.17.

As in the one-dimensional case, each object $a \in \mathbf{G}_0$ determines an interpretation ω_a , over \mathcal{T}_a , determined by the requirement: $\omega_a(*) = a$, $\omega_a(\mathbf{G}_i^*) =$

\mathbf{G}_i for $i = 1, 2$ and similarly for the face maps. We would like to show that the ω_a are objects in the 2-groupoid associated with $\mathcal{T}_{\mathbf{G}}$ over \mathcal{T} , namely:

Proposition 3.3.2. *For every object $a \in \mathbf{G}_0$, the interpretation $\omega_a : \mathcal{T}_{\mathbf{G}} \rightarrow \mathcal{T}_a$ is 2-stable. In particular, $\langle \mathcal{T}_{\mathbf{G}}, \Gamma \rangle$ is a 2-internal cover of \mathcal{T} .*

Proof. We need to show that over some parameter u , each admissible groupoid \mathbf{H} is equivalent to $\omega_a(\mathbf{H})$, over some parameters from \mathcal{T} . Let u be any element of $\underline{\mathbf{Hom}}_{\mathbf{G}^*}^L(*, a)_0$ (it is consistent that such a u exists: for any model M of \mathcal{T} such that $a \in \mathbf{G}_0(M)$, $M \circ \omega_a$ is a model of $\mathcal{T}_{\mathbf{G}}$ for which this set is non-empty).

Let $f : D^r \rightarrow \mathbf{G}^*$ be a map, and assume first that for some $i \in [r]$, $b = f(i) \in \mathbf{G}_0$. By the second item of Prop 3.2.14, $\mathbf{H} = \underline{\mathbf{Hom}}_{\mathbf{G}^*}^L(*, f)$ is equivalent (over no additional parameters) to $\underline{\mathbf{Hom}}_{\mathbf{G}^*}^L(*, b)$, so we may assume that $f = b$. We may also assume that b is in the same connected component as a , because otherwise \mathbf{H} is empty. According to Cor 3.2.15, it follows that \mathbf{H} is equivalent $\underline{\mathbf{Hom}}_{\mathbf{G}^*}^L(*, a)$. Again according to (a dual version of) Cor 3.2.15, the fixed element u determines an equivalence from \mathbf{H} to $\omega_a(\mathbf{H}) = \underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, a)$.

The remaining case is when f is the constant map $*$, so that $\mathbf{H} = \underline{\mathbf{Hom}}_{\mathbf{G}^*}^L(*, *)$, and $\omega_a(\mathbf{H}) = \underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, a)$. The same argument as above shows that both are equivalent to $\underline{\mathbf{Hom}}_{\mathbf{G}^*}^L(*, a)$ over u . \square

We would like to prove that the association $a \mapsto \omega_a$ is the object part of an assignment that recovers (up to equivalence) \mathbf{G} . To do that, we need to define the 2-groupoid which is the target of this assignment. This will be the analog of $I_{\mathcal{T}^*/\mathcal{T}}$ from the one-dimensional case (§2.2.1).

Definition 3.3.3. Let \mathcal{T}^* be a stable expansion of a theory \mathcal{T} , with admissible family of distinguished internal covers Γ . The 2-groupoid associated to this datum is defined as follows:

- (1) Objects are 2-stable interpretations of $\langle \mathcal{T}^*, \Gamma \rangle$ in \mathcal{T} , over \mathcal{T} .
- (2) If x, y are two objects as above, a morphism $u : x \rightarrow y$ is given by a bi-interpretation $u_{\mathcal{T}'} : x(\mathcal{T}') \rightarrow y(\mathcal{T}')$ over \mathcal{T} , for each admissible internal cover $\mathcal{T}' \in \Gamma$. These bi-interpretations are given with isomorphisms $c_i : u_{\mathcal{T}_2} \circ x(i) \rightarrow y(i) \circ u_{\mathcal{T}_1}$ for every admissible interpretation $i : \mathcal{T}_1 \rightarrow \mathcal{T}_2$ between admissible covers $\mathcal{T}_1, \mathcal{T}_2$ in Γ (uniformly in families).

The isomorphisms are required to satisfy: $c_{j \circ i} = y(j)(c_i) \circ c_j(x(i))$ for admissible interpretations $i : \mathcal{T}_1 \rightarrow \mathcal{T}_2$, $j : \mathcal{T}_2 \rightarrow \mathcal{T}_3$ as above (these make sense since, by definition, the c_i are definable maps in $y(\mathcal{T}_2)$, $y(j)$ is an interpretation of $y(\mathcal{T}_2)$ in $y(\mathcal{T}_3)$ and c_j is a map between interpretations of $x(\mathcal{T}_2)$, so can be applied to definable sets of the form $x(i)$.)

- (3) The 2-morphisms with edges $u : x \rightarrow y$, $v : y \rightarrow z$ and $w : x \rightarrow z$ are given by isomorphisms $v \circ u \rightarrow w$, all over \mathcal{T} .

- (4) The “2-composition” of the 2-morphisms $\alpha : v \circ u \rightarrow w$, $\beta : s \circ v \rightarrow r$ and $\gamma : r \circ u \rightarrow t$ is given by

$$s \circ w \xrightarrow{s \cdot \alpha^{-1}} s \circ (v \circ u) = (s \circ v) \circ u \xrightarrow{\beta \cdot u} r \circ u \xrightarrow{\gamma} t,$$

where \cdot stands for pointwise application (or *horizontal composition*) as above.

Applying the definition with \mathcal{T} replaced by \mathcal{T}_A , for a \mathcal{T} -structure A , we obtain a 2-groupoid for each such structure A , which we denote $I^2(A) = I^2_{\mathcal{T}^*/\mathcal{T}}(A)$.

Using the equivalence between internal covers and definable groupoids, this can be described in terms of definable groupoids. We give an explicit description in 3.3.6 below.

Remark 3.3.4. Let \mathcal{C} be the category of definable groupoids in \mathcal{T} , with weak equivalences as morphisms. We may form its bi-category of *cospan*s for this category, as in [9, 0084]. By Cor 2.2.13, it is equivalent (as a bi-category) to the category of internal covers and bi-interpretations. The 2-category in Def 3.3.3 can be viewed as the *Duskin nerve* ([9, 009T]) of this bi-category (this is clear from the description in [9, 00A1]). In particular, it follows that this is indeed a 2-category. \square

Proposition 3.3.5. *Let \mathbf{G} be a definable 2-groupoid in a theory \mathcal{T} , and let $\mathcal{T}_{\mathbf{G}} = \langle \mathcal{T}_{\mathbf{G}}, \Gamma \rangle$ be the corresponding 2-internal cover. The association $a \mapsto \omega_a$ extends to a map $\omega : \mathbf{G}(A) \rightarrow I^2_{\mathcal{T}_{\mathbf{G}}/\mathcal{T}}(A)$ of 2-groupoids, compatible with extensions of the structure A .*

Proof. Prop. 3.3.2 shows that for all $a \in \mathbf{G}_0(A)$, ω_a is indeed an object of $I^2(A)$. Given $t : a \rightarrow b$ in $\mathbf{G}_1(A)$, we define $\omega_t : \omega_a \rightarrow \omega_b$ as follows: Let \mathbf{H} be an admissible groupoid. As in the proof of 3.3.2, we may assume that $\mathbf{H} = \mathbf{H}_c = \underline{\mathbf{Hom}}_{\mathbf{G}^*}^L(*, c)$ for some $c \in \mathbf{G}_0(A)$, so that $\omega_a(\mathbf{H}) = \underline{\mathbf{Hom}}_{\mathbf{G}}^L(a, c)$ and $\omega_b(\mathbf{H}) = \underline{\mathbf{Hom}}_{\mathbf{G}}^L(b, c)$. By Cor 3.2.15, t induces an (admissible) equivalence from $\omega_a(\mathbf{H})$ to $\omega_b(\mathbf{H})$, which we take to be $\omega_t(\mathbf{H})$. Our definition (and construction) ensures the compatibility under admissible maps $\mathbf{H} \rightarrow \mathbf{H}'$.

Similarly, let $\alpha \in \mathbf{G}_2(A)$, with edges $r : a \rightarrow b$, $s : b \rightarrow c$ and $t : a \rightarrow c$. We need to construct an isomorphism (over \mathcal{T}) from $\omega_s \circ \omega_r$ to ω_t . Consider the map $f : D^2 \rightarrow \mathbf{G}$ determined by α . We have $f(01) = r$, $f(12) = s$ and $f(02) = t$, so that for each object $d \in \mathbf{G}_0(A)$, the equivalence $\omega_r(\mathbf{H}_d) : \omega_a(\mathbf{H}_d) \rightarrow \omega_b(\mathbf{H}_d)$ is given by $\underline{\mathbf{Hom}}^L(f \circ h_{01}, d)$, where $h_{01} : [1] \rightarrow [2]$ is the inclusion (and similarly for s, t). Hence, $\underline{\mathbf{Hom}}^L(f, d)$ represents the composition $\omega_s \circ \omega_r$, and restriction to ω_t provides the required map.

This completes the construction of ω . The proof that this is a map of 2-groupoids (i.e., that it commutes with composition) is similar to the above, using D^4 in place of D^3 , and the fact that it commutes with extension of scalars is obvious. \square

3.3.6. Our main goal is to prove that the map ω constructed in Prop 3.3.5 is a weak equivalence. Similarly to the 1-dimensional case, it will generally only be true in a model. As a preparation, we consider more explicitly the structure of I^2 from Def. 3.3.3, from a definable groupoid point of view.

Let ω_1 and ω_2 be two objects of $I_{\mathcal{T}^*/\mathcal{T}}^2$, i.e., 2-stable interpretations of \mathcal{T}^* in \mathcal{T} . An isomorphism over \mathcal{T} is given, according to Prop 2.2.10, by a family $\mathbf{K}(\mathbf{H})$ of groupoids in \mathcal{T} , for each admissible groupoid \mathbf{H} in \mathcal{T}^* , along with weak equivalences $u_i(\mathbf{H}) : \omega_i(\mathbf{H}) \xrightarrow{\sim} \mathbf{K}(\mathbf{H})$, all definable uniformly in \mathbf{H} . Given another admissible groupoid \mathbf{H}' , an admissible interpretation from $\mathcal{T}^*_{\mathbf{H}'}$ to $\mathcal{T}^*_{\mathbf{H}}$ is given, again by Prop 2.2.10, by an admissible groupoid \mathbf{X} and admissible maps $f : \mathbf{H} \rightarrow \mathbf{X}$ and $g : \mathbf{H}' \xrightarrow{\sim} \mathbf{X}$. According to the definition, we are provided with a definable isomorphism (realising the isomorphisms c_i , via Cor. 2.2.13)

$$t_{\mathbf{X},\mathbf{K}} : \mathbf{K}(\mathbf{H}) \circ_{\omega_1(\mathbf{H})} \omega_1(\mathbf{X}) \xrightarrow{\sim} \omega_2(\mathbf{X}) \circ_{\omega_2(\mathbf{H}')} \mathbf{K}(\mathbf{H}') \quad (6)$$

uniformly definable in \mathbf{X}, \mathbf{K} (and the associated embeddings), and restricting to the identity on $\omega_1(\mathbf{H}')$ and on $\omega_2(\mathbf{H})$. If \mathbf{Y} determines a map to $\mathcal{T}^*_{\mathbf{H}'}$ from $\mathcal{T}^*_{\mathbf{H}''}$ for a further groupoid \mathbf{H}'' , we have the maps

$$t_{\mathbf{X},\mathbf{K}} \otimes_{\omega_1(\mathbf{H}')} \mathbf{1}_{\omega_1(\mathbf{Y})} : \mathbf{K}(\mathbf{H}) \circ_{\omega_1(\mathbf{H})} \omega_1(\mathbf{X}) \circ_{\omega_1(\mathbf{H}')} \omega_1(\mathbf{Y}) \xrightarrow{\sim} \omega_2(\mathbf{X}) \circ_{\omega_2(\mathbf{H}')} \mathbf{K}(\mathbf{H}') \circ_{\omega_1(\mathbf{H}')} \omega_1(\mathbf{Y}) \quad (7)$$

and

$$\mathbf{1}_{\omega_2(\mathbf{X})} \otimes_{\omega_2(\mathbf{H}')} t_{\mathbf{Y},\mathbf{K}} : \omega_2(\mathbf{X}) \circ_{\omega_2(\mathbf{H}')} \mathbf{K}(\mathbf{H}') \circ_{\omega_1(\mathbf{H}')} \omega_1(\mathbf{Y}) \xrightarrow{\sim} \omega_2(\mathbf{X}) \circ_{\omega_2(\mathbf{H}')} \omega_2(\mathbf{Y}) \circ_{\omega_2(\mathbf{H}'')} \mathbf{K}(\mathbf{H}'') \quad (8)$$

the groupoid $\mathbf{X} \circ_{\mathbf{H}'} \mathbf{Y}$ represents the composition of interpretations, and $\omega_i(\mathbf{X} \circ_{\mathbf{H}'} \mathbf{Y}) = \omega_i(\mathbf{X}) \circ_{\omega_i(\mathbf{H}')} \omega_i(\mathbf{Y})$ (canonical identification), since ω_i is an interpretation. Under this identification, we require that

$$(\mathbf{1}_{\omega_2(\mathbf{X})} \otimes_{\omega_2(\mathbf{H}')} t_{\mathbf{Y},\mathbf{K}}) \circ (t_{\mathbf{X},\mathbf{K}} \otimes_{\omega_1(\mathbf{H}')} \mathbf{1}_{\omega_1(\mathbf{Y})}) = t_{(\mathbf{X} \circ_{\mathbf{H}'} \mathbf{Y}),\mathbf{K}}. \quad (9)$$

Finally, a 2-morphism is determined by a natural isomorphism between two maps as above (one a composition, which we already understand), so it is enough to describe those. Let ω_1, ω_2 and \mathbf{K} be as above, and let \mathbf{L} represent another morphism. A natural isomorphism is then given by a uniform family of isomorphisms $\alpha_{\mathbf{H}} : \mathbf{K}(\mathbf{H}) \rightarrow \mathbf{L}(\mathbf{H})$ over $\omega_i(\mathbf{H})$, which intertwine the maps $t_{\mathbf{X},\mathbf{K}}$ and $t_{\mathbf{X},\mathbf{L}}$ whenever \mathbf{X} represents an interpretation. The 2-composition of three such suitable maps is described as in Def 3.3.3, with composition replaced by pushouts as appropriate.

Remark 3.3.7. By definition, internality means that there is a non-empty definable *set* (i.e., a 0-groupoid) of isomorphisms between the internal sorts and sorts of the base theory. Similarly, the structure described above includes the description of a \mathcal{T}^* -definable 1-groupoid $\mathbf{Iso}_{\mathcal{T}}(\tilde{\mathcal{T}}^8, \tilde{\mathcal{T}})$ of weak equivalences between admissible covers $\tilde{\mathcal{T}}^*$ of \mathcal{T}^* and covers $\tilde{\mathcal{T}}$ of \mathcal{T} (non-empty for some $\tilde{\mathcal{T}}$ if \mathcal{T}^* is 2-internal). In terms of groupoids, the families \mathbf{K}

as in §3.3.6 are the objects, and the morphisms are the natural isomorphisms α . Furthermore, this groupoid itself is admissible. \square

Example 3.3.8. Let $\mathcal{T}^* = \mathcal{T}_{\mathbf{G}}$ as in Prop 3.3.5, and let $\omega_1 = \omega_a$ and $\omega_2 = \omega_b$ for some $a, b \in \mathbf{G}_0(A)$. Let $f : a \rightarrow b$ be a morphism in $\mathbf{G}(A)$ (identified with the corresponding map from D^1). Given an admissible groupoid $\mathbf{H} = \mathbf{H}_d = \underline{\mathbf{Hom}}^L(d, *)$, we let $\mathbf{K}_f(d) = \mathbf{K}_f(\mathbf{H}_d) = \underline{\mathbf{Hom}}_{\mathbf{G}}^L(d, f)$, with the canonical maps from $\omega_a(\mathbf{H}_d) = \underline{\mathbf{Hom}}^L(d, a)$ and $\omega_b(\mathbf{H}_d) = \underline{\mathbf{Hom}}^L(d, b)$ (these are weak equivalences by Prop 3.2.14).

To give \mathbf{K} the structure of an isomorphism from ω_a to ω_b , we need to supply the isomorphisms (6). If $\mathbf{H}' = \mathbf{H}_c = \underline{\mathbf{Hom}}^L(c, *)$ is another admissible groupoid in $\mathcal{T}_{\mathbf{G}}$, an admissible isomorphism from \mathbf{H}_c to \mathbf{H}_d is given by a groupoid $\mathbf{X}_g = \underline{\mathbf{Hom}}^L(g, *)$, with $g : c \rightarrow d$ in \mathbf{G} . Seeing as $\omega_x(\mathbf{X}_g) = \underline{\mathbf{Hom}}^L(g, x)$ for all x , such a structure consists of a definable family of maps

$$t_{g,f} : \underline{\mathbf{Hom}}^L(d, f) \circ_{\underline{\mathbf{Hom}}^L(d,a)} \underline{\mathbf{Hom}}^L(g, a) \xrightarrow{\sim} \underline{\mathbf{Hom}}^L(g, b) \circ_{\underline{\mathbf{Hom}}^L(c,b)} \underline{\mathbf{Hom}}^L(c, f)$$

Recall that t is the identity on objects, so we only need to define it on morphisms. Let $\langle u, v \rangle$ represent a morphism of $\underline{\mathbf{Hom}}^L(d, f) \circ_{\underline{\mathbf{Hom}}^L(d,a)} \underline{\mathbf{Hom}}^L(g, a)$. Let $h : c \rightarrow a$ be the domain of v . By the Kan property, there is a morphism $w \in \underline{\mathbf{Hom}}^L(c, f)$ whose domain is h . Then u, v, w form an element of $\Lambda_3^3(\mathbf{G})$, so composition provides a fourth face $y \in \underline{\mathbf{Hom}}^L(g, b)_1$. We let $t_{g,f}(u \otimes v) = y \otimes w$. If w' is a different choice in place of w , then $w' \circ w^{-1}$ is in $\underline{\mathbf{Hom}}^L(c, b)$, so the result represents the same morphism of $\underline{\mathbf{Hom}}^L(g, b) \circ_{\underline{\mathbf{Hom}}^L(c,b)} \underline{\mathbf{Hom}}^L(c, f)$. It is clear that t is well defined on the class $u \otimes v$, and uniformly definable in g, f .

To prove the identity (9), assume we are given another morphism $g' : c' \rightarrow c$, corresponding to an admissible interpretation represented by $\mathbf{Y} = \underline{\mathbf{Hom}}^L(g', *)$. Let $v' \in \underline{\mathbf{Hom}}^L(g', a) = \omega_1(\mathbf{Y})$. Proceeding with the notation above, we need to determine the image of $y \otimes w \otimes v'$ in $\underline{\mathbf{Hom}}^L(g, b) \circ_{\underline{\mathbf{Hom}}^L(c,b)} \underline{\mathbf{Hom}}^L(g', b) \circ_{\underline{\mathbf{Hom}}^L(c',b)} \underline{\mathbf{Hom}}^L(c', f)$. As above, it is given by $y \otimes y' \otimes w'$, where $y' \in \underline{\mathbf{Hom}}^L(g', b) = \omega_2(\mathbf{Y})$ and $w' \in \underline{\mathbf{Hom}}^L(c', f) = \mathbf{K}_f(\mathbf{H}_{c'})$ represent the other two faces of a partial simplex with faces w and v' (this other simplex can be visualised as attached to the previous one at the face w). On the other hand, $\mathbf{X} \circ_{\mathbf{H}_c} \mathbf{Y}$ was identified (as in Cor. 3.2.17) with $\underline{\mathbf{Hom}}^L(g \circ g', *)$, for any composition $g \circ g'$. After choosing such a composition h , $v \otimes v'$ is identified with an element of $\underline{\mathbf{Hom}}^L(h, a)$ and $y \otimes y'$ with an element of $\underline{\mathbf{Hom}}^L(h, b)$, so that they become two faces of the simplex with vertices a, b, c, d , the other two being u and w' , so that $u \otimes w' = t_{h,f}(v \otimes v', y \otimes y')$, as required.

Assume now that we are given a map $\gamma : D^2 \rightarrow \mathbf{G}$ corresponding to an element $w \in \mathbf{G}_2$, with edges $f = \gamma(01)$, $g = \gamma(12)$ and $h = \gamma(02)$. Given

element $u \in \mathbf{K}_f(\mathbf{H}_c) = \underline{\mathbf{Hom}}^L(c, f)$ and $v \in \mathbf{K}_g(\mathbf{H}_c) = \underline{\mathbf{Hom}}^L(c, g)$ (for an arbitrary $c \in \mathbf{G}_0$), the 2-composition applied to u, v and w provides an element of $\mathbf{K}_h(\mathbf{H}_c) = \underline{\mathbf{Hom}}^L(c, h)$. This process assembles into a family of isomorphisms $\alpha_\gamma : \mathbf{K}_g(\mathbf{H}_c) \circ \mathbf{K}_f(\mathbf{H}_c) \rightarrow \mathbf{K}_h(\mathbf{H}_c)$, definable uniformly in γ and c . This completes the description (and a reformulation of the proof) of the map in Prop. 3.3.5 in terms of definable groupoids. \square

We are now ready to prove our main result.

Theorem 3.3.9. *Let \mathbf{G} be a 2-groupoid defined in a theory \mathcal{T} , and let $\mathcal{T}_{\mathbf{G}}$ be the associated theory (and admissible covers), as in Def. 3.3.1. Then $\mathcal{T}_{\mathbf{G}}$ is a 2-internal cover of \mathcal{T} , and for every model M of \mathcal{T} , the 2-groupoid of M -points $I_{\mathcal{T}_{\mathbf{G}}/\mathcal{T}}^2(M)$ is weakly equivalent to $\mathbf{G}(M)$.*

Proof. The fact that $\mathcal{T}_{\mathbf{G}}$ is a 2-internal cover is Prop. 3.3.2. The map from \mathbf{G} to $I_{\mathcal{T}_{\mathbf{G}}/\mathcal{T}}^2$ was constructed in Prop. 3.3.5, and described in terms of groupoids in Ex. 3.3.8. We will use this description to show that the map is a weak equivalence, using Remark 3.2.8. We assume that we are working over M , and proceed by considering the possible dimensions $0 \leq k \leq 2$.

$k = 0$: We need to show that any 2-stable interpretation ω of $\mathcal{T}_{\mathbf{G}}$ in \mathcal{T} admits a coherent collection of bi-interpretations as in Def. 3.3.3.2 to some ω_a .

Since ω is an interpretation over \mathcal{T} , $\omega(\mathbf{G}^*)$ is a definable 2-groupoid in \mathcal{T} , containing \mathbf{G} , with the inclusion a weak equivalence. The proof now proceeds exactly as the proof of Prop. 3.3.2, with $\omega(\mathbf{G}^*)$ in place of \mathbf{G}^* .

$k = 1$: This is the main case, which can be viewed as a definable version of the Yoneda lemma. Let $a, b \in \mathbf{G}_0$, and assume we are given an equivalence from ω_a to ω_b . Hence, for every $c \in \mathbf{G}_0$ (some parameters), we are given a groupoid $\mathbf{K}(c)$ in \mathcal{T} and weak equivalences $\underline{\mathbf{Hom}}^L(c, a) = \omega_a(\mathbf{H}_c) \rightarrow \mathbf{K}(c)$ and $\underline{\mathbf{Hom}}^L(c, b) = \omega_b(\mathbf{H}_c) \rightarrow \mathbf{K}(c)$, uniformly in c , along with structure maps (6)

$$t_{g, \mathbf{K}} : \mathbf{K}(d) \circ_{\underline{\mathbf{Hom}}^L(d, a)} \underline{\mathbf{Hom}}^L(g, a) \rightarrow \underline{\mathbf{Hom}}^L(g, b) \circ_{\underline{\mathbf{Hom}}^L(c, b)} \mathbf{K}(c)$$

(all notation as in Ex. 3.3.8, except \mathbf{K} is no longer known to be of the given form). We identify $\omega_a(\mathbf{H}_c), \omega_b(\mathbf{H}_c)$ with their images in $\mathbf{K}(c)$.

In particular, we have the identity morphism $\mathbf{1}_a$ of a as an object $\mathbf{1}_a \in \omega_a(\mathbf{H}_a)$, and by weak equivalence, an object $f : a \rightarrow b$ in $\omega_b(\mathbf{H}_a) \subseteq \mathbf{K}(a)$, along with a morphism $u : \mathbf{1}_a \rightarrow f$ in $\mathbf{K}(a)$. We will show that \mathbf{K} is isomorphic to \mathbf{K}_f , by a unique isomorphism.

To do that, let $c \in \mathbf{G}_0$ be an arbitrary object, and let v be a morphism of $\mathbf{K}_f(c) = \underline{\mathbf{Hom}}^L(c, f)$ (so a 2-morphism of \mathbf{G}). Denote by $g \in \underline{\mathbf{Hom}}^L(c, a)_0$ the domain of v . Then v can also be viewed as a morphism in $\underline{\mathbf{Hom}}^L(g, b)$, and on the other hand, we have the canonical morphism w from g to $\mathbf{1}_a$ in $\underline{\mathbf{Hom}}^L(g, a)$. Applying $t_{g, \mathbf{K}}$

to the morphism $u \otimes w \in \mathbf{K}(a) \circ_{\underline{\mathbf{Hom}}^L(a,a)} \underline{\mathbf{Hom}}^L(g, a)$, we may write $t_{g,\mathbf{K}}(u \otimes w)$ as $v \otimes x$ for a unique $x \in \mathbf{K}(c)$, which we take to be the image of v . By construction this map commutes with the structure maps t , and is unique with this property.

$k = 2$: We need to show that each isomorphism $\alpha : \mathbf{K}_g \circ \mathbf{K}_f \rightarrow \mathbf{K}_h$ with $f : a \rightarrow b$, $g : b \rightarrow c$ and $h : a \rightarrow c$ arises from a unique $\gamma : D^2 \rightarrow \mathbf{G}$, with boundary f, g, h (as in the end of Ex. 3.3.8). Uniqueness was already shown in the part $k = 1$. For existence, we apply $\alpha_b : \mathbf{K}_g(b) \circ \mathbf{K}_f(b) \rightarrow \mathbf{K}_h(b)$ to the element $\mathbf{1}_g \otimes \mathbf{1}_f$ (where $\mathbf{1}_g$ is the identity morphism of the object g of $\underline{\mathbf{Hom}}^L(b, c)$, viewed as an element of \mathbf{G}_2 , and similarly for f), to obtain an element $\gamma_b \in \mathbf{K}_h(b)_1$, again viewed as a 2-morphism of \mathbf{G} . It is clear that the map α coincides with α_γ on the given maps, and then, again by the uniqueness statement, that $\alpha = \alpha_\gamma$ globally. \square

3.3.10. *Recovering an definable 2-groupoid.* The main statement of classical internality starts with the assumption of internality, and produces a definable (non-empty) groupoid from it. The general outline of this construction was recalled in §2.2, and in Cor. 2.2.4 we indicated how this construction is useful in the description of definable sets in the cover.

In our approach, the construction of the (2-) groupoid is almost tautological: we defined a groupoid (or a 2-groupoid) associated to every stable expansion, and by definition, it is an internal cover if the groupoid is non-empty. However, we still need to show that the groupoid is equivalent to a definable one, which we sketch below. The other part, describing the (admissible) 1-groupoids in the cover in terms of suitable definable fibrations in the base, is more involved, and we postpone most of the work here to future work.

To recover definability, assume \mathcal{T}^* is a 2-internal cover of \mathcal{T} , and let $\omega : \mathcal{T}^* \rightarrow \mathcal{T}_A$ be a 2-stable interpretation. For simplicity we assume that Γ , the collection of admissible covers, consists of one definable family. As in §3.3.6, we have a fixed parameter u_0 and a uniform family $\mathbf{K} = \mathbf{K}_c$ of groupoids in \mathcal{T}^* defined over u_0 , along with (uniformly definable) weak equivalences $f_c : \mathbf{X}_c \xrightarrow{\sim} \mathbf{K}_c$ and $g_c : \omega(\mathbf{X}_c) \rightarrow \mathbf{K}_c$ for \mathbf{X}_c members of Γ (note that c ranges over a definable set in \mathcal{T} by assumption). Like in Remark 3.3.7, as u_0 varies, we obtain a family $\mathbf{K}_{u,c}$ of objects of a definable 1-groupoid \mathbf{P}_c , along with a map $\mathbf{P}_c \rightarrow \underline{\mathbf{Iso}}(\mathbf{X}, \omega(\mathbf{X}))$ for each member \mathbf{X} of Γ . Furthermore, \mathbf{P}_c itself is also in Γ . Applying the above map to $\mathbf{X} = \mathbf{P}_{c'}$, we obtain a family of definable maps of 1-groupoids $a : \mathbf{P}_c \rightarrow \underline{\mathbf{Iso}}(\mathbf{P}_{c'}, \omega_c(\mathbf{P}_{c'}))$.

The 2-groupoid \mathbf{G} is constructed as follows: \mathbf{G}_0 is the definable set of parameters c as above. Each groupoid \mathbf{P}_c will be isomorphic to $\underline{\mathbf{Hom}}^L(*, c)$ in the corresponding \mathbf{G}^* . Let c, d be two elements of \mathbf{G}_0 . Given an object u of \mathbf{P}_c , the map a above produces a groupoid \mathbf{K}_u as an object of $\underline{\mathbf{Iso}}(\mathbf{P}_d, \omega_c(\mathbf{P}_d))$, along with weak equivalences $f : \mathbf{P}_d \xrightarrow{\sim} \mathbf{K}_u$ and $g : \omega_c(\mathbf{P}_d) \rightarrow \mathbf{K}_u$. Let $\mathbf{Q}_{u,v}$ be the set of morphisms in \mathbf{K}_u with domain $f(v)$.

We set the morphisms from c to d to be the definable types space of $\mathbf{Q}_{u,v}$ over \mathcal{T} (this is definable in \mathcal{T} by stability of the embedding). Note that each such type includes, in particular, the information of the object of $\omega_c(\mathbf{P}_d)$ which is the codomain of any realisation (as an element of $\mathbf{Q}_{u,v}$).

Similarly, assume e is another element of \mathbf{G}_0 , and w an object of \mathbf{P}_e . The elements of \mathbf{G}_2 with vertices c, d, e are defined as the types over \mathcal{T} of triples $\mathbf{Q}_{u,v} \times \mathbf{Q}_{v,w} \times \mathbf{Q}_{u,w}$ (over all such u, v, w). The 2-composition is defined similarly, by considering 4-tuples. We skip the details of the construction, as well as the proof that the map determined by a is a weak equivalence. We mention also that this description can, in principle, be used to give an equivalent combinatorial definition of 2-internality (similar to the original definition of internality), but I could not find one sufficiently pleasant to write.

3.4. Questions. I mention a few natural questions that I hope to address in the future.

3.4.1. Structure of admissible internal covers. The definable version of the 2-groupoid \mathbf{G} associated to a 2-internal cover \mathcal{T}^* of \mathcal{T} was only sketched above. Assuming it is properly described, it still needs to be seen that $\mathcal{T}_{\mathbf{G}}$ and \mathcal{T}^* are, in some sense, equivalent. It will also be useful to describe the admissible covers (in either) as suitably defined “higher local systems” on \mathbf{G} . Both questions require that we name the precise closure properties on the collection of admissible covers: we already assumed that they are closed under finite inverse limit and definable mapping spaces, but it is not clear, for example, if some closure under quantifiers is required.

3.4.2. Lax interpretations. We had not run into the questions above because we required to objects of the 2-groupoid I^2 to be actual interpretations. This works well in the example of $\mathcal{T}_{\mathbf{G}}$, but for general expansions it might make more sense to consider a larger class of “lax interpretations” that preserve only the admissible covers (possibly up to weak equivalence).

3.4.3. Internal covers of $\mathcal{T}_{\mathbf{G}}$. The requirement for introducing admissible covers was motivated above. However, it might still be the case in the case of $\mathcal{T}_{\mathbf{G}}$, essentially all internal covers are the ones described (up to covers that come from the base \mathcal{T}). Again, stating this precisely requires clarifying the structure of the collection of admissible covers.

3.4.4. Relation to analyzability. The 2-groupoid \mathbf{G}^* in the theory \mathcal{T} provides an example of a 2-analyzable set over \mathcal{T} . Can we describe (combinatorially) which 2-analyzable covers occur in this way?

3.4.5. Generalization to higher dimensions. This is rather clear: one continues by induction, defining an $i+1$ -groupoid associated to a stable expansion by taking into account i -internal covers, and then defining the expansion to be an $i+1$ -cover if this groupoid is non-empty. However, some of the proofs given above will be difficult to generalize, and it would be interesting to look for

a smoother way. In any case, this only applies to each finite level, and it does not seem reasonable to expect a generalization to arbitrary ∞ -groupoids.

3.4.6. *Structure at $*$* . We did not consider the structure of the groupoid $\mathbf{Hom}_{\mathbf{G}^*}^L(*, *)$ definable in $\mathcal{T}_{\mathbf{G}}$. On top of the groupoid structure, composition gives it a structure of a monoidal category up to homotopy (i.e., the homotopy category is monoidal). It also acts on all the admissible covers, so it is really a higher analog of the binding group. However, we did not consider what could be a version of the Galois correspondence or of descent, as in the 1-dimensional case.

REFERENCES

- [1] John Goodrick, Byunghan Kim, and Alexei Kolesnikov. “Type-amalgamation properties and polygroupoids in stable theories”. In: *Journal of Mathematical Logic* 15.1 (2015), pp. 1550004, 45. ISSN: 0219-0613. DOI: 10.1142/S021906131550004X (cit. on p. 2).
- [2] Levon Haykazyan and Rahim Moosa. “Functoriality and uniformity in Hrushovski’s groupoid-cover correspondence”. In: *Annals of Pure and Applied Logic* 169.8 (2018), pp. 705–730. ISSN: 0168-0072. DOI: 10.1016/j.apal.2018.03.005 (cit. on p. 2).
- [3] Ehud Hrushovski. “Computing the Galois group of a linear differential equation”. In: *Differential Galois theory*. Ed. by Teresa Crespo and Zbigniew Hajto. Banach Center Publications 58. May 28–June 1, 2001. Warsaw: Polish Academy of Sciences Institute of Mathematics, 2002, pp. 97–138 (cit. on p. 1).
- [4] Ehud Hrushovski. “Groupoids, imaginaries and internal covers”. In: *Turkish Journal of Mathematics* 36.2 (2012), pp. 173–198. ISSN: 1300-0098 (cit. on pp. 1–3, 5, 6, 9, 15).
- [5] A. Joyal. “Quasi-categories and Kan complexes”. In: *Journal of Pure and Applied Algebra* 175.1-3 (2002), pp. 207–222. ISSN: 0022-4049. DOI: 10.1016/S0022-4049(02)00135-4 (cit. on p. 12).
- [6] Moshe Kamensky. “A categorical approach to internality”. In: *Models, logics, and higher-dimensional categories. Proceedings of the meeting held at the Université de Montréal, Montréal, QC, June 18–20, 2009* (Centre de Recherches Mathématiques, Université de Montréal, June 2009). Ed. by Bradd Hart, Thomas G. Kucera, Anand Pillay, Philip J. Scott, and Robert A. G. Seely. Vol. 53. CRM Proceedings & Lecture Notes. A tribute to the work of Mihály Makkai. Providence, RI: American Mathematical Society, 2011, pp. 139–156. ISBN: 978-0-8218-7281-9. arXiv: 1012.3185 [math.LO] (cit. on p. 4).
- [7] Moshe Kamensky and Anand Pillay. “Interpretations and differential Galois extensions”. In: *International Mathematics Research Notices. IMRN* (Apr. 2016). DOI: 10.1093/imrn/rnw019. arXiv: 1409.5915 [math.AG] (cit. on pp. 3, 5).

- [8] Jacob Lurie. *Higher topos theory*. Annals of Mathematics Studies 170. Princeton University Press, Princeton, NJ, 2009. xviii+925. ISBN: 978-0-691-14049-0. DOI: 10.1515/9781400830558 (cit. on pp. 2, 3, 12, 13, 15–17).
- [9] Jacob Lurie. *Kerodon*. <https://kerodon.net>. 2018 (cit. on pp. 12, 14, 15, 21).
- [10] Saunders Mac Lane. *Categories for the working mathematician*. 2nd ed. Graduate Texts in Mathematics 5. New York: Springer-Verlag, 1998. xii+314. ISBN: 0-387-98403-8 (cit. on p. 4).
- [11] Bruno Poizat. “Une théorie de Galois imaginaire”. In: *The Journal of Symbolic Logic* 48.4 (1983), pp. 1151–1170. ISSN: 0022-4812 (cit. on p. 1).
- [12] Paul Wang. “Extending Hrushovski’s groupoid-cover correspondence using simplicial groupoids”. In: *Annals of Pure and Applied Logic* 172.7 (2021), p. 102970. ISSN: 0168-0072. DOI: 10.1016/j.apal.2021.102970. URL: <https://www.sciencedirect.com/science/article/pii/S0168007221000282> (cit. on p. 2).
- [13] Boris I. Zil’ber. “Totally categorical theories: structural properties and the nonfinite axiomatizability”. In: *Model theory of algebra and arithmetic*. Proceedings of the Conference on Applications of Logic to Algebra and Arithmetic (Karpacz, September 1–7, 1979). Ed. by L. Pacholski and J. Wierzejewski. Lecture Notes in Mathematics 834. Berlin: Springer, 1980, pp. 381–410. ISBN: 3-540-10269-8 (cit. on p. 1).

DEPARTMENT OF MATH, BEN-GURION UNIVERSITY, BE’ER-SHEVA, ISRAEL
Email address: <mailto:kamensky.bgu@gmail.com>
URL: <https://www.math.bgu.ac.il/~kamenskm>