

DISPERSIVE DECAY BOUND OF SMALL DATA SOLUTIONS TO KAWAHARA EQUATION IN A FINITE TIME SCALE

JONGWON LEE

ABSTRACT. In this article, we prove that small localized data yield solutions to Kawahara type equation which have linear dispersive decay on a finite time. We use the similar method used to derive the dispersive decay bound of the solutions to the KdV equation, with some steps being simpler. This result is expected to be the first result of the small data global bounds of the fifth-order dispersive equations with quadratic nonlinearity.

1. INTRODUCTION

A global behavior of a small data solution of a nonlinear dispersive equation has drawn attentions of many authors. It might result in the finite or infinite time linear decay bound, or the linear or modified scattering. The goal of this paper is to find a dispersive decay bound of a small data solution to the Kawahara equation:

$$(KW) \quad u_t - u_{5x} - uu_x = 0.$$

We also consider the case of the modified Kawahara equation

$$(mKW) \quad u_t - u_{5x} = cu^2u_x,$$

with $c = \pm 1$, which is the special case of the fifth-order KdV equation

$$(5KdV) \quad u_t - u_{5x} = c_1uu_{3x} + c_2u_xu_{2x} + c_3u^2u_x.$$

The equation (KW) arises from the study of capillary waves on a shallow layer and magneto-sound propagation in plasmas, see [Kaw72].

The local and global Cauchy problem of the fifth-order nonlinear dispersive equations such as (KW), (mKW), (5KdV) has been extensively studied. Kenig et al. proved the local well-posedness of the general class of dispersive equations

$$\partial_t u + \partial_x^{2j+1} u + P(u, \partial_x u, \dots, \partial_x^{2j} u) = 0$$

where P is a polynomial without constant or linear terms, on the space $H^s(\mathbb{R}) \cap L^2(|x|^m dx)$ with s, m being a sufficiently large positive integer. ([KPV94a],[KPV94b]) Kwon [Kwo08] proved the local well-posedness of (5KdV) in H^s with $s > 5/2$ without the weighted L^2 structure. Kenig and Pilod [KP12] proved that the equations (5KdV) with $\beta = 2\alpha$ and (KW) are globally well-posed in H^s with $s \geq 2$. Very recently, Bringmann et al. [BKV19] proved that (5KdV) with $c_1 = 10, c_2 = 20, c_3 = 30$ is globally well-posed in $H^{-1}(\mathbb{R})$. For the summary of the known result of the Cauchy problem of (KW), we refer to [KAT13].

Also, the small data global analysis of nonlinear dispersive equations has been spotlighted for a long time. For instance, [Caz03] introduces some classical scattering results for defocusing

NLS, together with asymptotic completeness. It also mentions that the one-dimensional cubic NLS

$$(3\text{NLS}) \quad iu_t + u_{xx} = \lambda|u|^2u, \quad \lambda = \pm 1$$

does not show a linear scattering. Rather, the modified scattering results for (3NLS) and similar types of equations are proved by many authors with different settings. Ifrim and Tataru [IT15] proved the global dynamics of (3NLS) by testing the wave packets. Adopting this idea of Ifrim and Tataru, Harrop-Griffiths [Har16] proved the modified scattering and asymptotic completeness results for the mKdV equation, and Okamoto [Oka+18] proved the global behavior of solutions to the fifth-order mKdV type equation, without proving asymptotic completeness, using the same idea as above. Okamoto later proved [Oka19] the asymptotic behavior of solutions to the higher order KdV-type equation with critical nonlinearity. Here the nonlinearity is said to be critical in the sense that the small data solution shows the modified scattering behavior at that order of nonlinearity.

In fact, Deift and Zhou [DZ93] proved the small data asymptotics of mKdV first, taking advantage of the complete integrability of the equation. Hayashi and Naumkin [HN99; HN01; HN06] proved the similar result without using the complete integrability, which helped them extend the result for nonlinearities with time-dependent coefficients. However, their results required an initial data to have zero mean. The work of Harrop-Griffiths [Har16] is meaningful in that it requires neither the integrability of the equation, nor the zero mean of the initial datum.

Unlike the mKdV-type equations, the global dynamics of KdV-type equations have not been studied so much. The main difficulty of the global analysis of KdV-type equations is that the nonlinearities are less perturbative than those of the equations on the mKdV hierarchy; the nonlinearities of mKdV-type equations are cubic, whereas those of KdV-type equations are quadratic. The first result regarding the nonlinear dispersive equation with a quadratic nonlinearity is due to [IT17], which discovered that the solution to the Benjamin-Ono equation

$$u_t + Hu_{xx} = uu_x$$

has linear dispersive decay bound in an almost global time scale $|t| \sim e^{\frac{\epsilon}{\epsilon}}$ where ϵ is the size of an initial datum. A global result for KdV equation is discovered fairly recently, by Ifrim et al., [IKT19] which stated that the linear dispersive decay bound may break down beyond the quartic time scale $|t| \sim \epsilon^{-3}$ where ϵ is the size of an initial datum. They proved that the result is optimal, i.e. the time scale cannot be extended further, using the inverse scattering theory. No other meaningful results are known about the global dynamics of KdV-type equations. The main goal of this paper is to show the first result regarding the global behavior of the fifth-order dispersive equations with quadratic nonlinearities.

1.1. Main result. First we state the main result of this paper:

Theorem 1.1. *Consider the Cauchy problem*

$$\begin{cases} u_t - u_{5x} = \pm u^m u_x, \\ u(0, x) = u_0(x), \end{cases} \quad m = 1, 2.$$

Suppose that the initial datum u_0 satisfies

$$\|u_0\|_{\dot{B}_{2,\infty}^{-\frac{1}{2}}} + \|xu_0\|_{\dot{H}^{\frac{1}{2}}} \leq \epsilon \ll 1,$$

then we have the following pointwise bound

$$|\partial_x^k u(t, x)| \lesssim \epsilon t^{-\frac{1}{8} - \frac{k}{4}} \langle x \rangle^{-\frac{3}{8} + \frac{k}{4}}, \quad k = 0, 1, 2, 3$$

provided $|t| \ll \epsilon^{-\frac{5}{5-2m}}$.

Remark 1.2. Note that the case $m = 1$ corresponds to (KW), and $m = 2$ to (mKW).

Remark 1.3. The time scale in (KW) is worse than that in the KdV equation, which is even worse than that in the Benjamin-Ono equation. This can be deduced heuristically as follows: The group velocity of the fifth-order linear equation is given as $\omega(\xi) = -5\xi^4$, so the high frequency part of the solution travels much faster towards to the negative x -axis than in the second or third-order equations. However, the possible emergence of nonlinear “bulk”, such as solitons and dispersive shocks, interrupts such linear dispersive decay. The faster the linear part travels, the earlier the nonlinear bulk emerges.

The time scale in (mKW) is better than that of the KdV and (KW). This is because the nonlinearity becomes cubic, making it more perturbative. However, such time scale is worse than that of the mKdV equation, as the mKdV equation has modified scattering and global dispersive decay bound; see [Har16]. Such discrepancy arises from the same heuristics used in the comparison between (KW), KdV, and Benjamin-Ono equations. One may also compare this result with [Oka+18] or [Oka19], in which case the nonlinearities become more perturbative than (mKW).

Remark 1.4. The two norms used in the main result are inspired from [IKT19]. As seen in the later analysis, the $\dot{B}_{2,\infty}^{-\frac{1}{2}}$ norm helps properly control the low frequency, and the growth of the high frequency norms is slow enough that it does not disrupt our analysis. The $\dot{H}^{\frac{1}{2}}$ norm exactly does the opposite. It does the control of the high frequency, and guarantees the mild growth of the low frequency part.

Let us explain the overall strategies of this paper. Mainly we focus on the case of (KW), since the case of (mKW) is entirely similar. Basically we are going to follow the argument of [IKT19], but the case becomes much simpler, thanks to the higher dispersive effect on high frequencies. In the Section 2, we are going to list some definitions, notations, and well-known theorems that will be used in the paper. Then in the Section 3 we find the pointwise decay bound of the linear solution under the same initial data assumption with Theorem 1.1. The argument used in that section will be also used in the last section, with some modifications for nonlinearities. In the Section 4, we establish a bootstrap assumption motivated from the linear analysis, and list the main propositions of the Section 5 and further. The Sections 5, 6, and 7 are dedicated for proving the propositions listed in the Section 4, which are the key part of proving the main theorem. In the appendix, we briefly sketch the proof of the main theorem in the case of (mKW).

2. PRELIMINARIES

2.1. Definitions and Notations. Let $1 \leq p < \infty$. $L^p(X)$ be the set of measurable function on X with the norm $\|f\|_{L^p(X)} = (\int_X |f|^p d\mu)^{1/p}$ on the measure space (X, μ) .

If $p = \infty$, then $\|f\|_{L^\infty(X)} = \inf\{M > 0 : |f(x)| \leq M \text{ a.e.}\}$. If $X = \mathbb{R}$ and μ is the Lebesgue measure, then we write $L^p(X) = L^p$ for simplicity, in either case $p < \infty$ or $p = \infty$.

We use the time-dependent Japanese bracket $\langle x \rangle = (x^2 + t^{\frac{1}{5}})^{\frac{1}{2}}$.

We define the spatial Fourier transform of $f \in \mathcal{S}(\mathbb{R})$ on \mathbb{R} by

$$\hat{f}(\xi) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) e^{-ix\xi} dx,$$

and its inverse transform by

$$f^\vee(x) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(\xi) e^{-ix\xi} d\xi.$$

Both transforms can be extended to $\mathcal{S}'(\mathbb{R})$, the space of tempered distributions, by duality.

Let A, B be two nonnegative quantities. If there exists $C > 0$ such that $A \leq CB$, then we denote by $A \lesssim B$. If A need not be nonnegative and $|A| \lesssim B$, then we denote $A = O(B)$. If $A \lesssim B$ and $B \lesssim A$, then we denote $A \sim B$. If the implied constant C depends on some parameters a, b, c, \dots , then we denote $A \lesssim_{a,b,c,\dots} B$, $A = O_{a,b,c,\dots}(B)$, $A \sim_{a,b,c,\dots} B$ respectively.

Let $\Psi \in \mathcal{S}(\mathbb{R})$ with $\psi = \widehat{\Psi}$ is even, supported on $[-2, 2]$, and $\psi = 1$ on $[-1, 1]$. Also let $\psi_{\leq N}(\xi) = \psi(\frac{\xi}{N})$ and $\Psi_{\leq N} = (\psi_{\leq N})^\vee$ for $N > 0$. Then define the Littlewood-Paley projection of $f \in \mathcal{S}'(\mathbb{R})$ by $P_{\leq N} f(x) = (\Psi_{\leq N} * f)(x)$, $P_N = P_{\leq N} - P_{\leq N/2}$. Also define $P_{< N} := P_{\leq N} - P_N$, $P_{\geq N} = 1 - P_{< N}$, $P_{> N} = 1 - P_{\leq N}$. By a slight abuse of notation, we denote a Fourier multiplier whose symbol is supported on $\{|\xi| \sim N\}$ by P_N . In particular, if $N = 2^j \in 2^{\mathbb{Z}}$, then we denote $P_N := P_j$ and similarly for $P_{\leq N}$, $P_{< N}$, $P_{> N}$, $P_{\geq N}$.

Now define Sobolev and Besov spaces. For $s \in \mathbb{R}$ and $1 < p < \infty$, the inhomogeneous Sobolev space $W^{s,p}$ and its homogeneous counterpart $\dot{W}^{s,p}$ are defined as the subspace of $\mathcal{S}'(\mathbb{R})$ under the norm

$$\|f\|_{W^{s,p}} = \left\| \left((1 + \xi^2)^{s/2} \hat{f} \right)^\vee \right\|_{L^p}, \quad \|f\|_{\dot{W}^{s,p}} = \left\| \left(|\xi|^s \hat{f} \right)^\vee \right\|_{L^p},$$

respectively. Recall the Littlewood-Paley inequality

$$\|f\|_{W^{s,p}} \sim_{s,p} \|P_{\leq 0} f\|_{L^p} + \left\| \left(\sum_{j \geq 1} 2^{2js} |P_j f|^2 \right)^{1/2} \right\|_{L^p}$$

and its homogeneous counterpart

$$\|f\|_{\dot{W}^{s,p}} \sim_{s,p} \left\| \left(\sum_{j \in \mathbb{Z}} 2^{2js} |P_j f|^2 \right)^{1/2} \right\|_{L^p}.$$

Motivated by the above Littlewood-Paley inequalities, we define the inhomogeneous and homogeneous Besov spaces, denoted as $B_{p,q}^s$ and $\dot{B}_{p,q}^s$, respectively, is defined as the subspace of $\mathcal{S}'(\mathbb{R})$ under the norm

$$\|f\|_{B_{p,q}^s} = \left(\|P_{\leq 1} f\|_{L^p}^q + \sum_{j \geq 1} 2^{qjs} \|P_j f\|_{L^p}^q \right)^{1/q},$$

$$\|f\|_{\dot{B}_{p,q}^s} = \left(\sum_{j \in \mathbb{Z}} 2^{qjs} \|P_j f\|_{L^p}^q \right)^{1/q},$$

with $1 \leq q < \infty$ and the natural modification for $q = \infty$.

2.2. Useful theorems. In this subsection, some of the theorems are presented for the later use. I omit the proofs here, since the proofs can be found in many literatures.

Theorem 2.1 (Bernstein's inequality). *Let $1 \leq p \leq q \leq \infty$. Then one has*

$$\|P_N f\|_{L^q} \lesssim_{p,q} N^{\frac{1}{p}-\frac{1}{q}} \|P_N f\|_{L^p}.$$

The same result holds when P_N is replaced by $P_{\leq N}$.

Also we have the Gagliardo-Nirenberg inequality, a refined version of the Sobolev embedding:

Theorem 2.2 (Gagliardo-Nirenberg inequality). *Let $1 < p < q \leq \infty$ and $s > 0$ be such that*

$$\frac{1}{q} = \frac{1}{p} - \frac{\theta s}{d}$$

for some $0 < \theta < 1$. Then for any $u \in W^{s,p}(\mathbb{R}^d)$ we have

$$\|u\|_{L^q(\mathbb{R}^d)} \lesssim_{d,p,q,s} \|u\|_{L^p(\mathbb{R}^d)}^{1-\theta} \|u\|_{W^{s,p}(\mathbb{R}^d)}^\theta.$$

For the proof, see for example [Tao06].

Throughout the paper, a special case of Theorem 2.2, namely

$$\|u\|_{L^\infty(\mathbb{R})} \lesssim \|u\|_{L^2(\mathbb{R})}^{1/2} \|u\|_{\dot{H}^1(\mathbb{R})}^{1/2},$$

will be used frequently.

Let M be one-dimensional Hardy-Littlewood maximal operator defined as

$$Mf(x) = \sup_{r>0} \frac{1}{2r} \int_{x-r}^{x+r} |f(y)| dy.$$

Then we have following pointwise bound by [FO00] :

Lemma 2.3. *Let f be a function whose Fourier support is in the annulus $|\xi| \sim R$. Then for any $x \in \mathbb{R}$ and $s \in \mathbb{R}$,*

$$||D|^s f(x)| \lesssim R^s Mf(x).$$

Moreover, if f is not necessarily frequency-localized, then

$$||D|^s P_N f(x)| \lesssim N^s Mf(x).$$

In both cases, if s is an integer, then replacing $|D|$ by ∂_x is acceptable. Also, if $s \geq 0$, then the annulus $|\xi| \sim R$ can be replaced by the disk $|\xi| \lesssim R$.

Remark 2.4. The above lemma implies that a polynomially decaying pointwise bound is stable under the frequency restriction, since if $g_s(x) = \langle x \rangle^{-s}$ for $s \geq 0$, then $Mg_s(x) \lesssim g_s(x)$. Thus, we can apply the bootstrap bounds of u appearing in the later sections freely to the frequency-localized pieces of u .

3. LINEAR ANALYSIS

First we consider the decay estimate of the solution of linear fifth-order KdV equation

$$(Lin) \quad u_t - u_{5x} = 0.$$

We introduce the time-dependent linear operator $L(t) := x + 5t\partial_x^4$, so that if u solves (Lin) with initial datum u_0 then Lu also solves the same equation with initial datum xu_0 .

Now we state the basic decay bound for the solution of (Lin), which is given as below:

Proposition 3.1. *Let u be a solution of (Lin) with initial datum $u_0 \in L^1$. Then u satisfies the uniform decay bound*

$$\|u(t)\|_{L^\infty} \lesssim t^{-1/5} \|u_0\|_{L^1}.$$

Proof. Solving the equation via spatial Fourier transform gives

$$u(t, x) = \int_{\mathbb{R}} \int_{\mathbb{R}} e^{i\xi^5 t + i\xi(x-y)} u_0(y) dy d\xi.$$

We change coordinates $\xi \mapsto t^{-1/5}\eta$ so that

$$u(t, x) = t^{-1/5} \int_{\mathbb{R}} \int_{\mathbb{R}} e^{i\eta^5 + i\eta t^{-1/5}(x-y)} u_0(y) dy d\eta = t^{-1/5} A(t^{-1/5}x) * u_0(x),$$

where $A(x)$ is an oscillatory integral

$$A(x) := \int_{\mathbb{R}} e^{i\eta^5 + i\eta x} d\eta.$$

Then A is bounded by standard stationary phase argument. Now the result follows from Young's inequality. \square

To obtain a more refined version of the above estimate, we need the following lemma due to [Dur14] and [Fuj07]:

Lemma 3.2. *$A(x)$ in the proof of Proposition 2.1 satisfy the following asymptotic bound:*

$$A(x) \lesssim \langle x \rangle^{-3/8} e^{-Cx_+^{5/4}},$$

where C is an appropriate positive constant.

From the above lemma, it follows that solutions with initial datum u_0 satisfying $\|u_0\|_{L^1} \leq 1$ and $\text{supp} u_0$ bounded has the following decay bound for $t \gtrsim 1$:

$$|u(t, x)| \lesssim t^{-1/8} \langle x \rangle^{-3/8} e^{-Cx_+^{5/4} t^{-1/4}}, \quad |u_x(t, x)| \lesssim t^{-3/8} \langle x \rangle^{-1/8} e^{-C'x_+^{5/4} t^{-1/4}}$$

Now we aim to relax the compact support assumption to a decay estimate, in the similar way with [IKT19]. Our goal of this section is to prove the following proposition:

Proposition 3.3. *Let $t > 0$. Assume that a function u satisfies:*

$$(3.1) \quad \|u\|_{\dot{B}_{2,\infty}^{-1/2}} + \|L(t)u\|_{\dot{H}^{1/2}} \leq 1.$$

Then it also satisfies the bound

$$(3.2) \quad t^{\frac{1}{8} + \frac{k}{4}} \langle x \rangle^{\frac{3}{8} - \frac{k}{4}} |\partial_x^k u| \lesssim 1$$

for $k = 0, 1, 2, 3$. Moreover, in the elliptic region E we have the better bound

$$(3.3) \quad t^{\frac{k}{4}} \langle x \rangle^{1 - \frac{k}{4}} |\partial_x^k u(x)| \lesssim \log(t^{-\frac{1}{5}} \langle x \rangle).$$

This is an analogous result of the Lemma 2.2 in [IKT19]. The only difference with it is the linear operator $L(t)$, where $L(t) = x - 3t\partial_x^2$ in [IKT19] and $L(t) = x + 5t\partial_x^4$ here. The steps of the proof will basically follow the idea of [IKT19]. First we may rescale to $t = 1$ since $\|u\|_{\dot{B}_{2,\infty}^{-1/2}} + \|L(t)u\|_{\dot{H}^{1/2}}$ is invariant under the scaling $u(\lambda^5, x) \mapsto \lambda u(\lambda^5, \lambda x)$. Also we will split the real line into the self-similar region $S = \{|x| \lesssim 1\}$ (would be $\{|x| \lesssim t^{1/5}\}$ in general), the elliptic region $E = \{x \gg 1\}$ (would be $\{x \gg t^{1/5}\}$ in general), and the hyperbolic region $H = \{-x \gg 1\}$. Also we split the elliptic and hyperbolic regions into dyadic components, namely $A_R = \{\langle x \rangle \sim R \gg 1\}$ and $A_1 = \{\langle x \rangle \lesssim 1\} = S$, $A_R^H = A_R \cap H$, $A_R^E = A_R \cap E$ for $R \gg 1$. Therefore the proof is reduced to showing that

$$(3.4) \quad \|\partial_x^k u\|_{L^\infty(A_R)} \lesssim R^{-\frac{3}{8} + \frac{k}{4}} \log R$$

for each $R \geq 1$ and $k = 0, 1, 2, 3$ (and the corresponding counterpart for the elliptic region).

3.1. Localized bounds of a linear solution. Now we present the following low frequency bound, which is the same as in [IKT19]:

Lemma 3.4. *If (3.1) holds, we have*

$$(3.5) \quad \|Lu\|_{L^2(A_R)} \lesssim R^{1/2}.$$

Proof. The proof is almost identical to the case of [IKT19], but I present the details here to clarify the ideas.

First we split u at some frequency cut-off M , where M is to be chosen later:

$$u = u_{<M} + u_{\geq M}.$$

Then the Besov bound from (3.1) tells that $\|u_M\|_{L^2(\mathbb{R})} \lesssim M^{1/2}$. Hence

$$\|u_{<M}\|_{L^2(\mathbb{R})} \leq \sum_{N < M} \|u_N\|_{L^2(\mathbb{R})} \lesssim \sum_{N < M} N^{1/2} \sim M^{1/2}$$

where the sum runs over all dyadic numbers $< M$. Therefore one has

$$(3.6) \quad \|Lu_{<M}\|_{L^2(A_R)} \leq \|xu_{<M}\|_{L^2(A_R)} + \|\partial_x^4 u_{<M}\|_{L^2(\mathbb{R})} \lesssim RM^{1/2} + M^{9/2}.$$

On the other hand,

$$Lu_{\geq M} = P_{\geq M}Lu + [L, P_{\geq M}]u = P_{\geq M}Lu + [x, P_{\geq M}]u,$$

where the last equality follows since $[\partial_x^4, P_{\geq M}] = 0$ (note that any two Fourier multipliers commute). Now we can estimate

$$\|P_{\geq M}Lu\|_{L^2(A_R)} \leq \|(1 - \psi(\xi/M))\widehat{Lu}\|_{L^2(\mathbb{R})} \lesssim M^{-1/2} \|Lu\|_{\dot{H}^{1/2}(\mathbb{R})} \leq M^{-1/2},$$

with ψ being a bump function supported in $|\xi| \lesssim 1$. Also we have

$$\|[x, P_{\geq M}]u\|_{L^2(A_R)} \leq \|M^{-1}\psi'(\xi/M)\hat{u}\|_{L^2(\mathbb{R})} \lesssim M^{-1/2}$$

by (3.1). This gives

$$(3.7) \quad \|Lu_{\geq M}\|_{L^2(A_R)} \lesssim M^{-1/2}.$$

Now we are going to optimize the bound, compromising between (3.6) and (3.7) by choosing the appropriate M . Note that the bound in (3.6) becomes smaller if M is small and the same holds for the bound in (3.7) if M is large. Hence the bound is optimized when the two bounds

are comparable. If $RM^{1/2} \sim M^{-1/2}$, then $M \sim R^{-1}$, and the resulting bound becomes $R^{1/2}$. Also if $M^{9/2} \sim M^{-1/2}$, then $M \sim 1$, so the resulting bound becomes R , and the former is better, so we choose $M = R^{-1}$. Hence (3.5) follows. \square

Remark 3.5. The above argument works in both cases $R = 1$ and $R > 1$.

Lemma 3.6. *If (3.1) holds, we have*

$$(3.8) \quad \|\partial_x^k u\|_{L^2(A_R)} \lesssim R^{\frac{1}{8} + \frac{k}{4}}, \quad k = 0, 1, 2, 3, 4.$$

Proof. Again we split $u = u_{<M} + \sum_{\lambda \geq M} u_\lambda$ and choose M later as before. The bound on $u_{<M}$ follows by:

$$(3.9) \quad \|\partial_x^k u_{<M}\|_{L^2(A_R)} \lesssim M^k \|u_{<M}\|_{L^2(\mathbb{R})} \lesssim M^{k + \frac{1}{2}}$$

where the last inequality follows from (3.1). Now consider the high frequency bound. The bound of Lu_λ can be computed in the same way with $Lu_{\geq M}$ in Lemma 2.4, and the bound of u_λ directly follows from (3.1) so that

$$(3.10a) \quad \|Lu_\lambda\|_{L^2} \lesssim \lambda^{-1/2},$$

$$(3.10b) \quad \|u_\lambda\|_{L^2} \lesssim \lambda^{1/2},$$

$$(3.10c) \quad \|u_{\lambda,x}\|_{L^2} \lesssim \lambda^{3/2}.$$

Now integrating by parts gives

$$\int \chi_R u_{\lambda,2x}^2 dx = -\frac{1}{5} \int \chi_R u_\lambda Lu_\lambda + \frac{1}{5} \int \chi_R x u_\lambda^2 + \int (\chi_R'' u_{\lambda,2x} + 2\chi_R' u_{\lambda,3x}) u_\lambda.$$

The first integral can be estimated via Cauchy-Schwarz inequality, which yields the bound 1 by (3.10a) and (3.10b), and the bound of the second integral follows directly from (3.10b), which yields $R\lambda$. To bound the third integral, we integrate by parts again to obtain:

$$\int (\chi_R'' u_{\lambda,2x} + 2\chi_R' u_{\lambda,3x}) u_\lambda = -\frac{1}{2} \int \chi_R^{(4)} u_\lambda^2 + 2 \int \chi_R'' u_{\lambda,x}^2.$$

Hence the third integral is bounded by $R^{-4}\lambda + R^{-2}\lambda^3$ due to (3.10b), (3.10c), and the fact $\|\partial_x^k \chi_R\|_{L^\infty} = O_{k,\chi}(R^{-k})$. This gives the preliminary bound

$$(3.11) \quad \|u_{\lambda,2x}\|_{L^2(A_R)} \lesssim 1 + R^{1/2}\lambda^{1/2} + R^{-1}\lambda^{3/2} := N.$$

Repeating the similar steps for the integral $\int \chi_R u_{\lambda,3x}^2 dx$ one gets the similar bound:

$$(3.12) \quad \|u_{\lambda,3x}\|_{L^2(A_R)} \lesssim R^{-1}N + \lambda^{1/4}N^{1/2} + R^{1/2}\lambda^{1/4}N^{1/2} := N'.$$

Now we are going to refine (3.10b) and (3.10c) using (3.11) and (3.12) so that the bound becomes summable with respect to λ . To do so we decompose

$$\chi_R u_\lambda = \partial_{x,\lambda}^{-2}(\chi_R u_{\lambda,2x}) - [\partial_{x,\lambda}^{-2}, \chi_R] u_{\lambda,2x},$$

where the antiderivative $\partial_{x,\lambda}^{-2}$ is localized at frequency λ , namely having the symbol $-\xi^{-2}\varphi(\xi/\lambda)$ with φ being a bump function supported on the annulus $|\xi| \sim 1$. Let ψ be the Fourier inversion

of $-\xi^{-2}\varphi(\xi)$, so that $\psi \in \mathcal{S}(\mathbb{R})$. Then the inversion of $-\xi^{-2}\varphi(\xi/\lambda)$ becomes $\lambda^{-1}\psi(\lambda x)$. Then one has

$$\begin{aligned} \|[\partial_{x,\lambda}^{-2}, \chi_R]u_{\lambda,2x}\|_{L^2(\mathbb{R})} &= \lambda^{-1} \left\| \int \psi(\lambda(x-y))(\chi_R(y) - \chi_R(x))u_{\lambda,2x}(y)dy \right\|_{L_x^2(\mathbb{R})} \\ &\lesssim \lambda^{-1} \| |\psi(\lambda \cdot)| * |u_{\lambda,2x}| \|_{L_x^2} \leq \lambda^{-2} \|\psi\|_{L_x^1} \|u_{\lambda,2x}\|_{L_x^2} \\ &\lesssim \lambda^{-2} N. \end{aligned}$$

Also

$$\|\partial_{x,\lambda}^{-2}(\chi_R u_{\lambda,2x})\|_{L^2(\mathbb{R})} \lesssim \lambda^{-2} \|u_{\lambda,2x}\|_{L^2(A_R)} \lesssim \lambda^{-2} N.$$

This yields a local bound for u_λ :

$$(3.13) \quad \|u_\lambda\|_{L^2(A_R)} \lesssim \lambda^{-2} N.$$

Now summing up (3.13) on the frequency range $\lambda \geq M$ gives

$$(3.14) \quad \|u_{\geq M}\|_{L^2(A_R)} \lesssim M^{-2} + R^{1/2}M^{-3/2} + R^{-1}M^{-1/2}.$$

Repeating the same argument above to the decomposition

$$(3.15) \quad \chi_R u_{\lambda,x} = \partial_{x,\lambda}^{-2}(\chi_R u_{\lambda,3x}) - [\partial_{x,\lambda}^{-2}, \chi_R]u_{\lambda,3x}$$

we also get the refined local bound for $u_{\lambda,x}$:

$$(3.16) \quad \|u_{\lambda,x}\|_{L^2(A_R)} \lesssim \lambda^{-2} N',$$

so that

$$(3.17) \quad \|\partial_x u_{\geq M}\|_{L^2(A_R)} \lesssim R^{-1}M^{-2} + R^{-2}M^{-\frac{1}{2}} + R^{\frac{1}{2}}M^{-\frac{7}{4}} + R^{\frac{3}{4}}M^{-\frac{3}{2}} + M^{-1}.$$

Now we want to optimize the bounds (3.9), (3.14), (3.17) by making them comparable. It turns out that $M = R^{1/4}$ is the case with the best bound, so that $M^{-2}N \sim R^{1/8}$. This gives

$$(3.18a) \quad \|u\|_{L^2(A_R)} \lesssim R^{1/8},$$

$$(3.18b) \quad \|u_x\|_{L^2(A_R)} \lesssim R^{3/8},$$

$$(3.18c) \quad \|u_{2x}\|_{L^2(A_R)} \lesssim R^{5/8}.$$

Here the bound (3.18c) follows from integration by parts below with (3.5), (3.18a), and (3.18b):

$$\int \chi_R u_{2x}^2 dx = -\frac{1}{5} \int \chi_R u L u + \frac{1}{5} \int \chi_R x u^2 + \int (\chi_R'' u_{2x} + 2\chi_R' u_{3x}) u.$$

Using the bound (3.5) and (3.18a) easily gives that

$$(3.19) \quad \|u_{4x}\|_{L^2(A_R)} \lesssim R^{9/8}.$$

Finally, it remains to find the bound of $\|u_{3x}\|_{L^2(A_R)}$. This can be done easily by integration by parts:

$$\int \chi_R u_{3x}^2 = - \int \chi_R' u_{3x} u_{2x} - \int \chi_R u_{4x} u_{2x} = \frac{1}{2} \int \chi_R'' u_{2x}^2 - \int \chi_R u_{4x} u_{2x}$$

so the bound follows from (3.18c) and (3.19). \square

3.2. Pointwise estimation on each region. Let $v := \chi_R u$. For notational convenience, we are going to restrict the norm to the region A_R , so we will write $L^2(A_R)$ simply as L^2 . Then v solves an equation of the form

$$(x + 5\partial_x^4)v = f$$

in A_R , where we control

$$(3.20) \quad \|\partial_x^k v\|_{L^2} \lesssim R^{\frac{1}{8} + \frac{k}{4}}, \quad 0 \leq k \leq 4$$

by (3.8), and

$$(3.21) \quad \|f\|_{\dot{H}^{1/2}} \lesssim 1, \quad \|f\|_{L^2} \lesssim R^{\frac{1}{2}}$$

which follows by (3.1), (3.5).

Now we turn to the pointwise estimate in each region. In the case of self-similar region, the result follows directly from Sobolev embedding (note that in this region just $\lesssim 1$ bound is enough, since $R = 1$.) Hence we concentrate on elliptic and hyperbolic regions.

First we see the elliptic region, namely $x \sim R$, which seems to be easier to deal with than the hyperbolic region. Split $f = f_{lo} + f_{hi}$, where the frequency scale is $R^{1/4}$. Here the operator $L = x + 5\partial_x^4$ is elliptic, so the leading part will then be $x^{-1}f_{lo}$. Then let $v_1 := v - x^{-1}f_{lo}$ be the remainder, which solves

$$Lv_1 = f_1 := f_{hi} - 5\partial_x^4(x^{-1}f_{lo}).$$

By (2) one has

$$\begin{aligned} \|f_{hi}\|_{L^2(\mathbb{R})} &\lesssim R^{-1/8} \|f\|_{\dot{H}^{1/2}} \lesssim R^{-1/8}, \\ \|\partial_x^4(x^{-1}f_{lo})\|_{L^2(\mathbb{R})} &\lesssim R^{-5} \|f_{lo}\|_{L^2(\mathbb{R})} + R^{-4} \|\partial_x f_{lo}\|_{L^2(\mathbb{R})} + R^{-3} \|\partial_x^2 f_{lo}\|_{L^2(\mathbb{R})} \\ &\quad + R^{-2} \|\partial_x^3 f_{lo}\|_{L^2(\mathbb{R})} + R^{-1} \|\partial_x^4 f_{lo}\|_{L^2(\mathbb{R})} \\ &\lesssim R^{-9/2} + R^{-31/8} + R^{-21/8} + R^{-11/8} + R^{-1/8} \\ &\lesssim R^{-1/8}, \end{aligned}$$

where for $k \geq 1$,

$$\|\partial_x^k f_{lo}\|_{L^2(\mathbb{R})} \lesssim R^{\frac{k}{4} - \frac{1}{8}} \|f_{lo}\|_{\dot{H}^{1/2}} \lesssim R^{\frac{k}{4} - \frac{1}{8}},$$

which yield

$$\|f_1\|_{L^2(\mathbb{R})} \lesssim R^{\frac{k}{4} - \frac{1}{8}}.$$

Hence we may integrate by parts in the following identity

$$\int_{\mathbb{R}} v_1 Lv_1 dx = \int_{\mathbb{R}} f_1 v_1 dx$$

and arrive at

$$\int_{\mathbb{R}} x |v_1|^2 dx + 5 \int_{\mathbb{R}} |\partial_x^2 v_1|^2 dx = \int_{\mathbb{R}} f_1 v_1 dx.$$

Then by Cauchy-Schwarz inequality,

$$R \|v_1\|_{L^2(\mathbb{R})}^2 + 5 \|\partial_x^2 v_1\|_{L^2(\mathbb{R})}^2 \lesssim \|f_1\|_{L^2(\mathbb{R})} \|v_1\|_{L^2(\mathbb{R})} \leq \delta R \|v_1\|_{L^2(\mathbb{R})}^2 + \frac{4}{\delta R} \|f_1\|_{L^2(\mathbb{R})}^2$$

for any $\delta > 0$, so letting $\delta > 0$ sufficiently small gives

$$R \|v_1\|_{L^2(\mathbb{R})}^2 + 5 \|\partial_x^2 v_1\|_{L^2(\mathbb{R})}^2 \lesssim R^{-1} \|f_1\|_{L^2(\mathbb{R})}^2.$$

Thus, using the bound on f_1 , we arrive at

$$\|\partial_x^k v_1\|_{L^2(\mathbb{R})} \lesssim R^{-\frac{9}{8} + \frac{k}{4}}, \quad k = 0, 2,$$

and using the v_1 equation shows that this bound also holds for $k = 4$, and by interpolation it still holds for $k = 1, 3$. To sum up, we have

$$\|\partial_x^k v_1\|_{L^2(\mathbb{R})} \lesssim R^{-\frac{9}{8} + \frac{k}{4}}, \quad k = 0, 1, 2, 3, 4.$$

Finally, by Gagliardo-Nirenberg inequality, one has

$$|\partial_x^k v_1| \lesssim R^{-1 + \frac{k}{4}}, \quad k = 0, 1, 2, 3,$$

which is exactly as needed.

Now we need to get the bound of $x^{-1}f_{lo}$. Proceeding as above, one has

$$\|\partial_x^k(x^{-1}f_{lo})\|_{L^2(\mathbb{R})} \lesssim R^{-\frac{9}{8} + \frac{k}{4}}$$

for $k = 1, 2, 3, 4$. Hence again Gagliardo-Nirenberg inequality gives

$$\|x^{-1}f_{lo}\|_{L^2(\mathbb{R})} \lesssim R^{-1 + \frac{k}{4}}, \quad k = 1, 2, 3.$$

For $k = 0$, we decompose f_{lo} further as

$$f_{lo} = f_{<R^{-1}} + \sum_{R^{-1} \leq \lambda \leq R^{1/4}} f_\lambda.$$

Then Bernstein's inequality and Littlewood-Paley characterization of Sobolev norms give

$$\begin{aligned} \|f_{<R^{-1}}\|_{L^\infty(\mathbb{R})} &\lesssim R^{-1/2} \|f_{<R^{-1}}\|_{L^2(\mathbb{R})} \lesssim 1, \\ \sum_{R^{-1} \leq \lambda \leq R^{1/4}} \|f_\lambda\|_{L^\infty(\mathbb{R})} &\lesssim \sum_{R^{-1} \leq \lambda \leq R^{1/4}} \lambda^{1/2} \|f_\lambda\|_{L^2(\mathbb{R})} \\ &\lesssim (\log R)^{1/2} \left(\sum_{R^{-1} \leq \lambda \leq R^{1/4}} \lambda \|f_\lambda\|_{L^2(\mathbb{R})}^2 \right)^{1/2} \\ &\lesssim (\log R)^{1/2} \|f\|_{\dot{H}^{1/2}} \lesssim (\log R)^{1/2}. \end{aligned}$$

where the inequality used in the penultimate line is just an elementary inequality

$$\sum_{k=1}^n a_k \leq (n \sum_{k=1}^n a_k^2)^{1/2}.$$

Hence one has

$$(3.22) \quad \|x^{-1}f_{lo}\|_{L^\infty} \lesssim R^{-1} (\log R)^{1/2},$$

which is exactly needed.

Finally we turn to the hyperbolic region. As in the third-order case, f might be worse at lower frequency (below $R^{\frac{1}{4}}$) than higher frequencies, due to the weight $|\xi|^{1/2}$. Hence we proceed by splitting f to low and high frequencies:

$$f = \tilde{\chi}_R f_{\leq R^{1/4}} + \tilde{\chi}_R f_{> R^{1/4}} := f_{lo} + f_{hi},$$

where $\tilde{\chi}_R$ is smooth, supported on $|x| \sim R$, and $\tilde{\chi}_R \chi_R = \chi_R$.

Then we simply treat the v equation as a hyperbolic evolution equation and use an energy estimate,

$$\frac{d}{dx} (-x|v|^2 + 5|v_{xx}|^2 - 10v_x v_{3x} + 2f_{lo}v) = -|v|^2 - 2f_{hi}v_x + 2f_{lo,x}v \leq -2f_{lo,x}v + 2f_{lo,x}v,$$

and integrating both sides gives

$$(3.23) \quad -x|v|^2 + 5|v_{xx}|^2 - 10v_x v_{3x} \lesssim \|f_{hi}\|_{L^2} \|v_x\|_{L^2} + \|f_{lo,x}\|_{L^2} \|v\|_{L^2} + |f_{lo}v|$$

whenever $x \in A_R^H$, so the term $|f_{lo}v|$ can be controlled via (3.22) and the Gagliardo-Nirenberg inequality

$$\|v\|_{L^\infty} \lesssim \|v\|_{L^2}^{\frac{1}{2}} \|v_x\|_{L^2}^{\frac{1}{2}} \lesssim R^{\frac{1}{4}},$$

and the other terms can be bounded by (3.20) and (3.21). This finally gives

$$(3.24) \quad -x|v|^2 + 5|v_{xx}|^2 - 10v_x v_{3x} \lesssim R^{\frac{1}{4}} (\log R)^{\frac{1}{2}}.$$

Now we have a task to deal with the extra $v_x v_{3x}$ term, which was absent in the third-order case. The problem is that, we should control the pointwise bound of $v_x v_{3x}$, either.

Now try a similar estimate as follows:

$$\begin{aligned} \frac{d}{dx}(-x|v_x|^2 + 5|v_{3x}|^2 + 2xvv_{xx} + 2f_{lo}v_{2x}) &\leq 2vv_{xx} + 2f_{hi}v_{3x} - 2f_{lo,x}v_{2x} \\ &\lesssim |x|^{-\frac{1}{2}}(-x|v|^2 + 5|v_{xx}|^2) + |f_{hi}v_{3x}| + |f_{lo,x}v_{2x}| \\ &\lesssim R^{-\frac{1}{2}}(R^{\frac{1}{4}}(\log R)^{\frac{1}{2}} + 10v_x v_{3x}) + |f_{hi}v_{3x}| + |f_{lo,x}v_{2x}|, \end{aligned}$$

and integrating both sides, together with Cauchy-Schwarz inequality gives

$$\begin{aligned} &-x|v_x|^2 + 5|v_{3x}|^2 + 2xvv_{xx} \\ &\lesssim R^{\frac{3}{4}}(\log R)^{\frac{1}{2}} + R^{-\frac{1}{2}}\|v_x\|_{L^2} \cdot \|v_{3x}\|_{L^2} + \|f_{hi}\|_{L^2} \|v_{3x}\|_{L^2} + \|f_{lo,x}\|_{L^2} \|v_{2x}\|_{L^2} + |f_{lo}v_{2x}| \\ &\lesssim R^{\frac{3}{4}}(\log R)^{\frac{1}{2}} + |f_{lo}v_{2x}| \end{aligned}$$

for $x \in A_R^H$. Here the term $|f_{lo}v_{2x}|$ can be treated in the same way as above, so one has

$$(3.25) \quad -x|v_x|^2 + 5|v_{3x}|^2 + 2xvv_{xx} \lesssim R^{\frac{3}{4}}(\log R)^{\frac{1}{2}}.$$

Now we obtained the system of two inequalities:

$$(3.26a) \quad -x|v|^2 + 5|v_{xx}|^2 - 10v_x v_{3x} \leq CR^{\frac{1}{4}}(\log R)^{\frac{1}{2}},$$

$$(3.26b) \quad -x|v_x|^2 + 5|v_{3x}|^2 + 2xvv_{xx} \leq CR^{\frac{3}{4}}(\log R)^{\frac{1}{2}}.$$

From (3.26a) and (3.26b) we have

$$\begin{aligned} -x|v|^2 + 5|v_{xx}|^2 &\leq CR^{\frac{1}{4}}(\log R)^{\frac{1}{2}} + 10v_x v_{3x} \leq CR^{\frac{1}{4}}(\log R)^{\frac{1}{2}} + \sqrt{\frac{5}{-x}}(-xv_x^2 + 5v_{3x}^2) \\ &\leq CR^{\frac{1}{4}}(\log R)^{\frac{1}{2}} + 2\sqrt{-5x}vv_{2x}, \end{aligned}$$

so that

$$|\sqrt{-x}v - \sqrt{5}v_{2x}| \lesssim R^{\frac{1}{8}}(\log R)^{\frac{1}{4}}$$

and in the same way one can show

$$|\sqrt{-x}v_x - \sqrt{5}v_{3x}| \lesssim R^{\frac{3}{8}}(\log R)^{\frac{1}{4}}.$$

Now it remains to show that

$$(3.27) \quad \|\partial_x^k v\|_{L^\infty(A_R)} \lesssim R^{-\frac{3}{8} + \frac{k}{4}}(\log R)^{\frac{1}{4}}, \quad k = 0, 1, 2, 3.$$

To deduce a contradiction, suppose that $\|v\|_{L^\infty(A_R)} \not\lesssim R^{-\frac{3}{8}}(\log R)^{\frac{1}{4}}$. Then there exists a sequence $\{R_n\} \uparrow \infty$ as $n \rightarrow \infty$ such that $\|v\|_{L^\infty(A_{R_n})} \geq 2^n R_n^{-\frac{3}{8}}(\log R_n)^{\frac{1}{4}}$. Now define the set $B_n := \{x :$

$|v(x)| \geq 2^{n-1} R_n^{-\frac{3}{8}} (\log R_n)^{\frac{1}{4}}$. Here v is assumed to be differentiable sufficiently many times, so by taking a connected component we may assume that $B_n = [a_n, b_n]$ with $a_n < b_n$. Note that $-\infty < a_n$ and $b_n < \infty$ because v is compactly supported.

Now observe by (3.26b) that

$$(3.28) \quad vv_{2x} \geq \frac{1}{-2x} \left(-x|v_x|^2 + 5|v_{3x}|^2 - CR_n^{\frac{3}{4}} (\log R_n)^{\frac{1}{2}} \right) \gtrsim 2^{n-1} R_n^{\frac{3}{4}} (\log R_n)^{\frac{1}{2}}$$

whenever $x \in B_n$ with n large enough. Similarly by (3.26a) one also has

$$(3.29) \quad v_x v_{3x} \geq \frac{1}{10} \left(-x|v|^2 + 5|v_{2x}|^2 - CR_n^{\frac{1}{4}} (\log R_n)^{\frac{1}{2}} \right) \gtrsim 2^{2(n-1)} R_n^{\frac{1}{4}} (\log R_n)^{\frac{1}{2}}$$

under the same assumption above. (3.28) shows that for sufficiently large n , v and v_{2x} always have the same sign on B_n , and the similar statement holds for v_x and v_{3x} .

First we may assume that $v > 0$ on B_n . (Otherwise we may replace v by $-v$.) Then $v \geq 2^{n-1} R_n^{-\frac{3}{8}} (\log R_n)^{\frac{1}{4}}$ and it is strictly convex on B_n since $v_{2x} > 0$. Also by continuity one must have $v(a_n) = v(b_n) = 2^{n-1} R_n^{-\frac{3}{8}} (\log R_n)^{\frac{1}{4}}$. However, this is a contradiction because strict convexity means $v(ta_n + (1-t)b_n) < tv(a_n) + (1-t)v(b_n) = 2^{n-1} R_n^{-\frac{3}{8}} (\log R_n)^{\frac{1}{4}}$ on R_n . Hence it follows that $\|v\|_{L^\infty} \lesssim R^{-\frac{3}{8}} (\log R)^{\frac{1}{4}}$. This proves (3.27), and plugging (3.27) into the inequality (3.23) and repeating the same argument removes the logarithm, and the Proposition 3.3 is proved.

Motivated from Proposition 3.3, we are going to make bootstrap assumptions

$$(3.30) \quad |u_{kx}(x, t)| \leq M\epsilon t^{-\frac{1}{8} - \frac{k}{4}} \langle x \rangle^{\frac{k}{4} - \frac{3}{8}}, \quad k = 0, 1, 2, 3,$$

where M is a large universal constant, independent of ϵ , to be chosen later, and $\epsilon \ll 1$ is small depending on M .

The overall process for nonlinear estimates is similar with that in [IKT19], sometimes being simpler. More precisely, the process may include energy estimates for both nonlinear and linearized equations, uniform Sobolev bounds for $L^{NL}u$ for some nonlinear operator L^{NL} , and finally the nonlinear dispersive bounds.

4. ENERGY ESTIMATES

The goal of this section is to establish the uniform $B_{2,\infty}^{-\frac{1}{2}}$ bound of u . Specifically, we have the following proposition:

Proposition 4.1. *Let u be a solution to (KW) with the smallness assumption of the initial data*

$$(4.1) \quad \|u_0\|_{\dot{B}_{2,\infty}^{-\frac{1}{2}}} + \|xu_0\|_{\dot{H}^{\frac{1}{2}}} \leq \epsilon \ll 1.$$

together with the bootstrap bound (3.30). Then one has

$$\|u(t)\|_{\dot{B}_{2,\infty}^{-\frac{1}{2}}} \lesssim \epsilon$$

for the time scale $|t| \ll_M \epsilon^{-\frac{5}{3}}$.

Proof. A direct energy estimate gives

$$\frac{d}{dt} \int (P_\lambda u)^2 = \frac{1}{2} \int P_\lambda u \cdot \partial_x P_\lambda (u^2) \lesssim \|P_\lambda u\|_{L^2} \lambda^{\frac{1}{2}} \| |D|^{\frac{1}{2}}(u^2) \|_{L^2}.$$

Also we may find the bound of $\| |D|^{\frac{1}{2}}(u^2) \|_{L^2}$ as follows: Interpolating the pointwise bounds $|u^2| \leq M^2 \epsilon^2 t^{-\frac{1}{4}} \langle x \rangle^{-\frac{3}{4}}$ and $|\partial_x(u^2)| \leq M^2 \epsilon^2 t^{-\frac{1}{2}} \langle x \rangle^{-\frac{1}{2}}$ gives $\| |D|^{\frac{1}{2}}(u^2) \| \lesssim M^2 \epsilon^2 t^{-\frac{3}{8}} \langle x \rangle^{-\frac{5}{8}}$, so that $\| |D|^{\frac{1}{2}}(u^2) \|_{L^2} \lesssim M^2 \epsilon^2 t^{-\frac{2}{5}}$. Hence it should be the case that

$$\|P_\lambda u\|_{L^2} \lesssim \|P_\lambda u_0\|_{L^2} + \lambda^{\frac{1}{2}} M^2 \epsilon^2 t^{\frac{3}{5}} \lesssim \epsilon \lambda^{\frac{1}{2}}$$

under the assumption of the time bound $M^2 \epsilon t^{\frac{3}{5}} \leq 1$.¹ Therefore, we get the desired bound

$$\|u(t)\|_{\dot{B}_{2,\infty}^{-\frac{1}{2}}} \lesssim \epsilon, \quad t \ll_M \epsilon^{-\frac{5}{3}}.$$

□

Remark 4.2. The reason why the direct energy estimate of the Besov bound for (KW) is possible is that it has a better spatial decay of the solution that it is directly square integrable. One may see that the above argument does not work for (5KdV) due to more derivatives in the nonlinearity and thus having worse spatial decay.

¹This is why the time scale for (KW) in Theorem 1.1 is restricted to $|t| \ll \epsilon^{-\frac{5}{3}}$.

5. BOUNDS FOR LINEARIZED EQUATION

Note that (KW) enjoys scaling symmetry $u(t, x) \mapsto \lambda^4 u(\lambda^5 t, \lambda x)$, which is critical in $\dot{H}^{-\frac{7}{2}}$, in the sense that the scaling preserves $\dot{H}^{-\frac{7}{2}}$ norm. The scaling vector field which generates the symmetry is given as $\mathcal{S} = 5t\partial_t + x\partial_x + 4$.

Define $\Lambda = \partial_x^{-1}\mathcal{S}$. Then by the commutator relation

$$[\mathcal{L}, \mathcal{S}] = 5\mathcal{L}, \quad [\mathcal{S}, \partial_x] = -\partial_x$$

one has

$$\mathcal{L}(\Lambda u) = \partial_x^{-1}(\mathcal{S} + 5)\mathcal{L}u = u\partial_x\Lambda u.$$

Hence, $w = \Lambda u$ solves the linearized equation

$$w_t - w_{5x} = uw_x.$$

However, the function $w = xu + u_{4x} + \frac{5}{2}u^2 + 3\partial_x^{-1}u$ contains the inverse derivative of u , which can cause problem if u is not mean-zero. Hence, we rather work with the function

$$w := L^{NL}u := xu + u_{4x} + \frac{5}{2}u^2,$$

which solves the inhomogeneous linearized equation

$$w_t + w_{5x} = uw_x + \frac{3}{2}u^2.$$

The main proposition of this section is as follows:

Proposition 5.1. *Let u be a solution to the fifth-order KdV equation (KW) which satisfies the smallness assumption 4.1 for the initial data, as well as the bootstrap assumption (3.30) for $t \ll \epsilon^{-\frac{5}{3}}$. Then the linearized equation*

$$(\text{LinKW}) \quad w_t - w_{5x} = uw_x.$$

is well-posed in $\dot{H}^{\frac{1}{2}}$ with uniform bounds

$$\|w(t)\|_{\dot{H}^{\frac{1}{2}}} \sim \|w(0)\|_{\dot{H}^{\frac{1}{2}}}, \quad t \ll_M \epsilon^{-\frac{5}{3}}.$$

Let $y = |D|^{\frac{1}{2}}w$, then y satisfies the equation

$$y_t - y_{5x} = -|D|^{\frac{1}{2}}(uH|D|^{\frac{1}{2}}y).$$

A direct calculation and frequency analysis as in [IKT19] gives

$$\frac{1}{2} \frac{d}{dt} \int y^2 = - \int |D|^{\frac{1}{2}}y \cdot uH|D|^{\frac{1}{2}}y = O(M\epsilon t^{-\frac{2}{5}}) \|y\|_{L^2}^2 - \int |D|^{\frac{1}{2}}y_{hi} \cdot u_{hi}H|D|^{\frac{1}{2}}y_{hi}$$

in the time scale $M^2\epsilon t^{\frac{3}{5}} \leq 1$. Here $y_{hi} := P_{hi}y_{hi}$ means a piece of y frequency-localized in the scale $|\xi| \gtrsim t^{-\frac{1}{5}}$.

Hence, we shall try a normal form analysis to cancel the high frequency term. For the equation (KW), It turns out that the normal form is given as $\frac{1}{5}B(\partial_x^{-1}u_{hi}, \partial_x^{-1}u_{hi})$ where B is a symmetric

bilinear Fourier multiplier whose symbol is $\hat{B}(\xi, \eta) = \frac{1}{\xi^2 + \xi\eta + \eta^2}$. Hence the energy correction term is given as $E'[y] = \frac{1}{5} \int |D|^{-\frac{1}{2}} y_{hi} B(\partial_x^{-1} u_{hi}, H|D|^{-\frac{1}{2}} y_{hi})$. Now a direct calculation gives²

$$\begin{aligned}
& \frac{d}{dt} E'[y] \\
&= D_1 + D_2 + D_3 + \int |D|^{-\frac{1}{2}} \partial_x^5 y_{hi} B(\partial_x^{-1} u_{hi}, H|D|^{-\frac{1}{2}} y_{hi}) \\
&+ \int |D|^{-\frac{1}{2}} y_{hi} B(\partial_x^4 u_{hi}, H|D|^{-\frac{1}{2}} y_{hi}) + \int |D|^{-\frac{1}{2}} y_{hi} B(\partial_x^{-1} u_{hi}, H|D|^{-\frac{1}{2}} \partial_x^5 y_{hi}) \\
&= D_1 + D_2 + D_3 - \int |D|^{-\frac{1}{2}} y_{hi} \partial_x^5 B(\partial_x^{-1} u_{hi}, H|D|^{-\frac{1}{2}} y_{hi}) \\
&+ \int |D|^{-\frac{1}{2}} y_{hi} B(\partial_x^4 u_{hi}, H|D|^{-\frac{1}{2}} y_{hi}) + \int |D|^{-\frac{1}{2}} y_{hi} B(\partial_x^4 u_{hi}, H|D|^{-\frac{1}{2}} \partial_x^5 y_{hi}) \\
&= D_1 + D_2 + D_3 + \int |D|^{-\frac{1}{2}} y_{hi} \partial_x (u_{hi} |D|^{\frac{1}{2}} y_{hi}) = D_1 + D_2 + D_3 + \int H|D|^{\frac{1}{2}} y_{hi} \cdot u_{hi} |D|^{\frac{1}{2}} y_{hi}
\end{aligned}$$

where the penultimate equality follows from the symbol calculation

$$\frac{(\xi + \eta)^5 - \xi^5 - \eta^5}{\xi^2 + \xi\eta + \eta^2} = 5\xi\eta(\xi + \eta).$$

Hence the last integral cancels out the "Bad" term of $\frac{1}{2} \frac{d}{dt} \int y^2$, and the other terms (D_1, D_2, D_3) are perturbative, as demonstrated below:

- (a) D_1 arises from the time derivative of the frequency scale truncation, i.e. $\partial_t P_{hi}$. Namely, $\partial_t(1 - \varphi(t^{\frac{1}{5}}\xi)) = -\frac{1}{5}t^{-\frac{4}{5}}\xi\varphi'(t^{\frac{1}{5}}\xi) = -t^{-1}\psi(t^{\frac{1}{5}}\xi)$ where $\psi(\eta) = \frac{1}{5}\eta\varphi'(\eta)$ is the Littlewood-Paley projection on the frequency scale ~ 1 . Thus the bound for D_1 is given as

$$\begin{aligned}
& \left| \int |D|^{-\frac{1}{2}} (\partial_t P_{hi}) y \cdot B(\partial_x^{-1} u_{hi}, H|D|^{-\frac{1}{2}} y_{hi}) \right| \lesssim t^{-\frac{9}{10}} \|y\|_{L^2} \|B(\partial_x^{-1} u_{hi}, H|D|^{-\frac{1}{2}} y_{hi})\|_{L^2} \\
& \lesssim t^{-\frac{9}{10}} \|y\|_{L^2} t^{\frac{2}{5}} \|\partial_x^{-1} u_{hi}\|_{L^\infty} \|H|D|^{-\frac{1}{2}} y_{hi}\|_{L^2} \lesssim M\epsilon t^{-\frac{2}{5}} \|y\|_{L^2}^2.
\end{aligned}$$

The rest terms, i.e. $\int |D|^{-\frac{1}{2}} y_{hi} \cdot B(\partial_x^{-1} (\partial_t P_{hi}) u, H|D|^{-\frac{1}{2}} y_{hi})$ and $\int |D|^{-\frac{1}{2}} y_{hi} \cdot B(\partial_x^{-1} u_{hi}, H|D|^{-\frac{1}{2}} (\partial_t P_{hi}) y)$ can be estimated in a similar fashion.

- (b) D_2 consists of the nonlinearity of u from (KW):

$$\left| \int |D|^{-\frac{1}{2}} y_{hi} B((u^2)_{hi}, H|D|^{-\frac{1}{2}} y_{hi}) \right| \lesssim t^{\frac{1}{10}} \|y\|_{L^2} t^{\frac{2}{5}} \|u^2\|_{L^\infty} t^{\frac{1}{10}} \|y\|_{L^2} \leq M\epsilon t^{-\frac{2}{5}}.$$

- (c) D_3 consists of the linearized counterpart of D_2 :

$$\begin{aligned}
& \left| \int (u H|D|^{\frac{1}{2}} y)_{hi} \cdot B(\partial_x^{-1} u_{hi}, H|D|^{-\frac{1}{2}} y_{hi}) \right| \\
&= \left| \int H y \cdot |D|^{\frac{1}{2}} \left(u P_{hi} B(\partial_x^{-1} u_{hi}, H|D|^{-\frac{1}{2}} y_{hi}) \right) \right| \lesssim \|y\|_{L^2} \| |D|^{\frac{1}{2}} \left(u P_{hi} B(\partial_x^{-1} u_{hi}, H|D|^{-\frac{1}{2}} y_{hi}) \right) \|_{L^2}^2.
\end{aligned}$$

²Here I intentionally dropped any insignificant constant multiplications.

Now the fractional chain rule gives:

$$\begin{aligned}
& \| |D|^{\frac{1}{2}} \left(u P_{hi} B(\partial_x^{-1} u_{hi}, H |D|^{-\frac{1}{2}} y_{hi}) \right) \|_{L^2} \\
& \lesssim \| |D|^{\frac{1}{2}} u \|_{L^\infty} \| P_{hi} B(\partial_x^{-1} u_{hi}, H |D|^{-\frac{1}{2}} y_{hi}) \|_{L^2} + \| u \|_{L^\infty} \| |D|^{\frac{1}{2}} P_{hi} B(\partial_x^{-1} u_{hi}, H |D|^{-\frac{1}{2}} y_{hi}) \|_{L^2} \\
& \lesssim M \epsilon t^{-\frac{3}{10}} \cdot t^{\frac{2}{5}} \| \partial_x^{-1} u_{hi} \|_{L^\infty} \| H |D|^{-\frac{1}{2}} y_{hi} \|_{L^2} + M \epsilon t^{-\frac{1}{5}} \cdot t^{\frac{3}{10}} \| \partial_x^{-1} u_{hi} \|_{L^\infty} \| H |D|^{-\frac{1}{2}} y_{hi} \|_{L^2} \\
& \lesssim M \epsilon t^{-\frac{2}{5}} \| y \|_{L^2}.
\end{aligned}$$

Hence we obtain the bound $M \epsilon t^{-\frac{2}{5}} \| y \|_{L^2}^2$. The other term $\int |D|^{-\frac{1}{2}} y_{hi} B(\partial_x^{-1} u_{hi}, H(u |D|^{\frac{1}{2}} y)_{hi})$ can be treated similarly after decomposing $(u |D|^{\frac{1}{2}} y)_{hi} = \partial_x(u H |D|^{-\frac{1}{2}} y)_{hi} - (u_x H |D|^{-\frac{1}{2}} y)_{hi}$. One may handle the former term by noting that the bilinear map $(u, v) \mapsto B(u, v_x)$ has symbol $\frac{i\eta}{\xi^2 + \xi\eta + \eta^2}$, so that $\| B(u, v_x) \|_{L^2} \lesssim t^{\frac{1}{5}} \| u \|_{L^\infty} \| v \|_{L^2}$, provided u and v are frequency supported at the scale $\gtrsim t^{-\frac{1}{5}}$.

Hence, we conclude that, if we let the corrected energy $E[y] = \frac{1}{2} \int y^2 + E'[y]$, then

$$\frac{d}{dt} E[y] \lesssim M \epsilon t^{-\frac{2}{5}} \| y \|_{L^2}^2 \sim M \epsilon t^{-\frac{2}{5}} E[y].$$

Hence Gronwall's inequality gives

$$E[y] \leq e^{CM \epsilon t^{\frac{3}{5}}} E[y_0] \lesssim E[y_0]$$

in the time scale $t \ll_M \epsilon^{-\frac{5}{3}}$.

Remark. The above analysis also works when the right hand side of (LinKW) is $u^k w_x$ with $k \geq 1$, with u_{hi} being replaced by $(u^k)_{hi}$. Namely, we may consider u^k as a single component, which makes no difference of the details. The only difference is the valid time scale of the analysis. For instance, if $k = 2$, which corresponds to the case of (5KdV) with quadratic terms gone, the time scale becomes ϵ^{-5} .

6. SOBOLEV BOUND OF $L^{NL}u$

In this section, we are going to prove the following proposition:

Proposition 6.1. *Let u be a solution to (KW) satisfying the smallness assumption (4.1) and the bootstrap assumption (3.30). Then $L^{NL}u := xu + u_{4x} + \frac{5}{2}u^2$ satisfies the bound*

$$\|L^{NL}u(t)\|_{\dot{H}^{\frac{1}{2}}} \lesssim \epsilon, \quad |t| \ll_M \epsilon^{-\frac{5}{3}}.$$

Unlike the [IKT19], we do not have to dive into a complicated normal form analysis, since we have a better bootstrap bound. Namely, we already proved that $\|u^2\|_{\dot{H}^{\frac{1}{2}}} \lesssim M^2\epsilon^2t^{-\frac{2}{5}}$ in the Section 4. Then it remains to see that the $H^{\frac{1}{2}}$ bound of the inhomogeneous term is perturbative enough that Proposition 5.1 is still valid to $L^{NL}u$.

Let $z := |D|^{\frac{1}{2}}L^{NL}u$. Then z solves the equation

$$z_t - z_{5x} = -|D|^{\frac{1}{2}}(uH|D|^{\frac{1}{2}}z + \frac{3}{2}u^2).$$

Hence one has:

$$\begin{aligned} \frac{d}{dt}E[z] &= -\frac{3}{2} \int z|D|^{\frac{1}{2}}(u^2) - \frac{3}{10} \int (u^2)_{hi}B(\partial_x^{-1}u_{hi}, H|D|^{-\frac{1}{2}}z_{hi}) \\ &\quad - \frac{3}{10} \int |D|^{-\frac{1}{2}}z_{hi}B(\partial_x^{-1}u_{hi}, H(u^2)_{hi}) + O(M\epsilon t^{-\frac{2}{5}})\|z\|_{L^2}^2, \end{aligned}$$

where $E[\cdot]$ is the corrected energy defined in the Section 5. Here most of the steps are identical to the Section 5, so it only remains to estimate the integrals including the inhomogeneous term. Namely,

$$\begin{aligned} \left| \int z|D|^{\frac{1}{2}}(u^2) \right| &\lesssim M^2\epsilon^2t^{-\frac{2}{5}}\|z\|_{L^2} \lesssim M^2\epsilon t^{-\frac{2}{5}}\|z\|_{L^2}^2 + M^2\epsilon^3t^{-\frac{2}{5}} \lesssim M^2\epsilon t^{-\frac{2}{5}}\|z\|_{L^2}^2 + \epsilon^2t^{-1}, \\ \left| \int (u^2)_{hi}B(\partial_x^{-1}u_{hi}, H|D|^{-\frac{1}{2}}z_{hi}) \right| \\ &\lesssim \|u^2\|_{\dot{H}^{\frac{1}{2}}}t^{\frac{2}{5}}\|\partial_x^{-1}u_{hi}\|_{L^\infty}\||D|^{-\frac{1}{2}}z_{hi}\|_{L^2} \lesssim M^3\epsilon^3t^{\frac{1}{10}}\|z\|_{L^2} \lesssim M^2\epsilon t^{-\frac{2}{5}}\|z\|_{L^2}^2 + M^4\epsilon^5t^{\frac{3}{5}} \\ &\lesssim M^2\epsilon t^{-\frac{2}{5}}\|z\|_{L^2}^2 + \epsilon^2t^{-\frac{6}{5}}, \end{aligned}$$

and the rest terms can be estimated similarly.

To sum up, one obtains the bound

$$\frac{d}{dt}E[z] \lesssim M^2\epsilon t^{-\frac{2}{5}}\|z\|_{L^2}^2 + \epsilon^2t^{-1}.$$

Hence Grönwall's inequality gives

$$\|z\|_{L^2}^2 \lesssim \epsilon^2e^{CM^2\epsilon t^{\frac{3}{5}}} \lesssim \epsilon^2$$

in the time scale $|t| \ll \epsilon^{-\frac{5}{3}}$. This completes the proof of Proposition 6.1.

7. THE KLAINERMAN-SOBOLEV ESTIMATES

In this section, we finish the proof of our main theorem, Theorem 1.1. For the sake of simplicity, we are going to rescale the problem to $t = 1$. Namely, given the equation

$$xu + 5tu_{4x} + \frac{5}{2}tu^2 = f, \quad f = L^{NL}u$$

let $\tilde{u}(x) = t^{\frac{4}{5}}u(t, xt^{\frac{1}{5}})$, $\tilde{f}(x) = t^{\frac{3}{5}}f(t, xt^{\frac{1}{5}})$. Then a direct calculation gives that \tilde{u} and \tilde{f} solve the same equation with $t = 1$, so that

$$(7.1) \quad (x + 5\partial_x^4)\tilde{u} + \frac{5}{2}\tilde{u}^2 = \tilde{f}.$$

Such scaling also gives the new bounds as follows:

$$(7.2) \quad \|\tilde{u}\|_{\dot{B}_{2,\infty}^{-\frac{1}{2}}} \lesssim \tilde{\epsilon} := \epsilon t^{\frac{3}{5}} \ll 1, \quad |\partial_x^k \tilde{u}(x)| \lesssim M\tilde{\epsilon}\langle x \rangle^{-\frac{3}{8} + \frac{k}{4}}, \quad k = 0, 1, 2, 3,$$

and

$$(7.3) \quad \|\tilde{f}\|_{\dot{H}^{\frac{1}{2}}} \lesssim \tilde{\epsilon}.$$

For notational convenience, we are going to drop the tilde notation.

Now most of the procedures are the same with the linear analysis, so I am going to skip most of the details, except the necessary nonlinear analysis.

Lemma 7.1. *Under the assumptions of (7.1), (7.2), (7.3), we have*

$$(7.4) \quad \|f\|_{L^2(A_R)} \lesssim \epsilon R^{\frac{1}{2}}.$$

Proof. As in the proof of Lemma 3.4, we split u at the frequency cutoff R^{-1} and compute:

$$f = P_{>R^{-1}}f + P_{<R^{-1}}Lu + P_{R^{-1}}(u^2).³$$

The first two terms can be estimated in the same way as Lemma 3.4. For the last term, the pointwise bootstrap bound (7.2) gives

$$|u^2| \lesssim \epsilon R^{-\frac{3}{4}},$$

so that

$$\|u^2\|_{L^2(A_R)} \lesssim \epsilon R^{-\frac{1}{4}},$$

which is better than we needed. □

³Here I intentionally dropped any insignificant constant coefficients.

Lemma 7.2. *Under the assumptions of (7.1), (7.2), (7.3), we have*

$$(7.5) \quad \|\partial_x^k u\|_{L^2(A_R)} \lesssim \epsilon R^{\frac{1}{8} + \frac{k}{4}}.$$

Proof. The low frequency case $\lambda \leq R^{\frac{1}{4}}$ can be dealt with in the same way in the linear case, so we focus on the high frequency case, $\lambda > R^{\frac{1}{4}}$. Again we split as

$$Lu_\lambda = P_\lambda L^{NL}u + \lambda^{-1}u_\lambda - P_\lambda(u^2),$$

and the first two terms can be estimated in the same way as in the linear case, so we again focus on the nonlinear terms. Using (7.2) and Lemma 2.3 gives

$$\|P_\lambda(u^2)\|_{L^2(A_R)} \lesssim \epsilon R^{-\frac{1}{4}}$$

and

$$\|\partial_x P_\lambda(u^2)\|_{L^2(A_R)} \lesssim \epsilon R^{\frac{1}{4}},$$

so that

$$\|P_\lambda(u^2)\|_{L^2(A_R)} \lesssim \epsilon \lambda^{-\frac{1}{2}}.$$

Now the argument is completed as in the linear case. \square

Now we may localize the solution u to the dyadic regions $|x| \sim R$. Let $v := \chi_R u$, then v solves the equation

$$(7.6) \quad (x + 5\partial_x^4)v + \frac{5}{2}uv = g,$$

with the bounds

$$(7.7) \quad \|\partial_x^k v\|_{L^2(A_R)} \lesssim \epsilon R^{\frac{1}{8} + \frac{k}{4}}, \quad k = 0, 1, 2, 3, 4,$$

and

$$(7.8) \quad \|f\|_{\dot{H}^{\frac{1}{2}}} \lesssim \epsilon, \quad \|g\|_{L^2(A_R)} \lesssim \epsilon R^{\frac{1}{2}}.$$

As in the linear case, we consider three different regions; self-similar, elliptic, and hyperbolic.

The self-similar region is entirely similar to the linear case, so we omit the details.

The elliptic region is also almost similar, but there is a subtle difference due to the presence of nonlinearity. More precisely, we again split $g = g_{lo} + g_{hi}$, where the frequency scale is $R^{\frac{1}{4}}$. Moreover, letting $v_1 := v - x^{-1}g_{lo}$ and

$$\tilde{L}^{NL}v_1 = (x + 5\partial_x^4)v_1 + uv_1 = g_{hi} - 5\partial_x^4(x^{-1}g_{lo}) := g_1.$$

The estimation of g_1 is the same as in the linear case, yielding the $\epsilon R^{-\frac{1}{8}}$ bound. The only difference is the estimation of integrals. To be more specific, we start from the equality

$$\int_{\mathbb{R}} v_1 \tilde{L}^{NL}v_1 dx = \int_{\mathbb{R}} g_1 v_1 dx$$

and integrating by parts gives

$$\int_{\mathbb{R}} x|v_1|^2 dx + 5 \int_{\mathbb{R}} |\partial_x^2 v_1|^2 dx = \int_{\mathbb{R}} g_1 v_1 dx + \int_{\mathbb{R}} uv_1^2 dx$$

The first integral on the right is routine, so we focus on the second integral including nonlinearity.

A direct calculation gives

$$\int_{\mathbb{R}} uv_1^2 dx = O(M\epsilon R^{-\frac{3}{8}})\|v_1\|_{L^2}^2,$$

so this bound can be absorbed into the left hand side provided R is large enough. Thus one may conclude

$$\|v_1\|_{L^2} \lesssim \epsilon R^{-\frac{9}{8}}, \quad \|\partial_x^2 v_1\|_{L^2} \lesssim \epsilon R^{-\frac{5}{8}}.$$

Now the same process as in the linear case gives the extra bounds

$$\|\partial_x^4 v_1\|_{L^2} \lesssim \epsilon R^{-\frac{1}{8}}, \quad \|\partial_x v_1\|_{L^2} \lesssim \epsilon R^{-\frac{7}{8}}, \quad \|\partial_x^3 v_1\|_{L^2} \lesssim \epsilon R^{-\frac{3}{8}}.$$

Hence Gagliardo-Nirenberg inequality gives

$$|\partial_x^k v_1| \lesssim \epsilon R^{-1+\frac{k}{4}}, \quad k = 0, 1, 2, 3.$$

The pointwise bound of $x^{-1}g_{lo}$ can be obtained in the same way as in the linear analysis, so we omit the details.

Lastly, let us turn to the hyperbolic region. Again we do a spatial energy estimate to obtain

$$\frac{d}{dx}(-x|v|^2 + 5|v_{2x}|^2 - 10v_x v_{3x} + 2g_{lo}v) = -|v|^2 + 5uvv_x - 2v_x g_{hi} + 2g_{lo,x}v.$$

Now integrating both sides and using Hölder's inequality gives

$$\begin{aligned} -x|v|^2 + 5|v_{2x}|^2 - 10v_x v_{3x} &\lesssim \|v_x\|_{L^2} \|g_{hi}\|_{L^2} + \|g_{lo,x}\|_{L^2} \|v\|_{L^2} + \sup_{x \in A_R^H} |g_{lo}v| + \int |uvv_x| dx \\ &\lesssim \epsilon^2 R^{\frac{1}{4}} + M\epsilon^3 R^{\frac{1}{8}} \lesssim \epsilon^2 R^{\frac{1}{4}}. \end{aligned}$$

A similar calculation gives

$$-x|v_x|^2 + 5|v_{3x}|^2 + 2xvv_{2x} \lesssim \epsilon^2 R^{\frac{3}{4}},$$

and the rest is now the same with the linear case. This completes the proof of the main theorem.

8. APPENDIX: THE CASE OF MODIFIED KAWAHARA EQUATION

In this appendix, we sketch the proof of the case $m = 2$ of the Theorem 1.1, namely (mKW). First, one may easily see that the argument in the Section 4 does work to prove Proposition 4.1 in the case of (mKW), under the time bound assumption $M^{\frac{3}{2}}\epsilon t^{\frac{1}{5}} \leq 1$. Here is the step where the time scale restriction $|t| \ll \epsilon^{-5}$ is needed.

Secondly, as mentioned in the remark at the end of the Section 5, one may prove the same bound from Proposition 5.1 with the same argument. Here we have a subtly different setting, and I state here for convenience.

The equation (mKW) enjoys the scaling symmetry $u(t, x) \mapsto \lambda^2 u(\lambda^5 t, \lambda x)$, so the scaling vector field is given as $\mathcal{S} = 5t\partial_t + x\partial_x + 2$, and one may see that $\Lambda u := \partial_x^{-1}\mathcal{S}u = xu + 5tu_{4x} + \frac{5}{3}tu^3 + \partial_x^{-1}u$ satisfies the linearized equation

$$w_t - w_{5x} = u^2 w_x.$$

Now the rest of the arguments are the same as in the Section 5.

The Sections 6 and 7 are again similar. In this case $L^{NL}u = xu + 5tu_{4x} + \frac{5}{3}tu^3 = \Lambda u - \partial_x^{-1}u$ solves the inhomogeneous equation

$$z_t - z_{5x} = u^2 z_x + \frac{2}{3}u^3,$$

and the energy estimate of the inhomogeneous equation is now routine.

When it comes to the Section 7, one may rescale the equation

$$xu + 5tu_{4x} + \frac{5}{3}tu^3 = f, \quad f = L^{NL}u$$

by letting $\tilde{u}(x) = t^{\frac{2}{5}}u(t, xt^{\frac{1}{5}})$ and $\tilde{f}(x) = t^{\frac{1}{5}}f(t, xt^{\frac{1}{5}})$. Also the localized equation becomes

$$(x + 5\partial_x^4)v + \frac{5}{3}u^2v = g.$$

In these settings, all the same steps work properly.

REFERENCES

- [Kaw72] Takuji Kawahara. “Oscillatory solitary waves in dispersive media”. In: *Journal of the physical society of Japan* 33.1 (1972), pp. 260–264.
- [DZ93] Percy Deift and Xin Zhou. “A steepest descent method for oscillatory Riemann–Hilbert problems. Asymptotics for the MKdV equation”. In: *Annals of Mathematics* (1993), pp. 295–368.
- [KPV94a] Carlos E Kenig, Gustavo Ponce, and Luis Vega. “Higher-order nonlinear dispersive equations”. In: *Proceedings of the American Mathematical Society* 122.1 (1994), pp. 157–166.
- [KPV94b] Carlos E Kenig, Gustavo Ponce, and Luis Vega. “On the hierarchy of the generalized KdV equations”. In: *Singular limits of dispersive waves*. Springer, 1994, pp. 347–356.
- [HN99] Nakao Hayashi and Pavel I Naumkin. “Large time behavior of solutions for the modified Korteweg–de Vries equation”. In: *International Mathematics Research Notices* 1999.8 (1999), pp. 395–418.
- [FO00] Kazumasa Fujiwara and Tohru Ozawa. “Remarks on bilinear estimates in the Sobolev spaces (Harmonic)”. In: *Math. Z* 235 (2000), pp. 173–194.
- [HN01] Nakao Hayashi and Pavel Naumkin. “On the modified Korteweg–de Vries equation”. In: *Mathematical Physics, Analysis and Geometry* 4.3 (2001), pp. 197–227.
- [Caz03] Thierry Cazenave. *Semilinear Schrödinger Equations*. Vol. 10. American Mathematical Soc., 2003.
- [HN06] Nakao Hayashi and Pavel I Naumkin. *Final state problem for Korteweg–de Vries type equations*. 2006.
- [Tao06] Terence Tao. *Nonlinear dispersive equations: local and global analysis*. 106. American Mathematical Soc., 2006.
- [Fuj07] Kazuyuki Fujii. “A Higher Order Non–Linear Differential Equation and a Generalization of the Airy Function”. In: *arXiv preprint arXiv:0712.2481* (2007).
- [Kwo08] Soonsik Kwon. “On the fifth-order KdV equation: local well-posedness and lack of uniform continuity of the solution map”. In: *Journal of Differential Equations* 245.9 (2008), pp. 2627–2659.
- [KP12] Carlos E Kenig and Didier Pilod. “Well-posedness for the fifth-order KdV equation in the energy space”. In: *arXiv preprint arXiv:1205.0169* (2012).
- [KAT13] TAKAMORI KATO. “GLOBAL WELL-POSEDNESS FOR THE KAWAHARA EQUATION WITH LOW REGULARITY.” In: *Communications on Pure & Applied Analysis* 12.3 (2013).
- [Dur14] Samuel O Durugo. “Higher-order airy functions of the first kind and spectral properties of the massless relativistic quartic anharmonic oscillator”. PhD thesis. Loughborough University, 2014.
- [IT15] Mihaela Ifrim and Daniel Tataru. “Global bounds for the cubic nonlinear Schrödinger equation (NLS) in one space dimension”. In: *Nonlinearity* 28.8 (2015), p. 2661.
- [Har16] Benjamin Harrop-Griffiths. “Long time behavior of solutions to the mKdV”. In: *Communications in Partial Differential Equations* 41.2 (2016), pp. 282–317.

- [IT17] Mihaela Ifrim and Daniel Tataru. “Well-posedness and dispersive decay of small data solutions for the Benjamin-Ono equation”. In: *arXiv preprint arXiv:1701.08476* (2017).
- [Oka+18] Mamoru Okamoto et al. “Long-time behavior of solutions to the fifth-order modified KdV-type equation”. In: *Advances in Differential Equations* 23.9/10 (2018), pp. 751–792.
- [BKV19] Bjoern Bringmann, Rowan Killip, and Monica Visan. “Global well-posedness for the fifth-order KdV equation in $H^{-1}(\mathbb{R})$ ”. In: *arXiv preprint arXiv:1912.01536* (2019).
- [IKT19] Mihaela Ifrim, Herbert Koch, and Daniel Tataru. “Dispersive decay of small data solutions for the KdV equation”. In: *arXiv preprint arXiv:1901.05934* (2019).
- [Oka19] Mamoru Okamoto. “Asymptotic behavior of solutions to a higher-order KdV-type equation with critical nonlinearity.” In: *Evolution Equations & Control Theory* 8.3 (2019).