

A BINARY QUADRATIC FORM APPROACH TO THE EQUATION $X^2 + (2k - 1)^Y = k^Z$

MAOHUA LE AND ANITHA SRINIVASAN

ABSTRACT. A conjecture of N. Terai states that for any integer $k > 1$, the equation $x^2 + (2k - 1)^y = k^z$ has only one solution, namely, $(x, y, z) = (k - 1, 1, 2)$. Using the theory of representations of integers by binary quadratic forms, we present methods to verify this conjecture in the case when $k \equiv 0 \pmod{4}$, with $2k - 1$ a prime power, and in the case when k is any odd integer. We present several values of k for which our method shows that the conjecture is true, while existing methods do not apply. However for $k \equiv 0 \pmod{4}$ our method does not always work.

AMS Subject Classification: 11D09, 11R29, 11E16. **Keywords:** binary quadratic forms, exponential diophantine equation, generalized Ramanujan-Nagell equation.

1. INTRODUCTION

N. Terai in [7] stated the following conjecture for a particular case of the generalized Ramanujan Nagell equation.

Conjecture 1.1. For any integer $k > 1$, the diophantine equation

$$(1.1) \quad X^2 + (2k - 1)^Y = k^Z, X, Y, Z \in \mathbb{N}$$

has only the solution $(X, Y, Z) = (k - 1, 1, 2)$.

The above conjecture has been proved only in some special cases. In [2], M.-J. Deng, J. Guo and A.-J. Xu verified Conjecture 1.1 when $k \equiv 3 \pmod{4}$ with $3 \leq k \leq 499$. Our method can verify the conjecture for any odd k , computational power being the only limitation. We also note here that to the best of our knowledge, there are no results that deal with the case when $k \equiv 1 \pmod{4}$.

N. Terai [7] dealt with $k \leq 30$ using methods that did not apply to $k = 12, 24$. Subsequently M. A. Bennett and N. Billerey [1] used the modular approach to solve these two cases. Our method proves the case $k = 12$ in a very straightforward manner. Interestingly, it does not work for $k = 24$.

Recently Mutlu, Le and Soydan [5] showed under the assumption of the GRH that if $k \equiv 0 \pmod{4}$ and $30 < k < 724$ with $2k - 1$ a

prime power, then Conjecture 1.1 is true. For several of these values our method applies easily. Moreover our method can be used for any (larger) value of k (such as $k = 1336$ given in section 3) satisfying these same conditions. The method for these values of k is based on the following theorem.

Theorem 1.2. *Let $k \equiv 0 \pmod{4}$ and $2k - 1$ be a prime power. Let $k - 1 = ab^2$ where a is square-free. If there exists a divisor d of ab such that z is even for all solutions (x, y, z) of*

$$(1.2) \quad d^2x^2 + (2k - 1)y^2 = k^z, x, y, z \in \mathbb{Z}, \gcd(x, y) = 1, z > 0,$$

then Conjecture 1.1 is true.

Remark 1.3. Observe that there are infinitely many solutions (x, y, z) to (1.2). This is because $((k - 1)/d, 1, 2)$ is a solution and we will see in Section 2 that all solutions can be divided into classes. Moreover it is easy to check whether all z are even as the least value z_1 of z in each class is bounded. Furthermore, all other solutions (x, y, z) satisfy $z = z_1 + z'_1 t$, where z'_1 is a fixed integer (for a given class) that divides the class number, and hence can be calculated.

The method for odd k ($\lambda = 1$) uses Lemma 2.4(4). If a solution to the equation $X^2 + (2k - 1)Y = k^Z$ exists, then in the case when Y is odd we have a solution to $x'^2 + (2k - 1)y'^2 = k^{z'}$ and in the case when Y is even, we have a solution to $x'^2 + (2k - 1)^2y'^2 = k^{z'}$. In both cases if the solution is the least solution in a class, then by Lemma 2.4, parts (3) and (4) and Lemma 2.2(4), z' divides the class number and we may verify for each divisor of the class number, whether for a solution (x', y') (if it exists) y' is a power of $(2k - 1)$. When the solution is not the least solution, by (2.7) we have a least solution, (x'_1, y'_1, z'_1) such that $y'_1 | (2k - 1)^Y$ and we may check whether this condition holds for each least solution. Hence we have the following result.

Theorem 1.4. *For a fixed odd positive integer k Conjecture 1.1 can be verified in a finite number of steps.*

The outline of the paper is as follows. In the next section we quote results from [3] and [4] that relate the solutions of two related equations that our theorems are based on. In Section 3 we prove our main theorems, and in the last section we present examples to illustrate the applications of these theorems, proving Conjecture 1.1 for several values of k , some that have resisted existing algorithms.

2. THE SOLUTIONS OF TWO RELATED EQUATIONS

Let D, D_1, D_2 be fixed odd positive integers with $D = D_1 D_2 > 3$ and $\gcd(D_1, D_2) = 1$. Let k be a fixed positive integer with $\gcd(D, k) = 1$. Let $w(k)$ be the number of distinct prime divisors of k and let

$$\lambda = \begin{cases} 2 & \text{if } k \text{ is even} \\ 1 & \text{if } k \text{ is odd.} \end{cases}$$

In this section we state results that are either given, or follow immediately from the results and proof methods in [3] and [4]. These results classify and connect the solutions of the equations

$$(2.1) \quad x'^2 + Dy'^2 = \lambda^2 k^{z'}, x', y', z' \in \mathbb{Z}, \gcd(x', y') = 1, z' > 0$$

and

$$(2.2) \quad D_1 x^2 + D_2 y^2 = k^z, x, y, z \in \mathbb{Z}, \gcd(x, y) = 1, z > 0.$$

Lemma 2.1. *If the congruence*

$$(2.3) \quad L^2 \equiv -D \pmod{\lambda^2 k}, 0 < L < \frac{\lambda^2 k}{2}, \gcd\left(\lambda^2 k, 2L, \frac{L^2 + D}{\lambda^2 k}\right) = 1$$

has solutions L , then it has exactly $r = 2^{w(k)-1}$ solutions.

We denote the r solutions above as $L_i, i = 1 \cdots L_r$.

Lemma 2.2. *A necessary and sufficient condition for (2.1) to have solutions is that (2.3) has solutions. Moreover the solutions of (2.1) satisfy the following.*

- (1) *For any fixed solution (x', y', z') of (2.1) there exists a unique solution L_j ($1 \leq j \leq r$) of (2.3) such that*

$$(2.4) \quad \frac{x'}{y'} \equiv \lambda' L_j \pmod{\lambda k}, \lambda' \in \{1, -1\}.$$

- (2) *All solutions of (2.1) can be divided into r disjoint equivalence classes C'_j , $1 \leq j \leq r$, where each class C'_j is non-empty and consists of solutions (x', y', z') of (2.1) that satisfy (2.4).*

- (3) *For each C'_j ($1 \leq j \leq r$), there is a unique least positive solution (x'_j, y'_j, z'_j) such that $z'_k \leq z'$, where z' runs through all solutions (x', y', z') in C'_j . Moreover every solution (x', y', z') of C'_j satisfies $z' = z'_j t, t \in \mathbb{N}$ and*

$$(2.5) \quad \frac{x' + y' \sqrt{-D}}{\lambda} = \pm \left(\frac{x'_j \pm y'_j \sqrt{-D}}{\lambda} \right)^t$$

for some positive integer t .

- (4) *The least solution (x'_j, y'_j, z'_j) of each class satisfies $z'_j | h(-4D)$, where $h(-4D)$ is the class number of binary quadratic primitive forms of discriminant $-4D$.*

Remark 2.3. Using Lemma 2.1 we may find all r primitive binary quadratic forms, F_j , of discriminant $-4D$ that represent $\lambda^2 k$ [6, Solution of Problem 2, page 120]. The order of F_j is precisely z'_j and it then follows that there is a primitive representation (α, β) such that $\alpha^2 + D\beta^2 = \lambda^2 k^{z'_j}$.

Lemma 2.4. *A necessary and sufficient condition for (2.2) to have solutions is that (2.1) has solutions. Moreover the solutions of (2.2) satisfy the following.*

- (1) *For any fixed solution (x, y, z) of (2.2) there exists a unique solution L_j ($1 \leq j \leq r$) of (2.3) such that*

$$(2.6) \quad \frac{D_1 x}{y} \equiv \lambda' L_j \pmod{\lambda k}, \lambda' \in \{1, -1\}.$$

- (2) *All solutions (x, y, z) of (2.2) can be divided into r equivalence classes, C_j , where each class C_j consists of solutions (x, y, z) of (2.2) that satisfy (2.6).*
- (3) *Some classes of (2.2) may be empty. In every fixed non-empty class C_j ($1 \leq j \leq r$), there is a unique least positive solution (x_{1j}, y_{1j}, z_{1j}) such that $z_{1j} \leq z$, where z runs through all solutions (x, y, z) of C_j . Moreover every solution (x, y, z) of C_j satisfies*

$$z = z_j + z'_j t,$$

for some integer $t \geq 0$, where (x'_j, y'_j, z'_j) is the least solution of the class C'_j of (2.1).

- (4) *If k is odd and C_j is non-empty, then $z'_j = 2z_j$ and every solution (x, y, z) of C_j satisfies*

$$(2.7) \quad x\sqrt{D_1} + y\sqrt{-D_2} = \pm(x_j\sqrt{D_1} \pm y_j\sqrt{-D_2})^{2t+1}$$

for some positive integer t .

Remark 2.5. Lemmas 2.3 and 2.4 show that each class of solutions of (2.2) corresponds to a class of solutions of (2.1). The correspondence is achieved via the integers L_j . These integers are distinct for different classes C'_j and the same for C_j and C'_j .

3. PROOFS OF MAIN THEOREMS

Lemma 3.1. *Let X, Y, Z be positive integers such that $X^2 + (2k-1)^Y = k^Z$. If $k-1 = ab^2$ with a square-free, then $X \equiv 0 \pmod{ab}$.*

Proof. The lemma follows on looking at the given equation modulo $k - 1$ which gives $X^2 \equiv 0 \pmod{ab^2}$. \square

Lemma 3.2. *If X, Y, Z is a solution of $X^2 + (2k - 1)^Y = k^Z$ with $k \equiv 0 \pmod{4}$ and $2k - 1$ a prime power, then Y and Z are odd.*

Proof. If $k \equiv 0 \pmod{4}$ it is easy to see (considering the equation mod 4) that Y is odd. In the case when $2k - 1$ is a prime power it is shown in [2, Lemma 2.6] that Z is odd. \square

The following lemma is key to our results and follows directly from part 3 of Lemma 2.4.

Lemma 3.3. *Let $k \equiv 0 \pmod{4}$ be a fixed positive integer. Then if the least solution (x_j, y_j, z_j) in each (non-empty) class of (2.2) and the least solution (x'_j, y'_j, z'_j) of the corresponding class of (2.1) both satisfy z_j and z'_j are even, then z is even for all solutions (x, y, z) of (2.2).*

3.1. Proof of Theorem 1.2. Let (X, Y, Z) be a solution of (1.1), where $k \equiv 0 \pmod{4}$ and $2k - 1$ is a prime power. Assume that d is a divisor of ab , that satisfies the hypothesis in the theorem. Then from Lemma 3.1 we have $d|X$ and hence $(X/d, (2k - 1)^{(Y-1)/2}, Z)$ is a solution of (1.2). From Lemma 3.2 we have that Z is odd, which contradicts the assumption of the theorem and hence Conjecture 1.1 is true. \square

3.2. Proof of Theorem 1.4. Suppose that $X^2 + (2k - 1)^Y = k^Z$ has a solution. If Y is odd this gives a solution to $x^2 + (2k - 1)y^2 = k^z$, namely $x = X$ and $y = (2k - 1)^{(Y-1)/2}$ and $z = Z$. Similarly if Y is even, we have a solution to $x^2 + (2k - 1)^2 y^2 = k^z$. Let $D = -4(2k - 1)$ if Y is odd and $D = -4(2k - 1)^2$ if Y is even. Now, if the solution (x, y, z) is the least solution, (x'_1, y'_1, z'_1) in a class, then by Lemma 2.2 part 4, $z'_1 | h(D)$. Note that we can find all these least solutions and therefore it may be verified if any y'_1 is a power of $2k - 1$. If the solution is not the least solution then by (2.7) we have

$$x + y\sqrt{-D} = \pm \left(x'_1 \pm y'_1 \sqrt{-D} \right)^l$$

for some integer l and least solution (x'_1, y'_1, z'_1) . In this case we note that each prime divisor of y'_1 must divide $2k - 1$. It is straightforward to verify whether this is the case for each least solution (x'_1, y'_1, z'_1) . \square

Next we present an algorithm for the implementation of Theorem 1.4. Recall that in this case we have $\lambda = 1$.

3.3. An algorithm for odd k . *Step 1.* Evaluate the class numbers $h = h(-4(2k - 1))$ and $h' = h(-4(2k - 1)^2)$.

Step 2. Determine all divisors of h and h' .

Step 3. Find all solutions (x, y, d) of (2.1) (with $D = 2k - 1$) where d divides h and $y > 1$.

Step 4. Find all solutions (x', y', d') of (2.1) (with $D = (2k - 1)^2$) where d' divides h' .

Step 5. If each y and y' has a prime divisor that does not divide $2k - 1$, Conjecture 1.1 is true.

4. EXAMPLES

Example 1: $k = 12$. We will show that $d = k - 1$ satisfies the assumption of Theorem 1.4 and hence Conjecture 1.1 is true for $k = 12$.

We have $2k - 1 = 23$ and $k - 1 = 11$. As $w(k) = 2$, equation (2.3) has two solutions, namely $L_1 = 7$ and $L_2 = 23$. Let $D = (2k - 1)(k - 1)^2$. The binary quadratic forms of discriminant $-4D$ representing $4k$ (see Remark 2.3) are $f_1 = (48, 14, 59)$ and $f_2 = (48, 46, 69)$ with orders $z'_1 = 6$ and $z'_2 = 18$. The least solutions of (2.1) are $(432, 65, 6)$ and $(9706934521, 66396265, 18)$ corresponding to $L_1 = 7$ and $L_2 = 23$ respectively (from (2.4)). By Lemma 2.4 part (3), a least solution (x, y, z) of (2.2) satisfies $z \leq 18$. Checking for all these values of z we get solutions $(1, 1, 2)$ and $(8870441293, 4089910213, 20)$ for equation (2.2). From (2.6) (with $D_1 = (k - 1)^2$) we see that both these solutions correspond to $L_2 = 23$ and hence are in the same class. It follows that the only least solution of (2.2) is $(1, 1, 2)$ corresponding to the class of $(9706934521, 66396265, 18)$ of (2.1). Hence by Lemma 3.3 all solutions of (1.2), for $d = k - 1$ satisfy z even and thus the hypothesis of Theorem 1.4 holds for $d = k - 1$ and hence Conjecture 1.1 is true.

Example 2: $k = 24$. As mentioned earlier, our method fails to work in this case. This is because there is no divisor d satisfying the hypothesis of Theorem 1.2.

We have $k - 1 = 23$. It is easy to see that $(7, 1, 1)$ is a least solution of (2.1). It follows that (2.1) represents all powers of k and thus the hypothesis of Theorem 1.2 does not hold for $d = 1$. For $d = 23$ we have $L_1 = 1$ and $L_2 = 31$ are the two solutions of (2.3). From Remark 2.3 the forms (of discriminant $-4(2k - 1)(k - 1)^2$) representing $4k$ are $(96, 62, 269)$ and $(96, 2, 259)$ that have orders 6 and 15 respectively. Therefore equation (2.1) with these exponents for k has the least solutions $(1727, 175, 6)$ and $(9795246139, 278142485, 15)$. Using (2.4) we see that the first solution corresponds to L_2 and the second to L_1 . Now we turn to the solutions of (2.2). By Lemma 2.4 part (3),

a least solution (x, y, z) here satisfies $z \leq 15$. Checking for all these values we get only one class of solutions for (2.2), represented by the least solution $(1, 1, 2)$. From (2.6) we identify that this class corresponds to $L_1 = 1$. The corresponding class of (2.1) has least solution $(9795246139, 278142485, 15)$. As the z value is odd here, the hypothesis of Lemma 3.3 is not satisfied and hence the hypothesis of Theorem 1.2 does not hold for $d = 23$. As there is no divisor d satisfying the hypothesis of Theorem 1.2, we are unable to prove the conjecture for $k = 24$.

Example 3: $k = 1336$. Here $2k - 1 = 2671$ is prime and $k - 1 = 3 \cdot 5 \cdot 89$. We take $d = 89$ in Theorem 1.2. As $w(k) = 2$ there are two solutions of (2.3) namely, $L_1 = 913$ and $L_2 = 1425$. The forms of discriminant $-4(2k - 1)89^2$ representing $4k$ are $(5344, 1826, 4115)$ and $(5344, 2850, 4339)$, both having order 92. By Remark 2.3 we find the two least solutions of (2.1) as $(a_1, b_1, 92)$ and $(a_2, b_2, 92)$ belonging to classes corresponding to L_1 and L_2 (we have omitted writing the values of a_i and b_i as they have several digits). Next we look for solutions of (2.2) for z up to 92 as by Lemma 2.4 part (3) we know that a least solution (x, y, z) satisfies $z \leq 92$. We obtain two classes of solutions of (2.2) namely $(x_1, y_1, 2)$ and $(x_2, y_2, 74)$ corresponding to L_2 and L_1 respectively. As the classes corresponding to L_1 and L_2 for (2.1) have least solutions (x', y', z') with z' even (namely 92), by Lemma 3.3 all solutions (x, y, z) of (1.2) with $d = 89$ have z even. Hence the hypothesis of Theorem 1.2 holds for $d = 89$ and therefore Conjecture 1.1 is true.

Example 4: $k = 96$. In this case our method fails. Here we have $k - 1 = 5 \cdot 19$. For $d = 1, 5, 19$ and $k - 1$, we see that there are coprime solutions (x, y) to $d^2x^2 + (2k - 1)y^2 = k^z$ with z odd. Thus Theorem 1.2 does not apply and we are unable to settle the conjecture in this case.

Next we consider odd values of k to illustrate Theorem 1.4 using the algorithm presented at the end of Section 3.

Example 5: $k = 531$. We have for the class numbers $h = h(-4(2k - 1)) = 26$ and $h' = h(-4(2k - 1)^2) = 530$. We verify for each divisor d of h whether (2.1) with $D = 2k - 1$ has a solution and obtain two classes of solutions $(a_1, b_1, 2)$ and $(a_2, b_2, 26)$. We observe that there are no solutions (x', y', d') of (2.1) with $D = (2k - 1)^2$ for any divisor d' of h' .

We see that b_1 and b_2 are co-prime with $2k - 1$. Hence Conjecture 1.1 holds.

Example 6: $k = 733$. Here $2k - 1 = 5 \cdot 23$. We have $h = 104$ and $h' = 584$. In the tables below we give the required primes p in Step 5 of the algorithm for odd k in Section 3.3 to conclude that Conjecture 1.1

d	4	8	26	52	104
p	61	7	79	61	7

TABLE 1. Solutions $x^2 + (2k - 1)y^2 = k^d$ with $p|y$

d'	146	292	584
p	29	29	7

TABLE 2. Solutions $x'^2 + (2k - 1)^2y'^2 = k^{d'}$ with $p|y'$

holds. More specifically, for each solution (x, y, d) (and (x', y', d')) of (2.1) the prime p satisfies $p|y$ ($p|y'$). It should be noted that $p \nmid (2k - 1)$.

REFERENCES

- [1] M. Bennett, N. Billerey, *Sums of two S -units via Frey-Hellegouarch curves*, Math. Comp. 305 (2017) 1375–1401.
- [2] M.-J. Deng, J. Guo, A.-J. Xu, *A note on the Diophantine equation $x^2 + (2c - 1)^m = c^n$* , Bull. Aust. Math. Soc. **98** (2018) 188–195.
- [3] Yasutsugu Fujita, Maohua Le, *Some exponential diophantine equations II: The equation $x^2 - Dy^2 = k^z$ for even k* , Mathematica Slovaca, **72**(2), 2022, 341–354.
- [4] Maohua Le, *Some exponential Diophantine equations I: The equation $D_1x^2 - D_2y^2 = \lambda k^z$* , J. Number Theory, bf 55, (1995), 209–221.
- [5] Elif Kızıldere Mutlu, Maohua Le, Gökhan Soydan, *A modular approach to the generalized Ramanujan-Nagell equation*, Indagationes Mathematicae, **33**(5), September 2022, 992–1000.
- [6] P. Ribenboim, *My Numbers, My Friends, Popular Lectures on Number Theory*, Springer-Verlag, 2000.
- [7] N. Terai, *A note on the diophantine equation $x^2 + q^m = c^n$* , Bull. Aust. Math. Soc. **90** (2014) 20–27.

INSTITUTE OF MATHEMATICS, LINGNAN NORMAL COLLEGE, ZHANGJIANG, GUANGDONG, 524048, CHINA

Email address: lemaohua2008@163.com

DEPARTAMENTO DE MÉTODOS CUANTITATIVOS, UNIVERSIDAD PONTIFICIA DE COMILLAS (ICADE), C/ ALBERTO AGUILERA, 23 - 28015 MADRID

Email address: asrinivasan@icade.comillas.edu