

ON Ext^1 FOR DRINFELD MODULES

DAWID EDMUND KĘDZIERSKI, PIOTR KRASOŃ

ABSTRACT. Let $A = \mathbb{F}_q[t]$ be the polynomial ring over a finite field \mathbb{F}_q and let ϕ and ψ be A -Drinfeld modules. In this paper we consider the group $\text{Ext}^1(\phi, \psi)$ with the Baer addition. We show that if $\text{rank}\phi > \text{rank}\psi$ then $\text{Ext}^1(\phi, \psi)$ has the structure of a \mathfrak{t} -module. We give complete formulas describing this structure. We also establish duality between Ext groups for \mathfrak{t} -modules and the corresponding adjoint \mathfrak{t}^σ -modules. Finally, we prove the existence of "Hom-Ext" six-term exact sequences for \mathfrak{t} -modules. As the category of \mathfrak{t} -modules is only additive (not abelian) this result is nontrivial.

1. INTRODUCTION

Drinfeld modules since their discovery [D74] gained a lot of attention in arithmetic algebraic geometry because of their numerous applications e.g. in class field theory, in Langlands conjectures, theory of automorphic forms or Diophantine geometry (see [G96], [Th04] or [F13]). There is a deep analogy between the theory of Drinfeld modules over function fields and theory of elliptic curves over number fields or more generally between \mathfrak{t} -modules and abelian varieties over number fields (see [BP09]). This analogy justifies various attempts to use the theory of Drinfeld modules in cryptology (see [G03], [N19]).

There are also some significant differences. For example the Mordell-Weil groups of \mathfrak{t} -modules are not finitely generated [P95]. The category of Drinfeld modules as well as the category of \mathfrak{t} -modules are not semisimple i.e. there are non-trivial extensions.

Remark 1.1. Since in this paper we consider Ext groups in several different categories, we indicate the appropriate category by a subscript τ , $\mathbb{F}_q[t]$ or σ .

Let $\text{Ext}_\tau^1(B, A)$ be the Bauer group of extensions of \mathfrak{t} -modules i.e. the group of exact sequences

$$(1.1) \quad 0 \rightarrow A \rightarrow M \rightarrow B \rightarrow 0$$

Date: October 18, 2022.

2020 Mathematics Subject Classification. 11G09, 18G50.

Key words and phrases. \mathfrak{t} -modules, Drinfeld modules, group of extensions, Hom-Ext exact sequence, biderivations.

with the usual addition known from homological algebra (cf. [M95]).

Let ϕ and ψ be Drinfeld modules. In this paper we study the group $\text{Ext}_\tau^1(\phi, \psi)$. It turns out that sometimes we can endow the extension group $\text{Ext}_\tau^1(A, B)$, for certain specific \mathfrak{t} -modules, with a \mathfrak{t} -module structure. In our study we apply the method (used in [PR03]) of expressing the elements of $\text{Ext}_\tau^1(\phi, \psi)$ by certain classes of biderivations. This idea was originally introduced by G. Hochschild in his thesis for the study of extensions (the first Hochschild cohomology groups) of associative algebras. It is worth mentioning that the concept of describing an extension space as a cokernel of a line map was used by C.M. Ringel for modules over path algebra of a quiver or more generally for K-species [R76], [R98]. The content of the paper is as follows. In Section 2 we recall basic definitions and properties of Drinfeld modules and \mathfrak{t} -modules. In Section 3 we identify the $\text{Ext}_\tau^1(\phi, \psi)$ for Drinfeld modules satisfying $\text{rk}\phi > \text{rk}\psi$ with some \mathbb{F}_q -subspace of the ring of skew polynomials $K\{\tau\}$. In Section 4 we compute an example which show how to endow $\text{Ext}_\tau^1(\phi, \psi)$ for the case $\text{rk}\phi > \text{rk}\psi$ with the structure of a \mathfrak{t} -module. Section 5 is devoted to the proof of general case i.e. that $\text{Ext}_\tau^1(\phi, \psi)$ is in the considered case a \mathfrak{t} -module. The exact calculations are long therefore we present them in the Appendix A. Notice that although the presented formulas are rather complicated it is easy to program them. Some consequences of the exact formulas are given then in the Appendix B. In Section 6 we generalize the results from Section 5 to certain \mathfrak{t} -modules that are products of Drinfeld modules. A brief discussion of the remaining cases i.e. $\text{rk}\phi \leq \text{rk}\psi$ is included in Section 7 where also, for a perfect field K , a duality between \mathfrak{t} -modules and \mathfrak{t}^σ -modules is established. The Duality Theorem (Theorem 7.2) allows one to describe $\text{Ext}_\tau^1(\phi, \psi)$ for $\text{rk}\phi < \text{rk}\psi$ as $\mathbb{F}_q(t)$ -module. In the Section 8 we prove directly the existence of pullbacks and pushouts in the category of \mathfrak{t} -modules. We also prove existence of the six-term " Hom – Ext " exact sequences, where the last map is surjective. As the category of \mathfrak{t} -modules is not an abelian category these are nontrivial results (cf. Example 8.1). We conclude with some consequences of the six-term exact sequences (Proposition 8.3 and Theorem 8.4).

2. PRELIMINARIES

Let p be a rational prime and $A = \mathbb{F}_q[t]$ the polynomial ring over the finite field with $q = p^m$ elements and let K be a field of characteristic p . We say that K is an A -**field** when we fix an F_q -linear homomorphism $\iota : A \rightarrow K$ with $\theta := \iota(t)$. Denote by $K\{\tau\}$ the ring of twisted polynomials

in τ with coefficients in K i.e. satisfying the additional relation $\tau x = x^q \tau$ for $x \in K$, see [G96].

In [A86] G. Anderson developed a generalization of the notion of a Drinfeld module called a t -module.

Definition 2.1. A d -dimensional \mathbf{t} -module over an A -field K is an \mathbb{F}_q -linear ring homomorphism

$$(2.1) \quad \Phi : \mathbb{F}_q[t] \rightarrow \mathrm{Mat}_d(K\{\tau\}),$$

such that $\Phi(t)$, as a polynomial in τ with coefficients in $\mathrm{Mat}_d(K)$ is of the following form

$$(2.2) \quad \Phi(t) = (\theta I_d + N)\tau^0 + M_1\tau^1 + \cdots + M_r\tau^r,$$

where I_d is the identity matrix and N is a nilpotent matrix.

In general, a \mathbf{t} -module over K is an algebraic group E defined over K and isomorphic over K to G_a^d together with a choice of \mathbb{F}_q -linear endomorphism $t : E \rightarrow E$ such that $d(t - \theta)^n \mathrm{Lie}(E) = 0$ for n sufficiently large. The choice of an isomorphism $E \cong G_a^d$ is equivalent to the choice of Φ . In order to indicate this choice of coordinates, we write $E = (G_a^d, \Phi)$.

Remark 2.1. Since the map Φ is a homomorphism of \mathbb{F}_q -algebras a \mathbf{t} -module is completely determined by the polynomial Φ_t i.e. by the image of t . The degree m of Φ_t is called the **rank** of Φ and it is denoted as $\mathrm{rk} \Phi$.

Definition 2.2. Let Φ and Ψ be two \mathbf{t} -modules of dimension d and e , respectively. A morphism $f : \Phi \rightarrow \Psi$ of \mathbf{t} -modules over K is a matrix $f \in \mathrm{Mat}_{e \times d}(K\{\tau\})$ such that

$$f\Psi(t) = \Phi(t)f.$$

In general, if $E = (G_a^d, \Phi)$ and $F = (G_a^e, \Psi)$, then a \mathbf{t} -module morphism $f : F \rightarrow E$ is a morphism of commutative algebraic groups $f : G_a^e \rightarrow G_a^d$ over K commuting with the action of A i.e:

$$f\Psi(t) = \Phi(t)f.$$

We also consider the **zero \mathbf{t} -module** of the form $0 : \mathbb{F}_q[t] \rightarrow 0$. Then the category of \mathbf{t} -modules becomes an $\mathbb{F}_q[t]$ -linear additive category and the notion of a short exact sequence (1.1) makes sense.

Dimension one \mathbf{t} -modules are called **Drinfeld** modules and the Drinfeld module $C : \mathbb{F}_q[t] \rightarrow K\{\tau\}$, given by the formula $C(t) = \theta + \tau$ is called the **Carlitz module**.

Every \mathfrak{t} -module $E = (G_a^d, \Phi)$ induces an $F_q[t]$ -module structure on K^d , where multiplication by t is given by evaluation of Φ_t , i.e.

$$t * x = \Phi_t(x) \quad \text{for } x \in K^d.$$

This $\mathbb{F}_q[t]$ -module is called the Mordell-Weil group of Φ and it is denoted as $\Phi(K^d)$. Similarly, each morphism $f : E = (G_a^d, \Phi) \rightarrow F = (G_a^e, \Psi)$ of \mathfrak{t} -modules induces a morphism $\Phi(f) : \Phi(K^d) \rightarrow \Psi(K^e)$ of $F_q[t]$ -modules. Then $\Phi(-)$ is a covariant inclusion functor from the category of \mathfrak{t} -modules to the category of $\mathbb{F}_q[t]$ -modules. In the Example 8.1 we will see that this functor is not full.

Recall [PR03], that each extension of an abelian \mathfrak{t} -module $\Phi : \mathbb{F}_q[t] \rightarrow \text{Mat}_d(K\{\tau\})$ by $\Psi : \mathbb{F}_q[t] \rightarrow \text{Mat}_e(K\{\tau\})$ can be determined by an \mathbb{F}_q -linear map $\delta : \mathbb{F}_q[t] \rightarrow \text{Mat}_{e \times d}(K\{\tau\})$ such that

$$(2.3) \quad \delta(ab) = \Psi(a)\delta(b) + \delta(a)\Phi(b) \quad \text{for all } a, b \in \mathbb{F}_q[t].$$

Such maps are called biderivations, and we will denote the \mathbb{F}_q -vector space of all biderivations by $\text{Der}(\Phi, \Psi)$. In the sequel we will, as usual, denote $\Phi_a := \Phi(a)$, $\Psi_a := \Psi(a)$ etc.. It is easy to check that the biderivation δ is uniquely determined, by the value $\delta(t) \in \text{Mat}_{e \times d}(K\{\tau\})$. Then the map $\delta \mapsto \delta(t)$ induces the \mathbb{F}_q -linear isomorphism of the vector spaces $\text{Der}(\Phi, \Psi)$ and $\text{Mat}_{e \times d}(K\{\tau\})$. Let $\delta^{(-)} : \text{Mat}_{e \times d}(K\{\tau\}) \rightarrow \text{Der}(\Phi, \Psi)$ be an \mathbb{F}_q -linear map defined by the following formula:

$$\delta^{(U)}(a) = U\Phi_a - \Psi_a U \quad \text{for all } a \in \mathbb{F}_q[t] \quad \text{and } U \in \text{Mat}_{e \times d}(K\{\tau\}).$$

The image of the map $\delta^{(-)}$ is denoted by $\text{Der}_{in}(\Phi, \Psi)$, and the elements of $\text{Der}_{in}(\Phi, \Psi)$ are called inner biderivations. In addition, both $\text{Ext}_\tau^1(\Phi, \Psi)$ and $\text{Der}(\Phi, \Psi)/\text{Der}_{in}(\Phi, \Psi)$ have $\mathbb{F}_q[t]$ -module structures. Then from [PR03, Lemma 2.1] there is an $\mathbb{F}_q[t]$ -module isomorphism

$$(2.4) \quad \text{Ext}_\tau^1(\Phi, \Psi) \cong \text{coker} \delta^{(-)} = \text{Der}(\Phi, \Psi)/\text{Der}_{in}(\Phi, \Psi).$$

Because the map $\delta \mapsto \delta(t)$ is an isomorphism, we will identify the coset $\delta + \text{Der}_{in}(\Phi, \Psi)$ with the coset $\delta(t) + \text{Der}_{in}(\Phi_t, \Psi_t)$ in $\text{Mat}_{e \times d}(K\{\tau\})$, where $\text{Der}_{in}(\Phi_t, \Psi_t) = \left\{ \delta(t) \mid \delta \in \text{Der}_{in}(\Phi, \Psi) \right\}$.

Remark 2.2. In what follows we adopt the following notation: $c^{(i)} := c^{q^i}$ is the evaluation of the Frobenius twist τ^i on an element $c \in K$. Moreover $c^{(0)} = c$.

3. EXTENSION OF DRINFELDS MODULES WITH T-MODULES STRUCTURES

In this section we study extensions of Drinfeld modules using biderivations. Our aim is to equip the extension space $\text{Ext}_\tau^1(\phi, \psi)$ with the natural

\mathbf{t} -module structure. This is generalization of M.A Papanikolas and N. Ramachandran results concerning the case where ψ is the Carlitz module. Further, we will give some basic properties of these \mathbf{t} -modules.

Lemma 3.1. *Let ϕ and ψ be Drinfeld modules, such that $\text{rk } \phi > \text{rk } \psi$. Then there is an \mathbb{F}_q -linear isomorphism*

$$\text{Ext}_\tau^1(\phi, \psi) \cong K\{\tau\}_{<\text{rk } \phi} := \left\{ w \in K\{\tau\} \mid \deg_\tau w < \text{rk } \phi \right\}$$

Proof. We will identify the biderivation δ with the polynomial $w(\tau) = \delta(t) \in K\{\tau\}$. Let $\phi_t = \theta + \sum_{i=1}^n a_i \tau^i$ and $\psi_t = \theta + \sum_{j=1}^m b_j \tau^j$, then $n > m$. From isomorphism (2.4) it suffices to show that $\text{Der}(\phi, \psi)/\text{Der}_{in}(\phi, \psi) \cong K\{\tau\}_{<\text{rk } \phi}$. To prove this, we consider generators of the space $\text{Der}_{in}(\phi, \psi)$ of the form $\delta^{(c\tau^k)}$ where $k = 0, 1, 2, \dots$ and $c \in K$. We have:

$$\begin{aligned} \delta^{(c\tau^k)}(t) &= c\tau^k \phi_t - \psi_t c\tau^k = c\tau^k \left(\theta + \sum_{i=1}^n a_i \tau^i \right) - \left(\theta + \sum_{j=1}^m b_j \tau^j \right) c\tau^k \\ &= c\theta^{(k)} \tau^k + \sum_{i=1}^n ca_i^{(k)} \tau^{i+k} - c\theta \tau^k - \sum_{j=1}^m c^{(k)} b_j \tau^{j+k} \\ (3.1) \quad &= c(\theta^{(k)} - \theta) \tau^k + \sum_{j=1}^m \left(ca_j^{(k)} - c^{(k)} b_j \right) \tau^{j+k} + \sum_{j=m+1}^n ca_j^{(k)} \tau^{i+k} \\ &= c(\theta^{(k)} - \theta) \tau^k + \sum_{j=k+1}^{m+k} \left(ca_{j-k}^{(k)} - c^{(j-k)} b_{j-k} \right) \tau^j + \sum_{j=m+k+1}^{n+k} ca_{j-k}^{(k)} \tau^j. \end{aligned}$$

Because $\deg_\tau \delta^{(c\tau^k)}(t) = n + k$ for $c \neq 0$, then each non-zero inner biderivation is given by a polynomial with τ -degree greater or equal to n . Therefore, if $w, \widehat{w} \in K\{\tau\}$, $w \neq \widehat{w}$ are such that $\deg_\tau w < n$ and $\deg_\tau \widehat{w} < n$, then w and \widehat{w} represent different cosets in $\text{Der}(\phi, \psi)/\text{Der}_{in}(\phi, \psi)$. On the other hand given a polynomial $w(\tau) = \sum_{i=1}^{n+l} a_i \tau^i \in K\{\tau\}$ with $l > 0$, one can find c_l (cf. (3.1)) such that the polynomial $w(\tau) - \delta^{(c_l \tau^l)}$ has τ -degree at most $l - 1$ and represents the same class as w in $\text{Der}(\phi, \psi)/\text{Der}_{in}(\phi, \psi)$. By downward induction we obtain:

$$\widetilde{w}(\tau) = w(\tau) - \delta^{(c_l \tau^l)} - \dots - \delta^{(c_1 \tau^1)} - \delta^{(c_0)}$$

where $\deg_\tau \widetilde{w} < n$ and both w and \widetilde{w} represent the same coset.

Therefore the isomorphism $\delta \mapsto \delta(t)$ induces the isomorphism of \mathbb{F}_q -vector spaces $\text{Der}(\phi, \psi)/\text{Der}_{in}(\phi, \psi) \cong K\{\tau\}_{<\text{rk } \phi}$. \square

Corollary 3.2. *Let $\delta \in K\{\tau\}_{<\text{rk } \phi}$ then δ corresponds to the extension:*

$$(3.2) \quad 0 \rightarrow \psi \rightarrow \Gamma \rightarrow \phi \rightarrow 0$$

where Γ is given by the matrix $\Gamma_t = \begin{bmatrix} \phi_t & 0 \\ \delta & \psi_t \end{bmatrix}$. Moreover, this correspondence is compatible with the Baer sum of extensions i.e. if $\delta_1, \delta_2 \in K\{\tau\}_{<\text{rk } \phi}$ correspond to $\Gamma_t^1 = \begin{bmatrix} \phi_t & 0 \\ \delta_1 & \psi_t \end{bmatrix}$ and $\Gamma_t^2 = \begin{bmatrix} \phi_t & 0 \\ \delta_2 & \psi_t \end{bmatrix}$ respectively then the Baer sum of extensions Γ^1 and Γ^2 corresponds to Γ such that $\Gamma_t = \begin{bmatrix} \phi_t & 0 \\ \delta_1 + \delta_2 & \psi_t \end{bmatrix}$. \square

Remark 3.1. Notice that $\text{Ext}_\tau^1(\phi, \psi)$ is a reflexive bilinear space with the bilinear product $\langle \delta_1, \delta_2 \rangle = \sum_{i=0}^{n-1} a_i b_i$ where $\delta_1 = a_0 + \dots + a_{n-1} \tau^{n-1}$ and $\delta_2 = b_0 + \dots + b_{n-1} \tau^{n-1}$.

Let ϕ and ψ be Drinfeld modules and let $r := \text{rk } \phi - \text{rk } \psi > 0$. Recall from [PR03] that the structure of the $\mathbb{F}_q[t]$ -module on $\text{Der}(\phi, \psi)/\text{Der}_{in}(\phi, \psi)$ is defined by the following formula:

$$(3.3) \quad a * \left(\delta + \text{Der}_{in}(\phi, \psi) \right) := \psi_a \delta + \text{Der}_{in}(\phi, \psi)$$

for $a \in \mathbb{F}_q[t]$ and $\delta \in \text{Der}(\phi, \psi)$.

Remark 3.2. For simplicity, we omit the notation $+\text{Der}_{in}(\phi, \psi)$ when considering the coset $\delta + \text{Der}_{in}(\phi, \psi)$ in the space $\text{Der}(\phi, \psi)/\text{Der}_{in}(\phi, \psi)$.

4. AN EASY BUT ESSENTIAL EXAMPLE

In this section we give an example to illustrate how to determine the structure of a \mathfrak{t} -module on the space of extensions of Drinfeld modules.

Example 4.1. Let $\phi_t = \theta + \tau^3$ and $\psi_t = \theta + \tau^2$. By Lemma 3.1

$$\text{Ext}_\tau^1(\phi, \psi) \cong K\{\tau\}_{<3} = \{c_0 + c_1\tau + c_2\tau^2 \mid c_i \in K\}.$$

In order to equip $\text{Ext}_\tau^1(\theta + \tau^3, \theta + \tau^2)$ with the \mathfrak{t} -module structure we will transfer the structure of the $F_q[t]$ -module from $\text{Ext}_\tau^1(\theta + \tau^3, \theta + \tau^2)$ to the space $K\{\tau\}_{<3}$ via the isomorphism from Lemma 3.1. Since each element of $K\{\tau\}$ is of the form $\sum_{i=0}^3 c_i \tau^i$, it is sufficient to determine the value of multiplication by t on the generators $c_i \tau^i$ for $i = 0, 1, 2$, where $c_i \in K$.

From (3.3) the value $t * (c_0)$ can be computed in the following way:

$$t * (c_0) = \psi_t \cdot c_0 = (\theta + \tau^2) \cdot c_0 = \theta c_0 + c_0^{(2)} \tau^2$$

Similarly,

$$t * (c_1 \tau) = \psi_t \cdot c_1 \tau = (\theta + \tau^2) \cdot c_1 \tau = \theta c_1 \tau + c_1^{(2)} \tau^3.$$

Because the polynomial $\theta c_0\tau + c_1^{(2)}\tau^3$ does not belong to $K\{\tau\}_{<3}$, then like in the proof of the Lemma 3.1 we reduce the term $c_1^{(2)}\tau^3$ by the generator

$$\delta^{(c)}(t) = c\phi_t - \psi_t c = c(\theta + \tau^3) - (\theta + \tau^2)c = -c^{(2)}\tau^2 + c\tau^3$$

for $c = c_1^{(2)}$. Then

$$t * (c_1\tau) = \theta c_0\tau + c_1^{(2)}\tau^3 - \delta^{(c)} = \theta c_1\tau + c_1^{(4)}\tau^2.$$

Next,

$$t * (c_2\tau^2) = \psi_t \cdot c_2\tau^2 = (\theta + \tau^2) \cdot c_2\tau^2 = \theta c_2\tau^2 + c_2^{(2)}\tau^4.$$

The term $c_1^{(2)}\tau^4$ can be reduced by the generator

$$\begin{aligned} \delta^{(c\tau)}(t) &= c\tau\phi_t - \psi_t c\tau = c\tau(\theta + \tau^3) - (\theta + \tau^2)c\tau \\ &= -c(\theta^{(1)} - \theta)\tau - c^{(2)}\tau^3 + c\tau^4, \end{aligned}$$

for $c = c_2^{(2)}$. Therefore

$$t * (c_2\tau^2) = t * (c_2\tau^2) - \delta^{(c\tau)} = c_1^{(2)}(\theta - \theta^{(1)})\tau + \theta c_2\tau^2 + c_1^{(4)}\tau^3.$$

The term $c_1^{(4)}\tau^3$ can be reduced by the generator $\delta^{(c)}$ for $c = c_1^{(4)}$. Hence

$$\begin{aligned} t * (c_2\tau^2) &= t * (c_2\tau^2) - \delta^{(c_1^{(2)}\tau)} - \delta^{(c_1^{(4)})} \\ &= c_1^{(2)}(\theta - \theta^{(1)})\tau + (\theta c_2 + c_2^{(6)})\tau^2. \end{aligned}$$

Now, choose the basis $e_0 = 1, e_1 = \tau, e_2 = \tau^2$ in $K\{\tau\}_{<3}$. In this basis the value $t * (c_0)$ has the following coordinates

$$t * (c_0) = \theta c_0 + c_0^{(2)}\tau^2 = [\theta c_0, 0, c_0^{(2)}].$$

Notice that $c_0^{(2)}$ is the value of the polynomial τ^2 at c_0 . Let $\tau^2|_{c_0} := c_0^{(2)}$.

Then we can express $t * (c_0)$ in the following form

$$t * (c_0) = [\theta, 0, \tau^2]_{|c_0}.$$

In a similar way we obtain:

$$\begin{aligned} t * c_0 &= c_0\theta + c_0^{(2)}\tau^2 = [\theta, 0, \tau^2]_{|c_0} \\ t * c_1\tau &= c_1\theta\tau + c_1^{(4)}\tau^2 = [0, \theta, \tau^4]_{|c_1} \\ t * c_2\tau^2 &= c_2^2(\theta - \theta^{(1)})\tau + (c_2\theta + c_2^{(6)})\tau^2 = [0, (\theta - \theta^{(1)})\tau^2, \theta + \tau^6]_{|c_2} \end{aligned}$$

Then the multiplication by t on $K\{\tau\}$ can be expressed by the following matrix:

$$(4.1) \quad \Pi_t = \begin{bmatrix} \theta & 0 & 0 \\ 0 & \theta & \theta - \theta^{(1)} \\ \tau^2 & \tau^4 & \theta + \tau^6 \end{bmatrix}$$

This matrix induces the homomorphism of \mathbb{F}_q -algebras

$$\Pi : \mathbb{F}_q[t] \longrightarrow \text{Mat}_3(K\{\tau\}),$$

such that

$$\Pi_t = \theta I_3 + \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & (\theta - \theta^{(1)}) \\ 1 & 0 & 0 \end{bmatrix} \tau^2 + \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \tau^4 + \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \tau^6.$$

Therefore Π gives rise to a \mathfrak{t} -module structure on $\text{Ext}_\tau^1(\theta + \tau^3, \theta + \tau^2)$.

5. $\text{Ext}_\tau^1(\phi, \psi)$ FOR DRINFELD MODULES ϕ AND ψ WITH $\text{rk } \phi > \text{rk } \psi$

We have the following:

Proposition 5.1. *Let ϕ and ψ be Drinfeld modules such that $\text{rk } \phi > \text{rk } \psi$. Then $\text{Ext}_\tau^1(\phi, \psi)$ has a natural structure of a \mathfrak{t} -module. This is given by the map $\Pi : \mathbb{F}_q[t] \longrightarrow M_{\text{rk } \phi}(k\{\tau\})$, where*

$$(5.1) \quad \Pi_t = \left[\begin{array}{c|ccc} \theta & 0 & \dots & 0 \\ \delta_1 & & & \\ \vdots & & & \\ \delta_{n-1} & & & \Pi_t^0 \end{array} \right].$$

At this moment, we will focus on the proof that the \mathfrak{t} -module structure comes from the matrix (5.1). Next we will see that the matrix Π_t^0 gives rise to a \mathfrak{t} -module structure on $\text{Ext}_{0,\tau}^1(\phi, \psi)$ and the vector $[\delta_1, \dots, \delta_{n-1}]^t$ determines the extension of $\text{Ext}_{0,\tau}^1(\phi, \psi)$ with $\text{Ext}_\tau^1(\phi, \psi)$ as the middle term.

Proof. We will describe an algorithm that allows us to determine the structure of the \mathfrak{t} -module on $\text{Ext}_\tau^1(\phi, \psi)$. The explicit formulas for \mathfrak{t} -module structure on $\text{Ext}_\tau^1(\phi, \psi)$ are given in the Appendix A.

From the Lemma 3.1 we know that

$$\text{Ext}_\tau^1(\phi, \psi) \cong \left\{ c_0 + c_1\tau + \dots + c_{\text{rk } \phi}\tau^{\text{rk } \phi} \mid c_i \in K \right\}$$

as F_q -vector space. In order to equip $\text{Ext}_\tau^1(\phi, \psi)$ with the \mathfrak{t} -module structure we will transfer the structure of the $F_q[t]$ -module from $\text{Ext}_\tau^1(\phi, \psi) = \text{Der}(\phi, \psi)/\text{Der}_{\text{in}}(\phi, \psi)$ to the space $K\{\tau\}_{<\text{rk } \phi}$ via the isomorphism from Lemma 3.1. Since each element of $K\{\tau\}$ is of the form $\sum_{i=0}^{\text{rk } \phi - 1} c_i\tau^i$, it is sufficient to determine the value of multiplication by t on the generators $c_i\tau^i$ for $i = 0, 1, \dots, \text{rk } \phi - 1$, where $c_i \in K$.

Next we will choose the coordinate system $e_i = \tau^i$ for $i = 0, 1, 2, \dots, \text{rk } \phi - 1$ in $K\{\tau\}_{<\text{rk } \phi}$ and see that the matrix Π_t of the multiplication map $t*(-) : K\{\tau\}_{<\text{rk } \phi} \longrightarrow K\{\tau\}_{<\text{rk } \phi}$ gives rise to the \mathfrak{t} -module structure on $\text{Ext}_\tau^1(\phi, \psi)$. This structure is given by (5.1).

Assume that $\phi_t = \theta + \sum_{i=1}^n a_i \tau^i$, $\psi_t = \theta + \sum_{j=1}^m b_j \tau^j$ and $r := n - m > 0$.

Then from the formula (3.3) we see that

$$t * (c_i \tau^i) = \psi_t \cdot c_i \tau^i \quad \text{for } c_i \in K \quad \text{and } i = 0, 1, 2, \dots, \text{rk } \phi - 1.$$

Suppose that $i \in \{0, 1, \dots, r - 1\}$. Then

$$t * (c_i \tau^i) = \left(\theta + \sum_{j=1}^m b_j \tau^j \right) c_i \tau^i = c_i \theta \tau^i + \sum_{j=1}^m b_j c_i^{(j)} \tau^{j+i} \in K\{\tau\}_{<\text{rk } \phi}$$

Hence

$$\begin{aligned} t * (c_i \tau^i) &= \left[\overbrace{0, \dots, 0}^{i-1}, \theta, b_1 c_i^{(1)}, b_2 c_i^{(2)}, \dots, b_m c_i^{(m)}, 0, \dots, 0 \right] = \\ &= \left[\underbrace{0, \dots, 0}_{i-1}, \theta, b_1 \tau, b_2 \tau^2, \dots, b_m \tau^m, 0, \dots, 0 \right]_{|c_i} = \end{aligned}$$

Therefore the first r column of the matrix Π_t satisfy our claim.

Consider the case $i \in \{r, r+1, \dots, \text{rk } \phi - 1\}$. In this situation the element $\psi_t \cdot c_i \tau^i$ has τ -degree bigger than $\text{rk } \phi - 1$, so like in the proof of the Lemma 3.1 we reduce the monomial with the biggest τ -degree by the generator $\delta^{(c\tau^k)} \in \text{Der}_{in}(\phi, \psi)$ and continue this procedure until we get the reduced polynomial belonging to $K\{\tau\}_{<\text{rk } \phi}$.

Then we perform downward induction from $n+i-r$ to n and see that at each step after the reduction we obtain polynomials satisfying the following properties:

- (i) the term at τ^0 of the reduced polynomial is equal to zero,
- (ii) each coefficient at τ^l of the reduced polynomial can be written as the evaluation at c_i of a skew polynomial $w_l(\tau)$,
- (iii) if $l \neq i$, then the skew polynomials $w_l(\tau)$ from (ii) have no free term,
- (iv) the skew polynomial $w_i(\tau)$ has a free term equal to θ .

From (i) we see that the matrix Π_t is of the form as claimed. On the other hand (ii) implies that after the reduction procedure is completed we can present the coefficients in the chosen coordinate system $e_i = \tau^i$, as the values of the skew polynomials in c_i . Write the matrix Π_t in the following form

$$\Pi_t = (\theta \cdot I + N) \tau^0 + \sum_{i=1}^{\text{finite}} A_i \tau^i, \quad \text{where } I, N, A_i \in \text{Mat}_{\text{rk } \phi}(K).$$

The condition (iii) implies that the matrix $N = 0$ and the condition (iv) implies that I is identity matrix. Therefore Π_t yields the \mathfrak{t} -module structure on $\text{Ext}_{\tau}^1(\phi, \psi)$

Induction: At the start of the induction we consider the following polynomial:

$$(5.2) \quad t * (c_i \tau^i) = c_i \theta \tau^i + \sum_{j=1}^m b_j c_i^{(j)} \tau^{j+i},$$

that satisfies conditions (i) – (iv) in an obvious way.

Suppose we have made k reductions that satisfy conditions (i) – (iv). Therefore the value $t * (c_i \tau^i)$ after reductions can be write in the following form:

$$w_{i-r-k+1}(c_i) \tau^{i-r-k+1} + w_{i-r-k+2}(c_i) \tau^{i-r-k+2} + \dots + w_{m+i-k}(c_i) \tau^{m+i-k},$$

where the polynomials $w_l(\tau)$ satisfy conditions (i) – (iv). We reduce the term $w_{m+i-k}(c_i) \tau^{m+i-k}$ by the generator $\delta^{(c\tau^{i-r-k})}$ where $c = \frac{w_{m+i-k}(c_i)}{a_n^{(i-r-k)}}$. Because all non-zero terms of $\delta^{(c\tau^{i-r-k})}$ have τ -degree ≥ 1 , then (i) is obvious. Recall that from (3.1) the generator $\delta^{(c\tau^{i-r-k})}$ can be rewritten in the following form:

$$\delta^{(c\tau^{i-r-k})}(t) = \sum_{j=i-r-k}^{m+i-k-1} p_j(c_i) \tau^j + \underbrace{ca_n^{(i-r-k)}}_{=w_{m+i-k}(c_i)} \tau^{m+i-k},$$

where the polynomials $p_l(\tau) \in K\{\tau\}$ satisfy the conditions (i) – (iii). Therefore after the reduction by the generator $\delta^{(c\tau^{i-r-k})}$ we obtain the following form of $t * (c_i \tau^i)$:

$$-p_{i-r-k}(c_i) \tau^{i-r-k} + \sum_{j=i-r-k+1}^{m+i-k-1} (w_j(c_i) - p_j(c_i)) \tau^j.$$

We put $\widehat{w_{i-r-k}}(\tau) = -p_{i-r-k}(\tau)$, $\widehat{w}_j(\tau) = w_j(\tau) - p_j(\tau)$ for $j = i - r - k + 1, \dots, m + i - k - 1$. Because the polynomials $p_l(\tau)$ and $w_l(\tau)$ satisfy the conditions (i) – (iii) and $w_i(\tau)$ satisfy (iv), then the polynomials $\widehat{w}_l(\tau)$ also satisfy these four conditions. This completes the induction step, and thus we proved that $\text{Ext}_{\tau}^1(\phi, \psi)$ has a \mathfrak{t} -module structure. \square

For $A \in \text{Mat}_{n_1 \times n_2}(K\{\tau\})$ let $dA \in \text{Mat}_{n_1 \times n_2}(K)$ be the constant term of A viewed as a polynomial in τ . For \mathfrak{t} -modules Φ and Ψ let

$$(5.3) \quad \text{Der}_0(\Phi, \Psi) = \{\delta \in \text{Der}(\Phi, \Psi) \mid d\delta(t) = 0\}.$$

Following [PR03] define $\text{Ext}_{0,\tau}^1(\Phi, \Psi) := \text{Der}_0(\Phi, \Psi) / \text{Der}_0(\Phi, \Psi) \cap \text{Der}_{in}(\Phi, \Psi)$. We have the following:

Lemma 5.2. *Let ϕ and ψ be the Drinfeld modules, such that $r = \text{rk } \phi - \text{rk } \psi > 0$. Then*

(i) there exists an isomorphism of $\mathbb{F}_q[t]$ -modules between $\text{Ext}_{0,\tau}^1(\phi, \psi)$

$$\text{and } K\{\tau\}_{\langle 1, \text{rk } \phi \rangle} = \left\{ \sum_{i=1}^{\text{rk } \phi - 1} c_i \tau^i \mid c_i \in K \right\},$$

(ii) $K\{\tau\}_{\langle 1, \text{rk } \phi \rangle}$ is an $\mathbb{F}_q[t]$ -submodule of $K\{\tau\}_{< \text{rk } \phi}$,

(iii) $\text{Ext}_{0,\tau}^1(\phi, \psi)$ is an $\mathbb{F}_q[t]$ -submodule of $\text{Ext}_{\tau}^1(\phi, \psi)$,

(iv) $\text{Ext}_{0,\tau}^1(\phi, \psi)$ has natural structure of a \mathfrak{t} -module.

Proof. The proof of part (i) is similar to the proof of Lemma 3.1. Recall, that $\text{Ext}_{0,\tau}^1(\phi, \psi) = \text{Der}_0(\phi, \psi) / \text{Der}_{in}(\phi, \psi) \cap \text{Der}_0(\phi, \psi)$ and $\delta^{(c\tau^k)} \in \text{Der}_0(\phi, \psi)$ for all $k \geq 0$. Because $\deg_{\tau}(\delta_t^{c\tau^k}) = \text{rk } \phi + k$ each biderivation $\delta \in \text{Der}_0(\phi, \psi)$ can be reduced as in the proof of the Lemma 3.1. This implies (i).

The vector space $K\{\tau\}_{\langle 1, \text{rk } \phi \rangle}$ has the $\mathbb{F}_q[t]$ -module structure coming from the same formulas $t * (c_k \tau^k)$ for $k = 1, 2, \dots, n-1$ as in the case $K\{\tau\}_{< \text{rk } \phi}$. Therefore the standard inclusion $K\{\tau\}_{\langle 1, \text{rk } \phi \rangle} \hookrightarrow K\{\tau\}_{< \text{rk } \phi}$ is the inclusion of $\mathbb{F}_q[t]$ -modules. This establishes (ii).

(iii) follows from (i) and (ii).

Part (iv) follows from the proof of the Proposition 5.1, where the matrix Π_t^0 from (5.1) induces the homomorphism $\Pi^0 : \mathbb{F}_q[t] \rightarrow M_{n-1}(K\{\tau\})$ giving rise to the \mathfrak{t} -module structure on $\text{Ext}_{0,\tau}^1(\phi, \psi)$. \square

The following theorem is a specialization of [PR03, Lemma 2.2]. However, we are able to describe the maps in an explicit way.

Theorem 5.3. *Let ϕ and ψ be Drinfeld modules, such that $r = \text{rk } \phi - \text{rk } \psi > 0$. Then there is a short exact sequence of \mathfrak{t} -modules*

$$(5.4) \quad 0 \longrightarrow \text{Ext}_{0,\tau}^1(\phi, \psi) \longrightarrow \text{Ext}_{\tau}^1(\phi, \psi) \longrightarrow \mathbb{G}_a \longrightarrow 0.$$

Proof. Let Π_t^0 and Π_t be the matrices giving the \mathfrak{t} -module structures for $\text{Ext}_{0,\tau}^1(\phi, \psi)$ and $\text{Ext}_{\tau}^1(\phi, \psi)$, respectively. Then the matrix

$$\begin{bmatrix} \overline{0 \dots 0} \\ I_{\text{rk } \phi - 1} \end{bmatrix}$$

induces the inclusion of \mathfrak{t} -modules $i : \text{Ext}_{0,\tau}^1(\phi, \psi) \hookrightarrow \text{Ext}_{\tau}^1(\phi, \psi)$. It is easy to check that $\text{coker}(i)$ is the trivial \mathfrak{t} -module \mathbb{G}_a . \square

Remark 5.1. Notice that the biderivation corresponding to the extension (5.4) comes from the first column of Π_t .

$$6. \text{Ext}_{\tau}^1 \left(\prod_{i=1}^n \phi_i, \prod_{j=1}^m \psi_j \right)$$

We have the following generalization of the Proposition 5.1: .

Theorem 6.1. *Assume that ϕ_i, ψ_j for $i = 1, \dots, n, j = 1, \dots, m$ are Drinfeld modules, such that $\text{rk } \phi_i > \text{rk } \psi_j$ for all i, j . Then $\text{Ext}_\tau^1\left(\prod_{i=1}^n \phi_i, \prod_{j=1}^m \psi_j\right)$ has a natural structure of a \mathfrak{t} -module.*

Proof. Let $\phi_i = \theta + \sum_{k=1}^{n_i} a_{i,k} \tau^k$ for $i = 1, \dots, n$ and $\psi_j = \theta + \sum_{k=1}^{m_j} b_{j,k} \tau^k$ for $j = 1, \dots, m$. We put

$$N = \max\{n_i \mid i = 1, \dots, n\} \quad \text{and} \quad M = \max\{m_j \mid j = 1, \dots, m\}.$$

Denote by $E_{j \times i}$ the $m \times n$ elementary matrix with the only one non-zero element equal to 1 at $j \times i$ place. The elements $\delta^{(c\tau^r E_{j \times i})}$ generate $\text{Der}_{in}\left(\prod_{i=1}^n \phi_i, \prod_{j=1}^m \psi_j\right)$, where

$$\delta^{(c\tau^r E_{i \times j})}(t) = E_{i \times j} \cdot \underbrace{\left(c(\theta^{(r)} - \theta)\tau^r + \sum_{l=1}^N c a_{i,l}^{(r)} \tau^{r+l} - \sum_{l=1}^M c^{(l)} b_{j,l} \tau^{l+r} \right)}_{\in \text{Der}_{in}(\phi_i, \psi_j)}$$

Then

$$\text{Der}_{in}\left(\prod_{i=1}^n \phi_i, \prod_{j=1}^m \psi_j\right) = \begin{bmatrix} \text{Der}_{in}(\phi_1, \psi_1) & \cdots & \text{Der}_{in}(\phi_n, \psi_1) \\ \text{Der}_{in}(\phi_1, \psi_2) & \cdots & \text{Der}_{in}(\phi_n, \psi_2) \\ \vdots & \ddots & \vdots \\ \text{Der}_{in}(\phi_1, \psi_m) & \cdots & \text{Der}_{in}(\phi_n, \psi_m) \end{bmatrix},$$

and therefore

$$(6.1) \quad \text{Ext}_\tau^1\left(\prod_{i=1}^n \phi_i, \prod_{j=1}^m \psi_j\right) \cong \begin{bmatrix} \text{Ext}_\tau^1(\phi_1, \psi_1) & \cdots & \text{Ext}_\tau^1(\phi_n, \psi_1) \\ \text{Ext}_\tau^1(\phi_1, \psi_2) & \cdots & \text{Ext}_\tau^1(\phi_n, \psi_2) \\ \vdots & \ddots & \vdots \\ \text{Ext}_\tau^1(\phi_1, \psi_m) & \cdots & \text{Ext}_\tau^1(\phi_n, \psi_m) \end{bmatrix},$$

as \mathbb{F}_q -linear spaces.

Moreover, for $a \in \mathbb{F}_q[t]$ and a biderivation $\delta \in \text{Der}\left(\prod_{i=1}^n \phi_i, \prod_{j=1}^m \psi_j\right)$ we

have

$$\begin{aligned}
a * \delta &= \left(\prod_{j=1}^m \psi_j \right) (a) \cdot \delta = \\
&= \begin{bmatrix} \psi_1(a) & 0 & \cdots & 0 \\ 0 & \psi_2(a) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \psi_m(a) \end{bmatrix} \cdot \begin{bmatrix} \delta_{1 \times 1}(\tau) & \cdots & \delta_{1 \times n}(\tau) \\ \delta_{2 \times 1}(\tau) & \cdots & \delta_{2 \times n}(\tau) \\ \vdots & \ddots & \vdots \\ \delta_{m \times 1}(\tau) & \cdots & \delta_{m \times n}(\tau) \end{bmatrix} = \\
&= \begin{bmatrix} \psi_1(a) \cdot \delta_{1 \times 1}(\tau) & \psi_1(a) \cdot \delta_{1 \times 2}(\tau) & \cdots & \psi_1(a) \cdot \delta_{1 \times n}(\tau) \\ \psi_2(a) \cdot \delta_{2 \times 1}(\tau) & \psi_2(a) \cdot \delta_{2 \times 2}(\tau) & \cdots & \psi_2(a) \cdot \delta_{2 \times n}(\tau) \\ \vdots & \vdots & \ddots & \vdots \\ \psi_m(a) \cdot \delta_{m \times 1}(\tau) & \psi_m(a) \cdot \delta_{m \times 2}(\tau) & \cdots & \psi_m(a) \cdot \delta_{m \times n}(\tau) \end{bmatrix} = \\
&= \begin{bmatrix} a * \delta_{1 \times 1}(\tau) & a * \delta_{1 \times 2}(\tau) & \cdots & a * \delta_{1 \times n}(\tau) \\ a * \delta_{2 \times 1}(\tau) & a * \delta_{2 \times 2}(\tau) & \cdots & a * \delta_{2 \times n}(\tau) \\ \vdots & \vdots & \ddots & \vdots \\ a * \delta_{m \times 1}(\tau) & a * \delta_{m \times 2}(\tau) & \cdots & a * \delta_{m \times n}(\tau) \end{bmatrix}.
\end{aligned}$$

Hence the isomorphism (6.1) preserves the structure of $\mathbb{F}_q[t]$ -modules. Because $\text{rk } \phi_i > \text{rk } \psi_j$, then $\text{Ext}_\tau^1(\phi_i, \psi_j)$ has a structure of \mathfrak{t} -module. Let $\Pi_{i \times j} : \mathbb{F}_q[t] \rightarrow \text{Mat}_{n_i}(K)\{\tau\}$ be the map, which gives the structure of the \mathfrak{t} -module on $\text{Ext}_\tau^1(\phi_i, \psi_j)$ in the base τ^l $l = 0, 1, \dots, n_i - 1$, same as in the proof of the Proposition 5.1.

Let $e_{i \times j}^l$ denote the element $\tau^l \cdot E_{i \times j}$. In $\mathbb{F}_q[t]$ -module

$$\text{Ext}_\tau^1 \left(\prod_{i=1}^n \phi_i, \prod_{j=1}^m \psi_j \right) \cong \begin{bmatrix} \text{Ext}_\tau^1(\phi_1, \psi_1) & \cdots & \text{Ext}_\tau^1(\phi_n, \psi_1) \\ \text{Ext}_\tau^1(\phi_1, \psi_2) & \cdots & \text{Ext}_\tau^1(\phi_n, \psi_2) \\ \vdots & \ddots & \vdots \\ \text{Ext}_\tau^1(\phi_1, \psi_m) & \cdots & \text{Ext}_\tau^1(\phi_n, \psi_m) \end{bmatrix},$$

we choose the following coordinate system:

$$\begin{aligned}
&\left(e_{1 \times 1}^l \right)_{l=0}^{n_1-1}, \left(e_{2 \times 1}^l \right)_{l=0}^{n_1-1}, \dots, \left(e_{m \times 1}^l \right)_{l=0}^{n_1-1}, \\
&\left(e_{1 \times 2}^l \right)_{l=0}^{n_2-1}, \left(e_{2 \times 2}^l \right)_{l=0}^{n_2-1}, \dots, \left(e_{m \times 2}^l \right)_{l=0}^{n_2-1}, \\
&\vdots \\
&\left(e_{1 \times n}^l \right)_{l=0}^{n_1-1}, \left(e_{2 \times n}^l \right)_{l=0}^{n_1-1}, \dots, \left(e_{m \times n}^l \right)_{l=0}^{n_n-1}.
\end{aligned}$$

Then the \mathfrak{t} -module structure on $\text{Ext}_\tau^1\left(\prod_{i=1}^n \phi_i, \prod_{j=1}^m \psi_j\right)$ is given by the map $\Pi : \mathbb{F}_q[t] \rightarrow \text{Mat}_z(K)\{\tau\}$, defined by

$$\Pi(t) = \begin{bmatrix} \left[\Pi_{1 \times i}(t)\right]_{i=1}^n & 0 & \cdots & 0 \\ 0 & \left[\Pi_{2 \times i}(t)\right]_{i=1}^n & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \left[\Pi_{m \times i}(t)\right]_{i=1}^n \end{bmatrix},$$

where $\left[\Pi_{j \times i}(t)\right]_{i=1}^n$ is the diagonal matrix with the elements $\Pi_{j \times 1}(t), \Pi_{j \times 2}(t), \dots, \Pi_{j \times n}(t)$ on the diagonal. \square

Corollary 6.2. *Assume that ϕ_i, ψ_j for $i = 1, \dots, n, j = 1, \dots, m$ are Drinfeld modules, such that $\text{rk } \phi_i > \text{rk } \psi_j$ for all i, j . Then there is a short exact sequence of \mathfrak{t} -modules*

$$0 \longrightarrow \text{Ext}_{0,\tau}\left(\prod_{i=1}^n \phi_i, \prod_{j=1}^m \psi_j\right) \longrightarrow \text{Ext}_\tau^1\left(\prod_{i=1}^n \phi_i, \prod_{j=1}^m \psi_j\right) \longrightarrow \mathbb{G}_a^{n \cdot m} \longrightarrow 0.$$

7. $\text{Ext}_\tau^1(\phi, \psi)$ FOR DRINFELD MODULES ϕ AND ψ WITH $\text{rk } \phi \leq \text{rk } \psi$

7.1. case $\text{rk } \phi = \text{rk } \psi$. This case is difficult to handle. We can express the generators $\delta^{(c\tau^k)}$, where $c \in K, k = 0, 1, \dots$, of the space $\text{Der}_{in}(\phi, \psi)$ in the following way:

$$\begin{aligned} \delta^{(c\tau^k)}(t) &= c\tau^k \phi_t - \psi_t c\tau^k = c\tau^k \left(\theta + \sum_{i=1}^n a_i \tau^i \right) - \left(\theta + \sum_{j=1}^n b_j \tau^j \right) c\tau^k \\ (7.1) \quad &= c(\theta^{(k)} - \theta)\tau^k + \sum_{j=k+1}^{n+k} \left(c a_{j-k}^{(k)} - c^{(j-k)} b_{j-k} \right) \tau^j. \end{aligned}$$

Notice that we cannot claim that the degree of the polynomial in the variable τ has degree $n + k$ which was important for the identification of the biderivation $c_j \tau^j$ with the polynomial $w(\tau) \in K\{\tau\}$. However excluding finite number of c 's in every degree k one can express a biderivation $c_j \tau^j$ as a polynomial in $K(\tau)$ of degree less than n . Once the $\mathbb{F}_q[t]$ -module structure is concerned at each stage of the reduction process a solution of a polynomial equation with the coefficients in the function field is required. There are efficient algorithms to do this (cf. [GS00]). However, it is unreasonable to expect that the roots of the polynomials that appear are expressed as polynomials in c_j . So, in general we do not obtain a \mathfrak{t} -module structure.

7.2. **case** $\text{rk } \phi < \text{rk } \psi$. Let K be a perfect field. Denote by σ the inverse map to τ . For the sake of simplicity we denote the value of $\sigma^k(c)$ as $c^{(-k)}$ for $c \in K$. Then the ring $K\{\sigma\}$ is the ring of adjoint twisted polynomials in K such that

$$(7.2) \quad \sigma x = x^{(-1)}\sigma \quad \text{for } x \in K,$$

(cf. [G95]). We can consider the \mathbf{t}^σ -module which we can define as in definition 2.1 by replacing τ with σ . Similarly, we define a morphism of the \mathbf{t}^σ -modules. Again the category of \mathbf{t}^σ -modules with the zero \mathbf{t}^σ -module attached is an additive, $\mathbb{F}_q[t]$ -linear category. Moreover, there is an isomorphism of $\mathbb{F}_q[t]$ -modules

$$\text{Ext}_\sigma^1(\Phi^{\text{ad}}, \Psi^{\text{ad}}) \cong \text{Der}(\Phi^{\text{ad}}, \Psi^{\text{ad}}) / \text{Der}_{in}(\Phi^{\text{ad}}, \Psi^{\text{ad}}),$$

where Φ^{ad} and Ψ^{ad} are \mathbf{t}^σ -modules.

Let $\omega(x) = \sum_{i=0}^n a_i x^i$ then $\omega(\tau) \in K\{\tau\}$ and $\omega^{\text{ad}} = \omega(\sigma) \in K\{\sigma\}$. Let $\phi(x) = \theta + \sum_{i=1}^n a_i x^i$ and $\psi(x) = \theta + \sum_{i=1}^m b_i x^i$ be polynomials in $K[x]$ then $\phi(\tau), \psi(\tau)$ are Drinfeld modules and $\phi(\sigma), \psi(\sigma)$ are Drinfeld \mathbf{t}^σ -modules.

Lemma 7.1. *Let $\phi(x) = \theta + \sum_{i=1}^n a_i x^i$ and $\psi(x) = \theta + \sum_{i=1}^m b_i x^i$ be polynomials in $K[x]$ such that $\deg \phi(x) > \deg \psi(x)$ and let the \mathbf{t} -module $\text{Ext}_\tau^1(\phi(\tau), \psi(\tau))$ be given by the map*

$$\Pi : \mathbb{F}_q[t] \longrightarrow \text{Mat}_n(K\{\tau\}).$$

Then the $\mathbb{F}_q[t]$ -module $\text{Ext}_\sigma^1(\phi(\sigma), \psi(\sigma))$ has a structure of \mathbf{t}^σ -module, given by the map

$$\widehat{\Pi} : \mathbb{F}_q[t] \longrightarrow \text{Mat}_n(K\{\sigma\}),$$

where the matrix $\widehat{\Pi}_t$ is obtained from the matrix Π_t , by replacing τ with σ and coefficients of the form $c^{(i)}$ with coefficients of the form $c^{(-i)}$.

Proof. Similarly as in the proof of Lemma 3.1 we reduce a biderivation $\delta \in \text{Der}(\phi(\sigma), \psi(\sigma))$ by the generators $\delta^{(c\sigma^k)}$ of $\text{Der}_{in}(\phi(\sigma), \psi(\sigma))$. As a result we obtain that

$$\text{Der}(\phi(\sigma), \psi(\sigma)) \cong K\{\sigma\}_{<n} := \{w(\sigma) \in K\{\sigma\} \mid \deg_\sigma(w) < n\}.$$

Let $c\sigma^k$ be a generator of $K\{\sigma\}_{<n}$. Then

$$(7.3) \quad t * c_i \sigma^i = \psi(\sigma) c_i \sigma^i = \theta c_i \sigma^i + \sum_{j=1}^m b_j c_i^{(-j)} \sigma^{j+i}.$$

Comparing the equality (7.3) with the value of $t * c_i \tau^i$ (see (5.2)), we conclude that $t * c_i \sigma^i$ can be obtained from $t * c_i \tau^i$, by replacing τ with σ and coefficients of the form $c_i^{(j)}$ with the coefficients of the form $c_i^{(-j)}$.

It is easy to see, that in the same way, we can obtain the generator $\delta^{(c\sigma^k)}$ from $\delta^{(c\tau^k)}$. Therefore after reduction process $t * c_i \sigma^i$ can be obtained from $t * c_i \tau^i$ by the previously described replacement. This proves the claim. \square

Example 7.1. Consider the following polynomials: $\phi(x) = \theta + ax^3$ and $\psi(x) = \theta + bx^2$ for fixed $a, b \in K$. Then the \mathfrak{t} -module structure on $\text{Ext}_\tau^1(\phi(\tau), \psi(\tau))$ is given by the following matrix:

$$\Pi_t = \begin{bmatrix} \theta & 0 & 0 \\ 0 & \theta & \frac{b}{a^{(1)}}(\theta - \theta^{(1)})\tau^2 \\ b\tau^2 & \frac{b \cdot b^{(2)}}{a^{(2)}}\tau^4 & \theta + \frac{b \cdot b^{(2)} \cdot b^{(4)}}{a^{(5)} \cdot a^{(2)}}\tau^6 \end{bmatrix}$$

Then the \mathfrak{t}^σ -module structure on $\text{Ext}_\sigma^1(\phi(\sigma), \psi(\sigma))$ is given by the matrix:

$$\widehat{\Pi}_t = \begin{bmatrix} \theta & 0 & 0 \\ 0 & \theta & \frac{b}{a^{(-1)}}(\theta - \theta^{(-1)})\sigma^2 \\ b\sigma^2 & \frac{b \cdot b^{(-2)}}{a^{(-2)}}\sigma^4 & \theta + \frac{b \cdot b^{(-2)} \cdot b^{(-4)}}{a^{(-5)} \cdot a^{(-2)}}\sigma^6 \end{bmatrix}$$

Let us note, that in this example we do not carry out calculations, because the matrix Π_t can be easily obtained using the explicit formulas described in Appendix A.

Consider the following maps:

$$(7.4) \quad (-)^\sigma : K\{\tau\} \rightarrow K\{\sigma\}; \quad \left(\sum_{i=0}^n a_i \tau^i \right)^\sigma = \sum_{i=0}^n a_i^{(-i)} \sigma^i,$$

$$(7.5) \quad (-)^\tau : K\{\sigma\} \rightarrow K\{\tau\}; \quad \left(\sum_{i=0}^n b_i \sigma^i \right)^\tau = \sum_{i=0}^n b_i^{(i)} \tau^i.$$

These maps are additive mutual inverses. We associate with $\Phi : \mathbb{F}_q[t] \rightarrow \text{Mat}_e(K\{\tau\})$ the adjoint homomorphism

$$(7.6) \quad \Phi^\sigma : \mathbb{F}_q[t] \rightarrow \text{Mat}_e(K\{\sigma\})$$

such that each matrix X_t is mapped to $\left[(X_t(\tau))^\sigma \right]^T$ i.e. the matrix entry $X_{i,j}(\tau)$ is mapped to $(X_{j,i}(\tau))^\sigma$. The inverse of $(-)^\sigma$ is given by the map that associates with $\Gamma : \mathbb{F}_q[t] \rightarrow \text{Mat}_e(K\{\sigma\})$ the following homomorphism:

$$(7.7) \quad \Gamma^\tau : \mathbb{F}_q[t] \rightarrow \text{Mat}_e(K\{\tau\}); \quad X_t \rightarrow \left[(X_t(\sigma))^\tau \right]^T$$

We have the following duality:

Theorem 7.2. *Assume that K is a perfect A -field. Let Φ and Ψ be \mathfrak{t} -modules. Then there exists an isomorphism of $\mathbb{F}_q[t]$ -modules:*

$$(7.8) \quad \text{Ext}_\tau^1(\Phi, \Psi) \cong \text{Ext}_\sigma^1(\Psi^\sigma, \Phi^\sigma)$$

Proof. Notice that

$$(7.9) \quad \text{Ext}_\tau^1(\Phi, \Psi) = \text{Der}(\Phi, \Psi)/\text{Der}_{in}(\Phi, \Psi); \quad t * \delta_t(\tau) = \Psi_t \cdot \delta_t(\tau)$$

and

$$(7.10) \quad \text{Ext}_\sigma^1(\Psi^\sigma, \Phi^\sigma) = \text{Der}(\Psi^\sigma, \Phi^\sigma)/\text{Der}_{in}(\Psi^\sigma, \Phi^\sigma); \quad t * \delta_t(\tau) = \delta_t^\sigma(\sigma)\Psi_t^\sigma$$

One readily verifies that $(-)^{\sigma}$ is well defined since it maps inner biderivations onto inner biderivations. Similarly, for $(-)^{\tau}$. Thus we have a bijective map (induced by $(-)^{\sigma}$) $\text{Ext}_\tau^1(\Phi, \Psi) \rightarrow \text{Ext}_\sigma^1(\Psi^\sigma, \Phi^\sigma)$. As

$$(t * c\tau^k)^\sigma = (\Psi_t c\tau^k)^\sigma = c^{(-k)}\sigma^k\Psi_t^\sigma = t * c^{(-k)}\sigma^k = t * (c\tau^k)^\sigma$$

we see that $(-)^{\sigma}$ is a homomorphism of $\mathbb{F}_q[t]$ -modules. \square

Lemma 7.1 and Theorem 7.2 allow one to compute easily the $\mathbb{F}_q[t]$ -module structure of $\text{Ext}_\tau(\phi, \psi)$ for Drinfeld modules satisfying $\text{rk } \phi < \text{rk } \psi$. We will see, that this $\mathbb{F}_q[t]$ -module structure comes from the \mathfrak{t}^σ -module structure in a natural way.

Example 7.2. Let $\psi_t = \theta + b\tau^2$ and $\phi_t = \theta + a\tau^3$, for fixed $a, b \in K$, be Drinfeld modules. Then Theorem 7.2 implies that there is an isomorphism of $F_q[t]$ -modules

$$\text{Ext}_\tau^1(\theta + b\tau^2, \theta + a\tau^3) \cong \text{Ext}_\sigma^1(\theta + a^{(-3)}\sigma^3, \theta + b^{(-2)}\sigma^2).$$

By Lemma 7.1 the adjoint \mathfrak{t}^σ -module structure on

$$\text{Ext}_\sigma^1(\theta + a^{(-3)}\sigma^3, \theta + b^{(-2)}\sigma^2)$$

is given by the matrix $\widehat{\Pi}_t$ obtained from matrix Π_t of

$$\text{Ext}_\tau^1(\theta + a^{(-3)}\tau^3, \theta + b^{(-2)}\tau^2),$$

by replacing τ with σ and coefficients of the form $c^{(i)}$ with coefficients of the form $c^{(-i)}$. Therefore $\text{Ext}_\tau^1(\theta + b\tau^2, \theta + a\tau^3)$ has the \mathfrak{t}^σ -module structure, given by the matrix

$$\begin{bmatrix} \theta & 0 & 0 \\ 0 & \theta & \frac{b^{(-2)}}{a^{(-4)}}(\theta - \theta^{(-1)})\sigma^2 \\ b^{(-2)}\sigma^2 & \frac{b^{(-2)} \cdot b^{(-4)}}{a^{(-5)}}\sigma^4 & \theta + \frac{b^{(-2)} \cdot b^{(-4)} \cdot b^{(-6)}}{a^{(-8)} \cdot a^{(-5)}}\sigma^6 \end{bmatrix}$$

The following theorem is a consequence of Theorem 5.3, Lemma 7.1 and Theorem 7.2,

Theorem 7.3. *Assume that K is a perfect A -field. Let ϕ and ψ be Drinfeld modules, such that $\text{rk } \phi < \text{rk } \psi$. There is a short exact sequence of \mathfrak{t}^σ -modules*

$$(7.11) \quad 0 \longrightarrow \text{Ext}_{0,\tau}(\phi, \psi) \longrightarrow \text{Ext}_\tau^1(\phi, \psi) \longrightarrow \mathbb{G}_a \longrightarrow 0.$$

Remark 7.3. In a similar way, one can prove " \mathfrak{t}^σ -version" of the Theorem 6.1 and Corollary 6.2 for the products of Drinfeld modules.

8. PUSHOUTS AND PULLBACKS FOR \mathfrak{t} -MODULES AND THEIR APPLICATIONS

In this section we study pullbacks and pushouts in the category of \mathfrak{t} -modules. For definitions and basic properties of these the reader is advised to consult [M78]. In what follows by $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ we mean the map of \mathfrak{t} -modules $(\psi, G_a^m) \hookrightarrow (X, G_a^{m+n})$ which on underlying group schemes is given by the injection $G_a^m \cong 0^n \times G_a^m \hookrightarrow G_a^{m+n}$. Similarly, by $\begin{bmatrix} 1 & 0 \end{bmatrix}$ we denote the map of \mathfrak{t} -modules $(X, G_a^{m+n}) \rightarrow (\psi, G_a^m)$ which on underlying group schemes is given by the surjection $G_a^{m+n} \rightarrow G_a^m \times 0^n \cong G_a^m$. So that in the category of $\mathbb{F}_q[t]$ -modules pullbacks and pushouts exist follows from the fact that this category is abelian. In the next theorem we show that they exist in the category of \mathfrak{t} -modules and can be nicely described in the language of biderivations.

Theorem 8.1. *Let*

$$\delta : \quad 0 \longrightarrow F \longrightarrow X \longrightarrow E \longrightarrow 0$$

be a short exact sequence of \mathfrak{t} -modules, given by the biderivation δ . Then

- (i) *for each morphism of \mathfrak{t} -modules $g : G \longrightarrow E$, the pull-back of δ by g is a \mathfrak{t} -module, given by the biderivation $\delta \cdot g$,*
- (ii) *for each morphism of \mathfrak{t} -modules $f : F \longrightarrow G$, the push-out of δ by f is a \mathfrak{t} -module, given by the biderivation $f \cdot \delta$.*

Proof. The sequence δ has the following form:

$$(8.1) \quad \delta : \quad 0 \longrightarrow F \xrightarrow{\begin{bmatrix} 0 \\ 1 \end{bmatrix}} X \xrightarrow{\begin{bmatrix} 1 & 0 \end{bmatrix}} E \longrightarrow 0,$$

where F (resp. E) is given by the map $\Psi : \mathbb{F}_q[t] \longrightarrow \text{Mat}_e(K\{\tau\})$ (resp. $\Phi : \mathbb{F}_q[t] \longrightarrow \text{Mat}_d(K\{\tau\})$) and X is given by the following block matrix:

$$\begin{bmatrix} \Phi & 0 \\ \delta & \Psi \end{bmatrix} : \mathbb{F}_q[t] \longrightarrow \text{Mat}_{e+d}(K\{\tau\}).$$

Proof of part (i) : Let G be given by $\Xi : \mathbb{F}_q[t] \longrightarrow \text{Mat}_r(K\{\tau\})$. We claim that the following diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & F & \xrightarrow{\begin{bmatrix} 0 \\ 1 \end{bmatrix}} & Y & \xrightarrow{\begin{bmatrix} 1 & 0 \end{bmatrix}} & G \longrightarrow 0 \\
& & \downarrow = & & \downarrow \begin{bmatrix} g & 0 \\ 0 & 1 \end{bmatrix} & & \downarrow g \\
0 & \longrightarrow & F & \xrightarrow{\begin{bmatrix} 0 \\ 1 \end{bmatrix}} & X & \xrightarrow{\begin{bmatrix} 1 & 0 \end{bmatrix}} & E \longrightarrow 0
\end{array}$$

is commutative with exact rows, where Y is given by the map:

$$\begin{bmatrix} \Xi & 0 \\ \delta \cdot g & \Psi \end{bmatrix} : \mathbb{F}_q[t] \longrightarrow \text{Mat}_{e+d}(K\{\tau\}).$$

It obvious, that the rows are exact, and it is easy to see, that the two squares are commutative. We will check that the middle vertical map is a morphism of \mathfrak{t} -modules.

$$\begin{bmatrix} g & 0 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} \Xi & 0 \\ \delta \cdot g & \Psi \end{bmatrix} = \begin{bmatrix} g\Xi & 0 \\ \delta \cdot g & \Psi \end{bmatrix} = \begin{bmatrix} \Phi & 0 \\ \delta & \Psi \end{bmatrix} \cdot \begin{bmatrix} g & 0 \\ 0 & 1 \end{bmatrix},$$

where $g\Xi = \Phi g$ because g is a morphism of \mathfrak{t} -modules.

It remains to show that the universal property for a pullback holds true. So, assume that there is a \mathfrak{t} -module \widehat{Y} with the morphisms of \mathfrak{t} -modules $\alpha : \widehat{Y} \longrightarrow G$ and $\beta = \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} : \widehat{Y} \longrightarrow X$ such that $g\alpha = \begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix}$.

We claim that there is a morphism of \mathfrak{t} -modules $\gamma = \begin{bmatrix} \gamma_1 \\ \gamma_2 \end{bmatrix} : \widehat{Y} \longrightarrow Y$ such that

$$\begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} \gamma_1 \\ \gamma_2 \end{bmatrix} = \alpha \quad \text{and} \quad \begin{bmatrix} g & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \gamma_1 \\ \gamma_2 \end{bmatrix} = \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix}.$$

It is easy to see, that $\gamma = \begin{bmatrix} \alpha \\ \beta_2 \end{bmatrix}$ satisfies the above conditions. Therefore Y is the pullback given by the biderivation $\delta \cdot g$.

Proof of part (ii) : Let G be given by $\Xi : \mathbb{F}_q[t] \longrightarrow \text{Mat}_r(K\{\tau\})$. It is easy to check that the following diagram:

$$\begin{array}{ccccccccc}
0 & \longrightarrow & F & \xrightarrow{\begin{bmatrix} 0 \\ 1 \end{bmatrix}} & X & \xrightarrow{\begin{bmatrix} 1 & 0 \end{bmatrix}} & E & \longrightarrow & 0 \\
& & \downarrow f & & \downarrow \begin{bmatrix} 1 & 0 \\ 0 & f \end{bmatrix} & & \downarrow = & & \\
0 & \longrightarrow & G & \xrightarrow{\begin{bmatrix} 0 \\ 1 \end{bmatrix}} & Y & \xrightarrow{\begin{bmatrix} 1 & 0 \end{bmatrix}} & E & \longrightarrow & 0
\end{array}$$

is commutative with exact rows, where Y is given by the following map:

$$\begin{bmatrix} \phi & 0 \\ f \cdot \delta & \Xi \end{bmatrix} : \mathbb{F}_q[t] \longrightarrow \text{Mat}_{e+d}(K\{\tau\}).$$

The proof of the universal property for a pushout is similar to that for a pullback presented in part (i). \square

Let Λ be a ring, $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ an exact sequence of Λ -modules and D a Λ -module. It is a standard result in homological algebra (cf. [HS71, Theorem 5.2]) that one has a six term exact sequence called Hom – Ext sequence in the second variable:

$$\begin{aligned}
(8.2) \quad & 0 \longrightarrow \text{Hom}_\Lambda(D, A) \xrightarrow{i \circ -} \text{Hom}_\Lambda(D, B) \xrightarrow{\pi \circ -} \text{Hom}_\Lambda(D, C) \longrightarrow \\
& \xrightarrow{\delta \circ -} \text{Ext}_\Lambda^1(D, A) \xrightarrow{-i \circ -} \text{Ext}_\Lambda^1(D, B) \xrightarrow{-\pi \circ -} \text{Ext}_\Lambda^1(D, C).
\end{aligned}$$

Dually, one has the following Hom – Ext sequence in the first variable:

$$\begin{aligned}
(8.3) \quad & 0 \longrightarrow \text{Hom}_\Lambda(C, D) \xrightarrow{-\circ \pi} \text{Hom}_\Lambda(B, D) \xrightarrow{-\circ i} \text{Hom}_\Lambda(A, D) \xrightarrow{-\circ \delta} \\
& \longrightarrow \text{Ext}_\Lambda^1(C, D) \xrightarrow{-\circ(-\pi)} \text{Ext}_\Lambda^1(B, D) \xrightarrow{-\circ(-i)} \text{Ext}_\Lambda^1(A, D).
\end{aligned}$$

These sequences in general can be continued by higher Ext bifunctors. However, if Λ is a P.I.D. (or more generally a Dedekind ring) the last maps in (8.2) and (8.3) are surjections [HS71, Corollary 5.7], [I59].

For exact sequences in the category of \mathfrak{t} -modules we obtain analogous exact sequences of $\mathbb{F}_q[t]$ -modules.

The following example shows that in general for \mathfrak{t} -modules (E, Φ) and (F, Ψ) we have $\text{Hom}_\tau(F, E) \subsetneq \text{Hom}_{\mathbb{F}_q[t]}(G_a(K), G_a(K))$.

Example 8.1. Let ϕ be a Drinfeld module of generic characteristic of rank r defined over $K = \mathbb{F}_q(t)$ i.e. $\phi_t = \sum_{i=0}^r a_i \tau^i$, $a_i \in \mathbb{F}_q[t]$. It is well-known that $\text{Hom}_\tau(\phi, \phi)$ is a projective $\mathbb{F}_q[t]$ -module of rank at most r (cf. [Th04]). However, the Mordell-Weil group $\phi(K) = K$ of ϕ as an $F_q[t]$ -module

is a direct sum of a finite torsion module and a free $F_q[t]$ -module on \aleph_0 generators. This shows that $\mathrm{Hom}_\tau(\phi, \phi) \subsetneq \mathrm{Hom}_{\mathbb{F}_q[t]}(G_a(K), G_a(K))$.

The following theorem gives an explicit description of the corresponding six term exact sequences:

Theorem 8.2. *Let*

$$\delta : 0 \longrightarrow F \xrightarrow{i} X \xrightarrow{\pi} E \longrightarrow 0$$

be a short exact sequence of \mathfrak{t} -modules given by the biderivation δ and let G be a \mathfrak{t} -module.

(i) *There is an exact sequence of $\mathbb{F}_q[t]$ -modules:*

$$\begin{aligned} 0 \longrightarrow \mathrm{Hom}_\tau(G, F) \xrightarrow{i \circ -} \mathrm{Hom}_\tau(G, X) \xrightarrow{\pi \circ -} \mathrm{Hom}_\tau(G, E) \longrightarrow \\ \xrightarrow{\delta \circ -} \mathrm{Ext}_\tau^1(G, F) \xrightarrow{-i \circ -} \mathrm{Ext}_\tau^1(G, X) \xrightarrow{-\pi \circ -} \mathrm{Ext}_\tau^1(G, E) \rightarrow 0. \end{aligned}$$

(ii) *There is an exact sequence of $\mathbb{F}_q[t]$ -modules:*

$$\begin{aligned} 0 \longrightarrow \mathrm{Hom}_\tau(E, G) \xrightarrow{-\circ \pi} \mathrm{Hom}_\tau(X, G) \xrightarrow{-\circ i} \mathrm{Hom}_\tau(F, G) \xrightarrow{-\circ \delta} \\ \longrightarrow \mathrm{Ext}_\tau^1(E, G) \xrightarrow{-\circ(-\pi)} \mathrm{Ext}_\tau^1(X, G) \xrightarrow{-\circ(-i)} \mathrm{Ext}_\tau^1(F, G) \rightarrow 0. \end{aligned}$$

Proof. We will give a proof of part (i). The proof for part (ii) is similar and is left to the reader,

Recall that $i = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$, $\pi = [1 \ 0]$ and \mathfrak{t} -modules F , E and X are given by the maps $\Psi : \mathbb{F}_q[t] \longrightarrow \mathrm{Mat}_e(K\{\tau\})$, $\Phi : \mathbb{F}_q[t] \longrightarrow \mathrm{Mat}_d(K\{\tau\})$ and

$$\begin{bmatrix} \Phi & 0 \\ \delta & \Psi \end{bmatrix} : \mathbb{F}_q[t] \longrightarrow \mathrm{Mat}_{e+d}(K\{\tau\}).$$

Assume that G is given by $\Xi : \mathbb{F}_q[t] \longrightarrow \mathrm{Mat}_r(K\{\tau\})$. The exactness at $\mathrm{Hom}_\tau(G, F)$, $\mathrm{Hom}_\tau(G, X)$ is obvious from the form of maps i and π .

Now we consider the exactness at $\mathrm{Hom}_\tau(G, E)$. Let $\begin{bmatrix} f_1 \\ f_2 \end{bmatrix} : G \longrightarrow X$ be a map of \mathfrak{t} -modules. Hence there is an equality:

$$\begin{bmatrix} f_1 \\ f_2 \end{bmatrix} \Xi = \begin{bmatrix} \Phi & 0 \\ \delta & \Psi \end{bmatrix} \begin{bmatrix} f_1 \\ f_2 \end{bmatrix}.$$

This implies

$$(8.4) \quad f_1 \Xi = \Phi f_1 \quad \text{and} \quad f_2 \Xi = \delta f_1 + \Psi f_2.$$

We will proof, that an exact sequence given by the biderivation:

$$\delta \circ \pi \circ \begin{bmatrix} f_1 \\ f_2 \end{bmatrix} = \delta \circ [1 \ 0] \circ \begin{bmatrix} f_1 \\ f_2 \end{bmatrix} = \delta \circ f_1 = \delta f_1$$

splits. Consider the following diagram:

$$\delta f_1 : \begin{array}{ccccccc} 0 & \longrightarrow & F & \xrightarrow{\begin{bmatrix} 0 \\ 1 \end{bmatrix}} & G \oplus F & \xrightarrow{\begin{bmatrix} 1 & 0 \end{bmatrix}} & G \longrightarrow 0, \\ & & \downarrow = & & \downarrow \begin{bmatrix} 1 & 0 \\ f_2 & 1 \end{bmatrix} & & \downarrow = \\ 0 & \longrightarrow & F & \xrightarrow{\begin{bmatrix} 0 \\ 1 \end{bmatrix}} & Y & \xrightarrow{\begin{bmatrix} 1 & 0 \end{bmatrix}} & G \longrightarrow 0 \end{array}$$

where the lower row is given by the biderivation δf_1 , ie. Y is defined by the map:

$$\begin{bmatrix} \Xi & 0 \\ \delta f_1 & \Psi \end{bmatrix} : \mathbb{F}_q[t] \longrightarrow \text{Mat}_{e+r}(K\{\tau\}).$$

It is easy to see, that this diagram is commutative, with exact rows. From (8.4) the middle vertical map is a morphism of \mathfrak{t} -modules that is also an isomorphism. Therefore the sequence δf_1 splits.

On the other hand, assume that for some morphism of \mathfrak{t} -modules $f_1 : G \longrightarrow E$ the sequence given by the biderivation δf_1 splits. Therefore $\delta f_1 \in \text{Der}_{in}(\Xi, \Psi)$, so there is $U \in \text{Mat}_{e+r}(K\{\tau\})$ such that

$$(8.5) \quad \delta f_1 = \delta^{(U)} = U\Xi - \Psi U.$$

Then $f = \begin{bmatrix} f_1 \\ U \end{bmatrix} : G \longrightarrow X$ is a morphism of \mathfrak{t} -modules. Indeed

$$\begin{bmatrix} f_1 \\ U \end{bmatrix} \Xi = \begin{bmatrix} f_1 \Xi \\ U \Xi \end{bmatrix} = \begin{bmatrix} \Phi f_1 \\ \delta f_1 + \Psi U \end{bmatrix} = \begin{bmatrix} \Phi & 0 \\ \delta & \Psi \end{bmatrix} \begin{bmatrix} f_1 \\ U \end{bmatrix}$$

where the second equality follows from (8.5) and the fact that $f_1 : G \longrightarrow E$ is a morphism of \mathfrak{t} -modules. Hence $\pi \circ f = \begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} f_1 \\ U \end{bmatrix} = f_1$, which shows exactness at $\text{Hom}_\tau(G, E)$.

For the exactness at $\text{Ext}_\tau^1(G, F)$ let $f : G \longrightarrow E$ be a morphism of \mathfrak{t} -modules. We will prove that the extension given by the biderivation $-i\delta f$ splits. Recall that δf determines the following short exact sequence:

$$0 \longrightarrow F \longrightarrow Y \longrightarrow G \longrightarrow 0,$$

where Y is defined by the map:

$$\begin{bmatrix} \Xi & 0 \\ \delta f & \Psi \end{bmatrix} : \mathbb{F}_q[t] \longrightarrow \text{Mat}_{r+e}(K\{\tau\}).$$

Then the biderivation $-i\delta f$ determines the following short exact sequence:

$$0 \longrightarrow X \longrightarrow \widehat{Y} \longrightarrow G \longrightarrow 0,$$

where \widehat{Y} is defined by the map:

$$\left[\begin{array}{c|cc} \Xi & 0 & 0 \\ \hline 0 & \Phi & 0 \\ -\delta f & \delta & \Psi \end{array} \right] : \mathbb{F}_q[t] \longrightarrow \text{Mat}_{r+e}(K\{\tau\}).$$

It is easy to check, that the following diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & X & \xrightarrow{\begin{bmatrix} 0 \\ 1 \end{bmatrix}} & Y & \xrightarrow{\begin{bmatrix} 1 & 0 \end{bmatrix}} & X \longrightarrow 0, \\ & & \downarrow = & & \downarrow \begin{bmatrix} 1 & 0 & 0 \\ -f & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} & & \downarrow = \\ 0 & \longrightarrow & X & \xrightarrow{\begin{bmatrix} 0 \\ 1 \end{bmatrix}} & G \oplus X & \xrightarrow{\begin{bmatrix} 1 & 0 \end{bmatrix}} & G \longrightarrow 0 \end{array}$$

is commutative and the map $\begin{bmatrix} 1 & 0 & 0 \\ -f & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ is an isomorphism of \mathfrak{t} -modules.

Therefore the sequence $-i\delta f$ splits.

Now consider the exact sequence:

$$\eta : 0 \longrightarrow F \longrightarrow Y \longrightarrow G \longrightarrow 0 \in \text{Ext}_\tau^1(G, F),$$

where Y is given by the map

$$\left[\begin{array}{c|cc} \Xi & 0 \\ \hline \eta & \Psi \end{array} \right] : \mathbb{F}_q[t] \longrightarrow \text{Mat}_{r+e}(K\{\tau\}),$$

and assume that the sequence

$$-i \circ \eta : 0 \longrightarrow X \longrightarrow \widehat{Y} \longrightarrow G \longrightarrow 0 \in \text{Ext}_\tau^1(G, F)$$

splits, where \widehat{Y} is defined by the map

$$\left[\begin{array}{c|cc} \Xi & 0 & 0 \\ \hline 0 & \Phi & 0 \\ -\eta & \delta & \Psi \end{array} \right] : \mathbb{F}_q[t] \longrightarrow \text{Mat}_{r+e}(K\{\tau\}).$$

Because $-i \circ \eta$ splits, then the biderivation $-i \circ \eta \in \text{Der}_{in}\left(\Xi, \begin{bmatrix} \Phi & 0 \\ \delta & \Psi \end{bmatrix}\right)$.

Hence, there is $U = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} \in \text{Mat}_{e+d \times r}(K\{\tau\})$ such that

$$\begin{bmatrix} 0 \\ -\eta \end{bmatrix} = \delta^{(U)} = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} \Xi - \begin{bmatrix} \Phi & 0 \\ \delta & \Psi \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} = \begin{bmatrix} u_1 \Xi - \Phi u_1 \\ u_2 \Xi - \delta u_1 - \Psi u_2 \end{bmatrix}.$$

Therefore u_1 is a morphism of \mathfrak{t} -modules and

$$\eta = \delta u_1 + \Psi u_2 - u_2 \Xi = \delta u_1 - \underbrace{(u_2 \Xi - \Psi u_2)}_{=\delta^{(u_2)}} = \delta u_1 - \delta^{(u_2)}.$$

Thus the biderivations η and δu_1 determine the same extension in $\text{Ext}_\tau^1(G, F)$, which shows the exactness at $\text{Ext}_\tau^1(G, F)$.

Now we consider exactness at $\text{Ext}_\tau^1(G, X)$. For $\eta \in \text{Ext}_\tau^1(G, F)$ there is an equality

$$-\pi \circ (-i \circ \eta) = \pi \circ i(\eta) = 0.$$

Hence the sequence given by the biderivation $-\pi \circ (-i \circ \eta)$ splits.

On the other hand assume, that $\eta = \begin{bmatrix} \eta_1 \\ \eta_2 \end{bmatrix}$ is the biderivation determining the exact sequence from $\text{Ext}_\tau^1(G, X)$, such that $-\pi \circ \eta$ gives a split sequence in $\text{Ext}_\tau^1(G, E)$. Then

$$-\pi \circ \eta = - \begin{bmatrix} 1 & 0 \end{bmatrix} \circ \begin{bmatrix} \eta_1 \\ \eta_2 \end{bmatrix} = -\eta_1 \in \text{Der}_{in}(\Xi, \Phi).$$

Thus there is $u \in \text{Mat}_{d \times r}(K\{\tau\})$ such that $-\eta_1 = \delta^{(u)} = u\Xi - \Phi u$. We put $U = \begin{bmatrix} u \\ 0 \end{bmatrix} \in \text{Mat}_{d+e \times r}(K\{\tau\})$. Then the inner biderivation $\delta^{(U)} \in \text{Der}_{in}\left(\Xi, \begin{bmatrix} \Phi & 0 \\ \delta & \Psi \end{bmatrix}\right)$ has the following form:

$$\delta^{(U)} = \begin{bmatrix} u \\ 0 \end{bmatrix} \Xi - \begin{bmatrix} \Phi & 0 \\ \delta & \Psi \end{bmatrix} \begin{bmatrix} u \\ 0 \end{bmatrix} = \begin{bmatrix} u\Xi - \Phi u \\ -\delta u \end{bmatrix} = \begin{bmatrix} \delta^{(u)} \\ -\delta u \end{bmatrix}.$$

Therefore

$$\begin{aligned} \eta + \delta^{(U)} &= \begin{bmatrix} -\delta^{(u)} \\ \eta_2 \end{bmatrix} + \begin{bmatrix} \delta^{(u)} \\ -\delta u \end{bmatrix} = \begin{bmatrix} 0 \\ \eta_2 - \delta u \end{bmatrix} = - \begin{bmatrix} 0 \\ 1 \end{bmatrix} \circ (\delta u - \eta_2) \\ &= -i \circ (\delta u - \eta_2). \end{aligned}$$

Thus the sequence corresponding to η is given by the biderivation $-i \circ (\delta u - \eta_2)$. This proves the exactness at $\text{Ext}_\tau^1(G, X)$. Now we will prove that the map: $\text{Ext}_\tau^1(G, X) \xrightarrow{-\pi \circ} \text{Ext}_\tau^1(G, E)$ is a surjection.

Let

$$(8.6) \quad \gamma: 0 \longrightarrow E \xrightarrow{\begin{bmatrix} 0 \\ 1 \end{bmatrix}} Y \xrightarrow{\begin{bmatrix} 1 & 0 \end{bmatrix}} G \longrightarrow 0,$$

be an element of $\text{Ext}_\tau^1(G, E)$ where Y is given by the following map:

$$(8.7) \quad \begin{bmatrix} \Xi & 0 \\ \gamma & \Psi \end{bmatrix}: \mathbb{F}_q[t] \rightarrow \text{Mat}_{e+r}(K\{\tau\}).$$

Then there exists the following commutative diagram with exact rows:

$$(8.8) \quad \begin{array}{ccccccc} 0 & \longrightarrow & X & \xrightarrow{\begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}} & Z & \xrightarrow{[1 \ 0]} & G \longrightarrow 0, \\ & & \downarrow & & \downarrow & & \downarrow = \\ 0 & \longrightarrow & E & \xrightarrow{-[1 \ 0]} & Y & \xrightarrow{\begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \end{bmatrix}} & G \longrightarrow 0 \\ & & & & \downarrow & & \\ & & & & \begin{bmatrix} 0 \\ 1 \end{bmatrix} & & \end{array}$$

where Z is given by the following map:

$$\begin{bmatrix} \Xi & 0 & 0 \\ -\gamma & \Phi & 0 \\ 0 & \delta & \Psi \end{bmatrix} : \mathbb{F}_q[t] \rightarrow M_{e+d+r}(K\{\tau\}).$$

Notice that

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \end{bmatrix} \begin{bmatrix} \Xi & 0 & 0 \\ -\gamma & \Phi & 0 \\ 0 & \delta & \Psi \end{bmatrix} = \begin{bmatrix} \Xi & 0 & 0 \\ \gamma & -\Phi & 0 \end{bmatrix} = \begin{bmatrix} \Xi & 0 \\ \gamma\Phi & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \end{bmatrix}.$$

Thus (8.8) is the diagram of maps of \mathfrak{t} -modules and $\begin{pmatrix} -\gamma \\ 0 \end{pmatrix} \rightarrow \gamma$, where $\begin{pmatrix} -\gamma \\ 0 \end{pmatrix}$ is the extension given by the upper row of (8.8). \square

Remark 8.2. In the Theorem 8.2 we assumed that the short exact sequence is given by the biderivation δ and in the result we obtained simple formula for the morphism from Hom_τ to Ext_τ^1 . In the case where the short exact sequence is not given by the biderivation, but it is isomorphic to a sequence given by the biderivation, the six-term exact sequences exist. The aforementioned isomorphism of short exact sequences induces the isomorphism of the corresponding six-term sequences.

We finish this section with the application of the six-term exact sequence.

Example 8.3. Let F (resp. E) be a Drinfeld module given by $\phi_t = \theta + \tau^3$ (resp. $\psi_t = \theta + \tau^2$) and consider the exact sequence of \mathfrak{t} -modules

$$(8.9) \quad 0 \rightarrow F \rightarrow X \rightarrow E \rightarrow 0,$$

where X is the extension given by the biderivation $\delta_t = 1 + \tau$ i.e. X is the \mathfrak{t} -module given by $\Gamma_t = \begin{bmatrix} \theta + \tau^2 & 0 \\ 1 + \tau & \theta + \tau^3 \end{bmatrix}$. Let C be the Drinfeld module

given by $\eta_t = \theta + \tau$. Since $\text{rk } F > \text{rk } C$ we have $\text{Hom}_\tau(F, C) = 0$ and from the six-term exact sequence we obtain the short exact sequence:

$$(8.10) \quad 0 \rightarrow \text{Ext}_\tau^1(E, C) \rightarrow \text{Ext}_\tau^1(X, C) \rightarrow \text{Ext}_\tau^1(F, C) \rightarrow 0$$

One readily verifies that $\text{Ext}_\tau^1(E, C)$ (resp. $\text{Ext}_\tau^1(F, C)$) is the \mathfrak{t} -module given by $\Psi_t = \begin{bmatrix} \theta & 0 \\ \tau & \theta + \tau^2 \end{bmatrix}$ (resp. $\Phi_t = \begin{bmatrix} \theta & 0 & 0 \\ \tau & \theta & \tau^2 \\ 0 & \tau & \theta \end{bmatrix}$).

We have:

$$(8.11) \quad \begin{aligned} \text{Ext}_\tau^1(X, C) &= \text{Der}(X, C) / \text{Der}_{\text{in}}(X, C) \\ &= K\{\tau\}^2 / \left\langle \delta \begin{bmatrix} c\tau^k & 0 \end{bmatrix}, \delta \begin{bmatrix} 0 & c\tau^k \end{bmatrix} \mid c \in K, k \in \mathbb{Z}_{\geq 0} \right\rangle, \end{aligned}$$

where

$$\begin{aligned} \delta \begin{bmatrix} c\tau^k & 0 \end{bmatrix} &= \begin{bmatrix} c\tau^k & 0 \end{bmatrix} \begin{bmatrix} \theta + \tau^2 & 0 \\ 1 + \tau & \theta + \tau^3 \end{bmatrix} - (\theta + \tau) \begin{bmatrix} c\tau^k & 0 \end{bmatrix} \\ &= \begin{bmatrix} \delta^{(c\tau^k)} & 0 \end{bmatrix} \quad \text{where } \delta^{(c\tau^k)} \in \text{Der}_{\text{in}}(E, C) \end{aligned}$$

$$\begin{aligned} \delta \begin{bmatrix} 0 & c\tau^k \end{bmatrix} &= \begin{bmatrix} 0 & c\tau^k \end{bmatrix} \begin{bmatrix} \theta + \tau^2 & 0 \\ 1 + \tau & \theta + \tau^3 \end{bmatrix} - (\theta + \tau) \begin{bmatrix} 0 & c\tau^k \end{bmatrix} \\ &= \begin{bmatrix} c\tau^k + c\tau^{k+1} & \delta^{(c\tau^k)} \end{bmatrix} \quad \text{where } \delta^{(c\tau^k)} \in \text{Der}_{\text{in}}(E, C) \end{aligned}$$

Thus first reducing the second coordinates by the elements $\delta \begin{bmatrix} 0 & c\tau^k \end{bmatrix}$ and then the first coordinates by $\delta \begin{bmatrix} c\tau^k & 0 \end{bmatrix}$ one can see that:

$$(8.12) \quad \text{Ext}_\tau^1(X, C) \cong \left\{ \begin{bmatrix} c_0 + c_1\tau & d_0 + d_1\tau + d_2\tau^2 \end{bmatrix} \mid c_i, d_i \in K \right\}$$

Enumerating the basis elements of (8.12) lexicographically i.e.

$$\begin{bmatrix} 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & \tau \end{bmatrix}, \begin{bmatrix} 0 & \tau^2 \end{bmatrix}, \begin{bmatrix} 1 & 0 \end{bmatrix}, \begin{bmatrix} 1 & \tau \end{bmatrix}$$

and computing $t * \begin{bmatrix} 0 & d_i \cdot \tau^i \end{bmatrix}$, $i = 0, 1, 2$ and $t * \begin{bmatrix} c_i \cdot \tau^i & 0 \end{bmatrix}$, $i = 0, 1$, in a similar to that in Section 4 way, one obtains that $\text{Ext}_\tau^1(X, C)$ is a \mathfrak{t} -module defined by the matrix:

$$(8.13) \quad \Omega_t = \begin{bmatrix} \theta & 0 & 0 & 0 & 0 \\ \tau & \theta & \tau^2 & 0 & 0 \\ 0 & \tau & \theta & 0 & 0 \\ 0 & 0 & -\tau & \theta & 0 \\ 0 & 0 & -\tau & \tau & \theta + \tau^2 \end{bmatrix}$$

Therefore

$$(8.14) \quad \begin{bmatrix} \Phi_t & 0 \\ \Delta_t & \Psi_t \end{bmatrix},$$

where $\Delta_t = \begin{bmatrix} 0 & 0 & -\tau \\ 0 & 0 & -\tau \end{bmatrix}$. Thus we showed that the exact sequence (8.10) is an extension of \mathfrak{t} -modules.

Change of δ_t in (8.9) to $\delta_t = 1 + \tau^3$ results in the change of Δ_t in (8.14).

For the new data $\Delta_t = \begin{bmatrix} 0 & 0 & -\tau \\ 0 & 0 & (\theta - \theta^{(1)})\tau + \tau^4 \end{bmatrix}$.

Similar arguments to that in the proof of Proposition 5.1, shows that the following assertions hold true.

Proposition 8.3. *Let $0 \rightarrow F \rightarrow X \rightarrow E \rightarrow 0$ be an exact sequence of \mathfrak{t} -modules, where F and E are Drinfeld modules, and let G be a Drinfeld module.*

(i) *If $\text{rk } G < \text{rk } F$ and $\text{rk } G < \text{rk } E$, then there is a short exact sequence of \mathfrak{t} -modules*

$$0 \rightarrow \text{Ext}_\tau^1(E, G) \rightarrow \text{Ext}_\tau^1(X, G) \rightarrow \text{Ext}_\tau^1(F, G) \rightarrow 0.$$

(i^D) *If $\text{rk } G > \text{rk } F$ and $\text{rk } G > \text{rk } E$, then there is a short exact sequence of \mathfrak{t} -modules*

$$0 \rightarrow \text{Ext}_\tau^1(G, F) \rightarrow \text{Ext}_\tau^1(G, X) \rightarrow \text{Ext}_\tau^1(G, E) \rightarrow 0.$$

As an immediate consequence of the Theorem 5.3 we obtain the following theorem:

Theorem 8.4. *Let $0 \rightarrow F \rightarrow X \rightarrow E \rightarrow 0$ be an exact sequence of \mathfrak{t} -modules, where F and E are Drinfeld modules, and let G be a Drinfeld module.*

(i) *If $\text{rk } G < \text{rk } F$ and $\text{rk } G < \text{rk } E$, then there is a short exact sequence of \mathfrak{t} -modules*

$$0 \rightarrow \text{Ext}_{0,\tau}(X, G) \rightarrow \text{Ext}_\tau^1(X, G) \rightarrow \mathbb{G}_a^2 \rightarrow 0.$$

(i^D) *If $\text{rk } G > \text{rk } F$ and $\text{rk } G > \text{rk } E$, then there is a short exact sequence of \mathfrak{t} -modules*

$$0 \rightarrow \text{Ext}_{0,\tau}(G, X) \rightarrow \text{Ext}_\tau^1(G, X) \rightarrow G_a^2 \rightarrow 0.$$

Remark 8.4. In the case where K is perfect, one can prove the corresponding " \mathfrak{t}^σ -versions" of the Proposition 8.3 and Theorem 8.4.

APPENDIX A. EXACT FORMULAS

Assume that ϕ and ψ are Drinfeld modules, such that $\phi_t = \theta + \sum_{i=1}^n a_i \tau^i$,

$\psi_t = \theta + \sum_{j=1}^m b_j \tau^j$ and $r := n - m > 0$. In this appendix we derive exact for-

mulas describing \mathfrak{t} -module structure on $\text{Ext}_\tau^1(\phi, \psi)$. This is done by transferring the $\mathbb{F}_q[t]$ -module structure from $\text{Ext}_\tau^1(\phi, \psi) = \text{Der}(\phi, \psi)/\text{Der}_{in}(\phi, \psi)$ to the space $K\{\tau\}_{<\text{rk } \phi}$, via the isomorphism from Lemma 3.1. Since each

element of $K\{\tau\}_{<\text{rk}\phi}$ is of the form $\sum_{i=0}^{n-1} c_i \tau^i$, it is sufficient to determine the values of multiplication by t on the generators $c_i \tau^i$ for $i = 0, 1, \dots, n-1$, where $c_i \in K$.

For the first r indices $i = 0, 1, \dots, r-1$ we have the following values:

$$(\mathbf{c}^i) \quad t * (c_i \tau^i) = \left(\theta + \sum_{j=1}^m b_j \tau^j \right) c_i \tau^i = c_i \theta \tau^i + \sum_{j=1}^m c_i^{(j)} b_j \tau^{j+i}$$

Now, we proceed with the computation of the value of $t * (c_{r+l} \tau^{r+l})$ where c_{r+l} is an element of field K . So, let us fix the natural number $l \in \{0, 1, \dots, m-1\}$. From the following computation

$$\begin{aligned} t * (c_{r+l} \tau^{r+l}) &= \left(\theta + \sum_{j=1}^m b_j \tau^j \right) c_{r+l} \tau^{r+l} = c_{r+l} \theta \tau^{r+l} + \sum_{j=1}^m c_{r+l}^{(j)} b_j \tau^{j+r+l} \\ (\text{A.1}) \quad &= c_{r+l} \theta \tau^{r+l} + \sum_{j=1}^{m-l-1} c_{r+l}^{(j)} b_j \tau^{j+r+l} + \sum_{j=m-l}^m c_{r+l}^{(j)} b_j \tau^{j+r+l} \end{aligned}$$

we see, that the monomials in the second sum have τ -degree greater or equal to n . Hence, like in the proof of Lemma 3.1 we will reduce these terms, using the generators of $\text{Der}_{in}(\phi, \psi)$. We do this in l steps. In each step we reduce the term with the highest exponent. Denote by $d_{r+l,k}$ the element of the field K such that the generator $\delta^{(d_{r+l,k} \tau^k)}$ will be used for reduction of the coefficient at τ^k for $k = m+l, m+l-1, \dots, 1, 0$. Note that during our reduction procedure we get a recursive relation for the elements $d_{r+l,k}$. However, this is enough to show that the $\mathbb{F}_q[t]$ -module structure on $K\{\tau\}_{<\text{rk}\phi}$ is in fact the \mathfrak{t} -module structure.

In order to simplify our formulas, we introduce the following notations:

$$(\text{A.2}) \quad \widehat{S_{\alpha,\beta}^\gamma} = \sum_{k=\alpha}^{\beta} \left(b_{\gamma-k} d_{n+l,n+k}^{(\gamma-k)} - a_{\gamma-k}^{(k)} d_{r+l,n+k} \right)$$

and

$$(\text{A.3}) \quad S_{\alpha,\beta}^\gamma = \sum_{k=\alpha}^{\beta} a_{\gamma-k}^{(k)} d_{r+l,n+k}.$$

We distinguish the following three cases: $m < r$, $m = r$ and $m > r$. In each of these cases we will carefully make the choice of the elements $d_{r+l,k}$. Recursively we obtain the final form of the value $t * (c_{r+l} \tau^{r+l})$.

A.1. **The case $c_1 : m < r$.** First we reduce the term $b_m c_{r+l}^{(m)} \tau^{n+l}$ using the generator

$$\begin{aligned} \delta^{(d_{r+l,n+l}\tau^l)}(t) &= d_{r+l,n+l}(\theta^{(l)} - \theta)\tau^l + \sum_{j=l+1}^{m+l} \left(a_{j-l}^{(l)} d_{r+l,n+l} - b_{j-l} d_{r+l,n+l}^{(j-l)} \right) \tau^j \\ &+ \sum_{j=m+l+1}^{n+l} a_{j-l}^{(l)} d_{r+l,n+l} \tau^j \end{aligned}$$

where $d_{r+l,n+l} = \frac{b_m c_{r+l}^{(m)}}{a_n^{(l)}}$.

Then we see that the coefficients at τ^{n+l-1} equal $b_{m-1} c_{r+l}^{(m-1)} - a_{n-1}^{(l)} d_{r+l,n+l}$. Therefore we will reduce this term with the generator $\delta^{(d_{r+l,n+l-1}\tau^{l-1})}(t)$, where

$$d_{r+l,n+l-1} = \frac{b_{m-1} c_{r+l}^{(m-1)} - a_{n-1}^{(l)} d_{r+l,n+l}}{a_n^{(l-1)}}.$$

Next, the coefficients at τ^{n+l-2} equal to

$$b_{m-2} c_{r+l}^{(m-2)} - a_{n-1}^{(l-1)} d_{r+l,n+l-1} - a_{n-2}^{(l)} d_{r+l,n+l},$$

can be reduced by the generator $\delta^{(d_{r+l,n+l-2}\tau^{l-2})}(t)$, with

$$d_{r+l,n+l-1} = \frac{b_{m-2} c_{r+l}^{(m-2)} - a_{n-1}^{(l-1)} d_{r+l,n+l-1} - a_{n-2}^{(l)} d_{r+l,n+l}}{a_n^{(l-2)}}.$$

It is easy to check that by continuing this process we get

$$(A.4) \quad d_{r+l,n+k} = \frac{b_{m-l+k} c_{r+l}^{(m-l+k)} - \sum_{i=k+1}^l a_{n+k-i}^{(i)} d_{r+l,n+i}}{a_n^{(k)}}$$

for $k = l, l-1, \dots, 1, 0$.

Now, we will determine the value of $t * (c_{r+l}\tau^{r+l})$. First, let's note that the smallest power of τ for $t * (c_{r+l}\tau^{r+l})$ before reduction (A.1) is $r+l$. Hence the elements $b_j c_{r+l}^{(j)}$ appear as coefficients at τ^j for $j > r+l$, and the element θc_{r+l} appears at τ^{r+l} . In the case of τ^j for $j < r+l$ there are only coefficients coming from the following generators:

$$(A.5) \quad \delta_l := \delta^{(d_{r+l,n+l}\tau^l)}, \delta_{l-1} := \delta^{(d_{r+l,n+l-1}\tau^{l-1})}, \dots, \delta_0 := \delta^{(d_{r+l,n})}.$$

Because the term $d_{r+l,n+k}(\theta^{(k)} - \theta)$ vanish for $k = 0$, then after reduction $t * (c_{r+l}\tau^{r+l})$ has no term at the power 0. For $j \in \{1, 2, \dots, l\}$ at τ^j appears one element $d_{r+l,n+j}(\theta - \theta^{(j)})$ from the generator δ^j and one element $d_{r+l,n+k}^{(j-k)} b_{j-k} - d_{r+l,n+k} a_{j-k}^{(k)}$ for each of the generators δ^k , $k = 0, 1, \dots, j-1$. Hence at τ^j we obtain the coefficient

$$(A.6) \quad d_{r+l,n+j}(\theta - \theta^{(j)}) + \widehat{S_{0,j-1}^j}.$$

In a similar way we see that the coefficient at τ^j for $j = l+1, l+2, \dots, m-1$ has the form $\widehat{S_{0,l}^j}$. In the case of τ^j for $j \in \{m, m+1, \dots, m+l\}$ the elements $d_{r+l, n+k}^{(j-k)} b_{j-k} - d_{r+l, n+k} a_{j-k}^{(k)}$ come from the generators $\delta_l, \delta_{l-1}, \dots, \delta_{j-m}$ and the elements $-d_{r+l, n+k} a_{j-k}^{(k)}$ come from the remaining generators i.e. $\delta_{j-m-1}, \dots, \delta_0$. Therefore at τ^j we obtain

$$(A.7) \quad \widehat{S_{j-m,l}^j} - S_{0, j-m-1}^j \quad \text{for } j = m, m+1, \dots, m+l.$$

For $j > m+l$ there are only elements of the form $-d_{r+l, n+k} a_{j-k}^{(k)}$ for all the generators. Thus the coefficients at τ^j have the following form:

$$\begin{aligned} -S_{0,l}^j & \quad \text{for } j = m+l+1, m+l+2, \dots, r+l-1, \\ c_{r+l}\theta - S_{0,l}^j & \quad \text{for } j = r+l, \\ b_{j-r-l} c_{r+l}^{(j-r-l)} - S_{0,l}^j & \quad \text{for } j = r+l+1, r+l+2, \dots, n-1. \end{aligned}$$

Therefore the values of $t * (c_{r+l} \tau^{r+l})$ for $l = 0, \dots, n-1-r$, after reduction, have the following form:

$$\begin{aligned} (\mathbf{c}_1^{r+l}) \quad t * (c_{r+l} \tau^{r+l}) &= \sum_{j=1}^l \left(d_{r+l, n+j} (\theta - \theta^{(j)}) + \widehat{S_{0, j-1}^j} \right) \tau^j + \sum_{j=l+1}^{m-1} \widehat{S_{0,l}^j} \tau^j \\ &+ \sum_{j=m}^{m+l} \left(\widehat{S_{j-m,l}^j} - S_{0, j-m-1}^j \right) \tau^j - \sum_{j=m+l+1}^{r+l-1} S_{0,l}^j \tau^j + (\theta c_{r+l} - S_{0,l}^{r+l}) \tau^{r+l} \\ &+ \sum_{j=r+l+1}^{n-1} \left(b_{j-r-l} c_{r+l}^{(j-r-l)} - S_{0,l}^j \right) \tau^j. \end{aligned}$$

We proved the following proposition.

Proposition A.1. *Let ϕ and ψ be Drinfeld modules, such that $r := \text{rk } \phi - \text{rk } \psi$. If $m < r$ then the \mathbb{F}_q -linear isomorphism from Lemma 3.1 is also an $\mathbb{F}_q[t]$ -module isomorphism, where the $\mathbb{F}_q[t]$ -module structure on $K\{\tau\}_{< \text{rk } \phi}$ is given by the formulas (\mathbf{c}^i) for $i = 0, 1, \dots, r-1$ and (\mathbf{c}_1^j) for $j = r, r+1, \dots, n-1$.*

A.2. The case \mathbf{c}_2 : $m = r$. Because the generators δ_k for $k = l, l-1, \dots, 0$ (cf. (A.5)) have coefficients at τ^j for $j \geq n$ of the form $d_{s,t} a_u^{(v)}$, the elements $d_{s,t}$ in this case will be given by the formula (A.4). Moreover, in this case, most of the calculations are completely analogous to the case \mathbf{c}_1 : $m < r$. The difference occurs for coefficients at τ^{r+l} . There, the element $d_{r+l, n+l} b_m - d_{r+l, n+l} a_m^{(l)}$ comes from δ_l and the elements $-d_{r+l, n+k} a_{j-k}^{(k)}$ come from the remaining generators: δ_k , $k = l-1, \dots, 0$. Hence the coefficient at τ^{r+l} has the form

$$c_{r+l}\theta + d_{r+l, n+l}^{(m)} b_m - S_{0,l}^j.$$

Therefore the value of $t * (c_{r+l}\tau^{r+l})$ after reduction has the following form:

$$\begin{aligned} (\mathbf{c}_2^{r+l}) \quad t * (c_{r+l}\tau^{r+l}) &= \sum_{j=1}^l \left(d_{r+l,n+1}(\theta - \theta^{(j)}) + \widehat{S_{0,j-1}^j} \right) \tau^j - \sum_{j=l+1}^m \widehat{S_{0,l}^j} \tau^j \\ &+ \sum_{j=m+1}^{r+l-1} \left(\widehat{S_{j-m,l}^j} = S_{0,j-m-1}^j \right) \tau^j + \left(c_{r+l}\theta + d_{r+l,n+l}^{(m)} b_m - S_{0,l}^j \right) \tau^{r+l} \\ &+ \sum_{j=r+l+1}^{n-1} \left(b_{j-r-l} c_{r+l}^{(j-r-l)} - S_{0,l}^j \right) \tau^j, \end{aligned}$$

where elements $d_{n+l,n+k}$ are the same as in the first case i.e. they are given by (A.4).

We obtained the following proposition:

Proposition A.2. *Let ϕ and ψ be Drinfeld modules, such that $r := \text{rk } \phi - \text{rk } \psi$. If $m = r$ then the \mathbb{F}_q -linear isomorphism from Lemma 3.1 is also $\mathbb{F}_q[t]$ -module isomorphism, where the $\mathbb{F}_q[t]$ -module structure on $K\{\tau\}_{<\text{rk } \phi}$ is given by the formulas (\mathbf{c}^i) for $i = 0, 1, \dots, r-1$ and (\mathbf{c}_2^j) for $j = r, r+1, \dots, n-1$.*

A.3. The case \mathbf{c}_3 : $m > r$. In this case the generators:

$$\delta^{(c\tau^{m-1})}, \delta^{(c\tau^{m-2})}, \dots, \delta^{(c\tau^r)}$$

have the coefficients at τ^j for $j \geq n$ of the form: $b_{r+l-*} d_{r+l,n+*}^{(r+l-*)} - a_{r+l-*}^{(k)} d_{r+l,n+*}$. Hence, these coefficients appear in the recursion for $d_{r+l,n+k}$ for some numbers k . Note, that for $l < r$ the elements $d_{r+l,n+k}$ will be the same as in two previous cases i.e. given by the formula (A.4). The difference will be occur in the case of $l \geq r$ for $d_{r+l,n+l-r}, d_{r+l,n+l-r-1}, \dots, d_{r+l,n}$. After reductions by the generators $\delta_l, \delta_{l-1}, \dots, \delta_{l-r+1}$ the highest power of τ is $l-r$, where the coefficient at τ^{l-r} is as follows:

$$b_{m-r} c_{r+l}^{(m-r)} - \sum_{i=l-r-1}^l a_{n+l-r-i}^{(i)} d_{r+l,n+i} + b_m d_{r+l,n+l}^{(m)}.$$

Then we put

$$d_{r+l,n+l-r} = \frac{b_{m-r} c_{r+l}^{(m-r)} - \sum_{i=l-r-1}^l a_{n+l-r-i}^{(i)} d_{r+l,n+i} + b_m d_{r+l,n+l}^{(m)}}{a_n^{(l-r)}}.$$

In each successive step there will be an additional element of the form $b_* d_{**}^{(*)}$, so we get the following form of recursion for $d_{r+l,n+k}$:

(A.8)

$$d_{r+l,n+k} = \frac{b_{m-l+k} c_{r+l}^{(m-l+k)} - \sum_{i=k+1}^l d_{r+l,n+i} a_{n+k-i}^{(i)} + \sum_{i=k+r}^l d_{r+l,n+i}^{(n+k-i)} b_{n+k-i}}{a_n^{(k)}}$$

for $k = l, l-1, \dots, 1, 0$.

Notice, that if $l < r$ the second sum in (A.8) disappears. Therefore this

formula describe $d_{n+l,n+k}$ in the both cases $l \geq r$ and $l < r$.

Determining the values for $t * (c_{r+l}\tau^{r+l})$ is similar to the previous cases. In particular, the coefficients at τ^j for $j < r+l$ will be the same as in the previous two cases. Differences occur for elements with exponent bigger than $r+l$. In addition, we have to distinguish between two sub-cases: $r+l > m$ and $r+l \leq m$. In each of these sub-cases the reduced form for t will be slightly different.

Sub-case $r+l > m$. For the monomial τ^{r+l} the elements $b_{r+l-*}d_{r+l,n+*}^{(r+l-*)} - a_{r+l-*}^{(k)}d_{r+l,n+*}$ appear in the coefficient. They come from the generators $\delta_l, \delta_{l-1}, \dots, \delta_{r+l-m}$. Additionally we have the elements $-a_{r+l-*}^{(k)}d_{r+l,n+*}$ coming from the remaining generators $\delta_{r+l-m-1}, \dots, \delta_0$. Hence at τ^{r+l} we obtain

$$\theta c_{r+l} + \widehat{S_{r+l-m,l}^{r+l}} - S_{0,r+l-m-1}^{r+l}.$$

In the case of τ^j for $j = r+l+1, \dots, m+l$ the situation is similar. There appear elements $b_{r+l-k}d_{r+l,n+k}^{(r+l-k)} - a_{r+l-k}^{(k)}d_{r+l,n+k}$ from generators $\delta_l, \dots, \delta_{j-m}$ and elements $-a_{r+l-k}^{(k)}d_{r+l,n+k}$ from the remaining generators: $\delta_{j-m-1}, \dots, \delta_0$. Therefore the coefficient at τ^j has the following form:

$$b_{j-r-l}c_{r+l}^{(j-r-l)} + \widehat{S_{j-m,l}^j} - S_{0,j-m-1}^j \quad \text{for } j = r+l+1, r+l+2, \dots, m+l.$$

For $j > m+l$ there are only elements of the form $-a_{r+l-k}^{(k)}d_{r+l,n+k}$ for all generators. Hence coefficient at τ^j equals $b_{j-r-l}c_{r+l}^{(j-r-l)} - S_{0,l}^j$ for $j > m+l$. Therefore the value of $t * (c_{r+l}\tau^{r+l})$ after reduction has the following form:

$$\begin{aligned} (\mathbf{c}_{3.1}^{r+l}) \quad t * (c_{r+l}\tau^{r+l}) &= \sum_{j=1}^l \left(d_{r+l,n+j}(\theta - \theta^{(j)}) + \widehat{S_{0,j-1}^j} \right) \tau^j + \sum_{j=l+1}^m \widehat{S_{0,l}^j} \tau^j \\ &+ \sum_{j=m+1}^{r+l-1} \left(\widehat{S_{j-m,l}^j} - S_{0,j-m-1}^j \right) \tau^j + \left(\theta c_{r+l} + \widehat{S_{r+l-m,l}^{r+l}} - S_{0,r+l-m-1}^{r+l} \right) \tau^{r+l} \\ &+ \sum_{j=r+l+1}^{m+l} \left(b_{j-r-l}c_{r+l}^{(j-r-l)} + \widehat{S_{j-m,l}^j} - S_{0,j-m-1}^j \right) \tau^j \\ &+ \sum_{j=m+l+1}^{n-1} \left(b_{j-r-l}c_{r+l}^{(j-r-l)} - S_{0,l}^j \right) \tau^j. \end{aligned}$$

Sub-case $r+l \leq m$. For $j = r+l, \dots, m$ only the elements $b_{r+l-*}d_{r+l,n+*}^{(r+l-*)} - a_{r+l-*}^{(k)}d_{r+l,n+*}$ coming from the generators $\delta_l, \delta_{l-1}, \dots, \delta_0$, appear at τ^j . Thus at τ^j we have the following coefficients:

$$\begin{aligned} c_{r+l}\theta + \widehat{S_{0,l}^{r+l}} & \quad \text{for } j = r+l, \\ b_{j-r-l}c_{r+l}^{(j-r-l)} + \widehat{S_{0,l}^j} & \quad \text{for } j = r+l+1, \dots, m. \end{aligned}$$

For $j = m+1, m+2, \dots, m+l$ the elements $b_{r+l-*}d_{r+l,n+*}^{(r+l-*)} - a_{r+l-*}^{(k)}d_{r+l,n+*}$ appear from generators $\delta_l, \delta_{l-1}, \dots, \delta_{j-m}$ and the elements $-a_{r+l-*}^{(k)}d_{r+l,n+*}$

from the other generators. Then the coefficient at τ^j is

$$b_{j-r-l}c_{r+l}^{(j-r-l)} + \widehat{S_{j-m,l}^j} - S_{0,j-m-1}^j \quad \text{for } j = m+1, m+2, \dots, m+l.$$

The coefficient at τ^j for $j > m+l$ is the same as in the former sub-case.

Therefore the value of $t * (c_{r+l}\tau^{r+l})$ after reduction has the following form:

$$\begin{aligned} (\mathbf{c}_{3.2}^{r+l}) \quad t * (c_{r+l}\tau^{r+l}) &= \sum_{j=1}^l \left(d_{r+l,n+j}(\theta - \theta^{(j)}) + \widehat{S_{0,j-1}^j} \right) \tau^j + \sum_{j=l+1}^m \widehat{S_{0,l}^j} \tau^j \\ &+ \sum_{j=m+1}^{r+l-1} \left(\widehat{S_{j-m,l}^j} - S_{0,j-m-1}^j \right) \tau^j + \left(c_{r+l}\theta + \widehat{S_{0,l}^{r+l}} \right) \tau^{r+l} \\ &+ \sum_{j=r+l+1}^m \left(b_{j-r-l}c_{r+l}^{(j-r-l)} + \widehat{S_{0,l}^j} \right) \tau^j + \sum_{j=m+1}^{m+l} \left(b_{j-r-l}c_{r+l}^{(j-r-l)} + \right. \\ &\left. \widehat{S_{j-m,l}^j} - S_{0,j-m-1}^j \right) \tau^j + \sum_{j=m+l+1}^{n-1} \left(b_{j-r-l}c_{r+l}^{(j-r-l)} - S_{0,l}^j \right) \tau^j, \end{aligned}$$

Proposition A.3. *Let ϕ and ψ be Drinfeld modules, such that $r := \text{rk } \phi - \text{rk } \psi$. If $m > r$ then the \mathbb{F}_q -linear isomorphism from Lemma 3.1 is also an $\mathbb{F}_q[t]$ -module isomorphism, where the $\mathbb{F}_q[t]$ -module structure on $K\{\tau\}_{<\text{rk } \phi}$ is given by the formulas (\mathbf{c}^i) for $i = 0, 1, \dots, r-1$ and if $r+l > m$ by formulas $(\mathbf{c}_{3.1}^{r+l})$ for $l = 0, 1, \dots, m-1$, and if $r+l \leq m$ by formulas $(\mathbf{c}_{3.2}^{r+l})$ for $l = 0, 1, \dots, m-1$.*

Using the calculated formulae in Propositions A.1, A.2, A.3 we can show that the $\mathbb{F}_q[t]$ -multiplication on the $K\{\tau\}_{<\text{rk } \phi}$ comes from the \mathfrak{t} -module structure.

Theorem A.4. *Let ϕ and ψ be Drinfeld modules, such that $r = \text{rk } \phi - \text{rk } \psi > 0$. Then $\text{Ext}_\tau^1(\phi, \psi)$ has natural structure of a \mathfrak{t} -module, given by the following matrix*

$$(A.9) \quad \Pi_t = \left[[t*\tau^0], [t*\tau^1], \dots, [t*\tau^{r-1}], [t*\tau^r], [t*\tau^{r+1}], \dots, [t*\tau^{n-1}] \right].$$

Proof. For the proof, it suffices to determine the matrix Π_t , defining the map $\Pi : F_q[t] \rightarrow M_n(K)\{\tau\}$, which gives the \mathfrak{t} -module structure on $\text{Ext}_\tau^1(\phi, \psi)$. We will carry out the reasoning in the case $m < r$. The other cases, are similar. Let $m < r$, then from Proposition A.1 we know the values of $t * (c_k\tau^k)$ for $k = 0, 1, \dots, n-1$.

In the space $K\{\tau\}_{<\text{rk } \phi}$, we choose the following basis: $e_i = \tau^i$ for $i = 0, 1, \dots, n-1$. In this basis, the element $t*(c_0)$ has the following coordinates

$$t * (c_0) = [c_0\theta, c_0^{(1)}b_1, c_0^{(2)}b_2, \dots, c_0^{(m)}b_m, 0, \dots, 0]$$

Notice that $c_0^{(k)}b_k$ is the value of the polynomial $b_k\tau^k$ at c_0 . Let $b_k\tau^k|_{c_0} := c_0^{(k)}b_k$ for $k = 1, \dots, m$. We can express $t * (c_0)$ in the following form:

$$t * (c_0) = [\theta, b_1\tau, b_2\tau^2, \dots, b_m\tau^m, 0, \dots, 0]|_{c_0}.$$

For simplicity, we introduce the following notation:

$$[t * \tau^0] := [\theta, b_1\tau, b_2\tau^2, \dots, b_m\tau^m, 0, \dots, 0]^t$$

for a vertical vector of polynomials, whose evaluation on c_0 are coordinates of $t * (c_0)$. In a similar way, we define $[t * \tau^k]$ for $k = 1, 2, \dots, n-1$:

$$[t * \tau^i] = \underbrace{[0, \dots, 0]}_{i-1}, \theta, b_1\tau, b_2\tau^2, \dots, b_m\tau^m, 0, \dots, 0]^t \quad \text{for } i = 1, 2, \dots, r-1.$$

Fix $l \in \{0, 1, \dots, m-1\}$ and define recursively the following polynomials:

(A.10)

$$d_{r+l, n+k}(\tau) = \frac{b_{m-l+k}\tau^{m-l+k} - \sum_{i=k+1}^l a_{n+k-i}^{(i)} d_{r+l, n+i}(\tau)}{a_n^{(k)}} \quad \text{for } k = l, l-1, \dots, 1, 0.$$

The evaluation of the polynomial $d_{r+l, n+k}(\tau)$ at c_{r+l} is equal to $d_{r+l, n+k}$. Then we can express $[t * \tau^{r+l}]$ in an explicit way:

$$\begin{aligned} [t * \tau^{r+l}] &= \\ &= \left[0, \left[(\theta - \theta^{(j)}) d_{r+l, n+j}(\tau) + \sum_{k=0}^{j-1} (b_{j-k} d_{r+l, n+k}^{(j-k)}(\tau) - a_{j-k}^{(k)} d_{r+l, n+k}(\tau)) \right]_{j=1}^l \right. \\ &\quad , \left[\sum_{k=0}^l (b_{j-k} d_{r+l, n+k}^{(j-k)}(\tau) - a_{j-k}^{(k)} d_{r+l, n+k}(\tau)) \right]_{j=l+1}^{m-1}, \\ &\quad , \left[\sum_{k=j-m}^l b_{j-k} d_{r+l, n+k}^{(j-k)}(\tau) - \sum_{k=0}^l a_{j-k}^{(k)} d_{r+l, n+k}(\tau) \right]_{j=m}^{m+l}, \\ &\quad , \left[- \sum_{k=0}^l d_{r+l, n+k} a_{j-k}^{(k)} \right]_{j=m+l+1}^{r+l-1}, \theta - \sum_{k=0}^l a_{r+l-k}^{(k)} d_{r+l, n+k}(\tau), \\ &\quad , \left. \left[b_{j-r-l} \tau^{j-r-l} - \sum_{k=0}^l a_{j-k}^{(k)} d_{r+l, n+k}(\tau) \right]_{j=r+l+1}^{n-1} \right]^t. \end{aligned}$$

Remark A.1. Notice that in the above formula the notation $[f(j)]_{j=u}^v$ means $[f(u), [f(u+1)], \dots, [f(v)]]$.

We claim that the \mathfrak{t} -module structure on $\text{Ext}^1(\phi, \psi)$ can be expressed by the map $\Pi : F_q[t] \longrightarrow M_n(K)\{\tau\}$, defined by

$$(A.11) \quad \Pi_t = \left[[t * \tau^0], [t * \tau^1], \dots, [t * \tau^{r-1}], [t * \tau^r], [t * \tau^{r+1}], \dots, [t * \tau^{n-1}] \right].$$

From the calculated form of Π_t , we observe that

- (i) The polynomials $d_{*,*}(\tau)$ used in the description of Π_t do not have constant terms.

(ii) If we write Π_t in the form $\Pi_t = (I\theta + N)\tau^0 + \sum_{i=1}^s A_i\tau^i$, then the matrix N vanishes.

Therefore Π_t defines the \mathfrak{t} -module structure on $\text{Ext}^1(\phi, \psi)$. Since the reasoning in the other two cases are analogous, we will only present the final forms of the corresponding matrices Π_t .

In the case $m = r$. We obtain the following coordinates

$$[t * \tau^i] = \left[\underbrace{0, \dots, 0}_{i-1}, \theta, b_1\tau, b_2\tau^2, \dots, b_m\tau^m, 0, \dots, 0 \right]^t \quad \text{for } i = 1, 2, \dots, r-1.$$

$$\begin{aligned} [t * \tau^{r+l}] &= \left[0, \left[(\theta - \theta^{(j)})d_{r+l, n+1}(\tau) + \sum_{k=0}^{j-1} \left(b_{j-k}d_{r+l, n+k}^{\{j-k\}}(\tau) - a_{j-k}^{(k)}d_{r+l, n+k}(\tau) \right) \right]_{j=1}^l, \right. \\ &\quad , \left[\sum_{k=0}^l \left(b_{j-k}d_{r+l, n+k}^{\{j-k\}}(\tau) - a_{j-k}^{(k)}d_{r+l, n+k}(\tau) \right) \right]_{j=l+1}^m, \\ &\quad , \left[\sum_{k=j-m}^l b_{j-k}d_{r+l, n+k}^{\{j-k\}}(\tau) - \sum_{k=0}^l a_{j-k}^{(k)}d_{r+l, n+k}(\tau) \right]_{j=m+1}^{r+l-1}, \\ &\quad , \left[\theta + b_m d_{r+l, n+l}^{\{m\}}(\tau) - \sum_{k=0}^l a_{r+l-k}^{(k)}d_{r+l, n+k}(\tau) \right], \\ &\quad , \left. \left[b_{j-r-l}\tau^{j-r-l} - \sum_{k=0}^l a_{j-k}^{(k)}d_{r+l, n+k}(\tau) \right]_{j=r+l+1}^{n-1} \right]^t, \end{aligned}$$

where the polynomials $d_{r+l, n+k}(\tau)$ are the same as before, see (A.10).

Therefore the \mathfrak{t} -module structure on $\text{Ext}^1(\phi, \psi)$ can be expressed by the map $\Pi : F_q[t] \rightarrow M_n(K)\{\tau\}$, defined by

$$(A.12) \quad \Pi_t = \left[[t*\tau^0], [t*\tau^1], \dots, [t*\tau^{r-1}], [t*\tau^r], [t*\tau^{r+1}], \dots, [t*\tau^{n-1}] \right].$$

In the case $m > r$. We obtain the following coordinates

$$[t * \tau^i] = \left[\underbrace{0, \dots, 0}_{i-1}, \theta, b_1\tau, b_2\tau^2, \dots, b_m\tau^m, 0, \dots, 0 \right]^t \quad \text{for } i = 1, 2, \dots, r-1.$$

To describe the remaining coordinates, as in the Proposition A.3, we must consider two sub-cases: $r+l \leq m$ or $r+l > m$.

In the sub-case $r + l \leq m$ we have:

$$\begin{aligned}
[t * \tau^{r+l}] &= \left[0, (\theta - \theta^{(j)}) \left[d_{r+l, n+j}(\tau) + \sum_{k=0}^{j-1} \left(b_{j-k} d_{r+l, n+k}^{\{j-k\}}(\tau) - a_{j-k}^{(k)} d_{r+l, n+k}(\tau) \right) \right]_{j=1}^l, \right. \\
&\quad , \left[\sum_{k=0}^l \left(b_{j-k} d_{r+l, n+k}^{\{j-k\}}(\tau) - a_{j-k}^{(k)} d_{r+l, n+k}(\tau) \right) \right]_{j=l+1}^{r+l-1}, \\
&\quad + \left[\theta + \sum_{k=0}^l \left(b_{r+l-k} d_{r+l, n+k}^{\{r+l-k\}}(\tau) - a_{r+l-k}^{(k)} d_{r+l, n+k}(\tau) \right) \right], \\
&\quad , \left[b_{j-r-l} \tau^{j-r-l} + \sum_{k=0}^l \left(b_{j-k} d_{r+l, n+k}^{\{j-k\}}(\tau) - a_{j-k}^{(k)} d_{r+l, n+k}(\tau) \right) \right]_{j=r+l+1}^m, \\
&\quad , \left[b_{j-r-l} \tau^{j-r-l} + \sum_{k=j-m}^l b_{j-k} d_{r+l, n+k}^{\{j-k\}}(\tau) - \sum_{k=0}^l a_{j-k}^{(k)} d_{r+l, n+k}(\tau) \right]_{j=m+1}^{m+l} \\
&\quad , \left. \left[b_{j-r-l} \tau^{j-r-l} - \sum_{k=0}^l a_{j-k}^{(k)} d_{r+l, n+k}(\tau) \right]_{j=m+l+1}^{n-1} \right]^t,
\end{aligned}$$

where the polynomials $d_{r+l, n+k}(\tau)$ are given recursively by the formula:

(A.13)

$$d_{r+l, n+k}(\tau) = \frac{b_{m-l+k} \tau^{m-l+k} - \sum_{i=k+1}^l a_{n+k-i}^{(i)} d_{r+l, n+i}(\tau) + \sum_{i=k+r}^l b_{n+k-i} d_{r+l, n+i}^{\{n+k-i\}}(\tau)}{a_n^{(k)}}$$

for $k = l, l-1, \dots, 1, 0$.

In the sub-case $r + l > m$ we have:

$$\begin{aligned}
[t * \tau^{r+l}] &= \left[0, \left[(\theta - \theta^{(j)}) d_{r+l, n+j}(\tau) + \sum_{k=0}^{j-1} \left(b_{j-k} d_{r+l, n+k}^{\{j-k\}}(\tau) - a_{j-k}^{(k)} d_{r+l, n+k}(\tau) \right) \right]_{j=1}^l, \right. \\
&\quad , \left[\sum_{k=0}^l \left(b_{j-k} d_{r+l, n+k}^{\{j-k\}}(\tau) - a_{j-k}^{(k)} d_{r+l, n+k}(\tau) \right) \right]_{j=l+1}^m, \\
&\quad , \left[\sum_{k=j-m}^l b_{j-k} d_{r+l, n+k}^{\{j-k\}}(\tau) - \sum_{k=0}^l a_{j-k}^{(k)} d_{r+l, n+k}(\tau) \right]_{j=m+1}^{r+l-1}, \\
&\quad , \left[\theta + \sum_{k=r+l-m}^l b_{r+l-k} d_{r+l, n+k}^{\{r+l-k\}}(\tau) - \sum_{k=0}^l a_{r+l-k}^{(k)} d_{r+l, n+k}(\tau) \right], \\
&\quad , \left[b_{j-r-l} \tau^{j-r-l} + \sum_{k=j-m}^l b_{j-k} d_{r+l, n+k}^{\{j-k\}}(\tau) - \sum_{k=0}^l a_{j-k}^{(k)} d_{r+l, n+k}(\tau) \right]_{j=r+l+1}^{m+l}, \\
&\quad , \left. \left[b_{j-r-l} \tau^{j-r-l} - \sum_{k=0}^l a_{j-k}^{(k)} d_{r+l, n+k}(\tau) \right]_{j=m+l+1}^{n-1} \right]^t,
\end{aligned}$$

where the polynomials $d_{r+l, n+k}(\tau)$ are given by A.13. \square

The Example 4.1 from section 4 is an example of a \mathfrak{t} -module $\text{Ext}^1(\psi, \phi)$ for the first case $m < r$. Now we give examples for the remaining two cases.

Example A.2. Let $\phi_t = \theta + \tau^4$ and $\psi_t = \theta + \tau^2$. Then $r = 2$ and $m = 2$ i.e. $m = r$. By Lemma 3.1 $\text{Ext}_\tau^1(\phi, \psi) \cong \{c_0 + c_1\tau + c_2\tau^2 + c_3\tau^3 \mid c_i \in K\}$. We have:

$$\begin{aligned} t * c_0 &= c_0\theta + c_0^{(2)}\tau^2 = [\theta, 0, \tau^2, 0]_{|c_0} \\ t * c_1 &= c_1\theta\tau + c_1^{(2)}\tau^3 = [0, \theta, 0, \tau^2]_{|c_1} \\ t * c_2 &= (c_2\theta + c_2^{(4)})\tau^2 = [0, 0, \theta + \tau^4, 0]_{|c_2} \\ t * c_3 &= c_3^{(2)}(\theta - \theta^{(1)})\tau + (c_3\theta + c_3^{(4)})\tau^3 = [0, (\theta - \theta^{(1)})\tau^2, 0, \theta + \tau^4]_{|c_3} \end{aligned}$$

and therefore

$$(A.14) \quad \Pi_t = \begin{bmatrix} \theta & 0 & 0 & 0 \\ 0 & \theta & 0 & (\theta - \theta^{(1)})\tau^2 \\ \tau^2 & 0 & \theta + \tau^4 & 0 \\ 0 & \tau^2 & 0 & \theta + \tau^4 \end{bmatrix}.$$

Example A.3. Let $\phi_t = \theta + \tau^5$ and $\psi_t = \theta + \tau^2$. Then $r = 3$ and $m = 2$ i.e. $m < r$. By Lemma 3.1 $\text{Ext}_\tau^1(\phi, \psi) \cong \{c_0 + c_1\tau + c_2\tau^2 + c_3\tau^3 + c_4\tau^4 \mid c_i \in K\}$. We have:

$$\begin{aligned} t * c_0 &= c_0\theta + c_0^{(2)}\tau^2 = [\theta, 0, \tau^2, 0, 0]_{|c_0}, \\ t * c_1 &= c_1\theta\tau + c_1^{(2)}\tau^3 = [0, \theta, 0, \tau^2, 0]_{|c_1}, \\ t * c_2 &= c_2\theta\tau^2 + c_2^{(2)}\tau^4 = [0, 0, \tau^2, 0, \tau^2]_{|c_2}, \\ t * c_3 &= c_3\theta\tau^3 + c_3^{(4)}\tau^2 = [0, 0, \tau^4, \theta, 0]_{|c_3}, \\ t * c_4 &= c_4^{(2)}(\theta - \theta^q)\tau + c_4^{(4)}\tau^3 + c_4\theta\tau^4 = [0, (\theta - \theta^{(1)})\tau^2, 0, \tau^4, \theta]_{|c_4} \end{aligned}$$

and therefore

$$(A.15) \quad \Pi_t = \begin{bmatrix} \theta & 0 & 0 & 0 & 0 \\ 0 & \theta & 0 & 0 & (\theta - \theta^{(1)})\tau^4 \\ \tau^2 & 0 & \theta & \tau^4 & 0 \\ 0 & \tau^2 & 0 & \theta & \tau^4 \\ 0 & 0 & \tau^2 & 0 & \theta \end{bmatrix}.$$

APPENDIX B. SOME CONSEQUENCES OF EXACT FORMULAS

Corollary B.1. *Let (K, v) be a valuation field. Assume that ϕ and ψ are Drinfeld's modules, such that $r := \text{rk } \phi - \text{rk } \psi > 0$ and $\phi_t = \theta + \sum_{i=1}^n a_i\tau^i$,*

$\psi_t = \theta + \sum_{j=1}^m b_j\tau^j$. If the following conditions

- (i) ψ has integer coefficients,
- (ii) $\phi - a_n\tau^n$ has integer coefficients,
- (iii) $v(a_n) < \min \left\{ v(b_1), q^i v(a_{n-i}), \frac{v(b_{m+1-i})}{q^{m-i}} \mid i = 1, 2, \dots, m-1 \right\}$,

hold true, then \mathfrak{t} -module $\text{Ext}_\tau^1(\phi, \psi)$ has integer coefficients.

Proof. First we will show, that the condition (iii) implies that the polynomials $d_{r+l, n+k}(\tau)$ have integer coefficients. The proof will be carried out for the case $m > r$. The other cases are similar. Fix $l \in \{0, 1, \dots, m-1\}$. We use the downward induction for $k = l, l-1, \dots, 1, 0$.

If $k = l$, then $d_{r+l, n+l}(\tau) = \frac{b_m}{a_n^{(l)}} \tau^m$. Because $v(a_n) < \frac{v(b_m)}{q^m}$, then

$$v(b_m) > q^m v(a_n) > q^l v(a_n) = v(a_n^{(l)}),$$

so the polynomial $d_{r+l, n+l}(\tau)$ has integer coefficients. Assume that the polynomials

$$d_{r+l, n+l}(\tau), d_{r+l, n+l-1}(\tau), \dots, d_{r+l, n+k+1}(\tau)$$

have integer coefficients. We have

$$\begin{aligned} d_{r+l, n+k}(\tau) &= \frac{b_{m-l+k} \tau^{m-l+k} - \sum_{i=k+1}^l a_{n+k-i}^{(i)} d_{r+l, n+i}(\tau) + \sum_{i=k+r}^l b_{n+k-i} d_{r+l, n+i}^{\{n+k-i\}}(\tau)}{a_n^{(k)}} \\ &= \frac{b_{m-l+k}}{a_n^{(k)}} \tau^{m-l+k} - \sum_{i=k+1}^l \frac{a_{n+k-i}^{(i)}}{a_n^{(k)}} d_{r+l, n+i}(\tau) + \sum_{i=k+r}^l \frac{b_{n+k-i}}{a_n^{(k)}} d_{r+l, n+i}^{\{n+k-i\}}(\tau). \end{aligned}$$

From the inductive hypotheses the polynomials $d_{r+l, n+i}(\tau)$ and $d_{r+l, n+i}^{\{n+k-i\}}(\tau)$ have integer coefficients. Thus, it is enough to show that the valuations of fractions

$$\frac{b_{m-l+k}}{a_n^{(k)}}, \quad \frac{a_{n+k-i}^{(i)}}{a_n^{(k)}}, \quad \frac{b_{n+k-i}}{a_n^{(k)}}$$

are positive. This follows from condition (iii). Hence the polynomials $d_{r+l, n+k}(\tau)$ have integer coefficients.

The claim follows now from the form of the matrix Π_t of the \mathfrak{t} -module $\text{Ext}_\tau^1(\phi, \psi)$ by using conditions (i) and (ii). \square

Corollary B.2. *Let ϕ and ψ be Drinfeld modules, such that $r = \text{rk } \phi - \text{rk } \psi > 0$. Then we have:*

$$\text{rk Ext}_\tau^1(\phi, \psi) = \begin{cases} 2m & \text{for } m \leq r \\ 3m & \text{for } m > r \end{cases}.$$

Proof. Claim follows immediately from the form of Π_t in the corresponding cases. \square

REFERENCES

- [A86] G. ANDERSON *t*-motives, Duke Math. J. 53 (1986), 457-502.
- [BP09] W. D. BROWNAWELL AND M. PAPANIKOLAS, *A rapid introduction to Drinfeld modules, t-modules, and t-motives*, Proceedings of the conference on t-motives: Hodge structures, transcendence and other motivic aspects, BIRS, Banff, Canada 2009.
- [D74] V. DRINFELD, *Elliptic modules*, Math. USSR Sb., 23 (1974) pp. 561- 592.
- [F13] Y. Z. FLICKER *Drinfeld Moduli Schemes and Automorphic Forms, The Theory of Elliptic Modules with Applications*, Springer 2013.

- [GS00] S. GAO, M. A. SHOKROLLAHI *Computing Roots of Polynomials over Function Fields of Curves* in Coding Theory and Cryptography. From Enigma and Geheimschreiber to Quantum Theory, Springer 2000, pp. 214-228.
- [G03] R. GILLARD, F. LEPREVOST, A. PANCHISHKIN, X-F. ROBLOT *Utilisation des modules de Drinfeld en cryptologie*, C. R. Acad. Sci. Paris, Ser. I 336 (2003), pp. 879-882.
- [G95] D. GOSS, *The Adjoint of the Carlitz Module and Fermat's Last Theorem*, Finite Fields and Their Applications **1**, (1995), pp.165-188.
- [G96] D. GOSS, *Basic Structures of Function Field Arithmetic*, Springer 1996.
- [HS71] P.J. HILTON, U. STAMMBACH *Course in Homological Algebra*, GTM 4, Springer 1971.
- [I59] T. ISHIKAWA, *On Dedekind Rings*, Journal of the Mathematical Society of Japan Vol. 11, No. 1, January, (1959), pp. 83-84.
- [M78] S. MACLANE, *Categories for Working Mathematician*, GTM 5, Springer 1978.
- [M95] S. MACLANE, *Homology*, (Classics in Mathematics), Springer 1995.
- [N19] A. K. NARAYANAN, *Polynomial Factorisation using Drinfeld Modules*, Newsletter of the London Mathematical Society, London Mathematical Society, 2019. hal-02368680.
- [PR03] M. A. PAPANIKOLAS, N. RAMACHANDRAN *A Weil-Barsotti formula for Drinfeld modules*, Journal of Number Theory 98, (2003), 407-431.
- [P95] B. POONEN, *Local height functions and the Mordell-Weil theorem for Drinfeld modules* Compositio Mathematica 97, (1995), pp.349-368.
- [R76] C.M. RINGEL, *Representations of K -species and bimodules* J. Algebra 41, (1976), 269-302.
- [R98] C.M. RINGEL, *Exceptional modules are tree modules*. Linear Algebra Appl. 275-276 (1998), 471-493.
- [Th04] D. THAKUR, *Function Field Arithmetic*, World Scientific, 2004.
- [W95] S.S. WOO, *Extensions of Drinfeld modules of rank 2 by the Carlitz module*, Bull. Korean Math. Soc. 32, (1995), pp. 251-257.

INSTITUTE OF MATHEMATICS, DEPARTMENT OF EXACT AND NATURAL SCIENCES,
UNIVERSITY OF SZCZECIN, UL. WIELKOPOLSKA 15, 70-451 SZCZECIN, POLAND
Email address: dawid.kedzierski@usz.edu.pl

INSTITUTE OF MATHEMATICS, DEPARTMENT OF EXACT AND NATURAL SCIENCES,
UNIVERSITY OF SZCZECIN, UL. WIELKOPOLSKA 15, 70-451 SZCZECIN, POLAND
Email address: piotrkras26@gmail.com