

ON THE ONE-SIDED BOUNDEDNESS OF THE LOCAL DISCREPANCY OF $\{n\alpha\}$ -SEQUENCES

JIANGANG YING AND YUSHU ZHENG

ABSTRACT. The main interest of this article is the one-sided boundedness of the local discrepancy of $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ on the interval $(0, c) \subset (0, 1)$ defined by

$$D_n(\alpha, c) = \sum_{j=1}^n 1_{\{\{j\alpha\} < c\}} - cn.$$

We focus on the special case $c \in (0, 1) \cap \mathbb{Q}$. Several necessary and sufficient conditions on α for $(D_n(\alpha, c))$ to be one-side bounded are derived. Using these, certain topological properties are given to describe the size of the set

$$O_c = \{\alpha \in \mathbb{R}^+ \setminus \mathbb{Q} : (D_n(\alpha, c)) \text{ is one-side bounded}\}.$$

1. INTRODUCTION

For $c \in (0, 1)$, an integer $n \geq 0$, and a sequence $\omega = (\omega_n : n \geq 1)$, the local discrepancy (sequence) of ω at c is defined to be

$$D_n(\omega, c) := \sum_{j=1}^n 1_{[0, c)}(\omega_j) - nc.$$

Given $\theta \in [0, 1)$ and $\alpha \in \mathbb{R}^+ \setminus \mathbb{Q}$, the sequence $\eta(\alpha, \theta) = (\eta_n(\alpha, \theta)) = (\{n\alpha + \theta\} : n \geq 1)$, where $\{\cdot\}$ denotes the fractional part, is called the sequence driven by irrational rotation with starting point θ and rotation parameter α . The boundedness of the local discrepancy (sequence) of this sequence

$$D_n(\eta, c) = D_n(\eta(\alpha, \theta), c), \quad n \geq 1,$$

has been a frequently discussed topic in discrepancy theory. The following well-known theorem was proved by Hecke [7], Ostrowski [10] (sufficiency) and Kesten [8] (necessity).

Theorem 1.1. *For any θ , the local discrepancy $(D_n(\eta, c))$ is bounded if and only if for some integer n , $\{n\alpha\} = c$.*

Theorem 1.1 gives a necessary and sufficient condition for $(D_n(\eta, c))$ to be bounded. Following this theorem, a natural question is, while $(D_n(\alpha, c))$ is unbounded, whether it is one-side bounded, i.e., bounded above or below. It was shown by Vera T Sós [12] that one can construct α and c , where c is not an integral multiple (mod 1) of α , such that $(D_n(\eta, c))$ is bounded from below (though unbounded by Theorem 1.1). Further results can be found in literature such as [2], [3], [6] and [4].

2010 *Mathematics Subject Classification.* Primary 11K06, 11K50.

Key words and phrases. irrational rotation, continued fraction, Diophantine approximation, discrepancy theory.

From now on, we will focus on the special case where $\theta = 0$ and $c = h/k \in (0, 1) \cap \mathbb{Q}$ ($h, k \in \mathbb{N}$ are coprime). Simply denote $D_n(\alpha, c) := D_n(\eta(\alpha, 0), c)$. We will use standard continued fraction notations. For $\alpha \in \mathbb{R}^+ \setminus \mathbb{Q}$, let $\alpha = [a_0; a_1 a_2 a_3 \cdots a_n \cdots]$ be the continued fraction of α , where $(a_n : n \geq 0)$ are by definition the partial quotients of α . The convergents $(p_n/q_n : n \geq -2)$ of α are defined by the following recursive equality:

$$\begin{cases} p_{-2} = 0, p_{-1} = 1, q_{-2} = 1, q_{-1} = 0; \\ p_n = a_n p_{n-1} + p_{n-2} \text{ and } q_n = a_n q_{n-1} + q_{n-2}, \text{ for } n \geq 0. \end{cases}$$

The main results in this article are stated as follows.

Theorem 1.2. *For $\alpha \in \mathbb{R}^+ \setminus \mathbb{Q}$, $c = h/k$, let $(a_n), (p_n/q_n)$ be the partial quotients and convergents of α respectively. Then the following statements are equivalent.*

- (1) $(D_n(\alpha, c))$ is bounded from above (resp. below), i.e. $\sup D_n(\alpha, c) < +\infty$ (resp. $\inf D_n(\alpha, c) > -\infty$);
- (2) there exists even (resp. odd) $m \geq -1$ such that for $n \geq 0$, $k \mid q_{m+2n}$;
- (3) there exists even (resp. odd) $m \geq -1$ such that (a_0, a_1, \dots, a_m) is of type- k and $k \mid a_{m+2n}$ for all $n \geq 1$.

(The definitions of tuples of type- k will be introduced in §4.)

Let $O = O_c$ be the set of $\alpha \in \mathbb{R}^+ \setminus \mathbb{Q}$ with $(D_n(\alpha, c))$ one-side bounded.

Theorem 1.3. (1) O is dense in \mathbb{R}^+ .

(2) O is of the 1st category.

(3) The Hausdorff dimension $\dim_H(O) \in (0, 1)$. Hence O is uncountable, of Lebesgue measure 0 and totally disconnected.

Let us review some other related works. In [1], the authors study the set

$$\mathcal{H} = \mathcal{H}(c) = \{\alpha \in \mathbb{R}^+ : D_n(\alpha, c) \geq 0 \text{ for all } n \geq 0\}$$

for $c = 1/k$. The characterizations of $\alpha \in \mathcal{H}$, via the partial quotients and convergents of α respectively, are obtained. Also, it is shown that \mathcal{H} is of positive Hausdorff dimension. If we set

$$\hat{O} = \hat{O}_c = \{\alpha \in O : (D_n(\alpha, c)) \text{ is bounded from below}\},$$

then it is direct from the definition that for $c = 1/k$, $\mathcal{H} \subset \hat{O}$. Comparing Theorem 1.2 and [1, Theorem 1], the relation between elements in \mathcal{H} and \hat{O} can be described as follows. For $\alpha \in \hat{O}$, let

$$m_1(\alpha) := \inf\{m \geq -1 : k \mid a_{m+2n} \text{ for all } n \geq 1\}.$$

The necessary and sufficient condition for α further in \mathcal{H} is $m_1(\alpha) = -1$.

In [11], some expressions for $\sup_{1 \leq n \leq q_m} D_n(\alpha, c)$ and $\inf_{1 \leq n \leq q_m} D_n(\alpha, c)$ are given respectively. From that, the sufficient condition in Theorem 1.2 (2) for D_n being bounded from above (resp. below) can be derived immediately. However, to obtain the necessary part, a more sophisticated analysis is needed. Besides the results in Theorem 1.2, in this paper we give a description on how the path $(D_n(\alpha, c))$ moves along with time n , that provides an intuition on the formulation of the one-sided boundedness.

The article is organized as follows. A recursive description of the moving mechanism of $(D_n(\alpha, c))$ is formulated in §2. Using this, we obtain in §3 a necessary and sufficient condition for α to have one-side bounded path via the convergents of α .

§4 is devoted to a pattern theory that leads to a necessary and sufficient condition via the partial quotients of α . As an application, in §5 we give certain topological properties to describe the size of the set of α with one-side bounded paths.

2. DESCRIPTION OF THE PATH

Fix $\alpha \in \mathbb{R}^+ \setminus \mathbb{Q}$. Let (p_n/q_n) and (a_n) be the convergents and partial quotients of α respectively and $I = (0, c)$, with $c = h/k \in (0, 1)$ ($h, k \in \mathbb{N}$ are coprime). The goal of this section is to give an intuitive description of the path $\xi_n := 1_{\{n\alpha \in I\}}$ and $D_n = \sum_{j=1}^n \xi_j - cn$.

It is well-known that $\left| \alpha - \frac{p_n}{q_n} \right| < \frac{1}{q_n q_{n+1}}$. The following lemma is an immediate consequence.

Lemma 2.1. *Each interval $\left(\frac{r}{q_n}, \frac{r+1}{q_n}\right)$, $r = 0, 1, \dots, q_n - 1$, contains exactly one point $\{j\alpha\}$ with $1 \leq j \leq q_n$. For each $1 \leq j \leq q_n$, there exists $0 \leq r < q_n$ such that*

$$\left\{ \{(lq_n + j)\alpha\} : 0 \leq l \leq \left\lfloor \frac{q_{n+1} - j}{q_n} \right\rfloor \right\} \subset \left(\frac{r}{q_n}, \frac{r+1}{q_n} \right).$$

Let λ_i^n ($i = 0, 1, \dots, q_n - 1$) be the unique integer $\in [1, q_n]$, such that

$$\lambda_i^n p_n \equiv i \pmod{q_n}.$$

In particular, $\lambda_0^n = q_n$. It follows from the basic properties of convergents that

$$\begin{cases} \text{when } n \text{ is odd, } \{\lambda_i^n \alpha\} \in \left(\frac{i-1}{q_n}, \frac{i}{q_n}\right) \forall i \in [1, q_n - 1] \text{ and } \{\lambda_0^n \alpha\} \in \left(\frac{q_n - 1}{q_n}, 1\right); \\ \text{when } n \text{ is even, } \{\lambda_i^n \alpha\} \in \left(\frac{i}{q_n}, \frac{i+1}{q_n}\right) \forall i \in [0, q_n - 1]. \end{cases}$$

When $k \mid q_n$, it is direct from Lemma 2.1 that

Proposition 2.2. *If $k \mid q_n$, then for $1 \leq j \leq q_n$ and $0 \leq l \leq \left\lfloor \frac{q_{n+1} - j}{q_n} \right\rfloor$,*

(1) (a) *when n is odd,*

$$\begin{cases} \xi_{\lambda_i^n} = 1, & \text{for } i \in [1, cq_n]; \\ \xi_{\lambda_i^n} = 0, & \text{for } i \in [cq_n + 1, q_n - 1] \cup \{0\}; \end{cases}$$

(b) *when n is even,*

$$\begin{cases} \xi_{\lambda_i^n} = 1, & \text{for } i \in [0, cq_n - 1]; \\ \xi_{\lambda_i^n} = 0, & \text{for } i \in [cq_n, q_n - 1]; \end{cases}$$

(2) $\xi_{lq_n + j} = \xi_j$; $D_{q_n} = 0$ and $D_{lq_n + j} = D_j$.

Now we turn to the case $k \nmid q_n$. Most ξ_j 's ($j = 1, \dots, q_{n+1}$) are clear by Lemma 2.1 except those j 's with $\{j\alpha\} \in \left(\frac{[cq_n]}{q_n}, \frac{[cq_n]+1}{q_n}\right)$. We only give the description of the path in the case n is even and it is analogous for the case n is odd. Note that when n is even, $\alpha > p_n/q_n$. We know that

$$\begin{aligned} j \in [1, q_{n+1}] \text{ and } \{j\alpha\} \in \left(\frac{[cq_n]}{q_n}, \frac{[cq_n]+1}{q_n}\right) \\ \iff j = \lambda_{[cq_n]}^n + lq_n, \text{ for some } 0 \leq l \leq \left\lfloor \frac{q_{n+1} - [cq_n]}{q_n} \right\rfloor. \end{aligned}$$

Note that

$$\left(\{(\lambda_{[cq_n]}^n + lq_n)\alpha\} : 0 \leq l \leq \left\lfloor \frac{q_{n+1} - \lambda_{[cq_n]}^n}{q_n} \right\rfloor \right)$$

is an increasing arithmetic sequence. Set

$$l_n := \inf \left\{ 0 \leq l \leq \left\lfloor \frac{q_{n+1} - \lambda_{[cq_n]}^n}{q_n} \right\rfloor : \{(\lambda_{[cq_n]}^n + lq_n)\alpha\} > c \right\},$$

with the convention that $\inf \emptyset = \infty$.

Proposition 2.3. *If n is even and $k \nmid q_n$, then for $1 \leq j \leq q_n$ and $0 \leq l \leq \left\lfloor \frac{q_{n+1} - j}{q_n} \right\rfloor$,*

$$(1) \quad \xi_{\lambda_{[cq_n]}^n} = 1 - 1_{\{l_n=0\}}, \text{ and}$$

$$\xi_{\lambda_i^n} = \begin{cases} 1, & \text{for } i \in [0, [cq_n] - 1]; \\ 0, & \text{for } i \in [[cq_n] + 1, q_n - 1]; \end{cases}$$

$$(2) \quad \xi_{lq_n + \lambda_{[cq_n]}^n} = 1 - 1_{\{l \geq l_n\}}; \text{ when } j \neq \lambda_{[cq_n]}^n, \quad \xi_{lq_n + j} = \xi_j; \text{ For the sequence } (D_n),$$

$$(a) \text{ when } l_n = 0, \quad D_{q_n} = -\left\{ \frac{hq_n}{k} \right\} \text{ and } D_{lq_n + j} = lD_{q_n} + D_j;$$

$$(b) \text{ when } l_n \geq 1, \quad D_{q_n} = 1 - \left\{ \frac{hq_n}{k} \right\} \text{ and}$$

$$D_{lq_n + j} = lD_{q_n} + D_j - 1_{\{l \geq l_n\}}(l - l_n + 1_{\{j \geq \lambda_{[cq_n]}^n\}}).$$

Proof. Most of the above results are straightforward. We only give an explanation for the value of D_{q_n} . It is simple to see from (1) that $|D_{q_n}| < 1$, and $D_{q_n} < 0$ if $l_n = 0$ while $D_{q_n} > 0$ if $l_n \geq 1$. Further note that

$$kD_{q_n} = k \sum_{j=1}^{q_n} \xi_j - hq_n \equiv -hq_n \pmod{k},$$

which leads to the conclusion. \square

By Proposition 2.3, the path of (D_n) has the following description. For n even and $k \nmid q_n$, define $\hat{D}^{(n)} := (\hat{D}_j^{(n)} : j = 1, \dots, q_n)$ (resp. $\check{D}^{(n)} := (\check{D}_j^{(n)} : j = 1, \dots, q_n)$) as:

$$\hat{D}_j^{(n)} = \sum_{i=1}^j \hat{\xi}_i - cj \quad \left(\text{resp. } \check{D}_j^{(n)} = \sum_{i=1}^j \check{\xi}_i - cj \right),$$

where for $i = 1, \dots, q_n$,

$$\hat{\xi}_{\lambda_i^n} = \begin{cases} 1, & \text{for } i \in [0, \lambda_{[cq_n]}^n]; \\ c - 1, & \text{for } i \in [\lambda_{[cq_n]}^n + 1, q_n - 1], \end{cases}$$

and $\check{\xi}_{\lambda_i^n} = \hat{\xi}_{\lambda_i^n} - 1_{\{i=[cq_n]\}}$. In particular,

$$\hat{D}_{q_n}^{(n)} = \check{D}_{q_n}^{(n)} + 1, \quad \hat{D}_{q_n}^{(n)} = 1 - \left\{ \frac{hq_n}{k} \right\} \text{ and } \check{D}_{q_n}^{(n)} = -\left\{ \frac{hq_n}{k} \right\}.$$

It follows from Proposition 2.3 that for $l \in [0, l_n)$,

$$(\xi_{lq_n + i} : i = 1, \dots, q_n) = (\hat{\xi}_j : i = 1, \dots, q_n),$$

and for $l \in [l_n, a_{n+1}]$,

$$(\xi_{lq_n+i} : i = 1, \dots, q_n \wedge (q_{n+1} - lq_n)) = (\check{\xi}_j : i = 1, \dots, q_n \wedge (q_{n+1} - lq_n)).$$

As a result, the path of (D_n) can be constructed by the translation of $\hat{D}^{(n)}$ and $\check{D}^{(n)}$. Precisely, we call the path $(D_n : n \in [(l-1)q_n + 1, lq_n])$ the l -th period before q_{n+1} , if $lq_n < q_{n+1}$. Then if $1 \leq l \leq l_n$, the l -th period is obtained by lifting the path, $(\hat{D}_j^{(n)} : 1 \leq j \leq q_n)$, $(l-1)D_{q_n}$ units up; if $l > l_n$, the l -th period is obtained by lifting the path, $(\check{D}^{(n)} : 1 \leq j \leq q_n)$, $(l_n \hat{D}_{q_n}^{(n)} + (l - l_n - 1)\check{D}_{q_n}^{(n)})$ units up. The path of $(D_n : 1 \leq n \leq q_{n+1})$ consists of a_{n+1} complete periods and an incomplete period which contains only q_{n-1} steps.

Now we directly give the analogous results for the case where n is odd and $k \nmid q_n$. Let

$$l_n := \inf \left\{ 0 \leq l \leq \left\lfloor \frac{q_{n+1} - \lambda_{[cq_n]+1}^n}{q_n} \right\rfloor : \{(lq_n + \lambda_{[cq_n]+1}^n)\alpha\} < c \right\},$$

where $\lambda_{q_n}^n := \lambda_0^n$, if $[cq_n] = q_n - 1$.

Proposition 2.3'. *If n is odd and $k \nmid q_n$, then for $1 \leq j \leq q_n$ and $0 \leq l \leq \left\lfloor \frac{q_{n+1}-j}{q_n} \right\rfloor$,*

$$(1) \quad \xi_{\lambda_{[cq_n]+1}^n} = 1_{\{l_n=0\}}, \text{ and}$$

$$\xi_{\lambda_i^n} = \begin{cases} 1, & \text{for } i \in [1, [cq_n]]; \\ 0, & \text{for } i \in [[cq_n] + 2, q_n - 1] \cup \{0\}; \end{cases}$$

$$(2) \quad \xi_{lq_n + \lambda_{[cq_n]+1}^n} = 1_{\{l \geq l_n\}}; \text{ when } j \neq \lambda_{[cq_n]+1}^n, \quad \xi_{lq_n+j} = \xi_j; \text{ For the sequence } (D_n),$$

$$(a) \text{ when } l_n = 0, \quad D_{q_n} = 1 - \left\{ \frac{hq_n}{k} \right\} \text{ and } D_{lq_n+j} = lD_{q_n} + D_j;$$

$$(b) \text{ when } l_n \geq 1, \quad D_{q_n} = - \left\{ \frac{hq_n}{k} \right\} \text{ and}$$

$$D_{lq_n+j} = lD_{q_n} + D_j + 1_{\{l \geq l_n\}}(l - l_n + 1_{\{j \geq \lambda_{[cq_n]+1}^n\}}).$$

In the next corollary, we compare the path (D_n) backwards from the q_{n+1} -th step and the q_{n-1} -th step (resp. the lq_n -th step and the q_n -th step) respectively, which is obtained directly from Proposition 2.3 (2) and will be frequently used in §3.

Corollary 2.4. *When n is even and $k \nmid q_n$,*

$$(1) \text{ for } j \in [0, q_{n-1}], \quad (D_0 := 0)$$

$$(D_{a_{n+1}q_n+j} - D_{q_{n+1}}) - (D_j - D_{q_{n-1}}) = \begin{cases} 1, & \text{if } 1 \leq l_n < \infty \text{ and } j < \lambda_{[cq_n]}^n \leq q_{n-1}; \\ 0, & \text{otherwise;} \end{cases}$$

$$(2) \text{ for } l \in [1, a_{n+1}] \text{ and } j \in [0, q_n],$$

$$(D_{(l-1)q_n+j} - D_{lq_n}) - (D_j - D_{q_n}) = \begin{cases} 1, & \text{if } 1 \leq l_n < \infty \text{ and } j < \lambda_{[cq_n]}^n \leq q_{n-1}; \\ 0, & \text{otherwise.} \end{cases}$$

3. ONE-SIDED BOUNDEDNESS

In this section, we will derive a necessary and sufficient condition on α with $(D_n(\alpha, c))$ one-side bounded. Define the level path of (D_n) as

$$\overline{D}_n := \sup\{D_j : 0 \leq j \leq n\} \text{ and } \underline{D}_n := \inf\{D_j : 0 \leq j \leq n\}.$$

Note that $D_n \in \frac{1}{k}\mathbb{N}$. Hence $\sup D_n < \infty$ (resp. $\inf D_n > -\infty$) if and only if \overline{D}_n increases (resp. \underline{D}_n decreases) only for finitely many times. First we give a useful lemma.

Lemma 3.1. (1) *If $D_{q_n} > 0$ for infinitely many n , then $\sup D_n = \infty$;*
 (2) *If $D_{q_n} < 0$ for infinitely many n , then $\inf D_n = -\infty$.*

Proof. We prove (1) only. Set

$$\lambda^n := \begin{cases} \lambda_{[cq_n]+1}^n, & \text{when } n \text{ is odd;} \\ \lambda_{[cq_n]}^n, & \text{when } n \text{ is even.} \end{cases}$$

Note that if $D_{q_n} > 0$, then $k \nmid q_n$ and $D_{q_n+j} > D_j$ for $j \in [1, q_{n-1} \wedge \lambda^n]$, and hence $\overline{D}_{q_{n+1}} > \overline{D}_{q_{n-1} \wedge \lambda^n}$. Now for the proof of $\sup D_n = \infty$, it suffices to show $\lambda^n \rightarrow \infty$, since $\lambda^n \rightarrow \infty$ implies \overline{D}_n increases for infinitely many times.

By definition, we readily check that when n is odd (resp. even), λ^n is the unique integer $\in [1, q_n]$, such that $\left\{\frac{\lambda^n p_n - 1}{q_n}\right\} < c < \left\{\frac{\lambda^n p_n}{q_n}\right\}$ (resp. $\left\{\frac{\lambda^n p_n}{q_n}\right\} < c < \left\{\frac{\lambda^n p_n + 1}{q_n}\right\}$). Fixing n , for $j = 1, \dots, \lambda^n$, $\left\{\frac{j p_N - 1}{q_N}\right\}$, $\left\{\frac{j p_N}{q_N}\right\}$ and $\left\{\frac{j p_N + 1}{q_N}\right\}$ all converge to $\{j\alpha\}$ as $N \uparrow \infty$. Hence, there exists $M > 0$, for $N > M$ and $j = 1, \dots, \lambda^n$,

$$\frac{1}{q_N} < \left\{\frac{j p_N}{q_N}\right\} < \frac{q_N - 1}{q_N} \text{ and } c \notin \left(\left\{\frac{j p_N - 1}{q_N}\right\}, \left\{\frac{j p_N + 1}{q_N}\right\}\right).$$

It follows that $\lambda^N > \lambda^n$, that leads to the conclusion. \square

A sequence (b_n) is called eventually a multiple of k if there exists N , such that for $n \geq N$, $k \mid b_n$. The next theorem gives a sufficient condition for the one-sided boundedness of (D_n) .

Theorem 3.2. *If (q_{2n}) (resp. (q_{2n+1})) is eventually a multiple of k , then $\sup D_n < \infty$ (resp. $\inf D_n > -\infty$) and $\inf D_n = -\infty$ (resp. $\sup D_n = \infty$).*

Proof. We only prove the case q_{2n+1} is eventually a multiple of k . Note that consecutive q_n must be coprime. So for n large enough, $k \nmid q_{2n}$. First we show $\inf D_n > -\infty$ and the following lemma is needed.

Lemma 3.3. (1) *When $k \mid q_n$, $\overline{D}_{q_{n+1}} = \overline{D}_{q_n}$ and $\underline{D}_{q_{n+1}} = \underline{D}_{q_n}$.*
 (2) *When n is even and $k \nmid q_n$, if $D_{q_{n+1}} \geq D_{q_{n-1}}$ and $D_{a_{n+1}q_n} \geq D_{q_n}$, then $\underline{D}_{q_{n+1}} = \underline{D}_{q_n}$.*

Proof. (1) is simple. For (2), by Corollary 2.4 (1), for $j \in [0, q_{n-1}]$

$$(3.1) \quad D_{a_{n+1}q_n+j} - D_j \geq D_{q_{n+1}} - D_{q_{n-1}} \geq 0.$$

In particular, taking $j = 0$, we get $D_{a_{n+1}q_n} \geq 0$. It follows from Proposition 2.3 (2a) that $l_n \geq 1$. Using Proposition 2.3 (2b) or the description of the path in §2,

$D_{lq_n} \geq D_{q_n}$ for $l = a_{n+1}$ is enough to ensure $D_{lq_n} \geq D_{q_n}$ for all $l = 1, \dots, a_{n+1}$. Hence by Corollary 2.4 (2), for $l \in [1, a_{n+1}]$ and $j \in [1, q_n]$,

$$D_{(l-1)q_n+j} - D_j \geq D_{lq_n} - D_{q_n} \geq 0.$$

Combined with (3.1), we readily see $\underline{D}_{q_{n+1}} = \underline{D}_{q_n}$. \square

Suppose $k \mid q_{2n-1}$ and $k \mid q_{2n+1}$. Then $\underline{D}_{q_{2n}} = \underline{D}_{q_{2n-1}}$ and $D_{q_{2n-1}} = D_{q_{2n+1}} = 0$. By Lemma 3.3, in order that $\underline{D}_{q_{2n+1}} = \underline{D}_{q_{2n}}$, it is enough to show $D_{a_{2n+1}q_{2n}} = 1$, since $D_{q_{2n}} < 1$. Using Proposition 2.3 (2), we readily see that $D_{q_{2n-1}} = D_{q_{2n+1}}$ implies $1 \leq l_{2n} < \infty$. By Corollary 2.4 (1), $D_{a_{2n+1}q_{2n}} = 1_{\{\lambda_{[cq_{2n}]}^{2n} \leq q_{2n-1}\}}$. Now it reduces to prove $\lambda_{[cq_{2n}]}^{2n} \leq q_{2n-1}$. We readily deduce from the definition that $\lambda_{[cq_{2n}]}^{2n}$ is the unique integer in $A := \{1 \leq j \leq q_{2n} : \{j\alpha\} < c\}$ such that

$$c - \{j\alpha\} = \min_{j \in A} (c - \{j\alpha\}).$$

Since $k \mid q_{2n-1}$, it is simple to see that $\lambda_{cq_{2n-1}}^{2n-1}$ also satisfies the above condition. Thus, $\lambda_{[cq_{2n}]}^{2n} = \lambda_{cq_{2n-1}}^{2n-1} \leq q_{2n-1}$ and we see that if $k \mid q_{2n-1}$ and $k \mid q_{2n+1}$, then $\underline{D}_{q_{2n+1}} = \underline{D}_{q_{2n}} = \underline{D}_{q_{2n-1}}$. It follows that if (q_{2n+1}) is eventually a multiple of k , then $\inf D_n > -\infty$.

Now we turn to the proof of $\sup D_n = \infty$.

Lemma 3.4. *For $\alpha \in \mathbb{R}^+ \setminus \mathbb{Q}$, let (p_n/q_n) and (a_n) be the convergents and partial quotients of α respectively. Then the following statements are equivalent for any $N \geq -1$ and $m \geq 2$.*

- (1) $m \mid q_{N+2n}$ for $n \geq 0$;
- (2) $m \mid q_N$ and $m \mid a_{N+2n}$ for $n \geq 1$.

Proof. (1) \Rightarrow (2): Suppose for $n \geq 0$, $m \mid q_{N+2n}$. In particular, $m \mid q_N$. Since consecutive q_n are coprime and

$$(3.2) \quad q_{N+2n+2} = a_{N+2n+2}q_{N+2n+1} + q_{N+2n},$$

$m \mid a_{N+2n}$ for all $n \geq 1$.

(2) \Rightarrow (1): Suppose $m \mid q_N$ and $m \mid a_{N+2n}$ for $n \geq 1$. Again using (3.2), we deduce inductively that $m \mid q_{N+2n}$ for $n \geq 0$. \square

It follows from Lemma 3.4 that the condition that q_{2n+1} is eventually a multiple of k implies that q_{2n} and k are coprime for large n , and a_{2n+1} is eventually a multiple of k . Note also that when q_{2n} and k are coprime, that $a_{2n+1} \geq k$ implies $l_{2n} \geq 1$. In fact, if $a_{2n+1} \geq k$, then

$$\alpha - \frac{p_{2n}}{q_{2n}} < \frac{1}{q_{2n}q_{2n+1}} < \frac{1}{kq_{2n}^2}.$$

It follows that

$$\lambda_{[cq_{2n}]}^{2n} \left(\alpha - \frac{p_{2n}}{q_{2n}} \right) < \frac{1}{kq_{2n}} \leq c - \frac{\lambda_{[cq_{2n}]}^{2n}}{q_{2n}} = \frac{h}{k} - \frac{\lambda_{[cq_{2n}]}^{2n}}{q_{2n}},$$

which implies $l_{2n} \geq 1$. Or equivalently, $D_{q_{2n}} > 0$ by Proposition 2.3 (2). Then $\sup D_n = \infty$ by Lemma 3.1. \square

Theorem 3.2 gives examples of α 's with $\sup D_n < \infty$ or $\inf D_n > -\infty$. Now we show that these α 's are exactly all the α 's with this condition.

Theorem 3.5. *If (q_{2n}) (resp. (q_{2n+1})) is not eventually a multiple of k , then $\sup D_n = \infty$ (resp. $\inf D_n = -\infty$).*

Proof. Suppose (q_{2n}) is not eventually a multiple of k . We shall prove $\sup D_n = \infty$. Note that $D_{q_n} \in \{-\frac{k-1}{k}, -\frac{k-2}{k}, \dots, \frac{k-2}{k}, \frac{k-1}{k}\}$, and $k \mid q_n$ implies $D_{q_{n-1}} = D_{q_{n+1}}$. It follows easily that there exist infinitely many n such that $k \nmid q_{2n}$ and $D_{q_{2n+1}} \geq D_{q_{2n-1}}$. By Corollary 2.4 (1), $D_{q_{2n+1}} > D_{q_{2n-1}}$ implies $\overline{D}_{q_{2n+1}} > \overline{D}_{q_{2n-1}}$. So we can focus on the case there exist infinitely many n such that $k \nmid q_{2n}$ and $D_{q_{2n+1}} = D_{q_{2n-1}}$. By Proposition 2.3 (2a), $D_{q_{2n+1}} = D_{q_{2n-1}}$ implies $1 \leq l_n < \infty$ and $D_{q_{2n}} > 0$. Hence, $\sup D_n = \infty$ follows from Lemma 3.1. \square

Now we have the following necessary and sufficient condition on α with D_n one-side bounded.

Theorem 3.6. *For $\alpha \in \mathbb{R}^+ \setminus \mathbb{Q}$, $c = h/k \in (0, 1)$, let (p_n/q_n) be the convergent of α . Then $\sup D_n < \infty$ (resp. $\inf D_n > -\infty$) if and only if (q_{2n}) (resp. (q_{2n+1})) is eventually a multiple of k .*

4. PATTERN DECOMPOSITION THEOREM

Theorem 3.6 gives a characterization of α with (D_n) one-side bounded, which is a condition imposed on convergents. Since the convergents are determined by the partial quotients, it is more natural to impose this condition equivalently on partial quotients.

From Lemma 3.4, we see that $\sup D_n < \infty$ (resp. $\inf D_n > -\infty$) is also equivalent to there exists even (resp. odd) $N > -1$ such that $k \mid q_N$ and $k \mid a_{N+2n}$ for $n \geq 1$. For an integer N , let $\pi_k(N)$ be the unique integer $\in \{0, 1, \dots, k-1\}$, such that $\pi_k(N) = N \pmod{k}$. In the following, we simply write π for π_k , since k is fixed. To answer the problem raised before, it suffices to characterize the $\pi(q_N)$ in terms of (a_0, a_1, \dots, a_N) .

Given an (ordered) tuple $M = (a_0, a_1, \dots, a_m)$ ($m \geq -1$) of non-negative integers, where m is called the length of M . When $m = -1$, M is the empty tuple $()$ (in the following, we will use the notation \mathcal{E} to represent the empty tuple). Define $(q_n^M : -2 \leq n \leq m)$ recursively by:

$$(4.1) \quad \begin{cases} q_{-2}^M = 1, q_{-1}^M = 0; \\ q_n^M = a_n q_{n-1}^M + q_{n-2}^M, \text{ for } 0 \leq n \leq m. \end{cases}$$

In other words, $(q_n^M : -2 \leq n \leq m)$ are the denominators of the convergents associated to $(a_n : 0 \leq n \leq m)$. What we are concerned is

$$\pi(q_{m-1}^M, q_m^M) := (\pi(q_{m-1}^M), \pi(q_m^M)).$$

Let's make a few definitions.

Definition 4.1. *For $M = (a_0, \dots, a_m)$,*

- (1) *A tuple with all elements $< k$ is called a pattern, and $\pi(M) := (\pi(a_0), \dots, \pi(a_m))$ is called the pattern of M .*
- (2) *The character of $M = (a_0, \dots, a_m)$ is defined by*

$$\kappa(M) = \pi(q_{m-1}^M, q_m^M).$$

- (3) *Operation of insertion: for tuples $M = (a_0, \dots, a_m)$, $N = (b_0, \dots, b_n)$, and $-1 \leq j \leq m$, define*

$$M \triangleleft_j N = (a_0, a_1, \dots, a_j, b_0, b_1, \dots, b_n, a_{j+1}, \dots, a_m),$$

i.e. the tuple induced by inserting N into M after a_j (or before a_0 if $j = -1$). The subscript j in \triangleleft_j may be omitted if we do not care where the tuple is inserted.

- (4) *If $M = M_0 \triangleleft N$, N is called a sub-tuple of M and proper if $N \neq M$. When N is a sub-tuple of M , we also say M contains N .*
- (5) *A non-empty tuple N is called elementary if N is null, in the sense that $\kappa(M) = \kappa(M \triangleleft_j N)$ for each tuple M and for each $-1 \leq j \leq m$, and minimal in the sense that N does not contain a proper null sub-tuple.*
- (6) *A tuple containing no elementary tuples is called prime.*

In particular, $\kappa(\mathcal{E}) = (1, 0)$. It is easy to see that $\kappa(M)$ depends only on the pattern $\pi(M)$. Hence as far as we are concerned, M is not different from $\pi(M)$.

Lemma 4.2. *Let $M = (a_0, \dots, a_m)$. If $m \geq (k^2 - 1)! - 1$, then M contains at least one elementary tuple with length $\leq (k^2 - 1)! - 1$.*

Proof. Note that since consecutive q_n^M are coprime, $\kappa(M) \neq (0, 0)$. Put

$$P_k = \{(N_1, N_2) : N_1, N_2 \in \{0, 1, \dots, k-1\} \text{ and } (N_1, N_2) \neq (0, 0)\}.$$

Then $\#P_k = k^2 - 1$. Denote $P_k = \{p_1, \dots, p_{k^2-1}\}$. Set

$$Q_k = \{(p_{\sigma(1)}, p_{\sigma(2)}, \dots, p_{\sigma(k^2-1)}) : \sigma \in S_{k^2-1}\},$$

where S_n is the collection of permutations on $\{1, 2, \dots, n\}$. Then $\#Q_k = (k^2 - 1)!$. For $a \in \mathbb{N}$, $p = (N_1, N_2) \in P_k$, define $T_a : P_k \rightarrow P_k$, $T_a((N_1, N_2)) = (N_2, \pi(aN_2 + N_1))$. It is easy to check T_a is a bijection. Hence for $q = (p_{\sigma(1)}, p_{\sigma(2)}, \dots, p_{\sigma(k^2-1)}) \in Q_k$,

$$T_a(q) := (T_a(p_{\sigma(1)}), T_a(p_{\sigma(2)}), \dots, T_a(p_{\sigma(k^2-1)})) \in Q_k.$$

For $M = (a_0, \dots, a_m)$ with $m \geq (k^2 - 1)! - 1$, fix any $q \in Q_k$ and consider the collection

$$C_k := (q, T_{a_0}(q), T_{a_1} \circ T_{a_0}(q), \dots, T_{a_m} \circ T_{a_{m-1}} \circ \dots \circ T_{a_0}(q)).$$

Since $\#C_k = m + 2 \geq (k^2 - 1)! + 1$, there exist $m_1 > m_2 \geq 0$, such that

$$T_{a_{m_1}} \circ \dots \circ T_{a_0}(q) = T_{a_{m_2}} \circ \dots \circ T_{a_0}(q).$$

Then it is easy to see $(a_{m_2+1}, a_{m_2+2}, \dots, a_{m_1})$ is elementary. \square

Corollary 4.3. (1) *There are only finite elementary or prime patterns.*

- (2) *For any $l \geq 0$, there exists $n \geq 1$, such that $\overbrace{(l, \dots, l)}^n$ is elementary.*

Proof. By Lemma 4.2, every pattern with length $\geq (k^2 - 1)!$ has a proper sub-tuple that is elementary. Hence, all the elementary or prime patterns have length

$\leq (k^2 - 1)! - 1$, which implies (1). For (2), we take $M = \overbrace{(l, \dots, l)}^N$ for some $N \geq (k^2 - 1)! - 1$. Then M contains an elementary tuple. Note that all the

sub-tuple of M is in the form $\overbrace{(l, \dots, l)}^n$. That concludes (2). \square

We now present a decomposition theorem.

Theorem 4.4. Let $M = (a_0, a_1, \dots, a_m)$.

- (1) For general M , there exists a prime tuple M_0 , and finite elementary tuples $(N_i : 1 \leq i \leq n)$ such that

$$(4.2) \quad M = M_0 \triangleleft N_1 \triangleleft N_2 \cdots \triangleleft N_n,$$

where the inserting operations are executed from left to right. This expression is called a prime decomposition of M .

- (2) $\kappa(M) = \kappa(M_0)$.

In general, the decomposition (4.2) is not unique, and neither is the prime tuple M_0 . The character of M is determined by any one of the prime tuples M_0 , which are hence called the prime tuples of M . A tuple (resp. pattern) M is called of type- k , if $\kappa(M) = (j, 0)$, for some $j = 1, \dots, k-1$. According to Corollary 4.3 (1), there are only finite prime patterns of type- k . In particular, \mathcal{E} is a prime pattern of type- k . Intuitively, all the patterns of type- k can be constructed by inserting finite elementary patterns one by one to an initial prime pattern of type- k .

Come back to partial quotients (a_n) . Combining previous results, we get

Theorem 4.5. $\sup D_n < \infty$ (resp. $\inf D_n > -\infty$) if and only if there exists even (resp. odd) $m \geq -1$ such that (a_0, a_1, \dots, a_m) is of type- k and $k \mid a_{m+2n}$ for all $n \geq 1$.

Example 4.6. In this example, we will explicitly calculate all the elementary patterns, prime patterns and prime patterns of type- k when $k = 2$.

Proposition 4.7. When $k = 2$,

- (1) A tuple M is elementary if and only if its pattern is one of the following 7 patterns (comma ‘,’s in pattern are omitted for simplicity): (00) , (111) , (0101) , (1010) , (011011) , (110110) , (101101) .
- (2) A tuple M is prime if and only if its pattern is one of the following 16 patterns: $\mathcal{E}, (0), (1), (11), (10), (01), (101), (110), (011), (010), (0110), (1101), (1011), (01101), (11011), (10110)$.

Proof. We will separate the proof into several steps. For the tuple $M = (a_0, \dots, a_m)$, we set $M^h = (a_0, \dots, a_h)$, for $-1 \leq h \leq m$. Our first step is to build the relation between $\kappa(M^h)$ and $\kappa(M^{h+1})$. Using (4.1), the relation is shown by the following two tables: for $-1 \leq h \leq m-1$,

- (1) when $\pi(a_{h+1}) = 1$,

$\kappa(M^h)$	$\kappa(M^{h+1})$
(11)	(10)
(10)	(01)
(01)	(11)

- (2) when $\pi(a_{h+1}) = 0$,

$\kappa(M^h)$	$\kappa(M^{h+1})$
(11)	(11)
(10)	(01)
(01)	(10)

Next we describe this relation algebraically. Digitize patterns first by defining a bijection $\Phi : \{(11), (10), (01)\} \rightarrow \mathbb{Z}/3$,

$$\Phi((11)) = 0, \quad \Phi((10)) = 1, \quad \Phi((01)) = 2.$$

Then the tables above can be written into

- (1) when $\pi(a_{h+1}) = 1$, $\kappa(M^{h+1}) = \Phi^{-1}(\Phi(\kappa(M^h)) + 1)$;
- (2) when $\pi(a_{h+1}) = 0$, $\kappa(M^{h+1}) = \Phi^{-1}(2\Phi(\kappa(M^h)))$.

To combine these two cases, we define $T_1, T_0 : \mathbb{Z}/3 \rightarrow \mathbb{Z}/3$, by

$$T_1(x) = x + 1 \pmod{3}, \quad T_0(x) = 2x \pmod{3}, \quad x \in \mathbb{Z}/3$$

and for any integer y , $T_y = T_{\pi(y)}$. With these notations, we have the parity relation

$$\kappa(M^{h+1}) = \Phi^{-1}(T_{a_{h+1}}(\Phi(\kappa(M^h)))) = \Phi^{-1} \circ T_{a_{h+1}} \circ \Phi(\kappa(M^h)), \quad h \geq -1,$$

or $\Phi(\kappa(M^{h+1})) = T_{a_{h+1}} \circ \Phi(\kappa(M^h))$. Hence it follows recursively that

$$(4.3) \quad \Phi(\kappa(M)) = T_{a_m} \circ T_{a_{m-1}} \circ \cdots \circ T_{a_0} \circ \Phi(\kappa(\mathcal{E})),$$

where $\Phi(\kappa(M)) = 0, 1$ or 2 is called the character value. Define two mappings T_M and $T_{\pi(M)}$ as

$$T_M = T_{a_m} \circ \cdots \circ T_{a_0} = T_{\pi a_m} \circ \cdots \circ T_{\pi a_0} = T_{\pi(M)},$$

and note that $\Phi(\kappa(\mathcal{E})) = 1$.

Lemma 4.8. *The character $\kappa(M)$ of a tuple M satisfies*

$$\Phi(\kappa(M)) = T_M \circ \Phi(\kappa(\mathcal{E})) = T_M(1).$$

Now the second step is to apply the relation to the case in Proposition 4.7. Note that for $N = (b_0, \dots, b_n)$ and $-1 \leq k \leq m$,

$$M \triangleleft_k N = (a_0, a_1, \dots, a_k, b_0, b_1, \dots, b_n, a_{k+1}, a_{k+2}, \dots, a_m).$$

Using the relation (4.3) above,

$$\begin{aligned} \Phi(\kappa(M \triangleleft_k N)) &= T_{a_m} \circ T_{a_{m-1}} \circ \cdots \circ T_{a_{k+1}} \circ T_{b_n} \circ T_{b_{n-1}} \\ &\quad \circ \cdots \circ T_{b_0} \circ T_{a_k} \circ T_{a_{k-1}} \circ \cdots \circ T_{a_0} (\Phi(\kappa(M^0))) \\ &= T_{(a_{k+1}, \dots, a_m)} \circ T_N \circ T_{(a_0, \dots, a_k)} \circ \Phi(\kappa(M^0)). \end{aligned}$$

Hence, it is easy to see that

Lemma 4.9. *The tuple N is null if and only if T_N is an identity.*

The third step is to use this lemma to prove Proposition 4.7. That is only a simple algebraic calculation by Lemma 4.9. We will do it according to the length of the tuples. Two tricks will make it easier. For $-1 \leq h \leq m-1$,

- (a) if M^h is elementary, M^{h+1} is neither elementary nor prime;
- (b) if M^h is not prime, M^{h+1} is neither elementary nor prime.

Now assume $M = (a_0, \dots, a_m)$,

- (i) when $m = -1$: \mathcal{E} is prime;
- (ii) when $m = 0$: (0), (1). Not null, all prime;
- (iii) when $m = 1$: (00), (01), (11), (10). Only (00) is elementary,

$$T_{(00)} = T_0 \circ T_0(x) = 2 \cdot (2x) = 4x \equiv x \pmod{3}.$$

The other three are prime.

- (iv) when $m = 2$: 6 patterns (010), (011), (110), (111), (100), (101) need to check. (010), (011), (110), (101) are prime. Only (111) is elementary, $T_{(111)} = T_1 \circ T_1 \circ T_1(x) = x + 1 + 1 + 1 = x + 3 \equiv x \pmod{3}$.

- (v) when $m = 3$: 8 patterns (1101), (1100), (1011), (1010), (0111), (0110), (0101), (0100) need to check. Among them, (1101), (1011), (0110) are prime and the following are elementary,
 $T_{(0101)} = T_1 \circ T_0 \circ T_1 \circ T_0(x) = 2(2x + 1) + 1 = 4x + 3 \equiv x \pmod{3}$,
 $T_{(1010)} = T_0 \circ T_1 \circ T_0 \circ T_1(x) = 2(2(x + 1) + 1) = 4x + 6 \equiv x \pmod{3}$.
- (vi) when $m = 4$: 6 patterns (11011), (11010), (10111), (10110), (01101), (01100) need to check. Among them, (11011), (10110), (01101) are prime and none is elementary.
- (vii) when $m = 5$: 6 patterns (110111), (110110), (101101), (101100), (011010), (011011) need to check. Among them, 3 patterns (110111), (101100), (011010) contain smaller elementary patterns, so they are neither elementary nor prime. The other three are elementary,
 $T_{(011011)} = T_1 \circ T_1 \circ T_0 \circ T_1 \circ T_1 \circ T_0(x) = 4x + 6 \equiv x \pmod{3}$.
 $T_{(110110)} = T_0 \circ T_1 \circ T_1 \circ T_0 \circ T_1 \circ T_1(x) = 4x + 12 \equiv x \pmod{3}$.
 $T_{(101101)} = T_1 \circ T_0 \circ T_1 \circ T_1 \circ T_0 \circ T_1(x) = 4x + 9 \equiv x \pmod{3}$.
- (viii) when $m \geq 6$: neither elementary nor prime.

Here only verification for 7 elementary patterns are presented. The verification of other patterns is similar, for example, $T_{(11)} = T_1 \circ T_1(x) = x + 2 \not\equiv x \pmod{3}$, so it is not null. That completes the proof. \square

By definition, a prime pattern M is of type-2 if and only if $\kappa(M) = (10)$, i.e., $\Phi(\kappa(M)) = 1$. Total 16 prime patterns in Proposition 4.7 may be classified according to their character values, which are calculated by the formula in Lemma 4.8:

$$\begin{aligned} \Phi(\kappa(\mathcal{E})) &= 1, \\ \Phi(\kappa(0)) &= 2, & \Phi(\kappa(1)) &= 2, \\ \Phi(\kappa(11)) &= 0, & \Phi(\kappa(10)) &= 1, & \Phi(\kappa(01)) &= 0, \\ \Phi(\kappa(101)) &= 2, & \Phi(\kappa(110)) &= 0, & \Phi(\kappa(011)) &= 1, & \Phi(\kappa(010)) &= 0, \\ \Phi(\kappa(0110)) &= 2, & \Phi(\kappa(1101)) &= 1, & \Phi(\kappa(1011)) &= 0, \\ \Phi(\kappa(01101)) &= 0, & \Phi(\kappa(11011)) &= 2, & \Phi(\kappa(10110)) &= 0. \end{aligned}$$

Proposition 4.10. *Let M_0 be any prime tuple of M . Then M is a tuple of type-2 if and only if $M_0 \in \{\mathcal{E}, (10), (011), (1101)\}$.*

5. SIZE OF O

Fix $c = h/k \in (0, 1)$. Recall that

$$O = O_c = \{\alpha \in \mathbb{R}^+ \setminus \mathbb{Q} : (D_n(\alpha, c)) \text{ is one-side bounded}\}.$$

We can readily derived from Theorem 4.5 that O is uncountable. To exhibit the application of Theorem 4.5 further, we shall use it to explore the size of O . We will obtain topological properties as shown in Theorem 1.3.

For any $m \geq 0$ and a tuple $B_m = (b_0, b_1, \dots, b_m)$, define

$$(5.1) \quad I(B_m) = \{\alpha \in \mathbb{R}^+ \setminus \mathbb{Q} : a_i(\alpha) = b_i, \text{ for } i = 0, 1, \dots, m\},$$

which is called a fundamental interval. It is known that $I(B_m)$ is an irrational interval, i.e. the intersection of an interval and $\mathbb{R}^+ \setminus \mathbb{Q}$.

Lemma 5.1. *For any tuple B_m , $I(B_m) \cap O \neq \emptyset$.*

Proof. Fix a tuple $B_m = (b_0, b_1, \dots, b_m)$. By Theorem 4.5, it suffices to construct a tuple of type- k with length $\geq m + 1$ and its first $m + 1$ numbers being b_0, b_1, \dots, b_m . We will prove this recursively.

The statement for $m = 1$ is already proved in Corollary 4.3 (2). Assume the above statement holds when $m = n$. When $m = n + 1$, for the first n of b_i 's, by inductive assumption, there exists a tuple $(d_i : 1 \leq i \leq r)$, such that $(b_0, \dots, b_n, d_1, \dots, d_r)$ is a tuple of type- k . For b_{n+1} , by Corollary 4.3 (2), there

exists $j > 0$ such that $\overbrace{(b_{n+1}, \dots, b_{n+1})}^j$ is elementary. Then

$$(b_0, \dots, b_n, \overbrace{b_{n+1}, \dots, b_{n+1}}^j, d_1, \dots, d_r)$$

is a tuple of type- k . That concludes the proof. \square

Corollary 5.2. *O is dense in \mathbb{R}^+ .*

Next, we will show that the Hausdorff dimension of O lies in $(0, 1)$. Define for any $m \geq -1$ and a tuple $B_m = (b_0, b_1, \dots, b_m)$, (recall that $B_{-1} = \mathcal{E}$)

$$(5.2) \quad F(B_m) := \{y \in \mathbb{R}^+ \setminus \mathbb{Q} : a_i(y) = b_i, \text{ for } i = 0, 1, \dots, m, \\ \text{and } k \mid a_{m+2n}(y) \text{ for any } n \geq 1\}.$$

It is known from Theorem 4.5 that

$$(5.3) \quad O = \bigcup_{m \geq 0} \bigcup_{\substack{b_0, b_1, \dots, b_m \in \mathbb{N}: \\ (b_0, \dots, b_m) \text{ is a tuple of type-}k}} F(b_0, b_1, \dots, b_m).$$

As shown in [5, Lemma 1], all subsets $F(B_m)$ have the same Hausdorff dimension. Hence it suffices to prove $\dim_H(F) \in (0, 1)$, where

$$F := F(\mathcal{E}) = \{\alpha \in \mathbb{R}^+ \setminus \mathbb{Q} : k \mid a_{2n+1}(\alpha) \text{ for any } n \geq 0\}.$$

Note that

$$F \supset \{\alpha \in \mathbb{R}^+ \setminus \mathbb{Q} : a_n(\alpha) \in \{k, 2k\} \text{ for any } n \geq 0\},$$

where the Hausdorff dimension of the set on the right hand side is known (cf. [5, Theorem 11]) and in particular it is positive. Hence $\dim_H(F) > 0$.

Now we give an estimation for upper bound.

Lemma 5.3. $\dim_H(F) < 1$.

Proof. F is the intersection of a decreasing sequence of sets $\{E_l : l \geq 0\}$, where

$$E_l = \{\alpha \in \mathbb{R}^+ \setminus \mathbb{Q} : k \mid a_{2n+1}(\alpha) \text{ for any } 0 \leq n \leq l\}.$$

Note that each E_l is a countable union of disjoint irrational intervals. More precisely, for $l \geq 0$ and $d_0, d_1, \dots, d_l \in k\mathbb{N}$, set

$$E(d_0, d_1, \dots, d_l) = \{\alpha \in (0, 1) \setminus \mathbb{Q} : a_{2n+1}(\alpha) = d_n \text{ for any } 0 \leq n \leq l\},$$

and then

$$(5.4) \quad E_l = \bigcup_{d_0, d_1, \dots, d_l \in k\mathbb{N}} E(d_0, d_1, \dots, d_l).$$

Since $E_l \supset F$, $\bigcup_{d_0, d_1, \dots, d_l \in k\mathbb{N}} E(d_0, d_1, \dots, d_l)$ is a countable cover of F . We will use the following fact, which can be easily derived by [9, (57)], to estimate the c -content of this cover for $c \in (0, 1)$. For $l \geq 1$ and $d_0, d_1, \dots, d_l \in k\mathbb{N}$,

$$(5.5) \quad \frac{1}{3d_l^2} < \frac{|E(d_0, d_1, \dots, d_l)|}{|E(d_0, d_1, \dots, d_{l-1})|} < \frac{2}{d_l^2}.$$

Repeatedly using (5.5), we have

$$|E(d_0, d_1, \dots, d_l)| < 2^l \prod_{r=1,2,\dots,l} \frac{1}{d_r^2}.$$

Following this, we get the estimation for the c -content of the above cover:

$$(5.6) \quad \begin{aligned} \sum_{d_0, d_1, \dots, d_l \in k\mathbb{N}} |E(d_0, d_1, \dots, d_l)|^c &< 2^{cl} \sum_{d_0, d_1, \dots, d_l \in k\mathbb{N}} \prod_{r=1,2,\dots,l} \frac{1}{d_r^{2c}} \\ &= \left[(2/k^2)^c \sum_{j \geq 1} \frac{1}{j^{2c}} \right]^l \leq \left(2^{-c} \sum_{j \geq 1} \frac{1}{j^{2c}} \right)^l. \end{aligned}$$

Let

$$g(c) = 2^{-c} \sum_{j \geq 1} \frac{1}{j^{2c}}, \quad c \in (1/2, 1].$$

It is easy to see that g is continuous and decreasing in $(1/2, 1]$, with $g(1) = \pi^2/12 < 1$ and $\lim_{c \rightarrow \frac{1}{2}^+} g(c) = \infty$. It follows that there exists a $c^* \in (0, 1)$ such that $g(c^*) < 1$,

which implies the c^* -content in (5.6) $< (g(c^*))^l < 1$. Since

$$\lim_{l \rightarrow \infty} \sup_{d_0, d_1, \dots, d_l \in k\mathbb{N}} |E(d_0, d_1, \dots, d_l)| = 0,$$

the c^* -Hausdorff measure $\mathcal{H}^{c^*}(F) \leq 1$ and $\dim_H(F) \leq c^* < 1$. \square

Now we have $\dim_H(F) \in (0, 1)$. It follows that

Theorem 5.4. $\dim_H(O) \in (0, 1)$. Hence, O is uncountable, of Lebesgue measure 0 and totally disconnected.

At last, we give a description of the size of O in the view of category.

Theorem 5.5. O is of 1st category.

Proof. We again use the same notation $F(B_m)$ defined in (5.2). By (5.3), O is covered by a countable union of $(F(B_m))$. It suffices to show that every $F(B_m)$ is nowhere dense. We have the following characterization of the closure of $F(B_m)$.

Lemma 5.6. For $B_m = (b_0, b_1, \dots, b_m)$,

$$\overline{F(B_m)} = F(B_m) \cup Q(B_m),$$

where

$$Q(B_m) = \{q \in \mathbb{Q} : q = [b_0; b_1 \cdots b_m d_{m+1} d_{m+2} \cdots d_{m+l}] \text{ for some } l \geq 1 \text{ and positive integers } d_{m+1}, \dots, d_{m+l}, \text{ where } k \mid d_{m+2n} \text{ for } 1 \leq n \leq [l/2]\}.$$

Proof. First we shall prove $\overline{F(B_m)} \supset Q(B_m)$. In fact, for any fixed $q = [b_0; b_1 \cdots b_m d_{m+1} \cdots d_{m+l}] \in Q(B_m)$, we can take for $r \geq 1$,

$$q_r = (b_0, \dots, b_m, d_{m+1}, \dots, d_{m+l}, rk, rk, \dots, rk, \dots).$$

It is easy to see $q_r \in F(B_m)$ and $q_r \rightarrow q$ as $r \uparrow \infty$ and hence $q \in \overline{F(B_m)}$.

For another direction, it suffices to prove that for any $\alpha \notin F(B_m) \cup Q(B_m)$, there exists an interval I containing α such that $I \cap F(B_m) = \emptyset$. Indeed, for such α , we can always find a fundamental interval associated to α satisfying the condition. This concludes the proof. \square

Proof of Theorem 5.5 continued. By Lemma 5.6, $\overline{F(B_m)} = F(B_m) \cup Q(B_m)$. Recall in the proof of Lemma 5.3, we have shown that $\dim_H(F(B_m)) < 1$. Since $Q(B_m) \subseteq \mathbb{Q}$, $\dim_H(Q(B_m)) = 0$. It follows that $\dim_H(\overline{F(B_m)}) < 1$. Hence $\overline{F(B_m)}$ is a totally disconnected closed set, which is then nowhere dense. \square

REFERENCES

- [1] M. Boshernitzan and D. Ralston. Continued fractions and heavy sequences. *Proceedings of the American Mathematical Society*, 137(10):3177–3185, 2009.
- [2] Y. Dupain. Intervalles a restes majorés pour la suite. *Acta Mathematica Academiae Scientiarum Hungaricae*, 29(3-4):289–303, 1977.
- [3] Y. Dupain and V. T Sós. On the one-sided boundedness of discrepancy-function of the sequence $\{n\alpha\}$. *Acta Arithmetica*, 37:363–374, 1980.
- [4] H. Furstenberg, H. Keynes, and L. Shapiro. Prime flows in topological dynamics. *Israel Journal of Mathematics*, 14(1):26–38, 1973.
- [5] I. J. Good. The fractional dimensional theory of continued fractions. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 37, pages 199–228. Cambridge University Press, 1941.
- [6] G. Halász. Remarks on the remainder in Birkhoff’s ergodic theorem. *Acta Mathematica Academiae Scientiarum Hungarica*, 28(3-4):389–395, 1976.
- [7] E. Hecke. Über analytische Funktionen und die Verteilung von Zahlen mod. eins. In *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, volume 1, pages 54–76. Springer, 1922.
- [8] H. Kesten. On a conjecture of Erdős and Szüs related to uniform distribution mod 1. *Acta Arithmetica*, 12:193–212, 1966.
- [9] A. I. Khinchin. *Continued fractions*, volume 525. P. Noordhoff, 1963.
- [10] A. Ostrowski. Bemerkungen zur Theorie der diophantischen Approximationen. In *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, volume 1, pages 77–98. Springer, 1922.
- [11] L. Roçadas and J. Schoißengeier. On the local discrepancy of $(n\alpha)$ -sequences. *Journal of Number Theory*, 131(8):1492–1497, 2011.
- [12] V. T Sós. On the discrepancy of the sequence $\{n\alpha\}$. *Coll. Math. Soc. J. Bolyai*, 13:359–367, 01 1976.

SCHOOL OF MATHEMATICS, FUDAN UNIVERSITY, SHANGHAI 200433, CHINA.

Email address: jgying@fudan.edu.cn

SHANGHAI CENTER FOR MATHEMATICAL SCIENCES, FUDAN UNIVERSITY, SHANGHAI 200433, CHINA.

Email address: yszheng666@gmail.com