

From Fourier series to infinite product representations of π and infinite-series forms for its positive powers

Jean-Christophe PAIN*

CEA, DAM, DIF, F-91297 ArpaJon, France
 Université Paris-Saclay, CEA, Laboratoire Matière en Conditions Extrêmes,
 91680 Bruyères-le-Châtel, France

October 18, 2022

Abstract

In this article, we derive, using Fourier series and multiple derivative of the function $\pi/\sin(\pi x)$, series representations for positive powers of π . We also show that the Euler-Wallis product can be easily obtained from the same formalism and deduce infinite products representations of π .

1 Introduction

Let us set $f(x) = \cos(\alpha x)$. The Fourier transform \mathcal{F} of f satisfies $\mathcal{F}(x) = f(x)$ in $[-\pi, \pi]$, $\mathcal{F}(x + 2\pi) = \mathcal{F}(x)$, \mathcal{F} is an odd function, \mathcal{C}^∞ and piecewise continuous. One has [1, 2]:

$$\mathcal{F}(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos(nx), \quad (1)$$

where

$$a_n = \frac{2}{\pi} \int_0^\pi \cos(\alpha t) \cos(nt) dt = \frac{(-1)^n}{\pi} \sin(\alpha\pi) \left(\frac{1}{\alpha + n} + \frac{1}{\alpha - n} \right). \quad (2)$$

Setting $x = 0$ and $x = \pi$, one gets respectively

$$\boxed{\frac{\pi}{\sin(\alpha\pi)} = \sum_{n=-\infty}^{\infty} \frac{(-1)^n}{\alpha + n}} \quad (3)$$

and

$$\boxed{\pi \cotan(\alpha\pi) = \sum_{n=-\infty}^{\infty} \frac{1}{\alpha + n}.} \quad (4)$$

*jean-christophe.pain@cea.fr

2 Infinite series representation for π^k with $k \geq 0$ involving multinomial coefficients

Starting from

$$\frac{\pi}{\sin(\pi x)} = \sum_{n=-\infty}^{\infty} \frac{(-1)^n}{x+n}, \quad (5)$$

with $x = 1/4$, one obtains

$$\pi = 2\sqrt{2} \left[1 + \sum_{n=-\infty}^{\infty} \frac{(-1)^n}{(1+4n)} \right]. \quad (6)$$

Using the Fàa di Bruno formula for the multiple derivative of a composite function (and in particular here of the inverse of a function):

$$\left(\frac{1}{f(x)} \right)^{(k)} = \frac{k!}{[f(x)]^{k+1}} \sum_{\substack{p_0+p_1+p_2+\dots+p_k=k \\ p_1+2p_2+\dots+kp_k=k}} \mathcal{C}(p_1, p_2, \dots, p_k) \prod_{i=0}^k [f^{(i)}(x)]^{p_i}, \quad (7)$$

with

$$\mathcal{C}(p_1, p_2, \dots, p_k) = \frac{(-1)^{k-p_0} (k-p_0)!}{\prod_{i=1}^k (i!)^{p_i} p_i!}. \quad (8)$$

$\mathcal{C}(p_1, p_2, \dots, p_k)$ can be expressed in terms of the multinomial coefficient

$$\mathcal{M}(p_1, p_2, \dots, p_k) = \frac{(p_1 + p_2 + \dots + p_k)!}{p_1! p_2! \dots p_k!} = \frac{(k-p_0)!}{p_1! p_2! \dots p_k!} \quad (9)$$

as

$$\mathcal{C}(p_1, p_2, \dots, p_k) = \frac{(-1)^{k-p_0}}{\prod_{i=1}^k (i!)^{p_i}} \mathcal{M}(p_1, p_2, \dots, p_k). \quad (10)$$

With $f(x) = \sin(\pi x)$, we get

$$f^k(x) = \pi^k \sin\left(\pi x + k\frac{\pi}{2}\right). \quad (11)$$

Thus, we obtain

$$\frac{\partial^k}{\partial x^k} \left[\frac{1}{\sin(\pi x)} \right] = \pi^k \frac{k!}{[\sin(\pi x)]^{k+1}} \sum_{\substack{p_0+p_1+p_2+\dots+p_k=k \\ p_1+2p_2+\dots+kp_k=k}} \mathcal{C}(p_1, p_2, \dots, p_k) \prod_{i=0}^k \left[\sin\left(\pi x + i\frac{\pi}{2}\right) \right]^{p_i} \quad (12)$$

and since

$$\frac{\partial^k}{\partial x^k} \left[\sum_{n=-\infty}^{\infty} \frac{(-1)^n}{x+n} \right] = (-1)^k k! \sum_{n=-\infty}^{\infty} \frac{(-1)^n}{(x+n)^{k+1}} \quad (13)$$

we get the final result

$$\boxed{\pi^{k+1} = \frac{(-1)^k}{\mathcal{B}_k(x)} \sum_{n=-\infty}^{\infty} \frac{(-1)^n}{(x+n)^{k+1}}} \quad (14)$$

with

$$\mathcal{B}_k(x) = \frac{1}{[\sin(\pi x)]^{k+1}} \sum_{\substack{p_0+p_1+p_2+\dots+p_k=k \\ p_1+2p_2+\dots+kp_k=k}} \mathcal{C}(p_1, p_2, \dots, p_k) \prod_{i=0}^k \left[\sin\left(\pi x + i\frac{\pi}{2}\right) \right]^{p_i}. \quad (15)$$

3 Particular cases

Let us first consider the case $k = 0$. All the p_i are equal to zero,

$$\mathcal{B}_0(x) = \frac{1}{\sin(\pi x)} \quad (16)$$

and we recover

$$\pi = \sin(\pi x) \sum_{n=-\infty}^{\infty} \frac{(-1)^n}{(x+n)}. \quad (17)$$

In the case $k = 1$, since $p_0 + p_1 = 1$ and $p_1 = 1$, we have necessarily $p_0 = 0$ and

$$\mathcal{B}_1(x) = \frac{\cos(\pi x)}{\sin^2(\pi x)} \quad (18)$$

yielding

$$\pi^2 = \frac{\sin^2(\pi x)}{\cos(\pi x)} \sum_{n=-\infty}^{\infty} \frac{(-1)^n}{(x+n)^2}. \quad (19)$$

For $k = 2$, since $p_0 + p_1 + p_2 = 2$ and $p_1 + 2p_2 = 2$, we have two possibilities:

- $p_0 = 0, p_1 = 2, p_2 = 0$, yielding the term

$$\frac{\cos^2(\pi x)}{\sin^3(\pi x)} \quad (20)$$

in the summation in \mathcal{B}_2 and

- $p_0 = 1, p_1 = 0, p_2 = 1$, responsible for the term

$$\frac{1}{2 \sin(\pi x)}, \quad (21)$$

leading to

$$\mathcal{B}_2(x) = \frac{1}{\sin^3(\pi x)} - \frac{1}{2 \sin(\pi x)}, \quad (22)$$

and finally

$$\pi^3 = \frac{2 \sin^3(\pi x)}{2 - \sin^2(\pi x)} \sum_{n=-\infty}^{\infty} \frac{(-1)^n}{(x+n)^3}. \quad (23)$$

4 Product formulas for π

Let us start with a simple proof of the so-called Euler-Wallis formula

$$\boxed{\frac{\sin(\pi\alpha)}{\pi\alpha} = \prod_{n=1}^{\infty} \left(1 - \frac{\alpha^2}{n^2}\right)}, \quad (24)$$

valid $\forall \alpha \in \mathbb{R} \setminus \mathbb{Z}^*$. Setting [3]

$$\mathcal{S}_n(x) = \frac{1}{x+n} + \frac{1}{x-n}, \quad (25)$$

one has, for $n \geq 1$,

$$\mathcal{S}_n(x) = \frac{2x}{x^2 - n^2} \quad (26)$$

and

$$|\mathcal{S}_n(x)| \leq \frac{2x_0}{x_0^2 - n^2} \quad (27)$$

if $|x| \leq x_0$ and $n > x_0$ with $x_0 > 0$. The series

$$\sum_{n=1}^{\infty} |\mathcal{S}_n(x)|^2 \quad (28)$$

converges normally in every segment included in $\mathbb{R} \setminus \mathbb{Z}^*$. The series

$$\sum_{n=1}^{\infty} \ln \left| 1 - \frac{x^2}{n^2} \right| \quad (29)$$

is therefore derivable term by term since

$$\frac{d}{dx} \ln \left| 1 - \frac{x^2}{n^2} \right| = \mathcal{S}_n(x). \quad (30)$$

Subsequently, since

$$\pi \cotan(\pi x) - \frac{1}{x} = \sum_{n=-\infty}^{\infty} \frac{1}{x+n} - \frac{1}{x} = \sum_{n=1}^{\infty} \mathcal{S}_n(x), \quad (31)$$

we get

$$\frac{d}{dx} \sum_{n=1}^{\infty} \ln \left| 1 - \frac{x^2}{n^2} \right| = \sum_{n=1}^{\infty} \mathcal{S}_n(x) = \pi \cotan(\pi x) - \frac{1}{x}. \quad (32)$$

Setting

$$g(x) = \prod_{n=1}^{\infty} \left(1 - \frac{x^2}{n^2}\right) \quad (33)$$

and

$$h(x) = \frac{\sin(\pi x)}{\pi x}, \quad (34)$$

we have that g and h are proportional in each interval of $\mathbb{R} \setminus \mathbb{Z}^*$. Since

$$h(0) = g(0) = 1, \quad (35)$$

we find that $h \equiv g$ over $] - 1, 1[$. Let us now introduce

$$g_N(x) = \prod_{n=1}^N \left(1 - \frac{x^2}{n^2}\right). \quad (36)$$

Then

$$\frac{g_N(x)}{g_{N+1}(x)} \rightarrow -\frac{(1+x)}{x} \quad (37)$$

which implies

$$x g(x) = -(x+1) g(x+1) \quad (38)$$

and

$$x h(x) = -(x+1) h(x+1) \quad (39)$$

and thus the equality of g and h on $] - 1, 1[$ implies that they are equal everywhere, and we have therefore, for any value of x :

$$\frac{\sin(\pi\alpha)}{\pi\alpha} = \prod_{n=1}^{\infty} \left(1 - \frac{\alpha^2}{n^2}\right). \quad (40)$$

The latter formula is in fact a typical example of application of the Weierstrass factorization theorem which states that every entire function can be represented as a (possibly infinite) product involving its zeroes [4, 5].

Let f be an entire function and let $\{a_n\}$ be the nonzero zeroes of f repeated according to multiplicity. Suppose f has a zero at $z = 0$ of order $m \geq 0$ (where order 0 means $f(0) \neq 0$). Then $\exists g$ an entire function and a sequence of integers $\{p_n\}$ such that

$$f(z) = z^m \exp[g(z)] \prod_{n=1}^{\infty} E_{p_n} \left(\frac{z}{a_n}\right) \quad (41)$$

where $E_n(y) = 1 - y$ if $n = 0$ and

$$E_n(y) = (1 - y) \exp \left(y + \frac{y^2}{2} + \cdots + \frac{y^n}{n} \right) \quad (42)$$

if $n = 1, 2, \dots$. It turns out that for $\sin(\pi x)$, the sequence $p_n = 1$ and the function $g(z) = \log(\pi z)$ work.

Thus Euler assumed that it must be possible to represent $\sin x$ as an infinite product of linear factors given by its roots [6, 7]. Using this procedure and knowing that the zeroes of $\sin x$ occur at $0, \pm\pi, \pm 2\pi$, Euler derived formula (40). Using the Euler-Wallis product (40), infinite series representations for π (or $1/\pi$) can be obtained. For instance, we get

$$\frac{1}{\pi} = \frac{\sqrt{2}}{4} \prod_{n=1}^{\infty} \left(1 - \frac{1}{16n^2}\right) \quad (43)$$

in the case $x = 1/4$ and

$$\frac{1}{\pi} = \frac{1}{2} \prod_{n=1}^{\infty} \left(1 - \frac{1}{4n^2}\right). \quad (44)$$

in the case $x = 1/2$. For $x = 1/5$, we have

$$\frac{1}{\pi} = \frac{2}{5} \frac{1}{\sqrt{3-\phi}} \prod_{n=1}^{\infty} \left(1 - \frac{1}{25n^2}\right) \quad (45)$$

which can be put in the form

$$\phi = 3 - \frac{4\pi^2}{25} \prod_{n=1}^{\infty} \left(1 - \frac{1}{25n^2}\right)^2 \quad (46)$$

or equivalently

$$\pi^2 = \frac{25}{4} (3 - \phi) \prod_{n=1}^{\infty} \left(\frac{25n^2}{25n^2 - 1}\right)^2, \quad (47)$$

and in the case $x = 1/10$, since $\sin(\pi/10) = 1/(2\phi)$, one finds

$$\frac{5}{\pi\phi} = \prod_{n=1}^{\infty} \left(1 - \frac{1}{100n^2}\right). \quad (48)$$

Similarly, for $x = 1/3$, we get

$$\frac{3\sqrt{3}}{2\pi} = \prod_{n=1}^{\infty} \left(1 - \frac{1}{9n^2}\right) \quad (49)$$

and for $x = 1/6$:

$$\frac{3}{\pi} = \prod_{n=1}^{\infty} \left(1 - \frac{1}{36n^2}\right). \quad (50)$$

We recall below a few other well-known infinite products, such as the Wallis formula [8]

$$\prod_{n=1}^{\infty} \left(\frac{2n}{2n-1}\right) \left(\frac{2n}{2n+1}\right) = \frac{\pi}{2} \quad (51)$$

and

$$\prod_{n=1}^{\infty} \left(1 - \frac{1}{(2n+1)^2}\right) = \frac{\pi}{4}, \quad (52)$$

as well as the Viète product:

$$\frac{\pi}{2} = \prod_{n=2}^{\infty} \frac{1}{\cos\left(\frac{\pi}{2^n}\right)} \quad (53)$$

where

$$\cos\left(\frac{\pi}{2^n}\right) = \frac{1}{2} \underbrace{\sqrt{2 + \sqrt{2 + \sqrt{2 + \dots}}}}_{(n-1) \text{ times}}, \quad (54)$$

and the two Eulerian products

$$\frac{\pi^2}{6} = \prod_{p \text{ prime}} \frac{p^2}{p^2 - 1} \quad (55)$$

and

$$\frac{\pi}{4} = \prod_{p \equiv 3 \pmod{4}} \left(\frac{p}{p+1} \right) \prod_{p \equiv 1 \pmod{4}} \left(\frac{p}{p-1} \right) = \prod_{p \text{ odd prime}} \frac{p}{p + (-1)^{\frac{p+1}{2}}}. \quad (56)$$

This is of course a non-exhaustive list, and other infinite products were derived, see for instance Ref. [9–11], or the more complicated product [12]:

$$\frac{\pi}{2} = \prod_{n=1}^{\infty} \left(\frac{1}{2n} \right)^{\frac{2}{2n-1}} \left[\prod_{k=1}^n \frac{(2k)^{2k}}{(2k-1)^{2k-1}} \right]^{\frac{4}{4n^2-1}}. \quad (57)$$

5 Conclusion

We obtained, using Fourier series expansions and multiple derivative of the function $\pi/\sin(\pi x)$, series representations for positive powers of π . Leaning on the fact that the Euler-Wallis product can be derived in a straightforward manner from the same formalism, we discussed infinite products representations of π .

References

- [1] G. P. Tolstov, *Fourier Series*, translated by R. Silverman, Englewood Cliffs, NJ: Prentice-Hall, Inc., 1962.
- [2] Kaplan, W. *Advanced Calculus*, 4th ed., Reading, MA: Addison-Wesley, p. 501, 1992.
- [3] A. Tissier, *Mathématiques générales - Agrégation interne de Mathématiques*, Bréal, Montreuil, 1991 [in french].
- [4] J. B. Conway, *Functions of one complex variable*, Springer-Verlag, New York, 1978.
- [5] T. B. Gamelin, *Complex analysis*, Springer-Verlag, New York, 2001.
- [6] O. Ciaurri, *Euler's Product Expansion for the Sine: An Elementary Proof*, Amer. Math. Monthly **122**, 693-695 (2015).
- [7] D. Meyer, A few notes on the Basel problem (2022), <https://davidmeyer.github.io/qc/basel.pdf>
- [8] J. Wallis, *Computation of π by successive interpolations*, in *Arithmetica Infinitorum*, Oxford, 1655; reprinted in *A Source Book in Mathematics, 1200-1800*, D. J. Struik, ed., Princeton University Press, Princeton, 1986, 244-253.
- [9] J. Sondow, *A faster product for π and a new integral for $\ln(\pi/2)$* , Amer. Math. Monthly **112**, 729-734 (2005).

- [10] J. Guillera and J. Sondow, *Double integrals and infinite products for some classical constants via analytic continuations of Lerch's transcendent*, *Ramanujan J.* **16**, 247-270 (2008).
- [11] J. Sondow and H. Yi, *New Wallis- and Catalan-type infinite products for π , e , and $\sqrt{2 + \sqrt{2}}$* , *Amer. Math. Monthly* **117**, 912-917 (2010).
- [12] <https://math.stackexchange.com/questions/4017806/an-infinite-product-for-frac-pi2>
- [13] J.-C. Pain, *Series representations for π^3 involving the golden ratio* (2022), <https://arxiv.org/pdf/2206.15281.pdf>
- [14] J.-C. Pain, *The fifth power of π : new series representation involving the golden ratio and an application in physics* (2022), <https://doi.org/10.5281/zenodo.6915559>
- [15] J.-C. Pain, *New series representations for any positive power of π from a relation involving trigonometric functions* (2022), <https://doi.org/10.48550/arXiv.2208.02624>
- [16] J. M. Borwein and P. B. Borwein, *Pi and the AGM*, John Wiley and Sons, New York, 1987.

A Proof of a relation previously used for deriving series for π^{k+2}

In previous works [13–15], we used the following relation attributed to Euler and mentioned as relation 13.a p. 382 in Ref. [16]. Using Eq. (4), we get

$$\pi \cotan(\pi x) - \pi \cotan(\pi a) = \sum_{n=-\infty}^{\infty} \left(\frac{1}{x-n} - \frac{1}{a-n} \right) = \sum_{n=-\infty}^{\infty} \frac{(a-x)}{(x-n)(a-n)}, \quad (58)$$

which is precisely the identity from which our main results in Refs. [14, 15] were derived. Of course, relation (58) can also be used to derive series representations of π . For instance, setting $a = 1/2$ and $x = 1/4$ yields the formula:

$$\pi = 2 \sum_{n=-\infty}^{\infty} \frac{1}{(2n-1)(4n-1)}. \quad (59)$$