

On Sylvester-type constructions of Hadamard matrices and their modifications

Dmitrii Zinoviev, Victor Zinoviev

November 2, 2022

Abstract

Using the ideas of concatenation construction of codes over the q -ary alphabet, we modify the known generalized Sylvester-type construction of the Hadamard matrices. The new construction is based on two collections of the Hadamard matrices. In particular this construction involves m Hadamard matrices of order k and k Hadamard matrices of order m . These matrices are not necessary different. As a result we obtain a Hadamard matrix of order km . The new construction gives many possibilities for construction of the new Hadamard matrices with different ranks and dimension of kernel.

1 Introduction

Let $E = \{0, 1, \dots, q - 1\}$. A q -ary block code \mathcal{C} is an arbitrary nonempty subset of E^n . We say that \mathcal{C} is a $(n, N, d)_q$ -code, where n is the *length* of the code, N is the number of codewords, i.e. the *cardinality* of \mathcal{C} and d is the *minimum Hamming distance*,

$$d = d(\mathcal{C}) = \min\{d(\mathbf{x}, \mathbf{y}) : \mathbf{x} \neq \mathbf{y}, \mathbf{x}, \mathbf{y} \in \mathcal{C}\}$$

where, for $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{y} = (y_1, \dots, y_n)$ from E^n ,

$$d(\mathbf{x}, \mathbf{y}) = |\{j : x_j \neq y_j, j = 1, \dots, n\}|.$$

A Hadamard matrix H of order n is an $n \times n$ matrix of $+1$'s and -1 's such that $HH^T = nI$, where I is the $n \times n$ identity matrix. We know that if a Hadamard matrix H of order n exists, then n is 1, 2 or a multiple of 4 [1]. Two Hadamard matrices are equivalent if one can be obtained from another one by permuting rows and/or columns and multiplying rows and/or columns by -1 . We can change the first row and column of H into $+1$'s and we obtain an equivalent Hadamard matrix which is called normalized. If $+1$'s are replaced by 0's and -1 's by 1's, and the complementary of all the rows are added, we obtain a binary $(n, 2n, n/2)$ -code which is called a (binary) Hadamard code and is denoted by H_n [1]. We always assume that the Hadamard matrix is normalized, so the corresponding Hadamard code contains the all-zero codeword.

Two structural properties of the binary codes (in particular, the Hadamard codes) are the rank and the dimension of its kernel. The rank of a binary code C is simply the dimension of the linear span, $\langle C \rangle$, of C . The kernel of a binary code C of length n is defined as [2]

$$\ker(C) = \{\mathbf{x} \in \mathbb{Z}_2^n : \mathbf{x} + C = C\}. \quad (1)$$

If the all-zero vector belongs to C , then $\ker(C)$ is a linear subcode of C . Note also that if C is linear, then $\ker(C) = C = \langle C \rangle$. We denote the rank of a binary code C as $\text{rank}(C)$, and the dimension of the kernel as $\dim(\ker(C))$. The rank and the dimension of the kernel can be used to distinguish between nonequivalent binary codes that contain the all-zero vector, since equivalent ones have the same rank and dimension of the kernel [3].

The goal of the present paper is to describe a new general construction of the binary Hadamard codes (or matrices). We extend our construction [4], to modify the known generalized Sylvester-type construction of Hadamard matrices. The new construction is based on two collections of the Hadamard matrices. In particular it involved m Hadamard matrices of order k and k Hadamard matrices of order m . These matrices are not necessarily different. As a result we obtain a Hadamard matrix of order km . This construction gives many possibilities for construction of new Hadamard matrices with different ranks and dimension of kernel.

2 New constructions

The purpose of this section is to describe a new construction of the Hadamard matrices, which modifies the known generalized Sylvester-type construction given by No and Song [5]. The new construction of the Hadamard matrices can be considered as a modification of the generalized Sylvester-type construction from the paper by No and Song [5].

First, recall that by the Sylvester construction of Hadamard matrices from the matrices $H_n = [h_{i,j}]$ and H_m , we mean matrix H_{mn} , denoted by $H_{mn} = H_n \otimes H_m$, obtained by replacing every element $h_{i,j}$ with the matrix $h_{i,j}H_m$ (clearly in this case $h_{i,j} \in \{\pm 1\}$), or with the matrix $h_{i,j} + H_m$ (clearly in this case $h_{i,j} \in \{0, 1\}$), where, for $H_m = [h_{r,k}]$ and any fixed pair i, j ,

$$h_{i,j} + H_m = [h_{i,j} + h_{r,k}].$$

In [5], No and Song suggested the following generalized Sylvester-type construction of Hadamard matrices. We give their result here.

Theorem 2.1. ([5]) *Suppose we have m Hadamard matrices B_1, B_2, \dots, B_m of order k (which are not necessary distinct) and a Hadamard matrix $C = [c_{i,j}]$ of order m where all matrices are over $\{0,1\}$. Then the matrix H ,*

$$H = \begin{bmatrix} c_{1,1} + B_1 & c_{1,2} + B_2 & \dots & c_{1,m} + B_m \\ c_{2,1} + B_1 & c_{2,2} + B_2 & \dots & c_{2,m} + B_m \\ \vdots & \vdots & \vdots & \vdots \\ c_{m,1} + B_1 & c_{m,2} + B_2 & \dots & c_{m,m} + B_m \end{bmatrix},$$

is a Hadamard matrix of order mk .

For a binary vector \mathbf{a} and $e \in \{0,1\}$ denote for the shortness the following vector

$$\mathbf{a} + e = \mathbf{a} + e(1, 1, \dots, 1).$$

Our modification of this construction can be stated in the following:

Theorem 2.2. *Suppose we have m Hadamard matrices A_1, A_2, \dots, A_m of order k (which are not necessary distinct) and k Hadamard matrices B_1, B_2, \dots, B_k of order m (which are also not necessary distinct). Let $\mathbf{a}_i^{(j)}$, $j = 1, 2, \dots, m$, $i = 1, 2, \dots, k$, denote the i -th row of A_j , let $B_u = [b_{r,s}^{(u)}]$, $u = 1, 2, \dots, k$, $r, s = 1, 2, \dots, m$ and $\mathbf{b}_r^{(u)}$ denote r -th row of B_u . Then the*

matrix H ,

$$H = \left[\begin{array}{cccc} \mathbf{a}_1^{(1)} + b_{1,1}^{(1)} & \mathbf{a}_1^{(2)} + b_{1,2}^{(1)} & \dots & \mathbf{a}_1^{(m)} + b_{1,m}^{(1)} \\ \mathbf{a}_2^{(1)} + b_{1,1}^{(2)} & \mathbf{a}_2^{(2)} + b_{1,2}^{(2)} & \dots & \mathbf{a}_2^{(m)} + b_{1,m}^{(2)} \\ \dots & \dots & \dots & \dots \\ \mathbf{a}_k^{(1)} + b_{1,1}^{(k)} & \mathbf{a}_k^{(2)} + b_{1,2}^{(k)} & \dots & \mathbf{a}_k^{(m)} + b_{1,m}^{(k)} \\ \hline \mathbf{a}_1^{(1)} + b_{2,1}^{(1)} & \mathbf{a}_1^{(2)} + b_{2,2}^{(1)} & \dots & \mathbf{a}_1^{(m)} + b_{2,m}^{(1)} \\ \mathbf{a}_2^{(1)} + b_{2,1}^{(2)} & \mathbf{a}_2^{(2)} + b_{2,2}^{(2)} & \dots & \mathbf{a}_2^{(m)} + b_{2,m}^{(2)} \\ \dots & \dots & \dots & \dots \\ \mathbf{a}_k^{(1)} + b_{2,1}^{(k)} & \mathbf{a}_k^{(2)} + b_{2,2}^{(k)} & \dots & \mathbf{a}_k^{(m)} + b_{2,m}^{(k)} \\ \hline \dots & \dots & \dots & \dots \\ \hline \mathbf{a}_1^{(1)} + b_{m,1}^{(1)} & \mathbf{a}_1^{(2)} + b_{m,2}^{(1)} & \dots & \mathbf{a}_1^{(m)} + b_{m,m}^{(1)} \\ \mathbf{a}_2^{(1)} + b_{m,1}^{(2)} & \mathbf{a}_2^{(2)} + b_{m,2}^{(2)} & \dots & \mathbf{a}_2^{(m)} + b_{m,m}^{(2)} \\ \dots & \dots & \dots & \dots \\ \mathbf{a}_k^{(1)} + b_{m,1}^{(k)} & \mathbf{a}_k^{(2)} + b_{m,2}^{(k)} & \dots & \mathbf{a}_k^{(m)} + b_{m,m}^{(k)} \end{array} \right],$$

is a Hadamard matrix of order mk .

We provide an independent proof of this result in terms of the matrices A_i and B_j :

Proof. Let \mathbf{h}_{i_1} and \mathbf{h}_{i_2} be two different rows of H . We have to consider three different cases.

(i) Both rows \mathbf{h}_{i_1} and \mathbf{h}_{i_2} belong to the same i -th shell of H , which is obtained from the elements $b_{i,r}^{(s)}$, where $s = 1, 2, \dots, k$ and $r = 1, 2, \dots, m$.

We have for his case

$$d(\mathbf{h}_{i_1} \mathbf{b}_{i_2}) = \sum_{r=1}^m \frac{k}{2} \cdot d(\mathbf{b}^{(s)}, \mathbf{b}^{(s')}) = \frac{km}{2}.$$

since $s \neq s'$.

(ii) Both rows \mathbf{h}_{i_1} and \mathbf{b}_{i_2} belong to the different i -th and i' -th shells of H ,

but have the same parameter s . In this case we obtain (taking into account, that $\mathbf{a}_s^{(j)} = \mathbf{a}_{s'}^{(j)}$)

$$\begin{aligned}
d(\mathbf{h}_{i_1}, \mathbf{b}_{i_2}) &= \sum_{r=1}^m d(\mathbf{a}_s^{(r)} + b_{i,r}^{(s)}, \mathbf{a}_s^{(r)} + b_{i',r}^{(s)}) \\
&= k \times \sum_{r=1}^m d(b_{i,r}^{(s)}, b_{i',r}^{(s)}) \\
&= k \times d(\mathbf{b}_i^{(s)}, \mathbf{b}_{i'}^{(s)}) \\
&= k \times \frac{m}{2} = \frac{km}{2}.
\end{aligned}$$

(iii) Both rows \mathbf{h}_{i_1} and \mathbf{b}_{i_2} belong to the different i -th and i' -th shells of H and correspond to the different rows $\mathbf{a}_s^{(r)}$ and $\mathbf{a}_{s'}^{(r)}$, $s \neq s'$. Now we have

$$\begin{aligned}
d(\mathbf{h}_{i_1}, \mathbf{b}_{i_2}) &= \sum_{r=1}^m d(\mathbf{a}_s^{(r)} + b_{i,r}^{(s)}, \mathbf{a}_{s'}^{(r)} + b_{i',r}^{(s')}) \\
&= k \times \sum_{r=1}^m d(\mathbf{a}_s^{(r)}, \mathbf{a}_{s'}^{(r)}) \\
&= k \times \frac{m}{2}.
\end{aligned}$$

Indeed, the second equality follows since the following equality is valid for any different s and s'

$$d(\mathbf{a}_s^{(r)}, \mathbf{a}_{s'}^{(r)}) = d(\mathbf{a}_s^{(r)}, \mathbf{a}_{s'}^{(r)} + (1, 1, \dots, 1)),$$

implying that

$$d(\mathbf{a}_s^{(r)} + b_{i,r}^{(s)}, \mathbf{a}_{s'}^{(r)} + b_{i',r}^{(s')}) = d(\mathbf{a}_s^{(r)}, \mathbf{a}_{s'}^{(r)})$$

□

3 Acknowledgements

The research was carried out at the IITP RAS within the framework of fundamental research on the topic mathematical foundations of the theory of

error-correcting codes and at the expense of the National Science Foundation of Bulgaria (NSFB) under project number 20-51-18002.

References

1. Assmus E. F. Jr., Key J. D., Designs and their codes. Cambridge University Press. Great Britain. 1992.
2. Bauer H., Ganter B., Hergert F., Algebraic techniques for nonlinear codes// *Combinatorica*. 1983. V. 3. No. 1, P. 21–33.
3. Fernández-Córdoba C., Vela C., Villanueva M., On \mathbb{Z}_2^s -linear Hadamard codes: kernel and partial classification// *Designs, Codes and Cryptography*. 2019. V. 87. No. 2-3, P. 417–435.
4. Villanueva M., Zinoviev V. A., Zinoviev D. V., On one construction of Hadamard matrices, *Problems of Information Transmission*, 2022, V. 58, No. 4.
5. No Jong-Seon, Song Hong-Yeop, Generalized Sylvester-Type Hadamard Matrices// ISIT. 2000, Sorrento, Italy, (June 25-30, 2000).