

\mathbb{A}^1 -INVARIANCE OF LOCALIZING INVARIANTS

VLADIMIR SOSNILO

ABSTRACT. Weibel proved that p -inverted K-theory is \mathbb{A}^1 -invariant on \mathbb{F}_p -schemes and K-theory with \mathbb{Z}/p -coefficients is \mathbb{A}^1 -invariant on $\mathbb{Z}[\frac{1}{p}]$ -schemes. We extend this result to all finitary localizing invariants of stable ∞ -categories. Along the way we study the Frobenius and Verschiebung endofunctors defined by Tabuada and give an easy proof of Stienstra’s projection formula.

CONTENTS

Introduction	1
1. Reminder on noncommutative motives and localizing invariants	4
2. Endomorphism categories	7
3. Nilpotent endomorphisms obstruct \mathbb{A}^1 -invariance	9
4. Witt vectors	12
5. Torsion in Nil-theories	16
References	19

INTRODUCTION

In the very foundation of motivic homotopy theory lies the idea of enforcing the affine line to be contractible, hence imposing the \mathbb{A}^1 -invariance property¹ on all cohomology theories of one’s interest. Concretely, a cohomology theory² E is said to be \mathbb{A}^1 -invariant if the projection map induces an equivalence

$$E(X) \simeq E(X \times \mathbb{A}^1)$$

for all schemes X . This is not as restrictive as it sounds at first, because many important cohomology theories satisfy \mathbb{A}^1 -property automatically on regular schemes. For example, this is true for K-theory, Chow groups, Morel-Levine’s algebraic cobordism in characteristic 0 (see [LM07]).

However, in general one is also interested in studying singular schemes. In this context K-theory is known not to be \mathbb{A}^1 -invariant—moreover, conjecturally \mathbb{A}^1 -invariance of K-theory on a scheme detects its regularity ([Vor79, Conjecture], [GH10]). As another example, the derived algebraic cobordism of Annala is not \mathbb{A}^1 -invariant (see [Ann22, Theorem 236]). This motivates the quest for *non- \mathbb{A}^1 -invariant motivic homotopy theory*.

0.1. Motivic theories. One approach is to replace \mathbb{A}^1 -invariance with the property of satisfying the projective bundle formula. This is done via *pbf-local sheaves with transfers* and *pbf-local \mathbb{P}^1 -spectra* in the recent works of Annala and Iwasa (see [AI22] and [AI22]). Another more naïve approach is to study *localizing invariants* (aka *noncommutative motives*) whose definition does not refer to any geometric properties at all but is nevertheless sensible enough to reflect many of them. Recall that a localizing invariant is a functor E from the ∞ -category of small idempotent complete stable ∞ -categories $\text{Cat}_\infty^{\text{perf}}$ to spectra Spt satisfying:

¹also called *homotopy invariance*

²i.e. a contravariant functor from the category of schemes

- for any Verdier sequence $\mathcal{A} \rightarrow \mathcal{B} \rightarrow \mathcal{C}$

$$E(\mathcal{A}) \rightarrow E(\mathcal{B}) \rightarrow E(\mathcal{C})$$

is a fiber sequence.

We say that a localizing invariant E is finitary if it preserves filtered colimits³. Any localizing invariant gives rise to a pbf-local \mathbb{P}^1 -spectrum. Given a commutative ring k we will also consider localizing invariants of k -linear stable ∞ -categories. We call them k -localizing invariants.

Many “motivic” properties are shared by pbf-local \mathbb{P}^1 -spectra and localizing invariants. To list a few, the Nisnevich descent [BKRS22, Corollary 4.1.2], the pro-cdh-descent for noetherian schemes [LT19, Theorem A.8], [BKRS22, Theorem C] hold for all localizing invariants. In the setting of pbf-local \mathbb{P}^1 -spectra the motives of Grassmanians $\mathrm{Gr}_n(S)$ and classifying stacks of vector bundles $\mathrm{Vect}_{n,S}$ can be computed as in the \mathbb{A}^1 -invariant theory [AI22, Corollary 4.2.5]. Annala and Iwasa also prove that K-theory can be obtained via a direct analogue of Snaith’s construction [AI22, Theorem 0.1.1] (the \mathbb{A}^1 -invariant analogue was previously proved in [GS09]).

0.2. Back to \mathbb{A}^1 -invariance. In many examples we have some control over the failure of \mathbb{A}^1 -invariance. In particular, this is illustrated by the following result of Weibel:

Theorem 0.3 ([Wei81, Corollary 3.3]). *$K(-)_{[\frac{1}{p}]}$ is \mathbb{A}^1 -invariant on \mathbb{F}_p -schemes. Moreover, $K(-)/l$ is \mathbb{A}^1 -invariant on $\mathbb{Z}[\frac{1}{l}]$ -schemes.*

Recently, Elmanto and Morrow have constructed a non- \mathbb{A}^1 -invariant version of motivic cohomology over schemes of characteristic p (see [EM22]). It is a well-behaved extension of the usual usual (meaning the one defined in [Spi18]) \mathbb{A}^1 -invariant motivic cohomology to non-regular schemes. The two theories coincide after inverting p , so the direct analogue of the above theorem holds. It follows from [Ann21, Theorem 4.22] that the *homological* version of the derived algebraic cobordism is also \mathbb{A}^1 -invariant away from the characteristic. So, it is not unreasonable to expect a similar theorem to hold even for a larger class of non- \mathbb{A}^1 -invariant motivic theories. We do not know whether this is true for pbf-local \mathbb{P}^1 -spectra of Annala-Iwasa under some assumptions, or for the derived algebraic cobordism. However, we prove that this is always true for all finitary localizing invariants:

Theorem 0.4 (Theorem 5.5). *Let k be a connective ring spectrum with $\pi_0(k)$ of characteristic p and E a k -localizing invariant which is finitary. Then $E(-)_{[\frac{1}{p}]}$ is \mathbb{A}^1 -invariant.*

We also generalize the other part of Weibel’s result:

Theorem 0.5 (Theorem 5.9). *Let k be a connective $\mathbb{S}[\frac{1}{l}]$ -algebra spectrum and E a finitary k -localizing invariant. Then $E(-)/l$ is \mathbb{A}^1 -invariant.*

In particular, this implies that Theorem 0.3 holds for all spectral schemes. In the context of dg-categories for E being the functor of nonconnective K-theory these theorems have been proved in [Tab17]. We note that the assumption of being finitary is necessary: for example, it follows from [Elm21, Theorem 3.1.1] that $TC \otimes \mathbb{Q}(-)$ is not \mathbb{A}^1 -invariant.

0.6. Outline of the proof. We follow the same steps as the original Weibel’s proof [Wei81]. Given a k -localizing invariant define

$$NE(\mathcal{C}) := \mathrm{Fib}(E(\mathcal{C} \otimes_k \mathbf{Perf}_{k[t]}) \rightarrow E(\mathcal{C}))$$

Similar to how it is done in [Tab17], we establish an action of *the ring of rational Witt vectors* $\mathbb{W}_0(k)$ on the homotopy groups of $NE(\mathcal{C})$. The main ingredients going into defining this action are the Almkvist’s isomorphism [Alm74]

$$\mathbb{Z} \oplus \mathbb{W}_0(k) \simeq K_0(\mathcal{E}nd(\mathbf{Perf}_k))$$

³In [BGT13] this property was included in the definition of a localizing invariant.

and an explicit isomorphism

$$\pi_i E(\mathcal{C}) \oplus \pi_{i+1} NE(\mathcal{C}) \simeq \pi_i E(\text{Nil}(\mathcal{C})).$$

Here $\mathcal{E}nd(\mathcal{C})$ is the stable ∞ -category of endomorphisms of \mathcal{C} and $\text{Nil}(\mathcal{C})$ is the subcategory of nilpotent endomorphisms. Having these, the action is induced by the tensor product functor

$$\mathcal{E}nd(\mathbf{Perf}_k) \times \text{Nil}(\mathcal{C}) \rightarrow \text{Nil}(\mathcal{C}).$$

Our main results then follow from a thorough analysis of the action, granted by the following steps:

- (1) There are Frobenius and Verschiebung endofunctors $\mathcal{F}_n, \mathcal{V}_n$ on $\text{Nil}(\mathcal{C})$ as well as $\mathcal{E}nd(\mathcal{C})$ (see [BGT15, Section 5]). We provide a different way of defining them, which allows to show that they are nicely compatible with the ring structure of $\mathbb{W}_0(k)$ and satisfy the projection formula with respect to the action of $\mathbb{W}_0(k)$ on $NE(\mathcal{C})$ (see Lemma 4.12). For $\mathcal{C} = \mathbf{Perf}_k$ they induce classical Frobenius and Verschiebung operations on $K_0(\mathcal{E}nd(\mathbf{Perf}_k)) \simeq \mathbb{Z} \oplus \mathbb{W}(\pi_0(k))$. This part allows to represent the action of $p \in \mathbb{W}_0(k)$ on $NE(\mathcal{C})$ by an explicit endofunctor of $\text{Nil}(\mathcal{C})$.
- (2) The action of $\mathbb{W}_0(k)$ on the (discrete) abelian group $\pi_i E(\text{Nil}(\mathcal{C}))$ is continuous. For $E = K$ this follows from the construction of K-theory. In general, we introduce a filtration

$$\dots \rightarrow \text{Nil}^{\leq n}(\mathcal{C}) \rightarrow \text{Nil}^{\leq n+1}(\mathcal{C}) \rightarrow \dots$$

of $\text{Nil}(\mathcal{C})$ such that

$$\underset{n}{\text{colim}} \text{Nil}^{\leq n}(\mathcal{C}) \simeq \text{Nil}(\mathcal{C})$$

and so that the action is continuous on the images of all $\pi_i E(\text{Nil}^{\leq n}(\mathcal{C}))$ by construction (see (5.1) and (5.3.1)). This allows to extend the action of $\mathbb{W}_0(k)$ to an action of the full ring of Witt vectors $\mathbb{W}(k)$ and deduce the main results from its properties.

The constructions in each of the steps are interesting on their own and appear to be completely new.

0.7. Structure of the paper. In Section 1 we remind the reader all they need to know about localizing invariants. In Section 2 we introduce the ∞ -categories of endomorphisms and of nilpotent endomorphisms, crucial for the main result. In Section 3 we study nil-theories $NE(-)$ that measure the failure of \mathbb{A}^1 -invariance of E . We show that $NE(\mathcal{C})$ can be recovered from the values of E on the ∞ -category of nilpotent endomorphisms in \mathcal{C} for any localizing invariant E . In Section 4 we remind the remind classical results about the ring of big Witt vectors. We then study the Frobenius and Verschiebung endofunctors of the ∞ -categories of endomorphisms and relate them to the classical Frobenius and Verschiebung operations on the ring of Witt vectors. Finally, in Section 5 we prove the main result.

Notations & Conventions.

- Throughout the paper we fix a connective commutative ring spectrum k .
- All spectral k -schemes are assumed quasi-compact and quasi-separated. They form a category Sch_k .
- For an ∞ -category \mathcal{C} we denote by \mathcal{C}^\simeq its maximal groupoid and by \mathcal{C}^ω the subcategory of compact objects.
- For ∞ -categories \mathcal{C}, \mathcal{D} , $\text{Fun}(\mathcal{C}, \mathcal{D})$ is the ∞ -category of functors between them. If \mathcal{C} and \mathcal{D} admit finite colimits $\text{Fun}^{\text{ex}}(\mathcal{C}, \mathcal{D})$ is the subcategory of those functors that preserve finite colimits. If \mathcal{C}, \mathcal{D} are also k -linear for some commutative ring spectrum k , we denote by $\text{Fun}_k^{\text{ex}}(\mathcal{C}, \mathcal{D})$ the ∞ -category of k -linear functors that preserve finite colimits.
- Given a small stable ∞ -category \mathcal{C} we use the following notation for the colimit-completion:

$$\widehat{\mathcal{C}} := \text{Fun}_{\text{ex}}(\mathcal{C}^{\text{op}}, \text{Spt}).$$

This is a stable (presentable) ∞ -category and it admits a fully faithful Yoneda embedding $\mathcal{C} \rightarrow \widehat{\mathcal{C}}$.

- For a stable ∞ -category \mathcal{C} and objects $x, y \in \mathcal{C}$ we denote by

$$\mathrm{maps}_{\mathcal{C}}(x, y)$$

the mapping spectrum in \mathcal{C} , while we keeping the notation

$$\mathrm{Maps}_{\mathcal{C}}(x, y)$$

for the mapping space.

- Given an \mathbb{E}_1 -ring R , \mathbf{Mod}_R is the ∞ -category of *right* R -modules, \mathbf{Perf}_R is the subcategory of perfect complexes.
- We denote the category of k -linear idempotent complete stable ∞ -categories by $\mathrm{Cat}_{\infty, k}^{\mathrm{perf}}$.

Acknowledgements. The results of the paper emerged after several busy conversations with Ryomei Iwasa, Marc Hoyois and Niklas Kipp. The author thanks them for sharing their ideas, wishes and insights. Toni Annala pointed out to the author that the Frobenius and Verschiebung endofunctors $\mathcal{F}_n, \mathcal{V}_n$ are induced by the pullback and pushforward functors along the maps $k[t] \xrightarrow{t \mapsto t^n} k[t]$. The author also wants to thank Denis-Charles Cisinski for a discussion about non-finitary localizing invariants. The author gladly acknowledges that he is a member of the SFB 1085 “Higher Invariants” funded by the Deutsche Forschungsgesellschaft (DFG).

1. REMINDER ON NONCOMMUTATIVE MOTIVES AND LOCALIZING INVARIANTS

In this section we recall some basic notions and facts from the theory of localizing invariants. We only discuss those results that are relevant for this paper. For a general theory of localizing invariants we refer to [BGT13]. We start from definitions:

Definition 1.1. *We say that a diagram in $\mathrm{Cat}_{\infty, k}^{\mathrm{perf}}$*

$$\mathcal{A} \xrightarrow{f} \mathcal{B} \xrightarrow{g} \mathcal{C}$$

*is an **exact sequence** of k -linear idempotent complete stable ∞ -categories if f is fully faithful, the composite $g \circ f$ is trivial and the induced functor*

$$\mathcal{B}/\mathcal{A} \rightarrow \mathcal{C}$$

is an equivalence up to idempotent completion.

Definition 1.2. *A functor*

$$E : \mathrm{Cat}_{\infty, k}^{\mathrm{perf}} \rightarrow \mathrm{Spt}$$

*is said to be a **k -localizing invariant** if for any exact sequence $\mathcal{A} \rightarrow \mathcal{B} \rightarrow \mathcal{C}$ of k -linear small idempotent complete stable ∞ -categories the sequence*

$$E(\mathcal{A}) \rightarrow E(\mathcal{B}) \rightarrow E(\mathcal{C})$$

is a fiber sequence.

1.2.1. Any k -localizing invariant gives rise to a cohomology theory on all k -schemes via assignment

$$X \mapsto E(\mathbf{Perf}_X)$$

for any $X \in \mathrm{Sch}_k$. We abuse the notation and write

$$E(X) := E(\mathbf{Perf}_X).$$

We also write

$$E(R) := E(\mathrm{Spec}(R))$$

when R is a k -algebra.

1.2.2. One of the most useful elementary properties of localizing invariants is the following result, sometimes called the additivity theorem. For a localizing invariant E and a fiber sequence

$$f \rightarrow g \rightarrow h$$

in $\mathrm{Fun}_k^{\mathrm{ex}}(\mathcal{C}, \mathcal{D})$ for some k -linear idempotent complete stable ∞ -categories \mathcal{C}, \mathcal{D} on has an equivalence of maps of spectra

$$E(g) \simeq E(f) + E(h) : E(\mathcal{C}) \rightarrow E(\mathcal{D}).$$

Example 1.3. *K-theory, topological Hochschild homology, topological Cyclic homology, TR, Blanc's topological K-theory, Selmer K-theory are k -localizing invariants for any k . Homotopy K-theory is a k -localizing invariant if k is a \mathbb{Z} -algebra.*

Proposition 1.4. *For any scheme X the pullback along the projection $\mathbb{P}_X^1 \xrightarrow{p} X$ and the pushforward along any section $X \xrightarrow{i} \mathbb{P}_X^1$ induce an equivalence*

$$E(\mathbf{Perf}_X) \oplus E(\mathbf{Perf}_X) \xrightarrow{\begin{pmatrix} i_* & p^* \end{pmatrix}} E(\mathbf{Perf}_{\mathbb{P}_X^1})$$

Proof. Follows from [Kha18, Theorem B]. □

Example 1.5. *Motivic cohomology of a fixed weight m does not extend to a k -localizing invariant for any k because there is no equivalence as in Proposition 1.4.*

Proposition 1.6. *Let E be a k -localizing invariant E . Then the square of spectra*

$$\begin{array}{ccc} E(\mathbb{P}_k^1) & \longrightarrow & E(\mathbb{A}_k^1) \\ \downarrow & & \downarrow \\ E(\mathbb{A}_k^1) & \longrightarrow & E(\mathbb{G}_{m,k}) \end{array}$$

is cartesian.

Proof. The diagram in question is obtained via applying E to the Nisnevich square

$$\begin{array}{ccc} \mathbb{G}_{m,k} & \longrightarrow & \mathbb{A}_k^1 \\ \downarrow & & \downarrow \\ \mathbb{A}_k^1 & \longrightarrow & \mathbb{P}_k^1, \end{array}$$

so the result follows from the étale excision ([BKRS22, Theorem 4.1.1], see also [TT90, Proposition 3.19]). □

1.7. Recall that $\mathrm{Cat}_{\infty,k}^{\mathrm{perf}}$ is symmetric monoidal in such a way that $\mathcal{C} \otimes_k \mathcal{D}$ corepresents functors $\mathcal{C} \times \mathcal{D}$ that are k -linear and exact in each variable. One can also see that $-\otimes_k \mathcal{D}$ left adjoint to $\mathrm{Fun}_k^{\mathrm{ex}}(\mathcal{D}, -)$ (see [Lur17, Theorem 4.5.2.1] and [Lur17, 4.8.1]). In fact, one can also explicitly compute $\mathcal{C} \otimes_k \mathcal{D}$ as the compact objects in $\mathrm{Fun}^{\mathrm{ex}}(\mathcal{C}^{\mathrm{op}}, \widehat{\mathcal{D}})$ (see [HSS17, Corollary 4.11]).

Lemma 1.8. *For any k -linear idempotent complete stable ∞ -category \mathcal{C}*

$$-\otimes_k \mathcal{C} : \mathrm{Cat}_{\infty,k}^{\mathrm{perf}} \rightarrow \mathrm{Cat}_{\infty,k}^{\mathrm{perf}}$$

preserves exact sequences and colimits.

Proof. See Lemma 9.35 of [BGT13]. □

1.8.1. *Non-commutative motives.* Tabuada, along with Bloomberg, Cisinski and Gepner has developed a theory of non-commutative motives based on localizing invariants. The idea to view localizing invariants as motives of small idempotent complete stable ∞ -categories comes quite naturally after one settles on an idea that localizing invariants are a good categorical analogue of a cohomology theory. Indeed, in a “category of motives” any suitable cohomology theory is expected to be representable by a motive and every motive should represent a cohomology theory.

However, one also expects there to be a “motive” functor

$$\mathrm{Cat}_{\infty}^{\mathrm{perf}} \rightarrow \mathrm{Fun}^{\mathrm{loc}}(\mathrm{Cat}_{\infty}^{\mathrm{perf}}, \mathrm{Spt})$$

such that $E(\mathcal{C}) \simeq \mathrm{maps}(M(\mathcal{C}), E)$ for any $E \in \mathrm{Fun}^{\mathrm{loc}}(\mathrm{Cat}_{\infty}^{\mathrm{perf}}, \mathrm{Spt})$. This is not easy to construct because of set-theoretic difficulties, so instead, one considers **finitary** localizing invariants.

Definition 1.9. *We say that a functor*

$$E : \mathrm{Cat}_{\infty, k}^{\mathrm{perf}} \rightarrow \mathrm{Spt}$$

is finitary if it preserves filtered colimits.

Example 1.10. *K-theory, topological Hochschild homology, homotopy K-theory, Blanc’s topological K-theory are finitary localizing invariants.*

We denote the ∞ -category of finitary k -localizing invariants $\mathcal{M}_{\mathrm{loc}, k}$.

Theorem 1.11 ([BGT13, Theorem 1.1]). *The embedding functor*

$$\mathcal{M}_{\mathrm{loc}, k} \rightarrow \mathrm{Fun}(\mathrm{Cat}_{\infty, k}^{\mathrm{perf}}, \mathrm{Spt})$$

admits a left adjoint. In particular, there is a functor

$$\mathcal{U}_{\mathrm{loc}, k} : \mathrm{Cat}_{\infty, k}^{\mathrm{perf}} \rightarrow \mathcal{M}_{\mathrm{loc}, k}$$

such that $E(\mathcal{C}) \simeq \mathrm{maps}(M(\mathcal{C}), E)$ for any $E \in \mathcal{M}_{\mathrm{loc}, k}$.

Remark 1.12. *Although all the constructions and the arguments in [BGT13] are done in the non-linear case, i.e. the case of localizing invariants, they work verbatim for k -localizing invariants. See [HSS17, 4,5] for details.*

Remark 1.13. *By construction, the ∞ -category $\mathcal{M}_{\mathrm{loc}, k}$ is presentable. In fact, it admits a presentable symmetric monoidal enhancement such that*

$$\mathcal{U}_{\mathrm{loc}, k} : \mathrm{Cat}_{\infty, k}^{\mathrm{perf}} \rightarrow \mathcal{M}_{\mathrm{loc}, k}$$

is symmetric monoidal.

1.14. **Smooth & proper categories.** Given $\mathcal{C} \in \mathrm{Cat}_{\infty, k}^{\mathrm{perf}}$ we can make sense of modules

Definition 1.15. *Given stable ∞ -categories \mathcal{C}, \mathcal{D} a \mathcal{C} - \mathcal{D} -bimodule is an exact functor*

$$\mathcal{C}^{\mathrm{op}} \otimes_k \mathcal{D} \rightarrow \mathrm{Spt}.$$

*For any \mathcal{C} there is a canonical \mathcal{C} - \mathcal{C} -bimodule given by the mapping spectrum functor, which we call canonical. A **perfect** \mathcal{C} - \mathcal{D} -bimodule is a bimodule which is a compact object of the functor category $\mathrm{Fun}^{\mathrm{ex}}(\mathcal{C}^{\mathrm{op}} \otimes_k \mathcal{D}, \mathrm{Spt})$.*

Definition 1.16. *For $\mathcal{C} \in \mathrm{Cat}_{\infty, k}^{\mathrm{perf}}$ we say:*

- \mathcal{C} is **smooth** if \mathcal{C} is perfect as a \mathcal{C} - \mathcal{C} -bimodule.
- \mathcal{C} is **proper** if the mapping spectra in \mathcal{C} are perfect k -modules.

Remark 1.17. *The notions of smoothness and properness are evidently self-dual.*

Remark 1.18. For a scheme $X \in \text{Sch}_k$ the smoothness and properness of \mathbf{Perf}_X correspond precisely to smoothness and properness of X .

Remark 1.19. For $\mathcal{C}, \mathcal{D} \in \text{Cat}_{\infty, k}^{\text{perf}}$ such that \mathcal{C} is smooth and proper, we have

$$\text{maps}_{\mathcal{M}_{\text{loc}, k}}(\mathcal{U}_{\text{loc}}(\mathcal{C}), \mathcal{U}_{\text{loc}}(\mathcal{D})) \simeq K(\mathcal{C}^{\text{op}} \otimes \mathcal{D}).$$

This is [HSS17, Theorem 5.25] (see also [BGT13, Theorem 9.36]).

Lemma 1.20. For $\mathcal{C}, \mathcal{D} \in \text{Cat}_{\infty, k}^{\text{perf}}$ such that \mathcal{C} is smooth, the functor

$$\text{Fun}^{\text{ex}}(\mathcal{C}^{\text{op}}, \mathcal{D}) \rightarrow \text{Fun}^{\text{ex}}(\mathcal{C}^{\text{op}}, \widehat{\mathcal{D}})$$

factors through the subcategory $\mathcal{C} \otimes \mathcal{D}$ (see 1.7).

Proof. We need to show that this functor lands in the subcategory of compact objects. The functor is natural in \mathcal{D} , so it suffices to show that the image of the identity functor along

$$\text{Fun}^{\text{ex}}(\mathcal{C}^{\text{op}}, \mathcal{C}^{\text{op}}) \rightarrow \text{Fun}^{\text{ex}}(\mathcal{C}^{\text{op}}, \widehat{\mathcal{C}^{\text{op}}}) \simeq \text{Fun}^{\text{ex}}(\mathcal{C}^{\text{op}} \otimes_k \mathcal{C}, \text{Spt})$$

is compact. The image is the canonical bimodule which is compact by smoothness of \mathcal{C} . \square

2. ENDOMORPHISM CATEGORIES

Definition 2.1. Let \mathcal{C} be a small k -linear idempotent complete stable ∞ -category. We introduce the following ∞ -categories:

(1) The stable ∞ -category of endomorphisms on objects of \mathcal{C} :

$$\mathcal{E}nd(\mathcal{C}) := \text{Fun}(B\mathbb{N}, \mathcal{C});$$

(2) The ∞ -category of nilpotent endomorphisms: $\text{Nil}(\mathcal{C})$ is defined to be the subcategory of $\mathcal{E}nd(\mathcal{C})$ consisting of endomorphisms $x \xrightarrow{p} x$ such that the colimit in $\widehat{\mathcal{C}}$

$$X[p^{-1}] := \text{colim}_{\rightarrow} X \xrightarrow{p} X \xrightarrow{p} X \xrightarrow{p} \dots$$

is trivial. It is a stable subcategory because the colimit functor

$$\text{Fun}(B\mathbb{N}, \mathcal{C}) \rightarrow \text{Fun}(B\mathbb{N}, \widehat{\mathcal{C}}) \xrightarrow{\text{colim}} \widehat{\mathcal{C}}$$

is exact.

Lemma 2.2. There is a canonical equivalence between $\mathcal{E}nd(\mathcal{C})$ and the subcategory of

$$\text{Fun}_k^{\text{ex}}(\mathcal{C}^{\text{op}}, \mathbf{Mod}_{k[t]})$$

consisting of functors $F : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Mod}_{k[t]}$ for which the composite

$$\mathcal{C}^{\text{op}} \rightarrow \mathbf{Mod}_{k[t]} \xrightarrow{\mathcal{U}} \mathbf{Mod}_k$$

is representable.

Proof. First we note that $k[t]$ is the free k -algebra on the monoid \mathbb{N} , so we may identify $\mathcal{E}nd(\mathcal{C})$ with $\text{Fun}_k^{\text{ex}}(\mathbf{Perf}_{k[t]}, \mathcal{C})$ (see also [BGT15, Proposition 3.8]). Under this identification taking the underlying object of an endomorphism corresponds to evaluating a functor at the tautological module $k[t]$ -module $k[t]$. Now consider the fully faithful composite

$$\text{Fun}_k^{\text{ex}}(\mathbf{Perf}_{k[t]}, \mathcal{C}) \rightarrow \text{Fun}_k^{\text{ex}}(\mathbf{Perf}_{k[t]}, \widehat{\mathcal{C}}) \simeq \text{Fun}_k^{\text{ex}}(\mathbf{Perf}_{k[t]} \times \mathcal{C}^{\text{op}}, \mathbf{Mod}_k) \simeq \text{Fun}_k^{\text{ex}}(\mathcal{C}^{\text{op}}, \mathbf{Mod}_{k[t]}).$$

It sends an exact functor $F : \mathbf{Perf}_{k[t]} \rightarrow \mathcal{C}$ to a functor

$$\mathcal{C}^{\text{op}} \rightarrow \mathbf{Mod}_{k[t]}$$

$$X \mapsto \text{maps}_{\mathcal{C}}(X, F(k[t])).$$

Here the action of $k[t]$ is induced by the action on $F(k[t])$. Forgetting the action we get a functor represented by $F(k[t])$. On the other hand, for any exact functor $\mathcal{C}^{\text{op}} \xrightarrow{G} \mathbf{Mod}_{k[t]}$ for which the composite

$$\mathcal{C}^{\text{op}} \xrightarrow{G} \mathbf{Mod}_{k[t]} \xrightarrow{u} \mathbf{Mod}_k$$

is representable by Y , the functor

$$\mathbf{Perf}_{k[t]} \rightarrow \mathcal{C}$$

sending $k[t]$ to Y with the induced action of $k[t]$ is in the preimage of G . □

The usage of the word *nilpotent* in the definition above is not accidental:

Lemma 2.3. *For $(P, \phi) \in \text{Nil}(\mathcal{C}) \subset \text{End}(\mathcal{C})$ there exists $n \in \mathbb{N}$ such that $\phi^n \simeq 0$.*

Proof. In view of the exact sequences

$$\text{Fib}(P \xrightarrow{\phi^n} P) \rightarrow P \xrightarrow{\phi} P$$

and the vanishing of $P[\phi^{-1}]$ we have

$$P = \varinjlim_n \text{Fib}(P \xrightarrow{\phi^n} P).$$

Now by compactness of $\mathcal{C} \subset \widehat{\mathcal{C}}$, we see that there exists an $n \in \mathbb{N}$ and a map $P \rightarrow \text{Fib}(P \xrightarrow{\phi^n} P)$ making the following commutative diagram commute:

$$\begin{array}{ccc} P & \longrightarrow & \text{Fib}(P \xrightarrow{\phi^n} P) \\ & \searrow \text{id} & \downarrow \\ & & P \xrightarrow{\phi^n} P \end{array}$$

In particular $\phi^n \simeq 0$. □

Remark 2.4. *Since $\mathbf{Perf}_{k[t]}$ is smooth, by Lemma 1.20 we have a fully faithful embedding*

$$\text{End}(\mathcal{C}) \rightarrow \mathcal{C} \otimes_k \mathbf{Perf}_{k[t]}.$$

We also have an embedding

$$\text{End}(\mathcal{C}) \rightarrow \text{Fun}_k^{\text{ex}}(\mathcal{C}^{\text{op}}, \mathbf{Mod}_{k[t]})$$

2.5. Tensoring endomorphisms. The symmetric monoidal structure on \mathbf{Perf}_k induces a sectionwise symmetric monoidal structure on

$$\text{Fun}_k^{\text{ex}}(\mathbf{Perf}_{k[t]}, \mathbf{Mod}_k) \simeq \mathbf{Mod}_{k[t]}$$

and on the subcategory

$$\text{End}(\mathbf{Perf}_k).$$

Now for \mathcal{C} being a k -linear stable ∞ -category $\text{End}(\mathcal{C})$ is $\text{End}(\mathbf{Perf}_k)$ -tensored. In particular, the biexact pairing

$$\otimes : \mathcal{C} \times \mathbf{Perf}_k \rightarrow \mathcal{C}$$

induces biexact pairings

$$\otimes : \text{Fun}_k^{\text{ex}}(\mathcal{C}^{\text{op}}, \mathbf{Mod}_{k[t]}) \times \mathbf{Mod}_{k[t]} \rightarrow \text{Fun}_k^{\text{ex}}(\mathcal{C}^{\text{op}}, \mathbf{Mod}_{k[t]})$$

and

$$\otimes : \text{End}(\mathcal{C}) \times \text{End}(\mathbf{Perf}_k) \rightarrow \text{End}(\mathcal{C})$$

$$(P, \phi), (Q, \psi) \mapsto (P \otimes Q, \phi \otimes \psi).$$

This also restricts to an action of $\text{End}(\mathbf{Perf}_k)$ on $\text{Nil}(\mathcal{C})$.

2.6. Sending an object $P \in \mathcal{C}$ to the pair $(P, 0)$ gives rise to an exact functor

$$\iota : \mathcal{C} \rightarrow \mathcal{E}nd(\mathcal{C})$$

which factors through an exact functor

$$\iota_{\text{nil}} : \mathcal{C} \rightarrow \mathcal{N}il(\mathcal{C}).$$

This functor admits a left inverse

$$\mathcal{E}nd(\mathcal{C}) \rightarrow \mathcal{C}$$

given by sending an endomorphism to its underlying module. We denote the respective cofibers of the maps

$$\mathcal{U}_{\text{loc},k}(\mathcal{C}) \rightarrow \mathcal{U}_{\text{loc},k}(\mathcal{E}nd(\mathcal{C}))$$

$$\mathcal{U}_{\text{loc},k}(\mathcal{C}) \rightarrow \mathcal{U}_{\text{loc},k}(\mathcal{N}il(\mathcal{C}))$$

in the ∞ -category NMot_k by $\mathcal{E}nd_0(\mathcal{C})$ and $\mathcal{N}il_0(\mathcal{C})$. We also may think of them as the images of idempotents

$$\pi : (P, \phi) \mapsto (P, \phi) - (P, 0).$$

Abusing the notation, we denote by $E(\mathcal{E}nd_0(\mathcal{C}))$ and $E(\mathcal{N}il_0(\mathcal{C}))$ the images of the corresponding idempotents on $E(\mathcal{E}nd(\mathcal{C}))$ and $E(\mathcal{N}il(\mathcal{C}))$ for any k -localizing invariant E (i.e. not necessarily finitary). This is justified by the fact that for a finitary E we have

$$E(\mathcal{E}nd_0(\mathcal{C})) = \text{maps}(\mathcal{E}nd_0(\mathcal{C}), E),$$

$$E(\mathcal{N}il_0(\mathcal{C})) = \text{maps}(\mathcal{N}il_0(\mathcal{C}), E).$$

Remark 2.7. *The image of $K_0(\mathcal{C})$ in $K_0(\mathcal{E}nd(\mathcal{C}))$ gives an ideal $\mathcal{J}_{\text{triv}}$, so $K_0(\mathcal{E}nd(\mathcal{C}))$ and $K_0(\mathcal{E}nd_0(\mathcal{C}))$ are both rings.*

3. NILPOTENT ENDOMORPHISMS OBSTRUCT \mathbb{A}^1 -INVARIANCE

Most of the paper is dedicated to studying the ∞ -categories $\mathcal{N}il(\mathcal{C})$ and $\mathcal{E}nd(\mathcal{C})$. This section explains the main reason we need to do it: as Proposition 3.4 shows, the ∞ -category $\mathcal{N}il(\mathcal{C})$ captures the failure of \mathbb{A}^1 -invariance.

Definition 3.1. *Let E be a functor*

$$\text{Cat}_{\infty,k}^{\text{perf}} \rightarrow \text{Spt}$$

For any $\mathcal{C} \in \text{Cat}_{\infty,k}^{\text{perf}}$ we define

$$NE(\mathcal{C}) := \text{Fib}(E(\mathcal{C} \otimes_k \mathbf{Perf}_{k[t]}) \rightarrow E(\mathcal{C} \otimes_k \mathbf{Perf}_k)),$$

where $k[t] \rightarrow k$ is the evaluation map.

3.1.1. $NE(-)$ promotes to a functor $\text{Cat}_{\infty,k}^{\text{perf}} \rightarrow \text{Spt}$. Equivalently $NE(\mathcal{C})$ can be defined as the cofiber of the map induced by the k -algebra structure:

$$E(\mathcal{C} \otimes_k \mathbf{Perf}_k) \rightarrow E(\mathcal{C} \otimes_k \mathbf{Perf}_{k[t]}).$$

Example 3.2. *The nonconnective K -theory spectrum gives rise to a functor $\text{Cat}_{\infty,k}^{\text{perf}} \rightarrow \text{Spt}$. We have $NK(\mathbf{Perf}_X) = 0$ if X is a regular noetherian scheme, however, in general it is non-trivial.*

Example 3.3. *Assume k is \mathbb{Z} -linear. Then given any functor $E : \text{Cat}_{\infty,k}^{\text{perf}} \rightarrow \text{Spt}$ one can define a functor*

$$EH(\mathcal{C}) := \underset{\Delta^{\text{op}}}{\text{colim}} E(\mathcal{C} \otimes_k \mathbf{Perf}_{k[\Delta \bullet]}).$$

This satisfies $NEH(\mathcal{C}) \simeq 0$ for all $\mathcal{C} \in \text{Cat}_{\infty,k}^{\text{perf}}$. See [Tab15, 5.1] and [LT19, 3.13].

Proposition 3.4. *For a k -linear idempotent complete stable ∞ -category \mathcal{C} the canonical embedding (see Remark 2.4) induces an equivalence*

$$\Psi : \mathrm{Nil}(\mathcal{C}) \rightarrow \mathrm{Ker}(\mathcal{C} \otimes_k \mathbf{Perf}_{k[t]} \rightarrow \mathcal{C} \otimes_k \mathbf{Perf}_{k[t,t^{-1}]}) .$$

Futhermore, the map ι_{nil} corresponds to the functor

$$\mathcal{C} \otimes_k \mathrm{ev}_* : \mathcal{C} \otimes_k \mathbf{Perf}_k \rightarrow \mathcal{C} \otimes_k \mathbf{Perf}_{k[t]}$$

under this equivalence. Here ev_* is the restriction of scalars functor along the evaluation map.

Proof. Ψ is fully faithful by construction, so we need to show that for any

$$(P, \phi) \in (\mathrm{Fun}^{\mathrm{ex}}(B\mathbb{N}, \widehat{\mathcal{C}}))^{\omega} \simeq \mathcal{C} \otimes_k \mathbf{Perf}_{k[t]}$$

such that $P[\phi^{-1}] \simeq 0$ we have that P belongs to $\mathcal{C} = (\widehat{\mathcal{C}})^{\omega}$.

First assume that $\phi \simeq 0$. The functor

$$\begin{aligned} \widehat{\mathcal{C}} &\rightarrow \mathrm{Fun}^{\mathrm{ex}}(B\mathbb{N}, \widehat{\mathcal{C}}) \\ X &\mapsto (X, 0) \end{aligned}$$

is fully faithful and preserves colimits, so it also reflects compact objects. In particular P should be compact.

In general, by Lemma 2.3 there exists an $n \in \mathbb{N}$ such that $\phi^n \simeq 0$. The pushforward along the map of rings $t \mapsto t^p$ yields a functor

$$\mathbf{Perf}_{k[t]} \rightarrow \mathbf{Perf}_{k[t]} .$$

This gives a functor

$$\mathcal{C} \otimes_k \mathbf{Perf}_{k[t]} \rightarrow \mathcal{C} \otimes_k \mathbf{Perf}_{k[t]}$$

which sends (P, ϕ) to $(P, \phi^n) \simeq (P, 0)$. By the above reasoning, P belongs to \mathcal{C} .

The second part of the claim follows from the construction. □

might be interesting but not used

Corollary 3.5. *There is an equivalence $\mathrm{Nil}(\mathcal{C}) \simeq \mathcal{C} \otimes_k \mathrm{Nil}(\mathbf{Perf}_k)$.*

Proof. By Proposition 3.4 we have a Verdier sequence

$$\mathrm{Nil}(\mathbf{Perf}_k) \rightarrow \mathbf{Perf}_{k[t]} \rightarrow \mathbf{Perf}_{k[t,t^{-1}]}$$

which by Lemma 1.8 induces a Verdier sequence

$$\mathcal{C} \otimes_k \mathrm{Nil}(\mathbf{Perf}_k) \rightarrow \mathcal{C} \otimes_k \mathbf{Perf}_{k[t]} \rightarrow \mathcal{C} \otimes_k \mathbf{Perf}_{k[t,t^{-1}]} .$$

Applying Proposition 3.4 again we get the desired equivalence. □

Proposition 3.6. *For a k -localizing invariant E we have a natural equivalence*

$$\Omega N E(\mathcal{C}) \simeq E(\mathrm{Nil}_0(\mathcal{C}))$$

of functors $\mathrm{Cat}_{\infty, k}^{\mathrm{perf}} \rightarrow \mathrm{Spt}$.

Proof. Denote the projection maps $\mathbb{P}_k^1 \rightarrow \mathrm{Spec}(k)$ and $\mathbb{A}_k^1 \rightarrow \mathrm{Spec}(k)$ by $pr_{\mathbb{P}^1}$ and $pr_{\mathbb{A}^1}$, respectively. We have maps given by embedding 0 and its complement into \mathbb{P}_k^1 :

$$\mathrm{Spec}(k) \xrightarrow{i_0} \mathbb{P}_k^1 \xleftarrow{j_0} \mathbb{A}_k^1 .$$

We also have an embedding given the complement to the section at infinity:

$$\mathbb{A}_k^1 \xrightarrow{j_{\infty}} \mathbb{P}_k^1 .$$

Consider the following diagram:

$$\begin{array}{ccccc}
 E(\mathcal{C} \otimes_k \mathbf{Perf}_{\mathbb{P}_k^1, \{0\}}) & \longrightarrow & E(\mathcal{C} \otimes_k \mathbf{Perf}_{\mathbb{P}_k^1}) & \xrightarrow{j_0^*} & E(\mathcal{C} \otimes_k \mathbf{Perf}_{\mathbb{A}_k^1}) \\
 \downarrow j & & \downarrow j_\infty^* & & \downarrow \\
 E(\mathbf{Nil}(\mathcal{C})) & \longrightarrow & E(\mathcal{C} \otimes_k \mathbf{Perf}_{\mathbb{A}_k^1}) & \longrightarrow & E(\mathcal{C} \otimes_k \mathbf{Perf}_{\mathbb{G}_{m,k}}).
 \end{array}$$

By Proposition 3.4 the rows in diagram are fiber sequences. By Lemma 1.8 $E(\mathcal{C} \otimes_k -)$ is a k -localizing invariant, so the right square is a pullback square by Proposition 1.6. This implies that the left vertical map j is an equivalence. It suffices to construct a horizontal map in the commutative diagram

$$\begin{array}{ccc}
 E(\mathcal{C}) & \xrightarrow{\quad} & E(\mathcal{C} \otimes_k \mathbf{Perf}_{\mathbb{P}_k^1, \{0\}}) \\
 & \searrow \iota_{\mathbf{nil}} & \downarrow j \\
 & & E(\mathbf{Nil}(\mathcal{C}))
 \end{array}$$

whose cofiber is $\Omega NE(\mathcal{C})$. It follows from the ‘‘furthermore’’ part of Proposition 3.4 that the direct image functor⁴ $i_{0,*} : \mathcal{C} \otimes_k \mathbf{Perf}_k \rightarrow \mathcal{C} \otimes_k \mathbf{Perf}_{\mathbb{P}_k^1}$ makes the diagram commutative. So we only need to show that its cofiber C is naturally equivalent to $\Omega NE(\mathcal{C})$.

The projective bundle formula of Proposition 1.4 yields that the map

$$E(\mathcal{C} \otimes_k \mathbf{Perf}_k) \oplus E(\mathcal{C} \otimes_k \mathbf{Perf}_k) \xrightarrow{\begin{pmatrix} i_{0,*} & pr_{\mathbb{P}^1}^* \end{pmatrix}} E(\mathcal{C} \otimes_k \mathbf{Perf}_{\mathbb{P}_k^1})$$

is an equivalence. To finish the proof, observe that the following diagram is commutative

$$\begin{array}{ccccc}
 E(\mathcal{C} \otimes_k \mathbf{Perf}_k) & \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} & E(\mathcal{C} \otimes_k \mathbf{Perf}_k) \oplus E(\mathcal{C} \otimes_k \mathbf{Perf}_k) & \xrightarrow{\begin{pmatrix} 0 & 1 \end{pmatrix}} & E(\mathcal{C} \otimes_k \mathbf{Perf}_k) \\
 \downarrow i_{0,*} & & \downarrow \begin{pmatrix} i_{0,*} & pr_{\mathbb{P}^1}^* \end{pmatrix} & & \downarrow pr_{\mathbb{A}^1}^* \\
 E(\mathcal{C} \otimes_k \mathbf{Perf}_{\mathbb{P}_k^1, \{0\}}) & \longrightarrow & E(\mathcal{C} \otimes_k \mathbf{Perf}_{\mathbb{P}_k^1}) & \xrightarrow{j_0^*} & E(\mathcal{C} \otimes_k \mathbf{Perf}_{\mathbb{A}_k^1}) \\
 \downarrow & & \downarrow & & \downarrow \\
 C & \longrightarrow & 0 & \longrightarrow & NE(\mathcal{C})
 \end{array}$$

and all its rows and columns are fiber sequences. □

Corollary 3.7. *Let E be any k -localizing invariant and \mathcal{C} be a k -linear stable ∞ -category. Then $\pi_i NE(\mathcal{C})$ admits an action of the ring $K_0(\mathcal{E}nd_0(\mathbf{Perf}_k))$ for any $i \in \mathbb{Z}$.*

Proof. By Proposition 3.6 $\pi_i NE(\mathcal{C}) \simeq E_{i-1}(\mathbf{Nil}_0(\mathcal{C}))$. Thus it suffices to construct the action of $K_0(\mathcal{E}nd_0(\mathbf{Perf}_k))$ on $\pi_i E(\mathbf{Nil}_0(\mathcal{C}))$. The action of $\mathcal{E}nd(\mathbf{Perf}_k)$ on $\mathbf{Nil}(\mathcal{C})$ (see 2.5)

$$\mathcal{E}nd(\mathbf{Perf}_k) \otimes \mathbf{Nil}(\mathcal{C}) \rightarrow \mathbf{Nil}(\mathcal{C})$$

by adjunction yields a functor

$$\mathcal{E}nd(\mathbf{Perf}_k) \xrightarrow{\simeq} \mathrm{Fun}_{e_x}(\mathbf{Nil}(\mathcal{C}), \mathbf{Nil}(\mathcal{C})) \xrightarrow{\simeq} \mathrm{End}_{\mathrm{Cat}_{\infty,k}^{\mathrm{perf}}}(\mathbf{Nil}(\mathcal{C})) \rightarrow \mathrm{End}_{\mathbb{Z}}(\pi_i E(\mathbf{Nil}(\mathcal{C}))).$$

By additivity theorem this descends into a map

$$K_0(\mathcal{E}nd(\mathbf{Perf}_k)) \rightarrow \mathrm{End}_{\mathbb{Z}}(\pi_i E(\mathbf{Nil}(\mathcal{C})))$$

⁴note that $i_{0,*}$ is well defined because i_0 is a regular embedding

which is a map of rings by construction. This action commutes with the idempotent given by sending $(P, \phi) \in \text{Nil}(\mathcal{C})$ to $(P, 0)$ (see 2.6), so we in fact have a map of rings

$$K_0(\mathcal{E}nd(k)) \rightarrow \text{End}_{\mathbb{Z}}(\pi_i E(\text{Nil}_0(\mathcal{C}))).$$

Finally, the ideal $\mathcal{J}_{\text{triv}}$ is sent to 0, so we obtain the desired action

$$K_0(\mathcal{E}nd_0(k)) \rightarrow \text{End}_{\mathbb{Z}}(\pi_i E(\text{Nil}_0(\mathcal{C}))).$$

□

4. WITT VECTORS

Assume k is a discrete ring. One can check that the map

$$\mathcal{E}nd(k) \rightarrow (1 + tk[[t]])^\times$$

given by sending a pair $(P, \phi) \in \mathcal{E}nd(k)$ to its **characteristic polynomial** $\det(1 - t\phi)$ is a group homomorphism. Here $\mathcal{E}nd(k)$ denotes the exact category of endomorphisms on finitely generated projective k -modules. In fact, the codomain of the map can be endowed with a ring structure in such a way that this map is also a ring homomorphism. This ring is called the **ring of Witt vectors** and is denoted by $\mathbb{W}(k)$. A trivial endomorphism has trivial characteristic polynomial, so we end up having a ring homomorphism

$$K_0(\mathcal{E}nd_0(k)) \rightarrow \mathbb{W}(k).$$

The image of this map consists of fractions of the form

$$\frac{1 + a_1 t + \dots + a_n t^n}{1 + b_1 t + \dots + b_l t^l}.$$

We call the subring spanned by the image the **ring of rational Witt vectors** and denote by $\mathbb{W}_0(k)$.

If k is an arbitrary connective ring spectrum and $\mathcal{E}nd(\mathbf{Perf}_k)$ is considered instead of $\mathcal{E}nd(k)$, the same story goes through, except that now the characteristic polynomial lands in power series over $\pi_0(k)$. The following is, in essence, a classical result of Almkvist [Alm74]. As stated, however, it follows from [BGT15, Theorem 1.12] (see also [BGT15, Theorem 1.9]):

Theorem 4.1 (Almkvist). *The characteristic polynomial map*

$$K_0(\mathcal{E}nd_0(\mathbf{Perf}_k)) \rightarrow \mathbb{W}(\pi_0(k))$$

is an isomorphism onto its image $\mathbb{W}_0(\pi_0(k))$. Moreover, it becomes an isomorphism after completing with respect to the topology defined by preimages of the ideals $(1 + t^N \pi_0(k)[[t]])$.

Example 4.2. $\mathbb{W}(\mathbb{F}_p)$ is isomorphic to the countable product of copies of \mathbb{Z}_p (see [Hes08, Proposition 10]).

Later we will make use of the following fact:

Lemma 4.3. *Assume R is a discrete $\mathbb{Z}[\frac{1}{l}]$ -algebra for $l \in \mathbb{Z}$. Then l is also invertible in $\mathbb{W}(R)$.*

Proof. Define $a_i \in \pi_0(k)$ inductively:

$$a_0 = 1, a_1 = -\frac{1}{l},$$

$$a_n = -\frac{1}{l} \left(\sum_{i_1 + \dots + i_k = n, i_k < n} a_{i_1} \dots a_{i_k} \right).$$

Now it is tedious to check that the power series

$$\mathbf{a} = \sum_{i \in \mathbb{N}} a_i t^i$$

belongs to $\mathbb{W}(\pi_0(k))$ and satisfies

$$\mathbf{a}^l = (1 - t).$$

□

4.4. Verschiebung and Frobenius. Any element x of the ring $\mathbb{W}(\pi_0(k))$ can be represented as an infinite product

$$\prod_{i \in \mathbb{N}} (1 - a_i t).$$

We define

$$\begin{aligned} V_n(1 - at) &= 1 - at^n \\ F_n(1 - at) &= 1 - a^n t. \end{aligned}$$

By additivity and continuity this extends to natural additive Verschiebung and Frobenius operations

$$V_n, F_n : \mathbb{W}(\pi_0(k)) \rightarrow \mathbb{W}(\pi_0(k))$$

for all $n \in \mathbb{N}$ (cf. [Gra78]).

These operations have been studied thoroughly over the past century and they have been shown to satisfy many interesting properties (see [Hes08], [Sti82]). These operations restrict to the subring $\mathbb{W}_0(\pi_0(k))$ and we use the same notation for the restricted operations. For our purposes the most important property is the following:

Lemma 4.5. *Assume R is a discrete \mathbb{F}_p -algebra. Then the equality*

$$V_{p^i} F_{p^i}(x) = p^i x$$

holds for all $x \in \mathbb{W}(R)$.

Proof. By continuity and additivity it suffices to show the equality for $x = (1 + at)$. In this case this is obvious from the definitions and the fact that taking the p^l -th power on R is additive. \square

Remark 4.6. *One has a nice description of the topology on $\mathbb{W}(R)$ from Theorem 4.1 using the Verschiebung operations. Recall that any element $x \in \mathbb{W}(R)$ can be uniquely represented as a product*

$$\prod_{i \in \mathbb{N}} (1 - a_i t^i) = \prod_{i \in \mathbb{N}} V_i(1 - a_i t).$$

If $x \in 1 + t^N R[[t]]$ then $a_i = 0$ for $i < N$. Hence x is a (possibly infinite) sum of $V_i(x_i)$ for $x_i \in \mathbb{W}(R)$ and $i \geq N$. Conversely, any such sum belongs to the ideal, so we have:

$$1 + t^N R[[t]] = \left\{ \prod_{i \geq N} V_i(x_i) \mid x_i \in \mathbb{W}(R) \right\}.$$

4.7. Categorical Frobenius & Verschiebung. A crucial observation about the operations V_n and F_n is that they are induced by explicit endofunctors of $\text{End}(k)$.

4.7.1. For a map of commutative ring spectra $f : R \rightarrow S$ one has an adjunction

$$\mathbf{Mod}_S \begin{array}{c} \xrightarrow{f_*} \\ \xleftarrow{f^!} \end{array} \mathbf{Mod}_R$$

The functor $f^!$ is called the **coinduction** functor. It can be described explicitly as follows:

$$f^!(M) := \text{maps}_R(S, M)$$

If S is perfect over R (for example, it is finite and flat), both f_* and $f^!$ restrict to the subcategories of perfect modules and we get an adjunction

$$\text{Perf}_S \begin{array}{c} \xrightarrow{f_*} \\ \xleftarrow{f^!} \end{array} \text{Perf}_R.$$

Remark 4.8. *If S is perfect over R , then the functor $f^!$ commutes with colimits. This follows from the equivalences*

$$f^! \simeq \text{maps}_R(S, P) \simeq S^\vee \otimes_R P$$

that make $f^!$ a left adjoint.

Construction 4.9. Fix a positive natural number l . Consider the k -linear map of algebras $p_l : k[t] \rightarrow k[t]$ that sends t to t^l . The functors $p_{l,*}$ and $p_l^!$ induce functors

$$\mathrm{Fun}_k^{\mathrm{ex}}(\mathcal{C}^{\mathrm{op}}, \mathbf{Mod}_{k[t]}) \rightarrow \mathrm{Fun}_k^{\mathrm{ex}}(\mathcal{C}^{\mathrm{op}}, \mathbf{Mod}_{k[t]}).$$

By Lemma 2.2 we may identify $\mathrm{End}(\mathcal{C})$ as a subcategory $\mathrm{Fun}_k^{\mathrm{ex}}(\mathcal{C}^{\mathrm{op}}, \mathbf{Mod}_{k[t]})$ of functors for which the composite

$$\mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{Mod}_{k[t]} \xrightarrow{u} \mathbf{Mod}_k$$

is representable. Now $p_{l,*}$ and $p_l^!$ preserve this subcategory because p_l is finite and free. We denote the restriction of $p_{l,*}$ to $\mathrm{End}(\mathcal{C})$ by \mathcal{F}_l and the restriction of $p_l^!$ by \mathcal{V}_l . We call these functors the categorical Frobenius and Verschiebung.

Remark 4.10. Unravelling the definitions we can describe the functors from Construction 4.9 explicitly as follows. Given any $(P, \phi) \in \mathrm{End}(\mathcal{C})$ we have

- $\mathcal{F}_l(P, \phi) \simeq (P, \phi^l)$
- $\mathcal{V}_l(P, \phi) \simeq \mathrm{maps}_{\mathbf{S}_{k[t]}}(k[t^{\frac{1}{l}}], P) \simeq (P^l, \begin{pmatrix} 0 & \cdots & 0 & (-1)^{l+1}\phi \\ 1 & \cdots & 0 & 0 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & 1 & 0 \end{pmatrix})$.

Proposition 4.11. For $\mathcal{C} = \mathbf{Perf}_k$ the induced maps

$$K_0(\mathrm{End}_0(\mathcal{C})) \rightarrow K_0(\mathrm{End}(\mathcal{C})) \xrightarrow{K_0(\mathcal{V}_n), K_0(\mathcal{F}_n)} K_0(\mathrm{End}(\mathcal{C})) \rightarrow K_0(\mathrm{End}_0(\mathcal{C}))$$

are equal to the Verschiebung and Frobenius maps under the isomorphism from Theorem 4.1.

Proof. By Remark 4.10 the functors \mathcal{F}_n and \mathcal{V}_n are the same as those defined in Section 5 of [BGT15]. The analogous functors on the exact category of endomorphisms of finitely generated projective modules over $\pi_0(k)$ were also defined by Grayson in [Gra78] and they were shown to induce the Verschiebung and Frobenius maps on the corresponding summand of K_0 . By Theorem 5.7 of [BGT15] the operations are compatible with the operations defined by Grayson. \square

4.11.1. Recall that

$$\mathrm{Fun}_k^{\mathrm{ex}}(\mathbf{Perf}_{k[t]}, \mathbf{Mod}_k) \simeq \mathbf{Mod}_{k[t]}$$

admits a sectionwise symmetric monoidal structure (see (2.5)). By construction, the functor $p_{l,*}$ is symmetric monoidal. The usual abstract nonsense implies that the adjoint functor $p_l^!$ is lax symmetric monoidal, i.e. there is a natural transformation

$$\mu_{X,Y} : p_l^!(X) \otimes p_l^!(Y) \rightarrow p_l^!(X \otimes Y).$$

Now let \mathcal{C} be a k -linear idempotent complete stable ∞ -category.

Lemma 4.12. There are natural equivalences for $X \in \mathrm{End}(\mathcal{C})$ and $Y \in \mathrm{End}(\mathbf{Perf}_k)$

- (1) $\mathcal{F}_l(X \otimes Y) \simeq \mathcal{F}_l(X) \otimes \mathcal{F}_l(Y)$
- (2) $X \otimes \mathcal{V}_l(Y) \simeq \mathcal{V}_l(X \otimes Y)$

Proof. The first equivalence follows from commutativity of the diagram

$$\begin{array}{ccc} \mathrm{Fun}_k^{\mathrm{ex}}(\mathcal{C}^{\mathrm{op}}, \mathbf{Mod}_{k[t]}) \times \mathbf{Mod}_{k[t]} & \xrightarrow{\otimes} & \mathrm{Fun}_k^{\mathrm{ex}}(\mathcal{C}^{\mathrm{op}}, \mathbf{Mod}_{k[t]}) \\ \downarrow p_{l,*} \times p_{l,*} & & \downarrow p_{l,*} \\ \mathrm{Fun}_k^{\mathrm{ex}}(\mathcal{C}^{\mathrm{op}}, \mathbf{Mod}_{k[t]}) \times \mathbf{Mod}_{k[t]} & \xrightarrow{\otimes} & \mathrm{Fun}_k^{\mathrm{ex}}(\mathcal{C}^{\mathrm{op}}, \mathbf{Mod}_{k[t]}) \end{array}$$

which is simply a manifestation of monoidality of $p_{l,*}$. To construct the second equivalence consider the natural transformation

$$\xi_{X,Y} : X \otimes p_i^!(Y) \xrightarrow{\eta_X \otimes \text{id}_{p_i^!(Y)}} p_i^!(p_*(X)) \otimes p_i^!(Y) \xrightarrow{\mu_{p_*(X),Y}} p_i^!(p_{l,*}(X) \otimes Y)$$

of functors

$$\mathbf{Mod}_{k[t]} \times \mathbf{Mod}_{k[t]} \longrightarrow \mathbf{Mod}_{k[t]}.$$

Notice that both functors commute with filtered colimits in both variables (see Remark 4.8). The natural transformation for $X = Y = (k, \text{id})$ is equivalent to the composite

$$\text{maps}_{k[t]}(k[t^{\frac{1}{t}}], k) \xrightarrow{\text{can} \otimes \text{id}} \text{maps}_{k[t]}(k[t^{\frac{1}{t}}], k) \otimes \text{maps}_{k[t]}(k[t^{\frac{1}{t}}], k) \xrightarrow{\times} \text{maps}_{k[t]}(k[t^{\frac{1}{t}}], k),$$

where *can* is induced by the ring map $k[t^{\frac{1}{t}}] \rightarrow k$ that sends the generator to the unit in k and the second map is the pointwise product in k . We observe that this composite is the identity. Now, the ∞ -category $\mathbf{Mod}_{k[t]}$ is generated under colimits and shifts by the object (k, id) , so this natural transformation is an equivalence in general. Now if X is a functor from $\text{Fun}^{\text{ex}}(\mathcal{C}, \mathbf{Mod}_{k[t]})$, we get a natural equivalence

$$X(C) \otimes p_i^!(Y) \simeq p_i^!(p_{l,*}(X(C)) \otimes Y)$$

of functors

$$\text{Fun}^{\text{ex}}(\mathcal{C}^{\text{op}}, \mathbf{Mod}_{k[t]}) \times \mathbf{Mod}_{k[t]} \longrightarrow \mathbf{Mod}_{k[t]}.$$

By construction, the restriction of the equivalence to the subcategories of endomorphisms gives

$$X \otimes V_l(Y) \simeq \mathcal{V}_l(F_l(X) \otimes Y).$$

□

Remark 4.13. *A quite involved proof of the relation Lemma 4.12(2) in K-theory appears in [Sti82] and later in [Tab17]. We are able to significantly simplify the proof by employing the adjunction between $p_{l,*}$ and $p_l^!$ and also obtain an equivalence on the level of ∞ -categories and not just an equality of K-theory classes.*

Corollary 4.14. *Assume k is an \mathbb{F}_p -algebra. Let \mathcal{C} be a small k -linear idempotent complete stable ∞ -category. Then multiplication by $p^l \in K_0(\text{End}_0(\mathbf{Perf}_k))$ on $K_0(\text{End}_0(\mathcal{C}))$ and $K_0(\text{Nil}_0(\mathcal{C}))$ are equal to the composites*

$$K_0(\text{End}_0(\mathcal{C})) \rightarrow K_0(\text{End}(\mathcal{C})) \xrightarrow{K_0(\mathcal{V}_{p^l} \circ \mathcal{F}_{p^l})} K_0(\text{End}(\mathcal{C})) \rightarrow K_0(\text{End}_0(\mathcal{C}))$$

and

$$K_0(\text{Nil}_0(\mathcal{C})) \rightarrow K_0(\text{Nil}(\mathcal{C})) \xrightarrow{K_0(\mathcal{V}_{p^l} \circ \mathcal{F}_{p^l})} K_0(\text{Nil}(\mathcal{C})) \rightarrow K_0(\text{Nil}_0(\mathcal{C}))$$

for any $l \in \mathbb{N}$.

Proof. If $\mathcal{C} = \mathbf{Perf}_k$, this follows from Proposition 4.11 and Lemma 4.5. In general, there is an equivalence

$$X \otimes V_{p^l}(Y) \simeq \mathcal{V}_{p^l}(F_{p^l}(X) \otimes Y)$$

for $X \in \text{End}(\mathcal{C})$ and $Y \in \text{End}(\mathbf{Perf}_k)$ by Lemma 4.12. In particular, for $Y = (k, \text{id})$ the following equalities hold in K_0 of either $\text{Nil}(\mathcal{C})$ or $\text{End}(\mathcal{C})$

$$p^l[X] = [X][\mathcal{V}_{p^l} \circ \mathcal{F}_{p^l}(k, \text{id})] = [X][\mathcal{V}_{p^l}(k, \text{id})] = [\mathcal{V}_{p^l} \circ \mathcal{F}_{p^l}(X)][(k, \text{id})] = [\mathcal{V}_{p^l} \circ \mathcal{F}_{p^l}(X)]$$

that prove the claim. □

5. TORSION IN NIL-THEORIES

This section is dedicated to proving the main results (Theorems 0.4 and 0.5). So far we never had to require our localizing invariants to be finitary. The reason we need it is to show that the action of $\mathbb{W}_0(\pi_0(k))$ constructed in Corollary 3.7 is continuous. This allows to extend the action to the action of the ring of Witt vectors $\mathbb{W}(\pi_0(k))$. The ring $\mathbb{W}(\pi_0(k))$ has many nice properties which the ring $\mathbb{W}_0(\pi_0(k))$ does not (for example, an integer l is invertible in $\mathbb{W}(\pi_0(k))$ whenever it is invertible in k). The $\mathbb{W}(\pi_0(k))$ -modules $NE(\mathcal{C})$ inherit these properties allowing to prove the main theorems.

5.1. Now we introduce a filtration on the ∞ -category $\text{Nil}(\mathcal{C})$. We start from the case $\mathcal{C} = \mathbf{Perf}_k$. Define $\text{Nil}^{\leq n}(\mathbf{Perf}_k)$ to be the stable ∞ -category of those $k[t]/(t^n)$ -modules that are perfect over k . Restricting scalars gives rise to canonical functors

$$\tau_{n,m} : \text{Nil}^{\leq n}(\mathbf{Perf}_k) \rightarrow \text{Nil}^{\leq m}(\mathbf{Perf}_k) \text{ for any } n \leq m$$

$$\tau_n : \text{Nil}^{\leq n}(\mathbf{Perf}_k) \rightarrow \text{Nil}(\mathbf{Perf}_k) \text{ for any } n.$$

Lemma 5.2. *Let $\{\mathcal{C}_i\}_{i \in I}$ be a filtered diagram of ∞ -categories and \mathcal{C} be an ∞ -category. A map $\text{colim}_I \mathcal{C}_i \rightarrow \mathcal{C}$ is an equivalence if any object of \mathcal{C} is an image of an object of \mathcal{C}_i for some i and for all $x, y \in \mathcal{C}_n$ the maps*

$$\begin{array}{c} \text{colim} \text{Maps}_{\mathcal{C}_i}(x, y) \rightarrow \mathcal{C} \\ \xrightarrow{I_{n,j}} \end{array}$$

are equivalences where $I_{n,j}$ is the category of objects of I under n .

Proof. This can be seen via expressing ∞ -categories as complete Segal spaces inside the functor category $\text{Fun}(\Delta^{\text{op}}, \text{Spc})$, where filtered colimits are computed sectionwise. See also the short note [Roz12]. \square

Lemma 5.3. *The canonical map*

$$\text{colim}_n \text{Nil}^{\leq n}(k) \rightarrow \text{Nil}(k)$$

is an equivalence.

Proof. Any object of $\text{Nil}(k)$ is the image of $\text{Nil}^{\leq k}(k)$ for big enough k . So it suffices to show that for any $U, V \in \text{Nil}^{\leq n}(k)$ the map

$$\text{colim}_m \text{Maps}_{\text{Nil}^{\leq m}(k)}(\tau_{n,m}U, \tau_{n,m}V) \rightarrow \text{Maps}_{\text{Nil}(k)}(\tau_n U, \tau_n V)$$

is an equivalence. Translating into the world of modules we need to show that the map

$$\text{colim}_m \text{Maps}_{k[x]/(x^m)}(U, V) \rightarrow \text{Maps}_{k[x]}(U, V)$$

is an equivalence. By adjunction it is the same as the map

$$\text{colim}_m \text{Maps}_{k[x]/(x^n)}(U \otimes_{k[x]/(x^m)} k[x]/(x^n), V) \rightarrow \text{Maps}_{k[x]}(U, V).$$

It suffices to show that the map of pro-objects

$$\{U\}_m \rightarrow \{U \otimes_{k[x]/(x^m)} k[x]/(x^n)\}_m,$$

is a weak equivalence (here the left-hand side is the constant pro-system). To compute the right hand-side we use the following free resolution for $k[x]/(x^n)$:

$$k[x]/(x^n) \leftarrow k[x]/(x^m) \xrightarrow{x^{m-n}} k[x]/(x^m) \xleftarrow{x^n} k[x]/(x^m) \xrightarrow{x^{m-n}} k[x]/(x^m) \xleftarrow{x^n} \dots$$

For $l > m + n$ we have a map of complexes

$$\begin{array}{ccccccccccc} k[x]/(x^n) & \xleftarrow{pr_{l,n}} & k[x]/(x^l) & \xleftarrow{x^{l-n}} & k[x]/(x^l) & \xleftarrow{x^l} & k[x]/(x^l) & \xleftarrow{x^{l-n}} & k[x]/(x^l) & \xleftarrow{\quad} & \dots \\ \downarrow = & & \downarrow pr_{l,m} & & \downarrow 0 & & \downarrow 0 & & \downarrow 0 & & \\ k[x]/(x^n) & \xleftarrow{pr_{m,n}} & k[x]/(x^m) & \xleftarrow{x^{m-n}} & k[x]/(x^m) & \xleftarrow{x^n} & k[x]/(x^m) & \xleftarrow{x^{m-n}} & k[x]/(x^m) & \xleftarrow{\quad} & \dots \end{array}$$

Now we can compute the map $U \otimes_{k[x]/(x^l)} k[x]/(x^n) \rightarrow U \otimes_{k[x]/(x^m)} k[x]/(x^n)$ as the totalization of the map of complexes

$$\begin{array}{ccccccc} U & \xleftarrow{x^{l-n}} & U & \xleftarrow{x^l} & U & \xleftarrow{x^{l-n}} & U & \xleftarrow{\quad} & \dots \\ \downarrow \text{id} & & \downarrow 0 & & \downarrow 0 & & \downarrow 0 & & \\ U & \xleftarrow{x^{m-n}} & U & \xleftarrow{x^n} & U & \xleftarrow{x^{m-n}} & U & \xleftarrow{\quad} & \dots \end{array}$$

Taking the brutal truncation of these complexes for all m we get a fiber sequence of pro-systems:

$$\{U\}_m \rightarrow \{U \otimes_{k[x]/(x^m)} k[x]/(x^n)\}_m \rightarrow \{(U \otimes_{k[x]/(x^m)} k[x]/(x^n))_{\geq 1}\}_m.$$

From the computation above we see that the morphisms in the pro-system on the right are eventually zero, so it has trivial limit. In particular, the map in question is a pro-equivalence. \square

5.3.1. For \mathcal{C} a k -linear idempotent complete stable ∞ -category we define

$$\text{Nil}^{\leq n}(\mathcal{C}) := \mathcal{C} \otimes_k \text{Nil}^{\leq n}(\mathbf{Perf}_k).$$

By Corollary 3.5 we have an equivalence

$$\text{Nil}(\mathcal{C}) \simeq \mathcal{C} \otimes_k \text{Nil}(\mathbf{Perf}_k)$$

so we have a filtration of $\text{Nil}(\mathcal{C})$ by $\text{Nil}^{\leq n}(\mathcal{C})$.

Corollary 5.4. *In the context above we have*

$$\text{colim}_{\rightarrow} \text{Nil}^{\leq n}(\mathcal{C}) \simeq \text{Nil}(\mathcal{C})$$

Proof. Follows from Lemma 1.8 and Lemma 5.3. \square

Now we are ready to state and prove our main result

Theorem 5.5. *Assume that $\pi_0(k)$ is an \mathbb{F}_p -algebra and E is a finitary k -localizing invariant. Then for any small k -linear idempotent complete stable ∞ -category \mathcal{C} we have*

$$NE(\mathcal{C})\left[\frac{1}{p}\right] \simeq 0.$$

Proof. It suffices to show that every element of $\pi_{i+1}NE(\mathcal{C}) \simeq \pi_i E(\text{Nil}_0(\mathcal{C}))$ is p -torsion (the equivalence is the content of Proposition 3.6). In other words, by Corollary 4.14 it suffices to show that for every $x \in \pi_i E(\text{Nil}_0(\mathcal{C}))$ there exists $l \in \mathbb{N}$ such that $(\mathcal{V}_{p^l} \circ \mathcal{F}_{p^l})(x)$ is in the image of the map

$$\pi_i E(\mathcal{L}_{\text{nil}}) : \pi_i E(\mathcal{C}) \rightarrow \pi_i E(\text{Nil}(\mathcal{C})).$$

Since E is finitary and by Corollary 5.4 we have

$$\pi_i E(\text{Nil}(\mathcal{C})) \simeq \text{colim}_{\rightarrow} \pi_i E(\text{Nil}^{\leq p^l}(\mathcal{C})).$$

Hence any element $x \in \pi_i E(\text{Nil}(\mathcal{C}))$ is the image of an element \bar{x} of $\pi_i E(\text{Nil}^{\leq p^l}(\mathcal{C}))$. Restricting scalars along maps in the commutative diagram of ring spectra

$$\begin{array}{ccc} k[t] & \xrightarrow{p^l} & k[t] \\ \downarrow t \rightarrow 0 & & \downarrow \\ k & \longrightarrow & k[t]/(t^{p^l}) \end{array}$$

we get a commutative diagram

$$\begin{array}{ccc} \mathcal{N}il^{\leq p^l}(\mathcal{C}) & \xrightarrow{\tau_{p^l}} & \mathcal{N}il(\mathcal{C}) \\ \downarrow \mathcal{U} & & \downarrow \mathcal{F}_{p^l} \\ \mathcal{C} & \xrightarrow{\iota_{\text{nil}}} & \mathcal{N}il(\mathcal{C}). \end{array}$$

Now

$$\mathcal{F}_{p^l}(x) = \mathcal{F}_{p^l}(\tau_{p^l}(\bar{x})) = \iota_{\text{nil}}(\mathcal{U}(\bar{x})).$$

It suffices to show that \mathcal{V}_{p^l} preserves the image of ι_{nil} which follows from the following lemma. \square

Lemma 5.6. *There is an equivalence of maps*

$$\mathcal{V}_i \circ \iota_{\text{nil}} \simeq i \cdot \iota_{\text{nil}} : E(\mathcal{C}) \rightarrow E(\mathcal{N}il(\mathcal{C})).$$

Proof. The base change theorem tells us that for all $i \in \mathbb{N} \setminus \{0\}$ we have commutative diagrams

$$\begin{array}{ccc} \mathbf{Mod}_k & \xrightarrow{u_i^!} & \mathbf{Mod}_{k[t]/(t^i)} \\ \downarrow a_{0,*} & & \downarrow a_{i,*} \\ \mathbf{Mod}_{k[t]} & \xrightarrow{p_i^!} & \mathbf{Mod}_{k[t]} \end{array}$$

induced by the pushout diagrams of ring spectra

$$\begin{array}{ccc} k[t] & \xrightarrow{p_i} & k[t] \\ \downarrow a_0 & & \downarrow a_i \\ k & \xrightarrow{u_i} & k[t]/(t^i) \end{array}$$

where the vertical maps are the obvious projections. By construction the composite $p_i^! a_{0,*}$ induces

$$\mathcal{V}_i \circ \iota_{\text{nil}} : \mathcal{N}il(\mathcal{C}) \rightarrow \mathcal{N}il(\mathcal{C})$$

The composite functors $a_i^! u_{i,*}$ fit into exact sequences

$$\begin{array}{ccccc} a_{0,*}(-) & \longrightarrow & a_i^! u_{i,*}(-) & \longrightarrow & a_{i-1}^! u_{i-1,*}(-) \\ \downarrow \simeq & & \downarrow \simeq & & \downarrow \simeq \\ a_{0,*}(-) & & a_{0,*}(-) \otimes_k k[t]/(t^i) & & a_{0,*}(-) \otimes_k k[t]/(t^{i-1}) \end{array}$$

which induce exact sequences of functors

$$\mathcal{C} \rightarrow \mathcal{N}il(\mathcal{C})$$

$$\iota_{\text{nil}} \rightarrow \mathcal{V}_i \circ \iota_{\text{nil}} \rightarrow \mathcal{V}_{i-1} \circ \iota_{\text{nil}}.$$

Applying additivity inductively we see that

$$\mathcal{V}_i \circ \iota_{\text{nil}} : E(\mathcal{C}) \rightarrow E(\mathcal{N}il(\mathcal{C})).$$

is equivalent to i copies of ι_{nil} . \square

5.7. Fix E a finitary k -localizing invariant and \mathcal{C} a k -linear idempotent complete stable ∞ -category. Arguing as in the proof of Theorem 5.5 one can see that $V_n(x)$ acts trivially on the image \mathbf{N}_n of $\pi_i E(\mathcal{N}il^{\leq n}(\mathcal{C}))$ in $\pi_i E(\mathcal{N}il_0(\mathcal{C}))$ for any $x \in \mathbf{W}_0(\pi_0(k))$. Indeed, for y in the image we have an equality in $\pi_i E(\mathcal{N}il_0(\mathcal{C}))$ (Lemma 4.12 and Lemma 5.6):

$$y \cdot V_n(x) = V_n(F_n(y) \cdot x) = V_n(0 \cdot x) = 0.$$

Now, any element of $\mathbf{W}(\pi_0(k))$ can be written as a product

$$\prod_{i \in \mathbb{N}} V_i(x_i)$$

for some $x_i \in \mathbf{W}_0(\pi_0(k))$ (see Remark 4.6), hence we can make sense of the action

$$\begin{aligned} \mathbf{W}(\pi_0(k)) \times \mathbf{N}_n &\rightarrow \mathbf{N}_n \\ \left(\prod_{i \in \mathbb{N}} V_i(x_i), y \right) &\mapsto \sum_{i \in \mathbb{N}} V_i(x_i) \cdot y = \sum_{0 \leq i \leq n} V_i(x_i) \cdot y, \end{aligned}$$

where $x_i \in \mathbf{W}_0(\pi_0(k))$. This action is continuous and extends the action of $\mathbf{W}_0(\pi_0(k)) \subset \mathbf{W}(\pi_0(k))$, so summarizing we have:

Lemma 5.8. *The action of Corollary 3.7*

$$\mathbf{W}_0(\pi_0(k)) \times \pi_i(E(\mathcal{N}il_0(\mathcal{C}))) \rightarrow \pi_i(E(\mathcal{N}il_0(\mathcal{C})))$$

is continuous.

Proof. We need to show that for any $x \in \pi_i(E(\mathcal{N}il_0(\mathcal{C})))$ there exists k such that $1 + t^k R[[t]] \subset \mathbf{W}(\pi_0(k))$ acts trivially on x . We have

$$\pi_i(E(\mathcal{N}il_0(\mathcal{C}))) \simeq \underset{n}{\operatorname{colim}} \mathbf{N}_n$$

in the category of $\mathbf{W}_0(\pi_0(k))$ -modules. Hence there exists n such that $x \in \mathbf{N}_n$. As explained above, the action on \mathbf{N}_n extends to a continuous action of $\mathbf{W}(\pi_0(k))$ with $1 + t^n R[[t]] \subset \mathbf{W}(\pi_0(k))$ acting trivially. \square

Now we can prove our second main result.

Theorem 5.9. *Assume l is invertible in $\pi_0(k)$ for some $l \in \mathbb{Z}$. Let E a finitary k -localizing invariant and \mathcal{C} a k -linear idempotent complete stable ∞ -category. Then*

- (1) l is invertible in $\pi_i NE(\mathcal{C})$;
- (2) E/l is \mathbb{A}^1 -invariant.

Proof. By Lemma 5.8 $\pi_i NE(\mathcal{C}) \simeq \pi_{i-1} E(\mathcal{N}il_0(\mathcal{C}))$ are continuous $\mathbf{W}(\pi_0(k))$ -modules. Now the first claim follows from Lemma 4.3. The second claim follows from the equivalence

$$NE/l(\mathcal{C}) \simeq N(E/l)(\mathcal{C}).$$

\square

REFERENCES

- [AI22] T. Annala and R. Iwasa, *Motivic spectra and universality of K-theory*, 2022
- [Alm74] G. Almkvist, *The Grothendieck ring of the category of endomorphisms*, Journal of Algebra **28** (1974), no. 3, pp. 375–388, <https://www.sciencedirect.com/science/article/pii/0021869374900477>
- [Ann21] T. Annala, *Algebraic Spivak’s theorem and applications*, 2021
- [Ann22] T. Annala, *Derived Algebraic Cobordism*, 2022
- [BGT13] A. J. Blumberg, D. Gepner, and G. Tabuada, *A universal characterization of higher algebraic K-theory*, Geom. Topol. **17** (2013), no. 2, pp. 733–838
- [BGT15] A. Blumberg, D. Gepner, and G. Tabuada, *K-theory of endomorphisms via noncommutative motives*, Transactions of the American Mathematical Society **368** (2015), no. 2, p. 1435–1465, <http://dx.doi.org/10.1090/TRAN/6507>

- [BKRS22] T. Bachmann, A. A. Khan, C. Ravi, and V. Sosnilo, *Categorical Milnor squares and K-theory of algebraic stacks*, *Selecta Mathematica* **28** (2022), no. 5, <http://dx.doi.org/10.1007/s00029-022-00796-w>
- [Elm21] E. Elmanto, *THH and TC are (very) far from being homotopy functors*, *Journal of Pure and Applied Algebra* **225** (2021), no. 8, p. 106640, <http://dx.doi.org/10.1016/j.jpaa.2020.106640>
- [EM22] E. Elmanto and M. Morrow, *Motivic cohomology of schemes*, in preparation
- [GH10] T. Geisser and L. Hesselholt, *On a conjecture of Vorst*, *Mathematische Zeitschrift* **270** (2010), no. 1–2, p. 445–452, <http://dx.doi.org/10.1007/s00209-010-0806-2>
- [Gra78] D. R. Grayson, *Grothendieck rings and Witt vectors*, *Communications in Algebra* **6** (1978), no. 3, pp. 249–255, preprint <https://doi.org/10.1080/00927877808822245>
- [GS09] D. Gepner and V. P. Snaith, *On the Motivic Spectra Representing Algebraic Cobordism and Algebraic K-Theory*, *Documenta Math.* (2009), no. 14, pp. 359–396
- [Hes08] L. Hesselholt, *Lecture notes on Witt vectors*, 2008
- [HSS17] M. Hoyois, S. Scherotzke, and N. Sibilla, *Higher traces, noncommutative motives, and the categorified Chern character*, *Adv. Math.* **309** (2017), pp. 97–154, <https://doi.org/10.1016/j.aim.2017.01.008>
- [Kha18] A. A. Khan, *Algebraic K-theory of quasi-smooth blow-ups and cdh descent*, 2018, to appear in *Annales Henri Lebesgue*
- [LM07] M. Levine and F. Morel, *Algebraic Cobordism*, Springer, 2007
- [LT19] Land and Tamme, *On the K-theory of pullbacks*, *Annals of Mathematics* **190** (2019), no. 3, p. 877, <http://dx.doi.org/10.4007/annals.2019.190.3.4>
- [Lur17] J. Lurie, *Higher Algebra*, September 2017, <http://www.math.harvard.edu/~lurie/papers/HA.pdf>
- [Roz12] N. Rozenblyum, *Filtered colimits of ∞ -categories*, 2012, <https://people.math.harvard.edu/~gaitsgde/GL/colimits.pdf>
- [Spi18] M. Spitzweck, *A commutative \mathbf{P}^1 -spectrum representing motivic cohomology over Dedekind domains*, *Mém. Soc. Math. Fr.* **157** (2018), preprint [arXiv:1207.4078](https://arxiv.org/abs/1207.4078)
- [Sti82] J. Stienstra, *Operations in the higher K-theory of endomorphisms*, no. 2, pp. 59–115, <https://books.google.de/books?id=Deb0fn093m4C>
- [Tab15] G. Tabuada, *A¹-homotopy theory of noncommutative motives*, *Journal of Noncommutative Geometry* **9** (2015), no. 3, p. 851–875
- [Tab17] ———, *A¹-homotopy invariance of algebraic K-theory with coefficients and du Val singularities*, *Annals of K-Theory* **2** (2017), no. 1, p. 1–25, <http://dx.doi.org/10.2140/AKT.2017.2.1>
- [TT90] R. W. Thomason and T. Trobaugh, *Higher algebraic K-theory of schemes and of derived categories*, *The Grothendieck Festschrift III*, *Progress in Mathematics*, vol. 88, Birkhäuser, 1990, pp. 247–435
- [Vor79] T. Vorst, *Localization of the K-Theory of Polynomial Extensions.*, *Mathematische Annalen* **244** (1979), pp. 33–54, <http://eudml.org/doc/163289>
- [Wei81] C. A. Weibel, *Mayer-Vietoris sequences and module structures on NK_** , pp. 466–493, <https://doi.org/10.1007/BFb0089534>

M309, UNIVERSITÄT REGENSBURG, UNIVERSITÄTSSTRASSE 31, 93053 REGENSBURG, GERMANY
 Email address: vsosnilo@gmail.com