

COLLAPSED MANIFOLDS WITH LOCAL RICCI BOUNDED COVERING GEOMETRY

XIAOCHUN RONG¹

ABSTRACT. For $\rho, v > 0$, we say that an n -manifold M satisfies local (ρ, v) -bound Ricci covering geometry, if Ricci curvature $\text{Ric}_M \geq -(n-1)$, and for all $x \in M$, $\text{vol}(B_\rho(\tilde{x})) \geq v > 0$, where \tilde{x} is an inverse image of x on the (local) Riemannian universal cover of the ρ -ball at x .

In this paper, we extend the nilpotent fiber bundle theorem of Cheeger-Fukaya-Gromov on a collapsed n -manifold M of bounded sectional curvature to M of a local (ρ, v) -bound Ricci covering geometry, and M is close to a non-collapsed Riemannian manifold of lower dimension. The nilpotent fiber bundle theorem significantly improves fiber bundle theorem in [Hu], and it strengthens a nilpotent fiber bundle seen from [NZ] and implies the torus bundle in [HW], which are obtained under additional local or global topological conditions, respectively.

Our construction of a nilpotent fibration requires a new proof for a result in [HKRX]: if an n -manifold M with local $(1, v)$ -bound Ricci covering geometry has diameter $< \epsilon(n, v)$, a constant depends on n and v , then M is diffeomorphic to an infra-nilmanifold. The proof in [HKRX] is to show that the Ricci flows produces an almost flat metric, thus the result follows from the Gromov's theorem on almost flat manifolds. The new proof is independent of the Gromov's theorem, thus has which as a corollary. If the first Betti number $b_1(M) = n$, then M satisfies a $(1, v)$ -bound Ricci covering geometry, thus M is diffeomorphic to a standard torus ([Co2]).

0. INTRODUCTION

A complete n -manifold M is called ϵ -collapsed, if every 1-ball on M has volume less than ϵ , while normalizing a bound on curvature to exclude a rescaling.

In this paper, we will extend the nilpotent fiber bundle theorem of Cheeger-Fukaya-Gromov on a collapsed n -manifold M of bounded sectional curvature (Theorem 0.2) to M of a local (ρ, v) -bound Ricci covering geometry (see below), and M is close, in the Gromov-Hausdorff distance, to a non-collapsed Riemannian manifold of lower dimension (Theorem B). The nilpotent fiber bundle theorem significantly improves fiber bundle theorem in [Hu] (Theorem 0.4), it much strengthens a nilpotent fiber bundle seen from [NZ] (Theorem 0.9) and implies the torus bundle in [HW] (Theorem 0.9), which are obtained under additional local and global topological restriction respectively.

¹A part of this work was done during the author's sabbatical leave in the fall of 2018 and spring of 2022 at Capital Normal University, partially supported by NSFC 11821101, BNSF Z19003, and a research fund from Capital Normal University.

Our construction of a nilpotent fibration requires a new proof for a result in [HKRX]: if an n -manifold M with local $(1, v)$ -bound Ricci covering geometry has diameter $< \epsilon(n, v)$, a constant depends on n and v , then M is diffeomorphic to an infra-nilmanifold (Theorem 0.3). The proof in [HKRX] is to show that the Ricci flows produces an almost flat metric, thus the result flows from the Gromov's theorem on almost flat manifolds. The new proof is independent of the Gromov's theorem (Theorem A), thus has which as a corollary (Corollary 0.5). Moreover, if $b_1(M) = n$ implies that M satisfies $(1, v)$ -bound Ricci covering geometry ((C1) in Theorem C), thus M is diffeomorphic to a standard torus (Theorem 0.9), the maximal first Betti number rigidity of almost non-negative Ricci curvature ([Co2]).

Let's start with some back ground. In the study of ϵ -collapsed manifolds with normalizing bound on sectional curvature, $|\sec_M| \leq 1$, the following almost flat manifolds theorem of Gromov has been a cornerstone:

Theorem 0.1. (Almost flat manifolds [Gr], [Ru]) *Given $n > 0$, there exist constants, $\epsilon(n), w(n) > 0$, such that if the scaling invariant of a compact n -manifold M , $(\max |\sec_M|)(\text{diam}(M))^2 < \epsilon(n)$, or equivalently,*

$$|\sec_M| \leq 1, \quad \text{diam}(M) < \epsilon(n),$$

then M is diffeomorphic to an infra-nilmanifold, N/Γ , where N is a simply connected nilpotent Lie group, $\Gamma < N \rtimes \text{Aut}(N)$ is a discrete subgroup such that $[\Gamma : \Gamma \cap N] \leq w(n)$.

In [Gr], Gromov proved that a bounded normal covering of M is diffeomorphic to a nilmanifold, and that M is diffeomorphic to an infra-nilmanifold was due to [Ru] via constructing a flat connection with a parallel torsion on M .

Starting with Theorem 0.1, Cheeger-Gromov ([CG1,2]), Fukaya ([Fu1-3]), and Cheeger-Fukaya-Gromov ([CFG]) investigated an ϵ -collapsed manifold M with $|\sec_M| \leq 1$, and the main discovery is a compatible local elementary nilpotent structures on M , every orbit has a positive dimension and points to all the collapsed directions (Theorem 1.2.2). Conversely, if a manifold admits such a compatible elementary nilpotent structures, then one constructs an one parameter family of invariant metrics g_ϵ on M ($\epsilon \in (0, 1]$) with $|\sec_{g_\epsilon}| \leq 1$ and any orbit collapses to a point, which implies that for any $x \in M$, $\text{vol}_{g_\epsilon}(B_1(x)) < \epsilon$ (Theorem 1.2.1, cf. [CR], [CG1], [Fu1]).

In the case that a collapsing is taking place with a bounded diameter (see blow), the nilpotent structure is pure, which has the following alternative formulation:

Theorem 0.2. ([Fu1,2], [CFG]) *Assume a sequence of compact n -manifolds converging in the Gromov-Hausdorff topology, $M_i \xrightarrow{\text{GH}} X$, such that*

$$|\sec_{M_i}| \leq 1, \quad \text{diam}(M_i) \leq d, \quad \dim(X) < n.$$

(0.2.1) (Nilpotent fiber bundle) *If X is a Riemannian manifold, then $d_{\text{GH}}(M_i, X) < \epsilon(X)$ (a constant depending on X) implies a smooth fiber bundle map, $f_i : M_i \rightarrow X$, with fiber an infra-nilmanifold and an affine structural group, satisfying*

(0.2.1.1) *f_i is an ϵ_i -Gromov-Hausdorff approximation (briefly GHA), $\epsilon_i \rightarrow 0$.*

(0.2.1.2) *f_i is an ϵ_i -Riemannian submersion i.e., for vector x_i orthogonal to a f_i -fiber, $e^{-\epsilon_i} \leq \frac{|df_i(\xi_i)|}{|\xi_i|} \leq e^{\epsilon_i}$.*

(0.2.1.3) *The second fundamental form of f_i -fibers, $|\Pi_{f_i}| \leq c(n)$.*

(0.2.2) (Singular nilpotent fibration) *If X is not a Riemannian manifold, then the frame bundle of M_i admits a smooth $O(n)$ -invariant fiber bundle map, $F_i \rightarrow (F(M_i), O(n)) \xrightarrow{f_i} (Y, O(n))$, with Y a $C^{1,\alpha}$ -Riemannian manifold, F_i a compact nilmanifold and an affine structure group, such that the following diagram commutes,*

$$\begin{array}{ccc} F_i \rightarrow (F(M_i), O(n)) & \xrightarrow{f_i} & (Y, O(n)) \\ \downarrow \text{proj}_i & & \downarrow \text{proj} \\ \bar{F}_i \rightarrow M_i & \xrightarrow{\bar{f}_i} & X = Y/O(n) \end{array}$$

where f_i satisfies (0.2.1.1)-(0.2.1.3), \bar{f}_i is the descending map of f_i and every $\bar{F}_i = \text{proj}_i(F_i)$ is diffeomorphic to an infra-nilmanifold.

For the existence of a mixed nilpotent structure on an ϵ -collapsed n -manifold with $|\text{sec}| \leq 1$ (but no bound on $\text{diam}(M)$), see Section 1.2, and for applications of the existence of nilpotent structures on collapsed manifolds with bounded sectional curvature, see [Fu3], [Ro1,2].

Let's turn to a collapsed manifold with bounded Ricci curvature. The weak curvature regularity bring in a complexity of underlying structures. For instance, it is unlikely to expect a classification on an almost Ricci flat manifolds (i.e., $|\text{Ric}_M| \cdot (\text{diam}(M))^2 < \epsilon$); partly because which contains all compact Ricci flat manifolds (comparing to Theorem 0.1), nor to expect any fibration structure as in (0.2.1), if replacing $|\text{sec}_M| \leq 1$ with $|\text{Ric}_M| \leq n - 1$ ([An]).

In view of the above, most work in recent investigation has been focusing on situations with various additional conditions ([CRX1,2], [DWY], [HKRX], [HRW], [HW1], [Ka], [NT1,2], [NZ], [PWY], etc.), in finding a nearby metric with bounded sectional curvature via local or global smoothing techniques ([Ab], [DWY], [Ha], [HW2], [Per], [PWY]), so as to obtain a underlying nilpotent structure from the work of Cheeger-Fukaya-Gromov.

Around 2016, the author proposed to investigate the class of collapsed manifolds satisfying a local (ρ, v) -bound Ricci covering geometry; which includes all manifolds in the above work, as well as ones admitting no nearby metric with bounded sectional curvature (Example 1.5). We conjectured that this class of collapsed manifolds should pose similar nilpotent structures (Conjecture 0.6). The proposal has been partially supported by recent construction of collapsing Calabi-Yau metrics ([GW], [GTZ1,2], [HSVZ], [Zh], etc), where a local (ρ, v) -bound Ricci covering geometry are satisfied on M , or a region in M away from compact subset of lower dimension.

For $\rho > 0$, we call the metric ball, $B_\rho(\tilde{x})$, a *local rewinding* of $B_\rho(x)$, $x \in M$, where $\pi : (\widetilde{B_\rho(x)}, \tilde{x}) \rightarrow (B_\rho(x), x)$ is the (incomplete) Riemannian universal cover. We call the number, $\text{vol}(B_\rho(x)) := \text{vol}(B_\rho(\tilde{x}))$, the local rewinding volume of $B_\rho(x)$. Clearly, $\widetilde{\text{vol}}(B_\rho(x)) = \text{vol}(B_\rho(x))$ if and only if $B_\rho(x)$ is simply connected. If $\text{vol}(B_\rho(x))$ is small and $\widetilde{\text{vol}}(B_\rho(x))$ is not small, we say that $B_\rho(x)$ is collapsed with local bounded covering geometry. We say that M satisfies local (ρ, v) -bound Ricci covering geometry if

$$\text{Ric}_M \geq -(n-1), \quad \widetilde{\text{vol}}(B_\rho(x)) \geq v > 0, \quad \forall x \in M,$$

or briefly, M satisfies local Ricci bounded covering geometry.

First, any collapsed n -manifold with $|\sec| \leq 1$ satisfies local (ρ, v) -bound covering geometry with ρ and v depending only on n ([CFG]). Secondly, if a collapsing is taking place along a nilpotent fibration i.e., each fiber collapses to a point while its normal slice is not collapsed, then the Riemannian universal cover of a local trivialization is not collapsed. Thirdly, a collapsed manifold with local/global (ρ, v) -bound Ricci covering geometry may admit no nearby metric of bounded sectional curvature (Example 1.5.1), or the present smoothing techniques fail to apply (Theorem 0.4).

The main goal of our investigation is to establish a similar nilpotent structure on a collapsed manifold with local (ρ, v) -bound Ricci covering geometry; whose regular (resp. singular) part coincides (resp. is different) with that of a nilpotent structure in [CFG] (Conjecture 0.6 below). Following a guide in the Cheeger-Fukaya-Gromov theory on collapsed manifolds with bounded sectional curvature, the first step is to classify maximally collapsed manifold (i.e., $\text{diam}(M)$ is small) with a local Ricci bounded covering geometry.

Theorem 0.3. (Maximally collapsed manifolds with Ricci bounded covering geometry, [HKRX]) *Given $n, v > 0$, there exist constants, $\epsilon(n, v), w(n) > 0$, such that if a compact n -manifold M satisfies*

$$\text{Ric}_M \geq -(n-1), \quad \text{vol}(B_1(\tilde{p})) \geq v, \quad \text{diam}(M) < \epsilon(n, v),$$

then M is diffeomorphic to an infra-nilmanifold, N/Γ , where \tilde{p} is a point in the Riemannian universal cover of M , N is a simply connected nilpotent Lie group, $\Gamma < N \rtimes \text{Aut}(N)$ is a discrete subgroup such that $[\Gamma : \Gamma \cap N] \leq w(n)$.

Note that for a maximally collapsed manifold M , the local Ricci bounded covering geometry is equivalent to that the Riemannian universal cover of M is not collapsed.

The proof in [HKRX] showed that the Ricci flows on M exists for a definite time which yields an almost flat metric, thus Theorem 0.3 follows from Theorem 0.1. The time estimate relies on the Perelman's pseudo-locality ([Per]) on the Riemannian universal cover \tilde{M} , which requires that every point $\tilde{x} \in \tilde{M}$, $\text{vol}(B_1(\tilde{x}))$ almost equals to $\text{vol}(\mathbb{B}_1^n(0))$ ([CM]), and which is verified based on the Cheeger-Colding's work on a non-collapsed Ricci limit space ([CC1-3]).

The next step is to generalize Theorem 0.2 to a collapsed manifold with local (ρ, v) -bound Ricci covering geometry; which has been partially done in the case that X is a Riemannian manifold (see (0.2.1)).

Theorem 0.4. (Fiber bundles, [Hu]) *Given $n, v > 0$, there exists a constant $\epsilon(n, v) > 0$ such that if compact n -manifold M and m -manifold N satisfy, $\forall x \in M$, $0 < \epsilon \leq \epsilon(n, v)$,*

$$\text{Ric}_M \geq -(n-1), \quad \widetilde{\text{vol}}(B_1(x)) \geq v, \quad |\sec_N| \leq 1, \quad \text{injrad}(N) \geq 1, \quad d_{\text{GH}}(M, N) < \epsilon,$$

then there is a smooth fiber bundle map, $f : M \rightarrow N$, which is also an $\Psi(\epsilon|n, v)$ -GHA, where $\Psi(\epsilon|n, v) \rightarrow 0$ as $\epsilon \rightarrow 0$ while n and v are fixed.

Note that smoothing methods mentioned in the above work unlikely apply under the conditions of Theorem 0.4. In the proof ([Hu]), f is constructed by gluing locally

defined (m, δ) -splitting maps ([CC1]) via the standard center of mass technique associate to a partition of unity on N , and the non-degeneracy of f relies on the non-degeneracy of any (n, δ) -splitting map ([CJN]).

Comparing with (0.2.1), the significant structural information missing in Theorem 0.4 is that a f -fiber should be diffeomorphic to an infra-nilmanifold, and the structural group should be reduced to an affine group; the two properties are expected and have been indispensable in applications ([Ro1]). Indeed, in the case that the Riemannian universal cover \tilde{M}_i is not collapsed, a singular nilpotent fibration with an affine structural group was obtained in [HKRX] by a smoothing method via Ricci flows, similar to the proof of Theorem 0.3.

In contrast to that (0.2.1) follows from Theorem 0.1, Theorem 0.3 fails to apply to f -fiber in Theorem 0.4 because the intrinsic metric on a f -fiber unlikely satisfies Theorem 0.3. Hence, to conclude that a fiber in Theorem 0.4 is diffeomorphic to an infra-nilmanifold and there is an affine structural group, one needs to find a new proof of Theorem 0.3 without Theorem 0.1.

We now begin to state main results in this paper.

Theorem A. *Theorem 0.3 can be proved independent of Theorem 0.1.*

A significance of Theorem A is that the proof naturally extends to a proof of a nilpotent fibration theorem (Theorem B). A bi-product from the proof of Theorem A is that one explicitly constructs a nearby left invariant almost flat metric (see completion of proof of Theorem 0.3').

A consequence of Theorem 0.1 is that the Riemannian universal covering of M of $|\sec_M| \leq 1$ and $\text{diam}(M) < \epsilon(n)$ is not collapsed. An elementary proof independent of Theorem 0.1 was recently given (see (8.4) in [Ro3]). Hence, a consequence of Theorem A is

Corollary 0.5. *Theorem 0.3 implies Theorem 0.1.*

Our approach to Theorem A is partially motivated by a new proof for Theorem 0.1 in [Ro4], which is by induction on dimension based on a recent criterion of nilmanifolds: a compact manifold M is a nilmanifold if and only if M admits an iterated principal circle bundles ([Na], [Be]). Because the inductive proof relies on smoothing techniques we are not allowed to use, we formulate an alternative criterion for our purpose (Theorem 2.1), which can be verified using tools available on manifolds of Ricci curvature bounded below (a brief outline is given at the end of introduction).

Theorem B. (Nilpotent fiber bundles) *Let the assumptions be as in Theorem 0.4. Then there is a smooth bundle map, $f : M \rightarrow N$, such that f is an $\Psi(\epsilon|n, v)$ -GHA, a f -fiber is diffeomorphic to an infra-nilmanifold, and the structural group is affine.*

Theorem B substantially improves Theorem 0.4 by asserting a fiber an infra-nilmanifold and an affine structural group.

We will present two proofs of Theorem B: one is to run the proof of Theorem A to a fiber (in a tubular neighborhood) in Theorem 0.4, and the other is to, by locally running the proof of Theorem A, construct a locally finite open cover for M , each supports an elementary N-structure (see Section 1.3), for which to apply gluing techniques in [CFG] to patch these elementary N-structures to a desired nilpotent fibration.

Theorem B extends (0.2.1) to collapsed manifolds with local (ρ, v) -bound Ricci covering geometry. Concerning a possible extension of (0.2.2), we recall the following conjecture ([Ro4], [HRW]):

Conjecture 0.6. Let a sequence of compact n -manifolds, $M_i \xrightarrow{\text{GH}} X$ (compact), with $\text{Ric}_{M_i} \geq -(n-1)$ and $\text{vol}(M_i) \rightarrow 0$.

(0.6.1) (Singular nilpotent fibration) Assume all $x_i \in M_i$ are local rewinding (ρ, ϵ) -Reifenberg points i.e., $d_{\text{GH}}(B_{i,s}(\tilde{x}_i), \mathbb{B}_s^n(0)) < s\epsilon$, $0 < s \leq \rho$, where $(\widetilde{B}_\rho(x_i), \tilde{x}_i) \rightarrow (B_\rho(x_i), x_i)$ is a (local) Riemannian universal covering map. Then for i large, there is a singular nilpotent fibration, $f_i : M_i \rightarrow X$, which satisfies the commutative diagram in (0.2.2).

(0.6.2) (Generalized singular nilpotent fibration) Assume M_i satisfies local (ρ, v) -bound Ricci covering geometry. Then for i large, there is a continuous map, $f_i : M_i \rightarrow X$, f_i is an ϵ_i -GHA, and for any $x \in X$, $f_i^{-1}(x)$ is finite quotient of an infra-nilmanifold, and at $x_i \in f_i^{-1}(x)$, any short geodesic loop is shortly homotopic to a nontrivial element in $\pi_1(f_i^{-1}(x))$.

Note that in the case that X is a Riemannian manifold, the conditions in (0.6.1) and (0.6.2) are equivalent, and Conjecture 0.6 holds in this case (Theorem B).

Roughly speaking, (0.6.2) says that the regular part of the singular nilpotent fibration coincides with one in Theorem 0.2, while the singular part is slightly ‘singular’ than one in Theorem 0.2 because a singular fiber is an infra-nilorbifold.

Next, we will give applications of Theorem B relating to recent work in [NZ] and [HW1,2] on a collapsed manifold M with Ricci curvature bounded in two sides or below, satisfying a local or global topological condition, respectively. In each work, mainly using analytic tools, it was proved that M satisfies a local or global bounded covering geometry, respectively. We will extend such bound covering property in a more general setting (Theorem C), and we will present a proof based on elementary tools in metric geometry.

The generalized Margulis lemma in [KW] asserts that there exist constants, $\epsilon(n), c(n) > 0$, such that for any point x in a complete n -manifold M of $\text{Ric}_M \geq -(n-1)$, the subgroup of $\pi_1(B_1(x), x)$, $\Gamma_x(\epsilon)$, generated by loops at x of length $< \epsilon \leq \epsilon(n)$ contains a nilpotent subgroup of index $\leq c(n)$ (Theorem 1.4.2). In particular, $\Gamma_x(\epsilon)$ contains a torsion-free nilpotent subgroup of finite index ([Ra]). The rank of a torsion-free nilpotent group is equal to the minimal number of generators, and the rank of $\Gamma_x(\epsilon)$ is defined as the rank of the torsion-free nilpotent subgroup of finite index. For the notions on pseudo-groups and their action, see Subsection 1.1 and Remark 1.1.5.

Let X be a Ricci limit space. According to [CoN], there is a unique integer k such that the subset of X consisting of points whose tangent cone is unique and is isomorphic to \mathbb{R}^k has a full renormalized measure. We will use the notation that $\dim(X) = k$.

Theorem C. (Local/global Ricci bounded covering geometry) *Let a sequence of compact n -manifolds, $(M_i, p_i) \xrightarrow{\text{GH}} (X, p)$, with $\text{Ric}_{M_i} \geq -(n-1)$, $\text{diam}(M_i) \leq d$.*

(C1) (Local rank estimate) *For $x_i \in M_i$, assume the following commutative dia-*

gram:

$$\begin{array}{ccc}
(\bar{B}_1(\tilde{x}_i), \Gamma_{x_i}(\epsilon)) & \xrightarrow{\text{eqGH}} & (\bar{B}_1(\tilde{x}), G) \\
\pi_i|_{B_1(\tilde{x}_i)} \downarrow & & \text{proj} \downarrow \\
\bar{B}_1(x_i) & \xrightarrow{\text{GH}} & B_1(x) = \bar{B}_1(\tilde{x})/G,
\end{array}$$

where $\pi_i : (\widetilde{B_2(x_i)}, \tilde{x}_i) \rightarrow (B_2(x_i), x_i)$ is a Riemannian universal cover map, $\Gamma_{x_i}(\epsilon)$ acts pseudollocally on $B_1(\tilde{x}_i) \subset \widetilde{B_2(x_i)}$, and G , the limit of $\Gamma_{x_i}(\epsilon)$, is a pseudo-group of isometries. Then $\text{rank}(\Gamma_{x_i}(\epsilon)) \leq \dim(G)$, and $\text{rank}(\Gamma_{x_i}(\epsilon)) = n - \dim(X)$ implies that $\text{vol}(B_1(\tilde{x}_i)) \geq v(X) > 0$.

(C2) (Global abelian rank estimate) Given the following commutative diagram,

$$\begin{array}{ccc}
(\tilde{M}_i, \tilde{p}_i, \Gamma_i) & \xrightarrow{\text{eqGH}} & (\tilde{X}, \tilde{p}, G) \\
\pi_i \downarrow & & \text{proj} \downarrow \\
(M_i, p_i) & \xrightarrow{\text{GH}} & (X = \tilde{X}/G, p),
\end{array}$$

the first Betti numbers, $b_1(M_i) - b_1(X) \leq \dim(G)$, and $b_1(M_i) - b_1(X) = n - \dim(X)$ implies that $\text{vol}(B_1(\tilde{p}_i)) \geq v(X) > 0$.

In (C1), the pseudo-group G contains a neighborhood of the identity that isomorphically embeds into a Lie group \hat{G} , and $\dim(G) := \dim(\hat{G})$.

Remark 0.7. (0.7.1) If one assumes that X is Riemannian manifold i.e., a collapsed M is d_{GH} -close to a non-collapsed Riemannian manifold, then one may state the non-collapsing on a local or global Riemannian universal cover in (C1) or (C2), without involving a GH-convergent sequence ([NZ], [HW1]).

(0.7.2) If X is a compact Riemannian m -manifold, the case of (C1) that $|\text{Ric}_M| \leq (n-1)$ was obtained in [NZ], and if X is an orbifold, (C2) was obtained in [HW2].

A consequence of Theorem A and (C1) is

Theorem 0.8. ([Co2]) *Given n , there exists $\delta(n) > 0$ such that if a compact n -manifold satisfies*

$$\text{Ric}_M \geq -(n-1), \quad \text{diam}(M) \leq \delta(n), \quad b_1(M) = n,$$

then M is diffeomorphic to a flat torus.

A consequence of Theorem B and (C1) is

Theorem 0.9. (Local maximal rank and nilpotent fiber bundles) *Assume that compact n -manifold M and m -manifold N ($m < n$) satisfy*

$$\text{Ric}_M \geq -(n-1), \quad |\text{sec}_N| \leq 1, \quad \text{injrads}(N) \geq 1, \quad d_{GH}(M, N) < \epsilon \leq \epsilon(n),$$

where $\epsilon(n)$ is the constant in Margulis lemma in [KW]. Then $\text{rank}(\Gamma_x(\epsilon)) \leq n - m$, and “=” for all $x \in M$ implies a smooth fiber bundle, $f : M \rightarrow N$, with fiber diffeomorphic to an infra-nilmanifold, and f is an $\Psi(\epsilon|n)$ -GHA.

The case of Theorem 0.9 that $|\text{Ric}_M| \leq n - 1$ can be easily seen from [NZ].

A consequence of Theorem B and (C2) is:

Theorem 0.10. (Generalized maximal first Betti number rigidity, [HW1]) *Assume that compact n -manifold M and m -manifold N ($m < n$) satisfy*

$$\text{Ric}_M \geq -(n-1), \quad |\text{sec}_N| \leq 1, \quad \text{injrads}(N) \geq 1, \quad d_{\text{GH}}(M, N) < \epsilon \leq \epsilon(n).$$

Then the first Betti numbers, $b_1(M) - b_1(N) \leq n - m$, and “=” implies a smooth torus fiber bundle, $f : M \rightarrow N$, with an affine structural group, and f is an $\Psi(\epsilon|n)$ -GHA.

Theorem 0.10 may be viewed as a bundle version of Theorem 0.8.

Observe that the case of Theorem 0.10 with bounded sectional curvature follows easily from Theorem 0.2. A key in the proof is to show that \tilde{M} is not collapsed, based on which a smoothing via Ricci flows applies ([HW2]).

We will conclude the introduction with an indication of our approach to Theorems A and B (see Section 2 for a detailed outline).

By the Gromov’s compactness, it is equivalent to consider a sequence of compact n -manifolds, $M_i \xrightarrow{\text{GH}} \text{pt}$, with local (ρ, v) -bound Ricci covering geometry, and show that for i large, M_i is diffeomorphic to an infra-nilmanifold (Theorem 0.3’).

The starting point in our proof is the generalized Margulis lemma in [KW] (Theorem 1.4.2), by which M_i has a bounded normal covering space, $M_{i,0}$, with $\pi_1(M_{i,0})$ a nilpotent group. The proof consists of two steps: (1) (the main work) we prove that $M_{i,0}$ is diffeomorphic to a nilmanifold, by exploring a sequence of sub-bundle structures on $M_{i,0}$ such that the adjacent orbit space is a principal tori bundle over the other (Theorems 2.1, 2.2). (2) we prove that M_i is diffeomorphic to an infra-nilmanifold: viewing \tilde{M}_i as a simply connected nilpotent Lie group, by the Malcév rigidity there is a $\Gamma_i = \pi_1(M_i)$ -action on \tilde{M}_i which preserves the sub-bundle structures. We will construct a Γ_i -conjugation on \tilde{M}_i via defining a left invariant metric on \tilde{M}_i , based on the geometry of a short basis in $\pi_1(M_{i,0})$, and applying a generalized stability of isometric Lie group actions (Lemma 3.5.1).

The sequence of sub-bundle structures on $M_{i,0}$ is constructed through a successively blowing up at every point in $M_{i,0}$, with rates obtained from a short basis at the point, and the nilpotency of $\pi_1(M_{i,0})$ guarantees that at each blow up $M_{i,0}$ is locally close (in the Gromov-Hausdorff distance) to a metric product of a flat torus and an Euclidean space (Theorem 2.4), which is used to construct a locally finite open cover on $M_{i,0}$, each carries an elementary T-structure. We will glue these elementary T-structures to a fiber bundle global ([CG2], [GK], [Pa1]), and the metric regularity required for the patching are that all points on the Riemannian universal cover are uniform (arbitrarily) Reifenberg (Lemma 2.3), which is achieved based on the Cheeger-Colding-Naber theory on the degeneration of metrics with Ricci curvature bounded below ([CC1-3], [Co1,2], [CN], [CJN]).

In the proof of Theorem B, by performing a similar successive blow ups as in the proof of Theorem A, we obtain a locally finite open cover on M with elementary N-structure, $f_\alpha : U_\alpha \rightarrow B_\alpha$, where f_α is a smooth trivial bundle map and $\Psi(\epsilon|n, v)$ -GHA, with fiber an infra-nilmanifold, such that on overlaps, elementary N-structures satisfy the patching criterion in [CFG]. Hence we are able glue elementary N-structures into a nilpotent fiber bundle with an affine structural group.

Finally, we pointed out the our proofs of Theorems A and B are mainly local arguments, hence they can be applied to a region in a collapsed manifold where a

local Ricci bounded covering geometry is satisfied (e.g., [GW], [GTZ1,2], [HSVZ], [Zh]).

The rest of the paper will be organized as follows:

In Section 1, we will review basic notions and results that will be used in the rest of the paper.

In Section 2, we will outline a proof of Theorem 0.3 that does not rely on Theorem 0.1.

In Section 3, we will prove Theorem A.

In Section 4, we will prove Theorem B.

In Section 5, we will prove Theorem C, Theorem 0.8 and Theorem 0.10.

1. PRELIMINARIES

We will review basic notions and properties that will be used in the rest of the paper. We will also present examples mentioned in the introduction.

1.1. Equivariant Gromov-Hausdorff convergence.

The reference of this subsection is [FY] (cf. [Ro2]).

Let X_i and X be compact length spaces. We say that X_i converge to X in the Gromov-Hausdorff topology, denoted by $X_i \xrightarrow{\text{GH}} X$, if there are a sequence $\epsilon_i \rightarrow 0$ and a sequence of maps, $h_i : X_i \rightarrow X$, such that $|d_X(h_i(x_i), h_i(x'_i)) - d_{X_i}(x_i, x'_i)| < \epsilon_i$ (ϵ_i -isometry), and for any $x \in X$, there is $x_i \in X_i$ such that $d_X(h_i(x_i), x) < \epsilon_i$ (ϵ_i -onto), and h_i is called an ϵ_i -Gromov-Hausdorff approximation, briefly, ϵ_i -GHA.

Assume that X_i admits a closed group Γ_i -action by isometries. Then $(X_i, \Gamma_i) \xrightarrow{\text{GH}} (X, \Gamma)$ means that there are a closed group of isometries Γ on X , a sequence $\epsilon_i \rightarrow 0$ and a sequence of triple of ϵ_i -GHA, (h_i, ϕ_i, ψ_i) , $h_i : X_i \rightarrow X$, $\phi_i : \Gamma_i \rightarrow \Gamma$ and $\psi_i : \Gamma \rightarrow \Gamma_i$, such that for all $x_i \in X_i, \gamma_i \in \Gamma_i$ and $\gamma \in \Gamma$,

$$(1.1.1) \quad d_X(h_i(x_i), \phi_i(\gamma_i)h_i(\gamma_i^{-1}(x_i))) < \epsilon_i, \quad d_X(h_i(x_i), \gamma^{-1}(h_i(\psi_i(\gamma)(x_i)))) < \epsilon_i,$$

where Γ_i and Γ are equipped with the induced metrics from X_i and X . We call (h_i, ϕ_i, ψ_i) an ϵ_i -equivariant GHA.

When X is not compact, then the above notion of equivariant convergence naturally extends to a pointed version $(h_i, \phi_i, \psi_i): h_i : B_{\epsilon_i^{-1}}(p_i) \rightarrow B_{\epsilon_i^{-1} + \epsilon_i}(p)$, $h_i(p_i) = p$, $\phi_i : \Gamma_i(\epsilon_i^{-1}) \rightarrow \Gamma(\epsilon_i^{-1} + \epsilon_i)$, $\phi_i(e_i) = e$, $\psi_i : \Gamma(\epsilon_i^{-1}) \rightarrow \Gamma_i(\epsilon_i^{-1} + \epsilon_i)$, $\psi_i(e) = e_i$, and (1.1.1) holds whenever the multiplications stay in the domain of h_i , where $\Gamma_i(R) = \{\gamma_i \in \Gamma_i, d_{X_i}(p_i, \gamma_i(p_i)) \leq R\}$.

Lemma 1.1.2. *Let $(X_i, p_i) \xrightarrow{\text{GH}} (X, p)$, where X_i is a complete locally compact length space. Assume that Γ_i is a closed group of isometries on X_i . Then there is a closed group G of isometries on X such that passing to a subsequence, $(X_i, p_i, \Gamma_i) \xrightarrow{\text{GH}} (X, p, G)$.*

Lemma 1.1.3. *Let $(X_i, p_i, \Gamma_i) \xrightarrow{\text{GH}} (X, p, \Gamma)$, where X_i is a complete locally compact length space and Γ_i is a closed subgroup of isometries. Then $(X_i/\Gamma_i, \bar{p}_i) \xrightarrow{\text{GH}} (X/\Gamma, \bar{p})$.*

Assume that the universal covering space, $\pi_i : (\tilde{X}_i, \tilde{p}_i) \rightarrow (X_i, p_i)$, exists, and let $\Gamma_i = \pi_1(X_i, p_i)$ denote the fundamental group.

Lemma 1.1.4. *Let $X_i \xrightarrow{\text{GH}} X$ be a sequence of compact length metric space. Then passing to a subsequence such that the following diagram commutes,*

$$\begin{array}{ccc} (\tilde{X}_i, \tilde{p}_i, \Gamma_i) & \xrightarrow{\text{GH}} & (\tilde{X}, \tilde{p}, \Gamma) \\ \downarrow \pi_i & & \downarrow \pi \\ (X_i, p_i) & \xrightarrow{\text{GH}} & (X, p). \end{array}$$

If X is compact and Γ/Γ_0 is discrete, then there is $\epsilon > 0$ such that the subgroup, $\Gamma_i(\epsilon)$, generated by elements with displacement bounded above by ϵ on $B_{2d}(\tilde{p}_i)$, is normal and for i large, $\Gamma_i/\Gamma_i(\epsilon) \cong^{\text{isom}} \Gamma/\Gamma_0$.

Remark 1.1.5. Inspecting proofs of Lemma 1.1.2-1.1.4, the whole group structure of Γ_i is not needed; the multiplication of elements in $\Gamma_i(\epsilon_i^{-1})$ is taken into account only if the product moves a point in $B_{\epsilon_i^{-1}}(p_i)$ and remained in the ball. We will call $\Gamma_i(R) = \{\gamma_i \in \Gamma_i, d(\gamma_i(p_i), p_i) \leq R\}$ is a pseudo-group, and the $\Gamma_i(R)$ -action (resp. the Γ_i -action) on $B_{\epsilon_i^{-1}}(p_i)$ a pseudo-action. Clearly, Lemma 1.1.2-1.1.4 hold when replacing Γ_i by $\Gamma_i(\epsilon_i^{-1})$ and the Γ_i -actions by pseudo-actions.

1.2. Patching atlas of elementary F-structures and N-structures.

In our proof of Theorem A, a basic tool is the patching technique in [CG2] on gluing an atlas of elementary F-structures on a manifold M to form a (mixed) F-structure (see below), and in the proof of Theorem B, we employ the generalized patching techniques in [CFG] to glue an atlas of elementary N-structures to obtain the desired nilpotent fiber bundle.

Indeed, we need only the simplest form of the above patching technique: patching local elementary T^k -structure by free T^k -actions (k is fixed) (resp. local nilpotent fiber bundles), which are C^1 -close on overlaps, to form an affine T^k -bundle i.e., T^k -fibers are orbits of local free T^k -actions which are commutative on the overlap (resp. to form a nilpotent fiber bundle on M). Hence the structure group of the bundle is affine.

Let M be a Riemannian n -manifold, and let $U \subseteq M$ be an open subset. An elementary N-structure on U , denoted by (U, \mathcal{N}) , is defined by an $O(n)$ -invariant fiber bundle on the frame bundle over U , $F \rightarrow (F(U), O(n)) \xrightarrow{f} (Y, O(n))$, where $F = N/\Lambda$ is a compact nilmanifold, and the $O(n)$ -action preserves F -fibers as well as the nilpotent structure on F . The $O(n)$ -invariance of f descends the F -bundle to a map, $\bar{f} : U \rightarrow X = Y/O(n)$, such that the following diagram commutes,

$$\begin{array}{ccc} F \rightarrow (F(U), O(n)) & \xrightarrow{f} & (Y, O(n)) \\ \pi \downarrow & & \text{proj} \downarrow \\ \bar{F} \rightarrow U & \xrightarrow{\bar{f}} & X = Y/O(n). \end{array}$$

We will call the π -image of F an orbit on U , and U decomposes into orbits. Note that the property that U admits a pure N-structure does not dependent on the

metric on M , because given M another metric, its frame bundle on U is equivalent to $(F(U), O(n))$. If $F = T^k$, then we will call this elementary N-structure an elementary F-structure, denoted by (U, \mathcal{F}) . If the T^k -bundle is also principal, we will call it an elementary T-structure; note that the free T^k -action on $F(U)$ descends to a T^k -action on U (which may not be free).

An N-structure on a manifold M is given by a collection of elementary N-structures, $(M, \mathcal{N}) = \{(U_i, \mathcal{N}_i)\}$, where $\{U_i\}$ form a locally finite open cover for M , which satisfy the following compatibility condition: if $U_i \cap U_j \neq \emptyset$, then $U_i \cap U_j$ is both \mathcal{N}_i - and \mathcal{N}_j -invariant, and on $F(U_i \cap U_j)$, the F_i -bundle is a sub-bundle of the F_j -bundle, or vice versa. When $U_i \cap U_j \neq \emptyset$, the F_i -bundle and F_j -bundle coincide, we will call \mathcal{N} a pure N-structure, and otherwise \mathcal{N} is called a mixed N-structure.

For any $x \in M$, we will call the orbit at x of any elementary N-structure \mathcal{N}_i containing x with the maximal dimension the N -orbit at x , denoted by \mathcal{O}_x . Clearly, M decomposes into disjoint N-orbits. The minimal dimension of N-orbits is called the rank of \mathcal{N} . A metric on M is called invariant, if for a canonical metric on $F(U_i)$, the induced metric on F_i is left invariant. If M admits an N-structure, then M always admits an invariant metric ([CR], [Ro2]).

Note that the case that a pure T^k -structure (T-structure) with every elementary T^k -structure defined by a free T^k -action form an affine T^k -bundle on M .

Theorem 1.2.1. *Let M be a manifold.*

(1.2.1.1) ([CG1]) *If M admits an F-structure of positive rank, then M admits one parameter family of invariant metrics, g_ϵ , such that $|\sec_{g_\epsilon}| \leq 1$, $\text{diam}(\mathcal{O}_x) \rightarrow 0$ as $\epsilon \rightarrow 0$, for all $x \in M$ (we refer such a collapse as long the F-structure).*

(1.2.1.2) ([CR]) *If M admits an N-structure of positive rank, then M admits one parameter family of invariant metrics, g_ϵ , such that $|\sec_{g_\epsilon}| \leq 1$ and $\text{diam}(\mathcal{O}_x) \rightarrow 0$ as $\epsilon \rightarrow 0$, for all $x \in M$.*

Theorem 1.2.2. *There exists a constant $\epsilon(n) > 0$ such that if M is a complete n -manifold satisfying $|\sec_M| \leq 1$ and $\text{vol}(B_1(x)) < \epsilon \leq \epsilon(n)$ for all $x \in M$, then*

(1.2.2.1) ([CFG]) *M admits a positive rank N-structure, \mathcal{N} , such that each \mathcal{N} -orbit has diameter $< \Psi(\epsilon)$, and there is a C^1 ϵ -close invariant metric, g_ϵ , with higher regularity.*

(1.2.2.2) ([CG2]) *The center part of the nilpotent structure of \mathcal{N} form an F-structure of positive rank.*

A proof of Theorem 1.2.2 is based on Theorem 0.2 and the patching techniques in [CG2]; which is an extension of the stability of compact Lie group actions ([Pa], [GK]).

To construct a nilpotent structure on a collapsed M in Theorem 1.2.2, one first covers M with locally bounded number of unit balls, and for each ball, apply a local version of Theorem 0.2 on an open subset, $B_1(x_i) \subset U_i \subset B_2(x_i)$, to obtain an elementary N-structure on U_i . Then one gets an open cover for M with elementary N-structure, $\{(U_i, \mathcal{N}_i)\}$, and on $U_i \cap U_j \neq \emptyset$, these elementary N-structures are C^1 -close in a suitable sense similar to that of a compact Lie group actions considered by [GK] and [Pa]. Extending the patching technique in [CG2], one can glue these elementary structures to form an N-structure ([CFG]).

In our proof the proof of Theorem 0.3, we perform blow-up operations on M a finite number times, and each time we blow up at any $x \in M$ with a scale from lengths a short basis at x . By assuming that $\pi_1(M)$ is nilpotent ([KW]), each

blow-up operation results in a manifold with a local finite open cover by elementary T^k -structures defined by free almost isometric T^k -actions (k is independent of points).

For our purpose, we will only state the gluing result (Lemmas 1.4 and 1.5 in [CG2]) for the case of constructing a T -structure on M by patching a collection of elementary T -structures which are C^1 -close on overlaps.

Theorem 1.2.3. (Gluing, [CG2]) *Let M be an n -manifold. Given $0 < \epsilon < 2\rho$, there exists a constant $\delta = \delta(n, \rho, \epsilon) > 0$ such that M admits a T^k -bundle with affine structural group, if M admits a collection of elementary T^k -structures, $\{(U_i, g_i, T_i^k)\}$ satisfying*

(1.2.3.1) *(U_i, T_i^k) is an elementary T -structure with a free T_i^k -action and invariant metric g_i such that $\{U_i^\rho\}$ is locally finite open cover for M and*

$$|\sec_{g_i}| \leq 1, \quad \text{injrads}(g_i) \geq \frac{1}{10},$$

where $U_i^\rho \subset U_i$ consisting of points distance $\geq \rho$ from $\partial\bar{U}_i$.

(1.2.3.2) *If $U_i^\rho \cap U_j^\rho \neq \emptyset$, then the T_i^k - and T_j^k -actions on $U_i^\rho \cap U_j^\rho$ is δ C^1 -close i.e.,*

$$e^{-\epsilon}g_i \leq g_j \leq e^\epsilon g_i, \quad \eta(t) = \max_{x \in U_i^\rho \cap U_j^\rho} \{d(t_i^{-1}t_j(x), x), t \in B_\rho(e)\} < \delta,$$

where T^k the product of unit circle and $B_\rho(r)$ the ρ -ball of centered at the identity element e .

Remark 1.2.4. The C^1 closeness condition in Theorem 1.2.3 is a modification of one in [GK] and [Pa] where the C^1 -closeness is for two actions of a compact Lie group on a manifold, here $U_i^\rho \cap U_j^\rho \neq \emptyset$ may neither T_i^k - nor T_j^k -invariant. The C^1 -closeness condition can be achieved on a collapsed manifold M in Theorem 1.2.2 via locally averaging the metric under the local almost isometric tori actions. Note that Theorem 1.2.3 can be applied on a collapsed M with Ricci curvature bounded below as long as U_i and U_j admit T^k -invariant metrics satisfying conditions in Theorem 1.2.3, even the invariant metrics may not be C^1 -close the induced metrics from M .

A prototype of our proof of Theorem 0.3' is seen by applying Theorem 1.2.3 to a sequence, $M_i \xrightarrow{\text{GH}} N$, where $|\sec_{M_i}| \leq 1$ and N is a compact Riemannian manifold. By Theorem 0.2, there is a fiber bundle map, $f_i : M_i \rightarrow N$, with fiber an infra-nilmanifold and affine structural group.

Let's apply the construction of an F-structure on M_i in [CG2]: M_i carries an F-structure \mathcal{F} of positive rank whose an orbit points the most collapsed directions. By the diameter bound (i.e., collapsing is taking place with a bounded diameter), \mathcal{F} is pure, and because N is a manifold, \mathcal{F} is polarized and is a fiber bundle with a flat fiber. Indeed, as pointed in [CFG] (p.330) that \mathcal{F} is a part in the centers of the infra-nilmanifold fibers, thus form a sub-bundle with fiber a flat manifold.

Inspecting the construction of F-structure in [CG2], one first gets local F-structure around each $x_i \in M_i$, by blowing up M_i by the inverse of $\ell(x_i) = \text{injrads}(M_i, x_i)$, and a limit,

$$(\ell(x_i)^{-1}M_i, x_i) \xrightarrow{\text{GH}} (Y, x),$$

is a non-compact flat manifold with a compact soul, a flat manifold F ; and an open subset at x_i that is diffeomorphic to a neighborhood of F in Y defines an elementary F-structure at x_i . By (0.2.1.3), the induced metrics on f_i -fibers are comparable, and in particular, one obtains a local finite open cover of M_i by $\{U_{i,j} \simeq T^{k_s} \times \mathbb{B}_{100}^{n-k_s}(0)\}$, and on overlaps the conditions Theorem 1.2.3 is satisfied, where $\mathbb{B}_r^m(0)$ denotes an m -dimensional Euclidean r -ball.

Corollary 1.2.5. *Let $f_i : M_i \rightarrow N$ be as in the above. Then there is a sub-bundle, $f_{i,s} : M_i \rightarrow B_{i,s}$, such that a $f_{i,s}$ -fiber is a flat manifold and affine structural group.*

In [CFG] (Part III, sections 5-8), Theorem 1.2.3 is generalized to glue elementary N-structures, which is more technique. Note that a gluing of elementary N-structures is carried out on local frame bundles (so one has to make gluing $O(n)$ -invariant), and the stability of a compact Lie group actions ([GK], [Pa]) cannot be applied because a (non-abelian) nilpotent Lie group is not compact. Instead, the stability of elementary N-structures also uses the Malcev's rigidity theorem for discrete co-compact subgroups of nilpotent Lie groups.

Given two elementary N-structures, (U_1, \mathcal{N}_1) and (U_2, \mathcal{N}_2) , with $U_1 \cap U_2 \neq \emptyset$. Assume that there is an open ball $B_r \subset U_1 \cap U_2 \neq \emptyset$ such that the \mathcal{N}_i -orbits on B_r is contained in $U_1 \cap U_2$. Then the two elementary N-structures are C^1 -close, if U_i admits an invariant Riemannian metric g_i with $|\sec_{g_i}| \leq 1$ such that the pullback metric \tilde{g}_1 and \tilde{g}_2 on the universal cover, $\widetilde{U_1 \cap U_2}$, are C^1 -close restricting to any ball always from the boundary of $\widetilde{U_1 \cap U_2}$.

Remark 1.2.6. The above C^1 -closeness is a simplest case for the gluing criterion in [CFG] (Part III, sections 5-8). Indeed, in our proof Theorem B, the orbits on U_i form a trivial fiber bundle over a unit Euclidean ball i.e., $f_i : U_i \rightarrow \mathbb{B}_1^m(0)$, thus $\Gamma = \pi_1(U_1) = \pi_1(U_2)$, which is the fundamental group of f_i -fiber.

To avoid a technique complexity, we will state the most general gluing criterion in [CFG], because our gluing charts are of the simplest form, a trivial bundle with fiber a fixed infra-nilmanifold, a gluing technique presented in Appendix 2 of [CFG] is all we need in the proof of Theorem B (Method 2).

1.3. Cheeger-Colding-Naber theory on Ricci limit spaces.

A length space, (X, p) , is called a Ricci limit space, if there is a sequence of pointed complete n -manifolds, $(M_i, p_i) \xrightarrow{\text{GH}} (X, p)$, with $\text{Ric}_{M_i} \geq -(n-1)$.

Observe that the most significant difference, from a limit space of a sequence of n -manifolds M_i with sectional curvature, $\sec_{M_i} \geq -1$, is a weak link between large scale geometry and small scale geometry; for instance, the tangent cone at $x \in X$ is not unique, and may not be a metric one.

Two basic tools in the Cheeger-Colding-Naber theory are two quantitative rigidity results on manifolds with Ricci curvature bounded below: the Splitting theorem of Cheeger-Gromoll, and the rigidity maximal volume rigidity.

Theorem 1.3.1. (Quantitative splitting, [CC1], [CoN]) *Given $n, \delta, \epsilon, L > 0$, there exists a constant such that if M is a complete n -manifold, $p, q, r \in M$ such that $E(p) = d(p, q_-) + d(p, q_+) - d(q_-, q_+) < \epsilon$, $d(p, q_{\pm}) \geq L$, then*

$$d_{\text{GH}}(B_R(p), B_R(X \times \mathbb{R})) < \Psi(\delta, \epsilon, L^{-1}|n, R),$$

where $0 < R \ll L$ and X is a compact length space.

Corollary 1.3.2. (Splitting of Ricci limit space) *Let $(M_i, p_i) \xrightarrow{\text{GH}} (X, p)$ such that $\text{Ric}_{M_i} \geq -(n-1)\delta_i \rightarrow 0$. If X contains a line, then X splits with a \mathbb{R}^1 -factor, $X = Y \times \mathbb{R}^1$.*

Theorem 1.3.3. (Volume convergence, [Co1], [CC1]) *Let $(M_i, p_i) \xrightarrow{\text{GH}} (X, p)$ be sequence of complete n -manifolds such that*

$$\text{Ric}_{M_i} \geq -(n-1), \quad \text{vol}(B_1(p_i)) \geq v > 0.$$

For any $M_i \ni x_i \rightarrow x \in X$ and any $R > 0$, $\text{vol}(B_R(x_i)) \rightarrow \text{Haus}^n(B_R(x))$ as $i \rightarrow \infty$, where Haus^n denotes the n -dimensional Hausdorff measure.

Theorem 1.3.4. (Almost volume balls are almost metric balls, [CC1]) *Given $n, \epsilon > 0$, there exists $\delta = \delta(n, \epsilon) > 0$ such that if a complete n -manifold M satisfies*

$$\text{Ric}_M \geq -(n-1)\delta, \quad \text{vol}(B_1(x)) \geq (1-\delta) \text{vol}(B_1^n(0)),$$

then $d_{\text{GH}}(B_r(x), B_r^n(0)) < \epsilon \cdot r$ for all $0 < r \leq 1$ (we refer x as and a $(n, \epsilon, 1)$ -Reifenberg point).

For any $q \in X$ and any $\epsilon_i \rightarrow 0$, passing to a subsequence we call the limit space, $(\epsilon_i^{-1}X, q) \xrightarrow{\text{GH}} (T_qX, q)$ a tangent cone of X at q . A tangent cone may depend on the subsequence. A point q is called regular, if T_qX is unique and is isometric to \mathbb{R}^k , $k \leq n$. We call $\mathcal{S} = X - \mathcal{R}$ the singular set.

Theorem 1.3.5. (Regularity/singularity) *Let $(M_i, p_i) \xrightarrow{\text{GH}} (X, p)$ with $\text{Ric}_{M_i} \geq -(n-1)$. Then*

(1.3.5.1) ([CC1]) *The set of regular points \mathcal{R} is dense in X .*

(1.3.5.2) ([CC1]) *If $\text{vol}(B_1(p_i)) \geq v > 0$, then $\dim_H(\mathcal{S}) \leq n-2$.*

(1.3.5.3) ([ChN]) *If $\text{vol}(B_1(p_i)) \geq v > 0$ and $\text{Ric}_{M_i} \leq n-1$, then \mathcal{R} is an open $C^{1,\alpha}$ -manifold, and $\dim_H(\mathcal{S}) \leq n-4$.*

Theorem 1.3.6. (Diffeomorphic stability, [CC2], [CoN]) (1.3.6.1) *Let $M_i \xrightarrow{\text{GH}} M_0$ such that M_i is a compact n -manifold with $\text{Ric}_{M_i} \geq -(n-1)$ and M_0 is a compact n -manifold. Then for i large, M_i is diffeomorphic to M_0 by a ϵ_i -GHA, $\epsilon_i \rightarrow 0$.*

(1.3.6.2) *Let $(M_i, p_i) \xrightarrow{\text{GH}} (M_0, p)$ such that M_i is a complete n -manifold with $\text{Ric}_{M_i} \geq -(n-1)$ and M_0 is a complete n -manifold. Given any $R > 0$, for i large $B_R(p_i)$ is diffeomorphic to $B_R(p)$ by a pointed ϵ_i -GHA.*

Theorem 1.3.7. (Limit groups are Lie groups, [CC2], [CoN]) *Let X be a Ricci limit space. Then the isometry group of X is a Lie group.*

We mention that in this paper, (1.3.5.2) and (1.3.5.3) are not used.

1.4. The Margulis Lemma.

In our proof of Theorem 0.3, we will perform a successive blow-ups on a suitable bounded covering space of M_i in Theorem 0.3' so that limit spaces are flat tori; which is based the generalized Margulis lemma in [KW], see Theorem 1.4.2.

Theorem 1.4.1. (Short generators estimate, [KW]) *Let M be a complete n -manifold with $\text{Ric}_M \geq -(n-1)$. For any $p \in M$, there is a subset of almost full measure, $A \subseteq B_1(p)$, such that for $q \in A$, the short generators for $\pi_1(M, q)$ with length $\leq R$ is bounded above by a constant $C(n, R)$.*

Theorem 1.4.1 is required in proving the following:

Theorem 1.4.2. (Generalized Margulis lemma, [KW]) *Given n , there exist constants, $\epsilon(n), c(n) > 0$, such that if M is a complete n -manifold with $\text{Ric}_M \geq -(n-1)$, then for any $x \in M$, the subgroup of $\pi_1(B_1(x), x)$ generated by loops of length $\leq \epsilon(n)$ contains a normal nilpotent subgroup of index $\leq c(n)$.*

We now give a brief description on a general approach to a Margulis lemma ([KW], [KPT], [FY], [XY]), assuming Theorem 1.4.1.

Let's start with an algebraic criterion for a finitely generated discrete group Γ to contain a nilpotent subgroup of a finite index:

(1.4.3.1) There is a descending sequence of normal subgroups,

$$\Gamma \triangleright \Gamma_1 \triangleright \cdots \triangleright \Gamma_s \triangleright \Gamma_{s+1} = e, \quad \Gamma_j / \Gamma_{j+1} \cong \mathbb{Z}^{k_j}, \quad 1 \leq j \leq s.$$

(1.4.3.2) The induced homomorphism via conjugation, $\rho_j : \Gamma \rightarrow \text{Aut}(\Gamma_j / \Gamma_{j+1})$, has a finite image.

Let $\Gamma_p(\epsilon)$ denote the subgroup generated by loops at a point p of length $< \epsilon$. By Theorem 1.4.1, one may assume that p is chosen such that the number of the Gromov short generators for $\Gamma_p(\epsilon)$ is bounded by $c(n)$. A basic property is that $\Gamma_p(\epsilon)$ contains a normal subgroup of bounded index such that the quotient group contains no element of uniformly small displacement on any open subset of the quotient space, which is based on that the isometry group of a Ricci limit space is a Lie group ([CC1], [CoN]). We now use $\Gamma(\epsilon)$ to denote its normal subgroup of bounded index.

With $0 < \epsilon \leq \epsilon(n)$, via a successively blowing ups argument one sees that the Gromov's short generators for $\Gamma(\epsilon)$ with norm in a decreasing order and $|\gamma_{1,1}| \leq \epsilon(n)$, are naturally divided in terms of gaps in their length (norm),

$$\gamma_{1,1}, \dots, \gamma_{1,k_1}, \quad \gamma_{2,k_1+1}, \dots, \gamma_{2,k_1+k_2}, \quad \dots, \quad \gamma_{s,k_{s-1}+1}, \dots, \gamma_{s,k_1+\dots+k_s},$$

such that

$$1 \leq \frac{|\gamma_{i,\alpha+k}|}{|\gamma_{i,\alpha}|} \leq c(n), \quad \frac{|\gamma_{i+k,\alpha}|}{|\gamma_{i,\beta}|} \rightarrow 0,$$

Let Γ_j denote the subgroup generated by short generators in blocks $s, s-1, \dots, j$.

The main technical work in a proof of Theorem 1.4.2 is to deform a short path from a point to another point so that the length distortion is proportional to the distance between the two points; once this is achieved, one is able to verify (1.4.3.1) and (1.4.3.2). Roughly, the deformation is by the gradient flows of certain functions constructed based on the geometry of $r^{-1}M$, where r is proportional to the norm of short generators in some block. Precisely, combining the holonomy distortion estimate and the norm gaps in blocks one is able to conclude that each Γ_j is normal in Γ , and the holonomy representation, $\phi_j : \Gamma \rightarrow \text{Aut}(\Gamma_j / \Gamma_{j+1})$, has a (bounded) finite image. Moreover, one can show that Γ_j / Γ_{j+1} contains a free abelian group of (bounded) finite index.

Let's analyze the gradient flows on manifolds with sectional (resp. Ricci) curvature bounded below.

(1.4.4.1) Assume that $\text{sec}_M \geq -1$. The main technique, a gradient push introduced in [KPT] (cf. [XY]), allows one deform a path between any two (m, δ) -strained points in the scale of $r^{-1}M$. A point p is called an (m, δ) -strained on $r^{-1}M$ if

there are $2m$ -points, (a_i^\pm, b_i^\pm) , with $d(p, \pm a_i), d(p, \pm b_i) \geq \rho > 0$, $\angle(a_i^+ p a_i^-) > \pi - \delta$ and $\angle(a_i^\pm p b_j^\pm) > \frac{\pi}{2} - \delta$. The gradient push is the gradient flows of the distance function at some point from (a_i^\pm, b_i^\pm) , and the distance distortion is exponentially proportional to the time of gradient flows. Note that along the gradient push one has to change distance functions from time to time, determined by each blow-up rate, the set of (m, δ) -strained point on $r^{-1}M$ is path connected, open and dense, hence one can further gradient push a path to a point which is not (m, δ) -strained (i.e., relative singular).

(1.4.4.2) Assume that $\text{Ric}_M \geq -(n-1)$. In general, there is no control from a large scale geometry on a small scale geometry (e.g., contrasts to $\text{sec}_M \geq -1$, the sizes of a geodesic triangle do not control angles), so the gradient-push technique is replaced by the so-called a zooming-in map in [KW], based on (k, δ) -splitting maps ([CC]). With a similar blow-up rate as in (1.4.1.1), one obtains a (k, δ) -splitting map, whose gradients are almost orthonormal in the L^2 -sense. The weak control on gradient leads to a possible vanishing of a gradient at some point, or a (k, δ) -splitting map on $B_1(p)$ may not be a (k, δ) -splitting map at $q \in B_1(p)$ in a smaller scale at q . As a result, for the holonomy estimate to hold, one has to exclude certain points.

Remark 1.4.5. In [CJN], it was proved that an (n, δ) -splitting map is everywhere non-degenerate. In our proof of Theorem A, we show that the points on the local Riemannian universal cover of any ρ -ball in M_i are uniform (n, δ) -Reifenberg (Lemma 2.3), which is used to check that an (k, δ) -splitting map is non-generate everywhere ([Hu]). Combining the transformation technique in [CJN], it is expected that the zooming in maps works at every point, hence a holonomy estimate should hold at every point, similar to (1.4.4.1). Because our proof of Theorem 0.3' does not require this property, we will not pursuit a proof in this paper.

1.5. Collapsing with/without local bounded Ricci covering geometry.

We will supply examples mentioned in the previous discussions.

Example 1.5.1. (No nearby metrics with bounded sectional curvature) Given any sequence, $M_i \xrightarrow{\text{GH}} X$, of compact n -manifolds such that

$$\text{Ric}_{M_i} \geq -(n-1), \quad \text{diam}(M_i) \leq d, \quad \text{vol}(M_i) \geq v > 0.$$

Then the metric product, $M_i \times \epsilon_i S^1 \xrightarrow{\text{GH}} X$ ($\epsilon_i \rightarrow 0$) satisfies the local $(1, v)$ -bounded Ricci covering geometry.

If X is not a topological manifold ([An], [Ot], [Per]), then $M_i \times \epsilon_i S^1$ admits no nearby metric with bounded sectional curvature. Otherwise, the collapsing structure on $M_i \times \epsilon_i S^1$ with respect to the nearby metric with bounded sectional curvature has to be the free S^1 -action (Theorem 0.2), thus the orbit space, $(M_i \times \epsilon_i S^1)/S^1 = M_i$, is homeomorphic to X ; a contradiction.

Example 1.5.2. (Collapsing with smooth limits, no fibration, [An]) If one removes the volume condition in Theorem 0.4, then there is no fiber bundle map.

Anderson ([An]) constructed one family of metrics g_δ on a compact n -manifold M with a nontrivial second homotopy group such that

$$(M, g_\delta) \xrightarrow{\text{GH}} T^{n-1} (\text{flat}), \quad 0 < \delta \leq \text{Ric}_{g_\delta} \leq 1.$$

There is no S^1 -fiber from (M, g_δ) to T^{n-1} ; otherwise all $\pi_q(M) = 0$ for $q \geq 2$.

Example 1.5.3. (Non-smooth/smooth limit vs. infra-nil/nilpotent fundamental groups)

Let (K^2, g) denote a Klein bottle, and let S^1 acts on K^2 by isometries with two exceptional S^1 -orbits with isotropy \mathbb{Z}_2 . Hence, $K^2/S^1 = [0, 1]$. Decomposes g_0 at each point as $g = ds^2 + g^\perp$, where ds^2 denote the restriction of g to the tangent space of S^1 -orbit, and g^\perp the orthogonal complement. For $\epsilon \in (0, 1]$, let $g_\epsilon = \epsilon^2 ds^2 + g^\perp$. Then $(K^2, g_\epsilon) \xrightarrow{\text{GH}} [0, 1]$ with $\text{sec}_{g_\epsilon} \equiv 0$, and $(K^2, \sqrt{\epsilon}g_\epsilon) \xrightarrow{\text{GH}} \text{pt}$ with $\text{sec}_{\sqrt{\epsilon}g_\epsilon} \equiv 0$. When blowing up by $\text{diam}(K^2, \sqrt{\epsilon}g_\epsilon)^{-1} \approx \epsilon^{-\frac{1}{2}}$, $(K^2, g_\epsilon) \xrightarrow{\text{GH}} [0, 1]$, while the double cover, $(T^2, g_\epsilon) \xrightarrow{\text{GH}} S^1$.

Example 1.5.4. (Collapsing Calabi-Yau metrics along a nilpotent fibration with singular fibers not topological manifolds). Let M be a compact Kähler manifold. Assume there is a map, $f : M \rightarrow X$, where X is a smooth manifold except a finite isolated singular points, $\{x_i\}$, and $f : M - \bigcup_i f^{-1}(x_i) \rightarrow X - \{x_i\}$ is smooth nilpotent fibration, $f^{-1}(x_i)$ is a not submanifold of dimension $\dim(M) - \dim(X)$. In [GW], [GTZ1,2], [HSVZ], and [Zh], a one-parameter family of Calabi-Yau metrics g_ϵ are constructed so that $(M, g_\epsilon) \xrightarrow{\text{GH}} X$. It can be checked that away from the singular fibers, sectional curvature is bounded, thus a local Ricci bounded covering geometry is satisfied.

Indeed, with some minor work, a local version of Theorem B apply to the above situation, $f : M - \bigcup_i f^{-1}(x_i) \rightarrow X - \{x_i\}$ is a nilpotent fiber bundle.

2. OUTLINE OF PROOF OF THEOREM A

In this section, we will outline our proof of Theorem A, with details supplied in Section 3.

By the Gromov's pre-compactness, Theorem 0.3 has the following equivalent form (which we will present a proof independent of Theorem 0.1):

Theorem 0.3'. *Let a sequence of compact n -manifolds, $M_i \xrightarrow{\text{GH}} \text{pt}$, such that*

$$\text{Ric}_{M_i} \geq -(n-1), \quad \text{vol}(B_1(\tilde{p}_i)) \geq v > 0, \quad \tilde{p}_i \in \tilde{M}_i.$$

Then for i large, M_i is diffeomorphic to an infra-nilmanifold.

A starting point is Theorem 1.4.2, by which we may assume that a bounded normal covering of M_i , $M_{i,0}$, with $\pi_1(M_{i,0})$ a nilpotent group. The main effort in our proof of Theorem 0.3' is to show that $M_{i,0}$ is diffeomorphic to a nilmanifold, which relies on the following fiber bundle criterion for a nilmanifold.

Theorem 2.1. (Nilmanifolds, a sequence of bundles) *A compact manifold M_0 is diffeomorphic to a nilmanifold if and only if M_0 carries a sequence of bundles:*

$$M_1 \rightarrow M_0 \rightarrow B_1, \quad M_2 \rightarrow M_0 \rightarrow B_2, \quad \dots, \quad M_{s+1} \rightarrow M_0 \rightarrow B_{s+1},$$

such that

(2.1.1) *Each M_i -fiber is a union of M_j -fibers for all $j \geq i$, $M_{s+1} = \{\text{pt}\}$.*

(2.1.2) *For each $1 \leq i \leq s$, the natural projection, $B_{i+1} \rightarrow B_i$, is a principal T^{k_i} -bundle.*

We will show that M_0 in Theorem 2.1 admits an iterated principal circle bundles:

$$S^1 \rightarrow M_0 \rightarrow M_1, \quad S^1 \rightarrow M_1 \rightarrow M_2, \quad \dots, \quad S^1 \rightarrow M_n \rightarrow \text{pt},$$

which, according to [Na] and [Be] (Theorem 3.1), implies that M_0 is diffeomorphic to a compact nilmanifold.

Our main work is the following result.

Theorem 2.2. *Let the assumptions be as in Theorem 0.3', and let $M_{i,0}$ be a (bounded) finite normal covering space of M_i such that $\pi_1(M_{i,0})$ is nilpotent. Then for i large $M_{i,0}$ admits a sequence of bundles satisfying Theorem 2.1 i.e. $M_{i,0}$ is diffeomorphic to a nilmanifold.*

A by-product of the proof of Theorem 2.2 is that we are able define, on $M_{i,0}$, a left-invariant metric, such that the normal covering transformations on $M_{i,0}$ by $\Gamma_i/\Gamma_{i,0}$ are almost isometries. Using the left invariant metric, we apply the center of mass technique (Lemma 3.5.1) to modify the normal covering transformations so that the $\Gamma_i/\Gamma_{i,0}$ action preserves the nilpotent structure on $M_{i,0}$ defined by the sub-bundles in Theorem 2.2.

As a preparation, we need some lemmas.

Lemma 2.3. (Euclidean limits and uniform Reifenberg points) *Let $M_i \xrightarrow{\text{GH}} \text{pt}$ with local (ρ, v) -bound Ricci covering geometry. Then up to a scaling of M_i , passing to a subsequence and for any $R > 1$ the following hold:*

$$\begin{array}{ccc} (\tilde{M}_i, \tilde{p}_i, \Gamma_i) & \xrightarrow{\text{eqGH}} & (\mathbb{R}^n, 0, \mathbb{R}^n) \\ \pi_i \downarrow & & \text{proj} \downarrow \\ M_i & \xrightarrow{\text{GH}} & \text{pt}, \end{array} \quad \left\{ \begin{array}{ll} \text{Ric}_{M_i} \geq -(n-1)\delta_i & \delta_i \rightarrow 0 \\ d_{\text{GH}}(B_{i,s}(\tilde{x}_i), B_s^n(0)) < s\epsilon_i & \epsilon_i \rightarrow 0 \\ \tilde{x}_i \in B_R(\tilde{p}_i) & 0 < s \leq R \end{array} \right. .$$

We will refer a maximal collapsing sequence in Lemma 2.3 as satisfying a uniform Reifenberg condition. From now on, we will always assume a sequence in Theorem 0.3' satisfying Lemma 2.3. Consequently, given any sequence, $r_i \rightarrow \infty$, and at any $\tilde{x}_i \in \tilde{M}_i$, the r_i -blow up limit is always \mathbb{R}^n i.e., $(r_i \tilde{M}_i, \tilde{x}_i) \xrightarrow{\text{GH}} (\mathbb{R}^n, 0)$.

We find that if $\pi_1(M_i)$ is nilpotent, then any blow-up limit of M_i must also be a product of a flat torus and an Euclidean space. This discovery leads to the following key geometric ingredients of Theorem 2.2.

Theorem 2.4. (Uniform local elementary T-structures) *Let a sequence, $M_i \xrightarrow{\text{GH}} \text{pt}$, be as in Lemma 2.3. Assume that $\Gamma_i = \pi_1(M_i)$ is nilpotent. Then passing to a subsequence, the following properties hold:*

(2.4.1) *There are integers, (k_1, \dots, k_s) , $m_j = \sum_{l=1}^j k_l$, $m_s = n$, such that for any $x_i \in M_i$, a short generators (ordered in lengths decreasing) for Γ_i at x_i , divides into groups,*

$$\gamma_{i,1}(x_i), \dots, \gamma_{i,m_1}(x_i), \gamma_{i,m_1+1}(x_i), \dots, \gamma_{i,m_2}(x_i), \dots, \gamma_{i,m_{s-1}+1}(x_i), \dots, \gamma_{i,m_s}(x_i),$$

$$\frac{l_j(x_i)}{l_{j'}(x_i)} \leq 1, \quad \frac{l_j(x_i)}{l_{j'}(x_i)} \rightarrow \infty, \quad j' > j, \quad 1 \leq j \leq s, \quad (i \rightarrow \infty),$$

where

$$\begin{cases} \ell_j(x_i) = \max\{|\gamma_{i,k}(x_i)|\}_{k=m_j+1}^{m_{j+1}} \\ l_j(x_i) = \min\{|\gamma_{i,k}(x_i)|\}_{k=m_j+1}^{m_{j+1}}. \end{cases}$$

(2.4.2) *There exists a constant $c = c(\{M_i\}) > 0$ such that the following properties hold: for any $\epsilon > 0$, there is $N(\epsilon) > 0$, such that for $i \geq N(\epsilon)$ and any $x_i \in M_i$,*

$$d_{\text{GH}}(B_{30\sqrt{n}}(x_i, \ell_j(x_i)^{-1}M_{i,0}), B_{30\sqrt{n}}(T^{k_j} \times \mathbb{R}^{m_{j-1}})) < \epsilon,$$

where $T^{k_j} \times \mathbb{R}^{m_j}$ is flat with T^{k_j} satisfying

$$\text{diam}(T^{k_j}) \leq \sqrt{k_j} \leq \sqrt{n}, \quad \text{inrad}(T^{k_j}) \geq c$$

Based on Theorem 2.4, we able to construct an atlas of elementary T -structures on suitable spaces, to which we apply Theorem 1.2.3 to construct, by induction on s , a sequence of sub-bundles on $M_{i,0}$ satisfying Theorem 2.1. Here is an outline of the proof:

Fixing $\epsilon > 0$ and $i > N(\epsilon)$ in Theorem 2.4, we construct a locally bounded open cover, $\{U_{i,\alpha}\}$,

$$(2.5.1) \quad B_{10\sqrt{n}}(x_{i,\alpha}, \ell_s(x_{i,\alpha})^{-1}M_{i,0}) \subset U_{i,\alpha} \subset B_{11\sqrt{n}}(x_{i,\alpha}, \ell_s(x_{i,\alpha})^{-1}M_{i,0}),$$

such that for any $x_i \in M_i$, $\{x_{i,\alpha}\} \cap B_{\ell_s(x_i)30\sqrt{n}}(x_i)$ is almost $\ell_s(x_i)10\sqrt{n}$ -separated and almost $\ell_s(x_i)10\sqrt{n}$ -dense.

(2.5.2) There is a diffeomorphism and ϵ -GHA (Theorem 1.3.6), $f_{i,\alpha} : U_{i,\alpha} \rightarrow T_\alpha^{k_s} \times \mathbb{B}_{10\sqrt{n}}^{m_s-1}(0_\alpha)$, where $U_{i,\alpha}$ (resp. $T^{k_s} \times \mathbb{B}_{10\sqrt{n}}^{m_s-1}(0_\alpha)$) is equipped with the induced metric on $\ell_s(x_{i,\alpha})^{-1}M_{i,0}$ (resp. the product of flat metrics). Moreover, a choice of a base point $e \in T^{k_s}$ and a canonical basis for $\pi_1(T^{k_s}, e)$ uniquely determines (via parallel translation) a Lie group structure on T^{k_s} , thus defines a free T^{k_s} -action on $U_{i,\alpha}$ with orbits $f_{i,\alpha}^{-1}(T_\alpha^{k_s} \times v)$, for any $v \in \mathbb{B}_{10\sqrt{n}}^{m_s-1}(0_\alpha)$.

(2.5.3) If $U_{i,\alpha} \cap U_{i,\beta} \neq \emptyset$, where $B_{\frac{1}{10}}(z_i) \subset U_{i,\alpha} \cap U_{i,\beta}$ for some z_i .

(2.5.4) There is a diffeomorphic ϵ -GHA, $\phi_{i,\alpha,\beta} : U_{i,\alpha,\beta} \rightarrow T^{k_s} \times \mathbb{B}_{30\sqrt{n}}^{m_s-1}(0)$, such that $\psi_{i,\alpha} = \phi_{i,\alpha,\beta} \circ f_{i,\alpha}^{-1} : T_\alpha^{k_s} \times \mathbb{B}_{10\sqrt{n}}^{m_s-1}(0) \rightarrow T^{k_s} \times \mathbb{B}_{30\sqrt{n}}^{m_s-1}(0)$ is an diffeomorphic embedding and 2ϵ -GHA.

It is obvious that the two T^{k_s} -actions on $U_{i,\alpha} \cap U_{i,\beta} \neq \emptyset$ are C^0 -close (with respects to either the original or the pull-back flat metric). Because the free T^{k_s} -action on $U_{i,\alpha}$ are isometric with respect to the pull-back metric from $T^{k_s} \times \mathbb{R}^{m_s-1}$, the C^1 -close condition in Theorem 1.2.3 is satisfied (Remark 1.2.4). Hence, we are able to glue the atlas of elementary T^{k_s} -structures, $\{(U_{i,\alpha}, T^{k_s})\}$, to form an affine T^{k_s} -bundle on $M_{i,0}$, $T^{k_s} \rightarrow M_{i,0} \xrightarrow{h_{s,i}} B_{i,s}$. Because $\Gamma_{i,0}$ is torsion-free nilpotent and $\Gamma_{i,s}$ is normal abelian subgroup, $\Gamma_{i,s}$ is contained in the center of $\Gamma_{i,0}$. Hence, the holonomy group is trivial i.e., the T^{k_s} -bundle is principal.

Next, we shall construct a principal T^{k_s-1} -bundle on $B_{i,s} = M_{i,0}/T^{k_s}$, where the free T^{k_s} -action on $M_{i,0}$ are almost isometries with respect to $\ell_s(x_{i,\alpha})^{-1}g_i$ around $x_{i,\alpha}$ (hence with respect to any $\ell_j(x_{i,\alpha})^{-1}g_i$, $j = 1, \dots, s-1$).

Fixing a probability measure on T^{k_s} , let $g_{i,s}$ denote the T^{k_s} -invariant metric obtained by averaging g_i under the T^{k_s} -action. Then for any fixed $R > 1$,

$$d_{\text{GH}}(B_R(x_i, \ell_s(x_i)^{-1}g_{i,s}), B_R(x_i, \ell_s(x_i)^{-1}g_i)) \rightarrow 0, \quad i \rightarrow \infty,$$

because $\frac{\ell_s(x_i)}{\ell_{s-1}(x_i)} \rightarrow 0$, all T^{k_s} -orbits on the above R -balls (with respect to both metrics) uniformly collapse to points. Hence, the above implies the following,

$$d_{\text{GH}}(B_R(\bar{x}_i, \ell_{s-1}(x_i)^{-1}\bar{g}_{i,s}), B_R(T^{k_{s-1}} \times \mathbb{R}^{m_{s-2}})) \rightarrow 0, \quad i \rightarrow \infty.$$

Consider the blow-up at the second fastest rate at any $x_{i,\beta}$, $\ell_{s-1}(x_{i,\beta})^{-1}M_{i,0}$, again by Theorem 2.4 we can construct a locally finite open cover for $M_{i,0}$, $\{V_{i,\beta}\}$, where similar to (2.5.1) $V_{i,\beta}$ is T^{k_s} -invariant open subset of $M_{i,0}$,

$$B_{10\sqrt{n}}(x_{i,\beta}, \ell_{s-1}(x_{i,\beta})^{-1}M_{i,0}) \subset V_{i,\beta} \subset B_{11\sqrt{n}}(x_{i,\beta}, \ell_{s-1}(x_{i,\beta})^{-1}M_{i,0}).$$

Lemma 2.6. *Let $B_{i,s} = M_{i,0}/T^{k_s}$ be equipped with $\bar{g}_{i,s}$. Then $B_{i,s}$ admits a locally bounded open cover, $\{\bar{V}_{i,\beta}\}$, and a diffeomorphic ϵ_i -GHA ($\epsilon_i \rightarrow 0$ as $i \rightarrow \infty$),*

$$f_{i,\beta} : \ell_{s-1}(x_{i,\beta})^{-1}\bar{V}_{i,\beta} \rightarrow T^{k_{s-1}} \times B_{10\sqrt{n}}(\mathbb{R}^{m_{s-2}}),$$

and $T^{k_{s-1}} \times B_{10\sqrt{n}}(\mathbb{R}^{m_{s-2}}) \subset T^{k_{s-1}} \times \mathbb{R}^{m_{s-2}}$ is a metric product of flat manifolds.

Consequently, we obtain a locally bounded open cover for $B_{i,s}$, $\{\bar{V}_{i,\beta}\}$, satisfying properties as in (2.5.2)-(2.5.4). By now we are able to apply Theorem 1.2.3, gluing the local elementary T -structures, $\{(\bar{V}_{i,\beta}, T^{k_{s-1}})\}$, on $B_{i,s}$, and obtain a principal T^{k_s} -bundle, $T^{k_s} \rightarrow B_{i,s} \xrightarrow{h_{i,s-1}} B_{i,s-1}$, and a fiber bundle, $M_{i,s} \rightarrow M_{i,0} \xrightarrow{h_{i,s-1} \circ h_{s,i}} B_{i,s-1}$, and each $M_{i,s}$ is a union of T^{k_s} -orbits.

Observe that if $s = 2$, then the above prove Theorem 2.2. For $s > 2$, we will proceed with induction on s to finish the proof of Theorem 2.2.

Remark 2.7. (2.7.1) Theorem 2.4 (both (2.4.1) and (2.4.2)) will be false, if one removes the assumption that $\pi_1(M_i)$ is nilpotent (Example 1.5.3).

(2.7.2) When restricting to the case with $|\text{sec}_{M_i}| \leq 1$, by Theorem 0.2 one sees a uniform bound on $\frac{\ell_j(x_i)}{\ell_j(y_i)}$ for any $x_i, y_i \in M_i$. Note that our proof of Theorem 2.2 does not required such a (uniform) bound, because our proof of Theorem 2.2 relies on local geometry (comparing the discussion at the end of Section 1.4).

(2.7.3) Let M_i be as in Theorem 0.3'. Combining Theorem 1.4.2 and (2.4.1), one sees that for any $x_i \in M_i$, in Theorem 0.3', the number of short generators at x_i is uniformly bounded above by a constant depending on n (comparing to Theorem 1.4.1 where the bound only holds at points which may not be specified).

3. PROOF OF THEOREM A

In this section, we will prove theorems/lemmas stated in Section 2.

3.1. Proof of Theorem 2.1.

Our proof of Theorem 2.1 uses the following criterion of a nilmanifold.

Theorem 3.1. (Nilmanifolds: iterated principal circle bundles, [Na], [Be]) *A compact manifold M is diffeomorphic to nilmanifold N/Γ if and only if M admits an iterated principal circle bundles:*

$$S^1 \rightarrow M \rightarrow M_1, \quad S^1 \rightarrow M_1 \rightarrow M_2, \quad \dots \quad S^1 \rightarrow M_n \rightarrow \text{pt.}$$

Proof of Theorem 2.1.

Given a nilmanifold, N/Γ , there is the descending sequence,

$$N \triangleright N_1 \triangleright \cdots \triangleright N_s \triangleright N_{s+1} = e, \quad \Gamma > \Gamma_1 > \cdots > \Gamma_s > \Gamma_{s+1} = e,$$

where $N_1 = [N, N]$, $N_i = [N, N_{i-1}]$, $i \geq 2$, and $\Gamma_i = N_i \cap \Gamma$, and $N_s \subseteq Z(N)$, the center of N . Then one obtains the desired iterated normal bundle over tori with trivial holonomy representation,

$$N_1/\Gamma_1 \rightarrow N/\Gamma \rightarrow T^{k_1} = (N/N_1)/(\Gamma/\Gamma_1), \quad \cdots, \quad e \rightarrow N_s/\Gamma_s \rightarrow T^{k_s}.$$

Let $M = N/\Gamma$, $M_i = N_i/\Gamma_i$, $i \geq 1$. Then the correspond sequence of sub-bundles are:

$$M_1 \rightarrow M \rightarrow B_1 = T^{k_1}, \quad M_2 \rightarrow M \rightarrow B_2, \quad \cdots, \quad \text{pt} \rightarrow M \rightarrow B_{s+1} = M,$$

where $B_j = (N/N_j)/(\Gamma/\Gamma_j)$, where $T^{k_j} \rightarrow B_j \rightarrow B_{j-1}$ is a principal bundle.

We now assume that M satisfies the conditions of Theorem 2.1. By Theorem 3.1, it suffices to show that M admits an iterated principal circle bundles. We proceed by induction on s , starting with $s = 1$, $T^{k_2} \rightarrow M \rightarrow T^{k_1}$. Because $T^{k_2} \rightarrow M \rightarrow T^{k_1}$ is a principal T^{k_2} -bundle, M can be expressed as iterated principal circle bundles.

Consider an iterated bundles over tori of length s on M in Theorem 3.1. Note that $M_s \rightarrow M \rightarrow B_s$ is a principal T^{k_s} -bundle. Consider

$$T^{k_s} \rightarrow M \rightarrow M/T^{k_s}.$$

It is easy to see that M/T^{k_s} again satisfies the conditions of Theorem 2.1 of length $s - 1$, and by induction M/T^{k_s} admits iterated circle bundles (Theorem 3.1). By now we can conclude that M admits iterated principal circle bundles. \square

3.2. Proof of Lemma 2.3 and Theorem 2.4.

Proof of Lemma 2.3.

Passing to a subsequence, we may assume the following commutative diagram (Lemma 1.1.3):

$$\begin{array}{ccc} (\tilde{M}_i, \tilde{p}_i, \Gamma_i) & \xrightarrow{\text{eqGH}} & (\tilde{X}, \tilde{p}, G) \\ \pi_i \downarrow & & \text{proj} \downarrow \\ (M_i, p_i) & \xrightarrow{\text{GH}} & \text{pt} = \tilde{X}/G. \end{array}$$

By (1.3.5.1), \tilde{X} contains a point where the tangent cone is unique and is isometric to \mathbb{R}^n . Because G acts transitively on \tilde{X} by isometries, every point has a unique tangent cone isometric to \mathbb{R}^n . Assume that $r_j \rightarrow \infty$ such that $(r_j \tilde{X}, \tilde{p}) \xrightarrow{\text{GH}} C_{\tilde{p}} \tilde{X} = \mathbb{R}^n$. Then for any $\tilde{x} \in \tilde{X}$, $(r_j \tilde{X}, \tilde{x}) \xrightarrow{\text{GH}} \mathbb{R}^n$. For each r_j , we may assume i_j large such that for all $i \geq i_j$, $d_{\text{GH}}((r_j \tilde{M}_i, \tilde{p}_i, \Gamma_i), (r_j \tilde{X}_i, \tilde{p})) < j^{-1}$, and $d_{\text{GH}}(r_j M_i, \text{pt}) < j^{-1}$. By Lemmas 1.1.2 and 1.1.3, a standard diagonal argument one gets the following commutative diagram,

$$\begin{array}{ccc} (r_j \tilde{M}_{i_j}, \tilde{p}_{i_j}, \Gamma_{i_j}) & \xrightarrow{\text{eqGH}} & (\mathbb{R}^n, \tilde{p}, \mathbb{R}^n) \\ \pi_i \downarrow & & \text{proj} \downarrow \\ (r_j M_{i_j}, p_{i_j}) & \xrightarrow{\text{GH}} & \text{pt} = \mathbb{R}^n/\mathbb{R}^n. \end{array}$$

Because $(r_j \tilde{M}_{i_j}, \tilde{p}_{i_j}) \xrightarrow{\text{GH}} (\mathbb{R}^n, 0)$ with $\text{Ric}_{r_j \tilde{M}_{i_j}} \geq -(n-1)r_j^{-2} \rightarrow 0$, for any $\epsilon > 0$, by Theorems 1.3.3 and 1.3.4, for i large we may assume that \tilde{p}_{i_j} is (R, ϵ) -Reifenberg i.e.,

$$d_{\text{GH}}(B_{i,s}(\tilde{p}_{i_j}), \mathbb{B}_s^n(0)) < \epsilon \cdot s, \quad 0 < s \leq R,$$

where $\mathbb{B}_s^n(0)$ denotes a s -ball in \mathbb{R}^n . Because $\pi_i(B_1(\tilde{p}_{i_j})) = M_{i_j} \xrightarrow{\text{GH}}$ pt, the above holds when replacing \tilde{p}_{i_j} by any point $\tilde{x}_{i_j} \in B_1(\tilde{p}_{i_j})$, hence all points on \tilde{M}_{i_j} are uniform (R, ϵ) -Reifenberg. Now let $\epsilon \rightarrow 0$, from the above one easily sees the desired result. \square

Our proof of Theorem 2.4 is divided into several lemmas.

We will first show that if $\pi_1(M_i)$ is nilpotent, then a blow up limit space is always a metric product of a flat torus and an Euclidean space.

Lemma 3.2.1. *Let $M_i \xrightarrow{\text{GH}}$ pt be as in Lemma 2.3. Assume that $\pi_1(M_i)$ is nilpotent. Then passing to a subsequence the following hold:*

(3.2.1.1)

$$\begin{array}{ccc} (\ell_1(M_i)^{-1} \tilde{M}_i, \tilde{p}_i, \Gamma_i) & \xrightarrow{\text{eqGH}} & (\mathbb{R}^n, 0, \mathbb{Z}^{k_1} \times \mathbb{R}^{n-k_1}) \\ \pi_i \downarrow & & \text{proj} \downarrow \\ (\ell_1(M_i)^{-1} M_i, p_i) & \xrightarrow{\text{GH}} & (T^{k_1}, e), \end{array} \quad \ell_1(M_i) = \text{diam}(M_i),$$

where the limit group, $\mathbb{Z}^{k_1} \times \mathbb{R}^{n-k_1}$, acts on \mathbb{R}^n by translations.

(3.2.1.2) Γ_i has a normal subgroup $\Gamma_{i,1}$, $\Gamma_{i,1} \rightarrow \mathbb{R}^{n-k_1}$, such that $\Gamma_i/\Gamma_{i,1} \cong \mathbb{Z}^{k_1}$.

(3.2.1.3) For $x_i \in M_i$, let $\{\gamma_{i,1,j}\}$ be a finite set of short generators for $\Gamma_{i,1}$ at x_i . Then

$$\ell_2(x_i) = \max_j \{d_{\ell_1(M_i)^{-1} \tilde{M}_i}(\tilde{x}_i, \gamma_{i,1,j}(\tilde{x}_i))\} \rightarrow 0, \quad i \rightarrow \infty.$$

Proof. (3.2.1.1) By Lemma 1.14 and Lemma 3.2.1, passing to a subsequence we may start with the following equivariant GH-convergence,

$$\begin{array}{ccc} (\ell_1(M_i)^{-1} \tilde{M}_i, \tilde{p}_i, \Gamma_{i,0}) & \xrightarrow{\text{eqGH}} & (\mathbb{R}^n, 0, G_1) \\ \pi_i \downarrow & & \text{proj} \downarrow \\ (\ell_1(M_i)^{-1} M_i, p_i) & \xrightarrow{\text{GH}} & (X_1, p). \end{array}$$

Because $\Gamma_{i,0}$ is nilpotent, the limit group G_1 is nilpotent. We claim that G_1 acts (freely) on \mathbb{R}^n by translations, thus $G_1 = \mathbb{Z}^{k_1} \times \mathbb{R}^{n-k_1}$ and $X_1 = \mathbb{R}^n/G_1 = T^{k_1}$.

We first show that G_1 acts freely. If the identity isotropy group at $0 \in \mathbb{R}^n$, $H_0 \neq e$. Because H_0 is a compact subgroup of a nilpotent group G_1 , $H_0 \subset Z(G_1)$, the center of G_1 , and G_1/H_0 acts isometrically on $\mathbb{R}^n/H_0 = \mathbb{R}^l \times C(S_1^{n-l-1}/H_0)$, where $C(S_1^{n-l-1}/H_0)$ is an Euclidean cone over S_1^{n-l-1}/H_0 . Consequently, $\mathbb{R}^n/G_1 = \mathbb{R}^l \times C(S_1^{n-l-1}/H_0)/(G_1/H_0)$ is not compact, a contradiction. Similarly, if $\mathbb{R}^1 < Z(G_1)$ acts not by translation, then $\mathbb{R}^n/\mathbb{R}^1 = \mathbb{R}^l \times Y$, and $\mathbb{R}^n/G_1 = \mathbb{R}^l \times Y/(G_1/\mathbb{R}^1)$ is not compact, a contradiction, where Y is non-compact manifold of $\text{sec}_Y \geq 0$ whose unique soul is the pole point. We then repeating the above to $\mathbb{R}^n/Z((G_1)_0) = \mathbb{R}^k$, on which $G_1/Z((G_1)_0)$ acts isometrically.

Because $G_1/Z((G_1)_0)$ is nilpotent, we repeat the above argument, and eventually get the desired result.

(3.2.1.2) Follows from (3.2.1.1) and Lemma 1.14.

(3.2.1.3) Follows from (3.2.1.2) and Lemma 1.14. \square

Lemma 3.2.2. *Let the assumptions be as in Lemma 3.2.1. Passing to a subsequence, we have*

$$(3.2.2.1) \quad \begin{array}{ccc} (\ell_2(x_i)^{-1} \tilde{M}_i, \tilde{x}_i, \Gamma_i) & \xrightarrow{\text{eqGH}} & (\mathbb{R}^n, 0, \mathbb{Z}^{k_2} \times \mathbb{R}^{n-k_1}) \\ \pi_i \downarrow & & \text{proj} \downarrow \\ (\ell_2(x_i)^{-1} M_i, x_i) & \xrightarrow{\text{GH}} & (T^{k_2} \times \mathbb{R}^{k_1}, e \times 0), \end{array}$$

where the limit group, $\mathbb{Z}^{k_2} \times \mathbb{R}^{n-k_1}$, acts on \mathbb{R}^n by translations.

(3.2.2.2) $\Gamma_{i,1}$ has a subgroup $\Gamma_{i,2}$ normal in Γ_i such that $\Gamma_{i,1}/\Gamma_{i,2} \cong \mathbb{Z}^{k_2}$.

(3.2.2.3) For $x_i \in M_i$, let $\{\gamma_{i,2,j}\}$ be a finite set of short generators for $\Gamma_{i,2}$ at x_i , and let $\ell_2(x_i) = \max\{d(\tilde{x}_i, \gamma_{i,2,j}(\tilde{x}_i))\}$. Then $\ell_2(x_i) \rightarrow 0$ as $i \rightarrow \infty$.

Proof. (3.2.2.1) Similarly, passing to a subsequence we may start with the following equivariant GH-convergence,

$$\begin{array}{ccc} (\ell_1(x_i)^{-1} \tilde{M}_i, \tilde{x}_i, \Gamma_{i,1}) & \xrightarrow{\text{eqGH}} & (\mathbb{R}^n, 0, G_2) \\ \pi_i \downarrow & & \text{proj} \downarrow \\ (\ell_1(x_i)^{-1} M_i, x_i) & \xrightarrow{\text{GH}} & (X_2 = \mathbb{R}^n/G_2, x). \end{array}$$

Because for any $\gamma_i \in \Gamma_i - \Gamma_{i,1}$, $d(\tilde{x}_i, \gamma_i(\tilde{x}_i)) \geq \text{inrad}(T^{k_1}) > 0$, $\Gamma_{i,1} \xrightarrow{\text{GH}} G_2$, which acts on the \mathbb{R}^{n-k_1} -factor of $\mathbb{R}^n = \mathbb{R}^{n-k_1} \times \mathbb{R}^{k_1}$. Because $\Gamma_{i,1}$ is nilpotent, G_2 is nilpotent. We claim that G_2 acts freely by translations on \mathbb{R}^{n-k_1} , thus $G_2 = \mathbb{Z}^{k_2} \times \mathbb{R}^{n-k_1-k_2}$ and $X_2 = \mathbb{R}^n/G_2 = T^{k_2} \times \mathbb{R}^{k_1}$.

Without loss of the generality, we may assume the isotropy group $H_0 \neq e$, $0 \in \mathbb{R}^n$. Because H_0 is compact, $H_0 \subset Z(G_2)$, the center of G_1 , and G_2/H_0 acts isometrically on $\mathbb{R}^n/H_0 = \mathbb{R}^l \times C(S_1^{n-l-1}/H_0)$, where $C(S_1^{n-l-1}/H_0)$ is an Euclidean cone over S_1^{n-l-1}/H_0 . Consequently, $\mathbb{R}^n/G_1 = \mathbb{R}^l \times C(S_1^{n-l-1}/H_0)/(G_1/H_0)$ is not compact, a contradiction. Similarly, if $\mathbb{R}^1 < Z(G_2)$ acts not by translation, then $\mathbb{R}^n/\mathbb{R}^1 = \mathbb{R}^l \times Y$, and $\mathbb{R}^n/G' = \mathbb{R}^l \times Y/(G_2/\mathbb{R}^1)$ is not compact, a contradiction, where Y is non-compact manifold of $\text{sec}_Y \geq 0$ whose unique soul is the pole point. We then repeating the above to $\mathbb{R}^n/Z((G_2)_0) = \mathbb{R}^k$, on which $G_2/Z((G_2)_0)$ acts isometrically.

Because $G_2/Z((G_2)_0)$ is nilpotent, we repeat the above argument, and eventually get the desired result.

By now, similar to the proof in Lemma 3.2.1, one gets (3.2.2.2) and (3.2.2.3). \square

Note that in Lemma 3.2.1, the flat torus T^{k_1} is independent of p_i . Consequently, for $\epsilon > 0$ small, there is $N(\epsilon) > 0$ such that at every point in M_i , the group of short generators for Γ_i , consisting of ones whose lengths are proportional to $\text{diam}(M_i)$, can be chosen having almost identical lengths; which are close to seen from parallel short basis for $\pi_1(T^{k_1})$ on T^{k_1} .

Note that in Lemma 3.2.2, a priori T^{k_2} may depend on x_i . We will show that k_2 is independent of x_i and every T^{k_2} has a bounded geometry.

Lemma 3.2.3. *Theorem 2.4 is true for $j = 1$ i.e., k_2 is independent of x_i , $\text{diam}(T^{k_2}) \leq \sqrt{n}$, and $\text{inrad}(T^{k_2}) \geq c > 0$, where $c = c(\{M_i\})$.*

Proof. Arguing by contradiction, given $c_j = 2^{-j} \rightarrow 0$, we will pick up a contradiction subsequence of M_i as follows:

(i) Fixing c_1 , assuming $\epsilon_1 > 0$ and a subsequence sequence (M_{i_1}, x_{i_1}) such that $B_{30\sqrt{n}}(x_{i_1}, \ell_1(x_{i_1})^{-1}M_{i_1})$ is at least ϵ_1 -away from any $B_{30\sqrt{n}}(e \times 0, T^k \times \mathbb{R}^{k_1})$, $\text{inrad}(T^k) \geq c_1$. Because T^k has a short basis of lengths ≤ 1 , $\text{diam}(T^k) \leq \sqrt{k} < \sqrt{n}$.

(ii) Fixing c_2 , assuming that there is $\epsilon_2 > 0$ and a subsequence of $\{(M_{i_1}, x_{i_1})\}$, (M_{i_2}, x_{i_2}) such that $B_{30\sqrt{n}}(x_{i_2}, \ell_1(x_{i_2})^{-1}M_{i_2})$ is at least ϵ_2 -away from any $B_{30\sqrt{n}}(e \times 0, T^k \times \mathbb{R}^{k_1})$, $\text{inrad}(T^k) \geq c_2$.

Iterating this process, fixing c_j , assuming that there is $\epsilon_j > 0$ and a subsequence of $\{(M_{i_{j-1}}, x_{i_{j-1}})\}$, (M_{i_j}, x_{i_j}) , such that $B_{30\sqrt{n}}(x_{i_j}, \ell_1(x_{i_j})^{-1}M_{i_j})$ is at least ϵ_j -away from any $B_{30\sqrt{n}}(e \times 0, T^k \times \mathbb{R}^{k_1})$, $\text{inrad}(T^k) \geq c_j$. For the sake of simple notion, we will use (M_{i_i}, x_{i_i}) to denote the diagonal subsequence from the above construction, which servers as a contradiction sequence.

Passing to a subsequence, by the proof of Lemma 3.2.2 we obtain

$$(\ell_1(x_{i_i})^{-1}M_{i_i}, x_{i_i}) \xrightarrow{\text{GH}} (T^k \times \mathbb{R}^{k_1}, e \times 0).$$

Fixing j_0 large so that $\text{inrad}(T^k) \geq c_{j_0} > 0$, for all i_i large such that

$$d_{\text{GH}}(B_{30\sqrt{n}}(x_{i_i}, \ell_1(x_{i_i})^{-1}M_{i_i}), B_{30\sqrt{n}}(T^k \times \mathbb{R}^{k_1}, e \times 0)) < \epsilon_{j_0},$$

we get a contradiction.

Finally we show that $k_2 := k$ in the above can be chosen independent of $x_i \in M_i$. Fixing $x_i \in M_i$, by the bounded geometry on T^k , it is clear that k_2 can be chosen for all points in a neighborhood of x_i ; hence the set of points with k_2 is open. Similarly, the set of points with k_2 is also close, hence we have proved that k_2 is independent of points in M_i . \square

Proof of Theorem 2.4.

Without loss of generality, we may assume that $M_i \xrightarrow{\text{GH}}$ pt satisfies Lemma 2.3. We will perform successive blow-ups on the sequence with a blow-up rate determined as in Lemmas 3.2.1-3.2.3, and we will proceed the proof by induction on the total number of the blow-ups, which is at most n .

By Lemma 3.2.2-3.2.3, we may assume that (2.4.1) is true for $j = 1, \dots, t-1$. If (2.4.2) is false on the t -th blow-up sequence, following the proof of Lemma 3.2.3 we get a contradiction. Moreover, $k = k_t$ is independent of $x_i \in M_i$. \square

3.3. Proof of Lemma 2.6.

Proof of Lemma 2.6.

We claim that given $\epsilon > 0$, there exists $N(\epsilon) > 0$ such that for $i \geq N(\epsilon)$, all $\bar{y}_i \in \bar{V}_{i,\alpha}(\bar{x}_i, \ell_{s-1}(x_i)^{-1}\bar{g}_{i,s})$ are uniform $(\epsilon, 1)$ -Reifenberg points, i.e.,

$$d_{\text{GH}}(B_r(\bar{y}_i, \ell_{s-1}(x_i, \alpha)^{-1}\bar{g}_{i,s}), \mathbb{B}_r^{n-k_s}(0)) < \epsilon r, \quad 0 < r \leq 1.$$

Assuming the claim, by the standard Reifenberg method in geometry measure theory ([CC1]) we conclude that $B_{10\sqrt{n}}(T^{k_s-1} \times \mathbb{R}^{m_s-2})$ is diffeomorphic to $\tilde{V}_{i,\alpha}$, thus by a similar construction of $\{U_{i,\alpha}\}$ for $M_{i,0}$, one gets a desired locally bounded open cover for $B_{i,s}$.

Arguing by contradiction, assuming that for some $\epsilon_0 > 0$, there is a contradicting sequence,

$$d_{\text{GH}}(B_{r_i}(\bar{y}_i, \ell_{s-1}(x_{i,\alpha})^{-1}\bar{g}_{i,s}), \mathbb{B}_{r_i}^{n-k_s}(0)) \geq \epsilon_0 r_i, \quad 0 < r_i \leq 1,$$

thus

$$(2.6.1) \quad d_{\text{GH}}(B_1(\bar{y}_i, r_i^{-1}\ell_{s-1}(x_{i,\alpha})^{-1}\bar{g}_{i,s}), \mathbb{B}_1^{n-k_s}(0)) \geq \epsilon_0.$$

Let $\text{proj}_{i,\alpha} : \tilde{V}_{i,\alpha} \rightarrow V_{i,\alpha} \rightarrow \bar{V}_{i,\alpha}$ denote the $\mathbb{R}_1^{k_s}$ -fiber projection map, where $\tilde{V}_{i,\alpha} \subset \widetilde{B_\rho(x_{i,\alpha})}$, and $\mathbb{R}_1^{k_s}$ denotes the universal covering group of $T_1^{k_s}$ which acts freely and isometrically on $\tilde{V}_{i,\alpha}$.

Fixing $R > 10$, consider the following diagram (passing to a subsequence if necessary):

$$(2.6.2) \quad \begin{array}{ccc} (\text{proj}_{i,\alpha}^{-1}(B_{i,1}) \cap B_R(\tilde{y}_i), \tilde{g}_{i,s}, \mathbb{R}_1^{k_s}) & \xrightarrow{\text{GH}} & (\mathbb{B}_1^{n-k_s}(0) \times \mathbb{B}_R^{k_s}(0), \mathbb{R}_2^{k_s}) \\ \text{proj}_{i,\alpha} \downarrow & & \text{proj} \downarrow \\ B_{i,1} = B_1(\bar{y}_i, r_i^{-1}\ell_{s-1}(x_{i,\alpha})^{-1}\bar{g}_{i,s}) & \xrightarrow{?} & \mathbb{B}_1^{n-k_s}(0), \end{array}$$

where the GH-convergence is originally with respect to \tilde{g}_i , which implies with respect to $\tilde{g}_{i,s}$, because the $R_1^{k_s+1}$ -action preserves the Euclidean metric which is ϵ_i -GH close to \tilde{g}_i with respect to any scale (Lemma 2.3).

We claim that there is a canonical identification between groups, $\mathbb{R}_1^{k_s+1}$ and $\mathbb{R}_2^{k_s+1}$, by which the ϵ_i -GH-convergence becomes an ϵ_i -equivariant $\Psi(\epsilon_i)$ -GHA with respect to the (pseudo) $\mathbb{R}_i^{k_s+1}$ -actions. Consequently, the eqGH-convergence in (2.6.2) descends to a GH-convergence on \mathbb{R}^{k_s} -orbit spaces, a contradiction to (2.6.1).

Recall that the $T_1^{k_s}$ -action on $V_{i,\alpha}$ is defined based on Theorem 2.4; from its proof one sees a canonical identification of the group of elements in a short basis at $x_{i,\alpha}$ with shortest lengths are identified with a canonical basis of the T^{k_s} -factor on the flat product, $\mathbb{B}_{10\sqrt{n}}^{n-k_s}(0) \times T^{k_s}$. By the Malcev's rigidity, this determines an identification between groups, $\mathbb{R}_1^{k_s}$ and $\mathbb{R}_2^{k_s}$.

Because the $\mathbb{R}_i^{k_s}$ -actions preserve metrics, by which the $\mathbb{R}_i^{k_s}$ -actions are completely determined. Hence a diffeomorphism and ϵ_i -GHA is also $\Psi(\epsilon_i)$ -equivariant, thus the claim has been verified. \square

3.4. $M_{i,0}$ is diffeomorphic to a nilmanifold.

Proof of Theorem 2.2.

Fixing small $\epsilon > 0$, without loss of the generality we may assume that $M_i \xrightarrow{\text{GH}} \text{pt}$ is chosen so that Theorem 2.4 holds.

We begin by summarizing the discussion following Theorem 2.4: with respect to the scaling functions, $\ell_s(\cdot)^{-1}$, $i \geq N(\epsilon)$, by Theorem 2.4 we are able to construct, on

$M_{i,0}$, a locally bounded open cover with an elementary T^{k_s} -structure, $\{(U_{i,\alpha}, T_\alpha^{k_s})\}$, where T^{k_s} acts freely on $U_{i,\alpha}$ and properties (2.5.1)–(2.5.4) hold. Equipped $U_{i,\alpha}$ with the pullback flat metrics, we are able to apply Theorem 1.2.3 to glue together the elementary T^{k_s} -structures, $\{(U_{i,\alpha}, T_\alpha^{k_s})\}$, to form a T^{k_s} -bundle, $T^{k_s} \rightarrow M_i \rightarrow B_{i,s}$, with affine structural group. Because $\Gamma_{i,0}$ is torsion-free nilpotent and the fiber fundamental group is a normal abelian subgroup, which has to be contained in the center of $\Gamma_{i,0}$, hence the structural group is trivial i.e., this is a principal T^{k_s} -bundle, $T^{k_s} \rightarrow M_i \xrightarrow{h_{i,s}} B_{i,s}$.

Next, with the scale function, $\ell_{s-1}(\cdot)^{-1}$, by (2.4.2) $B_{30\sqrt{n}}(x_{i,\alpha}, \ell_{s-1}(x_{i,\alpha})^{-1}M_i)$ is close to $B_{30\sqrt{n}}(T^{k_{s-1}} \times \mathbb{R}^{m_{s-2}})$ (note that a large radius, $30\sqrt{n}$, is for a gluing purpose later). By (2.4.1), any T^{k_s} -fiber in $B_{30\sqrt{n}}(x_{i,\alpha}, \ell_{s-1}(x_{i,\alpha})^{-1}M_i)$ is almost collapses to a point, as in (2.5.1) we may assume a T^{k_s} -invariant open subset $V_{i,\alpha}$ ϵ_i -close to $B_{10\sqrt{n}}(T^{k_{s-1}} \times \mathbb{R}^{n-k_s})$. By Lemma 2.6, similar to the construction of $\{(U_{i,\alpha}, T_\alpha^{k_s})\}$ for $M_{i,0}$, we construct a locally bounded open cover for $B_{i,s}$ with elementary $T^{k_{s-1}}$ -structures, $\{(\bar{V}_{i,\beta}, T_\beta^{k_{s-1}})\}$, which satisfies properties (2.5.1)–(2.5.4). Again by Theorem 1.2.3, we are able to glue together the elementary $T^{k_{s-1}}$ -structures to form a $T^{k_{s-1}}$ -bundle, $T^{k_{s-1}} \rightarrow B_{i,s} \xrightarrow{h_{i,s-1}} B_{i,s-1}$, which is principal because $\pi_1(B_{i,s})$ is torsion-free nilpotent. Clearly, the fiber bundle, $M_{i,s-1} \rightarrow M_{i,0} \xrightarrow{h_{i,s-1} \circ h_{i,s}} B_{i,s-1}$, has the principal T^{k_s} -bundle, $T^{k_s} \rightarrow M_{i,0} \xrightarrow{h_{i,s}} B_{i,s}$, as a sub-bundle, and each $M_{i,s-1}$ is a union of T^{k_s} -fibers.

We now process by induction on s (in the reverse order), assuming at up to local scales, $\ell_s(\cdot)^{-1}, \dots, \ell_j(\cdot)^{-1}$, we construct a sequence of bundles on $M_{i,0}$, and principal tori bundles on $B_{i,s}, B_{i,s-1}, \dots, B_{i,j}$, such that

$$\left\{ \begin{array}{ll} T^{k_s} \rightarrow M_i \xrightarrow{h_{i,s}} B_{i,s}, & \\ M_{i,s} \rightarrow M_i \xrightarrow{h_{i,s-1} \circ h_{i,s}} B_{i,s-1}, & T_i^{k_{s-1}} \rightarrow B_{i,s} \xrightarrow{h_{i,s-1}} B_{i,s-1}, \\ \dots, & \dots, \\ M_{i,j} \rightarrow M_i \xrightarrow{h_{i,j-1} \circ \dots \circ h_{i,s-1} \circ h_{i,s}} B_{i,j-1}, & T_i^{k_{j-1}} \rightarrow B_{i,j} \xrightarrow{h_{i,j-1}} B_{i,j-1}, \end{array} \right.$$

and each $M_{i,j-1}$ is a union of $M_{i,j}$ -fibers.

By Theorem 2.4 with the local scale number, $\ell_{i,j-1}(\cdot)^{-1}$, we observe the following properties:

$$d_{\text{GH}}(B_{10\sqrt{n}}(x_i, (\ell_{i,j-1}(x_i))^{-1}M_{i,0}), B_{10\sqrt{n}}(e \times 0, T_{x_i}^{k_{j-1}} \times \mathbb{R}^{m_{j-2}})) < \epsilon.$$

Now we are in a situation similar to the scale $\ell_s(\cdot)^{-1}$, where the fiber of the bundle, $M_{i,j} \rightarrow M_{i,0} \rightarrow B_{i,j}$, is collapsed to a point with respect to $\ell_{j-1}(\cdot)^{-1}$, thus (following the proof of Lemma 2.6) we may identify $B_{10\sqrt{n}}(T^{k_{j-1}} \times \mathbb{R}^{m_{j-2}})$ with an open subset of $B_{i,j}$, $h_{j,s}(W_{i,\gamma}) = \bar{W}_{i,\gamma}$, equipped with a flat metric. Consequently, we can construct a locally bounded open cover for $B_{i,j-1}$ with elementary $T^{k_{j-1}}$ -structures, $\{(\bar{W}_{i,\gamma}, T_\gamma^{k_{j-1}})\}$, which satisfies properties as in (2.5.1)–(2.5.4). Then by applying Theorem 1.2.3 we construct the desired bundle map, $T^{k_{j-1}} \rightarrow B_{i,j-1} \xrightarrow{h_{i,j-2}} B_{i,j-2}$, which is a principal $T^{k_{j-1}}$ -bundle because $\pi_1(B_j)$ is torsion-free nilpotent. Moreover, we extend the sequence of bundles, $M_{i,j-1} \rightarrow M_{i,0} \xrightarrow{h_{i,j-2} \circ \dots \circ h_{i,s}} B_{i,j-2}$. By the induction, the proof is complete. \square

3.5. Completion of Proof of Theorem 0.3’.

Let $M_i \xrightarrow{\text{GH}} \text{pt}$ be as in Theorem 0.3’. By Theorem 2.2, for i large, a bounded normal covering space, $M_{i,0}$, is diffeomorphic to a nilmanifold. To prove that M_i is diffeomorphic to an infra-nilmanifold, we shall use the stability of isometric Lie group actions in [Hu].

Consider a compact Lie group G acting effectively on a manifold M , if two G -actions on M are C^1 -close ([Pa]), then the two G -actions are conjugate via a diffeomorphism that is isotopy to id_M . According to [GK], if two G -actions are isometric with respect to two metrics, g_0, g_1 on M , with $|\text{sec}_{g_i}| \leq 1$ and the convexity radius of $g_0 \geq i_0 > 0$, then a C^0 -close of the two G -actions implies the C^1 -closeness. Moreover, one obtains a conjugation close to id_M constructed via the center of mass technique. The above has been extended to a non-compact nilpotent Lie group G in [CFG] (see discussion following Corollary 1.2.5.)

According to [Hu], the above stability result in [GK] holds if the condition, ‘ $|\text{sec}_{g_1}| \leq 1$ ’, is weakened to that $\text{Ric}_{g_1} \geq -(n-1)$ and all points in (M, g_1) are local rewinding (ρ, ϵ) -Reifenberg points.

Lemma 3.5.1. *Let M be an n -manifold equipped with two complete metrics, g_1 and g_0 , satisfying*

$$\text{Ric}_{g_1} \geq -(n-1), \quad |\text{sec}_{g_0}| \leq 1, \quad \text{inrad}(g_0) \geq i_0 > 0.$$

Then there exists a constant, $\epsilon(n, i_0) > 0$, such that if two isometric Lie group G -actions with respect to g and g_0 respectively, μ_1 and μ_0 , are ϵ -close i.e.,

$$\sup\{d_{g_0}(\mu_1(g^{-1}, \mu_0(g, x)), x), x \in M, g \in G\} < \epsilon, \quad 0 < \epsilon \leq \epsilon(n, i_0),$$

the two G -actions are conjugate via a diffeomorphism that is $\Psi(\epsilon|n, i_0)$ -isotopic to id_M .

Lemma 3.5.1 was stated in [Hu] for compact G ; it still holds without compactness of G because the conjugate map is constructed based on local geometry. Note that if G is not compact, then μ_1 and μ_0 are ϵ -close implies that the subgroup of G consisting of elements such that $\mu_1(g, \cdot) = \mu_0(g, \cdot)$ has a compact quotient.

Proof of Theorem 0.3’.

Consider $M_i \xrightarrow{\text{GH}} \text{pt}$ as in Theorem 0.3’, $\Gamma_i = \pi_1(M_i)$. By Theorem 1.4.1, Γ_i contains a nilpotent subgroup $\Gamma_{i,0}$ of index $\leq c(n)$. Let $M_{i,0} = \tilde{M}_i/\Gamma_{i,0}$. By Theorem 2.2, for i large $M_{i,0}$ admits a sequence of sub-bundles satisfying Theorem 2.1, thus there is a diffeomorphism, $f_i : N/\Gamma_{i,0} \rightarrow \tilde{M}/\Gamma_{i,0}$, preserving the sequence of sub-bundles, and the lifting map is an isomorphism, $\tilde{f}_i : N \rightarrow \tilde{M}_i$, of nilpotent Lie groups (determined by the sequence of sub-bundles).

By identifying \tilde{M}_i with N via \tilde{f}_i with the \tilde{f}_i^* -pullback metric, we may assume two Γ_i -actions μ_i on N , the other is obtained via the Malcév rigidity, $\Gamma_i < N \rtimes \text{Aut}(N)$, such that $\mu_1 = \mu_2$ on $\Gamma_{i,0}$, and μ_1 is the deck transformation of Γ_i , and $N/\mu_2(\Gamma_i)$ is an infra-nilmanifold. We shall apply Lemma 3.5.1 to slightly modify id_N to a Γ_i -conjugation, which descends to a diffeomorphism from M_i to N/Γ_i , an infra-nilmanifold.

To apply Lemma 3.5.1, a nearby ‘good’ metric (i.e., g_0) is required. We now define a left-invariant metric on N as follows: by Theorem 2.4 and following its

proof we may assume that a short basis for $\Gamma_{i,0}$ (at p_i) naturally divided into groups,

$$\begin{cases} \gamma_{i,1}, \dots, \gamma_{i,m_1}, \\ \gamma_{i,m_1+1}, \dots, \gamma_{i,m_2}, \\ \dots \\ \gamma_{i,m_{s-1}+1}, \dots, \gamma_{i,m_s=n}, \end{cases} \quad \begin{cases} \ell_1 = \max\{|\gamma_{i,1}|, \dots, |\gamma_{i,m_1}|\} \\ \ell_2 = \max\{|\gamma_{i,m_1+1}|, \dots, |\gamma_{i,m_2}|\} \\ \dots \\ \ell_s = \max\{|\gamma_{i,m_{s-1}+1}|, \dots, |\gamma_{i,m_s}|\} \end{cases}.$$

such that for $1 \leq j \leq s$, a blowup with ℓ_j^{-1} yields the following commutative diagram:

$$\begin{array}{ccc} (\ell_j^{-1}\tilde{M}_i, \tilde{p}_i, \Gamma_{i,0}) & \xrightarrow{\text{eqGH}} & (\mathbb{R}^n, 0, \mathbb{Z}^{k_j} \times \mathbb{R}^{n-m_{j-1}}) \\ \downarrow \pi_i & & \downarrow \text{proj} \\ (\ell_j^{-1}M_{i,0}, p_{i,0}) & \xrightarrow{\text{GH}} & (T^{k_j} \times \mathbb{R}^{n-m_{j-1}}, e \times 0). \end{array}$$

Assume that $\ell_j^{-1}\gamma_{i,m_j+1}, \dots, \ell_j^{-1}\gamma_{i,m_{j+1}}$ respectively converge to $\gamma_{m_j+1}, \dots, \gamma_{m_{j+1}}$, a canonical basis on flat T^{k_j} . Let $v_{i,k} = \gamma'_{i,k}(0)$ with $|\gamma'_{i,k}(0)| = \text{length}(\gamma_{i,k})$ (similarly, one gets $v_{m_j+1}, \dots, v_{m_{j+1}}$, a basis for $T_e T^{k_j}$). Then $\{v_{i,1}, \dots, v_{i,m_s=n}\}$ form a basis for the Lie algebra $\tilde{h}_i = T_{\tilde{p}_i}\tilde{M}_i = T_e N$. We then obtain a left invariant metric on N by specifying an inner product on a basis for $T_p\tilde{M}_i = T_e N$:

$$\tilde{g}_N(v_{i,k}, v_{i,l}) = \begin{cases} \ell_j(p_i)^2 \langle v_k, v_l \rangle & m_j + 1 \leq k, l \leq m_{j+1} \\ 0 & \text{otherwise} \end{cases}$$

where \langle, \rangle is the Euclidean inner product on flat T^{k_j} (roughly, we use the canonical basis on T^{k_j} to define the inner product for elements of \tilde{h}_i in the corresponding group, while elements of \tilde{h}_i in different groups are orthogonal).

From the above commutative diagram, it is clear that restricting to any R -ball at $e \in N$ with respect to \tilde{g}_N or \tilde{g}_i (i large), the two metrics are GH-close (to the limit Euclidean metric on $\mathbb{B}_R^n(0)$). Indeed, $\max |\sec_{\tilde{g}_N}| \rightarrow 0$ (otherwise, normalizing $\max |\sec_{\tilde{g}_N}| = 1$, one concludes that the left-invariant metric \tilde{g}_N is C^∞ -close to an Euclidean metric; a contradiction).

Let $\Gamma_i = \bigcup_{k=1}^s \alpha_k \Gamma_{i,0}$. For fixed $\alpha = \alpha_k$, $g \in N$, and any $\gamma \in \Gamma_{i,0}$,

$$\begin{aligned} d(\mu_2(\alpha\gamma)(\mu_1((\alpha\gamma)^{-1}, g), g) &= d(\mu_2(\alpha)\mu_2(\gamma)(\mu_1(\gamma^{-1})\mu_1(\alpha^{-1}g)), g) \\ &= d(\mu_2(\alpha)\mu_1(\alpha^{-1}g), g). \end{aligned}$$

Then $\{\mu_2(\gamma)^{-1}\mu_1(\gamma)(g), \gamma \in \Gamma_{i,0}\} \subset B_\rho(g)$, $\rho = \max\{d(\mu_2(\alpha_k)\mu_1(\alpha_k^{-1})(g), g), 1 \leq k \leq s\}$. We now specify a representative α_i as the projection of e to the coset, $\alpha_i \Gamma_{i,0}$ with respect to the \tilde{g}_N . Because (passing to a subsequence if necessary) the two metrics are GH-close and $\mu_2(\alpha)^{-1}\mu_1(\alpha)(e)$ can be very small (than the convexity radius of \tilde{g}_N), the finite set, $\{\mu_2(\alpha_k)^{-1}\mu_1(\alpha_k)(g)\} \subset N$, has a center of mass c_g .

By now we are able to apply Lemma 3.5.1 to $(N, \Gamma_i, \tilde{g}_N)$ and $(N, \Gamma_i, \tilde{g}_i)$, and conclude that to M_i is diffeomorphic to an infra-nilmanifold. \square

Corollary 3.5.2. *The finite group of isometries preserving the nilpotent structure of iterated bundles, thus preserves each sub-bundle. In particular, the structural group of each sub-bundles is affine.*

Remark 3.5.3. Note that the left-invariant metric on \tilde{M}_i is explicitly defined in terms of a short basis on $M_{i,0}$, which is different from ones via smoothing method.

Lemma 3.5.4. ((8.4) in [Ro4]) There are constants, $\epsilon(n), \delta(n) > 0$, such that if a compact n -manifold M satisfies

$$|\sec_M| \leq 1, \quad \text{diam}(M) < \epsilon < \epsilon(n),$$

then the injectivity radius of the Riemannian universal cover, $\text{inrad}(\tilde{M}) \geq \delta(n) > 0$.

Proof of Corollary 0.5.

By Lemma 3.5.4, an almost flat manifold M in Theorem 0.1, when normalizing metric by $|\sec_M| \leq 1$, satisfies the conditions of Theorem 0.3. Because the proofs of both Lemma 3.5.4 and Theorem 0.3 are independent of Theorem 0.1, Theorem 0.3 implies Theorem 0.1. \square

Together with Theorem 0.2, Lemma 3.5.4 has the following consequence, which is also independent of Theorem 0.1.

Corollary 3.5.5. ([CFG]) There exist constants, $\rho(n)$ and $v(n) > 0$, such that any complete n -manifold M with $|\sec_M| \leq 1$ satisfies a $(\rho(n), v(n))$ -bound covering geometry.

Proof. For any $x \in M$, without loss of generality we may assume that $\text{vol}(B_1(x))$ is small. Then by Theorem 0.2, we may assume the following commutative diagram,

$$\begin{array}{ccc} F \rightarrow (F(B_\rho(x)), O(n)) & \xrightarrow{\tilde{f}} & (Y, O(n)) \\ \text{proj} \downarrow & & \text{proj} \downarrow \\ B_\rho(x) & \xrightarrow{f} & B_\rho(x), \end{array}$$

where Y is a not collapsed (open) manifold. We now choose $\rho > 0$ such that \tilde{f} is a trivial fiber bundle i.e., $F(B_\rho(x))$ is diffeomorphic to $F \times Y$. Because Y is not collapsed, by Lemma 3.5.4 we conclude that the universal cover of $F(B_\rho(x))$ is not collapsed, thus $B_\rho(\tilde{x})$ is not collapsed. \square

4. PROOF OF THEOREM B

By a standard compactness, it suffices to consider a sequence, $M_i \xrightarrow{\text{GH}} N$, in Theorem B, and prove that for i large, there is a smooth bundle map and ϵ_i -GHA ($\epsilon_i \rightarrow 0$), $f_i : M_i \rightarrow N$, with fiber infra-nilmanifold and affine structural group.

We first observe a basic property on local fundamental groups.

Lemma 4.1. (The unique local fundamental group) *Let $M_i \xrightarrow{\text{GH}} N$ be as in Theorem B. Let $\delta = \min\{\rho, 2 \text{inrad}(N)\}$, and $0 < \epsilon \leq \epsilon(n)$, a constant in Theorem 1.4.2. For any $x_i \in M_i$ and i large, the isomorphic class,*

$$\Gamma_i(\epsilon) = \text{Im}[\pi_1(B_\epsilon(x_i), x_i) \rightarrow \pi_1(B_\delta(x_i), x_i)],$$

is independent of x_i .

Proof. Assume the following commutative diagram:

$$\begin{array}{ccc}
(B_\rho(\tilde{x}_i), \tilde{x}_i, \Gamma_{x_i}(\epsilon)) & \xrightarrow{\text{eqGH}} & (B_\rho(\tilde{x}), \tilde{x}, G) \\
\pi_i \downarrow & & \text{proj} \downarrow \\
(B_\rho(x_i), x_i) & \xrightarrow{\text{GH}} & (B_\rho(x), x).
\end{array}$$

In general, $\Gamma_{x_i}(\epsilon)$ contains a normal subgroup $\Gamma_i(\epsilon)$, of bounded index such that $\Gamma_i(\epsilon) \rightarrow G_0$, the identity component of G , where $\Gamma_i(\epsilon)$ is generated by a subset of short generators of $\Gamma_{x_i}(\epsilon)$ that moves every point in $B_{\frac{\rho}{2}}(\tilde{x}_i)$ uniformly small.

Because $B_\rho(x)$ is Riemannian and diffeomorphic to $\mathbb{B}_\rho^m(0)$, $G = G_0$ which (pseudodolly) acts freely, hence $\Gamma_{x_i}(\epsilon) \cong \Gamma_i(\epsilon)$ i.e., the isomorphism class of $\Gamma_{x_i}(\epsilon)$ is independent of points in $B_{\frac{\rho}{2}}(x_i)$. \square

Note that Lemma 4.1 is false if one replaces N with a length space, or if one removes the local (ρ, v) -bound Ricci covering geometry (Examples 1.5.3 and 1.5.2).

Proof of Theorem B (Method 1).

Let $M_i \xrightarrow{\text{GH}} N$ be as in Theorem B. By Theorem 0.4 for i large one may assume a smooth fiber bundle map, $F_i : M_i \xrightarrow{f_i} N$, which is an ϵ_i -GHA.

For any $x_i \in M_i$, by performing a successively blow up at x_i at rates determined by a short basis in $\Gamma_i(\epsilon)$ (Lemma 4.1), and following the proof of Theorem A (step by step), we conclude that a fiber F_i is diffeomorphic to an infra-nilmanifold, and the fiber bundle has a local trivialization compatible with the nilpotent structure on F_i (because the last blow-up at x_i yields a metric product limit, $X_1 \times \mathbb{R}^{n-m}$, $m = n - \dim(F_i)$). Because the infra-nilmanifold structure on F_i is determined a short basis for $\Gamma_i(\epsilon)$ at x_i , the structural group can be reduced to affine. \square

Indeed, Theorem B can also be proved without Theorem 0.4: fixing large i one may construct on M_i a local finite open cover by elementary N-structures such that each chart also carries an invariant metric determined by a short basis in the fiber so that on overlaps these invariant metrics are C^1 -close in terms of the gluing criterion on N-structures in [CFG] (see Remark 1.2.6). Our construction of an elementary N-structure and an invariant metrics follow the same arguments in the proof of Theorem A. Because the proof has its own interest, we will present it below.

Let's first consider a special case in Theorem B.

Lemma 4.2. *Let $M_i \xrightarrow{\text{GH}} N$ be as in Theorem B. Assume that for i large, $\Gamma_i(\epsilon)$ is nilpotent. Then there is a smooth fiber bundle map, $f_i : M_i \rightarrow N$, with fiber a nilmanifold and an affine structural group, and f_i is ϵ_i -GHA, $\epsilon_i \rightarrow 0$.*

Proof. We will only outline the proof because it imitates the proof of Theorem 2.2 (step-by-step) with obvious minor modifications.

Recall the proof of Theorem 2.2, given $M_i \xrightarrow{\text{GH}} \text{pt}$ with (ρ, v) -bound Ricci covering geometry such that $\Gamma_i = \pi_1(M_i)$ is nilpotent, we proved that, by via a successive blow-ups at rate depends on short basis at $x_i \in M_i$, M_i admits a sequence of bundle satisfying Theorem 2.1, hence M_i is diffeomorphic to a nilmanifold. The key ingredient is that each blow-ups is a metric product of flat $T^{k_j} \times \mathbb{R}^{m_j-1}$, such that T^{k_j} has a uniformly bounded geometry, which leads to a construction of a sequence of

bundles. For instance, the fastest rate of blow ups leads to a principal $T^{k_{s+1}}$ -bundle over M_i , and the slowest below-up leads to a $F_{i,1}$ -bundle on M_i over T^{k_1} .

We now consider $M_i \xrightarrow{\text{GH}} N$ with (ρ, v) -bound Ricci covering geometry such that $\Gamma_i(\epsilon)$ is nilpotent. For any $x_i \in M_i$, we perform a successive blow-ups at x_i with rates determined by a short basis for $\Gamma_i(\epsilon)$ at x_i . The fastest rate blow-up, $\ell_s(x_i)^{-1}M_i$,

$$\begin{array}{ccc} (\ell_s(x_i)^{-1}B_1(\tilde{x}_i), \tilde{x}_i, \Gamma_i(\epsilon)) & \xrightarrow{\text{eqGH}} & (\mathbb{R}^n, \tilde{x}, \mathbb{Z}^{k_{s+1}}) \\ \pi_i|_{B_1(\tilde{x}_i)} \downarrow & & \text{proj} \downarrow \\ (\ell_i(x_i)^{-1}B_1(x_i), x_i) & \xrightarrow{\text{GH}} & (T^{k_{s+1}} \times \mathbb{R}^{n-k_{s+1}}, e \times 0). \end{array}$$

which implies that for i large, M_i has locally finite an open cover by elementary $T^{k_{s+1}}$ -structures, thus by Theorem 1.2.3 we construct a bundle, $T^{k_{s+1}} \rightarrow M_i \rightarrow B_{i,s}$, with an affine structural group. Here differences from the proof of Theorem 2.2 are that this $T^{k_{s+1}}$ -bundle may not be principal, and that the last blow-up yields a T^{k_1} -bundle, $T^{k_1} \rightarrow B_1 \rightarrow N$, because the limit space of the last blow-up is $T^{k_1} \times \mathbb{R}^m$ (comparing to the proof of Theorem 2.2). also yields a bundle, $F_{i,1} \rightarrow M_i \rightarrow N$, with an affine structural group. Note that because $\pi_1(F_{i,1}) = \Gamma_i(\epsilon)$ is nilpotent, the restriction of a sequence of fiber bundles on M_i to $F_{i,1}$ satisfies Theorem 2.1 and therefore $F_{i,1}$ is diffeomorphic to an nilmanifold. \square

Note that the proof of Lemma 4.2 may not carry over if $\Gamma_i(\epsilon)$ is an infra-nil group, because it requires something like that there is a finite normal cover \hat{M}_i of M_i such that $\hat{\Gamma}_i(\epsilon)$ is a normal subgroup of $\Gamma_i(\epsilon)$. A priori, this property unlikely holds.

Observe that the above-desired property always holds over a local trivialization of the bundle with fiber infra-nilmanifold, hence we may apply the gluing construction in [CFG] to obtain a global fiber bundle with an affine structural group.

By the proof of Lemma 4.2 and the last part of the proof of Theorem 2.2 (using that $M_{i,0}$ is nilmanifold to show that M_i is infra-nilmanifold), when restricting to a local trivialization over $B_\rho(x) \subset N$, we obtain the following elementary N-structures.

Lemma 4.3. (Uniform elementary N-structures) *Let $M_i \xrightarrow{\text{GH}} N$ be as in Theorem B. For any $\delta > 0$, up to a scaling we may assume that any 2-ball on N is δ -close to $B_2^m(0)$ in C^3 -norm. When δ is suitably small, for i large the following hold:*

(4.3.1) *For any $x \in N$, $x_i \in M_i$, such that $x_i \rightarrow x$, there is open subset, $B_1(x_i) \subset U_i \subset B_2(x_i)$, and a fiber bundle map, $\phi_i : U_i \rightarrow B_1(x)$, which is an ϵ_i -GHA, $\epsilon_i \rightarrow 0$.*

(4.3.2) *A ϕ_i -fiber F_i is diffeomorphic to an infra-nilmanifold, equipped with a product metric determined by $\Gamma_i(\epsilon) = \pi_1(F_{i,1})$.*

(4.3.3) *U_i , equipped with the product of the invariant metric on $F_{i,1}$ in (4.3.2) and the Euclidean metric on $B_1(x)$ pullbacked by \exp_x^{-1} , satisfies the C^1 -closeness condition on overlaps.*

Proof of Theorem B (Method 2).

Let $M_i \xrightarrow{\text{GH}} N$ be as in Theorem B. By rescaling N and M_i with a suitable large constant, without loss of generality we may assume that $M_i \xrightarrow{\text{GH}} N$ satisfies the conditions of Lemma 4.3.

Let $\{x_\alpha\}$ be a $\frac{1}{4}$ -net on N . Fixing i large, for each α , applying Lemma 4.3 we obtain a locally finite open cover for M_i by atlas of elementary nilpotent structures, $\{(U_{i,\alpha}, \phi_{i,\alpha})\}$. For each α , let $V_{i,\alpha} = \phi_{i,\alpha}^{-1}(B_{\frac{1}{4}}(x_{i,\alpha}))$. Clearly, $\{(V_{i,\alpha}, \phi_{i,\alpha}|_{V_{i,\alpha}})\}$ also forms an atlas of elementary N-structures. We will give each V_i a left-invariant metric so that the C^1 -closeness of the gluing criterion in Appendix 2 in [CFG] holds (Remark 1.2.6).

In the proof of Theorem 0.3', we construct a left-invariant metric on $M_{i,0}$, completely determined by geometry from a short basis for Λ_i at x_i . Using the left-invariant metric, we construct a diffeomorphism, $M_i \rightarrow N_i/\Gamma$, hence a left-invariant metric on M_i (the pullback metric).

Similarly we construct, on a $\phi_{i,\alpha}$ -fiber F_i , a left-invariant metric determined by $\Gamma_i(\epsilon)$. By the local uniform short basis property in Lemma 3.2.3 (based on the uniform Reifenberg points on Riemannian universal cover), it is clear that fixing a left-invariant metric on F_i as the above and choosing $\delta > 0$ sufficiently small, equip $V_{i,\alpha}$ a product metric with the Euclidean metric on $\mathbb{B}_{\frac{1}{4}}^m(0)$. Then the following properties hold: if $V_{i,\alpha} \cap V_{i,\beta} \neq \emptyset$, then $B_{\frac{1}{20}}(z_i) \subset V_{i,\alpha} \cap V_{i,\beta}$ and $V_{i,\alpha}, V_{i,\beta} \subset U_{z_i}$ which satisfies Lemma 4.3, and it is clear that $V_{i,\alpha}$ and $V_{i,\beta}$ into the elementary N-structure, $\psi : U_{z_i}$, and these product metrics are ϵ_i -close, their pullback metrics on $\widetilde{V_{i,\alpha} \cap V_{i,\beta}}$ are C^1 -close (away from the boundary of $\widetilde{V_{i,\alpha} \cap V_{i,\beta}}$) in the sense of Appendix 2 in [CFG] (Remark 1.2.6)). Hence, we are able to glue the local trivial elementary N-structures on M_i to obtain a global pure N-structure, which is a fiber bundle with fiber infra-nilmanifold and affine structural group. \square

5. PROOF OF THEOREM C

The following result will be used in the proof of (C2); which is not required if one assumes that a Ricci limit space is a manifold.

Theorem 5.1. ([Wa]) *Given a sequence of complete n -manifolds, $(M_i, p_i) \xrightarrow{\text{GH}} (X, p)$ such that $\text{Ric}_{M_i} \geq -(n-1)$, then*

(5.1.1) *X is semi-locally simply connected (hence the universal cover \tilde{X} is simply connected) i.e., for any $x \in X$, there is $r_x > 0$ such that any loop at x that is contained in $B_{r_x}(x)$ is homotopically trivial in X ;*

(5.1.2) *assume $R > 0$ such that $\pi_1(M_i, p_i)$ is R -represented i.e., there is a set of generators and relations of bounded number with length $\leq R$. Then there is an onto homomorphism, $h_i : \pi_1(M_i, p_i) \rightarrow \pi_1(X, p)$.*

Remark 5.2. Note that (5.1.2) was originally stated for M_i with $\text{diam}(M_i) \leq R$. However, (5.1.2) can be seen from the proof in [Wa].

Proof of Theorem C.

(C1) We start with the following commutative diagram:

$$\begin{array}{ccc} (\bar{B}_1(\tilde{x}_i), \Gamma_{x_i}(\epsilon)) & \xrightarrow{\text{eqGH}} & (\bar{B}_1(\tilde{x}), G) \\ \downarrow \pi_i & & \downarrow \text{proj} \\ \bar{B}_1(x_i) & \xrightarrow{\text{GH}} & \bar{B}_1(x) = \bar{B}_1(\tilde{x})/G. \end{array}$$

By Theorem 1.4.2, $\Gamma_{x_i}(\epsilon)$ ($\epsilon < \epsilon(n)$) contains a nilpotent subgroup index $\leq c(n)$, thus $\Gamma_{x_i}(\epsilon)$ contains a torsion-free nilpotent subgroup, Λ_i , of finite index ([Ra]). We define $\text{rank}(\Gamma_{x_i}(\epsilon)) := \text{rank}(\Lambda_i)$. Let $(\bar{B}_1(\tilde{x}_i), \Lambda_i) \xrightarrow{\text{eqGH}} (\bar{B}_1(\tilde{x}), \hat{G})$, and we shall show that $\text{rank}(\Lambda_i) \leq \dim(\hat{G}) \leq \dim(G)$, where \hat{G} (resp. G) contains a neighborhood of the identity which isomorphically embeds into a Lie group ([Wa]). Without loss of generality, we may assume that $\text{rank}(\Lambda_i) = m \geq 1$.

Let $\gamma_{i,1}, \dots, \gamma_{i,m}$ denote a short basis for Λ_i in an order of length increasing. We first assume that the subgroup, $\Lambda_{i,s} \cong \mathbb{Z}$, generated by the first s elements is normal in $\Lambda_{i,s+1}$, $\langle \gamma_{i,s+1}, \Lambda_{i,s} \rangle \cong \mathbb{Z}^2$, and $\Lambda_{i,s+1}/\Lambda_{i,s} \cong \mathbb{Z}$. Let $\Lambda_{i,s} \rightarrow G_s$, and it suffices to show that $\text{rank}(\Lambda_{i,s}) \leq \dim(G_s)$. Because $o(\gamma_{i,1}) = \infty$ and $|\gamma_{i,1}| \rightarrow 0$, $\dim(G_s(\tilde{x})) \geq 1$, thus $\dim(G_s) \geq 1$. Assume that $\text{rank}(\Lambda_{i,s}) \leq \dim(G_s)$, and passing to a subsequence we may assume $(\bar{B}_1(\tilde{p}_i), \Lambda_{i,s}) \xrightarrow{\text{eqGH}} (\bar{B}_1(\tilde{p}), G_s)$. By Lemma 1.1.4, $(\bar{B}_1(\tilde{p}_i)/\Lambda_{i,s}, \Lambda_{i,s+1}/\Lambda_{i,s}) \xrightarrow{\text{eqGH}} (\bar{B}_1(\tilde{p})/G_s, H_s)$. Because H_s contains a short generator $\gamma_{i,s+1}$ such that $|\gamma_{i,s+1}| \rightarrow 0$ and $o(\gamma_{i,s+1}) = \infty$, $\dim(H_s(\tilde{x})) \geq 1$, thus $\dim(H_s) \geq 1$. Then

$$\begin{aligned} \text{rank}(\Lambda_{i,s+1}) &= \text{rank}(\Lambda_{i,s}) + 1 \leq \dim(G_s) + 1 \\ &\leq \dim(G_s) + \dim(H_s) = \dim(G_{s+1}). \end{aligned}$$

In general, we divide short generators for Λ_i into groups satisfying the following condition:

$$\gamma_{i,1}, \dots, \gamma_{i,k_1}, \quad \gamma_{i,k_1+1}, \dots, \gamma_{i,k_2}, \quad \dots, \quad \gamma_{i,k_{m-1}+1}, \dots, \gamma_{i,k_m},$$

such that $|\gamma_{i,m}| \rightarrow 0$ as $i \rightarrow \infty$, and

$$\left\{ \begin{array}{l} \text{rank}(\Lambda_{i,1}) = 1 \quad \Lambda_{i,1} = \langle \gamma_{i,1}, \dots, \gamma_{i,k_1} \rangle \\ \text{rank}(\Lambda_{i,2}) = 2 \quad \Lambda_{i,2} = \langle \gamma_{i,1}, \dots, \gamma_{i,k_1}, \gamma_{i,k_1+1}, \dots, \gamma_{i,k_2} \rangle \\ \dots \\ \text{rank}(\Lambda_{i,m}) = m \quad \Lambda_{i,m} = \langle \gamma_{i,1}, \dots, \gamma_{i,k_1}, \gamma_{i,k_1+1}, \dots, \gamma_{i,k_2}, \dots, \gamma_{i,k_m} \rangle, \end{array} \right.$$

every $\Lambda_{i,s}$ is normal in Λ_i , and $\Lambda_{i,s+1}/\Lambda_{i,s} \cong \mathbb{Z}$. Passing to a subsequence we may assume that $(\bar{B}_1(\tilde{x}_i), \Lambda_{i,j}) \xrightarrow{i \rightarrow \infty} (\bar{B}_1(\tilde{x}), G_j)$, $1 \leq j \leq m$. Then the above proof (replacing γ_{s+1} with γ_{k_s+1}) will go through and complete the proof.

We now assume that $n - \dim(X) = \text{rank}(\Lambda_i) \leq \dim(G) = \dim(B_1(\tilde{x})) - \dim(B_1(x)) \leq n - \dim(X)$. Then $\dim(B_1(\tilde{x})) = n$, thus we can apply Theorem 1.3.3, $\text{vol}(B_1(\tilde{x}_i)) \rightarrow \text{vol}(B_1(\tilde{x})) \geq v > 0$.

(C2) In the commutative diagram, by Theorem 1.3.7 $\text{Isom}(\tilde{X})$ is a Lie group, thus its a closed subgroup G is a Lie group and G/G_0 is a discrete group, where G_0 is the identity component of G . By Lemma 1.1.4, for any $0 < \epsilon \leq \epsilon(n)$, Γ_i contains a normal subgroup $\Gamma_i(\epsilon)$ such that $\Gamma_i/\Gamma_i(\epsilon) \cong G/G_0$.

By Theorem 5.1, the universal cover of X , \tilde{X}/H , where H is a normal subgroup generated by G_0 and isotropy subgroups ([EW]), is simply connected. Again by Lemma 1.1.4, we may assume a normal subgroup, $\Gamma'_i(\epsilon)$ such that $\Gamma'_i(\epsilon) \rightarrow H$ and $\Gamma_i/\Gamma'_i(\epsilon) \cong G/H$ (i large) satisfying the following commutative diagram:

$$\begin{array}{ccc} (\tilde{M}_i/\Gamma'_i(\epsilon), p'_i, \Gamma_i/\Gamma'_i(\epsilon)) & \xrightarrow{\text{eqGH}} & (\tilde{X}/H, p', G/H) \\ \pi_i \downarrow & & \text{proj} \downarrow \\ (M_i, p_i) & \xrightarrow{\text{GH}} & (X, p). \end{array}$$

By (5.1.2), we may assume the following commutative diagram:

$$\begin{array}{ccc} \pi_1(\tilde{M}_i/\Gamma'_i(\epsilon), p'_i) & \xrightarrow{\psi_i} & \pi_1(\tilde{X}/H, p') = e \\ (\pi_i)_* \downarrow & & \text{proj}_* \downarrow \\ \pi_1(M_i, p_i) & \xrightarrow{h_i} & \pi_1(X, p). \end{array}$$

Because h_i is an onto homomorphism, h_i induces an onto homomorphism,

$$\frac{\pi_1(M_i)}{[\pi_1(M_i), \pi_1(M_i)]} = H_1(M_i) \xrightarrow{h_i} H_1(X) = \frac{h_i(\pi_1(M_i))}{[h_i(\pi_1(M_i)), h_i(\pi_1(M_i))]}.$$

Thus we obtain the following commutative diagram,

$$\begin{array}{ccc} H_1(\tilde{M}_i/\Gamma'_i(\epsilon); \mathbb{Z}) & \xrightarrow{\psi_i} & H_1(\tilde{X}/H; \mathbb{Z}) = e \\ (\pi_i)_* \downarrow & & \text{proj}_* \downarrow \\ H_1(M_i; \mathbb{Z}) & \xrightarrow{h_i} & H_1(X; \mathbb{Z}), \end{array}$$

Hence, $\ker(h_i) = \text{im}((\pi_i)_*(H_1(\tilde{M}_i/\Gamma'_i(\epsilon); \mathbb{Z}))) = \Gamma_i(\epsilon)' / ([\Gamma_i, \Gamma_i] \cap \Gamma'_i(\epsilon))$. Then

$$\begin{aligned} b_1(M_i) - b_1(X) &= \text{rank}(\Gamma_i(\epsilon)' / ([\Gamma_i, \Gamma_i] \cap \Gamma'_i(\epsilon))) \\ &\leq \text{rank}(\Gamma_i(\epsilon)' / [\Gamma_i(\epsilon), \Gamma'_i(\epsilon)]). \end{aligned}$$

Because $\Gamma_i(\epsilon)/\Gamma'_i(\epsilon)$ is bounded for all i , as in (5.1.1) $\text{rank}(\Gamma_i(\epsilon)' / [\Gamma_i(\epsilon), \Gamma'_i(\epsilon)]) \leq \dim(G_0)$, and “ $b_1(M_i) - b_1(X) = n - m$ ” implies that $\dim(G_0) \geq m - n$, thus $\dim(B_1(\tilde{x})) = \dim(B_1(\tilde{x})) + \dim(G_0) \geq n - m + m = n$. By Theorem 1.3.3, $\text{vol}(B_1(\tilde{p}_i)) \rightarrow \text{vol}(B_1(\tilde{p})) \geq v > 0$. \square

Proof of Theorem 0.8.

We claim that the Riemannian universal cover of M is not collapsed, by Theorem A M is diffeomorphic to an infra-nilmanifold. Because $b_1(M) = n$, M is diffeomorphic to a standard torus.

We verify the claim via a contradiction argument; assuming a sequence of $M_i \xrightarrow{\text{GH}} \text{pt}$ such that $\text{Ric}_{M_i} \geq -(n-1)$ and $b_1(M_i) = n$. Then local fundamental group of a ball radius 1 is $\pi_1(M_i)$, thus M_i satisfies local maximal rank in (C1). Therefore the Riemannian universal cover of M_i is not collapsed. \square

Proof of Theorem 0.10.

By (C2), the Riemannian universal of M has a Ricci bounded covering geometry and thus we apply Theorem B to conclude a fiber bundle, $F \rightarrow M \rightarrow N$, with F diffeomorphic to an infra-nilmanifold of dimension $(n-m)$ and affine structure group. From the proof of (C2), we see that $b_1(M) - b_1(N) = n - m$ implies that $\pi_1(F) \cong \mathbb{Z}^{n-m}$, thus F is a torus. \square

Acknowledgment. The author would like to thank Shaosai Huang for some help comments on a draft of the present paper.

REFERENCES

- [Ab] U. Abresch, *Über das glätten Riemann'scher metriken*, Habilitationsschrift Rheinischen Friedrich-Wilhelms-Universität Bonn (1998).
- [An] M. T. Anderson, *Hausdorff perturbations of Ricci-Flat manifolds and the splitting theorem*, Duke Math. J. **68** (1992), 67-82.
- [Be] I. Belegradek, *Iterated circle bundles and infra-nilmanifolds*, arXiv:1805.06585 [math.GT].
- [CR] Q. Cai; X. Rong, *Collapsing construction with nilpotent structures*, Geom. Funct. Anal. **18** (2009), 1503-1524.
- [CM] F. Cavalletti; A. Mondino, *Almost Euclidean isoperimetric inequalities in spaces satisfying local Ricci curvature lower bounds*, International Mathematics Research Notices (2020), 1481-1510.
- [CC1] J. Cheeger; T. H. Colding, *Lower Bounds on Ricci Curvature and the Almost Rigidity of Warped Products*, Ann. of Math. **144** (Jul., 1996), no. 1, 189-237.
- [CC2] J. Cheeger; T. H. Colding, *On the structure of space with Ricci curvature bounded below I*, J. Diff. Geom. **46** (1997), 406-480.
- [CC3] J. Cheeger; T. H. Colding, *On the structure of spaces with Ricci curvature bounded below. III*, J. Diff. Geom. **54**. no. **1** (2000), 37-74.
- [CFG] J. Cheeger, K. Fukaya; M. Gromov, *Nilpotent structures and invariant metrics on collapsed manifolds*, J. Amer. Math. Soc. **5** (1992), 327-372.
- [CG1] J. Cheeger; D. Gromov, *Collapsing Riemannian manifolds while keeping their curvature bounded I*, J. Differential Geom. **23** (1986), 309-346.
- [CG2] J. Cheeger; D. Gromov, *Collapsing Riemannian manifolds while keeping their curvature bounded II*, J. Differential Geom. **32** (1990), 269-298.
- [CJN] J. Cheeger, W. Jiang; A. Naber, *Rectifiability of Singular Sets in Noncollapsed Spaces with Ricci Curvature bounded below*, Ann. of Math. **193** (2021), no. No. 3, 407-538.
- [ChN] J. Cheeger; A. Naber, *Regularity of Einstein manifolds and the codimension 4 conjecture*, Ann. of Math. **182** (2015), no. No. 3, 1093-1165.
- [CRX1] L. Chen, X. Rong; S. Xu, *Quantitative volume space form rigidity under lower Ricci curvature bound I*, J. Diff. Geom. **113** (2019), 222-272.
- [CRX2] L. Chen, X. Rong; S. Xu, *Quantitative volume space form rigidity under lower Ricci curvature bound II*, Trans. AMS **370** (2017), no. 6, 4509-4523.
- [Co1] T. H. Colding, *Large manifolds with positive Ricci curvature*, Invent. Math. **124** (1996), no. 1-3, 193-214.
- [Co2] T. H. Colding, *Ricci curvature and volume convergence*, Ann. of Math **145** (1997), no. 3, 477-501.
- [CoN] T. H. Colding; A. Naber, *Sharp Hölder continuity of tangent cones for spaces with a lower Ricci curvature bound and applications*, Ann. Math. **176** (2012), 1173-1229.
- [DWY] X. Dai, G. Wei; R. Ye, *Smoothing Riemannian metrics with Ricci curvature bounds*, Manuscripta Math **90** (1996), 49-61.
- [EW] J. Ennis; G. Wei, *Describing universal cover of a compact limit*, Diff. Geom. Appl **24** (2006), 554-562.
- [Fu1] K. Fukaya, *Collapsing Riemannian manifolds to ones with lower dimensions*, J. Differential Geom. **25** (1987), 139-156.
- [Fu2] K. Fukaya, *A boundary of the set of Riemannian manifolds with bounded curvature and diameters*, J. Differential Geom. **28** (1988), 1-21.
- [Fu3] K. Fukaya, *Collapsing Riemannian manifolds to ones with lower dimensions II*, J. Math. Soc. Japan **41** (1989), 333-356.
- [FY] K. Fukaya; T. Yamaguchi, *The fundamental groups of almost non-negatively curved manifolds*, Ann. of Math. **136** (1992), 253-333.
- [Gr] M. Gromov, *Almost flat manifolds*, J. Diff. Geom. **13** (1978), 231-241.
- [GTZ1] M. Gross; V. Tosatti; Y. Zhang, *Collapsing of abelian fibered Calabi-Yau manifolds*, Duke Math. J. **162** (2013), 517-551.
- [GTZ2] M. Gross; V. Tosatti; Y. Zhang, *Gromov-Hausdorff collapsing of Calabi-Yau manifolds*, Communication of Geometry and Analysis **24** (2016), 91-113.
- [GK] K. Grove; H. Karcher, *How to conjugate C^1 -close actions*, Math. Z **132** (1973), 11-20.
- [GW] M. Gross; P.M.H. Wilson, *Large complex structure limits of K3 surfaces*, J. Diff. Geom **55** (2000), 475-546.

- [Ha] R. Hamilton, *Three-manifolds with positive Ricci curvature*, J. Diff. Geom **17** (1982), 233-306.
- [HSVZ] H. Hein, S. Song, J. Viaclovsky; R. Zhang, *Nilpotent structures and collapsing Ricci-flat manifolds on $K3$ surfaces*, J. A.M.S. **35** (2022), 123-209.
- [Hu] H. Huang, *Fibrations, and stability for compact Lie group actions on manifolds with bounded local Ricci covering geometry*, Front. Math. China **15** (2020), 69-89.
- [HKRX] H. Huang, L. Kong, X. Rong; S. Xu, *Collapsed manifolds with Ricci bounded covering geometry*, Trans. A.M.S. **373** (2020), no. 11, 8039-8057.
- [HRW] S. Huang, X. Rong; B. Wang, *On the collapsing geometry of Ricci curvature bounded below and Ricci flow smoothing*, SIGMA (Special issue on scalar and Ricci curvature in honor of Misha Gromov on his 75th Birthday) **16** (2020), 1-25.
- [HW1] S. Huang; B. Wang, *Rigidity of first Betti number via Ricci flow smoothing*, arXiv:2004.09762.
- [HW2] S. Huang; B. Wang, *Ricci flow smoothing for locally collapsing manifolds*, arXiv:2008.09956.
- [Ka] V. Kapovitch, *Mixed curvature of almost flat manifolds*, Geom. & top. **25** (2021), 2017-2059.
- [KW] V. Kapovitch; B. Wilking, *Structure of fundamental groups of manifolds with Ricci curvature bounded below*, arXiv:1105.5955v2.
- [NZ] A. Naber; R. Zhang, *Topology and ϵ -regularity theorems on collapsed manifolds with Ricci curvature bounds*, Geom. Topol. **20** (2016), 2575-2664.
- [Na] M. Nakayama, *Seifert construction for nilpotent groups and application to S^1 -fibred nil-Bott Tower*, RIMS Kkyuroku Bessatsu **B39** (2013), 121-136.
- [Pa] R. Palais, *Equivalence of nearby differentiable actions of a compact group*, Bull. AMS. **67** (1961), 362-364.
- [Per] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications*, arXiv: math/0211159v1 (2002).
- [PWY] P. Petersen, G. Wei; R. Ye, *Controlled geometry via smoothing*, Comment. Math. Helv **74** (1999), 345-363.
- [Ra] M. S. Raghunathan, *Discrete Subgroups of Lie groups*, Springer-Verlag Berlin Heidelberg, New York (1972).
- [Ro1] X. Rong, *Collapsed manifolds with bounded curvature and applications*, Surveys in J. Diff. Geom. **XI** (2007), 1-23.
- [Ro2] X. Rong, *Convergence and collapsing theorems in Riemannian geometry*, Handbook of Geometric Analysis, Higher Education Press and International Press, Beijing-Boston (2010), no. II ALM 13, 193-298.
- [Ro3] X. Rong, *Manifolds with Ricci curvature and local rewinding volume bounded below*, Scientia Sinica Mathematica **28** (2018), 1-16.
- [Ro4] X. Rong, *A new proof of Gromov's theorem on almost flat manifolds*, ArXiv: 1906.03377v2 [math.DG] 15 Mar 2020.
- [Ru] E. Ruh, *Almost flat manifolds*, J. Diff. Geom. **17** (1982), 1-14.
- [Wa] J. Wang, *Ricci limit spaces are semi-locally simply connected*, arXiv:2104.02460v1.
- [XY] S. Xu; X. Yao, *Margulis lemma and Hurewicz fibration Theorem on Alexandrov spaces*, CCM (accepted) (2022).
- [Zh] Y. Zhang, *Collapsing of negative Kähler-Einstein metrics*, Mathematical Research Letters **22** (2015), 1843-1869.

MATHEMATICS DEPARTMENT, RUTGERS UNIVERSITY NEW BRUNSWICK, NJ 08903 USA
E-mail address: rong@math.rutgers.edu

MATHEMATICS DEPARTMENT, CAPITAL NORMAL UNIVERSITY, BEIJING, P.R.C.