

Generalized Positive Energy Representations of Groups of Jets

Niestijl, M.

December 1, 2022

Abstract

Let V be a finite-dimensional real vector space and K a compact simple Lie group with Lie algebra \mathfrak{k} . Consider the Fréchet-Lie group $G := J_0^\infty(V; K)$ of ∞ -jets at $0 \in V$ of smooth maps $V \rightarrow K$, with Lie algebra $\mathfrak{g} = J_0^\infty(V; \mathfrak{k})$. Let P be a Lie group and write $\mathfrak{p} := \text{Lie}(P)$. Let α be a smooth P -action on G . We study smooth projective unitary representations $\bar{\rho}$ of $G \rtimes_\alpha P$ that satisfy a so-called generalized positive energy condition. In particular, this class captures representations that are in a suitable sense compatible with a KMS state on the von Neumann algebra generated by $\bar{\rho}(G)$. We show that this condition imposes severe restrictions on the derived representation $d\bar{\rho}$ of $\mathfrak{g} \rtimes \mathfrak{p}$, leading in particular to sufficient conditions for $\bar{\rho}|_G$ to factor through $J_0^2(V; K)$, or even through K .

1 Introduction

This paper is concerned with projective representations of groups and Lie algebras of jets. Let K denote a compact simple Lie group with Lie algebra \mathfrak{k} and let V be a finite-dimensional real vector space. Then we consider the Fréchet-Lie group $G := J_0^\infty(V; K)$ with Lie algebra $\mathfrak{g} := J_0^\infty(V; \mathfrak{k}) \cong \mathbb{R}[[V^*]] \otimes \mathfrak{k}$. These consist of ∞ -jets at $0 \in V$ of smooth K - and \mathfrak{k} -valued functions, respectively. We are interested in smooth projective unitary representations of G which satisfy either a so-called positive energy, or a KMS condition.

To describe these, let P be a finite-dimensional Lie group with Lie algebra \mathfrak{p} . Assume that there is a smooth action α of P on G . A continuous projective unitary representation $\bar{\rho} : G \rtimes_\alpha P \rightarrow \text{PU}(\mathcal{H}_\rho)$ is of *positive energy* at the cone $\mathcal{C} \subseteq \mathfrak{p}$ if for every $p \in \mathcal{C}$, there is a strongly continuous homomorphic lift $t \mapsto e^{itH}$ of $t \mapsto \bar{\rho}(e^{tp})$ whose generator H satisfies $\text{Spec}(H) \geq 0$. We say that $\bar{\rho}$ is KMS at $p \in \mathfrak{p}$ if there is a normal state ϕ on the von Neumann algebra $\mathcal{M} := \bar{\rho}(G)''$ generated by $\bar{\rho}(G)$ such that ϕ satisfies the KMS condition for the one-parameter group $\mathbb{R} \rightarrow \text{Aut}(\mathcal{M})$, $t \mapsto \text{Ad}(\bar{\rho}(e^{tp}))$. As we shall see, these two seemingly very different classes of representations exhibit similar behavior in certain respects. In particular, they both give rise to so-called *generalized positive energy representations*, a notion which relaxes the positive energy condition and is introduced in Section 4 below. We study these generalized positive energy representations and thereby also those which satisfy either the positive energy or the KMS condition.

The motivation for looking at the positive energy and KMS representations of the group $G \rtimes_\alpha P$ originates in prior work by B. Janssens and K.H. Neeb, who studied in [JN21] a class of projective unitary representations of the group of compactly supported gauge transformations $\mathcal{G} := \Gamma_c(M; \text{Ad}(\mathcal{K}))$ of a principal K -bundle \mathcal{K} over M , where $\text{Ad}(\mathcal{K})$ denotes the corresponding adjoint bundle. Suppose that the Lie group P acts smoothly on \mathcal{K} by automorphisms of the principal bundle \mathcal{K} . This induces a smooth action of P on the infinite-dimensional Lie group \mathcal{G} . Their main result is:

Theorem 1.1. ([JN21, Theorem 7.19]):

Let $(\bar{\rho}, \mathcal{H})$ be a projective unitary representation of $\mathcal{G} \rtimes P$ which has a dense set of smooth rays and is of positive energy at the cone $\mathcal{C} \subseteq \mathfrak{p}$. If the cone \mathcal{C} has no fixed points in M , then there exists a 1-dimensional P -equivariantly embedded submanifold $S \subseteq M$ s.t. on the connected component \mathcal{G}_0 of the identity, the projective representation $\bar{\rho}$ factors through the restriction map $r : \mathcal{G}_0 \rightarrow \Gamma_c(S; \text{Ad}(\mathcal{K}))$.

Thus, if there are no fixed points in M for \mathcal{C} , then the problem of classifying the projective unitary positive energy representations of $\mathcal{G} \rtimes P$ is essentially reduced to the one-dimensional case, which has been extensively studied, see for example [PS86, Was98, Tan11, Kac90a, KR87, GW84, TL99]. Moreover, if there are no one-dimensional P -equivariantly embedded submanifolds in M , one is essentially reduced to the case where $\bar{\rho}$ factors through the germs at the fixed point set $\Sigma \subseteq M$ of the cone $\mathcal{C} \subseteq \mathfrak{p}$. In the present paper, we address the setting where fixed points *do* exist and where $\bar{\rho}$ actually factors through the germs at a *single* fixed point.

Thus, let $a \in M$ be a fixed point of the P -action on M and let $V := T_a(M)$. If a smooth projective unitary representation $\bar{\rho}$ of \mathcal{G} factors through the germs at $a \in M$, then the continuity of $\bar{\rho}$ implies that it must further factor through the ∞ -jets $J_a^\infty(\text{Ad}(\mathcal{K})) \cong J_0^\infty(V; K) = G$ at $a \in M$, as is shown in Section A of the appendix. This brings us to groups of jets and motivates the study of smooth projective unitary representations of $G \rtimes_\alpha P$. Clearly, any smooth projective unitary representation of $G \rtimes_\alpha P$ defines one of $\mathcal{G} \rtimes P$ via the jet homomorphism $j_a^\infty : \mathcal{G} \rightarrow J_a^\infty(\text{Ad}(\mathcal{K})) \cong G$. In this way, the present paper contributes to the understanding of positive energy and KMS-representations of gauge groups.

In relation to the unitary representation theory of gauge groups, let us also mention the papers [GGV77, AHeK78, PS76] and [Is76], in which unitary representations of gauge groups $C_c^\infty(M; K)$ are constructed which are non-local in the sense that they do not factor through the restriction map $C_c^\infty(M; K) \rightarrow C_c^\infty(N; K)$ for some proper submanifold $N \subseteq M$. When $\dim(M) \geq 3$, these are irreducible ([Wal87] and [AHeKT81]). They are also considered in [ADGV16] and [AHeKM⁺93]. Unitary representations of groups of jets have also been considered in [GG68] and [AT94].

Structure of the paper

The paper is divided in two parts. In Part I, we introduce both the (generalized) positive energy and the KMS condition. We start in Section 2 by briefly recalling the relation between continuous projective unitary representations, central extensions and the second Lie algebra cohomology $H_{\text{ct}}^2(\mathfrak{g}, \mathbb{R})$. We move on to consider positive energy representations in Section 3 and discuss some of their properties. In Section 4 we relax the positive energy condition and introduce the so-called *generalized positive energy* condition. We show that for a projective unitary representation which is of generalized positive energy, its kernel is related to a particular quadratic form canonically associated to the corresponding class in $H_{\text{ct}}^2(\mathfrak{g}, \mathbb{R})$. This observation will play a key role in Part II. In Section 5, we briefly recall the modular theory of von Neumann algebras and then proceed to define KMS representations. We consider a number of examples and discuss some of their properties, in particular making the important observation that KMS representations give rise to generalized positive energy ones. To some extent, this unifies the positive energy and KMS conditions, allowing their simultaneous study. We remark also that Part I is formulated in the general context of possibly infinite-dimensional locally convex Lie groups, which is the appropriate context within the larger program that studies the projective unitary representations of gauge groups.

In Part II, we return to the Fréchet-Lie group $G := J_a^\infty(V; K)$ of ∞ -jets at $0 \in V$. After fixing our notation, we discuss in Section 7 a normal-form problem for the \mathfrak{p} -action on $\mathfrak{g} = J_a^\infty(V; \mathfrak{k})$. Using the general observations made in Part I together with the normal-form results obtained in Section 7, we proceed in Section 8 with the study of (generalized) positive energy representations of the Lie algebra $\mathfrak{g} \rtimes_D \mathfrak{p}$, where $D : \mathfrak{p} \rightarrow \text{der}(\mathfrak{g})$ denotes \mathfrak{p} -action on \mathfrak{g} corresponding to α .

Overview of main results

Let us describe the main results of Section 8. We will first need to introduce some notation. We write \mathcal{X}_I for the Lie algebra of formal vector fields on V vanishing at the origin. The \mathfrak{p} -action D splits into a horizontal and a vertical part according to $D(p) = -\mathcal{L}_{\mathbf{v}(p)} + \text{ad}_{\sigma(p)}$, for some Lie algebra homomorphism $\mathbf{v} : \mathfrak{p} \rightarrow \mathcal{X}_I^{\text{op}}$ and a linear map $\sigma : \mathfrak{p} \rightarrow \mathfrak{g}$ satisfying the Maurer-Cartan equation

$$-\mathcal{L}_{\mathbf{v}(p_1)}\sigma(p_2) + \mathcal{L}_{\mathbf{v}(p_2)}\sigma(p_1) - \sigma([p_1, p_2]) + [\sigma(p_1), \sigma(p_2)] = 0, \quad \forall p_1, p_2 \in \mathfrak{p}.$$

For any $p \in \mathfrak{p}$, the formal vector field $\mathbf{v}(p)$ splits further as $\mathbf{v}(p) = \mathbf{v}_1(p) + \mathbf{v}_{\text{ho}}(p)$, into its linearization $\mathbf{v}_1(p)$ and its higher order part $\mathbf{v}_{\text{ho}}(p)$, which is a formal vector field on V vanishing up to first order at the origin. We denote by $\mathbf{v}_1(p) = \mathbf{v}_1(p)_s + \mathbf{v}_1(p)_n$ the Jordan decomposition of $\mathbf{v}_1(p)$ over \mathbb{C} . Let $\sigma_0(p)$ be the constant part of the formal power series $\sigma(p)$. Let $\Sigma_p \subseteq \mathbb{C}$ denote the additive subsemigroup of \mathbb{C} generated by $\text{Spec}(\mathbf{v}_1(p))$. Let $V_c^{\mathbb{C}}(p)$ denote the span in $V_{\mathbb{C}}$ of all generalized eigenspaces of $\mathbf{v}_1(p)$ corresponding to eigenvalues with zero real part. Set $V_c(p) := V_c^{\mathbb{C}}(p) \cap V$. If $\mathfrak{C} \subseteq \mathfrak{p}$ is a subset, define $V_c(\mathfrak{C}) := \bigcap_{p \in \mathfrak{C}} V_c(p)$, which we call the ‘center subspace of V associated to \mathfrak{C} ’, in analogy with the center manifold of a fixed point of a dynamical system. Let $V_c(\mathfrak{C})^\perp \subseteq V^*$ be its annihilator in V^* . If $\bar{\pi}$ is a continuous projective unitary representation of $\mathfrak{g} \rtimes_D \mathfrak{p}$, let $\mathfrak{C}(\bar{\pi})$ be the set of all points $p \in \mathfrak{p}$ for which $\bar{\pi}$ is of generalized positive energy at p . Write $R := \mathbb{R}[[V^*]] := \prod_{n=0}^{\infty} P^n(V)$ for the ring of formal power series on V , where $P^n(V)$ denotes the set of degree- n homogeneous polynomials on V .

The first main result concerns positive energy representations. It states that unless the spectrum of $\mathbf{v}_1(p)$ and $\sigma_0(p)$ happens to intersect non-trivially, any smooth projective unitary representation $\bar{\rho}$ of $G \rtimes_\alpha P$ which is of positive energy at $p \in \mathfrak{p}$ factors through the 2-jets $J_0^2(V; K) \rtimes_\alpha P$:

Theorem 8.1. *Let $\bar{\rho}$ be a smooth projective unitary representation of $G \rtimes_{\alpha} P$ which is of p.e. at $p \in \mathfrak{p}$. Assume that $\text{Spec}(\text{ad}_{\sigma_0(p)}) \cap \text{Spec}(\mathfrak{v}_1(p)) = \emptyset$. Then $\bar{\rho}$ factors through $J_0^2(V; K) \rtimes_{\alpha} P$. Moreover the image of $-\mathcal{L}_{\mathfrak{v}_1(p)} + \text{ad}_{\sigma_0(p)}$ in $P^2(V) \otimes \mathfrak{k} \subseteq J_0^2(V; K)$ is contained in $\ker \bar{\rho}$.*

The second main result determines restrictions imposed by the generalized positive energy condition. If $p \in \mathcal{C}(\bar{\pi})$, then unless possibly when the “non-resonance condition” $\text{Spec}(\text{ad}_{\sigma_0(p)}) \cap \Sigma_p = \emptyset$ is violated, it suffices to consider the case where all eigenvalues of $\mathfrak{v}_1(p)$ are purely imaginary. The precise statement is:

Theorem 8.3. *Let $\bar{\pi}$ be a continuous projective unitary representation of $\mathfrak{g} \rtimes_D \mathfrak{p}$. Let $\mathcal{C} \subseteq \mathcal{C}(\bar{\pi})$. Assume that $\text{Spec}(\text{ad}_{\sigma_0(p)}) \cap \Sigma_p = \emptyset$ for all $p \in \mathcal{C}$. Then $RV_c(\mathcal{C})^{\perp} \otimes \mathfrak{k} \subseteq \ker \bar{\pi}$ and hence $\bar{\pi}|_{\mathfrak{g}}$ factors through $\mathbb{R}[[V_c(\mathcal{C})^*]] \otimes \mathfrak{k}$.*

Since $\mathbb{R}[[V_c(\mathcal{C})^*]] \otimes \mathfrak{k} = \mathfrak{k}$ whenever $V_c(\mathcal{C}) = \{0\}$, Theorem 8.3 in particular gives sufficient conditions for $\bar{\pi}$ to factor through \mathfrak{k} , that depend only on the spectrum of σ_0 and $\mathfrak{v}_1(p)$. For the third main result, we consider the special case where \mathfrak{p} is non-compact and simple. In order to obtain generalized positive energy representations $\bar{\pi}$ of $\mathfrak{g} \rtimes_D \mathfrak{p}$ other than those for which $\bar{\pi}|_{\mathfrak{g}}$ factors through \mathfrak{k} , one has to consider the situation where $\mathcal{C}(\bar{\pi})$ is a pointed P -invariant convex cone:

Theorem 8.6. *Assume that \mathfrak{p} is non-compact and simple. Suppose that \mathfrak{v}_1 defines a non-trivial irreducible \mathfrak{p} -representation on V . Let $\bar{\pi}$ be a continuous projective unitary representation of $\mathfrak{g} \rtimes_D \mathfrak{p}$. Either $\mathcal{C}(\bar{\pi})$ is a pointed P -invariant convex cone or $\bar{\pi}|_{\mathfrak{g}}$ factors through \mathfrak{k} .*

Remark 1.2. If $\bar{\rho}$ is a smooth projective unitary representation of $G \rtimes_{\alpha} P$ which is of generalized positive energy at the cone $\mathcal{C} \subseteq \mathfrak{p}$, then its derived representation $d\bar{\rho}$ on the space of smooth vectors $\mathcal{H}_{\rho}^{\infty}$ is so, too. Moreover, as we shall see in Lemma 6.3 below, the exponential map of $G = J_0^{\infty}(V; K)$ restricts to a diffeomorphism from the pro-nilpotent ideal $\ker(\text{ev}_0 : J_0^{\infty}(V; \mathfrak{k}) \rightarrow \mathfrak{k})$ onto $\ker(\text{ev}_0 : J_0^{\infty}(V; K) \rightarrow K)$. Thus, the above results all have immediate analogous consequences on the group level.

Acknowledgements

This research is supported by the NWO grant 639.032.734 “Cohomology and representation theory of infinite-dimensional Lie groups”. I am grateful to be given the opportunity to pursue a PhD in mathematics, which was given to me by my supervisor Bas Janssens, whom I would also like to thank for his guidance during the process. I am also grateful for helpful discussions with Karl-Hermann Neeb, Tobias Diez and Lukas Miaskiowski.

Part I

Positive Energy and KMS Representations

2 Projective Representations and Central Extensions

In the following, the category of infinite-dimensional manifolds and smooth maps between them is defined in the Michal-Bastiani sense [Bas64, Mil84, Nee06]. This also defines the notion of a locally convex Lie group. Throughout the section, let G denote a locally convex Lie group which is *regular* in the sense of [Nee06, Def. II.5.2]. Then G in particular admits an exponential map $\exp_G : \mathfrak{g} \rightarrow G$, see e.g. [Nee06, Rem. II.5.3].

Definition 2.1.

- A (projective) unitary representation of G is said to be continuous if it is so w.r.t. the strong operator topology on $U(\mathcal{H}_{\rho})$.
- Let $(\rho, \mathcal{H}_{\rho})$ be a unitary representation of G on \mathcal{H}_{ρ} . A vector $\psi \in \mathcal{H}_{\rho}$ is called *smooth* if the orbit map $G \rightarrow \mathcal{H}_{\rho}$, $g \mapsto \rho(g)\psi$ is smooth. Denote by $\mathcal{H}_{\rho}^{\infty} \subseteq \mathcal{H}_{\rho}$ the subspace of smooth vectors. The representation ρ is called smooth if $\mathcal{H}_{\rho}^{\infty}$ is dense in \mathcal{H}_{ρ} .
- Let $(\bar{\rho}, \mathcal{H}_{\rho})$ be a projective unitary representation of G on \mathcal{H}_{ρ} . A ray $[\psi] \in P(\mathcal{H}_{\rho})$ is called *smooth* if the orbit map $G \rightarrow P(\mathcal{H}_{\rho})$, $g \mapsto \bar{\rho}(g)[\psi]$ is smooth. Denote by $P(\mathcal{H}_{\rho})^{\infty}$ the subspace of smooth rays. The projective representation $\bar{\rho}$ is called smooth if $P(\mathcal{H}_{\rho})^{\infty}$ is dense in $P(\mathcal{H}_{\rho})$.

Definition 2.2. If \mathcal{D} is a complex vector space, denote by $\mathcal{L}(\mathcal{D})$ the Lie algebra of linear operators on \mathcal{D} . If \mathcal{D} is a pre-Hilbert space with Hilbert space closure \mathcal{H} , we also write

$$\mathcal{L}^{\dagger}(\mathcal{D}) := \{ X \in \mathcal{L}(\mathcal{D}) : \mathcal{D} \subseteq \text{dom}(X^*) \text{ and } X^*\mathcal{D} \subseteq \mathcal{D} \}.$$

For $X \in \mathcal{L}^\dagger(\mathcal{D})$ set $X^\dagger := X^*|_{\mathcal{D}}$. Then $(X^\dagger)^\dagger = X$ and $(-)^\dagger$ endows $\mathcal{L}^\dagger(\mathcal{D})$ with an involution [Sch90, Ch. 2]. If \mathcal{D} is a pre-Hilbert space, define

$$\mathbf{U}(\mathcal{D}) := \{ u \in \mathcal{L}^\dagger(\mathcal{D}) : uu^\dagger = u^\dagger u = \text{id}_{\mathcal{D}} \}, \quad \mathbf{u}(\mathcal{D}) := \{ X \in \mathcal{L}^\dagger(\mathcal{D}) : X^\dagger + X = 0 \}.$$

Definition 2.3. Let \mathcal{D} be a complex pre-Hilbert space.

- A *unitary representation* of the Lie algebra \mathfrak{g} on \mathcal{D} is a Lie algebra homomorphism $\pi : \mathfrak{g} \rightarrow \mathbf{u}(\mathcal{D})$. A *projective unitary representation* is a Lie algebra homomorphism $\bar{\pi} : \mathfrak{g} \rightarrow \mathbf{pu}(\mathcal{D}) := \mathbf{u}(\mathcal{D})/i\mathbb{R}I$.
- A unitary representation π of \mathfrak{g} is called *continuous* if $\xi \mapsto \pi(\xi)\psi$ is continuous for any $\psi \in \mathcal{D}$. A projective unitary representation $\bar{\pi}$ is continuous if $\xi \mapsto \pi(\xi)[\psi]$ is continuous for every $[\psi] \in \mathbf{P}(\mathcal{D})$.
- A *unitary representation of G with domain $\mathcal{D} \subseteq \mathcal{H}$* is a homomorphism $\rho : G \rightarrow \mathbf{U}(\mathcal{D})$. A projective unitary representation with domain \mathcal{D} is a homomorphism $G \rightarrow \mathbf{PU}(\mathcal{D})$, where $\mathbf{PU}(\mathcal{D}) := \mathbf{U}(\mathcal{D})/\mathbb{T}I$.

Remark 2.4. Any unitary G -representation on \mathcal{H}_ρ defines a unitary \mathfrak{g} -representation $d\rho : \mathfrak{g} \rightarrow \mathbf{u}(\mathcal{H}_\rho^\infty)$ on \mathcal{H}_ρ^∞ by $d\rho(\xi)\psi := \frac{d}{dt}|_{t=0} \rho(e^{t\xi})\psi$. We will always consider elements of $d\rho(\mathfrak{g})$ as unbounded operators defined on the invariant domain \mathcal{H}_ρ^∞ . Projective unitary G -representations similarly define projective unitary \mathfrak{g} -representations on $\mathbf{P}(\mathcal{H}_\rho^\infty)$ by differentiation at the identity. If G is finite-dimensional, then \mathcal{H}_ρ^∞ is dense in \mathcal{H}_ρ for any continuous unitary representation ρ of G , by a result of Gårding [War72, Prop. 4.4.1.1].

A continuous projective unitary representation $\bar{\rho} : G \rightarrow \mathbf{PU}(\mathcal{H}_\rho)$ is equivalently given by a continuous central \mathbb{T} -extension \widehat{G} together with a unitary representation $\rho : \widehat{G} \rightarrow \mathbf{U}(\mathcal{H}_\rho)$ which satisfies $\rho(z) = zI$ for z in the central \mathbb{T} component. Of course, \widehat{G} is the pull-back of the central \mathbb{T} -extension $\mathbf{U}(\mathcal{H}_\rho) \rightarrow \mathbf{PU}(\mathcal{H}_\rho)$ along $\bar{\rho}$. We say that ρ *lifts* $\bar{\rho}$. Suppose $\bar{\rho}_1$ and $\bar{\rho}_2$ are two projective unitary representations, inducing by pull-back the lifts $\rho_1 : \widehat{G}_1 \rightarrow \mathbf{U}(\mathcal{H}_{\rho_1})$ and $\rho_2 : \widehat{G}_2 \rightarrow \mathbf{U}(\mathcal{H}_{\rho_1})$ of $\bar{\rho}_1$ and $\bar{\rho}_2$, respectively. Then $\bar{\rho}_1$ and $\bar{\rho}_2$ are unitarily equivalent if and only if there is an isomorphism $\Phi : \widehat{G}_1 \rightarrow \widehat{G}_2$ of central G -extensions and a unitary $U : \mathcal{H}_{\rho_1} \rightarrow \mathcal{H}_{\rho_2}$ such that $\rho_2(\Phi(x)) = U\rho_1(x)U^{-1}$ for all $x \in \widehat{G}_1$. Analogously, any projective unitary \mathfrak{g} -representation $\bar{\pi}$ can be lifted to a unitary representation $\pi : \widehat{\mathfrak{g}} \rightarrow \mathbf{u}(\mathcal{H}_\rho)$ of some central \mathbb{R} -extension $\widehat{\mathfrak{g}}$ of \mathfrak{g} . The continuous central extensions of \mathfrak{g} by \mathbb{R} are up to isomorphism classified by $H_{\text{ct}}^2(\mathfrak{g}, \mathbb{R})$, the continuous second Lie algebra cohomology with trivial coefficients [JN19, Def. 6.2, Prop. 6.3]. Thus, to study the projective unitary representations of \mathfrak{g} up to equivalence, one may first determine $H_{\text{ct}}^2(\mathfrak{g}, \mathbb{R})$, choose for each class $[\omega] \in H_{\text{ct}}^2(\mathfrak{g}, \mathbb{R})$ a representative ω and then proceed to determine the equivalence classes of unitary representations π of $\mathbb{R} \oplus_\omega \mathfrak{g}$ satisfying $\pi(1, 0) = iI$. We will also write $\mathbf{c} := (1, 0) \in \mathbb{R} \oplus_\omega \mathfrak{g}$ for the central generator.

Remark 2.5. In the literature, one encounters the notion of the *level* of a unitary representation π of $\mathbb{R}\mathbf{c} \oplus_\omega \mathfrak{g}$, which is the number $l \in \mathbb{R}$ such that $\pi(\mathbf{c}) = iI$ (see e.g. [PS86, sec. 9.3]). Let us briefly clarify how such representations are included in the program outlined above, even though $\pi(\mathbf{c}) = iI$ is always assumed. Simply notice that such a representation of level l factors through the map $\mathbb{R}\mathbf{c} \oplus_\omega \mathfrak{g} \rightarrow \mathbb{R}\mathbf{c} \oplus_{l\cdot\omega} \mathfrak{g}$ induced by multiplication by l on the central factor. The corresponding representation π_2 of $\mathbb{R}\mathbf{c} \oplus_{l\cdot\omega} \mathfrak{g}$ satisfies $\pi_2(\mathbf{c}) = iI$. Notice that $\mathbb{R}\mathbf{c} \oplus_\omega \mathfrak{g} \rightarrow \mathbb{R}\mathbf{c} \oplus_{l\cdot\omega} \mathfrak{g}$ is an isomorphism of Lie algebras whenever $l \neq 0$, but *not* as central extensions unless $l = 1$, because a morphism of central extensions is required to be the identity on the central component. For $1 \neq l \in \mathbb{R}$, the cocycles ω and $l \cdot \omega$ are *not* equivalent unless $[\omega] = 0$ in $H_{\text{ct}}^2(\mathfrak{g}, \mathbb{R})$.

Remark 2.6. If a projective unitary representation $\bar{\rho}$ of G is smooth, then the corresponding central \mathbb{T} -extension \widehat{G} is again a locally convex Lie group [JN19, Thm. 4.3]. Moreover, there is a similar correspondence between smooth projective unitary representations $\bar{\rho}$ of G and their lifts $\rho : \widehat{G} \rightarrow \mathbf{U}(\mathcal{H}_\rho)$, which are then again smooth [JN19, cor. 4.5, Thm. 7.3]. Moreover, we have $\mathbf{P}(\mathcal{H}_\rho)^\infty = \mathbf{P}(\mathcal{H}_\rho^\infty)$ by [JN19, Thm. 4.3].

3 Positive Energy Representations

Let us introduce the class of positive energy representations. After defining the notion, some immediate consequences are considered that will be relevant in Part II. Let G be a regular locally convex Lie group with Lie algebra \mathfrak{g} . If $c \in \mathbb{R}$, \mathcal{D} is a pre-Hilbert space and $X \in \mathcal{L}^\dagger(\mathcal{D})$ satisfies $X^\dagger = X$, we write $X \geq c$ if $\langle \psi, X\psi \rangle \geq c$ for every $\psi \in \mathcal{D}$.

Definition 3.1. Let $\mathcal{C} \subseteq \mathfrak{g}$ be a convex cone and \mathcal{D} be a pre-Hilbert space.

- A continuous unitary representation π of \mathfrak{g} on \mathcal{D} is said to be of *positive energy* (p.e.) at $\xi \in \mathfrak{g}$ if $-i\pi(\xi) \geq 0$. It is of p.e. at \mathcal{C} if it is of p.e. at every $\xi \in \mathcal{C}$. Write $\mathcal{C}(\pi) = \{\xi \in \mathfrak{g} : \pi \text{ is of p.e. at } \xi\}$.
- Let $\bar{\pi}$ be a continuous projective unitary representation of \mathfrak{g} on \mathcal{D} with lift $\pi : \widehat{\mathfrak{g}} \rightarrow \mathfrak{u}(\mathcal{D})$. Then $\bar{\pi}$ is of p.e. at ξ if there exists $\widehat{\xi} \in \mathcal{C}(\pi) \subseteq \widehat{\mathfrak{g}}$ covering ξ . Write $\mathcal{D}(\bar{\pi})$ for the set of all such ξ . We say that $\bar{\pi}$ is of p.e. at \mathcal{C} if $\mathcal{C} \subseteq \mathcal{C}(\bar{\pi})$.
- A smooth (projective) unitary representation of G on \mathcal{H}_ρ is said to be of p.e. at $\xi \in \mathfrak{g}$ if the corresponding derived (projective) unitary representation of \mathfrak{g} on \mathcal{H}_ρ^∞ is so. It is said to be of p.e. at \mathcal{C} if it is so at every $\xi \in \mathcal{C}$.

Remark 3.2. Let ρ be a smooth unitary representation of G . Then $\mathcal{C} := \mathcal{C}(d\rho)$ is always a closed, G -invariant convex cone. Consequently, its edge $\mathcal{C} \cap -\mathcal{C}$ and span $\mathcal{C} - \mathcal{C}$ are ideals in \mathfrak{g} . Moreover, if $\xi \in \mathcal{C} \cap -\mathcal{C}$ then $\xi \in \ker d\rho$, so by passing to the quotient $\mathfrak{g}/\ker d\rho$ one may always achieve that \mathcal{C} is pointed.

Next, we define the notion of a semibounded representation.

Definition 3.3. Let $\rho : G \rightarrow \mathfrak{U}(\mathcal{H})$ be a smooth unitary G -representation. Define the *momentum set* by:

$$I_\rho := \text{conv} \{ \xi \mapsto \langle \psi, -id\rho(\xi)\psi \rangle : \psi \in \mathcal{H}_\rho^\infty, \|\psi\| = 1 \} \subseteq \mathfrak{g}^*.$$

The representation ρ is said to be *semibounded* if W_ρ contains an interior point, where

$$W_\rho := \left\{ \xi \in \mathfrak{g} : \inf \text{Spec}(-i\overline{d\rho(\xi)}) > -\infty \right\}.$$

Remark 3.4. For finite-dimensional Lie groups, the class of semibounded representations has been subject to detailed study in [Nee00]. In particular, they are highest weight representations [Nee00, Def. X.2.9, Thm. X.3.9]. For a consideration of semibounded representations in the context of infinite-dimensional Lie groups, we refer to [Nee17] and [Nee10b].

In the finite-dimensional context, the semiboundedness condition turns out to be extremely restrictive, which in turn has consequences for arbitrary positive energy representations. The following result, Theorem 3.5, is based on the results in the monograph [Nee00].

Theorem 3.5. *Assume that G is connected and locally exponential. Take $\mathbf{d} \in \mathfrak{g}$ and let $\mathfrak{a} = \langle \mathbf{d} \rangle \triangleleft \mathfrak{g}$ be the closed ideal in \mathfrak{g} generated by \mathbf{d} . Assume that $\dim(\mathfrak{a}) < \infty$ and that \mathfrak{a} is stable, in the sense that $\text{Ad}_G(\mathfrak{a}) \subseteq \mathfrak{a}$. Let $A \triangleleft G$ be a connected normal Lie subgroup integrating \mathfrak{a} . Let \mathfrak{a}_n denote the maximal nilpotent ideal of \mathfrak{a} . Let (ρ, \mathcal{H}_ρ) be a smooth unitary G -representation which is of p.e. at $\mathbf{d} \in \mathfrak{g}$. Write $\mathfrak{h} := \mathfrak{a}/\ker d\rho$. The following assertions hold:*

1. $\mathfrak{a} = \mathcal{C} - \mathcal{C}$, where $\mathcal{C} \subseteq \mathfrak{g}$ is the closed G -invariant convex cone in \mathfrak{g} generated by \mathbf{d} .
2. $\mathcal{C} + \ker d\rho$ is a pointed, closed, generating and G -invariant convex cone in \mathfrak{h} . Thus $\mathcal{C} \cap -\mathcal{C} \subseteq \ker d\rho$.
3. $\rho|_A$ is semibounded.
4. Let \mathfrak{h}_n denote the maximal nilpotent ideal in \mathfrak{h} . Then \mathfrak{h}_n is a generalized Heisenberg algebra, meaning that $[\mathfrak{h}_n, \mathfrak{h}_n] \subseteq \mathfrak{z}(\mathfrak{h})$. Moreover, there exists a reductive Lie algebra \mathfrak{l} such that $\mathfrak{h} \cong \mathfrak{h}_n \rtimes \mathfrak{l}$.
5. $[\mathfrak{a}, [\mathfrak{a}_n, \mathfrak{a}_n]] \subseteq \ker d\rho$.

Proof. For the first point, let \mathfrak{a}' be the closure of $\mathcal{C} - \mathcal{C}$ in \mathfrak{g} . As \mathfrak{a}' is a closed ideal in \mathfrak{g} containing \mathbf{d} , we have $\mathfrak{a} \subseteq \mathfrak{a}'$. On the other hand, we know that $\text{Ad}_G(\mathbf{d}) \subseteq \mathfrak{a}$ because \mathfrak{a} is stable. Thus $\mathcal{C} \subseteq \mathfrak{a}$ and hence $\mathfrak{a}' \subseteq \mathfrak{a}$. So $\mathfrak{a}' = \mathfrak{a}$. In particular $\dim(\mathfrak{a}') < \infty$ and so $\mathcal{C} - \mathcal{C} = \mathfrak{a}' = \mathfrak{a}$. Next we prove the second statement. Take $\xi \in \mathcal{C} \cap -\mathcal{C}$. Then $d\rho(\xi) \geq 0$ and $d\rho(\xi) \leq 0$, in view of Remark 3.2, and hence $\text{Spec}(\overline{d\rho(\xi)}) = \{0\}$. As $d\rho(\xi)$ is essentially skew-adjoint, it follows that $\xi \in \ker d\rho$. Thus $\mathcal{C} + \ker d\rho$ is pointed in \mathfrak{h} . As \mathcal{C} is closed, G -invariant and convex, it is clear that the same holds for the cone $\mathcal{C} + \ker d\rho$ in \mathfrak{h} . The latter is also generating in \mathfrak{h} because $\mathfrak{a} = \mathcal{C} - \mathcal{C}$. Next we show that $\rho|_A$ is semibounded. As \mathfrak{a} is spanned by \mathcal{C} and $\dim \mathfrak{a} < \infty$, it follows that $\mathcal{C} \subseteq \mathfrak{a}$ has interior points. As $\mathcal{C} \subseteq W_\rho$, this implies that W_ρ has interior points. Hence $\rho|_A$ is semibounded. For the remaining points, we use the results in [Nee17]. We first show that \mathfrak{h} is *admissible*, in the sense of [Nee00, Def. VII.3.2]. Using the second point, the convex cone $\mathcal{C} \oplus \mathbb{R}_{\geq 0}$ in $\mathfrak{h} \oplus \mathbb{R}$ is closed, pointed, generating and $\text{Inn}(\mathfrak{h})$ -invariant. By [Nee00, Lem. VII.3.1, Def. VII.3.2] this implies that \mathfrak{h} is admissible. By [Nee00, Thm. VII.3.10], it follows that $[\mathfrak{h}_n, \mathfrak{h}_n] \subseteq \mathfrak{z}(\mathfrak{h})$ and that \mathfrak{h} contains a compactly embedded Cartan subalgebra \mathfrak{t} (where as in [Nee00, Def. VII.1.1], a subalgebra $\mathfrak{t} \subseteq \mathfrak{h}$ is called *compactly embedded* if $\overline{\langle e^{\text{ad}(\mathfrak{t})} \rangle}$ is compact in $\text{Aut}(\mathfrak{h})$). Using [Nee00, Lem. VII.2.26(iv)], we obtain that there exists some reductive Lie algebra \mathfrak{l} with $\mathfrak{h} \cong \mathfrak{h}_n \rtimes \mathfrak{l}$. Since $[\mathfrak{h}, [\mathfrak{h}_n, \mathfrak{h}_n]] = 0$ and $\mathfrak{h} = \mathfrak{a}/\ker d\rho$, it follows in particular that $[\mathfrak{a}, [\mathfrak{a}_n, \mathfrak{a}_n]] \subseteq \ker d\rho$. \square

For projective p.e. representations, this leads to:

Corollary 3.6. *Let G , \mathfrak{d} , \mathfrak{a} and \mathfrak{a}_n be as Theorem 3.5. Let $(\bar{\rho}, \mathcal{H}_\rho)$ be a smooth projective unitary representation of G . Suppose that $\bar{\rho}$ is of p.e. at $\mathfrak{d} \in \mathfrak{g}$. Then $[\mathfrak{a}, [\mathfrak{a}_n, \mathfrak{a}_n]] \subseteq \ker d\bar{\rho}$.*

Proof. Let $\rho : \widehat{G} \rightarrow \mathrm{U}(\mathcal{H}_\rho)$ be the lift of $\bar{\rho}$ to a central \mathbb{T} -extension \widehat{G} of G . Let $\widehat{\mathfrak{g}} := \mathrm{Lie}(\widehat{G})$. There exists some $\widehat{\mathfrak{d}} \in \widehat{\mathfrak{g}}$ s.t. $d\rho$ is of p.e. at $\widehat{\mathfrak{d}} \in \widehat{\mathfrak{g}}$. Let $\widehat{\mathfrak{a}}$ denote the ideal in $\widehat{\mathfrak{g}}$ generated by $\widehat{\mathfrak{d}}$ and let $\widehat{\mathfrak{a}}_n$ denote the maximal nilpotent ideal in $\widehat{\mathfrak{a}}$. Then $d\rho([\widehat{\mathfrak{a}}, [\widehat{\mathfrak{a}}_n, \widehat{\mathfrak{a}}_n]]) \subseteq \{0\}$ by Theorem 3.5. Thus $d\bar{\rho}([\mathfrak{a}, [\mathfrak{a}_n, \mathfrak{a}_n]]) = \{0\}$, where we used that the quotient map $\widehat{\mathfrak{g}} \rightarrow \mathfrak{g}$ projects $\widehat{\mathfrak{a}}$ and $\widehat{\mathfrak{a}}_n$ onto \mathfrak{a} and \mathfrak{a}_n , respectively. \square

The following simple lemma will also be useful.

Lemma 3.7. *Let $\bar{\rho} : G \rightarrow \mathrm{PU}(\mathcal{H}_\rho)$ be a continuous projective unitary representation of G which is of p.e. at the closed G -invariant convex cone \mathcal{C} . Let $\mathfrak{e} := \mathcal{C} \cap -\mathcal{C}$ and assume that $\dim(\mathfrak{e}) < \infty$. Then $\bar{\rho}|_{\mathfrak{e}}$ is continuous w.r.t. the norm-topology on $\mathrm{U}(\mathcal{H}_\rho)$.*

Proof. By restricting to \mathfrak{e} , we may assume that $\mathcal{C} = \mathfrak{g}$ and $\dim(\mathfrak{g}) < \infty$. Let $d\rho : \widehat{\mathfrak{g}} \rightarrow \mathfrak{u}(\mathcal{H}_\rho^\infty)$ be the lift of $d\bar{\rho}$. Identify $\widehat{\mathfrak{g}} \cong \mathbb{R} \oplus_\omega \mathfrak{g}$ for some 2-cocycle $\omega : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$. The assumptions imply that for every $\xi \in \mathfrak{g}$ there exists $E_\xi \in \mathbb{R}$ s.t. $-id\rho(\xi) \geq E_\xi$. As this holds in particular for both ξ and $-\xi$, $d\rho(\xi)$ is a bounded operator for any $\xi \in \mathfrak{g}$. As $\dim(\mathfrak{g}) < \infty$, one finds by choosing a basis (e_μ) of \mathfrak{g} that there exists $C > 0$ s.t. $\|d\rho(\xi)\| \leq C\|\xi\|$ where $\|\xi\| := \sup_\mu |\xi^\mu|$ if $\xi = \sum_\mu \xi^\mu e_\mu$. Thus $\xi \mapsto d\rho(\xi)$ is norm-continuous. This implies norm-continuity of $\bar{\rho}$ because $\mathcal{B}(\mathcal{H}_\rho) \rightarrow \mathcal{B}(\mathcal{H}_\rho)$, $T \mapsto e^T$ is norm-continuous and $\bar{\rho}(\exp(\xi)) = [e^{d\rho(\xi)}] \in \mathrm{PU}(\mathcal{H}_\rho)$ for $\xi \in \mathfrak{g}$. \square

4 Generalized Positive Energy Representations

Let G denote a regular locally convex Lie group with Lie algebra \mathfrak{g} . The class of p.e. representations can be generalized by relaxing the condition $-id\rho(\xi) \geq 0$ in Definition 3.1. We define a suitable relaxed notion, the *generalized positive energy condition*, and show that it can still be very restrictive. In Section 5, we will encounter a class of representations which are not of p.e. but are of generalized positive energy.

Definition 4.1. Let \mathcal{D} be a pre-Hilbert space with Hilbert space closure \mathcal{H} .

- A continuous unitary representation $\pi : \mathfrak{g} \rightarrow \mathfrak{u}(\mathcal{D})$ is of *generalized positive energy* (g.p.e.) at $\xi \in \mathfrak{g}$ if

$$\forall \psi \in \mathcal{D} : E_\psi(\pi, \xi) := \inf \{ \langle \psi, -i\pi(\mathrm{Ad}_g(\xi))\psi \rangle : g \in G \} > -\infty. \quad (1)$$

Write $\mathfrak{C}(\pi) := \{ \xi \in \mathfrak{g} : \pi \text{ is of g.p.e. at } \xi \}$. If $\mathfrak{C} \subseteq \mathfrak{g}$, we say that π is of g.p.e. at \mathfrak{C} if $\mathfrak{C} \subseteq \mathfrak{C}(\pi)$.

- Let $\bar{\pi}$ be a continuous projective unitary representation of \mathfrak{g} on \mathcal{D} with lift $\pi : \widehat{\mathfrak{g}} \rightarrow \mathfrak{u}(\mathcal{D})$. Let $\mathfrak{C}(\bar{\pi}) \subseteq \mathfrak{g}$ denote the image of $\mathfrak{C}(\pi) \subseteq \widehat{\mathfrak{g}}$ under the quotient map $\widehat{\mathfrak{g}} \rightarrow \mathfrak{g}$. Then $\bar{\pi}$ is said to be of generalized positive energy at $\xi \in \mathfrak{g}$ if $\xi \in \mathfrak{C}(\bar{\pi})$. Similarly, we say it is of g.p.e. at $\mathfrak{C} \subseteq \mathfrak{g}$ if $\mathfrak{C} \subseteq \mathfrak{C}(\bar{\pi})$.
- Let $\rho : G \rightarrow \mathrm{U}(\mathcal{D})$ be a smooth unitary representation of G with domain $\mathcal{D} \subseteq \mathcal{H}_\rho^\infty$ consisting of smooth vectors. Then ρ is said to be of g.p.e. at $\xi \in \mathfrak{g}$ if $E_\psi(d\rho, \xi) < \infty$ for all $\psi \in \mathcal{D} \subseteq \mathcal{H}_\rho^\infty$.
- Let $\bar{\rho} : G \rightarrow \mathrm{PU}(\mathcal{D})$ be a smooth projective unitary representation of G with domain $\mathrm{P}(\mathcal{D}) \subseteq \mathrm{P}(\mathcal{H}_\rho)^\infty$ consisting of smooth rays and with lift $\rho : \widehat{G} \rightarrow \mathrm{U}(\mathcal{D})$. Then $\bar{\rho}$ is of g.p.e. at $\xi \in \mathfrak{g}$ if there exists $\widehat{\xi} \in \mathfrak{C}(\rho)$ covering ξ .

Remark 4.2. If π is a (projective) continuous unitary representation of \mathfrak{g} , then the set $\mathfrak{C}(\pi) \subseteq \mathfrak{g}$ is always an Ad_G -invariant convex cone.

An important observation for the class of g.p.e. representations is the following one:

Lemma 4.3. *Let $\pi : \mathfrak{g} \rightarrow \mathfrak{u}(\mathcal{D})$ be a continuous unitary representation of \mathfrak{g} on the pre-Hilbert space \mathcal{D} . Let $\xi \in \mathfrak{C}(\pi)$. Suppose that $\eta \in \mathfrak{g}$ satisfies $[[\xi, \eta], \eta] \in \mathfrak{Z}(\mathfrak{g})$. Then for every $\psi \in \mathcal{D}$ we have:*

$$0 \leq \langle \psi, -i\pi([[\xi, \eta], \eta])\psi \rangle, \quad (2)$$

$$\langle \psi, -i\pi([\xi, \eta])\psi \rangle^2 \leq 2\langle \psi, -i\pi([\xi, \eta], \eta)\psi \rangle \left(\langle \psi, -i\pi(\xi)\psi \rangle - E_\psi(\pi, \xi) \right).$$

In particular, if $[[\xi, \eta], \eta] = 0$ then $\pi([\xi, \eta]) = 0$.

Proof. Let $\psi \in \mathcal{D}$. Then in particular $\langle \psi, -i\pi(e^{t\text{ad}_\eta}\xi)\psi \rangle \geq E_\psi(\pi, \xi)$ for all $t \in \mathbb{R}$. As $[[\xi, \eta], \eta] \in \mathfrak{Z}(\mathfrak{g})$, the third derivative $\gamma^{(3)} : \mathbb{R} \rightarrow \mathfrak{g}$ of the smooth path $\gamma : \mathbb{R} \rightarrow \mathfrak{g}$, $t \mapsto e^{t\text{ad}_\eta}\xi$ vanishes. From Taylor's formula (which holds for smooth maps between locally convex vector spaces by [Nee06, Prop. I.2.3]), it follows that $e^{t\text{ad}_\eta}\xi = \xi + t[\eta, \xi] + \frac{t^2}{2}[[\xi, \eta], \eta]$ for all $t \in \mathbb{R}$. Thus

$$\langle \psi, -i\pi(\xi)\psi \rangle + t\langle \psi, -i\pi([\eta, \xi])\psi \rangle + \frac{t^2}{2}\langle \psi, -i\pi([[\xi, \eta], \eta])\psi \rangle \geq E_\psi(\pi, \xi), \quad \forall t \in \mathbb{R}$$

The equations (2) follows from the fact that $at^2 + bt + c \geq 0$ for all $t \in \mathbb{R}$ if and only if $a, c \geq 0$ and $b^2 \leq 4ac$. In particular, if $[[\xi, \eta], \eta] = 0$ then $\langle \psi, -i\pi([\xi, \eta])\psi \rangle = 0$ for all $\psi \in \mathcal{D}$. As \mathcal{D} is a complex vector space, this implies by the polarization identity that $\pi([\xi, \eta]) = 0$. \square

In the projective context, this sets up a relation between $\ker \bar{\pi}$ and the class $[\omega] \in H_{\text{ct}}^2(\mathfrak{g}; \mathbb{R})$ defined by the corresponding central \mathbb{R} -extension $\widehat{\mathfrak{g}}$ of \mathfrak{g} . This is exploited in Section 8.

Proposition 4.4. *Let $\bar{\pi}$ be a continuous projective unitary \mathfrak{g} -representation on the pre-Hilbert space \mathcal{D} with lift $\pi : \widehat{\mathfrak{g}} \rightarrow \mathfrak{u}(\mathcal{D})$ for some continuous central \mathbb{R} -extension $\widehat{\mathfrak{g}}$ of \mathfrak{g} . Let ω represent the corresponding class in $H_{\text{ct}}^2(\mathfrak{g}, \mathbb{R})$. Let $\xi \in \mathfrak{C}(\bar{\pi})$. Suppose that $\eta \in \mathfrak{g}$ satisfies $[[\xi, \eta], \eta] = 0$. Then $\omega([\xi, \eta], \eta) \geq 0$ and*

$$\omega([\xi, \eta], \eta) = 0 \iff \bar{\pi}([\xi, \eta]) = 0.$$

Proof. Identify $\widehat{\mathfrak{g}}$ with $\mathbb{R} \oplus_\omega \mathfrak{g}$. Let $\widehat{\xi} \in \mathfrak{C}(\pi)$ and $\widehat{\eta} \in \widehat{\mathfrak{g}}$ be lifts of ξ and η , respectively. We have that $[[\widehat{\xi}, \widehat{\eta}], \widehat{\eta}] = \omega([\xi, \eta], \eta) \in \mathfrak{Z}(\widehat{\mathfrak{g}})$, because $[[\xi, \eta], \eta] = 0$. Using Lemma 4.3 it follows that $\omega([\xi, \eta], \eta) \geq 0$. If $\omega([\xi, \eta], \eta) = 0$, then $[[\widehat{\xi}, \widehat{\eta}], \widehat{\eta}] = 0$ and so Lemma 4.3 implies that $\pi([\widehat{\xi}, \widehat{\eta}]) = 0$. Hence $\bar{\pi}([\xi, \eta]) = 0$. Conversely, if $\bar{\pi}([\xi, \eta]) = 0$, then $i\omega([\xi, \eta], \eta) = [\pi([\xi, \eta]), \pi(\eta)] - \pi([\xi, \eta], \eta) = 0$, because $[[\xi, \eta], \eta] = 0$. \square

Remark 4.5. Notice in the setting of Proposition 4.4 that whenever $[[\xi, \eta], \eta] = 0$, the value of $\omega([\xi, \eta], \eta)$ does not depend on the choice of representative ω of the class $[\omega] \in H_{\text{ct}}^2(\mathfrak{g}, \mathbb{R})$.

In Part II, a particular special case of Proposition 4.4 is used extensively:

Corollary 4.6. *Let \mathfrak{p} and \mathfrak{g} be locally convex Lie algebras. Let $D : \mathfrak{p} \rightarrow \text{der}(\mathfrak{g})$ be a homomorphism for which the corresponding action $\mathfrak{p} \times \mathfrak{g} \rightarrow \mathfrak{g}$ is continuous. Let \mathcal{D} be a complex pre-Hilbert space and let $\bar{\pi} : \mathfrak{g} \rtimes_D \mathfrak{p} \rightarrow \mathfrak{pu}(\mathcal{D})$ be a continuous projective unitary representation of $\mathfrak{g} \rtimes_D \mathfrak{p}$ on \mathcal{D} . Let $[\omega] \in H_{\text{ct}}^2(\mathfrak{g} \rtimes_D \mathfrak{p}; \mathbb{R})$ be the corresponding class in $H_{\text{ct}}^2(\mathfrak{g} \rtimes_D \mathfrak{p}; \mathbb{R})$. Let $\eta \in \mathfrak{g}$, $p \in \mathfrak{C}(\bar{\pi}) \cap \mathfrak{p}$ and assume that $[D(p)\eta, \eta] = 0$. Then $\omega(D(p)\eta, \eta) \geq 0$ and $\omega(D(p)\eta, \eta) = 0 \iff \bar{\pi}(D(p)\eta) = 0$.*

5 KMS Representations

In the following, we introduce the class of KMS representations. We will see in particular that these give rise to generalized positive energy representations. Consequently, they can be studied using the results of Section 4. Its definition makes use of the modular theory of von Neumann algebras, which we recall first.

5.1 Modular Theory of von Neumann algebras

Let us recall the modular condition and the notion of a KMS state on a von Neumann algebra \mathcal{M} , whilst fixing our conventions and notation. We refer to [Tak03a, Ch. VIII], [BR87, Ch. 2.5] and [BR97, Ch. 5.3] for a detailed consideration of the modular theory of von Neumann algebras and of KMS states.

If \mathcal{M} is a von Neumann algebra, write \mathcal{M}_* for its pre-dual, equipped with the $\sigma(\mathcal{M}_*, \mathcal{M})$ -topology. Write $\mathcal{S}(\mathcal{M}) \subseteq \mathcal{M}_*$ for the set of normal states on \mathcal{M} . Further, if $\phi \in \mathcal{S}(\mathcal{M})$, write $\pi_\phi : \mathcal{M} \rightarrow \mathcal{B}(\mathcal{H}_\phi)$ for the GNS-representation of \mathcal{M} relative to ϕ . Write $\mathcal{M}_\phi := \pi_\phi(\mathcal{M})''$. Let $\Omega_\phi \in \mathcal{H}_\phi$ denote the canonical cyclic vector satisfying $\phi(x) = \langle \Omega_\phi, \pi_\phi(x)\Omega_\phi \rangle$ for all $x \in \mathcal{M}$. Whenever Ω_ϕ is separating for \mathcal{M}_ϕ , let S_ϕ denote the unique closed conjugate-linear operator satisfying $S_\phi x \Omega_\phi = x^* \Omega_\phi$ for all $x \in \mathcal{M}_\phi$. Let Δ_ϕ and J_ϕ be the unique self-adjoint and anti-unitary operators, respectively, such that $S_\phi = J_\phi \Delta_\phi^{\frac{1}{2}}$.

Definition 5.1. A map $\sigma : \mathbb{R} \rightarrow \text{Aut}(\mathcal{M})$ is said to be $\sigma(\mathcal{M}_*, \mathcal{M})$ -continuous if for every $x \in \mathcal{M}$, the map $\mathbb{R} \rightarrow \mathcal{M}$, $t \mapsto \sigma_t(x)$ is continuous w.r.t. the $\sigma(\mathcal{M}_*, \mathcal{M})$ -topology on \mathcal{M} .

Definition 5.2. Let $\phi \in \mathcal{S}(\mathcal{M})$ be a normal state. Let $\sigma : \mathbb{R} \rightarrow \text{Aut}(\mathcal{M})$ be a one-parameter group of automorphisms of \mathcal{M} . Define $\text{St} := \{z : z \in \mathbb{C}, 0 < \text{Im}(z) < 1\}$.

— ϕ is said to satisfy the *modular condition* for σ if the following two conditions are satisfied:

1. $\phi = \phi \circ \sigma_t$ for all $t \in \mathbb{R}$.
2. For every $x, y \in \mathcal{M}$, there exists a bounded continuous function $F_{x,y} : \overline{\text{St}} \rightarrow \mathbb{C}$ which is holomorphic on St and s.t. for every $t \in \mathbb{R}$:

$$\begin{aligned} F_{x,y}(t) &= \phi(\sigma_t(x)y), \\ F_{x,y}(t+i) &= \phi(y\sigma_t(x)). \end{aligned}$$

— ϕ is said to be *KMS* w.r.t. σ at inverse temperature $\beta > 0$ if it satisfies the modular condition for $t \mapsto \sigma_{-\beta t}$. In that case, we also say that ϕ is σ -KMS at inverse-temperature β . If $\beta = 1$ we simply say that ϕ is a σ -KMS state.

Remark 5.3.

1. Suppose that $\phi \in \mathcal{S}(\mathcal{M})$ is faithful. Then there exists a unique automorphism group $\sigma^\phi : \mathbb{R} \rightarrow \text{Aut}(\mathcal{M})$ for which ϕ satisfies the modular condition [Tak03a, Thm. VIII.1.2], [BR87, Thm. 2.5.14]. The automorphism group σ^ϕ is $\sigma(\mathcal{M}_*, \mathcal{M})$ -continuous. As ϕ is faithful, $\pi_\phi : \mathcal{M} \rightarrow \mathcal{M}_\phi$ is injective and hence a $*$ -isomorphism between \mathcal{M} and \mathcal{M}_ϕ [BR87, Thm. 2.4.24]. Thus one may identify \mathcal{M} with \mathcal{M}_ϕ via $\pi_\phi : \mathcal{M} \rightarrow \mathcal{M}_\phi \subseteq \mathcal{B}(\mathcal{H}_\phi)$. Finally, there is a unique conditional expectation $\mathcal{E} : \mathcal{M} \rightarrow \mathcal{M}^{\mathbb{R}}$ s.t. $\phi = \phi_0 \circ \mathcal{E}$, where $\mathcal{M}^{\mathbb{R}} := \{x \in \mathcal{M} : \sigma_t^\phi(x) = x \quad \forall t \in \mathbb{R}\}$ and $\phi_0 := \phi|_{\mathcal{M}^{\mathbb{R}}}$ [Tak03a, Thm. IX.4.2].
2. If ϕ is not necessarily faithful, then ϕ satisfies the modular condition for some $\sigma(\mathcal{M}_*, \mathcal{M})$ -continuous 1-parameter group $\sigma : \mathbb{R} \rightarrow \text{Aut}(\mathcal{M})$ of $*$ -automorphisms of \mathcal{M} if and only if $\Omega_\phi \in \mathcal{H}_\phi$ is separating for $\mathcal{M}_\phi = \pi_\phi(\mathcal{M})'' \subseteq \mathcal{B}(\mathcal{H}_\phi)$. In that case, there is a central projection $p \in \mathcal{Z}(\mathcal{M})$ such that $\phi(1-p) = 0$ and ϕ is faithful on $\mathcal{M}p$. Moreover, $\sigma_t(p) = p$ for all $t \in \mathbb{R}$ and $\sigma|_{\mathcal{M}p}$ is uniquely determined by the modular condition for ϕ [BR97, Thm. 5.3.10].
3. In particular, if \mathcal{M} is a factor and ϕ is KMS w.r.t. $\sigma : \mathbb{R} \rightarrow \text{Aut}(\mathcal{M})$, then necessarily $p = I$ and whence ϕ must be faithful. Consequently $\sigma = \sigma_t^\phi$ is necessary.
4. In the converse direction, given a $\sigma(\mathcal{M}_*, \mathcal{M})$ -continuous automorphism group $\sigma : \mathbb{R} \rightarrow \text{Aut}(\mathcal{M})$, there may be no, precisely one, or multiple states in $\mathcal{S}(\mathcal{M})$ that are KMS w.r.t. σ . The set of σ -KMS states in $\mathcal{S}(\mathcal{M})$ is considered in [BR97, Ch. 5.3.2]. In particular, if $\phi \in \mathcal{S}(\mathcal{M})$ is a faithful σ -KMS state and $\psi \in \mathcal{S}(\mathcal{M})$, then ψ is σ -KMS if and only if there is a (necessarily unique) positive operator T affiliated to $\mathcal{Z}(\mathcal{M})$ such that $\psi(x) = \phi(T^{\frac{1}{2}}xT^{\frac{1}{2}})$ for all $x \in \mathcal{M}$ [BR97, Prop. 5.3.29]. In [BEK80] and [BEK86], the set K_β of normal σ -KMS states at inverse temperature β is studied in the setting of C^* -dynamical systems.
5. As a consequence of the previous points, if \mathcal{M} is a factor and $\phi, \psi \in \mathcal{S}(\mathcal{M})$ are both σ -KMS, then $\phi = \psi$, so that two distinct normal states can not share the same modular automorphism group.

Remark 5.4. Suppose $\phi \in \mathcal{S}(\mathcal{M})$ is KMS w.r.t. $\sigma : \mathbb{R} \rightarrow \text{Aut}(\mathcal{M})$. Let $\sigma^\phi : \mathbb{R} \rightarrow \text{Aut}(\mathcal{M}_\phi)$ denote the modular automorphism group defined by the faithful state $\langle \Omega_\phi, \cdot \Omega_\phi \rangle$ on $\mathcal{M}_\phi = \pi_\phi(\mathcal{M})''$. It then holds true that $\sigma_t^\phi(\pi_\phi(x)) = \pi_\phi(\sigma_{-t}(x))$ for any $x \in \mathcal{M}$ and $t \in \mathbb{R}$. Indeed, by [BR97, cor. 5.3.4], the state $\langle \Omega_\phi, \cdot \Omega_\phi \rangle$ on \mathcal{M}_ϕ is KMS w.r.t. the unique automorphism group $\tau : \mathbb{R} \rightarrow \text{Aut}(\mathcal{M}_\phi)$ satisfying $\tau_t(\pi_\phi(x))\Omega_\phi = \pi_\phi(\sigma_t(x))\Omega_\phi$ for all $t \in \mathbb{R}$. Then $\sigma_t^\phi = \tau_{-t}$ by uniqueness of the modular automorphism group (and the minus sign in the definition of KMS states). As Ω_ϕ is separating for \mathcal{M}_ϕ , it follows that $\sigma_t^\phi(\pi_\phi(x)) = \pi_\phi(\sigma_{-t}(x))$.

Example 5.5 (Gibbs States). Let $\mathcal{M} = \mathcal{B}(\mathcal{H})$ and $\sigma_t(x) = e^{itH}xe^{-itH}$ for some self-adjoint operator H satisfying $Z_\beta := \text{Tr}(e^{-\beta H}) < \infty$ for some $\beta > 0$. Consider the normal state $\phi(x) = \frac{1}{Z_\beta} \text{Tr}(e^{-\beta H}x)$ on \mathcal{M} . Consider \mathcal{M} as a dense subspace of \mathcal{H}_ϕ via $x \mapsto \pi_\phi(x)\Omega_\phi$. The modular operators J_ϕ and Δ_ϕ satisfy:

$$\begin{aligned} J_\phi \pi_\phi(x)\Omega_\phi &= e^{-\beta \frac{H}{2}} \pi_\phi(x^*) e^{\beta \frac{H}{2}} \Omega_\phi, \\ \Delta_\phi \pi_\phi(x)\Omega_\phi &= e^{-\beta H} \pi_\phi(x) e^{\beta H} \Omega_\phi, \end{aligned}$$

for $x \in \mathcal{M}$. Moreover, $\sigma_t^\phi(x) = e^{-i\beta t H} x e^{i\beta t H} = \sigma_{-\beta t}(x)$. Thus ϕ satisfies the modular condition for $\sigma_{-\beta t}$ and is therefore KMS at inverse-temperature β w.r.t. σ_t .

Gibbs states $\phi(x) = \frac{1}{Z_\beta} \text{Tr}(e^{-\beta H}x)$ constitute the simplest class of examples of KMS states. We will encounter a variety of different KMS states in Section 5.2.2 below.

5.2 KMS representations

In the following, let G be a regular locally convex Lie group with Lie algebra \mathfrak{g} . Let $N \subseteq G$ be an embedded Lie subgroup.

Definition 5.6. Let (ρ, \mathcal{H}_ρ) be a unitary G -representation. Let $\mathcal{N} := \rho(N)'' \subseteq \mathcal{B}(\mathcal{H}_\rho)$ be the von Neumann algebra generated by $\rho(N)$. For $\phi \in \mathcal{N}_*$, define the function $\widehat{\phi} : N \rightarrow \mathbb{C}$ by $\widehat{\phi}(n) := \phi(\rho(n))$. Write $\mathcal{N}_*^\infty := \left\{ \phi \in \mathcal{N}_* : \widehat{\phi} \in C^\infty(N; \mathbb{C}) \right\}$ and set $\mathcal{S}(\mathcal{N})^\infty := \mathcal{S}(\mathcal{N}) \cap \mathcal{N}_*^\infty$.

- Let $\xi \in \mathfrak{g}$ and $\phi \in \mathcal{S}(\mathcal{N})$. We say that ϕ is *KMS-compatible* with (ρ, ξ, N) if $e^{t\xi} N e^{-t\xi} \subseteq N$ for all $t \in \mathbb{R}$ and ϕ is KMS w.r.t. the automorphism group $\mathbb{R} \rightarrow \text{Aut}(\mathcal{N})$ defined by $t \mapsto \text{Ad}(\rho(e^{t\xi}))$.
- Define $\text{KMS}(\rho, \xi, N) := \{ \phi \in \mathcal{S}(\mathcal{N}) : \phi \text{ is KMS-compatible with } (\rho, \xi, N) \}$. Similarly, let $\text{KMS}(\rho, \xi, N)^\infty := \text{KMS}(\rho, \xi, N) \cap \mathcal{S}(\mathcal{N})^\infty$.
- ρ is said to be *KMS at $\xi \in \mathfrak{g}$ relative to N* if $\text{KMS}(\rho, \xi, N) \neq \emptyset$. It is called *smoothly-KMS at ξ relative to N* if $\text{KMS}(\rho, \xi, N)^\infty \neq \emptyset$.

If the subgroup N is clear from the context, we drop N from the notation and simply write $\text{KMS}(\rho, \xi)$ and $\text{KMS}(\rho, \xi)^\infty$. We then also say that ρ is KMS at ξ if it is so relative to N .

Remark 5.7. For any fixed $\xi \in \mathfrak{g}$ satisfying $\text{Ad}(e^{t\xi})N \subseteq N$ for all $t \in \mathbb{R}$, one may as well consider the semidirect product $N \rtimes_\alpha \mathbb{R}$, where $\alpha : \mathbb{R} \rightarrow \text{Aut}(N)$ is defined by $\alpha_t := \text{Ad}(e^{t\xi})|_N$. Definition 5.6 additionally allows for the situation where ρ is KMS at multiple $\xi_I \in \mathfrak{g}$, relative to possibly distinct subgroups $N_I \subseteq G$, where $I \in \mathcal{I}$ for some indexing set \mathcal{I} . We will see an example of this in Example 5.20 below.

In the following, let (ρ, \mathcal{H}_ρ) be a unitary G -representation and $\mathcal{N} := \rho(N)'' \subseteq \mathcal{B}(\mathcal{H}_\rho)$ be the von Neumann algebra generated by $\rho(N)$. If $\phi \in \mathcal{S}(\mathcal{N})$, write $\pi_\phi : \mathcal{N} \rightarrow \mathcal{B}(\mathcal{H}_\phi)$ for the GNS-representation of \mathcal{N} relative to ϕ . Let $\Omega_\phi \in \mathcal{H}_\phi$ denote the canonical \mathcal{N} -cyclic vector satisfying $\phi(x) = \langle \Omega_\phi, \pi_\phi(x)\Omega_\phi \rangle$ for all $x \in \mathcal{N}$. Write $\rho_\phi := \pi_\phi \circ \rho : N \rightarrow \text{U}(\mathcal{H}_\phi)$ for the unitary N -representation on \mathcal{H}_ϕ . Define $\mathcal{N}_\phi := \rho_\phi(N)'' \subseteq \mathcal{B}(\mathcal{H}_\phi)$.

Lemma 5.8. *Let $\phi \in \mathcal{S}(\mathcal{N})$. Then $\widehat{\phi}$ is smooth $\iff \Omega_\phi \in \mathcal{H}_{\rho_\phi}^\infty$. In this case, $\mathcal{H}_{\rho_\phi}^\infty$ is dense, so ρ_ϕ is smooth.*

Proof. Assume that $\widehat{\phi}$ is smooth. Then $n \mapsto \langle \Omega_\phi, \rho_\phi(n)\Omega_\phi \rangle$ is smooth. By [Nee10a, Thm. 7.2], it follows $n \mapsto \rho_\phi(n)\Omega_\phi$ is smooth $N \rightarrow \mathcal{H}_\phi$. The converse direction is trivial. Assume that $\Omega_\phi \in \mathcal{H}_{\rho_\phi}^\infty$. As $\mathcal{H}_{\rho_\phi}^\infty$ is N -invariant and Ω_ϕ is cyclic for N , it follows that $\mathcal{H}_{\rho_\phi}^\infty$ is dense in \mathcal{H}_ϕ . \square

Consider the left action of G on $\mathcal{S}(\mathcal{N})$ defined by $(g.\phi)(x) := \phi(\rho(g)^{-1}x\rho(g))$ for $x \in \mathcal{N}$. Notice that this action leaves $\mathcal{S}(\mathcal{N})^\infty$ invariant.

Lemma 5.9. *Let $g \in G$ and $\xi \in \mathfrak{g}$. Then $\phi \in \text{KMS}(\rho, \xi, N) \iff g.\phi \in \text{KMS}(\rho, \text{Ad}_g(\xi), gNg^{-1})$.*

Proof. Write $\mathcal{N}_g := \rho(g)\mathcal{N}\rho(g)^{-1}$. Let $\phi \in \text{KMS}(\rho, \xi, N)$. As $e^{t\xi} N e^{-t\xi} \subseteq N$ it follows that $e^{t\text{Ad}_g(\xi)}$ normalizes gNg^{-1} for every $t \in \mathbb{R}$. Define the following automorphism groups:

$$\begin{aligned} \sigma^\xi : \mathbb{R} &\rightarrow \text{Aut}(\mathcal{N}), & \sigma^\xi &:= \text{Ad}(\rho(e^{t\xi})), \\ \eta^\xi : \mathbb{R} &\rightarrow \text{Aut}(\mathcal{N}_g), & \eta^\xi &:= \text{Ad}(\rho(e^{t\text{Ad}_g\xi})). \end{aligned}$$

In order to show $g.\phi \in \text{KMS}(\rho, \text{Ad}_g(\xi), gNg^{-1})$, we must verify that $g.\phi$ satisfies the modular condition for the automorphism group η_{-t}^ξ of \mathcal{N}_g . Notice that as isomorphisms $\mathcal{N}_g \rightarrow \mathcal{N}$ we have:

$$\sigma_t^\xi \circ \text{Ad}(\rho(g)^{-1}) = \text{Ad}(\rho(g)^{-1}) \circ \eta_t^\xi, \quad \forall t \in \mathbb{R}. \quad (3)$$

As $\phi \in \text{KMS}(\rho, \xi, N)$, we know that $\phi \circ \sigma_t^\xi = \phi$ for all $t \in \mathbb{R}$. It then follows immediately from (3) that

$$(g.\phi) \circ \eta_t^\xi = \phi \circ \text{Ad}(\rho(g)^{-1}) \circ \eta_t^\xi = \phi \circ \sigma_t^\xi \circ \text{Ad}(\rho(g)^{-1}) = \phi \circ \text{Ad}(\rho(g)^{-1}) = g.\phi, \quad \forall t \in \mathbb{R}.$$

Next, take $x, y \in \mathcal{N}_g$. Then $x = \rho(g)x'\rho(g)^{-1}$ and $y = \rho(g)y'\rho(g)^{-1}$ for some $x', y' \in \mathcal{N}$. Let the function $F_{x', y'} : \overline{\text{St}} \rightarrow \mathbb{C}$ be continuous and bounded, holomorphic on St and satisfy $F_{x', y'}(t) = \phi(\sigma_{-t}^\xi(x')y')$ and

$F_{x',y'}(t+i) = \phi(y'\sigma_{-t}^\xi(x'))$ for all $t \in \mathbb{R}$. Define $\tilde{F}_{x,y} : \overline{\mathbb{S}t} \rightarrow \mathbb{C}$ by $\tilde{F}_{x,y}(z) := F_{x',y'}(z)$. Then $\tilde{F}_{x,y}$ satisfies the conditions of Definition 5.2 for η_{-t}^ξ . Indeed, notice using Equation (3) that $\sigma_t^\xi(x') = \rho(g)^{-1}\eta_t^\xi(x)\rho(g)$. Thus

$$\begin{aligned}\tilde{F}_{x,y}(t) &= F_{x',y'}(t) = \phi(\sigma_{-t}^\xi(x')y') = \phi\left(\rho(g)^{-1}\eta_{-t}^\xi(x)y\rho(g)\right) = (g\cdot\phi)(\eta_{-t}^\xi(x)y), \\ \tilde{F}_{x,y}(t+i) &= F_{x',y'}(t+i) = \phi(y'\sigma_{-t}^\xi(x')) = \phi\left(\rho(g)^{-1}y\eta_{-t}^\xi(x)\rho(g)\right) = (g\cdot\phi)(y\eta_{-t}^\xi(x)).\end{aligned}$$

Thus $g\cdot\phi \in \text{KMS}(\rho, \text{Ad}_g(\xi), gNg^{-1})$. \square

Let $\phi \in \text{KMS}(\rho, \xi, N)$. Let α denote the smooth \mathbb{R} -action on N defined by $\alpha_t(n) := e^{t\xi}ne^{-t\xi}$. We extend ρ_ϕ to $N \rtimes_\alpha \mathbb{R}$ by setting $\rho_\phi(n, t) = \rho_\phi(n)\Delta_\phi^{-it}$. Define

$$\begin{aligned}\mathcal{N}^{\infty, \phi} &:= \{x \in \mathcal{N} : (n, t) \mapsto \rho_\phi(n, t)\pi_\phi(x)\Omega_\phi \text{ is smooth } N \rtimes_\alpha \mathbb{R} \rightarrow \mathcal{H}_\phi\}, \\ \mathcal{D}_\phi &:= \pi_\phi(\mathcal{N}^{\infty, \phi})\Omega_\phi \subseteq \mathcal{H}_{\rho_\phi}.\end{aligned}\tag{4}$$

Notice that $\mathcal{N}^{\infty, \phi}$ and \mathcal{D}_ϕ are invariant under the left N - and $N \rtimes_\alpha \mathbb{R}$ -actions, respectively.

Lemma 5.10. *If $\phi \in \text{KMS}(\rho, \xi, N)^\infty$, then $\mathcal{N}^{\infty, \phi}$ is SOT-dense in \mathcal{N} and \mathcal{D}_ϕ is dense in \mathcal{H}_ϕ . In particular, ρ_ϕ is smooth when considered as representation of $N \rtimes_\alpha \mathbb{R}$.*

Proof. Since $\phi \in \mathcal{S}(\mathcal{N})^\infty$, the vector Ω_ϕ is smooth for the N -action ρ_ϕ by Lemma 5.8. Let $m \in N$. Then

$$\rho_\phi(n, t)\rho_\phi(m)\Omega_\phi = \rho(n)\Delta_\phi^{-it}\rho_\phi(m)\Delta_\phi^{it}\Omega_\phi = \rho(n)\sigma_{-t}^\phi(\rho_\phi(m))\Omega_\phi = \rho_\phi(ne^{t\xi}me^{-t\xi})\Omega_\phi, \quad \forall n \in N, t \in \mathbb{R},$$

where the last equality follows by Remark 5.4. Thus $(n, t) \mapsto \rho_\phi(n, t)\rho_\phi(m)\Omega_\phi$ is smooth $N \rtimes_\alpha \mathbb{R} \rightarrow \mathcal{H}_\phi$ and so $\rho(m)\Omega_\phi \in \mathcal{N}^{\infty, \phi}$. Thus $\rho(N) \subseteq \mathcal{N}^{\infty, \phi}$ and $\rho_\phi(N)\Omega_\phi \subseteq \mathcal{D}_\phi$. Since $\rho(N)'' = \mathcal{N}$ and $\rho_\phi(N)\Omega_\phi$ is total for \mathcal{H}_ϕ , it follows that $\mathcal{N}^{\infty, \phi}$ is SOT-dense in \mathcal{N} and that \mathcal{D}_ϕ is dense in \mathcal{H}_ϕ . As \mathcal{D}_ϕ is contained in the set of $N \rtimes_\alpha \mathbb{R}$ -smooth vectors by definition, the final observation follows. \square

5.2.1 Restrictions Imposed by the KMS Condition

Let us next determine some consequences of the KMS condition. Most notably, we will show that representations ρ which are smoothly-KMS give rise to generalized positive energy representations ρ_ϕ on the GNS-Hilbert space \mathcal{H}_ϕ of the corresponding state ϕ .

We continue in the notation of Section 5.2. Fixing a Lie subgroup $N \subseteq G$ and some element $\xi \in \mathfrak{g}$ satisfying $\text{Ad}(e^{t\xi})N \subseteq N$ for all $t \in \mathbb{R}$, we may as well suppose that $G = N \rtimes_\alpha \mathbb{R}$ for some smooth \mathbb{R} -action α on N by automorphisms. Let $\mathfrak{g} := \text{Lie}(G)$, $\mathfrak{n} := \text{Lie}(N)$ and write $D \in \text{der}(\mathfrak{n})$ for the derivation on \mathfrak{n} corresponding to α . Thus $\mathfrak{g} = \mathfrak{n} \rtimes_D \mathbb{R}\mathbf{d}$, where $\mathbf{d} := 1 \in \mathbb{R}$ denotes the standard basis element. Assume that ρ is KMS at \mathbf{d} relative to N , and let $\phi \in \text{KMS}(\rho, \mathbf{d}, N)$. We extend the N -representation $\rho_\phi = \pi_\phi \circ \rho$ on the GNS-Hilbert space \mathcal{H}_ϕ to $G = N \rtimes_\alpha \mathbb{R}$ by setting $\rho_\phi(n, t) = \rho_\phi(n)\Delta_\phi^{-it}$. Define further $H_\phi := -\log \Delta_\phi = -id\rho_\phi(\mathbf{d})$.

A first observation is the following.

Proposition 5.11. *Let A be an Abelian Lie subgroup of N such that $\alpha_t(A) \subseteq A$ for all $t \in \mathbb{R}$. Then $\rho_\phi(\alpha_t(a)) = \rho_\phi(a)$ for every $t \in \mathbb{R}$ and $a \in A$. In particular, if \mathcal{N} is a factor then $\rho(\alpha_t(a)) = \rho(a)$ for every $t \in \mathbb{R}$ and $a \in A$.*

Proof. Let $\mathcal{A}_\phi := \rho_\phi(A)''$. Write again ϕ for the vector state $\langle \Omega_\phi, \cdot \Omega_\phi \rangle$ on \mathcal{N}_ϕ . Let $\psi := \phi|_{\mathcal{A}_\phi}$ denote its restriction to \mathcal{A}_ϕ . As A is \mathbb{R} -invariant, so is $\mathcal{A}_\phi \subseteq \mathcal{N}_\phi$. Thus, the modular automorphism group σ^ϕ of \mathcal{N}_ϕ leaves \mathcal{A}_ϕ invariant. As ϕ satisfies the modular condition for σ^ϕ , so does ψ for the automorphism group $t \mapsto \sigma_t^\phi|_{\mathcal{A}_\phi}$. Recall from Remark 5.3(2) that Ω_ϕ is separating for \mathcal{N}_ϕ . Hence it is so for \mathcal{A}_ϕ . In view of Remark 5.3(1) this implies that the modular automorphism group σ^ψ on \mathcal{A}_ϕ is uniquely determined by the modular condition. Thus $\sigma_t^\psi = \sigma_t^\phi|_{\mathcal{A}_\phi}$ for all $t \in \mathbb{R}$. As \mathcal{A}_ϕ is Abelian, we know by [BR97, Prop. 5.3.28] that $\sigma_t^\psi = \text{id}_{\mathcal{A}_\phi}$. Thus $\sigma_t^\phi|_{\mathcal{A}_\phi} = \text{id}_{\mathcal{A}_\phi}$. We know from Remark 5.4 that $\rho_\phi \circ \alpha_{-t} = \sigma_t^\phi \circ \rho_\phi$. Thus $\rho_\phi(\alpha_t(a)) = \rho_\phi(a)$ for all $a \in \mathcal{A}$ and $t \in \mathbb{R}$. If \mathcal{N} is a factor, then ϕ is faithful and π_ϕ is injective by Remark 5.3(1,3). Thus $\rho(\alpha_t(a)) = \rho(a)$ follows from $\rho_\phi(\alpha_t(a)) = \rho_\phi(a)$. \square

Let us illustrate Proposition 5.11 with the following noteworthy consequence for loop groups:

Corollary 5.12. *Let K be a compact 1-connected simple Lie group and $LK := C^\infty(S^1; K)$. Let α denote the \mathbb{T} -action on LK by rotations, with corresponding derivation $D := \frac{d}{d\theta}$ on $L\mathfrak{k}$. Consider the Lie group $G := LK \rtimes_\alpha \mathbb{T}$ with Lie algebra $\mathfrak{g} := L\mathfrak{k} \rtimes_D \mathbb{R}d$, where $d := 1 \in \mathbb{R}$. Suppose that the smooth unitary G -representation ρ is KMS at $d \in \mathfrak{g}$ relative to LK . Assume that $\rho(LK)''$ is a factor. Then $LK \subseteq \ker \rho$.*

Proof. Suppose $T \subseteq K$ is a maximal torus with Lie algebra \mathfrak{t} . Then $LT \subseteq LK$ is an Abelian α -invariant subgroup. By Proposition 5.11 it follows that $d\rho(DLt) = \{0\}$. As any $X \in \mathfrak{k}$ is contained in a maximal torus, it follows that $d\rho(\frac{df}{d\theta} \otimes X) = 0$ for any $f \in C^\infty(S^1)$ and $X \in \mathfrak{k}$. Consequently $d\rho(D\mathfrak{g}) = \{0\}$ and hence $D\mathfrak{g}_\mathbb{C} \subseteq \ker d\rho$, where we have extended $d\rho : \mathfrak{g} \rightarrow \mathcal{L}^1(\mathcal{H}_\rho^\infty)$ \mathbb{C} -linearly to the complexification $\mathfrak{g}_\mathbb{C}$. As $\ker d\rho$ is an ideal in $\mathfrak{g}_\mathbb{C}$ and $L\mathfrak{k}_\mathbb{C} = DL\mathfrak{k}_\mathbb{C} + [DL\mathfrak{k}_\mathbb{C}, DL\mathfrak{k}_\mathbb{C}]$, it follows that $L\mathfrak{k}_\mathbb{C} \subseteq \ker d\rho$. Notice that LK is connected because K is 1-connected. It is also locally exponential by [Nee01, Thm. II.1]. It follows that $LK \subseteq \ker \rho$. \square

Thus, one necessarily has to pass to a non-trivial central \mathbb{T} -extension \widehat{LK} of $LK \rtimes_\alpha \mathbb{T}$ to allow for interesting KMS-representations of \widehat{LK} that are smoothly-KMS at some $\widehat{d} \in \widehat{L\mathfrak{k}}$ covering $d \in L\mathfrak{k} \rtimes_D \mathbb{R}d$, as one may have expected from the positive energy analogue (which follows from [PS86, Thm 9.3.5]).

We now proceed with the observation that KMS representations give rise to generalized positive energy representations on the GNS-Hilbert space corresponding to the KMS state:

Theorem 5.13. *Let $\phi \in \text{KMS}(\rho, d, N)^\infty$. Let $x \in \mathcal{N}^{\phi, \infty}$ and assume $\psi := \pi_\phi(x)\Omega_\phi \in \mathcal{D}_\phi$ has unit norm. Then*

$$\langle \pi_\phi(x)\Omega_\phi, -id\rho_\phi(\text{Ad}_n(d))\pi_\phi(x)\Omega_\phi \rangle \geq -\log(\|\pi_\phi(x)\|^2) \quad \forall n \in N. \quad (5)$$

In particular the representation ρ_ϕ of $N \rtimes_\alpha \mathbb{R}$ with domain \mathcal{D}_ϕ is of generalized positive energy at $d \in n \rtimes_D \mathbb{R}d$.

Lemma 5.14. *Let $x \in \mathcal{N}$ be such that $0 \neq \psi := d\pi_\phi(x)\Omega_\phi \in \text{dom}(H_\phi)$. Then*

$$\frac{\langle \psi, H_\phi \psi \rangle}{\|\psi\|^2} \geq -\log\left(\frac{\|S_\phi \psi\|^2}{\|\psi\|^2}\right). \quad (6)$$

Proof. In view of the correlation lower bounds satisfied by KMS states, see e.g. [BR97, Thm. 5.3.15 (1) \implies (2)] or [FV77, Thm. II.4, (i) \implies (iii)], we have:

$$\langle \pi_\phi(x)\Omega_\phi, [H_\phi, \pi_\phi(x)]\Omega_\phi \rangle \geq -\|\pi_\phi(x)\Omega_\phi\|^2 \log\left(\frac{\|\pi_\phi(x)^*\Omega_\phi\|^2}{\|\pi_\phi(x)\Omega_\phi\|^2}\right).$$

Since $H_\phi\Omega_\phi = 0$, it follows that $\langle \pi_\phi(x)\Omega_\phi, [H_\phi, \pi_\phi(x)]\Omega_\phi \rangle = \langle \pi_\phi(x)\Omega_\phi, H_\phi\pi_\phi(x)\Omega_\phi \rangle$. The assertion follows. \square

Proof of Theorem 5.13:

Recall that $\mathcal{D}_\phi \subseteq \mathcal{H}_{\rho_\phi}^\infty$ and that $\mathcal{D}_\phi \subseteq \text{dom}(S_\phi)$, because the stronger condition $\mathcal{N}_\phi\Omega_\phi \subseteq \text{dom}(S_\phi)$ is satisfied. Let $n \in N$. Notice that $\|S_\phi\rho_\phi(n)\psi\| = \|\pi_\phi(x^*)\pi_\phi(n)^{-1}\Omega_\phi\| \leq \|\pi_\phi(x)\|$. Recalling that \mathcal{D}_ϕ is N -invariant, we can apply equation (6) to the vector $\rho_\phi(n)\psi$. Using $-id\rho(d) = -\log\Delta_\phi = H_\phi$ it follows that

$$\langle \psi, -id\rho_\phi(\text{Ad}_{n^{-1}}(d))\psi \rangle = \langle \rho_\phi(n)\psi, -id\rho_\phi(d)\rho_\phi(n)\psi \rangle \geq -\log(\|S_\phi\rho_\phi(n)\psi\|^2) \geq -\log(\|\pi_\phi(x)\|^2). \quad \square$$

As a consequence of Theorem 5.13, we find that the observations of Section 4 impose restrictions on KMS representations. Let us illustrate this with the following immediate consequence:

Corollary 5.15. *Let $\bar{\rho}$ be a smooth projective unitary representation of G on \mathcal{H}_ρ . Assume that $\mathcal{N} := \bar{\rho}(N)''$ is a factor. Let $\rho : \widehat{G} \rightarrow \text{U}(\mathcal{H}_\rho)$ be the lift of $\bar{\rho}$, for some central \mathbb{T} -extension \widehat{G} of G with Lie algebra $\widehat{\mathfrak{g}}$. Let $\widehat{N} \subseteq \widehat{G}$ cover N . Let ω represent the class in $H_{\text{ct}}^2(\mathfrak{g}, \mathbb{R})$ corresponding to $\widehat{\mathfrak{g}}$. Let $\xi \in \mathfrak{g}$ and suppose $\widehat{\xi} \in \widehat{\mathfrak{g}}$ covers ξ . Let $\phi \in \text{KMS}(\rho, \widehat{\xi}, \widehat{N})^\infty$. Assume that $\eta \in \mathfrak{g}$ satisfies $[[\xi, \eta], \eta] = 0$. Then $\omega([\xi, \eta], \eta) \geq 0$ and*

$$\omega([\xi, \eta], \eta) = 0 \iff d\bar{\rho}([\xi, \eta]) = 0.$$

Proof. Consider the representation ρ_ϕ of $\widehat{N} \rtimes \mathbb{R}$ on the GNS Hilbert space \mathcal{H}_ϕ , where \mathbb{R} acts on \widehat{N} by $\text{Ad}(e^{t\widehat{\xi}})|_{\widehat{N}}$ and where $\rho_\phi(1, t) = \Delta_\phi^{-it}$ for $t \in \mathbb{R}$. Let $\bar{\rho}_\phi$ be the corresponding projective unitary representation of $N \rtimes \mathbb{R}$ on \mathcal{H}_ϕ , where \mathbb{R} acts on N by $\text{Ad}(e^{t\xi})|_N$. By Theorem 5.13, ρ_ϕ is of g.p.e. at $d \in \widehat{n} \rtimes \mathbb{R}d$ and so $\bar{\rho}_\phi$ is of g.p.e. at d . It follows from Proposition 4.4 that $\omega([\xi, \eta], \eta) \geq 0$ and $\omega([\xi, \eta], \eta) = 0 \iff d\bar{\rho}_\phi([\xi, \eta]) = 0$. As \mathcal{N} is a factor, the KMS state $\phi \in \mathcal{S}(\mathcal{N})$ is faithful and the corresponding GNS-representation $\pi_\phi : \mathcal{N} \rightarrow \mathcal{B}(\mathcal{H}_\phi)$ is injective, by Remark 5.3(1,3). This implies that $\ker d\rho_\phi = \ker d\rho$. Thus $\omega([\xi, \eta], \eta) = 0 \iff d\bar{\rho}([\xi, \eta]) = 0$. \square

5.2.2 Some Examples of KMS Representations

Let us consider a variety of examples of KMS representations, thereby showing in various situations that a well-known σ -KMS state ϕ on a von Neumann algebra \mathcal{N} admits some underlying smooth structure. More precisely, we construct a unitary representation ρ of a (typically infinite-dimensional) Lie group G such that $\mathcal{N} = \rho(N)''$, $\phi \in \text{KMS}(\rho, \xi, N)^\infty$ and $\sigma_t = \text{Ad}(\rho(e^{t\xi}))$ for some $\xi \in \mathfrak{g}$ and Lie subgroup N of G . In particular, in this case the 1-parameter group σ on \mathcal{N} implements the \mathbb{R} -action $t \mapsto \text{Ad}(e^{t\xi})|_{\mathcal{N}}$ on the Lie subgroup N of G .

Let us begin with the simplest class of examples, which correspond to Gibbs states, as in Example 5.5.

Example 5.16. Take for N simply $N = G$. Let (ρ, \mathcal{H}) be an irreducible unitary G -representation. Then $\mathcal{N} = \mathcal{B}(\mathcal{H})$. Let $\xi \in \mathfrak{g}$ and define the self-adjoint operator $H := -i\overline{d\rho(\xi)}$. Let $\beta > 0$ and assume that $Z_\beta := \text{Tr}(e^{-\beta H}) < \infty$. Define the Gibbs state $\phi_\beta(x) := \frac{1}{Z_\beta} \text{Tr}(e^{-\beta H} x)$ for $x \in \mathcal{N}$. As in Example 5.5, we have $\sigma_{-t}^{\phi_\beta}(x) = e^{it\beta H} x e^{-it\beta H} = \rho(e^{t\beta\xi}) x \rho(e^{-t\beta\xi})$ for any $x \in \mathcal{N}$. Consequently, $\phi_\beta \in \text{KMS}(\rho, \beta\xi)$ and so ρ is a KMS representation at $\beta\xi \in \mathfrak{g}$. If in addition $\widehat{\phi_\beta} : G \rightarrow \mathbb{C}$ is smooth, then ρ is smoothly-KMS at $\beta\xi \in \mathfrak{g}$. By Lemma 5.9, ρ is also KMS at any element in the adjoint orbit of $\beta\xi$. In view of Example 5.5, the representation ρ_ϕ of $G \rtimes \mathbb{R}$ on $\mathcal{H}_\phi := \overline{\mathcal{B}(\mathcal{H}_\rho)}^{(-, -)^{\phi}}$ is given by $\rho_\phi(g, t)x\Omega_\phi = \rho(g)\rho(e^{t\beta\xi})x\rho(e^{-t\beta\xi})\Omega_\phi = \rho(g)\sigma_{-t}^{\phi_\beta}(x)\Omega_\phi$, where $\Omega_\phi := I \in \mathcal{B}(\mathcal{H}_\rho) \subseteq \mathcal{H}_\phi$ denotes the cyclic vector.

In fact, Proposition 5.17 below entails that any KMS representation ρ for which \mathcal{N} is a factor of type I is of the form described in Example 5.16.

Proposition 5.17. *Let $\xi \in \mathfrak{g}$ and $\beta > 0$. Suppose that $\rho|_{\mathcal{N}}$ is irreducible and that $\phi \in \text{KMS}(\rho, \beta\xi, N)$. Write $H := -i\overline{d\rho(\xi)}$. Then $Z_\beta := \text{Tr}(e^{-\beta H}) < \infty$ and $\phi(x) = \frac{1}{Z_\beta} \text{Tr}(e^{-\beta H} x)$.*

Proof. As $\rho|_{\mathcal{N}}$ is irreducible, it follows that $\mathcal{N} = \mathcal{B}(\mathcal{H}_\rho)$. Thus $\phi(x) = \text{Tr}(\delta x)$ for some $\delta \in L^1(\mathcal{H}_\rho)_+$ satisfying $\text{Tr}(\delta) = 1$, where $L^1(\mathcal{H}_\rho)$ denotes Banach space of trace-class operators on \mathcal{H}_ρ . Moreover, in view of Remark 5.3(3), we know that ϕ is faithful on \mathcal{N} . By assumption, ϕ satisfies the modular condition for the automorphism group $t \mapsto \text{Ad}(\rho(e^{-t\beta\xi})) =: \sigma_{-t\beta}$. On the other hand, as ϕ is faithful, there exists by Remark 5.3(1) a unique automorphism group σ_t^ϕ of \mathcal{N} for which ϕ satisfies the modular condition. It follows that $\sigma_{-t\beta} = \sigma_t^\phi$. When $\mathcal{N} = \mathcal{B}(\mathcal{H}_\rho)$ and $\phi(x) = \text{Tr}(\delta x)$, the modular automorphism group σ_t^ϕ corresponding to ϕ is $\sigma_t^\phi(x) = \delta^{it} x \delta^{-it}$. In view of $\sigma_t^\phi = \sigma_{-t\beta}$, it follows that $\delta^{it} x \delta^{-it} = \rho(e^{-t\beta\xi}) x \rho(e^{t\beta\xi})$ for every $x \in \mathcal{N}$. As $\mathcal{Z}(\mathcal{N}) = \mathbb{C}I$ and both $t \mapsto \delta^{it}$ and $t \mapsto \rho(e^{t\beta\xi})$ are strongly continuous unitary 1-parameter groups, it follows that there is some continuous homomorphism $c : \mathbb{R} \rightarrow \mathbb{T}$ such that $\delta^{it} = c(t)\rho(e^{-t\beta\xi}) = c(t)e^{-t\beta d\rho(\xi)}$ for all $t \in \mathbb{R}$. Thus there exists $\mu \in \mathbb{R}$ such that $\delta^{it} = e^{-it(\beta H + \mu I)}$ for all $t \in \mathbb{R}$; so $\log \delta = -(\beta H + \mu I)$. Since $\text{Tr}(\delta) = 1$, it follows that $Z_\beta = \text{Tr}(e^{-\beta H}) = \text{Tr}(e^{-(\beta H + \mu I)} e^{\mu I}) = \text{Tr}(\delta e^{\mu I}) = e^\mu \phi(1) = e^\mu < \infty$. It follows that

$$\frac{1}{Z_\beta} \text{Tr}(e^{-\beta H} x) = e^{-\beta\mu} \text{Tr}(e^{-\beta H} x) = \text{Tr}(e^{-(\beta H + \mu I)} x) = \text{Tr}(\delta x) = \phi(x), \quad \forall x \in \mathcal{B}(\mathcal{H}). \quad \square$$

For more interesting examples, one has to consider a Lie subgroup N of G which is not of type I, so that the von Neumann algebra \mathcal{N} need not be type I.

Example 5.18 (Powers' factors). Define $G_n := \prod_{k=1}^n \text{SU}(2)$ and let $\eta_n : G_n \hookrightarrow G_{n+1}$ be defined by

$$\eta_n : G_n \xrightarrow{\text{id} \times 1} G_n \times \text{SU}(2) = G_{n+1}.$$

Write $\mathfrak{g}_n := \text{Lie}(G_n)$ and $L(\eta_n) := \text{Lie}(\eta_n)$. The direct limit $G := \varinjlim_n (G_n, \eta_n)$ consists of sequences (u_k) in $\text{SU}(2)$ with $u_k = 1$ for all but finitely many values of k . It can be equipped with the structure of a regular Lie group that is modeled on the locally convex inductive limit $\mathfrak{g} := \varinjlim_n (\mathfrak{g}_n, L(\eta_n))$ [Glo05, Thm. 4.3] and has the exponential map $\exp_G = \varinjlim_n \exp_{G_n}$ [Glo05, Prop. 4.6]. Let $H := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and $\xi := iH \in \mathfrak{su}(2)$.

Consider the following \mathbb{R} -action α on G defined by $(\alpha_t(u))_k := e^{t\xi} u_k e^{-t\xi}$ for $u \in G$. The corresponding action $\mathbb{R} \times G \rightarrow G$ is smooth. Indeed, the restriction of α to $\mathbb{R} \times G_n$ yields a smooth action $\alpha^{(n)} : \mathbb{R} \times G_n \rightarrow G_n$ for every $n \in \mathbb{N}$. It follows from [Glo05, Thm. 4.6(c)] that $\varinjlim_n \alpha^{(n)} : \varinjlim_n (\mathbb{R} \times G_n) \rightarrow G$ is smooth. By [Glo05, Prop. 3.7] we further have $\varinjlim_n (\mathbb{R} \times G_n) = \mathbb{R} \times G$ as smooth manifolds. This shows that $\alpha : \mathbb{R} \times G \rightarrow G$ is smooth. Consider the Lie group $G^\sharp := G \rtimes_\alpha \mathbb{R}$ with Lie algebra $\mathfrak{g}^\sharp := \mathfrak{g} \rtimes_{\mathbb{D}} \mathbb{R}\mathbf{d}$, where $\mathbf{d} := (0, 1)$. Using the so-called Powers' factors, we define unitary representations ρ of G^\sharp which are smoothly-KMS at \mathbf{d} relative to $G \triangleleft G^\sharp$ and for which $\rho(G)''$ is a factor of type III $_\lambda$ for arbitrary $\lambda \in (0, 1)$. Define the finite-dimensional C^* -algebra $\mathcal{M}_n := \rho_n(G_n)'' = \bigotimes_{k=1}^n \mathcal{B}(\mathbb{C}^2)$ for every $n \in \mathbb{N}$. Let $\beta > 0$. Define the state $\phi(x) := \frac{1}{Z} \text{Tr}(e^{-\beta H} x)$ on $\mathcal{B}(\mathbb{C}^2)$, where $Z := \text{Tr}(e^{-\beta H}) = 2 \cosh(\beta)$. Let ϕ_n be the state on \mathcal{M}_n defined by

$\phi_n(x_1 \otimes \cdots \otimes x_n) = \prod_{k=1}^n \phi(x_k)$. The GNS-representation of $\mathcal{B}(\mathbb{C}^2)$ defined by ϕ is $\mathcal{H}_\phi := \mathcal{B}(\mathbb{C}^2)$ equipped with left $\mathcal{B}(\mathbb{C}^2)$ -action and the inner product $\langle a, b \rangle := \frac{1}{2} \text{Tr}(e^{-\beta H} a^* b)$. Similarly the GNS-representation of \mathcal{M}_n corresponding to ϕ_n is $\mathcal{H}_{\phi_n} := \bigotimes_{k=1}^n \mathcal{H}_\phi$. The isometric inclusions $\mathcal{H}_{\phi_n} \hookrightarrow \mathcal{H}_{\phi_{n+1}}, x \mapsto x \otimes 1$ define a directed system of Hilbert spaces, and the algebraic direct limit $\varinjlim_n \mathcal{H}_{\phi_n}$ becomes naturally a pre-Hilbert space. Let \mathcal{H} denote its Hilbert space closure. Let $\iota_n : \mathcal{H}_{\phi_n} \hookrightarrow \mathcal{H}$ denote the canonical inclusion. For every $n \in \mathbb{N}$, there is a $*$ -representation π_n of \mathcal{M}_n on \mathcal{H} defined by

$$\pi_n(x) \iota_m(\psi_1 \otimes \cdots \otimes \psi_m) := \iota_m(x_1 \psi_1 \otimes \cdots \otimes x_n \psi_n \otimes \psi_{n+1} \otimes \cdots \otimes \psi_m), \quad m \geq n.$$

Let $\mathcal{M}_\infty := \left(\bigcup_{n \in \mathbb{N}} \pi_n(\mathcal{M}_n) \right)''$. The vector $\Omega := 1 \otimes 1 \otimes \cdots \in \mathcal{H}$ is cyclic and separating for \mathcal{M}_∞ [Tak03b, XIV, Prop. 1.11], so \mathcal{H} may be identified with the GNS-representation of \mathcal{M}_∞ w.r.t. the state $\phi_\infty := \langle \Omega, \cdot \rangle$ on \mathcal{M}_∞ . Observe that ϕ_∞ satisfies $\phi_\infty(\iota_n(x)) = \phi_n(x)$ for all $n \in \mathbb{N}$ and $x \in \mathcal{M}_n$. The von Neumann algebra $(\mathcal{M}_\infty, \phi_\infty) =: \bigotimes_{k=1}^\infty (\mathcal{B}(\mathbb{C}^2), \phi)$ is the so-called Powers' factor with parameter $a := \frac{e^{-\beta}}{2 \cosh(\beta)} \in (0, \frac{1}{2})$, which is a factor of type III $_\lambda$ with $\lambda = e^{-2\beta} = \frac{a}{1-a} \in (0, 1)$ [Tak03b, XVIII, Theorem 1.1]. The modular automorphism group $\sigma_t^{\phi_\infty}$ on \mathcal{M}_∞ defined by ϕ_∞ is given by $\sigma_t^{\phi_\infty} = \bigotimes_{k=1}^\infty \text{Ad}(e^{-\beta t \xi})$ [Tak03b, XIV, Prop. 1.11], where $\bigotimes_{k=1}^\infty \text{Ad}(e^{\beta t \xi}) \in \text{Aut}(\mathcal{M}_\infty)$ satisfies $\bigotimes_{k=1}^\infty \text{Ad}(e^{\beta t \xi}) \circ \iota_n = \iota_n \circ \bigotimes_{k=1}^n \text{Ad}(e^{\beta t \xi})$ for all $t \in \mathbb{R}$ and $n \in \mathbb{N}$ and is defined from this condition by continuity, where we used [Tak03b, XIV, Thm. 1.13] and that $\phi \circ \text{Ad}(e^{\beta t \xi}) = \phi$ for all $t \in \mathbb{R}$. Consider the unitary representation $\rho : G \rtimes_\alpha \mathbb{R} \rightarrow \text{U}(\mathcal{H})$ defined by

$$\rho(u, \beta t) := \left(\bigotimes_k^\infty u_k \right) \circ \Delta_{\phi_\infty}^{-it}, \quad u \in G, t \in \mathbb{R}$$

which is well-defined because $u = (u_k) \in G$ is a sequence in $\text{SU}(2)$ with $u_k = 1$ for all k sufficiently large. Since $\rho(\beta t) = \Delta_{\phi_\infty}^{-it}$ and $\rho(G)'' = \mathcal{M}_\infty$, it follows that ρ is KMS at $\beta \mathbf{d} \in \mathfrak{g}^\sharp$ relative to $G \triangleleft G^\sharp$. To see that $\widehat{\phi}_\infty : G \rightarrow \mathbb{C}$ is smooth, it suffices to show that its restriction to G_n is smooth for every $n \in \mathbb{N}$, using the universal property of the smooth manifold structure on $G^\sharp = \varinjlim_n G_n^\sharp$ [Glo05, Thm. 3.1]. This is the case, as $\langle \Omega, \rho(u) \Omega \rangle = \prod_{k=1}^n \phi(u_k)$ for any $u \in G_n$, which is smooth $G_n \rightarrow \mathbb{C}$. Thus $\Omega \in \mathcal{H}_\rho^\infty$ and so ρ is smoothly-KMS at $\beta \mathbf{d} \in \mathfrak{g}^\sharp$ relative to $G \triangleleft G^\sharp$.

Example 5.19 (Standard real subspaces and Heisenberg representations). Let \mathcal{H} be a complex Hilbert space. Consider the real Heisenberg group $G := \text{H}(\mathcal{H}, \omega)$, where $\omega(v, w) = \text{Im} \langle v, w \rangle$. An \mathbb{R} -linear closed subspace $\mathcal{K} \subseteq \mathcal{H}$ is called *cyclic* if $\mathcal{K} + i\mathcal{K}$ is dense in \mathcal{H} . It is called *separating* if $\mathcal{K} \cap i\mathcal{K} = \{0\}$. A *standard subspace* is a closed \mathbb{R} -linear subspace $\mathcal{K} \subseteq \mathcal{H}$ which is both cyclic and separating. We show that any standard real subspace gives rise to a smooth KMS representation. Let $\mathcal{K} \subseteq \mathcal{H}$ be a standard real subspace. Write $\delta_{\mathcal{K}}$ for the corresponding modular operator on \mathcal{H} , which is generally unbounded, positive and self-adjoint, see e.g. [NO17, Sec. 3]. Then $t \mapsto \delta_{\mathcal{K}}^{it}$ is a strongly-continuous unitary 1-parameter group on \mathcal{H} satisfying in particular $\delta_{\mathcal{K}}^{it} \mathcal{K} \subseteq \mathcal{K}$. We first pass to the \mathbb{R} -smooth vectors \mathcal{K}^∞ to obtain a regular Lie group $\text{H}(\mathcal{K}^\infty, \omega) \rtimes \mathbb{R}$. We then construct a KMS representation thereof using second-quantization. The details are given below.

Let \mathcal{K}^∞ denote the set \mathbb{R} -smooth vectors in \mathcal{K} . Then \mathcal{K}^∞ is dense in \mathcal{K} and \mathbb{R} -invariant. It moreover carries a Fréchet topology which is finer than the one inherited as a subspace of \mathcal{K} and for which the action $\mathbb{R} \times \mathcal{K}^\infty \rightarrow \mathcal{K}^\infty$ is smooth [Nee10a, Thm. 4.4, Lem. 5.2]. As $\omega : \mathcal{K}^\infty \times \mathcal{K}^\infty \rightarrow \mathbb{R}$ is bilinear and continuous w.r.t. this topology, it is smooth. Thus the generalized Heisenberg group $N := \text{H}(\mathcal{K}^\infty, \omega)$ is a Lie group. (Notice that $\omega|_{\mathcal{K}^\infty}$ may be degenerate.) It is as a subgroup of G generated by \mathcal{K}^∞ . As \mathcal{K}^∞ is a Fréchet space, it is Mackey complete by [KM97, Thm. I.4.11], which implies using [Nee06, Thm. V.1.8] that N is regular. Write $\mathfrak{n} := \text{Lie}(N)$. By construction \mathbb{R} acts smoothly on N by $\delta_{\mathcal{K}}^{it}$, so that $N^\sharp := N \rtimes \mathbb{R}$ is a regular Lie group. Let $\mathfrak{n}^\sharp := \mathfrak{n} \rtimes \mathbb{R} \mathbf{d}$ denote its Lie algebra. We construct a representation of N^\sharp which is smoothly KMS at $\mathbf{d} \in \mathfrak{n}^\sharp$ relative to $N \triangleleft N^\sharp$. Let us recall the standard representation of $\text{H}(\mathcal{H}, \omega)$ on the Bosonic Fock space $\mathcal{F}(\mathcal{H})$. Equip the symmetric algebra $S^\bullet(\mathcal{H})$ with the inner product

$$\langle v_1 \cdots v_n, w_1 \cdots w_n \rangle = \sum_{\sigma \in S_n} \prod_{j=1}^n \langle v_j, w_{\sigma_j} \rangle. \quad (7)$$

Let $\mathcal{F}(\mathcal{H})$ denote the Hilbert space closure of $S^\bullet(\mathcal{H})$ and let $\Omega := 1 \in \mathcal{H}$ denote the vacuum vector. Then \mathcal{H} contains (and is generated by) the vectors $e^v := \sum_{n=0}^\infty \frac{1}{n!} v^n \in \mathcal{H}$ for $v \in \mathcal{H}$. There is a continuous irreducible unitary representation W of $\text{H}(\mathcal{H}, \omega)$ on $\mathcal{F}(\mathcal{H})$ satisfying $W(z, v) e^w = z e^{-\frac{1}{2} \|v\|^2 - \langle v, w \rangle} e^{v+w}$ for $v, w \in \mathcal{H}$ and $z \in \mathbb{T}$ [PS86, Sec. 9.5]. Moreover, any unitary $u \in \text{U}(\mathcal{H})$ extends canonically to a unitary $\mathcal{F}(u) \in \text{U}(\mathcal{F}(\mathcal{H}))$. We further have:

$$W(uv) = \mathcal{F}(u) W(v) \mathcal{F}(u)^{-1}, \quad \forall u \in \text{U}(\mathcal{H}), v \in \mathcal{H} \quad (8)$$

In view of (8), W and \mathcal{F} together define a representation ρ of the Lie group N^\sharp by $\rho(n, t) := W(n)\mathcal{F}(\delta_{\mathcal{K}}^{it})$. Let $\mathcal{N} := W(N)''$. As \mathcal{K} is a standard real subspace and \mathcal{K}^∞ is dense in \mathcal{K} , it follows that Ω is cyclic and separating for \mathcal{N} [NO17, Lem. 6.2]. Let ϕ denote the faithful vector state on \mathcal{N} defined by $\phi(x) = \langle \Omega, x\Omega \rangle$. Using [NO17, Prop. 6.10] we have $\Delta_\phi^{it} = \mathcal{F}(\delta_{\mathcal{K}}^{it})$ for all $t \in \mathbb{R}$. Consequently ρ is KMS at $-\mathbf{d} \in \mathfrak{n}^\sharp$ relative to $N \triangleleft N^\sharp$ (notice the minus sign in Definition 5.2). To see W is smoothly KMS, observe that $\widehat{\phi} : N \rightarrow \mathbb{C}$ is smooth because it is given by

$$\widehat{\phi}(z, v) = \langle \Omega, W(z, v)\Omega \rangle = ze^{-\frac{1}{2}\|v\|^2}. \quad (9)$$

The following provides an example where ρ is smoothly-KMS at various $\xi_I \in \mathfrak{g}$, relative to distinct subgroups $N_I \subseteq G$, where $I \in \mathcal{I}$ for some indexing set \mathcal{I} :

Example 5.20 (Bisognano-Wichmann and Möbius covariant nets).

Recall that $SU(1, 1)$ acts on S^1 . Explicitly, for $g = \begin{pmatrix} \alpha & \beta \\ \beta & \bar{\alpha} \end{pmatrix} \in SU(1, 1)$ with $\alpha, \beta \in \mathbb{C}$ satisfying $|\alpha|^2 - |\beta|^2 = 1$, define $g(z) := \frac{\alpha z + \beta}{\beta z + \bar{\alpha}}$ for $z \in \mathbb{C}$ with $|z| = 1$. With g as above, define the unitary action of $SU(1, 1)$ on the complex Hilbert space $V := L^2(S^1; \mathbb{C})$ by $(u(g)f)(z) := (\alpha - \bar{\beta}z)^{-1}f(g^{-1}(z))$. Let $V_{\mathbb{R}}$ denote the real vector space underlying V . Define the real Fréchet space $V_{\mathbb{R}}^\infty := C^\infty(S^1; \mathbb{C})$ and consider the symplectic vector space $(V_{\mathbb{R}}^\infty, \omega)$, where $\omega(v, w) := \text{Im}\langle v, w \rangle$ for $v, w \in V_{\mathbb{R}}^\infty$. Let $H(V_{\mathbb{R}}^\infty, \omega)$ be the corresponding real Heisenberg group. Consider the Lie group $G := H(V_{\mathbb{R}}^\infty, \omega) \rtimes SU(1, 1)$. Let $\mathbf{r} := \frac{i}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and $\mathbf{d} := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ denote the generators in $\mathfrak{su}(1, 1)$ of the rotation and the dilation subgroups in $SU(1, 1)$, respectively. Write \mathcal{I} for the set of intervals of S^1 , on which $SU(1, 1)$ acts naturally. For $I \in \mathcal{I}$, let $\xi_I \in \mathfrak{su}(1, 1)$ be the unique element in the adjoint orbit of \mathbf{d} fixing the endpoints of I . Define further the closed real subspace $V(I) := \{f \in L^2(S^1; \mathbb{C}) : \text{supp}(f) \subseteq I\}$ of $V_{\mathbb{R}}$, and set $V(I)^\infty := V(I) \cap V_{\mathbb{R}}^\infty$. Let $N_I := H(V(I)^\infty) \subseteq G$ be the corresponding closed subgroup of G . We construct a unitary representation ρ of G which is of p.e. at $\mathbf{r} \in \mathfrak{su}(1, 1)$ and which is KMS at $\xi_I \in \mathfrak{su}(1, 1)$ relative to N_I , for every $I \in \mathcal{I}$. The details are given below.

Let P be the orthogonal projection onto the space $PV := H_+^2(S^1; \mathbb{C})$ spanned by the non-negative Fourier modes. Let $P^\perp V$ denote its orthogonal complement in $L^2(S^1; \mathbb{C})$. Consider the complex structure \mathcal{J} on $V_{\mathbb{R}}$ defined by $\mathcal{J} := i(2P - I) = iP - iP^\perp$. Let $V_{\mathcal{J}}$ denote the corresponding complex Hilbert space, equipped with the inner product $\langle v, w \rangle_{\mathcal{J}} := \langle Pv, Pw \rangle_V + \langle P^\perp v, P^\perp w \rangle_V$, so that $V_{\mathcal{J}} \cong PV \oplus \overline{P^\perp V}$ as complex Hilbert spaces. Observe that $\text{Im}\langle v, w \rangle_{\mathcal{J}} = \text{Im}\langle v, w \rangle = \omega(v, w)$ for $v, w \in V_{\mathbb{R}}$, so $H(V_{\mathbb{R}}, \omega) = H(V_{\mathcal{J}}, \text{Im}\langle -, - \rangle_{\mathcal{J}})$ as real Lie groups. The $SU(1, 1)$ -action u on $L^2(S^1; \mathbb{C})$ commutes with P and hence also with \mathcal{J} . It consequently defines a continuous unitary representation of $SU(1, 1)$ on the complex Hilbert space $V_{\mathcal{J}} \cong PV \oplus \overline{P^\perp V}$ leaving both PV and $\overline{P^\perp V}$ invariant. It is by construction of p.e. at $\mathbf{r} \in \mathfrak{su}(1, 1)$. As in Example 5.19, let W denote the standard representation of $H(V_{\mathcal{J}}, \text{Im}\langle -, - \rangle_{\mathcal{J}})$ on $\mathcal{F}(V_{\mathcal{J}})$. Letting $SU(1, 1)$ act on $\mathcal{F}(V_{\mathcal{J}})$ by second quantization, we obtain a smooth unitary representation ρ of G on $\mathcal{F}(V_{\mathcal{J}})$ which is of p.e. at $\mathbf{r} \in \mathfrak{su}(1, 1)$. Explicitly, $\rho(v, u) = W(v)\mathcal{F}(u)$ for $v \in H(V_{\mathbb{R}}^\infty, \omega)$ and $u \in SU(1, 1)$. Now, it follows from [Was98, p. 497] that $V(I) \subseteq V_{\mathcal{J}}$ is a standard real subspace for any interval $I \in \mathcal{I}$. Let δ_I^{it} denote the corresponding modular 1-parameter group, as in Example 5.19. The assignment $I \mapsto V(I)$, called a *net of standard subspaces*, satisfies $I_1 \subseteq I_2 \implies V(I_1) \subseteq V(I_2)$ (*isotony*), $V_{g \cdot I} = u(g)V(I)$ for $g \in SU(1, 1)$ ($SU(1, 1)$ -*covariance*) and $I_1 \cap I_2 = \emptyset \implies V(I_2) \subseteq V(I_1)^\perp$ (*locality*). Recalling additionally that u is of p.e. at \mathbf{r} , it follows from [Lon08, Thm. 3.3.1], the Bisognano-Wichmann property of such nets, that $\delta_I^{it} = u(e^{-2\pi t \xi_I})$. Passing to the second quantization, let $\mathcal{N}_I := \rho(N_I)'' = W(V(I)^\infty)''$ denote the von Neumann algebra generated by $W(V(I)^\infty)$. By Example 5.19, we obtain that $\rho(e^{-2\pi t \xi_I}) = \mathcal{F}(u(e^{-2\pi t \xi_I})) = \mathcal{F}(\delta_I^{it}) = \Delta_I^{it}$, where Δ_I denotes the modular operator on $\mathcal{F}(V_{\mathcal{J}})$ defined from \mathcal{N}_I using the cyclic and separating vector $\Omega := 1 \in \mathcal{F}(V_{\mathcal{J}})$. Let $\phi = \langle \Omega, \cdot \Omega \rangle$ be the corresponding state on \mathcal{N}_I . Then (9) shows that $\widehat{\phi} : N_I \rightarrow \mathbb{C}$ is smooth. Thus ρ is smoothly-KMS at $2\pi \xi_I \in \mathfrak{su}(1, 1)$ relative to $N_I := H(V_I^\infty, \omega) \subseteq G$, for any $I \in \mathcal{I}$. For more details on the Bisognano-Wichmann property, see e.g. [Mor18] and [Mun01].

Part II

Generalized Positive Energy Representations of Jet Lie Groups and Algebras

We now depart from the general context of Part I. Using the observations made in Part I, we study projective unitary representations of jet Lie groups and algebras that are of generalized positive energy. Let us first fix our notation, which is kept throughout Part II.

6 Notation

Let V be a finite-dimensional real vector space and K a 1-connected compact simple Lie group with Lie algebra \mathfrak{k} . For any $n \in \mathbb{N}_{\geq 0}$, we denote by $P^n(V) \subseteq R$ the space of homogeneous polynomials on V of degree n . Let $R := \mathbb{R}[[V^*]] := \prod_{n=0}^{\infty} P^n(V)$ denote the ring of formal power series on V with coefficients in \mathbb{R} , equipped with the direct product topology. Let $I = (V^*)$ be the maximal ideal of R , containing those elements with vanishing constant term. We write $\text{ev}_0 : R \rightarrow \mathbb{R} \cong R/I$ for the corresponding quotient map. Let \mathfrak{g} be the R -module $\mathfrak{g} := R \otimes \mathfrak{k}$ of formal power series on V with coefficients in \mathfrak{k} . Then \mathfrak{g} is a topological Lie algebra with the Lie bracket defined by

$$[f \otimes X, g \otimes Y] := fg \otimes [X, Y], \quad f, g \in R, \quad X, Y \in \mathfrak{k}.$$

We also write fX instead of $f \otimes X$ for $f \in R$ and $X \in \mathfrak{k}$. Define $R_k := R/I^{k+1}$, $I_k := I/I^{k+1}$ and $\mathfrak{g}_k := \mathfrak{g}/(I^{k+1}\mathfrak{g})$ for $k \in \mathbb{N}_{\geq 0}$. Then $R = \varprojlim R_k$ and $\mathfrak{g} = \varprojlim \mathfrak{g}_k$ as topological vector spaces and Lie algebras, respectively. For $k \in \mathbb{N}_{\geq 0}$, let $G_k = J_0^k(V; K)$ be the unique 1-connected Lie group integrating the finite-dimensional Lie algebra \mathfrak{g}_k . Let $G := J_0^\infty(V; K) := \varprojlim G_k$ be the corresponding projective limit, which is a pro-Lie group with topological Lie algebra $\mathfrak{g} = \varprojlim \mathfrak{g}_k$. (See e.g. [HM07] for a detailed consideration of pro-Lie groups). Write \mathcal{X}_I for the Lie algebra of formal vector fields on V vanishing at the origin. Identify $\mathcal{X}_I \cong \text{der}(I)$ using the Lie derivative $\mathfrak{v} \mapsto \mathcal{L}_{\mathfrak{v}}$. Notice further that $\text{der}(I) \cong I \otimes V$. Define similarly $\mathcal{X}_{I_k} := \mathcal{X}_I/(I^{k+1}\mathcal{X}_I) \cong \text{der}(I_k)$.

Let \mathfrak{p} be a finite-dimensional Lie algebra acting on \mathfrak{g} by the homomorphism $D : \mathfrak{p} \rightarrow \text{der}(\mathfrak{g})$. Using the fact that all derivations of \mathfrak{k} are inner, by Whitehead's first Lemma [Jac79, III.7. Lem. 3], it follows from [Kac90b, Ex. 7.4] that $D(p)$ splits into a horizontal and vertical part according to $D(p) = -\mathcal{L}_{\mathfrak{v}(p)} + \text{ad}_{\sigma(p)}$, where $\mathfrak{v} : \mathfrak{p} \rightarrow \mathcal{X}_I^{\text{op}}$ is a homomorphism of Lie algebras and where $\sigma : \mathfrak{p} \rightarrow \mathfrak{g}$ is a linear map that necessarily satisfies the following Maurer-Cartan equation:

$$-\mathcal{L}_{\mathfrak{v}(p_1)}\sigma(p_2) + \mathcal{L}_{\mathfrak{v}(p_2)}\sigma(p_1) - \sigma([p_1, p_2]) + [\sigma(p_1), \sigma(p_2)] = 0, \quad \forall p_1, p_2 \in \mathfrak{p}. \quad (10)$$

Remark 6.1. As we shall see in Section 7.3 below, Equation (10) can be written as $\delta\sigma + \frac{1}{2}[\sigma, \sigma] = 0$ in the differential graded Lie algebra $(\bigwedge^\bullet \mathfrak{p}^*) \otimes \mathfrak{g}$, whose differential is that of the Chevalley-Eilenberg complex, where \mathfrak{g} is considered as \mathfrak{p} -module according to $p \mapsto -\mathcal{L}_{\mathfrak{v}(p)}$.

We will refer to D as a *lift* of the \mathfrak{p} -action on R to \mathfrak{g} and we call σ the *vertical twist* of the lift D . We remark also that $D(p)$ satisfies the following Leibniz rule:

$$D(p)(f\xi) = -\mathcal{L}_{\mathfrak{v}(p)}(f)\xi + fD(p)\xi, \quad f \in R, \xi \in \mathfrak{g}, \quad \forall p \in \mathfrak{p}. \quad (11)$$

We will denote by j^k various k -jet projections $R \rightarrow R_k$, $\mathfrak{g} \rightarrow \mathfrak{g}_k$ and $\mathcal{X}_I \rightarrow \mathcal{X}_{I_k}$. It should be clear from the context which map is being used. Also, we will freely identify the quotient $\mathfrak{g}_0 \cong \mathfrak{k}$ with the Lie subalgebra $\mathfrak{k} \subseteq \mathfrak{g}$ of formal power series having only a non-trivial constant term. Similarly, we identify $j^1\mathcal{X}_I = \mathcal{X}_{I_1} \cong \mathfrak{gl}(V)$ with the subalgebra $\mathfrak{gl}(V) \subseteq \mathcal{X}_I$ of linear vector fields on V .

A first observation is the fact that $G = J_0^\infty(V; K)$ is not just a pro-Lie group, but actually a regular Lie group modeled on the Fréchet space $\mathfrak{g} = J_0^\infty(V; \mathfrak{k})$.

Proposition 6.2. *Both G and $G \rtimes_\alpha P$ are regular BCH Fréchet-Lie groups.*

Proof. It is clear that the Lie algebras \mathfrak{g} and $\mathfrak{g} \rtimes_D \mathfrak{p}$ are Fréchet. For every $n \in \mathbb{N}$, the Lie group $G_n = J_0^n(V; K)$ is 1-connected, because K is so. Then also G is 1-connected, since $\pi_k(G) = \pi_k(\varprojlim_n G_n) = \varprojlim_n \pi_k(G_n)$ for

every $k \in \mathbb{N}$, see e.g. [Hat02, Prop. 4.67]. Thus G is the unique 1-connected pro-Lie group with $\text{Lie}(G) = \mathfrak{g}$, which is locally contractible by [HN09, Theorem 1.2]. Then [HN09, Theorem 1.3, Prop. 5.7] entails that G is a regular Lie group. As the action $\alpha : P \times G \rightarrow G$ is smooth, also $G \rtimes_{\alpha} P$ is a Lie group and it is regular by [Nee06, Thm. V.I.8], because both G and P are so. Both G and $G \rtimes_{\alpha} P$ are BCH because they carry the projective limit topology and all their finite-dimensional quotients are BCH. \square

Moreover, we have the following useful fact:

Lemma 6.3. *The exponential map $\exp_G : \mathfrak{g} \rightarrow G$ restricts to a diffeomorphism from $I \otimes \mathfrak{k} = \ker(\text{ev}_0 : \mathfrak{g} \rightarrow \mathfrak{k})$ onto $\ker(\text{ev}_0 : G \rightarrow K)$.*

Proof. For $k \in \mathbb{N}_{\geq 0}$, let $H_k := \ker(\text{ev}_0 : G_k \rightarrow K) \triangleleft G_k$ be the maximal nilpotent normal subgroup of G_k . Then $\mathfrak{h}_k := \text{Lie}(H_k) = \ker(\text{ev}_0 : \mathfrak{g}_k \rightarrow \mathfrak{k}) = (I_k \otimes \mathfrak{k}) \triangleleft \mathfrak{g}_k$. Write $H := \varprojlim_k H_k$ for the corresponding normal subgroup of G and $\mathfrak{h} = \varprojlim_k \mathfrak{h}_k$ for its Lie algebra. Let $k \in \mathbb{N}$. Notice that H_k is nilpotent and 1-connected. Consequently, its exponential map is a diffeomorphism $\exp_{H_k} : \mathfrak{h}_k \rightarrow H_k$ [CG90, Thm. 1.2.1]. Write $\log_{H_k} : H_k \rightarrow \mathfrak{h}_k$ for its inverse. If $m \geq k$, then $\exp_{H_k} \circ j^k = j^k \circ \exp_{H_m} : \mathfrak{h}_m \rightarrow H_k$ and consequently $\log_{H_k} \circ j^k = j^k \circ \log_{H_m} : H_m \rightarrow \mathfrak{h}_k$. Passing to the projective limit, we obtain the inverse $\log_H := \varprojlim_k \log_{H_k}$ of \exp_H . It is smooth because $H = \varprojlim_k H_k$ carries the projective limit topology and \log_{H_k} is smooth for every $k \in \mathbb{N}$. Thus $\exp_H : \mathfrak{h} \rightarrow H$ is a global diffeomorphism. \square

7 Normal Form Results

By choosing suitable local coordinates, one may attempt to simplify the vector fields $\mathbf{v}(p)$ and the vertical twist $\sigma(p)$ of the lift $D(p) = -\mathcal{L}_{\mathbf{v}(p)} + \text{ad}_{\sigma(p)}$ simultaneously. One might for example try to show that there are local coordinates in which the formal vector fields $\mathbf{v}(p)$ are linear for every $p \in \mathfrak{p}$ simultaneously, thereby linearizing the formal \mathfrak{p} -action. Similarly, one might aim to show that in suitable coordinates, $\sigma(p) \in \mathfrak{k} \subseteq R \otimes \mathfrak{k}$ is constant for all $p \in \mathfrak{p}$, so that σ is a Lie algebra homomorphism $\mathfrak{p} \rightarrow \mathfrak{k}$. In the following, this ‘normal form problem’ is considered. The results of Section 8 will depend on the availability of suitable normal forms, whose existence we study in the present section.

In Section 7.1, we briefly recall the transformation behavior of \mathbf{v} and σ under suitable automorphisms of \mathfrak{g} . We proceed in Section 7.2 to recollect some known results regarding normal forms for Lie algebras of vector fields with a common fixed point. Finally, we consider in Section 7.3 the vertical twist σ .

7.1 Transformation Behavior

Definition 7.1.

- A *formal diffeomorphism of V* is an automorphism h of R . An automorphism of \mathfrak{g} is said to be *horizontal* if it is of the form $h \otimes \text{id}_{\mathfrak{k}}$ for some $h \in \text{Aut}(R)$. We write $h.\xi$ or $h(\xi)$ instead of $(h \otimes \text{id}_{\mathfrak{k}})(\xi)$ for $\xi \in \mathfrak{g}$.
- A *gauge transformation* is an automorphism of \mathfrak{g} of the form $e^{\text{ad}_{\xi}}$ for some $\xi \in \mathfrak{g}$.

Remark 7.2. Any formal diffeomorphism $h \in \text{Aut}(R)$ preserves the maximal proper ideal I and is determined by its restriction $h|_{V^*}$, which can be regarded as an element of $I \otimes V$ for which $j^1 \tilde{h} \in V^* \otimes V \cong \mathfrak{gl}(V)$ is invertible. It is then a consequence of Borel’s Lemma [H03, Thm. 1.2.6] and the Inverse Function Theorem that for any automorphism h of R , there exist 0-neighborhoods $U, U' \subseteq V$ and a diffeomorphism $h_0 : U \rightarrow U'$ satisfying $h_0(0) = 0$ such that $h(j_0^{\infty}(f)) = j_0^{\infty}(f \circ h_0^{-1})$. Similarly, for $\xi \in \mathfrak{g}$ there exists $\eta \in C_c^{\infty}(V, \mathfrak{k})$ s.t. $j_0^{\infty}(\eta) = \xi$, where we have identified $\mathfrak{g} \cong J_0^{\infty}(V; \mathfrak{k})$. We then have $e^{\text{ad}_{\xi}} \circ j_0^{\infty} = j_0^{\infty} \circ e^{\text{ad}_{\eta}}$.

To determine the transformation behavior of $D : \mathfrak{p} \rightarrow \text{der}(\mathfrak{g})$, we have to consider the adjoint action of $\text{Aut}(\mathfrak{g})$ on $\text{der}(\mathfrak{g})$. Instead of considering arbitrary automorphisms of \mathfrak{g} , we will specialize to horizontal ones and to gauge transformations. For $h \in \text{Aut}(R)$ and $v \in \mathcal{X}_I^{op}$, we write $h.v$ for the action of $\text{Aut}(R)$ on \mathcal{X}_I^{op} obtained from the adjoint action of $\text{Aut}(R)$ on $\text{der}(R) \cong \mathcal{X}_I \cong \mathcal{X}_I^{op}$. The following two proofs are due to K.H. Neeb and B. Janssens. They appear in the presently unpublished article [JN].

Lemma 7.3 ([JN]). *Let $D \in \text{der}(\mathfrak{g})$ and $\xi \in \mathfrak{g}$. Then*

$$e^{\text{ad}_{\xi}} \circ D \circ e^{-\text{ad}_{\xi}} = D + \text{ad}(F(\text{ad}_{\xi})D\xi),$$

where $F(w) = -\int_0^1 e^{tw} dt = -\sum_{n=0}^{\infty} \frac{1}{(n+1)!} w^n$.

Proof. Let $k \in \mathbb{N}$ be arbitrary. Consider the continuous path $\gamma : I \rightarrow \text{der}(\mathfrak{g})$ defined by $\gamma(t) = e^{t\text{ad}_\xi} D e^{-t\text{ad}_\xi}$. Notice that $j^k \circ \gamma : I \rightarrow \text{der}(\mathfrak{g}_k)$ is smooth for all k and consequently so is γ . Moreover

$$\gamma'(t) = e^{t\text{ad}_\xi} [\text{ad}_\xi, D] e^{-t\text{ad}_\xi} = -e^{t\text{ad}_\xi} \text{ad}_{D\xi} e^{-t\text{ad}_\xi} = -\text{ad}(e^{t\text{ad}_\xi} D\xi),$$

where the last step uses that $\alpha \circ \text{ad}_\eta = \text{ad}_{\alpha(\eta)} \circ \alpha$ for any $\alpha \in \text{Aut}(\mathfrak{g})$. Thus

$$e^{\text{ad}_\xi} \circ D \circ e^{-\text{ad}_\xi} - D = \int_0^1 \gamma'(t) dt = - \int_0^1 \text{ad}(e^{t\text{ad}_\xi} D\xi) dt = -\text{ad}\left(\int_0^1 e^{t\text{ad}_\xi} dt\right)(D\xi) = \text{ad}(F(\text{ad}_\xi)D\xi). \quad \square$$

Proposition 7.4 ([JN]). *Let $h \in \text{Aut}(R) \subseteq \text{Aut}(\mathfrak{g})$, $\sigma, \xi \in \mathfrak{g}$ and $v \in \mathcal{X}_I$. Consider the derivation $D := -\mathcal{L}_v + \text{ad}_\sigma \in \text{der}(\mathfrak{g})$. Then*

$$\begin{aligned} h \circ D \circ h^{-1} &= -\mathcal{L}_{h.v} + \text{ad}(h.\sigma), \\ e^{\text{ad}_\xi} \circ D \circ e^{-\text{ad}_\xi} &= -\mathcal{L}_v + \text{ad}\left(e^{\text{ad}_\xi} \sigma + F(\text{ad}_\xi)(-\mathcal{L}_v \xi)\right). \end{aligned} \quad (12)$$

Proof. It is trivial that $h \circ \mathcal{L}_v \circ h^{-1} = \mathcal{L}_{h.v}$. Moreover, $h \circ \text{ad}_\sigma \circ h^{-1} = \text{ad}_{h.\sigma}$ is valid because $\alpha \circ \text{ad}_\sigma = \text{ad}_{\alpha(\sigma)} \circ \alpha$ for any $\alpha \in \text{Aut}(\mathfrak{g})$. Notice next that $F(\text{ad}_\xi)([\sigma, \xi]) = \sum_{n=1}^{\infty} \frac{1}{n!} \text{ad}_\xi^n \sigma = e^{\text{ad}_\xi} \sigma - \sigma$. It follows from Lemma 7.3:

$$h \circ D \circ h^{-1} = -\mathcal{L}_v + \text{ad}(\sigma) + \text{ad}(F(\text{ad}_\xi)(-\mathcal{L}_v \xi)) + \text{ad}(F(\text{ad}_\xi)[\sigma, \xi]) = -\mathcal{L}_v + \text{ad}\left(e^{\text{ad}_\xi} \sigma + F(\text{ad}_\xi)(-\mathcal{L}_v \xi)\right). \quad \square$$

Definition 7.5.

- Two homomorphisms $\mathbf{v}, \mathbf{w} : \mathfrak{p} \rightarrow \mathcal{X}_I^{op}$ are said to be *formally-equivalent* if there is a formal diffeomorphism $h \in \text{Aut}(R)$ such that $h.v(p) = \mathbf{w}(p)$ for all $p \in \mathfrak{p}$.
- Two linear maps $\sigma, \eta : \mathfrak{p} \rightarrow R \otimes \mathfrak{k}$ satisfying the Maurer-Cartan equation (10) are called *gauge-equivalent* if there is some $\xi \in \mathfrak{g}$ such that

$$\eta(p) = e^{\text{ad}_\xi} \sigma(p) + F(\text{ad}_\xi)(-\mathcal{L}_{\mathbf{v}(p)} \xi), \quad \forall p \in \mathfrak{p}. \quad (13)$$

In this case, we write $\sigma \sim \eta$ and say that σ and η are related by the gauge transformation e^{ad_ξ} .

7.2 Lie Algebras of Formal Vector Fields with a Common Fixed Point

The normal form problem for vector fields near a fixed point has been subject to extensive study. Let us first gather some relevant known results.

The case of a single vector field

Naturally, the special case which has been considered most is the case where \mathfrak{p} is simply \mathbb{R} , in which case one is looking for normal forms of dynamical systems near a fixed point, in the formal context. This case is already quite interesting. Let us recollect some relevant results. For more information, we refer to [Arn88].

Let \mathbf{v} be a vector field on V . Write $\mathbf{v} = \mathbf{v}_1 + \mathbf{v}_{\text{ho}}$, where $\mathbf{v}_1 = j_0^1(\mathbf{v}) \in \mathfrak{gl}(V) \subseteq \mathcal{X}_I$ is the linearization of \mathbf{v} at $0 \in V$ and $\mathbf{v}_{\text{ho}} \in \mathcal{X}_{I^2}$ is a vector field vanishing up to first order at $0 \in V$. Let $\mathbf{v}_1 = \mathbf{v}_{1,s} + \mathbf{v}_{1,n}$ be the Jordan decomposition of \mathbf{v}_1 over \mathbb{C} , where $\mathbf{v}_{1,s}$ is semisimple and $\mathbf{v}_{1,n}$ is nilpotent. Write $V_{\mathbb{C}} := V \otimes_{\mathbb{R}} \mathbb{C}$. Let $(e_j)_{j=1}^d$ be a basis of eigenvectors of $\mathbf{v}_{1,s}$ in $V_{\mathbb{C}}$ with dual basis $(x_j)_{j=1}^d$ of V^* . Let $(\mu_j)_{j=1}^d$ denote the corresponding eigenvalues.

Definition 7.6. Let $\mathbf{n} \in \mathbb{N}_{\geq 0}^d$ be a multi-index. A monomial vector field $x^{\mathbf{n}} \partial_{x_j}$ with $|\mathbf{n}| \geq 2$ is called *resonant* if $\langle \mathbf{n}, \boldsymbol{\mu} \rangle = \mu_j$, where $\langle \mathbf{n}, \boldsymbol{\mu} \rangle := \sum_{i=1}^d n_i \mu_i$. Identifying $\mathbf{v}_{1,s}$ with the linear vector field $\sum_{j=1}^d \mu_j x_j \partial_{x_j}$ on \mathbb{C}^d , this is equivalent to $[\mathbf{v}_{1,s}, x^{\mathbf{n}} \partial_{x_j}] = 0$.

Theorem 7.7 (Poincaré-Dulac Theorem [Dul04] [Arn88, ch.5]).

There exists $\mathbf{w}_{\text{ho}} \in \mathcal{X}_{I^2}$ which is a \mathbb{C} -linear combination of resonant monomials s.t. \mathbf{v} is formally equivalent to $\mathbf{w} = \mathbf{v}_1 + \mathbf{w}_{\text{ho}} \in \mathcal{X}_I$. In particular $[\mathbf{v}_{1,s}, \mathbf{w}_{\text{ho}}] = 0$ in \mathcal{X}_I .

Corollary 7.8 (Poincaré [Poi79]). *If there are no resonances, that is to say, if $\langle \mathbf{n}, \boldsymbol{\mu} \rangle \neq \mu_j$ for all $\mathbf{n} \in \mathbb{N}^d$ and $j \in \{1, \dots, d\}$, then the vector field \mathbf{v} can be formally linearized, so that \mathbf{v} is formally equivalent to the linear vector field \mathbf{v}_1 .*

The case of actions by a compact Lie group

For actions of compact Lie groups there is the following well-known result, see also [DK00, Ch. 2.2].

Theorem 7.9 (Bochner's Linearization Theorem [Boc45]).

Let $G \times M \rightarrow M$ be a smooth action of compact Lie group on a smooth manifold which has a fixed point $a \in M$. Then, in suitably chosen smooth local coordinates around the fixed point, the action is linear.

The case of actions by semisimple Lie algebras

Next, we move to Lie algebra representations by formal vector fields of semisimple Lie algebras. As nicely explained in [FM04] and first observed by Hermann in [Her68], in the formal setting the obstructions to being able to linearize a Lie algebra of vector fields simultaneously lie in various first Lie algebra cohomology groups $H^1(\mathfrak{p}, W)$ for suitable finite-dimensional \mathfrak{p} -modules W . In view of Whitehead's First Lemma [Jac79, III.7. Lem. 3], this results in:

Theorem 7.10 ([Her68]). *Let \mathfrak{p} be a semisimple Lie algebra and $\mathbf{v} : \mathfrak{p} \rightarrow \mathcal{X}_I^{op}$ be a Lie algebra homomorphism. Then \mathbf{v} is formally equivalent to its linearization $j^1\mathbf{v} : \mathfrak{p} \rightarrow \mathfrak{gl}(V) \subseteq \mathcal{X}_I^{op}$ around the origin.*

Remark 7.11. Corresponding statements of Theorem 7.10 in the setting of germs of smooth/analytic vector fields and diffeomorphisms have been proven in [GS68] and [FM04] under additional assumptions. They are false in general without suitable extra conditions, as was shown in [GS68].

7.3 Normal Form Results for the Vertical Twist

Let us next consider the vertical twist $\sigma : \mathfrak{p} \rightarrow \mathfrak{g}$ of the lift $D(p) = -\mathcal{L}_{\mathbf{v}(p)} + \text{ad}_{\sigma(p)}$ to \mathfrak{g} of the \mathfrak{p} -action $-\mathcal{L}_{\mathbf{v}(p)}$ on R , which has to satisfy the Maurer Cartan equation (10). We fix the horizontal part $\mathbf{v} : \mathfrak{p} \rightarrow \mathcal{X}_I^{op}$ and act by gauge transformations. The main results of this section are the following two theorems, whose proof comprises the remainder of the section. The reader who is eager to consider the projective unitary g.p.e. representations of \mathfrak{g} can proceed to Section 8 after reading Theorem 7.12 and Theorem 7.13 below.

Let us also remark that the methods used in this section to prove Theorem 7.12 and Theorem 7.13 were communicated to the author by B. Janssens and K.H. Neeb and appear in similar form in their presently unpublished work [JN], albeit in a more specific context. The author has placed their approach to a more general context and extracted the two theorems below.

Theorem 7.12. *Assume that \mathfrak{p} is semisimple. Let the linear map $\sigma : \mathfrak{p} \rightarrow \mathfrak{g}$ satisfy the Maurer-Cartan equation (10). Then σ is gauge-equivalent to $\sigma_0 := \text{ev}_0 \circ \sigma : \mathfrak{p} \rightarrow \mathfrak{k}$. If \mathfrak{p} has no non-trivial compact ideals, then σ is gauge-equivalent to 0.*

The next result concerns the case $\mathfrak{p} = \mathbb{R}$, in which case we identify \mathbf{v} with $\mathbf{v}(1) \in \mathcal{X}_I$ and σ with $\sigma(1) \in \mathfrak{g}$. In this case, the Maurer-Cartan equation (10) is trivially satisfied for any $\sigma \in \mathfrak{g}$ and $\mathbf{v} \in \mathcal{X}_I$.

Theorem 7.13. *Assume that $\mathfrak{p} = \mathbb{R}$. Let $\sigma \in \mathfrak{g}$ and $\mathbf{v} \in \mathcal{X}_I$. Let $\mathbf{v}_1 := j^1\mathbf{v} \in \mathfrak{gl}(V)$ be the linearization of \mathbf{v} at $0 \in V$. Assume w.l.o.g. that $\sigma_0 := \text{ev}_0(\sigma) \in \mathfrak{t}$ for some maximal torus $\mathfrak{t} \subseteq \mathfrak{k}$. The following assertions hold:*

1. *Assume that $\langle \mathbf{n}, \boldsymbol{\mu} \rangle \neq \alpha(\sigma_0)$ for any root $\alpha \in i\mathfrak{t}^*$ of \mathfrak{k} and $\mathbf{n} \in \mathbb{N}_{\geq}^d$ with $|\mathbf{n}| \geq 1$. Then σ is gauge-equivalent to some $\sigma' \in R \otimes \mathfrak{t}$.*
2. *If \mathbf{v}_1 is semisimple, then σ is gauge-equivalent to some $\nu \in R \otimes \mathfrak{k}$ satisfying $-\mathcal{L}_{\mathbf{v}_1}\nu + [\sigma_0, \nu] = 0$.*
3. *Suppose that $\mathbf{v} = \mathbf{v}_1$ is linear. Assume that $D = -\mathcal{L}_{\mathbf{v}} + \text{ad}_{\sigma}$ integrates to continuous $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ -action on \mathfrak{g} . Then σ is gauge-equivalent to $\sigma_0 \in \mathfrak{t}$. Moreover $\text{Spec}(\mathbf{v}_1) \cup \text{Spec}(\text{ad}_{\sigma}) \subseteq i\mathbb{Z}$.*

Remark 7.14. Suppose that $\mathcal{K} \rightarrow M$ is a principal fiber bundle with compact simple structure group K . Let $\alpha : \mathbb{T} \rightarrow \text{Aut}(\mathcal{K})$ be a smooth action on \mathcal{K} by bundle automorphisms. Suppose that $a \in M$ is a fixed point of the induced \mathbb{T} -action on M and set $V := T_a(M)$. By Theorem 7.9, the \mathbb{T} -action on M is linear in suitable local coordinates around $a \in M$. Passing to $J_a^\infty(M) \cong R$ and $J_a^\infty(\text{Ad}(\mathfrak{K})) \cong \mathfrak{g}$, one obtains a \mathbb{T} -action on both R and \mathfrak{g} . The corresponding derivations are given by $-\mathcal{L}_{\mathbf{v}}$ and $D = -\mathcal{L}_{\mathbf{v}} + \text{ad}_{\sigma}$ respectively, for some linear semisimple vector field \mathbf{v} on V and some $\sigma \in \mathfrak{g}$. This is the setting of the third item in Theorem 7.13, according to which we may further assume that $\sigma \in \mathfrak{t}$, where $\mathfrak{t} \subseteq \mathfrak{k}$ is a maximal torus, by acting with gauge transformations.

The remainder of this section is devoted to the proof of Theorem 7.12 and Theorem 7.13.

Reformulation Using Differential Graded Lie Algebras

In order to classify the equivalence classes of vertical twists $\sigma : \mathfrak{p} \rightarrow \mathfrak{g}$, we interpret equation (10) as the Maurer Cartan equation in the differential graded Lie algebra (DGLA) $L := L_R := (\bigwedge^\bullet \mathfrak{p}^*) \otimes \mathfrak{g}$. As a cochain complex, L is the Chevalley-Eilenberg complex associated to the \mathfrak{p} -module \mathfrak{g} , where \mathfrak{p} acts on \mathfrak{g} by $p.\psi = -\mathcal{L}_{v(p)}\psi$. Explicitly, the differential δ is given by

$$\delta(\alpha)(p_1, \dots, p_{k+1}) = \sum_i (-1)^{i+1} p_i \alpha(p_1, \dots, \widehat{p}_i, \dots, p_{k+1}) + \sum_{i < j} (-1)^{i+j} \alpha([p_i, p_j], p_1, \dots, \widehat{p}_i, \dots, \widehat{p}_j, \dots, p_{k+1}),$$

where as usual, the arguments with a caret are to be omitted. The graded Lie bracket on L is the unique bilinear map $[-, -] : L \times L \rightarrow L$ satisfying $[\alpha \otimes \sigma, \beta \otimes \psi] := (\alpha \wedge \beta) \otimes [\sigma, \psi]$ for $\alpha, \beta \in \bigwedge^\bullet \mathfrak{p}^*$ and $\sigma, \psi \in \mathfrak{g}$. Write $L^k := (\bigwedge^k \mathfrak{p}^*) \otimes \mathfrak{g}$ for the degree k -elements in L . Interpreting σ as a degree-1 element of L , equation (10) can now equivalently be written as the usual MC-equation $\delta\sigma + \frac{1}{2}[\sigma, \sigma] = 0$ in L .

Let us next reformulate the gauge-action (14) of \mathfrak{g} on the set of vertical twists, using the DGLA L . Consider the extended DGLA $L \rtimes \mathbb{R}D$, where D is a degree-1 element satisfying $[D, \sigma] = \delta(\sigma)$ for any $\sigma \in L$. Notice for $\xi \in \mathfrak{g} = L^0$ that $\delta(\xi)(p) = -\mathcal{L}_{v(p)}\xi$. We define the gauge-action of $L^0 = \mathfrak{g}$ on L by

$$\xi.\sigma = e^{\text{ad}(\xi)}(D + \sigma) - D = e^{\text{ad}(\xi)}(\sigma) + F(\text{ad}_\xi)(\delta(\xi)), \quad \xi \in \mathfrak{g}, \quad (14)$$

considered as an expression in $L \rtimes \mathbb{R}D$, where $F(w) = -\sum_{n=0}^{\infty} \frac{1}{(n+1)!} w^n = -\int_0^1 e^{tw} dt$. Let us check that the above is indeed well-defined, even though L is not a nilpotent DGLA. Since $G = \varprojlim_k G_k$ is a Lie group, it has an exponential map and so the automorphism $e^{\text{ad}_\xi} := \text{Ad}(e^\xi)$ on \mathfrak{g} is defined. Consequently, so is

$$F(\text{ad}_\xi)(-\mathcal{L}_{v(p)}\xi) = \int_0^1 e^{t\text{ad}_\xi}(\mathcal{L}_{v(p)}\xi) dt$$

for any $p \in \mathfrak{p}$. Thus the expression in equation (14) makes sense. Notice further that for $\sigma \in L^1$, the above reduces precisely to the transformation behavior (14) of the vertical twist. In accordance with Definition 7.5, we say that the MC-elements $\sigma, \sigma' \in L^1$ are gauge-equivalent if they satisfy $\sigma' = \xi.\sigma$ for some $\xi \in L^0$, in which case we write $\sigma \sim \sigma'$. Our goal is to study the MC-elements in L^1 up to gauge-equivalence.

Let $n \in \mathbb{N}_{\geq 0}$. Define analogously the following DGLAs, where we consider $P^n(V)$ as \mathfrak{p} -module by identifying $P^n(V)$ with I^n/I^{n+1} for $n \in \mathbb{N}_{\geq 0}$, so that $p.f = -\mathcal{L}_{v_1(p)}f$ for $p \in \mathfrak{p}$ and $f \in P^n(V)$:

$$\begin{aligned} L_I &:= \bigwedge^\bullet \mathfrak{p}^* \otimes (I \otimes \mathfrak{k}), & L_{R_n} &:= \bigwedge^\bullet \mathfrak{p}^* \otimes (R_n \otimes \mathfrak{k}), \\ L_{I_n} &:= \bigwedge^\bullet \mathfrak{p}^* \otimes (I_n \otimes \mathfrak{k}), & L_{P^n} &:= \bigwedge^\bullet \mathfrak{p}^* \otimes (P^n(V) \otimes \mathfrak{k}). \end{aligned}$$

Shifted DGLAs

It will be beneficial to split off the constants terms of the \mathfrak{k} -valued formal power series, because contrary to L_R , L_I is a projective limit of *nilpotent* DGLAs. We discuss next how this can be done.

For any MC-element $\chi \in L_{R_0}^1 = (\bigwedge^\bullet \mathfrak{p}^*) \otimes \mathfrak{k} \subseteq L_R$ of degree 1, define the "shifted" DGLA L_R^χ , which agrees with L_R as a graded Lie algebra but has a shifted differential given by $\delta_\chi(\sigma) := \delta(\sigma) + [\chi, \sigma]$. The differential δ_χ agrees with the Chevalley-Eilenberg differential of $(\bigwedge^\bullet \mathfrak{p}^*) \otimes \mathfrak{g}$ if \mathfrak{g} is considered as \mathfrak{p} -module with the twisted action $p.\sigma := -\mathcal{L}_{v(p)}\sigma + [\chi, \sigma]$. In particular, $\delta_\chi^2 = 0$. Let us write $R \otimes_\chi \mathfrak{k}$ for this module structure to distinguish it from the usual one on $\mathfrak{g} = R \otimes \mathfrak{k}$, which was given by $p.\xi = -\mathcal{L}_{v(p)}\xi$. Define also the extended DGLA $L_R^\chi \rtimes \mathbb{R}D_\chi$, where $[D_\chi, \sigma] = \delta_\chi(\sigma)$. Define in analogous fashion $L_{I_n}^\chi$, L_I^χ and $L_{P^n}^\chi$, where we have used that the \mathfrak{p} -action on $R \otimes_\chi \mathfrak{k}$ leaves $I_n \otimes \mathfrak{k}$ invariant for every n , so that $P^n(V) \otimes \mathfrak{k} \cong (I_n \otimes \mathfrak{k}) / (I_{n+1} \otimes \mathfrak{k})$ is naturally a \mathfrak{p} -module. The following is a standard result:

Lemma 7.15.

1. Let $\sigma \in L_R^1$. Then $\chi + \sigma$ is a MC-element in L_R if and only if σ is a MC-element in L_R^χ .
2. Let $\sigma, \sigma' \in L_R^\chi$ be degree-1 MC-elements. Then $\chi + \sigma \sim \chi + \sigma'$ in L_R if and only if $\sigma \sim \sigma'$ in L_R^χ .
3. Let $\psi \in L_R^1$ be a MC-element. Then $\psi = \chi + \sigma$ for some degree-1 MC-elements $\sigma \in L_I^\chi$ and $\chi \in L_{R_0}$.

Proof.

1. As χ is a MC-element and $[\sigma, \chi] = [\chi, \sigma]$ we have

$$\delta(\chi + \sigma) + \frac{1}{2}[\chi + \sigma, \chi + \sigma] = \delta(\sigma) + \frac{1}{2}[\sigma, \sigma] + [\chi, \sigma] = \delta_\chi(\sigma) + \frac{1}{2}[\sigma, \sigma].$$

2. Observe that $-F(\text{ad}_\xi)([\xi, \chi]) = e^{\text{ad}_\xi}(\chi) - \chi$. Consequently

$$F(\text{ad}_\xi)(\delta_\chi(\xi)) = F(\text{ad}_\xi)(\delta(\xi)) + F(\text{ad}_\xi)([\chi, \xi]) = F(\text{ad}_\xi)(\delta(\xi)) + e^{\text{ad}_\xi}(\chi) - \chi. \quad (15)$$

Thus, for any $\xi \in \mathfrak{g}$ we have

$$e^{\text{ad}_\xi}(\chi + \sigma) + F(\text{ad}_\xi)(\delta(\xi)) = \chi + \left(e^{\text{ad}_\xi}(\sigma) + F(\text{ad}_\xi)(\delta_\chi(\xi)) \right).$$

3. Since $R = R_0 \oplus I$ as vector space, we can write $\psi = \chi + \sigma$, where $\chi = j^0(\psi) \in L_{R_0}$ and $\sigma \in L_I^\chi$. As j^0 is a morphism of DGLAs, it is clear that $\chi = j^0(\psi)$ is a MC-element in $L_{R_0} \subseteq L_R$. By the first point it follows that σ is a MC-element in $L_I^\chi \subseteq L_R^\chi$. \square

Study of MC-Elements

In view of Lemma 7.15, let us first study the classification problem of gauge-orbits of MC-elements in $L_{R_0}^1$ and then, for each MC-element $\chi \in L_{R_0}^1$ consider the orbits in L_I^χ under the gauge-action.

Lemma 7.16.

1. Let $\chi : \mathfrak{p} \rightarrow \mathfrak{k}$ be linear. Then χ is a MC-element in $L_{R_0}^1$ if and only if it is a Lie algebra homomorphism. Thus if there are no homomorphisms $\mathfrak{p} \rightarrow \mathfrak{k}$, then any MC-element $\chi \in L_{R_0}^1 = \mathfrak{p}^* \otimes \mathfrak{k}$ is trivial.
2. The gauge-action of $X \in \mathfrak{k} = L_{R_0}^0$ on L_{R_0} is given by $X \cdot \chi = e^X \chi$.

Proof. Notice that $\delta(X) = 0$ for any $X \in \mathfrak{k} \subseteq \mathfrak{g}$, because $-\mathcal{L}_{v(p)}X = 0$ for any $p \in \mathfrak{p}$. So \mathfrak{p} acts trivially on $\mathfrak{k} = \mathfrak{g}_0 = j^0\mathfrak{g}$. Thus the Maurer-Cartan condition reads simply $\chi([p_1, p_2]) - [\chi(p_1), \chi(p_2)] = 0$ for all $p_1, p_2 \in \mathfrak{p}$, proving the first statement. The second statement follows at once from the definition (14), using once more that the \mathfrak{p} -action on \mathfrak{k} is trivial. \square

Next, we fix a homomorphism $\chi : \mathfrak{p} \rightarrow \mathfrak{k}$ and turn to the MC-elements of the twisted DGLAs L_I^χ . Consider the following diagram of DGLAs:

$$\begin{array}{ccccccc} & & L_I^\chi & & & & \\ & & \downarrow & \searrow & & & \\ \cdots & \longrightarrow & L_{I_{k+1}}^\chi & \longrightarrow & L_{I_k}^\chi & \longrightarrow & \cdots \longrightarrow L_{I_0}^\chi = \{0\} \end{array}$$

Any MC-element in L_I^χ projects to one in $L_{I_k}^\chi$ for any $k \in \mathbb{N}$, and all maps in the above diagram are equivariant w.r.t. the gauge-actions. Notice further that each $L_{I_k}^\chi$ is nilpotent. To study the MC-elements in L_I^χ , we consider lifts of MC-elements from $L_{I_k}^\chi$ to $L_{I_{k+1}}^\chi$, so as to solve the problem step-by-step. This can be done using the following central extension of nilpotent DGLAs, where $L_{I_0}^\chi = \{0\}$ is trivial:

$$0 \rightarrow L_{I_{k+1}}^\chi \rightarrow L_{I_k}^\chi \rightarrow L_{I_{k-1}}^\chi \rightarrow 0, \quad k \in \mathbb{N} \quad (16)$$

in combination with the following well-known result from deformation theory:

Lemma 7.17. *Let $0 \rightarrow K \rightarrow L \rightarrow M \rightarrow 0$ be a central extension of nilpotent DGLAs. Let $\sigma_M \in M^1$ be a MC-element.*

1. Suppose that $\sigma_L \in L$ projects to σ_M . Then $h := \delta\sigma_L + \frac{1}{2}[\sigma_L, \sigma_L] \in K^2$ is closed and $[h] \in H^2(K)$ is independent of the lift σ_L of σ_M . Moreover, there is some $\eta \in K^1$ such that $\sigma_L + \eta$ is a MC-element in L^1 if and only if $[h] = 0$ in $H^2(K)$.
2. If σ_L and σ'_L are two lifts of σ_M that are both MC-elements in L^1 , then $\Delta := \sigma'_L - \sigma_L \in K^1$ is closed. Conversely, if $\Delta \in K^1$ is closed and σ_L is a lift of σ_M which is a MC-element, then $\sigma'_L := \sigma_L + \Delta$ is also a lift of σ_M which is a MC-element. Moreover, the class $[\Delta] \in H^1(K)$ vanishes if and only if σ_L and σ'_L are related by a gauge transformation of some element $\xi \in K^0$.

3. If σ_L is any lift of σ_M which is a MC-element, then the map $\Delta \mapsto \sigma_L + \Delta$ induces a bijection between $H^1(K)$ and K^0 -orbits of MC-elements in L^1 lifting σ_M .

Proof.

1. It is clear that $h \in K^2$ as it projects to zero in M^2 . Since $\delta h = [\delta\sigma_L, \sigma_L]$ (by the graded Leibniz rule), we find using $\delta\sigma_L = h - \frac{1}{2}[\sigma_L, \sigma_L]$ that

$$\delta h = [h, \sigma_L] - \frac{1}{2}[[\sigma_L, \sigma_L], \sigma_L] = 0,$$

where the second term vanishes by the graded Jacobi identity and the first term vanishes because $h \in K$ is central. Thus h is closed. Suppose that σ'_L is some other lift of σ_M and define $h' := \delta\sigma'_L + \frac{1}{2}[\sigma'_L, \sigma'_L] \in K^2$. Then $\Delta := \sigma'_L - \sigma_L \in K$ lies in the center, so that $h' = h + \delta\Delta$. It follows that $[h] \in H^2(K)$ does not depend on the lift. If there is some $\eta \in K^1$ such that $\sigma_L + \eta$ is a MC-element in L^1 , then

$$0 = \delta(\sigma_L + \eta) + \frac{1}{2}[\sigma_L + \eta, \sigma_L + \eta] = \delta\eta + h.$$

Hence $[h] = 0$. Conversely, if $[h] = 0 \in H^2(K)$, then there exists $\eta \in K^1$ such that $h + \delta\eta = 0$. Then $\sigma_L + \eta$ is a MC-element, by the same computation.

2. Let σ'_L and σ_L be MC-elements in L^1 lifting σ_M . We have already noticed that $h' = h + \delta\Delta$, where $\Delta := \sigma'_L - \sigma_L \in K^1$. Since $h = h' = 0$ by assumption, it follows that $\delta\Delta = 0$. Conversely, suppose $\Delta \in K^1$ is closed and that σ_L is a MC-element projecting to σ_M . Then $\sigma'_L := \sigma_L + \Delta$ projects to σ_M as well. Also, σ'_L is a MC-element, because $\delta\sigma'_L + \frac{1}{2}[\sigma'_L, \sigma'_L] = \delta\sigma_L + \frac{1}{2}[\sigma_L, \sigma_L] + \delta\Delta = 0$. To see that $[\Delta] = 0$ in $H^1(K)$ if and only if σ_L and $\sigma'_L = \sigma_L + \Delta$ are related by a gauge transformation by some element $\xi \in K^0$, observation that if $\xi \in K^0$, then as ξ is central we have

$$\xi.\sigma_L = e^{\text{ad}\xi}(\sigma_L + D) - D = \sigma_L - [D, \xi] = \sigma_L - \delta\xi.$$

3. This is immediate from the previous point. □

Next, we apply Lemma 7.17 to the exact sequences (16).

Lemma 7.18. *For every sequence $(\xi_k)_{k \in \mathbb{N}}$ of degree-0 elements in L^X_I with $\xi_k \in P^k(V) \otimes_{\chi} \mathfrak{k}$ for every $k \in \mathbb{N}$, there exists $\eta \in I \otimes_{\chi} \mathfrak{k}$ such that $j^n(\eta.\sigma) = \xi_n.\xi_{n-1} \cdots \xi_1.\sigma$ for every $\sigma \in L^X_I$ and $n \in \mathbb{N}$.*

Proof. Consider the Lie subgroup $H := \ker(\text{ev}_0 : G \rightarrow K) \triangleleft G$ with Lie algebra $\mathfrak{h} := \ker(\text{ev}_0 : \mathfrak{g} \rightarrow \mathfrak{k}) = I \otimes \mathfrak{k}$. Similarly, for $n \in \mathbb{N}$ let $H_n := \ker(\text{ev}_0 : G_n \rightarrow K)$ and $\mathfrak{h}_n := \text{Lie}(H_n)$. Recall that the exponential map $\exp : \mathfrak{h} \rightarrow H$ is a global diffeomorphism, by Lemma 6.3. Write $\log : H \rightarrow \mathfrak{h}$ for its inverse. From $j^n \circ \exp = \exp \circ j^n : \mathfrak{h} \rightarrow H_n$ we obtain that $\log \circ j^n = j^n \circ \log : H \rightarrow \mathfrak{h}_n$ for any $n \in \mathbb{N}$. As $[\xi_k, I \otimes \mathfrak{k}] \subseteq I^{k+1} \otimes \mathfrak{k}$ and the ξ_k are of increasing order, we claim that the limit $\eta := \lim_{N \rightarrow \infty} \log\left(\prod_{k=1}^N e^{\xi_k}\right)$ exists in $I \otimes \mathfrak{k}$ w.r.t. the projective limit-topology, where k increases from *right to left* in the expression and $\eta_N := \log\left(\prod_{k=1}^N e^{\xi_k}\right)$. Indeed, to see this it suffices to show that for each $n \in \mathbb{N}$ the sequence $(j^n \eta_N)_{N=1}^{\infty}$ stabilizes for large enough values of N . This is the case because for $N \geq n$ we have:

$$j^n \eta_N = j^n \log\left(\prod_{k=1}^N e^{\xi_k}\right) = \log\left(\prod_{k=1}^N e^{j^n(\xi_k)}\right) = \log\left(\prod_{k=1}^n e^{j^n(\xi_k)}\right) = j^n \eta_n$$

where it was used that $j^n(\xi_k) = 0$ for all $k > n$, because $\xi_k \in I^k$. Thus $\eta = \lim_N \eta_N$ is well-defined and satisfies $j^n \eta = j^n \eta_n$ for all $n \in \mathbb{N}$. Let $\sigma \in L^X_I$. Using the fact that

$$\begin{aligned} \xi_{n+1}.\eta_n.\sigma &= e^{\text{ad}\xi_{n+1}}(\eta_n.\sigma + D_{\chi}) - D_{\chi} \\ &= e^{\text{ad}\xi_{n+1}}e^{\text{ad}(\eta_n)}(D_{\chi} + \sigma) - D_{\chi} = e^{\text{ad}\eta_{n+1}}(D_{\chi} + \sigma) - D_{\chi} = \eta_{n+1}.\sigma, \end{aligned}$$

it follows by induction that for any $n \in \mathbb{N}$, the equality $\eta_n.\sigma = \xi_n.\xi_{n-1} \cdots \xi_1.\sigma$ is valid. We thus get:

$$j^n(\eta.\sigma) = j^n(\eta_n.\sigma) = j^n(\xi_n.\xi_{n-1} \cdots \xi_1.\sigma), \quad \forall n \in \mathbb{N}. \quad \square$$

Proposition 7.19. *Assume that $H^1(\mathfrak{p}, P^k(V) \otimes_{\chi} \mathfrak{k}) = 0$ for every $k \in \mathbb{N}$. Then every degree-1 MC-element in L^X_I is gauge-equivalent to 0 in L^X_I .*

Proof. Fix a MC-element $\sigma \in L_I^\chi$. Recall that $j^n(\zeta.\sigma)$ is again a MC-element in $L_{I_n}^\chi$ for any $n \in \mathbb{N}$ and $\zeta \in L_I^\chi$ of degree 0. Notice also that $j^0\sigma = 0$. As $H^1(\mathfrak{p}, P^k(V) \otimes_\chi \mathfrak{k}) = 0$ for every $k \in \mathbb{N}$, it follows using Lemma 7.17 and the exact sequences (16), by induction on $n \in \mathbb{N}$, that we can find a sequence of degree-0 elements $(\xi_k)_{k \in \mathbb{N}}$ in L_I^χ with $\xi_k \in P^k(V) \otimes \mathfrak{k}$ such that $j^n(\xi_n.\xi_{n-1} \cdots \xi_1.\sigma) = 0$ for every $n \in \mathbb{N}$. It follows from Lemma 7.18 that there is some $\eta \in I \otimes_\chi \mathfrak{k}$ such that $j^n(\eta.\sigma) = \xi_n.\xi_{n-1} \cdots \xi_1.\sigma = 0$ in L_I^χ for all $n \in \mathbb{N}$. Thus $\sigma \sim 0$. \square

Lemma 7.20. *Let \mathfrak{p} be a semisimple Lie algebra with no nontrivial compact ideals. If \mathfrak{k} is a compact semisimple Lie algebra, then there are no non-trivial homomorphisms $\mathfrak{p} \rightarrow \mathfrak{k}$.*

Proof. Let $\chi : \mathfrak{p} \rightarrow \mathfrak{k}$ be a homomorphism. Then $\mathfrak{p}/\ker \chi$ is isomorphic to a subalgebra of \mathfrak{k} and is therefore compact. As \mathfrak{p} is semisimple and has no nontrivial compact ideals, it also has no non-trivial compact quotients. Thus $\mathfrak{p}/\ker \chi = \{0\}$ or equivalently $\mathfrak{p} = \ker \chi$, so χ is trivial. \square

Proposition 7.21. *Assume that \mathfrak{p} is semisimple. Let $\chi : \mathfrak{p} \rightarrow \mathfrak{k}$ be a homomorphism and let $\sigma' \in L_I^\chi$. Suppose that $\sigma := \chi + \sigma'$ is a degree-1 MC-element in L_R . Then σ is gauge-equivalent to χ in L_R .*

Proof. Since $H^1(\mathfrak{p}, P^k(V) \otimes_\chi \mathfrak{k}) = 0$ for all $k \in \mathbb{N}_{\geq 0}$ by Whitehead's Lemma [Jac79, III.7. Lem. 3], Proposition 7.19 implies that σ is equivalent to 0 in L_I^χ . Equivalently $\chi + \sigma$ is equivalent to χ in L_R . \square

Theorem 7.12. *Assume that \mathfrak{p} is semisimple. Let the linear map $\sigma : \mathfrak{p} \rightarrow \mathfrak{g}$ satisfy the Maurer-Cartan equation (10). Then σ is gauge-equivalent to $\sigma_0 := \text{ev}_0 \circ \sigma : \mathfrak{p} \rightarrow \mathfrak{k}$. If \mathfrak{p} has no non-trivial compact ideals, then σ is gauge-equivalent to 0.*

Proof. Since $\sigma \in \mathfrak{p}^* \otimes \mathfrak{g}$ is a MC-element in L_R , there is some degree-1 MC-element $\sigma' \in L_I^{\sigma_0}$ such that $\sigma = \sigma_0 + \sigma'$, by Lemma 7.15. Then Proposition 7.21 implies that σ is gauge-equivalent to σ_0 . By Lemma 7.16 we further know that $\sigma_0 : \mathfrak{p} \rightarrow \mathfrak{k}$ is a homomorphism of Lie algebras. Thus, if \mathfrak{p} has no non-compact ideals then σ_0 is trivial by Lemma 7.20. \square

We now prove Theorem 7.13.

Theorem 7.13. *Assume that $\mathfrak{p} = \mathbb{R}$. Let $\sigma \in \mathfrak{g}$ and $\mathbf{v} \in \mathcal{X}_I$. Let $\mathbf{v}_1 := j^1\mathbf{v} \in \mathfrak{gl}(V)$ be the linearization of \mathbf{v} at $0 \in V$. Assume w.l.o.g. that $\sigma_0 := \text{ev}_0(\sigma) \in \mathfrak{t}$ for some maximal torus $\mathfrak{t} \subseteq \mathfrak{k}$. The following assertions hold:*

1. *Assume that $\langle \mathbf{n}, \boldsymbol{\mu} \rangle \neq \alpha(\sigma_0)$ for any root $\alpha \in i\mathfrak{k}^*$ of \mathfrak{k} and $\mathbf{n} \in \mathbb{N}_{\geq}^d$ with $|\mathbf{n}| \geq 1$. Then σ is gauge-equivalent to some $\sigma' \in R \otimes \mathfrak{t}$.*
2. *If \mathbf{v}_1 is semisimple, then σ is gauge-equivalent to some $\nu \in R \otimes \mathfrak{k}$ satisfying $-\mathcal{L}_{\mathbf{v}_1}\nu + [\sigma_0, \nu] = 0$.*
3. *Suppose that $\mathbf{v} = \mathbf{v}_1$ is linear. Assume that $D = -\mathcal{L}_{\mathbf{v}} + \text{ad}_\sigma$ integrates to continuous $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ -action on \mathfrak{g} . Then σ is gauge-equivalent to $\sigma_0 \in \mathfrak{t}$. Moreover $\text{Spec}(\mathbf{v}_1) \cup \text{Spec}(\text{ad}_\sigma) \subseteq i\mathbb{Z}$.*

Proof.

1. Using Lemma 7.15, write $\sigma = \sigma_0 + \sigma'$, where $\sigma' \in L_I^{\sigma_0}$ is a degree-1 MC-element in the shifted DGLA $L_I^{\sigma_0}$. Passing to the complexification, observe that

$$P^n(V_{\mathbb{C}}) \otimes_{\sigma_0} (\mathfrak{k}/\mathfrak{t})_{\mathbb{C}} \cong \bigoplus_{\alpha} P^n(V_{\mathbb{C}}) \otimes_{\sigma_0} (\mathfrak{k}_{\mathbb{C}})_{\alpha}$$

as \mathfrak{p} -modules. The eigenvalues of $-\mathcal{L}_{\mathbf{v}_1} + \text{ad}_{\sigma_0}$ acting on $P^n(V) \otimes_{\sigma_0} (\mathfrak{k}_{\mathbb{C}})_{\alpha}$ are given by $\alpha(\sigma_0) - \langle \mathbf{n}, \boldsymbol{\mu} \rangle$, as \mathbf{n} ranges over the multi-indices $\mathbf{n} \in \mathbb{N}^d$ with $|\mathbf{n}| = n$, and α over the roots of \mathfrak{k} . Thus $-\mathcal{L}_{\mathbf{v}_1} + \text{ad}_{\sigma_0}$ is invertible on $P^n(V) \otimes_{\sigma_0} (\mathfrak{k}_{\mathbb{C}})_{\alpha}$. Consequently

$$H^0(\mathfrak{p}, P^n(V_{\mathbb{C}}) \otimes_{\sigma_0} (\mathfrak{k}_{\mathbb{C}})_{\alpha}) = H^1(\mathfrak{p}, P^n(V_{\mathbb{C}}) \otimes_{\sigma_0} (\mathfrak{k}_{\mathbb{C}})_{\alpha}) = 0$$

for any $n \in \mathbb{N}$ and root α , which in turn implies that $H^0(\mathfrak{p}, P^n(V) \otimes_{\sigma_0} \mathfrak{k}/\mathfrak{t}) = H^1(\mathfrak{p}, P^n(V) \otimes_{\sigma_0} \mathfrak{k}/\mathfrak{t}) = 0$. By the long exact sequence of cohomology groups associated to the short exact sequence

$$0 \rightarrow \left(\bigwedge^{\bullet} \mathfrak{p}^* \right) \otimes P^n(V) \otimes \mathfrak{t} \rightarrow \left(\bigwedge^{\bullet} \mathfrak{p}^* \right) \otimes P^n(V) \otimes_{\sigma_0} \mathfrak{k} \rightarrow \left(\bigwedge^{\bullet} \mathfrak{p}^* \right) \otimes P^n(V) \otimes_{\sigma_0} \mathfrak{k}/\mathfrak{t} \rightarrow 0,$$

it follows that for every $n \in \mathbb{N}$, the inclusion $\mathfrak{p}^* \otimes P^n(V) \otimes \mathfrak{t} \hookrightarrow P^n(V) \otimes_{\sigma_0} \mathfrak{k}$ induces an isomorphism $H^1(\mathfrak{p}, P^n(V) \otimes \mathfrak{t}) \cong H^1(\mathfrak{p}, P^n(V) \otimes_{\sigma_0} \mathfrak{k})$. It then follows using Lemma 7.17 by induction on $n \in \mathbb{N}$ that we can find elements $\xi_k \in P^k(V) \otimes_{\sigma_0} \mathfrak{k}$ s.t. $j^n(\xi_n.\xi_{n-1} \cdots \xi_1.\sigma') \in I_n \otimes \mathfrak{t}$ for every $n \in \mathbb{N}$, the gauge action taking place in $L_I^{\sigma_0}$. By Lemma 7.18 there exists $\eta \in I \otimes_{\sigma_0} \mathfrak{k}$ s.t. $j^n(\eta.\sigma') = j^n(\xi_n.\xi_{n-1} \cdots \xi_1.\sigma') \in I_n \otimes \mathfrak{t}$ for every $n \in \mathbb{N}$. Hence $\zeta := \eta.\sigma' \in I \otimes \mathfrak{t}$ and σ' is gauge-equivalent to ζ in $L_I^{\sigma_0}$. By Lemma 7.15 it follows that $\sigma = \sigma_0 + \sigma'$ is gauge-equivalent to $\sigma_0 + \zeta \in R \otimes \mathfrak{t}$.

2. If \mathbf{v}_1 is semisimple, then so is $-\mathcal{L}_{\mathbf{v}_1} + \text{ad}_{\sigma_0}$ as operator on $P^n(V) \otimes_{\sigma_0} \mathfrak{k}$. Thus

$$H^1(\mathfrak{p}, P^n(V) \otimes_{\sigma_0} \mathfrak{k}) = H^0(\mathfrak{p}, P^n(V) \otimes_{\sigma_0} \mathfrak{k}) = (P^n(V) \otimes_{\sigma_0} \mathfrak{k})^{\mathfrak{p}}.$$

By a similar argument as in the previous item it follows that σ' is gauge-equivalent to some $\zeta \in I \otimes_{\sigma_0} \mathfrak{k}$ in $L_I^{\sigma_0}$ that satisfies $-\mathcal{L}_{\mathbf{v}_1}(\zeta) + [\sigma_0, \zeta] = 0$. By Lemma 7.15 it follows that $\sigma_0 + \sigma' \sim \sigma_0 + \zeta$ in L_R . Notice that $\nu := \sigma_0 + \zeta$ satisfies $-\mathcal{L}_{\mathbf{v}_1}\nu + [\sigma_0, \nu] = 0$.

3. Observe first that any derivation $D' \in \text{der}(\mathfrak{g})$ satisfying $D'(I \otimes \mathfrak{k}) \subseteq (I \otimes \mathfrak{k})$ integrates to a unique 1-parameter group $t \mapsto e^{tD'}$ of automorphisms on \mathfrak{g} that leave the ideal $I \otimes \mathfrak{k} \subseteq \mathfrak{g}$ invariant. Indeed, this follows from fact that the corresponding statement is true for the finite-dimensional Lie algebra $\text{der}(\mathfrak{g}_n)$ for every $n \in \mathbb{N}$, where we use that \mathfrak{g} is the projective limit $\mathfrak{g} = \varprojlim_n \mathfrak{g}_n$. In particular this applies to the \mathbb{T} -action e^{tD} on \mathfrak{g} , which therefore leaves $I \otimes \mathfrak{k}$ invariant. It thus induces a continuous \mathbb{T} -action on $V^* \otimes \mathfrak{k} \cong (I \otimes \mathfrak{k}) / (I^2 \otimes \mathfrak{k})$, which integrates the linear operator $-\mathcal{L}_{\mathbf{v}_1} \otimes 1 + 1 \otimes \text{ad}_{\sigma_0}$ on $V^* \otimes \mathfrak{k}$. This implies that $\mathbf{v}_1 \in \mathfrak{gl}(V)$ and $\text{ad}_{\sigma_0} \in \text{der}(\mathfrak{k})$ integrate to continuous $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ -actions on V and \mathfrak{k} , respectively. As \mathbb{T} is compact it follows in particular that \mathbf{v}_1 is semisimple and that $\text{Spec}(\mathbf{v}_1) \cup \text{Spec}(\text{ad}_{\sigma_0}) \subseteq i\mathbb{Z}$. By Lemma 7.15 we know that there is some degree-1 MC-element $\sigma' \in L_I^{\sigma_0}$ such that $\sigma = \sigma_0 + \sigma'$. By the previous item, it follows that we may assume that σ' satisfies $-\mathcal{L}_{\mathbf{v}_1}\sigma' + [\sigma_0, \sigma'] = 0$, by acting with gauge transformations in $L_I^{\sigma_0}$ if necessary. Let $n \in \mathbb{N}$. The \mathbb{T} -action on $\mathfrak{g}_n = R_n \otimes \mathfrak{k}$ must be unitarizable because \mathbb{T} is compact, so that its generator $D_n := -\mathcal{L}_{\mathbf{v}_1} + [j^n \sigma, -] \in \text{der}(\mathfrak{g}_n)$ must be semisimple. Notice further that $-\mathcal{L}_{\mathbf{v}_1} + [\sigma_0, -]$ is semisimple on \mathfrak{g}_n whereas $[j^n \sigma', -]$ is nilpotent. Since $-\mathcal{L}_{\mathbf{v}_1}\sigma' + [\sigma_0, \sigma'] = 0$, the operators $-\mathcal{L}_{\mathbf{v}_1} + [\sigma_0, -]$ and $[j^n \sigma', -]$ on \mathfrak{g}_n commute. Thus $D_n = (-\mathcal{L}_{\mathbf{v}_1} + [\sigma_0, -]) + [j^n \sigma', -]$ is the Jordan decomposition of D_n . As D_n is semisimple, this implies that $[j^n \sigma', -] = 0$. Thus $j^n \sigma' \in \mathfrak{Z}(\mathfrak{g}_n)$, where $\mathfrak{Z}(\mathfrak{g}_n)$ denotes the center of \mathfrak{g}_n . As \mathfrak{k} is simple, we know that $\mathfrak{Z}(\mathfrak{g}_n) = P^n(V) \otimes \mathfrak{k} \subseteq \mathfrak{g}_n$. Thus $\sigma' \in I^{n-1} \otimes \mathfrak{k}$ for every $n \in \mathbb{N}$, where $I^0 := R$. As $\bigcap_{n \in \mathbb{N}} (I^{n-1} \otimes \mathfrak{k}) = \{0\}$, it follows that $\sigma' = 0$. Hence $\sigma = \sigma_0 \in \mathfrak{t}$. \square

8 Projective Unitary G.P.E. Representations of $(R \otimes \mathfrak{k}) \rtimes_D \mathfrak{p}$

Having obtained the normal form results Theorem 7.12 and Theorem 7.13, we now proceed with the study of continuous projective unitary representation of jet Lie groups and algebras that are of generalized positive energy.

Let us begin by briefly recalling the setting and our notation. We have that V is a finite-dimensional real vector space, $R = \mathbb{R}[[V^*]] := \prod_{n=0}^{\infty} P^k(V)$ is the ring of formal power series on V with coefficients in \mathbb{R} and equipped with the direct product topology. Moreover, \mathfrak{g} denotes the topological Lie algebra $\mathfrak{g} = R \otimes \mathfrak{k}$, where \mathfrak{k} is a compact simple Lie algebra and \mathfrak{p} is a finite-dimensional real Lie algebra acting on \mathfrak{g} by the homomorphism $D : \mathfrak{p} \rightarrow \text{der}(\mathfrak{g})$, which splits into a horizontal and a vertical part according to $D(p) = -\mathcal{L}_{\mathbf{v}(p)} + \text{ad}_{\sigma(p)}$, where $\mathbf{v} : \mathfrak{p} \rightarrow \mathcal{X}_I^{\text{op}}$ is a homomorphism and where the linear map $\sigma : \mathfrak{p} \rightarrow \mathfrak{g}$ satisfies the Maurer Cartan equation (10). Let P and K be the 1-connected Lie groups integrating \mathfrak{p} and K , respectively. For $n \in \mathbb{N}$ write $G_n := J_0^n(V; K)$, $G_n^{\sharp} := J_0^n(V; K) \rtimes P$ and $\mathfrak{g}_n^{\sharp} := \mathfrak{g}_n \rtimes_D \mathfrak{p}$. Define further $G := J_0^{\infty}(V; K) := \varprojlim_n G_n$, $G^{\sharp} := G \rtimes_{\alpha} P$ and $\mathfrak{g}^{\sharp} := \mathfrak{g} \rtimes_D \mathfrak{p} = \varprojlim_n \mathfrak{g}_n^{\sharp}$.

In the following, we are interested in understanding the extent to which the linearization \mathbf{v}_1 of \mathbf{v} and the values of $\sigma(p)$ at the origin already determine properties of the class of representations which are of g.p.e. at a given cone $\mathfrak{C} \subseteq \mathfrak{p}$. To describe the main results, we first have to introduce some more notation.

Define $\sigma_0 := \text{ev}_0 \circ \sigma : \mathfrak{p} \rightarrow \mathfrak{k}$ and let $\mathbf{v}_1 = j^1 \mathbf{v} : \mathfrak{p} \rightarrow \mathfrak{gl}(V)$ be the linearization of \mathbf{v} at the origin. For $p \in \mathfrak{p}$, the vector fields $\mathbf{v}(p)$ splits as $\mathbf{v}(p) = \mathbf{v}_1(p) + \mathbf{v}_{\text{ho}}(p)$ for some formal vector field $\mathbf{v}_{\text{ho}}(p) \in \mathcal{X}_{I^2}$ vanishing up to first order at the origin. Let $\mathbf{v}_1(p) = \mathbf{v}_1(p)_s + \mathbf{v}_1(p)_n$ be the Jordan decomposition of $\mathbf{v}_1(p)$ over \mathbb{C} . Let $V_c^{\mathbb{C}}(p)$ denote the span in $V_{\mathbb{C}}$ of all generalized eigenspaces of $\mathbf{v}_1(p)$ corresponding to eigenvalues with zero real part. Set $V_c(p) := V_c^{\mathbb{C}}(p) \cap V$. If $\mathfrak{C} \subseteq \mathfrak{p}$ is a subset, define $V_c(\mathfrak{C}) := \bigcap_{p \in \mathfrak{C}} V_c(p)$. We call $V_c(\mathfrak{C})$ the ‘center vector space associated \mathfrak{C} ’, in analogy with the center manifold of a fixed point of a dynamical system. Let $V_c(\mathfrak{C})^{\perp} \subseteq V^*$ denote the annihilator of $V_c(\mathfrak{C})$ in V^* . For any $p \in \mathfrak{p}$, let $\Sigma_p \subseteq \mathbb{C}$ denote the additive subsemigroup of \mathbb{C} generated by $\text{Spec}(\mathbf{v}_1(p))$. Recall from Definition 4.1 that for any continuous projective unitary representation $\bar{\pi}$ of $\mathfrak{g} \rtimes_D \mathfrak{p}$, the set $\mathfrak{C}(\bar{\pi})$ consists of all points $p \in \mathfrak{p}$ for which $\bar{\pi}$ is of generalized positive energy at p .

Let us describe the main results of this section. In the context of positive energy representations, we have:

Theorem 8.1. *Let $\bar{\rho}$ be a smooth projective unitary representation of $G \rtimes_{\alpha} P$ which is of p.e. at $p \in \mathfrak{p}$. Assume that $\text{Spec}(\text{ad}_{\sigma_0(p)}) \cap \text{Spec}(\mathfrak{v}_1(p)) = \emptyset$. Then $\bar{\rho}$ factors through $J_0^2(V; K) \rtimes_{\alpha} P$. Moreover the image of $-\mathcal{L}_{\mathfrak{v}_1(p)} + \text{ad}_{\sigma_0(p)}$ in $P^2(V) \otimes \mathfrak{k} \subseteq J_0^2(V; K)$ is contained in $\ker \bar{\rho}$.*

Remark 8.2. Notice that $\text{Spec}(\text{ad}_{\sigma_0(p)}) = \{ \alpha(\sigma_0(p)) : \alpha \in \Delta \} \cup \{0\}$ is a finite subset of $i\mathbb{R}$. In particular the condition $\text{Spec}(\mathfrak{v}_1(p)) \cap \text{Spec}(\text{ad}_{\sigma_0(p)}) = \emptyset$ is satisfied if \mathfrak{v}_1 does not contain purely imaginary eigenvalues.

This is complemented by the following results, which in particular give sufficient conditions for $\bar{\pi}$ to factor through \mathfrak{k} , because $\mathbb{R}[[V_c(\mathfrak{C})^*]] \otimes \mathfrak{k} = \mathfrak{k}$ whenever $V_c(\mathfrak{C}) = \{0\}$.

Theorem 8.3. *Let $\bar{\pi}$ be a continuous projective unitary representation of $\mathfrak{g} \rtimes_D \mathfrak{p}$. Let $\mathfrak{C} \subseteq \mathfrak{C}(\bar{\pi})$. Assume that $\text{Spec}(\text{ad}_{\sigma_0(p)}) \cap \Sigma_p = \emptyset$ for all $p \in \mathfrak{C}$. Then $RV_c(\mathfrak{C})^{\perp} \otimes \mathfrak{k} \subseteq \ker \bar{\pi}$ and hence $\bar{\pi}|_{\mathfrak{g}}$ factors through $\mathbb{R}[[V_c(\mathfrak{C})^*]] \otimes \mathfrak{k}$.*

Theorem 8.4. *Let $\mathfrak{t} \subseteq \mathfrak{k}$ be a maximal Abelian subalgebra. Let $\bar{\pi}$ be a continuous projective unitary representation of \mathfrak{g}^{\sharp} . Let $\mathfrak{C} \subseteq \mathfrak{C}(\bar{\pi})$. Assume for every $p \in \mathfrak{C}$ that $\sigma(p) \in R \otimes \mathfrak{t}$ and $[\mathfrak{v}_1(p)_s, \mathfrak{v}_{\text{ho}}(p)] = 0$. Then $RV_c(\mathfrak{C})^{\perp} \otimes \mathfrak{k} \subseteq \ker \bar{\pi}$ and hence $\bar{\pi}|_{\mathfrak{g}}$ factors through $\mathbb{R}[[V_c(\mathfrak{C})^*]] \otimes \mathfrak{k}$.*

To prove these results, we consider in Section 8.1 the second continuous Lie algebra cohomology $H_{\text{ct}}^2(\mathfrak{g} \rtimes_D \mathfrak{p}; \mathbb{R})$ so as to obtain particular representatives ω of cohomology classes therein. We proceed in Section 8.2 to show that any irreducible smooth projective unitary representation $G \rtimes_{\alpha} P$ factors through the finite-dimensional $G_n \rtimes P$ for some $n \in \mathbb{N}$. This gives us access techniques that are available for finite-dimensional Lie groups, and in particular to Corollary 3.6, which leads to Theorem 8.1. In Section 8.3 and Section 8.4 we study the kernel of the quadratic form $\xi \mapsto \omega(D(p)\xi, \xi)$. Recalling from Corollary 4.6 that

$$[D(p)\eta, \eta] = 0 \implies \left(\omega(D(p)\eta, \eta) = 0 \iff \bar{\pi}(D(p)\eta) = 0 \right), \quad \forall \eta \in \mathfrak{g},$$

this leads to an ideal in \mathfrak{g} contained in $\ker \bar{\pi}$ and is used to prove Theorem 8.3 and Theorem 8.4. These results are supplemented in Section 8.5 by a consideration of the special case where \mathfrak{p} is a simple non-compact Lie algebra, in which case Theorem 7.12 is available. This leads to the following.

Theorem 8.5. *Assume that \mathfrak{p} is non-compact and simple. Let $\bar{\pi}$ be a continuous projective unitary representation of $\mathfrak{g} \rtimes_D \mathfrak{p}$. Write $\mathfrak{C} := \mathfrak{C}(\bar{\pi}) \subseteq \mathfrak{p}$. Then $\bar{\pi}|_{\mathfrak{g}}$ factors through $\mathbb{R}[[V_c(\mathfrak{C})^*]] \otimes \mathfrak{k}$.*

Theorem 8.6. *Assume that \mathfrak{p} is non-compact and simple. Suppose that \mathfrak{v}_1 defines a non-trivial irreducible \mathfrak{p} -representation on V . Let $\bar{\pi}$ be a continuous projective unitary representation of $\mathfrak{g} \rtimes_D \mathfrak{p}$. Either $\mathfrak{C}(\bar{\pi})$ is a pointed P -invariant convex cone or $\bar{\pi}|_{\mathfrak{g}}$ factors through \mathfrak{k} .*

8.1 The Second Lie Algebra Cohomology $H_{\text{ct}}^2(\mathfrak{g} \rtimes_D \mathfrak{p}, \mathbb{R})$.

We next determine suitable representatives of classes in the second Lie algebra cohomology $H_{\text{ct}}^2(\mathfrak{g} \rtimes_D \mathfrak{p}, \mathbb{R})$, which classifies the continuous \mathbb{R} -central extensions of $\mathfrak{g} \rtimes \mathfrak{p} =: \mathfrak{g}^{\sharp}$ up to equivalence. As an intermediate step we first consider $H_{\text{ct}}^2(\mathfrak{g}, \mathbb{R})$, which is completely understood.

Define $\Omega_R^k := R \otimes \bigwedge^k V^*$, equipped with projective limit topology obtained from $\Omega_R^k = \varprojlim_n \Omega_{R_n}^k$, where $\Omega_{R_n}^k := R_n \otimes \bigwedge^k V^*$. This makes Ω_R^k into a Fréchet space. In particular $\Omega_R^0 = R$. Since $J_0^{\infty}(\Omega^k(V)) \cong \Omega_R^k$, we can define a continuous differential $d : \Omega_R^k \rightarrow \Omega_R^{k+1}$ by $dj_0^{\infty} \alpha := j_0^{\infty} d\alpha \in J_0^{\infty}(\Omega^{k+1}(V)) \cong \Omega_R^{k+1}$, which is indeed well-defined. Choosing a basis e_{μ} of V with dual basis dx^{μ} of V^* , the above differential d is on R given $df = \sum_{\mu} (\partial_{\mu} f) \otimes dx^{\mu}$ for $f \in R$.

Lemma 8.7. *Let E be a topological R -module and let $D : R \rightarrow E$ be a continuous derivation. Then there exists a unique continuous R -linear map $\bar{D} : \Omega_R^1 \rightarrow E$ such that $D = \bar{D} \circ d$.*

Proof. Let $\mathbb{R}[V^*]$ denote the ring of polynomial functions on V . As $\mathbb{R}[V^*] \subseteq R$, E is also a $\mathbb{R}[V^*]$ -module and $D|_{\mathbb{R}[V^*]} : \mathbb{R}[V^*] \rightarrow E$ is a derivation. Using the universal property of the Kähler differential forms $\Omega_{\mathbb{R}[V^*]}^1 \cong \mathbb{R}[V^*] \otimes V^*$, there is a unique $\mathbb{R}[V^*]$ -linear map $\bar{D} : \Omega_{\mathbb{R}[V^*]}^1 \rightarrow E$ such that $\bar{D} \circ d|_{\mathbb{R}[V^*]} = D|_{\mathbb{R}[V^*]}$. As $\Omega_{\mathbb{R}[V^*]}^1$ is dense in Ω_R^1 , it remains to extend \bar{D} continuously to the Fréchet space Ω_R^1 . Let $\alpha \in \Omega_R^1$ and let $(\alpha_n)_{n \in \mathbb{N}}$ be a sequence in $\Omega_{\mathbb{R}[V^*]}^1$ s.t. $\alpha_n \rightarrow \alpha$ in Ω_R^1 . We show that $D\alpha := \lim_{n \rightarrow \infty} \bar{D}\alpha_n$ exists and is independent of the approximating sequence (α_n) . Choose a basis $(dx_{\mu})_{\mu=1}^d$ of V^* . Write $\alpha_n = \sum_{\mu=1}^d f_{\mu}^{(n)} dx_{\mu}$ and $\alpha = \sum_{\mu=1}^d f_{\mu} dx_{\mu}$ for some unique $f_{\mu}^{(n)} \in \mathbb{R}[V^*]$ and $f_{\mu} \in R$. Then $f_{\mu}^{(n)} \rightarrow f_{\mu}$ in R for every μ and hence $\lim_{n \rightarrow \infty} \bar{D}\alpha_n = \sum_{\mu=1}^d \lim_{n \rightarrow \infty} f_{\mu}^{(n)} Ddx_{\mu} = \sum_{\mu=1}^d f_{\mu} Ddx_{\mu}$, which is independent of the approximating sequence

(α_n). It follows that \overline{D} extends to a continuous R -linear map $D : \Omega_R^1 \rightarrow E$, which satisfies $\overline{D} \circ d = D$ by construction. It is unique with these properties because its restriction to the dense subspace $\mathbb{R}[V^*] \subseteq R$ is so. \square

Remark 8.8. Lemma 8.7 entails that $d : R \rightarrow \Omega_R^1$ is the universal differential module R in the category of complete locally convex \mathbb{R} -modules, in the sense of [Mai02, Thm. 6]

Proposition 8.9. *Any class $[\omega] \in H_{\text{ct}}^2(\mathfrak{g}, \mathbb{R})$ has a unique representative of the form $\omega(\xi, \eta) = \lambda(\kappa(\xi, d\eta))$, where $\lambda \in \Omega_R^{1,*}$ is closed and continuous functional on Ω_R^1 and κ is the Killing form on \mathfrak{k} . (Closed meaning that $\lambda(dR) = 0$.) Conversely, any such λ defines a 2-cocycle representing some non-zero class in $H_{\text{ct}}^2(\mathfrak{g}, \mathbb{R})$. Consequently, the center of the universal central extension of \mathfrak{g} is (Ω_R^1/dR) .*

Proof. This is a special case of [Mai02, Theorem 16], seeing as $\mathfrak{g} = R \otimes \mathfrak{k}$, where R is a unital, associative and commutative Fréchet algebra and \mathfrak{k} is a simple Lie algebra. \square

Lemma 8.10. *Let ω be an extension of the 2-cocycle $\lambda(\kappa(\xi, d\eta))$ on \mathfrak{g} to a 2-cocycle on $\mathfrak{g}^\sharp = \mathfrak{g} \rtimes \mathfrak{p}$. Then λ is \mathfrak{p} -invariant in the sense that $\lambda(\mathcal{L}_{\mathfrak{v}(p)}\Omega_R^1) = 0$ for every $p \in \mathfrak{p}$.*

Proof. Take $\xi = f \otimes X$ and $\eta = g \otimes X$ for $f, g \in R$ and $0 \neq X \in \mathfrak{k}$. Notice that $[\xi, \eta] = 0$ in $R \otimes \mathfrak{k}$. Using the cocycle identity, this implies

$$\begin{aligned} \omega(D(p)\xi, \eta) + \omega(\xi, D(p)\eta) &= \omega(D(p), [\xi, \eta]) = 0 \\ \omega([\sigma(p), \xi], \eta) + \omega(\xi, [\sigma(p), \eta]) &= \omega(\sigma(p), [\xi, \eta]) = 0. \end{aligned}$$

Using $D(p)\xi = -\mathcal{L}_{\mathfrak{v}(p)}\xi + [\sigma(p), \xi]$ it follows that $0 = \omega(\mathcal{L}_{\mathfrak{v}(p)}\xi, \eta) + \omega(\xi, \mathcal{L}_{\mathfrak{v}(p)}\eta) = \lambda(\mathcal{L}_{\mathfrak{v}(p)}fdg)\kappa(X, X)$. As $\kappa(X, X) \neq 0$ and $RdR = \Omega_R^1$, this shows the claim. \square

Lemma 8.11. *Let $\omega : \mathfrak{g}^\sharp \times \mathfrak{g}^\sharp \rightarrow \mathbb{R}$ be a continuous 2-cocycle on $\mathfrak{g}^\sharp = \mathfrak{g} \rtimes_D \mathfrak{p}$. Then there exists $n \in \mathbb{N}$ and a 2-cocycle ω_n on \mathfrak{g}_n^\sharp such that $\omega(\xi, \eta) = \omega_n(j^n\xi, j^n\eta)$ for all $\xi, \eta \in \mathfrak{g}^\sharp$.*

Proof. Let $\omega : \mathfrak{g}^\sharp \times \mathfrak{g}^\sharp \rightarrow \mathbb{R}$ be a continuous 2-cocycle on \mathfrak{g}^\sharp . Choose norms $\| \cdot \|_n$ on the finite-dimensional Lie algebras \mathfrak{g}_n^\sharp s.t. the quotient maps $j^n : \mathfrak{g}_m^\sharp \rightarrow \mathfrak{g}_n^\sharp$ are contractive for any $n, m \in \mathbb{N}$ with $n \leq m$. The topology on $\mathfrak{g}^\sharp = \varprojlim \mathfrak{g}_n^\sharp$ is specified by the seminorms $\xi \mapsto \|j^n\xi\|_n$ for $n \in \mathbb{N}$ and $\xi \in \mathfrak{g}^\sharp$. As ω is continuous and the maps $\mathfrak{g}_m^\sharp \rightarrow \mathfrak{g}_n^\sharp$ are contractive for $n \leq m$, there exist $n \in \mathbb{N}$ such that $|\omega(\xi, \eta)| \leq \|j^n\xi\|_n \|j^n\eta\|_n$ for all $\xi, \eta \in \mathfrak{g}^\sharp$. As $j^n : \mathfrak{g}^\sharp \rightarrow \mathfrak{g}_n^\sharp$ is surjective, it follows that $\omega(\xi, \eta) = \omega_n(j^n\xi, j^n\eta)$ for a unique 2-cocycle ω_n on \mathfrak{g}_n^\sharp . \square

8.2 Factorization Through Finite Jets

In the context of smooth projective unitary representations $\overline{\rho}$ of the Lie group G^\sharp , it is no loss of generality to consider the case where $\overline{\rho}$ factors through the finite-dimensional Lie group G_n^\sharp for some $n \in \mathbb{N}$:

Theorem 8.12. *Let $\overline{\rho}$ be a smooth projective unitary representation of G^\sharp with lift $\rho : \widehat{G} \rightarrow \text{U}(\mathcal{H}_\rho)$ for some central \mathbb{T} -extension \widehat{G} of G^\sharp . Then ρ decomposes as a (possibly uncountable) direct sum $\rho = \bigoplus_{i \in \mathcal{I}} \rho_i$ s.t. for every $i \in \mathcal{I}$ there exists $n \in \mathbb{N}$ s.t. the projective unitary representations $\overline{\rho}_i$ associated to ρ_i factors through G_n^\sharp . In particular, if $\overline{\rho}$ is irreducible then it factors through G_n^\sharp for some $n \in \mathbb{N}$.*

Proof. Write $N_m^\sharp := \ker(j^{m+1} : G^\sharp \rightarrow G_{m+1}^\sharp)$ and $\mathfrak{n}_m := \ker(j^{m+1} : \mathfrak{g}^\sharp \rightarrow \mathfrak{g}_{m+1}^\sharp)$ for any $m \in \mathbb{N}_{\geq 0}$, so that $G_m^\sharp \cong G^\sharp/N_m^\sharp$ for any $m \in \mathbb{N}_{\geq 0}$. Notice that $N_m^\sharp \subseteq N_n^\sharp$ whenever $n \leq m$. Since $\overline{\rho}$ is a smooth projective representation, it follows from [JN19, Thm. 4.3] that \widehat{G} is a Lie group. It is moreover regular by [Nee06, Thm. V.I.8], because both G^\sharp and \mathbb{T} are so. Let $\widehat{\mathfrak{g}} := \text{Lie}(\widehat{G})$. Then $\widehat{\mathfrak{g}}$ is a central \mathbb{R} -extension of \mathfrak{g}^\sharp in the category of locally convex Lie algebras. Let the continuous 2-cocycle $\omega : \mathfrak{g}^\sharp \times \mathfrak{g}^\sharp \rightarrow \mathbb{R}$ represent the corresponding class in $H_{\text{ct}}^2(\mathfrak{g}^\sharp, \mathbb{R})$. By Lemma 8.11, there is some $n \in \mathbb{N}$ such that for all $\xi \in \mathfrak{g}^\sharp$ we have $j^n\xi = 0 \implies \omega(\xi, \eta) = 0$ for all $\eta \in \mathfrak{g}^\sharp$. Let \widehat{N}_n be the closed normal subgroup of \widehat{G} covering N_n^\sharp and let $\widehat{\mathfrak{n}}_n$ be its Lie algebra. Then \widehat{N}_n is a central \mathbb{T} -extension of N_n^\sharp integrating $\widehat{\mathfrak{n}}_n$. Since $\omega|_{\mathfrak{n}^\sharp \times \mathfrak{n}^\sharp} = 0$, the central \mathbb{R} -extension $\widehat{\mathfrak{n}}_n$ is trivial. Hence $\widehat{\mathfrak{n}}_n \cong \mathbb{R} \oplus \mathfrak{n}_n^\sharp$ as central \mathbb{R} -extensions of \mathfrak{n}_n^\sharp . As N_n^\sharp is regular and 1-connected, it follows from [Nee06, Thm. III.1.5] that there is a commutative diagram

$$\begin{array}{ccccc} \mathbb{R} & \longrightarrow & \mathbb{R} \times N_n^\sharp & \longrightarrow & N_n^\sharp \\ \downarrow \text{id} & & \downarrow \tilde{\phi} & & \downarrow \text{id} \\ \mathbb{T} & \longrightarrow & \widehat{N}_n & \longrightarrow & N_n^\sharp \end{array}$$

of locally convex regular Lie groups. Observe that $\tilde{\phi}$ is surjective and that $\ker \tilde{\phi} = \mathbb{Z}$. Thus $\widehat{N}_n \cong \mathbb{T} \times N_n^\sharp$ as central \mathbb{T} -extension of N_n^\sharp . Let $\phi : \mathbb{T} \times N_n^\sharp \rightarrow \widehat{N}_n$ realize the isomorphism. For any integer $m \geq n$, let $\mathcal{N}_m := \phi(\{1\} \times N_m^\sharp) \subseteq \widehat{N}_n \subseteq \widehat{G}$, which is a closed normal subgroup of \widehat{G} isomorphic to and covering N_n^\sharp . Then $\mathcal{N} := \{\mathcal{N}_m\}_{m \geq n}$ is a filter basis of (decreasing) closed normal subgroups of \widehat{G} satisfying $\varprojlim \mathcal{N} = \{1\}$, in the sense that for any 1-neighborhood U of \widehat{G} there exists $m \geq n$ such that $\mathcal{N}_m \subseteq U$. Indeed, since $G^\sharp = \varprojlim_m G_m^\sharp$ carries the projective limit topology and \widehat{G} is a locally trivial principal \mathbb{T} -bundle over G^\sharp [JN19, Thm. 4.3], it follows that any 1-neighborhood $U \subseteq \widehat{G}$ contains $\phi(I \times N_m^\sharp)$ for large enough m and some open 1-neighborhood $I \subseteq \mathbb{T}$. It now follows from [Nee10a, Thm.12.2] that ρ decomposes as a possibly uncountable direct sum $\rho \cong \bigoplus_{i \in \mathcal{I}} \rho_i$ such that for every $i \in \mathcal{I}$ there exists some $m \geq n$ with $\rho_i(\mathcal{N}_m) = \{1\}$, which implies that $\bar{\rho}_i(N_m^\sharp) = 1$. \square

Theorem 8.12 gives us access to techniques that are available for finite-dimensional Lie groups, and in particular to Corollary 3.6. This can be used to prove Theorem 8.1.

Theorem 8.1. *Let $\bar{\rho}$ be a smooth projective unitary representation of $G \rtimes_\alpha P$ which is of p.e. at $p \in \mathfrak{p}$. Assume that $\text{Spec}(\text{ad}_{\sigma_0(p)}) \cap \text{Spec}(\mathfrak{v}_1(p)) = \emptyset$. Then $\bar{\rho}$ factors through $J_0^2(V; K) \rtimes_\alpha P$. Moreover the image of $-\mathcal{L}_{\mathfrak{v}_1(p)} + \text{ad}_{\sigma_0(p)}$ in $P^2(V) \otimes \mathfrak{k} \subseteq J_0^2(V; K)$ is contained in $\ker \bar{\rho}$.*

To prove Theorem 8.1, it suffices by Theorem 8.12 to consider the case where $\bar{\rho}$ factors through the finite-dimensional Lie group G_k^\sharp for some $k \in \mathbb{N}$, which we thus assume. Write \mathfrak{a} for the ideal in \mathfrak{g}_k^\sharp generated by $p \in \mathfrak{p}$. Let $\mathfrak{a}_n \subseteq \mathfrak{a}$ denote the maximal nilpotent ideal in \mathfrak{a} . According to Corollary 3.6 we have $[\mathfrak{a}, [\mathfrak{a}_n, \mathfrak{a}_n]] \subseteq \ker d\bar{\rho}$. Recall that $I_k := I/I^{k+1}$.

Lemma 8.13. *Suppose that $V^* \otimes \mathfrak{k} \subseteq j^1 \mathfrak{a}_n$. Then $I_k \otimes \mathfrak{k} \subseteq \mathfrak{a}_n$.*

Proof. By assumption $V^* \otimes \mathfrak{k} \subseteq \mathfrak{a}_n + I_k^2 \otimes \mathfrak{k}$. As \mathfrak{k} is perfect it follows that

$$I_k^{l+1} \otimes \mathfrak{k} = [V^* \otimes \mathfrak{k}, I_k^l \otimes \mathfrak{k}] \subseteq \mathfrak{a}_n + [I_k^2 \otimes \mathfrak{k}, I_k^l \otimes \mathfrak{k}] = \mathfrak{a}_n + I_k^{l+2} \otimes \mathfrak{k}, \quad \forall l \in \mathbb{N}.$$

Thus it follows by induction that $V^* \otimes \mathfrak{k} \subseteq \mathfrak{a}_n + I_k^{l+1} \otimes \mathfrak{k}$ for all $l \in \mathbb{N}$. As $\bigcap_l (\mathfrak{a}_n + I_k^{l+1} \otimes \mathfrak{k}) = \mathfrak{a}_n$, it follows that $V^* \otimes \mathfrak{k} \subseteq \mathfrak{a}_n$ and hence $I_k \otimes \mathfrak{k} \subseteq \mathfrak{a}_n$. \square

Proof of Theorem 8.1. We may assume that $\bar{\rho}$ factors through G_k^\sharp for some $k \in \mathbb{N}$. It suffices to show that $d\bar{\rho}$ factors through \mathfrak{g}_2^\sharp and that the image of $-\mathcal{L}_{\mathfrak{v}_1(p)} + \text{ad}_{\sigma_0(p)}$ in $P^2(V) \otimes \mathfrak{k} \subseteq \mathfrak{g}_2^\sharp$ is contained in $\ker d\bar{\rho}$. By Corollary 3.6 we know that $[\mathfrak{a}, [\mathfrak{a}_n, \mathfrak{a}_n]] \subseteq \ker d\bar{\rho}$. Moreover $I_k^3 \otimes \mathfrak{k} = [I_k \otimes \mathfrak{k}, [I_k \otimes \mathfrak{k}, I_k \otimes \mathfrak{k}]]$, because \mathfrak{k} is perfect. To see that $d\rho$ factors through \mathfrak{g}_2^\sharp it thus suffices to show that $I_k \otimes \mathfrak{k} \subseteq \mathfrak{a}_n$. By Lemma 8.13 it is further sufficient to show that $V^* \otimes \mathfrak{k} \subseteq j^1(\mathfrak{a}_n)$. Write $D_1(p) := \mathcal{L}_{\mathfrak{v}_1(p)} + [\sigma_0(p), -]$. Notice that $j^1(D(p)\xi) = D_1(p)\xi$ for $\xi \in V^* \otimes \mathfrak{k}$. As $\text{Spec}(\text{ad}_{\sigma_0(p)})$ is a symmetric subset of $i\mathbb{R}$, the assumption $\text{Spec}(\mathfrak{v}_1(p)) \cap \text{Spec}(\text{ad}_{\sigma_0(p)}) = \emptyset$ implies that $\text{Spec}(D_1(p)) = \{ \mu + \nu : \mu \in \text{Spec}(\mathfrak{v}_1(p)), \nu \in \text{Spec}(\text{ad}_{\sigma_0(p)}) \}$ does not contain zero. Hence $D_1(p)$ is invertible on $V^* \otimes \mathfrak{k}$. Thus if $\eta \in V^* \otimes \mathfrak{k}$ is arbitrary, there exists $\xi \in V^* \otimes \mathfrak{k}$ such that $\eta = D_1(p)\xi$. Then $\eta = D_1(p)\xi = j^1(D(p)\xi) = j^1([p, \xi]) \in j^1(\mathfrak{a}_n)$. Thus $V^* \otimes \mathfrak{k} \subseteq j^1(\mathfrak{a}_n)$. We obtain that $I_k \otimes \mathfrak{k} \subseteq \mathfrak{a}_n$ and $I_k^3 \otimes \mathfrak{k} \subseteq \ker d\rho$, so $d\rho$ factors through \mathfrak{g}_2^\sharp . We may thus assume that $k = 2$. We then obtain

$$D_1(p)(P^2(V) \otimes \mathfrak{k}) = D(p)(I_k^2 \otimes \mathfrak{k}) = [p, [I_k \otimes \mathfrak{k}, I_k \otimes \mathfrak{k}]] \subseteq [\mathfrak{a}, [\mathfrak{a}_n, \mathfrak{a}_n]] \subseteq \ker d\bar{\rho}. \quad \square$$

8.3 The Case Where $\mathfrak{p} = \mathbb{R}$

We proceed the study of projective unitary representations $\bar{\pi}$ of $\mathfrak{g}^\sharp = \mathfrak{g} \rtimes_D \mathfrak{p}$ which are of generalized positive energy. We first specialize to the case where $\mathfrak{p} = \mathbb{R}$, aiming to consider its consequences for the general case afterwards.

As $\mathfrak{p} = \mathbb{R}$, we may as well identify \mathfrak{v} with $\mathfrak{v}(1) \in \mathcal{X}_I$, D with $D(1)$ and σ with $\sigma(1) \in \mathfrak{g}$. Recall that the derivation D is given by $D = -\mathcal{L}_{\mathfrak{v}} + [\sigma, -]$. Write $\mathfrak{v} = \mathfrak{v}_1 + \mathfrak{v}_{\text{ho}}$, where $\mathfrak{v}_1 := j^1 \mathfrak{v} \in \mathfrak{gl}(V)$ is the linearization of \mathfrak{v} at $0 \in V$ and where $\mathfrak{v}_{\text{ho}} \in \mathcal{X}_{I^2}$. Let $\mathfrak{v}_1 = \mathfrak{v}_{1,s} + \mathfrak{v}_{1,n}$ denote the Jordan decomposition of \mathfrak{v}_1 over \mathbb{C} . Write $V_c^\mathbb{C}$ for the span of the eigenspaces of \mathfrak{v}_1 whose corresponding eigenvalue has zero real part. Set $V_c := V_c^\mathbb{C} \cap V$. Write $\mathfrak{d} := (0, 1) \in \mathfrak{g}^\sharp = \mathfrak{g} \rtimes_D \mathbb{R}$. Let $V_c^\perp \subseteq V^*$ denote the annihilator of V_c in V^* , so $V_c^\perp \cong (V/V_c)^*$.

Theorem 8.14. *Let $\mathfrak{t} \subseteq \mathfrak{k}$ be a maximal Abelian subalgebra. Assume that $\sigma \in R \otimes \mathfrak{t} \subseteq \mathfrak{g}$ and $[\mathfrak{v}_{1,s}, \mathfrak{v}_{\text{ho}}] = 0$ in \mathcal{X}_I . Let $\bar{\pi}$ be a continuous projective unitary representation of \mathfrak{g}^\sharp on the pre-Hilbert space \mathcal{D} . Assume that $\bar{\pi}$ is of g.p.e. at $\mathfrak{d} \in \mathfrak{g} \rtimes_D \mathbb{R}\mathfrak{d}$. Then $R V_c^\perp \subseteq \ker \bar{\pi}$. Consequently, $\bar{\pi}|_{\mathfrak{g}}$ factors through $\mathbb{R}[[V_c^*]] \otimes \mathfrak{k}$. In particular, if $V_c = \{0\}$ then $\bar{\pi}|_{\mathfrak{g}}$ factors through \mathfrak{k} .*

Remark 8.15. By acting with formal diffeomorphisms if necessary, one may by Theorem 7.7 always bring \mathfrak{v} into a normal form, in the sense that $[\mathfrak{v}_{1,s}, \mathfrak{v}_{\text{ho}}] = 0$ in \mathcal{X}_I . Moreover, Theorem 7.13 provides sufficient conditions guaranteeing that σ is gauge equivalent to some element in $R \otimes \mathfrak{t}$.

Proof of Theorem 8.14

Let ω be a continuous 2-cocycle on \mathfrak{g}^\sharp that represents the class in $H_{ct}^2(\mathfrak{g}^\sharp, \mathbb{R})$ corresponding to the central \mathbb{R} -extension of \mathfrak{g}^\sharp obtained from $\bar{\pi}$ by pulling back $\mathfrak{u}(\mathcal{D}) \rightarrow \mathfrak{pu}(\mathcal{D})$ along $\bar{\pi}$. In view of Proposition 8.9 and Lemma 8.10, we may and do assume that ω satisfies $\omega(\xi, \eta) = \lambda(\kappa(\xi, d\eta))$ for any $\xi, \eta \in \mathfrak{g}$, where $\lambda : \Omega_R \rightarrow \mathbb{R}$ is continuous, \mathfrak{p} -invariant and closed. We write fX instead of $f \otimes X$ for $f \in R_{\mathbb{C}}$ and $X \in \mathfrak{k}_{\mathbb{C}}$. Let $\Delta \subseteq i\mathfrak{t}^*$ denote the set of roots of \mathfrak{k} . Finally, write $\mathfrak{h} := \mathfrak{t}_{\mathbb{C}} \subseteq \mathfrak{k}_{\mathbb{C}}$. Recall from Corollary 4.6 that

$$[D\eta, \eta] = 0 \implies \left(\omega(D\eta, \eta) = 0 \iff \bar{\pi}(D\eta) = 0 \right), \quad \forall \eta \in \mathfrak{g}. \quad (17)$$

Moreover, $\omega(D\eta, \eta) \geq 0$ whenever $[D(p)\eta, \eta] = 0$. In the present setting, this yields:

Proposition 8.16. *Fix $f \in R$. Then $\bar{\pi}(R\mathcal{L}_v f \otimes \mathfrak{k}) = \{0\} \iff \lambda(fd\mathcal{L}_v f) = 0$.*

Proof. For any $H \in \mathfrak{t}$, observe that $DfH = -\mathcal{L}_v fH$ because $\sigma \in R \otimes \mathfrak{t}$, so $[DfH, fH] = -[\mathcal{L}_v fH, fH] = 0$. Using (17) we obtain

$$-\kappa(H, H)\lambda(\mathcal{L}_v fdf) = 0 \iff \bar{\pi}(\mathcal{L}_v fH) = 0, \quad \forall H \in \mathfrak{t}. \quad (18)$$

Assume $\bar{\pi}(R\mathcal{L}_v f \otimes \mathfrak{k}) = \{0\}$. Then $\bar{\pi}(\mathcal{L}_v fH) = 0$ for any $H \in \mathfrak{t}$ and hence $\lambda(\mathcal{L}_v fdf) = 0$ by (18). Conversely, suppose $\lambda(\mathcal{L}_v fdf) = 0$. Then $\bar{\pi}(\mathcal{L}_v fH) = 0$ for all $H \in \mathfrak{t}$, by (18). Taking the commutator with $\bar{\pi}(gX_\alpha)$, where $g \in R$, $\alpha \in \Delta$ is a root and $X_\alpha \in (\mathfrak{k}_{\mathbb{C}})_\alpha$ is a corresponding root vector, it follows that

$$\bar{\pi}(g\mathcal{L}_v fX_\alpha) = 0 \quad \forall X_\alpha \in (\mathfrak{k}_{\mathbb{C}})_\alpha, g \in R. \quad (19)$$

Take $X_\alpha \in (\mathfrak{k}_{\mathbb{C}})_\alpha$ and $Y_{-\alpha} \in (\mathfrak{k}_{\mathbb{C}})_{-\alpha}$. Write $H_\alpha = [X_\alpha, Y_{-\alpha}]$. By taking commutators with $\bar{\pi}(1 \otimes Y_{-\alpha})$ in equation (19) we find that $\bar{\pi}(g\mathcal{L}_v fH_\alpha) = 0$. As $\mathfrak{h} = \sum_\alpha [(\mathfrak{k}_{\mathbb{C}})_\alpha, (\mathfrak{k}_{\mathbb{C}})_{-\alpha}]$, this shows together with equation (19) by linearity and the root space decomposition that $\bar{\pi}(R\mathcal{L}_v f \otimes \mathfrak{k}) = \{0\}$. \square

Define the quadratic form $q(f) := \lambda(\mathcal{L}_v fdf) = -\lambda(fd\mathcal{L}_v f)$ on R . Let $\mathcal{N} := \ker q$ denote its kernel. By Proposition 8.16, \mathcal{N} generates an ideal $J \otimes \mathfrak{k}$ on which $\bar{\pi}$ vanishes, where $J := R\mathcal{L}_v \mathcal{N}$.

Corollary 8.17. *Set $J := R\mathcal{L}_v \mathcal{N}$. Then $J \otimes \mathfrak{k} \subseteq \ker(\bar{\pi})$.*

Together with the fact that λ vanishes on exact forms and is \mathcal{L}_v -invariant, this puts severe restrictions on the representation $\bar{\pi}$ and leads to Theorem 8.14. Let us also remark the following:

Lemma 8.18. *The bilinear form $\beta_q(f, g) := \lambda(\mathcal{L}_v fdg)$ on R associated to q is symmetric, the quadratic form q is positive semi-definite and $\mathcal{N} = \{f \in R : \beta_q(f, g) = 0 \ \forall g \in R\}$.*

Proof. As λ is closed and \mathcal{L}_v -invariant, it follows that β is symmetric. To see that q is positive semi-definite, let $f \in R$ and $0 \neq H \in \mathfrak{t}$. Write $\eta := fH$ and notice that $[D\eta, \eta] = 0$. By Corollary 4.6 we know that $-\kappa(H, H)\lambda(\mathcal{L}_v fdf) = \omega(D\eta, \eta) \geq 0$. As κ is negative definite on \mathfrak{k} we obtain that q is positive semi-definite. It follows that $|\beta_q(f, g)|^2 \leq q(f)q(g)$, which implies $\mathcal{N} = \{f \in R : \beta_q(f, g) = 0 \ \forall g \in R\}$. \square

The following observation is also noteworthy, although it will not be used:

Lemma 8.19. *$\mathcal{N} \subseteq R$ is a subalgebra.*

Proof. Let $f, g \in \mathcal{N}$. Then using the Leibniz rule and Proposition 8.16 we obtain

$$\bar{\pi}(R\mathcal{L}_v(fg) \otimes \mathfrak{k}) = \bar{\pi}(fR\mathcal{L}_v g \otimes \mathfrak{k}) + \bar{\pi}(gR\mathcal{L}_v f \otimes \mathfrak{k}) \subseteq \{0\},$$

Applying Proposition 8.16 once more, we conclude that $fg \in \mathcal{N}$. \square

Lemma 8.20. *$\lambda \circ \mathcal{L}_{v_{1,s}} = 0$.*

Proof. As $\lambda : \Omega_{R_k}^1 \rightarrow \mathbb{R}$ is continuous, it factors through the finite-dimensional space $\Omega_{R_k}^1 = R_k \otimes V^*$ for some $k \in \mathbb{N}$. Notice that both $\mathcal{L}_{v_{1,n}}$ and $\mathcal{L}_{v_{ho}}$ are nilpotent on $\Omega_{R_k}^1 \otimes_{\mathbb{R}} \mathbb{C}$, whereas $\mathcal{L}_{v_{1,s}}$ is semisimple on it. Also $[\mathcal{L}_{v_{1,s}}, \mathcal{L}_{v_{1,n}} + \mathcal{L}_{v_{ho}}] = 0$ because $[v_{1,s}, v_{ho}] = [v_{1,s}, v_{1,n}] = 0$. Thus $\mathcal{L}_v = \mathcal{L}_{v_{1,s}} + (\mathcal{L}_{v_{1,n}} + \mathcal{L}_{v_{ho}})$ is the Jordan decomposition of \mathcal{L}_v acting on $\Omega_{R_k}^1 \otimes_{\mathbb{R}} \mathbb{C}$. Thus $\text{Im}(\mathcal{L}_{v_{1,s}}) \subseteq \text{Im}(\mathcal{L}_v)$ when $\mathcal{L}_{v_{1,s}}$ and \mathcal{L}_v are considered as operators on $\Omega_{R_k}^1 \otimes_{\mathbb{R}} \mathbb{C}$. As λ is \mathcal{L}_v -invariant, we know $\lambda \circ \mathcal{L}_v = 0$. Thus $\lambda \circ \mathcal{L}_{v_{1,s}} = 0$. \square

In particular, λ vanishes on the eigenspaces in $\Omega_{R_{\mathbb{C}}}^1$ of $\mathcal{L}_{v_{1,s}}$ corresponding to non-zero eigenvalues. We introduce some more notation. Let $E_{\mathbb{C}}$ denote the span of all eigenspaces in $R_{\mathbb{C}}$ of $\mathcal{L}_{v_{1,s}}$ corresponding to eigenvalues with non-zero real part. Define $E := E_{\mathbb{C}} \cap R$ and $E^n := E \cap I^n$.

Lemma 8.21. $E \subseteq \mathcal{N}$.

Proof. Let $\mu \in \text{Spec}(\mathcal{L}_{\mathbf{v}_{1,s}})$ with $\text{Re}(\mu) \neq 0$. Set $E_\mu := \ker(\mathcal{L}_{\mathbf{v}_{1,s}} - \mu I) \subseteq R_{\mathbb{C}}$. Suppose first that $\mu \in \mathbb{R}$. If $\psi \in E_\mu \cap R$ then because $\mathcal{L}_{\mathbf{v}}$ leaves the eigenspaces of $\mathcal{L}_{\mathbf{v}_{1,s}}$ invariant, the 1-form $\psi d\mathcal{L}_{\mathbf{v}}\psi$ is an eigenvector of $\mathcal{L}_{\mathbf{v}_{1,s}}$ with non-zero eigenvalue 2μ . By Lemma 8.20 it follows that $q(\psi) = 0$ and hence $\psi \in \mathcal{N}$. Thus $E_\mu \subseteq \mathcal{N}$. Next, suppose that μ is not real. Then also $\bar{\mu}$ is an eigenvalue of $\mathcal{L}_{\mathbf{v}_{1,s}}$. Write $W_{\mathbb{C}} := E_\mu \oplus E_{\bar{\mu}}$ and $W := W_{\mathbb{C}} \cap R$. Take $\psi \in W$ arbitrary. Then $\psi = \eta + \bar{\eta}$ for some $\eta \in E_\mu$ (and hence $\bar{\eta} \in E_{\bar{\mu}}$). As $\mu + \bar{\mu} = 2\text{Re}(\mu) \neq 0$ and $\mathcal{L}_{\mathbf{v}}$ leaves the eigenspaces of $\mathcal{L}_{\mathbf{v}_{1,s}}$ invariant, each of the 1-forms $\eta d\mathcal{L}_{\mathbf{v}}\eta$, $\eta d\mathcal{L}_{\mathbf{v}}\bar{\eta}$, $\bar{\eta} d\mathcal{L}_{\mathbf{v}}\eta$ and $\bar{\eta} d\mathcal{L}_{\mathbf{v}}\bar{\eta}$ are eigenvectors of $\mathcal{L}_{\mathbf{v}_{1,s}}$ with non-zero eigenvalue. Using Lemma 8.20 it follows that $q(\psi) = 0$ and hence $\psi \in \mathcal{N}$. Thus $W \subseteq \mathcal{N}$. As \mathcal{N} is a linear subspace, we have shown $E \subseteq \mathcal{N}$. \square

Lemma 8.22. $RE \subseteq R\mathcal{L}_{\mathbf{v}}E$.

Proof. Write $J := R\mathcal{L}_{\mathbf{v}}E$. As J is an ideal in R it suffices to show $E \subseteq J$. We claim that $E^n \subseteq J + E^{n+1}$ for every $n \in \mathbb{N}_{\geq 0}$. Indeed, take $\psi \in E^n$. As $\mathcal{L}_{\mathbf{v}_1}$ is invertible on E^n (which is true because $\mathcal{L}_{\mathbf{v}_1}$ is invertible on every finite-dimensional and $\mathcal{L}_{\mathbf{v}_1}$ -invariant subspace $E \cap P^k(V) \subseteq E$), there exists some $\eta \in E^n$ s.t. $\mathcal{L}_{\mathbf{v}_1}\eta = \psi$. Observe that $\mathcal{L}_{\mathbf{v}_{\text{ho}}}E \subseteq E$ because $[\mathcal{L}_{\mathbf{v}_{1,s}}, \mathcal{L}_{\mathbf{v}_{\text{ho}}}] = 0$. Also $\mathcal{L}_{\mathbf{v}_{\text{ho}}}I^n \subseteq I^{n+1}$, since $\mathbf{v}_{\text{ho}} \in \mathcal{X}_{I^2}$. Thus $\mathcal{L}_{\mathbf{v}_{\text{ho}}}E^n \subseteq E^{n+1}$. In particular $\mathcal{L}_{\mathbf{v}_{\text{ho}}}\eta \in E^{n+1}$. Then $\psi = \mathcal{L}_{\mathbf{v}_1}\eta = \mathcal{L}_{\mathbf{v}}\eta - \mathcal{L}_{\mathbf{v}_{\text{ho}}}\eta \in J + E^{n+1}$, as required. By induction it follows that $E = E^0 \subseteq J + E^n$ for every $n \in \mathbb{N}$. As $\bigcap_{n \in \mathbb{N}}(J + E^n) = J$, this implies $E \subseteq J$. \square

Proof of Theorem 8.14:

Using Lemma 8.21 we obtain $E \subseteq \mathcal{N}$. By Corollary 8.17, this implies $J \otimes \mathfrak{k} \subseteq \ker \bar{\pi}$, where $J = R\mathcal{L}_{\mathbf{v}}E$. By Lemma 8.22, we know $RE \subseteq J$. Notice that $E \cap V^* = V_c^\perp$, so in particular $RV_c^\perp \subseteq J$. Thus $RV_c^\perp \otimes \mathfrak{k} \subseteq \ker \bar{\pi}$. Notice that $R/(RV_c^\perp) \cong \mathbb{R}[[V_c^*]]$, because $V_c^* = V^*/V_c^\perp$. We conclude that $\bar{\pi}$ factors through the quotient $(R \otimes \mathfrak{k})/(RV_c^\perp \otimes \mathfrak{k}) \cong (\mathbb{R}[[V_c^*]] \otimes \mathfrak{k})$. \square

8.4 The Case of General \mathfrak{p}

Let us return to the case where P is a 1-connected finite-dimensional Lie group with Lie algebra \mathfrak{p} . Let us recall some of the notation introduced in Section 8.

Define $\sigma_0 := \text{ev}_0 \circ \sigma : \mathfrak{p} \rightarrow \mathfrak{k}$ and let $\mathbf{v}_1 = j^1 \mathbf{v} : \mathfrak{p} \rightarrow \mathfrak{gl}(V)$ be the linearization of \mathbf{v} at $0 \in V$. For $p \in \mathfrak{p}$, the vector fields $\mathbf{v}(p)$ splits as $\mathbf{v}(p) = \mathbf{v}_1(p) + \mathbf{v}_{\text{ho}}(p)$ for some formal vector field $\mathbf{v}_{\text{ho}}(p) \in \mathcal{X}_{I^2}$ vanishing up to first order at the origin. Let $\mathbf{v}_1(p) = \mathbf{v}_1(p)_s + \mathbf{v}_1(p)_n$ be the Jordan decomposition of $\mathbf{v}_1(p)$ over \mathbb{C} . Let $V_c^{\mathbb{C}}(p)$ denote the span in $V_{\mathbb{C}}$ of all generalized eigenspaces of $\mathbf{v}_1(p)$ corresponding to eigenvalues with zero real part. Set $V_c(p) := V_c^{\mathbb{C}}(p) \cap V$. If $\mathfrak{C} \subseteq \mathfrak{p}$ is a subset, define $V_c(\mathfrak{C}) := \bigcap_{p \in \mathfrak{C}} V_c(p)$. Let $V_c(\mathfrak{C})^\perp \subseteq V^*$ denote the annihilator of $V_c(\mathfrak{C})$ in V^* . Let $\Sigma_p \subseteq \mathbb{C}$ denote the additive subsemigroup of \mathbb{C} generated by $\text{Spec}(\mathbf{v}_1(p))$. For any continuous projective unitary representation $\bar{\pi}$ of $\mathfrak{g} \rtimes_D \mathfrak{p}$, the set $\mathfrak{C}(\bar{\pi})$ consists of all points $p \in \mathfrak{p}$ for which $\bar{\pi}$ is of generalized positive energy at p .

We use Theorem 8.14 combined with suitable normal form results to prove Theorem 8.3 and Theorem 8.4.

Lemma 8.23. *Let W be a finite-dimensional real vector space and let $W_i \subseteq W$ be a collection of linear subspaces, where $i \in \mathcal{I}$ for some indexing set \mathcal{I} . Then $(\bigcap_{i \in \mathcal{I}} W_i)^\perp = \text{Span}_{i \in \mathcal{I}} W_i^\perp$.*

Proof. Notice first that $\bigcap_{i \in \mathcal{I}} W_i^\perp = [\text{Span}_{i \in \mathcal{I}} W_i]^\perp$. Applying this observation to the subspaces $W_i^\perp \subseteq W^*$, we obtain that $\bigcap_{i \in \mathcal{I}} W_i = \bigcap_{i \in \mathcal{I}} (W_i^\perp)^\perp = [\text{Span}_{i \in \mathcal{I}} W_i^\perp]^\perp$, where we also used that $(W_i^\perp)^\perp \cong W_i$ for any $i \in \mathcal{I}$. Taking annihilators, this implies $(\bigcap_{i \in \mathcal{I}} W_i)^\perp = \text{Span}_{i \in \mathcal{I}} W_i^\perp$. \square

Theorem 8.3. *Let $\bar{\pi}$ be a continuous projective unitary representation of $\mathfrak{g} \rtimes_D \mathfrak{p}$. Let $\mathfrak{C} \subseteq \mathfrak{C}(\bar{\pi})$. Assume that $\text{Spec}(\text{ad}_{\sigma_0(p)}) \cap \Sigma_p = \emptyset$ for all $p \in \mathfrak{C}$. Then $RV_c(\mathfrak{C})^\perp \otimes \mathfrak{k} \subseteq \ker \bar{\pi}$ and hence $\bar{\pi}|_{\mathfrak{g}}$ factors through $\mathbb{R}[[V_c(\mathfrak{C})^*]] \otimes \mathfrak{k}$.*

Proof. Let $p \in \mathfrak{C}$. By Theorem 7.7, there is a formal vector field $\mathbf{w}_{\text{ho}}(p) \in \mathcal{X}_{I^2}$ satisfying $[\mathbf{v}_1(p)_s, \mathbf{w}_{\text{ho}}(p)] = 0$ s.t. $\mathbf{v}(p)$ is formally equivalent to $\mathbf{w}(p) := \mathbf{v}_1(p) + \mathbf{w}_{\text{ho}}(p)$. If $h \in \text{Aut}(R) \subseteq \text{Aut}(\mathfrak{g})$ is a formal diffeomorphism s.t. $\mathbf{w}(p) = h.\mathbf{v}(p)$, then h leaves the constant part $\sigma_0(p)$ of $\sigma(p)$ fixed, so $\text{ev}_0(h.\sigma(p)) = \text{ev}_0(\sigma(p)) = \sigma_0(p)$. Thus, we may assume that $[\mathbf{v}_1(p)_s, \mathbf{w}_{\text{ho}}(p)] = 0$ and $\text{Spec}(\text{ad}_{\sigma_0(p)}) \cap \Sigma_p = \emptyset$. By acting with gauge transformations, we may by Theorem 7.13 further assume that $\sigma \in R \otimes \mathfrak{t}$, where \mathfrak{t} is a maximal torus containing $\sigma_0(p)$. By Theorem 8.14, it follows that $RV_c(p)^\perp \subseteq \ker \bar{\pi}$. The above holds for all $p \in \mathfrak{C}$, so $\text{Span}_{p \in \mathfrak{C}} RV_c(p)^\perp \subseteq \ker \bar{\pi}$. By Lemma 8.23 we know $\text{Span}_{p \in \mathfrak{C}} (V_c(p)^\perp) = V_c(\mathfrak{C})^\perp$, so that $R/(\text{Span}_{p \in \mathfrak{C}} RV_c(p)^\perp) \cong \mathbb{R}[[V_c(\mathfrak{C})^*]]$. \square

Theorem 8.4. *Let $\mathfrak{t} \subseteq \mathfrak{k}$ be a maximal Abelian subalgebra. Let $\bar{\pi}$ be a continuous projective unitary representation of \mathfrak{g}^\sharp . Let $\mathfrak{C} \subseteq \mathfrak{C}(\bar{\pi})$. Assume for every $p \in \mathfrak{C}$ that $\sigma(p) \in R \otimes \mathfrak{t}$ and $[\mathbf{v}_1(p)_s, \mathbf{v}_{\text{ho}}(p)] = 0$. Then $RV_c(\mathfrak{C})^\perp \otimes \mathfrak{k} \subseteq \ker \bar{\pi}$ and hence $\bar{\pi}|_{\mathfrak{g}}$ factors through $\mathbb{R}[[V_c(\mathfrak{C})^*]] \otimes \mathfrak{k}$.*

Proof. By Theorem 8.14 it follows that $\text{Span}_{p \in \mathfrak{C}} RV_c(p)^\perp = RV_c(\mathfrak{C})^\perp \subseteq \ker \bar{\pi}$. \square

8.5 The Case Where \mathfrak{p} is Semisimple.

Let us consider the special case where \mathfrak{p} is simple. Let P be a 1-connected Lie group with $\text{Lie}(P) = \mathfrak{p}$. In this case, suitable normal form theorems for \mathbf{v} and σ are available (see Theorem 7.10 and Theorem 7.12). In particular, $\mathbf{v} : \mathfrak{p} \rightarrow \mathcal{X}_I^{op}$ is always formally equivalent to its linearization \mathbf{v}_1 at $0 \in V$. Similarly, by Theorem 7.12, the vertical twist $\sigma : \mathfrak{p} \rightarrow \mathfrak{g}$ is gauge-equivalent to some Lie algebra homomorphism $\sigma_0 : \mathfrak{p} \rightarrow \mathfrak{k}$. In particular, if \mathfrak{p} is not compact then we may assume that $\sigma = 0$ by acting with gauge transformations if necessary, for in that case there are no homomorphisms $\mathfrak{p} \rightarrow \mathfrak{k}$ (because \mathfrak{k} is compact, see Lemma 7.20). Combined with Theorem 8.4 we immediately obtain Theorem 8.5 below, where $V_c(\mathfrak{C}) := \bigcap_{p \in \mathfrak{C}} V_c(p)$.

Theorem 8.5. *Assume that \mathfrak{p} is non-compact and simple. Let $\bar{\pi}$ be a continuous projective unitary representation of $\mathfrak{g} \rtimes_D \mathfrak{p}$. Write $\mathfrak{C} := \mathfrak{C}(\bar{\pi}) \subseteq \mathfrak{p}$. Then $\bar{\pi}|_{\mathfrak{g}}$ factors through $\mathbb{R}[[V_c(\mathfrak{C})^*]] \otimes \mathfrak{k}$.*

Let $\mathfrak{p} = \mathfrak{k}_0 \oplus \mathfrak{p}_0$ be a Cartan decomposition of \mathfrak{p} , so that \mathfrak{k}_0 and \mathfrak{p}_0 are the $+1$ and -1 eigenspaces of a Cartan involution θ , respectively [Kna96, cor. 6.18]. Let $\mathfrak{a}_0 \subseteq \mathfrak{p}_0$ be a maximal Abelian subalgebra of \mathfrak{p}_0 . According to the Iwasawa decomposition [Kna96, Prop. 6.43], \mathfrak{p} decomposes as $\mathfrak{p} \cong \mathfrak{k}_0 \oplus \mathfrak{a}_0 \oplus \mathfrak{n}_0$, where $\mathfrak{n}_0 \subseteq \mathfrak{p}$ is nilpotent. For $p \in \mathfrak{p}$ we write $p = p_e + p_h + p_n$ for the corresponding decomposition of p , where $p_e \in \mathfrak{k}_0, p_h \in \mathfrak{a}_0$ and $p_n \in \mathfrak{n}_0$. Then $\text{Spec}(\text{ad}_{p_e}) \subseteq i\mathbb{R}$, $\text{Spec}(\text{ad}_{p_h}) \subseteq \mathbb{R}$ and ad_{p_n} is nilpotent [Kna96, Lem. 6.45]. Moreover, \mathfrak{a}_0 is contained in a Cartan subalgebra of \mathfrak{p}_0 [Kna96, cor. 6.47].

Proposition 8.24. *Suppose that \mathfrak{p} is simple and that the \mathfrak{p} -representation \mathbf{v}_1 on V is non-trivial and irreducible. Let $\mathfrak{C} \subseteq \mathfrak{p}$ be an Ad_P -invariant convex cone and let $V_c(\mathfrak{C}) := \bigcap_{p \in \mathfrak{C}} V_c(p)$. Assume that \mathfrak{C} contains some $p \in \mathfrak{C}$ with non-trivial hyperbolic part. Then $V_c(\mathfrak{C}) = \{0\}$.*

Proof. Notice first that as P is 1-connected, the \mathfrak{p} -action $\mathbf{v}_1 : \mathfrak{p} \rightarrow \mathfrak{gl}(V)$ integrates to a continuous representation of P on V . As $\mathfrak{C}(\bar{\pi})$ is Ad_P -invariant, the subspace $V_c(\mathfrak{C})$ is P -invariant. Thus either $V_c(\mathfrak{C}) = \{0\}$ or $V_c(\mathfrak{C}) = V$, so it suffices to show $V_c(\mathfrak{C}) \neq V$. By assumption $p_h \neq 0$. In view of Cartan's unitary trick, see e.g. [Kna01, V. Prop. 5.3], the image of elements in \mathfrak{a}_0 in any finite-dimensional representation are semisimple and have real spectrum. Thus $\text{Spec}(\mathbf{v}_1(p_h)) \subseteq \mathbb{R}$. As \mathfrak{p} is simple and \mathbf{v}_1 is a non-trivial \mathfrak{p} -representation by assumption, it follows that \mathbf{v}_1 is injective. As $\mathbf{v}_1(p_h) \in \mathfrak{gl}(V)$ is semisimple, there exists $0 \neq v \in V$ s.t. $\mathbf{v}_1(p_h)v = \mu v$ for some $0 \neq \mu \in \mathbb{R}$. Thus $0 \neq v \notin V_c(\mathfrak{C})$. Hence $V_c(\mathfrak{C}) \neq V$ and so $V_c(\mathfrak{C}) = \{0\}$. \square

Theorem 8.6. *Assume that \mathfrak{p} is non-compact and simple. Suppose that \mathbf{v}_1 defines a non-trivial irreducible \mathfrak{p} -representation on V . Let $\bar{\pi}$ be a continuous projective unitary representation of $\mathfrak{g} \rtimes_D \mathfrak{p}$. Either $\mathfrak{C}(\bar{\pi})$ is a pointed P -invariant convex cone or $\bar{\pi}|_{\mathfrak{g}}$ factors through \mathfrak{k} .*

Remark 8.25. Notice that if \mathfrak{p} is simple and \mathcal{C} is a closed Ad_P -invariant convex cone which is not pointed, then $\mathcal{C} \cap -\mathcal{C} = \mathfrak{p}$ and hence $\mathcal{C} = \mathfrak{p}$.

Proof of Theorem 8.6: Write $\mathfrak{C} := \mathfrak{C}(\bar{\pi})$. Recall that \mathfrak{C} is an Ad_P -invariant convex cone. Consequently, the edge $\mathfrak{e} := \bar{\mathfrak{C}} \cap -\bar{\mathfrak{C}}$ of the closure $\bar{\mathfrak{C}}$ of \mathfrak{C} is an ideal in \mathfrak{p} . Assume that \mathfrak{C} is not pointed. Then neither is $\bar{\mathfrak{C}}$. As \mathfrak{p} is simple, it follows that $\mathfrak{e} = \mathfrak{p}$ and hence $\bar{\mathfrak{C}} = \mathfrak{p}$. Thus \mathfrak{C} is a dense convex cone in the finite-dimensional real vector space \mathfrak{p} , which implies that $\mathfrak{C} = \mathfrak{p}$. As \mathfrak{p} is non-compact, it contains some hyperbolic element. Thus, so does \mathfrak{C} . By Proposition 8.24 it follows that $V_c(\mathfrak{C}) = \{0\}$ and hence Theorem 8.5 implies that $\bar{\pi}$ factors through \mathfrak{k} . \square

Thus if \mathcal{C} is an Ad_P -invariant convex cone which is not pointed, then \mathfrak{g} admits no continuous projective unitary representations which are of g.p.e. at $\mathcal{C} \subseteq \mathfrak{p}$ other than those which factor through \mathfrak{k} . On the other hand, we know by [Pan81, cor. 2.3] that if \mathfrak{p} is simple, then a non-trivial pointed closed and P -invariant convex cone exists in \mathfrak{p} if and only if \mathfrak{p} is of *hermitian type*, meaning that $\dim(\mathfrak{z}(\mathfrak{k}_0)) = 1$, where $\mathfrak{p} = \mathfrak{k}_0 \oplus \mathfrak{p}_0$ is a Cartan decomposition of \mathfrak{p} and where \mathfrak{k}_0 is the Lie algebra of a compact Lie group. In fact, the proof of [Pan81, cor. 2.3] clarifies that a closed P -invariant convex cone is pointed if and only if it intersects $\mathfrak{z}(\mathfrak{k}_0) \cong \mathbb{R}$ non-trivially. In particular, if \mathfrak{p} is simple and of hermitian type, then up to a sign there is a unique non-trivial closed P -invariant cone that is pointed.

Let us shift our attention to positive energy representations, in which case a different argument is available.

Lemma 8.26. *Suppose that P is a non-compact simple connected linear Lie group. If $(\sigma, \mathcal{H}_\sigma)$ is a unitary P -representation that is norm-continuous, then σ is trivial.*

Proof. As simple real lie groups are of type I [HC53, Thm. 7], we may assume that σ is irreducible. As P is a connected, linear and simple Lie group, every irreducible P -representation is admissible [Kna01, Thm. 8.1]. Thus if $K_{\mathfrak{p}} \subseteq P$ is a maximal compact Lie subgroup of P , then every irreducible $K_{\mathfrak{p}}$ -representation occurs with finite multiplicity in $\sigma|_{K_{\mathfrak{p}}}$. If σ is norm continuous, then $\sigma|_{K_{\mathfrak{p}}}$ can contain only finitely many irreducible $K_{\mathfrak{p}}$ -representations. Thus we conclude that \mathcal{H}_{σ} must be finite-dimensional. But there are no non-trivial homomorphisms from a non-compact simple Lie group into the compact $SU(n)$ by Lemma 7.20. \square

Proposition 8.27. *Suppose that P is non-compact simple, connected and linear. Suppose that the P -action on V is irreducible and non-trivial. Let $\bar{\rho}$ be a continuous projective unitary representation of G which is of p.e. at $\mathcal{C} := \mathfrak{p}$. Then $\bar{\rho}|_G$ factors through K .*

Proof. By Theorem 8.12 it suffices to consider the case where $\bar{\rho}$ factors through G_k for some $k \in \mathbb{N}$. Whitehead's Second Lemma, [Jac79, III.9. Lem. 6], we know that $H^2(\mathfrak{p}, \mathbb{R}) = \{0\}$. Using in addition that P is simply connected, it follows that $\bar{\rho}|_P$ lifts to a continuous unitary representation $\sigma : P \rightarrow U(\mathcal{H}_{\rho})$ of P , so that $\bar{\rho}(p) = [\sigma(p)]$ in $PU(\mathcal{H}_{\rho})$ for all $p \in P$. By Lemma 3.7, the fact that σ is of p.e. at $\mathcal{C} = \mathfrak{p}$ implies that it is norm-continuous. It follows from Lemma 8.26 that $\sigma = \text{id}_{\mathcal{H}}$. Thus $\bar{\rho}(\alpha_p(g)) = \bar{\rho}(g)$ for all $g \in G$ and $p \in P$. It follows that $d\bar{\rho}$ vanishes on $D(\mathfrak{p})\mathfrak{g}$. As \mathfrak{p} acts irreducibly and non-trivially on V , it follows that $D(\mathfrak{p})\mathfrak{g} = I \otimes \mathfrak{k}$. Thus $I \otimes \mathfrak{k} \subseteq \ker d\bar{\rho}$. This implies that $\bar{\rho}|_{G_k}$ factors through K . \square

The following provides a simple example of a projective p.e. representation $\bar{\rho}$ of $G_1 \rtimes P$ s.t. $\bar{\rho}|_{G_1}$ does not factor through K .

Example 8.28. Let $P = \text{Mp}(2, \mathbb{R})$ be the double-cover of $SL(2, \mathbb{R})$. Let P act on $V := \mathbb{R}^2$ via the defining action of $SL(2, \mathbb{R})$. We consider a trivial vertical twist, so that the \mathfrak{p} -action on $\mathfrak{g} = R \otimes \mathfrak{k}$ is given by $D(p) = -\mathcal{L}_{v(p)}$. In this case the generator p_0 of rotations generates the unique (up to a sign) pointed, closed and P -invariant cone \mathcal{C} in \mathfrak{p} . Explicitly, $v(p_0) = y\partial_x - x\partial_y$. Let us construct a non-trivial continuous projective unitary representation of $G_1 \rtimes P \cong (V^* \otimes \mathfrak{k}) \rtimes (K \times P)$ that is of p.e. at the cone $\mathcal{C} \subseteq \mathfrak{p}$. Write $W := V^* \otimes \mathfrak{k}$.

We begin by specifying a suitable 2-cocycle on $V^* \otimes \mathfrak{k} \subseteq \mathfrak{g}_1$. Notice that $(\wedge^2 V)^{\mathfrak{p}} \cong \mathbb{R}$ is one-dimensional.

Let $0 \neq \lambda \in (\wedge^2 V)^{\mathfrak{p}}$ and consider it as a \mathfrak{p} -invariant bilinear map $V^* \times V^* \rightarrow \mathbb{R}$. To be consistent with Proposition 8.9, let us write $\lambda(fdg)$ instead of $\lambda(f, g)$ for $f, g \in V^*$. Let $x, y \in V^*$ be the usual basis of V^* . Then λ is fully specified by the number $\lambda(ydx)$. If $\lambda(ydx) > 0$, then the quadratic form $q(v) := \lambda(\mathcal{L}_{v(p_0)}v, dv)$ is positive-definite, because $q(ax + by) = (a^2 + b^2)\lambda(ydx)$ for $a, b \in \mathbb{R}$. Let ω be the unique symplectic form on W satisfying $\omega(vX, wY) := \lambda(vdw)\kappa(X, Y)$ for $X, Y \in \mathfrak{k}$ and $v, w \in V^*$. Then $\omega(D(p_0)\xi, \xi) \geq 0$ for every $\xi \in W$ (recalling that κ is *negative* definite). Let $H(W, \omega)$ be the corresponding Heisenberg group. Let L_{\pm} be the $\pm i$ -eigenspaces in $W_{\mathbb{C}}$ of the complex structure $\mathcal{J} := D(p_0)$ on $W_{\mathbb{C}}$, so that $W_{\mathbb{C}} = L_- \oplus L_+$. The \mathfrak{p} -invariance of λ ensures that $\mathcal{J}^*\omega = \omega$. Indeed, extend ω \mathbb{C} -bilinearly to $W_{\mathbb{C}}$. As λ is \mathfrak{p} -invariant, it follows that $\omega(\mathcal{J}\xi, \eta) + \omega(\xi, \mathcal{J}\eta) = 0$ for all $\xi, \eta \in W_{\mathbb{C}}$, which implies that $L_{\pm} \subseteq W_{\mathbb{C}}$ are \mathcal{J} -invariant Lagrangian subspaces for ω . Then $\mathcal{J}^*\omega = \omega$ follows from $\mathcal{J}^*\omega(w_+, w_-) = \omega(iw_+, -iw_-) = \omega(w_+, w_-)$ for $w_{\pm} \in L_{\pm}$. Notice further that $\omega(\mathcal{J}\xi, \xi) \geq 0$ holds for all $\xi \in W$, by construction. Equip L_+ with the positive definite hermitian form defined by $\langle v, w \rangle_{L_+} := -2i\omega(\bar{v}, w)$ for $v, w \in L_+$. For each $n \in \mathbb{N}$, equip the symmetric algebra $S^n(L_+)$ with the inner product satisfying

$$\langle v_1 \cdots v_n, w_1 \cdots w_n \rangle := \sum_{\sigma \in S_n} \prod_{k=1}^n \langle v_{\sigma_k}, w_k \rangle_{L_+}, \quad v_k, w_k \in L_+.$$

Let $\mathcal{F} := \overline{S^{\bullet}(L_+)}$ be the Hilbert space closure, where $S^{\bullet}(L_+) = \bigoplus_{n=0}^{\infty} S^n(L_+)$. The metaplectic representation ρ of $H(W, \omega) \rtimes \text{Mp}(W, \omega)$, with $\rho(z) = zI$ on the central \mathbb{T} component, can be realized on the Fock space \mathcal{F} , where $\text{Mp}(W, \omega)$ denotes the metaplectic group [Nee00, Thm X.3.3]. Notice that $SL(2, \mathbb{R}) \hookrightarrow \text{Sp}(W, \omega)$ because λ is \mathfrak{p} -invariant. By pulling back the metaplectic representation we obtain a continuous unitary representation of $H(W, \omega) \rtimes P$ which is of p.e. at \mathcal{C} and does not factor through K .

Appendix

A From Germs to Jets

Let $\mathcal{K} \rightarrow M$ be locally trivial bundle of Lie groups with typical fiber a finite-dimensional Lie group G with Lie algebra \mathfrak{g} . Write $\mathfrak{K} \rightarrow M$ for the corresponding Lie algebra bundle. The following justifies the claim made in Section 1 that any continuous projective unitary representation of $\Gamma_c(\mathcal{K})$ which factors through the germs at a point $a \in M$ actually factors through the ∞ -jets $J_a^\infty(\mathcal{K})$ at $a \in M$. The group $\Gamma_c(\mathcal{K})$ is a locally exponential Lie group modeled on the LF-Lie algebra $\Gamma_c(\mathfrak{K})$ [JN17, Prop. 2.3].

Let $U \subseteq \mathbb{R}^d$ be an open neighborhood of the origin. Let $C_{\text{flat}}^\infty(U)$ denote the kernel of the ∞ -jet projection

$$j_0^\infty : C_c^\infty(U) \rightarrow J_0^\infty(C_c^\infty(U)) \cong \mathbb{R}[[x_1, \dots, x_d]]$$

at $0 \in U$. In the following we show well-known fact that the closure $C_c^\infty(U \setminus \{0\})$ in $C_c^\infty(U)$ is $C_{\text{flat}(a)}^\infty(U)$. As a consequence, we deduce that if a continuous projective unitary representation of the Lie algebra $\Gamma_c(\mathfrak{K})$ factors through the germs at a point $a \in M$, then it factors through the ∞ -jets $J_a^\infty(\mathfrak{K})$ at $a \in M$. In turn, this implies a group level-analogue.

If $K \subset U$ is a compact set, write $C_K^\infty(U)$ for the subspace of $C_c^\infty(U)$ consisting of functions on U with support in K . Then $C_K^\infty(U)$ is the projective limit $C_K^\infty(U) = \varprojlim_n C_K^n(U)$ of the Banach spaces $C_K^n(U)$, which we equip with the norm $\|f\|_{C_K^n(U)} := \sup_{|k| \leq n} \|D^k f\|_{C_K(U)}$, where the supremum runs over all multi-indices $k \in \mathbb{N}_{\geq 0}^d$ with $|k| \leq n$. Then $C_c^\infty(U) := \varinjlim C_K^\infty(U)$ is the corresponding locally convex inductive limit. See e.g. [Rud91, Theorem 6.5] for a description of this topology. For $r > 0$, write $B_r := \{x \in \mathbb{R}^d : \|x\| \leq r\}$ for the closed ball centered at $0 \in \mathbb{R}^d$ with radius r .

Lemma A.1. *The closure of $C_c^\infty(U \setminus \{0\})$ in $C_c^\infty(U)$ is $C_{\text{flat}(a)}^\infty(U)$.*

Proof. As $C_{\text{flat}(a)}^\infty(U) \subseteq C_c^\infty(U)$ is closed and $C_c^\infty(U \setminus \{0\}) \subseteq C_{\text{flat}(a)}^\infty(U)$, it follows that $\overline{C_c^\infty(U \setminus \{0\})} \subseteq C_{\text{flat}(a)}^\infty(U)$. It remains to show the reverse inclusion. Fix $f \in C_{\text{flat}(a)}^\infty(U) \subseteq C_c^\infty(U)$. We show $f \in \overline{C_c^\infty(U \setminus \{0\})}$. Let $K_0 \subseteq M$ be a relatively compact open subset such that $\text{supp } f \subseteq K_0$. Set $K := \overline{K_0}$. We may assume that $0 \in K_0$, for otherwise $f \in C_c^\infty(U \setminus \{0\})$ and we are done. By [Mal67, Lem 4.2], we can find for any multi-index $k \in \mathbb{N}_{\geq 0}^d$ some constant $C_k > 0$ depending only on the multi-index k s.t. for any $0 < r < 1$ with $B_{2r} \subseteq K_0$, there exists a smooth function $\psi_r \in C_c^\infty(\mathbb{R}^d)$ s.t. $\psi_r \geq 0$, $\psi_r|_{B_r} = 0$, $\psi_r|_{(\mathbb{R}^d \setminus B_{2r})} = 1$ and $\sup_{x \in \mathbb{R}^d} |D^k \psi_r(x)| \leq C_k r^{-|k|}$. In particular $f\psi_r \in C_c^\infty(U \setminus \{0\})$ and $\text{supp } f\psi_r \subseteq K$. Moreover observe that $\text{supp}(1 - \psi_r) \subseteq B_{2r}$ and $\|(1 - \psi_r)\|_{C_{B_{2r}}^n(\mathbb{R}^d)} \lesssim r^{-n}$ for some constant depending on $n \in \mathbb{N}_{\geq 0}$, where we used that $0 < r < 1$. On the other hand, suppose that $\alpha \in \mathbb{N}_{\geq 0}^d$ is a multi-index. Since $j_0^\infty(D^\alpha f) = 0$, it follows from Taylor's Theorem that $\|D^\alpha f\|_{C(B_{2r})} \lesssim r^l$ for arbitrary $l \in \mathbb{N}_{\geq 0}$, with a constant depending on f, α and l but not on r . Thus $\|f\|_{C^n(B_{2r})} \lesssim r^l$ for arbitrary $n, l \in \mathbb{N}_{\geq 0}$. In particular $\|f\|_{C^n(B_{2r})} \lesssim r^{n+1}$. Combining the previous observations, we obtain that

$$\|f - f\psi_r\|_{C^n(K)} = \|f(1 - \psi_r)\|_{C^n(K)} = \|f(1 - \psi_r)\|_{C^n(B_{2r})} \lesssim \|f\|_{C^n(B_{2r})} \|(1 - \psi_r)\|_{C^n(B_{2r})} \lesssim r,$$

the constants depending only on f and n but not on r . This shows that $f\psi_r \rightarrow f$ in $C_K^\infty(U)$ as $r \rightarrow 0$. Thus $f\psi_r \rightarrow f$ in $C_c^\infty(U)$. Since $f\psi_r \in C_c^\infty(U \setminus \{0\})$ for every r , we conclude that $f \in \overline{C_c^\infty(U \setminus \{0\})}$. \square

If $a \in M$, define the spaces of smooth section of \mathcal{K} and \mathfrak{K} which are flat at $a \in M$:

$$\Gamma_{\text{flat}(a)}(\mathcal{K}) := \ker \left(j_a^\infty : \Gamma_c(\mathcal{K}) \rightarrow J_a^\infty(\mathcal{K}) \right), \quad \Gamma_{\text{flat}(a)}(\mathfrak{K}) := \ker \left(j_a^\infty : \Gamma_c(\mathfrak{K}) \rightarrow J_a^\infty(\mathfrak{K}) \right).$$

Proposition A.2 below clarifies the apparent ambiguity in the topology on $J_a^\infty(\mathfrak{K})$, for which two candidates are available.

Proposition A.2. *Let $a \in M$. The projective limit topology on $J_a^\infty(\mathfrak{K}) := \varprojlim_k J_a^k(\mathfrak{K})$ coincides with the quotient topology obtained from $J_a^\infty(\mathfrak{K}) \cong \Gamma_c(\mathfrak{K})/\Gamma_{\text{flat}(a)}(\mathfrak{K})$.*

Proof. The continuous k -jet projections $j_a^k : \Gamma_c(\mathfrak{K}) \rightarrow J_a^k(\mathfrak{K})$ at $a \in M$ all descend to continuous maps $\Gamma_c(\mathfrak{K})/\Gamma_{\text{flat}(a)}(\mathfrak{K}) \rightarrow J_a^k(\mathfrak{K})$. By the universal property of the projective limit, they induce a continuous map $\Phi : \Gamma_c(\mathfrak{K})/\Gamma_{\text{flat}(a)}(\mathfrak{K}) \rightarrow J_a^\infty(\mathfrak{K})$. Using Borel's Lemma [Hö3, Thm. 1.2.6], it is not hard to check that this map is bijective. It remains to show it is an open map, which follows immediately from the Open Mapping Theorem [Rud91, cor. 2.12] because $\Gamma_c(\mathfrak{K})/\Gamma_{\text{flat}(a)}(\mathfrak{K})$ and $J_a^\infty(\mathfrak{K})$ are both Fréchet spaces and Φ is bijective and continuous. \square

Proposition A.3. *Let $a \in M$.*

- *The closure of $\Gamma_c(M \setminus \{a\}; \mathfrak{K})$ in $\Gamma_c(M; \mathfrak{K})$ is $\Gamma_{\text{flat}(a)}(\mathfrak{K})$.*
- *The closure of $\Gamma_c(M \setminus \{a\}; \mathcal{K})$ in $\Gamma_c(M; \mathcal{K})$ is $\Gamma_{\text{flat}(a)}(\mathcal{K})$.*

Proof. By a partition of unity argument, we may assume that the bundle $\mathfrak{K} \rightarrow M$ is trivial, that $M \subseteq \mathbb{R}^d$ is open neighborhood of $0 \in \mathbb{R}^d$ and that $a = 0$. Then $\Gamma_c(M; \mathfrak{K}) \cong C_c^\infty(M; \mathfrak{K})$. The claim now follows from Lemma A.1. Notice for the second assertion that $\Gamma_{\text{flat}(a)}(M, \mathcal{K})$ is a locally exponential, being an embedded closed Lie subgroup of the locally exponential Lie group $\Gamma_c(M, \mathcal{K})$. The result is then immediate from the previous point. \square

Proposition A.4. *Let $a \in M$.*

1. *Let $\bar{\pi} : \Gamma_c(M; \mathfrak{K}) \rightarrow \mathcal{L}^\dagger(\mathcal{D})$ be a continuous projective unitary representation on the pre-Hilbert space \mathcal{D} . Assume that $\bar{\pi}$ vanishes on $\Gamma_c(M \setminus \{a\}; \mathfrak{K})$. Then $\bar{\pi}$ factors continuously through $J_a^\infty(\mathfrak{K})$.*
2. *Let $\bar{\rho} : \Gamma_c(M; \mathcal{K}) \rightarrow \text{PU}(\mathcal{H})$ be a continuous projective unitary representation of $\Gamma_c(M; \mathcal{K})$. Assume that $\bar{\rho}$ vanishes on $\Gamma_c(M \setminus \{a\}; \mathcal{K})$. Then $\bar{\rho}$ factors through $\Gamma_{\text{flat}(a)}(M, \mathcal{K})$.*

Proof. For the first point, notice by continuity that $\bar{\pi}$ must also vanish on the closure of $\Gamma_c(M \setminus \{a\}; \mathfrak{K})$ in $\Gamma_c(M; \mathfrak{K})$, which by Proposition A.3 equals $\Gamma_{\text{flat}(a)}(M; \mathfrak{K})$. Thus $\Gamma_c(M; \mathfrak{K})$ factors continuously through the quotient space $J_a^\infty(\mathfrak{K}) \cong \Gamma_c(M; \mathfrak{K})/\Gamma_{\text{flat}(a)}(M; \mathfrak{K})$, where Proposition A.2 was used. The second point is proven similarly using Proposition A.3. \square

References

- [ADGV16] Sergio Albeverio, Bruce Driver, Maria Gordina, and A. Vershik. Equivalence of the brownian and energy representations. *Journal of Mathematical Sciences*, 219:612–630, 12 2016.
- [AHeK78] Sergio Albeverio and Raphael Hø egh Krohn. The energy representation of Sobolev-Lie groups. *Compositio Math.*, 36(1):37–51, 1978.
- [AHeKM⁺93] Sergio A. Albeverio, Raphael J. Hø egh Krohn, Jean A. Marion, Daniel H. Testard, and Bruno S. Torrèsani. *Noncommutative distributions*, volume 175 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker, Inc., New York, 1993. Unitary representation of gauge groups and algebras.
- [AHeKT81] S. Albeverio, R. Hø egh Krohn, and D. Testard. Irreducibility and reducibility for the energy representation of the group of mappings of a Riemannian manifold into a compact semisimple Lie group. *J. Functional Analysis*, 41(3):378–396, 1981.
- [Arn88] V. I. Arnol'd. *Geometrical methods in the theory of ordinary differential equations*, volume 250 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, New York, second edition, 1988. Translated from the Russian by Joseph Szucs [József M. Szücs].
- [AT94] S. Albeverio and B. Torrèsani. Some remarks on representations of jet groups and gauge groups. *J. Math. Phys.*, 35(9):4897–4908, 1994.
- [Bas64] Andrée Bastiani. Applications différentiables et variétés différentiables de dimension infinie. *J. Analyse Math.*, 13:1–114, 1964.
- [BEK80] Ola Bratteli, George A. Elliott, and Akitaka Kishimoto. The temperature state space of a C^* -dynamical system. I. *Yokohama Math. J.*, 28(1-2):125–167, 1980.
- [BEK86] Ola Bratteli, George A. Elliott, and Akitaka Kishimoto. The temperature state space of a C^* -dynamical system. II. *Ann. of Math. (2)*, 123(2):205–263, 1986.
- [Boc45] S. Bochner. Compact groups of differentiable transformations. *Ann. of Math. (2)*, 46:372–381, 1945.
- [BR87] Ola Bratteli and Derek W. Robinson. *Operator algebras and quantum statistical mechanics. 1*. Texts and Monographs in Physics. Springer-Verlag, New York, second edition, 1987. C^* - and W^* -algebras, symmetry groups, decomposition of states.

- [BR97] Ola Bratteli and Derek W. Robinson. *Operator algebras and quantum statistical mechanics. 2.* Texts and Monographs in Physics. Springer-Verlag, Berlin, second edition, 1997. Equilibrium states. Models in quantum statistical mechanics.
- [CG90] Lawrence J. Corwin and Frederick P. Greenleaf. *Representations of nilpotent Lie groups and their applications. Part I*, volume 18 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1990. Basic theory and examples.
- [DK00] J. J. Duistermaat and J. A. C. Kolk. *Lie groups*. Universitext. Springer-Verlag, Berlin, 2000.
- [Dul04] H. Dulac. Recherches sur les points singuliers des équations différentielles. *Journ. École Polyt. Série 2*, 9:1–25, 1904.
- [FM04] Rui Loja Fernandes and Philippe Monnier. Linearization of Poisson brackets. *Lett. Math. Phys.*, 69:89–114, 2004.
- [FV77] M. Fannes and A. Verbeure. Correlation inequalities and equilibrium states. II. *Comm. Math. Phys.*, 57(2):165–171, 1977.
- [GG68] I. M. Gelfand and M. I. Graev. Representations of quaternion groups over locally compact fields and function fields. *Funkcional. Anal. i Priložen*, 2(1):20–35, 1968.
- [GGV77] I. M. Gelfand, M. I. Graev, and A. M. Veršik. Representations of the group of smooth mappings of a manifold X into a compact Lie group. *Compositio Math.*, 35(3):299–334, 1977.
- [Glo05] Helge Glockner. Fundamentals of direct limit Lie theory. *Compos. Math.*, 141(6):1551–1577, 2005.
- [GS68] Victor W. Guillemin and Shlomo Sternberg. Remarks on a paper of Hermann. *Trans. Amer. Math. Soc.*, 130:110–116, 1968.
- [GW84] Roe Goodman and Nolan R. Wallach. Structure and unitary cocycle representations of loop groups and the group of diffeomorphisms of the circle. *J. Reine Angew. Math.*, 347:69–133, 1984.
- [Hö03] Lars Hörmander. *The analysis of linear partial differential operators. I.* Classics in Mathematics. Springer-Verlag, Berlin, 2003. Distribution theory and Fourier analysis, Reprint of the second (1990) edition [Springer, Berlin; MR1065993 (91m:35001a)].
- [Hat02] Allen Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002.
- [HC53] Harish-Chandra. Representations of a semisimple Lie group on a Banach space. I. *Trans. Amer. Math. Soc.*, 75:185–243, 1953.
- [Her68] Robert Hermann. The formal linearization of a semisimple Lie algebra of vector fields about a singular point. *Trans. Amer. Math. Soc.*, 130:105–109, 1968.
- [HM07] Karl H. Hofmann and Sidney A. Morris. *The Lie theory of connected pro-Lie groups*, volume 2 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2007. A structure theory for pro-Lie algebras, pro-Lie groups, and connected locally compact groups.
- [HN09] K. H. Hofmann and K.-H. Neeb. Pro-Lie groups which are infinite-dimensional Lie groups. *Math. Proc. Cambridge Philos. Soc.*, 146(2):351–378, 2009.
- [Ism76] R. S. Ismagilov. Unitary representations of the group $C_0^\infty(X, G)$, $G = \text{SU}_2$. *Mat. Sb. (N.S.)*, 100(142)(1):117–131, 166, 1976.
- [Jac79] Nathan Jacobson. *Lie algebras*. Dover Publications, Inc., New York, 1979. Republication of the 1962 original.
- [JN] Bas Janssens and Karl-Hermann Neeb. Positive energy representations of gauge groups II: Conformal completion of space-time. To appear.
- [JN17] Bas Janssens and Karl-Hermann Neeb. Covariant central extensions of gauge Lie algebras. In *Stochastic geometric mechanics*, volume 202 of *Springer Proc. Math. Stat.*, pages 101–114. Springer, Cham, 2017.

- [JN19] Bas Janssens and Karl-Hermann Neeb. Projective unitary representations of infinite-dimensional Lie groups. *Kyoto J. Math.*, 59(2):293–341, 2019.
- [JN21] Bas Janssens and Karl-Hermann Neeb. Positive energy representations of gauge groups I: Localization. *In preparation*, 2021.
- [Kac90a] Victor G. Kac. *Infinite-dimensional Lie algebras*. Cambridge University Press, Cambridge, third edition, 1990.
- [Kac90b] Victor G. Kac. *Infinite-dimensional Lie algebras*. Cambridge University Press, Cambridge, third edition, 1990.
- [KM97] Andreas Kriegl and Peter W. Michor. *The convenient setting of global analysis*, volume 53 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1997.
- [Kna96] Anthony W. Knaapp. *Lie groups beyond an introduction*, volume 140 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 1996.
- [Kna01] Anthony W. Knaapp. *Representation theory of semisimple groups*. Princeton Landmarks in Mathematics. Princeton University Press, Princeton, NJ, 2001. An overview based on examples, Reprint of the 1986 original.
- [KR87] V. G. Kac and A. K. Raina. *Bombay lectures on highest weight representations of infinite-dimensional Lie algebras*, volume 2 of *Advanced Series in Mathematical Physics*. World Scientific Publishing Co., Inc., Teaneck, NJ, 1987.
- [Lon08] R Longo. Lectures on conformal nets, November 2008.
- [Mai02] Peter Maier. Central extensions of topological current algebras. In *Geometry and analysis on finite- and infinite-dimensional Lie groups*, volume 55 of *Banach Center Publ.*, pages 61–76. Polish Acad. Sci. Inst. Math., Warsaw, 2002.
- [Mal67] B. Malgrange. *Ideals of differentiable functions*, volume 3 of *Tata Institute of Fundamental Research Studies in Mathematics*. Tata Institute of Fundamental Research, Bombay; Oxford University Press, London, 1967.
- [Mil84] J. Milnor. Remarks on infinite-dimensional Lie groups. In *Relativity, groups and topology, II (Les Houches, 1983)*, pages 1007–1057. North-Holland, Amsterdam, 1984.
- [Mor18] Vincenzo Morinelli. The Bisognano-Wichmann property on nets of standard subspaces, some sufficient conditions. *Ann. Henri Poincaré*, 19(3):937–958, 2018.
- [Mun01] J. Mund. The Bisognano-Wichmann theorem for massive theories. *Ann. Henri Poincaré*, 2(5):907–926, 2001.
- [Nee00] Karl-Hermann Neeb. *Holomorphy and convexity in Lie theory*, volume 28 of *De Gruyter Expositions in Mathematics*. Walter de Gruyter & Co., Berlin, 2000.
- [Nee01] Karl-Hermann Neeb. Borel-Weil theory for loop groups. In *Infinite dimensional Kähler manifolds (Oberwolfach, 1995)*, volume 31 of *DMV Sem.*, pages 179–229. Birkhäuser, Basel, 2001.
- [Nee06] Karl-Hermann Neeb. Towards a Lie theory of locally convex groups. *Jpn. J. Math.*, 1(2):291–468, 2006.
- [Nee10a] Karl-Hermann Neeb. On differentiable vectors for representations of infinite dimensional Lie groups. *J. Funct. Anal.*, 259(11):2814–2855, 2010.
- [Nee10b] Karl-Hermann Neeb. Semibounded representations and invariant cones in infinite dimensional Lie algebras. *Confluentes Math.*, 2(1):37–134, 2010.
- [Nee17] Karl-Hermann Neeb. Bounded and semibounded representations of infinite dimensional Lie groups. In *Representation theory—current trends and perspectives*, EMS Ser. Congr. Rep., pages 541–563. Eur. Math. Soc., Zürich, 2017.
- [NO17] Karl-Hermann Neeb and Gestur Ólafsson. Antiunitary representations and modular theory. In *50th Seminar “Sophus Lie”*, volume 113 of *Banach Center Publ.*, pages 291–362. Polish Acad. Sci. Inst. Math., Warsaw, 2017.

- [Pan81] Stephen M. Paneitz. Invariant convex cones and causality in semisimple Lie algebras and groups. *J. Functional Analysis*, 43(3):313–359, 1981.
- [Poi79] Henri Poincaré. *Sur les propriétés des fonctions définies par les équations aux différences partielles*. Gauthier-Villars, 1879.
- [PS76] K. R. Parthasarathy and K. Schmidt. A new method for constructing factorisable representations for current groups and current algebras. *Comm. Math. Phys.*, 50(2):167–175, 1976.
- [PS86] Andrew Pressley and Graeme Segal. *Loop groups*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 1986. Oxford Science Publications.
- [Rud91] Walter Rudin. *Functional analysis*. International Series in Pure and Applied Mathematics. McGraw-Hill, Inc., New York, second edition, 1991.
- [Sch90] Konrad Schmüdgen. *Unbounded operator algebras and representation theory*, volume 37 of *Operator Theory: Advances and Applications*. Birkhäuser Verlag, Basel, 1990.
- [Tak03a] M. Takesaki. *Theory of operator algebras. II*, volume 125 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2003. Operator Algebras and Non-commutative Geometry, 6.
- [Tak03b] M. Takesaki. *Theory of operator algebras. III*, volume 127 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2003. Operator Algebras and Non-commutative Geometry, 8.
- [Tan11] Yoh Tanimoto. Ground state representations of loop algebras. *Ann. Henri Poincaré*, 12(4):805–827, 2011.
- [TL99] Valerio Toledano Laredo. Positive energy representations of the loop groups of non-simply connected Lie groups. *Comm. Math. Phys.*, 207(2):307–339, 1999.
- [Wal87] Nolan R. Wallach. On the irreducibility and inequivalence of unitary representations of gauge groups. *Compositio Math.*, 64(1):3–29, 1987.
- [War72] Garth Warner. *Harmonic analysis on semi-simple Lie groups. I*. Die Grundlehren der mathematischen Wissenschaften, Band 188. Springer-Verlag, New York-Heidelberg, 1972.
- [Was98] Antony Wassermann. Operator algebras and conformal field theory. III. Fusion of positive energy representations of $LSU(N)$ using bounded operators. *Invent. Math.*, 133(3):467–538, 1998.