

# VALUATIONS AND NONZERO TORSION IN MODULE OF DIFFERENTIALS

SARASIJ MAITRA AND VIVEK MUKUNDAN

ABSTRACT. Let  $(R, \mathfrak{m}_R, \mathbb{k})$  be a one-dimensional complete local reduced  $\mathbb{k}$ -algebra over a field of characteristic zero. Berger conjectured that  $R$  is regular if and only if the universally finite module of differentials  $\Omega_R$  is torsion free. When  $R$  is a domain, we prove the conjecture in several cases. Our techniques are primarily reliant on making use of the valuation semi-group of  $R$ . First, we establish a method of verifying the conjecture by analyzing the valuation semi-group of  $R$  and orders of units of the integral closure of  $R$ . We also prove the conjecture in the case when certain monomials are missing from the monomial support of the defining ideal of  $R$ . These monomials are based on the smallest power of  $\mathfrak{m}_R$  that is contained within the conductor ideal. This also generalizes a previous result of [CnGW98].

## 1. INTRODUCTION

Let  $X$  be a (affine or projective) curve over a field  $\mathbb{k}$  of characteristic zero. The classification of circumstances when  $X$  is smooth is of importance to both algebraists and geometers. Though there are other smoothness characterizations, the one involving the differential sheaf  $\Omega_{X,x}$ , where  $x$  is a closed point on  $X$ , has eluded researchers. Consider the completion of the stalk of the structure sheaf  $\mathcal{O}_X$  at  $x$  and denote this by  $\widehat{\mathcal{O}_{X,x}}$ . R.W. Berger conjectured that  $X$  is smooth at  $x$  if and only if the sheaf of differentials  $\widehat{\mathcal{O}_{X,x}} \otimes \Omega_{X,x}$  is torsion free.

Let  $R = \widehat{\mathcal{O}_{X,x}} \cong \mathbb{k}[[X_1, \dots, X_n]]/I$  with  $I \subseteq (X_1, \dots, X_n)^2$ . For a curve, the differential sheaf mentioned above is  $R \otimes \Omega_{X,x}$  where  $\Omega_{X,x}$  is the module of Kähler differentials and  $R \otimes \Omega_{X,x}$  is the universally finite module of differentials, henceforth denoted by  $\Omega_R$ . For the purposes of this article, we can define the universally finite module of differentials  $\Omega_R$  to be the module whose presentation is as follows:

$$R^{\mu(I)} \xrightarrow{\mathcal{J}} R^n \rightarrow \Omega_R \rightarrow 0$$

where  $\mu(I)$  is the minimal set of generators of  $I$  and  $\mathcal{J}$  is the Jacobian matrix on a generating set for  $I$ , considered as a matrix over  $R$ . For further details, we refer the reader to the excellent resource [Kun86]. One of the first pieces of evidence on the smoothness of the curve  $X$  via the torsion freeness of its differential sheaf was studied by E. Kunz ([Kun61]) when the ring is of characteristic  $p$  and the base field is imperfect. Berger ([Ber63]) in 1963 conjectured the following for perfect fields:

**Conjecture A** (R. W. Berger [Ber63]). Let  $\mathbb{k}$  be a perfect field and let  $R$  be a reduced one-dimensional analytic  $\mathbb{k}$ -algebra. Then  $R$  is regular if and only if the torsion submodule  $\tau(\Omega_R)$  of  $\Omega_R$  vanishes.

---

2010 *Mathematics Subject Classification*. Primary: 13A15. Secondary: 13H05.

*Key words and phrases*. module of differentials, Berger Conjecture, reduced curves.

It follows from the above presentation that if  $R$  is regular (i.e.,  $n = 1$ ), then  $\Omega_R$  is free. Hence it is the converse direction that is of interest. Thus we assume  $n \geq 2$  throughout this article.

A direct generalization of this conjecture for higher dimensional surfaces does not hold (see for example [Mat67, Example Pg 160], [Sch70, Satz 9.4]). But with added constraints on  $R$  (such as Cohen-Macaulayness of  $R$ ), a generalized version does exist due to R. Hübl and D. Arapura independently ([H90, Ara89]). There have been numerous instances in which Conjecture A has been established by a variety of different approaches (see [Sch70], [HK71], [Her78] [BG80], [Ulr81] [Koc83], [HW84], [HW86], [Ber88], [Poh89], [Güt90], [Iso91], [Poh91], [CnGW98], [CK99], [HMM21], [Mai22a]). An excellent exposition on the earlier ones among the above listed results can be found in [Ber94]. An Artinian variant of the conjecture, named the Artin-Berger conjecture, was explored in [CnGW98] recovering some of the earlier cases.

Recently, the authors of this article along with C. Huneke [HMM21] demonstrated a new approach for explicitly constructing torsion in  $\Omega_R$ , where  $R$  is assumed to be a domain with residue field  $\mathbb{k}$  of characteristic 0 and embedding dimension  $n$ . In that article, we provided a strategy for first constructing an over-ring  $S$  such that  $R \subseteq S \subseteq \overline{R} = \mathbb{k}[[t]]$ , and embedding dimension of  $S$  is  $n + s$ . Here  $\overline{R}$  denotes the integral closure of  $R$  in its field of fractions  $\text{Quot}(R)$ . We then demonstrated that constructing “sufficiently many” torsion elements in  $\Omega_S$  enables us to pull back a torsion element to  $\Omega_R$ . In Proposition 3.3 of this article, we show that

$$\lambda(\Omega_S) \geq ns + \binom{s}{2}$$

(where  $\lambda(M)$  denotes the length of a module  $M$ ), which is one less than the required number of torsion elements meeting the above sufficiency criteria. This settles [HMM21, Remark 4.10].

Identifying  $R$  within  $\overline{R} = \mathbb{k}[[t]]$  allows us to write  $R = \mathbb{k}[[\alpha_1 t^{a_1}, \dots, \alpha_n t^{a_n}]]$  where  $\alpha_i$ 's are units in  $\overline{R}$ . The valuation semi-group of  $R$ , namely  $v(R)$ , is critical for our methods throughout this article. In fact, the precise valuations  $a_i$  and the valuation of the conductor ideal,  $\mathfrak{C}_R = R :_{\text{Quot}(R)} \overline{R}$ , are critical in our work of constructing torsion elements. Moreover, the “order” of the unit  $\alpha_d$  in the description of  $x_d = \alpha_d t^{a_d}$  also plays a role. Given  $\alpha_d = \sum_{i=0}^{\infty} u_{di} t^i$ , a unit in  $\mathbb{k}[[t]]$ , its order is defined to be

$$o(\alpha_d) = v(\alpha_d - u_{d0}) = \min_{i \geq 1} \{i \mid u_{ji} \neq 0\}.$$

We state our first result here which is quite technical, but we illustrate with examples that the conditions are often satisfied (see Theorem 4.4 and Theorem 4.6).

**Theorem B.** Let  $R = \mathbb{k}[[\alpha_1 t^{a_1}, \dots, \alpha_n t^{a_n}]] \subseteq \overline{R} = \mathbb{k}[[t]]$  be as above with  $\mathfrak{C}_R = (t^{c_R})\overline{R}$  and  $\mathbb{k}$ , an algebraically closed field of characteristic zero. Then  $\tau(\Omega_R)$  is nonzero in the following cases:

- (1) the valuations  $a_n, a_{n-1}$  satisfy  $a_n + a_{n-1} \geq c_R$ .
- (2) there exist  $i, d$  such that  $o(\alpha_d) + a_i \geq c_R$ ,  $o(\alpha_d) < \infty$  and any one of the following statements holds.
  - (a)  $o(\alpha_d) \leq o(\alpha_i) < \infty$ ,
  - (b)  $d \geq i$ .

It is possible to get hold of a torsion element by mapping  $\Omega_R$  to a rank one torsion free module (see Section 2.2). However, proving that it is non-zero is the most challenging

part. Our discussion in Proposition 4.7 provides an approach towards achieving this goal, by going modulo a monomial ideal. We implement this method and obtain our second main result. This is the broadest class where we establish the truth of Berger's Conjecture (see Theorem 4.9).

**Theorem C.** Let  $R = \mathbb{k}[[X_1, \dots, X_n]]/I$  be a one-dimensional local domain where  $\mathbb{k}$  is algebraically closed of characteristic zero and  $I \subseteq (X_1, \dots, X_n)^2$ . Let  $x_i$  denote the image of  $X_i$  in  $R$  and let the conductor be  $\mathfrak{C}_R$ . Suppose  $\mathfrak{m}^N \subseteq \mathfrak{C}_R$  and  $x_1^{N-1}, x_1^{N-2}x_2, N \geq 2$ , are not in the monomial support ideal of  $I$ . Then the torsion submodule  $\tau(\Omega_R)$  is nonzero.

The monomial support ideal of  $I$  is the same as  $\text{Mono}(I)$  appearing in [PUV07]. The theorem recovers the main theorem of [CnGW98]. We also present other interesting examples that satisfy the hypothesis of the above theorem.

This raises the question if condition (2) in Theorem C can be relaxed. So far our methods have been unable to resolve this. Getting a complete answer even in the case when  $\mathfrak{C}_R = \mathfrak{m}^N$  for some large enough  $N$  would be interesting. The methods used in this article are innovative, in that, they explicitly compute torsion elements in  $\Omega_R$ , whereas most of the previous approaches to the Conjecture A have been unable to do so.

**Acknowledgements.** Both authors are extremely grateful to C. Huneke for insightful talks about the article's content. Numerous results in this article are ascribed to the concepts he imparted to us. Additionally, we are obliged to D. Arapura for providing us with a geometric viewpoint on Berger's Conjecture.

## 2. SETTING AND PRELIMINARIES

Throughout this article we will assume that  $\mathbb{k}$  is an algebraically closed field of characteristic 0, and  $(R, \mathfrak{m}_R, \mathbb{k})$  is a one-dimensional complete equicharacteristic local  $\mathbb{k}$ -algebra which is a domain with embedding dimension  $n$  ( $\mu_R(\mathfrak{m}_R) = n$ ). Choosing  $t$  to be a uniformizing parameter of the integral closure  $\overline{R}$  of  $R$ , we may assume that  $\overline{R} = \mathbb{k}[[t]]$ . As a result, we may write  $R = \mathbb{k}[[\alpha_1 t^{a_1}, \dots, \alpha_n t^{a_n}]]$  where  $\alpha_i$ 's are units in  $\overline{R}$  and  $a_1 < a_2 < \dots < a_n$ . Using the technique as in the first paragraph of the proof of [HMM21, Theorem 3.1], we can alternatively write  $R = \mathbb{k}[[t^{a_1}, \alpha_2 t^{a_2}, \dots, \alpha_n t^{a_n}]]$ . We briefly explain the process. First we multiply by a nonzero element of  $\mathbb{k}$  to assume that the constant term of  $\alpha_1$  is 1. Let  $s = \beta t$  where  $\beta^{a_1} = \alpha_1$ . Such a  $\beta$  exists due to Hensel's Lemma [Eis95, Theorem 7.3]. The characteristic of  $\mathbb{k}$  being zero is important here. Notice that under this change of variables,  $\mathbb{k}[[t]] = \mathbb{k}[[s]]$  and  $s = \beta t = t + \beta_1 t^2 + \beta_2 t^3 + \dots$  where  $\beta = 1 + \sum_{i=1}^{\infty} \beta_i t^i$ . Notice that  $s^{a_1} = (\beta t)^{a_1} = \alpha_1 t^{a_1}$ . Also  $\alpha_i t^{a_i} = \alpha'_i s^{a_i}$ . Then  $R = \mathbb{k}[[\alpha_1 t^{a_1}, \dots, \alpha_n t^{a_n}]] \cong \mathbb{k}[[s^{a_1}, \alpha'_2 s^{a_2}, \dots, \alpha'_n s^{a_n}]]$ .

We also define the valuation  $v$  on  $\overline{R}$  given by  $v(p(t)) = a$  if  $p(t) = t^a \alpha$  where  $\alpha$  is a unit in  $\mathbb{k}[[t]]$  (also called  $\text{ord}(-)$ , see for example [SH06, Example 6.7.5]). Let  $v(R)$  denote the valuation semi-group of  $R$ .

The conductor  $\mathfrak{C}_R$  is the largest common ideal of  $R$  and its integral closure,  $\overline{R}$ . Since  $\overline{R} = \mathbb{k}[[t]]$ , we have that  $\mathfrak{C}_R = (t^i)_{i \geq c_R} \overline{R}$  where  $c_R$  is the least integer such that  $t^{c_R-1} \notin R$ , and  $t^{c_R+i} \in R$  for all  $i \geq 0$ . The number  $c_R$  is characterized as the least valuation in  $\mathfrak{C}_R$ . It is clear from this discussion that there cannot be any element  $r \in R$ , such that  $v(r) = c_R - 1$ . Since  $\overline{R}$  is finitely generated over  $R$  ([SH06, Theorem 4.3.4]),  $\mathfrak{C}_R \neq 0$ , and it is never all of  $R$  unless  $R$  is regular.

We define an epimorphism

$$\begin{aligned}\Phi : \mathbb{k}[[X_1, \dots, X_n]] &\twoheadrightarrow R \\ \Phi(X_i) &= \alpha_i t^{a_i} \text{ for } 1 \leq i \leq n.\end{aligned}$$

We denote the kernel of  $\Phi$  by  $I = (f_1, \dots, f_m)$  and hence have the natural isomorphism  $R \cong \mathbb{k}[[X_1, \dots, X_n]]/I$ . Since  $\text{edim } R = n$ ,  $I$  is contained in  $\mathfrak{m}_R^2$  where  $\mathfrak{m}_R = (X_1, \dots, X_n)$ , the maximal ideal of  $\mathbb{k}[[X_1, \dots, X_n]]$ . Such rings are called *analytic  $\mathbb{k}$ -algebras*. We will interchangeably use  $\alpha_i t^{a_i}$  for  $x_i$ , the images of  $X_i$  in the quotient  $\mathbb{k}[[X_1, \dots, X_n]]/I$ .

## 2.1. Universally Finite Module of Differentials.

**Definition 2.1.** Let  $R$  be an analytic one-dimensional  $\mathbb{k}$ -algebra as above, which is a domain. Let  $I = (f_1, \dots, f_m)$  where  $f_j \in P = \mathbb{k}[[X_1, \dots, X_n]]$ ,  $n \geq 2$ . We assume that  $I \subseteq \mathfrak{m}_P^2$  where  $\mathfrak{m}_P = (X_1, \dots, X_n)$ . Then the *universally finite module of differentials over  $\mathbb{k}$* , denoted by  $\Omega_R$ , has a (minimal) presentation given as follows:

$$R^m \xrightarrow{\begin{bmatrix} \frac{\partial f_j}{\partial x_i} \end{bmatrix}} R^n \rightarrow \Omega_R \rightarrow 0$$

where  $\begin{bmatrix} \frac{\partial f_j}{\partial x_i} \end{bmatrix}$  is the Jacobian matrix of  $I$ , with entries in  $R$ .

We refer the reader to the excellent resource [Kum86] for more information. Let  $\tau(\Omega_R)$  denote the torsion submodule of  $\Omega_R$ .

**2.2. Computing Torsion.** It is well-known that  $\text{rank}_R(\Omega_R) = \text{rank}_R(\Omega_{\overline{R}}) = 1$  ( $R$  and  $\overline{R}$  have the same fraction field) and  $\Omega_{\overline{R}}$  is torsion-free. Let  $f : \Omega_R \rightarrow \Omega_{\overline{R}}$  be the natural map induced by the inclusion  $R \hookrightarrow \overline{R}$ . Recall that the torsion submodule of  $\Omega_R$  is the kernel of the homomorphism  $\Omega_R \rightarrow \Omega_R \otimes Q$  where  $Q = \text{Quot}(R)$ . Since  $\Omega_{\overline{R}}$  is torsion-free, we have  $\Omega_{\overline{R}} \hookrightarrow \Omega_R \otimes Q$ . Thus  $\ker f$  is the torsion submodule of  $\Omega_R$  and hence  $\tau(\Omega_R) = \ker f$ . The map  $f$  acts as follows:

$$f(dx_i) = \frac{dx_i}{dt} dt.$$

Since  $\Omega_{\overline{R}}$  is isomorphic to  $\overline{R}$ ,  $\Omega_R$  surjects to an  $R$ -submodule  $\sum_{i=1}^n R \frac{dx_i}{dt}$  of  $\overline{R} = \mathbb{k}[[t]]$ . Thus the torsion submodule,  $\tau(\Omega_R)$ , consists of the column vectors  $[r_1 \dots r_n]^t$  such that  $\sum_{i=1}^n r_i \frac{dx_i}{dt} = 0$ . Evidently,  $\tau(\Omega_R)$  is nonzero precisely when the tuples  $[r_1 \dots r_n]^t$  are not in the image of the presentation matrix (Jacobian matrix of  $I$ ) of  $\Omega_R$ , all entries written in terms of the uniformizing parameter  $t$ .

One of the main issues while constructing torsion is to check whether the column vectors  $[r_1, \dots, r_n]^t$  is nonzero in  $\Omega_R$ . One technique is to show that this column vector is nonzero in  $\Omega_{R/J}$  for some suitable ideal  $J$  in  $R$  which will then imply that it is nonzero in  $\Omega_R$ . Though we have a more general theorem for this purpose in Proposition 4.7, a version which we use in some of the results in the sequel is the following theorem.

**Theorem 2.2.** [CnGW98, Proposition 2.6] Suppose  $R = \mathbb{k}[[\alpha_1 t^{a_1}, \dots, \alpha_n t^{a_n}]]$  with maximal ideal  $\mathfrak{m}_R = (x_1, \dots, x_n)$ . Then the torsion  $\tau(\Omega_{R/\mathfrak{m}_R^2})$  has  $\binom{n}{2}$   $\mathbb{k}$ -linearly independent torsion elements  $\overline{x_i} d\overline{x_j}$  for  $i < j$ . Here  $\overline{x_i}$  denotes the image of  $x_i$  in  $R/\mathfrak{m}_R^2$ .

We often avoid using the  $\overline{\phantom{x}}$  for notational convenience.

**2.3. Quasi-homogeneous rings.** These rings form an important class which satisfies the Berger conjecture. A ring  $R$  is said to be quasi-homogeneous, if there is a surjection  $\Omega_R \rightarrow \mathfrak{m}_R$ . Kunz and Ruppert in [KR77] showed that such rings are completions of graded rings. Scheja ([Sch70]) showed that the torsion submodule of the module of differentials of such rings are always nonzero. In fact, he showed that  $\mu(\tau(\Omega_R)) \geq \binom{\text{edim } R}{2}$  when  $R$  is quasi-homogeneous.

**2.4. The extension  $R[\frac{\mathfrak{C}_R}{x_1}]$ .** The basic facts about the finite birational extension  $S = R[\frac{\mathfrak{C}_R}{x_1}]$  of  $R$  are discussed in detail in [HMM21, Section 4].

We write  $\mathfrak{C}_R/x_1$  to denote the set of elements of the form  $c/x_1$  where  $c \in \mathfrak{C}_R$ . Using [HK71, Proposition 2.9] and [HMM21, Lemma 4.1], we get that  $\mathfrak{C}_R/x_1$  is generated as an  $R$ -submodule of  $\overline{R}$  (in fact, it is the ideal  $(t^{c-a_1}\overline{R})$ ) by  $t^{b_i}$  with  $b_1 < \dots < b_s$  where  $b_i \in [c - a_1, c - 1] \setminus v(R)$ . For further details, see [HMM21, Notation 4.2]. It turns out that  $\mathfrak{C}_R/x_1$  is the conductor of the ring  $S$ . In other words the conductor  $\mathfrak{C}_S$  equals the ideal  $(t^{c-a_1}\overline{R})$ . A complete description of the ring  $S$  is as follows:

**Theorem 2.3.** [HMM21, Theorem 4.6] Let  $\mathfrak{C}_R \subseteq \mathfrak{m}_R^2$ . Set  $P = \mathbb{k}[[X_1, \dots, X_n]]$ . Choose  $S, b_1, \dots, b_s$  as described above. Then there exists a presentation of  $S$  as follows:

$$S = R\left[\frac{\mathfrak{C}_R}{x_1}\right] = \frac{\mathbb{k}[[X_1, \dots, X_n, T_1, \dots, T_s]]}{\ker \Phi + (X_i T_j - g_{ij}(X_1, \dots, X_n), T_k T_l - h_{kl}(X_1, \dots, X_n))_{\substack{1 \leq i \leq n, 1 \leq j \leq s \\ 1 \leq k \leq l \leq s}}}$$

where  $g_{ij}(X_1, \dots, X_n), h_{kl}(X_1, \dots, X_n) \in (X_1, \dots, X_n)^2 P$  for all  $i, j, k, l$ .

Moreover,  $g_{ij}(x_1, \dots, x_n), h_{kl}(x_1, \dots, x_n) \in \mathfrak{C}_R$ . In particular, when  $\mathfrak{C}_R \subseteq \mathfrak{m}_R^2$ , it follows that  $\text{edim } S = n + s$ .

### 3. FINER COMPUTATIONS OF TORSION ELEMENTS IN $\Omega_S$

One of the main results in this section is the existence of torsion elements in  $\Omega_S$ . We show that we can in fact construct  $ns + \binom{s}{2}$   $\mathbb{k}$ -linearly independent torsion elements in  $\Omega_S$  which completes [HMM21, Remark 4.10].

The following theorem establishes that once  $ns + \binom{s}{2} + 1$  torsion elements are constructed with entries in  $\mathfrak{m}_S$ , then we can pull back a nonzero torsion element from  $\Omega_S$  to  $\Omega_R$ .

**Theorem 3.1.** [HMM21, Theorem 4.9] Let  $R = \mathbb{k}[[\alpha_1 t^{a_1}, \dots, \alpha_n t^{a_n}]]$  with conductor  $\mathfrak{C}_R \subseteq \mathfrak{m}_R^2$ . Construct  $S = R\left[\frac{\mathfrak{C}_R}{x_1}\right] = R[T_1, \dots, T_s]$  as in Theorem 2.3. Let  $\tau(\Omega_R), \tau(\Omega_S)$  represent the torsion submodules of  $\Omega_R, \Omega_S$  respectively. If  $\lambda(\tau(\Omega_S)) \geq ns + \binom{s}{2} + 1$  and all these torsion elements have non-units in the last  $s$  rows (corresponding to  $dT_1, \dots, dT_s$ ), then a  $\mathbb{k}$ -linear combination of these torsion elements can be pulled back to a nonzero torsion element in  $\tau(\Omega_R)$ . In particular,  $\tau(\Omega_R) \neq 0$ .

**Remark 3.2.** An obvious consequence of this theorem is that if  $\lambda(\tau(\Omega_S)) \geq ns + \binom{s}{2} + k, k \geq 1$  and all these torsions have non-units in the last  $s$  rows, then we can pull back  $k$  torsion elements to  $\Omega_R$ , each of which are  $\mathbb{k}$ -linear combinations of the  $ns + \binom{s}{2} + k$  torsion elements. Notice that these  $k$  torsion elements are  $\mathbb{k}$ -linearly independent in  $\Omega_R$  as they are  $\mathbb{k}$ -linearly independent in  $\Omega_S$ . Thus  $\lambda(\tau(\Omega_R)) \geq k$ .

It is easy to see that we can construct  $\binom{s}{2}$  torsion elements, namely  $\Gamma_{ij} := b_j T_j dT_i - b_i T_i dT_j$ . This can be clearly seen as  $T_i$  are monomials  $t^{b_i}$  in  $S$ . Now Proposition 3.3 shows that we can indeed construct  $ns$  torsion elements in  $\Omega_S$  too.

If  $x_i$  is a monomial  $t^{a_i}$  to begin with, then we can construct the torsion element  $a_i x_i dT_j - b_j T_j dx_i$ . This is clearly nonzero in  $\Omega_S$ . But often  $x_i$  are not of this form and hence the need for the following theorem.

**Proposition 3.3.** Let  $R = \mathbb{k}[\alpha_1 t^{a_1}, \dots, \alpha_n t^{a_n}]$  with conductor  $\mathfrak{C}_R$ . Construct  $S = R \left[ \frac{\mathfrak{C}_R}{x_1} \right] = R[T_1, \dots, T_s]$ . Then

$$(3.1) \quad [x_i, T_j] = \begin{bmatrix} -a_i x_i \partial f_{ij} / \partial x_1 \\ \vdots \\ b_j (T_j + f_{ij}(\underline{x}) + \sum_{k=1}^s \delta_k T_k) - a_i x_i \partial f_{ij} / \partial x_i \\ \vdots \\ -a_i x_i (1 + \delta_j) \\ \vdots \\ -a_i x_i \delta_s \end{bmatrix} \begin{array}{c} \text{basis} \\ \vdots \\ dx_1 \\ \vdots \\ dx_i \\ \vdots \\ dT_j \\ \vdots \\ dT_s \end{array}$$

where  $f_{ij} \in (x_1, \dots, x_n)$  and  $\delta_r \in \mathbb{k}$ , are  $\mathbb{k}$ -linearly independent nonzero torsion elements in  $\tau(\Omega_S)$ .

*Proof.* We follow the process of ‘‘monomialization’’ as in [HMM21, Theorem 3.1] (see Section 2). Let  $\beta = \alpha_i^{1/a_i}$  and  $s = \beta t$  (since the characteristic of  $\mathbb{k}$  is 0, by Hensel’s lemma [Eis95, Theorem 7.3], there exists such a  $\beta \in \overline{R}$ ). Thus  $R \cong \mathbb{k}[\alpha'_1 s^{a_1}, \alpha'_2 s^{a_2}, \dots, s^{a_i}, \dots, \alpha'_n s^{a_n}]$ . Under this isomorphism (change of variables), let the new variables be  $x'_1, \dots, x'_n, T'_1, \dots, T'_s$ . The valuations do not change under this isomorphism. Since  $T'_1, \dots, T'_s \in \mathfrak{C}_S$ , we assume that, without loss of generality, under another change of variables,  $T'_1, \dots, T'_s$  are monomials  $s^{b_1}, \dots, s^{b_s}$ . Notice that under this change of variables  $x'_i = s^{a_i}$ . Clearly, for  $j = 1, \dots, s$  we have

$$(3.2) \quad b_j T'_j d(x'_i) - a_i x'_i d(T'_j)$$

is a nonzero torsion element of  $\Omega_S$  (the nonzero part can be obtained by showing that  $\overline{b_j T'_j d(x'_i) - a_i x'_i d(T'_j)} \equiv (b_j - a_i) \overline{T'_j d x'_i}$  in  $\Omega_{S/\mathfrak{m}_S^2}$ , which is nonzero using [CnGW98, Proposition 2.6]).

Now we convert it back to the original variables  $x_1, \dots, x_n, T_1, \dots, T_s$ . Under the change of variables,  $x_i$  is mapped to  $x'_i$ . So we just need to convert  $T'_j$  back to  $T_j$ . We have  $T'_j = s^{b_j} = \beta^{b_j} t^{b_j} = t^{b_j} + \sum_k \gamma_k t^{b_j+k}$ . Since  $\mathfrak{C}_S = (t^{c_R - a_1}) \overline{R}$  and  $c_R - a_1$  is smaller than  $b_j$ , we can replace  $t^{b_j+k}$  as a function of  $x_1, \dots, x_n, T_1, \dots, T_s$ . Thus we have

$$T'_j = t^{b_j} + \sum_k \gamma_k t^{b_j+k} = T_j + g_{ij}(x_1, \dots, x_n, T_1, \dots, T_s).$$

Due to the nature of the defining ideal of  $S$ ,

$$(3.3) \quad g_{ij}(x_1, \dots, x_n, T_1, \dots, T_s) = f_{ij}(x_1, \dots, x_n) + \sum_{k>j}^s \delta_{ik} T_k$$

where  $\delta_{ik} \in \mathbb{k}$ . In the latter equality, only  $T_k$  with  $k > j$  appear because the only nonzero terms which can appear in  $g_{ij}(\underline{x}, \underline{T})$  are those which have valuations more than that of  $T_j$ .

Therefore

$$dT'_j = dT_j + \sum_{u=1}^n \frac{\partial f_{ij}}{\partial x_u} dx_u + \sum_{k>j}^s \delta_{ik} dT_k$$

Putting the above equation in (3.2) we have

$$(3.4) \quad [x_i, T_j] = \begin{bmatrix} -a_i x_i \partial f_{ij} / \partial x_1 \\ \vdots \\ b_j(T_j + f_{ij}(\underline{x}) + \sum_{k>j}^s \delta_{ik} T_k) - a_i x_i \partial f_{ij} / \partial x_i \\ \vdots \\ -a_i x_i \\ -a_i x_i \delta_{ij+1} \\ \vdots \\ -a_i x_i \delta_{is} \end{bmatrix} \begin{array}{l} \overbrace{\hspace{1cm}}^{\text{basis}} \\ dx_1 \\ \vdots \\ dx_i \\ \vdots \\ dT_j \\ dT_{j+1} \\ \vdots \\ dT_s \end{array}.$$

Next we show that  $[x_i, T_j], 1 \leq i \leq n, 1 \leq j \leq s$  are  $\mathbb{k}$ -linearly independent. First let  $f_{ij}(x_1, \dots, x_n) = \sum_k \beta_{ijk} x_k + f'_{ij}(x_1, \dots, x_n)$  where  $f'_{ij}(x_1, \dots, x_n) \in \mathfrak{m}_R^2, \beta_{ijk} \in \mathbb{k}$ . Now consider the torsion elements  $[x_i, T_j]$  in  $\Omega_{S/\mathfrak{m}_S^2}$ . These would take the form

$$(3.5) \quad \begin{bmatrix} -a_i x_i \beta_{ij1} \\ \vdots \\ -a_i x_i \beta_{iji-1} \\ b_j(T_j + \sum_{k \neq i} \beta_{ijk} x_k + \sum_{k>j}^s \delta_{ijk} T_k) \\ -a_i x_i \beta_{iji+1} \\ \vdots \\ -a_i x_i \beta_{ijn} \\ 0 \\ \vdots \\ 0 \\ -a_i x_i \\ \vdots \\ -a_i x_i \delta_{ijs} \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \\ (b_j - a_i)T_j + \sum_{k>j}^s \delta_{ijk}(b_j - a_i)T_k - (b_j + a_i) \sum_{k \leq i} \beta_{ijk} x_k \\ -(b_j + a_i)x_i \beta_{iji+1} \\ \vdots \\ -(b_j + a_i)x_i \beta_{ijn} \\ 0 \\ \vdots \\ 0 \end{bmatrix},$$

where the equality is due to the fact that  $x_1 x_i, \dots, x_{i-1} x_i, x_i T_1, \dots, x_i T_s \in \mathfrak{m}_S^2$  (the latter containment implies that  $x_j dx_i = -x_i dx_j, j \leq i$  and  $x_i dT_j = -T_j dx_i$  in  $\Omega_{S/\mathfrak{m}_S^2}$ ). So to show that  $[x_i, T_j], 1 \leq i \leq n, 1 \leq j \leq s$  are  $\mathbb{k}$ -linearly independent, it is enough to show that these torsions are  $\mathbb{k}$ -linearly independent in  $\Omega_{S/\mathfrak{m}_S^2}$ .

We first show that  $[x_i, T_j], 1 \leq j \leq s$ , for a fixed  $i$ , are  $\mathbb{k}$ -linearly independent. It is known that  $x_k dx_l, T_j dx_i, 1 \leq k < l \leq n, 1 \leq j \leq s$  are  $\mathbb{k}$ -linearly independent in  $\Omega_{S/\mathfrak{m}_S^2}$  [CnGW98, Proposition 2.6, Corollary 2.7]. Now writing the coefficients of part of the ordered basis



$x_k dx_l, T_j dx_i, 1 \leq k < l \leq n, 1 \leq j \leq s$  in (3.5), we get a  $s \times \binom{n}{2} + ns$  matrix

$$\begin{bmatrix} [x_i, T_1] \\ [x_i, T_2] \\ [x_i, T_3] \\ \vdots \\ [x_i, T_s] \end{bmatrix} = \begin{bmatrix} 0 & \cdots & 0 & -(b_1 + a_i)\beta_{i11} & \cdots & -(b_1 + a_i)\beta_{i1n} & 0 & \cdots & 0 & 0 & \cdots & 0 & b_1 - a_i & \delta_{i12}(b_1 - a_i) & \cdots & \delta_{i1s}(b_1 - a_i) & 0 & \cdots & 0 \\ 0 & \cdots & 0 & -(b_2 + a_i)\beta_{i21} & \cdots & -(b_1 + a_i)\beta_{i2n} & 0 & \cdots & 0 & 0 & \cdots & 0 & 0 & b_2 - a_i & \cdots & \delta_{i2s}(b_2 - a_i) & 0 & \cdots & 0 \\ 0 & \cdots & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 & 0 & 0 & \cdots & \delta_{i3s}(b_3 - a_i) & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots \\ 0 & \cdots & 0 & -(b_s + a_i)\beta_{is1} & \cdots & -(b_1 + a_i)\beta_{isn} & 0 & \cdots & 0 & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 & b_s - a_i & 0 & \cdots & 0 \end{bmatrix} \begin{bmatrix} x_1 dx_2 \\ \vdots \\ x_{n-1} dx_n \\ T_1 dx_1 \\ \vdots \\ T_s dx_n \end{bmatrix}$$

Assume the first block is  $A_i$  which is a  $s \times \binom{n}{2}$  matrix, and the second block is  $[0 \cdots 0 B_i 0 \cdots 0]$  which is a  $s \times ns$  matrix, where  $B_i$  is an upper triangular  $s \times s$  matrix. The columns of  $A_i$  correspond to the ordered basis  $x_k dx_l, 1 \leq k < l \leq n$  and that of  $[0 \cdots 0 B_i 0 \cdots 0]$  correspond to the ordered basis  $T_j dx_i, 1 \leq j \leq s$ . Clearly the matrix  $B_i$  has rank  $s$  as  $a_i, b_1, \dots, b_s$  are distinct and nonzero. Since the rows are  $\mathbb{k}$ -linearly independent,

$$[x_i, T_j] = \left( (b_j - a_i)T_j + \sum_{k>j}^s \delta_{ijk}(b_j - a_i)T_k - (b_j + a_i) \sum_{k \leq i} \beta_{ijk} x_k \right) dx_i - \sum_{k=i+1}^n a_i x_i \beta_{ijk} dx_k$$

are  $\mathbb{k}$ -linearly independent for a fixed  $i$  and  $1 \leq j \leq s$ .

If we consider all the torsion elements  $[x_i, T_j]$  for arbitrary  $i, j$ , then we get the following matrix

$$\begin{bmatrix} [x_1, T_1] \\ \vdots \\ [x_n, T_s] \end{bmatrix} = \begin{bmatrix} A_1 & B_1 & 0 & 0 & \cdots & 0 \\ A_2 & 0 & B_2 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ A_n & 0 & 0 & 0 & \cdots & B_n \end{bmatrix} \begin{bmatrix} x_1 dx_2 \\ \vdots \\ x_{n-1} dx_n \\ T_1 dx_1 \\ \vdots \\ T_s dx_n \end{bmatrix}$$

where each of the  $B_i$  are upper triangular invertible  $s \times s$  matrix. Thus the rows are  $\mathbb{k}$ -linearly independent and hence  $[x_i, T_j]$ 's are  $\mathbb{k}$ -linearly independent.  $\square$

**Theorem 3.4.** With the hypothesis as in Proposition 3.3, we have

$$\lambda(\tau(\Omega_S)) \geq ns + \binom{s}{2}.$$

*Proof.* We observe that  $\Gamma_{ij} = b_j T_j dT_i - b_i T_i dT_j$  and  $[x_i, T_j]$  are  $\mathbb{k}$ -linearly independent: suppose

$$0 = \sum_{i,j} k_{ij} [x_i, T_j] + \sum_{r,s} k'_{rs} \Gamma_{rs}, k_{ij}, k'_{rs} \in \mathbb{k}.$$

Now let  $J = \langle x_1, \dots, x_n \rangle + \langle T_1, \dots, T_s \rangle^2$ . Then in  $\Omega_{S/J}$ ,  $[x_i, T_j] = 0$  (using (3.5)). Thus in  $\Omega_{S/J}$ , we have  $0 = \sum_{r,s} k'_{rs} \Gamma_{rs}$ . Since  $\Gamma_{rs}$  are  $\mathbb{k}$ -linearly independent ([CnGW98, Proposition 2.6]), we have  $k'_{rs} = 0$ . Thus we have  $0 = \sum_{i,j} k_{ij} [x_i, T_j]$  in  $\Omega_S$ . Hence  $k_{ij} = 0$  by Proposition 3.3. Thus  $\lambda(\tau(\Omega_S)) \geq ns + \binom{s}{2}$ .  $\square$

**Remark 3.5.** In [HMM21, Theorem 4.11], we established that  $\tau(\Omega_R)$  is nonzero if  $S = R \left[ \frac{\mathfrak{c}_R}{x_1} \right]$  is quasi-homogeneous. Recall that quasi-homogeneous means that there is a surjection  $\Omega_R \rightarrow \mathfrak{m}_R$ . In fact, the proof mentioned above proceeded by showing that there are  $\binom{n+s}{2}$   $\mathbb{k}$ -linearly independent torsion elements in  $\Omega_S$  and none of these have units in the last  $s$  rows. However, notice that  $\binom{n+s}{2} = ns + \binom{s}{2} + \binom{n}{2}$ . Thus we can pull back  $\binom{n}{2}$  torsions to  $\Omega_R$  using Theorem 3.1 and Remark 3.2, i.e.,  $\lambda(\tau(\Omega_R)) \geq \binom{n}{2}$ .



4. VALUE SEMI-GROUP OF  $R$  AND NONZERO TORSION

In this section, we provide partial answers to Berger's conjecture by carefully observing the value semi-group of  $R$  as well as order of the units  $\alpha_j$  in  $\mathbb{k}[[t]]$ .

We begin by stating a useful test of checking the nonvanishing of torsion in  $\Omega_R$ . First notice that units in  $\mathbb{k}[[t]]$  are power series of the form  $\sum_{i=0}^{\infty} u_i t^i$  where  $u_0 \neq 0$ .

**Notation 4.1.** Let  $\alpha_j = \sum_{i=0}^{\infty} u_{ji} t^i \in \mathbb{k}[[t]]$  be a unit. We denote

$$o(\alpha_j) = v(\alpha_j - u_{j0}) = \min_{i \geq 1} \{i \mid u_{ji} \neq 0\},$$

where  $v(\cdot)$  denotes the order valuation (also called  $\text{ord}(\cdot)$ , see for example [SH06, Example 6.7.5]). If  $\alpha_j = u_{j0} \in \mathbb{k}$ , then notice that  $o(\alpha_j) = \infty$ .

**Theorem 4.2.** Let the conductor of  $R$  be  $\mathfrak{C}_R = (t^{c_R})\overline{R}$ . Suppose there exists  $i, d$  such that  $o(\alpha_d) + a_i \geq c_R$  and  $o(\alpha_d) < \infty$ . Then the torsion  $\tau(\Omega_R)$  is nonzero if any one of the following statements holds.

- (1)  $o(\alpha_d) \leq o(\alpha_i) < \infty$ ,
- (2)  $d \geq i$ .

*Proof.* As in [HMM21, Theorem 3.1], we write  $R = \mathbb{k}[[t^{a_1}, \alpha_2 t^{a_2}, \dots, \alpha_n t^{a_n}]]$  with conductor  $\mathfrak{C}_R = (t^{c_R})\overline{R}$ . Notice that the  $o(\alpha_i)$  remain the same by applying the above change of variables. For (1), observe that  $o(\alpha_i) + a_i \geq o(\alpha_d) + a_i \geq c_R$ . Now

$$x_i = \alpha_i t^{a_i} = \alpha_{i0} t^{a_i} + t^{o(\alpha_i) + a_i} b$$

where  $b \in \overline{R}$ . Since  $o(\alpha_i) + a_i \geq c_R$ ,  $t^{o(\alpha_i) + a_i} b \in \mathfrak{C}_R \subseteq R$ . Thus,  $t^{a_i} = \frac{1}{\alpha_{i0}} (x_i - t^{o(\alpha_i) + a_i} b) \in R$ . Thus, we get that  $R = k[[t^{a_1}, \alpha_2 t^{a_2}, \dots, t^{a_i}, \dots, \alpha_n t^{a_n}]]$ . The proof is now complete using [HMM21, Remark 3.3].

For (2), first observe that by our setup,  $o(\alpha_d) + a_d \geq o(\alpha_d) + a_i \geq c_R$ . Apply the same argument as above to see that  $R = k[[t^{a_1}, \alpha_2 t^{a_2}, \dots, t^{a_d}, \dots, \alpha_n t^{a_n}]]$ . This finishes the proof.  $\square$

The following is an example of a non-quasi-homogeneous ring  $R$  where the above result can be quickly applied.

**Example 4.3.** Let  $R = \mathbb{k}[[t^8 + t^9, t^9 + t^{15}, t^{12} + t^{20}, t^{14}]] = \mathbb{k}[[t^8(1+t), t^9(1+t^6), t^{12}(1+t^8), t^{14}]]$ . Macaulay2 computations show that the conductor is  $(t^{c_R})\overline{R} = (t^{20})\overline{R}$ . With  $i = 3, d = 3$ , we see that  $a_i + o(\alpha_d) = 12 + 8 = 20 = c_R$ . Thus by the above theorem, the torsion  $\tau(\Omega_R)$  is nonzero. Moreover, Macaulay 2 computations show that  $v(\text{trace}(\Omega_R)) = 16$  where  $\text{trace}(\cdot)$  refers to the trace ideal (see [Lin17] for details). Hence,  $R$  is not quasi-homogeneous [Mai22a, Conclusion 2].

The second part of the next result appears as [Mai22b, Theorem 4.9].

**Theorem 4.4.** Suppose the valuations  $a_1, \dots, a_n$  of  $x_1, \dots, x_n$  of  $R$  satisfies one of the following:

- (1)  $a_1 + a_n \geq c_R$ ,
- (2)  $a_{n-1} + a_n \geq c_R + a_1$

Then the torsion  $\tau(\Omega_R)$  is nonzero.

*Proof.* Throughout the proof, we can assume that  $\mathfrak{C}_R \subseteq \mathfrak{m}_R^2$  as otherwise the conclusion follows from [HMM21, Theorem 3.1]. Further applying the same ‘‘monomialization’’ as in the proof of Theorem 4.2, we can assume that  $R = k[[t^{a_1}, \alpha_2 t^{a_2}, \dots, \alpha_n t^{a_n}]]$ .

(1): Consider  $S = R \left[ \frac{\mathfrak{C}_R}{x_1} \right]$  with conductor  $\mathfrak{C}_S = (t^{c-a_1})\overline{R}$ . Since  $a_n \geq c - a_1$ ,  $x_n$ , by change of variables, can be made into monomial  $t^{a_n}$ . Thus  $t^{a_n} = x'_n = x_n - \sum \delta_i T_i - g(\underline{x})$ ,  $\delta_i \in k$ . Then  $S = k[[x_1, \dots, x_{n-1}, x'_n, T_1, \dots, T_s]] = k[[t^{a_1}, \alpha_2 t^{a_2}, \dots, \alpha_{n-1} t^{a_{n-1}}, t^{a_n}, t^{b_1}, \dots, t^{b_s}]]$ . Clearly  $\tau = a_n x'_n dx_1 - a_1 x_1 dx'_n$  is a nonzero torsion in  $\Omega_S$ . Hence (using the notations as in Section 3)  $\tau, [x_i, T_j], \Gamma_{ij}$  are  $ns + \binom{s}{2} + 1$  torsion elements in  $\Omega_S$ . If we show that these torsion elements are  $k$ -linearly independent, then the result follows from Theorem 3.1.

Consider

$$(4.1) \quad \gamma\tau + \sum k_{ij}[x_i, T_j] + \sum k'_{ij}\Gamma_{ij} = 0.$$

First let  $J = \langle x_1, \dots, x'_n \rangle + \mathfrak{m}_S^2$ ; then in  $\Omega_{S/J}$ , (4.1) takes the form  $\sum k'_{ij}\Gamma_{ij} = 0$ . But these are  $k$ -linearly independent and hence  $k'_{ij} = 0$ . Now let  $J_1 = \langle x'_n \rangle + \mathfrak{m}_S^2$ . Then in  $\Omega_{S/J_1}$ ,  $\tau = 0$  as  $x'_n \in J_1$  and  $dx'_n = 0$  in  $\Omega_{S/J_1}$ . Thus (4.1) now becomes  $\sum_{i \neq n} k_{ij}[x_i, T_j] = 0$ . Since we already proved  $[x_i, T_j]$ 's are  $k$ -linearly independent, we have  $k_{ij} = 0$  for  $i \neq n, 1 \leq j \leq s$ . Now (4.1) becomes  $\gamma\tau + \sum_j k_{nj}[x'_n, T_j] = 0$ . Now let  $J_2 = \langle x_1 \rangle + \mathfrak{m}_S^2$ . Then in  $\Omega_{S/J_2}$ , we have again  $\tau = 0$ , but  $0 \neq [x'_n, T_j] = a_n x'_n dT_j - b_j T_j dx'_n = (a_n + b_j)x'_n dT_j$ . Thus we have  $k_{nj} = 0$  as well. We now have  $\gamma\tau = 0$  which implies  $\gamma = 0$ . Thus  $\tau, [x_i, T_j], \Gamma_{ij}$  are  $k$ -linearly independent.

(2): In  $\Omega_{\overline{R}} \cong k[[t]]$  where  $\overline{R} = k[[t]]$ , we have

$$(4.2) \quad \begin{aligned} & \frac{dx_n}{dt} \left( \frac{dx_1}{dt} \right) - \frac{dx_1}{dt} \left( \frac{dx_n}{dt} \right) = 0 \\ & \left( a_n \alpha_n t^{a_n-1} + t^{a_n} \frac{d(\alpha_n)}{dt} \right) \left( \frac{dx_1}{dt} \right) - a_1 t^{a_1-1} \left( \frac{dx_n}{dt} \right) = 0 \end{aligned}$$

Multiplying (4.2) by  $\alpha_{n-1} t^{a_{n-1}-a_1+1}$ , we get

$$(4.3) \quad \begin{aligned} & \left( a_n \alpha_n \alpha_{n-1} t^{a_n-1+a_{n-1}-a_1+1} + \alpha_{n-1} t^{a_n+a_{n-1}-a_1+1} \frac{d\alpha_n}{dt} \right) \frac{dx_1}{dt} - a_1 \alpha_{n-1} t^{a_1-1+a_{n-1}-a_1+1} \frac{dx_n}{dt} = 0 \\ & \left( a_n \alpha_n \alpha_{n-1} t^{a_n+a_{n-1}-a_1} + \alpha_{n-1} t^{a_n+a_{n-1}-a_1+1} \frac{d\alpha_n}{dt} \right) \frac{dx_1}{dt} - a_1 \alpha_{n-1} t^{a_n-1} \frac{dx_n}{dt} = 0 \\ & \left( a_n \alpha_n \alpha_{n-1} t^{a_n+a_{n-1}-a_1} + \alpha_{n-1} t^{a_n+a_{n-1}-a_1+1} \frac{d\alpha_n}{dt} \right) \frac{dx_1}{dt} - a_1 x_{n-1} \frac{dx_n}{dt} = 0 \end{aligned}$$

The order valuation of  $u = a_n \alpha_n \alpha_{n-1} t^{a_n+a_{n-1}-a_1} + \alpha_{n-1} t^{a_n+a_{n-1}-a_1+1} \frac{d\alpha_n}{dt}$  is at least  $a_n + a_{n-1} - a_1$ . Since  $a_n + a_{n-1} - a_1 \geq c_R$ , we get that  $u \in \mathfrak{C}_R \subseteq R$ . Thus (4.3), as a member of  $\Omega_R$ , takes the form

$$(4.4) \quad \begin{bmatrix} u \\ 0 \\ \vdots \\ 0 \\ a_1 x_{n-1} \end{bmatrix}$$

Since  $\mathfrak{C}_R \subseteq \mathfrak{m}_R^2$ , consider the torsion element (4.4) in  $\Omega_{R/\mathfrak{m}_R^2}$ . It takes the form  $a_1 x_{n-1} d(x_n)$  which is nonzero due to [CnGW98, Proposition 2.6, Corollary 2.7].  $\square$

**Remark 4.5.** An immediate consequence of the above theorem is that, henceforth, we can always assume  $a_n < c_R - a_1$ . This in turn results in the simplification of the description of  $[x_i, T_j]$  (3.5) in  $\Omega_{S/\mathfrak{m}_S^2}$ . First, consider the description of  $f_{ij}(x_1, \dots, x_n)$  in (3.3). The valuations of the terms in  $f_{ij}(x_1, \dots, x_n)$  are more than that of  $T_j$ , or more than  $c_R - a_1$ . Since  $a_n < c_R - a_1$ , no linear terms can appear in  $f_{ij}(x_1, \dots, x_n)$ . Thus  $f_{ij}(x_1, \dots, x_n) \in \mathfrak{m}_R^2$  and hence  $\beta_{ijk} = 0$ . Thus (3.5) takes the form

$$(4.5) \quad \left[ \begin{array}{c} 0 \\ \vdots \\ 0 \\ (b_j - a_i)T_j + \sum_{k>j}^s \delta_{ik}(b_j - a_i)T_k \\ 0 \\ \vdots \\ 0 \end{array} \right] \begin{array}{c} \overbrace{\hspace{1cm}}^{\text{basis}} \\ dx_1 \\ \vdots \\ dx_{i-1} \\ dx_i \\ dx_{i+1} \\ \vdots \\ dT_s \\ \vdots \end{array}$$

in  $\Omega_{S/\mathfrak{m}_S^2}$ . This simpler form will help us to prove  $\mathbb{k}$ -linear independence of the torsion elements.

**Theorem 4.6.** Suppose the valuations  $a_1, \dots, a_n$  of  $x_1, \dots, x_n$  of  $R$  satisfy  $a_n + a_{n-1} \geq c_R$ , then the torsion  $\tau(\Omega_R)$  is nonzero.

*Proof.* Throughout the proof, we can assume that  $\mathfrak{C}_R \subseteq \mathfrak{m}_R^2$  as otherwise the conclusion follows from [HMM21, Theorem 3.1].

Consider  $S = R \left[ \frac{\mathfrak{C}_R}{x_1} \right]$  with conductor  $\mathfrak{C}_S = (t^{c_R - a_1})\overline{R}$ . If  $x_n \in \mathfrak{C}_S$ , then  $a_n \geq c_S = c_R - a_1$  and hence we are done by Theorem 4.4(1).

So it is enough to assume  $x_n \notin \mathfrak{C}_S$ . Since  $a_{n-1} + a_n \geq c_S + a_1 = (c_R - a_1) + a_1$ , we use the same steps as in Theorem 4.4(2) and end up at the torsion element (4.4), now in  $\Omega_S$  (we put zeros in the last  $s$  rows corresponding to  $dT_1, \dots, dT_s$ ). This torsion element is nonzero as it is nonzero in  $\Omega_{S/(\mathfrak{C}_S + \mathfrak{m}_S^2)}$ .

Let  $\tau$  denote this torsion element. Next we show that  $\tau, \mathfrak{r}_{ij} = [x_i, T_j], \Gamma_{ij} = [T_i, T_j]$  (as in Section 3) are  $\mathbb{k}$ -linearly independent. In the description of  $\tau$ , let  $u = f_u(x_1, \dots, x_n) + \sum_{k=1}^s \gamma_k T_k, \gamma_k \in \mathbb{k}$ . Since  $x_n \notin \mathfrak{C}_S$  and  $u \in \mathfrak{C}_S$ , we have  $f_u(x_1, \dots, x_n) \in \mathfrak{m}_R^2$ . Thus  $\tau$  in  $\Omega_{S/\mathfrak{m}_S^2}$ ,

will take the form

$$\tau = \begin{bmatrix} \sum_{k=1}^s \gamma_k T_k \\ 0 \\ \vdots \\ 0 \\ a_1 x_{n-1} \\ 0 \\ \vdots \\ 0 \end{bmatrix} \begin{array}{c} \overbrace{\hspace{1.5cm}}^{\text{basis}} \\ dx_1 \\ dx_2 \\ \vdots \\ dx_{n-1} \\ dx_n \\ dT_1 \\ \vdots \\ dT_s \end{array} .$$

Let  $J = \mathfrak{m}_S^2 + \langle T_1, \dots, T_s \rangle$ . In  $\Omega_{S/J}$ , the above torsion takes the form  $a_1 x_{n-1} dx_n$  which is nonzero by [CnGW98, Proposition 2.6]. Now suppose  $\tau = \sum_{i,j} k_{ij} [x_i, T_j] + \sum_{i,j} k'_{ij} \Gamma_{ij}$ ,  $k_{ij}, k'_{ij} \in \mathbb{k}$ . In  $\Omega_{S/J}$ ,  $[x_i, T_j] = \Gamma_{ij} = 0$  (see (4.5)) and this contradicts the fact that  $\tau$  is nonzero in  $\Omega_{S/J}$ . Thus  $\tau$  is  $\mathbb{k}$ -linearly independent with  $[x_i, T_j], \Gamma_{ij}$ . Now the result follows from [HMM21, Theorem 4.9].  $\square$

**Proposition 4.7. (Monomial Case)** Let  $K = (m_1, \dots, m_s)$  be a monomial ideal minimally generated by monomials  $m_i = X_1^{b_{i1}} \cdots X_n^{b_{in}}$  in  $\mathbb{k}[[X_1, \dots, X_n]]$  and  $R' = \mathbb{k}[[X_1, \dots, X_n]]/K$ . Suppose  $m = X_1^{b_1} \cdots X_n^{b_n} \notin K$  where  $b_u \geq 1$  for some  $1 \leq u \leq n$ . Then  $mdx_u = 0$  in  $\Omega_{R'}$  if and only if  $X_u^{b_u+1} \in K$  or  $\frac{\partial(mX_u)}{\partial X_i} \in K$  for all  $1 \leq i \neq u \leq n$ . Here  $x_i$  denotes the image of  $X_i$  in  $R'$ .

*Proof.* First, let  $mdx_u = 0$ . Without loss of generality we assume that  $u = 1, i = 2$ . Suppose by contradiction,  $X_1^{b_1+1} \notin K$  and  $\frac{mX_1}{X_2} \notin K$ . Here we replaced  $\frac{\partial(mX_u)}{\partial X_i} = \frac{\partial(mX_1)}{\partial X_2}$  by  $\frac{mX_1}{X_2}$ .

Consider the monomial ideal  $L = (X_1^{b_1+1} X_2^{b_2}) + K = (X_1^{b_1+1} X_2^{b_2}, m_1, \dots, m_s)$ . First suppose that  $m_i$ , for some  $1 \leq i \leq s$ , divides  $X_1^{b_1+1} X_2^{b_2}$ . Then  $m_i = X_1^u X_2^v$  where  $u \leq b_1 + 1, v \leq b_2$ . If  $u < b_1 + 1, v \leq b_2$ , then  $m \in (m_i) \subseteq K$ , a contradiction. Now suppose  $u = b_1 + 1, v < b_2$ , then  $\frac{mX_1}{X_2} \in (m_i) \subseteq K$ , a contradiction. Therefore, in these cases, none of the  $m_i$ 's divide  $X_1^{b_1+1} X_2^{b_2}$ . Finally if  $u = b_1 + 1, v = b_2$ , then  $m_i = X_1^{b_1+1} X_2^{b_2}$  and hence  $K = L$ . Let  $(X_1^{b_1+1} X_2^{b_2}, m_1, \dots, m_l), l \leq s$  be a minimal generating set for  $L$ . Also, clearly  $\frac{mX_1}{X_2} = X_1^{b_1+1} X_2^{b_2-1} X_2^{b_3} \cdots X_n^{b_n} \notin L$ .

Now since  $mdx_1 = 0$  in  $\Omega_{R'}$ , we also have  $mdx_1 = 0$  in  $\Omega_{\mathbb{k}[[X_1, \dots, X_n]]/L}$ . Thus

$$mdx_1 = rd(x_1^{b_1+1} x_2^{b_2}) + \sum_{i=1}^l r_i dm_i, r, r_i \in \mathbb{k}[[X_1, \dots, X_n]]/L$$

(here we have again used  $x_i$  to denote the images in the suitable quotient). Therefore,

$$(4.6) \quad m = r \frac{\partial(x_1^{b_1+1} x_2^{b_2})}{\partial x_1} + \sum_{i=1}^l r_i \frac{\partial m_i}{\partial x_1} = r(b_1 + 1)x_1^{b_1} x_2^{b_2} + \sum_{i=1}^l r_i b_{i1} x_1^{b_{i1}-1} x_2^{b_{i2}} \cdots x_n^{b_{in}}.$$

$$(4.7) \quad 0 = r \frac{\partial(x_1^{b_1+1} x_2^{b_2})}{\partial x_k} + \sum_{i=1}^l r_i \frac{\partial m_i}{\partial x_k} \text{ for } 2 \leq k \leq n.$$

First we analyze (4.6). Suppose  $\frac{\partial m_i}{\partial x_1}$  divides  $m$ , then we have  $b_{i1} - 1 \leq b_1$  and  $b_{ik} \leq b_k$  for  $2 \leq k \leq n$ . If  $b_{i1} - 1 < b_1$ , then  $m \in K$  which leads to a contradiction. Therefore,  $b_{i1} = b_1 + 1$ . Now if  $b_{i2} < b_2$ , then  $\frac{mX_1}{X_2} \in K$  which is again a contradiction. Thus  $b_{i2} = b_2$ . But now this forces  $m_i \in (X_1^{b_1+1}X_2^{b_2})$  which is also contradiction as  $(X_1^{b_1+1}X_2^{b_2}, m_1, \dots, m_l)$  is a minimal generating set for  $L$ . Thus  $\frac{\partial m_i}{\partial x_1}$  does not divide  $m$ . This shows that  $m_i$ 's have the property that either  $b_{i1} > b_1 + 1$  or  $b_{ik} > b_k$  for some  $2 \leq k \leq n$ .

From the above discussion and (4.6), we have  $r = \frac{1}{b_1+1}x_3^{b_3} \dots x_n^{b_n}$ . Using  $k = 2$  in (4.7), we have

$$\begin{aligned} 0 &= r \frac{\partial(x_1^{b_1+1}x_2^{b_2})}{\partial x_2} + \sum_{i=1}^s r_i \frac{\partial m_i}{\partial x_2} \\ &= rb_2 x_1^{b_1+1} x_2^{b_2-1} + \sum_{i=1}^s r_i b_{i2} x_1^{b_{i1}} x_2^{b_{i2}-1} \dots x_n^{b_{in}} \\ &= \frac{b_2}{b_1+1} \frac{mX_1}{x_2} + \sum_{i=1}^s r_i b_{i2} x_1^{b_{i1}} x_2^{b_{i2}-1} \dots x_n^{b_{in}} \end{aligned}$$

By our choice of  $m_i$  (preceeding discussion on choice of  $b_{ik}$ ), we have that the term  $\frac{b_2}{b_1+1} \frac{mX_1}{x_2}$  cannot cancel with any term in the sum  $\sum_{i=1}^s r_i b_{i2} x_1^{b_{i1}} x_2^{b_{i2}-1} \dots x_n^{b_{in}}$ . Thus  $\frac{b_2}{b_1+1} \frac{mX_1}{x_2} = 0$  or in other words,  $\frac{b_2}{b_1+1} \frac{mX_1}{X_2} \in L$ . Since  $\frac{b_2}{b_1+1} \in \mathbb{k}$ , this forces  $\frac{mX_1}{X_2} \in L$ , which leads to a contradiction. Thus we have either  $X_1^{b_u+1} \in K$  or  $\frac{\partial(mX_u)}{\partial X_i} \in K$ .

Conversely, suppose  $\frac{\partial(mX_u)}{\partial X_i} \in K$  for all  $1 \leq i \neq u \leq n$ . This implies  $mX_u \in K$  and hence  $d(mX_u) = 0$  in  $\Omega_{R'}$ . Hence  $(b_u + 1)mdx_u + \sum_{i=1, i \neq u}^n b_i \frac{mX_u}{x_i} dx_i \in KdR' + dK$ . Using the assumption, we have  $(b_u + 1)mdx_u \in KdR' + dK$ . Also, clearly, if  $X_u^{b_u+1} \in K$ , then  $mdx_u = 0$ , completing the proof.  $\square$

In the following, we use the notion of monomial support of an ideal which is the same as  $\text{Mono}(I)$  in [PUV07]. The ideal  $\text{Mono}(I)$  is defined to be the smallest monomial ideal containing  $I$  and is generated by the monomial support of a generating set for  $I$ .

**Example 4.8.** If  $I = (X^3 - YZ, X^2Y - Z^2, XZ - Y^2) \subseteq S = \mathbb{k}\langle\langle X, Y, Z \rangle\rangle$ , then  $\text{Mono}(I) = (X^3, YZ, X^2Y, Z^2, XZ, Y^2)$ . Notice that  $I$  is the defining ideal of  $\mathbb{k}\langle\langle t^3, t^4, t^5 \rangle\rangle$ . Clearly, we have  $\tau = 3xdz - 5zdx \in \tau(\Omega_{S/I})$ . We can now use Proposition 4.7 with  $K = \text{Mono}(I)$  and  $mdx_u = xdz$  to see that  $\tau$  is nonzero in  $\Omega_{S/\text{Mono}(I)}$ . Hence, it is nonzero in  $\Omega_{S/I}$ .

**Theorem 4.9.** Let  $R = \mathbb{k}\langle\langle \alpha_1 t^{a_1}, \dots, \alpha_n t^{a_n} \rangle\rangle \cong \mathbb{k}\langle\langle X_1, \dots, X_n \rangle\rangle / I$  with conductor  $\mathfrak{C}_R = (t^{c_R})\overline{R}$ . Let  $\tau(\Omega_R)$  denote the torsion submodule of  $\Omega_R$  and  $x_i$  denote the images of  $X_i$  in  $R$ . Let  $N \geq 2$  be the integer such that  $\mathfrak{m}^N \subseteq \mathfrak{C}_R$  and suppose that  $x_1^{N-1}, x_1^{N-2}x_2 \notin \text{Mono}(I)$  where  $I$  is the defining ideal of  $R$ . Then the torsion submodule  $\tau(\Omega_R)$  is nonzero.

We can always choose  $N$  to be the least integer satisfying all the conditions in the hypothesis. For example, if  $k$  is the least integer such that  $\mathfrak{m}^k \subseteq \mathfrak{C}_R$  and  $k \leq N$ , then  $x_1^{k-1}, x_1^{k-2}x_2 \notin \text{Mono}(I)$  (as  $x_1^{n-1}, x_1^{N-2}x_2 \notin \text{Mono}(I)$ ) as well. Thus, we will assume  $N$  is least henceforth. The hypothesis now guarantees that  $x_1^{N-1} \notin \mathfrak{C}_R$ . Before we go to the proof of the theorem, we present a small lemma which helps us reduce to the case that  $x_1^{N-2}x_2 \notin \mathfrak{C}_R$ .

**Lemma 4.10.** Under the conditions of the above theorem, if  $x_1^{N-2}x_2 \in \mathfrak{C}_R$ , then the torsion  $\tau(\Omega_R)$  is nonzero.

*Proof.* Since  $x_1^{N-2}x_2 \in \mathfrak{C}_R$ , we have  $t^{(N-2)a_1+a_2} \in \mathfrak{C}_R$ . Thus there exists  $f(\underline{x})$  such that  $t^{(N-2)a_1+a_2} = x_1^{N-2}x_2 - f(\underline{x})$ . Since the valuation  $v(x_1^{N-2}x_2) = a_1(N-2) + a_2$ , the valuation of  $f(\underline{x})$  is more than  $(N-2)a_1 + a_2$ . Now we know that  $a_1(t^{a_1})dt^{(N-2)a_1+a_2} - ((N-2)a_1 + a_2)t^{(N-2)a_1+a_2}dt^{a_1} = 0$ . Thus we have the torsion element

$$\begin{aligned} & a_1x_1d(x_1^{N-2}x_2 - f(\underline{x})) - ((N-2)a_1 + a_2)x_1^{N-2}x_2 - f(\underline{x}) dx_1 \\ &= a_1x_1 \left( (N-2)x_1^{N-3}x_2dx_1 + x_1^{N-2}dx_2 - \sum_i \frac{\partial f}{\partial x_i} dx_i \right) - ((N-2)a_1 + a_2)x_1^{N-2}x_2 - f(\underline{x}) dx_1 \\ &= (N-2)a_1x_1^{N-2}x_2dx_1 + a_1x_1^{N-1}dx_2 - \sum_i a_1x_1 \frac{\partial f}{\partial x_i} dx_i - ((N-2)a_1 + a_2)x_1^{N-2}x_2 - f(\underline{x}) dx_1 \\ &= (-a_2x_1^{N-2}x_2 + f(\underline{x})) dx_1 + a_1x_1^{N-1}dx_2 - \sum_i a_1x_1 \frac{\partial f}{\partial x_i} dx_i. \end{aligned}$$

Let  $J = \langle x_3, \dots, x_n \rangle + \text{Mono}\langle p \in R \mid v(p) > (N-2)a_1 + a_2 \rangle + \text{Mono}(I)$  (we treat the last part as an ideal in  $R$ ; we keep writing  $\text{Mono}(I)$  for notational convenience) where  $I$  is the defining ideal of  $R$ . Note that  $J$  is a monomial ideal not containing  $x_1^{N-1}, x_1^{N-2}x_2$  as these are not in  $\text{Mono}(I)$  and its valuations are less than or equal to  $(N-2)a_1 + a_2$ . Also notice that  $f(\underline{x}) \in J$  as the valuation of  $f(\underline{x})$  is more than  $(N-2)a_1 + a_2$ . Now in  $\Omega_{R/J}$ , we show that the above torsion element takes the form  $(-a_2x_1^{N-2}x_2) dx_1 + a_1x_1^{N-1}dx_2$ . First,  $dx_i = 0$  for  $i \geq 3$  and hence it is enough to show that  $\sum_{i=1}^2 a_1x_1 \frac{\partial f}{\partial x_i} dx_i = 0$  in  $\Omega_{R/J}$ .

Since  $f(\underline{x})$  has valuation more than  $(N-2)a_1 + a_2$ ,  $a_1x_1 \frac{\partial f}{\partial x_1}$  is either 0 or has valuation more than  $a_1 + (N-3)a_1 + a_2 = (N-2)a_1 + a_2$ . Thus it vanishes in  $R/J$ . So all that remains is  $a_1x_1 \frac{\partial f}{\partial x_2}$  which is either 0 or has valuation more than  $a_1 + (N-2)a_1 = (N-1)a_1 = v(x_1^{N-1})$ . Since  $x_1^{N-1} \notin \text{Mono}(I)$  and  $x_i \in J$  for  $i \geq 3$ , we need to only focus on the case when one of the terms in the product  $a_1x_1 \frac{\partial f}{\partial x_2}$  is possibly  $x_1^{N-2}x_2$ . This in turn shows that  $f(\underline{x})$  has a term  $\beta x_1^{N-3}x_2^2, \beta \in \mathbb{k}$ . But the valuation shows that  $x_1^{N-3}x_2^2 \in J$ . Hence  $a_1x_1 \frac{\partial f}{\partial x_1} dx_1 + a_1x_1 \frac{\partial f}{\partial x_2} dx_2 = 0 + \beta a_1x_1d(x_1^{N-3}x_2^2) = 0$  in  $\Omega_{R/J}$ , i.e.,  $\sum_i a_1x_1 \frac{\partial f}{\partial x_i} dx_i = 0$  in  $\Omega_{R/J}$  as well.

Now in  $\Omega_{R/J}$ , the torsion element  $(-a_2x_1^{N-2}x_2) dx_1 + a_1x_1^{N-1}dx_2$  simplifies to  $\omega = -(a_2 + a_1(N-1))x_1^{N-2}x_2dx_1$ . This is because  $x_1^{N-1}x_2 \in J$  and hence  $x_1^{N-1}dx_2 + (N-1)x_1^{N-2}x_2dx_1 = 0$  in  $\Omega_{R/J}$ . Now Proposition 4.7 can be applied with  $m = x_1^{N-2}x_2$  to see that  $\omega$  is nonzero.  $\square$

*Proof of Theorem 4.9.* Due to the previous lemma, we assume  $x_1^{N-1}, x_1^{N-2}x_2 \notin \mathfrak{C}_R$ . Also, we can safely assume that we have ‘‘monomialized’’  $x_1$  so that  $R = k[[t^{a_1}, \alpha_2 t^{a_2}, \dots, \alpha_n t^{a_n}]]$ .

Consider

$$\begin{aligned} & \frac{dx_2}{dt} \left( \frac{dx_1}{dt} \right) - \frac{dx_1}{dt} \left( \frac{dx_2}{dt} \right) = 0 \\ & (\alpha_2 a_2 t^{a_2-1} + t^{a_2} \frac{\partial \alpha_2}{\partial t}) \left( \frac{dx_1}{dt} \right) - a_1 t^{a_1-1} \left( \frac{dx_2}{dt} \right) = 0 \end{aligned}$$

Multiply both sides by  $t^{(N-2)a_1+1}$  to get

$$(4.8) \quad \begin{aligned} & (\alpha_2 a_2 t^{(N-2)a_1+a_2} + t^{(N-2)a_1+a_2+1} \frac{\partial \alpha_2}{\partial t}) \left( \frac{dx_1}{dt} \right) - a_1 t^{(N-1)a_1} \left( \frac{dx_2}{dt} \right) = 0 \\ & (a_2 x_2 x_1^{N-2} + t^{(N-2)a_1+a_2+1} \frac{\partial \alpha_2}{\partial t}) \left( \frac{dx_1}{dt} \right) - a_1 x_1^{N-1} \left( \frac{dx_2}{dt} \right) = 0 \end{aligned}$$

We have  $S = R[T_1, \dots, T_s]$  and  $\mathfrak{C}_S = (t^{c-a_1})\overline{R}$ . Since  $(N-2)a_1 + a_2 \geq c - a_1$  ( $\mathfrak{m}^N \subseteq \mathfrak{C}_R$ ), we have that the above torsion element is a member of  $\tau(\Omega_S)$ .

The above torsion element takes the form:

$$\tau = \begin{bmatrix} a_2 x_2 x_1^{N-2} + t^{(N-2)a_1+a_2+1} \frac{\partial \alpha_2}{\partial t} \\ -a_1 x_1^{N-1} \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

For the defining ideal  $L$  of  $S$ , let the monomial support be denoted by  $\text{Mono}(L)$ .

We first show that  $\tau$  is nonzero in  $\Omega_S$ . Let  $J' = \langle x_3, \dots, x_n, T_1, \dots, T_s \rangle + \text{Mono}(L) + \text{Mono}(p \in S \mid v(p) > (N-2)a_1 + a_2)$ . It is a monomial ideal. Now consider  $\tau$  in  $\Omega_{S/J'}$ . Since the valuation of  $t^{(N-2)a_1+a_2+1} \frac{\partial \alpha_2}{\partial t}$  is more than  $(N-2)a_1 + a_2$ , it will be of the form

$$\overline{\tau} = \begin{bmatrix} a_2 x_1^{N-2} x_2 \\ -a_1 x_1^{N-1} \\ 0 \\ \vdots \\ 0 \end{bmatrix} \in \Omega_{S/J'}.$$

By hypothesis,  $x_1^{N-2} x_2, x_1^{N-1}$  are not in the monomial support of the defining ideal of  $R$ . Now the rest of the defining ideal of  $S$  is of the form  $x_i T_j - g_{ij}(\underline{x}), T_i T_j - h_{ij}(\underline{x})$ . The valuation of  $x_i T_j$  is  $a_i + b$  where  $b \geq c_R - a_1$ . Since  $x_1^{N-1} \notin \mathfrak{C}_R$ , the valuation  $c_R > (N-1)a_1$ . Thus for  $i \geq 2$ , we have  $a_i + b \geq a_i + c_R - a_1 > a_i + (N-2)a_1 \geq a_2 + (N-2)a_1 = v(x_1^{N-2} x_2) > v(x_1^{N-1})$ . Thus the monomial support of  $g_{ij}(\underline{x}), i \geq 2$  can never have  $x_1^{N-2} x_2, x_1^{N-1}$  as terms. Also, the valuation of  $x_1 T_j$  is at least  $c_R > (N-2)a_1 + a_2$  (as  $x_1^{N-2} x_2 \notin \mathfrak{C}_R$ ). Thus  $g_{1j}(\underline{x})$  cannot have  $x_1^{N-2} x_2, x_1^{N-1}$  as terms as well. Applying similar arguments to analyze the elements  $T_i T_j - h_{ij}(\underline{x})$ , we conclude that  $x_1^{N-2} x_2, x_1^{N-1}$  cannot appear in the monomial support of  $L$ . Thus these terms do not lie in  $J'$  too.

Since  $x_1^{N-1} x_2 \in J'$ , we have  $d(x_1^{N-1} x_2) = 0$  in  $\Omega_{S/J'}$ . Thus we have  $(N-1)x_1^{N-2} x_2 dx_1 = -x_1^{N-1} dx_2$ . Using this equality in the above torsion element we have

$$(4.9) \quad \begin{aligned} \overline{\tau} &= (a_2 x_1^{N-2} x_2) dx_1 - a_1 x_1^{N-1} dx_2 \\ &= (a_2 x_1^{N-2} x_2) dx_1 + a_1 (N-1) x_1^{N-2} x_2 dx_1 \\ &= (a_2 + a_1 (N-1)) x_1^{N-2} x_2 dx_1. \end{aligned}$$

Clearly  $(a_2 + a_1(N-1)) \neq 0$ , and  $x_1^{N-2} x_2 \notin J'$ . Using Proposition 4.7, we get that the above torsion element is nonzero in  $\Omega_{S/J'}$  and hence  $\tau$  is nonzero in  $\Omega_S$ .



Next we show that  $\tau$  is  $\mathbb{k}$ -linearly independent with the torsion elements  $[x_i, T_j], \Gamma_{ij}$  (using the notations as in Section 3). Suppose

$$(4.10) \quad \tau = \sum_{i,j} k_{ij}[x_i, T_j] + \sum_{i,j} k'_{ij}\Gamma_{ij}$$

where  $k_{ij}, k'_{ij} \in \mathbb{k}$ .

Let  $J'' = \langle x_1 \rangle + \mathfrak{m}_S^2$ . Then in  $\Omega_{S/J''}$ , the torsion element  $\tau$  (4.8) is zero. Since  $[x_1, T_j] = a_1 x_1 dT_j - b_j T_j dx_1$ , this element is also zero in  $\Omega_{S/J''}$ . Thus in  $\Omega_{S/J''}$ , Equation (4.10) is of the form  $0 = \sum_{i \geq 2, j \geq 1} k_{ij} \overline{[x_i, T_j]} + \sum_{i,j} k'_{ij} \overline{\Gamma_{ij}}$ . Now consider the description of  $\overline{[x_i, T_j]}, i \geq 2, j \geq 1$  in (4.5). Following the same proof as in Theorem 3.4, it is easy to see that  $\overline{[x_i, T_j]}, i \geq 2, j \geq 1$  and the  $\overline{\Gamma_{ij}}$  are still  $\mathbb{k}$ -linearly independent in  $\Omega_{S/J''}$ . It follows that  $k_{ij} = 0$  for  $i \geq 2, j \geq 1$  and also  $k'_{ij} = 0$  for  $1 \leq i, j \leq s$ . Thus we have  $\tau = \sum_{j \geq 1} k_{ij}[x_1, T_j]$  in  $\Omega_S$ . Now consider the ideal  $J' = \langle x_3, \dots, x_n, T_1, \dots, T_s \rangle + \text{Mono}(L) + \text{Mono}(p \in S \mid v(p) > (N-2)a_1 + a_2)$  from the previous part of the proof. In  $\Omega_{S/J'}$ ,  $\tau$  is nonzero which we already proved above. But  $\overline{[x_1, T_j]} = 0$  in  $\Omega_{S/J'}$  as  $T_j \in J'$ . Thus we arrive at  $\tau = 0$  in  $\Omega_{S/J'}$ , a contradiction. This proves the  $\mathbb{k}$ -linear independence of  $\tau, [x_i, T_j], \Gamma_{ij}$ . Thus,  $\lambda(\tau(\Omega_S)) \geq ns + \binom{s}{2} + 1$ .

Hence, a  $\mathbb{k}$ -linear combination of  $\tau, [x_i, T_j], \Gamma_{ij}$  can be pulled back to a nonzero torsion element in  $\Omega_R$  by [HMM21, Theorem 4.9].  $\square$

The above theorem recovers [CnGW98, Theorem 2.13].

**Corollary 4.11.** [CnGW98, Theorem 2.13] Under the conditions of Theorem 4.9, if  $\mathfrak{m}^3 \subseteq \mathfrak{C}_R$ , then the torsion  $\tau(\Omega_R)$  is nonzero.

*Proof.* Since  $\mathfrak{m}^3 \subseteq \mathfrak{C}_R$ , we only need to check if  $x_1^2, x_1 x_2 \notin \text{Mono}(I)$  to use the above theorem. This is equivalent to checking if  $X_1^2 - f(\underline{X}), X_1 X_2 - g(\underline{X}) \in I$  or  $x_1^2 = f(\underline{x}), x_1 x_2 = g(\underline{x})$  in  $R$ .

We have  $v(x_1^2) = 2a_1, v(x_1 x_2) = a_1 + a_2$ . Also, note that  $v(f)$  or  $v(g)$  is of the form  $\sum_{i=1}^n j_i a_i, j_i \in \mathbb{N}$ . Since  $I \subseteq \mathfrak{m}^2$ ,  $f(\underline{X})$  and  $g(\underline{X})$  cannot have any linear terms in  $X_i$ . Thus, either  $j_i \geq 2$  for some  $i \geq 2$  or there exist at least two values of  $i$ , say  $\{i_1, i_2\} \neq \{1, 2\}$ , such that  $j_i \geq 1$ . Since we have  $a_1 < a_2 < \dots < a_n$ , it is clear that the valuations  $2a_1, a_1 + a_2$  cannot be attained in either of these situations. Hence,  $x_1^2, x_1 x_2 \notin \text{Mono}(I)$ .  $\square$

**Example 4.12.** Let  $R = \mathbb{k}[[t^{22}, t^{23} + t^{27}, t^{24} + t^{27}, t^{25} + t^{27}, t^{26} + t^{27}]] \cong \mathbb{k}[[x, y, z, w, u]]$ . Here  $\mathfrak{C}_R = (t^{110})\overline{R}$  and hence  $\mathfrak{m}^5 \subseteq \mathfrak{C}_R$ . M2 computations show that  $x^4, x^3 y \notin \text{Mono}(I)$ . Thus the torsion submodule  $\tau(\Omega_R) \neq 0$  by the above theorem.

**Example 4.13.** Let  $R = \mathbb{k}[[t^{30}, t^{31} + t^{36}, t^{32} + t^{36}, t^{33} + t^{36}, t^{34} + t^{36}]] \cong \mathbb{k}[[x, y, z, w, u]]$ . Here  $\mathfrak{C}_R = (t^{180})\overline{R}$  and hence  $\mathfrak{m}^6 \subseteq \mathfrak{C}_R$ . M2 computations show that  $x^5, x^4 y \notin \text{Mono}(I)$ . Thus the torsion submodule  $\tau(\Omega_R) \neq 0$  by the above theorem.

## REFERENCES

- [Ara89] Donu Arapura. Smoothable varieties with torsion free canonical sheaf. *Manuscripta Math.*, 63(1):17–20, 1989.
- [Ber63] Robert Berger. Differentialmoduln eindimensionaler lokaler Ringe. *Math. Z.*, 81:326–354, 1963.
- [Ber88] Robert W. Berger. On the torsion of the differential module of a curve singularity. *Arch. Math. (Basel)*, 50(6):526–533, 1988.
- [Ber94] Robert W Berger. Report on the Torsion of Differential Module of an Algebraic Curve. In *Algebraic Geometry and its Applications*, pages 285–303. Springer, 1994.

- [BG80] Ragnar-Olaf Buchweitz and Gert-Martin Greuel. The Milnor number and deformations of complex curve singularities. *Invent. Math.*, 58(3):241–281, 1980.
- [CK99] Guillermo Cortiñas and Fabiana Krongold. Artinian algebras and differential forms. *Communications in Algebra*, 27(4):1711–1716, 1999.
- [CnGW98] Guillermo Cortiñas, Susan C. Geller, and Charles A. Weibel. The Artinian Berger conjecture. *Math. Z.*, 228(3):569–588, 1998.
- [Eis95] David Eisenbud. *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. With a view toward algebraic geometry.
- [Güt90] Klaus Güttes. Zum Torsionsproblem bei Kurvensingularitäten. *Archiv der Mathematik*, 54(5):499–510, May 1990.
- [H90] Reinhold Hübl. A note on the torsion of differential forms. *Arch. Math. (Basel)*, 54(2):142–145, 1990.
- [Her78] Jürgen Herzog. Ein Cohen-Macaulay-Kriterium mit Anwendungen auf den Konormalenmodul und den Differentialmodul. *Math. Z.*, 163(2):149–162, 1978.
- [HK71] Jürgen Herzog and Ernst Kunz. Die Wertehalbgruppe eines lokalen Rings der Dimension 1. In *Die Wertehalbgruppe eines lokalen Rings der Dimension I*, pages 3–43. Springer, 1971.
- [HMM21] Craig Huneke, Sarasij Maitra, and Vivek Mukundan. Torsion in differentials and berger’s conjecture. *Research in the Mathematical Sciences*, 8(4):1–15, 2021.
- [HW84] Jürgen Herzog and Roland Waldi. Differentials of linked curve singularities. *Archiv der Mathematik*, 42:335–343, 1984.
- [HW86] Jürgen Herzog and Rolf Waldi. Cotangent functors of curve singularities. *Manuscripta Math.*, 55(3-4):307–341, 1986.
- [Iso91] Satoru Isogawa. On Berger’s conjecture about one-dimensional local rings. *Arch. Math. (Basel)*, 57(5):432–437, 1991.
- [Koc83] Josef Koch. *Über die Torsion des Differentialmoduls von Kurvensingularitäten*, volume 5 of *Regensburger Mathematische Schriften [Regensburg Mathematical Publications]*. Universität Regensburg, Fachbereich Mathematik, Regensburg, 1983.
- [KR77] Ernst Kunz and Walter Ruppert. Quasihomogene Singularitäten algebraischer Kurven. *Manuscripta Math.*, 22(1):47–61, 1977.
- [Kun61] Ernst Kunz. Differentialformen inseparabler algebraischer Funktionenkörper. *Math. Z.*, 76:56–74, 1961.
- [Kun86] Ernst Kunz. *Kähler differentials*. Advanced Lectures in Mathematics. Friedr. Vieweg & Sohn, Braunschweig, 1986.
- [Lin17] Haydee Lindo. Trace ideals and centers of endomorphism rings of modules over commutative rings. *J. Algebra*, 482:102–130, 2017.
- [Mai22a] Sarasij Maitra. Partial trace ideals and Berger’s conjecture. *Journal of Algebra*, 598:1–23, 2022.
- [Mai22b] Sarasij Maitra. *Partial Trace Ideals, The Conductor and Berger’s Conjecture*. PhD thesis, University of Virginia, Charlottesville, VA, 2022.
- [Mat67] Tadayuki Matsuoka. On the torsion submodule of a module of type  $(F_1)$ . *J. Sci. Hiroshima Univ. Ser. A-I Math.*, 31:151–160, 1967.
- [Poh89] Thomas Pohl. Torsion des Differentialmoduls von Kurvensingularitäten mit maximaler Hilbertfunktion. *Archiv der Mathematik*, 52(1):53–60, 1989.
- [Poh91] Thomas Pohl. Differential modules with maximal torsion. *Arch. Math. (Basel)*, 57(5):438–445, 1991.
- [PUV07] Claudia Polini, Bernd Ulrich, and Marie A. Vitulli. The core of zero-dimensional monomial ideals. *Adv. Math.*, 211(1):72–93, 2007.
- [Sch70] Günter Scheja. Differentialmoduln lokaler analytischer Algebren, Schriftenreihe Math. Inst. Univ. Fribourg, Univ. Fribourg, Switzerland, 1970.
- [SH06] Irena Swanson and Craig Huneke. *Integral closure of ideals, rings, and modules*, volume 336 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2006.
- [Ulr81] Bernd Ulrich. Torsion des Differentialmoduls und Kotangentenmodul von Kurvensingularitäten. *Arch. Math. (Basel)*, 36(6):510–523, 1981.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY, UT, USA  
*Email address:* maitra@math.utah.edu

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY DELHI, HAUZ KHAS, INDIA.  
*Email address:* vmukunda@iitd.ac.in