

THE HURWITZ MAP AND HARMONIC WAVE FUNCTIONS FOR INTEGER AND HALF-INTEGERS ANGULAR MOMENTUM

SERGIO A. HOJMAN^{1,2,3;(a)}, EDUARDO NAHMAD-ACHAR^{4;(b)},
AND ADOLFO SÁNCHEZ-VALENZUELA^{5;(c)}

November 19, 2022

ABSTRACT. Harmonic wave functions for integer and half-integer angular momentum are given in terms of the Euler angles (ϕ, θ, ψ) that define a rotation in $SO(3)$, and the Euclidean norm r in \mathbb{R}^3 , keeping the usual meaning of the spherical coordinates (r, ϕ, θ) . They form a Hilbert (super)-space decomposed in the form $\mathcal{H} = \mathcal{H}_0 \oplus \mathcal{H}_1$. Following a classical work by Schwinger, 2-dimensional harmonic oscillators are used to produce raising and lowering operators that change the total angular momentum eigenvalue of the wave functions in half units. The nature of the representation space \mathcal{H} is approached from the double covering group homomorphism $SU(2) \rightarrow SO(3)$ and the topology involved is taken care of by using the Hurwitz map $H : \mathbb{R}^4 \rightarrow \mathbb{R}^3$. It is shown how to reconsider H as a 2-to-1 group map, $G_0 = \mathbb{R}^+ \times SU(2) \rightarrow \mathbb{R}^+ \times SO(3)$, translating into an assignment $(z_1, z_2) \mapsto (r, \phi, \theta, \psi)$ in terms of two complex variables (z_1, z_2) , under the appropriate identification of \mathbb{R}^4 with \mathbb{C}^2 . It is shown how the Lie algebra of G_0 is coupled with the two Heisenberg Lie algebras of the 2-dimensional (Schwinger's) harmonic oscillators generated by the operators $\{z_1, z_2, \bar{z}_1, \bar{z}_2\}$ and their adjoints. The whole set of operators close either into a 9-dimensional Lie algebra or into an 8-dimensional Lie superalgebra. The wave functions in \mathcal{H} can also be written in terms polynomials in the complex coordinates (z_1, z_2) and their complex conjugates (\bar{z}_1, \bar{z}_2) and the representations are explicitly constructed via highest weight (or lowest weight) vector representations for G_0 .

INTRODUCTION AND SUMMARY OF RESULTS

In 1965, Julian Schwinger introduced a set of quantum operators based on the Heisenberg commutation relations from which he also produced the Lie algebra generators of the quantum theory of angular momentum (see [6]). Schwinger's operators had the property of transforming states of a given *total angular momentum* j (or *spin value* s), into states having total angular momentum $j \pm 1/2$ (or spin value $s \pm 1/2$). Later on, in 1979, he wrote in [7]:

(a) sergio.hojman@uai.cl; (b) nahmad@nucleares.unam.mx; (c) adolfo@cimat.mx

(1) Departamento de Ciencias, Facultad de Artes Liberales, Universidad Adolfo Ibáñez, Santiago, Chile.

(2) Departamento de Física, Facultad de Ciencias, Universidad de Chile, Santiago, Chile.

(3) Centro de Recursos Educativos Avanzados, CREA, Santiago, Chile.

(4) Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Apdo. Postal 70-543, Mexico City CP 04510, México.

(5) Centro de Investigación en Matemáticas A.C. Unidad Mérida Km 5.5 Carretera Sierra Papacal - Chuburná Puerto. Mérida, Yucatán. CP 97302, México.

“ *The recent recognition of the existence and possible utility of transformations between particles of different statistics* [For a mathematically oriented review see L. Corwin, Y. Ne’eman, and S. Sternberg, *Rev. Modern Phys.* 47 (1975), 573; D.Z. Freedman, P. Van Nieuwenhuizen, and S. Ferrara, *Phys. Rev. D* 13 (1976), 3214; S. Deser and B. Zumino, *Phys. Lett. B* 62 (1976), 335.] *had its origin in the properties of certain two-dimensional dual models, although the concept was clearly prefigured in the simple unification of all spins and statistics by means of multispinor sources and fields* [J. Schwinger, *Phys. Rev.* 152 (1966), 1219]. [*Offstage voice: “All right, wise guy! Then why didn’t you do it first ?”*] *The question is well put and I cannot answer it* [There is a relevant discussion in J. Hadamard, “*The Psychology of Invention in the Mathematical Field*,” Princeton Univ. Press, Princeton, 1945], *but it may yet be appropriate, and useful, to indicate the elementary kinematical basis for such transformations and, hopefully, to assist in the redirection of these efforts toward more modest, physical, goals.*”

The purpose of this work is show how Schwinger’s ideas produce a Hilbert space (superspace, actually) of harmonic wave functions describing both, integer and half-integer angular momentum states from a unified mathematical point of view and having an easy physical interpretation. The produced wave functions depend on the Euler angles (ϕ, θ, ψ) that define a rotation in $SO(3)$, and on the Euclidean norm r in \mathbb{R}^3 , keeping the usual meaning of the spherical coordinates (r, ϕ, θ) . They can also be written in terms of two complex variables (z_1, z_2) and their complex conjugates (\bar{z}_1, \bar{z}_2) .

The nature of the given Hilbert space is approached from the double covering group homomorphism $SU(2) \rightarrow SO(3)$ and the corresponding representation theory of these groups. The topology involved is taken care of by using the Hurwitz map $H : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ as in Hage-Hassan and Kibler [2]. Under appropriate identifications, $\mathbb{R}^4 = \text{Dom } H$ is viewed as Hamilton’s quaternions \mathbb{H} which are in turn identified with the domain \mathbb{C}^2 of the complex variables (z_1, z_2) .

Our approach consists of changing the point of view of H and considering it as a 2-to-1 group map, $G_0 = \mathbb{R}^+ \times SU(2) \rightarrow \mathbb{R}^+ \times SO(3)$, which translates into a correspondence $\mathbb{C}^2 \ni (z_1, z_2) \mapsto (r, \phi, \theta, \psi) \in \mathbb{R}^+ \times SO(3)$, whenever $\begin{pmatrix} \bar{z}_1/\sqrt{r} & -\bar{z}_2/\sqrt{r} \\ z_2/\sqrt{r} & z_1/\sqrt{r} \end{pmatrix} \in SU(2)$, and $|z_1|^2 + |z_2|^2 = r > 0$. In particular, we show how to write z_1 and z_2 in terms of $(\sqrt{r}, \phi/2, \theta/2, \psi/2)$. The Lie algebra generators, $\{\mathbf{L}, \mathbf{L}_x, \mathbf{L}_y, \mathbf{L}_z\}$ of $\mathfrak{g} = \text{Lie}(G_0)$ find simple expressions in terms of the monomials $\{z_1, z_2, \bar{z}_1, \bar{z}_2\}$ and the partial derivatives with respect to these variables. A direct computation shows that,

$$\mathbf{L}_x^2 + \mathbf{L}_y^2 + \mathbf{L}_z^2 = \mathbf{L}^2 + \mathbf{L} + \frac{r}{4} \Delta, \quad \text{where} \quad \Delta = \left(\frac{\partial^2}{\partial z_1 \partial \bar{z}_1} + \frac{\partial^2}{\partial z_2 \partial \bar{z}_2} \right),$$

is the Laplace operator in the domain $\mathbb{R}^4 \simeq \mathbb{C}^2$. It turns out that the wave functions f produced through our approach satisfy $\Delta f = 0$, and they can be written as, $f(r, \phi, \theta, \psi) = r^{n+1/2} e^{i\frac{\psi}{2}} \Psi(\theta, \phi)$ if the eigenvalue j of the total angular momentum operator \mathbf{L} is $n + 1/2$, or $f(r, \phi, \theta, \psi) = r^n \Psi(\theta, \phi)$ if $j = n$ ($n \in \mathbb{N}$). Furthermore, $\Psi(\theta, \phi) \in \oplus_{m=-j}^{m=j} \langle j, m \rangle$, where $\langle j, m \rangle$ stands for the space of harmonic functions f on the 2-sphere satisfying $\mathbf{L}f = jf$ and $\mathbf{L}_z = mf$; *ie*, the usual spherical harmonics.

In terms of the complex variables $(z_1, z_2, \bar{z}_1, \bar{z}_2)$, the wave functions hereby produced are obtained in one of two ways as follows: either, get first the highest weight vector $|j, +j\rangle = (z_1 \bar{z}_2)^j$ (if $j = n \in \mathbb{N}$) or $|j, +j\rangle = z_1 (z_1 \bar{z}_2)^{\frac{2j-1}{2}}$ (if $j = n + 1/2$, with $n \in \mathbb{N} \cup \{0\}$), and then move from it by successive application of the ladder operator $\mathbf{L}_- = \mathbf{L}_x - i\mathbf{L}_y$ which preserves the eigenvalue j but lowers the eigenvalue m in one unit at each step, until reaching a non-zero scalar multiple of the lowest weight vector $|j, -j\rangle$, characterized by $\mathbf{L}_- |j, -j\rangle = 0$; or else, produce first the lowest weight vector $|j, -j\rangle = (\bar{z}_1 z_2)^j$ (if $j = n \in \mathbb{N}$) or $|j, -j\rangle = z_2 (\bar{z}_1 z_2)^{\frac{2j-1}{2}}$ (if $j = n + 1/2$, with $n \in \mathbb{N} \cup \{0\}$), and then move from it by successive application of the ladder operator $\mathbf{L}_+ = \mathbf{L}_x + i\mathbf{L}_y$ which preserves the eigenvalue j but raises the eigenvalue m in one unit at each step, until reaching a non-zero scalar multiple of the highest weight vector $|j, +j\rangle$, characterized by $\mathbf{L}_+ |j, +j\rangle = 0$.

The whole picture is completed by showing how the Lie algebra of G_0 is coupled with two Heisenberg Lie algebras of 2-dimensional harmonic oscillators. Following Schwinger, the 2-dimensional harmonic oscillators are used to produce raising and lowering operators that change the total angular momentum eigenvalue of the wave functions in half units. It is explained how the complete set of Lie algebra operators of \mathfrak{g} together with the generators of the two Heisenberg Lie algebras (raising and lowering operators) can close either into a 9-dimensional Lie algebra or into an 8-dimensional Lie superalgebra.

The Hilbert space where the produced wave functions *live* can be naturally decomposed into a direct sum $\mathcal{H} = \mathcal{H}_0 \oplus \mathcal{H}_1$, where \mathcal{H}_0 contains all the states having integer angular momentum (or *space of bosons*) and \mathcal{H}_1 contains all the states having half-integer angular momentum (or *space of fermions*), thus providing an explicit realization of a *superspace* in which both, the Lie algebra and the Lie superalgebra are faithfully represented.

We mention in passing that mathematical related work in a similar direction was done in the mid 70's of the last century by M. Kashiwara and M. Vergne following ideas introduced by R. Howe (see [3] and [4]). It turns out that their approach is adapted to a joint action of a given symplectic and a given orthogonal group, which is a clear indicative of supersymmetry (see [1] and [3]). In Schwinger's approach —on which we focus this work— the coexistence between a symplectic group and an orthogonal group comes from the coupling of two harmonic oscillators with Lie algebra \mathfrak{g}_1 on which the usual angular momentum operators with Lie algebra \mathfrak{g}_0 act, giving rise either to a 9-dimensional Lie algebra or to an 8-dimensional Lie superalgebra, as explained in §9 below.

1. SCHWINGER'S HEISENBERG-ANGULAR-MOMENTUM LIE ALGEBRA

Schwinger's Lie algebra is defined in terms of the 2-dimensional quantum harmonic oscillator operators, a_x, a_y , and their adjoints, a_x^\dagger, a_y^\dagger , satisfying the Heisenberg commutation relations,

$$[a_x, a_x^\dagger] = 1, \quad [a_y, a_y^\dagger] = 1.$$

Then define,

$$J_+ = a_x^\dagger a_y, \quad J_- = a_y^\dagger a_x,$$

$$J_z = \frac{1}{2}(a_x^\dagger a_x - a_y^\dagger a_y), \quad J = \frac{1}{2}(a_x^\dagger a_x + a_y^\dagger a_y).$$

It follows that,

$$[J_z, J_\pm] = \pm J_\pm, \quad [J_+, J_-] = 2J_z,$$

and

$$[J, J_\pm] = 0, \quad [J, J_z] = 0.$$

It is not difficult to see that,

$$\begin{aligned} [J, a_x] &= -\frac{1}{2} a_x, & [J, a_x^\dagger] &= \frac{1}{2} a_x^\dagger, & [J, a_y] &= -\frac{1}{2} a_y, & [J, a_y^\dagger] &= \frac{1}{2} a_y^\dagger, \\ [J_z, a_x] &= -\frac{1}{2} a_x, & [J_z, a_x^\dagger] &= \frac{1}{2} a_x^\dagger, & [J_z, a_y] &= \frac{1}{2} a_y, & [J_z, a_y^\dagger] &= -\frac{1}{2} a_y^\dagger. \end{aligned}$$

And also that,

$$\begin{aligned} [J_+, a_x] &= -a_y, & [J_+, a_x^\dagger] &= 0, & [J_+, a_y] &= 0, & [J_+, a_y^\dagger] &= a_x^\dagger, \\ [J_-, a_x] &= 0, & [J_-, a_x^\dagger] &= a_y^\dagger, & [J_-, a_y] &= -a_x, & [J_-, a_y^\dagger] &= 0. \end{aligned}$$

We shall give a Lie algebra, realized in terms of operators adapted to the geometry and the symmetry involved, which fully contains the subalgebra of the J 's and the a 's, but includes an additional Heisenberg algebra.

2. THE HURWITZ MAP

From the point of view of Lie's theory, Schwinger's operators may transform representation spaces of the 3-dimensional rotation group $SO(3)$ (*ie*, states whose *total angular momentum* j (or *spin value* s) lies in \mathbb{Z}), into representation spaces of its *spin group* or *double covering group*, $SU(2)$, (*ie*, into states whose *total angular momentum* is $j \pm 1/2$ (or whose *spin value* is $s \pm 1/2$), and viceversa.

Even though the Lie groups $SU(2)$ and $SO(3)$ are locally isomorphic, they are topologically different. To unveil the spin representations of $SO(3)$ one needs to consider the double covering map $SU(2) \rightarrow SO(3)$. One way to take the topology involved into account is as in Hage-Hassan and Kibler (see [2]) by considering the Hurwitz map, $H : \mathbb{R}^4 \ni u \mapsto H(u) = x \in \mathbb{R}^3$, given by,

$$\begin{aligned} x_1 &= H_1(u_1, u_2, u_3, u_4) = 2(u_1 u_3 + u_2 u_4), \\ x_2 &= H_2(u_1, u_2, u_3, u_4) = 2(u_2 u_3 - u_1 u_4), \\ x_3 &= H_3(u_1, u_2, u_3, u_4) = u_1^2 + u_2^2 - u_3^2 - u_4^2, \end{aligned}$$

which has the property that,

$$r = \sqrt{x_1^2 + x_2^2 + x_3^2} = u_1^2 + u_2^2 + u_3^2 + u_4^2 = R^2.$$

Thus, H maps 3-spheres centered at the origin in \mathbb{R}^4 , onto 2-spheres centered at the origin in \mathbb{R}^3 . The Hurwitz map can also be described as a map $H : \mathbb{C}^2 \rightarrow \mathbb{R}^3$, using the complex variables,

$$z_1 = u_1 + i u_2 \quad \text{and} \quad z_2 = u_3 + i u_4,$$

so that,

$$x_1 = z_1\bar{z}_2 + \bar{z}_1z_2, \quad x_2 = -i(z_1\bar{z}_2 - \bar{z}_1z_2), \quad x_3 = z_1\bar{z}_1 - z_2\bar{z}_2.$$

That is,

$$\begin{aligned} x_1 &= H_1(z_1, z_2) = 2 \operatorname{Re}(z_1\bar{z}_2), \\ x_2 &= H_2(z_1, z_2) = 2 \operatorname{Im}(z_1\bar{z}_2), \\ x_3 &= H_3(z_1, z_2) = |z_1|^2 - |z_2|^2 \end{aligned}$$

One possible symmetry group associated to the Hurwitz map is the *real* Lie group,

$$G = \left\{ \begin{pmatrix} \bar{z}_1 & -\bar{z}_2 \\ z_2 & z_1 \end{pmatrix} \mid \begin{pmatrix} z_1 \\ z_2 \end{pmatrix} \in \mathbb{C}^2, \text{ and } R^2 = |z_1|^2 + |z_2|^2 > 0 \right\},$$

whose connected component to the identity G_0 is isomorphic to $\mathbb{R}^+ \times SU(2)$, where \mathbb{R}^+ stands for the multiplicative group of positive real numbers, obtained when $R = \sqrt{r} > 0$. The Lie group $SU(2)$ can be identified with the unit quaternions. It acts on 3-spheres in $\mathbb{R}^4 \simeq \mathbb{H}$ via left (or right) multiplication in \mathbb{H} . Then the Hurwitz map transforms 3-spheres in \mathbb{R}^4 into 2-spheres in \mathbb{R}^3 , and \mathbb{R}^+ acts either in \mathbb{R}^4 or in \mathbb{R}^3 by changing the spheres radii.

3. THE LIE GROUP $SU(2)$ AND ITS LIE ALGEBRA $\mathfrak{su}(2)$

To fix the notation we recall that the special unitary group $SU(2)$ is defined as,

$$SU(2) = \{A \in \operatorname{Mat}_{2 \times 2}(\mathbb{C}) \mid A^*A = \mathbb{1}_2, \text{ and } \det A = 1\},$$

where $A^* = (\bar{A})^t$ stands for the conjugate transpose matrix of A and $\mathbb{1}_2$ is the 2×2 unit matrix. Thus, $A \in \operatorname{Mat}_{2 \times 2}(\mathbb{C})$ belongs to $SU(2)$ if and only if it is invertible and its inverse A^{-1} is equal to A^* . That is,

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} = \frac{1}{\det A} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} = \begin{pmatrix} \bar{a} & \bar{c} \\ \bar{b} & \bar{d} \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}^*.$$

Since $\det A = 1$ for any $A \in SU(2)$, it follows that,

$$d = \bar{a}, \quad b = -\bar{c}, \quad \text{and} \quad \det A = |a|^2 + |c|^2 = 1.$$

Therefore,

$$SU(2) \simeq \left\{ \begin{pmatrix} a & -\bar{c} \\ c & \bar{a} \end{pmatrix} \in \operatorname{Mat}_{2 \times 2}(\mathbb{C}) \mid a, c \in \mathbb{C} \text{ and } |a|^2 + |c|^2 = 1 \right\},$$

making clear the fact that $SU(2)$ is actually diffeomorphic to the unit 3-sphere S^3 in $\mathbb{R}^4 \simeq \mathbb{C}^2$.

The Lie algebra $\mathfrak{su}(2)$ of $SU(2)$ is the *real* vector space,

$$\mathfrak{su}(2) \simeq \{X \in \operatorname{Mat}_{2 \times 2}(\mathbb{C}) \mid X + X^* = 0, \text{ and } \operatorname{Tr} X = 0\}.$$

It is well known that $\mathfrak{su}(2)$ is isomorphic to \mathbb{R}^3 via,

$$\mathfrak{su}(2) \ni X \iff X = i \begin{pmatrix} z & x - iy \\ x + iy & -z \end{pmatrix}, \quad \text{with } \begin{pmatrix} x \\ y \\ z \end{pmatrix} \in \mathbb{R}.$$

A convenient set of linearly independent generators for $\mathfrak{su}(2)$ over the real field \mathbb{R} is, $\{i\sigma_1, i\sigma_2, i\sigma_3\}$, where, the σ_i 's are the Pauli matrices, so that,

$$i\sigma_1 = i \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad i\sigma_2 = i \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad i\sigma_3 = i \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

and,

$$[i\sigma_1, i\sigma_2] = -2i\sigma_3, \quad [i\sigma_2, i\sigma_3] = -2i\sigma_1, \quad [i\sigma_3, i\sigma_1] = -2i\sigma_2.$$

Clearly,

$$\mathfrak{su}(2) \ni X \Rightarrow \det X = x^2 + y^2 + z^2.$$

4. THE DOUBLE COVERING MAP $SU(2) \rightarrow SO(3)$

For each $A \in SU(2)$, the assignment $X \mapsto AXA^*$ defines a transformation $\rho(A) : \mathfrak{su}(2) \rightarrow \mathfrak{su}(2)$ which preserves the quadratic form, $\det X = x^2 + y^2 + z^2$, since $A^*A = \mathbb{1}_2$ and $\det(AXA^*) = \det(XA^*A) = \det X$. Therefore, $\rho(A) \in SO(3)$.

It can be proved (and it is well known) that ρ is a 2 to 1 map, and in fact, a local diffeomorphism. This is usually expressed by saying that ρ induces the exact sequence of groups,

$$\{\mathbb{1}_2\} \hookrightarrow \mathbb{Z}_2 \hookrightarrow SU(2) \twoheadrightarrow SO(3) \twoheadrightarrow \{\mathbb{1}_3\},$$

where, $\mathbb{Z}_2 = \{\pm\mathbb{1}_2\}$ as a subgroup of $SU(2)$, and $\rho : SU(2) \twoheadrightarrow SO(3)$ is actually a *group homomorphism*, as it satisfies,

$$\rho(A_1 A_2) = \rho(A_1) \rho(A_2), \quad \text{for any } A_1, A_2 \in SU(2),$$

and it is straightforward to rewrite $\rho(A_1)$ and $\rho(A_2)$ as 3×3 matrices with real entries acting (in that order) on column vectors $\begin{pmatrix} x \\ y \\ z \end{pmatrix} \in \mathbb{R}^3$, whenever, $X = i \begin{pmatrix} z & x - iy \\ x + iy & -z \end{pmatrix} \in \mathfrak{su}(2)$.

The nature of the double covering map $SU(2) \ni A \mapsto \rho(A) \in SO(3)$ can be understood through the following observations:

1. Take $A = \begin{pmatrix} e^{-i\phi} & 0 \\ 0 & e^{i\phi} \end{pmatrix} \in SU(2)$. Then,

$$\rho(A)X = AXA^* = i \begin{pmatrix} z & e^{-2i\phi}(x - iy) \\ e^{2i\phi}(x + iy) & -z \end{pmatrix},$$

corresponds to a counterclockwise rotation, $R_z(2\phi)$, of the x - y plane (performed around the z -axis in \mathbb{R}^3) by an angle of 2ϕ .

2. Take $A = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \in SU(2)$. Then,

$$\rho(A)X = A X A^* = i \begin{pmatrix} z \cos 2\theta - x \sin 2\theta & z \sin 2\theta + x \cos 2\theta - iy \\ z \sin 2\theta + x \cos 2\theta + iy & -(z \cos 2\theta - x \sin 2\theta) \end{pmatrix},$$

corresponds to a counterclockwise rotation, $R_y(2\theta)$, of the z - x plane (performed around the y -axis in \mathbb{R}^3) by an angle of 2θ .

3. In order to parametrize a given rotation in $SO(3)$ through the Euler angles (ϕ, θ, ψ) consistently with the given parametrization of $SU(2)$ and the given identification of \mathbb{R}^3 with $\mathfrak{su}(2) = \text{Lie } SU(2)$, we shall adhere ourselves to the following convention: In passing from a given x - y - z (fixed) frame in \mathbb{R}^3 to a (moving) X - Y - Z frame (assuming the most general situation in which the directions defined by z and Z are linearly independent), we fix first the orientation of the *line of nodes* ℓ defined by the intersection of the x - y plane with the X - Y plane, in such a way that that the positive direction of ℓ coincides with the direction of $\hat{z} \times \hat{Z}$, where \hat{z} and \hat{Z} are unit vectors along the corresponding z and Z axes. Then, the three rotations defined by the Euler angles (ϕ, θ, ψ) to go from the x - y - z frame to the X - Y - Z frame, must be performed as follows:

- (1) First make a counterclockwise rotation by an angle ϕ around the z -axis, to take the x -axis into the positive direction of the line of nodes.
- (2) Then make a counterclockwise rotation by an angle θ around the positive direction of the line of nodes, to make the z -axis coincide with the Z -axis.
- (3) Finally, make a counterclockwise rotation by an angle ψ around the Z -axis to make the positive direction of the line of nodes, coincide with the positive direction defined by the X -axis.
- (4) Due to the symmetry under rotations in the x - y plane, the positive direction of the line of nodes can be labeled either by x' or by y' . All that matters is that the *auxiliary x' - y' - z frame must have* the same orientation as the x - y - z frame does. We shall take the positive direction of the line of nodes to be the auxiliary y' -axis, so that the x' -axis will have the direction of $\hat{y}' \times \hat{z}$, where \hat{y}' and \hat{z} are unit vectors along the positive direction of the line of nodes and the z -axis, respectively.

4. Using the double covering group homomorphism $\rho : SU(2) \rightarrow SO(3)$, we may write $A \in SU(2)$ in the form,

$$\begin{aligned} A &= \begin{pmatrix} e^{-i\frac{\phi}{2}} & 0 \\ 0 & e^{i\frac{\phi}{2}} \end{pmatrix} \begin{pmatrix} \cos \frac{\theta}{2} & -\sin \frac{\theta}{2} \\ \sin \frac{\theta}{2} & \cos \frac{\theta}{2} \end{pmatrix} \begin{pmatrix} e^{-i\frac{\psi}{2}} & 0 \\ 0 & e^{i\frac{\psi}{2}} \end{pmatrix} \\ &= \begin{pmatrix} e^{-i\frac{(\psi+\phi)}{2}} \cos \frac{\theta}{2} & -e^{-i\frac{(\psi-\phi)}{2}} \sin \frac{\theta}{2} \\ e^{i\frac{(\psi-\phi)}{2}} \sin \frac{\theta}{2} & e^{i\frac{(\psi+\phi)}{2}} \cos \frac{\theta}{2} \end{pmatrix}, \end{aligned}$$

and obtain, under $\rho(A)$, the desired rotation $R_z(\psi) \circ R_y(\theta) \circ R_z(\phi) \in SO(3)$ defined by the Euler angles (ϕ, θ, ψ) .

5. THE HURWITZ MAP IN TERMS OF THE FOUR VARIABLES (r, θ, ϕ, ψ)

Consider the 4-dimensional *real* Lie group,

$$G = \left\{ \begin{pmatrix} \bar{z}_1 & -\bar{z}_2 \\ z_2 & z_1 \end{pmatrix} \in \text{Mat}_{2 \times 2}(\mathbb{C}) \mid \begin{pmatrix} z_1 \\ z_2 \end{pmatrix} \in \mathbb{C}^2, \text{ and } R^2 = |z_1|^2 + |z_2|^2 > 0 \right\}.$$

For any $g = \begin{pmatrix} \bar{z}_1 & -\bar{z}_2 \\ z_2 & z_1 \end{pmatrix} \in G$, choose a positive real number $R = \sqrt{r} \in \mathbb{R}^+$, so as to have g in the *identity component* G_0 of G , and write,

$$g = \sqrt{r} \begin{pmatrix} \bar{z}_1/\sqrt{r} & -\bar{z}_2/\sqrt{r} \\ z_2/\sqrt{r} & z_1/\sqrt{r} \end{pmatrix}, \quad \text{where,} \quad \frac{1}{\sqrt{r}} \begin{pmatrix} \bar{z}_1 & -\bar{z}_2 \\ z_2 & z_1 \end{pmatrix} \in SU(2).$$

Thus,

$$\frac{1}{\sqrt{r}} \begin{pmatrix} \bar{z}_1 & -\bar{z}_2 \\ z_2 & z_1 \end{pmatrix} = \begin{pmatrix} e^{-i\frac{(\psi+\phi)}{2}} \cos \frac{\theta}{2} & -e^{-i\frac{(\psi-\phi)}{2}} \sin \frac{\theta}{2} \\ e^{i\frac{(\psi-\phi)}{2}} \sin \frac{\theta}{2} & e^{i\frac{(\psi+\phi)}{2}} \cos \frac{\theta}{2} \end{pmatrix} \in SU(2),$$

and,

$$\begin{aligned} z_1 &= \sqrt{r} e^{i\frac{\psi}{2}} e^{i\frac{\phi}{2}} \cos \frac{\theta}{2} \\ z_2 &= \sqrt{r} e^{i\frac{\psi}{2}} e^{-i\frac{\phi}{2}} \sin \frac{\theta}{2}. \end{aligned}$$

In particular, using these expressions for z_1 and z_2 , the cartesian coordinates x_1, x_2 and x_3 —given by the Hurwitz map, $x_i = H_i(z_1, z_2)$ ($1 \leq i \leq 3$)— can be written in terms of the spherical coordinates r, θ and ϕ , in the usual way,

$$x_1 = r \sin \theta \cos \phi, \quad x_2 = r \sin \theta \sin \phi, \quad x_3 = r \cos \theta.$$

Also, H yields the following expressions for r and the Euler angles (θ, ϕ, ψ) in terms of the complex variables z_1 and z_2 :

$$\begin{aligned} H^*r &= |z_1|^2 + |z_2|^2, & H^*\theta &= \cos^{-1} \left(\frac{|z_1|^2 - |z_2|^2}{|z_1|^2 + |z_2|^2} \right), \\ H^*\phi &= \tan^{-1} \left(-i \frac{z_1 \bar{z}_2 - \bar{z}_1 z_2}{z_1 \bar{z}_2 + \bar{z}_1 z_2} \right), & H^*\psi &= \tan^{-1} \left(-i \frac{z_1 z_2 - \bar{z}_1 \bar{z}_2}{z_1 z_2 + \bar{z}_1 \bar{z}_2} \right), \end{aligned}$$

where, H^* stands for the *pullback* of H , so that H^*f means $f \circ H$ for any smooth function f . Thus, for example, $H^*r = r(H(z_1, z_2)) = |z_1|^2 + |z_2|^2$, etc. In particular, H itself gets described as a map changing two specific local coordinate charts; say, $(z_1, z_2, \bar{z}_1, \bar{z}_2) \mapsto (r, \theta, \phi, \psi)$.

6. THE LIE ALGEBRA \mathfrak{g} OF THE SYMMETRY GROUP G ASSOCIATED TO THE HURWITZ MAP

Since $G_0 \simeq \mathbb{R}^+ \times SU(2)$, it is immediate to see that the Lie algebra \mathfrak{g} of G gets isomorphic to the *real* 4-dimensional space,

$$\mathfrak{g} = \text{Span}_{\mathbb{R}}\{ \mathbb{1}_2 \} \oplus \mathfrak{su}(2).$$

Its complexification $\mathfrak{g}_{\mathbb{C}}$ is isomorphic to the Lie algebra $\mathfrak{gl}_2(\mathbb{C})$ of complex 2×2 matrices. Geometrically, we may realize the Lie algebras \mathfrak{g} and $\mathfrak{g}_{\mathbb{C}}$ in terms of real and complex 4-dimensional vector spaces of left G -invariant vector fields, respectively. A convenient set of linearly independent generators for either \mathfrak{g} , or for $\mathfrak{g}_{\mathbb{C}}$, is $\{\mathbf{L}, \mathbf{L}_z, \mathbf{L}_+, \mathbf{L}_-\}$, where,

$$\begin{aligned}\mathbf{L} &= \frac{1}{2} \left(z_1 \frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_2} + \bar{z}_1 \frac{\partial}{\partial \bar{z}_1} + \bar{z}_2 \frac{\partial}{\partial \bar{z}_2} \right), \\ \mathbf{L}_z &= \frac{1}{2} \left(z_1 \frac{\partial}{\partial z_1} - z_2 \frac{\partial}{\partial z_2} - \bar{z}_1 \frac{\partial}{\partial \bar{z}_1} + \bar{z}_2 \frac{\partial}{\partial \bar{z}_2} \right), \\ \mathbf{L}_+ &= \left(z_1 \frac{\partial}{\partial z_2} - \bar{z}_2 \frac{\partial}{\partial \bar{z}_1} \right), \quad \text{and} \quad \mathbf{L}_- = - \left(\bar{z}_1 \frac{\partial}{\partial \bar{z}_2} - z_2 \frac{\partial}{\partial z_1} \right).\end{aligned}$$

It is a straightforward matter to verify that \mathbf{L} commutes with all the operators and that,

$$[\mathbf{L}_z, \mathbf{L}_{\pm}] = \pm \mathbf{L}_{\pm}, \quad \text{and} \quad [\mathbf{L}_+, \mathbf{L}_-] = 2\mathbf{L}_z.$$

These operators may act, *for example*, on the space of complex polynomials in the variables $\{z_1, z_2, \bar{z}_1, \bar{z}_2\}$.

7. COMPUTATIONS WITH THE ANGULAR MOMENTUM COMPLEX OPERATORS

Take the Lie algebra generators $\{\mathbf{L}, \mathbf{L}_z, \mathbf{L}_+, \mathbf{L}_-\}$ as given before, and define the operators \mathbf{L}_x and \mathbf{L}_y through,

$$\mathbf{L}_+ = \mathbf{L}_x + i \mathbf{L}_y, \quad \text{and} \quad \mathbf{L}_- = \mathbf{L}_x - i \mathbf{L}_y.$$

Therefore,

$$\mathbf{L}_x = \frac{1}{2} (\mathbf{L}_+ + \mathbf{L}_-), \quad \text{and} \quad \mathbf{L}_y = \frac{1}{2i} (\mathbf{L}_+ - \mathbf{L}_-).$$

It turns out that,

$$[\mathbf{L}_x, \mathbf{L}_y] = i \mathbf{L}_z, \quad [\mathbf{L}_z, \mathbf{L}_x] = i \mathbf{L}_y, \quad [\mathbf{L}_y, \mathbf{L}_z] = i \mathbf{L}_x.$$

Clearly,

$$\mathbf{L}_+ \mathbf{L}_- = \mathbf{L}_x^2 + \mathbf{L}_y^2 - i [\mathbf{L}_x, \mathbf{L}_y] = \mathbf{L}_x^2 + \mathbf{L}_y^2 + \mathbf{L}_z.$$

Whence,

$$\mathbf{L}_x^2 + \mathbf{L}_y^2 + \mathbf{L}_z^2 = \mathbf{L}_+ \mathbf{L}_- + \mathbf{L}_z^2 - \mathbf{L}_z.$$

Proposition. A direct computation shows that,

$$\mathbf{L}_x^2 + \mathbf{L}_y^2 + \mathbf{L}_z^2 = \mathbf{L}^2 + \mathbf{L} + (|z_1|^2 + |z_2|^2) \left(\frac{\partial^2}{\partial z_1 \partial \bar{z}_1} + \frac{\partial^2}{\partial z_2 \partial \bar{z}_2} \right),$$

where,

$$\left(\frac{\partial^2}{\partial z_1 \partial \bar{z}_1} + \frac{\partial^2}{\partial z_2 \partial \bar{z}_2} \right) = \frac{1}{4} \left(\frac{\partial^2}{\partial u_1^2} + \frac{\partial^2}{\partial u_2^2} + \frac{\partial^2}{\partial u_3^2} + \frac{\partial^2}{\partial u_4^2} \right),$$

is the Laplace operator in \mathbb{R}^4 .

Proof: This is a straightforward computation. \square

Remark. What turns out to be useful is **not to project the operators** $\{\mathbf{L}, \mathbf{L}_z, \mathbf{L}_\pm\}$ onto \mathbb{R}^3 , **but to write them in terms of** the four variables (r, θ, ϕ, ψ) **using the Hurwitz map**. Thus the significant result is the following:

Proposition. Using,

$$\begin{aligned} H^*r &= |z_1|^2 + |z_2|^2, & H^*\theta &= \cos^{-1} \left(\frac{|z_1|^2 - |z_2|^2}{|z_1|^2 + |z_2|^2} \right), \\ H^*\phi &= \tan^{-1} \left(-i \frac{z_1 \bar{z}_2 - \bar{z}_1 z_2}{z_1 \bar{z}_2 + \bar{z}_1 z_2} \right), & H^*\psi &= \tan^{-1} \left(-i \frac{z_1 z_2 - \bar{z}_1 \bar{z}_2}{z_1 z_2 + \bar{z}_1 \bar{z}_2} \right), \end{aligned}$$

together with the covering map $\mathbb{R}^+ \times SU(2) \rightarrow \mathbb{R}^+ \times SO(3)$, to write,

$$\begin{aligned} z_1 &= \sqrt{r} e^{i\frac{\psi}{2}} e^{i\frac{\phi}{2}} \cos \frac{\theta}{2}, \\ z_2 &= \sqrt{r} e^{i\frac{\psi}{2}} e^{-i\frac{\phi}{2}} \sin \frac{\theta}{2}, \end{aligned} \quad \text{when,} \quad \begin{pmatrix} \bar{z}_1 & -\bar{z}_2 \\ z_2 & z_1 \end{pmatrix} \in G_0 = \mathbb{R}^+ \times SU(2),$$

the chain rule,

$$\begin{aligned} \frac{\partial}{\partial z_1} &= \frac{\partial H^*r}{\partial z_1} \frac{\partial}{\partial r} + \frac{\partial H^*\theta}{\partial z_1} \frac{\partial}{\partial \theta} + \frac{\partial H^*\phi}{\partial z_1} \frac{\partial}{\partial \phi} + \frac{\partial H^*\psi}{\partial z_1} \frac{\partial}{\partial \psi}, \\ \frac{\partial}{\partial z_2} &= \frac{\partial H^*r}{\partial z_2} \frac{\partial}{\partial r} + \frac{\partial H^*\theta}{\partial z_2} \frac{\partial}{\partial \theta} + \frac{\partial H^*\phi}{\partial z_2} \frac{\partial}{\partial \phi} + \frac{\partial H^*\psi}{\partial z_2} \frac{\partial}{\partial \psi}, \\ \frac{\partial}{\partial \bar{z}_1} &= \frac{\partial H^*r}{\partial \bar{z}_1} \frac{\partial}{\partial r} + \frac{\partial H^*\theta}{\partial \bar{z}_1} \frac{\partial}{\partial \theta} + \frac{\partial H^*\phi}{\partial \bar{z}_1} \frac{\partial}{\partial \phi} + \frac{\partial H^*\psi}{\partial \bar{z}_1} \frac{\partial}{\partial \psi}, \\ \frac{\partial}{\partial \bar{z}_2} &= \frac{\partial H^*r}{\partial \bar{z}_2} \frac{\partial}{\partial r} + \frac{\partial H^*\theta}{\partial \bar{z}_2} \frac{\partial}{\partial \theta} + \frac{\partial H^*\phi}{\partial \bar{z}_2} \frac{\partial}{\partial \phi} + \frac{\partial H^*\psi}{\partial \bar{z}_2} \frac{\partial}{\partial \psi}, \end{aligned}$$

yields,

$$\frac{\partial}{\partial z_1} = \sqrt{r} e^{-i\frac{\psi}{2}} e^{-i\frac{\phi}{2}} \left(\cos \frac{\theta}{2} \frac{\partial}{\partial r} - \frac{1}{r} \sin \frac{\theta}{2} \frac{\partial}{\partial \theta} - \frac{i}{2r \cos \frac{\theta}{2}} \left(\frac{\partial}{\partial \phi} + \frac{\partial}{\partial \psi} \right) \right)$$

$$\frac{\partial}{\partial z_2} = \sqrt{r} e^{-i\frac{\psi}{2}} e^{i\frac{\phi}{2}} \left(\sin \frac{\theta}{2} \frac{\partial}{\partial r} + \frac{1}{r} \cos \frac{\theta}{2} \frac{\partial}{\partial \theta} + \frac{i}{2r \sin \frac{\theta}{2}} \left(\frac{\partial}{\partial \phi} - \frac{\partial}{\partial \psi} \right) \right).$$

$$\frac{\partial}{\partial \bar{z}_1} = \sqrt{r} e^{i\frac{\psi}{2}} e^{i\frac{\phi}{2}} \left(\cos \frac{\theta}{2} \frac{\partial}{\partial r} - \frac{1}{r} \sin \frac{\theta}{2} \frac{\partial}{\partial \theta} + \frac{i}{2r \cos \frac{\theta}{2}} \left(\frac{\partial}{\partial \phi} + \frac{\partial}{\partial \psi} \right) \right).$$

$$\frac{\partial}{\partial \bar{z}_2} = \sqrt{r} e^{i\frac{\psi}{2}} e^{-i\frac{\phi}{2}} \left(\sin \frac{\theta}{2} \frac{\partial}{\partial r} + \frac{1}{r} \cos \frac{\theta}{2} \frac{\partial}{\partial \theta} - \frac{i}{2r \sin \frac{\theta}{2}} \left(\frac{\partial}{\partial \phi} - \frac{\partial}{\partial \psi} \right) \right).$$

Moreover,

$$\begin{aligned} \mathbf{L} &= \frac{1}{2} \left(z_1 \frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_2} + \bar{z}_1 \frac{\partial}{\partial \bar{z}_1} + \bar{z}_2 \frac{\partial}{\partial \bar{z}_2} \right) = r \frac{\partial}{\partial r}, \\ \mathbf{L}_z &= \frac{1}{2} \left(z_1 \frac{\partial}{\partial z_1} - z_2 \frac{\partial}{\partial z_2} - \bar{z}_1 \frac{\partial}{\partial \bar{z}_1} + \bar{z}_2 \frac{\partial}{\partial \bar{z}_2} \right) = -i \frac{\partial}{\partial \phi}, \\ \mathbf{L}_+ &= \left(z_1 \frac{\partial}{\partial z_2} - \bar{z}_2 \frac{\partial}{\partial \bar{z}_1} \right) = e^{i\phi} \left(\frac{\partial}{\partial \theta} + i \frac{\cos \theta}{\sin \theta} \left(\frac{\partial}{\partial \phi} - \frac{1}{\cos \theta} \frac{\partial}{\partial \psi} \right) \right), \\ \mathbf{L}_- &= - \left(\bar{z}_1 \frac{\partial}{\partial \bar{z}_2} - z_2 \frac{\partial}{\partial z_1} \right) = -e^{-i\phi} \left(\frac{\partial}{\partial \theta} - i \frac{\cos \theta}{\sin \theta} \left(\frac{\partial}{\partial \phi} - \frac{1}{\cos \theta} \frac{\partial}{\partial \psi} \right) \right). \end{aligned}$$

Finally, since,

$$\mathbf{L}_x = \frac{1}{2} (\mathbf{L}_+ + \mathbf{L}_-), \quad \text{and} \quad \mathbf{L}_y = \frac{1}{2i} (\mathbf{L}_+ - \mathbf{L}_-),$$

it follows that,

$$\mathbf{L}_x = i \left(\sin \phi \frac{\partial}{\partial \theta} + \cos \phi \frac{\cos \theta}{\sin \theta} \left(\frac{\partial}{\partial \phi} - \frac{1}{\cos \theta} \frac{\partial}{\partial \psi} \right) \right),$$

and,

$$\mathbf{L}_y = -i \left(\cos \phi \frac{\partial}{\partial \theta} - \sin \phi \frac{\cos \theta}{\sin \theta} \left(\frac{\partial}{\partial \phi} - \frac{1}{\cos \theta} \frac{\partial}{\partial \psi} \right) \right).$$

Proof: This is again a straightforward computation \square

8. THE COMPLETE SET OF SCHWINGER'S OPERATORS

Following Schwinger [6] and [7], introduce the following set of *eight operators*:

$$\begin{aligned} \mathbf{j}_1^{++} &= z_1 & \mathbf{j}_1^{+-} &= \bar{z}_1 & \mathbf{j}_1^{-+} &= \frac{\partial}{\partial z_2} & \mathbf{j}_1^{--} &= \frac{\partial}{\partial z_1} \\ \mathbf{j}_2^{++} &= \bar{z}_2 & \mathbf{j}_2^{+-} &= z_2 & \mathbf{j}_2^{-+} &= \frac{\partial}{\partial \bar{z}_1} & \mathbf{j}_2^{--} &= \frac{\partial}{\partial \bar{z}_2} \end{aligned}$$

We shall refer to them collectively as “*the \mathbf{j}_*^{**} -operators*”, or simply as “*the \mathbf{j}_*^{**} 's*”. Just like the operators from the set $\text{Span}\{\mathbf{L}, \mathbf{L}_z, \mathbf{L}_+, \mathbf{L}_-\} \simeq \mathfrak{gl}_2(\mathbb{C})$, the \mathbf{j}_*^{**} 's may also act on the space of homogeneous polynomials in the variables $(z_1, z_2, \bar{z}_1, \bar{z}_2)$, to produce polynomials in the same variables.

Commutators of the operators \mathbf{L}_* and \mathbf{j}_*^{} .** A direct computation shows that the commutators between the \mathbf{L}_* 's and the \mathbf{j}_*^{**} 's are given by,

$$[\mathbf{L}, \mathbf{j}_1^{++}] = \frac{1}{2} \mathbf{j}_1^{++}, \quad [\mathbf{L}, \mathbf{j}_1^{+-}] = \frac{1}{2} \mathbf{j}_1^{+-}, \quad [\mathbf{L}, \mathbf{j}_1^{-+}] = -\frac{1}{2} \mathbf{j}_1^{-+}, \quad [\mathbf{L}, \mathbf{j}_1^{--}] = -\frac{1}{2} \mathbf{j}_1^{--},$$

$$[\mathbf{L}, \mathbf{j}_2^{++}] = \frac{1}{2} \mathbf{j}_2^{++}, \quad [\mathbf{L}, \mathbf{j}_2^{+-}] = \frac{1}{2} \mathbf{j}_2^{+-}, \quad [\mathbf{L}, \mathbf{j}_2^{-+}] = -\frac{1}{2} \mathbf{j}_2^{-+}, \quad [\mathbf{L}, \mathbf{j}_2^{--}] = -\frac{1}{2} \mathbf{j}_2^{--},$$

$$[\mathbf{L}_z, \mathbf{j}_1^{++}] = \frac{1}{2} \mathbf{j}_1^{++}, \quad [\mathbf{L}_z, \mathbf{j}_1^{+-}] = -\frac{1}{2} \mathbf{j}_1^{+-}, \quad [\mathbf{L}_z, \mathbf{j}_1^{-+}] = \frac{1}{2} \mathbf{j}_1^{-+}, \quad [\mathbf{L}_z, \mathbf{j}_1^{--}] = -\frac{1}{2} \mathbf{j}_1^{--},$$

$$[\mathbf{L}_z, \mathbf{j}_2^{++}] = \frac{1}{2} \mathbf{j}_2^{++}, \quad [\mathbf{L}_z, \mathbf{j}_2^{+-}] = -\frac{1}{2} \mathbf{j}_2^{+-}, \quad [\mathbf{L}_z, \mathbf{j}_2^{-+}] = \frac{1}{2} \mathbf{j}_2^{-+}, \quad [\mathbf{L}_z, \mathbf{j}_2^{--}] = -\frac{1}{2} \mathbf{j}_2^{--},$$

$$[\mathbf{L}_+, \mathbf{j}_1^{++}] = 0, \quad [\mathbf{L}_+, \mathbf{j}_1^{+-}] = -\mathbf{j}_2^{++}, \quad [\mathbf{L}_+, \mathbf{j}_1^{-+}] = 0, \quad [\mathbf{L}_+, \mathbf{j}_1^{--}] = -\mathbf{j}_1^{-+},$$

$$[\mathbf{L}_+, \mathbf{j}_2^{++}] = 0, \quad [\mathbf{L}_+, \mathbf{j}_2^{+-}] = \mathbf{j}_1^{++}, \quad [\mathbf{L}_+, \mathbf{j}_2^{-+}] = 0, \quad [\mathbf{L}_+, \mathbf{j}_2^{--}] = \mathbf{j}_2^{-+},$$

$$[\mathbf{L}_-, \mathbf{j}_1^{++}] = \mathbf{j}_2^{+-}, \quad [\mathbf{L}_-, \mathbf{j}_1^{+-}] = 0, \quad [\mathbf{L}_-, \mathbf{j}_1^{-+}] = -\mathbf{j}_1^{--}, \quad [\mathbf{L}_-, \mathbf{j}_1^{--}] = 0,$$

$$[\mathbf{L}_-, \mathbf{j}_2^{++}] = -\mathbf{j}_1^{+-}, \quad [\mathbf{L}_-, \mathbf{j}_2^{+-}] = 0, \quad [\mathbf{L}_-, \mathbf{j}_2^{-+}] = \mathbf{j}_2^{--}, \quad [\mathbf{L}_-, \mathbf{j}_2^{--}] = 0.$$

These results show that the subspace generated by the \mathbf{j}_*^{**} 's gets decomposed into four copies of the 2-dimensional irreducible representation of $\mathfrak{gl}_2(\mathbb{C})$:

1. On $\text{Span}\{\mathbf{j}_1^{++}, \mathbf{j}_2^{+-}\}$, we have,

$$\mathbf{L} = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \mathbf{L}_z = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathbf{L}_+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \mathbf{L}_- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

2. On $\text{Span}\{\mathbf{j}_1^{-+}, \mathbf{j}_1^{--}\}$,

$$\mathbf{L} = -\frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \mathbf{L}_z = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathbf{L}_+ = -\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \mathbf{L}_- = -\begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

3. On $\text{Span}\{\mathbf{j}_2^{++}, \mathbf{j}_1^{+-}\}$,

$$\mathbf{L} = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \mathbf{L}_z = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathbf{L}_+ = - \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \mathbf{L}_- = - \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

4. On $\text{Span}\{\mathbf{j}_2^{-+}, \mathbf{j}_2^{--}\}$,

$$\mathbf{L} = -\frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \mathbf{L}_z = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathbf{L}_+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \mathbf{L}_- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

Write,

$$W_+ = W_{+1/2}^{(1++|2+-)} \oplus W_{-1/2}^{(1-+|1--)} = \text{Span}\{\mathbf{j}_1^{++}, \mathbf{j}_2^{+-}\} \oplus \text{Span}\{\mathbf{j}_1^{-+}, \mathbf{j}_1^{--}\},$$

$$W_- = W_{+1/2}^{(2++|1+-)} \oplus W_{-1/2}^{(2-+|2--)} = \text{Span}\{\mathbf{j}_2^{++}, \mathbf{j}_1^{+-}\} \oplus \text{Span}\{\mathbf{j}_2^{-+}, \mathbf{j}_2^{--}\}.$$

Observe that the four subspaces $W_{+1/2}^{(1++|2+-)}$, $W_{-1/2}^{(1-+|1--)}$, $W_{+1/2}^{(2++|1+-)}$ and $W_{-1/2}^{(2-+|2--)}$ are 2-dimensional irreducible representations for \mathfrak{gl}_2 on which the operator \mathbf{L} acts as the scalar $j = \pm 1/2$, as indicated by the subindex in the notation used: $W_j^{(*|*)}$. On each of these 2-dimensional subspaces, the operator \mathbf{L}_z acts diagonally in the given ordered bases as $\text{diag}(+1/2, -1/2)$. Both, W_+ and W_- are symplectic vector spaces and the given decompositions $W_{+1/2}^{(*|*)} \oplus W_{-1/2}^{(*'|*)}$ correspond to the totally isotropic 2-dimensional subspaces $W_j^{(*|*)}$, defined by their corresponding symplectic forms with which one may define 5-dimensional Heisenberg Lie algebras as in [6] and [7] (see §9 below).

Observation. Based on these observations, one may deduce the following possible correspondences with the original Schwinger operators; namely,

$$\begin{aligned} a_x^\dagger &\leftrightarrow \mathbf{j}_1^{++}, & a_y^\dagger &\leftrightarrow \mathbf{j}_2^{+-}, & a_x &\leftrightarrow \mathbf{j}_1^{--}, & a_y &\leftrightarrow \mathbf{j}_1^{-+}, & \text{or,} \\ a_x^\dagger &\leftrightarrow \mathbf{j}_2^{++}, & a_y^\dagger &\leftrightarrow \mathbf{j}_1^{+-}, & a_x &\leftrightarrow \mathbf{j}_2^{--}, & a_y &\leftrightarrow \mathbf{j}_2^{-+}. \end{aligned}$$

9. THE LIE ALGEBRA VS THE LIE SUPERALGEBRA APPROACH

Assuming that the operators \mathbf{L}_* and \mathbf{j}_*^{**} act on an appropriate Hilbert space of functions, \mathcal{H} , one may consider the **Lie algebra**,

$$\mathfrak{g} = \langle \text{Id} \rangle \oplus \mathfrak{gl}_2 \oplus W_+ \oplus W_-,$$

where $\langle \text{Id} \rangle$ stands for the one-dimensional subspace generated by the identity operator $\text{Id} : \mathcal{H} \rightarrow \mathcal{H}$. In this case, the subspaces,

$$\mathfrak{h}_+ = W_+ \oplus \langle \text{Id} \rangle, \quad \text{and} \quad \mathfrak{h}_- = W_- \oplus \langle \text{Id} \rangle,$$

get closed under the Lie bracket $[\cdot, \cdot]$ which is the ordinary commutator of the input operators, and each \mathfrak{h}_* is a 5-dimensional Heisenberg algebra. Moreover,

$[\mathfrak{h}_+, \mathfrak{h}_-] = 0$. Observe that one requires the symplectic forms of W_+ and W_- in order to close the algebraic structure via Lie brackets (commutators).

On the other hand, one may consider the **Lie superalgebra**,

$$\mathfrak{g}/\langle \text{Id} \rangle \simeq \mathfrak{g}' = \mathfrak{gl}_2 \oplus W_+ \oplus W_-$$

where the algebraic structure for pairs of elements in $W_+ \oplus W_-$ gets closed with anticommutators taking values in \mathfrak{gl}_2 . The underlying \mathbb{Z}_2 -graded vector space of this Lie superalgebra is,

$$\mathfrak{g}' = \mathfrak{g}'_0 \oplus \mathfrak{g}'_1, \quad \text{where,} \quad \begin{cases} \mathfrak{g}'_0 = \mathfrak{gl}_2, \\ \mathfrak{g}'_1 = W_+ \oplus W_-. \end{cases}$$

Since $\mathfrak{g}'_1 = W_+ \oplus W_-$ may be decomposed in terms of various copies of the fundamental 2-dimensional irreducible representation of $\mathfrak{g}'_0 = \mathfrak{gl}_2$, we may apply the results from [5] to conclude that **all the anticommutators of the Lie superalgebra \mathfrak{g}' vanish**.

10. THE REPRESENTATION SPACE AND THE WAVE FUNCTIONS

For either the Lie algebra \mathfrak{g} or the Lie superalgebra \mathfrak{g}' , we shall take the representation space to be the L^2 -completion with respect to the hermitian product given by,

$$\langle \psi_1, \psi_2 \rangle = \int_{\mathbb{R}^4} \psi_1(z_1, z_2) \overline{\psi_2(z_1, z_2)} e^{-\frac{1}{2}(|z_1|^2 + |z_2|^2)} dz_1 dz_2 d\bar{z}_1 d\bar{z}_2,$$

in the vector space generated by the harmonic polynomials that result from $(\mathbf{L}_+)^k (\mathbf{j}_1^{+-} \mathbf{j}_2^{+-})^n |0, 0\rangle$, together with those that result from $(\mathbf{L}_+)^k \mathbf{j}_2^{+-} (\mathbf{j}_1^{+-} \mathbf{j}_2^{+-})^n |0, 0\rangle$ ($0 \leq k \leq n$ and $n \in \mathbb{N}$), and the base state space $\langle 0, 0 \rangle$ being generated by the constant functions.

Proposition. The polynomials $(\mathbf{L}_+)^k (\mathbf{j}_1^{+-} \mathbf{j}_2^{+-})^n |0, 0\rangle$ and $(\mathbf{L}_+)^k \mathbf{j}_2^{+-} (\mathbf{j}_1^{+-} \mathbf{j}_2^{+-})^n |0, 0\rangle$ (resp, $(\mathbf{L}_-)^k (\mathbf{j}_2^{++} \mathbf{j}_1^{++})^n |0, 0\rangle$ and $(\mathbf{L}_-)^k \mathbf{j}_1^{++} (\mathbf{j}_2^{++} \mathbf{j}_1^{++})^n |0, 0\rangle$) are linear combinations of monomials of the form $\bar{z}_2^k z_1^{m-k}$ and $\bar{z}_1^k z_2^{m-k}$ ($0 \leq k \leq m$, $m = n$ or $m = n+1$, and $n \in \mathbb{N}$). In particular, these monomials f satisfy, $\Delta f = 0$, where,

$$\Delta = \frac{\partial^2}{\partial z_1 \partial \bar{z}_1} + \frac{\partial^2}{\partial z_2 \partial \bar{z}_2}.$$

11. EXAMPLES

Define $|0, 0\rangle = 1$ and leave the normalization questions aside, for the moment. Define the $j = 1/2$ kets by,

$$\begin{aligned} |1/2, +1/2\rangle &= \mathbf{j}_1^{++} |0, 0\rangle = z_1 |0, 0\rangle = z_1, \\ |1/2, -1/2\rangle &= \mathbf{j}_2^{+-} |0, 0\rangle = z_2 |0, 0\rangle = z_2, \end{aligned}$$

to get the $j = 1/2$ harmonic functions,

$$f_{1/2,+1/2}(r, \theta, \phi, \psi) = \sqrt{r} e^{i\frac{\psi}{2}} e^{i\frac{\phi}{2}} \cos \frac{\theta}{2},$$

$$f_{1/2,-1/2}(r, \theta, \phi, \psi) = \sqrt{r} e^{i\frac{\psi}{2}} e^{-i\frac{\phi}{2}} \sin \frac{\theta}{2}.$$

In terms of the complex coordinates $(z_1, z_2, \bar{z}_1, \bar{z}_2)$, we have,

$$f_{1/2,+1/2}(z_1, z_2, \bar{z}_1, \bar{z}_2) = z_1,$$

$$f_{1/2,-1/2}(z_1, z_2, \bar{z}_1, \bar{z}_2) = z_2.$$

This pair corresponds to the 2-dimensional *spin representation*. The spinor $\begin{pmatrix} z_1 \\ z_2 \end{pmatrix} = \begin{pmatrix} f_{1/2,+1/2} \\ f_{1/2,-1/2} \end{pmatrix}$ may be acted on by the *gauge transformation*, $\begin{pmatrix} z_1 \\ z_2 \end{pmatrix} \mapsto e^{-i\frac{\psi}{2}} \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}$ to eliminate its dependence on ψ . Note indeed that,

$$\mathbf{L}f_{1/2,+1/2} = \frac{1}{2}f_{1/2,+1/2}, \quad \text{and} \quad \mathbf{L}f_{1/2,-1/2} = \frac{1}{2}f_{1/2,-1/2},$$

whereas,

$$\mathbf{L}_z f_{1/2,+1/2} = \frac{1}{2}f_{1/2,+1/2}, \quad \text{and} \quad \mathbf{L}_z f_{1/2,-1/2} = -\frac{1}{2}f_{1/2,-1/2}.$$

On the other hand,

$$\mathbf{L}_- z_1 = z_2, \quad \text{and} \quad \mathbf{L}_+ z_2 = z_1.$$

That is,

$$\begin{aligned} \mathbf{L}_- f_{1/2,+1/2} &= f_{1/2,-1/2}; & ie, & \quad \mathbf{L}_- |1/2, +1/2\rangle = |1/2, -1/2\rangle, \\ \mathbf{L}_+ f_{1/2,-1/2} &= f_{1/2,+1/2}; & ie, & \quad \mathbf{L}_+ |1/2, -1/2\rangle = |1/2, +1/2\rangle. \end{aligned}$$

Notation: Let $\langle j, m \rangle$ stand for the subspace generated by the wave functions $|j, m\rangle$ —also denoted by $f_{j,m}$ — satisfying,

$$\mathbf{L} |j, m\rangle = j |j, m\rangle, \quad \text{and} \quad \mathbf{L}_z |j, m\rangle = m |j, m\rangle.$$

In what follows, we shall use the well known relations,

$$\begin{aligned} \mathbf{L}_+ |j, m\rangle &= \sqrt{(j-m)(j+m+1)} |j, m+1\rangle, \\ \mathbf{L}_- |j, m\rangle &= \sqrt{(j+m)(j-m+1)} |j, m-1\rangle, \end{aligned}$$

Proposition. $\mathbf{j}_1^{+-} \langle 1/2, -1/2 \rangle \subset \langle 1, -1 \rangle$ and $\mathbf{j}_2^{++} \langle 1/2, +1/2 \rangle \subset \langle 1, +1 \rangle$.

Proof: Since $\langle 1/2, -1/2 \rangle = \langle z_2 \rangle$, it follows that, $\mathbf{j}_1^{+-} z_2 = \bar{z}_1 z_2$. Therefore,

$$\mathbf{L}(\bar{z}_1 z_2) = \bar{z}_1 z_2, \quad \text{and} \quad \mathbf{L}_z(\bar{z}_1 z_2) = -\bar{z}_1 z_2.$$

Similarly, since $\langle 1/2, +1/2 \rangle = \langle z_1 \rangle$, it follows that, $\mathbf{j}_2^{++} z_1 = z_1 \bar{z}_2$, and,

$$\mathbf{L}(z_1 \bar{z}_2) = z_1 \bar{z}_2, \quad \text{and} \quad \mathbf{L}_z(z_1 \bar{z}_2) = z_1 \bar{z}_2.$$

Proposition.

$$\begin{aligned} |1, -1\rangle &= \mathbf{j}_1^{+-} z_2 = \mathbf{j}_1^{+-} |1/2, -1/2\rangle = \mathbf{j}_1^{+-} (\mathbf{j}_2^{+-} |0, 0\rangle) = \bar{z}_1 z_2, \\ |1, +1\rangle &= \mathbf{j}_2^{++} z_1 = \mathbf{j}_2^{++} |1/2, +1/2\rangle = \mathbf{j}_2^{++} (\mathbf{j}_1^{++} |0, 0\rangle) = z_1 \bar{z}_2. \end{aligned}$$

Proof: Straightforward computations show that,

$$\mathbf{L}(\bar{z}_1 z_2) = \bar{z}_1 z_2 \Rightarrow j = +1, \quad \text{and} \quad \mathbf{L}(z_1 \bar{z}_2) = z_1 \bar{z}_2 \Rightarrow j = +1,$$

$$\mathbf{L}_z(\bar{z}_1 z_2) = -\bar{z}_1 z_2 \Rightarrow m = -1, \quad \text{and} \quad \mathbf{L}_z(z_1 \bar{z}_2) = z_1 \bar{z}_2 \Rightarrow m = +1.$$

Straightforward computations also show that,

$$\mathbf{L}_+(\bar{z}_1 z_2) = |z_1|^2 - |z_2|^2, \quad \mathbf{L}_+(|z_1|^2 - |z_2|^2) = -2z_1 \bar{z}_2, \quad \mathbf{L}_+(-2z_1 \bar{z}_2) = 0,$$

$$\mathbf{L}_-(z_1 \bar{z}_2) = |z_2|^2 - |z_1|^2, \quad \mathbf{L}_-(|z_2|^2 - |z_1|^2) = -2\bar{z}_1 z_2, \quad \mathbf{L}_-(-2\bar{z}_1 z_2) = 0. \quad \square$$

In particular,

$$\begin{aligned} |z_1|^2 - |z_2|^2 &= r \cos \theta = \mathbf{L}_+ |1, -1\rangle = \sqrt{2} |1, 0\rangle \\ &\Rightarrow |1, 0\rangle = \frac{r}{\sqrt{2}} \cos \theta, \\ -2z_1 \bar{z}_2 &= -r e^{-i\phi} \sin \theta = (\mathbf{L}_+)^2 |1, -1\rangle = \sqrt{2} \mathbf{L}_+ |1, 0\rangle = 2 |1, -1\rangle \\ &\Rightarrow |1, -1\rangle = -\frac{r}{2} e^{-i\phi} \sin \theta. \end{aligned}$$

Therefore, we may write,

$$\begin{pmatrix} f_{1,+1}(r, \theta, \phi) \\ f_{1,0}(r, \theta, \phi) \\ f_{1,-1}(r, \theta, \phi) \end{pmatrix} = \begin{pmatrix} r/2 e^{i\phi} \sin \theta \\ r/\sqrt{2} \cos \theta \\ -r/2 e^{-i\phi} \sin \theta \end{pmatrix}.$$

Observe that up to an overall scalar multiple of r , the triple $\begin{pmatrix} f_{1,+1} \\ f_{1,0} \\ f_{1,-1} \end{pmatrix}$ yields the

triple of spherical harmonics $\begin{pmatrix} Y_1^{+1}(\theta, \phi) \\ Y_1^0(\theta, \phi) \\ Y_1^{-1}(\theta, \phi) \end{pmatrix}$ as expected.

Summary. One may summarize the results obtained in passing from $j = 1/2$ to $j = 1$ by means of the following ‘starting’ diagram:

$$\begin{array}{ccccc} \langle 1, -1 \rangle & \xrightarrow{\mathbf{L}_+} & \langle 1, 0 \rangle & \xleftarrow{\mathbf{L}_-} & \langle 1, +1 \rangle \\ \mathbf{j}_1^{+-} \uparrow & & \nearrow \alpha & & \nwarrow \beta \\ \langle 1/2, -1/2 \rangle & & & & \langle 1/2, +1/2 \rangle \\ & & \mathbf{j}_2^{+-} \searrow & & \nearrow \mathbf{j}_1^{++} \\ & & \langle 0, 0 \rangle & & \end{array}$$

where, the maps α and β can be determined using the commutation relations between the \mathbf{L} 's and the \mathbf{j}_*^{**} 's:

$$\alpha = \mathbf{L}_+ \circ \mathbf{j}_1^{+-} = -\mathbf{j}_2^{++} + \mathbf{j}_1^{+-} \circ \mathbf{L}_+,$$

$$\beta = \mathbf{L}_- \circ \mathbf{j}_2^{++} = -\mathbf{j}_1^{+-} + \mathbf{j}_2^{++} \circ \mathbf{L}_-.$$

12. GENERALIZATION

Proposition. Straightforward computations show that,

$$\begin{aligned} |n, -n\rangle &= (\mathbf{j}_1^{+-} \mathbf{j}_2^{+-})^n |0, 0\rangle = (\bar{z}_1 z_2)^n \\ |n, +n\rangle &= (\mathbf{j}_2^{++} \mathbf{j}_1^{++})^n |0, 0\rangle = (z_1 \bar{z}_2)^n \\ |n+1/2, -(n+1/2)\rangle &= \mathbf{j}_2^{+-} (\mathbf{j}_1^{+-} \mathbf{j}_2^{+-})^n |0, 0\rangle = z_2 (\bar{z}_1 z_2)^n \\ |n+1/2, +(n+1/2)\rangle &= \mathbf{j}_1^{++} (\mathbf{j}_2^{++} \mathbf{j}_1^{++})^n |0, 0\rangle = z_1 (z_1 \bar{z}_2)^n \end{aligned}$$

with,

$$\begin{aligned} \mathbf{L}(\bar{z}_1 z_2)^n &= n (\bar{z}_1 z_2)^n, & \mathbf{L}(z_1 \bar{z}_2)^n &= n (z_1 \bar{z}_2)^n, \\ \mathbf{L}(\bar{z}_1^n z_2^{n+1}) &= \left(n + \frac{1}{2}\right) \bar{z}_1^n z_2^{n+1}, & \mathbf{L}(z_1^{n+1} \bar{z}_2^n) &= \left(n + \frac{1}{2}\right) z_1^{n+1} \bar{z}_2^n, \\ \mathbf{L}_z(\bar{z}_1 z_2)^n &= -n (\bar{z}_1 z_2)^n, & \mathbf{L}_z(z_1 \bar{z}_2)^n &= +n (z_1 \bar{z}_2)^n, \\ \mathbf{L}_z(\bar{z}_1^n z_2^{n+1}) &= -\left(n + \frac{1}{2}\right) \bar{z}_1^n z_2^{n+1}, & \mathbf{L}_z(z_1^{n+1} \bar{z}_2^n) &= +\left(n + \frac{1}{2}\right) z_1^{n+1} \bar{z}_2^n. \quad \square \end{aligned}$$

Proposition. For, for each positive integer n , the $2n+1$ -tuples f obtained from $\oplus_{m=-j}^{+j} \langle j, m \rangle$ having $j = n+1/2$, can be decomposed in the form,

$$f(r, \phi, \theta, \psi) = r^{n+1/2} e^{i\psi/2} \Psi(\theta, \phi), \quad \Psi(\theta, \phi) \in \bigoplus_{m=-j}^{+j} \langle j, m \rangle.$$

In particular, the gauge transformation $f \mapsto e^{-i\psi/2} f$ eliminates the ψ dependence. On the other hand, the $2n+1$ -tuples f obtained from $\oplus_{m=-j}^{+j} \langle j, m \rangle$ having $j = n$, do not depend on ψ , and can be decomposed in the form

$$f(r, \phi, \theta) = r^n \Psi(\theta, \phi), \quad \Psi(\theta, \phi) \in \bigoplus_{m=-j}^{+j} \langle j, m \rangle.$$

Lemma.

$$\begin{aligned} \mathbf{L}_+ (\mathbf{j}_1^{+-} \mathbf{j}_2^{+-}) &= (\mathbf{j}_1^{+-} \mathbf{j}_2^{+-}) \mathbf{L}_+ + (\mathbf{j}_1^{+-} \mathbf{j}_1^{++} - \mathbf{j}_2^{++} \mathbf{j}_2^{+-}) \\ &= (\mathbf{j}_1^{+-} \mathbf{j}_2^{+-}) \mathbf{L}_+ + (|z_1|^2 - |z_2|^2) \text{Id} \\ \mathbf{L}_- (\mathbf{j}_2^{++} \mathbf{j}_1^{++}) &= (\mathbf{j}_2^{++} \mathbf{j}_1^{++}) \mathbf{L}_- + (\mathbf{j}_2^{++} \mathbf{j}_2^{+-} - \mathbf{j}_1^{+-} \mathbf{j}_1^{++}) \\ &= (\mathbf{j}_2^{++} \mathbf{j}_1^{++}) \mathbf{L}_- + (|z_2|^2 - |z_1|^2) \text{Id} \end{aligned}$$

Corollary. Since $[\mathbf{L}_+, \mathbf{j}_2^{+-}] = \mathbf{j}_1^{++}$ and $[\mathbf{L}_-, \mathbf{j}_1^{++}] = \mathbf{j}_2^{+-}$, it follows that,

$$\mathbf{L}_+ \mathbf{j}_2^{+-} = \mathbf{j}_2^{+-} \mathbf{L}_+ + \mathbf{j}_1^{++},$$

implying that,

$$\begin{aligned} \mathbf{L}_+ \mathbf{j}_2^{+-} (\mathbf{j}_1^{+-} \mathbf{j}_2^{+-}) &= \mathbf{j}_2^{+-} \mathbf{L}_+ (\mathbf{j}_1^{+-} \mathbf{j}_2^{+-}) + \mathbf{j}_1^{++} (\mathbf{j}_1^{+-} \mathbf{j}_2^{+-}) \\ &= (\mathbf{j}_2^{+-} \mathbf{j}_1^{+-} \mathbf{j}_2^{+-}) \mathbf{L}_+ + (|z_2|^2 - |z_1|^2) \mathbf{j}_2^{+-} + |z_1|^2 \mathbf{j}_2^{+-} \\ &= (\mathbf{j}_2^{+-} \mathbf{j}_1^{+-} \mathbf{j}_2^{+-}) \mathbf{L}_+ + |z_2|^2 \mathbf{j}_2^{+-} \\ &= \mathbf{j}_2^{+-} (\mathbf{j}_1^{+-} \mathbf{j}_2^{+-}) \mathbf{L}_+ + (|z_2|^2 z_2) \text{Id}. \end{aligned}$$

Corollary. One can:

- (1) Go from $\langle 0, 0 \rangle$ to $\langle j, -j \rangle$ using $(\mathbf{j}_1^{+-}\mathbf{j}_2^{+-})^n$ if $j = n$,
 (or using $\mathbf{j}_2^{+-}(\mathbf{j}_1^{+-}\mathbf{j}_2^{+-})^n$ if $j = n + 1/2$),
 and then move from $\langle j, -j \rangle$ to $\langle j, +j \rangle$ by applying successively \mathbf{L}_+ ; or,
- (2) Go from $\langle 0, 0 \rangle$ to $\langle j, +j \rangle$ using $(\mathbf{j}_2^{++}\mathbf{j}_1^{++})^n$ if $j = n$,
 (or using $\mathbf{j}_1^{++}(\mathbf{j}_2^{++}\mathbf{j}_1^{++})^n$ if $j = n + 1/2$),
 and then move from $\langle j, +j \rangle$ to $\langle j, -j \rangle$ by applying successively \mathbf{L}_- .

In the first case, each application of \mathbf{L}_+ changes the eigenvalue m of \mathbf{L}_z by adding 1 in each step, whereas in the second case, each application of \mathbf{L}_- changes m by subtracting 1 in each step. In particular,

- (3) Application of the operators \mathbf{j}_*^{++} and \mathbf{j}_*^{+-} to any of the above states raise the eigenvalue j by a $1/2$ step, whereas the operators \mathbf{j}_*^{-+} and \mathbf{j}_*^{--} lower j by a $1/2$ step.

13. CONCLUDING REMARKS

We have given a new interpretation of the classical Hurwitz map $H : \mathbb{R}^4 \rightarrow \mathbb{R}^3$, providing an assignment $\mathbb{C}^2 \ni (z_1, z_2) \mapsto (r, \phi, \theta, \psi)$ that realizes the double covering map of groups, $G_0 = \mathbb{R}^+ \times SU(2) \rightarrow \mathbb{R}^+ \times SO(3)$, keeping the real 3-dimensional euclidean space interpretation of both, the radius $r = \sqrt{x_1^2 + x_2^2 + x_3^2}$, and the Euler angles (ϕ, θ, ψ) that parametrize rotations in the group $SO(3)$. The interpretation only requires that $\begin{pmatrix} \bar{z}_1/\sqrt{r} & -\bar{z}_2/\sqrt{r} \\ z_2/\sqrt{r} & z_1/\sqrt{r} \end{pmatrix} \in SU(2)$, whenever $r = |z_1|^2 + |z_2|^2$. The Lie algebra generators, $\{\mathbf{L}, \mathbf{L}_x, \mathbf{L}_y, \mathbf{L}_z\}$ of $\mathfrak{g} = \text{Lie}(G_0)$, written in terms of the complex variables $\{z_1, z_2, \bar{z}_1, \bar{z}_2\}$, satisfy,

$$\mathbf{L}_x^2 + \mathbf{L}_y^2 + \mathbf{L}_z^2 = \mathbf{L}^2 + \mathbf{L} + \frac{r}{4} \Delta,$$

where Δ is the Laplace operator in the domain $\mathbb{R}^4 \simeq \mathbb{C}^2$. We use these facts to produce a Hilbert space \mathcal{H} of harmonic wave functions—depending on the four real variables (r, ϕ, θ, ψ) —with which one can describe either integer or half-integer angular momentum states. The angular momentum operators $\{\mathbf{L}_x, \mathbf{L}_y, \mathbf{L}_z\}$ coincide with their usual expressions in the spherical coordinates (r, ϕ, θ) when they act on states having integer angular momentum, but they show an additional term with a derivative depending on the angle ψ when acting on half-integer angular momentum states. The total angular momentum operator \mathbf{L} gets expressed as $r \partial_r$. Finally, following Schwinger (see [6] and [7]), the 4-dimensional Lie algebra of G_0 is coupled with two Heisenberg Lie algebras of 2-dimensional harmonic oscillators generated by $\{z_1, z_2, \bar{z}_1, \bar{z}_2\}$ and their adjoints thus producing raising and lowering operators that change the total angular momentum in half-units. These two Heisenberg Lie algebras intersect in the identity operator $\text{Id}_{\mathcal{H}}$. All the operators close either into a 9-dimensional Lie algebra or, leaving out only the operator $\text{Id}_{\mathcal{H}}$, into an 8-dimensional Lie superalgebra. Both algebraic structures can be faithfully represented in the Hilbert space \mathcal{H} obtained by the L^2 -completion of the span of the subspace of complex polynomials in the complex variables $(z_1, z_2, \bar{z}_1, \bar{z}_2)$ with the usual Hermitian product having the Hermite weight factor $\exp(-1/2(|z_1|^2 + |z_2|^2))$. Moreover, \mathcal{H}

can be decomposed in the form $\mathcal{H} = \mathcal{H}_0 \oplus \mathcal{H}_1$, where \mathcal{H}_0 contains all the states having integer angular momentum (or *space of bosons*) and \mathcal{H}_1 contains all the states having half-integer angular momentum (or *space of fermions*). The closed subspaces \mathcal{H}_0 and \mathcal{H}_1 are obtained by producing first the highest weight vector $|j, +j\rangle = (z_1 \bar{z}_2)^j$ (if j is integer, with $j \geq 1$) or $|j, +j\rangle = z_1 (z_1 \bar{z}_2)^{\frac{2j-1}{2}}$ (if j is half-integer, with $j \geq 1/2$), and then move from it by successive application of the ladder operator \mathbf{L}_- which preserves the eigenvalue j but lowers the eigenvalue m in one unit at each step, until reaching the lowest weight vector $|j, -j\rangle$, characterized by $\mathbf{L}_- |j, -j\rangle = 0$, or by doing the analogue constructions producing first the lowest weight vectors $|j, -j\rangle$. Similarly, closed subspaces can be selected by producing first a lowest weight vector $|j, -j\rangle$.

Acknowledgements. Two of the authors acknowledge supports received by their research grants. EN-A acknowledges partial support by DGAPA-UNAM under project IN100323. OA-SV acknowledges CONACYT Grant # A1-S-45886.

REFERENCES

- [1] Corwin, L., Ne’eman, Y., and Sternberg, S., *Graded Lie algebras in mathematics and physics (Bose-Fermi symmetry)*, Rev. Mod. Phys. **47** (1975), 573-603.
- [2] Hage Hassan, M., and Kibler, M., *On Hurwitz Transformations*, Proceedings of the Workshop “Le problème de factorisation de Hurwitz: approche historique, solutions, applications en physique”, Eds., A. Ronveaux and D. Lambert (Facultés Universitaires N.D. de la Paix, Namur) (1991 <http://hal.in2p3.fr/in2p3-00115773/>), 1-29.
- [3] Howe, R., *Dual Pairs in Physics: Harmonic Oscillators, Photons, Electrons, and Singletons*, Lectures in Applied Math. **21** (1985), 179-207.
- [4] Kashiwara, M. and Vergne, M., *On the Segal-Shale-Weil Representations and Harmonic Polynomials*, Inventiones math. **44** (1978), 1-47.
- [5] Salgado, G., and Sánchez-Valenzuela, O.A., *Lie superalgebras over \mathfrak{gl}_2* , Communications in Algebra **39** (2011), 2114-2136.
- [6] Schwinger, J., *On Angular Momentum*, Quantum Theory of Angular Momentum (Biedenharn, L.C. and Van Dam, H., eds.), Academic Press, New York, 1965, pp. 229-279.
- [7] Schwinger, J., *Multispinor Basis of Fermi-Bose Transformations*, Annals of Physics **119** (1979), 192-237.