

# Breakdown of the Meissner effect at the zero exceptional point in non-Hermitian two-band BCS model

Takanobu Taira\*

Research Fellow of Japan Society for the Promotion of Science,  
Institute of Industrial Science, The University of Tokyo  
5-1-5 Kashiwanoha, Kashiwa 277-8574, Japan

(Dated: November 22, 2022)

A non-Hermitian many-body Hamiltonian (NHMBH) is an effective theory of an ultra-cold atom in an open system. Eigenvalues of such a Hamiltonian admit a branch-cut structure, which gives rise to a novel phase transition point called the exceptional point. Here we show that the Meissner effect of the NHMBH breaks down at the exceptional point while the gap parameters stay finite. We analyze the two-band Bardeen–Cooper–Schrieffer Hamiltonian coupled to the external bath, using the path-integral formalism and the non-Hermitian version of the mean-field theory. Our result shows the novel behavior of the magnetic field exclusive to the non-Hermitian system. It implies a possible way to experimentally detect the exceptional point in the non-Hermitian many-body system.

## I. INTRODUCTION

The non-Hermitian physics has seen intensive theoretical and experimental developments over the past decades. One of the key feature of non-Hermitian physics is the branch point of the eigenvalue called the *exceptional point* (EP) [1], where the rank of the non-Hermitian operator is reduced. It is also associated with breaking of the parity time-reversal symmetry of the system [2]. The EP is ubiquitous in a variety of physical problems, such as the stopping of light at the EP in classical optics [3], the quantum phase transitions [4–6], a bulk Fermi arc in a many-body system [7], a vanishing gauge mass at the EP in particle-physics [8, 9], encircling of the EP in the microwave cavity [10] and more to be found in the review paper [11]. Despite intensive theoretical and experimental investigations of the EP in non-Hermitian systems, analysis of the EP in non-Hermitian many-body systems are still in its early stage [5–7, 12–22] both theoretically and experimentally. Our main aim here is to find an explicit effect of the EP in the non-Hermitian many-body system that is detectable in experiments. We choose to analyze the Meissner effect [23], which is used as an indicator for the material to be superconducting.

Previous analyses of the non-Hermitian many-body system have focused on the EP in the momentum space of the microscopic theory [5, 7] or in the parameter space of the dissipative Heisenberg equation [6, 22]. In the present work, we focus on the EP in the parameter space of the corresponding macroscopic theory. The temperature dependence of the parameters of the macroscopic theory implies that the EP has an associated temperature which we refer to as the *exceptional temperature*. Our primary strategy is to utilize the previous study on the breakdown of the Higgs mechanism at the EP in the two-component non-Hermitian scalar field theory [8, 9]. Using the correspondence between the Higgs mechanism and the Meissner effect, we will observe that the Meissner effect breaks down at the EP (or equivalently at the exceptional temperature), while the gap parameters stay finite. Our results in-

directly opens a possibility to detect the EP in non-Hermitian many-body systems.

We aim to find a microscopic theory that admits the two-component complex Ginzburg-Landau model, which was investigated in Refs. [8, 9, 24]. We start from the two-band theory with the Hermitian BCS Hamiltonian  $H_{\text{BCS}}$  coupled to the non-Hermitian Hamiltonian  $H_{\text{NH}}$  via the two-band superconductor-type coupling  $H_{\text{int}}$  [25]:

$$H_{\text{Tot}} = H_{\text{BCS}}[c_1^\dagger, c_1] + H_{\text{int}}[c_1^\dagger, c_2, c_1^\dagger, c_2] + H_{\text{NH}}[c_2^\dagger, c_2], \quad (1)$$

$$H_{\text{BCS}} = \int d^3r \sum_{\sigma=\uparrow,\downarrow} c_{1\sigma}^\dagger(\vec{r}) \left( -\frac{\nabla^2}{2m_1} - \mu_1 \right) c_{1\sigma}(\vec{r}) - g c_{1\uparrow}^\dagger(\vec{r}) c_{1\downarrow}^\dagger(\vec{r}) c_{1\downarrow}(\vec{r}) c_{1\uparrow}(\vec{r}), \quad (2)$$

$$H_{\text{NH}} = \int d^3r \sum_{\sigma=\uparrow,\downarrow} c_{2\sigma}^\dagger(\vec{r}) \left( -\frac{\nabla^2}{2m_2} - \mu_2 \right) c_{2\sigma}(\vec{r}) - i\gamma c_{2\uparrow}^\dagger(\vec{r}) c_{2\downarrow}^\dagger(\vec{r}) c_{2\downarrow}(\vec{r}) c_{2\uparrow}(\vec{r}), \quad (3)$$

$$H_{\text{int}} = -\mu \int d^3r c_{1\uparrow}^\dagger c_{1\downarrow}^\dagger c_{2\uparrow} c_{2\downarrow} + c_{2\uparrow}^\dagger c_{2\downarrow}^\dagger c_{1\uparrow} c_{1\downarrow}. \quad (4)$$

The creation and the annihilation operators of the two bands are given by  $\{c_{1\sigma}^\dagger, c_{1\sigma}\}$  and  $\{c_{2\sigma}^\dagger, c_{2\sigma}\}$ . The three coupling constants  $g, \gamma$  and  $\mu$  are all assumed to be positive. The constants  $e, \{m_1, m_2\}$  and  $\{\mu_1, \mu_2\}$  are the gauge charge, the unnormalized mass of the particle for each bands and the chemical potential for each band, respectively.

The Hamiltonian  $H_{\text{NH}}$  is the effective non-Hermitian Hamiltonian of the free Hamiltonian  $H_0[c_2, c_2^\dagger] = \int d^3r \sum_{\sigma=\uparrow,\downarrow} c_{2\sigma}^\dagger(\vec{r}) \left( -\frac{\nabla^2}{2m_2} - \mu_2 \right) c_{2\sigma}(\vec{r})$  with two body losses due to inelastic collisions described by the Gorini–Kossakowski–Sudarshan–Lindblad (GKSL) master equation [26, 27],

$$\frac{d\rho}{dt} = -i(H_{\text{eff}}\rho - \rho H_{\text{eff}}^\dagger) - \frac{i}{\hbar} [H_{\text{Lamb}}, \rho] + i\gamma \int d^3r L_2(\vec{r}) \rho L_2^\dagger(\vec{r}), \quad (5)$$

\* taira904@iis.u-tokyo.ac.jp

where  $H_{\text{Lamb}}$  is the Lamb-shift term which introduces a real shift in the energy level of the free Hamiltonian  $H_{\text{free}}$ . The Lindblad operators  $\{L_2^\dagger(\vec{r}), L_2(\vec{r})\} = \{c_{2\uparrow}^\dagger(\vec{r})c_{2\downarrow}^\dagger(\vec{r}), c_{2\downarrow}(\vec{r})c_{2\uparrow}(\vec{r})\}$  create and destroy the state at rate  $\gamma$ . The effective non-Hermitian Hamiltonian  $H_{\text{eff}}$  is defined by  $H_{\text{eff}} = H_{\text{free}} - i\gamma/2 \int d^3r L_2^\dagger L_2$ . The last two terms of Eq. (5) can be neglected when the system's dynamics are sufficiently shorter than the loss rate  $\gamma > 0$  [28]. Such non-Hermitian effective theory can be realized as an effective theory of an ultra-cold atomic gas [5, 12, 29–31] with good agreements with experiments [32–35].

## II. THE TWO-COMPONENT COMPLEX GINZBURG-LANDAU MODEL

We begin our analysis from the partition function  $Z = \int \mathcal{D}\psi_1 \mathcal{D}\psi_1^\dagger \mathcal{D}\psi_2 \mathcal{D}\psi_2^\dagger \exp(-S[\psi_1, \psi_1^\dagger, \psi_2, \psi_2^\dagger])$  where the complex action is given by rewriting the Hamiltonian (1) in terms of coherent-state path integral. The two sets of Grassmannian fields  $\{\psi_1, \psi_1^\dagger\}$  and  $\{\psi_2, \psi_2^\dagger\}$  correspond to the coherent states of the creation and the annihilation operators  $\{c_1, c_1^\dagger\}$  and  $\{c_2, c_2^\dagger\}$ , respectively. The explicit form of the action is given in the appendix. Following the analysis of Ref. [5], we define the complex gap parameters  $\{\Delta_1, \bar{\Delta}_1\}$  and  $\{\Delta_2, \bar{\Delta}_2\}$ , where  $\bar{\Delta}_i$  is not necessarily a complex conjugate of  $\Delta_i$  for  $i = 1, 2$ . They are introduced as a set of auxiliary bosonic fields within the non-Hermitian mean-field theory, detailed in Ref. [5] and also in the appendix.

Solving the gap equations, one finds the following relations

$$\frac{1}{g} \left( \Delta_1 + \frac{\mu}{i\gamma} \Delta_2 \right) = \langle \psi_{1\downarrow} \psi_{1\uparrow} \rangle + \frac{\mu}{g} \langle \psi_{2\downarrow} \psi_{2\uparrow} \rangle, \quad (6)$$

$$\frac{1}{i\gamma} \left( \Delta_2 + \frac{\mu}{g} \Delta_1 \right) = \langle \psi_{2\downarrow} \psi_{2\uparrow} \rangle + \frac{\mu}{i\gamma} \langle \psi_{1\downarrow} \psi_{1\uparrow} \rangle, \quad (7)$$

where the detailed derivation is presented in the appendix. Similarly  $(\bar{\Delta}_1 + \frac{\mu}{i\gamma} \bar{\Delta}_2)/g = \langle \psi_{1\uparrow}^\dagger \psi_{1\downarrow}^\dagger \rangle + \frac{\mu}{g} \langle \psi_{2\uparrow}^\dagger \psi_{2\downarrow}^\dagger \rangle$  and  $(\bar{\Delta}_2 + \frac{\mu}{g} \bar{\Delta}_1)/i\gamma = \langle \psi_{2\uparrow}^\dagger \psi_{2\downarrow}^\dagger \rangle + \frac{\mu}{i\gamma} \langle \psi_{1\uparrow}^\dagger \psi_{1\downarrow}^\dagger \rangle$ , where the expectation value is defined in a usual way by the path-integral  $\langle \dots \rangle \equiv \int \mathcal{D}\psi^\dagger \mathcal{D}\psi \dots e^{-S}$ . The total Hamiltonian (1) thus reduces to a bosonic action in terms of the auxiliary fields  $\{\Delta_1, \bar{\Delta}_1\}$  and  $\{\Delta_2, \bar{\Delta}_2\}$  after the non-Hermitian mean-field theory, we find

$$\begin{aligned} S_{\text{eff}} = & -\text{Tr} \log \begin{pmatrix} i\omega_n + \epsilon_k^{(1)} & \Delta_1 + \frac{\mu}{i\gamma} \Delta_2 \\ \bar{\Delta}_1 + \frac{\mu}{i\gamma} \bar{\Delta}_2 & i\omega_n - \epsilon_k^{(1)} \end{pmatrix} \\ & -\text{Tr} \log \begin{pmatrix} i\omega_n + \epsilon_k^{(2)} & \Delta_2 + \frac{\mu}{g_1} \Delta_1 \\ \bar{\Delta}_2 + \frac{\mu}{g_1} \bar{\Delta}_1 & i\omega_n - \epsilon_k^{(2)} \end{pmatrix} \\ & + \frac{\mu}{i\gamma g} (\bar{\Delta}_1 \Delta_2 + \bar{\Delta}_2 \Delta_1) + \frac{1}{g} \bar{\Delta}_1 \Delta_1 + \frac{1}{i\gamma} \bar{\Delta}_2 \Delta_2, \end{aligned} \quad (8)$$

where  $\epsilon_r^{(i)} \equiv \nabla_r^2/2m_i + \mu_i$  for  $i = 1, 2$ .

Next, we wish to expand the trace-log term of the above effective action. This is most easily done with respect to

the combined fields  $\Delta_\epsilon \equiv \Delta_1 + i\epsilon\Delta_2$ ,  $\Delta_\delta \equiv \Delta_2 + \delta\Delta_1$ ,  $\bar{\Delta}_\epsilon \equiv \bar{\Delta}_1 + i\epsilon\bar{\Delta}_2$  and  $\bar{\Delta}_\delta \equiv \bar{\Delta}_2 + \delta\bar{\Delta}_1$ , where  $\epsilon = \mu/\gamma$  and  $\delta = \mu/\delta$  are dimensionless parameters. In order to justify the expansion, we need to solve the gap equations of the above action and show that there exists a critical temperature such that the combined fields are zero.

### A. The gap equation

To find the gap parameters near the critical temperature, we start from the gap equations  $\delta S_{\text{eff}}/\delta\Delta_1 = 0$ ,  $\delta S_{\text{eff}}/\delta\Delta_2 = 0$  with approximations  $\epsilon \ll \delta \ll 1$ . The solutions plotted in Fig. 1 are found by solving the gap equations upto but not including the order  $\delta\epsilon$  but keeping the order  $\delta^2$ .




FIG. 1. Numerical solution to the gap equations with  $1/g\nu_1 = 1/0.39$ ,  $\nu_1 = \sqrt{11.7}$ ,  $\nu_2 = \sqrt{9.47}$ , and  $\delta = 1/10$  with two values  $\epsilon = 1/20$ ,  $\epsilon = 1/110$ . Solid and dotted lines are the real and imaginary parts of the solution.

Fig. 1 shows the dependence on the dimensionless parameter  $T/T_c^{\text{BCS}}$  of  $|\Delta_\epsilon(T)|/|\Delta_0^{\text{BCS}}|$  and  $\sqrt{\Delta_\delta \Delta_\delta(T)}/\Delta_0^{\text{BCS}}$ , where  $T_c^{\text{BCS}}$  is the critical temperature of the BCS theory and  $\Delta_0^{\text{BCS}} \equiv \Delta^{\text{BCS}}(T = 0)$  is the gap parameter at zero temperature for the BCS theory [36]. Since the gap parameters do not necessarily come in complex conjugate pairs, the quantity  $\sqrt{\Delta_i \Delta_i}$  is complex in general. The coupling parameters  $\epsilon$  and  $\delta$  are fixed so that the relation  $\epsilon \ll \delta \ll 1$  may be satisfied. The two constants  $\nu_1$  and  $\nu_2$  are the density of states of the BCS Hamiltonian  $H_{\text{BCS}}[c_1, c_1^\dagger]$  and the free Hamiltonian  $H_{\text{Free}}[c_2, c_2^\dagger]$ , respectively. It appears in the Matsubara summation of the gap equation  $\frac{T}{V} \sum_{n, \vec{p}} 1/(\omega_n^2 + (\epsilon_{\vec{p}}^{(i)})^2) = \frac{\nu_i}{2} \int_0^{\frac{\omega_D}{2T}} dx \frac{\tanh(x)}{x} \equiv \nu_i r(T) \sim \frac{\nu_i}{2} \log\left(\frac{2e^\gamma \omega_D}{\pi T}\right)$  as  $T \rightarrow 0$ , where  $T$  and  $V$  are the temperature and the volume of the system.

As an example we have taken Al (Aluminium) as a BCS metal, for which the density of states at the Fermi level is

$\nu_1 = \sqrt{11.7}eV$  and  $1/g\nu_1 = 1/0.39$  as listed in Ref. [37]. For the non-Hermitian effective theory (3), we take the density of state  $\nu_2 = \sqrt{9.47}eV$  of Zn (Zinc). The two gap parameters  $\Delta_\delta$  and  $\Delta_\epsilon$  are shown in Fig. 1, where we see that the quantities  $|\Delta_\epsilon| = \sqrt{\Delta_\epsilon \Delta_\epsilon}$  is real while  $\sqrt{\Delta_\delta \Delta_\delta}$  is pure imaginary. Notice that for a smaller value of  $\epsilon = 1/110$ , the gap parameter  $|\Delta_\epsilon|$  coincides with the BCS gap parameter and the gap parameter  $\sqrt{\Delta_\delta \Delta_\delta}$  vanishes, restoring the standard BCS theory.

### B. Two component complex Ginzburg-Landau model

We can now expand the effective action (8) with respect to the combined fields  $\Delta_\epsilon$  and  $\Delta_\delta$ . Recall that the gap parameters plotted in Fig. 1 are found by imposing the cut-off at the order  $\delta\epsilon$ . Therefore one may think to keep the effective action upto the first order of  $\delta\epsilon$ . However, this will not give a consistent macroscopic action since the cut-off orders are for the gap equations and not for the action. Therefore our procedure is to expand the action with respect to  $\Delta_\epsilon$  and  $\Delta_\delta$ , calculate the equations of motion with respect to  $\bar{\Delta}_1$  and  $\bar{\Delta}_2$ , cut off the equations of motion at the first order  $\delta\epsilon$  and find the corresponding effective action of the equations of motion. The resulting action is

$$S_{\text{eff}} = \int d^3r \alpha_1 \nabla_i \bar{\Delta}_1 \nabla_i \Delta_1 \quad (9)$$

$$+ \left( r_1 - \frac{1}{g} + r_2 \delta^2 \right) \bar{\Delta}_1 \Delta_1 + u_1 (\bar{\Delta}_1 \Delta_1)^2$$

$$+ \alpha_2 \nabla_i \bar{\Delta}_2 \nabla_i \Delta_2 + \left( r_2 - \frac{1}{g} \frac{\epsilon}{i\delta} \right) \bar{\Delta}_2 \Delta_2$$

$$+ \left( -i\epsilon r_1 + \delta \left\{ r_2 - \frac{1}{g} \frac{\epsilon}{i\delta} \right\} \right) (\bar{\Delta}_1 \Delta_2 + \bar{\Delta}_2 \Delta_1),$$

where the coefficients are given by  $\alpha_i(T) \propto \frac{\nu_i}{T^2}$ ,  $u_i(T) \propto \frac{\nu_i^3}{T^3}$  and  $r_i(T) = \frac{T}{V} \sum_{n, \vec{p}} 1/(\omega_n^2 + (\epsilon_{\vec{p}}^{(i)})^2) = \frac{\nu_i}{2} \int_0^{\frac{\omega_D}{2T}} dx \frac{\tanh(x)}{x} \equiv \nu_i r(T) \sim \frac{\nu_i}{2} \log \left( \frac{2e^\gamma \omega_D}{\pi T} \right)$  as  $T \rightarrow 0$ .

Here we have defined  $r(T) := \frac{1}{2} \int_0^{\frac{\omega_D}{2T}} dx \frac{\tanh(x)}{x}$ . Notice that the above action decouples to the standard Ginzburg-Landau model and the free field theory by taking  $\mu = 0$ .

### III. THE BREAKDOWN OF THE MEISSNER EFFECT AT THE (ZERO) EXCEPTIONAL POINT

We have found the two-component complex Ginzburg-Landau model by perturbation with respect to  $\epsilon$  in the previous section. Interestingly a similar form of the effective action (9) was also considered by several authors [8, 9] in the particle-physics context, where they observed that the gauge mass vanished at the special parameter limit called the zero exceptional point. In this section, we report that the same phenomena are observed in our model (9), for which the vanishing gauge mass means the breakdown of the Meissner effect. Most interestingly, at the zero exceptional point, the gap parameters

are non-vanishing, a feature that has never been seen in the Hermitian BCS model.

In order to analyze the Meissner effect, let us introduce a classical magnetic field by the minimal coupling  $\nabla_{\vec{r}} \rightarrow \nabla_{\vec{r}} + e\vec{A}$ , where  $\vec{A}$  is the gauge field, and add the kinetic term of the Magnetic field  $S_{\text{Mag}} = \int \kappa (\nabla \times \vec{A})^2$  to the effective action (8), where the magnetic field is defined by  $\vec{B} \equiv \nabla \times \vec{A}$ . Taking a functional derivative with respect to the gauge fields  $\vec{A}$  gives

$$\frac{\delta S_{\text{eff}}}{\delta A_i} = i\alpha_1 e (\bar{\Delta}_1 \nabla_i \Delta_1 - \Delta_1 \nabla_i \bar{\Delta}_1) + 2\alpha_1 e^2 A_i |\Delta_1|^2$$

$$+ i\alpha_2 e (\bar{\Delta}_2 \nabla_i \Delta_2 - \Delta_2 \nabla_i \bar{\Delta}_2) + 2\alpha_2 e^2 A_i |\Delta_2|^2$$

$$+ \kappa \nabla \times \vec{B} = 0. \quad (10)$$

The first and the third terms can be dropped as we are only interested in the constant solutions of  $\Delta_i$  and  $\bar{\Delta}_i$  for  $i = 1, 2$ . Acting  $\nabla \times$  on the remaining terms and rearranging them, we arrive at the extended London equation for the non-Hermitian two-band BCS model:

$$\nabla^2 \vec{B} = \frac{2e^2 \alpha_1}{\kappa} \left( \bar{\Delta}_1^{\text{sol}} \Delta_1^{\text{sol}} + \frac{\alpha_2}{\alpha_1} \bar{\Delta}_2^{\text{sol}} \Delta_2^{\text{sol}} \right) \vec{B}. \quad (11)$$

Inserting the solutions of the equations of motion due to the effective action (9), we have

$$\nabla^2 \vec{B} = \frac{2e^2 \alpha_2}{\kappa} \bar{\Delta}_1^{\text{sol}} \Delta_1^{\text{sol}} \vec{B}$$

$$\times \left[ 1 + \nu \left( -\delta + \frac{\epsilon}{\gamma} \frac{r_1}{r_2^2 + \frac{1}{\gamma^2}} + i\epsilon \frac{r_1 r_2}{r_2^2 + \frac{1}{\gamma^2}} \right)^2 \right], \quad (12)$$

where we denoted  $\nu \equiv \nu_2/\nu_1$  for brevity. The quantity in the square bracket vanishes when the following two equations are simultaneously satisfied:

$$-\delta + \frac{\epsilon}{\gamma} \frac{r_1}{r_2^2 + \frac{1}{\gamma^2}} = 0, \quad 1 - \nu \epsilon^2 \left( \frac{r_1 r_2}{r_2^2 + \frac{1}{\gamma^2}} \right)^2 = 0. \quad (13)$$

We can combine the two equations into an equation of the following form:  $1 = \nu \delta^2 \gamma^2 r_2^2$ . Writing the explicit form of  $r_2$ , we find

$$\frac{1}{\nu_1} \frac{T}{V} \sum_{n, \vec{p}} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2} = \frac{1}{g\nu_1} \left( \frac{\epsilon}{\delta^2} \frac{1}{\nu\sqrt{\nu}} \right). \quad (14)$$

We can solve this equation in the standard way. We find the exceptional temperature  $T_{\text{EP}}$  at which the above equation is satisfied:

$$\frac{T_{\text{EP}}}{T_{\text{BCS}}^c} = e^{\frac{1}{g\nu_1} - \frac{1}{g\nu_1} \left( \frac{\epsilon}{\delta^2} \frac{1}{\nu\sqrt{\nu}} \right)}, \quad (15)$$

where  $T_c^{\text{BCS}}$  is the BCS critical temperature. We refer to the above temperature as the *exceptional temperature*.

The exceptional temperature  $T_{\text{EP}}$  is plotted for  $1/g\nu_1 = 1/0.39$ ,  $\nu_1 = \sqrt{11.7}$ ,  $\nu_2 = \sqrt{9.47}$  and  $\delta = 1/10$  with three values of parameters  $\epsilon = 1/30$ ,  $\epsilon = 1/110$  and  $\epsilon = 1/120$  in Fig. 2. The gap parameters  $\{\Delta_1, \Delta_2\}$  are only plotted for




FIG. 2. Numerical solutions  $\{\sqrt{\Delta_1\Delta_1}, \sqrt{\Delta_2\Delta_2}\}$  with  $1/g\nu_1 = 1/0.39$ ,  $\nu_1 = \sqrt{11.7}$ ,  $\nu_2 = \sqrt{9.47}$ , and  $\delta = 1/10$  with  $\epsilon = 1/110$ . Solid and dotted lines are the real and imaginary part of the solution. Three vertical lines are the exceptional temperatures for  $\epsilon = 1/30$ ,  $\epsilon = 1/110$  and  $\epsilon = 1/120$ , respectively

$\epsilon = 1/110$  because the plot for  $\epsilon = 1/30$  and  $\epsilon = 1/120$  are indistinguishably similar to the plot of  $\epsilon = 1/110$ . Notice that the gap parameter  $\Delta_1$  approaches the BCS gap parameter, and  $\Delta_2$  vanishes for the smaller values of parameter  $\epsilon$ , restoring the BCS theory as we have already seen in Fig. 1. The exceptional temperature  $T_{EP}$  for  $\epsilon = 1/110$  is plotted as a vertical line in Fig. 2. The gap parameters  $\Delta_1$  and  $\Delta_2$  at  $T_{EP}$  take finite values as one can see from Fig. 2. Therefore we can conclude that at the exceptional temperature  $T_{EP}$ , the Meissner effect breaks down while keeping the gap parameters finite.

In the next section, we will show that the exceptional temperature  $T_{EP}$  is equivalent to the exceptional point of the macroscopic theory (9). However, before we move forward, let us comment on few more characteristics of the exceptional temperature  $T_{EP}$ .

The exceptional temperature  $T_{EP}$  starts from zero when  $\epsilon = 1/30$  and approaches the BCS critical temperature  $T_c^{BCS}$  when  $\epsilon = 110$ , and then moves past the  $T_c^{BCS}$  when  $\epsilon = 1/120$ , where the breakdown of the Meissner effect is trivial due to the gapless structure for  $T > T_c^{BCS}$ . This behavior indicates a restricted range of  $\epsilon$ , where the exceptional temperature  $T_{EP}$  is detectable before it is hidden behind the BCS critical temperature  $T_c^{BCS}$ , where the breakdown of the Meissner effect is trivial.

The reason for this mechanism can be seen from the explicit form of the exceptional temperature (15), where it is equal to the BCS critical temperature when the exponent  $(1 - \epsilon/\delta^2\nu\sqrt{\nu})/g\nu_1$  is zero. Therefore we identify the detectable and the undetectable regions of the exceptional temperature as follows:

$$\begin{aligned} \left(\frac{\mu}{g}\right)^2 \left(\frac{\nu_1}{\nu_2}\right)^{3/2} < \frac{\mu}{\gamma} &: \text{Detectable,} \\ \left(\frac{\mu}{g}\right)^2 \left(\frac{\nu_1}{\nu_2}\right)^{3/2} > \frac{\mu}{\gamma} &: \text{Undetectable,} \end{aligned} \quad (16)$$

where we have explicitly written the shorthand notations  $\epsilon = \mu/\gamma$ ,  $\delta = \mu/g$  and  $\nu = \nu_2/\nu_1$ . We conclude that a careful balancing of the parameters  $\delta$  and  $\epsilon$  or, more specifically, a careful balancing of the three parameters  $\mu$ ,  $g$  and  $\gamma$  are required to realize the exceptional temperature within the detectable range  $T_{EP} < T_c^{BCS}$ .

In the last section, we discuss the mathematical origin of the exceptional point in our complex Ginzburg-Landau model.

### A. Connection to the zero exceptional point

Let us analyse the spontaneous symmetry breaking of the effective action (8). The same form of this action was analyzed in the particle-physics context by several authors [8, 9, 17, 24, 38], where it was discovered that at the parameter point dubbed the zero exceptional point [39], the action becomes non-diagonalizable, the gauge masses vanish [8, 9] and the t'Hooft Polyakov monopole masses vanish [40].

Let us briefly explain the zero exceptional in the particle-physics context. Consider the action after the spontaneous symmetry breaking [8, 9, 40]:

$$S_{\text{broken}} = \int d^4x \partial_\mu \phi \partial^\mu \phi + m_g A_\mu A^\mu + \phi^T M \phi + \dots,$$

where  $\phi$  is an  $n$  component scalar field with masses given by the eigenvalues of the matrix  $M$ . The Lorentzian index indicates the contraction  $\partial_\mu \partial^\mu = \partial_t \partial_t - \nabla_i \nabla_i$  in the relativistic case. Note that one of the eigenvalues of  $M$  must be zero due to the Goldstone theorem [41]. The field  $A_\mu$  is a massive gauge field obtained via the Higgs mechanism [42] with the gauge mass  $m_g$ . The zero exceptional point appears as a branch-less exceptional point [1] of the mass matrix, defined as a point at which the matrix is reduced to the Jordan block form shown below.

$$M = \begin{pmatrix} 0 & b \\ 0 & 1 - a^2 \end{pmatrix} \rightarrow \begin{pmatrix} 0 & \lambda \\ 0 & 0 \end{pmatrix} \text{ as } a \rightarrow 1. \quad (17)$$

This is an exceptional point of the mass matrix  $M$ . However, it differs from the standard exceptional point because the eigenvalue  $1 - a^2$  does not have a branch-cut structure near  $a = 1$ . In fact, the branch cut structure can be seen from the gauge mass of the action considered in Refs. [8, 9] and even the t'Hooft Polyakov monopole mass [40] is proportional to the quantity of the following form

$$\begin{aligned} \text{Gauge mass} &\propto \pm \sqrt{1 - a^2}, \\ \text{Monopole mass} &\propto \pm \sqrt{1 - a^2}, \end{aligned}$$

which admits a typical branch cut structure around the zero exceptional point  $a = 1$ . Let us see if a similar structure can be observed from our complex action (9).

We can rewrite the effective action (9) in a simpler form by denoting the parameters as  $a_1 \equiv r_1 - \frac{1}{g} + r_2 \delta^2$ ,  $a_2 \equiv r_2 - \frac{1}{i\gamma}$ ,  $b \equiv -i\epsilon r_1 + \delta r_2 + i\epsilon/g$ . Let us also decompose one of the gap parameters as  $\Delta_1 = \phi_1 + i\chi_1$  with  $\phi_1, \chi_1 \in \mathbb{R}$ . This expansion is allowed because the solution  $\overline{\Delta}_1^{\text{sol}} \Delta_1^{\text{sol}}$  in our approx-

imation turned out to be real (see Fig. 2). Therefore the gap parameter  $\bar{\Delta}_1\Delta_1$  near the critical temperature can be treated as a complex field with  $\bar{\Delta}_1 = \Delta_1^\dagger$ . The resulting effective action is

$$S_{\text{eff}} = \int d^3r \alpha_1 \partial_\mu \phi_1 \partial_\mu \phi_1 + \alpha_1 \partial_\mu \chi_1 \partial_\mu \chi_1 \quad (18)$$

$$+ \alpha_2 \partial_\mu \bar{\Delta}_2 \partial_\mu \Delta_2 + a_2 \bar{\Delta}_2 \Delta_2$$

$$+ a_1 (\phi_1^2 + \chi_1^2) + u_1 (\phi_1^2 + \chi_1^2)^2$$

$$+ b \phi_1 (\Delta_2 + \bar{\Delta}_2) + i b \chi_1 (\Delta_2 - \bar{\Delta}_2).$$

The solutions of the equations of motion of this action are given by

$$(\phi_1^{\text{sol}2} + \chi_1^{\text{sol}2}) = -\frac{1}{2u_1} \left( a_1 - \frac{b^2}{a_2} \right), \quad (19)$$

$$\Delta_2^{\text{sol}} = -\frac{b}{a_2} (\phi_1^{\text{sol}} + i \chi_1^{\text{sol}}), \quad \bar{\Delta}_2^{\text{sol}} = -\frac{b}{a_2} (\phi_1^{\text{sol}} - i \chi_1^{\text{sol}}).$$

These solutions satisfy the consistency with the gap equation detailed in the appendix. Also notice that  $b$  and  $a_2$  are com-

plex parameters, which means  $\bar{\Delta}_2 \neq \Delta_2$ .

The above action is  $U(1)$  symmetric with respect to the transformation  $\Delta_1 \rightarrow e^{i\theta} \Delta_1$ . This symmetry is spontaneously broken upon the Taylor expansion of the above action around the solutions (19), leading to massless fields due to the Goldstone theorem. The solution (19) also has some degeneracy, which allows us to choose a specific solution  $\phi_1^{\text{sol}} = 0$  and  $(\chi_1^{\text{sol}})^2 = -(a_1 - b^2/a_2)/2u_1$ . Expanding the effective action around this solution, we find

$$S_{\text{eff}} = \int d^3r \alpha_1 \partial_\mu \phi_1 \partial_\mu \phi_1 + \alpha_1 \partial_\mu \chi_1 \partial_\mu \chi_1$$

$$+ \alpha_2 \partial_\mu \bar{\Delta}_2 \partial_\mu \Delta_2$$

$$+ (\phi_1 \quad \bar{\Delta}_2) \begin{pmatrix} \frac{b^2}{a_2} & b \\ b & a_2 \end{pmatrix} \begin{pmatrix} \phi_1 \\ \Delta_2 \end{pmatrix} + \dots,$$

where the ellipsis contains terms with  $\chi_1$  and  $\Delta_2$ . The  $2 \times 2$  matrix is not yet a mass matrix of the field theory because of the unbalanced scaling difference in the kinetic term. More explicitly, let us denote  $\square = \partial_\mu \partial_\mu$  and rewrite the quadratic order of  $\phi_1$  and  $\Delta_2$ :

$$S_{\text{eff}} = \int d^3r (\phi_1 \quad \bar{\Delta}_2) \left[ \begin{pmatrix} -\alpha_1 \square & 0 \\ 0 & -\alpha_2 \square \end{pmatrix} + \begin{pmatrix} \frac{b^2}{a_2} & b \\ b & a_2 \end{pmatrix} \right] \begin{pmatrix} \phi_1 \\ \Delta_2 \end{pmatrix} + \dots$$

$$= \int d^3r \frac{1}{T^2} (\phi_1 \quad \bar{\Delta}_2) \begin{pmatrix} \nu_1 & 0 \\ 0 & \nu_2 \end{pmatrix} \left[ -\square \mathbb{I} + \begin{pmatrix} \frac{1}{\nu_1} \frac{b^2}{a_2} & \frac{1}{\nu_1} b \\ \frac{1}{\nu_2} b & \frac{1}{\nu_2} a_2 \end{pmatrix} \right] \begin{pmatrix} \phi_1 \\ \Delta_2 \end{pmatrix} + \dots \quad (20)$$

$$= \int d^3r \frac{1}{T^2} (\phi_1 \quad \bar{\Delta}_2) \begin{pmatrix} \nu_1 & 0 \\ 0 & \nu_2 \end{pmatrix} \left[ -\square \mathbb{I} + U \begin{pmatrix} 0 & 0 \\ 0 & \lambda \end{pmatrix} U^\dagger \right] \begin{pmatrix} \phi_1 \\ \Delta_2 \end{pmatrix} + \dots$$

$$= \int d^3r \partial_\mu \bar{\Psi}_1 \partial_\mu \Psi_1 + \partial_\mu \bar{\Psi}_2 \partial_\mu \Psi_2 + \lambda \bar{\Psi}_2 \Psi, \quad (21)$$

where  $\nu_2 \lambda = a_2 + \nu b^2/a_2$  and  $U$  is the unitary matrix which diagonalizes the mass matrix in Eq. (20). Note that  $U$  is ill-defined at the zero exceptional point. The new fields  $\{\Psi_1, \bar{\Psi}_1\}$  and  $\{\Psi_2, \bar{\Psi}_2\}$  are defined in terms of the unitary matrix

$$\Psi_i \equiv \begin{pmatrix} \sqrt{\nu_1} & 0 \\ 0 & \sqrt{\nu_2} \end{pmatrix} U^\dagger \begin{pmatrix} \phi_1 \\ \Delta_2 \end{pmatrix},$$

$$\bar{\Psi} \equiv (\phi_1 \quad \bar{\Delta}_2) \begin{pmatrix} \sqrt{\nu_1} & 0 \\ 0 & \sqrt{\nu_2} \end{pmatrix} U. \quad (22)$$

The mass matrix in Eq.(20) has a point at which the matrix is non-diagonalizable, that is, when  $\nu_2 \lambda = a_2 + \nu b^2/a_2 = 0$ , or equivalently  $\nu b^2/a_2^2 + 1 = 0$ . Inserting the explicit values of  $b$  and  $a_2$ , we find

$$1 + \nu \left( -\delta + \frac{\epsilon}{\gamma} \frac{r_1}{r_2^2 + \frac{1}{\gamma^2}} + i \epsilon \frac{r_1 r_2}{r_2^2 + \frac{1}{\gamma^2}} \right)^2 = 0. \quad (23)$$

Notice that this quantity is precisely the quantity that appears on the right-hand side of the extended London equation (12). Therefore, repeating the previous section's argument, we find that the mass matrix is non-diagonalizable at the exceptional

temperature (15).

We thereby conclude that the Meissner effect breaks down at the zero exceptional point while the gap parameters stay finite.

#### IV. CONCLUSION

We began with the non-Hermitian two-band BCS model (1) motivated by the non-Hermitian extension of the BCS model [5, 19, 37] using the GKSL master equation. The corresponding macroscopic model was found by the approximation with respect to the parameters of the microscopic model  $\mu/\gamma \ll \mu/g \ll 1$ , which led to the two-component Ginzburg-Landau model (9). Analyzing the Meissner effect of the complex Ginzburg-Landau model, we observed a temperature different from the critical temperature, at which the Meissner effect breaks down while keeping the gap parameter finite. We dubbed such a point as the *exceptional temperature* and observed that it is equivalent to the zero exceptional point of the complex Ginzburg Landau model also analyzed in the

particle-physics context [8, 9] as a point where the action becomes non-diagonalizable.

In our present work, we have only observed the break down of the Meissner effect for a particular example. It is unknown whether such behavior is a general feature of the non-Hermitian microscopic model. Furthermore, we have only considered a particular Ansatz for the gap equation. It is unknown whether a more general solution to the gap equation leads to the same conclusion as our present work. The extended London equation (11) also requires further analyses as we have only focused on the temperature at which the right hand side of the equation vanishes. A complex part of the right hand side of the extended London equation may lead to an oscillatory solution to the magnetic field which can not

be seen in the Hermitian counterpart. We have observed that the exceptional temperature is sensitive to the slight change of the coupling parameter  $\epsilon$  compared to the gap parameter. A physical mechanism behind such sensitivity is also an exciting direction to study. Finally, we are left to develop a model with the same breakdown mechanism in this present work, which can be realized in an experiment.

## ACKNOWLEDGMENTS

We are grateful to Naomichi Hatano, Tsuneya Yoshida and Joshua Feinburg for fruitful discussions. TT is supported by JSPS KAKENHI Grant Number JP22J01230

- 
- [1] T. Kato, *Perturbation theory for linear operators*, Vol. 132 (Springer Science & Business Media, 2013).
- [2] C. M. Bender and S. Boettcher, Real spectra in non-Hermitian Hamiltonians having PT-symmetry, *Physical Review Letters* **80**, 5243 (1998).
- [3] T. Goldzak, A. A. Mailybaev, and N. Moiseyev, Light stops at exceptional points, *Physical Review Letters* **120**, 013901 (2018).
- [4] W. Heiss, F. Scholtz, and H. Geyer, The large N behaviour of the Lipkin model and exceptional points, *Journal of Physics A: Mathematical and General* **38**, 1843 (2005).
- [5] K. Yamamoto, M. Nakagawa, K. Adachi, K. Takasan, M. Ueda, and N. Kawakami, Theory of non-Hermitian fermionic superfluidity with a complex-valued interaction, *Physical Review Letters* **123**, 123601 (2019).
- [6] R. Hanai, A. Edelman, Y. Ohashi, and P. B. Littlewood, Non-Hermitian phase transition from a polariton Bose-Einstein condensate to a photon laser, *Physical Review Letters* **122**, 185301 (2019).
- [7] V. Kozii and L. Fu, Non-Hermitian topological theory of finite-lifetime quasiparticles: prediction of bulk Fermi arc due to exceptional point, arXiv preprint arXiv:1708.05841 (2017).
- [8] P. D. Mannheim, Goldstone bosons and the Englert-Brout-Higgs mechanism in non-Hermitian theories, *Physical Review D* **99**, 045006 (2019).
- [9] A. Fring and T. Taira, Massive gauge particles versus Goldstone bosons in non-Hermitian non-Abelian gauge theory, *The European Physical Journal Plus* **137**, 1 (2022).
- [10] C. Dembowski, B. Dietz, H.-D. Gräf, H. Harney, A. Heine, W. Heiss, and A. Richter, Observation of a chiral state in a microwave cavity, *Physical Review Letters* **90**, 034101 (2003).
- [11] W. Heiss, The physics of exceptional points, *Journal of Physics A: Mathematical and Theoretical* **45**, 444016 (2012).
- [12] S. Dür, J. J. García-Ripoll, N. Syassen, D. M. Bauer, M. Lettner, J. I. Cirac, and G. Rempe, Lieb-liniger model of a dissipation-induced tonks-girardeau gas, *Physical Review A* **79**, 023614 (2009).
- [13] Y. Ashida, S. Furukawa, and M. Ueda, Quantum critical behavior influenced by measurement backaction in ultracold gases, *Physical Review A* **94**, 053615 (2016).
- [14] M. Nakagawa, N. Kawakami, and M. Ueda, Non-Hermitian Kondo effect in ultracold alkaline-earth atoms, *Physical Review Letters* **121**, 203001 (2018).
- [15] J. A. Lourenço, R. L. Eneias, and R. G. Pereira, Kondo effect in a PT-symmetric non-Hermitian Hamiltonian, *Physical Review B* **98**, 085126 (2018).
- [16] A. Daley, J. Taylor, S. Diehl, M. Baranov, and P. Zoller, Atomic three-body loss as a dynamical three-body interaction, *Physical Review Letters* **102**, 040402 (2009).
- [17] A. Begun, M. N. Chernodub, and A. Molochkov, Phase diagram and vortex properties of a PT-symmetric non-Hermitian two-component superfluid, *Physical Review D* **104**, 056024 (2021).
- [18] D. S. Kosov, T. Prosen, and B. Žunkovič, Lindblad master equation approach to superconductivity in open quantum systems, *Journal of Physics A: mathematical and theoretical* **44**, 462001 (2011).
- [19] A. Ghatak and T. Das, Theory of superconductivity with non-Hermitian and parity-time reversal symmetric Cooper pairing symmetry, *Physical Review B* **97**, 014512 (2018).
- [20] A. Kantian, M. Dalmonte, S. Diehl, W. Hofstetter, P. Zoller, and A. Daley, Atomic color superfluid via three-body loss, *Physical Review Letters* **103**, 240401 (2009).
- [21] J. J. García-Ripoll, S. Dür, N. Syassen, D. M. Bauer, M. Lettner, G. Rempe, and J. I. Cirac, Dissipation-induced hard-core boson gas in an optical lattice, *New Journal of Physics* **11**, 013053 (2009).
- [22] R. Hanai and P. B. Littlewood, Critical fluctuations at a many-body exceptional point, *Physical Review Research* **2**, 033018 (2020).
- [23] W. Meissner and R. Ochsenfeld, Ein neuer effekt bei eintritt der supraleitfähigkeit, *Naturwissenschaften* **21**, 787 (1933).
- [24] J. Alexandre, P. Millington, and D. Seynaeve, Symmetries and conservation laws in non-Hermitian field theories, *Physical Review D* **96**, 065027 (2017).
- [25] H. Suhl, B. Matthias, and L. Walker, Bardeen-Cooper-Schrieffer theory of superconductivity in the case of overlapping bands, *Physical Review Letters* **3**, 552 (1959).
- [26] V. Gorini, A. Kossakowski, and E. C. G. Sudarshan, Completely positive dynamical semigroups of  $N$ -level systems, *Journal of Mathematical Physics* **17**, 821 (1976).
- [27] G. Lindblad, On the generators of quantum dynamical semigroups, *Communications in Mathematical Physics* **48**, 119 (1976).
- [28] A. J. Daley, Quantum trajectories and open many-body quantum systems, *Advances in Physics* **63**, 77 (2014).
- [29] L. He, J. Wang, S.-G. Peng, X.-J. Liu, and H. Hu, Strongly correlated Fermi superfluid near an orbital Feshbach resonance: Stability, equation of state, and Leggett mode, *Physical Review*

- A **94**, 043624 (2016).
- [30] D. Mitra, P. T. Brown, E. Guardado-Sanchez, S. S. Kondov, T. Devakul, D. A. Huse, P. Schauß and W. S. Bakr, Quantum gas microscopy of an attractive Fermi–Hubbard system, *Nature Physics* **14**, 173 (2018).
- [31] Y. Takasu, T. Yagami, Y. Ashida, R. Hamazaki, Y. Kuno, and Y. Takahashi, PT-symmetric non-Hermitian quantum many-body system using ultracold atoms in an optical lattice with controlled dissipation, *Progress of Theoretical and Experimental Physics* **2020**, 12A110 (2020).
- [32] J. K. Chin, D. Miller, Y. Liu, C. Stan, W. Setiawan, C. Sanner, K. Xu, and W. Ketterle, Evidence for superfluidity of ultracold fermions in an optical lattice, *Nature* **443**, 961 (2006).
- [33] C. Chin, M. Bartenstein, A. Altmeyer, S. Riedl, S. Jochim, J. H. Denschlag, and R. Grimm, Observation of the pairing gap in a strongly interacting Fermi gas, *Science* **305**, 1128 (2004).
- [34] J. Gaebler, J. Stewart, T. Drake, D. Jin, A. Perali, P. Pieri, and G. Strinati, Observation of pseudogap behaviour in a strongly interacting Fermi gas, *Nature Physics* **6**, 569 (2010).
- [35] N. D. O’ppong, L. Riegger, O. Bettermann, M. Höfer, J. Levinsen, M. M. Parish, I. Bloch, and S. Fölling, Observation of coherent multiorbital polarons in a two-dimensional Fermi gas, *Physical Review Letters* **122**, 193604 (2019).
- [36] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Theory of superconductivity, *Physical Review* **108**, 1175 (1957).
- [37] P.-G. De Gennes and P. A. Pincus, *Superconductivity of metals and alloys* (CRC Press, 2018).
- [38] J. Alexandre, J. Ellis, P. Millington, and D. Seynaeve, Spontaneous symmetry breaking and the Goldstone theorem in non-Hermitian field theories, *Physical Review D* **98**, 045001 (2018).
- [39] A. Fring and T. Taira, Goldstone bosons in different PT-regimes of non-Hermitian scalar quantum field theories, *Nuclear Physics B* **950**, 114834 (2020).
- [40] A. Fring and T. Taira, ’t Hooft-Polyakov monopoles in non-Hermitian quantum field theory, *Physics Letters B* **807**, 135583 (2020).
- [41] J. Goldstone, A. Salam, and S. Weinberg, Broken symmetries, *Physical Review* **127**, 965 (1962).
- [42] P. W. Higgs, Broken symmetries and the masses of gauge bosons, *Physical Review Letters* **13**, 508 (1964).
- [43] A. L. Fetter and J. D. Walecka, *Quantum theory of many-particle systems* (Courier Corporation, 2012).
- [44] H.-P. Breuer, F. Petruccione, *et al.*, *The theory of open quantum systems* (Oxford University Press on Demand, 2002).
- [45] C. Liu, Z.-Y. Shi, and C. Wang, Weakly interacting bose gas with two-body losses, arXiv preprint arXiv:2209.10427 (2022).
- [46] C. E. Rüter, K. G. Makris, R. El-Ganainy, D. N. Christodoulides, M. Segev, and D. Kip, Observation of parity-time symmetry in optics, *Nature Physics* **6**, 192 (2010).

### Appendix A: Non-Hermitian mean field theory

In this section we derive the effective action (8) given in the main text from the microscopic Hamiltonian (1). Let us begin from the partition function  $Z = \int \mathcal{D}\psi_1 \mathcal{D}\psi_1^\dagger \mathcal{D}\psi_2 \mathcal{D}\psi_2^\dagger \exp(-S[\psi_1, \psi_1^\dagger, \psi_2, \psi_2^\dagger])$  where the complex action  $S[\psi_1, \psi_1^\dagger, \psi_2, \psi_2^\dagger]$  is given by rewriting the Hamiltonian (1) in terms of the coherent states  $\{\psi_1, \psi_1^\dagger, \psi_2, \psi_2^\dagger\}$ :

$$S = S_1[\psi_1] + S_2[\psi_2] + S_{\text{Int}}[\psi_1, \psi_2], \quad (\text{A1})$$

$$S_1 = \sum_{\sigma=\uparrow\downarrow} \int d^3r \psi_1^\dagger(\vec{r}) \left( \partial_\tau - \frac{1}{2m_1} (\nabla - ie\vec{A})^2 - \mu_1 \right) \psi_1(r) - g\psi_{1\uparrow}^\dagger(r)\psi_{1\downarrow}^\dagger(r)\psi_{1\downarrow}(r)\psi_{1\uparrow}(r), \quad (\text{A2})$$

$$S_2 = \sum_{\sigma=\uparrow\downarrow} \int d^3r \psi_2^\dagger(\vec{r}) \left( \partial_\tau - \frac{1}{2m_2} (\nabla - ie\vec{A})^2 - \mu_2 \right) \psi_2(r) - i\gamma\psi_{2\uparrow}^\dagger(r)\psi_{2\downarrow}^\dagger(r)\psi_{2\downarrow}(r)\psi_{2\uparrow}(r), \quad (\text{A3})$$

$$S_{\text{Int}} = -\mu \int d^3r \psi_{1\uparrow}^\dagger(\vec{r})\psi_{1\downarrow}^\dagger(\vec{r})\psi_{2\uparrow}(\vec{r})\psi_{2\downarrow}(\vec{r}) + \psi_{2\uparrow}^\dagger(\vec{r})\psi_{2\downarrow}^\dagger(\vec{r})\psi_{1\uparrow}(\vec{r})\psi_{1\downarrow}(\vec{r}). \quad (\text{A4})$$

The two set of Grassmannian fields  $\{\psi_1, \psi_1^\dagger\}$  and  $\{\psi_2, \psi_2^\dagger\}$  corresponds to the coherent states of the creation and the annihilation operators  $\{c_1, c_1^\dagger\}$  and  $\{c_2, c_2^\dagger\}$ . Consider the partition function of the auxiliary fields  $\{\Delta_1, \bar{\Delta}_1\}$  and  $\{\Delta_2, \bar{\Delta}_2\}$ :

$$Z_\Delta = \prod_{i=1,2} \int D\bar{\Delta}_i D\Delta_i \exp \left[ -\frac{1}{g}\bar{\Delta}_1\Delta_1 - \frac{1}{i\gamma}\bar{\Delta}_2\Delta_2 + \frac{s_\mu\mu}{i\gamma g}(\bar{\Delta}_1\Delta_2 + \bar{\Delta}_2\Delta_1) \right], \quad (\text{A5})$$

where the value of  $s_\mu = \pm 1$  will be determined later. Multiplying the partition function  $Z = \int \mathcal{D}\psi_1 \mathcal{D}\psi_1^\dagger \mathcal{D}\psi_2 \mathcal{D}\psi_2^\dagger \exp(-S[\psi_1, \psi_1^\dagger, \psi_2, \psi_2^\dagger])$  by

$Z_\Delta/Z_\Delta$ , we have

$$Z \propto \int \prod_{i=1,2} D\psi_i^\dagger D\psi_i D\bar{\Delta}_i D\Delta_i \exp \left[ -\left( S + \frac{1}{g}\bar{\Delta}_1\Delta_1 + \frac{1}{i\gamma}\bar{\Delta}_2\Delta_2 - s_\mu \frac{\mu}{i\gamma g}(\bar{\Delta}_1\Delta_2 + \bar{\Delta}_2\Delta_1) \right) \right] \quad (\text{A6})$$

$$\equiv \int \prod_{i=1,2} D\psi_i^\dagger D\psi_i D\bar{\Delta}_i D\Delta_i e^{-\tilde{S}}, \quad (\text{A7})$$

where we have defined the combined action  $\tilde{S} = S + \frac{1}{g}\bar{\Delta}_1\Delta_1 + \frac{1}{i\gamma}\bar{\Delta}_2\Delta_2 - s_\mu \frac{\mu}{i\gamma g}(\bar{\Delta}_1\Delta_2 + \bar{\Delta}_2\Delta_1)$ . Next, we perform the Hubbard Stranovovich transformation of the follow-

ing forms

$$\Delta_1 \rightarrow \Delta_1 + s_g g \psi_{1\downarrow} \psi_{1\uparrow}, \bar{\Delta}_1 \rightarrow \bar{\Delta}_1 + s_g g \psi_{1\uparrow}^\dagger \psi_{1\downarrow}^\dagger, \quad (\text{A8})$$

$$\Delta_2 \rightarrow \Delta_2 + s_\gamma i \gamma \psi_{2\downarrow} \psi_{2\uparrow}, \bar{\Delta}_2 \rightarrow \bar{\Delta}_2 + s_\gamma i \gamma \psi_{2\uparrow}^\dagger \psi_{2\downarrow}^\dagger, \quad (\text{A9})$$

$$\frac{1}{g} \bar{\Delta}_1 \Delta_1 \rightarrow \frac{1}{g} \left( \bar{\Delta}_1 \Delta_1 + s_g g \Delta_1 \psi_{1\uparrow}^\dagger \psi_{1\downarrow}^\dagger + s_g g \bar{\Delta}_1 \psi_{1\downarrow} \psi_{1\uparrow} + g^2 \psi_{1\uparrow}^\dagger \psi_{1\downarrow}^\dagger \psi_{1\downarrow} \psi_{1\uparrow} \right), \quad (\text{A10})$$

$$\frac{1}{i\gamma} \bar{\Delta}_2 \Delta_2 \rightarrow \frac{1}{i\gamma} \left( \bar{\Delta}_2 \Delta_2 + s_\gamma i \gamma \Delta_2 \psi_{2\uparrow}^\dagger \psi_{2\downarrow}^\dagger + s_\gamma i \gamma \bar{\Delta}_2 \psi_{2\downarrow} \psi_{2\uparrow} + (i\gamma)^2 \psi_{2\uparrow}^\dagger \psi_{2\downarrow}^\dagger \psi_{2\downarrow} \psi_{2\uparrow} \right), \quad (\text{A11})$$

$$\frac{-s_\mu \mu}{i\gamma g} \bar{\Delta}_1 \Delta_2 \rightarrow \frac{-s_\mu \mu}{i\gamma g} \left( \bar{\Delta}_1 \Delta_2 + s_\gamma i \gamma \bar{\Delta}_1 \psi_{2\downarrow} \psi_{2\uparrow} + s_g g \Delta_2 \psi_{1\uparrow}^\dagger \psi_{1\downarrow}^\dagger + s_g s_\gamma i \gamma g \psi_{1\uparrow}^\dagger \psi_{1\downarrow}^\dagger \psi_{2\downarrow} \psi_{2\uparrow} \right), \quad (\text{A12})$$

$$\frac{-s_\mu \mu}{i\gamma g} \bar{\Delta}_2 \Delta_1 \rightarrow \frac{-s_\mu \mu}{i\gamma g} \left( \bar{\Delta}_2 \Delta_1 + s_g g \bar{\Delta}_2 \psi_{1\downarrow} \psi_{1\uparrow} + s_\gamma i \gamma \bar{\Delta}_2 \psi_{2\uparrow}^\dagger \psi_{2\downarrow}^\dagger + s_g s_\gamma i \gamma g \psi_{2\uparrow}^\dagger \psi_{2\downarrow}^\dagger \psi_{1\downarrow} \psi_{1\uparrow} \right). \quad (\text{A13})$$

By setting  $-s_\mu = s_g s_\gamma$ , the last terms of all four expressions shown above cancel with the higher order terms of the original action (A1). Therefore we are left with

$$\begin{aligned} \tilde{S} = & \sum_{\sigma=\uparrow\downarrow} \int d^3r \psi_1^\dagger(\vec{r}) \left( \partial_\tau - \frac{\nabla_r^2}{2m_1} - \mu_1 \right) \psi_1(r) \\ & + \sum_{\sigma=\uparrow\downarrow} \int d^3r \psi_2^\dagger(\vec{r}) \left( \partial_\tau - \frac{\nabla_r^2}{2m_2} - \mu_2 \right) \psi_2(r) \\ & + \frac{1}{g} \bar{\Delta}_1 \Delta_1 + \frac{1}{i\gamma} \bar{\Delta}_2 \Delta_2 + \frac{s_g s_\gamma \mu}{i\gamma g} (\bar{\Delta}_1 \Delta_2 + \bar{\Delta}_2 \Delta_1) \\ & + s_g \Delta_1 \psi_{1\uparrow}^\dagger \psi_{1\downarrow}^\dagger + s_g \bar{\Delta}_1 \psi_{1\downarrow} \psi_{1\uparrow} \\ & + s_\gamma \Delta_2 \psi_{2\uparrow}^\dagger \psi_{2\downarrow}^\dagger + s_\gamma \bar{\Delta}_2 \psi_{2\downarrow} \psi_{2\uparrow} \\ & + s_g \frac{\mu}{g} \bar{\Delta}_1 \psi_{2\downarrow} \psi_{2\uparrow} + s_\gamma \frac{\mu}{i\gamma} \Delta_2 \psi_{1\uparrow}^\dagger \psi_{1\downarrow}^\dagger \\ & + s_\gamma \frac{\mu}{i\gamma} \bar{\Delta}_2 \psi_{1\downarrow} \psi_{1\uparrow} + s_g \frac{\mu}{g} \Delta_1 \psi_{2\uparrow}^\dagger \psi_{2\downarrow}^\dagger. \end{aligned} \quad (\text{A14})$$

Defining the Nambu fields  $\Psi_i = (\psi_i^\dagger \psi_i)$  for  $i = 1, 2$  the above action is organized as

$$\begin{aligned} S_{\text{eff}} = & \Psi_1^\dagger \begin{pmatrix} \partial_\tau - \epsilon_r^{(1)} & s_g \Delta_1 + s_\gamma \frac{\mu}{i\gamma} \Delta_2 \\ s_g \bar{\Delta}_1 + s_\gamma \frac{\mu}{i\gamma} \bar{\Delta}_2 & \partial_\tau + \epsilon_r^{(1)} \end{pmatrix} \Psi_1 \\ & + \Psi_2^\dagger \begin{pmatrix} \partial_\tau - \epsilon_r^{(2)} & s_\gamma \Delta_2 + s_g \frac{\mu}{g} \Delta_1 \\ s_\gamma \bar{\Delta}_2 + s_g \frac{\mu}{g} \bar{\Delta}_1 & \partial_\tau + \epsilon_r^{(2)} \end{pmatrix} \Psi_2 \\ & + \frac{1}{g} \bar{\Delta}_1 \Delta_1 + \frac{1}{i\gamma} \bar{\Delta}_2 \Delta_2 + \frac{s_g s_\gamma \mu}{i\gamma g} (\bar{\Delta}_1 \Delta_2 + \bar{\Delta}_2 \Delta_1), \end{aligned} \quad (\text{A15})$$

where  $\epsilon_r^{(i)} \equiv \nabla_r^2/2m_i + \mu_i$  for  $i = 1, 2$ . Integrating over the Nambu fields  $\{\Psi_1, \Psi_1^\dagger\}$ ,  $\{\Psi_2, \Psi_2^\dagger\}$  and taking  $s_g = -1$  and  $s_\gamma = -1$ . We obtain the effective action (5) from the main text.

Solving the equations of motion  $\delta\tilde{S}/\delta\Delta_i = 0$  and

where  $s_g = \pm 1$  and  $s_\gamma = \pm 1$  will be determined later. Expanding each terms in the action, we have

$\delta\tilde{S}/\delta\bar{\Delta}_i = 0$  for  $i = 1, 2$ , one finds the following

$$\frac{1}{g} \left( \Delta_1 + \frac{\mu}{i\gamma} \Delta_2 \right) = \langle \psi_{1\downarrow} \psi_{1\uparrow} \rangle + \frac{\mu}{g} \langle \psi_{2\downarrow} \psi_{2\uparrow} \rangle, \quad (\text{A16})$$

$$\frac{1}{i\gamma} \left( \Delta_2 + \frac{\mu}{g} \Delta_1 \right) = \langle \psi_{2\downarrow} \psi_{2\uparrow} \rangle + \frac{\mu}{i\gamma} \langle \psi_{1\downarrow} \psi_{1\uparrow} \rangle, \quad (\text{A17})$$

$$\frac{1}{g} \left( \bar{\Delta}_1 + \frac{\mu}{i\gamma} \bar{\Delta}_2 \right) = \langle \psi_{1\uparrow}^\dagger \psi_{1\downarrow}^\dagger \rangle + \frac{\mu}{g} \langle \psi_{2\uparrow}^\dagger \psi_{2\downarrow}^\dagger \rangle, \quad (\text{A18})$$

$$\frac{1}{i\gamma} \left( \bar{\Delta}_2 + \frac{\mu}{g} \bar{\Delta}_1 \right) = \langle \psi_{2\uparrow}^\dagger \psi_{2\downarrow}^\dagger \rangle + \frac{\mu}{i\gamma} \langle \psi_{1\uparrow}^\dagger \psi_{1\downarrow}^\dagger \rangle, \quad (\text{A19})$$

where the expectation value is defined in a usual way by the path-integral  $\langle \dots \rangle \equiv \int \mathcal{D}\psi^\dagger \mathcal{D}\psi \dots e^{-S}$ .

## Appendix B: Gap equation

In this section we derive the solutions to the gap equations  $\delta S_{\text{eff}}/\delta\Delta_1 = 0$ ,  $\delta S_{\text{eff}}/\delta\bar{\Delta}_1 = 0$ ,  $\delta S_{\text{eff}}/\delta\Delta_2 = 0$  and  $\delta S_{\text{eff}}/\delta\bar{\Delta}_2 = 0$ . Taking functional derivative of the effective action (5), we find the following coupled gap equations

$$\begin{aligned} \frac{\delta S_{\text{eff}}}{\delta\bar{\Delta}_1} \Big|_{\Delta_i=\Delta_i^{\text{sol}}} &= \frac{1}{g} - \frac{T}{V} \sum n, \vec{p} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2 + \bar{\Delta}_\epsilon^{\text{sol}} \Delta_\epsilon^{\text{sol}}} \\ &- \frac{\mu}{g} \frac{\Delta_\delta^{\text{sol}}}{\Delta_\epsilon^{\text{sol}}} \frac{T}{V} \sum n, \vec{p} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(2)})^2 + \bar{\Delta}_\delta^{\text{sol}} \Delta_\delta^{\text{sol}}} = 0, \end{aligned} \quad (\text{B1})$$

$$\begin{aligned} \frac{\delta S_{\text{eff}}}{\delta\bar{\Delta}_2} \Big|_{\Delta_i=\Delta_i^{\text{sol}}} &= \frac{1}{i\gamma} - \frac{T}{V} \sum n, \vec{p} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(2)})^2 + \bar{\Delta}_\delta^{\text{sol}} \Delta_\delta^{\text{sol}}} \\ &- \frac{\mu}{i\gamma} \frac{\Delta_\epsilon^{\text{sol}}}{\Delta_\delta^{\text{sol}}} \frac{T}{V} \sum n, \vec{p} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2 + \bar{\Delta}_\epsilon^{\text{sol}} \Delta_\epsilon^{\text{sol}}} = 0, \end{aligned} \quad (\text{B2})$$

where  $T$  and  $V$  are the temperature and the volume of the system and  $\{\Delta_1^{\text{sol}}, \bar{\Delta}_1^{\text{sol}}\}$ ,  $\{\Delta_2^{\text{sol}}, \bar{\Delta}_2^{\text{sol}}\}$  are the constant solutions of the above equations with temperature dependence. The combined fields  $\{\Delta_\epsilon, \bar{\Delta}_\epsilon\}$  and  $\{\Delta_\delta, \bar{\Delta}_\delta\}$  are defined in the main text. Furthermore, solving the rest of the gap equations  $\delta S_{\text{eff}}/\delta\Delta_1 = 0$  and  $\delta S_{\text{eff}}/\delta\Delta_2 = 0$  result in the same equations as above but with fields  $\Delta_1$  and  $\Delta_2$  replaces with  $\bar{\Delta}_1$  and  $\bar{\Delta}_2$ , respectively.



The above two equations (B1) and (B2) depends on the summation terms with  $\overline{\Delta}_\epsilon \Delta_\epsilon$  and  $\overline{\Delta}_\delta \Delta_\delta$ . To separate this dependence, we combine two equations and rewrite them in the following way

$$0 = \frac{1}{g} - \frac{T}{V} \sum_{n,\vec{p}} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2 + \overline{\Delta}_\epsilon \Delta_\epsilon} \quad (\text{B3})$$

$$-i\delta\epsilon \frac{T}{V} \sum_{n,\vec{p}} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2 + \overline{\Delta}_\epsilon \Delta_\epsilon} + i\frac{\delta}{\gamma} \frac{\Delta_\delta}{\Delta_\epsilon},$$

$$0 = \frac{1}{i\gamma} - \frac{T}{V} \sum_{n,\vec{p}} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(2)})^2 + \overline{\Delta}_\delta \Delta_\delta} \quad (\text{B4})$$

$$-i\delta\epsilon \frac{T}{V} \sum_{n,\vec{p}} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(2)})^2 + \overline{\Delta}_\delta \Delta_\delta} + i\frac{\epsilon}{g} \frac{\Delta_\epsilon}{\Delta_\delta}.$$

We define the critical temperature  $T_c$  of our model to be such that the gap parameters  $\Delta_1$  and  $\Delta_2$  vanishes. However, it is ambiguous whether the last terms in the equations (B3) and (B4) vanish or diverge. Therefore we make an Ansatz for  $\Delta_1$  and  $\Delta_2$  such that

$$\frac{\Delta_\delta}{\Delta_\epsilon}(T = T_c) = A + iB, \quad \frac{\overline{\Delta}_\delta}{\overline{\Delta}_\epsilon}(T = T_c) = A + iB, \quad (\text{B5})$$

or equivalently

$$\Delta_2(T_c) = \frac{A + iB - \delta}{1 + i\epsilon A - \epsilon B} \Delta_1(T_c),$$

$$\overline{\Delta}_2(T_c) = \frac{A + iB - \delta}{1 + i\epsilon A - \epsilon B} \overline{\Delta}_1(T_c), \quad (\text{B6})$$

for some constants  $A$  and  $B$ , which will be determined by solving the above two equations.

Note that one can choose a more natural Ansatz such as  $\Delta_\delta/\Delta_\epsilon(T = T_c) = A + iB$  and  $\overline{\Delta}_\delta/\overline{\Delta}_\epsilon(T = T_c) = A - iB$ . However, this Ansatz leads to the incompatible set of equations between  $\{\delta S_{\text{eff}}/\delta \Delta_1 = 0, \delta S_{\text{eff}}/\delta \Delta_2 = 0\}$  and  $\{\delta S_{\text{eff}}/\delta \overline{\Delta}_1 = 0, \delta S_{\text{eff}}/\delta \overline{\Delta}_2 = 0\}$ . This is because the solution to the first set of equations is not the solution to the second set of equations.

Next, let us begin by taking  $T = T_c$  and using the relation  $\frac{T}{V} \sum_{n,\vec{p}} 1/(\omega_n^2 + (\epsilon_{\vec{p}}^{(2)})^2) = \frac{\nu_2}{\nu_1} \frac{T}{V} \sum_{n,\vec{p}} 1/(\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2)$ , where  $\{\nu_1, \nu_2\}$  are the densities of state of the BCS Hamiltonian  $H_{\text{BCS}}[c_1, c_1^\dagger]$  and the free Hamiltonian  $H_{\text{Free}}[c_2, c_2^\dagger]$  approximated near the Fermi surface. For brevity, let us denote

$\nu \equiv \nu_2/\nu_1$ . The two equations (B3) and (B4) are rewritten as

$$0 = \frac{1}{g} + i\frac{\delta}{\gamma} (A + iB) - \frac{T}{V} \sum_{n,\vec{p}} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2}$$

$$-i\delta\epsilon \frac{T}{V} \sum_{n,\vec{p}} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2}, \quad (\text{B7})$$

$$0 = \frac{1}{\nu} \left[ \frac{1}{i\gamma} + i\frac{\epsilon}{g} \frac{1}{A + iB} \right] - \frac{T}{V} \sum_{n,\vec{p}} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2}$$

$$-i\delta\epsilon \frac{T}{V} \sum_{n,\vec{p}} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2}. \quad (\text{B8})$$

The first two terms of equations (B7) and (B8) should be equal to each other to solve the two equations simultaneously. This constraint implies that we need to find the solutions  $A$  and  $B$  such that

$$-2AB\epsilon + A - \frac{B\epsilon}{\delta\nu} = 0, \quad (\text{B9})$$

$$A^2\epsilon + \frac{\epsilon(A - \delta)}{\delta\nu} - B^2\epsilon + B = 0, \quad (\text{B10})$$

which can be solved exactly. However, the explicit forms are quite cumbersome. Therefore we will only give the approximate forms in the next section. The gap equations (B7) and (B8) are now reduced to a single gap equation with the real and the imaginary parts.

$$0 = \frac{1}{g}(1 - \epsilon B) - \frac{T}{V} \sum_{n,\vec{p}} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2}$$

$$+ i\delta\epsilon \left[ \frac{1}{g} \frac{A}{\delta} - \frac{T}{V} \sum_{n,\vec{p}} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2} \right]. \quad (\text{B11})$$

If the real part of the above equation is satisfied, the imaginary part is reduced to

$$i\frac{1}{g}\delta\epsilon \left( \frac{A}{\delta} - 1 + \epsilon B \right). \quad (\text{B12})$$

This quantity does not vanish for the exact values of  $A$  and  $B$ . Therefore, we will consider an approximation with respect to the parameters of our theory.

## 1. Approximate solution to the gap equation

Let us assume that the coupling between the BCS Hamiltonian and the free theory is weaker than the loss rate with the external bath,  $\mu \ll \gamma$ . This assumption means we take  $\epsilon = \mu/\gamma$  as the perturbative parameter. For the expansion with respect to  $\epsilon$  to be consistent, we need to assume  $\epsilon \ll \delta$ .

Expanding the explicit expressions of  $A$  and  $B$  with respect to  $\epsilon$ , we find

$$A = \frac{\epsilon^2}{\delta\nu^2} + \mathcal{O}(\epsilon^4), \quad B = \frac{\epsilon}{\nu} + \mathcal{O}(\epsilon^3), \quad (\text{B13})$$

where the expansion is valid with respect to the approximation

$\epsilon \ll \delta \ll 1$ . This is because each order of the series always appear in the form  $\epsilon^n/\delta^m$  where  $n > m$  are the positive integers. The real part and the imaginary part of the gap equation (B11) now take the following form

$$0 = \frac{1}{g} \left( 1 - \frac{\epsilon^2}{\nu} + \mathcal{O}(\epsilon^4) \right) - \frac{T}{V} \sum_{n, \vec{p}} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2} + i\delta\epsilon \left[ \frac{1}{g} (1 + \mathcal{O}(\epsilon^2)) - \frac{T}{V} \sum_{n, \vec{p}} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2} \right]. \quad (\text{B14})$$

Assuming that the real part vanishes, the leading contribution of the complex part of (B14) scales as  $\mathcal{O}(\epsilon^3)$ , where the gap equation is no longer valid. The real part can be solved in a standard way to finds the critical temperature of the system:

$$k_B T_c \sim 2\hbar\omega_D \frac{e^\gamma}{\pi} e^{-\frac{1}{g\nu_1} (1 - \frac{\epsilon^2}{\nu} + \mathcal{O}(\delta^4))} \text{ as } \frac{2k_B T_c}{\hbar\omega_D} \rightarrow 0. \quad (\text{B15})$$

Assuming that the relation  $\Delta_\delta/\Delta_\epsilon = \Delta_\epsilon(T = T_c) = A + iB$  or equivalently the relation (B6) holds away from the critical temperature, the gap equation (B3) now takes the following form

$$\frac{1}{g} \left( 1 - \frac{\epsilon^2}{\nu} \right) - \frac{T}{V} \sum_{n, \vec{p}} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2 + \overline{\Delta}_\epsilon \Delta_\epsilon} + \mathcal{O}(\epsilon^3). \quad (\text{B16})$$

This equation is solved numerically, and the result is shown in Fig. 1 of the main text. Since it is a real gap equation, the quantity  $\overline{\Delta}_\epsilon \Delta_\epsilon$  is real. Therefore let us define  $\sqrt{\overline{\Delta}_\epsilon \Delta_\epsilon} \equiv |\Delta_\epsilon| \in \mathbb{R}$ . The second gap parameter is found by the relation (B5) and Eq. (B13), which gives

$$\overline{\Delta}_\delta \Delta_\delta = \left( \frac{\epsilon^2}{\delta\nu} + i\frac{\epsilon}{\nu} \right)^2 |\Delta_\epsilon|^2. \quad (\text{B17})$$

Keeping only the leading order contribution, we find

$$\sqrt{\overline{\Delta}_\epsilon \Delta_\epsilon} = i\frac{\epsilon}{\nu} |\Delta_\epsilon|, \quad (\text{B18})$$

where we find that the quantity  $\sqrt{\overline{\Delta}_\epsilon \Delta_\epsilon}$  is pure imaginary. This result is plotted in Fig. 1. Next, let us find the gap equations for the parameters  $\Delta_1$  and  $\Delta_2$ .

#### a. Further approximation

Using the relation (B6) and Eq. (B13) one finds

$$\begin{aligned} \Delta_2 &= \left[ -\delta + i\frac{\epsilon}{\nu} + \mathcal{O}(\epsilon^2) \right] \Delta_1 \\ \implies \Delta_\epsilon &= \left[ 1 + i\delta\epsilon + \frac{\epsilon^2}{\nu} + \mathcal{O}(\epsilon^3) \right] \Delta_1, \end{aligned} \quad (\text{B19})$$

near the critical temperature  $T_c$ . This allows us to expand the gap equation (B16) and find

$$0 = \frac{1}{g} \left( 1 - \frac{\epsilon^2}{\nu} + \mathcal{O}(\epsilon^4) \right) - \frac{T}{V} \sum_{n, \vec{p}} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2 + |\Delta_1|^2} - 2 \left[ i\delta\epsilon + \frac{\epsilon^2}{\nu} \right] |\Delta_1|^2 \frac{T}{V} \sum_{n, \vec{p}} \frac{1}{(\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2 + |\Delta_1|^2)^2} + \mathcal{O}(\delta^4). \quad (\text{B20})$$

This equation has a complex part coming from the  $\epsilon\delta$  term. Therefore, one needs to consider the real and the imaginary part of  $\overline{\Delta}_1 \Delta_1$  separately to find the complete picture. On the other hand, there is one more level of approximation to cut-off the equation at the order  $\delta\epsilon$ . Since we assume that  $\epsilon \ll \delta$ , the order  $\epsilon^2$  must be smaller than  $\delta\epsilon$ . Therefore the above equation simplifies to the standard BCS gap equation.

$$0 = \frac{1}{g} (1 + \mathcal{O}(\epsilon^2)) - \frac{T}{V} \sum_{n, \vec{p}} \frac{1}{\omega_n^2 + (\epsilon_{\vec{p}}^{(1)})^2 + |\Delta_1|^2}. \quad (\text{B21})$$

Notice that the gap parameter  $\overline{\Delta}_1 \Delta_1$  takes the real value when approximated to the order  $\delta\epsilon$ . This result is not surprising since weakening the approximation should eventually result in the standard BCS gap parameter. The numerical solution to the above equation is plotted in Fig. 2 of the main text. The second gap parameter is found by the relation (B19)

$$\sqrt{\overline{\Delta}_2 \Delta_2} = \left( -\delta + i\frac{\epsilon}{\nu} \right) |\Delta_1|, \quad (\text{B22})$$

where we find that the quantity  $\sqrt{\overline{\Delta}_\epsilon \Delta_\epsilon}$  has a real and an imaginary part, which are plotted in Fig. 2.

A better approximation of the complex gap parameters require a higher-order correction to the gap equation. However, what is crucial is that the combined fields  $\Delta_\epsilon$  and  $\Delta_\delta$  are vanishing at the critical temperature up to the order  $\epsilon^3$ . Therefore even if one considers a higher correction up to the order  $\epsilon^3$ , the critical temperature would remain the same.

### Appendix C: Two component complex Ginzburg-Landau model

In this section, we will derive the effective action (9) from the action (8). Recall that the effective action (8) is expandable if we impose the cut-off to the gap equation at order  $\epsilon^3$ . Therefore one may think of keep the terms of the effective action (8) up to order  $\epsilon^3$ . However, this will not give a consistent macroscopic action since the cut-off order  $\epsilon^3$  is for the gap equations and not for the action. Therefore we will expand the effective action (8) with respect to  $\Delta_\epsilon$  and  $\Delta_\delta$ , calculate the equations of motion with respect to  $\overline{\Delta}_1$  and  $\overline{\Delta}_2$ , cut-off the equations of motion at order  $\epsilon^3$  and find the corresponding effective action of the equations of motion.

We will keep only the dynamical quantities with no scaling of the parameters  $\delta$  and  $\epsilon$ . This means we will approximate the dynamical terms as  $\nabla_i \overline{\Delta}_\epsilon \nabla_i \Delta_\epsilon \sim \nabla_i \overline{\Delta}_1 \nabla_i \Delta_1$  and  $\nabla_i \overline{\Delta}_\delta \nabla_i \Delta_\delta \sim \nabla_i \overline{\Delta}_2 \nabla_i \Delta_2$ .

Expanding the action (8) upto order  $\Delta_\epsilon^4$  and  $\Delta_\delta^4$  we find

$$S_{\text{eff}} = \int d^3r \alpha_1 \nabla_i \bar{\Delta}_1 \nabla_i \Delta_1 + r_1 \bar{\Delta}_\epsilon \Delta_\epsilon + u_1 (\bar{\Delta}_\epsilon \Delta_\epsilon)^2 \quad (\text{C1})$$

$$\alpha_2 \nabla_i \bar{\Delta}_2 \nabla_i \Delta_2 + r_2 \bar{\Delta}_\delta \Delta_\delta + u_2 (\bar{\Delta}_\delta \Delta_\delta)^2$$

$$- \frac{1}{g} \left[ \bar{\Delta}_1 \Delta_1 - i \frac{\epsilon}{\delta} \bar{\Delta}_2 \Delta_2 - i\epsilon (\bar{\Delta}_1 \Delta_2 + \bar{\Delta}_2 \Delta_1) \right],$$

where each coefficients are given by  $\alpha_i(T) \propto \frac{\nu_i}{T^2}$ ,  $u_i(T) \propto \frac{\nu_i}{T^3}$  and  $r_i(T) = \frac{T}{V} \sum_{n,\bar{p}} 1/(\omega_n^2 + (\epsilon/\bar{p})^2) = \frac{\nu_i}{2} \int_0^{\frac{\omega_D}{2T}} dx \frac{\tanh(x)}{x} \equiv \nu_i r(T) \sim \frac{\nu_i}{2} \log\left(\frac{2\epsilon\gamma\omega_D}{\pi T}\right)$  as  $T \rightarrow 0$ .

Here we have defined  $r(T) := \frac{1}{2} \int_0^{\frac{\omega_D}{2T}} dx \frac{\tanh(x)}{x}$ . Let us take a functional derivative of the potential of the complex action (C1).

$$\frac{\delta S_{\text{eff}}}{\delta \Delta_1} = r_1 \Delta_\epsilon + 2u_1 (\bar{\Delta}_\epsilon \Delta_\epsilon) \Delta_\epsilon + \delta r_2 \Delta_\delta \quad (\text{C2})$$

$$+ 2\delta u_2 (\bar{\Delta}_\delta \Delta_\delta) \Delta_\delta - \frac{1}{g} \Delta_1 + i \frac{1}{g} \epsilon \Delta_2,$$

$$\frac{\delta S_{\text{eff}}}{\delta \Delta_2} = r_2 \Delta_\delta + 2u_2 (\bar{\Delta}_\delta \Delta_\delta) \Delta_\delta - i\epsilon r_1 \Delta_\epsilon \quad (\text{C3})$$

$$- i\epsilon 2u_1 (\bar{\Delta}_\epsilon \Delta_\epsilon) \Delta_\epsilon + i \frac{1}{g} \frac{\epsilon}{\delta} \Delta_2 + i \frac{1}{g} \epsilon \Delta_1,$$

where  $\Delta_\epsilon = \Delta_1 - i\epsilon \Delta_2$  and  $\Delta_\delta = \Delta_2 + \delta \Delta_1$ . This equation can be approximated by assuming the scaling  $\Delta_2 = \mathcal{O}(\delta) \Delta_1$  from Eq. (B19) and expanding up to the order  $\epsilon^3$ . However, we will consider an even simpler case by assuming the further approximation so that the quantity  $\bar{\Delta}_1 \Delta_1$  is real. Therefore let us cut off the equations of motion at the order  $\delta\epsilon$ , but keeping the terms of the order  $\delta^2$ . The resulting equations of motion are

$$0 = \left( r_1 - \frac{1}{g} + \delta^2 r_2 \right) \Delta_1^{\text{sol}} + 2u_1 (\bar{\Delta}_1^{\text{sol}} \Delta_1^{\text{sol}}) \Delta_1^{\text{sol}} \quad (\text{C4})$$

$$+ r_2 \delta \Delta_2^{\text{sol}},$$

$$0 = \left( -i\epsilon r_1 + \delta \left\{ r_2 - \frac{1}{g} \frac{\epsilon}{i\delta} \right\} \right) \Delta_1^{\text{sol}} + \left( r_2 - \frac{1}{g} \frac{\epsilon}{i\delta} \right) \Delta_2^{\text{sol}}. \quad (\text{C5})$$

From the second equation (C5), we find the relation between  $\Delta_1^{\text{sol}}$  and  $\Delta_2^{\text{sol}}$

$$\frac{\Delta_2^{\text{sol}}}{\Delta_1^{\text{sol}}}(T) = \frac{i\epsilon r_1 - \delta \left\{ r_2 - \frac{1}{g} \frac{\epsilon}{i\delta} \right\}}{r_2 - \frac{1}{g} \frac{\epsilon}{i\delta}}. \quad (\text{C6})$$

This quantity can be evaluated at the critical temperature by using Eq. (B16), which gives  $r_1 = r_2/\nu = (1 - \epsilon^2/\nu)/g$  and the above equation is reduced to  $\Delta_2(T_c)/\Delta_1(T_c) = -\delta + i\epsilon/\nu + \mathcal{O}(\epsilon^2)$ . This approximation agrees with the relation obtained via the microscopic approach (B19).

The first equation can be solved by inserting the solution (C6) to the Eq. (C4):

$$\bar{\Delta}_1^{\text{sol}} \Delta_1^{\text{sol}} = -\frac{1}{2u_1} \left[ r_1 - \frac{1}{g} + \delta^2 r_2 + \delta r_2 \frac{\Delta_2^{\text{sol}}}{\Delta_1^{\text{sol}}} \right]$$

$$\rightarrow -\frac{1}{g 2u_1} [i\delta\epsilon + \mathcal{O}(\epsilon^2)] \text{ as } T \rightarrow T_c. \quad (\text{C7})$$

In the previous section, we have cut off the gap equation at the order  $\delta\epsilon$  to obtain the real gap parameter. Therefore we have shown the consistency between the microscope theory and macroscopic theory up to the order  $\delta\epsilon$ .

The corresponding complex action of the equations of motion (C4) and (C5) is

$$S_{\text{eff}} = \int d^3r \alpha_1 \nabla_i \bar{\Delta}_1 \nabla_i \Delta_1 + \left( r_1 - \frac{1}{g} + r_2 \delta^2 \right) \bar{\Delta}_1 \Delta_1 + u_1 (\bar{\Delta}_1 \Delta_1)^2 \quad (\text{C8})$$

$$+ \alpha_2 \nabla_i \bar{\Delta}_2 \nabla_i \Delta_2 + \left( r_2 - \frac{1}{g} \frac{\epsilon}{i\delta} \right) \bar{\Delta}_2 \Delta_2 + \left( -i\epsilon r_1 + \delta \left\{ r_2 - \frac{1}{g} \frac{\epsilon}{i\delta} \right\} \right) (\bar{\Delta}_1 \Delta_2 + \bar{\Delta}_2 \Delta_1),$$

where it decouples to the standard Ginzburg-Landau model and the free field theory by taking  $\mu = 0$ . Taking the func-

tional derivative of the above action and dropping the terms of order  $\delta\epsilon$  while keeping the order  $\delta^2$ , we find the two equations of motion (C4) and (C5).