

STRONG BARRIERS FOR WEIGHTED QUASILINEAR EQUATIONS

TAKANOBU HARA

ABSTRACT. In potential theory, use of barriers is one of the most important techniques. We construct strong barriers for weighted quasilinear elliptic operators. There are two applications: (i) solvability of Poisson-type equations with boundary singular data, and (ii) a geometric version of Hardy inequality. Our construction method can be applied to a general class of divergence form elliptic operators on domains with rough boundary.

1. INTRODUCTION

Let $\Omega \subsetneq \mathbb{R}^n$ ($n \geq 1$) be an open set with nonempty boundary, and let $1 < p < \infty$. We consider elliptic differential equations of the type

$$(1.1) \quad \begin{cases} -\operatorname{div} \mathcal{A}(x, \nabla u(x)) = f(x) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where $\operatorname{div} \mathcal{A}(x, \nabla \cdot)$ is a weighted (p, w) -Laplacian type elliptic operator, w is a doubling weight on \mathbb{R}^n which admit a p -Poincaré inequality (see (2.1) and (2.2) for detail), and f is a locally bounded function on Ω . The most simple example of w is $w \equiv 1$, and the reason for considering weighted equations will be explained later. The precise assumptions on $\mathcal{A}: \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ are as follows: For each $z \in \mathbb{R}^n$, $\mathcal{A}(\cdot, z)$ is measurable, for each $x \in \Omega$, $\mathcal{A}(x, \cdot)$ is continuous, and there exists $1 \leq L < \infty$ such that

$$(1.2) \quad \mathcal{A}(x, z) \cdot z \geq w(x)|z|^p,$$

$$(1.3) \quad |\mathcal{A}(x, z)| \leq Lw(x)|z|^{p-1},$$

$$(1.4) \quad (\mathcal{A}(x, z_1) - \mathcal{A}(x, z_2)) \cdot (z_1 - z_2) > 0,$$

$$(1.5) \quad \mathcal{A}(x, tz) = t|t|^{p-2}\mathcal{A}(x, z)$$

for all $x \in \Omega$, $z, z_1, z_2 \in \mathbb{R}^n$, $z_1 \neq z_2$ and $t \in \mathbb{R}$. When $\mathcal{A}(x, z) = w(x)|z|^{p-2}z$, the operator $\operatorname{div} \mathcal{A}(x, \nabla u)$ is called as the (p, w) -Laplacian. In particular, if $\mathcal{A}(x, z) = z$, then the operator coincides with the classical Laplacian.

The aim of this paper is to provide an existence result of weak solutions to (1.1) for boundary singular data. The study of Eq. (1.1) has long history, and the standard approach to solve this problem is the variational method and its generalization (see, [34]). However, this method yields only solutions with finite energy. On the other hand, we can confirm existence of infinite energy solutions for

Date: December 23, 2022.

2020 Mathematics Subject Classification. 35J92; 35J25; 31C15; 31C45.

Key words and phrases. potential theory, Hardy inequality, p -Laplacian, quasilinear elliptic equation, boundary value problem, boundary regularity.

boundary singular data by considering Poisson's equation (in the classical sense) or ordinary differential equations. To obtain such solutions, we divide the problem into three steps: (i) find finite energy solutions to approximate problems, (ii) prove compactness of solutions, and (iii) derive a uniform bound for solutions. The part (i) is trivial in our setting, and we can find sufficient results for (ii) in prior work (e.g., [9, 23, 39]). This sequel from [20, 21] proposes a new perspective for (iii). More precisely, we construct supersolutions called strong barriers.

Supersolutions are an effective tool to control boundary behavior of solutions. For concrete problems, direct calculations for δ often yield sharp estimates (see, e.g., [38, 35, 10, 1, 15]). Unfortunately, this method can not be applied to general elliptic equations on domains with rough boundary. When conditions of the form (1.2)-(1.5) have to be considered, such as in applications to homogenization problems (e.g., [6]), it is needed to consider a construction method of supersolutions itself.

Ancona [4] defined strong barriers for linear elliptic operators and constructed them under capacity density type conditions. In addition, a Hardy inequality was proved as one application of them. Since other proofs by Lewis [33] and Wannebo [40], many authors have proved more general Hardy-type inequalities under capacity density conditions (see, §2). However, another application of strong barriers, to the Dirichlet problem of the type (1.1), seems not to be discussed sufficiently.

We construct strong barriers for quasilinear operators (Theorem 5.1) and apply the result to (1.1) (Theorem 6.3). Specifically, we make auxiliary functions by a boundary Hölder estimate in the De Giorgi-Nash-Moser theory and construct a global function by gluing them. These results can be regarded as extensions of [4, Remark 6.2] or [21, Corollary 4.4] as well as analogs of known facts for the p -Laplacian on C^2 domains (see, [17, Problem 6.6], [35, Theorem 1]). In addition, we prove a Hardy-type inequality (Corollary 5.3) by combining the results and the Picone inequality (see [2]). For connection with prior work on Hardy-type inequalities, we consider weighted operators borrowing a framework in [23] (see also [7, 28] for recent progress). Known results for Hardy-type inequalities that seem to be particularly relevant to this study will be discussed in §2. Throughout the paper, we assume only (1.2)-(1.5), (2.1)-(2.2) and (p, w) -capacity density conditions. The quantitative statements in results are new even for unweighted linear equations (compare with [4, Remark 5.2]). Our method seems to work for Cheeger differential equations on metric measure spaces (see, [11, 8], [7, Chapter B.2], [19]). On the other hand, there are large gaps in its direct application to the minimizers of variational problems. Another problem is that assumption in the results is clearly not optimal for interior regularity of solutions. These problems are left for future work.

Organization of the paper. In §2, we confirm of our problem and pick up related results on weighted Sobolev spaces. In §3, we define weak solutions to (1.1) and prove a Kato-type inequality. In §4, we states regularity estimates that will be used in §5. In §5, we construct strong barriers for (1.1). In §6, we apply the result in §5 to (1.1) and achieve our goals.

Notation. Let $\Omega \subsetneq \mathbb{R}^n$ be an open set.

- $\mathbf{1}_E(x)$:= the indicator function of a set E .
- $C_c^\infty(\Omega)$:= the set of all infinitely-differentiable functions with compact support in Ω .

- $L^p(\Omega; \mu) :=$ the L^p space with respect to a measure μ on Ω .
- $f_+ := \max\{f, 0\}$ and $f_- := -\min\{f, 0\}$.

For a closed set $\Gamma \subset \mathbb{R}^n$, we denote by δ_Γ the distance from Γ . If $\Gamma = \mathbb{R}^n \setminus \Omega$, we write $\delta_{\mathbb{R}^n \setminus \Omega}$ as δ simply. For a ball $B = B(x, R) = \{y: \text{dist}(x, y) < R\}$ and $\lambda > 0$, we denote $B(x, \lambda R)$ by λB . The letters c and C denote various constants with and without indices.

2. SETTING AND RELATED WORK

2.1. Admissible weights. Let $1 < p < \infty$ be a fixed number. A function $w \in L^1_{\text{loc}}(\mathbb{R}^n; dx)$ is called the *weight* if $w(x) > 0$ a.e. in \mathbb{R}^n . We write $w(E) = \int_E w dx$ for a Lebesgue measurable set $E \subset \mathbb{R}^n$. Throughout the below, we assume that w satisfying the doubling condition

$$(2.1) \quad w(2B) \leq C_D w(B)$$

and the p -Poincaré inequality

$$(2.2) \quad \int_B |u - u_B| dw \leq C_P \text{diam}(B) \left(\int_{\lambda B} |\nabla u|^p dw \right)^{1/p}, \quad \forall u \in C_c^\infty(\mathbb{R}^n),$$

where B is an arbitrary ball in \mathbb{R}^n , $f_B := w(B)^{-1} \int_B$, $u_B := \int_B u dw$ and C_D , C_P and $\lambda \geq 1$ are constants. A weight w satisfying (2.1) and (2.2) said to be *p -admissible*.

It is well-known that (2.1) and (2.2) yield the following Sobolev type inequality:

$$(2.3) \quad \left(\int_B |u|^{\chi p} dw \right)^{1/\chi p} \leq C \text{diam}(B) \left(\int_B |\nabla u|^p dw \right)^{1/p}, \quad \forall u \in C_c^\infty(B).$$

where C and $\chi > 1$ are constants depending only on p , C_D , C_P and λ . For detail, we refer to [23, Chapter 20] and the references cited therein.

Muckenhoupt A_p -weights are one typical example of p -admissible weights ([23, Chapter 15]). The power function $|x|^\mu$ is an A_p -weight on \mathbb{R}^n if and only if $-n < \mu < n(p-1)$. It seems to be known conventionally that if Ω is a bounded Lipschitz domain, then $w(x) = \delta(x)^\mu$ is a p -admissible weight on \mathbb{R}^n for $-1 < \mu < p-1$. Finer results can be found in [24, 25, 12, 13] and [28, Chapter 10]. Roughly speaking, if Γ is an s -dimensional set with $0 < s < n$, then $w(x) = \delta_\Gamma(x)^\mu$ is an A_p -weight on \mathbb{R}^n for $-(n-s) < \mu < (n-s)(p-1)$.

2.2. Sobolev spaces and capacities. The weighted Sobolev space $H^{1,p}(\Omega; w)$ is the closure of $C^\infty(\Omega)$ with respect to the norm

$$\|u\|_{H^{1,p}(\Omega; w)} := \left(\int_\Omega |u|^p + |\nabla u|^p dw \right)^{1/p},$$

where ∇u is the gradient of u in the sense of \mathbb{R}^n . The corresponding local space $H^{1,p}_{\text{loc}}(\Omega; w)$ is defined in the usual manner. We denote by $H^{1,p}_0(\Omega; w)$ the closure of $C_c^\infty(\Omega)$ in $H^{1,p}(\Omega; w)$. It is well-known that if $u, v \in H^{1,p}_0(\Omega; w)$, then $\min\{u, v\} \in H^{1,p}_0(\Omega; w)$.

Let $O \subset \mathbb{R}^n$ be open, and let $K \subset O$ be compact. The (p, w) -capacity $\text{cap}_{p,w}(K, O)$ of the condenser (K, O) is defined by

$$(2.4) \quad \text{cap}_{p,w}(K, O) := \inf \left\{ \|\nabla u\|_{L^p(O; w)}^p : u \geq 1 \text{ on } K, u \in C_c^\infty(O) \right\}.$$

For a boundary point $\xi \in \partial\Omega$ (more generally, for $\xi \in \mathbb{R}^n \setminus \Omega$), we consider the following (p, w) -capacity density condition: There exists $\gamma > 0$ such that

$$(2.5) \quad \frac{\text{cap}_{p,w}(\overline{B(\xi, R)} \setminus \Omega, B(\xi, 2R))}{\text{cap}_{p,w}(\overline{B(\xi, R)}, B(\xi, 2R))} \geq \gamma.$$

If (2.5) holds for all small $R > 0$, then ξ is a regular point of the corresponding Dirichlet problem (see Lemma 4.3 below, [23, Theorem 6.31] and the references cited therein). Sufficient conditions for (2.5) via exterior corkscrew-type conditions can be found in [23, Theorem 6.31] and [7, Corollary 11.25]. In particular, every boundary point of a ball satisfies (2.5) for all $R > 0$.

When (2.5) holds for all $\xi \in \mathbb{R}^n \setminus \Omega$ and $R > 0$, the set $\mathbb{R}^n \setminus \Omega$ is called as *uniformly (p, w) -fat*. This property is closely relates to Hardy-type inequalities. The study of Hardy-type inequalities under capacity density conditions started with Ancona's work [3, 4]. Lewis [33] developed theory of self-improvement property of uniformly $(p, 1)$ -fat sets and proved p -Hardy inequalities as a consequence. Wannebo [40] proved higher order Hardy-type inequalities using another approach. Mikkonen [36] studied nonlinear potential theory with p -admissible weights and proved the self-improvement property of (p, w) -fat sets using a new method. Björn, MacManus and Shanmugalingam [8] extended the results to the metric measure space setting. The equivalence between pointwise Hardy inequalities and uniformly fatness was established by several authors' contributions (see, [18, 29, 31, 30]). We refer to [28, Chapter 6] and [5, Chapter 3] for further related work. However, Ancona's proof in [4] is not very similar to any of the others.

The study of cases where (2.5) holds on a part of the boundary is more limited. The most similar known result to Corollary 5.3 below is [32, Proposition 5.4]. They are using Wannebo's idea. In [14], another form of Hardy inequality and its applications are discussed under assumptions on Sobolev extensibility of domains.

3. QUASILINEAR ELLIPTIC OPERATOR

Assume that $\mathcal{A}: \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ satisfies (1.2)-(1.5). For simplicity, we denote the following extended function $\overline{\mathcal{A}}$ by \mathcal{A} again:

$$\overline{\mathcal{A}}(x, z) = \begin{cases} \mathcal{A}(x, z) & x \in \Omega, \\ w(x)|z|^{p-2}z & \text{otherwise,} \end{cases} \quad z \in \mathbb{R}^n.$$

Let $f \in L^1_{\text{loc}}(\Omega)$. A function $u \in H^{1,p}_{\text{loc}}(\Omega; w)$ is called weak (super-, sub-)solution to $-\text{div } \mathcal{A}(x, \nabla u) = f$ in Ω if

$$(3.1) \quad \int_{\Omega} \mathcal{A}(x, \nabla u) \cdot \nabla \varphi \, dx = (\geq, \leq) \int_{\Omega} \varphi f \, dx$$

for any nonnegative $\varphi \in C_c^\infty(\Omega)$.

If u is a weak supersolution to $-\text{div } \mathcal{A}(x, \nabla u) = 0$ in Ω , then, by the Riesz representation theorem, there is a unique Radon measure $\nu[u]$ in Ω such that

$$\int_{\Omega} \mathcal{A}(x, \nabla u) \cdot \nabla \varphi \, dx = \int_{\Omega} \varphi \, d\nu[u]$$

for any $\varphi \in C_c^\infty(\Omega)$. The measure $\nu[u]$ is called the *Riesz measure* of u . We refer to [23, Chapter 21] for further detail.

By (1.2), if u is a solution to $-\operatorname{div} \mathcal{A}(x, \nabla u) = f$ in Ω , then its truncation $\min\{u, k\}$ is a supersolution to $-\operatorname{div} \mathcal{A}(x, \nabla u) = \min\{f, 0\}$ in Ω . The monotonicity condition (1.4) yields the following more general result.

Lemma 3.1. *Assume that u and v are weak supersolutions to $-\operatorname{div} \mathcal{A}(x, \nabla u) = 0$ in Ω . Then,*

$$(3.2) \quad \int_{\Omega} \varphi \, d\nu[\min\{u, v\}] \geq \int_{\Omega} \varphi \mathbf{1}_{\{u \leq v\}} \, d\nu[u] + \int_{\Omega} \varphi \mathbf{1}_{\{u > v\}} \, d\nu[v]$$

for any nonnegative $\varphi \in C_c^\infty(\Omega)$. Moreover, if u_1, \dots, u_k are weak supersolutions to $-\operatorname{div} \mathcal{A}(x, \nabla u) = 0$ in Ω , and if there is a Radon measure σ such that $\nu[u_k] \ll \sigma$ for all k , then

$$(3.3) \quad \nu[\min\{u_1, \dots, u_k\}] \geq \min\{f_1, \dots, f_k\} \sigma \quad \text{in } \Omega$$

in the sense of distributions, where f_k is the Radon-Nikodým derivative of $\nu[u_k]$ with respect to σ .

Proof. Note that $u, v \in H_{\text{loc}}^{1,p}(\Omega; w)$. For $\epsilon > 0$, consider the functions

$$\Phi_\epsilon(u - v) := \frac{\epsilon}{(u - v)_+ + \epsilon} \quad \text{and} \quad \Psi_\epsilon(u - v) := 1 - \Phi_\epsilon(u - v).$$

These functions are globally Lipschitz continuous with respect to $u - v$; therefore, $\Phi_\epsilon(u - v), \Psi_\epsilon(u - v) \in H_{\text{loc}}^{1,p}(\Omega; w)$. We also note that $\Phi_\epsilon(u - v)(x) \rightarrow \mathbf{1}_{\{u \leq v\}}(x)$ for all x . Fix $\varphi \in C_c^\infty(\Omega)$. Then,

$$\begin{aligned} & \int_{\Omega} \varphi \, d\nu[\min\{u, v\}] \\ &= \int_{\Omega} \varphi \Phi_\epsilon(u - v) \, d\nu[\min\{u, v\}] + \int_{\Omega} \varphi \Psi_\epsilon(u - v) \, d\nu[\min\{u, v\}]. \end{aligned}$$

By the definition of $\nu[\min\{u, v\}]$, we have

$$\begin{aligned} \int_{\Omega} \varphi \Phi_\epsilon(u - v) \, d\nu[\min\{u, v\}] &= \int_{[u \leq v]} \mathcal{A}(x, \nabla u) \cdot \nabla(\varphi \Phi_\epsilon(u - v)) \, dx \\ &\quad + \int_{[u > v]} \mathcal{A}(x, \nabla v) \cdot \nabla(\varphi \Phi_\epsilon(u - v)) \, dx. \end{aligned}$$

Note that $(\mathcal{A}(x, \nabla v) - \mathcal{A}(x, \nabla u)) \cdot \nabla \Phi_\epsilon(u - v) \geq 0$ by (1.4). Therefore,

$$\int_{[u > v]} \mathcal{A}(x, \nabla v) \cdot \nabla(\varphi \Phi_\epsilon(u - v)) \, dx + I_\epsilon \geq \int_{[u > v]} \mathcal{A}(x, \nabla u) \cdot \nabla(\varphi \Phi_\epsilon(u - v)) \, dx,$$

where

$$I_\epsilon := \int_{[u > v]} (\mathcal{A}(x, \nabla u) - \mathcal{A}(x, \nabla v)) \cdot \nabla \varphi \Phi_\epsilon(u - v) \, dx.$$

Adding the two inequalities, we get

$$(3.4) \quad \int_{\Omega} \varphi \Phi_\epsilon(u - v) \, d\nu[\min\{u, v\}] + I_\epsilon \geq \int_{\Omega} \varphi \Phi_\epsilon(u - v) \, d\nu[u].$$

Similarly, since

$$\begin{aligned} \int_{\Omega} \varphi \Psi_\epsilon(u - v) \, d\nu[\min\{u, v\}] &= \int_{[u \leq v]} \mathcal{A}(x, \nabla u) \cdot \nabla(\varphi \Psi_\epsilon(u - v)) \, dx \\ &\quad + \int_{[u > v]} \mathcal{A}(x, \nabla v) \cdot \nabla(\varphi \Psi_\epsilon(u - v)) \, dx \end{aligned}$$

and $(\mathcal{A}(x, \nabla u) - \mathcal{A}(x, \nabla v)) \cdot \nabla \Psi_\epsilon(u - v) \geq 0$, we have

$$(3.5) \quad \int_{\Omega} \varphi \Psi_\epsilon(u - v) d\nu[\min\{u, v\}] + II_\epsilon \geq \int_{\Omega} \varphi \Psi_\epsilon(u - v) d\nu[v],$$

where

$$II_\epsilon := \int_{[u \leq v]} (\mathcal{A}(x, \nabla v) - \mathcal{A}(x, \nabla u)) \cdot \nabla \varphi \Psi_\epsilon(u - v) dx.$$

Combining (3.4) and (3.5), we obtain

$$\int_{\Omega} \varphi d\nu[\min\{u, v\}] + I_\epsilon + II_\epsilon \geq \int_{\Omega} \varphi \Phi_\epsilon(u - v) d\nu[u] + \int_{\Omega} \varphi \Psi_\epsilon(u - v) d\nu[v].$$

Take the limit $\epsilon \rightarrow 0$. By the dominated convergence theorem, the right-hand side of this inequality goes to the right-hand of (3.2). By the same reason, $I_\epsilon, II_\epsilon \rightarrow 0$. Therefore, the former statement holds. The latter statement is a consequence of induction. \square

4. REGULARITY ESTIMATES

By standard techniques in the De Giorgi-Nash-Moser theory, the Sobolev inequality (2.3) yields the following global L^∞ estimate and weak Harnack inequality. The proofs are combinations of [27, p.63, Lemma B.2], [17, Theorem 8.18] and [23, Theorems 3.59 and 7.46]. See also [37, Chapter 3.1.0].

Lemma 4.1. *Let $u \in H^{1,p}(\Omega; w)$ be a weak subsolution to $-\operatorname{div} \mathcal{A}(x, \nabla u) = f$ in Ω . Let $F = (\operatorname{diam}(\Omega)^p \|f_+/w\|_{L^\infty(\Omega)})^{1/(p-1)}$. Then,*

$$\operatorname{ess\,sup}_{\Omega} u \leq \sup_{\partial\Omega} u + CF_+,$$

where $\sup_{\partial\Omega} u := \inf\{k \in \mathbb{R}^n : (u - k)_+ \in H_0^{1,p}(\Omega; w)\}$ and C is a constant depending only on p, C_D and $\{C_P, \lambda\}$. In particular, there is a constant $c_1 = c_1(p, C_D, \{C_P, \lambda\})$ such that if $\|f_+/w\|_{L^\infty(\Omega)} \leq c_1 \operatorname{diam}(\Omega)^p$, then $\operatorname{ess\,sup}_{\Omega} u \leq \sup_{\partial\Omega} u + 1/4$.

Lemma 4.2. *Let $u \in H^{1,p}(2B; w)$ be a nonnegative weak supersolution to $-\operatorname{div} \mathcal{A}(x, \nabla u) = f$ in $2B$, and let $F_- = (R^p \|f_-/w\|_{L^\infty(2B)})^{1/(p-1)}$. Then, (i) for each $0 < s < \chi(p-1)$, there exists a constant C depending only on $p, C_D, \{C_P, \lambda\}, L$ and s such that*

$$(4.1) \quad \left(\int_B u^s dw \right)^{1/s} \leq C \left(\operatorname{ess\,inf}_B u + F_- \right).$$

(ii) there exists a constant C depending only on $p, C_D, \{C_P, \lambda\}$ and L such that

$$(4.2) \quad R^{p-1} \int_B |\nabla u|^{p-1} dw \leq C \left(\operatorname{ess\,inf}_B u + F_- \right)^{p-1}.$$

It is well-known that Lemma 4.2 yields a local Hölder estimate (see, e.g., [17, Theorem 8.24]). Below, we always assume that f/w is locally bounded. Under this assumption, any weak solution u to $-\operatorname{div} \mathcal{A}(x, \nabla u) = f$ can be regarded as a locally Hölder continuous function.

We use the exterior condition (2.5) via the following boundary Hölder estimate. The proof below is the same as [17, pp. 206-209] (see also [16] and [37]).

Lemma 4.3. *Let B be a ball centered at $\xi \in \partial\Omega$ with radius R . Assume that (2.5) holds. Let $u \in H^{1,p}(\Omega; w) \cap L^\infty(\Omega)$ satisfy $-\operatorname{div} \mathcal{A}(x, \nabla u) = f$ in $\Omega \cap 4B$. Let $F_\pm = (R^p \|f_\pm/w\|_{L^\infty(2B)})^{1/(p-1)}$. Then,*

$$\operatorname{osc}_{\Omega \cap B} u \leq \left(1 - \frac{\gamma^{1/(p-1)}}{C}\right) \operatorname{osc}_{\Omega \cap 4B} u + \frac{\gamma^{1/(p-1)}}{C} \operatorname{osc}_{\partial\Omega \cap B} u + F_+ + F_-,$$

where C is a constant depending only on p , C_D , $\{C_P, \lambda\}$ and L and $\operatorname{osc} := \sup - \inf$.

Proof. We first assume that u is a nonnegative supersolution to $-\operatorname{div} \mathcal{A}(x, \nabla u) = f$ in $\Omega \cap 4B$. Let $m = \inf_{\partial\Omega \cap B} u$, and let

$$u_m^-(x) = \begin{cases} \min\{u(x), m\} & x \in \Omega, \\ m & \text{otherwise.} \end{cases}$$

Note that $0 \leq u_m^- \leq m$ and $u_m^- = m$ on $\overline{B} \setminus \Omega$. Take $\eta \in C_c^\infty(2B)$ such that $0 \leq \eta \leq 1$ in $2B$, $\eta = 1$ on B and $|\nabla \eta| \leq C/R$. Then, (2.5) gives

$$m^p \gamma \leq \frac{m^p \operatorname{cap}_{p,w}(\overline{B} \setminus \Omega, 2B)}{\operatorname{cap}_{p,w}(\overline{B}, 2B)} \leq CR^p \int_{2B} |\nabla(u_m^- \eta)|^p dw.$$

By the product rule, $\nabla(u_m^- \eta) = \nabla u_m^- \eta + u_m^- \nabla \eta$ a.e. Therefore,

$$(4.3) \quad \int_{2B} |\nabla(u_m^- \eta)|^p dw \leq C \left(R^{-p} \int_{2B} (u_m^-)^p d\mu + \int_{2B} |\nabla u_m^-|^p \eta^p dw \right).$$

By Lemma 4.2 (i), the former term on the right-hand side is estimated by

$$(4.4) \quad \int_{2B} (u_m^-)^p dw \leq m \int_{2B} (u_m^-)^{p-1} dw \leq Cm \left(\inf_B u_m^- + F_- \right)^{p-1}.$$

Consider the test function $(m - u_m^-) \eta^p$. Since u_m^- is a supersolution to $-\operatorname{div} \mathcal{A}(x, \nabla u) = \min\{f, 0\}$ in Ω , we have

$$\begin{aligned} & \int_{\Omega} \mathcal{A}(x, \nabla u_m^-) \cdot \nabla(m - u_m^-) \eta^p dx + p \int_{\Omega} \mathcal{A}(x, \nabla u_m^-) \cdot \nabla \eta \eta^{p-1} (m - u_m^-) dx \\ & \geq \int_{\Omega} (m - u_m^-) \eta^p \frac{\min\{f, 0\}}{w} dw. \end{aligned}$$

By (1.2) and (1.3), this inequality yields

$$\begin{aligned} \int_{2B} |\nabla u_m^-|^p \eta^p dw &= \int_{2B} |\nabla(m - u_m^-)|^p \eta^p dw \\ &\leq Cm \left(R^{-1} \int_{2B} |\nabla u_m^-|^{p-1} dw + F_-^{p-1} R^{-p} w(2B) \right). \end{aligned}$$

By Lemma 4.2 (ii), the former term on the right-hand side is estimated by

$$(4.5) \quad R^{-1} \int_{2B} |\nabla u_m^-|^{p-1} dw \leq CR^{-p} \left(\inf_B u_m^- + F_- \right)^{p-1}.$$

Combining (4.3), (4.4) and (4.5), we obtain

$$(4.6) \quad m \leq \frac{C}{\gamma^{1/(p-1)}} \left(\inf_B u_m^- + F_- \right).$$

Let $M(R) = \sup_{\Omega \cap B(\xi, R)} u$ and $m(R) = \inf_{\Omega \cap B(\xi, R)} u$. Applying (4.6) to $M(4R) - u$ and $u - m(4R)$, we obtain

$$(4.7) \quad \begin{aligned} M(4R) - \sup_{\partial\Omega \cap B} u &\leq \frac{C}{\gamma^{1/(p-1)}} (M(4R) - M(R) + F_+), \\ \inf_{\partial\Omega \cap B} u - m(4R) &\leq \frac{C}{\gamma^{1/(p-1)}} (m(R) - m(4R) + F_-). \end{aligned}$$

Adding the two inequalities, we arrive at the desired estimate. \square

Using (4.7) in Lemma 4.3 iteratively, we get the following lemma.

Lemma 4.4. *Let B be a ball centered at $\xi \in \partial\Omega$ with radius R_0 . Assume that (2.5) holds for all $0 < R \leq R_0$. Then, there are positive constants c_2 and $\theta \in (0, 1)$ depending only on p , C_D , $\{C_P, \lambda\}$, L and γ such that if $u \in H_0^{1,p}(\Omega; w)$ is a nonnegative bounded weak subsolution to $-\operatorname{div} \mathcal{A}(x, \nabla u) = f$ in $\Omega \cap B$, and if $\|f_+/w\|_{L^\infty(\Omega)} \leq c_2 R_0^{-p} (\sup_{\Omega \cap B} u)^{p-1}$, then*

$$\sup_{\Omega \cap \theta B} u \leq \frac{1}{4} \sup_{\Omega \cap B} u.$$

5. CONSTRUCTION OF STRONG BARRIERS

Theorem 5.1. *Let $\Gamma \subset \partial\Omega$ be a closed set, and assume that (2.5) holds for all $\xi \in \Gamma$ and $R > 0$. Then, there exists a nonnegative function $s_\Gamma \in H_{\text{loc}}^{1,p}(\Omega; w) \cap C(\Omega)$ satisfying*

$$(5.1) \quad -\operatorname{div} \mathcal{A}(x, \nabla s_\Gamma) \geq c_H \frac{s_\Gamma(x)^{p-1}}{\delta_\Gamma(x)^p} w(x) \quad \text{in } \Omega$$

and

$$(5.2) \quad \delta_\Gamma(x)^\alpha \leq s_\Gamma(x) \leq 30 \delta_\Gamma(x)^\alpha$$

for all $x \in \Omega$, where, c_H and $\alpha > 0$ are constants depending only on p , C_D , $\{C_P, \lambda\}$, L and γ .

Lemma 5.2. *Let B be a ball centered at $\xi \in \partial\Omega$ with radius R_0 . Assume that (2.5) holds for all $0 < R \leq R_0$. Let $c_3 := \min\{c_1, c_2\}$, where c_1 and c_2 are constants in Lemmas 4.1 and 4.4. Then, there exists a function $u_B \in H_{\text{loc}}^{1,p}(\Omega; w) \cap C(\Omega)$ satisfying*

$$(5.3) \quad -\operatorname{div} \mathcal{A}(x, \nabla u_B) = c_3 R_0^{-p} w \quad \text{in } \Omega \cap B,$$

$$(5.4) \quad \frac{1}{4} \leq u_B \leq \frac{5}{4} \quad \text{in } \Omega,$$

$$(5.5) \quad u_B = 1 \quad \text{on } \Omega \setminus B, \quad u_B \leq \frac{1}{2} \quad \text{on } \Omega \cap \theta B,$$

where θ is a constant in Lemma 4.4.

Proof. Existence of the function follows from the standard argument in [34, p.177]. Take $\eta_B \in C^\infty(\overline{\Omega})$ such that

$$\eta_B = \frac{1}{4} \text{ on } \overline{\Omega \cap B/2}, \quad \eta_B = 1 \text{ on } \overline{\Omega} \setminus B, \quad \frac{1}{4} \leq \eta_B \leq 1 \text{ in } \overline{\Omega}.$$

Consider the Dirichlet problem

$$\begin{cases} -\operatorname{div} A_B(x, \nabla v_B) = c_3 R_0^{-p} w & \text{in } \Omega \cap B, \\ v_B \in H_0^{1,p}(\Omega \cap B; w), \end{cases}$$

where

$$A_B(x, z) := \mathcal{A}(x, z + \nabla \eta_B(x)).$$

By (2.3), the right-hand side belongs to the dual of $H_0^{1,p}(\Omega \cap B; w)$. Therefore, by the Minty-Browder theorem, there exists a unique solution $u_B \in \eta_B + H_0^{1,p}(\Omega \cap B; w)$ to (5.3). By Lemma 4.2, u_B is continuous in $\Omega \cap B$. Meanwhile, if $x \in \Omega \cap \partial B$, then u_B is continuous at x by Lemma 4.3. Consequently, u_B is continuous in Ω . By the comparison principle, $u_B \geq 1/4$ in Ω . By Lemma 4.1, $u_B \leq \sup_{\partial\Omega} u + 1/4 \leq 5/4$ in Ω . The latter bound in (5.5) follows from Lemma 4.4. \square

Proof of Theorem 5.1. First, we construct a function s by the following two steps. (i) For $k \in \mathbb{Z}$, set $R_k = (\theta/2)^k$ and $E_k = \{x \in \Omega : \delta_\Gamma(x) \leq R_k\}$. Choosing $\{\xi_j\}_{j \in J_k} \subset \Gamma$, we construct a locally finite covering $\{B(\xi_j, \theta R_k)\}_{j \in J_k}$ of E_{k+1} . Note that

$$E_{k+1} \subset D_k := \Omega \cap \bigcup_{j \in J_k} B(\xi_j, R_k) \subset E_k.$$

Using Lemma 5.2, we define a function $v_k \in H_{\text{loc}}^{1,p}(\Omega; w) \cap C(\Omega)$ by

$$v_k = \inf_{j \in J_k} u_{B(\xi_j, R_k)}.$$

By (5.4) and (5.5), we have

$$(5.6) \quad \frac{1}{4} \leq v_k \leq \frac{5}{4} \text{ in } \Omega$$

and

$$(5.7) \quad v_k = 1 \text{ on } \Omega \setminus D_k, \quad v_k \leq \frac{1}{2} \text{ on } E_{k+1}.$$

Moreover, by Lemma 3.1, we have

$$(5.8) \quad -\operatorname{div} \mathcal{A}(x, \nabla v_k) \geq c_3 R_k^{-p} w \quad \text{in } D_k.$$

(ii) Define a function s on Ω by

$$s(x) = \inf_{k \in \mathbb{Z}} \left\{ \left(\frac{3}{4} \right)^k \tilde{v}_k(x) \right\}, \quad \text{where } \tilde{v}_k(x) = \begin{cases} v_k(x) & \text{if } x \in E_k, \\ +\infty & \text{otherwise.} \end{cases}$$

By (5.6) and the inequality $(3/4)^6 \leq 1/5 < (3/4)^5$,

$$\left(\frac{3}{4} \right)^{k-6} v_{k-6}(x) \geq \left(\frac{3}{4} \right)^{k-6} \frac{1}{4} \geq \left(\frac{3}{4} \right)^k \frac{5}{4} \geq \left(\frac{3}{4} \right)^k v_k(x)$$

for any $x \in E_k$. Therefore,

$$(5.9) \quad s(x) = \min \left\{ \left(\frac{3}{4} \right)^{k-5} v_{k-5}(x), \dots, \left(\frac{3}{4} \right)^k v_k(x) \right\}$$

for all $x \in E_k \setminus E_{k+1}$. In particular, $s \in H_{\text{loc}}^{1,p}(\Omega; w) \cap C(\Omega)$.

Next, we claim that

$$(5.10) \quad -\operatorname{div} \mathcal{A}(x, \nabla s) \geq \left(\frac{3}{4} \right)^{k(p-1)} c_3 R_{k-5}^{-p} w$$

in an open neighborhood of $E_k \setminus E_{k+1}$. By Lemma 3.1 and (5.8), this inequality holds in $D_k \setminus E_{k+1}$. Meanwhile, by (5.7), we have

$$v_{k-1} \leq \frac{1}{2} < \frac{3}{4}v_k \quad \text{in } E_k \setminus D_k.$$

By continuity of v_k and v_{k-1} , we can take an open set O_k such that $v_{k-1} < (3/4)v_k$ in O_k . Note that

$$s(x) = \min \left\{ \left(\frac{3}{4} \right)^{k-5} v_{k-5}(x), \dots, \left(\frac{3}{4} \right)^{k-1} v_{k-1}(x) \right\}$$

for all $x \in O_k$. Since v_{k-5}, \dots, v_{k-1} satisfy (5.8) in D_{k-1} , (5.10) holds in $O_k \cap D_{k-1}$ by the same reason as above. The open set $(D_k \setminus E_{k+1}) \cup (O_k \cap D_{k-1})$ has the desired property.

Finally, we consider pointwise behavior of s . It follows from (5.9) that

$$(5.11) \quad \frac{1}{4} \left(\frac{3}{4} \right)^k \leq s(x) \leq \frac{5}{4} \left(\frac{4}{3} \right)^5 \left(\frac{3}{4} \right)^k$$

for all $x \in E_k \setminus E_{k+1}$. By this inequality and the definition of R_k , the right-hand side of (5.10) is estimated from below by

$$\left(\frac{3}{4} \right)^{k(p-1)} c_3 R_{k-5}^{-p} \geq s^{p-1} \frac{c_H}{\delta_\Gamma(x)^p},$$

where $c_H = \left\{ (5/4) (4/3)^5 \right\}^{1-p} (\theta/2)^{6p} c_3$. Hence, s satisfies (5.1). Take $\alpha > 0$ such that $3/4 = (\theta/2)^\alpha$. Then, (5.11) gives

$$\frac{1}{4} \delta_\Gamma(x)^\alpha \leq s(x) \leq \frac{5}{4} \left(\frac{4}{3} \right)^6 \delta_\Gamma(x)^\alpha \leq \frac{30}{4} \delta_\Gamma(x)^\alpha$$

for all $x \in \Omega$. Thus, the function $s_\Gamma := 4s$ has the desired properties. \square

Corollary 5.3. *Assume that w is a p -admissible weight on \mathbb{R}^n . Let $\Gamma \subset \partial\Omega$ be a nonempty closed set, and assume that (2.5) holds for all $\xi \in \Gamma$ and $R > 0$. Let $c_H = c_H(p, C_D, \{C_P, \lambda\}, L = 1, \gamma) > 0$ be the constant in Theorem 5.1. Then,*

$$c_H \int_\Omega \frac{|\varphi|^p}{\delta_\Gamma^p} dw \leq \int_\Omega |\nabla \varphi|^p dw$$

for all $\varphi \in C_c^\infty(\Omega)$.

Proof. Applying Theorem 5.1 to $\mathcal{A}(x, z) = w(x)|z|^{p-2}z$, we get a nonnegative function $s_\Gamma \in H_{\text{loc}}^{1,p}(\Omega; w) \cap C(\Omega)$ satisfying

$$-\Delta_{p,w} s_\Gamma \geq c_H \frac{s_\Gamma(x)^{p-1}}{\delta_\Gamma(x)^p} w(x) \quad \text{in } \Omega.$$

Take $\varphi \in C_c^\infty(\Omega)$ such that $\varphi \geq 0$. Then,

$$c_H \int_\Omega \frac{\varphi^p}{\delta_\Gamma^p} dw = c_H \int_\Omega \frac{\varphi^p}{s_\Gamma^{p-1}} \frac{s_\Gamma^{p-1}}{\delta_\Gamma^p} dw \leq \int_\Omega \varphi^p \frac{d\nu[s_\Gamma]}{s_\Gamma^{p-1}},$$

where $\nu[s_\Gamma]$ is the Riesz measure of s_Γ with respect the (p, w) -Laplacian. Applying the Picone inequality ([2, Theorem 1.1]) to the right-hand side (for detail, see e.g. [22, Lemma 3.2]), we get

$$\int_{\Omega} \varphi^p \frac{d\nu[s_\Gamma]}{s_\Gamma^{p-1}} \leq \int_{\Omega} |\nabla \varphi|^p dw.$$

Combining the two inequalities, we obtain the desired inequality. \square

Remark 5.4. If $\mathbb{R}^n \setminus \Omega$ is uniformly (p, w) -fat, then

$$c_H \int_{\Omega} \frac{|\varphi|^p}{\delta^p} dw \leq \int_{\Omega} |\nabla \varphi|^p dw$$

for all $\varphi \in C_c^\infty(\Omega)$. This inequality was first proved in [36, Theorem 8.15] with an additional assumption on w . Later, Keith and Zhong [26] proved the self-improvement property of p -admissible weights, and the assumption was dropped. We note that our proof is not via the self-improvement property of (p, w) -fat sets and does not use of this result.

6. APPLICATIONS TO DIRICHLET PROBLEMS

Lemma 6.1. *Assume that there exists a function s_Γ on Ω satisfying (5.1) and (5.2). Let $h: (0, \infty) \rightarrow (0, \infty)$ be a continuously differentiable nondecreasing concave function such that*

$$g(s) := \int_0^s h(t) \frac{dt}{t} < \infty$$

for some $s > 0$. Then, $v(x) := g(s_\Gamma(x)) \in H_{\text{loc}}^{1,p}(\Omega; w) \cap C(\Omega)$ is a nonnegative weak supersolution to

$$-\operatorname{div} \mathcal{A}(x, \nabla v) \geq c_H \frac{h(\delta_\Gamma(x)^\alpha)^{p-1}}{\delta_\Gamma(x)^p} w(x) \quad \text{in } \Omega.$$

Moreover, $g(\delta_\Gamma(x)^\alpha) \leq v(x) \leq g(30 \delta_\Gamma(x)^\alpha)$ for all $x \in \Omega$.

Proof. For simplicity, we denote s_Γ by s . Let $' = \frac{d}{dt}$. By assumption, $h'(t)t \leq h(t)$ for all $t \in (0, \infty)$; therefore, g is increasing and concave on $(0, \infty)$. Fix a nonnegative function $\varphi \in C_c^\infty(\Omega)$. Since $g'' \leq 0$, we have

$$\int_{\Omega} \mathcal{A}(x, \nabla s) \cdot \nabla (g'(s)^{p-1}) \varphi dx \leq 0.$$

By the chain rule, (1.5) and (5.1), we get

$$\begin{aligned} \int_{\Omega} \mathcal{A}(x, \nabla g(s)) \cdot \nabla \varphi dx &= \int_{\Omega} \mathcal{A}(x, g'(s) \nabla s) \cdot \nabla \varphi dx \\ &\geq \int_{\Omega} \mathcal{A}(x, \nabla s) \cdot \nabla (g'(s)^{p-1} \varphi) dx \\ &\geq c_H \int_{\Omega} \frac{s^{p-1}}{\delta_\Gamma^p} (g'(s)^{p-1} \varphi) dw. \end{aligned}$$

Since h is nondecreasing, (5.2) gives

$$s^{p-1} g'(s)^{p-1} = h(s)^{p-1} \geq h(\delta_\Gamma^\alpha)^{p-1}.$$

The latter statement is a consequence of the monotonicity of g . \square

Proposition 6.2. *Let Ω be a bounded open set, and let $\Gamma \subset \partial\Omega$ be a nonempty closed set satisfying (2.5) for all $\xi \in \Gamma$ and $R > 0$. Let $f \in L^\infty_{\text{loc}}(\Omega)$, and assume that there exists a function h satisfying the assumption in Lemma 6.1 and*

$$|f(x)| \leq c_H \frac{h(\delta_\Gamma(x)^\alpha)^{p-1}}{\delta_\Gamma(x)^p} w(x)$$

for almost every $x \in \Omega$, where c_H and α are constants in Theorem 5.1. Then, there exists a nonnegative unique weak solution $u \in H_{\text{loc}}^{1,p}(\Omega; w) \cap C(\Omega)$ to (1.1) in the sense that (i) the zero extension of u belongs to $H_{\text{loc}}^{1,p}(\mathbb{R}^n \setminus \Gamma; w)$, and (ii) $\lim_{x \rightarrow \xi} u(x) = 0$ for all $\xi \in \Gamma$.

Proof. Using Theorem 5.1 and Lemma 6.1, we get a nonnegative supersolution $v(x) = g(s(x))$ to $-\text{div } \mathcal{A}(x, \nabla u) = |f|$ in Ω . Set $D_k = \{x \in \Omega : \text{dist}(x, \Gamma) > 1/k\}$, and consider the sequence of weak solutions $\{v_k\}_{k=1}^\infty \subset H_0^{1,p}(\Omega; w) \cap C(\Omega)$ to

$$\begin{cases} -\text{div } \mathcal{A}(x, \nabla v_k) = |f| \mathbf{1}_{D_k} & \text{in } \Omega, \\ v_k = 0 & \text{on } \partial\Omega. \end{cases}$$

Since Ω is bounded, the right-hand side belongs to the dual of $H_0^{1,p}(\Omega; w)$. By the comparison principle in [21, Theorem 3.5],

$$(6.1) \quad 0 \leq v_k(x) \leq v(x)$$

for all $x \in \Omega$. Also, consider the sequence of weak solutions $\{u_k\}_{k=1}^\infty \subset H_0^{1,p}(\Omega; w) \cap C(\Omega)$ to

$$\begin{cases} -\text{div } \mathcal{A}(x, \nabla u_k) = f \mathbf{1}_{D_k} & \text{in } \Omega, \\ u_k = 0 & \text{on } \partial\Omega. \end{cases}$$

By the comparison principle for weak solutions, we have

$$-v_k(x) \leq u_k(x) \leq v_k(x)$$

for all $x \in \Omega$; therefore,

$$(6.2) \quad |u_k(x)| \leq v(x)$$

for all $x \in \Omega$. Fix $j \geq 1$, and take a nonnegative function $\eta \in C^\infty(\mathbb{R}^n)$ such that $\eta \equiv 1$ on $\overline{D_j}$ and $\eta \equiv 0$ on $\Omega \setminus D_{j+1}$. Testing the equation of u_k with $u_k \eta^p$, we get

$$\int_\Omega \mathcal{A}(x, \nabla u_k) \cdot \nabla u_k \eta^p dx + p \int_\Omega \mathcal{A}(x, \nabla u_k) \cdot \nabla \eta \eta^{p-1} u_k dx = \int_\Omega f u_k \eta^p dx.$$

By (1.2) and (1.3), this inequality yields

$$(6.3) \quad \int_{D_j} |\nabla u_k|^p dw \leq C \int_{D_{j+1}} |u_k|^p dw + C \left\| \frac{f}{w} \right\|_{L^\infty(D_{j+1})}^{p/(p-1)} w(D_{j+1}).$$

By (6.2) and assumption on f , the right-hand side is bounded with respect to k . Meanwhile, Lemma 4.2 yields a local Hölder estimate of u_k . Therefore, for each $j \geq 1$, there are constants C_j and $\alpha_j \in (0, 1)$ independent of k such that

$$\|u_k\|_{C^{\alpha_j}(\overline{D_j})} + \|\nabla u_k\|_{L^p(D_j, w)} \leq C_j.$$

Take a subsequence of $\{u_k\}_{k=1}^\infty$ and a function u on Ω such that $u_k \rightarrow u$ locally uniformly in Ω and $\nabla u_k \rightharpoonup \nabla u$ weakly in $L^p(D_j; w)$ for all j . Let η be the function

defined before. By the product rule, we have

$$\int_{\Omega} |\nabla(u_k \eta)|^p dw \leq C \left(\int_{D_{j+1}} |\nabla u_k|^p \eta^p dw + \int_{D_{j+1}} |u_k|^p |\nabla \eta|^p dw \right).$$

The left-hand side belong to $H_0^{1,p}(\Omega; w)$, and thus, $u\eta \in H_0^{1,p}(\Omega; w)$. This implies that the zero extension of u belongs to $H_{\text{loc}}^{1,p}(\mathbb{R}^n \setminus \Gamma; w)$. Using the test function $u\eta - u_k\eta \in H_0^{1,p}(\Omega; w)$, we obtain

$$(6.4) \quad \int_{\Omega} \mathcal{A}(x, \nabla u_k) \cdot \nabla(u - u_k)\eta dx = - \int_{\Omega} \mathcal{A}(x, \nabla u_k) \cdot \nabla \eta (u - u_k) dx + \int_{\Omega} f(u - u_k)\eta dx.$$

Take the limit $k \rightarrow \infty$. The latter term on the right-hand side goes to zero. Moreover, by (1.3), the former term is estimated by

$$\begin{aligned} & \left| \int_{\Omega} \mathcal{A}(x, \nabla u_k) \cdot \nabla \eta (u - u_k) dx \right| \\ & \leq C \left(\int_{D_{j+1}} |\nabla u_j|^p dw \right)^{(p-1)/p} \left(\int_{D_{j+1}} |u - u_k|^p dw \right)^{1/p}. \end{aligned}$$

The right-hand side goes to zero because $u_k \rightarrow u$ uniformly in D_{j+1} . Thus, the left-hand side of (6.4) goes to zero. Meanwhile, by the weak convergence of ∇u_j in $L^p(D_{j+1}; w)$, we have

$$(6.5) \quad \int_{\Omega} \mathcal{A}(x, \nabla u) \cdot \nabla(u - u_k)\eta dx \rightarrow 0.$$

Combining (6.4), (6.5) and (1.4), we find that

$$\int_{D_j} \mathcal{A}(x, \nabla u) - \mathcal{A}(x, \nabla u_k) \cdot \nabla(u - u_k)\eta dx \rightarrow 0.$$

It follows from [23, Lemma 3.73] that u satisfies $-\text{div } \mathcal{A}(x, \nabla u) = f$ in Ω . Interior regularity of u follows from Lemma 4.2. If $\xi \in \Gamma$, then u is continuous at ξ by the upper bound (6.2).

Let $u, v \in H_{\text{loc}}^{1,p}(\Omega; w) \cap C(\Omega)$ be weak solutions to (1.1) satisfying the Dirichlet boundary condition in the statement. Assume that $u(x) > v(x)$ for some $x \in \Omega$. Then, $D = \{x \in \Omega: u(x) > v(x) + \epsilon\}$ is a nonempty bounded open set for some $\epsilon > 0$. If $\text{dist}(\overline{D}, \Gamma) = 0$, then there is a boundary point $\xi \in \overline{D} \cap \Gamma$. This contradicts to assumption because $(u-v)(\xi) = 0$ and $\inf_D (u-v) \geq \epsilon > 0$. Therefore, $\text{dist}(\overline{D}, \Gamma) > 0$ and $u, v \in H^{1,p}(D; w)$. By density,

$$\int_D (\mathcal{A}(x, \nabla u) - \mathcal{A}(x, \nabla v)) \cdot \nabla \varphi dx = 0$$

for all $\varphi \in H_0^{1,p}(D; w)$. Testing this equation with $\varphi = u - v - \epsilon$ and using (1.4), we find that $u = v$ on D . This contradicts to the assumption. \square

Theorem 6.3. *Assume that Ω is a bounded open set and that $\mathbb{R}^n \setminus \Omega$ is uniformly (p, w) -fat. Let $f \in L_{\text{loc}}^{\infty}(\Omega)$, and assume that there are constants $K > 0$ and $0 < \beta \leq \alpha$ such that*

$$|f(x)| \leq K \delta(x)^{\beta(p-1)-p} w(x)$$

for almost every $x \in \Omega$, where α is a positive number in Theorem 5.1. Then, there exists a unique weak solution $u \in H_{\text{loc}}^{1,p}(\Omega; w) \cap C(\bar{\Omega})$ to (1.1). Moreover,

$$(6.6) \quad |u(x)| \leq CK^{1/(p-1)}\delta(x)^\beta$$

for all $x \in \Omega$, where $C = c_H^{1-p}30^{\beta/\alpha}(\alpha/\beta)$.

Proof. Applying to Proposition 6.2 to $\Gamma = \partial\Omega$ and $h(t) = (K/c_H)^{1/(p-1)}t^{\beta/\alpha}$, we obtain the desired unique weak solution u . The upper bound (6.6) follows from (6.2). \square

ACKNOWLEDGMENTS

This work was supported by JST CREST (doi:10.13039/501100003382) Grant Number JPMJCR18K3 and JSPS KAKENHI (doi:10.13039/501100001691) Grant Number JP17H01092.

REFERENCES

- [1] H. Aikawa, T. Kilpeläinen, N. Shanmugalingam, and X. Zhong. Boundary Harnack principle for p -harmonic functions in smooth Euclidean domains. *Potential Anal.*, 26(3):281–301, 2007.
- [2] W. Allegretto and Y. X. Huang. A Picone’s identity for the p -Laplacian and applications. *Nonlinear Anal.*, 32(7):819–830, 1998.
- [3] A. Ancona. Une propriété des espaces de Sobolev. *C. R. Acad. Sci. Paris Sér. I Math.*, 292(9):477–480, 1981.
- [4] A. Ancona. On strong barriers and an inequality of Hardy for domains in \mathbf{R}^n . *J. London Math. Soc. (2)*, 34(2):274–290, 1986.
- [5] A. A. Balinsky, W. D. Evans, and R. T. Lewis. *The analysis and geometry of Hardy’s inequality*. Universitext. Springer, Cham, 2015.
- [6] A. Bensoussan, J.-L. Lions, and G. Papanicolaou. *Asymptotic analysis for periodic structures*, volume 5 of *Studies in Mathematics and its Applications*. North-Holland Publishing Co., Amsterdam-New York, 1978.
- [7] A. Björn and J. Björn. *Nonlinear potential theory on metric spaces*, volume 17 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2011.
- [8] J. Björn, P. MacManus, and N. Shanmugalingam. Fat sets and pointwise boundary estimates for p -harmonic functions in metric spaces. *J. Anal. Math.*, 85:339–369, 2001.
- [9] L. Boccardo and F. Murat. Almost everywhere convergence of the gradients of solutions to elliptic and parabolic equations. *Nonlinear Anal.*, 19(6):581–597, 1992.
- [10] M. Borsuk and V. Kondratiev. *Elliptic boundary value problems of second order in piecewise smooth domains*, volume 69 of *North-Holland Mathematical Library*. Elsevier Science B.V., Amsterdam, 2006.
- [11] J. Cheeger. Differentiability of Lipschitz functions on metric measure spaces. *Geom. Funct. Anal.*, 9(3):428–517, 1999.
- [12] R. G. Durán and F. López García. Solutions of the divergence and analysis of the Stokes equations in planar Hölder- α domains. *Math. Models Methods Appl. Sci.*, 20(1):95–120, 2010.
- [13] B. Dyda, L. Ihnatsyeva, J. Lehrbäck, H. Tuominen, and A. V. Vähäkangas. Muckenhoupt A_p -properties of distance functions and applications to Hardy-Sobolev-type inequalities. *Potential Anal.*, 50(1):83–105, 2019.
- [14] M. Egert, R. Haller-Dintelmann, and J. Rehberg. Hardy’s inequality for functions vanishing on a part of the boundary. *Potential Anal.*, 43(1):49–78, 2015.
- [15] S. Filippas, V. Maz’ya, and A. Tertikas. Critical Hardy-Sobolev inequalities. *J. Math. Pures Appl. (9)*, 87(1):37–56, 2007.
- [16] R. Gariepy and W. P. Ziemer. A regularity condition at the boundary for solutions of quasi-linear elliptic equations. *Arch. Rational Mech. Anal.*, 67(1):25–39, 1977.
- [17] D. Gilbarg and N. S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [18] P. Hajlasz. Pointwise Hardy inequalities. *Proc. Amer. Math. Soc.*, 127(2):417–423, 1999.

- [19] T. Hara. Wolff potential estimates for Cheeger p -harmonic functions. *Collect. Math.*, 69(3):407–426, 2018.
- [20] T. Hara. Quasilinear elliptic equations with sub-natural growth terms in bounded domains. *NoDEA Nonlinear Differential Equations Appl.*, 28(6):Paper No. 62, 2021.
- [21] T. Hara. Trace inequalities of the Sobolev type and nonlinear Dirichlet problems. *Calc. Var. Partial Differential Equations*, 61(6):Paper No. 216, 2022.
- [22] T. Hara and A. Seesanea. Existence of minimal solutions to quasilinear elliptic equations with several sub-natural growth terms. *Nonlinear Anal.*, 197:111847, 21, 2020.
- [23] J. Heinonen, T. Kilpeläinen, and O. Martio. *Nonlinear potential theory of degenerate elliptic equations*. Dover Publications, Inc., Mineola, NY, 2006. Unabridged republication of the 1993 original.
- [24] T. Horiuchi. The imbedding theorems for weighted Sobolev spaces. *J. Math. Kyoto Univ.*, 29(3):365–403, 1989.
- [25] T. Horiuchi. The imbedding theorems for weighted Sobolev spaces. II. *Bull. Fac. Sci. Ibaraki Univ. Ser. A*, (23):11–37, 1991.
- [26] S. Keith and X. Zhong. The Poincaré inequality is an open ended condition. *Ann. of Math. (2)*, 167(2):575–599, 2008.
- [27] D. Kinderlehrer and G. Stampacchia. *An introduction to variational inequalities and their applications*, volume 88 of *Pure and Applied Mathematics*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1980.
- [28] J. Kinnunen, J. Lehrbäck, and A. Vähäkangas. *Maximal function methods for Sobolev spaces*, volume 257 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2021.
- [29] J. Kinnunen and O. Martio. Hardy’s inequalities for Sobolev functions. *Math. Res. Lett.*, 4(4):489–500, 1997.
- [30] R. Korte, J. Lehrbäck, and H. Tuominen. The equivalence between pointwise Hardy inequalities and uniform fatness. *Math. Ann.*, 351(3):711–731, 2011.
- [31] J. Lehrbäck. Pointwise Hardy inequalities and uniformly fat sets. *Proc. Amer. Math. Soc.*, 136(6):2193–2200, 2008.
- [32] J. Lehrbäck, H. Tuominen, and A. V. Vähäkangas. Self-improvement of uniform fatness revisited. *Math. Ann.*, 368(3-4):1439–1464, 2017.
- [33] J. L. Lewis. Uniformly fat sets. *Trans. Amer. Math. Soc.*, 308(1):177–196, 1988.
- [34] J.-L. Lions. *Quelques méthodes de résolution des problèmes aux limites non linéaires*. Dunod; Gauthier-Villars, Paris, 1969.
- [35] M. Marcus and I. Shafir. An eigenvalue problem related to Hardy’s L^p inequality. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 29(3):581–604, 2000.
- [36] P. Mikkonen. On the Wolff potential and quasilinear elliptic equations involving measures. *Ann. Acad. Sci. Fenn. Math. Diss.*, (104):71, 1996.
- [37] E. W. Stredulinsky. *Weighted inequalities and degenerate elliptic partial differential equations*, volume 1074 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1984.
- [38] P. Tolksdorf. On the Dirichlet problem for quasilinear equations in domains with conical boundary points. *Comm. Partial Differential Equations*, 8(7):773–817, 1983.
- [39] N. S. Trudinger and X.-J. Wang. On the weak continuity of elliptic operators and applications to potential theory. *Amer. J. Math.*, 124(2):369–410, 2002.
- [40] A. Wannebo. Hardy inequalities. *Proc. Amer. Math. Soc.*, 109(1):85–95, 1990.

Email address: takanobu.hara.math@gmail.com

GRADUATE SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY, KITA 14, NISHI 9, KITA-KU, SAPPORO, HOKKAIDO, 060-0814, JAPAN