

INEQUALITIES FOR WEIGHTED SPACES WITH VARIABLE EXPONENTS

PABLO ROCHA

Abstract

In this article we obtain an "off-diagonal" version of the Fefferman-Stein vector-valued maximal inequality on weighted Lebesgue spaces with variable exponents. As an application of this result and the atomic decomposition developed in [12] we prove, for certain exponents $q(\cdot)$ in $\mathcal{P}^{\log}(\mathbb{R}^n)$ and certain weights ω , that the Riesz potential I_α , with $0 < \alpha < n$, can be extended to a bounded operator from $H_\omega^{p(\cdot)}(\mathbb{R}^n)$ into $L_\omega^{q(\cdot)}(\mathbb{R}^n)$, for $\frac{1}{p(\cdot)} := \frac{1}{q(\cdot)} + \frac{\alpha}{n}$.

1 Introduction

The theory of variable exponent Hardy spaces $H^{p(\cdot)}(\mathbb{R}^n)$ was developed independently by Nakai and Sawano in [15] and by Cruz-Urbe and Wang in [5]. Both theories prove equivalent definitions in terms of maximal operators using different approaches. In [15, 5], one of their main goals is the atomic decomposition of elements in $H^{p(\cdot)}$, as an application of the atomic decomposition they proved that singular integrals are bounded on $H^{p(\cdot)}$. Later in [18], the author jointly with Urciuolo proved the $H^{p(\cdot)} - L^{q(\cdot)}$ boundedness of certain generalized Riesz potentials and the $H^{p(\cdot)} - H^{q(\cdot)}$ boundedness of Riesz potential via the infinite atomic and molecular decomposition developed in [15]. In [17], the author gave another proof of the results obtained in [18], but by using the finite atomic decomposition developed in [5].

Recently, Kwok-Pun Ho in [12] developed the weighted theory for variable Hardy spaces on \mathbb{R}^n . He established the atomic decompositions for the weighted Hardy spaces with variable exponents $H_\omega^{p(\cdot)}(\mathbb{R}^n)$ and also revealed some intrinsic structures of atomic

2020 *Mathematics Subject Classification*. Primary 42B30; Secondary 42B25, 42B35, 46E30.

Key words and phrases: Fefferman-Stein inequalities, weighted variable Hardy spaces, atomic decomposition, Riesz potential.

decomposition for Hardy type spaces. His results generalize the infinite atomic decompositions obtained in [8, 1, 22, 15].

The main results of this work are contained in Theorems 4.1 and 5.1 below. The first result concerns to an "off-diagonal" version of the Fefferman-Stein vector-valued maximal inequality on weighted variable Lebesgue spaces (originally due to Harboure, Macías and Segovia [10]). Our second result states, for certain exponent functions $q(\cdot)$ and certain weights ω , the boundedness of Riesz potential I_α from $H_\omega^{p(\cdot)}$ into $L_\omega^{p(\cdot)}$ where $0 < \alpha < n$ and $\frac{1}{p(\cdot)} := \frac{1}{q(\cdot)} + \frac{\alpha}{n}$. This result is achieved via the off-diagonal Fefferman-Stein inequality (Theorem 4.1), the atomic decomposition established in [12], and the following additional assumption: for every cube $Q \subset \mathbb{R}^n$

$$(1) \quad \|\chi_Q\|_{L_\omega^{q(\cdot)}} \approx |Q|^{-\alpha/n} \|\chi_Q\|_{L_\omega^{p(\cdot)}}.$$

If $\omega \equiv 1$ and $q(\cdot)$ is log-Hölder continuous (locally and at infinite), then (1) holds. This case was proved in [18]. In this article we will give non trivial examples of power weights satisfying (1). So, (1) is an admissible hypothesis.

This paper is organized as follows. Section 2 gives the definition of weighted variable Lebesgue spaces $L_\omega^{p(\cdot)}$ and weighted variable Hardy spaces $H_\omega^{p(\cdot)}$ and some of their preliminary results. Section 3 presents some basics properties of the set of weights $\mathcal{W}_{p(\cdot)}$ used to define $H_\omega^{p(\cdot)}$ and some results about the size of the cubes in the $L_\omega^{p(\cdot)}$ -norm. The off-diagonal version of the Fefferman-Stein inequality on weighted variable Lebesgue spaces is established in Section 4. The $H_\omega^{p(\cdot)} - L_\omega^{q(\cdot)}$ boundedness of Riesz potential I_α is proved in Section 5. Finally, Section 6 gives non trivial examples of power weights satisfying (1).

Notation. The symbol $A \lesssim B$ stands for the inequality $A \leq cB$ for some constant c . The symbol $A \approx B$ stands for $B \lesssim A \lesssim B$. We denote by $Q(z, r)$ the cube centered at $z \in \mathbb{R}^n$ with side length r . Given a cube $Q = Q(z, r)$, we set $kQ = Q(z, kr)$ and $l(Q) = r$. For a measurable subset $E \subseteq \mathbb{R}^n$ we denote by $|E|$ and χ_E the Lebesgue measure of E and the characteristic function of E respectively. Given a real number $s \geq 0$, we write $[s]$ for the integer part of s . As usual we denote with $\mathcal{S}(\mathbb{R}^n)$ the space of smooth and rapidly decreasing functions and with $\mathcal{S}'(\mathbb{R}^n)$ the dual space. If β is the multiindex $\beta = (\beta_1, \dots, \beta_n)$, then $|\beta| = \beta_1 + \dots + \beta_n$.

2 Preliminaries

Let $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function. Given a measurable set E , let

$$p_-(E) = \operatorname{ess\,inf}_{x \in E} p(x), \quad \text{and} \quad p_+(E) = \operatorname{ess\,sup}_{x \in E} p(x).$$

When $E = \mathbb{R}^n$, we will simply write $p_- := p_-(\mathbb{R}^n)$ and $p_+ := p_+(\mathbb{R}^n)$.

Given a measurable function f on \mathbb{R}^n , define the modular ρ associated with $p(\cdot)$ by

$$\rho(f) = \int_{\mathbb{R}^n} |f(x)|^{p(x)} dx.$$

We define the variable Lebesgue space $L^{p(\cdot)} = L^{p(\cdot)}(\mathbb{R}^n)$ to be the set of all measurable functions f such that, for some $\lambda > 0$, $\rho(f/\lambda) < \infty$. This becomes a quasi normed space when equipped with the Luxemburg norm

$$\|f\|_{L^{p(\cdot)}} = \inf \{ \lambda > 0 : \rho(f/\lambda) \leq 1 \}.$$

Given a weight ω , i.e.: a locally integrable function on \mathbb{R}^n such that $0 < \omega(x) < \infty$ almost everywhere, we define the weighted variable Lebesgue space $L_\omega^{p(\cdot)}$ as the set of all measurable functions $f : \mathbb{R}^n \rightarrow \mathbb{C}$ such that $\|f\omega\|_{L^{p(\cdot)}} < \infty$. If $f \in L_\omega^{p(\cdot)}$, we define its "norm" by

$$\|f\|_{L_\omega^{p(\cdot)}} := \|f\omega\|_{L^{p(\cdot)}}.$$

LEMMA 2.1. *Given a measurable function $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ with $0 < p_- \leq p_+ < \infty$ and a weight ω , then for every $s > 0$,*

$$\|f\|_{L_\omega^{p(\cdot)}}^s = \| |f|^s \|_{L_\omega^{p(\cdot)/s}}.$$

PROOF. The condition $p_+ < \infty$ implies that $|\{x : p(x) = \infty\}| = 0$. Then, for every $s > 0$,

$$\begin{aligned} \| |f|^s \|_{L_\omega^{p(\cdot)/s}} &= \inf \left\{ \lambda > 0 : \int \left(\frac{|f(x)| \omega(x)}{\lambda^{1/s}} \right)^{p(x)} dx \leq 1 \right\} \\ &= \inf \left\{ \mu^s > 0 : \int \left(\frac{|f(x)| \omega(x)}{\mu} \right)^{p(x)} dx \leq 1 \right\} = \|f\|_{L_\omega^{p(\cdot)}}^s. \end{aligned}$$

□

For a measurable function $p(\cdot) : \mathbb{R}^n \rightarrow [1, \infty)$, its conjugate function $p'(\cdot)$ is defined by $\frac{1}{p(x)} + \frac{1}{p'(x)} = 1$. We have the following generalization of Hölder's inequality and an equivalent expression for the $L_\omega^{p(\cdot)}$ -norm.

LEMMA 2.2. (*Hölder's inequality*) *Let $p(\cdot) : \mathbb{R}^n \rightarrow [1, \infty)$ be a measurable function and ω be a locally integrable function such that $0 < \omega(x) < \infty$ almost everywhere. Then, there exists a constant $C > 0$ such that*

$$\int_{\mathbb{R}^n} |f(x)g(x)| dx \leq C \|f\|_{L_\omega^{p(\cdot)}} \|g\|_{L_\omega^{p'(\cdot)}}.$$

PROOF. The lemma follows from [6, Lemma 3.2.20].

□

PROPOSITION 2.3. Let $p(\cdot) : \mathbb{R}^n \rightarrow [1, \infty)$ be a measurable function and ω be a locally integrable function such that $0 < \omega(x) < \infty$ almost everywhere. Then

$$\|f\|_{L_\omega^{p(\cdot)}} \approx \sup \left\{ \int_{\mathbb{R}^n} |f(x)g(x)| dx : \|g\|_{L_{\omega^{-1}}^{p'(\cdot)}} \leq 1 \right\}.$$

PROOF. The proposition follows from [6, Corollary 3.2.14]. \square

Now, we briefly present the basics of weighted variable Hardy spaces theory and recall the atomic decomposition developed by K.-P. Ho in [12].

We introduce the weights used in [12] to define weighted Hardy spaces with variable exponents.

DEFINITION 2.4. Let $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function with $0 < p_- \leq p_+ < \infty$. We define $\mathcal{W}_{p(\cdot)}$ as the set of all weights ω such that

(i) there exists $0 < p_* < \min\{1, p_-\}$ such that $\|\chi_Q\|_{L_{\omega^{p_*}}^{p(\cdot)/p_*}} < \infty$, and

$$\|\chi_Q\|_{L_{\omega^{-p_*}}^{(p(\cdot)/p_*)'}} < \infty, \text{ for all cube } Q;$$

(ii) there exist $\kappa > 1$ and $s > 1$ such that Hardy-Littlewood maximal

$$\text{operator } M \text{ is bounded on } L_{\omega^{-\kappa/s}}^{(sp(\cdot))'/\kappa}.$$

REMARK 2.5. In [12, Definition 2.3], the author considers $p_* = \min\{1, p_-\}$. We observe that the whole theory is still valid if we take $0 < p_* < \min\{1, p_-\}$ instead of $p_* = \min\{1, p_-\}$. This slight modification allows us to compare the sets $\mathcal{W}_{q(\cdot)}$ and $\mathcal{W}_{p(\cdot)}$ related by $\frac{1}{p(\cdot)} - \frac{1}{q(\cdot)} = \frac{\alpha}{n}$ with $0 < \alpha < n$ (see Proposition 3.1 below).

For a measurable function $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ such that $0 < p_- \leq p_+ < \infty$ and $\omega \in \mathcal{W}_{p(\cdot)}$, in [12] the author give a variety of distinct approaches, based on differing definitions, all lead to the same notion of weighted variable Hardy space $H_\omega^{p(\cdot)}$.

We recall some terminologies and notations from the study of maximal functions. Given $N \in \mathbb{N}$, define

$$\mathcal{F}_N = \left\{ \varphi \in \mathcal{S}(\mathbb{R}^n) : \sum_{|\beta| \leq N} \sup_{x \in \mathbb{R}^n} (1 + |x|)^N |\partial^\beta \varphi(x)| \leq 1 \right\}.$$

For any $f \in \mathcal{S}'(\mathbb{R}^n)$, the grand maximal function of f is given by

$$\mathcal{M}f(x) = \sup_{t > 0} \sup_{\varphi \in \mathcal{F}_N} |(\varphi_t * f)(x)|,$$

where $\varphi_t(x) = t^{-n} \varphi(t^{-1}x)$.

DEFINITION 2.6. Let $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$, $0 < p_- \leq p_+ < \infty$, and $\omega \in \mathcal{W}_{p(\cdot)}$. The weighted variable Hardy space $H_\omega^{p(\cdot)}(\mathbb{R}^n)$ is the set of all $f \in \mathcal{S}'(\mathbb{R}^n)$ for which $\|\mathcal{M}f\|_{L_\omega^{p(\cdot)}} < \infty$. In this case we define $\|f\|_{H_\omega^{p(\cdot)}} := \|\mathcal{M}f\|_{L_\omega^{p(\cdot)}}$.

DEFINITION 2.7. Let $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$, $0 < p_- \leq p_+ < \infty$, $p_0 > 1$, and $\omega \in \mathcal{W}_{p(\cdot)}$. Fix an integer $d \geq 1$. A function $a(\cdot)$ on \mathbb{R}^n is called a $\omega - (p(\cdot), p_0, d)$ atom if there exists a cube Q such that

- $a_1)$ $\text{supp}(a) \subset Q$,
- $a_2)$ $\|a\|_{L^{p_0}} \leq \frac{|Q|^{\frac{1}{p_0}}}{\|\chi_Q\|_{L_\omega^{p(\cdot)}}}$,
- $a_3)$ $\int x^\beta a(x) dx = 0$ for all $|\beta| \leq d$.

Next, we introduce two indices, which are related to the intrinsic structure of the atomic decomposition of $H_\omega^{p(\cdot)}$. Given $\omega \in \mathcal{W}_{p(\cdot)}$, we write

$$s_{\omega, p(\cdot)} := \inf \left\{ s \geq 1 : M \text{ is bounded on } L_{\omega^{-1/s}}^{(sp(\cdot))'} \right\}$$

and

$$\mathbb{S}_{\omega, p(\cdot)} := \left\{ s \geq 1 : M \text{ is bounded on } L_{\omega^{-\kappa/s}}^{(sp(\cdot))'/\kappa} \text{ for some } \kappa > 1 \right\}.$$

Then, for every fixed $s \in \mathbb{S}_{\omega, p(\cdot)}$, we define

$$\kappa_{\omega, p(\cdot)}^s := \sup \left\{ \kappa > 1 : M \text{ is bounded on } L_{\omega^{-\kappa/s}}^{(sp(\cdot))'/\kappa} \right\}.$$

The index $\kappa_{\omega, p(\cdot)}^s$ is used to measure the left-openness of the boundedness of M on the family $\left\{ L_{\omega^{-\kappa/s}}^{(sp(\cdot))'/\kappa} \right\}_{\kappa > 1}$. The index $s_{\omega, p(\cdot)}$ is related to the vanishing moment condition and the index $\kappa_{\omega, p(\cdot)}^s$ is related to the size condition of the atoms (see [12, Theorems 5.3 and 6.3]).

The following three results are crucial in obtaining the $H_\omega^{p(\cdot)} - L_\omega^{q(\cdot)}$ boundedness of the Riesz potential. The first is a supporting result [12, Lemma 5.4], the second is a combination of [12, Theorem 6.2] and [16, Theorem 3.1], the last one was proved in [13, Proposition 2.1].

PROPOSITION 2.8. ([12, Lemma 5.4]) Let $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function with $0 < p_- \leq p_+ < \infty$ and $\omega \in \mathcal{W}_{p(\cdot)}$. Let $s \in \mathbb{S}_{\omega, p(\cdot)}$ and $\{\lambda_j\}_{j \in \mathbb{N}}$ be a sequence of scalars. If $r > (\kappa_{\omega, p(\cdot)}^s)'$, $\{b_j\}_{j \in \mathbb{N}}$ is a sequence in L^r such that, for every j , $\text{supp}(b_j)$ is contained in a cube Q_j and

$$\|b_j\|_{L^r} \leq A_j |Q_j|^{1/r},$$

where the A_j 's are positive scalars for all $j \in \mathbb{N}$, then

$$\left\| \sum_{j \in \mathbb{N}} \lambda_j b_j \right\|_{L_{\omega^{1/s}}^{sp(\cdot)}} \leq C \left\| \sum_{j \in \mathbb{N}} A_j |\lambda_j| \chi_{Q_j} \right\|_{L_{\omega^{1/s}}^{sp(\cdot)}}$$

for some $C > 0$ independent of $\{\lambda_j\}_{j \in \mathbb{N}}$, $\{b_j\}_{j \in \mathbb{N}}$ and $\{A_j\}_{j \in \mathbb{N}}$.

THEOREM 2.9. *Let $1 < p_0 < \infty$, $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function with $0 < p_- \leq p_+ < \infty$ and $\omega \in \mathcal{W}_{p(\cdot)}$. Then, for every $s > 1$ fixed, and every $f \in H_\omega^{p(\cdot)}(\mathbb{R}^n) \cap L^s(\mathbb{R}^n)$ there exist a sequence of scalars $\{\lambda_j\}$, a sequence of cubes $\{Q_j\}$ and $\omega - (p(\cdot), p_0, \lfloor ns_{\omega, p(\cdot)} - n \rfloor)$ atoms a_j supported on Q_j such that*

$$(2) \quad \left\| \sum_j \left(\frac{|\lambda_j|}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \right)^\theta \chi_{Q_j} \right\|_{L_\omega^{p(\cdot)/\theta}}^{1/\theta} \leq C \|f\|_{H_\omega^{p(\cdot)}}, \text{ for all } 0 < \theta < \infty,$$

and $f = \sum_j \lambda_j a_j$ converges in $L^s(\mathbb{R}^n)$.

PROOF. The existence of a such atomic decomposition as well as the validity of inequality in (2) are guaranteed by [12, Theorem 6.2]. The construction of a such atomic decomposition is the same that the given for the classical Hardy spaces (see [19, Chapter III]). So, following the proof in [16, Theorem 3.1], we obtain the convergence in $L^s(\mathbb{R}^n)$. \square

We define the set $\mathcal{S}_0(\mathbb{R}^n)$ by

$$\mathcal{S}_0(\mathbb{R}^n) = \left\{ \varphi \in \mathcal{S}(\mathbb{R}^n) : \int x^\beta \varphi(x) dx = 0, \text{ for all } \beta \in \mathbb{N}_0^n \right\}.$$

We say that an exponent function $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ such that $0 < p_- \leq p_+ < \infty$ belongs to $\mathcal{P}^{\log}(\mathbb{R}^n)$, if there exist two positive constants C and C_∞ such that $p(\cdot)$ satisfies the local log-Hölder continuity condition, i.e.:

$$|p(x) - p(y)| \leq \frac{C}{-\log(|x - y|)}, \quad |x - y| \leq \frac{1}{2},$$

and is log-Hölder continuous at infinity, i.e.:

$$|p(x) - p_\infty| \leq \frac{C_\infty}{\log(e + |x|)}, \quad x \in \mathbb{R}^n,$$

for some $p_- \leq p_\infty \leq p_+$.

THEOREM 2.10. *([13, Proposition 2.1]) Let $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $0 < p_- \leq p_+ < \infty$. If $\omega \in \mathcal{W}_{p(\cdot)}$, then $\mathcal{S}_0(\mathbb{R}^n) \subset H_\omega^{p(\cdot)}(\mathbb{R}^n)$ densely.*

3 Auxiliary Results

In this section we prove some basics properties of the set $\mathcal{W}_{p(\cdot)}$ and some results about the size of the cubes in the $L_\omega^{p(\cdot)}$ -norm.

PROPOSITION 3.1. *Let $0 < \alpha < n$ and let $q(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function such that $0 < q_- \leq q_+ < \infty$. If $\omega \in \mathcal{W}_{q(\cdot)}$ and $\frac{1}{p(\cdot)} := \frac{1}{q(\cdot)} + \frac{\alpha}{n}$, then $\omega \in \mathcal{W}_{p(\cdot)}$. Moreover, $s_{\omega, p(\cdot)} \leq s_{\omega, q(\cdot)} + \frac{\alpha}{n}$.*

PROOF. By Definition 2.4, $\omega \in \mathcal{W}_{q(\cdot)}$ if and only if

(i) there exists $0 < q_* < \min\{1, q_-\}$ such that $\|\chi_Q\|_{L_{\omega}^{q(\cdot)/q_*}} < \infty$, and

$$\|\chi_Q\|_{L_{\omega^{-q_*}}^{(q(\cdot)/q_*)'}} < \infty, \text{ for all cube } Q;$$

(ii) there exist $\kappa > 1$ and $s > 1$ such that Hardy-Littlewood maximal

operator M is bounded on $L_{\omega^{-\kappa/s}}^{(sq(\cdot))' / \kappa}$.

We define $\frac{1}{p_*} := \frac{1}{q_*} + \frac{\alpha}{n}$. Since $\frac{1}{p(\cdot)} - \frac{1}{q(\cdot)} = \frac{1}{p_*} - \frac{1}{q_*}$, it follows that $0 < p_* < \min\{1, p_-\}$ and $\frac{p_*}{q_*} \left(\frac{p(\cdot)}{p_*}\right)' = \left(\frac{q(\cdot)}{q_*}\right)'$. So, from Lemma 2.1 and (i) above, we have

$$\|\chi_Q\|_{L_{\omega^{-p_*}} \left(\frac{p(\cdot)}{p_*}\right)'} = \|\chi_Q\|_{L_{\omega^{-q_*}} \frac{p_*}{q_*} \left(\frac{p(\cdot)}{p_*}\right)'} = \|\chi_Q\|_{L_{\omega^{-q_*}} \left(\frac{q(\cdot)}{q_*}\right)'} < \infty.$$

Being $p(\cdot) \leq q(\cdot)$, by [4, Corollary 2.48], Lemma 2.1 and (i), we have

$$\begin{aligned} \|\chi_Q\|_{L_{\omega^{p_*}}^{q_*/p_*}} &= \|\chi_Q \omega^{p_*}\|_{L^{p(\cdot)/p_*}}^{q_*/p_*} \leq (1 + |Q|)^{q_*/p_*} \|\chi_Q \omega^{p_*}\|_{L^{q(\cdot)/p_*}}^{q_*/p_*} \\ &= (1 + |Q|)^{q_*/p_*} \|\chi_Q\|_{L_{\omega^{q_*}}^{q(\cdot)/q_*}} < \infty. \end{aligned}$$

From (ii) follows that the maximal operator M is bounded on $L_{\omega^{-1/s}}^{(sq(\cdot))'}$, with $s > 1$. We fix $r > s + \frac{\alpha}{n}$, and define $q_0 := \frac{r}{s}$ and $\frac{1}{p_0} := \frac{1}{q_0} + \frac{\alpha}{nr}$. It is clear that $\frac{r}{p_0} > 1$, $\frac{q_0}{p_0} > 1$ and $\frac{p_0}{q_0} \left(\frac{r}{p_0} p(\cdot)\right)' = \left(\frac{r}{q_0} q(\cdot)\right)' = (sq(\cdot))'$. Thus, for $\tilde{s} := \frac{r}{p_0}$ and $\tilde{\kappa} := \frac{q_0}{p_0}$, we obtain

$$\|Mf\|_{L_{\omega^{-\tilde{\kappa}/\tilde{s}}}^{(\tilde{s}p(\cdot))'/\tilde{\kappa}}} = \|Mf\|_{L_{\omega^{-1/s}}^{(sq(\cdot))'}} \leq C\|f\|_{L_{\omega^{-1/s}}^{(sq(\cdot))'}} = C\|f\|_{L_{\omega^{-\tilde{\kappa}/\tilde{s}}}^{(\tilde{s}p(\cdot))'/\tilde{\kappa}}}.$$

So, $\omega \in \mathcal{W}_{p(\cdot)}$ and $s_{\omega, p(\cdot)} \leq s_{\omega, q(\cdot)} + \frac{\alpha}{n}$. \square

The following necessary condition is due to Cruz-Uribe, Fiorenza and Neugebauer (see [3, Theorem 1.5]). It should be compared to the Muckenhoupt \mathcal{A}_p condition from the study of weighted norm inequalities (see [9, Chapter 7]).

LEMMA 3.2. *Given a weight ω and an exponent function $p(\cdot) : \mathbb{R}^n \rightarrow (1, \infty)$ such that the Hardy-Littlewood maximal operator is bounded on $L_w^{p(\cdot)}(\mathbb{R}^n)$, then, there exists a positive constant C such that for every cube $Q \subset \mathbb{R}^n$,*

$$(3) \quad \|\chi_Q\|_{L_w^{p(\cdot)}} \|\chi_Q\|_{L_{\omega^{-1}}^{p'(\cdot)}} \leq C|Q|.$$

DEFINITION 3.3. Given an exponent function $p(\cdot) : \mathbb{R}^n \rightarrow (1, \infty)$ and a weight ω , we write $\omega \in \mathcal{A}_{p(\cdot)}$ if ω satisfies (3).

LEMMA 3.4. Let $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function with $0 < p_- \leq p_+ < \infty$. If $\omega \in \mathcal{W}_{p(\cdot)}$, then, for every cube $Q \subset \mathbb{R}^n$,

$$\|\chi_{2Q}\|_{L_\omega^{p(\cdot)}} \approx \|\chi_Q\|_{L_\omega^{p(\cdot)}}.$$

PROOF. By the order preserving property of the norm $\|\cdot\|_{L_\omega^{p(\cdot)}}$, we have that

$$(4) \quad \|\chi_Q\|_{L_\omega^{p(\cdot)}} \leq \|\chi_{2Q}\|_{L_\omega^{p(\cdot)}}.$$

On the other hand, since $\omega \in \mathcal{W}_{p(\cdot)}$, the maximal operator is bounded on $L_{\omega^{-1/s}}^{(sp(\cdot))'}$. Then, by Lemma 3.2 (exchanging the roles of $p(\cdot)$ and $p'(\cdot)$), (4) above, and Hölder's inequality applied to $|Q| = \int \chi_Q(x) dx$ (see Lemma 2.2), we have that

$$\begin{aligned} \|\chi_{2Q}\|_{L_\omega^{p(\cdot)}}^{1/s} &= \|\chi_{2Q}\|_{L_{\omega^{1/s}}^{sp(\cdot)}} \leq C|Q| \|\chi_{2Q}\|_{L_{\omega^{-1/s}}^{(sp(\cdot))'}}^{-1} \leq C|Q| \|\chi_Q\|_{L_{\omega^{-1/s}}^{(sp(\cdot))'}}^{-1} \\ &\leq C \|\chi_Q\|_{L_{\omega^{1/s}}^{sp(\cdot)}} = C \|\chi_Q\|_{L_\omega^{p(\cdot)}}^{1/s}. \end{aligned}$$

This completes the proof. \square

A weight ω satisfies the *reverse Hölder inequality* with exponent $s > 1$, denoted by $\omega \in RH_s$, if there exists a constant $C > 0$ such that for every cube Q ,

$$\left(\frac{1}{|Q|} \int_Q [\omega(x)]^s dx \right)^{\frac{1}{s}} \leq C \frac{1}{|Q|} \int_Q \omega(x) dx;$$

the best possible constant is denoted by $[\omega]_{RH_s}$. We observe that if $\omega \in RH_s$, then by Hölder's inequality, $\omega \in RH_t$ for all $1 < t < s$, and $[\omega]_{RH_t} \leq [\omega]_{RH_s}$.

LEMMA 3.5. Let $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $0 < p_- \leq p_+ < \infty$, $\gamma \in \mathbb{R}$, and let $f \in L_{loc}^1(\mathbb{R}^n)$ be a function such that $|f(x)| \lesssim (1 + |x|)^\gamma$. Then $\|f\|_{L^{p(\cdot)}(\mathbb{R}^n)} \approx \|f\|_{L^{p_\infty}(\mathbb{R}^n)}$.

PROOF. We take $s > 1/p_-$. Then, by applying [7, Lemma 2.7] with $\omega \equiv 1$, we get $\|f\|_{L^{p(\cdot)}}^{1/s} = \| |f|^{1/s} \|_{L^{sp(\cdot)}} \approx \| |f|^{1/s} \|_{L^{sp_\infty}} = \|f\|_{L^{p_\infty}}^{1/s}$. \square

LEMMA 3.6. Let $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $1 < p_- \leq p_+ < \infty$ and $p_\infty = p_-$, and let ω be a weight such that $\omega(x) \lesssim (1 + |x|)^\gamma$ for some $\gamma \in \mathbb{R}$. If $\omega \in RH_{p_+}$, then for every cube Q we have

$$(5) \quad \|\chi_Q\|_{L_\omega^{p(\cdot)}} \approx \begin{cases} [\omega^{p_-}(Q)]^{1/p_-}, & \text{if } \ell(Q) > 1 \\ [\omega^{p_-(Q)}(Q)]^{1/p_-(Q)}, & \text{if } \ell(Q) \leq 1 \end{cases}.$$

PROOF. Applying Lemma 3.5, with $f = \chi_Q \omega$, we obtain

$$(6) \quad \|\chi_Q\|_{L_\omega^{p(\cdot)}} = \|\chi_Q \omega\|_{L^{p(\cdot)}} \approx [\omega^{p_-}(Q)]^{1/p_-},$$

for all cube Q .

From (6) and since $1 < p_- \leq p_-(Q)$, by Hölder's inequality, we have

$$\|\chi_Q\|_{L_\omega^{p(\cdot)}}^{p_-} \approx \int_Q [\omega(x)]^{p_-} dx \leq [\omega^{p_-(Q)}(Q)]^{\frac{p_-}{p_-(Q)}} \|\chi_Q\|_{\left(\frac{p_-(Q)}{p_-}\right)'} \leq [\omega^{p_-(Q)}(Q)]^{\frac{p_-}{p_-(Q)}},$$

if $\ell(Q) \leq 1$. So,

$$(7) \quad \|\chi_Q\|_{L_\omega^{p(\cdot)}} \lesssim [\omega^{p_-(Q)}(Q)]^{\frac{1}{p_-(Q)}},$$

holds for every cube Q with $\ell(Q) \leq 1$.

On the other hand, $\omega \in RH_{p_+}$ and $p_-(Q) \leq p_+$, then $\omega \in RH_{p_-(Q)}$ and

$$\left(\frac{1}{|Q|} \int_Q [\omega(x)]^{p_-(Q)} dx \right)^{1/p_-(Q)} \leq C|Q|^{-1} \int_Q \omega(x) dx \leq C|Q|^{-1} \|\chi_Q \omega\|_{L^{p(\cdot)}} \|\chi_Q\|_{L^{p'(Q)}},$$

[15, Lemma 2.2] gives $\|\chi_Q\|_{L^{p'(Q)}} \approx |Q|^{1/p'_+(Q)}$ if $\ell(Q) \leq 1$, being $p'_+(Q) = (p_-(Q))'$, we obtain

$$(8) \quad [\omega^{p_-(Q)}(Q)]^{1/p_-(Q)} \lesssim \|\chi_Q\|_{L_\omega^{p(\cdot)}}, \quad \text{if } \ell(Q) \leq 1.$$

Finally, (6), (7) and (8) give (5). \square

COROLLARY 3.7. *Let $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $0 < p_- \leq p_+ < \infty$ and $p_\infty = p_-$, and let $s > 1/p_-$. If $\omega(x) \lesssim (1 + |x|)^\gamma$ for some $\gamma \in \mathbb{R}$ and $\omega^{1/s} \in RH_{sp_+}$, then (5) holds.*

PROOF. Since $(sp(\cdot))_+ = sp_+$, $(sp(\cdot))_- = sp_-$ and $(sp(\cdot))_-(Q) = sp_-(Q)$, Lemma 3.6, applied to $sp(\cdot)$ and $\omega^{1/s}$, gives

$$\|\chi_Q\|_{L_\omega^{p(\cdot)}}^{1/s} = \|\chi_Q\|_{L_{\omega^{1/s}}^{sp(\cdot)}} \approx \begin{cases} [\omega^{p_-(Q)}]^{1/sp_-}, & \text{if } \ell(Q) > 1 \\ [\omega^{p_-(Q)}(Q)]^{1/sp_-(Q)}, & \text{if } \ell(Q) \leq 1 \end{cases}.$$

So, the lemma follows. \square

LEMMA 3.8. *Let $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $1 < p_- \leq p_+ < \infty$ and $p_\infty = p_+$. If $\omega \in RH_{p_+}$, then for every cube Q we have*

$$(9) \quad \|\chi_Q\|_{L_\omega^{p(\cdot)}} \approx \begin{cases} [\omega^{p_+}(Q)]^{1/p_+}, & \text{if } \ell(Q) > 1 \\ [\omega^{p_+(Q)}(Q)]^{1/p_+(Q)}, & \text{if } \ell(Q) \leq 1 \end{cases}.$$

PROOF. By hypothesis $p_\infty = p_+$, thus there exists $C_\infty > 0$ such that

$$0 \leq p_+ - p(x) \leq \frac{C_\infty}{\log(e + |x|)}, \text{ for all } x \in \mathbb{R}^n.$$

Let $\lambda = [\omega^{p_+}(Q)]^{1/p_+}$, by applying [3, Lemma 2.8] with $t = 2/p_-$, we have

$$\begin{aligned} \int_Q \left(\frac{\omega(x)}{\lambda} \right)^{p(x)} dx &\leq C_{2/p_-} \int_Q \left(\frac{\omega(x)}{\lambda} \right)^{p_+} dx + \int_Q \frac{dx}{(e + |x|)^{2n}} \\ &\leq C_{2/p_-} + \int_{\mathbb{R}^n} \frac{dx}{(e + |x|)^{2n}} =: M < \infty. \end{aligned}$$

So, from the definition of the "norm" $\|\cdot\|_{L_\omega^{p(\cdot)}}$ we have

$$(10) \quad \|\chi_Q\|_{L_\omega^{p(\cdot)}} \leq \max\{M^{1/p_-}, M^{1/p_+}\} [\omega^{p_+}(Q)]^{1/p_+}, \text{ for all cube } Q.$$

Now, if $\ell(Q) > 1$, the condition $\omega \in RH_{p_+}$, Hölder's inequality and [15, Lemma 2.2] give

$$\begin{aligned} \left(\frac{1}{|Q|} \int_Q [\omega(x)]^{p_+} dx \right)^{1/p_+} &\leq \frac{C}{|Q|} \int_Q \omega(x) dx \\ &\leq \frac{C}{|Q|} \|\chi_Q \omega\|_{L^{p(\cdot)}} \|\chi_Q\|_{L^{p'(\cdot)}} \leq \frac{C}{|Q|} \|\chi_Q \omega\|_{L^{p(\cdot)}} |Q|^{1-\frac{1}{p_+}}. \end{aligned}$$

Consequently,

$$(11) \quad [\omega^{p_+}(Q)]^{1/p_+} \leq C \|\chi_Q\|_{L_\omega^{p(\cdot)}}, \text{ if } \ell(Q) > 1.$$

Then, (10) and (11) give

$$(12) \quad \|\chi_Q\|_{L_\omega^{p(\cdot)}} \approx [\omega^{p_+}(Q)]^{1/p_+}, \text{ if } \ell(Q) > 1.$$

Now, we study the case $\ell(Q) \leq 1$. It is easy to check that

$$0 \leq p_+(Q) - p(x) \leq \frac{C}{\log(e + |x|)}, \text{ for all } x \in Q.$$

Since $1 < p_+(Q) \leq p_+$, we have that $\omega \in RH_{p_+(Q)}$. Then, by reasoning as above, we get

$$(13) \quad \|\chi_Q\|_{L_\omega^{p(\cdot)}} \approx [\omega^{p_+(Q)}(Q)]^{1/p_+(Q)}, \text{ if } \ell(Q) \leq 1.$$

Finally, (12) and (13) give (9). □

COROLLARY 3.9. *Let $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $0 < p_- \leq p_+ < \infty$ and $p_\infty = p_+$, and let $s > 1/p_-$. If $\omega^{1/s} \in RH_{sp_+}$, then (9) holds.*

PROOF. The proof follows from Lemmas 2.1 and 3.8. □

4 Off-diagonal Fefferman-Stein inequality

We apply extrapolation techniques to obtain an "off-diagonal" version of the Fefferman-Stein vector-valued maximal inequality on $L_\omega^{p(\cdot)}$. The following result generalizes Theorem 3.1 obtained in [12].

THEOREM 4.1. *Let $0 \leq \alpha < n$, $1 < u < \infty$ and let $q(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function with $0 < q_- \leq q_+ < \infty$. If $\omega \in \mathcal{W}_{q(\cdot)}$, then for $\frac{1}{p(\cdot)} := \frac{1}{q(\cdot)} + \frac{\alpha}{n}$ and any $r > s_{\omega, q(\cdot)} + \frac{\alpha}{n}$,*

$$(14) \quad \left\| \left(\sum_{j \in \mathbb{N}} (M_{\frac{\alpha}{r}} f_j)^u \right)^{1/u} \right\|_{L_{\omega^{1/r}}^{r q(\cdot)}} \lesssim \left\| \left(\sum_{j \in \mathbb{N}} |f_j|^u \right)^{1/u} \right\|_{L_{\omega^{1/r}}^{r p(\cdot)}},$$

holds for all sequences of bounded measurable functions with compact support $\{f_j\}_{j=1}^\infty$.

PROOF. Given $r > s_{\omega, q(\cdot)} + \frac{\alpha}{n}$, from the definition of $s_{\omega, q(\cdot)}$, we have $s > s_{\omega, q(\cdot)}$ such that $s + \frac{\alpha}{n} < r$ and M is bounded on $L_{\omega^{-1/s}}^{(sq(\cdot))'(\mathbb{R}^n)}$. Define

$$\mathcal{F} = \left\{ \left(\left(\sum_{j=1}^K (M_{\frac{\alpha}{r}} f_j)^u \right)^{1/u}, \left(\sum_{j=1}^K |f_j|^u \right)^{1/u} \right) : K \in \mathbb{N}, \{f_j\}_{j=1}^K \subset L_{comp}^\infty \right\},$$

where L_{comp}^∞ denotes the set of bounded functions with compact support.

Let $q_0 = \frac{r}{s}$ and let p_0 be defined by $\frac{1}{p_0} := \frac{1}{q_0} + \frac{\alpha}{nr}$. Since $r > s + \frac{\alpha}{n}$ we have that $1 < p_0 < \frac{nr}{\alpha}$. From [14, Theorem 3] follows that there exists an universal constant $C > 0$ such that for any $(F, G) \in \mathcal{F}$ and any $v \in \mathcal{A}_1$ (for the definition of the \mathcal{A}_1 class, the reader may refer to [9, Chapter 7])

$$(15) \quad \int [F(x)]^{q_0} v(x) dx \leq C \left(\int [G(x)]^{p_0} [v(x)]^{p_0/q_0} dx \right)^{q_0/p_0},$$

since $v \in \mathcal{A}_1$ implies that $v^{1/q_0} \in \mathcal{A}_{p_0, q_0}$ (for the definition of the $\mathcal{A}_{p, q}$ class, see [14, inequality (1.1)]). On the other hand, by Proposition 2.3, we have

$$(16) \quad \|F\|_{L_{\omega^{1/r}}^{r q(\cdot)}}^{q_0} = \|F^{q_0}\|_{L_{\omega^{1/s}}^{sq(\cdot)}} \leq C \sup \left\{ \int_{\mathbb{R}^n} |[F(x)]^{q_0} g(x)| dx : \|g\|_{L_{\omega^{-1/s}}^{(sq(\cdot))'}} \leq 1 \right\}$$

for some constant $C > 0$.

Let \mathcal{R} be the operator defined on $L_{\omega^{-1/s}}^{(sq(\cdot))'}$ by

$$\mathcal{R}g(x) = \sum_{k=0}^{\infty} \frac{M^k g(x)}{2^k \|M\|_{L_{\omega^{-1/s}}^{(sq(\cdot))'}}^k},$$

where, for $k \geq 1$, M^k denotes k iterations of the Hardy-Littlewood maximal operator M , $M^0 = M$, and $\|M\|_{L_{\omega^{-1/s}}^{(sq(\cdot))}'}$ is the operator norm of the maximal operator M on $L_{\omega^{-1/s}}^{(sq(\cdot))}'$. It follows immediately from this definition that:

- (i) if g is non-negative, $g(x) \leq \mathcal{R}g(x)$ a.e. $x \in \mathbb{R}^n$;
- (ii) $\|\mathcal{R}g\|_{L_{\omega^{-1/s}}^{(sq(\cdot))}' } \leq 2\|g\|_{L_{\omega^{-1/s}}^{(sq(\cdot))}'}$;
- (iii) $\mathcal{R}g \in \mathcal{A}_1$ with $[\mathcal{R}g]_{\mathcal{A}_1} \leq 2\|M\|_{L_{\omega^{-1/s}}^{(sq(\cdot))}'}$.

Since F is non-negative, we can take the supremum in (16) over those non-negative g only. For any fixed non-negative $g \in L_{\omega^{-1/s}}^{(sq(\cdot))}'$, by (i) above we have that

$$(17) \quad \int [F(x)]^{q_0} g(x) dx \leq \int [F(x)]^{q_0} (\mathcal{R}g)(x) dx.$$

Then (iii) and (15), and Hölder's inequality yield

$$(18) \quad \begin{aligned} \int [F(x)]^{q_0} (\mathcal{R}g)(x) dx &\leq C \left(\int [G(x)]^{p_0} [(\mathcal{R}g)(x)]^{p_0/q_0} dx \right)^{q_0/p_0} \\ &\leq C \|G^{p_0}\|_{L_{\omega^{p_0/r}}^{rp(\cdot)/p_0}}^{q_0/p_0} \|(\mathcal{R}g)^{p_0/q_0}\|_{L_{\omega^{-p_0/r}}^{(rp(\cdot)/p_0)'}}^{q_0/p_0} \\ &= C \|G\|_{L_{\omega^{1/r}}^{rp(\cdot)}}^{q_0} \|\mathcal{R}g\|_{L_{\omega^{-q_0/r}}^{q_0 \left(\frac{rp(\cdot)}{p_0}\right)'}} \end{aligned}$$

being $\frac{1}{rp(\cdot)} - \frac{1}{rq(\cdot)} = \frac{1}{p_0} - \frac{1}{q_0}$ and $q_0 = \frac{r}{s}$, we have $\frac{p_0}{q_0} \left(\frac{rp(\cdot)}{p_0}\right)' = \left(\frac{r}{q_0}q(\cdot)\right)' = (sq(\cdot))'$, so

$$= C \|G\|_{L_{\omega^{1/r}}^{rp(\cdot)}}^{q_0} \|\mathcal{R}g\|_{L_{\omega^{-1/s}}^{(sq(\cdot))}'}$$

now, (ii) gives

$$\leq C \|G\|_{L_{\omega^{1/r}}^{rp(\cdot)}}^{q_0} \|g\|_{L_{\omega^{-1/s}}^{(sq(\cdot))}'}$$

Thus, (17) and (18) lead to

$$(19) \quad \int [F(x)]^{q_0} g(x) dx \leq C \|G\|_{L_{\omega^{1/r}}^{rp(\cdot)}}^{q_0},$$

for all non-negative g with $\|g\|_{L_{\omega^{-1/s}}^{(sq(\cdot))}' } \leq 1$. Then, (16) and (19) give (14) for all finite sequences $\{f_j\}_{j=1}^K \subset L_{comp}^\infty$. Finally, by passing to the limit, we obtain (14) for all infinite sequences $\{f_j\}_{j=1}^\infty \subset L_{comp}^\infty$. \square

COROLLARY 4.2. *Let $0 < \alpha < n$, $q(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function with $0 < q_- \leq q_+ < \infty$ and $\omega \in \mathcal{W}_{q(\cdot)}$. If $\frac{1}{p(\cdot)} := \frac{1}{q(\cdot)} + \frac{\alpha}{n}$ and $\|\chi_Q\|_{L_\omega^{q(\cdot)}} \approx |Q|^{-\alpha/n} \|\chi_Q\|_{L_\omega^{p(\cdot)}}$ for every cube Q , then for any sequence of scalars $\{\lambda_j\}_{j \in \mathbb{N}}$, any family of cubes $\{Q_j\}_{j \in \mathbb{N}}$, and any $\theta \in (0, \infty)$ fixed we have*

$$\left\| \sum_{j \in \mathbb{N}} \left(\frac{|\lambda_j| \chi_{Q_j}}{\|\chi_{Q_j}\|_{L_\omega^{q(\cdot)}}} \right)^\theta \right\|_{L_{\omega^\theta}^{q(\cdot)/\theta}}^{1/\theta} \lesssim \left\| \sum_{j \in \mathbb{N}} \left(\frac{|\lambda_j| \chi_{Q_j}}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \right)^\theta \right\|_{L_{\omega^\theta}^{p(\cdot)/\theta}}^{1/\theta}.$$

PROOF. Since $\|\chi_Q\|_{L_\omega^{q(\cdot)}} \approx |Q|^{-\alpha/n} \|\chi_Q\|_{L_\omega^{p(\cdot)}}$ for every cube Q we obtain

$$\begin{aligned} \left\| \sum_j \left(\frac{|\lambda_j| \chi_{Q_j}}{\|\chi_{Q_j}\|_{L_\omega^{q(\cdot)}}} \right)^\theta \right\|_{L_\omega^{q(\cdot)/\theta}}^{1/\theta} &= \left\| \left\{ \sum_j \left(\frac{|\lambda_j| \chi_{Q_j}}{\|\chi_{Q_j}\|_{L_\omega^{q(\cdot)}}} \right)^\theta \right\}^{1/\theta} \right\|_{L_\omega^{q(\cdot)}} \\ &\lesssim \left\| \left\{ \sum_j \left(\frac{|\lambda_j| |Q_j|^{\alpha/n} \chi_{Q_j}}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \right)^\theta \right\}^{1/\theta} \right\|_{L_\omega^{q(\cdot)}}, \end{aligned}$$

it is easy to check that $|Q_j|^{\alpha/n} \chi_{Q_j}(x) \leq M_{\frac{\alpha\theta}{N}}(\chi_{Q_j})^{\frac{N}{\theta}}(x)$ for all j and all $N \in \mathbb{N}$, so

$$\begin{aligned} &\lesssim \left\| \left\{ \sum_j \left(\frac{|\lambda_j| M_{\frac{\alpha\theta}{N}}(\chi_{Q_j})^{\frac{N}{\theta}}}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \right)^\theta \right\}^{1/\theta} \right\|_{L_\omega^{q(\cdot)}} = \left\| \left\{ \sum_j \left(\frac{|\lambda_j|^\theta M_{\frac{\alpha\theta}{N}}(\chi_{Q_j})^N}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}^\theta} \right) \right\}^{1/\theta} \right\|_{L_\omega^{q(\cdot)}} \\ &= \left\| \left\{ \sum_j \left(\frac{|\lambda_j|^\theta M_{\frac{\alpha\theta}{N}}(\chi_{Q_j})^N}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}^\theta} \right) \right\}^{1/N} \right\|_{L_\omega^{Nq(\cdot)/\theta}}^{N/\theta}, \end{aligned}$$

taking N such that $N/\theta > s_{\omega, q(\cdot)} + \frac{\alpha}{n}$, by Theorem 4.1, we get

$$\lesssim \left\| \left\{ \sum_j \left(\frac{|\lambda_j|^\theta \chi_{Q_j}}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}^\theta} \right) \right\}^{1/N} \right\|_{L_\omega^{Np(\cdot)/\theta}}^{N/\theta} = \left\| \sum_j \left(\frac{|\lambda_j| \chi_{Q_j}}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \right)^\theta \right\|_{L_\omega^{p(\cdot)/\theta}}^{1/\theta}.$$

This completes the proof. \square

5 Weighted variable estimates for Riesz potential

Let $0 < \alpha < n$. The *Riesz potential* of order α is the fractional operator I_α defined by

$$(20) \quad I_\alpha f(x) = \int_{\mathbb{R}^n} f(y) |x - y|^{\alpha-n} dy, \quad x \in \mathbb{R}^n,$$

$f \in \mathcal{S}(\mathbb{R}^n)$. A well known result of Sobolev gives the boundedness of I_α from $L^p(\mathbb{R}^n)$ into $L^q(\mathbb{R}^n)$ for $1 < p < \frac{n}{\alpha}$ and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$. In [2] C. Capone, D. Cruz Uribe and A. Fiorenza extend this result to the case of Lebesgue spaces with variable exponents $L^{p(\cdot)}$. In [20] E. Stein and G. Weiss used the theory of harmonic functions of several variables to prove that these operators are bounded from $H^1(\mathbb{R}^n)$ into $L^{\frac{n}{n-\alpha}}(\mathbb{R}^n)$. In [21], M. Taibleson and G. Weiss obtained the boundedness of the Riesz potential I_α from the Hardy spaces $H^p(\mathbb{R}^n)$ into $H^q(\mathbb{R}^n)$, for $0 < p < 1$ and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$. These results were extended to the context of variable Hardy spaces by M. Urciuolo and the author in [18, 17].

In this section we will prove that the Riesz potential I_α is bounded from weighted variable Hardy spaces into weighted variable Lebesgue spaces. The main tools that we will use are Theorem 4.1, Corollary 4.2 and Theorem 2.9.

THEOREM 5.1. *Let $0 < \alpha < n$, $q(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $0 < q_- \leq q_+ < \infty$, and $\omega \in \mathcal{W}_{q(\cdot)}$. If $\frac{1}{p(\cdot)} := \frac{1}{q(\cdot)} + \frac{\alpha}{n}$ and $\|\chi_Q\|_{L_\omega^{q(\cdot)}} \approx |Q|^{-\alpha/n} \|\chi_Q\|_{L_\omega^{p(\cdot)}}$ for every cube Q , then the Riesz potential I_α given by (20) can be extended to a bounded operator $H_\omega^{p(\cdot)}(\mathbb{R}^n) \rightarrow L_\omega^{q(\cdot)}(\mathbb{R}^n)$.*

PROOF. Let $\omega \in \mathcal{W}_{q(\cdot)}$, by Definition 2.4, there exists $0 < \theta < 1$ such that $\frac{1}{\theta} \in \mathbb{S}_{\omega, q(\cdot)}$. Now, we take $q_0 > \frac{n}{n-\alpha}$ such that $q_0 > \theta \left(\kappa_{\omega, q(\cdot)}^{1/\theta} \right)'$, and define $\frac{1}{p_0} := \frac{1}{q_0} + \frac{\alpha}{n}$. By Proposition 3.1, for $\frac{1}{p(\cdot)} = \frac{1}{q(\cdot)} + \frac{\alpha}{n}$, we have that $\mathcal{W}_{q(\cdot)} \subset \mathcal{W}_{p(\cdot)}$ and $s_{\omega, p(\cdot)} \leq s_{\omega, q(\cdot)} + \frac{\alpha}{n}$. So, given $f \in S_0(\mathbb{R}^n)$, from Theorem 2.9 (taking $s = p_0$) and since one can always choose atoms with additional vanishing moment, we have that there exist a sequence of real numbers $\{\lambda_j\}_{j=1}^\infty$, a sequence of cubes $Q_j = Q(z_j, r_j)$ centered at z_j with side length r_j and $\omega - (p(\cdot), p_0, [ns_{\omega, q(\cdot)} + \alpha - n])$ atoms a_j supported on Q_j , satisfying

$$(21) \quad \left\| \sum_j \left(\frac{|\lambda_j|}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \right)^\theta \chi_{Q_j} \right\|_{L_\omega^{p(\cdot)/\theta}}^{1/\theta} \lesssim \|f\|_{H_\omega^{p(\cdot)}},$$

and $f = \sum_j \lambda_j a_j$ converges in $L^{p_0}(\mathbb{R}^n)$. By Sobolev's Theorem we have that I_α is bounded from $L^{p_0}(\mathbb{R}^n)$ into $L^{q_0}(\mathbb{R}^n)$, so

$$|I_\alpha f(x)| \leq \sum_j |\lambda_j| |I_\alpha a_j(x)|, \quad a.e. x \in \mathbb{R}^n.$$

Then

$$(22) \quad \|I_\alpha f\|_{L_\omega^{q(\cdot)}} \lesssim \left\| \sum_j |\lambda_j| \chi_{2Q_j} \cdot I_\alpha a_j \right\|_{L_\omega^{q(\cdot)}} + \left\| \sum_j |\lambda_j| \chi_{\mathbb{R}^n \setminus 2Q_j} \cdot I_\alpha a_j \right\|_{L_\omega^{q(\cdot)}} = J_1 + J_2,$$

where $2Q_j = Q(z_j, 2r_j)$. To estimate J_1 , we again apply Sobolev's Theorem and obtain

$$\|(I_\alpha a_j)^\theta\|_{L_\omega^{q_0/\theta}(2Q_j)} = \|I_\alpha a_j\|_{L_\omega^{q_0}(2Q_j)}^\theta \lesssim \|a_j\|_{L^{p_0}}^\theta \lesssim \frac{|Q_j|^{p_0/\theta}}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)/\theta}}^\theta} \lesssim \frac{|2Q_j|^{q_0/\theta}}{\|\chi_{2Q_j}\|_{L_\omega^{q(\cdot)/\theta}}^\theta},$$

where θ , q_0 and p_0 are given above. The last inequality follows from the condition $\|\chi_Q\|_{L_\omega^{q(\cdot)}} \approx |Q|^{-\alpha/n} \|\chi_Q\|_{L_\omega^{p(\cdot)}}$ assumed for every cube Q and Lemma 3.4 applied to the exponent $q(\cdot)$. Since $\frac{1}{\theta} \in \mathbb{S}_{\omega, q(\cdot)}$ and $q_0 > \theta \left(\kappa_{\omega, q(\cdot)}^{1/\theta} \right)'$, we apply the θ -inequality and Proposition 2.8 with $b_j = (\chi_{2Q_j} \cdot I_\alpha(a_j))^\theta$ and $A_j = \|\chi_{2Q_j}\|_{L_\omega^{q(\cdot)/\theta}}^{-1}$ to obtain

$$J_1 \lesssim \left\| \sum_j (|\lambda_j| \chi_{2Q_j} \cdot I_\alpha a_j)^\theta \right\|_{L_\omega^{q(\cdot)/\theta}}^{1/\theta} \lesssim \left\| \sum_j \left(\frac{|\lambda_j|}{\|\chi_{2Q_j}\|_{L_\omega^{q(\cdot)/\theta}}} \right)^\theta \chi_{2Q_j} \right\|_{L_\omega^{q(\cdot)/\theta}}^{1/\theta}.$$

Being $\chi_{2Q_j} \leq M(\chi_{Q_j})^2$, by Theorem 4.1, we have

$$J_1 \lesssim \left\| \left\{ \sum_j \left(\frac{|\lambda_j|^{\theta/2}}{\|\chi_{2Q_j}\|_{L_\omega^{q(\cdot)}}^{\theta/2}} M(\chi_{Q_j}) \right)^2 \right\}^{1/2} \right\|_{L_\omega^{2q(\cdot)/\theta}}^{2/\theta} \lesssim \left\| \sum_j \left(\frac{|\lambda_j|}{\|\chi_{Q_j}\|_{L_\omega^{q(\cdot)}}} \right)^\theta \chi_{Q_j} \right\|_{L_\omega^{q(\cdot)/\theta}}^{1/\theta}.$$

Corollary 4.2 and (21) give

$$(23) \quad J_1 \lesssim \left\| \sum_j \left(\frac{|\lambda_j|}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \right)^\theta \chi_{Q_j} \right\|_{L_\omega^{p(\cdot)/\theta}}^{1/\theta} \lesssim \|f\|_{H_\omega^{p(\cdot)}}.$$

Now, we estimate J_2 . Let $d = \lfloor ns_{\omega, q(\cdot)} + \alpha - n \rfloor$, and let $a_j(\cdot)$ be a $\omega - (p(\cdot), p_0, d)$ atom supported on the cube $Q_j = Q(z_j, r_j)$. In view of the moment condition of $a_j(\cdot)$ we obtain

$$I_\alpha a_j(x) = \int_{Q_j} (|x - y|^{\alpha-n} - q_d(x, y)) a(y) dy, \quad \text{for all } x \notin 2Q_j,$$

where q_d is the degree d Taylor polynomial of the function $y \rightarrow |x - y|^{\alpha-n}$ expanded around z_j . By the standard estimate of the remainder term of the Taylor expansion, there exists ξ between y and z_j such that

$$||x - y|^{\alpha-n} - q_d(x, y)| \leq C|y - z_j|^{d+1}|x - \xi|^{-n+\alpha-d-1},$$

for any $y \in Q_j$ and any $x \notin 2Q_j$, since $|x - \xi| \geq \frac{|x - z_j|}{1 + \sqrt{n}}$, we get

$$||x - y|^{\alpha-n} - q_d(x, y)| \leq Cr^{d+1}|x - z_j|^{-n+\alpha-d-1},$$

this inequality and the condition (a2) allow us to conclude that

$$(24) \quad \begin{aligned} |I_\alpha a_j(x)| &\lesssim \|a_j\|_1 r^{d+1} |x - z_j|^{-n+\alpha-d-1} \\ &\lesssim |Q_j|^{1-\frac{1}{p_0}} \|a_j\|_{p_0} r^{d+1} |x - z_j|^{-n+\alpha-d-1} \lesssim \frac{r^{n+d+1}}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} |x - z_j|^{-n+\alpha-d-1} \\ &\lesssim \frac{\left[M_{\frac{\alpha n}{n+d+1}}(\chi_{Q_j})(x) \right]^{\frac{n+d+1}{n}}}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}}, \quad \text{for all } x \notin 2Q_j. \end{aligned}$$

We put $r = \frac{n+d+1}{n}$, thus (24) leads to

$$J_2 \lesssim \left\| \left\{ \sum_j \frac{|\lambda_j|}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \left[M_{\frac{\alpha}{r}}(\chi_{Q_j}) \right]^r \right\}^{1/r} \right\|_{L_\omega^{rq(\cdot)/r}}^r.$$

Since

$$r = \frac{n + \lfloor n s_{\omega, q(\cdot)} + \alpha - n \rfloor + 1}{n} > s_{\omega, q(\cdot)} + \frac{\alpha}{n},$$

to apply Theorem 4.1, with $u = r$, we obtain

$$J_2 \lesssim \left\| \left\{ \sum_j \frac{|\lambda_j|}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \chi_{Q_j} \right\}^{1/r} \right\|_{L_{\omega^{1/r}}^{rp(\cdot)}}^r = \left\| \sum_j \frac{|\lambda_j|}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \chi_{Q_j} \right\|_{L_\omega^{p(\cdot)}}.$$

Being $0 < \theta < 1$, the θ -inequality and (21) give

$$(25) \quad J_2 \lesssim \left\| \sum_j \left(\frac{|\lambda_j|}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \right)^\theta \chi_{Q_j} \right\|_{L_{\omega^\theta}^{p(\cdot)/\theta}}^{1/\theta} \lesssim \|f\|_{H_\omega^{p(\cdot)}}.$$

Hence, (22), (23) and (25) yield

$$\|I_\alpha f\|_{L_\omega^{q(\cdot)}} \lesssim \|f\|_{H_\omega^{p(\cdot)}}, \text{ for all } f \in \mathcal{S}_0(\mathbb{R}^n).$$

Finally, by taking into account that $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$, $0 < p_- \leq p_+ < \infty$ and $\omega \in \mathcal{W}_{p(\cdot)}$, the theorem follows from Theorem 2.10. \square

6 Example: power weights

For $\gamma > -n$, let $w_\gamma(x) = |x|^\gamma$ with $x \in \mathbb{R}^n \setminus \{0\}$. From the estimates obtained in [9, Example 7.1.6, pp. 505], we observe that

$$(26) \quad \omega_\gamma \in \bigcap_{t>1} RH_t, \text{ if } \gamma \geq 0$$

and

$$(27) \quad \omega_\gamma \in \bigcap_{1 < t < n/|\gamma|} RH_t, \text{ if } -n < \gamma < 0.$$

Moreover, for $0 \leq \alpha < n$, $0 < p < \frac{n}{\alpha}$, $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ and each cube Q we have that

$$(28) \quad [\omega_\gamma^p(Q)]^{-1/p} [\omega_\gamma^q(Q)]^{1/q} = [\omega_{\gamma p}(Q)]^{-1/p} [\omega_{\gamma q}(Q)]^{1/q} \approx |Q|^{-\alpha/n}, \text{ for every } \gamma \geq 0$$

and

$$[\omega_\gamma^p(Q)]^{-1/p} [\omega_\gamma^q(Q)]^{1/q} \approx |Q|^{-\alpha/n}, \text{ for every } -n < \gamma < 0 \text{ with } q < \frac{n}{|\gamma|}.$$

Let $q(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $0 < q_\infty = q_- \leq q_+ < \infty$, and let $p(\cdot)$ be defined by $\frac{1}{p(\cdot)} := \frac{1}{q(\cdot)} + \frac{\alpha}{n}$. If $\gamma \geq 0$, then, by (28) and Corollary 3.7 (first applied to $q(\cdot)$ and then to $p(\cdot)$), we get

$$\|\chi_Q\|_{L_{\omega_\gamma}^{q(\cdot)}} \approx |Q|^{-\alpha/n} \|\chi_Q\|_{L_{\omega_\gamma}^{p(\cdot)}}.$$

It is clear that $q'(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$, $0 < q'_- \leq q'_+ = q'_\infty < \infty$, $q'_+ = (q_-)'$ and $q'_+(Q) = (q_-(Q))'$. Then for every $0 \leq \gamma < \frac{n}{q'_+}$, by (26) and Corollary 3.7; and (27) and Corollary 3.9 (applied to ω_γ and $q(\cdot)$; and $\omega_{-\gamma}$ and $q'(\cdot)$ respectively), we obtain that

$$\|\chi_Q\|_{L_{\omega_\gamma}^{q(\cdot)}} \|\chi_Q\|_{L_{\omega_{-\gamma}}^{(q(\cdot))'}} \approx \left\{ \begin{array}{ll} [\omega_\gamma^{q_-(Q)}]^{1/q_-} [\omega_{-\gamma}^{q'_+(Q)}]^{1/q'_+}, & \text{if } \ell(Q) > 1 \\ [\omega_\gamma^{q_-(Q)}(Q)]^{1/q_-(Q)} [\omega_{-\gamma}^{q'_+(Q)}(Q)]^{1/q'_+(Q)}, & \text{if } \ell(Q) \leq 1 \end{array} \right\} \approx |Q|,$$

for all cube $Q \subset \mathbb{R}^n$, where the last estimate follows from [9, Example 7.1.6, pp. 505]. In particular, for $s > 1/q_-$ and $0 \leq \gamma < \frac{sn}{(sq(\cdot))'_+}$, we have

$$\|\chi_Q\|_{L_{\omega_{\gamma/s}}^{sq(\cdot)}} \|\chi_Q\|_{L_{\omega_{-\gamma/s}}^{(sq(\cdot))'}} \approx |Q|.$$

Since $\omega_{\pm\gamma}^{1/s} = \omega_{\pm\gamma/s}$, by [3, Theorem 1.5] we have that the maximal operator M is bounded on $L_{\omega_\gamma^{-1/s}}^{(sq(\cdot))'}$. By the left-openness property of the set $\mathcal{A}_{(sq(\cdot))'}$ (see definition 3.3 and [11, Theorem 3.3]), it follows that there exists a constant $\kappa > 1$ such that the maximal operator M is bounded on $L_{\omega_\gamma^{-\kappa/s}}^{(sq(\cdot))'/\kappa}$. So, $\omega_\gamma \in \mathcal{W}_{q(\cdot)}$ and $\|\chi_Q\|_{L_{\omega_\gamma}^{q(\cdot)}} \approx |Q|^{-\alpha/n} \|\chi_Q\|_{L_{\omega_\gamma}^{p(\cdot)}}$ for every cube $Q \subset \mathbb{R}^n$.

References

- [1] H. Q. BUI, Weighted Hardy spaces, Math. Nachr. 103 (1981), 45-62.
- [2] C. CAPONE, D. CRUZ-URIBE and A. FIORENZA, The fractional maximal operator and fractional integral on variable L^p spaces, Rev. Mat. Iberoam. 23 (3) (2007), 743-770.
- [3] D. CRUZ-URIBE, A. FIORENZA and C. NEUGEBAUER, Weighted norm inequalities for the maximal operator on variable Lebesgue spaces, J. Math. Anal. Appl. 394 (2012), 744-760.
- [4] D. CRUZ-URIBE and A. FIORENZA, Variable Lebesgue Spaces, Birkhäuser, 2013.
- [5] D. CRUZ-URIBE and D. WANG, Variable Hardy Spaces, Indiana Univ. Math. J. 63 (2) (2014), 447-493.
- [6] L. DIENING, P. HARJULEHTO, P. HÄSTÖ and M. RUŽIČKA, Lebesgue and Sobolev spaces with Variable Exponents, Springer, 2011.
- [7] L. DIENING and P. HÄSTÖ, Muckenhoupt weights in variable exponent spaces, Preprint (2010).

- [8] J. GARCÍA-CUERVA, Weighted H^p spaces, *Dissertations Math.* 162 (1979), 1-63.
- [9] L. GRAFAKOS, *Classical Fourier Analysis*, 3rd edition, Graduate Texts in Mathematics, 249, Springer New York, 2014.
- [10] E. HARBOURE, R. MACÍAS and C. SEGOVIA, Extrapolation results for classes of weights, *Amer. J. Math.* 110 no. 3 (1988), 383-397.
- [11] K.-P. HO, Singular integral operators, John-Nirenberg inequalities and Triebel-Lizorkin type spaces on weighted Lebesgue spaces with variable exponents, *Rev. Un. Mat. Argentina* 57 (1) (2016), 85-101.
- [12] K.-P. HO, Atomic decompositions of weighted Hardy spaces with variable exponents, *Tohoku Math. J.* 69 (2) (2017), no. 3, 383-413.
- [13] K.-P. HO, Sublinear operators on weighted Hardy spaces with variable exponents, *Forum Math.* 31 (3) (2019), 607-617.
- [14] B. MUCKENHOUPT and R. WHEEDEN, Weighted norm inequalities for fractional integrals, *Trans. Amer. Math. Soc.*, 192 (1974), 261-274.
- [15] E. NAKAI and Y. SAWANO, Hardy spaces with variable exponents and generalized Campanato spaces, *J. Funct. Anal.* 262 (2012), 3665-3748.
- [16] P. ROCHA, A note on Hardy spaces and bounded linear operators, *Georgian Math. J.* 25 (1) (2018), 73-76.
- [17] P. ROCHA, Boundedness of generalized Riesz potentials on the variable Hardy spaces, *J. Aust. Math. Soc.* 104 (2018), 255-273.
- [18] P. ROCHA and M. URUIOLO, Fractional type integral operators on variable Hardy spaces, *Acta Math. Hung.* 143 (2) (2014), 502-514.
- [19] E. STEIN, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton University Press, 1993.
- [20] E. STEIN and G. WEISS, On the theory of harmonic functions of several variables I: The theory of H^p spaces, *Acta Math.* 103 (1960), 25-62.
- [21] M. H. TAIBLESON and G. WEISS, The molecular characterization of certain Hardy spaces, *Astérisque* 77 (1980), 67-149.
- [22] J.-O. STRÖMBERG and A. TORCHINSKY, *Weighted Hardy Spaces*, Lecture Notes in Mathematics, 1381, Springer-Verlag, Berlin, 1989.

DEPARTAMENTO DE MATEMÁTICA
UNIVERSIDAD NACIONAL DEL SUR
BAHÍA BLANCA, 8000 BUENOS AIRES
ARGENTINA

E-mail address: pablo.rocha@uns.edu.ar