

Note: a counterexample to a conjecture of Gilmer
which would imply the union-closed conjecture

David Ellis*

22nd November 2022

Abstract

In this very short note, we give a counterexample to a recent conjecture of Gilmer which would have implied the union-closed conjecture.

1 Introduction

We say a family \mathcal{F} of sets is *union-closed* if $A \cup B \in \mathcal{F}$ whenever $A, B \in \mathcal{F}$. The celebrated Union-Closed conjecture of Frankl states that if $n \in \mathbb{N}$ and $\mathcal{F} \neq \{\emptyset\}$ is a union-closed family of subsets of $\{1, 2, \dots, n\}$, then there exists $i \in \{1, 2, \dots, n\}$ such that at least half of the sets in \mathcal{F} contain i . Gilmer [2] recently obtained a breakthrough on this conjecture, proving that for any union-closed family \mathcal{F} of subsets of $\{1, 2, \dots, n\}$ with $\mathcal{F} \neq \{\emptyset\}$, there exists an element i contained in at least $0.01|\mathcal{F}|$ of the sets in \mathcal{F} . (We refer the reader to the survey of Bruhn and Schaudt [1] for survey of work prior to Gilmer's, on the problem.)

Gilmer's proof is an elegant entropy argument. At the end of his paper, Gilmer makes the following (information-theoretic) conjecture which would immediately imply the union-closed conjecture.

Conjecture 1 (Gilmer). *Let A, B be i.i.d. samples from a distribution over a family of subsets of $\{1, 2, \dots, n\}$. Assume that $\text{Prob}[i \in A] < \frac{1}{2}$ for all i , and that $H(A) > 0$. Then*

$$H(A \cup B) + D(A \cup B \| A) > H(A).$$

Here, $H(A)$ denotes the entropy of A ; recall that for a probability distribution $p = (p_x)_{x \in X}$ over a finite set X , the entropy of p is defined by

$$H(p) = \sum_{x \in X} p_x \log_2(1/p_x).$$

*School of Mathematics, University of Bristol, UK. Email: david.ellis@bristol.ac.uk

Also, $D(q||p)$ denotes the *Kullback-Leibler divergence* of q from p , for probability distributions $(p_x)_{x \in X}$ and $(q_x)_{x \in X}$ over a finite set X ; recall that this is defined by

$$D(q||p) = \sum_{x \in X} q_x \log_2(q_x/p_x).$$

The purpose of this very short note is to give a (simple) counterexample to this conjecture (with $n = 2$). Another counterexample (for large n) was independently and simultaneously obtained by Sawin.

2 The counterexample

As usual, we write $[n] := \{1, 2, \dots, n\}$, and for a set S , we write $\mathcal{P}(S)$ for the power-set of S . We first note that, writing p for the distribution of A in Gilmer's conjecture, and q for the distribution of $A \cup B$, the left-hand side of the conjectured inequality is equal to

$$\sum_{x \subset [n]} q_x \log_2(1/q_x) + \sum_{x \subset [n]} q_x \log_2(q_x/p_x) = \sum_{x \subset [n]} q_x \log_2(1/p_x),$$

and therefore Gilmer's conjecture is equivalent to

$$\sum_{x \subset [n]} q_x \log_2(1/p_x) - \sum_{x \subset [n]} p_x \log_2(1/p_x) > 0. \quad (1)$$

We first give a probability distribution p on $\mathcal{P}([2])$ such that, if A and B are i.i.d. samples from p , then $\text{Prob}[1 \in A] = \text{Prob}[2 \in A] = \frac{1}{2}$, but writing q for the distribution of $A \cup B$, the quantity on the left-hand side of (1) satisfies

$$\sum_{x \subset [n]} q_x \log_2(1/p_x) - \sum_{x \subset [n]} p_x \log_2(1/p_x) < -0.04. \quad (2)$$

The distribution p is as follows:

$$p(\emptyset) = p(\{1, 2\}) = x, \quad p(\{1\}) = p(\{2\}) = \frac{1}{2} - x, \quad (3)$$

where (for concreteness) we take $x = 0.3$. (We use the variable x to make the construction more readable.)

It remains to observe that an arbitrarily small perturbation (p' , say) of the probability distribution p satisfies the hypotheses of Gilmer's conjecture and yet also has the above quantity (2) being negative; indeed, we may replace p by the distribution p' defined by

$$p'(\emptyset) = x, \quad p'(\{1, 2\}) = x - 2\epsilon, \quad p'(\{1\}) = p'(\{2\}) = \frac{1}{2} + \epsilon - x,$$

for ϵ a sufficiently small positive number.

We now check that we do indeed have

$$\sum_{x \subset [2]} q_x \log_2(1/p_x) - \sum_{x \subset [2]} p_x \log_2(1/p_x) < -0.04,$$

for the probability distribution p defined in (3) above. Indeed, we have

$$q_\emptyset = \text{Prob}[A = B = \emptyset] = x^2,$$

and

$$\begin{aligned} q_{\{1\}} &= \text{Prob}[A = \emptyset \text{ and } B = \{1\}] + \text{Prob}[A = \{1\} \text{ and } B = \emptyset] + \text{Prob}[A = B = \{1\}] \\ &= 2x\left(\frac{1}{2} - x\right) + \left(\frac{1}{2} - x\right)^2 \\ &= x - 2x^2 + \frac{1}{4} - x + x^2 \\ &= \frac{1}{4} - x^2, \end{aligned}$$

and by symmetry $q_{\{2\}} = q_{\{1\}} = \frac{1}{4} - x^2$. Since q is a probability distribution, we have

$$q_{\{1,2\}} = 1 - q_\emptyset - q_{\{1\}} - q_{\{2\}} = 1 - x^2 - 2\left(\frac{1}{4} - x^2\right) = \frac{1}{2} + x^2.$$

Substituting in the above values, we have

$$\begin{aligned} &\sum_{x \subset [2]} q_x \log_2(1/p_x) - \sum_{x \subset [2]} p_x \log_2(1/p_x) \\ &= \left(\frac{1}{2} + 2x^2 - 2x\right) \log_2(1/x) + \left(-\frac{1}{2} - 2x^2 + 2x\right) \log_2(1/(\frac{1}{2} - x)), \end{aligned}$$

which is indeed less than -0.04 when $x = 0.3$, as claimed.

References

- [1] H. Bruhn and O. Schaudt. The journey of the union-closed sets conjecture. *Graphs and Combinatorics*, 31(6):2043–2074, 2015.
- [2] J. Gilmer. A constant lower bound for the union-closed sets conjecture. Preprint. arXiv:2211.09055.