

# MONOTONE MESHFREE METHODS FOR LINEAR ELLIPTIC EQUATIONS IN NON-DIVERGENCE FORM VIA NONLOCAL RELAXATION

QIHAO YE AND XIAOCHUAN TIAN

**ABSTRACT.** We design a monotone meshfree finite difference method for linear elliptic equations in the non-divergence form on point clouds via a nonlocal relaxation method. The key idea is a novel combination of a nonlocal integral relaxation of the PDE problem with a robust meshfree discretization on point clouds. Minimal positive stencils are obtained through a local  $l_1$ -type optimization procedure that automatically guarantees the stability and, therefore, the convergence of the meshfree discretization for linear elliptic equations. A major theoretical contribution is the existence of consistent and positive stencils for a given point cloud geometry. We provide sufficient conditions for the existence of positive stencils by finding neighbors within an ellipse (2d) or ellipsoid (3d) surrounding each interior point, generalizing the study for Poisson's equation by Seibold in 2008. It is well-known that wide stencils are in general needed for constructing consistent and monotone finite difference schemes for linear elliptic equations. Our result represents a significant improvement in the stencil width estimate for positive-type finite difference methods for linear elliptic equations in the near-degenerate regime (when the ellipticity constant becomes small), compared to previously known works in this area. Numerical algorithms and practical guidance are provided with an eye on the case of small ellipticity constant. At the end, we present numerical results for the performance of our method in both 2d and 3d, examining a range of ellipticity constants including the near-degenerate regime.

## 1. INTRODUCTION

In this work, we consider numerical approximations to the second-order elliptic equations in non-divergence form

$$(1.1) \quad \begin{cases} -Lu(\mathbf{x}) := -\sum_{i,j=1}^d a^{ij}(\mathbf{x})\partial_{ij}u(\mathbf{x}) = f(\mathbf{x}) & \mathbf{x} \in \Omega \\ u(\mathbf{x}) = g(\mathbf{x}) & \mathbf{x} \in \partial\Omega, \end{cases}$$

for a bounded domain  $\Omega \subset \mathbb{R}^d$ .  $A(\mathbf{x}) = (a^{ij}(\mathbf{x}))_{i,j=1}^d$  is a bounded and measurable matrix-valued function and is assumed to be symmetric and positive definite satisfying the uniform ellipticity condition

$$(1.2) \quad \lambda|\boldsymbol{\xi}|^2 \leq \boldsymbol{\xi}^T A(\mathbf{x})\boldsymbol{\xi} \leq \Lambda|\boldsymbol{\xi}|^2 \quad \forall \boldsymbol{\xi} \in \mathbb{R}^d, \forall \mathbf{x} \in \Omega$$

---

2020 *Mathematics Subject Classification.* Primary 65N06, 65N12, 65N35, 35B50, 35J15; Secondary 65R20, 35J70, 45A05.

*Key words and phrases.* meshfree method, minimal positive stencil, monotone scheme, discrete maximum principle, linear elliptic PDEs, near-degenerate, non-divergence form, nonlocal relaxation, linear minimization problem, point cloud, super-convergence.

for positive constants  $\lambda$  and  $\Lambda$ . Notice that by dividing both sides of the first equation in (1.1) by  $\Lambda$ , we can assume without loss of generality that  $\Lambda = 1$  and the ratio  $\varrho := \lambda/\Lambda = \lambda \leq 1$ .

Linear elliptic equations in non-divergence form are ubiquitous in science and engineering. They arise in many applications such as stochastic optimal control, materials science, and mathematical finance [21], and they are also recognized as the linearization of fully nonlinear PDEs such as the Monge-Ampère equation [34]. Notice that if the coefficient matrix  $A$  is differentiable, then the non-divergence form equation (1.1) can be recast into a divergence form convection-diffusion equation. However, when  $A$  is not differentiable, such reformulation no longer exists. PDE theories for non-divergence form elliptic equations are well-established in the literature. Existence, uniqueness and regularity theories are established for different notions of solutions including classical solutions, strong solutions, and viscosity solutions [9, 26, 40, 50].

In terms of numerical methods, non-divergence form elliptic PDEs are much less discussed than the divergence form PDEs because of a lack of variational formulation aforementioned. Discussions on finite element methods can be found in [18, 19, 32, 44, 47, 53] and the references therein. We pursue the direction of positive-type finite difference method for elliptic PDEs which has guaranteed convergence as a result of consistency and monotonicity/stability. Such pursuit dates back to Motzkin and Wasow [43], Kuo and Trudinger [31], and Kocan [30]. What was found in these work was that consistent and positive-type finite difference schemes exist for a given elliptic operator, but the stencil size grows with  $\varrho \rightarrow 0$ , and therefore the so-called wide stencil is a necessary feature of monotone finite difference methods even in the case of linear elliptic equations. Wide-stencil methods are later developed also for fully nonlinear elliptic PDEs [23, 49, 48, 46], and a few recent developments on monotone finite difference methods for nonlinear elliptic PDEs are found in [20, 22, 29]. For the linear elliptic PDEs, Kocan [30] gives an estimate of the stencil width for the existence of positive-type finite difference method, and it grows linearly with  $\varrho^{-1}$  in 2d and superlinearly with  $\varrho^{-1}$  in 3d, which severely impacts the practical use of such scheme for small ratio  $\varrho$ . In a more recent work [41], a positive-type finite difference scheme with anisotropic stencils is discussed but their stencil width estimate in 2d is also at the scale of  $\varrho^{-1}$ .

Our work is inspired by the recent development of meshfree methods for nonlocal models [15, 16, 35, 55]. The fundamental new idea of numerical approximation to (1.1) is a continuous nonlocal relaxation of the PDE problem followed by asymptotically compatible discretizations. More specifically, we first define a family of nonlocal operators  $\{\mathcal{L}_\delta\}_{\delta>0}$  as continuous approximations to the elliptic operator  $L$ , and then seek for robust discretizations  $\{\mathcal{L}_\delta^h\}_{\delta>0, h>0}$  by which the discretization parameter  $h$  is allowed to be proportional to the nonlocal length parameter  $\delta$ , i.e.,  $h = \mathcal{O}(\delta)$ , for the convergence to the elliptic problem (1.1). In our work,  $\mathcal{L}_\delta u(\mathbf{x})$  is defined by an integral over an elliptical region depending on  $A(\mathbf{x})$  where  $\delta > 0$  denotes the bound of the semi-major axis of it. We note that similar integral approximation to the linear elliptic operator is used in [47], while a non-robust discretization is employed so that a stronger relation,  $h = o(\delta)$ , is needed for convergence. In contrast, the idea of asymptotically compatible schemes [54, 55] could significantly improve the efficiency of numerical methods via nonlocal relaxation. The basic idea of the mesh-free discretization is an  $l_1$ -type local optimization method for obtaining minimal

positive stencils on point clouds. The major theoretical contribution (cf. Theorem 3.9) is an estimate of the elliptical searching region, parametrized by  $\delta > 0$ , for guaranteeing the existence of consistent and positive stencils. There are two main takeaways from Theorem 3.9. First, for a fixed ratio  $\varrho > 0$ ,  $\delta$  can be chosen as a constant multiple of  $h$ , leading to discretizations of asymptotically compatible type. Second, for small  $\varrho > 0$ , our theorem guarantees the existence of positive-type finite difference method within a stencil width of  $\mathcal{O}(\varrho^{-1/2})$ , which is a substantial improvement of the known theoretical results in [30, 41]. In terms of numerical algorithms, the size of the searching region determines the computational complexity of the local minimizations problems, and the near-optimal searching area estimate allows a practical assembly process in both 2d and 3d. In addition, the  $l_1$ -type minimization leads to minimal stencils with at most six points in 2d and ten points in 3d, thus the resulting linear system is sparse and can be solved efficiently by iterative methods. One notable feature of our approach is that it is the first numerical method, as far as we are aware, that has been successfully implemented in both 2d and 3d for the solution of linear elliptic equations with a wide range of  $\varrho \in (0, 1]$ , including the near-degenerate regime when  $\varrho \ll 1$ . We emphasize that our approach takes an innovative path that could have a potential impact on how meshfree and particle methods can be designed to enhance their mathematical properties as well as practical performances. Indeed, many traditional meshfree methods and data analysis algorithms unanimously suffer the restrictive condition  $h = o(\delta)$  on the two length-scales  $h$  and  $\delta$ , see e.g., [3, 8, 25, 36]. Our work exemplifies the possible approach to improving the performance of meshfree and particle methods via nonlocal relaxation and robust discretizations. Extension of the approach to other interesting problems will be considered in the further.

The rest of the paper is organized as follows. We first define a family of  $\delta > 0$  parameterized nonlocal integral approximations to eq. (1.1) in Section 2. The consistency errors of the continuum nonlocal model to the elliptic equation with respect to  $\delta$  are discussed. In Section 3, we present the meshfree discretization based on the nonlocal regularized problem. An  $l_1$ -type local optimization method is proposed in search of positive stencils in an elliptical neighborhood surrounding each node. A major theoretical result concerning the neighborhood criteria for guaranteeing the existence of consistent and positive stencils is presented in Theorem 3.9. Convergence of numerical solutions in terms of the discretization parameter  $h$  is then followed by mononicity and the relation of  $\delta$  and  $h$  in Theorem 3.9. In Section 4, we discuss point cloud generation and management, the assembling process, and provide complexity analysis and practical guidance for the implementation of the numerical method. Since the feasibility condition in Section 4 contains an implicit constant  $c = c(d) > 0$ , we estimate the constant numerically in this section and provide the searching neighborhood estimate that can be used in practice for  $d = 2$  and  $d = 3$ . Finally, we present the 2d and 3d numerical results in Section 5, and make conclusion and further discussions in Section 6.

## 2. NONLOCAL RELAXATION TO ELLIPTIC EQUATIONS

In this section, we discuss the nonlocal integral approximation to eq. (1.1) on which our numerical methods are based.

**2.1. Nonlocal elliptic operators in non-divergence form.** Nonlocal models have gained much interest in recent years [1, 5, 14]. In [12, 15], nonlocal Laplace

operator with a parameter dependence on  $\delta > 0$  is used as an approximation to the classical Laplace operator  $\Delta$  (when  $A(\mathbf{x}) = I$ ). The nonlocal Laplace operator is given by

$$\tilde{\mathcal{L}}_\delta u(\mathbf{x}) = \int_{B_\delta(\mathbf{0})} \frac{1}{\delta^{d+2}} \gamma\left(\frac{|\mathbf{y}|}{\delta}\right) (u(\mathbf{x} + \mathbf{y}) - u(\mathbf{x})) d\mathbf{y}$$

where  $\gamma$  is a nonnegative kernel with

$$(2.1) \quad \int_{B_1(\mathbf{0})} |\mathbf{y}|^2 \gamma(|\mathbf{y}|) d\mathbf{y} = 2d.$$

It can be shown that as  $\delta \rightarrow 0$ ,  $\tilde{\mathcal{L}}_\delta u(\mathbf{x}) \rightarrow \Delta u(\mathbf{x})$  for a sufficiently smooth function  $u$ . Here we consider a more general nonlocal elliptic operator that approximates the classical elliptic operator  $L$  in eq. (1.1) in the  $\delta \rightarrow 0$  limit. Following [47], we define the nonlocal elliptic operator parameterized by  $\delta$  as

$$(2.2) \quad \mathcal{L}_\delta u(\mathbf{x}) = \int_{\mathcal{E}_\delta^\mathbf{x}(\mathbf{0})} \frac{1}{\delta^{d+2}} \gamma\left(\frac{|M(\mathbf{x})^{-1}\mathbf{y}|}{\delta}\right) \det(M(\mathbf{x}))^{-1} (u(\mathbf{x} + \mathbf{y}) - u(\mathbf{x})) d\mathbf{y}$$

where  $M(\mathbf{x}) := (A(\mathbf{x}))^{1/2}$  is a positive definite matrix and  $\mathcal{E}_\delta^\mathbf{x}(\mathbf{z})$  denotes an ellipse with definition

$$(2.3) \quad \mathcal{E}_\delta^\mathbf{x}(\mathbf{z}) := \{\mathbf{y} \in \mathbb{R}^d : M(\mathbf{x})^{-1}(\mathbf{y} - \mathbf{z}) \in B_\delta(\mathbf{0})\}.$$

Notice that by our assumption on  $A(\mathbf{x})$ ,  $\mathcal{E}_\delta^\mathbf{x}(\mathbf{z})$  is an elliptical region centered




FIGURE 1. An illustration of  $\mathcal{E}_\delta^\mathbf{x}(\mathbf{z})$  in 2d.

at  $\mathbf{z}$  with semi-axes being  $\{\delta\sqrt{\lambda_i(\mathbf{x})}\}_{i=1}^d$  where  $\lambda_i(\mathbf{x})$  denotes the  $i$ -th smallest eigenvalue of  $A(\mathbf{x})$ . By our assumption,  $\varrho \leq \lambda_1(\mathbf{x}) \leq \dots \leq \lambda_d(\mathbf{x}) \leq 1$ . Figure 1 shows a 2d example where the semi-major axis of the ellipse is  $\delta\lambda_2^{1/2}(\mathbf{x}) \leq \delta$  and the semi-minor axis is  $\delta\lambda_1^{1/2}(\mathbf{x}) \geq \delta\sqrt{\varrho}$ . For convenience, we define

$$(2.4) \quad \rho_\delta(\mathbf{x}, \mathbf{y}) = \frac{1}{\delta^{d+2}} \gamma\left(\frac{|M(\mathbf{x})^{-1}\mathbf{y}|}{\delta}\right) \det(M(\mathbf{x}))^{-1}$$

and simply write  $\mathcal{L}_\delta u(\mathbf{x}) = \int_{\mathcal{E}_\delta^\mathbf{x}(\mathbf{0})} \rho_\delta(\mathbf{x}, \mathbf{y}) (u(\mathbf{x} + \mathbf{y}) - u(\mathbf{x})) d\mathbf{y}$ .

In the next, we show the consistency between  $\mathcal{L}_\delta$  and  $L$  on sufficiently smooth functions. The first result asserts that  $\mathcal{L}_\delta u$  agrees with  $Lu$  for all polynomials up to the third order, and the second result gives pointwise truncation error for sufficiently smooth functions. Similar calculations can be found in [47], here we present them for completeness and for the convenience of analyzing our numerical method in Section 3.

**Lemma 2.1.** *Let  $\mathcal{P}_p(\mathbb{R}^d)$  denote the space of all polynomials up to order  $p$  and  $\boldsymbol{\alpha} = (\alpha_1, \alpha_2, \dots, \alpha_d) \in (\mathbb{Z}^+ \cup \{0\})^d$  with  $|\boldsymbol{\alpha}| = \sum_i \alpha_i$ . Then for any  $\mathbf{x} \in \mathbb{R}^d$ ,*

$$(2.5) \quad \int_{\mathcal{E}_\delta^\mathbf{x}(\mathbf{0})} \rho_\delta(\mathbf{x}, \mathbf{y}) \mathbf{y}^\alpha d\mathbf{y} = \mathbf{0}$$

if  $|\boldsymbol{\alpha}|$  is an odd number and

$$(2.6) \quad \mathcal{L}_\delta u(\mathbf{x}) = Lu(\mathbf{x}) \quad \forall u \in \mathcal{P}_3(\mathbb{R}^d).$$

*Proof.* Consider  $\mathbf{x} \in \mathbb{R}^d$  fixed. If  $|\boldsymbol{\alpha}|$  is an odd number we have

$$\begin{aligned} \int_{\mathcal{E}_\delta^\mathbf{x}(\mathbf{0})} \rho_\delta(\mathbf{x}, \mathbf{y}) \mathbf{y}^\alpha d\mathbf{y} &= \int_{\mathcal{E}_\delta^\mathbf{x}(\mathbf{0})} \frac{1}{\delta^{d+2}} \gamma \left( \frac{|M(\mathbf{x})^{-1}\mathbf{y}|}{\delta} \right) \det(M(\mathbf{x}))^{-1} \mathbf{y}^\alpha d\mathbf{y} \\ &= \int_{B_\delta(\mathbf{0})} \frac{1}{\delta^{d+2}} \gamma \left( \frac{|\mathbf{y}|}{\delta} \right) (M(\mathbf{x})\mathbf{y})^\alpha d\mathbf{y} \\ &= 0. \end{aligned}$$

This last equality above is due to symmetry of the integration domain and anti-symmetry of the integrand.

Now it is easy to see that  $\mathcal{L}_\delta u(\mathbf{x}) = Lu(\mathbf{x}) = 0$  when  $u$  is a constant or  $u(\mathbf{z}) = (\mathbf{z} - \mathbf{x})^\alpha$  with  $|\boldsymbol{\alpha}|$  being an odd number. On the other hand, notice that

$$\int_{B_\delta(\mathbf{0})} \frac{1}{\delta^{d+2}} \gamma \left( \frac{|\mathbf{y}|}{\delta} \right) y_i y_j d\mathbf{y} = 0 \quad i \neq j$$

and

$$\int_{B_\delta(\mathbf{0})} \frac{1}{\delta^{d+2}} \gamma \left( \frac{|\mathbf{y}|}{\delta} \right) y_i^2 d\mathbf{y} = \frac{1}{d} \int_{B_\delta(\mathbf{0})} \frac{1}{\delta^{d+2}} \gamma \left( \frac{|\mathbf{y}|}{\delta} \right) |\mathbf{y}|^2 d\mathbf{y} = 2 \quad \forall i.$$

We have

$$\begin{aligned} \int_{\mathcal{E}_\delta^\mathbf{x}(\mathbf{0})} \rho_\delta(\mathbf{x}, \mathbf{y}) (\mathbf{y} \otimes \mathbf{y}) d\mathbf{y} &= \int_{\mathcal{E}_\delta^\mathbf{x}(\mathbf{0})} \frac{1}{\delta^{d+2}} \gamma \left( \frac{|M(\mathbf{x})^{-1}\mathbf{y}|}{\delta} \right) \det(M(\mathbf{x}))^{-1} (\mathbf{y} \otimes \mathbf{y}) d\mathbf{y} \\ &= \int_{B_\delta(\mathbf{0})} \frac{1}{\delta^{d+2}} \gamma \left( \frac{|\mathbf{y}|}{\delta} \right) (M(\mathbf{x})\mathbf{y} \otimes M(\mathbf{x})\mathbf{y}) d\mathbf{y} \\ &= M(\mathbf{x}) \int_{B_\delta(\mathbf{0})} \left( \frac{1}{\delta^{d+2}} \gamma \left( \frac{|\mathbf{y}|}{\delta} \right) (\mathbf{y} \otimes \mathbf{y}) d\mathbf{y} \right) M(\mathbf{x}) \\ &= 2(M(\mathbf{x}))^2 = 2A(\mathbf{x}), \end{aligned}$$

and thus  $\mathcal{L}_\delta u(\mathbf{x}) = Lu(\mathbf{x})$  when  $u(\mathbf{z}) = (\mathbf{z} - \mathbf{x})^\alpha$  with  $|\boldsymbol{\alpha}| = 2$ . Since  $\mathcal{L}_\delta(\mathbf{x})$  agrees with  $Lu(\mathbf{x})$  for  $u(\mathbf{z}) = (\mathbf{z} - \mathbf{x})^\alpha$  with  $|\boldsymbol{\alpha}| \leq 3$ , eq. (2.6) is true.  $\square$

We now consider an open bounded domain  $\Omega \subset \mathbb{R}^d$ . For  $\mathbf{x} \in \Omega$  near the boundary of  $\Omega$ , the definition in eq. (2.2) requires the values of  $u$  outside  $\Omega$ . Therefore nonlocal equations on bounded domains are usually accompanied by volumetric constraints ([15]) imposed over a boundary layer surrounding  $\Omega$ . In our case, we need to define the boundary interaction layer as

$$\Omega_{\mathcal{I}_\delta} = \{\mathbf{x} \in \mathbb{R}^d \setminus \Omega : \text{dist}(\mathbf{x}, \partial\Omega) < \delta\}.$$

We then denote the extended domain  $\Omega_\delta := \Omega \cup \Omega_{\mathcal{I}_\delta}$ . The consistency of  $\mathcal{L}_\delta$  to  $L$  is indicated in the following lemma.

**Lemma 2.2.** *Let  $\mathcal{L}_\delta u$  be defined by (2.2) and  $C > 0$  being a generic constant. Let  $\delta_0 > 0$  be a fixed number.*

(1) *If  $u \in C^2(\overline{\Omega_{\delta_0}})$ , then  $|\mathcal{L}_\delta u(\mathbf{x}) - Lu(\mathbf{x})| \rightarrow 0$  as  $\delta \rightarrow 0$  for all  $\mathbf{x} \in \Omega$ .*

(2) If  $u \in C^{k,\alpha}(\overline{\Omega_{\delta_0}})$  for  $k = 2$  or  $3$  and  $\alpha \in (0, 1]$ , then

$$|\mathcal{L}_\delta u(\mathbf{x}) - Lu(\mathbf{x})| \leq C|u|_{C^{k,\alpha}(\overline{\Omega_{\delta_0}})} \delta^{k-2+\alpha}$$

for all  $\mathbf{x} \in \Omega$  and  $\delta \leq \delta_0$ .

*Proof.* Notice that

$$u(\mathbf{x} + \mathbf{y}) - u(\mathbf{x}) = \int_0^1 \frac{d}{dt} u(\mathbf{x} + t\mathbf{y}) dt = \int_0^1 \nabla u(\mathbf{x} + t\mathbf{y}) \cdot \mathbf{y} dt.$$

Therefore

$$\begin{aligned} \mathcal{L}_\delta u(\mathbf{x}) &= \int_{\mathcal{E}_\delta^{\mathbf{x}}(\mathbf{0})} \rho_\delta(\mathbf{x}, \mathbf{y}) \mathbf{y}^T \int_0^1 \nabla u(\mathbf{x} + t\mathbf{y}) dt d\mathbf{y} \\ &= \int_{\mathcal{E}_\delta^{\mathbf{x}}(\mathbf{0})} \rho_\delta(\mathbf{x}, \mathbf{y}) \mathbf{y}^T \cdot \int_0^1 (\nabla u(\mathbf{x} + t\mathbf{y}) - \nabla u(\mathbf{x})) dt d\mathbf{y} \end{aligned}$$

where we have used  $\int_{\mathcal{E}_\delta^{\mathbf{x}}(\mathbf{0})} \rho_\delta(\mathbf{x}, \mathbf{y}) \mathbf{y} d\mathbf{y} = \mathbf{0}$  for the second line above. Then use

$$\nabla u(\mathbf{x} + t\mathbf{y}) - \nabla u(\mathbf{x}) = \int_0^1 D^2 u(\mathbf{x} + s\mathbf{y}) \cdot t\mathbf{y} ds,$$

we have

$$\mathcal{L}_\delta u(\mathbf{x}) = \int_{\mathcal{E}_\delta^{\mathbf{x}}(\mathbf{0})} \rho_\delta(\mathbf{x}, \mathbf{y}) (\mathbf{y} \otimes \mathbf{y}) : \int_0^1 t \int_0^1 D^2 u(\mathbf{x} + s\mathbf{y}) ds dt d\mathbf{y}$$

for all  $\mathbf{x} \in \Omega$ . On the other hand, from calculations in Lemma 2.1, one can show for  $\mathbf{x} \in \Omega$ ,

$$\begin{aligned} Lu(\mathbf{x}) &= \int_{\mathcal{E}_\delta^{\mathbf{x}}(\mathbf{0})} \rho_\delta(\mathbf{x}, \mathbf{y}) \frac{(\mathbf{y} \otimes \mathbf{y})}{2} : D^2 u(\mathbf{x}) d\mathbf{y} \\ &= \int_{\mathcal{E}_\delta^{\mathbf{x}}(\mathbf{0})} \rho_\delta(\mathbf{x}, \mathbf{y}) (\mathbf{y} \otimes \mathbf{y}) : \int_0^1 t \int_0^1 D^2 u(\mathbf{x}) ds dt d\mathbf{y}. \end{aligned}$$

For  $u \in C^2(\overline{\Omega_{\delta_0}})$ , we have  $D^2 u(\mathbf{x} + s\mathbf{y}) \rightarrow D^2 u(\mathbf{x})$  as  $\delta \rightarrow 0$  since  $\mathbf{y} \in \mathcal{E}_\delta^{\mathbf{x}}(\mathbf{0})$ . Therefore case (1) holds. For  $u \in C^{2,\alpha}(\overline{\Omega_{\delta_0}})$ , we have

$$|D^2 u(\mathbf{x} + s\mathbf{y}) - D^2 u(\mathbf{x})|_\infty \leq |u|_{C^{2,\alpha}(\overline{\Omega_{\delta_0}})} |s\mathbf{y}|^\alpha.$$

So

$$\begin{aligned} |\mathcal{L}_\delta u(\mathbf{x}) - Lu(\mathbf{x})| &\leq C|u|_{C^{2,\alpha}(\overline{\Omega_{\delta_0}})} \int_{\mathcal{E}_\delta^{\mathbf{x}}(\mathbf{0})} \rho_\delta(\mathbf{x}, \mathbf{y}) |\mathbf{y}|^{2+\alpha} d\mathbf{y} \\ &\leq C|u|_{C^{2,\alpha}(\overline{\Omega_{\delta_0}})} \delta^\alpha \int_{\mathcal{E}_\delta^{\mathbf{x}}(\mathbf{0})} \rho_\delta(\mathbf{x}, \mathbf{y}) |\mathbf{y}|^2 d\mathbf{y} \\ &= C|u|_{C^{2,\alpha}(\overline{\Omega_{\delta_0}})} \delta^\alpha \int_{B_\delta(\mathbf{0})} \frac{1}{\delta^{d+2}} \gamma_\delta \left( \frac{|\mathbf{y}|}{\delta} \right) \mathbf{y}^T A(\mathbf{x}) \mathbf{y} d\mathbf{y} \\ &\leq C|u|_{C^{2,\alpha}(\overline{\Omega_{\delta_0}})} \delta^\alpha \int_{B_\delta(\mathbf{0})} \frac{1}{\delta^{d+2}} \gamma_\delta \left( \frac{|\mathbf{y}|}{\delta} \right) |\mathbf{y}|^2 d\mathbf{y} \\ &= 2dC|u|_{C^{2,\alpha}(\overline{\Omega_{\delta_0}})} \delta^\alpha. \end{aligned}$$

Finally, if  $u \in C^{3,\alpha}(\overline{\Omega_{\delta_0}})$ , then we can write

$$\begin{aligned} & \mathcal{L}_\delta u(\mathbf{x}) - Lu(\mathbf{x}) \\ &= \int_{\mathcal{E}_\delta^{\mathbf{x}}(\mathbf{0})} \rho_\delta(\mathbf{x}, \mathbf{y})(\mathbf{y} \otimes \mathbf{y}) : \int_0^1 t \int_0^1 (D^2 u(\mathbf{x} + st\mathbf{y}) - D^2 u(\mathbf{x})) ds dt d\mathbf{y} \\ &= \int_{\mathcal{E}_\delta^{\mathbf{x}}(\mathbf{0})} \rho_\delta(\mathbf{x}, \mathbf{y})(\mathbf{y} \otimes \mathbf{y} \otimes \mathbf{y}) : \int_0^1 t^2 \int_0^1 s \int_0^1 D^3 u(\mathbf{x} + rst\mathbf{y}) dr ds dt d\mathbf{y} \\ &= \int_{\mathcal{E}_\delta^{\mathbf{x}}(\mathbf{0})} \rho_\delta(\mathbf{x}, \mathbf{y})(\mathbf{y} \otimes \mathbf{y} \otimes \mathbf{y}) : \int_0^1 t^2 \int_0^1 s \int_0^1 (D^3 u(\mathbf{x} + rst\mathbf{y}) - D^3 u(\mathbf{x})) dr ds dt d\mathbf{y}. \end{aligned}$$

Therefore, by the same reasoning as before, we have

$$|\mathcal{L}_\delta u(\mathbf{x}) - Lu(\mathbf{x})| \leq C|u|_{C^{3,\alpha}(\overline{\Omega_{\delta_0}})} \int_{\mathcal{E}_\delta^{\mathbf{x}}(\mathbf{0})} \rho_\delta(\mathbf{x}, \mathbf{y}) |\mathbf{y}|^{3+\alpha} d\mathbf{y} \leq 2dC|u|_{C^{3,\alpha}(\overline{\Omega_{\delta_0}})} \delta^{1+\alpha}.$$

□

### 3. MESHFREE DISCRETIZATION

Meshfree methods have been widely used in simulations of scientific problems. There are many existing meshfree approaches for solving PDEs on scattered datasets, such as vortex methods [7], smooth particle hydrodynamics (SPH) [27], radial basis functions (RBF)[6], moving least squares (MLS)/generalized moving least squares (GMLS) [33, 42, 59], and reproducing kernel particle method (RKPM) [38]. The key idea in MLS and GMLS is a local fitting of data using least squares approximation. They can also be written as a weighted  $l_2$ -type local optimization under certain reproducing conditions. In [51, 11], weighted  $l_1$ -type optimization was discussed for the sparsity of stencils. It was shown that using  $l_1$ -type optimization, the number of nonzero weights generated is at most the number of constraints in the reproducing condition. Such property is important to keep the linear system sparse, especially when the elliptic problem is nearly degenerate ( $\varrho \ll 1$ ), in the case of which the searching region becomes large (cf. Theorem 3.9). For the rest of this section, we discuss the meshfree method for solving eq. (1.1) based on the nonlocal relaxation and the convergence of the numerical method.

**3.1. Optimization based meshfree discretization.** Our numerical method is inspired by the meshfree finite difference method presented in [51] for solving the classical Poisson equation. The focus here is on the generation of positive stencils which lead to monotone schemes. The desirability of positive stencils was observed in other meshless methods, see e.g., [13, 37], although there was no guarantee of positive stencils in these works. We now present a reformulation of the meshfree method in [51] as a nonlocal relaxation method on which a generalization to elliptic equations is based. Given a point cloud  $X = \{\mathbf{x}_i\} \subset \mathbb{R}^d$  with  $h$  being its associated fill distance to be defined later, it is proposed in [51] to discretize the Laplace operator by

$$(3.1) \quad \Delta u(\mathbf{x}_i) \approx \Delta_h u(\mathbf{x}_i) = \sum_{\mathbf{x}_j \in B_\delta(\mathbf{x}_i)} \beta_{j,i}(u(\mathbf{x}_j) - u(\mathbf{x}_i)),$$

where the weights  $\{\beta_{j,i}\}$  are determined by the following linear minimization problem in order to achieve the so-called minimal positive stencils:

$$(3.2) \quad \begin{aligned} \{\beta_{j,i}\} &= \operatorname{argmin} \sum_j \frac{\beta_{j,i}}{W(|\mathbf{x}_j - \mathbf{x}_i|)} \\ \text{s.t. } \beta_{j,i} &\geq 0 \text{ and } \Delta u(\mathbf{x}_i) = \Delta_h u(\mathbf{x}_i) \forall u \in \mathcal{P}_p(\mathbb{R}^d) \end{aligned}$$

In [51], the polynomial space  $\mathcal{P}_p(\mathbb{R}^d)$  is taken to be  $\mathcal{P}_2(\mathbb{R}^d)$  with  $p = 2$ . The parameter  $\delta$  in eq. (3.1) is determined in relation to  $h$  such that the feasible set of the minimization problem is non-empty. The weight function  $W(r)$  is suggested in [51] as  $W(r) = r^{-\alpha}$  for  $\alpha > 0$ . It is not hard to see that when we choose the nonlocal kernel function  $\gamma(r) = Cr^{-\alpha}\chi_{\{|r|<1\}}$  that satisfies eq. (2.1), then by letting  $\beta_{j,i} = \frac{1}{\delta^{d+2}}\gamma\left(\frac{|\mathbf{x}_j - \mathbf{x}_i|}{\delta}\right)\omega_{j,i}$ , the minimization problem (3.2) is equivalent to

$$(3.3) \quad \begin{aligned} \{\omega_{j,i}\} &= \operatorname{argmin} \sum_j \omega_{j,i} \\ \text{s.t. } \omega_{j,i} &\geq 0 \text{ and } \tilde{\mathcal{L}}_\delta u(\mathbf{x}_i) = \tilde{\mathcal{L}}_\delta^h u(\mathbf{x}_i) \forall u \in \mathcal{P}_p(\mathbb{R}^d). \end{aligned}$$

where  $\tilde{\mathcal{L}}_\delta^h$  is the nonlocal approximate operator defined by

$$(3.4) \quad \tilde{\mathcal{L}}_\delta^h u(\mathbf{x}_i) = \sum_{\mathbf{x}_j \in B_\delta(\mathbf{x}_i)} \frac{1}{\delta^{d+2}} \gamma\left(\frac{|\mathbf{x}_j - \mathbf{x}_i|}{\delta}\right) \omega_{j,i} (u(\mathbf{x}_j) - u(\mathbf{x}_i)).$$

Based on this observation, we propose the discretization of nonlocal elliptic operator (2.2) as

$$(3.5) \quad \mathcal{L}_\delta^h u(\mathbf{x}_i) = \sum_{\mathbf{x}_j \in \mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i)} \rho_\delta(\mathbf{x}_i, \mathbf{x}_j - \mathbf{x}_i) \omega_{j,i} (u(\mathbf{x}_j) - u(\mathbf{x}_i)).$$

where  $\rho_\delta(\mathbf{x}_i, \mathbf{x}_j - \mathbf{x}_i) = \frac{1}{\delta^{d+2}} \gamma\left(\frac{|M(\mathbf{x}_i)^{-1}(\mathbf{x}_j - \mathbf{x}_i)|}{\delta}\right) \det(M(\mathbf{x}_i))^{-1}$ . The weights  $\{\omega_{j,i}\}$  in eq. (3.5) are solved by the minimization problem

$$(3.6) \quad \begin{aligned} \{\omega_{j,i}\} &= \operatorname{argmin} \sum_j \omega_{j,i} \\ \text{s.t. } \omega_{j,i} &\geq 0 \text{ and } \mathcal{L}_\delta^h u(\mathbf{x}_i) = \mathcal{L}_\delta u(\mathbf{x}_i) \forall u \in \mathcal{P}_p(\mathbb{R}^d). \end{aligned}$$

The well-posedness (3.6) is guaranteed only if the feasible set is non-empty. We will discuss in Section 3.4 the neighborhood criteria for non-emptiness of the feasible set

$$(3.7) \quad S_{\delta,h,p}(\mathbf{x}_i) := \{\{\omega_{j,i}\} : \omega_{j,i} \geq 0 \text{ and } \mathcal{L}_\delta^h u(\mathbf{x}_i) = \mathcal{L}_\delta u(\mathbf{x}_i) \forall u \in \mathcal{P}_p(\mathbb{R}^d)\}.$$

**3.2. Boundary treatment.** For an open and bounded domain  $\Omega \subset \mathbb{R}^d$ , we take a point cloud  $X = \{\mathbf{x}_i\}_{i=1}^M \subset \Omega_\delta$  and define its associated fill distance

$$(3.8) \quad h := \sup_{\mathbf{x} \in \Omega_\delta} \min_{1 \leq i \leq M} |\mathbf{x} - \mathbf{x}_i|$$

following the convention in [59]. Assume that  $\{\mathbf{x}_i\}_{i=1}^N \subset \Omega$ . For  $\mathbf{x}_i \in \Omega$  near the boundary of  $\partial\Omega$ , the definition in eq. (3.5) may require the value of  $u(\mathbf{x}_j)$  for  $\mathbf{x}_j \in \Omega_{\mathcal{I}_\delta}$ . Therefore, extensions of the boundary values from  $\partial\Omega$  to  $\Omega_{\mathcal{I}_\delta}$  are needed. However, it is usually hard to find an easy way to do the extension, especially in higher dimensions, to guarantee a second-order convergence rate for nonlocal solutions. We propose an alternative way for the boundary treatment.






FIGURE 2. Illustration of the projection

For  $\mathbf{x}_i \in \Omega$  and  $\mathbf{x}_j \in \mathcal{E}_\delta^{\mathbf{x}_i}$ , we define  $\overline{\mathbf{x}}_j = \mathbf{x}_j$  if  $\mathbf{x}_j \in \overline{\Omega}$ , otherwise  $\overline{\mathbf{x}}_j \in \partial\Omega$  is defined as the projection of  $\mathbf{x}_j$  onto  $\partial\Omega$  such that the line from  $\mathbf{x}_i$  to  $\overline{\mathbf{x}}_j$  is contained in  $\overline{\Omega}$ . Notice that the projected point  $\overline{\mathbf{x}}_j$  depends on both  $\mathbf{x}_j$  and  $\mathbf{x}_i$ . Here for notational convenience, we have omitted the dependence on  $i$  and simply denoted the projected point as  $\overline{\mathbf{x}}_j$ . See Figure 2 as an illustration of the projection. We then define the approximate operator associated with  $\delta$ ,  $h$  and  $\Omega$  as

$$(3.9) \quad \mathcal{L}_{\delta,\Omega}^h u(\mathbf{x}_i) = \sum_{\mathbf{x}_j \in \mathcal{E}_\delta^{\mathbf{x}_i}} \rho_\delta(\mathbf{x}_i, \overline{\mathbf{x}}_j - \mathbf{x}_i) \omega_{j,i} (u(\overline{\mathbf{x}}_j) - u(\mathbf{x}_i))$$

where  $\{\omega_{j,i}\}$  is solved from

$$(3.10) \quad \{\omega_{j,i}\} = \underset{\{\omega_{j,i}\} \in \overline{S}_{\delta,h,p}(\mathbf{x}_i)}{\operatorname{argmin}} \sum_j \omega_{j,i}$$

where  $\overline{S}_{\delta,h,p}(\mathbf{x}_i)$  is the feasible set defined as

$$(3.11) \quad \overline{S}_{\delta,h,p}(\mathbf{x}_i) := \{\{\omega_{j,i}\} : \omega_{j,i} \geq 0 \text{ and } \mathcal{L}_{\delta,\Omega}^h u(\mathbf{x}_i) = \mathcal{L}_\delta u(\mathbf{x}_i) \forall u \in \mathcal{P}_p(\mathbb{R}^d)\}.$$

We will address in Section 3.4 the feasibility of the minimization problem (3.10).

With the definition of the discrete operator  $\mathcal{L}_{\delta,\Omega}^h$ , we define the discrete problem as to find a function  $u_\delta^h : \{\mathbf{x}_i\}_{i=1}^N \cup \partial\Omega \rightarrow \mathbb{R}$  such that

$$(3.12) \quad \begin{cases} -\mathcal{L}_{\delta,\Omega}^h u_\delta^h(\mathbf{x}_i) = f(\mathbf{x}_i) & \mathbf{x}_i \in \Omega \\ u_\delta^h(\mathbf{x}) = g(\mathbf{x}) & \mathbf{x} \in \partial\Omega \end{cases}$$

**3.3. Convergence analysis.** We first provide a truncation error analysis for the discrete operator  $\mathcal{L}_{\delta,\Omega}^h$  and then the convergence is followed by the mononicity of the numerical scheme. In this subsection, the errors are presented in terms of  $\delta$ , which denotes the upper bound of the semi-major axis of the elliptical neighborhood of each point. Following the neighborhood criteria in Section 3.4, the convergence errors are finally presented in terms of  $h$ .

**Lemma 3.1.** *Take a point cloud  $X = \{\mathbf{x}_i\}_{i=1}^M \subset \Omega_\delta$  with  $\{\mathbf{x}_i\}_{i=1}^N \subset \Omega$ . Assume also that  $\overline{S}_{\delta,h,p}(\mathbf{x}_i)$  is not empty and  $C > 0$  is a generic constant.*

- (1) *If  $p \geq 2$  and  $u \in C^2(\overline{\Omega})$ , then  $|\mathcal{L}_{\delta,\Omega}^h u(\mathbf{x}_i) - Lu(\mathbf{x}_i)| \rightarrow 0$  as  $\delta \rightarrow 0$  for all  $\mathbf{x}_i \in \Omega$ .*

- (2) If  $p \geq 2$  and  $u \in C^{2,\alpha}(\overline{\Omega})$  for  $\alpha \in (0, 1]$ , then  $|\mathcal{L}_{\delta,\Omega}^h u(\mathbf{x}_i) - Lu(\mathbf{x}_i)| \leq C|u|_{C^{2,\alpha}(\overline{\Omega})} \delta^\alpha$  for all  $\mathbf{x}_i \in \Omega$ .
- (3) If  $p \geq 3$  and  $u \in C^{3,\alpha}(\overline{\Omega})$  for  $\alpha \in (0, 1]$ , then  $|\mathcal{L}_{\delta,\Omega}^h u(\mathbf{x}_i) - Lu(\mathbf{x}_i)| \leq C|u|_{C^{3,\alpha}(\overline{\Omega})} \delta^{1+\alpha}$  for all  $\mathbf{x}_i \in \Omega$ .

*Proof.* Consider a fixed  $\mathbf{x}_i \in \Omega$ . The proof follows closely from the proof of Lemma 2.2 by noticing that  $\{\omega_{j,i}\} \in \overline{S}_{\delta,h,p}(\mathbf{x}_i)$  implies that

$$(3.13) \quad \int_{\mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{0})} \rho_\delta(\mathbf{x}_i, \mathbf{y}) \mathbf{y}^\alpha d\mathbf{y} = \sum_{\mathbf{x}_j \in \mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i)} \rho_\delta(\mathbf{x}_i, \overline{\mathbf{x}}_j - \mathbf{x}_i) (\overline{\mathbf{x}}_j - \mathbf{x}_i)^\alpha \omega_{j,i}$$

for  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_d) \in (\mathbb{Z}^+ \cup \{0\})^d$  with  $|\alpha| = \sum_i \alpha_i \leq p$ .  $\square$

**Remark 3.2.** In practice, we often observe superconvergence for  $p = 2$ , which is likely due to symmetry. When  $p = 2$  and  $u \in C^{3,\alpha}(\overline{\Omega})$ , a more precise error estimate for  $\mathbf{x}_i \in \Omega$  is given by

$$|\mathcal{L}_{\delta,\Omega}^h u(\mathbf{x}_i) - Lu(\mathbf{x}_i)| \leq C \left( |u|_{C^3(\overline{\Omega})} T_3(\mathbf{x}_i) + |u|_{C^{3,\alpha}(\overline{\Omega})} \delta^{1+\alpha} \right)$$

where  $T_3(\mathbf{x}_i) = \max_{|\alpha|=3} \left| \sum_{\mathbf{x}_j \in \mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i)} \rho_\delta(\mathbf{x}_i, \overline{\mathbf{x}}_j - \mathbf{x}_i) (\overline{\mathbf{x}}_j - \mathbf{x}_i)^\alpha \omega_{j,i} \right|$

**Lemma 3.3** (Discrete maximum principle). *Let  $\Omega \subset \mathbb{R}^d$  be an open, bounded and simply connected domain. Take a point cloud  $X = \{\mathbf{x}_i\}_{i=1}^M \subset \Omega_\delta$  with  $\{\mathbf{x}_i\}_{i=1}^N \subset \Omega$ . Assume that there exists  $\mathbf{x}_i \in \Omega$  such that  $\mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i) \cap \Omega^c \neq \emptyset$ . If  $u \in C(\{\mathbf{x}_i\}_{i=1}^N \cup \partial\Omega)$  and  $\mathcal{L}_{\delta,\Omega}^h u(\mathbf{x}_i) \geq 0$  for all  $\mathbf{x}_i \in \Omega$ , then*

$$\max_{\mathbf{x}_i \in \Omega} u(\mathbf{x}_i) \leq \max_{\mathbf{x} \in \partial\Omega} u(\mathbf{x}).$$

*Proof.* First notice that  $\max_{\mathbf{x} \in \partial\Omega} u(\mathbf{x})$  is well-defined since  $\partial\Omega$  is a closed set and  $u$  is continuous on  $\partial\Omega$ . Assume that  $\max_{\mathbf{x}_i \in \Omega} u(\mathbf{x}_i) > \max_{\mathbf{x} \in \partial\Omega} u(\mathbf{x})$ , then there exists  $\mathbf{x}_k \in \Omega$  such that

$$u(\mathbf{x}_k) = \max_{\mathbf{x}_i \in \Omega} u(\mathbf{x}_i) \geq u(\mathbf{x}) \quad \forall \mathbf{x} \in \{\mathbf{x}_i\}_{i=1}^N \cup \partial\Omega.$$

Therefore

$$\mathcal{L}_{\delta,\Omega}^h u(\mathbf{x}_k) \leq 0.$$

By the assumption, we must have  $\mathcal{L}_{\delta,\Omega}^h u(\mathbf{x}_k) = 0$  and  $u(\mathbf{x}_j) = u(\mathbf{x}_k)$  for  $\mathbf{x}_j \in \mathcal{E}_\delta^{\mathbf{x}_k}(\mathbf{x}_k) \cap \Omega$ . Continue this process we can show that  $u$  is constant on  $\{\mathbf{x}_i\}_{i=1}^N \subset \Omega$ . Choose  $\mathbf{x}_i \in \Omega$  such that there exists  $\mathbf{x}_j \in \mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i) \cap \Omega^c$ , so  $\overline{\mathbf{x}}_j \in \partial\Omega$ . However, since we can argue that  $\mathcal{L}_{\delta,\Omega}^h u(\mathbf{x}_i) = 0$ , it implies  $u(\mathbf{x}_i) = u(\overline{\mathbf{x}}_j)$ , which is contradiction to  $\max_{\mathbf{x}_i \in \Omega} u(\mathbf{x}_i) > \max_{\mathbf{x} \in \partial\Omega} u(\mathbf{x})$ .  $\square$

**Theorem 3.4.** *Take a point cloud  $X = \{\mathbf{x}_i\}_{i=1}^M \subset \Omega_\delta$  and assume that  $\overline{S}_{\delta,h,p}(\mathbf{x}_i)$  is not empty. Let  $u$  and  $u_\delta^h$  be the solutions to eqs. (1.1) and (3.12) respectively.*

- (1) If  $p \geq 2$  and  $u \in C^{2,\alpha}(\overline{\Omega})$  for  $\alpha \in (0, 1]$ , then

$$\max_{\mathbf{x}_i \in \Omega} |u(\mathbf{x}_i) - u_\delta^h(\mathbf{x}_i)| \leq C|u|_{C^{2,\alpha}(\overline{\Omega})} \delta^\alpha.$$

- (2) If  $p \geq 3$  and  $u \in C^{3,\alpha}(\overline{\Omega})$  for  $\alpha \in (0, 1]$ , then

$$\max_{\mathbf{x}_i \in \Omega} |u(\mathbf{x}_i) - u_\delta^h(\mathbf{x}_i)| \leq C|u|_{C^{3,\alpha}(\overline{\Omega})} \delta^{1+\alpha}.$$

**Remark 3.5.** Using the truncation error estimate in Remark 3.2, one can show that if  $p = 2$  and  $u \in C^{3,\alpha}(\overline{\Omega})$ , then

$$\max_{\mathbf{x}_i \in \Omega} |u(\mathbf{x}_i) - u_\delta^h(\mathbf{x}_i)| \leq C(|u|_{C^3(\overline{\Omega})} \tau + |u|_{C^{3,\alpha}(\overline{\Omega})} \delta^{1+\alpha}).$$

where

$$\tau = \max_{\mathbf{x}_i \in \Omega} T_3(\mathbf{x}_i)$$

In practice,  $\tau$  might be a very small number depending on the point cloud. Therefore, superconvergence may be observed.

*Proof of Theorem 3.9.* We only show the proof for the first case and the second case can be similarly shown. Denote  $e_\delta^h(\mathbf{x}) = u(\mathbf{x}) - u_\delta^h(\mathbf{x})$  for  $\mathbf{x} \in \{\mathbf{x}_i\}_{i=1}^N \cup \partial\Omega$  and  $T_\delta^h(\mathbf{x}_i) = \mathcal{L}_{\delta,\Omega}^h u(\mathbf{x}_i) - Lu(\mathbf{x}_i)$  for  $\mathbf{x}_i \in \Omega$ . Notice that  $\mathcal{L}_{\delta,\Omega}^h e_\delta^h(\mathbf{x}_i) = T_\delta^h(\mathbf{x}_i)$  for  $\mathbf{x}_i \in \Omega$ . By Lemma 3.1, we have

$$K := \max_{\mathbf{x}_i \in \Omega} |T_\delta^h(\mathbf{x}_i)| \leq C|u|_{C^{2,\alpha}(\overline{\Omega})} \delta^\alpha.$$

Take  $\mathbf{x}^* \in \Omega$  such that  $\Omega \subset B_R(\mathbf{x}^*)$  for some  $R > 0$ . Then define  $\Phi(\mathbf{x}) = (\mathbf{x} - \mathbf{x}^*)^T (\mathbf{x} - \mathbf{x}^*) / (2d)$ , we have

$$\mathcal{L}_{\delta,\Omega}^h \Phi(\mathbf{x}) = A(\mathbf{x}) : D^2 \Phi(\mathbf{x}) = \frac{1}{d} \sum_{i=1}^d a_{ii}(\mathbf{x}) = \frac{1}{d} \sum_{i=1}^d \mathbf{e}_i^T A(\mathbf{x}) \mathbf{e}_i \geq \lambda,$$

where  $\mathbf{e}_i \in \mathbb{R}^d$  is the unit vector with the  $i$ -th component equal 1. Therefore we have

$$\mathcal{L}_{\delta,\Omega}^h \left( \frac{K}{\lambda} \Phi + e_\delta^h \right) (\mathbf{x}_i) \geq 0 \quad \forall \mathbf{x}_i \in \Omega,$$

and by Lemma 3.3, we have

$$\begin{aligned} \max_{\mathbf{x}_i \in \Omega} e_\delta^h(\mathbf{x}_i) &\leq \max_{\mathbf{x}_i \in \Omega} \left( \frac{K}{\lambda} \Phi(\mathbf{x}_i) + e_\delta^h(\mathbf{x}_i) \right) \leq \max_{\mathbf{x} \in \partial\Omega} \left( \frac{K}{\lambda} \Phi(\mathbf{x}) + e_\delta^h(\mathbf{x}) \right) \\ &= \frac{K}{\lambda} \max_{\mathbf{x} \in \partial\Omega} \Phi(\mathbf{x}) \leq \frac{K R^2}{\lambda 2d} \leq \frac{C R^2}{2\lambda d} |u|_{C^{2,\alpha}(\overline{\Omega})} \delta^\alpha. \end{aligned}$$

Similar estimates can be done for  $-e_\delta^h(\mathbf{x}_i)$  and therefore the proof is complete.  $\square$

**3.4. Neighborhood criteria.** In this subsection, we will discuss the neighborhood criteria that guarantee positive stencils. We only discuss the case  $p = 2$  in this subsection. The case  $p = 3$  is much harder to characterize which will be left for future work.

First of all, there is a sufficient criterion for positive stencils for solving the Laplace equation, and it is presented as a cone condition in [51] for  $d = 2$  or  $d = 3$ . For any  $\mathbf{x}_i \in \Omega$  and unit vector  $\mathbf{v} \in \mathbb{R}^d$ , we define an associated cone  $\mathcal{C}_\delta^{\mathbf{v}}(\mathbf{x}_i)$  in  $B_\delta(\mathbf{x}_i)$  by

$$(3.14) \quad \mathcal{C}_\delta^{\mathbf{v}}(\mathbf{x}_i) := \left\{ \mathbf{x} \in B_\delta(\mathbf{x}_i) : \mathbf{x}^T \mathbf{v} \geq \frac{1}{\sqrt{1 + \sigma_d}} |\mathbf{x}|^2 \right\}$$

where  $\sigma_d = \sqrt{2} - 1$  (a cone with total opening angle  $45^\circ$ ) for  $d = 2$  and  $\sigma_d = \sqrt{(3 - \sqrt{6})}/6$  (a cone with total opening angle  $33.7^\circ$ ) for  $d = 3$ . With a rephrasing of words, we quote the result in [51, Theorems 9 and 10] in the following lemma.

**Lemma 3.6** (Theorems 9 and 10 in [51]). *Take a point cloud  $X = \{\mathbf{x}_i\}_{i=1}^M \subset \Omega_\delta \subset \mathbb{R}^d$  and let  $\mathbf{x}_i \in \Omega$  be fixed. If for any unit vector  $\mathbf{v} \in \mathbb{R}^d$ ,  $\mathcal{C}_\delta^{\mathbf{v}}(\mathbf{x}_i) \cap X \setminus \{\mathbf{x}_i\} \neq \emptyset$ , then the feasible set to problem (3.3) with  $p = 2$  is not empty.*

To discuss the neighborhood criteria for our problem, we first notice that for  $\mathbf{x}_i \in \Omega$ , one can define a one-to-one mapping between  $B_\delta(\mathbf{x}_i)$  and  $\mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i)$  by

$$T_i \mathbf{x} = \mathbf{x}_i + M(\mathbf{x}_i)(\mathbf{x} - \mathbf{x}_i) \quad \mathbf{x} \in B_\delta(\mathbf{x}_i).$$

The inverse of  $T_i$  is then given by

$$T_i^{-1} \mathbf{x} = \mathbf{x}_i + M(\mathbf{x}_i)^{-1}(\mathbf{x} - \mathbf{x}_i) \quad \mathbf{x} \in \mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i).$$

We also denote  $T_i(D) = \{\mathbf{y} = T_i \mathbf{x} : \mathbf{x} \in D\}$  and  $T_i^{-1}(D) = \{\mathbf{y} = T_i^{-1} \mathbf{x} : \mathbf{x} \in D\}$  for any set  $D \subset \mathbb{R}^d$ .




FIGURE 3. Illustration of  $T_i$  and  $T_i^{-1}$

**Lemma 3.7.** *Take a point cloud  $X = \{\mathbf{x}_i\}_{i=1}^M \subset \Omega_\delta \subset \mathbb{R}^d$  and let  $\mathbf{x}_i \in \Omega$  be fixed. For any  $\mathbf{x}_j \in \mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i) \cap X \setminus \{\mathbf{x}_i\}$ , we write  $\tilde{\mathbf{x}}_j = T_i^{-1} \mathbf{x}_j$ . Let  $\mathcal{L}_\delta^h u(\mathbf{x}_i)$  be defined by eq. (3.5) and*

$$(3.15) \quad \tilde{\mathcal{L}}_\delta^h u(\mathbf{x}_i) := \sum_{\tilde{\mathbf{x}}_j \in B_\delta(\mathbf{x}_i)} \frac{1}{\delta^{d+2}} \gamma \left( \frac{|\tilde{\mathbf{x}}_j - \mathbf{x}_i|}{\delta} \right) \tilde{\omega}_{j,i}(u(\tilde{\mathbf{x}}_j) - u(\mathbf{x}_i)).$$

The following statements are equivalent.

- (1) There exists  $\{\tilde{\omega}_{j,i} \geq 0\}$  such that  $\tilde{\mathcal{L}}_\delta^h u(\mathbf{x}_i) = \tilde{\mathcal{L}}_\delta u(\mathbf{x}_i) \forall u \in \mathcal{P}_2(\mathbb{R}^d)$ .
- (2) There exists  $\{\omega_{j,i} \geq 0\}$  such that  $\mathcal{L}_\delta^h u(\mathbf{x}_i) = \mathcal{L}_\delta u(\mathbf{x}_i) \forall u \in \mathcal{P}_2(\mathbb{R}^d)$ .

*Proof.* By definition, we see that for any  $\mathbf{x}_j \in \mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i) \iff \tilde{\mathbf{x}}_j \in B_\delta(\mathbf{x}_i)$ . We show that (1) and (2) are equivalent by letting  $\tilde{\omega}_{j,i} = \det(M(\mathbf{x}_i))^{-1} \omega_{j,i}$ . Assume that (2) is true, then since  $\mathcal{L}_\delta u = 0$  if  $u \in \mathcal{P}_1(\mathbb{R}^d)$ , we have

$$\begin{aligned} \mathbf{0} &= \sum_{\mathbf{x}_j \in \mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i)} \frac{1}{\delta^{d+2}} \gamma \left( \frac{|M(\mathbf{x}_i)^{-1}(\mathbf{x}_j - \mathbf{x}_i)|}{\delta} \right) \det(M(\mathbf{x}_i))^{-1} \omega_{j,i} M(\mathbf{x}_i)^{-1}(\mathbf{x}_j - \mathbf{x}_i) \\ &= \sum_{\tilde{\mathbf{x}}_j \in B_\delta(\mathbf{x}_i)} \frac{1}{\delta^{d+2}} \gamma \left( \frac{|\tilde{\mathbf{x}}_j - \mathbf{x}_i|}{\delta} \right) \det(M(\mathbf{x}_i))^{-1} \omega_{j,i} (\tilde{\mathbf{x}}_j - \mathbf{x}_i). \end{aligned}$$

By letting  $\tilde{\omega}_{j,i} = \det(M(\mathbf{x}_i))^{-1} \omega_{j,i}$ , we see that  $\tilde{\mathcal{L}}_\delta^h u(\mathbf{x}_i) = 0 = \tilde{\mathcal{L}}_\delta u(\mathbf{x}_i) \forall u \in \mathcal{P}_1(\mathbb{R}^d)$ . Next by using  $\mathcal{L}_\delta^h u(\mathbf{x}_i) = \mathcal{L}_\delta u(\mathbf{x}_i) = A(\mathbf{x}_i) : D^2 u(\mathbf{x}_i)$  for  $u \in \mathcal{P}_2(\mathbb{R}^d)$ , we

have

$$\begin{aligned}
2 &= M(\mathbf{x}_i)^{-1}(2A(\mathbf{x}_i))M(\mathbf{x}_i)^{-1} \\
&= M(\mathbf{x}_i)^{-1} \sum_{\mathbf{x}_j \in \mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i)} \frac{1}{\delta^{d+2}} \gamma \left( \frac{|M(\mathbf{x}_i)^{-1}(\mathbf{x}_j - \mathbf{x}_i)|}{\delta} \right) \det(M(\mathbf{x}_i))^{-1} \omega_{j,i}((\mathbf{x}_j - \mathbf{x}_i) \otimes (\mathbf{x}_j - \mathbf{x}_i)) M(\mathbf{x}_i)^{-1} \\
&= \sum_{\tilde{\mathbf{x}}_j \in \mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i)} \frac{1}{\delta^{d+2}} \gamma \left( \frac{|\tilde{\mathbf{x}}_j - \mathbf{x}_i|}{\delta} \right) \tilde{\omega}_{j,i}(\tilde{\mathbf{x}}_j - \mathbf{x}_i) \otimes (\tilde{\mathbf{x}}_j - \mathbf{x}_i).
\end{aligned}$$

This implies that  $\tilde{\mathcal{L}}_\delta^h u(\mathbf{x}_i) = \tilde{\mathcal{L}}_\delta u(\mathbf{x}_i) = \Delta u(\mathbf{x}_i)$  when  $u \in \mathcal{P}_2(\mathbb{R}^d)$ . Therefore (2) implies (1). Similarly, we can show (1) also implies (2).  $\square$

The following result is an implication of Lemma 3.6 and Lemma 3.7.

**Corollary 3.8.** *Take a point cloud  $X = \{\mathbf{x}_i\}_{i=1}^M \subset \Omega_\delta \subset \mathbb{R}^d$  and let  $\mathbf{x}_i \in \Omega$  be fixed. If for any unit vector  $\mathbf{v} \in \mathbb{R}^d$ ,  $T_i(\mathcal{C}_\delta^{\mathbf{v}}(\mathbf{x}_i)) \cap X \setminus \{\mathbf{x}_i\} \neq \emptyset$ , then  $S_{\delta,h,2}(\mathbf{x}_i)$  and  $\bar{S}_{\delta,h,2}(\mathbf{x}_i)$  are not empty.*

*Proof.* First of all, from Lemma 3.6 and Lemma 3.7, it is easy to see that  $T_i(\mathcal{C}_\delta^{\mathbf{v}}(\mathbf{x}_i)) \cap X \setminus \{\mathbf{x}_i\} \neq \emptyset$  for all unit vector  $\mathbf{v} \in \mathbb{R}^d$  implies  $S_{\delta,h,2}(\mathbf{x}_i)$  (as defined in eq. (3.7)) is not empty. Now if we define a new point cloud  $\bar{X}$  by replacing all  $\mathbf{x}_j \in X \cap \mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i) \setminus \{\mathbf{x}_i\}$  with  $\bar{\mathbf{x}}_j$  in  $X$ . Then since  $\bar{\mathbf{x}}_j$  lies on the line connecting  $\mathbf{x}_i$  and  $\mathbf{x}_j$ , we see that

$$T_i(\mathcal{C}_\delta^{\mathbf{v}}(\mathbf{x}_i)) \cap X \setminus \{\mathbf{x}_i\} \neq \emptyset \implies T_i(\mathcal{C}_\delta^{\mathbf{v}}(\mathbf{x}_i)) \cap \bar{X} \setminus \{\mathbf{x}_i\} \neq \emptyset,$$

and the latter implies  $\bar{S}_{\delta,h,2}(\mathbf{x}_i)$  is not empty by the same reasoning. See an illustration of the sets  $T_i(\mathcal{C}_\delta^{\mathbf{v}}(\mathbf{x}_i)) \cap X \setminus \{\mathbf{x}_i\}$  and  $T_i(\mathcal{C}_\delta^{\mathbf{v}}(\mathbf{x}_i)) \cap \bar{X} \setminus \{\mathbf{x}_i\}$  in Figure 4.  $\square$




FIGURE 4. Illustration of Corollary 3.8.

Although Corollary 3.8 is a complete characterization of a sufficient condition for the well-posedness of eq. (3.10). It is hard to use in practice. In the following, we proceed to give a sufficient condition that is easy to use in the case  $d = 2$ . We leave the proof of the following theorem in Appendix A.

**Theorem 3.9.** *Let  $d = 2$  or  $d = 3$  and  $h$  be the fill distance defined in eq. (3.8). Let  $\lambda_1 = \lambda_1(\mathbf{x}_i)$  denote the smallest eigenvalue of  $A(\mathbf{x}_i)$ . Then there exists a constant  $c = c(d) > 0$  depending only on  $d$  such that if*

$$\delta \geq ch(\lambda_1)^{-1/2},$$

*then  $S_{\delta,h,2}(\mathbf{x}_i)$  and  $\bar{S}_{\delta,h,2}(\mathbf{x}_i)$  are not empty. Since  $\lambda_1 \leq \varrho$ , this implies the existence of positive stencils given  $\delta \geq ch(\varrho)^{-1/2}$ .*

**Remark 3.10.** *Notice that for a given point  $\mathbf{x}$ , the elliptical searching region surrounding  $\mathbf{x}$  has semi-axes  $\{\delta\sqrt{\lambda_i(\mathbf{x})}\}_{i=1}^d$  where  $\lambda_i(\mathbf{x})$  denotes the  $i$ -th smallest eigenvalue of  $A(\mathbf{x}_i)$ . The estimate in Theorem 3.9 is near-optimal in the sense that the semi-minor axis  $\delta\sqrt{\lambda_1(\mathbf{x})}$  of the searching neighborhood can be made proportional to  $h$  asymptotically for the existence of positive stencils. By Theorem 3.9, we can choose an elliptical neighborhood of  $\mathbf{x}$  whose volume is proportional to  $h^d(\prod_2^d \varrho_i)^{-1/2}$ , where  $\varrho_i = \varrho_i(\mathbf{x}) := \lambda_1(\mathbf{x})/\lambda_i(\mathbf{x}) \geq \varrho$ . This implies that the number of points within the searching neighborhood of  $\mathbf{x}$  is proportional to  $\varrho_2^{-1/2} \leq \varrho^{-1/2}$  in 2d and  $(\varrho_2\varrho_3)^{-1/2} \leq \varrho^{-1}$  in 3d. However, we do not have an explicit estimate of the constant  $c = c(d)$  in Theorem 3.9. In practice, we estimate this constant numerically which is described in detail in Section 4.3.*

Combining Theorem 3.9 with Theorem 3.4, one can take  $\delta = ch\varrho^{-1/2}$  and then the convergence rate is given in terms of  $h$ . This is summarized in the following Corollary.

**Corollary 3.11.** *Let  $\delta = ch\varrho^{-1/2}$  where  $c = c(d)$  is the constant in Theorem 3.9, then with a generic constant  $C > 0$ ,*

$$(3.16) \quad \max_{\mathbf{x}_i \in \Omega} |u(\mathbf{x}_i) - u_\delta^h(\mathbf{x}_i)| \leq C|u|_{C^{2+k,\alpha}(\bar{\Omega})} \varrho^{-(k+\alpha)/2} h^{k+\alpha}.$$

*for  $k = 0$  or  $k = 1$  and  $\alpha \in (0, 1]$ .*

#### 4. ALGORITHM DESIGN & COMPLEXITY ANALYSIS

In this section, we explain our algorithms in detail, mainly focusing on point cloud generation and matrix assembly.

**4.1. Point cloud generation.** In order to perform numerical experiments, some criteria need to be given on the point cloud geometry. For this, we first need to define two geometric quantities with respect to point clouds in addition to the fill distance defined in eq. (3.8). For a point cloud  $X = \{\mathbf{x}_i\}_{i=1}^M \subset \Omega_\delta$ , we define the separation distance  $\zeta$  as

$$(4.1) \quad \zeta := \frac{1}{2} \min_{1 \leq i < j \leq M} |\mathbf{x}_i - \mathbf{x}_j|.$$

In addition, for the points  $\{\mathbf{x}_i\}_{i=1}^N$  inside  $\Omega$ , we denote  $\kappa$  the minimum distance to the boundary, i.e.,

$$(4.2) \quad \kappa := \min_{1 \leq i \leq N} \text{dist}(\mathbf{x}_i, \partial\Omega).$$

With these geometric quantities, we now define proper point clouds to be used in numerical experiments.

**Definition 4.1.** Let  $X = \{\mathbf{x}_i\}_{i=1}^M \subset \Omega_\delta$  be a point cloud with its geometric quantities  $h$ ,  $\zeta$  and  $\kappa$  defined in eqs. (3.8), (4.1) and (4.2), respectively. Given a set of positive constants  $\{c_h, c_\zeta, c_\kappa\}$ , we say  $X$  is a proper point cloud (with respect to the constants  $\{c_h, c_\zeta, c_\kappa\}$ ) if it satisfies the following conditions:

- (i)  $h \leq c_h \left( \frac{|\Omega_\delta|}{|X|} \right)^{1/d}$ ;
- (ii)  $\zeta \geq c_\zeta h$ ;
- (iii)  $\kappa \geq c_\kappa h$ .

Notice that  $|\Omega_\delta|$  denotes the  $d$ -dimensional Lebesgue measure of  $\Omega_\delta$  and  $|X| = M$ .

**Remark 4.2.** Condition (ii) in Definition 4.1 essentially requires the point cloud to be quasi-uniform ([59]), and condition (iii) requires a certain distance from the interior points to the boundary set so that interior points would not be too close to the boundary points after the mapping described in Section 3.2 (this allows numerically solving eq. (3.10)). Notice that if (ii) is satisfied, then there exists  $C = C(c_\zeta, d) > 0$  such that  $h \leq C \left( \frac{|\Omega_\delta|}{|X|} \right)^{1/d}$ . In practice, we impose condition (i) with a chosen constant  $c_h > 0$  to have explicit control over the fill distance. In our numerical experiments in Section 5, we take  $c_h = 1$ ,  $c_\zeta = 0.175$ , and  $c_\kappa = 0.25$ .

**Remark 4.3.** Since the domain  $\Omega_\delta$  may be irregular, in practice, we always generate point clouds on a larger bounding box of  $\Omega_{\delta_0}$  for  $\delta \in (0, \delta_0]$ , as indicated by figs. 5a to 5d. The formula for the fill distance in eq. (3.8) and condition (i) Definition 4.1 are then modified accordingly. In practice,  $\delta_0$  is determined by the largest discretization parameter as well as the ratio  $\rho$  as indicated by neighborhood estimate in Section 3.4 and Section 4.3.

To generate a proper point cloud, we first initialize a random point cloud using the Quasi-Monte Carlo method [45] (see fig. 5a), then adjust this point cloud to make it proper. Adjustment contains three steps in each loop:

- Step 1. add points until  $h$  satisfies condition (i) (see fig. 5b);
- Step 2. map points until  $\kappa$  satisfies condition (iii) (see fig. 5c);
- Step 3. merge points until  $\zeta$  satisfies condition (ii) (see fig. 5d).

Adjustment stops when the conditions in Definition 4.1 are satisfied. In practice, it usually takes a few loops to make the point cloud proper.

**Remark 4.4.** One can also use the Quasi-Monte Carlo method for generating the initial point cloud and perform only step 2 without the adding and merging steps. Our adjustment algorithm provides explicit control over the fill distance and the separation distance and it leads to smaller fill distances for the same number of interior points compared with point clouds without adjustment. This is a trade-off situation, namely, one can save memory by using extra time adjusting the initialized point cloud, or vice versa.

**4.2. Matrix assembly.** The major effort in matrix assembly is the generation of the weights  $\{\omega_{j,i}\}$  defined in eq. (3.10). Notice that with respect to each point cloud and coefficient matrix  $A(\mathbf{x})$ , we need to solve  $N$  number of linear minimization problems to get the weights where  $N$  denotes the number of interior points. For each  $\mathbf{x}_i \in \Omega$ , we first need to find all the points inside the searching area, i.e., the domain  $\mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i)$  for a given a  $\delta > 0$ , then solve the linear minimization problem to get the stencil.




FIGURE 5. The process of proper point cloud generation. The grey circular domain is  $\Omega$ . The square domain is a bounding box of  $\Omega_{\delta_0}$  for some  $\delta_0 > 0$ . (A): initialize a point cloud by the Quasi-Monte Carlo method. (B): use the Voronoi diagram [57, 58] for the calculation of the fill distance, and then add the green points to the point cloud so that  $h$  satisfies condition (i). (C): map points near the boundary of  $\Omega$  to the interior so that  $\kappa$  satisfies condition (iii). (D): merge points whose distances are less than  $2c_\zeta h$  so that  $\zeta$  satisfies condition (ii). Notice that after merging of points, the fill distance may increase, as a result, the adjustment loop may be needed again.

We now describe the process of finding points inside an elliptical searching area. Notice that the proper point cloud is quasi-uniform, an easy way to accelerate this procedure is dividing the domain into same-size axis-aligned blocks [51, 59]. We call these blocks voxels. Alternatively, point clouds can also be managed by k-d trees [4, 59]. To search neighbors in the given elliptical area  $\mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i)$ , we first compute which voxel contains the current point  $\mathbf{x}_i$ , then search all the neighboring voxels that intersect non-trivially with  $\mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i)$ . If a voxel that intersects non-trivially with the searching area is contained in that area, then we add all the points in the voxel to the result set; otherwise, points in the voxel need to be checked one by



one. The intersection algorithm of voxels with ellipses/ellipsoids is crucial and we now describe it below.

Let  $\mathcal{H}$  denote a  $d$ -dimensional (hyper)rectangle and  $\mathcal{E}$  a  $d$ -dimensional ellipsoid.  $\mathcal{H}$  and  $\mathcal{E}$  are both open sets. We present an intersection detection algorithm in Algorithm 1 that distinguishes the following cases:

- Case 1.*  $\mathcal{H}$  does not intersect with  $\mathcal{E}$ ;
- Case 2.*  $\mathcal{H}$  is contained in  $\mathcal{E}$ ;
- Case 3.*  $\mathcal{H}$  intersects with  $\mathcal{E}$  but is not contained in  $\mathcal{E}$ .

---

**Algorithm 1:** Intersection detection of hyperrectangles with ellipsoids.

---

```

function intersection( $\mathcal{H}$ ,  $\mathcal{E}$ )
  if the center of  $\mathcal{H}$  is inside  $\mathcal{E}$  then
    if all the vertices of  $\mathcal{H}$  are inside  $\bar{\mathcal{E}}$  then
      | return Case 2;
    else
      | return Case 3;
  else if the center of  $\mathcal{E}$  is inside  $\mathcal{H}$  then
    | return Case 3;
  else
    if any one of the faces of  $\mathcal{H}$  intersects with  $\mathcal{E}$  then
      | return Case 3;
    else
      | return Case 1;

```

---

Notice that in Algorithm 1, the most time-consuming part is the intersection detection of faces of  $\mathcal{H}$  with  $\mathcal{E}$ . In 2d, the faces of a rectangle are line segments. Intersection detection of a line segment with an ellipse is relatively easy to carry out. One can first find the intersection (if exists) of the underlying line with the ellipse, which is a line segment (see fig. 6a as an illustration), by solving a quadratic equation. Then the intersection of two line segments can be easily checked. In 3d, to check whether a face of a 3d rectangle intersects with an ellipsoid, we first find the intersection area (if exists) of the underlying plane with the ellipse. Then since the intersection area (see fig. 6b as an illustration) is an ellipse, the problem is then reduced to the intersection detection of two-dimensional rectangles with ellipses. This can be further extended to higher dimensions, and a  $d$ -dimensional intersection problem can be reduced to a  $d-1$ -dimensional problem by this reasoning. Let  $Q_I(d)$  denote the complexity of the intersection algorithm in  $d$  dimensions. Notice that a  $d$ -dimensional hyperrectangle has  $2d$  faces, we can then deduce the recurrence relation

$$Q_I(d) \leq cdQ_I(d-1) \quad \text{with} \quad Q_I(1) = \mathcal{O}(1).$$

for some constant  $c > 0$  independent of  $d$ . Finally, the recurrence relation leads to

$$(4.3) \quad Q_I(d) = \mathcal{O}(c^d d!).$$

**Remark 4.5.** *The recursive algorithm for face-ellipsoid intersection detection gives a complexity (eq. (4.3)) that grows quickly with dimension. In this work, we only consider  $d = 2$  or  $d = 3$  so that  $Q_I(2)$  or  $Q_I(3)$  can be treated as constants. It*




FIGURE 6. Intersection illustration

will be of future interest to explore better intersection detection algorithms in higher dimensions.

**Remark 4.6.** The complexity for testing whether a hyperrectangle is contained in an ellipsoid is less than  $Q_I(d)$ . In fact, since a hyperrectangle has  $2^d$  vertices, this gives the complexity  $\mathcal{O}(2^d d^2)$ . Notice that when the ellipsoid is not aligned with the axes, we need to do the mapping with complexity  $\mathcal{O}(d^2)$  first, and then check with complexity  $\mathcal{O}(d)$  for each vertex.

For  $\mathbf{x}_i \in \Omega$ , let  $q(\mathbf{x}_i)$  denote the number of points in the searching area  $\mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i)$ , then the corresponding searching process needs  $\mathcal{O}(q(\mathbf{x}_i))$  intersection detections. For point clouds managed by k-d trees, it can be shown that  $\mathcal{O}(q(\mathbf{x}_i) \log M)$  intersection detections are needed for such range query [59]. Once we find all the points inside  $\mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i)$ , we proceed to solve the linear minimization problem eq. (3.10). We adopt the simplex method [10], which in average has a linear complexity in  $q(\mathbf{x}_i)$  when the dimension is fixed [52].

Combining the above discussions, when the dimension  $d$  is fixed, the total average complexity of finding a stencil for a given interior point  $\mathbf{x}_i$  is  $\mathcal{O}(q(\mathbf{x}_i))$ . Note that by the quasi-uniform assumption (condition (2) in Definition 4.1), we have  $q(\mathbf{x}_i) \leq C \frac{|\mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i)|}{h^d}$  for some  $C > 0$ . The volume  $|\mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i)|$  depends on  $\delta$  and the coefficient matrix  $A(\mathbf{x}_i)$ . In practice, we take  $h$  to be proportional to  $\delta$ , and therefore  $\frac{|\mathcal{E}_\delta^{\mathbf{x}_i}(\mathbf{x}_i)|}{h^d} = \mathcal{O}(1)$  considering  $A(\mathbf{x})$  to be fixed. As a result, the total complexity of searching for neighbors near a given point can be considered as a constant. In the near degenerate case, i.e.,  $\rho \ll 1$ ,  $q(\mathbf{x}_i)$  may grow with the decrease of  $\rho$  as mentioned in Remark 3.10.

Traversing all  $N$  number of interior points, we can get all the weights  $\{\omega_{j,i}\}$  to complete the matrix assembly process. Therefore, the whole complexity of assembling a matrix is given by  $\mathcal{O}(N)$  for a fixed problem. In addition, notice that the weights generation process is embarrassingly parallelizable, the actual computational time can be further reduced by parallelization.

**Remark 4.7.** From the  $l_1$  type minimization, we get a minimal positive stencil, and therefore the assembled matrix is sparse. It is also recommended that a reindexing process be applied to the point cloud to reduce the bandwidth of the assembled matrix. The simplest way to do this is to sort all the interior points by coordinates so that the index distance between two close points is not too large.

**Remark 4.8.** One may encounter memory issues using exact solvers when the linear system gets large. Iterative methods can be used in the case of large and sparse linear systems. We use the biconjugate gradient stabilized method (BiCGSTAB) [56] to approximately solve the sparse linear system when  $N$  is large.

**4.3. Searching area estimate.** Theorem 3.9 does not specify the constant  $c = c(d) > 0$  which determines the searching neighborhoods. Here we discuss how to determine the searching neighborhoods in practice. For a given fill distance  $h$ , we let  $\delta = ch(\varrho)^{-1/2}$ , where the determination of  $c > 0$  is described below. Then the searching neighborhood of a point  $\mathbf{x}$  is the domain  $\mathcal{E}_\delta^\mathbf{x}(\mathbf{x})$ .

We now discuss the choice of  $c > 0$  in practice. We first discuss the 2d case, and then use the 2d result to approximately estimate the searching area in 3d. Without loss of generality, we fix  $\mathbf{x}_i \in \Omega$  and assume that

$$A(\mathbf{x}_i) = \begin{pmatrix} \varrho & 0 \\ 0 & 1 \end{pmatrix}.$$

According to Lemma A.1, we need to find the smallest radius of the inscribed circles of the domains  $\{T_i(\mathcal{C}_\delta^\mathbf{v}(\mathbf{x}_i))\}_{\mathbf{v} \in \mathbb{R}^2, |\mathbf{v}|=1}$ . The problem is a rescaling of the case  $\delta = 1$ , as illustrated by fig. 7. Therefore we only need to consider the case  $\delta = 1$




FIGURE 7. An illustration of  $T_i(\mathcal{C}_\delta^\mathbf{v}(\mathbf{x}_i))$  and its inscribed circle as a rescaling from the  $\delta = 1$  case.

and find  $r(\varrho) := \min_{\mathbf{v} \in \mathbb{R}^2, |\mathbf{v}|=1} T_i(\mathcal{C}_1^\mathbf{v}(\mathbf{x}_i))$ . The detailed procedure for finding  $r(\varrho)$  numerically is provided in Appendix B. One may choose  $c = \max_{\varrho \in (0,1]} \sqrt{\varrho}/r(\varrho)$  and then by letting  $\delta = ch(\varrho)^{-1/2}$  we have the desired relation  $h \leq \delta r$ . In practice, we find that the  $\sqrt{\varrho}/r(\varrho)$  is a bit smaller with smaller  $\varrho > 0$  using the estimate of  $r(\varrho)$ . Since smaller constant  $c$  leads to smaller searching neighborhood, we therefore suggest taking different  $c$  for different values of  $\varrho \in (0, 1]$ . In particular, in 2d, by numerical approximations, we have the following estimate of  $c$ ,

$$c = c_{2d}(\varrho) := 2.836\chi_{(0,0.01]}(\varrho) + 2.901\chi_{(0.01,0.1]}(\varrho) + 3.614\chi_{(0.1,1]}(\varrho).$$

In 3d, it is difficult to estimate the radius of the inscribed ball in  $T_i(\mathcal{C}_\delta^\mathbf{v}(\mathbf{x}_i))$ . Therefore, we only take the intersecting ellipses of a given ellipsoid with the three planes that go through its principal axes, and perform the 2d estimate described above to obtain an estimate of the constant  $c > 0$  in 3d. The result is given as follows.

$$c = c_{3d}(\varrho) := 3.623\chi_{(0,0.01]}(\varrho) + 3.776\chi_{(0.01,0.1]}(\varrho) + 4.450\chi_{(0.1,1]}(\varrho).$$

In practice, we find that solutions often exist for even smaller searching areas and this means that one may take even smaller values of  $c > 0$  to further decrease the computational cost. We suggest taking  $c = \frac{1}{\sqrt{3}}c_{2d}(\varrho) \approx 0.58c_{2d}(\varrho)$  in 2d and  $c = \frac{1}{\sqrt[3]{18}}c_{3d} \approx 0.38C_{3d}(\varrho)$  in 3d first, and if no solution exists resetting  $c = c_{2d}(\varrho)$  in 2d and  $c = c_{3d}(\varrho)$  in 3d. This procedure could reduce the number of points in a searching neighborhood by a large factor.

## 5. NUMERICAL RESULTS

In this section, we report the results of numerical experiments for the study of the numerical accuracy of our method. We present 2d numerical results in Section 5.1 and 3d numerical results in Section 5.2.

**5.1. 2d numerical tests.** We test our numerical algorithm in 2d using two domains. The first domain is a unit disk given by  $\{x_1^2 + x_2^2 < 1\}$ , and the second domain is an L-shaped domain given by  $(-1, 1)^2 \setminus [0, 1]^2$ . The nonlocal kernel function is chosen as  $\gamma(r) = Cr^{-3}\chi_{\{|r|<1\}}$  so that eq. (2.1) is satisfied. We implement the numerical algorithm with  $p = 2$ . Smooth manufactured solutions are used in our tests with the right-hand side of eq. (1.1) computed based on them.

**5.1.1. Tests for continuous coefficient matrices.** We first test our algorithm for continuous coefficient matrices. Our baseline is  $A_0(\mathbf{x}) = I$  with  $\varrho = 1$ . A list of coefficient matrices used in numerical experiments is given below.

#	$A(\mathbf{x})$	$\varrho$
1	$\begin{pmatrix} 1 - 0.5 x_1  & 0 \\ 0 & 0.25 + 0.25 x_2  \end{pmatrix}$	0.2500
2	$\frac{1}{2.21} \begin{pmatrix} 2 -  x_1  & 0.5 \\ 0.5 & 0.5 + 0.5 x_2  \end{pmatrix}$	0.0864
3	$\begin{pmatrix} 1 - 0.5 x_1  & 0 \\ 0 & 0.025 + 0.025 x_2  \end{pmatrix}$	0.0250
4	$\begin{pmatrix} 1 - 0.5 x_1  & 0 \\ 0 & 0.0025 + 0.0025 x_2  \end{pmatrix}$	0.0025
5	$\frac{1}{2.001} \begin{pmatrix} 2 -  x_1(0.5 - x_2)  & 0.025 \\ 0.025 & 0.01 + 0.0025x_1 \exp(x_2) \end{pmatrix}$	0.0014

Notice that the value  $\varrho$  is computed approximately in the domain  $[-1, 1]^2$ , which contains both the unit disk and the L-shaped domain as subsets. Numerical results are presented in figs. 8 to 10. We observe second-order convergence in  $h$  for all cases, which is better than the theoretical analysis in Theorem 3.4 for  $p = 2$ . This superconvergence phenomenon may likely be due to the cancellation of terms as mentioned in Remarks 3.2 and 3.5. In figs. 8 and 9, we test our method on two manufactured solutions  $u_1^{(2d)}(x_1, x_2) = x_1x_2 + \cos(x_1)\exp(x_2)$  and  $u_2^{(2d)}(x_1, x_2) = (x_1 + x_2)^4 \cos(x_1(x_1 + 2x_2))$ . The numerical errors in these graphs grow as  $\varrho$  becomes smaller as predicted by theory. In some very special cases, the numerical errors may behave differently as  $\varrho \rightarrow 0$ , and one example is with  $u_3^{(2d)}(x_1, x_2) = x_1^2 + \sin(x_2)\exp(x_2^2 - 1)$  illustrated by fig. 10. The reason for this abnormal behavior is because the elliptic operator degenerates to  $\partial_{x_1}^2$  as  $\varrho \rightarrow 0$  by our choices of  $A(\mathbf{x})$

and the exact solution in this case is a second order polynomial in  $x_1$  which can be exactly reproduced by our method.




FIGURE 8. 2d tests on the unit disk domain with continuous coefficient matrices




FIGURE 9. 2d tests on the L-shaped domain with continuous coefficient matrices




FIGURE 10. 2d tests for a separable function

5.1.2. *Tests for discontinuous coefficient matrices.* We now show numerical results for discontinuous coefficient matrices. Notice that when  $A(\mathbf{x})$  is discontinuous, the elliptic equation in the non-divergence form cannot be recast into a variational form. Therefore, non-variational methods are especially important in this case. We divide the computational domains into smaller blocks and define piecewise constant coefficient matrices with respect to the blocks. More specifically, for  $n \in \mathbb{N}$ , we divide the domain  $[-1, 1]^2$  into  $(2n + 1)^2$  blocks, and define the corresponding piecewise constant coefficient matrix

$$A_\psi(\mathbf{x}, n) := (B_\psi(\mathbf{x}, n) + B_\psi^T(\mathbf{x}, n) + 4I)/8,$$

where  $B_\psi(\mathbf{x}, n)$  is generated by `mt19937`[39] (a pseudorandom number generator) with seed  $\psi(\mathbf{x}, n) := \text{round}(x_1 * n) * 2 + \text{round}(x_2 * n) * 3 \bmod 2^{32}$ . Here  $\text{round}(x)$  maps  $x$  to the closest integer. A list of coefficient matrices used in our experiments is given below.

#	$A(\mathbf{x})$	$\varrho$	discription
6	$A_\psi(\mathbf{x}, 10^{10})$	0.2500	dense blocks
7	$A_\psi(\mathbf{x}, 10^4)$	0.2500	medium blocks
8	$A_\psi(\mathbf{x}, 10^0)$	0.2500	loose blocks

In addition, we have the last example matrix:

$$A_9(\mathbf{x}) = \begin{cases} A_2(\mathbf{x}), & x_1 < 0 \\ A_3(\mathbf{x}), & \text{otherwise} \end{cases} \quad \text{with} \quad \varrho = 0.0250.$$

Numerical results are presented in figs. 11 and 12. We observe similar second-order convergence in  $h$  for all cases.




FIGURE 11. 2d tests on the unit disk domain with discontinuous coefficient matrices

5.2. **3d numerical tests.** 3d numerical tests are performed over the unit sphere given by  $\{x_1^2 + x_2^2 + x_3^2 < 1\}$  and a 3d L-shaped domain given by  $(-1, 1)^3 \setminus [0, 1] \times [-1, 1] \times [0, 1]$ . The nonlocal kernel function is chosen to be the same one as in the 2d case. We again test our algorithm for smooth manufactured solutions and  $p = 2$ .




FIGURE 12. 2d tests on the L-shaped domain with discontinuous coefficient matrices

5.2.1. *Tests for continuous coefficient matrices.* For the test on continuous coefficient matrices, we use the following list of coefficient matrices. Notice again that our baseline case is  $A_0(\mathbf{x}) = I$  with  $\varrho = 1$ .

#	$A(\mathbf{x})$	$\varrho$
1	$\begin{pmatrix} 1 - 0.5 x_1  & 0 & 0 \\ 0 & 0.5 - 0.25 x_2  & 0 \\ 0 & 0 & 0.25 + 0.25 x_3  \end{pmatrix}$	0.2500
2	$\frac{1}{2.21} \begin{pmatrix} 2 -  x_1  & 0 & 0.5 \\ 0 & 0.5 + 0.5 x_2  & 0 \\ 0.5 & 0 & 1 - 0.5 x_3  \end{pmatrix}$	0.0864
3	$\begin{pmatrix} 1 - 0.5 x_1  & 0 & 0 \\ 0 & 0.05 - 0.025 x_2  & 0 \\ 0 & 0 & 0.025 + 0.025 x_3  \end{pmatrix}$	0.0250
4	$\begin{pmatrix} 1 - 0.5 x_1  & 0 & 0 \\ 0 & 0.005 - 0.0025 x_2  & 0 \\ 0 & 0 & 0.0025 + 0.0025 x_3  \end{pmatrix}$	0.0025
5	$\frac{1}{2.001} \begin{pmatrix} 2 -  x_1(0.5 - x_2)  & -0.02 & 0.005 \\ -0.02 & 0.005 + 0.005 x_1 + x_3  & -0.001 \\ 0.005 & -0.001 & 0.01 + 0.0025x_2 \exp(x_3) \end{pmatrix}$	0.0014

Here the value  $\varrho$  is computed approximately in the domain  $[-1, 1]^3$ . Numerical results are presented in figs. 13 and 14 for the two manufactured solutions  $u_1^{(3d)}(x_1, x_2, x_3) = x_1x_2 + x_1x_3 + x_2x_3 + \cos(x_1) \exp(x_2 + x_3)$  and  $u_2^{(3d)}(x_1, x_2, x_3) = (x_1 + x_2 + x_3)^4 \cos(x_1(x_1 + 2x_2 + 2x_3))$ . We observe similar second-order convergence in  $h$  for all cases and numerical errors grow as  $\varrho$  becomes smaller.

5.2.2. *Tests for discontinuous coefficient matrices.* We now show numerical results for discontinuous coefficient matrices. Again, we divide the computational domains into smaller blocks and define piecewise constant coefficient matrices with respect to the blocks. More specifically, for  $n \in \mathbb{N}$ , we divide the domain  $[-1, 1]^3$  into  $(2n + 1)^3$  blocks, and define the corresponding piecewise constant coefficient matrix

$$A_\psi(\mathbf{x}, n) := (B_\psi(\mathbf{x}, n) + B_\psi^T(\mathbf{x}, n) + 4I)/10,$$

where  $B_\psi(\mathbf{x}, n)$  is generated by `mt19937` with seed  $\psi(\mathbf{x}, n) := \text{round}(x_1 * n) * 2 + \text{round}(x_2 * n) * 3 + \text{round}(x_3 * n) * 5 \pmod{2^{32}}$ . A list of coefficient matrices used in




FIGURE 13. 3d tests on the unit sphere domain with continuous coefficient matrices




FIGURE 14. 3d tests on the 3d L-shaped domain with continuous coefficient matrices

our experiments is given below.

#	$A(\mathbf{x})$	$\varrho$	discription
6	$A_\psi(\mathbf{x}, 10^{10})$	0.1847	dense blocks
7	$A_\psi(\mathbf{x}, 10^4)$	0.1847	medium blocks
8	$A_\psi(\mathbf{x}, 10^0)$	0.1847	loose blocks

In addition, we have the last example matrix:

$$A_9(\mathbf{x}) = \begin{cases} A_2(\mathbf{x}), & x_1 < 0 \\ A_3(\mathbf{x}), & \text{otherwise} \end{cases} \quad \text{with} \quad \varrho = 0.0250.$$

Numerical results are presented in figs. 15 and 16. We observe similar second-order convergence in  $h$  for all cases.

## 6. CONCLUSION

In this paper, we have presented a monotone meshfree finite difference method for linear elliptic equations in non-divergence form via integral relaxation. Minimal positive stencils are found through an  $l_1$ -type minimization problem within a local elliptical searching neighborhood of each point in a meshfree point cloud. For the






FIGURE 15. 3d tests on the unit sphere domain with discontinuous coefficient matrices




FIGURE 16. 3d tests on the 3d L-shaped domain with discontinuous coefficient matrices

treatment of Dirichlet boundary conditions, a mapping strategy near the boundary is incorporated into the numerical scheme. Convergence is guaranteed by the consistency and monotonicity of the scheme and efficient solvers can be designed by the sparsity of the resulting linear system. It is essential to characterize the shape and size of the elliptical searching neighborhood for the guarantee of positive stencils. Our theoretical result improves the previously known result for the stencil sizes when  $\varrho$ , the ratio between the smallest and the largest eigenvalues of the coefficient matrix, is a very small number. More precisely, our theory predicts that within an elliptical region with semi-major axis proportional to  $\varrho^{-1/2}h$ , we are able to find a positive stencil. The searching region determines the size of the  $l_1$ -type minimization problem, and therefore the efficiency of our algorithm. Our theory predicts that the number of points within the searching neighborhood grows with  $1/\varrho$  with a rate not worse than  $\mathcal{O}(\varrho^{-1/2})$  in 2d and  $\mathcal{O}(\varrho^{-1})$  in 3d.

We present algorithms for point cloud management and matrix assembly. We also give practical guidance for finding the elliptical searching neighborhood and present numerical experiments. Numerical tests are presented in both 2d and 3d for several different domains and coefficient matrices, including the near degenerate cases when  $\varrho \ll 1$ . While theoretical convergence in  $h$  for the numerical method (when the polynomial order  $p = 2$ ) is only first order, we observe second-order

convergence in all cases for manufactured smooth solutions. The super-convergence is likely due to the cancellation of odd order terms for the stencils obtained from the  $l_1$ -type minimization problem. A rigorous explanation for this phenomenon is still an open question.

Our current study focuses on the case of  $p = 2$  and  $d \in \{2, 3\}$  with Dirichlet boundary conditions. Future work includes higher order methods, problems in higher dimensions, and Neumann boundary value problems. Our convergence theorem assumes that the exact solution exists at least in  $C^2$ . Further question on the convergence of the method to viscosity solutions can be discussed, following the approaches discussed in [2, 17, 44]. Extending the study to surface PDEs is also a natural direction for future research. While we only test our algorithm for smooth manufactured solutions, adaptive methods will be useful when solutions display singularity. For adaptive point cloud management, some data structures that support fast insertion and deletion may be needed, for instance, R-tree [28] and scapegoat k-d tree [24]. The topic of monotone schemes for solving PDEs has a long history in numerical analysis. While our new ideas, inspired by the recent development of nonlocal modeling and meshfree methods, are presented for the linear elliptic equations, extending them to other types of PDEs is also possible for future research.

#### ACKNOWLEDGEMENT

This research is supported in part by NSF DMS-2111608. UC San Diego Research IT Services and Academic Technology Services are acknowledged for providing the Research Cluster computing resource for the numerical simulations in this work. The authors would like to thank Qiang Du, Xiaobing Feng, Nathaniel Trask and Zhen Zhang for their helpful discussions on the subject.

#### APPENDIX A. PROOF OF THEOREM 3.9

We show the proof of Theorem 3.9 in Appendix A. We begin with some useful lemmas before proving the theorem.

**Lemma A.1.** *Let  $r(\mathbf{v}, \mathbf{x}_i)$  denote the radius of the inscribed ball in  $T_i(\mathcal{C}_\delta^y(\mathbf{x}_i))$  and  $h$  be the fill distance associated with  $X = \{\mathbf{x}_i\}_{i=1}^M \subset \Omega_\delta$ . If*

$$h < \min_{\mathbf{v} \in \mathbb{R}^d, |\mathbf{v}|=1} r(\mathbf{v}, \mathbf{x}_i),$$

*then  $S_{\delta, h, 2}(\mathbf{x}_i)$  and  $\overline{S}_{\delta, h, 2}(\mathbf{x}_i)$  are not empty.*

*Proof.* Notice that by the definition of the fill distance in eq. (3.8), there are no holes with a radius larger than  $h$ . Suppose  $S_{\delta, h, 2}(\mathbf{x}_i)$  or  $\overline{S}_{\delta, h, 2}(\mathbf{x}_i)$  is empty, then by Corollary 3.8, there exists  $\mathbf{v}$  such that  $T_i(\mathcal{C}_\delta^y(\mathbf{x}_i))$  contains no point in  $X \setminus \{\mathbf{x}_i\}$ . Therefore the inscribed ball in  $T_i(\mathcal{C}_\delta^y(\mathbf{x}_i))$  is a hole with radius larger than  $h$  by the assumption, which gives a contradiction.  $\square$

From the lemma above, our goal is then to get a lower bound for

$$\min_{\mathbf{v} \in \mathbb{R}^d, |\mathbf{v}|=1} r(\mathbf{v}, \mathbf{x}_i)$$

for each  $\mathbf{x}_i \in \Omega$ . We first present a result in 2d which will also be useful for the 3d estimates. In 2d, we assume that  $\mathcal{C}_\delta^y(\mathbf{x}_i)$  is a cone with total opening angle  $2\phi$ . In

addition, without loss of generality, we fix  $\mathbf{x}_i \in \Omega$  and assume that

$$(A.1) \quad A(\mathbf{x}_i) = \begin{pmatrix} \varrho & 0 \\ 0 & 1 \end{pmatrix}.$$

From symmetry, we only need to consider  $\mathbf{v}(\theta) = (\cos(\theta), \sin(\theta))$  for  $\theta \in [0, \frac{\pi}{2}]$ .

**Lemma A.2.** *Consider  $d = 2$  and  $A(\mathbf{x}_i)$  given by eq. (A.1). Assume that  $\mathcal{C}_\delta^{\mathbf{v}}(\mathbf{x}_i)$  is a cone with total opening angle  $2\phi$  for  $\phi \in (0, \frac{\pi}{8}]$ , and  $r(\mathbf{v}, \mathbf{x}_i)$  denote the radius of the inscribed circle in  $T_i(\mathcal{C}_\delta^{\mathbf{v}}(\mathbf{x}_i))$ . In addition, let  $\mathbf{v}(\theta) = (\cos(\theta), \sin(\theta))$  for  $\theta \in [0, \frac{\pi}{2}]$ . Then, there exists a constant  $K = K(\phi) > 0$  such that*

$$\min_{\theta \in [0, \frac{\pi}{2}]} r(\mathbf{v}(\theta), \mathbf{x}_i) \geq K\delta\sqrt{\varrho}.$$

*Proof.* We try to fit a cone in  $T_i(\mathcal{C}_\delta^{\mathbf{v}(\theta)}(\mathbf{x}_i))$  and then find the inscribed circle in the cone. First, notice that for a cone with a radius  $R$  and total opening angle  $\alpha \in (0, \pi)$ , the radius of the inscribed circle is given by the formula

$$\frac{\sin(\alpha/2)}{1 + \sin(\alpha/2)} R \geq \frac{1}{2} \sin(\alpha/2) R.$$

Notice that  $\frac{1}{2} \sin(\alpha/2) R$  increases with  $\alpha \in (0, \pi)$  and  $R$ . Now for  $\theta \in [0, \frac{\pi}{2}]$ , we let  $\Gamma(\theta)$  denote the opening angle of  $T_i(\mathcal{C}_\delta^{\mathbf{v}(\theta)}(\mathbf{x}_i))$  and define

$$R(\theta) := \min_{\varphi \in [\theta - \phi, \theta + \phi]} \delta \sqrt{\varrho \cos^2(\varphi) + \sin^2(\varphi)} = \min_{\varphi \in [\theta - \phi, \theta + \phi]} \delta \sqrt{\varrho + (1 - \varrho) \sin^2(\varphi)},$$

then it is easy to see that a cone with radius  $R(\theta)$  and total opening angle  $\Gamma(\theta)$  is contained in  $T_i(\mathcal{C}_\delta^{\mathbf{v}(\theta)}(\mathbf{x}_i))$ . Therefore we have

$$\min_{\theta \in [0, \frac{\pi}{2}]} r(\mathbf{v}(\theta), \mathbf{x}_i) \geq \min_{\theta \in [0, \frac{\pi}{2}]} \frac{1}{2} \sin(\Gamma(\theta)/2) R(\theta).$$

By calculation we have

$$\Gamma(\theta) = \begin{cases} \arctan(\sqrt{\varrho^{-1}} \tan(\theta + \phi)) - \arctan(\sqrt{\varrho^{-1}} \tan(\theta - \phi)), & \theta \in [0, \pi/2 - \phi], \\ \pi + \arctan(\sqrt{\varrho^{-1}} \tan(\theta + \phi)) - \arctan(\sqrt{\varrho^{-1}} \tan(\theta - \phi)), & \theta \in (\pi/2 - \phi, \pi/2], \end{cases}$$

and

$$R(\theta) = \begin{cases} \delta\sqrt{\varrho}, & \theta \in [0, \phi], \\ \delta\sqrt{\varrho + (1 - \varrho) \sin^2(\theta - \phi)}, & \theta \in [\phi, \pi/2]. \end{cases}$$

For  $\theta \in [0, \phi]$ ,  $\Gamma(\theta)$  decreases and  $R(\theta) = \delta\sqrt{\varrho}$ , so

$$\min_{\theta \in [0, \phi]} \frac{1}{2} \sin(\Gamma(\theta)/2) R(\theta) = \frac{1}{2} \sin(\Gamma(\phi)/2) \delta\sqrt{\varrho} \geq \frac{\sin(\phi)}{2} \delta\sqrt{\varrho}$$

where we have used  $2\phi \leq \Gamma(\phi) \leq \pi/2$ .

For  $\theta \in [\pi/2 - 2\phi, \pi/2]$ ,  $\Gamma(\theta)$  decreases and  $R(\theta)$  increases, so

$$\begin{aligned} & \min_{\theta \in [\pi/2 - 2\phi, \pi/2]} \frac{1}{2} \sin(\Gamma(\theta)/2) R(\theta) \\ & \geq \frac{1}{2} \sin(\Gamma(\pi/2)/2) R(\pi/2 - 2\phi) \geq \frac{\delta}{2} \sin(\Gamma(\pi/2)/2) \sin(\pi/8) \\ & = \frac{\sin(\pi/8)\delta}{2} \frac{\sqrt{\varrho}}{\sqrt{\varrho + \tan^2(\pi/2 - \phi)}} \geq \frac{\sin(\pi/8)}{2\sqrt{1 + \tan^2(3\pi/8)}} \delta\sqrt{\varrho}. \end{aligned}$$

where we have used  $\phi \leq \pi/8$  and  $\Gamma(\pi/2) = 2 \arctan(\sqrt{\varrho} \cot(\pi/2 - \phi))$ .

Now for  $\theta \in [\phi, \pi/2 - 2\phi]$ , we use the formulas for  $\Gamma(\theta)$  and  $R(\theta)$  to compute  $\frac{1}{2} \sin(\Gamma(\theta)/2)R(\theta)$ . Denote

$$\alpha = \arctan(\sqrt{\varrho^{-1}} \tan(\theta + \phi)) \quad \text{and} \quad \beta = \arctan(\sqrt{\varrho^{-1}} \tan(\theta - \phi)).$$

Use the formula

$$\sin\left(\frac{\alpha - \beta}{2}\right) = \sqrt{\frac{1 - \cos(\alpha - \beta)}{2}} = \sqrt{\frac{1 - \cos(\alpha)\cos(\beta) - \sin(\alpha)\sin(\beta)}{2}},$$

and the fact that

$$\begin{aligned} \sin(\alpha) &= \frac{\sqrt{\varrho^{-1}} \tan(\theta + \phi)}{\sqrt{1 + \varrho^{-1} \tan^2(\theta + \phi)}}, \quad \cos(\alpha) = \frac{1}{\sqrt{1 + \varrho^{-1} \tan^2(\theta + \phi)}}, \\ \sin(\beta) &= \frac{\sqrt{\varrho^{-1}} \tan(\theta - \phi)}{\sqrt{1 + \varrho^{-1} \tan^2(\theta - \phi)}}, \quad \cos(\beta) = \frac{1}{\sqrt{1 + \varrho^{-1} \tan^2(\theta - \phi)}}, \end{aligned}$$

we can obtain the formula for  $\frac{1}{2} \sin(\Gamma(\theta)/2)R(\theta)$  where  $\theta \in [\phi, \pi/2 - 2\phi]$ . In particular, denote  $g_1(\theta) = \tan^2(\theta - \phi)$ ,  $g_2(\theta) = \tan^2(\theta + \phi)$ , and  $g_3(\theta) = \sin^2(\theta - \phi)$ , we fine

$$\begin{aligned} & \frac{1}{2} \sin(\Gamma(\theta)/2)R(\theta) \\ &= \frac{\delta}{2\sqrt{2}} \sqrt{\frac{\sqrt{(\varrho + g_1(\theta))(\varrho + g_2(\theta))} - \varrho - \sqrt{g_1(\theta)g_2(\theta)}}{\sqrt{(\varrho + g_1(\theta))(\varrho + g_2(\theta))}}} (\varrho + (1 - \varrho)g_3(\theta)) \\ &= \frac{\delta\sqrt{\varrho}}{2} \sqrt{\frac{((g_1(\theta))^{1/2} - (g_2(\theta))^{1/2})^2 (\varrho + (1 - \varrho)g_3(\theta))}{\sqrt{(\varrho + g_1(\theta))(\varrho + g_2(\theta))} (\sqrt{(\varrho + g_1(\theta))(\varrho + g_2(\theta))} + \varrho + \sqrt{g_1(\theta)g_2(\theta)})}} \\ &=: \frac{\delta\sqrt{\varrho}}{2} G(\varrho, \theta). \end{aligned}$$

Notice that  $G(\varrho, \theta)$  defined in the above is a continuous function on  $(\varrho, \theta) \in [0, 1] \times [\phi, \pi/2 - 2\phi]$ . Therefore it attains a minimum value at some  $(\varrho^*, \theta^*) \in [0, 1] \times [\phi, \pi/2 - 2\phi]$ . Next we show that we must have  $G(\varrho^*, \theta^*) > 0$ . Indeed, if  $\varrho^* > 0$ , then it is easy to see that  $G(\varrho^*, \theta^*) > 0$ . Now if  $\varrho^* = 0$ , then

$$\begin{aligned} G(0, \theta) &= \sqrt{\frac{((g_1(\theta))^{1/2} - (g_2(\theta))^{1/2})^2 g_3(\theta)}{2g_1(\theta)g_2(\theta)}} \\ &= (\tan(\theta + \phi) - \tan(\theta - \phi)) \frac{\sin(\theta - \phi)}{2 \tan(\theta - \phi) \tan(\theta + \phi)} \\ &= (\tan(\theta + \phi) - \tan(\theta - \phi)) \frac{\cos(\theta - \phi)}{2 \tan(\theta + \phi)} > 0 \end{aligned}$$

for any  $\theta \in [\phi, \pi/2 - 2\phi]$ . Therefore, we can take

$$K(\phi) = \min \left\{ \frac{\sin(\phi)}{2}, \frac{\sin(\pi/8)}{2\sqrt{1 + \tan^2(3\pi/8)}}, G(\varrho^*, \theta^*)/2 \right\} > 0$$

for the claim to be true.  $\square$

*Proof of Theorem 3.9.* Let  $\lambda_j = \lambda_j(\mathbf{x}_i)$  denotes the  $j$ -th smallest eigenvalue of  $A(\mathbf{x}_i)$ . For  $d = 2$ , we apply Lemma A.2 with  $\phi = \pi/8$  on a rescaled ellipse of  $T_i(B_\delta(\mathbf{x}_i))$ , we get

$$\min_{\theta \in [0, \frac{\pi}{2}]} r(\mathbf{v}(\theta), \mathbf{x}_i) \geq C_1 \delta \sqrt{\lambda_2} \sqrt{\frac{\lambda_1}{\lambda_2}} = C_1 \delta \sqrt{\lambda_1},$$

where  $C_1 = K(\pi/8)$ .

Now consider  $d = 3$ . First of all, we can assume without loss of generality that  $\lambda_3 = 1$  by the method of rescaling. By the discussions at the beginning of Section 3.4,  $\mathcal{C}_\delta^{\mathbf{v}}(\mathbf{x}_i)$  is a 3d cone with total opening angle  $33.7^\circ$  for a given unit vector  $\mathbf{v} \in \mathbb{R}^3$ . Let  $P_{\mathbf{v}} \subset \mathbb{R}^3$  be a 2d plane that contains the vector  $\mathbf{v}$ , then we see that  $P_{\mathbf{v}} \cap B_\delta(\mathbf{x}_i)$  is a circular domain and  $P_{\mathbf{v}} \cap \mathcal{C}_\delta^{\mathbf{v}}(\mathbf{x}_i)$  is a 2d cone with total opening angle  $33.7^\circ$ . With the transform  $T_i$ , we see that  $T_i(P_{\mathbf{v}} \cap B_\delta(\mathbf{x}_i))$  is a 2d ellipse and  $T_i(P_{\mathbf{v}} \cap \mathcal{C}_\delta^{\mathbf{v}}(\mathbf{x}_i))$  is a section of the ellipse. Therefore, the 2d calculations in Lemma A.2 can be applied. Notice that for each  $P_{\mathbf{v}}$ , there exists  $\rho_1$  and  $\rho_2$  with  $\lambda_1 \leq \rho_1 \leq \rho_2 \leq 1$  such that the lengths of the semi-axes of the ellipse  $T_i(P_{\mathbf{v}} \cap B_\delta(\mathbf{x}_i))$  are given by  $\sqrt{\rho_1}$  and  $\sqrt{\rho_2}$ . We can then rescale the the ellipse  $T_i(P_{\mathbf{v}} \cap B_\delta(\mathbf{x}_i))$  and use Lemma A.2 with  $\phi = 33.7^\circ/2$  we see that the radius of the inscribed circle in  $T_i(P_{\mathbf{v}} \cap \mathcal{C}_\delta^{\mathbf{v}}(\mathbf{x}_i))$  has a lower bound

$$\tilde{C}(\delta\sqrt{\rho_2})\sqrt{\frac{\rho_1}{\rho_2}} \geq \tilde{C}\delta\sqrt{\lambda_1},$$

where  $\tilde{C} = K(33.7^\circ/2) > 0$ . Notice that  $P_{\mathbf{v}}$  is an arbitrary plane that contains  $\mathbf{v}$ , and in addition, the average length of the line segments that connect  $T_i(\mathbf{x}_i)$  and the edge of  $T_i(P_{\mathbf{v}} \cap \mathcal{C}_\delta^{\mathbf{v}}(\mathbf{x}_i))$  is at the same scale for different plane  $P_{\mathbf{v}}$ . Therefore, it can be shown that there exists  $C_2 > 0$  such that

$$\min_{\theta \in [0, \frac{\pi}{2}]} r(\mathbf{v}(\theta), \mathbf{x}_i) \geq C_2 \delta \sqrt{\lambda_1}.$$

At last, by Lemma A.1 and the above discussions, there exists  $C = C(d) > 0$  such that whenever  $h \leq C\delta\sqrt{\lambda_1(\mathbf{x}_i)}$ ,  $S_{\delta,h,2}(\mathbf{x}_i)$  and  $\bar{S}_{\delta,h,2}(\mathbf{x}_i)$  are not empty. Therefore taking  $c(d) = 1/C(d)$  the conclusion is true. By our assumption,  $\varrho \leq \lambda_1(\mathbf{x}_i)$  for all  $\mathbf{x}_i$ , and therefore  $\delta \geq ch(\varrho)^{-1/2}$  implies the existence of positive stencils.  $\square$

## APPENDIX B. INSCRIBED CIRCLE SEARCH ALGORITHM

We show the details of finding the radius of the inscribed circle in  $T_i(\mathcal{C}_1^{\mathbf{v}}(\mathbf{x}_i))$  that contained in an ellipse given by  $x^2/\varrho + y^2 = 1$ . Let  $(x_0, y_0)$  be the center of the inscribed circle with radius  $r$ , then it can only sit on the angle bisector of  $T_i(\mathcal{C}_1^{\mathbf{v}}(\mathbf{x}_i))$ . Let  $(x_1, y_1)$  and  $(x_2, y_2)$  represent the two corner points of  $T_i(\mathcal{C}_1^{\mathbf{v}}(\mathbf{x}_i))$ . Then by some elementary calculations, there exists some  $t > 0$  such that

$$(x_0, y_0) = t \left( \sqrt{x_2^2 + y_2^2}x_1 + \sqrt{x_1^2 + y_1^2}x_2, \sqrt{x_2^2 + y_2^2}y_1 + \sqrt{x_1^2 + y_1^2}y_2 \right)$$

and

$$r = t|x_1y_2 - y_1x_2|.$$

To determine  $t > 0$ , we find the closest point to the circle center on the ellipse and choose  $t > 0$  such that the point is also on the circle. The closest point to the circle

center on the ellipse can be found by the minimization problem

$$\min_{\theta} \|(\sqrt{\varrho} \cos(\theta), \sin(\theta)) - (x_0, y_0)\|_2^2,$$

which can be solved by, e.g., Newton's method. At last, one may use a numerical method, e.g., the bisection method, to determine  $t > 0$  such that the point is also on the circle.

#### REFERENCES

1. Fuensanta Andreu-Vaillo, José M Mazón, Julio D Rossi, and J Julián Toledo-Melero, *Nonlocal diffusion problems*, no. 165, American Mathematical Soc., Providence, 2010.
2. Guy Barles and Panagiotis E Souganidis, *Convergence of approximation schemes for fully nonlinear second order equations*, *Asymptotic analysis* **4** (1991), no. 3, 271–283.
3. J Thomas Beale and Andrew Majda, *Vortex methods. I. convergence in three dimensions*, *Mathematics of computation* **39** (1982), no. 159, 1–27.
4. Jon Louis Bentley, *Multidimensional binary search trees used for associative searching*, *Communications of the ACM* **18** (1975), no. 9, 509–517.
5. Claudia Bucur and Enrico Valdinoci, *Nonlocal diffusion and applications*, vol. 20, Springer, New York, 2016.
6. Martin D Buhmann, *Radial basis functions: Theory and implementations*, vol. 12, Cambridge university press, Cambridge, 2003.
7. Georges-Henri Cottet, Petros D Koumoutsakos, et al., *Vortex methods: theory and practice*, vol. 8, Cambridge university press Cambridge, Cambridge, 2000.
8. Katy Craig and Andrea Bertozzi, *A blob method for the aggregation equation*, *Mathematics of computation* **85** (2016), no. 300, 1681–1717.
9. Michael G Crandall, Hitoshi Ishii, and Pierre-Louis Lions, *User's guide to viscosity solutions of second order partial differential equations*, *Bulletin of the American mathematical society* **27** (1992), no. 1, 1–67.
10. George B Dantzig, *Origins of the simplex method*, *A history of scientific computing*, 1990, pp. 141–151.
11. Oleg Davydov and Robert Schaback, *Minimal numerical differentiation formulas*, *Numerische Mathematik* **140** (2018), no. 3, 555–592.
12. Marta D'Elia, Qiang Du, Christian Glusa, Max Gunzburger, Xiaochuan Tian, and Zhi Zhou, *Numerical methods for nonlocal and fractional models*, *Acta Numerica* **29** (2020), 1–124.
13. Leszek Demkowicz, Andrzej Karafiat, and Tadeusz Liszka, *On some convergence results for FDM with irregular mesh*, *Computer Methods in Applied Mechanics and Engineering* **42** (1984), no. 3, 343–355.
14. Qiang Du, *Nonlocal modeling, analysis, and computation*, vol. 94, SIAM, Philadelphia, 2019.
15. Qiang Du, Max Gunzburger, Richard B Lehoucq, and Kun Zhou, *Analysis and approximation of nonlocal diffusion problems with volume constraints*, *SIAM review* **54** (2012), no. 4, 667–696.
16. Yiming Fan, Huaiqian You, Xiaochuan Tian, Xiu Yang, Xingjie Li, Naveen Prakash, and Yue Yu, *A meshfree peridynamic model for brittle fracture in randomly heterogeneous materials*, *Computer Methods in Applied Mechanics and Engineering* **339** (2022), 115340.
17. Xiaobing Feng, Roland Glowinski, and Michael Neilan, *Recent developments in numerical methods for fully nonlinear second order partial differential equations*, *SIAM Review* **55** (2013), no. 2, 205–267.
18. Xiaobing Feng, Lauren Hennings, and Michael Neilan, *Finite element methods for second order linear elliptic partial differential equations in non-divergence form*, *Mathematics of Computation* **86** (2017), no. 307, 2025–2051.
19. Xiaobing Feng, Michael Neilan, and Stefan Schnake, *Interior penalty discontinuous galerkin methods for second order linear non-divergence form elliptic pdes*, *Journal of Scientific Computing* **74** (2018), no. 3, 1651–1676.
20. Chris Finlay and Adam Oberman, *Improved accuracy of monotone finite difference schemes on point clouds and regular grids*, *SIAM Journal on Scientific Computing* **41** (2019), no. 5, A3097–A3117.

21. Wendell H Fleming and Halil Mete Soner, *Controlled markov processes and viscosity solutions*, vol. 25, Springer Science & Business Media, New York, 2006.
22. Brittany D Froese, *Meshfree finite difference approximations for functions of the eigenvalues of the Hessian*, *Numerische Mathematik* **138** (2018), no. 1, 75–99.
23. Brittany D Froese and Adam M Oberman, *Convergent finite difference solvers for viscosity solutions of the elliptic Monge–Ampère equation in dimensions two and higher*, *SIAM Journal on Numerical Analysis* **49** (2011), no. 4, 1692–1714.
24. Igal Galperin and Ronald L Rivest, *Scapegoat trees.*, *SODA*, vol. 93, 1993, pp. 165–174.
25. Nicolás García Trillos, Moritz Gerlach, Matthias Hein, and Dejan Slepčev, *Error estimates for spectral convergence of the graph laplacian on random geometric graphs toward the laplace–beltrami operator*, *Foundations of Computational Mathematics* **20** (2020), no. 4, 827–887.
26. David Gilbarg and Neil S Trudinger, *Elliptic partial differential equations of second order*, vol. 224, Springer, Berlin, 1977.
27. Robert A Gingold and Joseph J Monaghan, *Smoothed particle hydrodynamics: Theory and application to non-spherical stars*, *Monthly notices of the royal astronomical society* **181** (1977), no. 3, 375–389.
28. Antonin Guttman, *R-trees: A dynamic index structure for spatial searching*, *Proceedings of the 1984 ACM SIGMOD international conference on Management of data*, 1984, pp. 47–57.
29. Brittany Froese Hamfeldt and Jacob Lesniewski, *Convergent finite difference methods for fully nonlinear elliptic equations in three dimensions*, *Journal of Scientific Computing* **90** (2022), no. 1, 1–36.
30. Maciej Kocan, *Approximation of viscosity solutions of elliptic partial differential equations on minimal grids*, *Numerische Mathematik* **72** (1995), no. 1, 73–92.
31. Hung Ju Kuo and Neil S Trudinger, *Linear elliptic difference inequalities with random coefficients*, *Mathematics of computation* **55** (1990), no. 191, 37–53.
32. Omar Lakkis and Tristan Pryer, *A finite element method for second order nonvariational elliptic problems*, *SIAM Journal on Scientific Computing* **33** (2011), no. 2, 786–801.
33. Peter Lancaster and Kes Salkauskas, *Surfaces generated by moving least squares methods*, *Mathematics of computation* **37** (1981), no. 155, 141–158.
34. Nam Q Le, Hiroyoshi Mitake, and Hung V Tran, *Dynamical and geometric aspects of hamilton-jacobi and linearized Monge–Ampère equations*, Springer, New York, 2017.
35. Yu Leng, Xiaochuan Tian, Nathaniel Trask, and John T Foster, *Asymptotically compatible reproducing kernel collocation and meshfree integration for nonlocal diffusion*, *SIAM Journal on Numerical Analysis* **59** (2021), no. 1, 88–118.
36. Zhen Li and Zuoqiang Shi, *A convergent point integral method for isotropic elliptic equations on a point cloud*, *Multiscale Modeling & Simulation* **14** (2016), no. 2, 874–905.
37. TJ Liszka, CAM Duarte, and WW Tworzydło, *Hp-Meshless cloud method*, *Computer Methods in Applied Mechanics and Engineering* **139** (1996), no. 1-4, 263–288.
38. Wing Kam Liu, Y Chen, S Jun, JS Chen, T Belytschko, C Pan, RA Uras, and CT Chang, *Overview and applications of the reproducing kernel particle methods*, *Archives of Computational Methods in Engineering* **3** (1996), no. 1, 3–80.
39. Makoto Matsumoto and Takuji Nishimura, *Mersenne twister: a 623-dimensionally equidistributed uniform pseudo-random number generator*, *ACM Transactions on Modeling and Computer Simulation (TOMACS)* **8** (1998), no. 1, 3–30.
40. Antonino Maugeri, Dian K Palagachev, and Lubomira Softova, *Elliptic and parabolic equations with discontinuous coefficients*, *Mathematical Research* (2000).
41. Jean-Marie Mirebeau, *Minimal stencils for discretizations of anisotropic pdes preserving causality or the maximum principle*, *SIAM Journal on Numerical Analysis* **54** (2016), no. 3, 1582–1611.
42. Davoud Mirzaei, Robert Schaback, and Mehdi Dehghan, *On generalized moving least squares and diffuse derivatives*, *IMA Journal of Numerical Analysis* **32** (2012), no. 3, 983–1000.
43. Theodore S Motzkin and Wolfgang Wasow, *On the approximation of linear elliptic differential equations by difference equations with positive coefficients*, *Journal of Mathematics and Physics* **31** (1952), no. 1-4, 253–259.
44. Michael Neilan, Abner J Salgado, and Wujun Zhang, *Numerical analysis of strongly nonlinear pdes*, *Acta Numerica* **26** (2017), 137–303.
45. Harald Niederreiter, *Quasi-monte carlo methods and pseudo-random numbers*, *Bulletin of the American mathematical society* **84** (1978), no. 6, 957–1041.

46. R. Nochetto, Dimitrios Ntoggas, and Wujun Zhang, *Two-scale method for the Monge–Ampère equation: convergence to the viscosity solution*, *Mathematics of Computation* **88** (2019), no. 316, 637–664.
47. Ricardo H Nochetto and Wujun Zhang, *Discrete ABP estimate and convergence rates for linear elliptic equations in non-divergence form*, *Foundations of Computational Mathematics* **18** (2018), no. 3, 537–593.
48. ———, *Pointwise rates of convergence for the Ollier–Prüssner method for the Monge–Ampère equation*, *Numerische Mathematik* **141** (2019), no. 1, 253–288.
49. Adam M Oberman, *Wide stencil finite difference schemes for the elliptic monge-ampere equation and functions of the eigenvalues of the hessian*, *Discrete and Continuous Dynamical Systems-B* **10** (2008), no. 1, 221.
50. Mikhail V Safonov, *Nonuniqueness for second-order elliptic equations with measurable coefficients*, *SIAM Journal on Mathematical Analysis* **30** (1999), no. 4, 879–895.
51. Benjamin Seibold, *Minimal positive stencils in meshfree finite difference methods for the poisson equation*, *Computer Methods in Applied Mechanics and Engineering* **198** (2008), no. 3-4, 592–601.
52. Steve Smale, *On the average number of steps of the simplex method of linear programming*, *Mathematical programming* **27** (1983), no. 3, 241–262.
53. Iain Smears and Endre Suli, *Discontinuous galerkin finite element approximation of nondivergence form elliptic equations with cordes coefficients*, *SIAM Journal on Numerical Analysis* **51** (2013), no. 4, 2088–2106.
54. Xiaochuan Tian and Qiang Du, *Asymptotically compatible schemes and applications to robust discretization of nonlocal models*, *SIAM J. Numerical Analysis* **52** (2014), 1641–1665.
55. Nathaniel Trask, Huaqian You, Yue Yu, and Michael L Parks, *An asymptotically compatible meshfree quadrature rule for nonlocal problems with applications to peridynamics*, *Computer Methods in Applied Mechanics and Engineering* **343** (2019), 151–165.
56. Henk A Van der Vorst, *Bi-cgstab: A fast and smoothly converging variant of bi-cg for the solution of nonsymmetric linear systems*, *SIAM Journal on scientific and Statistical Computing* **13** (1992), no. 2, 631–644.
57. Georges Voronoi, *Nouvelles applications des paramètres continus à la théorie des formes quadratiques. deuxième mémoire. recherches sur les parallélogrammes primitifs.*, *Journal für die reine und angewandte Mathematik (Crelles Journal)* **1908** (1908), no. 134, 198–287.
58. ———, *Nouvelles applications des paramètres continus à la théorie des formes quadratiques. premier mémoire. sur quelques propriétés des formes quadratiques positives parfaites.*, *Journal für die reine und angewandte Mathematik (Crelles Journal)* **1908** (1908), no. 133, 97–102.
59. Holger Wendland, *Scattered data approximation*, vol. 17, Cambridge university press, Cambridge, 2004.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SAN DIEGO, CA 92093, UNITED STATES

*Email address:* q8ye@ucsd.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SAN DIEGO, CA 92093, UNITED STATES

*Email address:* xctian@ucsd.edu