

BEYOND THE BROKEN TETRAHEDRON

AUGUST CHEN AND BJARNE SCHÜLKE

ABSTRACT. Here we consider the hypergraph Turán problem in uniformly dense hypergraphs as was suggested by Erdős and Sós. Given a 3-graph F , the uniform Turán density $\pi_{\bullet}(F)$ of F is defined as the supremum over all $d \in [0, 1]$ for which there is an F -free uniformly d -dense 3-graph, where uniformly d -dense means that every linearly sized sub-hypergraph has density at least d . Recently, Glebov, Král', and Volec and, independently, Reiher, Rödl, and Schacht proved that $\pi_{\bullet}(K_4^{(3)-}) = \frac{1}{4}$, solving a conjecture by Erdős and Sós. Reiher, Rödl, and Schacht also characterised all 3-graphs with $\pi_{\bullet}(F) = 0$. Very recently, Bucić, Cooper, Král', Mohr, and Correia determined the uniform Turán density of 3-uniform tight cycles (of length at least 5), answering a question by Reiher. These hypergraphs and their blow-ups are the only hypergraphs for which the uniform Turán density is known.

In this work, we determine the uniform Turán density of the 3-graph on five vertices that is obtained from $K_4^{(3)-}$ by adding an additional vertex whose link forms a matching on the vertices of $K_4^{(3)-}$. Further, we point to two natural intermediate problems on the way to determining $\pi_{\bullet}(K_4^{(3)})$ and solve the first of these.

§1. INTRODUCTION

Turán's paper [18] in which he shows that complete, balanced, $(r - 1)$ -partite graphs maximise the number of edges among all K_r -free graphs is often seen as the starting point of extremal combinatorics. Already in this paper, he asked for similar results for hypergraphs. More than 80 years later, still little is known in this regard.

A k -uniform hypergraph (or k -graph) $H = (V, E)$ consists of a vertex set V and an edge set $E \subseteq V^{(k)} = \{e \subseteq V : |e| = k\}$. The extremal number for $n \in \mathbb{N}$ and a k -uniform hypergraph F is the maximum number of edges in an F -free k -uniform hypergraph on n vertices and it is denoted by $\text{ex}(n, F)$. The Turán density of F is $\pi(F) = \lim_{n \rightarrow \infty} \frac{\text{ex}(n, F)}{\binom{n}{k}}$ (this limit was shown to exist by monotonicity in [12]). The aforementioned result by Turán was later generalised to the Erdős-Stone-Simonovits theorem [7, 9], which determines the Turán density of any graph F to be $\frac{\chi(F)-2}{\chi(F)-1}$ (where $\chi(F)$ is the chromatic number of F).

For hypergraphs, only few (non-trivial) Turán densities are known, one prominent example being the Fano plane [2, 5, 10, 14]. The Turán density of the complete 3-uniform hypergraph

Key words and phrases. Hypergraphs, Turán density.

on four vertices, $K_4^{(3)}$, is still not known, although it is widely conjectured to be $5/9$. In fact, the problem is even open for the 3-uniform hypergraph that consists of three edges on four vertices, denoted by $K_4^{(3)-}$. For more on the hypergraph Turán problem, we refer to the survey by Keevash [13].


In this work, we consider a variant of the hypergraph Turán problem that was suggested by Erdős and Sós [6, 8] who asked for the maximum density of F -free hypergraphs H if we require H to be uniformly dense. More precisely, given $d \in [0, 1]$ and $\eta > 0$, a 3-uniform hypergraph $H = (V, E)$ is called *uniformly (d, η) -dense* if for every $U \subseteq V$, we have $|U^{(3)} \cap E| \geq d \binom{|U|}{3} - \eta |V|^3$. Then the uniform Turán density of a 3-uniform hypergraph F is defined as

$$\pi_{\bullet}(F) = \sup\{d \in [0, 1] : \text{for every } \eta > 0 \text{ and } n \in \mathbb{N} \text{ there is an } \\ F\text{-free, uniformly } (d, \eta)\text{-dense 3-graph } H \text{ with } |V(H)| \geq n\}.$$

Erdős and Sós specifically asked to determine $\pi_{\bullet}(K_4^{(3)})$ and $\pi_{\bullet}(K_4^{(3)-})$. Similarly as with the original Turán density, these problems turned out to be very difficult. Only recently, Glebov, Král', and Volec [11] and Reiher, Rödl, and Schacht [16] independently solved the latter, showing that $\pi_{\bullet}(K_4^{(3)-}) = 1/4$ and thus confirming a conjecture by Erdős and Sós. We refer to Reiher's survey [15] for a full description of the landscape of extremal problems in uniformly dense hypergraphs. Further, Reiher, Rödl, and Schacht [17] *characterised* all 3-uniform hypergraphs F with $\pi_{\bullet}(F) = 0$. Very recently, Bucić, Cooper, Král', Mohr, and Correia [4] determined the uniform Turán density of all tight 3-uniform cycles of length at least five. Note that by supersaturation $\pi_{\bullet}(F) = \pi_{\bullet}(F(t))$ for every 3-uniform hypergraph F and $t \in \mathbb{N}$, where $F(t)$ is the t -blow-up of F , the hypergraph obtained from F by replacing every vertex of F by t copies of itself. We say that F' is a blow-up of F if $F' \subseteq F(t)$ for some $t \in \mathbb{N}$. Thus, by determining $\pi_{\bullet}(F)$ for some 3-uniform hypergraph F , also the uniform Turán density of any blow-up of F is determined. Although there are more results on other variants of the Turán density (see [1, 15] and the references therein, and [3]) and some partial results for π_{\bullet} of some hypergraphs [1], as far as we know, (blow-ups of) the $K_4^{(3)-}$, hypergraphs F with $\pi_{\bullet}(F) = 0$, and tight cycles of length at least five are the only hypergraphs for which π_{\bullet} is known.

In this work, we add another example, namely F_{\star} , which we define as the hypergraph on five vertices with edge set $\{123, 124, 134, 125, 345\}$, see Figure 1.1. In other words, F_{\star} is the 3-uniform hypergraph obtained from a $K_4^{(3)-}$ by adding an additional vertex whose link is a matching.

Theorem 1.1. *We have $\pi_{\bullet}(F_{\star}) = 1/4$.*

FIGURE 1.1. The hypergraph F_* .

One potential difficulty in moving from determining $\pi_{\bullet}(K_4^{(3)-})$ to $\pi_{\bullet}(K_4^{(3)})$, which has been described as perhaps the most urgent problem in this area [15], is that there are no 3-uniform hypergraphs F with $K_4^{(3)-} \subsetneq F \subsetneq K_4^3$ that could serve as intermediate steps. However, Reiher, Rödl, and Schacht reduced the problem of determining the uniform Turán density of any hypergraph to a problem in *reduced hypergraphs* (see Theorem 2.4). The terminology of reduced hypergraphs gives rise to two natural problems ‘between’ determining $\pi_{\bullet}(K_4^{(3)-})$ and $\pi_{\bullet}(K_4^{(3)})$. Our second result solves the first of these. Since the notation needed is not the lightest, we postpone the introduction of the notation to Section 2 and ask the reader unfamiliar with it to read it first before continuing here.

Suppose $\mathcal{A} = (I, \mathcal{P}^{ij}, \mathcal{A}^{ijk})$ is a (d, \bullet) -dense reduced hypergraph and (λ, φ) is a reduced map from $K_4^{(3)-}$ (with vertex set $[4]$ and apex at vertex 1) to \mathcal{A} . Let $\lambda(j) = i_j$ and $\varphi(jk) = \alpha_{jk}$ for $jk \in [4]^{(2)}$. Now note that if $\alpha_{23}\alpha_{24}\alpha_{34} \in E(\mathcal{A}^{i_2i_3i_4})$ (that is, if there is an edge in $\mathcal{A}^{i_2i_3i_4}$ which intersects $\{\alpha_{23}, \alpha_{24}, \alpha_{34}\}$ in three vertices) then (λ, φ) would actually be a reduced map from $K_4^{(3)}$ to \mathcal{A} . If \mathcal{A} is (d, \bullet) -dense for any $d > 0$, then $\mathcal{A}^{i_2i_3i_4}$ does contain edges (which intersect $\{\alpha_{23}, \alpha_{24}, \alpha_{34}\}$ in zero or more vertices). This leads to the question which density d we need to guarantee that there is an edge in $\mathcal{A}^{i_2i_3i_4}$ which intersects $\{\alpha_{23}, \alpha_{24}, \alpha_{34}\}$ in at least one (or in at least two) vertices. In this sense, moving from the problem for $K_4^{(3)-}$ to the problem for K_4^3 can be described as ‘gluing’ first one, then two, and eventually three of the vertices in the constituent $\mathcal{A}^{i_2i_3i_4}$. The setting in \mathcal{A} with one vertex ‘glued’ is shown in Figure 1.2.

Our second result states that an intersection in at least one vertex (or the first gluing) is already guaranteed if the density is larger than $1/4$ (which is clearly best possible). More precisely, we prove the following.

Theorem 1.2. *For $\varepsilon > 0$, there is some $M \in \mathbb{N}$ such that in every $(\frac{1}{4} + \varepsilon, \bullet)$ -dense reduced hypergraph $\mathcal{A} = (I, \mathcal{P}^{ij}, \mathcal{A}^{ijk})$, there are indices $i_1, i_2, i_3, i_4 \in I$ and vertices $\alpha_{jk} \in \mathcal{P}^{i_ji_k}$ for $jk \in [4]^{(2)}$, as well as vertices $\alpha'_{23} \in \mathcal{P}^{i_2i_3}$ and $\alpha'_{24} \in \mathcal{P}^{i_2i_4}$, such that $\alpha_{13}\alpha_{14}\alpha_{34} \in E(\mathcal{A}^{i_1i_3i_4})$, $\alpha_{12}\alpha_{13}\alpha_{23} \in E(\mathcal{A}^{i_1i_2i_3})$, $\alpha_{12}\alpha_{14}\alpha_{24} \in E(\mathcal{A}^{i_1i_2i_4})$, and $\alpha'_{23}\alpha'_{24}\alpha_{34} \in E(\mathcal{A}^{i_2i_3i_4})$.*

Here, α_{34} corresponds to the vertex resulting from gluing (see Figure 1.2).




FIGURE 1.2. The reduced image of $K_4^{(3)-}$ with an additional edge and one vertex ‘glued’ (α_{34}).

It would be interesting to determine the optimal density d such that we find such a structure with two vertices glued, i.e., only requiring one of the vertices α'_{23} and α'_{24} .

The proofs of Theorem 1.1 and Theorem 1.2 are very similar and could be formulated as one. For clarity however, we give the key steps separately in Section 3 (from Lemma 3.3 onward) and Section 4, respectively.

§2. PRELIMINARIES

When proving $\pi_*(K_4^{(3)-}) = \frac{1}{4}$, Reiher, Rödl, and Schacht [16] used the hypergraph regularity lemma. After “regularising” a hypergraph, the problem reduces to an extremal problem in the reduced hypergraph. We will follow this approach and use the terminology introduced in [16] and laid out systematically by Reiher in his survey [15].

For a hypergraph H , we denote the number of its edges by $e(H) = |E(H)|$ and the number of its vertices by $v(H) = |V(H)|$. Given a set I , we denote by $I^{(\ell)}$ the set of all ℓ -element subsets of I . When considering such subsets, we often omit the parentheses and write for instance ijk for an element of $I^{(3)}$, instead of $\{ijk\}$. For a hypergraph H , we define the shadow of H as $\partial H = \{e \setminus x : x \in e \in E(H)\}$.

Definition 2.1. *Let I be a set of indices, for all $ij \in I^{(2)}$, let \mathcal{P}^{ij} be a set of vertices which are disjoint for $ij \neq i'j'$, and for $ijk \in I^{(3)}$, let \mathcal{A}^{ijk} be a 3-partite 3-graph with partition classes \mathcal{P}^{ij} , \mathcal{P}^{ik} , and \mathcal{P}^{jk} . Then the $\binom{I}{2}$ -partite 3-graph \mathcal{A} with vertex set $V(\mathcal{A}) = \bigcup_{ij \in I^{(2)}} \mathcal{P}^{ij}$ and edge set $E(\mathcal{A}) = \bigcup_{ijk \in I^{(3)}} E(\mathcal{A}^{ijk})$ is called a reduced hypergraph with index set I , vertex classes \mathcal{P}^{ij} and constituents \mathcal{A}^{ijk} .*

For brevity, we often simply write ‘let $\mathcal{A} = (I, \mathcal{P}^{ij}, \mathcal{A}^{ijk})$ be a reduced hypergraph’ instead of ‘let \mathcal{A} be a reduced hypergraph with index set I , vertex classes \mathcal{P}^{ij} and constituents \mathcal{A}^{ijk} ’.

Next, let us take a look at how the density condition in our host hypergraphs translates to a density condition in the respective reduced hypergraphs.

Definition 2.2. Let $\mathcal{A} = (I, \mathcal{P}^{ij}, \mathcal{A}^{ijk})$ be a reduced hypergraph and let $d \in [0, 1]$ be a real number. If $e(\mathcal{A}^{ijk}) \geq d|\mathcal{P}^{ij}||\mathcal{P}^{ik}||\mathcal{P}^{jk}|$ holds for all $ijk \in I^{(3)}$, we say that \mathcal{A} is $(d, \bullet\bullet)$ -dense.

The following definition of a reduced map describes the setting in which a hypergraph F can be embedded into a host hypergraph H (by using the hypergraph regularity method) in terms of the reduced hypergraph of H .

Definition 2.3. A reduced map from a 3-graph F to a reduced hypergraph $\mathcal{A} = (I, \mathcal{P}^{ij}, \mathcal{A}^{ijk})$ is a pair (λ, φ) such that

- (1) $\lambda : V(F) \rightarrow I$ and $\varphi : \partial F \rightarrow V(\mathcal{A})$;
- (2) for $uv \in \partial F$, we have $\lambda(u) \neq \lambda(v)$ and $\varphi(uv) \in \mathcal{P}^{\lambda(u)\lambda(v)}$;
- (3) if $uvw \in E(F)$, then $\varphi(uv)\varphi(uw)\varphi(vw) \in E(\mathcal{A}^{\lambda(u)\lambda(v)\lambda(w)})$.

If there is a reduced map from F to \mathcal{A} , we say that \mathcal{A} contains a reduced image of F , otherwise \mathcal{A} is called F -free.

Now the following theorem due to Reiher, Rödl, and Schacht (and appearing in this formulation in Reiher’s survey [15]) reduces the problem of determining $\pi_{\bullet\bullet}(F)$ to a problem about reduced hypergraphs.

Theorem 2.4. For a 3-graph F , we have

$$\pi_{\bullet\bullet}(F) = \sup\{d \in [0, 1] : \text{For every } M \in \mathbb{N}, \text{ there is a } (d, \bullet\bullet)\text{-dense, } F\text{-free reduced hypergraph } \mathcal{A} \text{ with index set } [M]\}.$$

§3. PROOF OF THEOREM 1.1

Proof of Theorem 1.1. Since $K_4^{(3)-} \subseteq F_\star$, we have $1/4 = \pi_{\bullet\bullet}(K_4^{(3)-}) \leq \pi_{\bullet\bullet}(F_\star)$.

Due to Theorem 2.4, it is thus enough to show that for every $\varepsilon > 0$, there is some large $M \in \mathbb{N}$ such that every $(\frac{1}{4} + \varepsilon, \bullet\bullet)$ -dense reduced hypergraph with index set $[M]$ contains a reduced image of F_\star . Let thus $\varepsilon > 0$ be given, without loss of generality $\varepsilon \ll 1$, and choose $M, M_*, m \in \mathbb{N}$, and $\delta > 0$ such that

$$M^{-1} \ll M_*^{-1} \ll m^{-1} \ll \delta \ll \varepsilon \ll 1. \quad (3.1)$$

Then let \mathcal{A} be a $(\frac{1}{4} + \varepsilon, \bullet\bullet)$ -dense reduced hypergraph with index set $[M]$, vertex classes \mathcal{P}^{ij} and constituents \mathcal{A}^{ijk} . We will show that \mathcal{A} contains a reduced image of F_\star .

We begin with a ‘cleaning process’ (until Lemma 3.2) similar to the one in [16]. However, we will allow for a bit more ‘robustness’. Consider any $ijk \in [M]^{(3)}$ with $i < j < k$. We define the bipartite graph Q_{jk}^i on the vertex set $\mathcal{P}^{ij} \cup \mathcal{P}^{ik}$ by placing an edge between $w \in \mathcal{P}^{ij}$ and $v \in \mathcal{P}^{ik}$ if there are at least $\varepsilon^2 |\mathcal{P}^{jk}|$ vertices u in \mathcal{P}^{jk} such that $wvu \in E(\mathcal{A}^{ijk})$. Similarly, we define the bipartite graph Q_{ij}^k on the vertex set $\mathcal{P}^{ik} \cup \mathcal{P}^{jk}$ by placing an edge between $v \in \mathcal{P}^{ik}$ and $w \in \mathcal{P}^{jk}$ if there are at least $\varepsilon^2 |\mathcal{P}^{ij}|$ vertices u in \mathcal{P}^{ij} such that $uvw \in E(\mathcal{A}^{ijk})$. We write $N_{Q_{jk}^i}(x)$ and $d_{Q_{jk}^i}(x)$ for the neighbourhood and the degree of a vertex x in the graph Q_{jk}^i , respectively, and similarly we define $N_{Q_{ij}^k}(x)$ and $d_{Q_{ij}^k}(x)$.

Lemma 3.1. *For any $ijk \in [M]^{(3)}$ with $i < j < k$, we have*

$$\sum_{v \in \mathcal{P}^{ik}} d_{Q_{jk}^i}(v) d_{Q_{ij}^k}(v) \geq \left(\frac{1}{4} + \frac{\varepsilon}{2}\right) |\mathcal{P}^{ij}| |\mathcal{P}^{jk}| |\mathcal{P}^{ik}|.$$

Proof. We use a double-counting argument. For $v \in \mathcal{P}^{ik}$, set $T(v) = \{w_1 v w_2 \in E(\mathcal{A}^{ijk})\}$ and $S(v) = \{w_1 v w_2 \in E(\mathcal{A}^{ijk}) : vw_1 \in E(Q_{jk}^i), vw_2 \in E(Q_{ij}^k)\}$,

$$\begin{aligned} S_1(v) &= \{w_1 v w_2 \in E(\mathcal{A}^{ijk}) : w_1 \in \mathcal{P}^{ij}, vw_1 \notin E(Q_{jk}^i)\}, \text{ and} \\ S_2(v) &= \{w_1 v w_2 \in E(\mathcal{A}^{ijk}) : w_2 \in \mathcal{P}^{jk}, vw_2 \notin E(Q_{ij}^k)\}. \end{aligned}$$

Note that by double-counting, we have the following.

$$\begin{aligned} \left(\frac{1}{4} + \varepsilon\right) |\mathcal{P}^{ij}| |\mathcal{P}^{jk}| |\mathcal{P}^{ik}| &\leq \sum_{v \in \mathcal{P}^{ik}} |T(v)| \leq \sum_{v \in \mathcal{P}^{ik}} |S(v)| + |S_1(v)| + |S_2(v)| \\ &\leq \sum_{v \in \mathcal{P}^{ik}} d_{Q_{jk}^i}(v) d_{Q_{ij}^k}(v) + 2\varepsilon^2 |\mathcal{P}^{ij}| |\mathcal{P}^{jk}| |\mathcal{P}^{ik}|. \end{aligned}$$

Since $\varepsilon \ll 1$, the assertion follows. \square

Given $ijk \in [M]^{(3)}$ with $i < j < k$, by Lemma 3.1 and the Cauchy-Schwartz inequality, we have either

$$\sum_{v \in \mathcal{P}^{ik}} d_{Q_{jk}^i}(v)^2 \geq \left(\frac{1}{4} + \frac{\varepsilon}{2}\right) |\mathcal{P}^{ij}|^2 |\mathcal{P}^{ik}|, \quad (3.2)$$

in which case we colour ijk blue, and if this is not true, we must have

$$\sum_{v \in \mathcal{P}^{ik}} d_{Q_{ij}^k}(v)^2 \geq \left(\frac{1}{4} + \frac{\varepsilon}{2}\right) |\mathcal{P}^{jk}|^2 |\mathcal{P}^{ik}|,$$

in which case we colour ijk red.

By (3.1) and by Ramsey’s theorem, there is a subset of M_* indices, without loss of generality $[M_*]$, such that all 3-subsets of this set are coloured with the same colour, say blue.

There is one more step in the ‘cleaning process’, for which we apply Ramsey’s theorem to a different colouring of the 3-subsets of $[M_*]$. For $ijk \in [M_*]^{(3)}$ with $i < j < k$ and $r \in \mathbb{N}$, let

$$S_{jk}^i(r) = \{x \in \mathcal{P}^{ik} : d_{Q_{jk}^i}(x) \geq (\frac{1}{2} + r\delta)|\mathcal{P}^{ij}|\}.$$

Since ijk was coloured in blue, (3.2) entails

$$(\frac{1}{4} + \frac{\varepsilon}{2})|\mathcal{P}^{ij}|^2|\mathcal{P}^{ik}| \leq \sum_{v \in \mathcal{P}^{ik}} d_{Q_{jk}^i}(v)^2 \leq (\frac{1}{2} + \delta)^2|\mathcal{P}^{ij}|^2|\mathcal{P}^{ik}| \setminus |S_{jk}^i(1)| + |\mathcal{P}^{ij}|^2|S_{jk}^i(1)|.$$

Because of (3.1), we infer that $|S_{jk}^i(1)| \geq \delta|\mathcal{P}^{ik}|$.

Clearly the $S_{jk}^i(r)$ ’s are decreasing and $S_{jk}^i(r) = \emptyset$ for $r > \frac{1}{2\delta}$, so we may colour ijk by the maximum $r \in [\frac{1}{2\delta}]$ (w.l.o.g., we assume that $\frac{1}{2\delta}$ is an integer) for which $|S_{jk}^i(r)| \geq \delta|\mathcal{P}^{ik}|$.

Again by Ramsey’s theorem, as $m \ll M_*$ and after potentially reordering the indices, we can suppose that $[m]$ is a blue clique in the first coloring and an r_* -coloured clique in the second coloring, for some $r_* \in [\frac{1}{2\delta}]$. The following is the conclusion of our cleaning.

(\star) Every $ijk \in [m]^{(3)}$ with $i < j < k$ satisfies (3.2) and $|S_{jk}^i(r_*)| \geq \delta|\mathcal{P}^{ik}| > |S_{jk}^i(r_* + 1)|$.

Now, for all $i \in [m]$, define the graph $Q^i = \bigcup_{1 \leq i < j < k \leq m} Q_{jk}^i$. We need the following auxiliary result, which is a robust version of a result in [16].

Lemma 3.2. *For $i, j_1, j_2, k \in [m]$ with $i < j_1 < j_2 < k$, and any $x \in S_{j_1 k}^i(r_*) \cap S_{j_2 k}^i(r_*)$, there are at least $\delta|\mathcal{P}^{ij_1}||\mathcal{P}^{ij_2}|$ triangles xyz in Q^i with $y \in \mathcal{P}^{ij_2}, z \in \mathcal{P}^{ij_1}$.*

Proof. Let A_{j_1} be the neighbors of x in \mathcal{P}^{ij_1} , and let A_{j_2} be the neighbors of x in \mathcal{P}^{ij_2} . By definition of x (and S_{jk}^i), we have that $|A_{j_1}| \geq (\frac{1}{2} + r_*\delta)|\mathcal{P}^{ij_1}|, |A_{j_2}| \geq (\frac{1}{2} + r_*\delta)|\mathcal{P}^{ij_2}|$.

Also, denote $B_{j_1} = \mathcal{P}^{ij_1} \setminus A_{j_1}$ and $B_{j_2} = \mathcal{P}^{ij_2} \setminus A_{j_2}$. Now let $C = B_{j_2} \cap S_{j_1 j_2}^i(r_* + 1)$. By (\star), we know $|C| < \delta|\mathcal{P}^{ij_2}|$, and by definition of $S_{j_1 j_2}^i(r_* + 1)$, we have $d_{Q_{j_1 j_2}^i}(y) < (\frac{1}{2} + (r_* + 1)\delta)|\mathcal{P}^{ij_1}|$ for all $y \in B_{j_2} \setminus C$.

Note that any edge between A_{j_1} and A_{j_2} gives rise to a triangle of the kind the statement asks for, so suppose for the sake of contradiction that there are less than $\delta|\mathcal{P}^{ij_1}||\mathcal{P}^{ij_2}|$ edges between A_{j_1} and A_{j_2} .

Then, when setting $d'_{Q_{j_1 j_2}^i}(y) = |N_{Q_{j_1 j_2}^i}(y) \cap A_{j_1}|$ for all $y \in A_{j_2}$, we have $\sum_{y \in A_{j_2}} d'_{Q_{j_1 j_2}^i}(y) < \delta|\mathcal{P}^{ij_1}||\mathcal{P}^{ij_2}|$. Since each $d'_{Q_{j_1 j_2}^i}(y) \leq |\mathcal{P}^{ij_1}|$, we also know $\sum_{y \in A_{j_2}} (d'_{Q_{j_1 j_2}^i}(y))^2 \leq \delta|\mathcal{P}^{ij_1}|^2|\mathcal{P}^{ij_2}|$.

By our conditions on $|A_{j_1}|$, we have $d_{Q_{j_1 j_2}^i}(y) - d'_{Q_{j_1 j_2}^i}(y) \leq |B_{j_1}| \leq (\frac{1}{2} - r_* \delta) |\mathcal{P}^{ij_1}|$. Hence

$$\begin{aligned} \sum_{y \in A_{j_2}} d_{Q_{j_1 j_2}^i}(y)^2 &= \sum_{y \in A_{j_2}} (d_{Q_{j_1 j_2}^i}(y) - d'_{Q_{j_1 j_2}^i}(y) + d'_{Q_{j_1 j_2}^i}(y))^2 \\ &\leq |A_{j_2}| \left(\frac{1}{2} - r_* \delta \right)^2 |\mathcal{P}^{ij_1}|^2 + \delta |\mathcal{P}^{ij_1}|^2 |\mathcal{P}^{ij_2}| + 2 \left(\frac{1}{2} - r_* \delta \right) \delta |\mathcal{P}^{ij_1}|^2 |\mathcal{P}^{ij_2}| \\ &< |A_{j_2}| \left(\frac{1}{2} - r_* \delta \right)^2 |\mathcal{P}^{ij_1}|^2 + 2\delta |\mathcal{P}^{ij_1}|^2 |\mathcal{P}^{ij_2}|. \end{aligned}$$

Now recalling (\star) and splitting the sum yields

$$\begin{aligned} \left(\frac{1}{4} + \frac{\varepsilon}{2} \right) |\mathcal{P}^{ij_1}|^2 |\mathcal{P}^{ij_2}| &\leq \sum_{x \in \mathcal{P}^{ij_2}} d_{Q_{j_1 j_2}^i}(x)^2 \\ &\leq \sum_{x \in B_{j_2} \setminus C} d_{Q_{j_1 j_2}^i}(x)^2 + \sum_{x \in A_{j_2}} d_{Q_{j_1 j_2}^i}(x)^2 + \sum_{x \in C} d_{Q_{j_1 j_2}^i}(x)^2 \\ &\leq |B_{j_2}| \left(\frac{1}{2} + (r_* + 1)\delta \right)^2 |\mathcal{P}^{ij_1}|^2 + |A_{j_2}| \left(\frac{1}{2} - r_* \delta \right)^2 |\mathcal{P}^{ij_1}|^2 + 3\delta |\mathcal{P}^{ij_1}|^2 |\mathcal{P}^{ij_2}|. \end{aligned}$$

Since $|A_{j_2}| + |B_{j_2}| = |\mathcal{P}^{ij_2}|$ and $|B_{j_2}| \leq (\frac{1}{2} - r_* \delta) |\mathcal{P}^{ij_2}|$, the right-hand side is maximised when $|B_{j_2}| = (\frac{1}{2} - r_* \delta) |\mathcal{P}^{ij_2}|$ and $|A_{j_2}| = (\frac{1}{2} + r_* \delta) |\mathcal{P}^{ij_2}|$. Cancelling $|\mathcal{P}^{ij_1}|^2 |\mathcal{P}^{ij_2}|$ entails

$$\begin{aligned} \frac{1}{4} + \frac{\varepsilon}{2} &\leq \left(\frac{1}{2} - r_* \delta \right) \left(\frac{1}{2} + (r_* + 1)\delta \right)^2 + \left(\frac{1}{2} + r_* \delta \right) \left(\frac{1}{2} - r_* \delta \right)^2 + 3\delta \\ &\leq \left(\frac{1}{2} - r_* \delta \right) \left(\frac{1}{2} + r_* \delta \right)^2 + \left(\frac{1}{2} + r_* \delta \right) \left(\frac{1}{2} - r_* \delta \right)^2 + 6\delta \\ &< \frac{1}{4} + 6\delta. \end{aligned}$$

Since $\delta \ll \varepsilon$, this yields a contradiction. \square

In our proof, we will iteratively prepare a number of ‘rows’, each of which is a subgraph of a Q^r , by finding many triangles that are ‘compatible’ with each other. By definition of Q^r , for each edge in $e \in E(Q^r)$, there exists a small subset X in some vertex class of $Q^{r'}$ for some $r' > r$, such that every $x \in X$ together with e forms an edge in \mathcal{A} . After preparing many rows, we will show that some specific of these ‘projections’ have to intersect. From this, the reduced image of F_\star will arise (see Figure 3.1).

The following lemma encapsulates the preparation of one row which is ‘paid for’ by shrinking the index set.

Lemma 3.3. *Let $I \subseteq [m]$ with $|I| > \frac{2}{\delta^2}$, $m \in I$, and $r = \min I$. Then there are $J \subseteq I \setminus r$, with $m \in J$ and $|J| \geq \delta^2 |I|$, $x \in \mathcal{P}^{rm}$, $y \in \mathcal{P}^{r'r}$, where $r' = \min J$, and $z_j \in \mathcal{P}^{rj}$ for all $j \in J \setminus \{r', m\}$ such $xz_j y$ forms a triangle in Q^r .*

Proof. First, we find $x \in \mathcal{P}^{rm}$. As $|S_{jm}^r(r_*)| \geq \delta |\mathcal{P}^{rm}|$ for all $j \in I \setminus \{m, r\}$, it follows that there is some set $I' \subseteq I \setminus \{r, m\}$ such that $|I'| \geq \delta(|I| - 2)$ and some $x \in \bigcap_{j \in I'} S_{jm}^r(r_*)$.

Let $r' = \min I'$; next, we find $y \in \mathcal{P}^{rr'}$. For any $j \in I'$, let A_j be the set of neighbours of x in \mathcal{P}^{rj} . Since $x \in S_{jm}^r(r_*) \cap S_{r'm}^r(r_*)$, by Lemma 3.2, there are at least $\delta|\mathcal{P}^{rj}||\mathcal{P}^{rr'}|$ edges in Q^r between A_j and $A_{r'}$. For $j \in I' \setminus \{r'\}$, let D_j be the set of vertices in $A_{r'}$ that are connected by some edge in Q^r to a vertex in A_j . Then, $|D_j| \geq \frac{\delta|\mathcal{P}^{rj}||\mathcal{P}^{rr'}|}{|\mathcal{P}^{rj}|} = \delta|\mathcal{P}^{rr'}|$ and double-counting yields that there is some vertex $y \in A_{r'}$ that lies in D_j for at least $\frac{\delta|\mathcal{P}^{rr'}|(|I'|-1)}{|\mathcal{P}^{rr'}|} = \delta(|I'|-1)$ such $j \in I'$. Denote the set of such $j \in I'$ by I'' and let $J = I'' \cup \{m, r'\}$.

Now $r' = \min J$ and for all $j \in I'' = J \setminus \{r', m\}$, we have that $y \in D_j$. Hence, there is some vertex $z_j \in A_j$ such that $z_j y$ is an edge in Q^r . Since $z_j \in A_j$ and $y \in A_{r'}$, we know that xz_j and xy are also edges in Q^r . Thus, for all $j \in J \setminus \{r', m\}$, $xz_j y$ is a triangle in Q^r . The lower bound on $|J|$ follows as $|J| \geq \delta(|I'|-1) + 2 \geq \delta^2(|I|-2) - \delta + 2 \geq \delta^2|I|$. \square

Now, we apply Lemma 3.3 iteratively, in each step preparing a row and shrinking the index set. First, let $I_0 = [m]$, and $r_1 = \min I_0$. Lemma 3.3 provides some $I_1 \subseteq I_0 \setminus r_1$ with $m \in I_1$ and $|I_1| \geq \delta^2|I_0|$, $x_1 \in \mathcal{P}^{r_1 m}$, $y_1 \in \mathcal{P}^{r_1 r_2}$, where $r_2 = \min I_1$, and $z_{1j} \in \mathcal{P}^{r_1 j}$ for all $j \in I_1 \setminus \{r_2, m\}$ such that $x_1 z_{1j} y_1$ forms a triangle in Q^{r_1} .

Next, we apply Lemma 3.3 again, but this time to I_1 . Generally, assume that for some $i \in [\frac{2}{\varepsilon^2}]$, we have defined sets $I_0, \dots, I_i \subseteq [m]$, indices $r_1, \dots, r_{i+1} \in [m]$, and for all $j \in [i]$ and $k \in I_j \setminus \{r_{j+1}, m\}$, vertices $x_j \in \mathcal{P}^{r_j m}$, $y_j \in \mathcal{P}^{r_j r_{j+1}}$, and $z_{jk} \in \mathcal{P}^{r_j k}$ such that the following holds. For all $j \in [i]$, we have $I_j \subseteq I_{j-1} \setminus r_j$, $m \in I_j$, $r_{j+1} = \min I_j$, and $|I_j| \geq \delta^2|I_{j-1}|$, and further for all $k \in I_j \setminus \{r_{j+1}, m\}$, $x_j z_{jk} y_j$ forms a triangle in Q^{r_j} . Then applying Lemma 3.3 to I_i yields $I_{i+1} \subseteq I_i \setminus r_{i+1}$ with $m \in I_{i+1}$ and $|I_{i+1}| \geq \delta^2|I_i|$, $x_{i+1} \in \mathcal{P}^{r_{i+1} m}$, $y_{i+1} \in \mathcal{P}^{r_{i+1} r_{i+2}}$, where $r_{i+2} = \min I_{i+1}$, and $z_{i+1j} \in \mathcal{P}^{r_{i+1} j}$ for all $j \in I_{i+1} \setminus \{r_{i+2}, m\}$ such that $x_{i+1} z_{i+1j} y_{i+1}$ forms a triangle in $Q^{r_{i+1}}$.

Note that we can perform this process $\frac{2}{\varepsilon^2} + 1$ times since in the last step (and thus in all steps prior) we have that $|I_{\frac{2}{\varepsilon^2}}| \geq (\delta^2)^{\frac{2}{\varepsilon^2}} m > \frac{2}{\delta^2}$ (using the hierarchy (3.1)) and can thus we can still construct the set $I_{\frac{2}{\varepsilon^2}+1}$ with $|I_{\frac{2}{\varepsilon^2}+1}| \geq 3$.

After this iteration procedure, let the maximum element of $I_{\frac{2}{\varepsilon^2}+1} \setminus m$ be m' . For all $i \in [\frac{2}{\varepsilon^2} + 1]$, since $I_{\frac{2}{\varepsilon^2}+1} \subseteq I_i$, there is some triangle $x_i z_{im'} y_i$ in Q^{r_i} and we set $e_i = x_i z_{im'}$. Note that e_i is between $\mathcal{P}^{r_i m}$ and $\mathcal{P}^{r_i m'}$. Since $e_i \in E(Q^{r_i})$, we know that there is a set $H(e_i) \subseteq \mathcal{P}^{m'm}$ with $|H(e_i)| \geq \varepsilon^2 |\mathcal{P}^{m'm}|$ such that $x_i z_{im'} v \in E(\mathcal{A}^{r_i m' m})$ for every $v \in H(e_i)$. Thus, there are at least three indices $i, j, k \in [\frac{2}{\varepsilon^2} + 1]$ such that $H(e_i) \cap H(e_j) \cap H(e_k) \neq \emptyset$. Without loss of generality, assume $i < j < k$.

Let $v \in H(e_i) \cap H(e_j) \cap H(e_k)$ and note that $r_k > r_j \geq r_{i+1}$ and that since $|I_{\frac{2}{\varepsilon^2}+1}| \geq 3$ and $m = \max I_{\frac{2}{\varepsilon^2}+1}$, $m' = \max I_{\frac{2}{\varepsilon^2}+1} \setminus m$, we know $r_k < m'$. Thus, there are $z_{im'} \in \mathcal{P}^{r_i m'}$ and $z_{ir_k} \in \mathcal{P}^{r_i r_k}$ such that $x_i z_{im'} y_i$ is a triangle in Q^{r_i} and $z_{ir_k} y_i \in E(Q^{r_i})$ (we do not




FIGURE 3.1. F_\star embedded across several prepared rows in the reduced hypergraph \mathcal{A} . Blue lines represent edges in Q^r .

need the other edges in the respective triangle). Now these edges imply the existence of vertices $u_1 \in \mathcal{P}^{r_{i+1}m}$, $u_2 \in \mathcal{P}^{r_{i+1}m'}$, $u_3 \in \mathcal{P}^{r_{i+1}r_k}$ such that $x_i y_i u_1 \in E(\mathcal{A}^{r_i r_{i+1} m})$, $z_{im'} y_i u_2 \in E(\mathcal{A}^{r_i r_{i+1} m'})$, and $z_{ir_k} y_i u_3 \in E(\mathcal{A}^{r_i r_{i+1} r_k})$. Together with the edges $x_i z_{im'} v \in E(\mathcal{A}^{r_i m' m})$ and $x_k z_{km'} v \in E(\mathcal{A}^{r_k m' m})$, this yields a reduced embedding of F_\star . This iteration process and the resulting embedding is illustrated in Figure 3.1.

Indeed, as shown in Figure 3.2, the edges mentioned show that when setting $\lambda(1) = r_i$, $\lambda(2) = r_{i+1}$, $\lambda(3) = m$, $\lambda(4) = m'$, and $\lambda(5) = r_k$ and $\varphi(12) = y_i$, $\varphi(13) = x_i$, $\varphi(14) = z_{im'}$, $\varphi(15) = z_{ir_k}$, $\varphi(23) = u_1$, $\varphi(24) = u_2$, $\varphi(25) = u_3$, $\varphi(34) = v$, $\varphi(35) = x_k$, and $\varphi(45) = z_{km'}$, then (λ, φ) is a reduced embedding of F_\star into \mathcal{A} .

□

§4. PROOF OF THEOREM 1.2

Proof. Let $\varepsilon > 0$ be given, without loss of generality, $\varepsilon \ll 1$, and choose $M, M_*, m \in \mathbb{N}$ and $\delta > 0$ such that

$$M^{-1} \ll M_*^{-1} \ll m^{-1} \ll \delta \ll \varepsilon \ll 1. \quad (4.1)$$

Then let \mathcal{A} be a $(\frac{1}{4} + \varepsilon)$ -dense reduced hypergraph with index set $[M]$, vertex classes \mathcal{P}^{ij} and constituents \mathcal{A}^{ijk} . We need to show that there are four indices $i_1, i_2, i_3, i_4 \in [M]$ and




FIGURE 3.2. The reduced image of F_* in the reduced hypergraph \mathcal{A} . As before, the gray lines between two indices i and j represent the vertex class \mathcal{P}^{ij} .

vertices $x_{\alpha\beta} \in \mathcal{P}^{i\alpha i\beta}$ for $\alpha\beta \in [4]^{(2)}$ as well as $x'_{23} \in \mathcal{P}^{i2i3}$ and $x'_{24} \in \mathcal{P}^{i2i4}$ such that $x_{12}x_{13}x_{23} \in E(\mathcal{A}^{i1i2i3})$, $x_{12}x_{14}x_{24} \in E(\mathcal{A}^{i1i2i4})$, $x_{13}x_{14}x_{34} \in E(\mathcal{A}^{i1i3i4})$, and $x'_{23}x'_{24}x_{34} \in E(\mathcal{A}^{i2i3i4})$.

We proceed as in the last section until Lemma 3.3. Instead of this lemma, we need the following variant.

Lemma 4.1. *Let $m_1, m_2 \in \mathbb{N}$ with $m_1 \geq \frac{m_2+1}{\delta^{m_2}}$ and let $I \subseteq [m]$ with $|I| \geq m_1$, $m \in I$ and $r = \min I$. Then there are $J \subseteq I \setminus r$ with $m \in J$ and $|J| \geq m_2$, $x \in \mathcal{P}^{rm}$ and $z_k \in \mathcal{P}^{rk}$ for all $k \in J \setminus \{r', m\}$, where $r' = \min J$, such that for all $jk \in (J \setminus m)^{(2)}$ with $j < k$, there is some $y \in \mathcal{P}^{rj}$ such that xz_ky forms a triangle in Q^r .*

Proof. We will choose the z_k iteratively and with each choice the set of considered indices will shrink. As $|S_{jm}^r(r_*)| \geq \delta|\mathcal{P}^{rm}|$ for all $j \in I \setminus \{m, r\}$, it follows that there is some set $I' \subseteq I \setminus \{r, m\}$ such that $|I'| \geq \delta(|I| - 2)$ and some $x \in \bigcap_{j \in I'} S_{jm}^r(r_*)$. For any $k \in I'$, let A_k be the set of neighbours of x in \mathcal{P}^{rk} . Since for all $jk \in I'^{(2)}$ with $j < k$, we have $x \in S_{km}^r(r_*) \cap S_{jm}^r(r_*)$, by Lemma 3.2, there are at least $\delta|\mathcal{P}^{rk}||\mathcal{P}^{rj}|$ edges in Q^r between A_k and A_j . Let now $k_1 = \max I'$ and let D_j be the set of vertices in \mathcal{P}^{rk_1} which lie in a triangle between A_{k_1} and A_j . Note that $|D_j| \geq \delta|\mathcal{P}^{rk_1}|$. Therefore, there is some set $I'_1 \subseteq I' \setminus k_1$ with $|I'_1| \geq \delta(|I'| - 1)$ and some $z_{k_1} \in \bigcap_{j \in I'_1} D_j$. Set $I_1 = I'_1 \cup \{k_1, m\}$ and note that $|I_1| \geq \delta^2|I|$ and that for all $j \in I_1$ with $j < k_1$, there is some $y \in \mathcal{P}^{rj}$ such that $xz_{k_1}y$ forms a triangle in Q^r .

Now suppose that we have found indices $m > k_1 > \dots > k_i$ in I for some $i < m_2$, a set $I_i \subseteq I \setminus r$ with $\{m, k_1, \dots, k_i\} \subseteq I_i$ and $|I_i| \geq \delta^{i+1}|I|$, and vertices $z_k \in \mathcal{P}^{rk}$ for all $k \in \{k_1, \dots, k_i\}$ such that for all $k \in \{k_1, \dots, k_i\}$ and $j \in I_i$ with $j < k$, there is

some $y \in \mathcal{P}^{rj}$ such that xz_ky forms a triangle in Q^r . Then let $I'_{i+1} = I_i \setminus \{m, k_1, \dots, k_i\}$ and note that $|I'_{i+1}| \geq \delta^{i+1}m_1 - (i+1) \geq 1$. Further, let $k_{i+1} = \max I'_{i+1}$ and for $j \in I'_{i+1} \setminus k_{i+1}$, let D'_j be the set of vertices in $\mathcal{P}^{rk_{i+1}}$ which lie in a triangle between $A_{k_{i+1}}$ and A_j . As before, we have that $|D'_j| \geq \delta|\mathcal{P}^{rk_{i+1}}|$ and thus, there is some set $I''_{i+1} \subseteq I'_{i+1} \setminus k_{i+1}$ and some $z_{k_{i+1}} \in \bigcap_{j \in I''_{i+1}} D'_j$ (if $I''_{i+1} = \emptyset$, pick $z_{k_{i+1}} \in \mathcal{P}^{rk_{i+1}}$ arbitrarily). Set $I_{i+1} = I''_{i+1} \cup \{m, k_1, \dots, k_{i+1}\}$ and note that

$$|I_{i+1}| \geq \delta(|I_i| - (i+2)) + i+2 \geq \delta|I_i| \geq \delta^{i+2}|I|,$$

and that for all $k \in \{k_1, \dots, k_{i+1}\}$ and $j \in I_{i+1}$ with $j < k$, there is some $y \in \mathcal{P}^{rj}$ such that xz_ky forms a triangle in Q^r .

Eventually, we set $J = \{m, k_1, \dots, k_{m_2}\}$ and note that J , x , and z_k for $k \in J$ have the asserted properties. \square

Similarly as in the last section, we will now conclude the proof by an iteration process, this time repeatedly applying Lemma 4.1. Let $m_1, \dots, m_{\frac{1}{\varepsilon^2}+1} \in \mathbb{N}$ with

$$m^{-1} = m_1^{-1} \ll m_2^{-1} \ll \dots \ll m_{\frac{1}{\varepsilon^2}+1} \ll \delta, \quad (4.2)$$

and note that this is compatible with the hierarchy (4.1). Set $I_0 = [m]$, $r_1 = 1$ (note that we will use I_i here in a different sense than in the proof of Lemma 4.1 but essentially in the same sense as in the proof of Theorem 1.1). Generally, assume that we have defined, for some $i \in [\frac{1}{\varepsilon^2}]$, sets $I_0, \dots, I_i \subseteq [m]$, indices $r_1, \dots, r_{i+1} \in [m]$, and for each $i' \in [i]$ and $k \in I_{i'} \setminus \{r_{i'+1}, m\}$, vertices $x_{i'} \in \mathcal{P}^{r_{i'}m}$ and $z_{i'k} \in \mathcal{P}^{r_{i'}k}$ such that the following holds. For all $i' \in [i]$, we have $I_{i'} \subseteq I_{i'-1} \setminus r_{i'}$, $m \in I_{i'}$, $r_{i'+1} = \min I_{i'}$, and $|I_{i'}| \geq m_{i'}$, and further for all $jk \in (I_{i'} \setminus m)^{(2)}$ with $j < k$, there is some $y \in \mathcal{P}^{r_{i'}j}$ such that $x_{i'}z_{i'k}y$ forms a triangle in $Q^{r_{i'}}$.

Then we apply Lemma 4.1 with I_i instead of I , and m_i and m_{i+1} in place of m_1 and m_2 , respectively. This yields a set $I_{i+1} \subseteq I_i \setminus r_{i+1}$ with $m \in I_{i+1}$ and $|I_{i+1}| \geq m_{i+1}$, a vertex $x_{i+1} \in \mathcal{P}^{r_{i+1}m}$, and for all $k \in I_{i+1} \setminus \{r_{i+2}, m\}$, where $r_{i+2} = \min I_{i+1}$, vertices $z_{i+1k} \in \mathcal{P}^{r_{i+1}k}$ such that for all $jk \in (I_{i+1} \setminus m)^{(2)}$ with $j < k$, there is some $y \in \mathcal{P}^{r_{i+1}j}$ such that $x_{i+1}z_{i+1k}y$ forms a triangle in $Q^{r_{i+1}}$.

Due to (4.2), we can proceed in this manner until $i = \frac{1}{\varepsilon^2} + 1$, that is, we can construct a set $I_{\frac{1}{\varepsilon^2}+1}$ with $|I_{\frac{1}{\varepsilon^2}+1}| \geq m_{\frac{1}{\varepsilon^2}+1}$ in this manner. Let $m' = \max I_{\frac{1}{\varepsilon^2}+1} \setminus m$. During the iteration, we found for each $i \in [\frac{1}{\varepsilon^2} + 1]$ vertices x_i and $z_{im'}$ such that, in particular, $x_i z_{im'} \in E(Q^{r_i})$. We set $e_i = x_i z_{im'}$. By definition of Q^{r_i} , there is a set $H(e_i) \subseteq \mathcal{P}^{m'm}$ with $|H(e_i)| \geq \varepsilon^2 |\mathcal{P}^{m'm}|$ such that $x_i z_{im'} v \in E(\mathcal{A}^{r_i m' m})$ for every $v \in H(e_i)$. Thus there are two indices $i, j \in [\frac{1}{\varepsilon^2} + 1]$ such that $H(e_i) \cap H(e_j) \neq \emptyset$. W.l.o.g., assume $i < j$.

Let $v \in H(e_i) \cap H(e_j)$ and let $y \in \mathcal{P}^{r_i r_j}$ such that $x_i z_{im'} y$ forms a triangle in Q_{r_i} (such a y exists due to the choice of x_i and $z_{im'}$). The definition of Q^{r_i} implies that




FIGURE 4.1. $K_4^{(3)-}$ with one vertex glued embedded in the reduced hypergraph \mathcal{A} that we obtain, following the procedure above.

there are vertices $u_1 \in \mathcal{P}^{r_j m}$ and $u_2 \in \mathcal{P}^{r_j m'}$ such that $x_i y u_1 \in E(\mathcal{A}^{r_i r_j m})$ and $z_{im'} y u_2 \in E(\mathcal{A}^{r_i r_j m'})$. Since we also have the edges $x_i z_{im'} v \in E(\mathcal{A}^{r_i m' m})$ and $x_j z_{jm'} v \in E(\mathcal{A}^{r_j m' m})$, the indices r_i, r_j, m' , and m and the vertices $x_i, z_{im'}, y, x_j, u_1, z_{jm'}, u_2$, and v form the desired configuration. This is illustrated in Figure 4.1. \square

Acknowledgements. The second author thanks Christian Reiher and Mathias Schacht for fruitful discussions and for teaching him [16].

REFERENCES

- [1] J. Balogh, F. C. Clemen, and B. Lidický, *Hypergraph Turán problems in ℓ_2 -norm*, Surveys in combinatorics 2022, London Math. Soc. Lecture Note Ser., vol. 481, Cambridge Univ. Press, Cambridge, 2022, pp. 21–63. MR4421399 $\uparrow 1$
- [2] L. Bellmann and Chr. Reiher, *Turán's theorem for the Fano plane*, Combinatorica **39** (2019), no. 5, 961–982, DOI 10.1007/s00493-019-3981-8. MR4039597 $\uparrow 1$
- [3] S. Berger, S. Piga, Chr. Reiher, V. Rödl, and M. Schacht, *Turán density of cliques of order five in 3-uniform hypergraphs with quasirandom links*, posted on 2022, DOI 10.48550/ARXIV.2206.07354. $\uparrow 1$
- [4] M. Bucić, J. W. Cooper, D. Král, S. Mohr, and D. M. Correia, *Uniform Turán density of cycles*, posted on 2021, DOI 10.48550/ARXIV.2112.01385. $\uparrow 1$
- [5] D. De Caen and Z. Füredi, *The maximum size of 3-uniform hypergraphs not containing a Fano plane*, J. Combin. Theory Ser. B **78** (2000), no. 2, 274–276, DOI 10.1006/jctb.1999.1938. MR1750899 $\uparrow 1$

- [6] P. Erdős, *Problems and results on graphs and hypergraphs: similarities and differences*, Mathematics of Ramsey theory, Algorithms Combin., vol. 5, Springer, Berlin, 1990, pp. 12–28, DOI [10.1007/978-3-642-72905-8_2](https://doi.org/10.1007/978-3-642-72905-8_2). MR1083590 [↑1](#)
- [7] P. Erdős and M. Simonovits, *A limit theorem in graph theory*, Studia Sci. Math. Hungar. **1** (1966), 51–57. MR205876 [↑1](#)
- [8] P. Erdős and V. T. Sós, *On Ramsey-Turán type theorems for hypergraphs*, Combinatorica **2** (1982), no. 3, 289–295, DOI [10.1007/BF02579235](https://doi.org/10.1007/BF02579235). MR698654 [↑1](#)
- [9] P. Erdős and A. H. Stone, *On the structure of linear graphs*, Bull. Amer. Math. Soc. **52** (1946), 1087–1091, DOI [10.1090/S0002-9904-1946-08715-7](https://doi.org/10.1090/S0002-9904-1946-08715-7). MR18807 [↑1](#)
- [10] Z. Füredi and M. Simonovits, *Triple systems not containing a Fano configuration*, Combin. Probab. Comput. **14** (2005), no. 4, 467–484, DOI [10.1017/S0963548305006784](https://doi.org/10.1017/S0963548305006784). MR2160414 [↑1](#)
- [11] R. Glebov, D. Král', and J. Volec, *A problem of Erdős and Sós on 3-graphs*, Israel J. Math. **211** (2016), no. 1, 349–366, DOI [10.1007/s11856-015-1267-4](https://doi.org/10.1007/s11856-015-1267-4). MR3474967 [↑1](#)
- [12] G. Katona, T. Nemetz, and M. Simonovits, *On a problem of Turán in the theory of graphs*, Mat. Lapok **15** (1964), 228–238 (Hungarian, with English and Russian summaries). MR172263 [↑1](#)
- [13] P. Keevash, *Hypergraph Turán problems*, Surveys in combinatorics 2011, London Math. Soc. Lecture Note Ser., vol. 392, Cambridge Univ. Press, Cambridge, 2011, pp. 83–139. MR2866732 [↑1](#)
- [14] P. Keevash and B. Sudakov, *The Turán number of the Fano plane*, Combinatorica **25** (2005), no. 5, 561–574, DOI [10.1007/s00493-005-0034-2](https://doi.org/10.1007/s00493-005-0034-2). MR2176425 [↑1](#)
- [15] Chr. Reiher, *Extremal problems in uniformly dense hypergraphs*, European J. Combin. **88** (2020), 103117, 22, DOI [10.1016/j.ejc.2020.103117](https://doi.org/10.1016/j.ejc.2020.103117). MR4111729 [↑1](#), [1](#), [2](#), [2](#)
- [16] Chr. Reiher, V. Rödl, and M. Schacht, *On a Turán problem in weakly quasirandom 3-uniform hypergraphs*, J. Eur. Math. Soc. (JEMS) **20** (2018), no. 5, 1139–1159, DOI [10.4171/JEMS/784](https://doi.org/10.4171/JEMS/784). MR3790065 [↑1](#), [2](#), [3](#), [3](#), [4](#)
- [17] ———, *Hypergraphs with vanishing Turán density in uniformly dense hypergraphs*, J. Lond. Math. Soc. (2) **97** (2018), no. 1, 77–97, DOI [10.1112/jlms.12095](https://doi.org/10.1112/jlms.12095). MR3764068 [↑1](#)
- [18] P. Turán, *Eine Extremalaufgabe aus der Graphentheorie*, Mat. Fiz. Lapok **48** (1941), 436–452 (Hungarian, with German summary). MR18405 [↑1](#)

CALIFORNIA INSTITUTE OF TECHNOLOGY

Email address: augustc@alumni.caltech.edu

CALIFORNIA INSTITUTE OF TECHNOLOGY

Email address: schuelke@caltech.edu