

# Mean-field phase transitions in TGFT quantum gravity

Luca Marchetti,<sup>1,\*</sup> Daniele Oriti,<sup>1,2,†</sup> Andreas G. A. Pithis,<sup>1,2,‡</sup> and Johannes Thürigen<sup>3,4,§</sup>

<sup>1</sup>*Arnold Sommerfeld Center for Theoretical Physics, Ludwig-Maximilians-Universität München, Theresienstr. 37, 80333 München, Germany, EU*

<sup>2</sup>*Munich Center for Quantum Science and Technology (MCQST), Schellingstr. 4, 80799 München, Germany, EU*

<sup>3</sup>*Mathematisches Institut der Westfälischen Wilhelms-Universität, Einsteinstr. 62, 48149 Münster, Germany, EU*

<sup>4</sup>*Institut für Physik/Institut für Mathematik, Humboldt-Universität zu Berlin, Unter den Linden 6, 10099 Berlin, Germany, EU*

Controlling the continuum limit and extracting effective gravitational physics is a shared challenge for quantum gravity approaches based on quantum discrete structures. The description of quantum gravity in terms of tensorial group field theory (TGFT) has recently led to much progress in its application to phenomenology, in particular cosmology. This application relies on the assumption of a phase transition to a nontrivial vacuum (condensate) state describable by mean-field theory, an assumption that is difficult to corroborate by a full RG flow analysis due to the complexity of the relevant TGFT models. Here we demonstrate that this assumption is justified due to the specific ingredients of realistic quantum geometric TGFT models: combinatorially non-local interactions, matter degrees of freedom and Lorentz group data together with the encoding of micro-causality. This greatly strengthens the evidence for the existence of a meaningful continuum gravitational regime in group-field and spin foam quantum gravity, the phenomenology of which is amenable to explicit computations in a mean-field approximation.

The main challenge for quantum gravity, in all its candidate formulations [1–17], is to show the existence of a regime of the fundamental theory which reproduces gravitational physics together with new observable consequences. This is highly non-trivial because we expect the new quantum gravity effects to become dominant at rather extreme energy and length scales. Still, early-universe cosmology, black holes and several astrophysical phenomena might provide many testing grounds for quantum gravity, also thanks to the recent improvement of observational techniques [18]. In quantum gravity formulations based on fundamentally discrete quantum structures, the challenge of extracting an effective continuum gravitational physics is intertwined with that of controlling the continuum limit of the fundamental quantum dynamics, i.e. its renormalization group flow and its continuum phase structure.

Black hole [19, 20] and cosmological [21–24] physics have been studied in the framework of tensorial group field theory (TGFT) [14–16]. This is a quantum field theory (QFT) generating spacetime geometries from discrete geometric building blocks given by combinatorially non-local interactions. This framework provides also a completion or reformulation of the quantum dynamics encoded in lattice gravity path integrals and spin foam models [9, 10, 25] (a covariant counterpart of canonical loop quantum gravity [12, 13]).

The underlying assumption in all these works is that there is a condensate phase, the dynamics of which can be captured by a mean-field approximation. However, this assumption is difficult to corroborate by a full RG flow analysis due to the complexity of the relevant TGFT models. On the other hand, this formal complexity of the fundamental quantum dynamics is the result of trying to incorporate the conditions required for a geometric in-

terpretation at the discrete level, some seed of the causal structure we expect to emerge in the continuum approximation, and appropriate matter degrees of freedom.

In this letter we show that an appropriate mean-field description of a condensate phase generically exists in TGFT and that the quantum geometric and physical ingredients even improve the mean-field theory behaviour. To this aim, we apply a Landau-Ginzburg analysis, in particular the Ginzburg criterion, to models with (1) non-local interactions [26, 27], (2) additional matter degrees of freedom [28] as well as (3) the Lorentz group  $SL(2, \mathbb{C})$  together with the implementation of geometricity constraints [29]. We further use the notion of effective dimension coming from renormalization group results [30, 31] compatible with the findings here.

Our results thus strongly support the existence of a meaningful continuum gravitational regime in TGFT quantum gravity as well as the closely related spin foam models and lattice quantum gravity, the phenomenology of which is accessible to explicit computations in a mean-field approximation. They also confirm, thus, how the TGFT formulation provides new powerful tools for tackling the difficult problem of the continuum limit in lattice quantum gravity and the quantum dynamics of spin networks states in the canonical loop quantum gravity approach.

## MEAN-FIELD TGFT

TGFT is a field theory which perturbatively generates a sum over lattices to which group-theoretic data are associated, which encode their discrete geometry [15, 16]. The field  $\Phi : G^r \rightarrow \mathbb{R}$  excites  $r$  elements  $\mathbf{g} = (g^1, \dots, g^r)$  in a Lie group  $G$ . From the Feynman diagrams one obtains

cell complexes due to combinatorially non-local interactions  $\prod_i^n \Phi(\mathbf{g}_i)$  in which the field arguments convolute pairwise, one  $g_i^a$  with one other  $g_j^b$ . This is encoded by an  $r$ -valent interaction graph  $\gamma$  with  $n_\gamma = n$  vertices and  $r \cdot n/2$  edges  $(i, a; j, b)$ . A generic action is thus

$$S[\Phi] = \int_{G^r} d\mathbf{g} \Phi(\mathbf{g}) \mathcal{K} \Phi(\mathbf{g}) + \sum_\gamma \frac{\lambda_\gamma}{n_\gamma} \text{Tr}_\gamma [\Phi] \quad (1)$$

where  $\mathcal{K}$  is a kinetic operator,  $\sum_\gamma$  is over a given set of interaction graphs  $\gamma$  and  $\text{Tr}_\gamma$  defines the resulting pairwise convolutions of each  $\gamma$  with kernels  $\mathcal{V}$ ,

$$\text{Tr}_\gamma [\Phi] = \int_{G^{r \cdot n_\gamma}} \prod_{i=1}^{n_\gamma} d\mathbf{g}_i \prod_{(i,a;j,b)} \mathcal{V}(g_i^a, g_j^b) \prod_{i=1}^{n_\gamma} \Phi(\mathbf{g}_i). \quad (2)$$

If in each convolution  $a = b$ , this integer  $a$  labels the edges yielding an  $r$ -coloured graph (related to tensorial symmetry [17, 32]). The resulting Feynman diagrams are then dual to  $r$ -dimensional simplicial pseudo-manifolds.

To associate to the simplicial structures a discrete geometry (in terms of connection variables), additional geometricity constraints on the group elements (called closure and simplicity constraints) are necessary [15, 16]. With such constraints in place, the TGFT amplitude on a given simplicial lattice takes the form of a lattice gravity path integral (or, equivalently, of a spin foam state sum) [10, 25] (different models correspond to different implementations of such constraints i.e. a choice of kinetic operator  $\mathcal{K}$  and interaction convolutions  $\mathcal{V}$ ). TGFT thus differs significantly from local QFT in technical details and interpretation. Still, QFT methods can be adapted to this peculiar quantum gravity framework.

Mean-field theory provides an approximation of the full QFT partition function and thus an effective description of the phase structure [33]. One considers Gaussian fluctuations around a non-vanishing vacuum solution, e.g. for a constant mean-field  $\Phi_0$ . This is a (real) solution to the TGFT equation of motion [28]

$$\mathcal{K} \Phi + \sum_\gamma \frac{\lambda_\gamma}{n_\gamma} \sum_v \text{Tr}_{\gamma \setminus v} (\Phi) = 0 \quad (3)$$

where the trace is over the graph  $\gamma \setminus v$  obtained by deleting the vertex  $v$  from  $\gamma$  and  $\sum_v$  runs over all its vertices  $v$ .

The validity of a mean-field description of phase transitions with diverging correlation length  $\xi$ , can be checked via the Ginzburg criterion. The ratio

$$Q := \frac{\int_{\Omega_\xi} d\mathbf{g} C(\mathbf{g})}{\int_{\Omega_\xi} d\mathbf{g} \Phi_0^2} \quad (4)$$

compares correlations  $C(\mathbf{g})$  of Gaussian fluctuations with the vacuum  $\Phi_0^2$  both averaged up to the correlation length scale  $\xi$ , that is integrated over a suitable domain  $\Omega_\xi$  [28]. If fluctuations remain small towards the phase transition, i.e.  $Q \ll 1$  when  $\xi \rightarrow \infty$ , mean-field theory is valid.





double-trace		$\hat{\mathcal{X}} = 4(2 \prod_{c=1}^4 \delta_{j_c 0} + 1)$
$n = 4$ melonic		$\hat{\mathcal{X}} = 4(\prod_c \delta_{j_c 0} + \prod_{b \neq c} \delta_{j_b 0} + \delta_{j_c 0})$
$n = 4$ necklace		$\hat{\mathcal{X}} = 4(\prod_c \delta_{j_c 0} + \delta_{j_1 0} \delta_{j_2 0} + \delta_{j_3 0} \delta_{j_4 0})$
simplicial		$\hat{\mathcal{X}} = 5 \sum_{i=0}^4 \prod_{k \neq i} \delta_{j_{(ik)} 0}$

TABLE I. Examples of non-local interaction graphs for  $r = 4$  and the resulting operator  $\hat{\mathcal{X}}_\gamma$  [28]. Additional to the usual legs (red) of interacting fields  $\Phi$ , green half-edges represent the pairwise convolution of group arguments  $g_i^a$  [34].

In local  $d$ -dimensional  $\Phi_d^4$  scalar field theory,  $Q \sim \xi^{4-d}$  such that the mean-field description is valid only beyond the critical dimension  $d_{\text{crit}} = 4$ . In the following, we explain how this is affected by the various features of TGFT models, progressively including all main ingredients for realistic quantum gravity models.

## 1) COMBINATORIAL NON-LOCALITY

The effect of combinatorially non-local interactions is most transparent for a simplified TGFT model. We choose  $\mathcal{K} = \sum_{c=1}^r -\Delta_c + \mu$  with  $\Delta_c$  being the Laplacian on  $G$  and Dirac delta convolutions  $\mathcal{V} = \delta$ . Then the equation of motions (3) for the constant field  $\Phi_0$  is

$$\left( \mu + \sum_\gamma \lambda_\gamma V_G^{r \frac{n_\gamma - 2}{2}} \Phi_0^{n_\gamma - 2} \right) \Phi_0 = 0 \quad (5)$$

where  $V_G$  is the (formal) volume of  $G$ . Such factors arise due to the combinatorial non-locality and need regularization if  $G$  is not compact. For the example of this section we take  $G = \mathbb{R}^{d_c}$  and regularize it to the  $d_c$ -torus  $G_L = \mathbb{T}_L^{d_c}$  with radii  $L/2\pi$  such that  $V_L \equiv V_{G_L} = L^{d_c}$ .

For a single interaction  $\gamma$ , the vacuum equation Eq. (5) reduces to  $(V_L^{r/2} \Phi_0)^{n_\gamma - 2} = -\frac{\mu}{n_\gamma \lambda_\gamma}$  which has real solution for  $\mu < 0$ . The correlation function of Gaussian fluctuations around such  $\Phi_0$  in momentum space is [28]

$$\hat{C}(\mathbf{j}) = \frac{1}{\frac{1}{V_L^2} \sum_c \text{Cas}_{j_c} + \mu - \mu \hat{\mathcal{X}}_\gamma(\mathbf{j})} = \frac{1}{\frac{1}{V_L^2} \sum_c \text{Cas}_{j_c} + b_j} \quad (6)$$

where  $j_c$  labels representations of  $G$  and  $\mathbf{j} \equiv (j_1, \dots, j_r)$ . On a compact (ified) group these are countable, here  $j_c \in \mathbb{Z}^{d_c}$ . Moreover,  $\hat{\mathcal{X}}_\gamma(\mathbf{j})$  is a sum over products of Kronecker deltas specific to  $\gamma$  (see Tab. I) which gives rise to an effective mass  $b_j = \mu(1 - \hat{\mathcal{X}}_\gamma(\mathbf{j}))$ . Consequently, cor-

relations expand in various multiplicities of zero modes,

$$C(\mathbf{g}) = \frac{1}{V_L^r} \sum_{s=0}^r \sum_{(c_1, \dots, c_s)} \sum_{\substack{j_{c_1}, \dots, j_{c_s} = 0 \\ j_{c_{s+1}}, \dots, j_{c_r} \neq 0}} \text{tr}_j \left[ \hat{C}(\mathbf{j}) \bigotimes_{c=1}^r D^{j_c}(g_c) \right], \quad (7)$$

with representation matrices  $D^{j_c}$ , here  $D^{j_c}(e^{i\theta}) = e^{i\theta \cdot j_c}$ . Since  $D^0(g) = 1$ , each  $s$ -fold zero-mode  $(c_1, \dots, c_s)$  contribution depends only on the other  $r - s$  group variables and the effective mass reduces to a number  $b_j = b_{c_1, \dots, c_s}$ .

The correlation length  $\xi$  can be obtained from the second moment or asymptotic behaviour of  $C(\mathbf{g})$  [28, 29]. It sets the characteristic scale beyond which correlations decay exponentially and diverges as  $\xi^2 \sim -1/\mu$  at criticality, i.e. when  $\mu \rightarrow 0$ . For large cut-off  $L$ , integrating  $C(\mathbf{g})$  up to  $\xi$  in each parameter  $\theta_c^a$  yields [28]

$$\int_{\Omega_\xi} d\mathbf{g} C(\mathbf{g}) = \sum_{s=s_0}^r \left( \frac{\xi}{L} \right)^{d_G s} \sum_{(c_1, \dots, c_s)} \frac{1}{b_{c_1 \dots c_s}} \quad (8)$$

where  $s_0$  is the minimal number of zero modes, that is deltas in  $\hat{\mathcal{X}}$ . Interaction graphs  $\gamma$  have multiple edges in general; it is the maximal multiplicity  $s_{\max}$  occurring in  $\gamma$  which determines  $s_0$ ; in the model here  $s_0 = r - s_{\max}$ .

Taking the ratio (4) with integrated  $\Phi_0^2$  gives

$$Q_L = \left( \frac{\lambda_\gamma}{-\mu} \right)^{\frac{2}{n_\gamma - 2}} \sum_{s=s_0}^r \frac{f_s^\gamma}{-\mu} \left( \frac{\xi}{L} \right)^{d_G(s-r)} \quad (9)$$

where we abbreviate the coefficients in the polynomial  $f_s^\gamma := -\mu \sum_{(c_1, \dots, c_s)} b_{c_1, \dots, c_s}^{-1}$ . Removing the cut-off  $L$ , only the  $s_0$ -fold zero modes of the interaction  $\gamma$  survive, yielding large- $\xi$  asymptotics (using  $-\mu \sim \xi^{-2}$ )

$$Q \underset{\xi \rightarrow \infty}{\sim} \bar{\lambda}_\gamma^{\frac{2}{n_\gamma - 2}} f_{s_0}^\gamma \xi^{\frac{2n_\gamma}{n_\gamma - 2} - d_G(r-s_0)} \quad (10)$$

with  $\bar{\lambda}_\gamma = L^{d_G(r-s_0)(n_\gamma-2)/2} \lambda_\gamma$ . This is the  $Q \sim \xi^{d_{\text{crit}} - d}$  result of local QFT with  $d_{\text{crit}} = \frac{2n_\gamma}{n_\gamma - 2}$ . The effect of the non-local interaction given by the graph  $\gamma$  with  $s_0$  minimal zero modes is a reduction of the configuration space dimension  $d_G \cdot r$  to an effective dimension

$$d = d_{\text{eff}} := d_G(r - s_0). \quad (11)$$

A typical model of 4d quantum gravity has  $r = 4$ , group dimension at least  $d_G = 3$  (e.g.  $G = \text{SU}(2)$  models) and quartic, quintic or higher-order interactions. Though the configuration space is then at least 12-dimensional and  $d_{\text{crit}} = 4$  or  $10/3$  respectively, the effective dimension  $d_{\text{eff}}$  might be smaller, e.g. for  $r - s_0 = 1$  in the case of  $\gamma$  with no multiple edges like the simplicial interaction (see Tab. I). Adding gauge invariance can shift  $s_0 \rightarrow s_0 + 1$  and thus reduce  $d_{\text{eff}}$  even further [28]. On the other hand, if there are edges of high enough multiplicities, e.g. melonic interactions with  $s_0 = 1$ ,  $d_{\text{eff}}$  is larger than  $d_{\text{crit}}$  such

that  $Q \ll 1$  and mean-field theory is a valid description of phase transitions.<sup>1</sup> Thus, combinatorial non-locality of TGFTs affects the detailed mean-field behaviour but does not spoil the very applicability of mean-field theory.

A special case are models where  $G$  is compact and no thermodynamic limit is applicable. Then, there is a fixed finite volume  $L^{d_G}$  (no  $L \rightarrow \infty$  limit) and it is the  $r$ -fold zero mode which dominates in the IR, i.e. at small momentum scale  $k \sim 1/\xi$  [31]. In the current language this is the  $s = r$  mode and thus  $d_{\text{eff}} = 0$ . That is, also in TGFT one has [30, 36] the standard result that a QFT on compact domain is effectively zero dimensional in the IR and does not allow for phase transitions [38].

## 2) MATTER DEGREES OF FREEDOM

Adding matter degrees of freedom in a TGFT model increases the effective dimension [28]. We extend the group field to  $\Phi(\phi, \mathbf{g})$  with  $d_\phi$  local degrees of freedom  $\phi = (\phi_1, \dots, \phi_{d_\phi}) \in \mathbb{R}^{d_\phi}$  with point-like interactions  $\int d\phi \text{Tr}_\gamma[\Phi](\phi)$  while the convolutions of  $\mathbf{g}_i$  remain as in (2). This generates free scalar fields minimally coupled to the discrete geometry. Such fields can be used as reference frames which can allow to retrieve the dynamics of quantum geometry in relational terms [39–42]. Furthermore, this is of course a necessary ingredient of realistic models of quantum gravity coupled to elementary particle fields.

In the Landau-Ginzburg analysis such local degrees of freedom simply add to the dimension. Gaussian correlations (6) extend to

$$\hat{C}(\mathbf{p}, \mathbf{j}) = \frac{1}{\alpha(\mathbf{j}) \sum_{a=1}^{d_\phi} p_a^2 + \frac{1}{V_L} \sum_c \text{Cas}_{j_c=1}^{d_G} + b_j} \quad (12)$$

where the momenta  $p_a$  of  $\phi_a$  can couple to group representations  $\mathbf{j}$  via a function  $\alpha(\mathbf{j})$ . However, only modes with  $p_a \approx 0$  are relevant upon integration over  $\phi_a$  so that the Ginzburg  $Q$  (10) receives no contribution from  $\alpha(\mathbf{j})$  and simply gets an additional factor  $\xi^{-d_\phi}$  [28]. Accordingly, the effective dimension is

$$d_{\text{eff}} = d_\phi + d_G(r - s_0). \quad (13)$$

Therefore, even if  $d_G(r - s_0) \leq d_{\text{crit}}$ , the Ginzburg criterion is still satisfied when there are  $d_\phi > d_{\text{crit}} - d_G(r - s_0)$  matter fields. Even in the special case of compact  $G$  without thermodynamic large-volume limit, mean-field

<sup>1</sup> The extreme case  $s_0 = 0$  occurs for multi-trace interactions like the double trace in Tab. I. Then  $d_{\text{eff}} = d_G r$  like in local QFT and the theory indeed turns out to be a local theory in disguise [35]. While multi-trace couplings are generated along the RG flow in general, they are suppressed in the cyclic melonic [31, 36] and necklace [37] potential approximation.

phase transitions are possible if  $d_\phi > d_{\text{crit}}$ . In this way, adding such matter degrees of freedom not only makes these models more realistic from a physical point of view, but also always improves their mean-field behaviour.

### 3) HYPERBOLIC GEOMETRY

Realistic models of quantum gravity involve a Lie group with hyperbolic geometry related to the Lorentz group. The latter is a crucial ingredient to properly implement micro-causality, but of course brings new challenges, in particular infinite volume factors.

One such model is the Lorentzian Barrett-Crane TGFT model [43] generating triangulations formed by space-like tetrahedra. This is a TGFT on the non-compact curved group  $G = \text{SL}(2, \mathbb{C})$  subject to closure and simplicity constraints. Due to the restriction to space-like tetrahedra the group domain is effectively  $r = 4$  copies of the 3-hyperboloid  $\text{H}^3 = \text{SL}(2, \mathbb{C})/\text{SU}(2)$ . Due to the interplay with combinatorial non-locality, infinite volume factors do not cancel and need to be regularized (see e.g. (5)). We do so in the Cartan decomposition  $\text{SL}(2, \mathbb{C}) \cong \text{SU}(2) \times \text{A}^+ \times \text{SU}(2)$  via a cut-off  $L$  on the diagonal Cartan subgroup

$$\text{A}^+ = \{e^{\frac{\sigma_3}{2} \frac{\eta}{a}} | \eta \in \mathbb{R}_+\} \rightarrow \text{A}_L^+ := \{e^{\frac{\sigma_3}{2} \frac{\eta}{a}} | 0 \leq \eta < L\}. \quad (14)$$

Here  $a$  is the curvature scale, i.e. the skirt radius of the group's hyperbolic part  $\text{H}^3$ . Using the Haar measure on  $\text{SU}(2)$  and  $\sinh^2(\eta/a) d\eta/a$  for  $\text{A}^+$ , one finds the regularized volume of  $\text{SL}(2, \mathbb{C})$  to be

$$V_L = \frac{1}{4} \left( \sinh\left(\frac{2L}{a}\right) - \frac{2L}{a} \right) \underset{L \rightarrow \infty}{\sim} \frac{1}{8} e^{\frac{2L}{a}}. \quad (15)$$

The relative magnitude of Gaussian fluctuations around the mean-field  $\Phi_0$  at large cut-off  $L$  is [29]

$$Q_L = \frac{\sum_{s=s_0}^r \frac{f_s^\gamma}{-\mu} V_\xi^s V_L^{1-s}}{\left(\frac{-\mu}{\lambda_\gamma}\right)^{\frac{2}{n_\gamma-2}} V_\xi^r V_L^{1-r-2\frac{n_\gamma-1}{n_\gamma-2}}}. \quad (16)$$

Removing the regularization, the minimal number  $s_0$  of zero modes dominates such that

$$Q_L \underset{L \rightarrow \infty}{\sim} \bar{\lambda}_\gamma^{\frac{2}{n_\gamma-2}} (-\mu)^{-\frac{n_\gamma}{n_\gamma-2}} f_{s_0}^\gamma V_\xi^{-(r-s_0)} \quad (17)$$

with rescaling  $\bar{\lambda}_\gamma = V_L^{\frac{n_\gamma-2}{2}(r-s_0)+n_\gamma-1} \lambda_\gamma$ . Due to the hyperbolic geometry of the configuration space, the correlation length scales  $\xi \underset{\mu \rightarrow 0}{\sim} \frac{1}{a\mu}$  [29]. Thus,

$$Q \underset{\xi \rightarrow \infty}{\sim} \bar{\lambda}_\gamma^{\frac{2}{n_\gamma-2}} f_{s_0}^\gamma (a\xi)^{\frac{n_\gamma}{n_\gamma-2}} e^{-2(r-s_0)\frac{\xi}{a}}. \quad (18)$$

This is an exponential suppression such that the Ginzburg criterion  $Q \ll 1$  is satisfied regardless of the order  $n_\gamma$  and zero modes  $s_0$  of the interaction.




FIG. 1. The effective dimension (20) for  $s_0 = r - 1$  flowing from three at small  $\xi$  to infinity.

A complementary way to understand this is by using the notion of a scale-dependent effective dimension  $d_{\text{eff}}$  [31]. One can still write (17) as a power function but with scale-dependent inverse power

$$d_{\text{eff}}(\xi) := -\frac{\partial \log F(\xi)}{\partial \log \xi}, \text{ here } F(\xi) = f_{s_0}^\gamma V_\xi^{-(r-s_0)}. \quad (19)$$

Then, the general form  $Q \propto \mu^{-d_{\text{crit}}/2} \xi^{-d_{\text{eff}}(\xi)}$  is valid on all scales  $\xi$ , with the resulting effective dimension

$$d_{\text{eff}}(\xi) = (r - s_0) \frac{\cosh\left(\frac{2\xi}{a}\right) - 1}{\frac{a}{2\xi} \sinh\left(\frac{2\xi}{a}\right) - 1} \quad (20)$$

flowing from  $d_{\text{eff}} = 3(r - s_0)$  in the UV (small  $\xi$ )<sup>2</sup> to  $d_{\text{eff}} \rightarrow \infty$  in the IR. Thus, mean-field theory can describe the phase transition because the theory's effective dimension blows up towards the IR and thereby supersedes any possible critical dimension.

This result for the Barrett-Crane TGFT model suggests that mean-field theory is valid for any TGFT on hyperbolic spaces irrespective of the dimension  $d_G \cdot r$  of the field domain or of combinatorics  $n_\gamma$ ,  $s_0$ . Including  $d_\phi$  matter fields adds again a power  $\xi^{-d_\phi}$  [29] but the exponential suppression still dominates. A hyperbolic sector arises also for models with compact  $G$  but Lorentzian integrals implicit in the kernel  $\mathcal{K}$  [44]. Similarly, one expects this exponential decay to remain also for TGFT models of quantum gravity that also contain light- and time-like tetrahedra like in the complete Lorentzian Barrett-Crane model [45]. Thus, it seems to be a generic feature of TGFT quantum gravity that phase transitions towards a non-vanishing (non-perturbative) vacuum state exist which can be self-consistently described using mean-field theory. Since such a state is typically highly populated by TGFT quanta, this indeed makes a compelling case for an interesting continuum geometric approximation.

<sup>2</sup> Small  $\xi$  is equivalent to large curvature  $a$  here. Thus, this limit also shows consistency with the flat case (11) with  $d_G = 3$  [29]. This is because only the hyperbolic part  $\text{H}^3$  contributes here.

## CONCLUSIONS

The key result of this work is that it is possible to understand some part of the phase structure of TGFT with straightforward QFT methods. The full theory space is very involved and largely out of reach for explicit control due to the intricate interplay between the tensorial nature of the field, the Lie group domain and its geometry, combinatorially non-local interactions, as well as geometricity constraints. Still, considering Gaussian fluctuations around a constant background vacuum field, we find that these features of TGFT can be controlled and even work in favour of a mean-field description: (1) The general  $\xi^{d_{\text{crit}}-d}$  asymptotic scaling towards the IR remains even with non-locality, only with a modified effective dimension  $d = d_{\text{eff}}$ . (2) Coupling local degrees of freedom on  $\mathbb{R}^{d_\phi}$  adds to the dimension  $d_{\text{eff}} \rightarrow d_{\text{eff}} + d_\phi$ . Most importantly, (3) the holonomies of the Lorentz group produce an exponential suppression of the fluctuation size such that we can always find a transition towards a phase which is self-consistently described in terms of mean-field theory regardless of the critical dimension.

Crucially, this provides evidence for the existence of a sensible continuum limit in TGFT quantum gravity as well as in the closely related spin foam models and lattice quantum gravity (where renormalization and continuum limit are the main outstanding challenges, see e.g. [11, 46]), directly rooted in the Lorentz group and thus the causal structure it enforces, which is reminiscent of results of the causal dynamical triangulations approach [5, 47–49]. Therefore, our results impact directly on a wide set of quantum gravity approaches based on discrete structures comprising lattice quantum gravity and canonical loop quantum gravity.

The exponential-suppression effect allows for an educated guess of the phase structure even beyond the Gaussian approximation. Renormalization group calculations of local QFT on  $H^3$  in the local-potential approximation have shown that the flowing effective potential freezes around the curvature scale  $a$  [38] such that the entire phase diagram is already described by the two phases of the mean-field regime. Using the floating-point method one can identify the Wilson-Fisher fixed point but it is pushed to infinity when  $(ka)^{-1} \sim \xi/a \rightarrow \infty$ , where  $k$  denotes the RG scale [38]. This agrees with our result that  $d_{\text{eff}}$  flows to infinity in this regime such that non-trivial phase structure at finite dimension vanishes. We can thus expect also for TGFT<sup>3</sup> that hyperbolic geometry yields a universal phase structure already captured by the two phases around the Gaussian fixed point.

---

<sup>3</sup> Higher orders in a derivative expansion such as the anomalous dimension may uncover further structure as in [31]. But all fixed points found occur only up to some dimension. Thus, we expect them to vanish when  $d_{\text{eff}}$  blows up in the IR.

All in all, our results greatly strengthen the evidence for the existence of a continuum regime in TGFT quantum gravity, the phenomenology of which is tractable by explicit computations in a mean-field approximation. This is all the more important given how difficult it is, due to their analytic and combinatorial complexity, to establish the same fact via a complete RG analysis of the relevant quantum geometric TGFT (and spin foam) models.

## ACKNOWLEDGEMENTS

The work of L. Marchetti was funded by Fondazione Angelo Della Riccia. D. Oriti and A. Pithis acknowledge funding from DFG research grants OR432/3-1 and OR432/4-1. A. Pithis is grateful for the generous financial support from the MCQST via the seed funding Aost 862983-4. J. Thürigen’s research was funded by DFG grant number 418838388 and Germany’s Excellence Strategy EXC 2044–390685587, Mathematics Münster: Dynamics–Geometry–Structure.

---

\* luca.marchetti@physik.lmu.de

† daniele.oriti@physik.lmu.de

‡ andreas.pithis@physik.lmu.de

§ johannes.thuerigen@uni-muenster.de

- [1] R. Blumenhagen, D. Lüst, and S. Theisen, *Basic concepts of string theory*, Theoretical and Mathematical Physics (Springer, Heidelberg, Germany, 2013).
- [2] R. Percacci, *An Introduction to Covariant Quantum Gravity and Asymptotic Safety*, 100 Years of General Relativity, Vol. 3 (World Scientific, 2017).
- [3] M. Reuter and F. Saueressig, *Quantum Gravity and the Functional Renormalization Group: The Road towards Asymptotic Safety* (Cambridge University Press, 2019).
- [4] A. Bonanno, A. Eichhorn, H. Gies, J. M. Pawłowski, R. Percacci, M. Reuter, F. Saueressig, and G. P. Vacca, *Front. in Phys.* **8**, 269 (2020), arXiv:2004.06810 [gr-qc].
- [5] J. Ambjorn, A. Goerlich, J. Jurkiewicz, and R. Loll, *Phys. Rept.* **519**, 127 (2012), arXiv:1203.3591 [hep-th].
- [6] H. W. Hamber, *Quantum gravitation: The Feynman path integral approach* (Springer, Berlin, 2009).
- [7] S. Surya, *Living Rev. Rel.* **22**, 5 (2019), arXiv:1903.11544 [gr-qc].
- [8] C. Rovelli, *Quantum gravity*, Cambridge Monographs on Mathematical Physics (Univ. Pr., Cambridge, UK, 2004).
- [9] C. Rovelli, *PoS QGQS2011*, 003 (2011), arXiv:1102.3660 [gr-qc].
- [10] A. Perez, *Living Rev. Rel.* **16**, 3 (2013), arXiv:1205.2019 [gr-qc].
- [11] S. K. Asante, B. Dittrich, and S. Steinhaus, (2022), arXiv:2211.09578 [gr-qc].
- [12] A. Ashtekar and J. Lewandowski, *Class. Quant. Grav.* **21**, R53 (2004), arXiv:gr-qc/0404018.
- [13] T. Thiemann, *Modern Canonical Quantum General Relativity*, Cambridge Monographs on Mathematical Physics

- (Cambridge University Press, 2007).
- [14] L. Freidel, *Int. J. Theor. Phys.* **44**, 1769 (2005), arXiv:hep-th/0505016.
  - [15] D. Oriti, in *Foundations of Space and Time: Reflections on Quantum Gravity* (2011) pp. 257–320, arXiv:1110.5606 [hep-th].
  - [16] S. Carrozza, *Tensorial methods and renormalization in Group Field Theories*, Ph.D. thesis, Orsay, LPT (2013), arXiv:1310.3736 [hep-th].
  - [17] R. G. Gurău, *Random tensors* (Oxford University Press, 2017).
  - [18] A. Addazi *et al.*, *Prog. Part. Nucl. Phys.* **125**, 103948 (2022), arXiv:2111.05659 [hep-ph].
  - [19] D. Oriti, D. Pranzetti, and L. Sindoni, *Phys. Rev. Lett.* **116**, 211301 (2016), arXiv:1510.06991 [gr-qc].
  - [20] D. Oriti, D. Pranzetti, and L. Sindoni, *Phys. Rev. D* **97**, 066017 (2018), arXiv:1801.01479 [gr-qc].
  - [21] S. Gielen, D. Oriti, and L. Sindoni, *Phys. Rev. Lett.* **111**, 031301 (2013), 1303.3576.
  - [22] A. G. A. Pithis and M. Sakellariadou, *Universe* **5**, 147 (2019), arXiv:1904.00598 [gr-qc].
  - [23] D. Oriti (2021) arXiv:2112.02585 [gr-qc].
  - [24] L. Marchetti and D. Oriti, *JCAP* **07**, 004 (2022), arXiv:2112.12677 [gr-qc].
  - [25] M. Finocchiaro and D. Oriti, *Class. Quant. Grav.* **37**, 015010 (2020), arXiv:1812.03550 [gr-qc].
  - [26] A. G. A. Pithis and J. Thürigen, *Phys. Rev. D* **98**, 126006 (2018), 1808.09765.
  - [27] A. G. A. Pithis, *Aspects of quantum gravity*, Ph.D. thesis, King's Coll. London (2019), arXiv:1903.07735 [gr-qc].
  - [28] L. Marchetti, D. Oriti, A. G. A. Pithis, and J. Thürigen, *JHEP* **21**, 201 (2021), arXiv:2110.15336 [gr-qc].
  - [29] L. Marchetti, D. Oriti, A. G. A. Pithis, and J. Thürigen, *arXiv* (2022), arXiv:2209.04297 [gr-qc].
  - [30] A. G. A. Pithis and J. Thürigen, *Phys. Lett. B* **816**, 136215 (2021), arXiv:2007.08982 [hep-th].
  - [31] A. G. A. Pithis and J. Thürigen, *JHEP* **12**, 159 (2020), arXiv:2009.13588 [hep-th].
  - [32] V. Bonzom, R. Gurau, and V. Rivasseau, *Phys. Rev. D* **85**, 084037 (2012), arXiv:1202.3637 [hep-th].
  - [33] P. Kopietz, L. Bartosch, and F. Schütz, *Introduction to the functional renormalization group*, Vol. 798 (Springer Heidelberg, 2010).
  - [34] J. Thürigen, *Math Phys Anal Geom* **24**, 19 (2021), 2102.12453.
  - [35] V. Rivasseau, *EPL* **111**, 60011 (2015), arXiv:1507.04190 [hep-th].
  - [36] S. Carrozza and V. Lahoche, *Class. Quant. Grav.* **34**, 115004 (2017), arXiv:1612.02452 [hep-th].
  - [37] S. Carrozza, V. Lahoche, and D. Oriti, *Phys. Rev. D* **96**, 066007 (2017), arXiv:1703.06729 [gr-qc].
  - [38] D. Benedetti, *Journal of Statistical Mechanics: Theory and Experiment* **1501**, P01002 (2015), 1403.6712.
  - [39] D. Oriti, L. Sindoni, and E. Wilson-Ewing, *Class. Quant. Grav.* **33**, 224001 (2016), arXiv:1602.05881 [gr-qc].
  - [40] Y. Li, D. Oriti, and M. Zhang, *Class. Quant. Grav.* **34**, 195001 (2017), arXiv:1701.08719 [gr-qc].
  - [41] S. Gielen, *Universe* **4**, 103 (2018), arXiv:1808.10469 [gr-qc].
  - [42] L. Marchetti and D. Oriti, *JHEP* **05**, 025 (2021), arXiv:2008.02774 [gr-qc].
  - [43] A. F. Jercher, D. Oriti, and A. G. A. Pithis, *JCAP* **01**, 050 (2022), arXiv:2112.00091 [gr-qc].
  - [44] M. Finocchiaro, Y. Jeong, and D. Oriti, *Class. Quant. Grav.* **39**, 135014 (2022), arXiv:2012.11536 [gr-qc].
  - [45] A. F. Jercher, D. Oriti, and A. G. A. Pithis, *Phys. Rev. D* **106**, 066019 (2022), arXiv:2206.15442 [gr-qc].
  - [46] J. Ambjorn, J. Gizbert-Studnicki, A. Görlich, J. Jurkiewicz, and R. Loll, *Front. in Phys.* **8**, 247 (2020), arXiv:2002.01693 [hep-th].
  - [47] J. Ambjorn, J. Jurkiewicz, and R. Loll, *Phys. Rev. Lett.* **93**, 131301 (2004), arXiv:hep-th/0404156.
  - [48] S. Jordan and R. Loll, *Phys. Lett. B* **724**, 155 (2013), arXiv:1305.4582 [hep-th].
  - [49] J. Ambjorn (2022) arXiv:2209.06555 [hep-lat].