

# AN INVERSE SPECTRAL PROBLEM FOR A FRACTIONAL SCHRÖDINGER OPERATOR

MOURAD CHOULLI

ABSTRACT. We establish that the potential appearing in a fractional Schrödinger operator is uniquely determined by an internal spectral data.

## 1. INTRODUCTION

Let  $\mathcal{M} = (\mathcal{M}, \mathbf{g})$  be a smooth connected compact Riemannian manifold with boundary  $\partial\mathcal{M}$  of dimension  $n \geq 2$ . Recall that the Laplace-Beltrami operator associated to the metric tensor  $\mathbf{g} = (\mathbf{g}_{k\ell})$  is given in local coordinates by

$$\Delta_{\mathbf{g}} = \frac{1}{\sqrt{|\mathbf{g}|}} \sum_{k,\ell=1}^n \frac{\partial}{\partial x_k} \left( \sqrt{|\mathbf{g}|} \mathbf{g}^{k\ell} \frac{\partial}{\partial x_\ell} \cdot \right),$$

where  $(\mathbf{g}^{k\ell})$  is the matrix inverse of  $(\mathbf{g}_{k\ell})$  and  $|\mathbf{g}|$  is the determinant of  $\mathbf{g}$ .

The following notation will be useful in the sequel

$$H_D^\theta(\mathcal{M}) = \{u \in H^\theta(\mathcal{M}); u|_{\partial\mathcal{M}} = 0\}, \quad \theta > 1/2.$$

We denote by  $A$  the realization on  $L^2(\mathcal{M})$  of the negative Laplace-Beltrami operator with Dirichlet boundary conditions, that is,  $D(A) = H_D^2(\mathcal{M})$  and  $A = -\Delta_{\mathbf{g}}$ .

As  $A$  is positive definite self-adjoint operator with compact resolvent its spectrum is reduced to a sequence of eigenvalues counted according to their multiplicity:

$$0 < \lambda_1 < \lambda_2 \leq \dots \lambda_k \leq \dots \quad \text{and} \quad \lambda_k \rightarrow \infty \text{ as } k \rightarrow \infty.$$

Moreover there exists  $(\phi_k)_{k \geq 1}$  an orthonormal basis of  $L^2(\mathcal{M})$  consisting of eigenfunctions or more precisely, each  $\phi_k$  is an eigenfunction for the eigenvalue  $\lambda_k$ .

Let  $0 < s \leq 1$ . We recall that the fractional power  $A^s$  is defined as follows

$$A^s u = \sum_{k \geq 1} \lambda_k^s (u | \phi_k) \phi_k, \quad u \in D(A^s).$$

Here and henceforth the symbol  $(\cdot | \cdot)$  denotes the natural scalar product of  $L^2(\mathcal{M})$ .

---

2010 *Mathematics Subject Classification.* 35R30.

*Key words and phrases.* Compact Riemannian manifold with boundary, internal spectral data, source-to-solution operator.

The author is supported by the grant ANR-17-CE40-0029 of the French National Research Agency ANR (project MultiOnde).

Following [4, Theorem 1] (see also [8, Theorem 2.1]) we have

$$D(A^s) := \mathcal{H}^s = \begin{cases} H^{2s}(\mathcal{M}), & 0 < s < 1/4, \\ \left\{ u \in H^{1/2}(\mathcal{M}); \int_{\mathcal{M}} \zeta^{-1}(x)|u(x)|^2 dV < \infty \right\}, & s = 1/4, \\ H_D^{2s}(\mathcal{M}), & 1/4 < s \leq 1, \end{cases}$$

where  $dV = \sqrt{|\mathbf{g}|} dx^1 \dots dx^n$  denotes the Riemannian measure associated to  $\mathbf{g}$  and  $\zeta$  is the distance to the boundary  $\partial\mathcal{M}$ .

It is worth noticing that, contrary to  $A$ , the operator  $A^s$ ,  $0 < s < 1$ , is nonlocal.

Set

$$L_+^\infty(\mathcal{M}) = \{\mathfrak{q} \in L^\infty(\mathcal{M}, \mathbb{R}); \mathfrak{q} \geq 0\}$$

For  $\mathfrak{q} \in L_+^\infty(\mathcal{M})$  we denote by  $A_{\mathfrak{q}}$  the realization on  $L^2(\mathcal{M})$  of the operator  $A^s + \mathfrak{q}$  with domain  $D(A_{\mathfrak{q}}) = \mathcal{H}^s$ .

As  $\mathcal{H}^s$  is compactly embedded in  $L^2(\mathcal{M})$  and  $A_{\mathfrak{q}}$  is self-adjoint we derive that the spectrum of  $A_{\mathfrak{q}}$ , denoted by  $\sigma(A_{\mathfrak{q}})$ , consists in a nondecreasing sequence  $(\mu_k^{\mathfrak{q}})$  converging to  $\infty$ . Furthermore to each  $\mu_k^{\mathfrak{q}}$  we can associate an eigenfunction  $\varphi_k^{\mathfrak{q}}$  in such a way that  $(\varphi_k^{\mathfrak{q}})$  form an orthonormal basis of  $L^2(\mathcal{M})$ .

Let  $\Omega_0, \Omega_1$  and  $\Omega'$  be three nonempty open subsets of  $\text{Int}(\mathcal{M})$  chosen in such a way that  $\Omega_j \setminus \overline{\Omega'} \neq \emptyset$ ,  $j = 0, 1$ . Set  $\Omega = \Omega_0 \cup \Omega_1$  and consider the following subset of  $L_+^\infty(\mathcal{M}) \times L_+^\infty(\mathcal{M})$

$$\mathcal{Q} = \{(\mathfrak{q}_1, \mathfrak{q}_2) \in L_+^\infty(\mathcal{M}) \times L_+^\infty(\mathcal{M}); \text{supp}(\mathfrak{q}_1 - \mathfrak{q}_2) \subset \Omega'\}.$$

**Theorem 1.1.** *Let  $(\mathfrak{q}_1, \mathfrak{q}_2) \in \mathcal{Q}$ . If*

$$\mu_k^{\mathfrak{q}_1} = \mu_k^{\mathfrak{q}_2}, \quad \varphi_k^{\mathfrak{q}_1}|_{\Omega} = \varphi_k^{\mathfrak{q}_2}|_{\Omega}, \quad k \geq 1,$$

*then  $\mathfrak{q}_1 = \mathfrak{q}_2$ .*

As  $\mu_k^{\mathfrak{q}_j + \mu} = \mu_k^{\mathfrak{q}_j} + \mu$ , for any  $k \geq 1$ ,  $\mu \in \mathbb{R}$  and  $j = 1, 2$ , it is clear that we can remove the condition  $\mathfrak{q}_j \geq 0$  in Theorem 1.1.

The case  $s = 1$  was considered in [7] when  $(\mathcal{M}, \mathbf{g})$  is a smooth connected compact Riemannian manifold without boundary. The authors established in [7, Corollary 2] that the internal spectral data  $(\lambda_k^{\mathfrak{g}}, \phi_k^{\mathfrak{g}}|_{\Omega})$ , where  $(\lambda_k^{\mathfrak{g}}, \phi_k^{\mathfrak{g}})$  is the sequence of eigenvalues and the corresponding eigenfunctions of  $-\Delta_{\mathfrak{g}}$  and  $\Omega$  is an open subset of  $\text{Int}(\mathcal{M})$ , determines uniquely  $\mathbf{g}$  up to isometry.

There is a wide literature devoted to inverse spectral problems for Schrödinger operators and magnetic Schrödinger operators involving boundary spectral data. We refer to the recent work [1] and references therein.

## 2. PROOF OF THEOREM 1.1

We give the proof in several steps. The key step is based on the density result in Lemma 2.1 which is inspired by the proof of [6, Theorem 1.1].

The resolvent of  $A_{\mathfrak{q}}$ ,  $\mathfrak{q} \in L_+^\infty(\mathcal{M})$ , is denoted by  $R_{\mathfrak{q}}$ :

$$R_{\mathfrak{q}} : \mu \in \rho(A_{\mathfrak{q}}) \mapsto R_{\mathfrak{q}}(\mu) = (A_{\mathfrak{q}} - \mu)^{-1} \in \mathcal{B}(L^2(\mathcal{M}), \mathcal{H}^s),$$

where  $\rho(A_q) = \mathbb{C} \setminus \sigma(A_q)$  is the resolvent set of  $A_q$ .

We can then assert that, for each  $f \in L^2(\mathcal{M})$  and  $\mu \in \rho(A_q)$ ,  $u = R_q(\mu)f \in \mathcal{H}^s$  is the unique variational solution of the equation

$$(2.1) \quad (A^s + \mathfrak{q} - \mu)u = f.$$

That is, we have

$$(2.2) \quad \left( A^{s/2}u | A^{s/2}v \right) + ((\mathfrak{q} - \mu)u | v) = (f | v), \quad v \in \mathcal{H}^{s/2}.$$

As  $\varphi_k^q = R_q(0) (-\mathfrak{q}\varphi_k^q + \mu_k^q\varphi_k^q) \in \mathcal{H}^s$ ,  $k \geq 1$ , we have

$$\begin{aligned} (u | \mu_k^q \varphi_k^q) &= (u | A^s \varphi_k^q) + (u | \mathfrak{q} \varphi_k^q) \\ &= \left( A^{s/2}u | A^{s/2} \varphi_k^q \right) + (u | \mathfrak{q} \varphi_k^q). \end{aligned}$$

This and (2.2) imply

$$(u | \mu_k^q \varphi_k^q) = (f | \varphi_k^q) - ((\mathfrak{q} - \mu)u | \varphi_k^q) + (u | \mathfrak{q} \varphi_k^q)$$

from which we derive in a straightforward manner

$$(2.3) \quad (u | \varphi_k^q) = (\mu_k^q - \mu)^{-1} (f | \varphi_k^q).$$

Hence

$$(2.4) \quad R_q(\mu)f = \sum_{k \geq 1} (\mu_k^q - \mu)^{-1} (f | \varphi_k^q) \varphi_k^q.$$

Note that since  $\min_{k \geq 1} |\mu_k^q - \mu| = \text{dist}(\mu, \sigma(A_q))$  it follows from (2.4) the following well-known resolvent estimate

$$(2.5) \quad \|R_q(\mu)\|_{\mathcal{B}(L^2(\mathcal{M}))} \leq \frac{1}{\text{dist}(\mu, \sigma(A_q))}.$$

We now make an observation that will be used in the sequel.

According to Weyl's asymptotic formula we have  $\lambda_k = O(k^{2/n})$  (e.g. [2]). On the other hand we infer from the min-max principle that  $\lambda_k^s \leq \mu_k^q \leq \lambda_k^s + m$ , for every  $q \in L^\infty(\mathcal{M})$  satisfying  $0 \leq q \leq m$ . We derive that  $\mu_k^q = O(k^{2s/n})$ , uniformly with respect to  $q \in L^\infty(\mathcal{M})$  satisfying  $0 \leq q \leq m$ .

For notational convenience  $R_q(0)$  is simply denoted in the rest of this section by  $R_q$ . Also, we consider  $L^2(\Omega_0)$  as a subset of  $L^2(\mathcal{M})$ :

$$L^2(\Omega_0) = \{f \in L^2(\mathcal{M}); \text{supp}(f) \subset \Omega_0\}$$

and for  $\mathfrak{q} \in L^\infty_+(\mathcal{M})$  define the source-to-solution operator  $\Sigma_{\mathfrak{q}}$  as follows

$$\Sigma_{\mathfrak{q}} : L^2(\Omega_0) \rightarrow L^2(\Omega_1) : f \mapsto R_{\mathfrak{q}}f|_{\Omega_1}.$$

Pick  $\mathfrak{q}_j \in L^\infty(\mathcal{M})$  satisfying  $0 \leq \mathfrak{q}_j \leq m$ ,  $j = 1, 2$ . Since

$$\Sigma_{\mathfrak{q}_j} f = \sum_{k \geq 1} (\mu_k^{\mathfrak{q}_j})^{-1} (f | \varphi_k^{\mathfrak{q}_j}) \varphi_k^{\mathfrak{q}_j}|_{\Omega_1}, \quad j = 1, 2,$$

we have

$$\Sigma_{\mathfrak{q}_1} f - \Sigma_{\mathfrak{q}_2} f = I_1 + I_2 + I_3, \quad f \in L^2(\Omega_0),$$

where

$$\begin{aligned} I_1 &= \sum_{k \geq 1} \left[ (\mu_k^{\mathfrak{q}_1})^{-1} - (\mu_k^{\mathfrak{q}_2})^{-1} \right] (f|_{\varphi_k^{\mathfrak{q}_1}}) \varphi_k^{\mathfrak{q}_1}|_{\Omega_1}, \\ I_2 &= \sum_{k \geq 1} (\mu_k^{\mathfrak{q}_2})^{-1} (f|_{\varphi_k^{\mathfrak{q}_1}} - \varphi_k^{\mathfrak{q}_2}) \varphi_k^{\mathfrak{q}_1}|_{\Omega_1}, \\ I_3 &= \sum_{k \geq 1} (\mu_k^{\mathfrak{q}_2})^{-1} (f|_{\varphi_k^{\mathfrak{q}_2}}) (\varphi_k^{\mathfrak{q}_1}|_{\Omega_1} - \varphi_k^{\mathfrak{q}_2}|_{\Omega_1}). \end{aligned}$$

If  $\|f\|_{L^2(\mathcal{M})} = 1$  then

$$(2.6) \quad \|I_1\|_{L^2(\Omega_1)} \leq C \sum_{k \geq 1} k^{-4s/n} |\mu_k^{\mathfrak{q}_1} - \mu_k^{\mathfrak{q}_2}|$$

and

$$(2.7) \quad \|I_{2+j}\|_{L^2(\Omega_1)} \leq C \sum_{k \geq 1} k^{-2s/n} \|\varphi_k^{\mathfrak{q}_1} - \varphi_k^{\mathfrak{q}_2}\|_{L^2(\Omega_j)}, \quad j = 0, 1.$$

Here and henceforth  $C > 0$  is a generic constant only depending on  $\mathcal{M}$ ,  $\mathfrak{g}$  and  $m$ .

Define

$$\mathbf{d}(\mathfrak{q}_1, \mathfrak{q}_2) = \sum_{k \geq 1} \left[ k^{-4s/n} |\mu_k^{\mathfrak{q}_1} - \mu_k^{\mathfrak{q}_2}| + k^{-2s/n} \|\varphi_k^{\mathfrak{q}_1} - \varphi_k^{\mathfrak{q}_2}\|_{L^2(\Omega)} \right]$$

and suppose that  $\mathbf{d}(\mathfrak{q}_1, \mathfrak{q}_2) < \infty$ . Then inequalities (2.6) and (2.7) yield

$$(2.8) \quad \|\Sigma_{\mathfrak{q}_1} - \Sigma_{\mathfrak{q}_2}\|_{\mathcal{B}(L^2(\Omega_0), L^2(\Omega_1))} \leq C \mathbf{d}(\mathfrak{q}_1, \mathfrak{q}_2).$$

An immediate consequence of (2.8) is

**Proposition 2.1.** *Let  $\mathfrak{q}_1, \mathfrak{q}_2 \in L_+^\infty(\mathcal{M})$ . If*

$$\mu_k^{\mathfrak{q}_2} = \mu_k^{\mathfrak{q}_1}, \quad \varphi_k^{\mathfrak{q}_2}|_{\Omega} = \varphi_k^{\mathfrak{q}_1}|_{\Omega} \quad k \geq 1,$$

then  $\Sigma_{\mathfrak{q}_1} = \Sigma_{\mathfrak{q}_2}$ .

Next, we state a result borrowed to [11, Theorem 1.5] which still holds for  $H^{2s}$ -solutions. We point out that the reflexion extension argument used in the proof of [11, Theorem 1.5] is contained in [10, Theorem 3.1].

**Theorem 2.1.** *Pick  $V$  a nonempty open subset of  $\text{Int}(\mathcal{M})$ . If  $u \in \mathcal{H}^s$  satisfies  $u = A^s u = 0$  in  $V$  then  $u = 0$ .*

Theorem 2.1 will serve to establish the following density result.

**Lemma 2.1.** *Let  $\omega_0, \omega_1$  be two nonempty open subsets of  $\text{Int}(\mathcal{M})$  so that  $V = \omega_0 \setminus \bar{\omega}_1 \neq \emptyset$ . For every  $q \in L_+^\infty(\mathcal{M})$  the set*

$$\mathcal{S}_q = \{u = R_q f|_{\omega_1}; f \in C_0^\infty(\omega_0)\}$$

is dense in  $L^2(\omega_1)$ .

*Proof.* Let  $g \in L^2(\omega_1)$ , extended by 0 outside  $\omega_1$ , satisfy  $(g|u) = 0$  for any  $u \in \mathcal{S}_q$ . Equivalently we have

$$(2.9) \quad (g|R_q f) = 0, \quad f \in C_0^\infty(\omega_0).$$

If  $v = R_q g$  then

$$(g|R_q f) = \sum_{k \geq 1} (\mu_k^q)^{-1} (\varphi_k^q | f) (g | \varphi_k^q) = (R_q g | f) = (v | f), \quad f \in C_0^\infty(\omega_0).$$

Hence (2.9) yields

$$(v | f) = 0, \quad f \in C_0^\infty(\omega_0).$$

Therefore  $v = 0$  in  $\omega_0$ . Using  $A^s v = g - qv$  and  $\text{supp}(g) \subset \omega_1$  we obtain  $v = A^s v = 0$  in  $V$ . Theorem 2.1 implies  $v = 0$  and hence  $g = 0$ .  $\square$

*Completion of the proof of Theorem 1.1.* Let  $(q_1, q_2) \in \mathcal{Q}$  satisfying

$$\mu_k^{q_1} = \mu_k^{q_2}, \quad \varphi_k^{q_1}|_\Omega = \varphi_k^{q_2}|_\Omega, \quad k \geq 1.$$

For  $u_j = R_{q_j} f$ ,  $f \in L^2(\Omega_0)$ , we have

$$A^s u_1 + q_1 u_1 = A^s u_2 + q_2 u_2.$$

Hence

$$A^s(u_1 - u_2) + q_1(u_1 - u_2) = (q_2 - q_1)u_2.$$

If  $v_1 = R_{q_1} g$ ,  $g \in C_0^\infty(\Omega_1)$ , then

$$((u_1 - u_2) | A^s v_1) + ((u_1 - u_2) | q_1 v_1) = ((q_2 - q_1) u_2 | v_1).$$

Using that  $A^s v_1 + q_1 v_1 = g$  we find

$$((u_1 - u_2) | g) = ((q_2 - q_1) u_2 | v_1).$$

It follows from Proposition 2.1 that  $u_1 - u_2 = 0$  in  $\Omega_1$ . This and the fact that  $\text{supp}(g) \subset \Omega_1$  imply

$$((q_2 - q_1) u_2 | v_1) = 0.$$

In other words we proved

$$(2.10) \quad ((q_2 - q_1) R_{q_2} f | R_{q_1} g) = 0, \quad f \in C_0^\infty(\Omega_0), \quad g \in C_0^\infty(\Omega_1).$$

As  $\text{supp}(q_2 - q_1) \subset \Omega'$  and  $\{R_{q_1} g|_{\Omega'}; g \in C_0^\infty(\Omega_1)\}$  is dense  $L^2(\Omega')$  by Lemma 2.1 we find by applying (2.10) to a sequence  $(g_k)$  in  $C_0^\infty(\Omega_1)$  so that  $R_{q_1} g_k|_{\Omega'}$  converges to 1 in  $L^2(\Omega')$

$$((q_2 - q_1) R_{q_2} f | 1) = (R_{q_2} f | q_2 - q_1) = 0, \quad f \in C_0^\infty(\Omega_0).$$

Applying once again Lemma 2.1 (with  $\omega_0 = \Omega_0$  and  $\omega_1 = \Omega'$ ) we finally obtain  $q_2 = q_1$ .  $\square$

The preceding proof contains the following result

**Theorem 2.2.** *Let  $(q_1, q_2) \in \mathcal{Q}$ . If  $\Sigma_{q_1} = \Sigma_{q_2}$ , then  $q_1 = q_2$ .*

Let  $(-\Delta)^s = \text{Op}(|\xi|^{2s})$ ,  $0 < s < 1$ , be the fractional power of  $-\Delta$  acting as an operator on  $\mathbb{R}^n$ . Pick  $D$  a bounded domain of  $\mathbb{R}^n$  and consider the following problem

$$(2.11) \quad ((-\Delta)^s + q)u = 0 \text{ in } D, \quad u|_{D_e} = f,$$

where  $D_e = \mathbb{R}^n \setminus \overline{D}$ .

When  $D$  and  $q$  are sufficiently smooth the authors define in [6] an analogue of the so-called Dirichlet-to-Neumann map as follows

$$\Lambda_q : f \in H^{s+\beta}(D_e) \mapsto (-\Delta u(f))^s|_{D_e} \in H^{-s+\beta}(D_e),$$

where  $u(f)$  is the variational solution of (2.11) and  $\max(0, s - 1/2) < \beta < 1/2$ .

Let  $W_1, W_2$  be two open subsets of  $D_e$ . It is proved in [6, Theorem 1.1] that  $f \in C_0^\infty(W_1) \mapsto \Lambda_q f|_{W_2}$  determines uniquely  $q$ .

A generalization of this result was obtained in [5] when  $-\Delta$  is substituted by an elliptic operator with variable coefficients.

We refer to [9, Theorem 1.2] to a stability inequality corresponding to the uniqueness result in [6].

### 3. COMMENTS

Fix  $\Omega$  a nonempty open subset of  $\text{Int}(\mathcal{M})$  and let  $\Omega_0$  and  $\Omega_1$  be two nonempty open subsets so that  $\Omega_0 \cup \Omega_1 = \Omega$ . Let  $q_1, q_2 \in L_+^\infty(\mathcal{M})$ . If the assumption

$$(3.1) \quad \mu_k^{q_1} = \mu_k^{q_2}, \quad \varphi_k^{q_1}|_\Omega = \varphi_k^{q_2}|_\Omega, \quad k \geq 1,$$

holds then we derive from formula (2.4)

$$(3.2) \quad R_{q_1}(-\mu)f|_{\Omega_1} = R_{q_2}(-\mu)f|_{\Omega_1}, \quad \Re\mu > 0, \quad f \in C_0^\infty(\Omega_0).$$

This means that the internal spectral data  $(\lambda_k^q, \varphi_k^q|_\Omega)$  does not only determine the source-to-solution operator  $\Sigma_q$  but the whole family of source-to-solution operators defined as follows

$$\Sigma_q(\mu) : f \in C_0^\infty(\Omega_0) \mapsto R_q(-\mu)f|_{\Omega_1}, \quad \Re\mu > 0,$$

It turns out that the knowledge of the family of source-to-solution operators  $(\Sigma_q(\mu))_{\Re\mu > 0}$  determines the internal data  $(\lambda_k^q, \varphi_k^q|_\Omega)$ . Precisely, we have the following result.

**Proposition 3.1.** *Suppose  $\Omega_0 \cap \Omega_1 \neq \emptyset$ . Let  $q_1, q_2 \in L_+^\infty(\mathcal{M})$  satisfying*

$$\Sigma_{q_1}(\mu) = \Sigma_{q_2}(\mu), \quad \Re\mu > 0.$$

*Then there exists  $(\hat{\varphi}_k^{q_j})$  an orthonormal basis of eigenfunctions of  $A_{q_j}$ ,  $j = 1, 2$ , such that we have*

$$(3.3) \quad \mu_k^{q_1} = \mu_k^{q_2}, \quad \hat{\varphi}_k^{q_1}|_\Omega = \hat{\varphi}_k^{q_2}|_\Omega, \quad k \geq 1.$$

*Proof.* Noting that

$$\int_0^{+\infty} e^{-\mu t} e^{-\mu_k^{q_j} t} dt = (\mu + \mu_k^{q_j})^{-1}, \quad k \geq 1, \quad j = 1, 2, \quad \Re\mu > 0,$$

we obtain that  $\mu \in \{z \in \mathbb{C}; \Re z > 0\} \mapsto R_{q_j}(-\mu)f|_{\Omega_1}$  is the Laplace transform of the mapping

$$(3.4) \quad T_{q_j}(t)f = \sum_{k \geq 1} e^{-\mu_k^{q_j} t} (f|\varphi_k^{q_j}) \varphi_k^{q_j}|_{\Omega_1}, \quad j = 1, 2, f \in C_0^\infty(\Omega_0), t > 0.$$

Therefore we have

$$(3.5) \quad T_{q_1}(t)f = T_{q_2}(t)f, \quad t > 0, f \in C_0^\infty(\Omega_0).$$

Observe that the mapping

$$f \in L^2(\Omega) \mapsto S_{q_j}(t)f = \sum_{k \geq 1} e^{-\mu_k^{q_j} t} (f|\varphi_k^{q_j}) \varphi_k^{q_j}, \quad t \geq 0,$$

is nothing but the semi-group generated by  $-A_{q_j}$ ,  $j = 1, 2$ .

Let  $(\tilde{\mu}_k^{q_j})$  be the sequence of distinct eigenvalues of  $A_{q_j}$  and  $(\tilde{\varphi}_{k,1}^{q_j}, \dots, \varphi_{k,m_k^j}^{q_j})$  be an orthonormal basis of the eigenspace associated to  $\tilde{\mu}_k^{q_j}$ ,  $k \geq 1$  and  $j = 1, 2$ .

In light of these new definitions we rewrite (3.4) in the form

$$T_{q_j}(t)f = \sum_{k \geq 1} e^{-\tilde{\mu}_k^{q_j} t} \sum_{\ell=1}^{m_k^j} (f|\tilde{\varphi}_{k,\ell}^{q_j}) \tilde{\varphi}_{k,\ell}^{q_j}|_{\Omega_1}, \quad j = 1, 2, f \in C_0^\infty(\Omega_0), t > 0.$$

The uniqueness of Dirichlet series yields  $\tilde{\mu}_k^{q_1} = \tilde{\mu}_k^{q_2}$ ,  $k \geq 1$ , and

$$\sum_{\ell=1}^{m_k^1} (f|\tilde{\varphi}_{k,\ell}^{q_1}) \tilde{\varphi}_{k,\ell}^{q_1}|_{\Omega_1} = \sum_{\ell=1}^{m_k^2} (f|\tilde{\varphi}_{k,\ell}^{q_2}) \tilde{\varphi}_{k,\ell}^{q_2}|_{\Omega_1}, \quad k \geq 1, f \in C_0^\infty(\Omega_0).$$

We derive in a straightforward manner from these identities

$$\sum_{\ell=1}^{m_k^1} \tilde{\varphi}_{k,\ell}^{q_1}|_{\Omega_0} \otimes \tilde{\varphi}_{k,\ell}^{q_1}|_{\Omega_1} = \sum_{\ell=1}^{m_k^2} \tilde{\varphi}_{k,\ell}^{q_2}|_{\Omega_0} \otimes \tilde{\varphi}_{k,\ell}^{q_2}|_{\Omega_1}, \quad k \geq 1.$$

Upon substituting  $(\tilde{\varphi}_{k,\ell}^{q_2})_{1 \leq \ell \leq m_k^2}$  by another orthonormal basis of eigenfunctions we infer from [3, Lemma 2.3] that  $m_k^1 = m_k^2 = m_k$ ,  $k \geq 1$ , and

$$\tilde{\varphi}_{k,\ell}^{q_1}|_{\Omega} = \tilde{\varphi}_{k,\ell}^{q_2}|_{\Omega}, \quad 1 \leq \ell \leq m_k, k \geq 1.$$

In other words we proved that (3.3) holds.

It is worth noticing that conditions (i) and (iii) of [3, Lemma 2.3] hold in our case as a consequence of the uniqueness of continuation property of Theorem 2.1.  $\square$

**Acknowledgement.** I would like to thank the referee for his valuable comments.

#### REFERENCES

- [1] M. Bellassoued, M. Choulli, D. Dos Santos Ferreira, Y. Kian and P. Stefanov, A Borg-Levinson theorem for magnetic Schrödinger operators on a Riemannian manifold, *Ann. Inst. Fourier* 17 (6) (2021), 2471-2517. [2](#)
- [2] P. H. Bérard, *Spectral geometry : direct and inverse problems*, Lecture Notes in Mathematics, 1207, Springer-Verlag, Berlin, 1986. [3](#)
- [3] B. Canuto and O. Kavian, Determining coefficients in a class of heat equations via boundary measurements, *SIAM J. Math. Anal.* 32 (5) (2001), 963-986. [7](#)

- [4] D. Fujiwara, The asymptotic formula for the trace of Green operators of elliptic operators on compact manifold, Proc. Japan Acad. 43 (1967), 426-428. [2](#)
- [5] T. Ghosh, H.-Y. Lin and J. Xiao, The Calderón problem for variable coefficients nonlocal elliptic operators, Commun. Part. Diff. Equat. 42 (12), (2017), 1923-1961. [6](#)
- [6] T. Ghosh, M. Salo, Mikko and G. Uhlmann, The Calderón problem for the fractional Schrödinger equation, Anal. PDE 13 (2) (2020), 455-475. [2](#), [6](#)
- [7] T. Helin, M. Lassas, L. Oksanen and T. Saksala, Correlation based passive imaging with a white noise source, J. Math. Pures Appl. (9) 116 (2018), 132-160. [2](#)
- [8] T. Nambu, Characterization of the domain of fractional powers of a class of elliptic differential operators with feedback boundary conditions. J. Differential Equations 136 (2) (1997), 294-324. [2](#)
- [9] A. Rüländ and M. Salo, The fractional Calderón problem: low regularity and stability, Non-linear Anal., 193:111529, 56, (2020). [6](#)
- [10] P. R. Stinga and C. Zhang, Harnack's inequality for fractional nonlocal equations, Discrete Contin. Dyn. Syst. 33 (7) (2013), 3153-3170. [4](#)
- [11] H. Yu, Unique continuation for fractional orders of elliptic equations, Ann. PDE 3 (2) (2017), Paper No. 16, 21 pp. [4](#)

UNIVERSITÉ DE LORRAINE

*Email address:* mourad.choulli@univ-lorraine.fr