

ESTIMATES CONTROLLING A FUNCTION BY ONLY ITS RADIAL DERIVATIVE AND APPLICATIONS TO STABLE SOLUTIONS OF ELLIPTIC EQUATIONS

XAVIER CABRÉ

ABSTRACT. We establish two new estimates which control a function in L^1 by only the L^1 norm of its radial derivative. While the interior estimate holds for all superharmonic functions, the boundary version is much more delicate. It requires the function to be a stable solution of a semilinear elliptic equation with a nonnegative, nondecreasing, and convex nonlinearity. As an application, our estimates provide quantitative proofs of two results established in [Cabré, Figalli, Ros-Oton, and Serra, *Acta Math.* 224 (2020)] by contradiction-compactness proofs. We recall that this work proved the Hölder regularity of stable solutions to semilinear elliptic equations in the optimal range of dimensions $n \leq 9$.

1. INTRODUCTION

The recent paper [CFRS] by Figalli, Ros-Oton, Serra, and the author established the Hölder regularity of stable solutions to semilinear elliptic equations in the optimal range of dimensions $n \leq 9$. It concerned the semilinear elliptic equation

$$-\Delta u = f(u) \quad \text{in } \Omega \subset \mathbb{R}^n, \quad (1.1)$$

where Ω is a bounded domain, $f \in C^1(\mathbb{R})$, and $u : \bar{\Omega} \subset \mathbb{R}^n \rightarrow \mathbb{R}$. Note that (1.1) is the first variation of the functional

$$\mathcal{E}[u] := \int_{\Omega} \left(\frac{|\nabla u|^2}{2} - F(u) \right) dx, \quad (1.2)$$

where $F' = f$. Let us consider the second variation of \mathcal{E} at u , given by

$$\frac{d^2}{d\varepsilon^2} \Big|_{\varepsilon=0} \mathcal{E}[u + \varepsilon\xi] = \int_{\Omega} \left(|\nabla \xi|^2 - f'(u)\xi^2 \right) dx.$$

One says that u is a *stable* solution of equation (1.1) in Ω if the second variation at u is nonnegative, namely, if

$$\int_{\Omega} f'(u)\xi^2 dx \leq \int_{\Omega} |\nabla \xi|^2 dx \quad \text{for all } \xi \in C^1(\bar{\Omega}) \text{ with } \xi|_{\partial\Omega} \equiv 0. \quad (1.3)$$

The author was supported by grants MTM2017-84214-C2-1-P, RED2018-102650-T, and PID2021-123903NB-I00 funded by MCIN/AEI/10.13039/501100011033 and by ERDF “A way of making Europe”, as well as by the Spanish State Research Agency through the Severo Ochoa and María de Maeztu Program for Centers and Units of Excellence in R&D (CEX2020-001084-M).

Note that the stability of u is defined within the class of functions $u + \varepsilon\xi$ agreeing with u on the boundary $\partial\Omega$. Stability is equivalent to assuming the nonnegativeness of the first Dirichlet eigenvalue in Ω for the linearized operator of (1.1) at u , namely $-\Delta - f'(u)$.

If a *local minimizer* of \mathcal{E} exists (that is, a minimizer for small perturbations having same boundary values), then it will be a stable solution as well.

See the monograph [D] by Dupaigne for more information and results on stable solutions.

The following is the result of [CFRS] on interior regularity. Here, and throughout the paper, by “*dimensional constant*” we mean a constant that depends only on n .

Theorem 1.1. [CFRS, Theorem 1.2]. *Let $u \in C^\infty(\overline{B_1})$ be a stable solution of $-\Delta u = f(u)$ in $B_1 \subset \mathbb{R}^n$, for some nonnegative function $f \in C^1(\mathbb{R})$.*

Then,

$$\|\nabla u\|_{L^{2+\gamma}(B_{1/2})} \leq C\|u\|_{L^1(B_1)} \quad (1.4)$$

for some dimensional constants $\gamma > 0$ and C . In addition,

$$\|u\|_{C^\alpha(\overline{B_{1/2}})} \leq C\|u\|_{L^1(B_1)} \quad \text{if } n \leq 9, \quad (1.5)$$

where $\alpha > 0$ and C are dimensional constants.

The proof of the Hölder estimate (1.5) starts from the bound

$$\int_{B_\rho} r^{2-n} u_r^2 dx \leq C \int_{B_{3\rho/2} \setminus B_\rho} r^{2-n} |\nabla u|^2 dx, \quad (1.6)$$

for all $\rho < 2/3$, which holds for $3 \leq n \leq 9$ and all stable solutions. It is easily proved in Lemma 2.1 of [CFRS] using the stability condition (1.3) with the choice of test function $\xi = x \cdot \nabla u r^{(2-n)/2} = u_r r^{1+(2-n)/2}$. Here and throughout the paper we will use the notation

$$r = |x| \quad \text{and} \quad u_r(x) = \frac{x}{|x|} \cdot \nabla u(x). \quad (1.7)$$

To conclude Hölder regularity from (1.6),¹ it suffices to replace the gradient $|\nabla u|^2$ integrated on the annulus in the right-hand side by the radial derivative u_r , also in an annulus. This was done in [CFRS] using the following result.

Lemma 1.2. [CFRS, Lemma 3.1]. *Let $u \in C^2(B_2)$ be a stable solution of $-\Delta u = f(u)$ in $B_2 \subset \mathbb{R}^n$, with f locally Lipschitz and nonnegative. Assume that*

$$\int_{B_1} |\nabla u|^2 dx \geq \delta \int_{B_2} |\nabla u|^2 dx \quad (1.8)$$

¹At this point one could think to replace the origin on the left-hand side of (1.6) by any other point y , and hence to have the same inequality with the new left-hand side $\int_{B_\rho(y)} r_y^{2-n} u_{r_y}^2 dx$ for all points y in a certain ball around the origin, where $r_y(x) = |x - y|$ and u_{r_y} is the radial derivative centered at y . Averaging in y one could think that $r^{2-n} u_r^2$ on the left-hand side of (1.6) could then be replaced by $r^{2-n} |\nabla u|^2$. This, given the right-hand side of (1.6) in an annulus, would lead to Hölder regularity. However, as shown in Appendix E of [C] with some concrete examples, such averaging procedure cannot work for $n \geq 3$, due to the presence of the weight r^{2-n} .

for some $\delta > 0$.

Then, there exists a constant C_δ , depending only on n and δ , such that

$$\int_{B_{3/2}} |\nabla u|^2 dx \leq C_\delta \int_{B_{3/2} \setminus B_1} u_r^2 dx. \quad (1.9)$$

Note the *doubling assumption* (1.8) made on $|\nabla u|^2$ for the validity of the result. The proof of the lemma was based on a contradiction-compactness argument. Compactness in the $W^{1,2}$ norm followed from the higher integrability bound (1.4) (obtained in $B_{3/2}$ instead of $B_{1/2}$). The contradiction came from analyzing the validity of (1.9) in the limiting case $u_r \equiv 0$. This is simple, since if $u_r \equiv 0$, then u is 0-homogeneous and, thus, u is a superharmonic function for the Laplace-Beltrami operator on the unit sphere (recall that $f \geq 0$). This yields u to be constant, and hence $|\nabla u| \equiv 0$.

In the current paper we succeed to avoid such a contradiction-compactness proof combining estimate (1.6) with two more ingredients. The first one is [CFRS, Proposition 2.5], which claims that stable solutions satisfy the bound

$$\|\nabla u\|_{L^2(B_{1/2})} \leq C \|u\|_{L^1(B_1)}.$$

This estimate was proved in [CFRS] in a quantitative way by using, among other ideas, the test function $\xi = |\nabla u| \zeta$, with ζ a simple cut-off function, in the stability inequality (1.3). Now, one covers the annulus on the right-hand side of (1.6) by balls and uses the previous estimate in each ball. We conclude that it suffices to bound the L^1 norm of the solution in an annulus —instead of the L^2 norm of its gradient in (1.6)— by only its radial derivative.

This is the content of the first result of the current paper. It is a new inequality that will be established in a quantitative way and that will suffice to complete the proof of (1.5) in Theorem 1.1, thus avoiding the use of Lemma 1.2 and its contradiction-compactness proof.

Theorem 1.3. *Let $u \in C^\infty(\overline{B_1})$ be superharmonic in $B_1 \subset \mathbb{R}^n$.*

Then, there exists a constant t , which depends on u , such that

$$\|u - t\|_{L^1(B_1 \setminus B_{1/2})} \leq C \|u_r\|_{L^1(B_1 \setminus B_{1/2})} \quad (1.10)$$

and

$$\|u - t\|_{L^1(B_1)} \leq C \|u_r\|_{L^1(B_1)} \quad (1.11)$$

for some dimensional constant C .

The estimates hold for all superharmonic functions, but still we could not find them in previous literature.²

Note also that, with respect to (1.9), we have replaced L^2 norms by L^1 norms.

In addition, the new estimates do not require a doubling assumption, unlike (1.9) —which required (1.8). This will simplify the Hölder regularity proof of [CFRS], as

²That superharmonicity suffices is in contrast with the much more delicate case of boundary regularity, treated next, where the analogue estimate will crucially require u to be a stable solution of a semilinear equation for a certain class of nonlinearities.

presented in all detail in [C]. More importantly, we will establish Theorem 1.3 with a quantitative proof. In particular, this will allow to quantify the Hölder exponent α in Theorem 1.1. It will use only estimate (1.10) in an annulus —it will not use estimate (1.11) in balls, which is stated here as additional information and possible future use.

In fact, regarding estimates (1.10) and (1.11), one could wonder about their validity for all functions u , and not only superharmonic ones. For $n \geq 2$, both inequalities would be false.³

Let us finally point out that having two different proofs of the regularity result of [CFRS] is of interest for possible future use in other frameworks. For instance, the compactness proof of [CFRS] has been extended to the p -Laplacian in [CMS]. Instead, Theorem 1.3 (and Theorem 1.1) have been extended to stable solutions for operators with variable coefficients by Erneta [E1].

Remark 1.4. Our quantitative proof of Theorem 1.1 allows for an easy extension of the result to the case $f \geq -K$ for some positive constant K —note that Theorem 1.1 assumed $f \geq 0$. The resulting estimates in this case are those of Theorem 1.1 with the right-hand sides replaced by $C(\|u\|_{L^1(B_1)} + K)$, where C is still a dimensional constant. The necessary changes in the proofs are explained in [C]. We point out that it was not clear how to obtain such a result for $f \geq -K$ by extending the proof in [CFRS], due to the compactness argument used in that paper.

Very recently, and independently of our work, Fa Peng [P] has weakened the hypothesis $f \geq 0$ in [CFRS] to allow $f \geq A \min(0, t) - K$ for positive constants A and K . This improves our requirement $f \geq -K$. He needs, however, to replace the L^1 norm of u on the right-hand side of (1.5) by the $W^{1,2}$ norm of u .

The proof of Theorem 1.3 uses a harmonic replacement together with the maximum principle. This allows us to reduce the problem to the case when u is harmonic —for this reduction it will be crucial that we work with L^1 instead of L^2 norms. But when $\Delta u = 0$, we confront a Neumann problem for the homogeneous Laplace equation (since u_r , restricted to any sphere, may be thought as the Neumann data), for which an estimate on spheres similar to the one of the theorem is not surprising and, in fact, easy to prove.

Let us now turn into the much more delicate case of boundary regularity. The main result in [CFRS] established a C^α estimate up to the boundary, in domains of class C^3 and in dimensions $n \leq 9$, for stable solutions of the problem

$$\begin{cases} -\Delta u = f(u) & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

³This can be seen with the family of functions $u^\varepsilon(x) := \varphi(|x|/\varepsilon)x_n/|x|$, where $\varepsilon \in (0, 1)$, φ is nonnegative, vanishes in $(0, 1/4)$, and is identically 1 in $(1/2, \infty)$. Then, u_r^ε has support in $B_{\varepsilon/2}$ and hence its $L^1(B_1)$ norm is bounded above by $C\varepsilon^{n-1}$. Instead, to control the $L^1(B_1 \setminus B_{1/2})$ norm of $u^\varepsilon - t^\varepsilon$ by below, if $t^\varepsilon \leq 0$ (the case $t^\varepsilon \geq 0$ is done similarly) we may integrate only in $\{1/2 < |x| < 1, x_n > 0\}$. In this set $|u^\varepsilon - t^\varepsilon| = u^\varepsilon - t^\varepsilon \geq u^\varepsilon = x_n/|x|$, and thus the $L^1(B_1 \setminus B_{1/2})$ norm is larger than a positive constant independent of ε .

The key boundary ingredient was the following result, Theorem 6.1 from [CFRS] in the case $\theta = 0$ of half-balls. To state it, let us introduce the notation

$$\mathbb{R}_+^n = \{x \in \mathbb{R}^n : x_n > 0\}, \quad B_\rho^+ = \mathbb{R}_+^n \cap B_\rho, \quad \text{and} \quad \partial^0 B_\rho^+ = \{x_n = 0\} \cap \partial B_\rho^+.$$

Theorem 1.5. [CFRS, Theorem 6.1]. *Let $u \in C^\infty(\overline{B_1^+})$ be a nonnegative stable solution of $-\Delta u = f(u)$ in $B_1^+ \subset \mathbb{R}^n$, with $u = 0$ on $\partial^0 B_1^+$. Assume that $f \in C^1(\mathbb{R})$ is nonnegative, nondecreasing, and convex.*

Then,

$$\|u\|_{C^\alpha(\overline{B_{1/2}^+})} \leq C \|u\|_{L^1(B_1^+)} \quad \text{if } n \leq 9, \quad (1.12)$$

where $\alpha > 0$ and C are dimensional constants.

It is important here to assume the solution u to be nonnegative —an assumption that will hold when applying the theorem to a problem in all of Ω with zero Dirichlet boundary conditions, since $f \geq 0$. Note also that the result requires more assumptions on the nonlinearity than the interior Theorem 1.1.

As in the interior case, the proof of Theorem 1.5 given in [CFRS] requires a boundary analogue of Lemma 1.2 on control of ∇u by only its radial derivative u_r . It is here where [CFRS] used a contradiction-compactness argument (Step 2 of the proof of Proposition 6.3 in that paper) in addition to a new delicate result. To explain the new result, it is interesting to first look at the extreme case in which $u_r \equiv 0$ —this can be seen as a first validity test for a boundary version of estimate (1.9). In fact, such case had to be analyzed also in the contradiction-compactness proof of [CFRS]. In contrast with the interior case, “bad news” arise here. Indeed, there exist nonzero nonnegative superharmonic functions u in B_1^+ which vanish on $\partial^0 B_1^+$ and are 0-homogeneous (i.e., for which $u_r \equiv 0$) —obviously, they are not smooth at the origin, but belong to $W^{1,2}(B_1^+)$ when $n \geq 3$. An example is given by the function

$$u(x) = \frac{x_n}{|x|}.$$

This is the key reason behind the fact that our new boundary result, Theorem 1.6 below, does not hold in the class of superharmonic functions which are smooth up to the flat boundary, as we will show in all detail in Remark 3.1.

As a consequence, the contradiction in [CFRS] could not come from the limiting function being superharmonic —as in the interior case—, but from a stronger property. Namely, Theorem 4.1 of [CFRS] established that the class of $W^{1,2}$ functions u for which there exists a nonnegative, nondecreasing, and convex nonlinearity f (with f blowing-up, perhaps, at the value $\sup u$) such that u is a stable solution of $-\Delta u = f(u)$, is closed under the L^1 -convergence of the functions u in compact sets. Note that the class does not correspond to stable solutions of only one equation, but of all equations with such nonlinearities. This is a very strong result, of interest by itself, which will not be needed in the new quantitative proof given in [C] using the result of the current paper. The work [CFRS] used this compactness result to find that, in the extreme case discussed above, the limiting function u would be not only superharmonic, but also a solution of an equation $-\Delta u = g(u)$ for some

nonlinearity g with the properties listed above. This gave a contradiction, since then $g(u)$ would be 0-homogeneous (being u_r identically 0), while $-\Delta u$ would be -2 -homogeneous.

The compactness Theorem 4.1 of [CFRS] will not be needed in our quantitative proof. As a consequence, it will be simpler to implement our proof in other frameworks than that of [CFRS]. This is the case of stable solutions for operators with variable coefficients, as treated in [E2].

The new quantitative proof of boundary Hölder regularity (described in detail in [C]) uses the main and most delicate result of the current paper —more precisely, estimate (1.13) below, in a half-annulus. Its statement is the following.

Theorem 1.6. *Let $u \in C^\infty(\overline{B_1^+})$ be a nonnegative stable solution of $-\Delta u = f(u)$ in $B_1^+ \subset \mathbb{R}^n$, with $u = 0$ on $\partial^0 B_1^+$. Assume that $f \in C^1(\mathbb{R})$ is nonnegative, nondecreasing, and convex.*

Then,

$$\|u\|_{L^1(B_1^+ \setminus B_{1/2}^+)} \leq C \|u_r\|_{L^1(B_1^+ \setminus B_{1/2}^+)} \quad (1.13)$$

and

$$\|u\|_{L^1(B_1^+)} \leq C \|u_r\|_{L^1(B_1^+)} \quad (1.14)$$

for some dimensional constant C .

Several comments are in order.

First, in Remark 3.1 we will show that, unlike the interior case of Theorem 1.3, estimate (1.13) does not hold in the larger class of nonnegative superharmonic functions in B_1^+ which are smooth up to the boundary of B_1^+ and vanish on $\partial^0 B_1^+$.

In Remark 3.3 we will prove that, at least for $n \geq 3$, Theorem 1.6 also holds when replacing (1.13), or (1.14), by the stronger bound

$$\|u\|_{L^1(B_1^+)} \leq C \|u_r\|_{L^1(B_1^+ \setminus B_{1/2}^+)} \quad (\text{if } n \geq 3).$$

The proof of Theorem 1.6 is rather delicate and required new ideas. It is completely different than that of the analogue interior estimate, Theorem 1.3. It uses not only the semilinear equation for u , but also the stability condition. Furthermore, it requires the nonlinearity to satisfy $f \geq 0$, $f' \geq 0$, and $f'' \geq 0$. Note that these are the exactly same assumptions needed in the contradiction-compactness proof in [CFRS]. This fact came partly as a surprise, since our quantitative proof and the one of [CFRS] are based in very different ingredients, as described next.

The proof of Theorem 1.6 will be quantitative. It starts from the identity

$$-2\Delta u + \Delta(x \cdot \nabla u) = -f'(u) x \cdot \nabla u, \quad (1.15)$$

which is easily checked by computing $\Delta(x \cdot \nabla u)$. After multiplying (1.15) by an appropriate test function and integrating by parts in a half-annulus, it is easily seen that the left-hand side bounds the L^1 norm of u by below, modulus an error which is admissible for (1.13): the L^1 norm of $x \cdot \nabla u = ru_r$ in a half-annulus. The main difficulty is how to control the right-hand side of (1.15) by above, in terms of only u_r . This is delicate, due to the presence of $f'(u)$. We will explain the details in

Section 3. The proof involves introducing the functions $u_\lambda(x) := u(\lambda x)$, interpreting the radial derivative $x \cdot \nabla u(\lambda x)$ as $\frac{d}{d\lambda} u_\lambda(x)$, considering a “ u_λ -version” of (1.15), and averaging such version in λ to reduce the number of derivatives falling on the solution u through the proof arguments. This will be the crucial and key new idea in the proof: to average the inequalities also with respect to the variable λ .

Plan of the paper. In Section 2 we prove our interior estimate, Theorem 1.3. Section 3 is devoted to the boundary bound, Theorem 1.6. The paper contains also two Appendices (the first one includes a new interpolation inequality), with results that are needed in Section 3.

2. THE RADIAL DERIVATIVE CONTROLS THE FUNCTION IN L^1

In this section we prove, as claimed in Theorem 1.3, that the L^1 norm in an annulus (respectively, in a ball) of any superharmonic function can be controlled, up to an additive constant, by the L^1 norm of its radial derivative, taken also in the annulus (respectively, in the ball). For this, we first need a similar result for harmonic functions, with norms taken now on a sphere.

Lemma 2.1. *Let $v \in C^\infty(\overline{B}_1)$ solve $\Delta v = 0$ in $B_1 \subset \mathbb{R}^n$. Then,*

$$\|v - v(0)\|_{L^\infty(\partial B_1)} \leq 2n^{3/2} \|v_r\|_{L^\infty(\partial B_1)} \quad (2.1)$$

and

$$\|v - t\|_{L^1(\partial B_1)} \leq 2n^{3/2} \|v_r\|_{L^1(\partial B_1)}, \quad (2.2)$$

where $t := \inf \{\bar{t} : |\{v > \bar{t}\} \cap \partial B_1| \leq |\partial B_1|/2\}$ is the median of v .

The previous estimates should not be surprising. Indeed, we are controlling the solution to a boundary value problem for an homogeneous equation by its Neumann data $v_r|_{\partial B_1}$. Alternatively, the flux $v_r|_{\partial B_1}$ can be considered as a first-order elliptic integro-differential operator acting on the function $v|_{\partial B_1}$. This operator is a kind of “half-Laplacian”, since we extend the data $v|_{\partial B_1}$ harmonically in the ball—the half-Laplacian corresponds to the harmonic extension in a half-space. Anyhow, these comments will not be needed in the following elementary proof.

Proof of Lemma 2.1. We start proving (2.1). From it, we will easily deduce (2.2) by duality.

Let

$$w(x) := x \cdot \nabla v(x) = r v_r \quad \text{for } x \in \overline{B}_1.$$

We claim that

$$|w(x)| \leq 2n^{3/2} \|v_r\|_{L^\infty(\partial B_1)} r \quad \text{for } x \in \overline{B}_1; \quad (2.3)$$

recall that $r = |x|$. From this, (2.1) will clearly follow since, for all $\sigma \in \partial B_1$ we have

$$v(\sigma) - v(0) = \int_0^1 v_r(r\sigma) dr = \int_0^1 r^{-1} w(r\sigma) dr.$$

Now, to prove (2.3), note that w is harmonic in B_1 and agrees with v_r on ∂B_1 . Consequently,

$$\|w\|_{L^\infty(B_1)} \leq \|v_r\|_{L^\infty(\partial B_1)}. \quad (2.4)$$

It follows that (2.3) holds whenever $|x| = r \geq 1/2$.

Assume now that $x \in B_{1/2}$. Note that

$$|w(x)| = |w(x) - w(0)| \leq \|\nabla w\|_{L^\infty(B_{1/2})} r. \quad (2.5)$$

Using that the partial derivatives w_i of w are harmonic and also (2.4), we see that

$$\begin{aligned} |w_i(x)| &= \frac{1}{|B_{1/2}|} \left| \int_{B_{1/2}(x)} w_i dx \right| = \frac{1}{|B_{1/2}|} \left| \int_{\partial B_{1/2}(x)} w \nu^i d\mathcal{H}^{n-1} \right| \\ &\leq \frac{|\partial B_{1/2}|}{|B_{1/2}|} \|v_r\|_{L^\infty(\partial B_1)} = 2n \|v_r\|_{L^\infty(\partial B_1)}. \end{aligned}$$

Hence $|\nabla w(x)| \leq 2n^{3/2} \|v_r\|_{L^\infty(\partial B_1)}$, which combined with (2.5), gives (2.3) when $r < 1/2$.

Next, we establish (2.2) by a duality argument. First note that the value t , defined in the statement of Lemma 2.1, is finite and well-defined. Replacing v by $v - t$ we may assume that $t = 0$, and therefore that $0 = \inf \{\bar{t} : |\{v > \bar{t}\} \cap \partial B_1| \leq |\partial B_1|/2\}$. It follows that $|\{v > 0\} \cap \partial B_1| \leq |\partial B_1|/2$ and that $|\{v > \bar{t}\} \cap \partial B_1| > |\partial B_1|/2$ for all $\bar{t} < 0$. As a consequence, $|\{v \leq \bar{t}\} \cap \partial B_1| < |\partial B_1|/2$ for all $\bar{t} < 0$, and thus $|\{v < 0\} \cap \partial B_1| \leq |\partial B_1|/2$.

Let now $\text{sgn}(v) := v/|v|$ where $v \neq 0$. Since $|\{v > 0\} \cap \partial B_1| \leq |\partial B_1|/2$ and $|\{v < 0\} \cap \partial B_1| \leq |\partial B_1|/2$, it is possible to extend $\text{sgn}(v)$ to $\{v = 0\} \cap \partial B_1$, still taking values ± 1 and in such a way that we have $\int_{\partial B_1} \text{sgn}(v) d\mathcal{H}^{n-1} = 0$.⁴ In addition, it will also hold that $|v| = v \text{sgn}(v)$ on ∂B_1 .

We now define the functions

$$g_k(x) := \int_{\partial B_1} \text{sgn}(v)(y) \rho_k(|x - y|) d\mathcal{H}^{n-1}(y),$$

where $\{\rho_k = \rho_k(|\cdot|)\}$ is a sequence of smooth mollifiers on ∂B_1 . We have that $g_k \in C^\infty(\partial B_1)$, $|g_k| \leq 1$, and $\int_{\partial B_1} g_k d\mathcal{H}^{n-1} = 0$ since $\text{sgn}(v)$ has zero average on ∂B_1 . In addition, since $|v| = v \text{sgn}(v)$ a.e. on ∂B_1 , it holds that

$$\int_{\partial B_1} |v| d\mathcal{H}^{n-1} = \lim_k \int_{\partial B_1} v g_k d\mathcal{H}^{n-1}. \quad (2.6)$$

Now, since g_k is smooth and has zero average on ∂B_1 , we can uniquely solve the problem

$$\begin{cases} \Delta \varphi_k = 0 & \text{in } B_1 \\ \partial_r \varphi_k = g_k & \text{on } \partial B_1 \end{cases}$$

⁴This follows from the fact that, given a measurable set $A \subset \partial B_1$ (which we take here to be $\{v = 0\} \cap \partial B_1$) and $\theta \in (0, 1)$, there exists a measurable subset $B \subset A$ with $|B| = \theta|A|$. This follows from the continuity with respect to ρ of the quantity $|A \cap B_\rho(1, 0, \dots, 0)|$.

by imposing, additionally, $\varphi_k(0) = 0$. By (2.1), we have $\|\varphi_k\|_{L^\infty(\partial B_1)} \leq 2n^{3/2}$. We conclude that

$$\begin{aligned} \left| \int_{\partial B_1} v g_k d\mathcal{H}^{n-1} \right| &= \left| \int_{\partial B_1} v \partial_r \varphi_k d\mathcal{H}^{n-1} \right| = \left| \int_{B_1} \nabla v \cdot \nabla \varphi_k dx \right| = \left| \int_{\partial B_1} v_r \varphi_k d\mathcal{H}^{n-1} \right| \\ &\leq 2n^{3/2} \int_{\partial B_1} |v_r| d\mathcal{H}^{n-1}. \end{aligned}$$

This, together with (2.6), concludes the proof of (2.2). \square

We can now give the

Proof of Theorem 1.3. Since

$$\|u_r\|_{L^1(B_1 \setminus B_{1/2})} = \frac{1}{1/2} \int_{1/2}^1 \left(\int_{\partial B_t} \frac{1}{2} |u_r| d\mathcal{H}^{n-1} \right) dt,$$

there exists $\rho \in [1/2, 1]$ such that $\|u_r\|_{L^1(B_1 \setminus B_{1/2})} = \frac{1}{2} \int_{\partial B_\rho} |u_r| d\mathcal{H}^{n-1}$.

Let v be the harmonic function in B_ρ which agrees with u on ∂B_ρ . We have $v \in C^\infty(\overline{B_\rho})$. Decompose the radial derivatives $u_r = u_r^+ - u_r^-$ and $v_r = v_r^+ - v_r^-$ in their positive and negative parts. Since $v \leq u$ in B_ρ and the functions agree on the boundary, we have $v_r \geq u_r$ on ∂B_ρ . As a consequence, $v_r^- \leq u_r^-$ on ∂B_ρ . Since, in addition $0 = \int_{B_\rho} \Delta v dx = \int_{\partial B_\rho} v_r d\mathcal{H}^{n-1} = \int_{\partial B_\rho} v_r^+ d\mathcal{H}^{n-1} - \int_{\partial B_\rho} v_r^- d\mathcal{H}^{n-1}$, it follows that

$$\begin{aligned} 2\|u_r\|_{L^1(B_1 \setminus B_{1/2})} &= \int_{\partial B_\rho} |u_r| d\mathcal{H}^{n-1} \geq \int_{\partial B_\rho} u_r^- d\mathcal{H}^{n-1} \\ &\geq \int_{\partial B_\rho} v_r^- d\mathcal{H}^{n-1} = \int_{\partial B_\rho} v_r^+ d\mathcal{H}^{n-1} = \frac{1}{2} \int_{\partial B_\rho} |v_r| d\mathcal{H}^{n-1}. \end{aligned}$$

Now, since $\rho \in [1/2, 1]$, we can rescale the estimate (2.2) in Lemma 2.1 and apply it to the harmonic function v to deduce

$$C \int_{\partial B_\rho} |v_r| d\mathcal{H}^{n-1} \geq \int_{\partial B_\rho} |v - t| d\mathcal{H}^{n-1} = \int_{\partial B_\rho} |u - t| d\mathcal{H}^{n-1}$$

for some value t and a dimensional constant C , where in the last equality we used that $v = u$ on ∂B_ρ . Therefore, by the previous two estimates, the theorem will be proven once we establish that

$$\|u - t\|_{L^1(B_1 \setminus B_{1/2})} \leq C \left(\int_{\partial B_\rho} |u - t| d\mathcal{H}^{n-1} + \|u_r\|_{L^1(B_1 \setminus B_{1/2})} \right) \quad (2.7)$$

and

$$\|u - t\|_{L^1(B_{1/2})} \leq C \left(\int_{\partial B_\rho} |u - t| d\mathcal{H}^{n-1} + \|u_r\|_{L^1(B_1)} \right). \quad (2.8)$$

This is simple. For both estimates, we will use that

$$(u - t)(s\sigma) = (u - t)(\rho\sigma) - \int_s^\rho u_r(r\sigma) dr$$

for every $s \in (0, 1)$ and $\sigma \in S^{n-1}$.

Now, to check (2.7) we take $s \in (1/2, 1)$ and note that

$$s^{n-1}|(u-t)(s\sigma)| \leq 2^{n-1}\rho^{n-1}|(u-t)(\rho\sigma)| + 2^{n-1} \int_{1/2}^1 r^{n-1}|u_r(r\sigma)| dr.$$

Integrating in $\sigma \in S^{n-1}$, and then in $s \in (1/2, 1)$, we conclude (2.7).

Finally, to prove (2.8) we take $s \in (0, 1/2)$ and note that (since $\rho \geq 1/2 > s$)

$$\begin{aligned} s^{n-1}|(u-t)(s\sigma)| &\leq \rho^{n-1}|(u-t)(\rho\sigma)| + \int_s^\rho r^{n-1}|u_r(r\sigma)| dr \\ &\leq \rho^{n-1}|(u-t)(\rho\sigma)| + \int_0^1 r^{n-1}|u_r(r\sigma)| dr. \end{aligned}$$

Integrating in $\sigma \in S^{n-1}$, and then in $s \in (0, 1/2)$, we conclude (2.8) by using also (2.7). \square

3. THE RADIAL DERIVATIVE CONTROLS THE FUNCTION IN L^1 UP TO THE BOUNDARY

In this section we establish our main result, Theorem 1.6. This will be much more delicate than the interior case of Theorem 1.3, which holds assuming only that the function u is superharmonic. Instead, in the boundary setting, superharmonicity is not enough for the theorem to hold, as the following remark shows. Hence, within the proof we will need to use the semilinear equation satisfied by u and, in fact, also the stability of u .

With $\mathbb{R}_+^n = \{x \in \mathbb{R}^n : x_n > 0\}$ and $r = |x|$, throughout this section we use the notation

$$B_\rho^+ = \mathbb{R}_+^n \cap B_\rho, \quad A_{\rho, \bar{\rho}} := \{\rho < r < \bar{\rho}\}, \quad A_{\rho, \bar{\rho}}^+ := \{x_n > 0, \rho < r < \bar{\rho}\},$$

and

$$\partial^0 \Omega = \{x_n = 0\} \cap \partial \Omega \quad \text{and} \quad \partial^+ \Omega = \mathbb{R}_+^n \cap \partial \Omega$$

for an open set $\Omega \subset \mathbb{R}_+^n$.

Remark 3.1. For $n \geq 2$, the estimate (1.13) of Theorem 1.6, controlling the function in L^1 in a half-annulus by its radial derivative in L^1 (or even in L^∞), also in a half-annulus, cannot hold within the class of nonnegative superharmonic functions which are smooth in $\overline{\mathbb{R}_+^n}$ and vanish on $\partial \mathbb{R}_+^n$.

Indeed, for $\delta \in (0, 1)$, consider

$$u^\delta(x) := \frac{x_n}{|(x', x_n + \delta)|} \quad \text{for } x = (x', x_n) \in \mathbb{R}^{n-1} \times \mathbb{R}, x_n \geq 0.$$

The function u^δ is nonnegative and smooth in $\overline{\mathbb{R}_+^n}$, vanishes on $\{x_n = 0\}$, and, as a simple computation shows,⁵ is superharmonic in \mathbb{R}_+^n for $n \geq 2$. At the same time,

⁵One may start from $\Delta u^\delta = x_n \Delta \varphi + 2\varphi_{x_n}$, where $\varphi = (|x|^2 + 2\delta x_n + \delta^2)^{-1/2}$.

in the half-annulus it satisfies

$$|\partial_r u^\delta| = \frac{x_n}{r|(x', x_n + \delta)|^3} \delta(x_n + \delta) \leq C\delta \quad \text{in } A_{1/2,1}^+$$

for some constant C independent of δ . By taking δ small enough, this shows that the estimate (1.13) of Theorem 1.6 cannot hold within this class of functions.

Note also that in the limiting case $\delta = 0$, we are exhibiting a nonnegative superharmonic function $u^0 = x_n/r$ which belongs to $W^{1,2}(B_1^+)$ for $n \geq 3$, vanishes a.e. on $\{x_n = 0\}$, and is zero homogeneous (i.e., $\partial_r u^0 \equiv 0$).

When proving Theorem 1.6, we will need the following lemma. It follows easily from results of [CFRS] which were proved, there, in a quantitative manner.

Lemma 3.2. *Let $u \in C^\infty(\overline{B_1^+})$ be a nonnegative stable solution of $-\Delta u = f(u)$ in $B_1^+ \subset \mathbb{R}^n$, with $u = 0$ on $\partial^0 B_1^+$. Assume that $f \in C^1(\mathbb{R})$ is nonnegative and nondecreasing. Let $0 < \rho_1 < \rho_2 < \rho_3 < \rho_4 \leq 1$.*

Then,

$$\|\nabla u\|_{L^{2+\gamma}(A_{\rho_2, \rho_3}^+)} \leq C_{\rho_i} \|u\|_{L^1(A_{\rho_1, \rho_4}^+)} \quad (3.1)$$

and

$$\|D^2 u\|_{L^1(A_{\rho_2, \rho_3}^+)} \leq C_{\rho_i} \|u\|_{L^1(A_{\rho_1, \rho_4}^+)} \quad (3.2)$$

for some dimensional constant $\gamma > 0$ and some constant C_{ρ_i} depending only on n , ρ_1 , ρ_2 , ρ_3 , and ρ_4 .

Proof. First of all, by [CFRS, Propositions 2.5 and 5.5] (and a covering of the half-annulus by balls) it suffices to prove the estimates of the lemma with the L^1 norm of u on their right-hand sides replaced by the L^2 norm of ∇u (in a slightly smaller annulus). This new estimates will follow from a simple covering and scaling argument combined with the next interior and boundary estimates.

For (3.1) we use the $W^{1,2+\gamma}$ estimate of Propositions 2.4 in [CFRS] for the interior balls, and of Propositions 5.2 in [CFRS] for the boundary balls.

Next, to show (3.2), let $\nu = \nabla u/|\nabla u|$ in the set $\{\nabla u \neq 0\}$ and $\nu = 0$ in $\{\nabla u = 0\}$. Recall the definition of the quantity \mathcal{A} in (2.7) of [CFRS]. Take an orthonormal basis of \mathbb{R}^n where the last basic vector is ν . Since $D^2 u$ is a symmetric matrix and \mathcal{A}^2 contains all the squares of its elements except for those in the last column (which also appear in the last row), we conclude that, up to a multiplicative constant, \mathcal{A}^2 controls all the squares u_{ij}^2 in the Hessian matrix except for $u_{\nu\nu}^2$. As a consequence, writing $u_{\nu\nu}$ as Δu minus the sum of all other diagonal elements, we see that

$$|D^2 u| \leq |\Delta u| + C\mathcal{A} \quad \text{a.e. in } B_1^+, \quad (3.3)$$

for some dimensional constant C .

Now, choose a nonnegative function $\zeta \in C_c^\infty(B_{1/2})$ with $\zeta \equiv 1$ in $B_{1/4}$. Then, since $-\Delta u \geq 0$ in B_1^+ , we have

$$\int_{B_{1/4}} |\Delta u| dx \leq - \int_{B_{1/2}} \Delta u \zeta dx = \int_{B_{1/2}} \nabla u \cdot \nabla \zeta dx \leq C \|\nabla u\|_{L^2(B_{1/2})}. \quad (3.4)$$

This bound, (3.3), and [CFRS, Lemma 2.3] lead to the interior estimate needed to prove (3.2) (after a standard covering and scaling argument to replace the balls $B_{1/4}$ and $B_{1/2}$ in (3.4) by balls B_{r_1} and B_{r_2} , respectively, for any $r_1 < r_2$).

To complete the proof of (3.2), we need the analogue Hessian boundary estimate. Since $-\Delta u \geq 0$ in B_1^+ , with ζ as above we see that

$$\begin{aligned} \|\Delta u\|_{L^1(B_{1/4}^+)} &\leq \int_{B_{1/2}^+} (-\Delta u) \zeta \, dx = - \int_{\partial^0 B_{1/2}^+} u_\nu \zeta \, d\mathcal{H}^{n-1} + \int_{B_{1/2}^+} \nabla u \cdot \nabla \zeta \, dx \\ &\leq C \|\nabla u\|_{L^2(B_1^+)}, \end{aligned} \quad (3.5)$$

where we have used [CFRS, Lemma 5.3] (rescaled) to control u_ν . This bound, (3.3), and estimate (5.14) of [CFRS] for \mathcal{A} give the boundary estimate needed to prove (3.2) after a covering and scaling argument. \square

Towards the proof of Theorem 1.6, to control the L^1 norm of a stable solution in a half-annulus by its radial derivative in L^1 , the starting idea is to use the equation

$$-2\Delta u + \Delta(x \cdot \nabla u) = -f'(u) x \cdot \nabla u, \quad (3.6)$$

after multiplying it against an appropriate test function and integrating it in the half-annulus. We will see that this easily yields a lower bound for the integral of the left-hand side which is appropriate for our purposes. The difficulty is how to control the integral of the right-hand side, by above, in terms of only $x \cdot \nabla u = ru_r$. As we will explain later in Footnote 6, we wish to use the stability condition to deal with the factor $f'(u)$ in (3.6). However, through the simplest approach this would force to control the L^2 norm of $\nabla(x \cdot \nabla u)$ —which is not at our hands since we do not have L^2 control on the full Hessian of u (but only the L^1 control of Lemma 3.2).

To proceed in a similar manner but reducing the number of derivatives falling on u , for $\lambda > 0$ we consider the functions

$$u_\lambda(x) := u(\lambda x)$$

and note that $\frac{d}{d\lambda} u_\lambda(x) = x \cdot \nabla u(\lambda x) = \lambda^{-1} x \cdot \nabla u_\lambda(x)$. Using this and noticing that $\lambda^{-2} \Delta u_\lambda = -f(u_\lambda)$, we have

$$\begin{aligned} -2\lambda^{-3} \Delta u_\lambda + \lambda^{-2} \Delta(\lambda^{-1} x \cdot \nabla u_\lambda) &= \frac{d}{d\lambda} (\lambda^{-2} \Delta u_\lambda) \\ &= -\frac{d}{d\lambda} f(u_\lambda) = -f'(u_\lambda) \lambda^{-1} x \cdot \nabla u_\lambda. \end{aligned} \quad (3.7)$$

Evaluating the first and last expressions at $\lambda = 1$, we recover (3.6). However, it will be crucial (see the later Footnote 6) to use instead the equality between the first and third expressions, after integrating them not only in x , but also in λ . Integrating in λ will be essential to reduce the number of derivatives falling on u . In addition, the monotonicity and convexity of f will allow to use the stability condition appropriately.

This will be understood in all detail going through the following proof.

Proof of Theorem 1.6. By rescaling, we may suppose that we have a stable solution in B_6^+ instead of B_1^+ .

We choose a nonnegative smooth function ζ with compact support in the annulus $A_{4,5}$ and such that $\zeta \equiv 1$ in $A_{4.1,4.9}$. Then, the function $\xi := x_n \zeta$ satisfies

$$\begin{aligned} \xi &\geq 0 \text{ in } A_{4,5}^+, \quad \xi = 0 \text{ on } \partial^0 A_{4,5}^+, \\ \xi &= \xi_\nu = 0 \text{ on } \partial^+ A_{4,5}^+, \quad \text{and} \quad \xi = x_n \text{ in } A_{4.1,4.9}^+. \end{aligned}$$

The proof starts from the identity

$$\begin{aligned} 2\lambda^{-3} \int_{A_{4,5}^+} (-\Delta u_\lambda) \xi \, dx + \lambda^{-2} \int_{A_{4,5}^+} \Delta(\lambda^{-1}x \cdot \nabla u_\lambda) \xi \, dx \\ = -\frac{d}{d\lambda} \int_{A_{4,5}^+} f(u_\lambda) \xi \, dx, \end{aligned} \quad (3.8)$$

which follows from the equality between the first and third expressions in (3.7). In a first step, we will bound the left-hand side of (3.8) by below. But the subtle part of the proof, where we use the stability of the solution, will be the second step. It will bound the right-hand side of (3.8) by above, but only after averaging it in λ . For this, we will use that

$$\int_1^{1.1} \left(-\frac{d}{d\lambda} \int_{A_{4,5}^+} f(u_\lambda) \xi \, dx \right) d\lambda = \int_{A_{4,5}^+} (f(u) - f(u_{1.1})) \xi \, dx, \quad (3.9)$$

where

$$u_{1.1} := u_{11/10} = u((11/10) \cdot).$$

Step 1: We prove that

$$\begin{aligned} 2\lambda^{-3} \int_{A_{4,5}^+} (-\Delta u_\lambda) \xi \, dx + \lambda^{-2} \int_{A_{4,5}^+} \Delta(\lambda^{-1}x \cdot \nabla u_\lambda) \xi \, dx \\ \geq c \|u\|_{L^1(A_{4.7,4.8}^+)} - C \|u_r\|_{L^1(A_{3,6}^+)} \end{aligned} \quad (3.10)$$

for every $\lambda \in [1, 1.1]$, where c and C are positive dimensional constants.

To bound by below the second integral in (3.10) is easy. Since ξ and ξ_ν vanish on $\partial^+ A_{4,5}^+$, and both ξ and $x \cdot \nabla u_\lambda$ vanish on $\partial^0 A_{4,5}^+$, we deduce that

$$\int_{A_{4,5}^+} \Delta(x \cdot \nabla u_\lambda) \xi \, dx = \int_{A_{4,5}^+} x \cdot \nabla u_\lambda \Delta \xi \, dx \geq -C \|u_r\|_{L^1(A_{4.5,5.5}^+)} \quad (3.11)$$

since $x \cdot \nabla u_\lambda(x) = \lambda x \cdot \nabla u(\lambda x) = \lambda r u_r(\lambda x)$.

Next, to control the first integral in (3.10), given any $\rho_1 \in (4.1, 4.2)$ and $\rho_2 \in (4.8, 4.9)$ we consider the solution φ of

$$\begin{cases} -\Delta \varphi = 1 & \text{in } A_{\rho_1, \rho_2}^+ \\ \varphi = 0 & \text{on } \partial^0 A_{\rho_1, \rho_2}^+ \\ \varphi_\nu = 0 & \text{on } \partial^+ A_{\rho_1, \rho_2}^+. \end{cases} \quad (3.12)$$

Note that $\varphi \geq 0$ by the maximum principle. In addition, as shown in Appendix B, we have $|\nabla\varphi| \leq C$ in A_{ρ_1, ρ_2}^+ for some dimensional constant C . This bound yields $c\varphi \leq x_n = \xi$ in A_{ρ_1, ρ_2}^+ for some small dimensional constant $c > 0$. Since, in addition, $-\Delta u_\lambda$, ξ , and φ are all nonnegative, it follows that (for positive constants c and C that may differ from line to line)

$$\begin{aligned}
\int_{A_{4,5}^+} (-\Delta u_\lambda) \xi \, dx &\geq c \int_{A_{\rho_1, \rho_2}^+} (-\Delta u_\lambda) \varphi \, dx \\
&= -c \int_{\partial^+ A_{\rho_1, \rho_2}^+} (u_\lambda)_\nu \varphi \, d\mathcal{H}^{n-1} + c \int_{A_{\rho_1, \rho_2}^+} u_\lambda \, dx \\
&\geq -C \int_{\partial^+ B_{\rho_1}^+} |(u_\lambda)_r| \, d\mathcal{H}^{n-1} - C \int_{\partial^+ B_{\rho_2}^+} |(u_\lambda)_r| \, d\mathcal{H}^{n-1} + c \|u\|_{L^1(A_{\lambda\rho_1, \lambda\rho_2}^+)} \\
&\geq -C \int_{\partial^+ B_{\rho_1}^+} |(u_\lambda)_r| \, d\mathcal{H}^{n-1} - C \int_{\partial^+ B_{\rho_2}^+} |(u_\lambda)_r| \, d\mathcal{H}^{n-1} + c \|u\|_{L^1(A_{4.7, 4.8}^+)}
\end{aligned}$$

since $\lambda\rho_1 \leq 1.1 \cdot 4.2 \leq 4.7$ and $4.8 \leq \lambda\rho_2$. Finally, integrating first in $\rho_1 \in (4.1, 4.2)$ and then in $\rho_2 \in (4.8, 4.9)$, we arrive at

$$\int_{A_{4,5}^+} (-\Delta u_\lambda) \xi \, dx \geq -C \|u_r\|_{L^1(A_{4.1, 6}^+)} + c \|u\|_{L^1(A_{4.7, 4.8}^+)},$$

since $1.1 \cdot 4.9 \leq 6$.

This and (3.11) establish the claim of Step 1.

Step 2: We prove that, for every $\varepsilon \in (0, 1)$,

$$\int_{A_{4,5}^+} (f(u) - f(u_{1.1})) \xi \, dx \leq C(\varepsilon \|u\|_{L^1(A_{3,6}^+)} + \varepsilon^{-1-2\frac{2+\gamma}{\gamma}} \|u_r\|_{L^1(A_{3,6}^+)}) \quad (3.13)$$

for some dimensional constants $\gamma > 0$ and C —with γ being the exponent in Lemma 3.2.

By convexity of f we have $f(u) - f(u_{1.1}) \leq f'(u)(u - u_{1.1})$. We now use that $\xi = 0$ on $\partial A_{4,5}^+$ and that $u - u_{1.1} = 0$ on $\partial^0 A_{3.9, 5.1}^+$ in order to take advantage, twice, of the stability of u . Taking a function $\phi \in C_c^\infty(A_{3.9, 5.1})$ with $\phi = 1$ in $A_{4,5}$, and

since ξ and $f'(u)$ are nonnegative, we deduce

$$\int_{A_{4,5}^+} (f(u) - f(u_{1.1}))\xi \, dx \leq \int_{A_{4,5}^+} f'(u)(u - u_{1.1})\xi \, dx \quad (3.14)$$

$$\begin{aligned} &\leq \left(\int_{A_{4,5}^+} f'(u)\xi^2 \, dx \right)^{1/2} \left(\int_{A_{4,5}^+} f'(u)(u - u_{1.1})^2 \, dx \right)^{1/2} \\ &\leq \left(\int_{A_{4,5}^+} |\nabla \xi|^2 \, dx \right)^{1/2} \left(\int_{A_{3.9,5.1}^+} f'(u)((u - u_{1.1})\phi)^2 \, dx \right)^{1/2} \\ &\leq C \left(\int_{A_{3.9,5.1}^+} |\nabla((u - u_{1.1})\phi)|^2 \, dx \right)^{1/2} \\ &\leq C \|\nabla(u - u_{1.1})\|_{L^2(A_{3.9,5.1}^+)}, \end{aligned} \quad (3.15)$$

where in the last bound we have used Poincaré's inequality in $A_{3.9,5.1}^+$ for functions vanishing on $\partial^0 A_{3.9,5.1}^+$. The previous chain of inequalities, which are a crucial part of the proof, have used the stability of the solution twice.⁶

Next, by the $W^{1,2+\gamma}$ estimate (3.1) (rescaled to hold in B_6^+ instead of B_1^+) and taking $q := \frac{2(1+\gamma)}{2+\gamma}$, we have

$$\begin{aligned} \|\nabla(u - u_{1.1})\|_{L^2(A_{3.9,5.1}^+)} &\leq \|\nabla(u - u_{1.1})\|_{L^{2+\gamma}(A_{3.9,5.1}^+)}^{\frac{1}{q}} \|\nabla(u - u_{1.1})\|_{L^1(A_{3.9,5.1}^+)}^{\frac{1}{q'}} \\ &\leq C \|\nabla u\|_{L^{2+\gamma}(A_{3.9,5.1,1.1}^+)}^{\frac{1}{q}} \|\nabla(u - u_{1.1})\|_{L^1(A_{3.9,5.1}^+)}^{\frac{1}{q'}} \\ &\leq C \|u\|_{L^1(A_{3,6}^+)}^{\frac{1}{q}} \|\nabla(u - u_{1.1})\|_{L^1(A_{3.9,5.1}^+)}^{\frac{1}{q'}} \\ &\leq \varepsilon \|u\|_{L^1(A_{3,6}^+)} + C\varepsilon^{-\frac{q'}{q}} \|\nabla(u - u_{1.1})\|_{L^1(A_{3.9,5.1}^+)} \end{aligned} \quad (3.16)$$

for all $\varepsilon \in (0, 1)$. Now, by the interpolation inequality in cubes of Proposition A.1 applied with $p = 1$, we claim that

$$\begin{aligned} &\|\nabla(u - u_{1.1})\|_{L^1(A_{3.9,5.1}^+)} \\ &\leq C\varepsilon^{1+\frac{q'}{q}} \|D^2(u - u_{1.1})\|_{L^1(A_{3.8,5.2}^+)} + C\varepsilon^{-1-\frac{q'}{q}} \|u - u_{1.1}\|_{L^1(A_{3.8,5.2}^+)} \\ &\leq C\varepsilon^{1+\frac{q'}{q}} \|D^2 u\|_{L^1(A_{3.8,5.8}^+)} + C\varepsilon^{-1-\frac{q'}{q}} \|u - u_{1.1}\|_{L^1(A_{3.8,5.2}^+)}. \end{aligned} \quad (3.17)$$

⁶Note first that a simple Cauchy-Schwarz argument to control the right-hand side of (3.14) would not work, since we do not have control on the integral of $f'(u)^2$. On the other hand, we would be in trouble if we had performed the above chain of inequalities starting from (3.6) instead of (3.7). Indeed, in such case, $f'(u)(u - u_{1.1})\xi$ in (3.14) would be replaced by $f'(u)x \cdot \nabla u \xi$. Hence, proceeding as we have done above, the final quantity appearing in (3.15) would be the L^2 norm of $\nabla(x \cdot \nabla u)$. But recall that we do not have control on the L^2 norm of the full Hessian of u .

To see the first inequality, one covers the half-annulus $A_{3,9,5,1}^+$ (except for a set of measure zero) by cubes of sufficiently small side-length to be contained in $A_{3,8,5,2}^+$. One then applies Proposition A.1 with $p = 1$, after rescaling it and renaming ε , in each of these cubes, and finally one adds up all the inequalities.

From (3.17) and the Hessian bound (3.2) (rescaled to hold in B_6^+ instead of B_1^+) we conclude

$$\|\nabla(u - u_{1.1})\|_{L^1(A_{3,9,5,1}^+)} \leq C\varepsilon^{1+\frac{q'}{q}}\|u\|_{L^1(A_{3,6}^+)} + C\varepsilon^{-1-\frac{q'}{q}}\|u - u_{1.1}\|_{L^1(A_{3,8,5,2}^+)}.$$

Putting together this last bound with (3.15) and (3.16), and using again (3.2), we arrive at

$$\int_{A_{4,5}^+} (f(u) - f(u_{1.1}))\xi \, dx \leq C\varepsilon\|u\|_{L^1(A_{3,6}^+)} + C\varepsilon^{-1-2\frac{2+\gamma}{\gamma}}\|u - u_{1.1}\|_{L^1(A_{3,8,5,2}^+)}.$$

At the same time, since $\frac{d}{d\lambda}u_\lambda(x) = x \cdot \nabla u(\lambda x) = r u_r(\lambda x)$, we have

$$\begin{aligned} \|u - u_{1.1}\|_{L^1(A_{3,8,5,2}^+)} &= \int_{A_{3,8,5,2}^+} dx \left| \int_1^{1.1} d\lambda r u_r(\lambda x) \right| \\ &\leq C \int_1^{1.1} d\lambda \int_{A_{3,8,5,2,1.1}^+} dy |u_r(y)| \leq C\|u_r\|_{L^1(A_{3,6}^+)}. \end{aligned}$$

The last two bounds establish Step 2.

Step 3: Conclusion.

We integrate (3.8) in $\lambda \in (1, 1.1)$ and use (3.9), as well as (3.10) and (3.13) in the statements of Steps 1 and 2. We obtain the bound

$$\|u\|_{L^1(A_{4,7,4,8}^+)} \leq C(\varepsilon\|u\|_{L^1(A_{3,6}^+)} + \varepsilon^{-1-2\frac{2+\gamma}{\gamma}}\|u_r\|_{L^1(A_{3,6}^+)}) \quad (3.18)$$

for all $\varepsilon \in (0, 1)$. It is now simple to conclude the desired estimate

$$\|u\|_{L^1(A_{3,6}^+)} \leq C\|u_r\|_{L^1(A_{3,6}^+)}. \quad (3.19)$$

Indeed, we first show that

$$\|u\|_{L^1(A_{3,6}^+)} \leq C(\|u\|_{L^1(A_{4,7,4,8}^+)} + \|u_r\|_{L^1(A_{3,6}^+)}). \quad (3.20)$$

To prove this, use that

$$u(s\sigma) = u(t\sigma) - \int_s^t u_r(r\sigma) \, dr \quad (3.21)$$

for $s \in (3, 6)$, $t \in (4.7, 4.8)$, and $\sigma \in S^{n-1}$. We deduce

$$(s/6)^{n-1}|u(s\sigma)| \leq (t/4)^{n-1}|u(t\sigma)| + \int_3^6 (r/3)^{n-1}|u_r(r\sigma)| \, dr.$$

Integrating in $\sigma \in S^{n-1}$, and then in $s \in (3, 6)$ and in $t \in (4.7, 4.8)$, we conclude (3.20).

Now, we use (3.20) to bound the right-hand side of (3.18). In the resulting inequality we choose $\varepsilon \in (0, 1)$ small enough so that the constant $C\varepsilon$ multiplying $\|u\|_{L^1(A_{4.7,4.8}^+)}$ satisfies $C\varepsilon \leq 1/2$. We deduce

$$\|u\|_{L^1(A_{4.7,4.8}^+)} \leq C\|u_r\|_{L^1(A_{3,6}^+)},$$

that, together with (3.20), yields (3.19).

Next, to show $\|u\|_{L^1(B_6^+)} \leq C\|u_r\|_{L^1(B_6^+)}$, by (3.19) it suffices to show $\|u\|_{L^1(B_3^+)} \leq C\|u_r\|_{L^1(B_6^+)}$. Using again (3.19), it is indeed enough to show

$$\|u\|_{L^1(B_3^+)} \leq C \left(\|u\|_{L^1(A_{3,6}^+)} + \|u_r\|_{L^1(B_6^+)} \right).$$

This is easily shown using (3.21), with a similar argument as above or as in the interior case of Theorem 1.3.

The proven bounds hold for a solution in B_6^+ . By rescaling we conclude the estimates of the theorem for solutions in B_1^+ . \square

For $n \geq 3$, the estimate that we have just proven can be improved as follows.

Remark 3.3. When $n \geq 3$, Theorem 1.6 also holds when replacing $\|u\|_{L^1(A_{1/2,1}^+)}$, in the left-hand side of (1.13), by $\|u\|_{L^1(B_1^+)}$.

To show this, and given the theorem, it suffices to prove

$$\|u\|_{L^1(B_{1/2}^+)} \leq C\|u\|_{L^1(A_{1/2,1}^+)}.$$

But notice that, by the Poincaré inequality for functions vanishing on $\partial^0 B_{1/2}^+$, we have that

$$\|u\|_{L^1(B_{1/2}^+)} \leq C\|\nabla u\|_{L^2(B_{1/2}^+)} \leq C\|\nabla u\|_{L^2(B_{4/6}^+)}. \quad (3.22)$$

Now, we use the estimate

$$\int_{B_\rho^+} |\nabla u|^2 dx \leq C_\lambda \int_{A_{\rho,\lambda\rho}^+} |\nabla u|^2 dx \quad \text{if } n \geq 3, \quad (3.23)$$

where $\rho \in (0, 1/\lambda)$, $\lambda > 1$ is given, and C_λ is a constant depending only on n and λ . Such estimate follows from Lemma 6.2 in [CFRS], applied with $\Omega = B_1^+$ and $\theta = 0$. Indeed, take $\psi \in C_c^\infty(B_\lambda)$ to be a nonnegative radially nonincreasing function with $\psi \equiv 1$ in B_1 . For $\rho \in (0, 1/\lambda)$, set $\psi_\rho(x) := \psi(x/\rho)$. Note that $|\nabla \psi_\rho| \leq C_\lambda/\rho$ (where C_λ depends only on λ) and that $\nabla \psi_\rho$ vanishes outside of the annulus $A_{\rho,\lambda\rho}$. Choosing $\eta = \psi_\rho$ in the inequality in [CFRS, Lemma 6.2] with $\theta = 0$, for $n \geq 3$ we immediately deduce (3.23).

Thus, we can bound the last quantity in (3.22) computed in a ball by the L^2 norm $C\|\nabla u\|_{L^2(A_{4/6,5/6}^+)}$ computed in an annulus. Finally, by (3.1) we know that

$$\|\nabla u\|_{L^2(A_{4/6,5/6}^+)} \leq C\|u\|_{L^1(A_{1/2,1}^+)},$$

which leads to our claim.

APPENDICES

APPENDIX A. TWO SIMPLE INTERPOLATION INEQUALITIES IN CUBES

In contrast with [CFRS], the new quantitative proof of the interior result Theorem 1.1 given in [C] does not use the $W^{1,2+\gamma}$ bound—but just the $W^{1,2}$ estimate. This simplification is accomplished thanks to a new interpolation inequality in cubes, which has been used also in our boundary regularity proof of the previous section.

The inequality concerns the quantity $\int |\nabla u|^{p-1} |D^2 u| dx$. It is stated in cubes, it does not assume any boundary values, and it is extremely simple to be established. Indeed, in contrast with other interpolation results for functions with no prescribed boundary values, such as [GT, Theorem 7.28], proving our inequality in a cube of \mathbb{R}^n will be immediate once we prove it in dimension one.⁷

We conceived the inequality recently, in our work [CMS, Proposition A.3], in order to extend the results of [CFRS] to the $(p+1)$ -Laplacian operator.⁸ We do not know if the interpolation inequality has appeared somewhere before.

Proposition A.1. *Let $Q = (0, 1)^n \subset \mathbb{R}^n$, $p \geq 1$, and $u \in C^2(\overline{Q})$.*

Then, for every $\varepsilon \in (0, 1)$,

$$\int_Q |\nabla u|^p dx \leq C_p \left(\varepsilon \int_Q |\nabla u|^{p-1} |D^2 u| dx + \frac{1}{\varepsilon^p} \int_Q |u|^p dx \right), \quad (\text{A.1})$$

where C_p is a constant depending only on n and p .

Proof. We first prove it for $n = 1$, in the interval $(0, \delta)$ for a given $\delta \in (0, 1)$. Let $u \in C^2([0, \delta])$. Let $x_0 \in [0, \delta]$ be such that $|u'(x_0)| = \min_{[0, \delta]} |u'|$. For $0 < y < \frac{\delta}{3} < \frac{2\delta}{3} < z < \delta$, since $(u(z) - u(y))/(z - y)$ is equal to u' at some point, we deduce that $|u'(x_0)| \leq 3\delta^{-1}(|u(y)| + |u(z)|)$. Integrating this inequality first in y and later in z , we see that $|u'(x_0)| \leq 9\delta^{-2} \int_0^\delta |u| dx$. Raising this inequality to the power $p \in [1, \infty)$ we get

$$|u'(x_0)|^p \leq 9^p \delta^{-p-1} \int_0^\delta |u|^p dx. \quad (\text{A.2})$$

Now, for $x \in (0, \delta)$, integrating $(|u'|^p)'$ in the interval with end points x_0 and x , we deduce

$$|u'(x)|^p \leq p \int_0^\delta |u'|^{p-1} |u''| dx + |u'(x_0)|^p.$$

⁷Instead, the interpolation inequality [GT, Theorem 7.28] for functions with arbitrary boundary values requires a result on Sobolev seminorms for extension operators (i.e., operators extending functions of n variables to a larger domain).

⁸The proof that we give here in cubes is exactly the same as that of [CMS]. However, [CMS] claimed the same interpolation inequality in balls instead of cubes, and this made the proof (and perhaps the statement) in [CMS] not to be correct. Indeed, proving inequality (A.5) of [CMS] contains a mistake: it fails when, given $\varepsilon \in (0, 1)$, one has $|I| \ll \varepsilon$ —as it occurs later in the proof in \mathbb{R}^n when 1d sections of the ball are very small compared to ε . Despite this, all other results of [CMS] remain correct since, within Step 1 of the proof of Theorem 1.1 in [CMS], the interpolation inequality may be applied in small enough cubes covering the ball $B_{1/2}$ (as we do in the current paper) instead of applying it in the whole ball.

Combining this inequality with (A.2) and integrating in $x \in (0, \delta)$, we obtain

$$\int_0^\delta |u'|^p dx \leq p \delta \int_0^\delta |u'|^{p-1} |u''| dx + 9^p \delta^{-p} \int_0^\delta |u|^p dx. \quad (\text{A.3})$$

We come back now to the statement of the proposition in dimension one and let $u \in C^2([0, 1])$. For any given integer $k > 1$ we divide $(0, 1)$ into k disjoint intervals of length $\delta = 1/k$. Since for (A.3) we did not require any specific boundary values for u , we can use the inequality in each of these intervals of length $\delta = 1/k$ (instead of in $(0, \delta)$), and then add up all the inequalities, to deduce

$$\int_0^1 |u'|^p dx \leq \frac{p}{k} \int_0^1 |u'|^{p-1} |u''| dx + (9k)^p \int_0^1 |u|^p dx. \quad (\text{A.4})$$

Since $0 < \varepsilon < 1$, there exists an integer $k > 1$ such that $\frac{1}{\varepsilon} \leq k < \frac{2}{\varepsilon}$. This and (A.4) establish the proposition in dimension one.

Finally, for $u \in C^2([0, 1]^n)$, denote $x = (x_1, x') \in \mathbb{R} \times \mathbb{R}^{n-1}$. Using (A.1) with $n = 1$ for every x' , we get

$$\begin{aligned} \int_Q |u_{x_1}|^p dx &= \int_{(0,1)^{n-1}} dx' \int_0^1 dx_1 |u_{x_1}(x)|^p \\ &\leq C_p \varepsilon \int_{(0,1)^{n-1}} dx' \int_0^1 dx_1 |u_{x_1}(x)|^{p-1} |u_{x_1 x_1}(x)| \\ &\quad + C_p \varepsilon^{-p} \int_{(0,1)^{n-1}} dx' \int_0^1 dx_1 |u(x)|^p \\ &= C_p \left(\varepsilon \int_Q |u_{x_1}(x)|^{p-1} |u_{x_1 x_1}(x)| dx + \varepsilon^{-p} \int_Q |u(x)|^p dx \right). \end{aligned}$$

Since the same inequality holds for the partial derivatives with respect to each variable x_k instead of x_1 , adding up the inequalities we conclude (A.1). \square

APPENDIX B. GRADIENT ESTIMATE FOR A MIXED BOUNDARY VALUE PROBLEM

Here we show that the solution to (3.12), i.e.,

$$\begin{cases} -\Delta \varphi = 1 & \text{in } A_{\rho_1, \rho_2}^+ \\ \varphi = 0 & \text{on } \partial^0 A_{\rho_1, \rho_2}^+ \\ \varphi_\nu = 0 & \text{on } \partial^+ A_{\rho_1, \rho_2}^+, \end{cases} \quad (\text{B.1})$$

with $\rho_1 \in (4.1, 4.2)$ and $\rho_2 \in (4.8, 4.9)$ —that we used in Step 1 of the proof of Theorem 1.6—satisfies $|\nabla \varphi| \leq C$ in A_{ρ_1, ρ_2}^+ for some dimensional constant C . Note that the equation is posed in a nonsmooth domain and that this is a mixed Dirichlet-Neumann problem.

To establish this gradient estimate, we first need to control the L^∞ norm of φ . For this, recall that, by the maximum principle, $\varphi \geq 0$. Multiplying the equation in

(B.1) by φ^β , with $\beta \geq 1$, and integrating by parts, we get

$$\int_{A_{\rho_1, \rho_2}^+} |\nabla(\varphi^{(\beta+1)/2})|^2 dx \leq \frac{(\beta+1)^2}{4\beta} \int_{A_{\rho_1, \rho_2}^+} \varphi^\beta dx. \quad (\text{B.2})$$

We now proceed with a standard iteration procedure, as in the proof of Theorem 8.15 in [GT]. To bound by below the left-hand side of (B.2), we use the Sobolev inequality for functions vanishing on $\partial^0 A_{\rho_1, \rho_2}^+$ and also that the averaged L^p norms of φ are increasing in p and, hence, greater or equal than $\|\varphi\|_{L^1}$. Note also that $\|\varphi\|_{L^1} = \|\nabla\varphi\|_{L^2}^2$ is a positive universal quantity —the solution can be found variationally and thus belongs to $W^{1,2}$. In this way, we obtain the recurrence inequality $\|\varphi\|_{L^{\chi\cdot\beta}} \leq (C\beta)^{1/\beta} \|\varphi\|_{L^\beta}$ with $\chi = n/(n-2) > 1$. As in the proof of Theorem 8.15 in [GT], this leads to an L^∞ bound for φ .

Now, a simple way to prove the gradient estimate consists of considering the function $\tilde{\varphi} := \varphi + x_n^2/2$, which is harmonic and vanishes on the flat boundary. Reflecting it oddly, it becomes a harmonic function in the full annulus A_{ρ_1, ρ_2} (a smooth domain) and has $\tilde{\varphi}_r = |x_n|x_n/r$ as Neumann data on its boundary. Since this data is a $C^{1,\alpha}$ function, we can now apply global Schauder's regularity theory for the Neumann problem. We use Theorems 6.30 and 6.31 in [GT] to get a $C^{2,\alpha}$ estimate for $\tilde{\varphi}$ up to the boundary —and, in particular, a global gradient estimate. The $C^{2,\alpha}$ bound depends only on the $C^{1,\alpha}$ norm of the Neumann data, the L^∞ norm of $\tilde{\varphi}$ (that we already controlled), and the smoothness of the annulus A_{ρ_1, ρ_2} (which is uniform in ρ_1 and ρ_2 due to the constraints we placed on these parameters).

REFERENCES

- [C] Cabré, Xavier. A quantitative proof of the Hölder regularity of stable solutions to semilinear elliptic equations. *arXiv:2205.11352 [math.AP]*.
- [CFRS] Cabré, Xavier; Figalli, Alessio; Ros-Oton, Xavier; Serra, Joaquim. Stable solutions to semilinear elliptic equations are smooth up to dimension 9. *Acta Math.* 224 (2020), 187–252.
- [CMS] Cabré, Xavier; Miraglio, Pietro; Sanchón, Manel. Optimal regularity of stable solutions to nonlinear equations involving the p-Laplacian. *Adv. Calc. Var.* (2020), <https://doi.org/10.1515/acv-2020-0055>.
- [D] Dupaigne, Louis. *Stable Solutions of Elliptic Partial Differential Equations*. Chapman & Hall/CRC Monogr. Surv. Pure Appl. Math. 143, CRC Press, Boca Raton, 2011.
- [E1] Erneta, Iñigo U. Stable solutions to semilinear elliptic equations for operators with variable coefficients. *arXiv:2206.01572 [math.AP]*.
- [E2] Erneta, Iñigo U. Boundary regularity for stable solutions to semilinear elliptic equations. *Forthcoming*.
- [GT] Gilbarg, David; Trudinger, Neil S. *Elliptic partial differential equations of second order*. Reprint of the 1998 edition. Classics in Mathematics. Springer-Verlag, Berlin, 2001.
- [L] Leoni, Giovanni. *A first course in Sobolev spaces*. Graduate Studies in Mathematics, 105. American Mathematical Society, Providence, RI, 2009.
- [P] Peng, Fa. The boundedness of stable solutions to semilinear elliptic equations with linear lower bound on nonlinearities. *Preprint*.

X.C.^{1,2,3} — ¹ICREA, PG. LLUIS COMPANYYS 23, 08010 BARCELONA, SPAIN & ²UNIVERSITAT POLITÈCNICA DE CATALUNYA, DEPARTAMENT DE MATEMÀTIQUES AND IMTECH, DIAGONAL 647, 08028 BARCELONA, SPAIN & ³CENTRE DE RECERCA MATEMÀTICA, EDIFICI C, CAMPUS BELLATERRA, 08193 BELLATERRA, SPAIN

Email address: `xavier.cabre@upc.edu`