

# A note on graphs with purely imaginary per-spectrum\*

Hitesh Wankhede<sup>†</sup>      Ranveer Singh<sup>‡</sup>

November 24, 2022

## Abstract

In 1983, Borowiecki and Józwiak posed an open problem of characterizing graphs with purely imaginary per-spectrum. The most general result, although a partial solution, was given in 2004 by Yan and Zhang, who show that if a graph contains no subgraph which is an even subdivision of  $K_{2,3}$ , then it has purely imaginary per-spectrum. Zhang and Li in 2012 proved that such graphs are planar and admit a pfaffian orientation. In this article, we describe how to construct graphs with purely imaginary per-spectrum having a subgraph which is an even subdivision of  $K_{2,3}$  (planar and nonplanar) using coalescence of rooted graphs.

Key words: Permanental Polynomial, Bipartite Graphs, Theta Graphs, Coalescence.

AMS subject classification (2020): 05C05, 05C31, 05C50, 05C75, 05C76.

## 1 Introduction

We consider simple and undirected graphs. Let  $V(G)$  and  $E(G)$  denote the vertex set and the edge set of a graph  $G$ , respectively. If two vertices  $i, j \in V(G)$  are adjacent, then we write  $i \sim j$ . The *adjacency matrix*  $A(G) = (a_{ij})$  of a graph  $G$  with  $V(G) = \{1, \dots, n\}$  is the  $n \times n$  matrix in which  $a_{ij} = 1$  if  $i \sim j$ , and 0 otherwise. The *determinant* and the *permanent* of  $A(G)$ , are defined as

$$\det(A(G)) = \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) \prod_{i=1}^n a_{i, \sigma(i)},$$

$$\operatorname{per}(A(G)) = \sum_{\sigma \in S_n} \prod_{i=1}^n a_{i, \sigma(i)},$$

respectively, where  $S_n$  is the set of all permutation of the set  $\{1, 2, \dots, n\}$  and  $\operatorname{sgn}(\sigma)$  is the signature of the permutation  $\sigma$ . The *characteristic polynomial* and the *permanental polynomial* of graph  $G$  are defined as

$$\phi(G, x) = \det(xI - A(G)),$$

$$\pi(G, x) = \operatorname{per}(xI - A(G)),$$

respectively, where  $I$  is the identity matrix of order  $n$ . The set of all the roots (counting multiplicities) of  $\phi(G, x)$  and  $\pi(G, x)$  is called the *spectrum* and the *per-spectrum* of the graph  $G$ , respectively. Let  $\sigma(G)$  and  $\sigma_p(G)$  denote the spectrum and per-spectrum of a graph  $G$ , respectively. Two graphs are *cospectral* if they have the same spectrum and *per-cospectral* if they have the same per-spectrum. Both the spectrum and the per-spectrum are graph invariants. Because if two graphs are isomorphic, then they must necessarily have the same spectrum and the per-spectrum. The converse need not be true.

\*This work is supported by Department of Science and Technology (Govt. of India) through project DST/04/2019/002676.

<sup>†</sup>hitesh.wankhede@students.iiserpune.ac.in, hiteshwankhede9@gmail.com, Department of Computer Science and Engineering, Indian Institute of Technology Indore, India.

<sup>‡</sup>ranveer@iiti.ac.in, Department of Computer Science and Engineering, Indian Institute of Technology Indore, India.

A graph is said to be *determined by its spectrum (per-spectrum)*, if any other graph which is cospectral (per-cospectral) to it is also isomorphic. Haemers and van Dam have conjectured that ‘Almost all graphs are determined by their spectrum’ [12, 23]. Much work has been done on studying graphs using the characteristic polynomial and its roots in Spectral Graph Theory, but very less using the permanental polynomial in comparison [17]. The eigenvalues of a matrix are the roots of its characteristic polynomial, but the familiar Linear Algebra is not much helpful when studying the permanent as it is a combinatorial function. However, much work has been done on finding bounds on the permanent [3, 26]. Further, the determinant and the permanent look quite similar by definition, but computationally they are extremely different. The determinant can be computed in polynomial time using LUP decomposition, while computing permanent is a #P-complete problem [22]. The ‘Computation of Determinant vs. Permanent’ is a significant problem in Computational Complexity Theory [1, 2]. This computational dichotomy is also explored using the general notion of immanants in [8]. Some numerical evidence suggests that the permanental polynomial might be better than the characteristic polynomial in distinguishing graphs [9, 18]. This tradeoff between computational hardness and the graph characterizing power is an interesting area to study. For an excellent survey on permanental polynomial, we refer to [17]. For some recent work on permanental polynomial, we refer to [27, 13, 14, 15, 16, 24].

Let  $K_{m,n}$  denote the *complete bipartite graph* with partite sets of size  $m$  and  $n$ , respectively. In case  $m = 1$ , we get a *star graph* with  $n$  pendent vertices. Let  $K_n$ ,  $P_n$ , and  $C_n$  denote the *complete graph*, the *path graph*, and the *cycle graph* on  $n$  vertices, respectively. Note that the spectrum is always real, but the per-spectrum may not be real. H. Sachs (1978) had noted that if  $G$  is a tree such that  $\sigma(G) = \{\lambda_1, \lambda_2, \dots, \lambda_n\}$ , then  $\sigma_p(G) = \{i\lambda_1, i\lambda_2, \dots, i\lambda_n\}$ . Borowiecki (1985) [4] generalized this observation by showing that a bipartite graph  $G$  contains no  $C_k$  for any  $k \equiv 0 \pmod{4}$  if and only if  $\sigma(G) = \{\lambda_1, \lambda_2, \dots, \lambda_n\}$  and  $\sigma_p(G) = \{i\lambda_1, i\lambda_2, \dots, i\lambda_n\}$ . Thus, if  $G_1$  and  $G_2$  are two bipartite graphs without  $C_k$  for any  $k \equiv 0 \pmod{4}$ , then  $G_1$  and  $G_2$  are cospectral if and only if they are per-cospectral. In other words, the permanental polynomial is not any more useful than the characteristic polynomial in characterization of such graphs. Earlier, Borowiecki and Józwiak (1983) had posed the following open problem.

**Problem 1.1.** [6] *Characterize graphs with purely imaginary per-spectrum.*<sup>1</sup>

Let  $\mathcal{G}$  denote the class of graphs with purely imaginary per-spectrum. The *subdivision* of an edge  $(i, j)$  is the operation of replacing  $(i, j)$  with a path  $(i, k, j)$  through a new vertex  $k$ . Let  $G = G_0, G_1, G_2, \dots, G_k$  be a sequence of graphs such that for each  $i$ ,  $G_i$  can be obtained from  $G_{i-1}$  by subdividing an edge twice. Then,  $G_k$  is said to be an even subdivision of  $G$ . Yan and Zhang (2004) [25] have given the most general but a partial solution to this problem. They showed that if  $G$  is a bipartite graph containing no subgraph which is an even subdivision of  $K_{2,3}$ , then there exists an orientation  $G^e$  with skew-adjacency matrix  $A(G^e)$  such that  $\pi(G, x) = \det(xI - A(G^e))$ . The converse was proved by Zhang and Li (2012) [27] along with showing that such graphs are planar and the orientation is pfaffian. They also give a linear time algorithm to determine whether a given bipartite graph contains no even subdivision of  $K_{2,3}$ . Since the skew-adjacency matrix  $A(G^e)$  is skew-symmetric matrix, it has purely imaginary eigenvalues. It follows that the class of graphs containing no subgraph of an even subdivision of  $K_{2,3}$  is a subset of  $\mathcal{G}$ . Thus, we have

$$\text{Class of trees} \subseteq \text{Class of bipartite graphs without } C_k \text{ for any } k \equiv 0 \pmod{4} \subseteq \text{Class of graphs without a subgraph which is an even subdivision of } K_{2,3} \subseteq \mathcal{G}.$$

In this article, we wish to study the following problem.

**Problem 1.2.** *Let  $G$  be a bipartite graph containing a subgraph which is an even subdivision of  $K_{2,3}$ . When does it admit purely imaginary per-spectrum?*

The rest of paper is organised as follows. In Section 2, we recall some definitions, known results, and compute permanental polynomial of rooted trees. In Section 3, we describe a construction of graphs with purely imaginary per-spectrum and demonstrate it for some special cases.

---

<sup>1</sup>Zero is also considered to be a purely imaginary number. The per-spectrum is purely imaginary if each of the roots of the permanental polynomial is purely imaginary.

## 2 Preliminaries

The permanent polynomial of a graph  $G$  can also be computed using its Sachs subgraphs, that is, subgraphs whose components are either edges or cycles of length at least 3.

**Proposition 2.1.** [5] *Let  $G$  be a graph on  $n$  vertices with  $\pi(G, x) = \sum_{k=0}^n b_k x^{n-k}$ . Then,*

$$b_k = (-1)^k \sum_{U_k} 2^{c(U_k)},$$

where the summation is taken over all Sachs subgraphs  $U_k$  on  $k$  vertices of  $G$ , and  $c(U_k)$  denotes the number of components of  $U_k$  which are cycles.

As a consequence, we get the following.

**Proposition 2.2.** [5, 4, 7] *A graph  $G$  is bipartite if and only if  $b_k = 0$  for each odd  $k \in \{0, 1, 2, \dots, n\}$*

Next, we show that graphs with purely imaginary per-spectrum are bipartite.

**Proposition 2.3.** *If  $G \in \mathcal{G}$ , then  $G$  is bipartite.*

*Proof.* Let  $\sigma_p(G) = \{i\lambda_1, i\lambda_2, \dots, i\lambda_n\}$ , then  $\pi(G, x) = (x - i\lambda_1)(x - i\lambda_2) \dots (x - i\lambda_n)$ . If  $k$  is odd, then  $b_k$  has the factor  $i$ . But since  $\pi(G, x)$  has all real coefficients,  $b_k$  must be zero whenever  $k$  is odd. Proof follows from Proposition 2.2.  $\square$

Zhang, Liu and Li give a characterization of bipartite graphs in terms of the per-spectrum.

**Theorem 2.4.** [28] *Let  $G$  be a graph. Then,*

1.  $\pi(G, x)$  has no negative root.
2. If all the roots of  $\pi(G, x)$  are real, then  $G$  is empty.
3. If  $G$  is bipartite, then  $\pi(G, x)$  has no real root except 0.
4.  $G$  is bipartite if and only if the per-spectrum is symmetric with respect to the real and the imaginary axes.

As a consequence of Proposition 2.3 and Theorem 2.4(4), a graph  $G \in \mathcal{G}$  if and only if its permanent polynomial has the form

$$\pi(G, x) = x^{n-2k}(x^2 + c_1)(x^2 + c_2) \dots (x^2 + c_k),$$

where  $c_i > 0$  for each  $i$  for some  $k$ . Let  $G - S$  denote the induced subgraph on the vertex set  $V(G) - S$  for some  $S \subseteq V(G)$ . In case  $S = \{r\}$ , we denote it by  $G - r$ .

Next, we recall a recursive formula to compute  $\pi(G, x)$  of a graph  $G$ .

**Lemma 2.5.** [5] *Let  $G$  be a graph,  $u \in V(G)$ , and  $\Gamma_u(G)$  denote the set of cycles passing through  $u$ . Then,*

$$\pi(G, x) = x\pi(G - u, x) + \sum_{v \sim u} \pi(G - u - v, x) + 2 \sum_{C \in \Gamma_u(G)} (-1)^{|V(C)|} \pi(G - V(C), x),$$

where  $V(C)$  is the set of vertices in cycle  $C$ .

Let  $(T, r)$  denote a tree  $T$  rooted at the vertex  $r$ . Let  $u_1, u_2, \dots, u_k$  denote the children of the root  $r$ , and  $(T_1, u_1), (T_2, u_2), \dots, (T_k, u_k)$  denote the corresponding rooted subtrees for some  $k \geq 0$ . We write  $(T, r) = \mathcal{R}(T_1, T_2, \dots, T_k, r)$ . In case  $T_1 = T_2 = \dots = T_k$ , let  $T' := T_i$  and  $u' := u_i$  for each  $i = 1, \dots, k$ , then we write  $(T, r) = \mathcal{R}(T'^{(k)}, r)$ . Next, we give a recursive formula to compute permanent polynomial of rooted trees.

**Proposition 2.6.** Consider a rooted tree  $(T, r) = \mathcal{R}(T_1, T_2, \dots, T_k, r)$ . Then,

$$\pi(T, x) = x\pi(T - r, x) + \sum_{i=1}^k \left( \pi(T_i - u_i, x) \prod_{j=1, j \neq i}^k \pi(T_j, x) \right).$$

In case  $(T, r) = \mathcal{R}(T^{(k)}, r)$ , then the expression simplifies to

$$\pi(T, x) = x\pi(T - r, x) + k (\pi(T' - u', x)\pi(T', x)^{k-1}).$$

*Proof.* First, note that  $T - r = \cup_{i=1}^k T_i$ . Hence, for each  $i = 1, \dots, k$ , we have

$$\pi(T - r, x) = \prod_{i=1}^k \pi(T_i, x), \text{ and } \pi(T - r - u_i, x) = \pi(T_i - u_i, x) \prod_{j=1, j \neq i}^k \pi(T_j, x).$$

Using Lemma 2.5, we get

$$\pi(T, x) = x\pi(T - r, x) + \sum_{i=1}^k \pi(T - r - u_i, x).$$

Proof follows by substitutions, and the second statement follows as a corollary.  $\square$

The permenental polynomial of star graphs and path graphs can be computed directly using the following proposition.

**Proposition 2.7.** 1.  $\pi(K_{1,n}, x) = x^{n-1}(x^2 + n)$

2.  $\pi(P_n, x) = x\pi(P_{n-1}) + \pi(P_{n-2})$  where  $\pi(P_0) = 1$ ,  $\pi(P_1) = x$ . Equivalently,

$$\pi(P_n, x) = \sum_{m=0}^{\lfloor n/2 \rfloor} \binom{n-m}{n-2m} x^{n-2m}.$$

*Proof.* For the star graph, substitute  $k = n$  and  $T_i = K_1$ , and for the path graph, substitute  $k = 1$  and  $T_i = P_{n-1}$  in Proposition 2.6. The recurrence relation of the permenental polynomial of the path graph can be solved either using generating series or using Proposition 2.1.  $\square$

We say that a rooted tree  $(T, r)$  is *starlike* if  $(T, r) = \mathcal{R}(K_{1,l}^{(k)}, r)$ , and *pathlike* if  $(T, r) = \mathcal{R}(P_{l+1}^{(k)}, r)$  for some  $l \geq 0$ . See Figure 1a, 1b. Note that for a rooted starlike tree  $T$ , the subtrees  $(K_{1,l}, u')$  are rooted at the central vertex of degree  $l$ . So consequently  $\pi(T', x) = x^{l-1}(x^2 + l)$ , and  $\pi(T' - u') = \pi(lK_1, x) = x^l$ , where  $lK_1$  denotes  $l$  copies of  $K_1$ . For a rooted pathlike tree  $T$ , we have  $\pi(T', x) = \pi(P_{l+1})$ , and  $\pi(T' - u') = \pi(P_l)$ .




Figure 1

Let  $u, v$  be two vertices, and  $a, b, c \geq 1$ . A *theta graph*  $\Theta_{a,b,c}(u, v)$  is a graph on  $a + b + c + 2$  vertices such that there are three disjoint paths from  $u$  to  $v$  of lengths  $a + 1$ ,  $b + 1$  and  $c + 1$ . See Figure 1c. Any




Figure 2:  $\mathcal{B}$ : Bipartite Graphs,  $\mathcal{P}$ : Planar Bipartite Graphs,  $\mathcal{G}$ : Graphs with purely imaginary per-spectrum,  $\mathcal{K}$ : Graphs without a subgraph which is an even subdivision of  $K_{2,3}$ .

theta graph is a homeomorphic image of  $K_{2,3}$  as it can be obtained by repeated subdivisions of  $K_{2,3}$ <sup>2</sup>. For example, we have  $\Theta_{1,1,1} = K_{2,3}$ . An even subdivision of  $K_{2,3}$  is a theta graph with all odd parameters. Note that after removing a vertex of degree 3 from a theta graph, we get a pathlike tree. Hence, we can obtain a formula for  $\pi(\Theta_{a,b,c}, x)$  in terms of the permanental polynomial of path graphs using Lemma 2.5.

Consider two rooted graphs  $(G_1, r_1)$  and  $(G_2, r_2)$ . By identifying  $r_1$  in  $G_1$  with  $r_2$  in  $G_2$ , we construct a new graph  $G_1 \cdot G_2$  called the *coalescence* of  $G_1$  and  $G_2$ . Schwenk (1974) proved the following relation which holds for both the characteristic and the permanental polynomial.

**Lemma 2.8.** [19] *Let  $(G_1, r_1)$  and  $(G_2, r_2)$  be two rooted graphs, then*

$$\pi(G_1 \cdot G_2, x) = \pi(G_1, x)\pi(G_2 - r_2, x) + \pi(G_1 - r_1, x)\pi(G_2, x) - x\pi(G_1 - r_1, x)\pi(G_2 - r_2, x).$$

### 3 Construction

In this section, we describe a construction of graphs with purely imaginary per-spectrum and demonstrate it for some special cases. First, we discuss some examples.

The graphs containing no subgraph which is an even subdivision of  $K_{2,3}$  are planar, but not all planar graphs are in  $\mathcal{G}$ . The smallest example of a planar graph not in  $\mathcal{G}$  is  $K_{2,3}$  itself. The smallest planar graph in  $\mathcal{G}$  containing an even subdivision of  $K_{2,3}$  is a graph on 8 vertices (denote this by  $G_8$ ) obtained by subdividing any edge of  $K_{2,4}$  twice. The smallest nonplanar graph in  $\mathcal{G}$  containing an even subdivision of  $K_{2,3}$  is a graph on 11 vertices (denote this by  $G_{11}$ ) obtained as follows. Consider  $K_{3,3}$  and  $v \in V(K_{3,3})$ . Take two edges adjacent to  $v$  and subdivide each of them twice. Now add a new vertex  $u$  and draw an edge between  $u$  and  $v$ . The graph  $G_{11}$  serves as a motivation for our construction using coalescence. The smallest nonplanar graph, which is not in  $\mathcal{G}$  is  $K_{3,3}$ . See Figure 2 and 3 for these graphs. Their permanental polynomial and per-spectrum are as follows.

- $\pi(K_{2,3}, x) = x(x^4 + 6x^2 + 12)$ ,  
 $\sigma_p(K_{2,3}) = \{0, z, \bar{z}, iz, i\bar{z}\}$  where  $z = 0.79 + 2.10i$  (approximated to two decimals),
- $\pi(G_8, x) = x^2(x^6 + 10x^4 + 33x^2 + 36)$ ,  
 $\sigma_p(G_8) = \{0, 0, \pm 2i, \pm\sqrt{3}i, \pm\sqrt{3}i\}$ ,
- $\pi(K_{3,3}, x) = x^6 + 9x^4 + 36x^2 + 36$ ,  
 $\sigma_p(K_{3,3}) = \{-1.96i, 1.96i, y, \bar{y}, iy, i\bar{y}\}$  where  $y = 0.76 + 2.10i$  (approximated to two decimals),
- $\pi(G_{11}, x) = x(x^{10} + 14x^8 + 72x^6 + 168x^4 + 172x^2 + 56)$ ,  
 $\sigma_p(G_{11}) = \{0, \pm i\sqrt{4 + \sqrt{2}}, \pm i\sqrt{2 + \sqrt{2}}, \pm i\sqrt{4 - \sqrt{2}}, \pm i\sqrt{2}, \pm i\sqrt{2 - \sqrt{2}}\}$ .

Next, we consider the coalescence of a rooted graph and a rooted tree.

<sup>2</sup>A recursive formula to compute the characteristic polynomial of a theta graph in one of the parameters was given by Sciriha and Fiorini [20].




Figure 3: (Left to right)  $K_{2,4}$ ,  $G_8$ ,  $K_{3,3}$  and  $G_{11}$ . By subdividing the blue edge twice in  $K_{2,4}$  we obtain  $G_8$ . By subdividing the blue and the red edge twice in  $K_{3,3}$ , and drawing an edge between their common vertex and the newly added vertex, we obtain  $G_{11}$ .

**Theorem 3.1.** *Let  $(G_1, r_1)$  be a rooted graph and  $(T, r_2) = \mathcal{R}(T^{(k)}, r_2)$  be a rooted tree. Their coalescence  $G = G_1 \cdot T$  has purely imaginary per-spectrum if and only if the polynomial*

$$\mathcal{H}(G_1, T) := \pi(G_1, x)\pi(T', x) + k\pi(G_1 - r_1, x)\pi(T' - u', x)$$

*has purely imaginary roots.*

*Proof.* Applying Lemma 2.8, we get

$$\begin{aligned} \pi(G, x) &= \pi(G_1, x)\pi(T - r_2, x) + \pi(G_1 - r_1, x)\pi(T, x) - x\pi(G_1 - r_1, x)\pi(T - r_2, x) \\ &= \pi(G_1, x)\pi(T - r_2, x) + \pi(G_1 - r_1, x)(\pi(T, x) - x\pi(T - r_2, x)) \\ &= \pi(G_1, x)\pi(T', x)^k + \pi(G_1 - r_1, x)(k\pi(T' - u', x)\pi(T', x)^{k-1}) \\ &= \pi(T', x)^{k-1}(\pi(G_1, x)\pi(T', x) + k\pi(G_1 - r_1, x)\pi(T' - u', x)). \end{aligned}$$

In the third step, we use Proposition 2.6. We already know that a tree has purely imaginary per-spectrum. Hence, we can ignore the common factor  $\pi(T', x)^{k-1}$ .  $\square$

In the remaining section, we take different examples of graph  $G_1$ , in particular,  $K_{2,3}$  rooted at a degree 3 vertex,  $K_{2,3}$  rooted at a degree 2 vertex, and a rooted  $K_{3,3}$ . As for the tree, we take  $T$  to be a rooted starlike or a rooted pathlike tree.

Consider the case when  $G_1 = K_{2,3}$ , and  $T$  is starlike.

**Theorem 3.2.** *The coalescence of  $(K_{2,3}, r_1)$  such that  $\text{degree}(r_1) = 3$  and rooted starlike tree  $\mathcal{R}(K_{1,l}^{(k)}, r_2)$  has purely imaginary per-spectrum if and only if any of the following holds:*

1.  $l + k \geq 4$  when  $l \leq 3$ ,
2.  $k \geq 2l - 4$  when  $l \geq 4$ .

*Proof.* We have  $\pi(K_{2,3}, x) = x(x^4 + 6x^2 + 12)$  and  $\pi(K_{1,3}, x) = x^2(x^2 + 3)$ . Now consider the polynomial

$$\begin{aligned} \mathcal{H}(G_1, T) &= \pi(G_1, x)\pi(T', x) + k\pi(G_1 - r_1, x)\pi(T' - u', x) \\ &= \pi(K_{2,3}, x)\pi(K_{1,l}, x) + k\pi(K_{1,3}, x)\pi(lK_1, x) \\ &= x(x^4 + 6x^2 + 12)x^{l-1}(x^2 + l) + kx^2(x^2 + 3)x^l \\ &= x^l(x^6 + (l + k + 6)x^4 + (6l + 3k + 12)x^2 + 12l). \end{aligned}$$

This polynomial has purely imaginary roots if and only if  $l + k \geq 4$  when  $l \leq 3$  and  $k \geq 2l - 4$  when  $l \geq 4$ .<sup>3</sup> Proof follows by Theorem 3.1.  $\square$

**Example 1.** Let  $k = n$  and  $l = 0$ . Then we get a star  $T = K_{1,n}$  is rooted at the central vertex. Let  $G_n$  be the coalescence of  $K_{2,3}$  (rooted at a vertex of degree 3) and  $K_{1,n}$  (rooted at a vertex of degree  $n$ ), then it has purely imaginary per-spectrum if and only if  $n \geq 4$ .

<sup>3</sup>This is verified using SageMath software [21].

**Example 2.** Let  $k = 1$  and  $l = n - 1$ , then we get a star  $T = K_{1,n}$  rooted a pendant vertex. Let  $G_n$  be the coalescence of  $K_{2,3}$  (rooted at a vertex of degree 3) and  $K_{1,n}$  (rooted at a vertex of degree 1), then it has purely imaginary roots if and only if  $n = 4$ .

**Remark 3.3.** In case when the rooted tree  $T = \mathcal{R}(P_{l+1}^{(k)}, r_1)$  is pathlike, substituting  $\pi(T', x) = \pi(P_{l+1}, x)$  and  $\pi(T' - u') = \pi(P_l, x)$ , we get

$$\mathcal{H}(G_1, T) = x((x^4 + 6x^2 + 12)\pi(P_{l+1}, x) + kx(x^2 + 3)\pi(P_l, x)).$$

There is no closed form for the set of all pairs  $(k, l)$  such that the polynomial has purely imaginary roots. But the first few  $l$ 's and the corresponding  $k$ 's for which this happens are given as follows.

1.  $k \geq 4$  when  $l = 0$ ,
2.  $k \geq 3$  when  $l = 1, 2, 3, 4$ ,
3.  $k \geq 13$  when  $l = 7$ .

Next, we make a degree 2 vertex of  $K_{2,3}$  a root.

**Theorem 3.4.** *The coalescence of  $(K_{2,3}, r_1)$  such that  $\text{degree}(r_1) = 2$  with a rooted starlike tree  $\mathcal{R}(K_{1,l}^{(k)}, r_2)$  has purely imaginary per-spectrum if and only if  $l + k \geq 4$  and  $l \leq 2$ .*

*Proof.* We have  $\pi(K_{2,3}, x) = x(x^4 + 6x^2 + 12)$  and  $\pi(C_4, x) = (x^2 + 2)^2$ . Now consider the polynomial

$$\begin{aligned} \mathcal{H}(G_1, T) &= \pi(G_1, x)\pi(T', x) + k\pi(G_1 - r_1, x)\pi(T' - u', x) \\ &= \pi(K_{2,3}, x)\pi(K_{1,l}, x) + k\pi(C_4, x)\pi(lK_1, x) \\ &= x(x^4 + 6x^2 + 12)x^{l-1}(x^2 + l) + k(x^2 + 2)^2x^l \\ &= x^l(x^6 + (l + k + 6)x^4 + (6l + 4k + 12)x^2 + (12l + 4k)). \end{aligned}$$

This polynomial has purely imaginary roots if and only if  $l + k \geq 4$  and  $l \leq 2$ . Proof follows by Theorem 3.1.  $\square$

**Example 3.** Let  $k = n$  and  $l = 0$ , then we get a star  $K_{1,n}$  rooted at the central vertex. Let  $G_n$  be the coalescence of  $K_{2,3}$  (rooted at a vertex of degree 2) and  $K_{1,n}$  (rooted at a vertex of degree  $n$ ). It has purely imaginary per-spectrum if and only if  $n \geq 4$ .

**Remark 3.5.** Similarly, in case when the rooted tree  $T = \mathcal{R}(P_{l+1}^{(k)}, r_1)$  is pathlike, we get

$$\mathcal{H}(G_1, T) = x(x^4 + 6x^2 + 12)\pi(P_{l+1}, x) + k(x^2 + 2)^2\pi(P_l, x).$$

There is no closed form for the set of all pairs  $(k, l)$  such that the polynomial has purely imaginary roots. But the first few  $l$ 's and the corresponding  $k$ 's for which this happens are given as follows.

1.  $k \geq 4$  when  $l = 0$ ,
2.  $k \geq 3$  when  $l = 1, 2$ ,
3.  $k = 3$  when  $l = 3$ ,
4.  $k \geq 19$  when  $l = 5$ ,
5.  $k \geq 9$  when  $l = 6$ .

**Remark 3.6.** The smallest connected bipartite graph in  $\mathcal{G}$  which has an even subdivision of  $K_{2,3}$  is the graph  $G_8$  (see Figure 3), and there are no other such graphs on 8 vertices in  $\mathcal{G}$ . On 9 vertices, there are 8 graphs in  $\mathcal{G}$  which contain an even subdivision of  $K_{2,3}$  (see Figure 4). Note that 7 of them can be obtained using the single coalescence of either  $K_{2,3}$  or  $\Theta_{3,1,1}$  with a starlike or a pathlike tree. The remaining example requires coalescences at two different roots.

The graph $G_1$	Coalescence of $G_1$ with rooted tree
$K_{2,3}$ rooted at a degree 3 vertex	
$K_{2,3}$ rooted at a degree 2 vertex	
$\Theta_{3,1,1}$ rooted at a degree 3 vertex	
$\Theta_{3,1,1}$ rooted at a degree 2 vertex	
$\Theta_{3,1,1}$ with two root vertices of degree 3 each	

Figure 4: Connected bipartite graphs on 9 vertices in  $\mathcal{G}$  which have an even subdivision of  $K_{2,3}$  as a subgraph

Next, we consider the case when  $G_1$  is  $K_{3,3}$  to produce nonplanar graphs with purely imaginary per-spectrum.

**Theorem 3.7.** *The coalescence of  $K_{3,3}$  rooted at any vertex with a rooted starlike tree  $T = \mathcal{R}(K_{1,l}^{(k)}, r_2)$  has purely imaginary per-spectrum if and only if  $(k, l) \in \{(5, 1), (6, 0), (7, 0)\}$ . The coalescence of  $K_{3,3}$  rooted at any vertex with a rooted pathlike tree  $T = \mathcal{R}(P_{l+1}^{(k)}, r_2)$  has purely imaginary per-spectrum if and only if  $(k, l) \in \{(5, 1), (6, 0), (7, 0)\}$ .*

*Proof.* First recall the useful facts  $\pi(K_{3,3}, x) = x^6 + 9x^4 + 36x^2 + 36$  and  $\pi(K_{2,3}, x) = x(x^4 + 6x^2 + 12)$ . When the tree is starlike, we have

$$\begin{aligned}
\mathcal{H}(G_1, T) &= \pi(G_1, x)\pi(T', x) + k\pi(G_1 - r_1, x)\pi(T' - u', x) \\
&= \pi(K_{3,3}, x)\pi(K_{1,l}, x) + k\pi(K_{2,3}, x)\pi(lK_1, x) \\
&= (x^6 + 9x^4 + 36x^2 + 36)x^{l-1}(x^2 + l) + kx(x^4 + 6x^2 + 12)x^l \\
&= x^{l-1}(x^8 + (k+l+9)x^6 + (6k+9l+36)x^4 + (12k+36l+36)x^2 + 36l).
\end{aligned}$$

This polynomial has purely imaginary roots if and only if  $(k, l) \in \{(5, 1), (6, 0), (7, 0)\}$ . When the tree is pathlike, we have

$$\begin{aligned}
\mathcal{H}(G_1, T) &= \pi(G_1, x)\pi(T', x) + k\pi(G_1 - r_1, x)\pi(T' - u', x) \\
&= \pi(K_{3,3}, x)\pi(P_{l+1}, x) + k\pi(K_{2,3}, x)\pi(P_l, x) \\
&= (x^6 + 9x^4 + 36x^2 + 36)\pi(P_{l+1}, x) + kx(x^4 + 6x^2 + 12)\pi(P_l, x).
\end{aligned}$$

This polynomial has purely imaginary roots if and only if  $(k, l) \in \{(5, 1), (6, 0), (7, 0)\}$ . Proof follows by Theorem 3.1.  $\square$

## 4 Conclusion

There are various ways in which one can make use of the construction idea. The graph  $G_1$  in Theorem 3.1 can be taken to be any theta graph with odd parameters or any graph containing it. One can also consider multiple coalescences (see [10]) or a different definition of the rooted product (see neighbourhood rooted product [11]). The second graph  $T$  need not be only a starlike or a pathlike tree. For example, the coalescence of the cycle  $C_6$  and  $K_{2,3}$  (rooted at a degree 3 vertex) has purely imaginary per-spectrum, but it is not obtainable using the construction described in this article. Accounting for such cases is the subject of future study.



## References

- [1] Manindra Agrawal. Determinant versus permanent. In *Proceedings of the 25th International Congress of Mathematicians, ICM 2006*, volume 3, pages 985–997, 2006.
- [2] Frank Ban. Permanent vs. determinant. 2014.
- [3] RB Bapat. Recent developments and open problems in the theory of permanents. *The Mathematics student*, 76(1):55, 2007.
- [4] Mieczysław Borowiecki. On spectrum and per-spectrum of graphs. *Publ. Inst. Math.(Beograd)*, 38:31–33, 1985.
- [5] Mieczysław Borowiecki and T Józwiak. Computing the permanental polynomial of a multigraph. *Discussiones Mathematicae*, 5:9–16, 1982.
- [6] Mieczysław Borowiecki and Tadeusz Józwiak. A note on characteristic and permanental polynomials of multigraphs. In *Graph theory*, pages 75–78. Springer, 1983.
- [7] Andries E Brouwer and Willem H Haemers. *Spectra of graphs*. Springer Science & Business Media, 2011.
- [8] Radu Curticapean. A full complexity dichotomy for immanant families. In *Proceedings of the 53rd Annual ACM SIGACT Symposium on Theory of Computing*, pages 1770–1783, 2021.
- [9] Matthias Dehmer, Frank Emmert-Streib, Bo Hu, Yongtang Shi, Monica Stefu, and Shailesh Tripathi. Highly unique network descriptors based on the roots of the permanental polynomial. *Information Sciences*, 408:176–181, 2017.
- [10] CD Godsil and BD McKay. A new graph product and its spectrum. *Bulletin of the Australian Mathematical Society*, 18(1):21–28, 1978.
- [11] Chris D Godsil and Brendan D McKay. Constructing cospectral graphs. *Aequationes Mathematicae*, 25(1):257–268, 1982.
- [12] Willem H Haemers. Are almost all graphs determined by their spectrum. *Notices of the South African Mathematical Society*, 47:42–45, 2016.
- [13] Wei Li. Graphs whose characteristic and permanental polynomials have coefficients of the same magnitude. *Discrete Mathematics*, 339(8):2127–2135, 2016.
- [14] Wei Li. On the skew-permanental polynomials of orientation graphs. *Discrete Applied Mathematics*, 208:79–87, 2016.
- [15] Wei Li. The graphs whose permanental polynomials are symmetric. *Discussiones Mathematicae: Graph Theory*, 38(1), 2018.
- [16] Wei Li. On the matching and permanental polynomials of graphs. *Discrete Applied Mathematics*, 302:16–23, 2021.
- [17] Wei Li, Shunyi Liu, Tingzeng Wu, and Heping Zhang. On the permanental polynomials of graphs. *Graph Polynomials*, pages 101–121, 2016.
- [18] Shunyi Liu and Jinjun Ren. Enumeration of copermanental graphs. *arXiv preprint arXiv:1411.0184*, 2014.
- [19] Allen J Schwenk. Computing the characteristic polynomial of a graph. In *Graphs and combinatorics*, pages 153–172. Springer, 1974.
- [20] Irene Sciriha and Stanley Fiorini. On the characteristic polynomial of homeomorphic images of a graph. *Discrete Mathematics*, 174(1-3):293–308, 1997.

- [21] The Sage Developers. *SageMath, the Sage Mathematics Software System (Version 9.7)*, 2022. <https://www.sagemath.org>.
- [22] Leslie G Valiant. The complexity of computing the permanent. *Theoretical computer science*, 8(2):189–201, 1979.
- [23] Edwin R Van Dam and Willem H Haemers. Which graphs are determined by their spectrum? *Linear Algebra and its applications*, 373:241–272, 2003.
- [24] Tingzeng Wu and Hong-Jian Lai. Constructing graphs which are permanental cospectral and adjacency cospectral. *Graphs and Combinatorics*, 34(6):1713–1721, 2018.
- [25] Weigen Yan and Fuji Zhang. On the permanental polynomials of some graphs. *Journal of mathematical chemistry*, 35(3):175–188, 2004.
- [26] Fuzhen Zhang. An update on a few permanent conjectures. *Special Matrices*, 4(1), 2016.
- [27] Heping Zhang and Wei Li. Computing the permanental polynomials of bipartite graphs by pfaffian orientation. *Discrete Applied Mathematics*, 160(13-14):2069–2074, 2012.
- [28] Heping Zhang, Shunyi Liu, and Wei Li. A note on the permanental roots of bipartite graphs. *Discussiones Mathematicae: Graph Theory*, 34(1), 2014.