

# Fuzzy hyperspheres via confining potentials and energy cutoffs

Gaetano Fiore

Dip. di Matematica e applicazioni, Università di Napoli “Federico II”,  
& INFN, Sezione di Napoli,  
Complesso Universitario M. S. Angelo, Via Cintia, 80126 Napoli, Italy

## Abstract

We simplify and complete the construction of fully  $O(D)$ -equivariant fuzzy spheres  $S_\Lambda^d$ , for all dimensions  $d \equiv D - 1$ , initiated in [G. Fiore, F. Pisacane, J. Geom. Phys. 132 (2018), 423-451]. This is based on imposing a suitable energy cutoff on a quantum particle in  $\mathbb{R}^D$  subject to a confining potential well  $V(r)$  with a very sharp minimum on the sphere of radius  $r = 1$ ; the cutoff and the depth of the well diverge with  $\Lambda \in \mathbb{N}$ . We show that the irreducible representation  $\pi_\Lambda$  of  $O(D+1)$  on the space of harmonic homogeneous polynomials of degree  $\Lambda$  in the Cartesian coordinates of  $\mathbb{R}^{D+1}$  (which we express via trace-free completely symmetric projections) is isomorphic to the Hilbert space  $\mathcal{H}_\Lambda$  of the particle, as a reducible representation of  $O(D) \subset O(D+1)$ . Moreover, we show that the algebra of observables is isomorphic to  $\pi_\Lambda(Uso(D+1))$ . As  $\Lambda$  diverges (commutative limit) so does the dimension of  $\mathcal{H}_\Lambda$ , and we recover ordinary quantum mechanics on the sphere  $S^d$ ; more formally, we have a fuzzy quantization of a coadjoint orbit of  $O(D+1)$  that goes to the classical phase space  $T^*S^d$ .

## 1 Introduction

Noncommutative space(time) algebras are introduced and studied with various motivations, notably to provide an arena for regularizing ultraviolet (UV) divergences in quantum field theory (QFT) (see e.g. [1]), reconciling Quantum Mechanics and General Relativity in a satisfactory Quantum Gravity (QG) theory (see e.g. [2]), unifying fundamental interactions (see e.g. [3, 4]). Noncommutative Geometry (NCG) [5, 8, 6, 7] has become a sophisticated framework that develops the whole machinery of differential geometry on noncommutative spaces. Fuzzy spaces are particularly appealing noncommutative spaces: a fuzzy space is a sequence  $\{\mathcal{A}\}_{n \in \mathbb{N}}$  of *finite-dimensional* algebras such that  $\mathcal{A}_n \xrightarrow{n \rightarrow \infty} \mathcal{A} \equiv$  algebra of regular functions on an ordinary manifold, with  $\dim(\mathcal{A}_n) \xrightarrow{n \rightarrow \infty} \infty$ . They have raised a big interest in the high energy physics community as a non-perturbative technique in quantum field theory (QFT) based on a finite discretization of space(time) alternative to the lattice one: the main advantage is that the algebras  $\mathcal{A}_n$  can carry representations of Lie groups (not only

of discrete ones). They can be used also for internal (e.g. gauge) degrees of freedom (see e.g. [9]), or as a new tool in string and  $D$ -brane theories (see e.g. [10, 11]). The first and seminal fuzzy space is the 2-dimensional Fuzzy Sphere (FS) of Madore and Hoppe [12, 13], where  $\mathcal{A}_n \simeq M_n(\mathbb{C})$ , which is generated by coordinates  $x^i$  ( $i = 1, 2, 3$ ) fulfilling

$$[x^i, x^j] = \frac{2i}{\sqrt{n^2-1}} \varepsilon^{ijk} x^k, \quad r^2 := x^i x^i = 1, \quad n \in \mathbb{N} \setminus \{1\} \quad (1)$$

(sum over repeated indices is understood); they are obtained by the rescaling  $x^i = 2L_i/\sqrt{n^2-1}$  of the elements  $L_i$  of the standard basis of  $so(3)$  in the unitary irreducible representation (irrep)  $(\pi^l, V^l)$  of dimension  $n = 2l+1$ , i.e. where  $V^l$  is the eigenspace of the Casimir  $\mathbf{L}^2 = L_i L_i$  with eigenvalue  $l(l+1)$ . Ref. [14, 15] first proposed a QFT based on it. Each matrix in  $M_n$  can be expressed as a polynomial in the  $x^i$  that can be rearranged as the expansion in spherical harmonics of an element of  $C(S^2)$  truncated at level  $n$ . Unfortunately, such a nice feature is not shared by the fuzzy spheres of dimension  $d > 2$ : the product of two spherical harmonics is not a combination of spherical harmonics, but an element in a larger algebra  $\mathcal{A}_n$ . Fuzzy spheres of dimension  $d = 4$ , any  $d \geq 3$  were first introduced respectively in [16] and [17]; other versions in  $d = 3, 4$  or  $d \geq 3$  have been proposed in [18, 19, 20, 21].

The Hilbert space of a (zero-spin) quantum particle on  $S^d$  carries a *reducible* representation of  $O(D)$ ; this decomposes into irreducible representations (irreps) as follows

$$\mathcal{L}^2(S^d) = \bigoplus_{l=0}^{\infty} V_D^l, \quad (2)$$

where the carrier space  $V_D^l$  is an eigenspace of the quadratic Casimir  $\mathbf{L}^2$  with eigenvalue

$$E_l := l(l+D-2). \quad (3)$$

For  $D = 3$  and  $l \in \mathbb{N}_0$   $V_3^l \equiv V^l$ . On the contrary, the mentioned fuzzy hyperspheres (including the Madore-Hoppe FS) are either based on sequences of irreps of  $Spin(D)$  (so that  $r^2$ , which is proportional to  $\mathbf{L}^2$ , is identically 1) parametrized by  $n$  [12, 13, 16, 17, 18, 19], or on sequences of reducible representations that are the direct sums of small bunches of such irreps [20, 21]. In either case, even excluding the  $n$  for which the associated representation of  $O(D)$  is only *projective*, the carrier space does not go to (2) in the limit  $n \rightarrow \infty$ ; we think this makes the interpretation of these fuzzy spheres as fuzzy configuration spaces (and of the  $x^i$  as spatial coordinates) questionable. For the Madore-Hoppe FS such an interpretation is even more difficult, because relations (1) are equivariant under  $SO(3)$ , but not under the whole  $O(3)$ , e.g. not under parity  $x^i \mapsto -x^i$ , while the ordinary sphere  $S^2$  is; on the contrary, all the other mentioned fuzzy spheres are  $O(D)$ -equivariant, because the commutators  $[x^i, x^j]$  are Snyder-like [1], i.e. proportional to angular momentum components  $L_{ij}$ .

The purpose of this work is to complete the construction [22, 23] of new, fully  $O(D)$ -equivariant fuzzy approximations of spheres  $S^d$  of arbitrary dimension  $d = D - 1 \in \mathbb{N}$ , thought as configuration spaces, simplifying and uniformizing (with respect to  $D$ ) the construction procedure; in the commutative limit the involved  $O(D)$ -representation goes to (2).

We recall the procedure starting from the general underlying philosophy [24, 22, 25]. Consider a quantum theory  $\mathcal{T}$  with Hilbert space  $\mathcal{H}$ , algebra of observables on  $\mathcal{H}$  (or with a

domain dense in  $\mathcal{H}$ )  $\mathcal{A} \equiv \text{Lin}(\mathcal{H})$ , Hamiltonian  $H \in \mathcal{A}$ . For any subspace  $\overline{\mathcal{H}} \subset \mathcal{H}$  preserved by the action of  $H$ , let  $\overline{P} : \mathcal{H} \mapsto \overline{\mathcal{H}}$  be the associated projection and

$$\overline{\mathcal{A}} \equiv \text{Lin}(\overline{\mathcal{H}}) = \{\overline{A} \equiv \overline{P}A\overline{P} \mid A \in \mathcal{A}\};$$

the observable  $\overline{A} \equiv \overline{P}A\overline{P} \in \overline{\mathcal{A}}$  will have the same physical interpretation as  $A$ . By construction  $\overline{H} = \overline{P}H = H\overline{P}$ . The projected Hilbert space  $\overline{\mathcal{H}}$ , algebra of observables  $\overline{\mathcal{A}}$  and Hamiltonian  $\overline{H}$  provide a new quantum theory  $\overline{\mathcal{T}}$ . If  $\overline{\mathcal{H}}, H$  are invariant under some group  $G$ , then  $\overline{P}, \overline{\mathcal{A}}, \overline{H}, \overline{\mathcal{T}}$  will be as well. In general, the relations among the generators of  $\overline{\mathcal{A}}$  differ from those among the generators of  $\mathcal{A}$ . In particular, if the theory  $\mathcal{T}$  is based on commuting coordinates  $x^i$  (commutative space) this will be in general no longer true for  $\overline{\mathcal{T}}$ :  $[\overline{x}^i, \overline{x}^j] \neq 0$ , and we have generated a quantum theory on a NC space.

A physically relevant instance of the above projection mechanism occurs when  $\overline{\mathcal{H}}$  is the subspace of  $\mathcal{H}$  characterized by energies  $E$  below a certain cutoff,  $E \leq \overline{E}$ ; then  $\overline{\mathcal{T}}$  is a *low-energy effective approximation* of  $\mathcal{T}$ . What it can be useful for? If  $\overline{\mathcal{A}}$  contains all the observables corresponding to measurements that we can *really* perform with the experimental apparati at our disposal, and the initial state of the system belongs to  $\overline{\mathcal{H}}$ , then neither the dynamical evolution ruled by  $H$ , nor any measurement can map it out of  $\overline{\mathcal{H}}$ , and we can replace  $\mathcal{T}$  by the effective theory  $\overline{\mathcal{T}}$ . Moreover, if at  $E > \overline{E}$  we even expect new physics not accountable by  $\mathcal{T}$ , then  $\overline{\mathcal{T}}$  may also help to figure out a new theory  $\mathcal{T}'$  valid for all  $E$ .

For an ordinary (for simplicity, zero-spin) quantum particle in the Euclidean (configuration) space  $\mathbb{R}^D$  it is  $\mathcal{H} = \mathcal{L}^2(\mathbb{R}^d)$ . Fixed a Hamiltonian  $H(x, p)$ , by standard wisdom the dimension of  $\overline{\mathcal{H}}$  is

$$\dim(\overline{\mathcal{H}}) \approx \text{Vol}(\mathcal{B}_{\overline{E}})/h^D,$$

where  $h$  is the Planck constant and  $\mathcal{B}_{\overline{E}} \equiv \{(x, p) \in \mathbb{R}^{2D} \mid H(x, p) \leq \overline{E}\}$  is the classical phase space region with energy below  $\overline{E}$ . If  $H$  consists only of the kinetic energy  $T$ , then this is infinite (fig. 1 left). If  $H = T + V$ , with a confining potential  $V$ , then this is finite (fig. 1 right) at least for sufficiently small  $\overline{E}$ , and also the classical region  $v_{\overline{E}} \subset \mathbb{R}^D$  in configuration space determined by the condition  $V \leq \overline{E}$  is bounded. In the sequel we rescale  $x, p, H, V$  so that they are dimensionless and, denoting by  $\Delta$  the Laplacian in  $\mathbb{R}^D$ , we can write

$$H = -\Delta + V. \tag{4}$$

The ‘dimensional reduction’  $\mathbb{R}^D \rightsquigarrow S^d$  of the configuration space is obtained:

1. Assuming in addition that  $V = V(r)$  depends only on the distance  $r$  from the center of the sphere  $S^d \subset \mathbb{R}^D$  and has a very sharp minimum, parametrized by a very large  $k \equiv V''(1)/4$ , on the sphere  $S^d$  of equation  $r = 1$ , see fig. 2.
2. Choosing  $\overline{E}$  so low that all radial quantum excitations are ‘frozen’, i.e. excluded from  $\overline{\mathcal{H}} \subset \mathcal{H}$ ; this makes  $H$  coincide with the Laplacian  $\mathbf{L}^2$  on  $S^d$ , up to terms  $O(1/\sqrt{k})$ .
3. Making both  $k, \overline{E}$  depend on, grow and diverge with a natural number  $\Lambda$ . Thereby we rename  $\overline{\mathcal{H}}, \overline{P}, \overline{\mathcal{A}}$  as  $\mathcal{H}_\Lambda, P_\Lambda, \mathcal{A}_\Lambda$ .




Figure 1: Classical phase space regions with energy  $E \leq \bar{E}$

As  $H$  is  $O(D)$ -invariant, so are  $P_\Lambda$ ,  $\bar{H} = P_\Lambda H$ , and the projected theory is  $O(D)$ -equivariant.

Technical details are given in sections 2, 3. Section 2 fixes the notation and contains preliminaries partly developed in [23]. The representation-theoretical results of section 3, which deserve attention also on their own, allow to explicitly characterize the space  $V_D^l$  as the space of harmonic homogeneous polynomials of degree  $l$  in the Cartesian coordinates  $x^i$  of  $\mathbb{R}^D$  restricted to the sphere  $S^d$ ; we determine such polynomials constructing the trace-free completely symmetric projection of  $(\mathbb{R}^D)^{\otimes l}$  and applying it to the polynomials of degree  $l$  in  $x^i$ . The actions of the  $L_{hk}$  and of the multiplication operators  $x^i \cdot$  on such polynomials can be expressed by general formulae valid for all  $D, l$ ; this allows to avoid the rather complicated actions of the  $L_{hk}$  on spherical harmonics (which also span  $V_D^l$ ) used in [23]. It turns out that both  $\mathcal{H}_\Lambda, V_{D+1}^\Lambda$  decompose into irreps of  $O(D)$  as follows:

$$\mathcal{H}_\Lambda \simeq V_{D+1}^\Lambda = \bigoplus_{l=0}^{\Lambda} V_D^l, \quad (5)$$

The second equality shows that, in contrast with the mentioned fuzzy hyperspheres, we recover (2) in the limit  $\Lambda \rightarrow \infty$ . The first equality suggests that the unitary irrep of the  $*$ -algebra  $\mathcal{A}_\Lambda$  on  $\mathcal{H}_\Lambda$  is isomorphic to the irrep  $\pi_\Lambda$  of  $Uso(D+1)$  on  $V_{D+1}^\Lambda$ , what we in fact prove in section 5 (this had been conjectured in [22, 23]). The relations fulfilled by  $\bar{x}^i, \bar{L}_{hk}$  are determined in section 4: the commutators  $[\bar{x}^i, \bar{x}^j]$  are also Snyder-like [1], i.e. are proportional to  $\bar{L}_{hk}/k$ , with a proportionality factor that is the same constant on all of  $\mathcal{H}_\Lambda$ , except on the  $l = \Lambda$  component of the latter, where it is a slightly different constant.  $\bar{x}^i$  generate the whole  $\mathcal{A}_\Lambda$ . The square distance  $\bar{\mathbf{x}}^2 \equiv \bar{x}^i \bar{x}_i$  is a function of  $\mathbf{L}^2$  only, such that almost all its spectrum is very close to 1 and goes to 1 in the limit  $\Lambda \rightarrow \infty$ . In section 6 we show in which sense  $\mathcal{H}_\Lambda, \mathcal{A}_\Lambda$  go to  $\mathcal{H}, \mathcal{A}$  as  $\Lambda \rightarrow \infty$ , in particular how one can recover  $f \cdot \in C(S^d) \subset \mathcal{A}$ , the multiplication operator of wavefunctions in  $\mathcal{L}^2(S^d)$  by a continuous function  $f$ , as the strong limit of suitable sequence  $f_\Lambda \in \mathcal{A}_\Lambda$  (again, this had been only conjectured in [23]). In section 7 we discuss our results and possible developments in comparison with the literature; in particular, we point out that our pair  $(\mathcal{H}_\Lambda, \mathcal{A}_\Lambda)$  can be seen as a fuzzy quantization of a coadjoint orbit of  $O(D)$  that can be identified with the cotangent space  $T^*S^d$ , the classical phase space over the  $d$ -dimensional sphere. Finally, we have concentrated most proofs in the appendix 8.

## 2 General setting

We choose a set of real Cartesian coordinates  $x := (x^1, \dots, x^D)$  of  $\mathbb{R}^D$  and abbreviate  $\partial_i \equiv \partial/\partial x^i$ . We normalize  $x^i, \partial_i$  and  $H$  itself so as to be dimensionless. Then we can express  $r^2 := \mathbf{x}^2 \equiv x_i x^i$ ,  $\Delta := \partial_i \partial^i$  (sum over repeated indices understood), where actually  $x_i = x^i$  and  $\partial^i = \partial_i$  because the coordinates are real and Cartesian. The self-adjoint operators  $x^i, -i\partial_i$  on  $\mathcal{L}^2(\mathbb{R}^D)$  fulfill the canonical commutation relations

$$[x^i, x^j] = 0, \quad [-i\partial_i, -i\partial_j] = 0, \quad [x^i, -i\partial_j] = i\delta_j^i, \quad (6)$$

which are equivariant under all orthogonal transformations  $Q$  (including parity  $Q = -I$ )

$$x^i \mapsto x'^i = Q_j^i x^j, \quad Q^{-1} = Q^T. \quad (7)$$

All scalars  $S$ , in particular  $S = \Delta, r^2, V, H$ , are invariant. This implies  $[S, L_{ij}] = 0$ , where

$$L_{ij} := i(x^j \partial_i - x^i \partial_j) \quad (8)$$

are the angular momentum components associated to  $x$ . These generate rotations of  $\mathbb{R}^D$ , i.e.

$$[iL_{ij}, v^h] = v^i \delta_j^h - v^j \delta_i^h \quad (9)$$

hold for the components  $v^h$  of all vector operators, in particular  $v^h = x^h, \partial_h$ , and close the commutation relations of  $so(D)$ ,

$$[iL_{ij}, iL_{hk}] = i(L_{ik}\delta_{hj} - L_{jk}\delta_{hi} - L_{ih}\delta_{kj} + L_{jh}\delta_{ki}). \quad (10)$$

The  $D$  derivatives  $\partial_i$  make up a globally defined basis for the linear space of smooth vector fields on  $\mathbb{R}^D$ . As the  $L_{ij}$  are vector fields tangent to all spheres  $r = \text{const}$ , the set  $B = \{\partial_r, L_{ij} \mid i < j\}$  ( $\partial_r := \partial/\partial r$ ) is an alternative complete set that is singular for  $r = 0$ , but globally defined elsewhere; for  $D = 2$  it is a basis, while for  $D > 2$  it is redundant, because of the relations

$$\varepsilon^{i_1 i_2 i_3 \dots i_D} x^{i_1} L_{i_2 i_3} = 0. \quad (11)$$

This redundancy (unavoidable if  $S^d$  is not parallelizable) will be no problem for our purposes.

We shall assume that  $V(r)$  has a very sharp minimum at  $r = 1$  with very large  $k = V''(1)/4 > 0$ , and fix  $V_0 := V(1)$  so that the ground state has zero energy, i.e.  $E_0 = 0$  (see fig. 2). We choose an energy cutoff  $\bar{E}$  fulfilling first of all the condition

$$V(r) \simeq V_0 + 2k(r-1)^2 \quad \text{if } r \text{ fulfills } V(r) \leq \bar{E}, \quad (12)$$

so that we can neglect terms of order higher than two in the Taylor expansion of  $V(r)$  around 1 and approximate the potential as a harmonic one in the classical region  $v_{\bar{E}} \subset \mathbb{R}^D$  determined by the condition  $V(r) \leq \bar{E}$ . By (12),  $v_{\bar{E}}$  is approximately the spherical shell  $|r-1| \leq \sqrt{\frac{\bar{E}-V_0}{2k}}$ ; when both  $\bar{E}-V_0$  and  $k$  diverge, while their ratio goes to zero, then  $v_{\bar{E}}$  reduces to the unit sphere  $S^d$ . We expect that in this limit the dimension of  $\mathcal{H}_{\bar{E}}$  diverges, and we recover standard quantum mechanics on  $S^d$ . As we shall see, this is the case.




Figure 2: Three-dimensional plot of  $V(r)$  in dimension  $D = 2$ .




Figure 3: Two-dimensional plot of  $V(r)$ , including the energy-cutoff.

Of course, the eigenfunctions of  $H$  can be more easily determined in terms of polar coordinates  $r, \theta_1, \dots, \theta_d$ , recalling that the Laplacian in  $D$  dimensions decomposes as follows

$$\Delta = \partial_r^2 + (D-1)\frac{1}{r}\partial_r - \frac{1}{r^2}\mathbf{L}^2, \quad (13)$$

(see section 3.1) where  $\mathbf{L}^2 := L_{ij}L_{ij}/2$  is the square angular momentum (in normalized units), i.e. the quadratic Casimir of  $Uso(D)$  and the Laplacian on the sphere  $S^d$ ;  $\mathbf{L}^2$  can be expressed in terms of angles  $\theta_a$  and derivatives  $\partial/\partial\theta^a$  only. The eigenvalues of  $\mathbf{L}^2$  are  $l(l+D-2)$ , see section 3.1; we denote by  $V_D^l$  the  $\mathbf{L}^2 = E_l$  eigenspace within  $\mathcal{L}^2(S^d)$ . Replacing the Ansatz  $\psi = f(r)Y_l(\boldsymbol{\theta})$ , with  $f(r) = r^{-d/2}g(r)$ ,  $Y_l \in V_D^l$ ,  $\boldsymbol{\theta} \equiv (\theta_1, \dots, \theta_d)$ , transforms the Schrödinger PDE  $H\psi = E\psi$  into the Fuchsian ODE in the unknown  $g(r)$ <sup>1</sup>

$$-g''(r) + \left[ \frac{[D^2 - 4D + 3 + 4l(l+D-2)]}{4r^2} + V(r) \right] g(r) = Eg(r). \quad (14)$$

The requirement  $\psi \in \mathcal{L}^2(\mathbb{R}^D)$  implies that  $g$  belongs to  $\mathcal{L}^2(\mathbb{R}^+)$ , in particular goes to zero as  $r \rightarrow \infty$ . The self-adjointness of  $H$  implies that it must be  $f(0) = 0$ ; this is compatible [23] with Fuchs theorem provided  $r^2V(r) \xrightarrow{r \rightarrow 0^+} T \in \mathbb{R}^+$  [what is in turn compatible with (12)]. Since  $V(r)$  is very large outside the thin spherical shell  $v_{\bar{E}}$  (a neighbourhood of  $S^d$ ),

<sup>1</sup>The present treatment of the equations in the case  $D = 2$  is slightly different from the one adopted in Ref. [22], where the independent radial variable had been changed according to  $r \mapsto \rho \equiv \ln r$ ; however this gives the same wavefunctions at lowest order in  $\rho \simeq r-1$ .

$g, f, \psi$  become negligibly small there, and, by condition (12), the lowest eigenvalues  $E$  are at leading order those of the 1-dimensional harmonic oscillator approximation [23] of (14)

$$-g''(r) + g(r)k_l(r - \tilde{r}_l)^2 = \tilde{E}_l g(r), \quad (15)$$

which is obtained neglecting terms  $O((r-1)^3)$  in the Taylor expansions of  $1/r^2, V(r)$  about  $r=1$ . Here

$$\begin{aligned} \tilde{r}_l &:= 1 + \frac{b(l,D)}{3b(l,D)+2k}, & \tilde{E}_l &:= E - V_0 \frac{2b(l,D)[k+b(l,D)]}{3b(l,D)+2k}, \\ k_l &:= 2k + 3b(l,D), & b(l,D) &:= \frac{D^2-4D+3+4l(l+D-2)}{4}. \end{aligned} \quad (16)$$

The (Hermite functions) square-integrable solutions of (15)

$$g_{n,l}(r) = M_{n,l} e^{-\frac{\sqrt{k_l}}{2}(r-\tilde{r}_l)^2} \cdot H_n \left( (r - \tilde{r}_l) \sqrt[4]{k_l} \right) \quad \text{with } n \in \mathbb{N}_0$$

(here  $M_{n,l}$  are normalization constants and  $H_n$  are the Hermite polynomials) lead to

$$f_{n,l}(r) = \frac{M_{n,l}}{r^{\frac{d}{2}}} e^{-\frac{\sqrt{k_l}}{2}(r-\tilde{r}_l)^2} \cdot H_n \left( (r - \tilde{r}_l) \sqrt[4]{k_l} \right) \quad \text{with } n \in \mathbb{N}_0. \quad (17)$$

The corresponding ‘eigenvalues’ in (15)  $\tilde{E}_{n,l} = (2n+1)\sqrt{k_l}$  lead to energies

$$E_{n,l} = (2n+1)\sqrt{k_l} + V_0 + \frac{2b(l,D)[k+b(l,D)]}{3b(l,D)+2k}.$$

As said, we fix  $V_0$  requiring that the lowest one  $E_{0,0}$  be zero; this implies

$$V_0 = -\sqrt{k_0} - \frac{2b(0,D)[k+b(0,D)]}{3b(0,D)+2k} = -\sqrt{2k} - b(0,D) - \frac{3b(0,D)}{2\sqrt{2k}} + O\left(k^{-\frac{1}{2}}\right),$$

and the expansions of  $E_{n,l}$  and  $\tilde{r}_l$  at leading order in  $k$  become

$$E_{n,l} = l(l+D-2) + 2n\sqrt{2k} + O\left(k^{-\frac{1}{2}}\right), \quad \tilde{r}_l = 1 + \frac{b(l,D)}{2k} + O\left(k^{-2}\right). \quad (18)$$

$E_{0,l}$  coincide at lowest order with the desired eigenvalues  $E_l$  of the Laplacian  $\mathbf{L}^2$  on  $S^d$ , while if  $n > 0$   $E_{n,l}$  diverge as  $k \rightarrow \infty$ ; to exclude all states with  $n > 0$  (i.e., to ‘freeze’ radial oscillations; then all corresponding classical trajectories are circles) we impose the cutoff

$$E_{n,l} \leq \bar{E}(\Lambda) := \Lambda(\Lambda+D-2) < 2\sqrt{2k}, \quad \Lambda \in \mathbb{N}. \quad (19)$$

The right inequality is satisfied prescribing a suitable dependence  $k(\Lambda)$ , e.g.  $k(\Lambda) = [\Lambda(\Lambda+D-2)]^2$ ; the left one is satisfied setting  $n=0$  and  $l \leq \Lambda$ . Abbreviating  $f_l \equiv f_{0,l}$ , we end up with eigenfunctions and associated energies (at leading order in  $1/\Lambda$ )

$$\psi_l(r, \boldsymbol{\theta}) = f_l(r) Y_l(\boldsymbol{\theta}), \quad H\psi_l = E_l \psi_l, \quad l = 0, 1, \dots, \Lambda. \quad (20)$$

Thus  $\mathcal{H}_\Lambda$  decomposes into irreps of  $O(D)$  (and eigenspaces of  $\mathbf{L}^2, H$ ) as follows

$$\mathcal{H}_\Lambda = \bigoplus_{l=0}^{\Lambda} \mathcal{H}_\Lambda^l, \quad \mathcal{H}_\Lambda^l := f_l(r) V_D^l. \quad (21)$$

We can express the projections  $P_\Lambda^l : \mathcal{H}_\Lambda \rightarrow \mathcal{H}_\Lambda^l$  as the following polynomials in  $\bar{\mathbf{L}}^2$ :

$$P_\Lambda^l = \prod_{n=0, n \neq l}^\Lambda \frac{\bar{\mathbf{L}}^2 - E_n}{E_l - E_n}. \quad (22)$$

In the commutative limit  $\Lambda \rightarrow \infty$  the spectrum  $\{E_{0,l}\}_{l=0}^\Lambda$  of  $\bar{H}$  goes to the whole spectrum  $\{E_l\}_{l \in \mathbb{N}_0}$  of  $\mathbf{L}^2$ . If  $\phi, \phi' \in \mathcal{H} \equiv \mathcal{L}^2(\mathbb{R}^D)$  can be factorized into radial parts  $f(r), f'(r)$  and angular parts  $T, T' \in \mathcal{L}^2(S^d)$ , i.e.  $\phi = f T, \phi' = f' T'$ , then so can be their scalar product:

$$(\phi, \phi') := \int_{\mathbb{R}^D} d^D x \phi^*(x) \phi'(x) = \langle T, T' \rangle \int_0^\infty dr r^d f^*(r) f'(r). \quad (23)$$

Here we have denoted by  $\langle \cdot, \cdot \rangle$  the scalar product of  $\mathcal{L}^2(S^d)$ ,

$$\langle T, T' \rangle := \int_{S^d} d\alpha T^* T', \quad (24)$$

where  $d\alpha$  is the  $O(D)$ -invariant measure on  $S^{d2}$ . Assume that  $\mathcal{B} := \{Y_l^{\mathbf{m}}\}_{(l, \mathbf{m}) \in \mathfrak{l}}$ , is an orthonormal basis of  $\mathcal{L}^2(S^d)$  consisting of eigenvectors of  $\mathbf{L}^2$  (e.g. spherical harmonics),

$$\mathbf{L}^2 Y_l^{\mathbf{m}} = E_l Y_l^{\mathbf{m}}, \quad \langle Y_{l'}^{\mathbf{m}'}, Y_l^{\mathbf{m}} \rangle = \delta^{\mathbf{m}\mathbf{m}'} \delta_{ll'}; \quad (25)$$

here  $\mathbf{m} \in \mathfrak{l}$  is a (multi-)index<sup>3</sup> labelling the elements of an orthonormal basis  $\mathcal{B}_l \equiv \{Y_l^{\mathbf{m}}\}_{\mathbf{m} \in \mathfrak{l}_l}$  of  $V_D^l$ , and  $\mathfrak{l} := \{(l, \mathbf{m}) \mid l \in \mathbb{N}_0, \mathbf{m} \in \mathfrak{l}_l\}$ . Then, by appropriate choices of the normalization constants<sup>4</sup> of (17), one obtains as orthonormal bases respectively of  $\mathcal{H}$  and  $\mathcal{H}_\Lambda^l$

$$\mathcal{B}' := \{\psi_{n,l}^{\mathbf{m}} := f_{n,l}(r) Y_l^{\mathbf{m}}, \mid n \in \mathbb{N}_0, \mathbf{l} \in \mathcal{I}\}, \quad \mathcal{B}'_l := \{\psi_{0,l}^{\mathbf{m}} = f_l(r) Y_l^{\mathbf{m}}\}_{\mathbf{m} \in \mathfrak{l}_l}.$$

The projection  $\tilde{P}_\Lambda^l : \mathcal{H} \rightarrow \mathcal{H}_\Lambda^l$  acts by

$$\left(\tilde{P}_\Lambda^l \phi\right)(x) = \sum_{\mathbf{m} \in \mathfrak{l}_l} \psi_{0,l}^{\mathbf{m}}(x) (\psi_{0,l}^{\mathbf{m}}, \phi) = \sum_{\mathbf{m} \in \mathfrak{l}_l} \psi_{0,l}^{\mathbf{m}}(x) \int_{\mathbb{R}^D} d^D x' \psi_{0,l}^{\mathbf{m}*}(x') \phi(x').$$

If  $\phi$  has the form  $\phi(r, \boldsymbol{\theta}) = \Theta_j(\boldsymbol{\theta}) \phi(r)$ , with  $\Theta_j \in V_D^j$ , then by (23), (25) this simplifies to

$$\left(\tilde{P}_\Lambda^l \phi\right)(r, \boldsymbol{\theta}) = \delta_{lj} \Theta_j(\boldsymbol{\theta}) f_l(r) \int_0^\infty r^d dr f_l^*(r) \phi(r), \quad (26)$$

which is zero if  $l \neq j$ , has the same angular dependence  $\Theta_j(\boldsymbol{\theta})$  as  $\phi$  if  $l = j$ .

In next section we provide an explicit characterization of elements  $\Theta_l \in V_D^l$  as polynomials in the coordinates  $t^i$  of points of the unit sphere  $S^d$ , which fulfill the relation

$$t^i t_i = 1 \quad (\text{sum over } i), \quad (27)$$

rather than as combinations of spherical harmonics  $Y_l^{\mathbf{m}}(\boldsymbol{\theta})$ ,  $\mathbf{m} \in \mathfrak{l}_l$ .

<sup>2</sup>In terms of the angles  $\boldsymbol{\theta}$ ,  $d\alpha = [\sin^{d-1}(\theta_d) \sin^{d-2}(\theta_{d-1}) \cdots \sin(\theta_2)] d\theta_1 d\theta_2 \cdots d\theta_d$ .

<sup>3</sup>If  $D = 2, 3$ , then  $\mathfrak{l} \subset \mathbb{Z}$ ; more precisely  $\mathfrak{l}_0 = \{0\}$ , while for  $l > 0$  it is  $\mathfrak{l}_l = \{+, -\}$  if  $D = 2$ ,  $\mathfrak{l}_l = \{-l, 1-l, \dots, l\}$  if  $D = 3$ . If  $D > 3$ , then  $\mathfrak{l} \subset \mathbb{Z}^{D-2}$ .

<sup>4</sup>Choosing them positive, one easily finds  $M_l = \sqrt[8]{k_l} / \sqrt[4]{\pi}$ . In fact, by (25), (23) normalizing  $\psi_{0,l}^{\mathbf{m}}$  amounts to  $1 = \int_0^\infty dr r^d f_l^2(r) = \int_0^\infty dr g_l^2(r) \simeq M_l^2 \int_{-\infty}^\infty dr e^{-\sqrt{k_l}(r-\tilde{r}_l)^2}$ ; here  $\simeq$  (due to the shift  $0 \mapsto -\infty$  of the left integration extreme) means equality up to terms of the order  $1/\sqrt{2k_l} e^{-\sqrt{2k_l}}$ , which has zero asymptotic expansion in  $1/\sqrt{2k_l}$ , see [22].



### 3 Representations of $O(D)$ via polynomials in $x^i, t^i$

The differential operator  $\mathbf{L}^2$  can be expressed as

$$\mathbf{L}^2 = \eta(D-2+\eta) - r^2\Delta, \quad \eta := x^i\partial_i. \quad (28)$$

The ‘dilatation operator’  $\eta$  and the Laplacian  $\Delta$  fulfill

$$\eta x^i = x^i(\eta + 1), \quad \eta\partial_i = \partial_i(\eta - 1), \quad (29)$$

$$\Delta x^i = x^i\Delta + 2\partial^i, \quad \Delta r^2 = r^2\Delta + 4\eta + 2D. \quad (30)$$

In particular, the action of  $\eta$  on monomials in the  $x^i$  amounts to multiplication by their total degree. In terms of polar coordinates it is  $\eta = r\partial_r$ , which replaced in (28) gives (13).

Let  $\mathbb{C}[x^1, \dots, x^D]$  be the space of complex polynomial functions on  $\mathbb{R}^D$  and, for all  $l \in \mathbb{N}_0$ , let  $W_D^l$  be the subspace of homogeneous ones of degree  $l$ . The monomials of degree  $l$   $x^{i_1}x^{i_2}\dots x^{i_l} \in W_D^l$  can be reordered in the form  $(x^1)^{l_1}\dots(x^D)^{l_D}$  and make up a basis of  $W_D^l$ :

$$\mathcal{B}_{W_D^l} := \left\{ (x^1)^{l_1}\dots(x^D)^{l_D} \mid (l_1, \dots, l_D) \in \mathbb{N}_0, \sum_{i=1}^D l_i = l \right\}, \quad \dim(W_D^l) = \binom{D+l-1}{l};$$

the dimension of  $W_D^l$  is the number of elements of  $\mathcal{B}_{W_D^l}$ . Clearly  $W_D^l$  carries a representation of  $O(D)$  as well as  $Uso(D)$ , but this is reducible if  $l \geq 2$ ; in fact, the subspace  $r^2W_D^{l-2} \subset W_D^l$  manifestly carries a smaller representation. We denote by  $\check{V}_D^l$  the ‘‘trace-free’’ component of  $W_D^l$ , namely the subspace such that  $W_D^l = \check{V}_D^l \oplus r^2W_D^{l-2}$ . As a consequence,

$$\dim(\check{V}_D^l) = \dim(W_D^l) - \dim(W_D^{l-2}) = \frac{(l+D-3)\dots(l+1)}{(D-2)!}(D+2l-2). \quad (31)$$

$\check{V}_D^l$  carries the irreducible representation (irrep)  $\pi_D^l$  of  $Uso(D)$  and  $O(D)$  characterized by the highest eigenvalue of  $\mathbf{L}^2$  within  $W_D^l$ , namely  $E_l$  (the eigenvalues of all other  $[D/2-1]$  Casimirs are determined by  $l$ ). Abbreviating  $X_{l,\pm}^{hk} := (x^h \pm ix^k)^l$ , this can be easily shown observing that for all  $h, k \in \{1, \dots, D\}$   $X_{l,\pm}^{hk} \in W_D^l$  are annihilated by  $\Delta$  and are eigenvectors of  $\mathbf{L}^2$  with that eigenvalue; moreover, they are eigenvectors of  $L_{hk}$  with eigenvalue  $\pm l$ . Hence  $X_{l,+}^{hk}, X_{l,-}^{hk}$  can be used as the highest and lowest weight vectors of  $\check{V}_D^l$ <sup>5</sup>. Since all the  $L_{ij}$  commute with  $\Delta$ ,  $\check{V}_D^l$  can be characterized also as the subspace of  $W_D^l$  which is annihilated by  $\Delta$ . A

<sup>5</sup>In fact, in terms of Cartesian coordinates, using (28), (30) we immediately find the following commutation relations among operators of multiplication  $(x^h \pm ix^k)$  and differential operators

$$\begin{aligned} (\partial_h + i\partial_k)(x^h + ix^k) &= (x^h + ix^k)(\partial_h + i\partial_k), & (\partial_h - i\partial_k)(x^h - ix^k) &= (x^h - ix^k)(\partial_h - i\partial_k) \\ \Delta(x^h \pm ix^k) &= (x^h \pm ix^k)\Delta + 2(\partial_h \pm i\partial_k), & \Delta X_{l,\pm}^{hk} &= X_{l,\pm}^{hk}\Delta + 2lX_{l-1,\pm}^{hk}(\partial_h + i\partial_k), \\ L_{hk}(x^h \pm ix^k) &= (x^h \pm ix^k)(L_{hk} \pm 1), & L_{hk}(L_{hj} \pm iL_{kj}) &= (L_{hj} \pm iL_{kj})[L_{hk} \pm 1]. \end{aligned} \quad (32)$$

Consequently, we obtain the following functions at the rhs as results of the operator actions on functions at

complete set in  $\check{V}_D^l$  consists of trace-free homogeneous polynomials  $X_l^{i_1 i_2 \dots i_l}$ , which we will obtain below applying the completely symmetric trace-free projector  $\mathcal{P}^l$  to the  $x^{i_1} x^{i_2} \dots x^{i_l}$ 's.

We slightly enlarge  $\mathbb{C}[x^1, \dots, x^D]$  introducing as new generators  $r, r^{-1}$  subject to the relations  $r^2 = x^i x_i$  (sum over  $i$ ),  $rr^{-1} = 1$ . Inside this enlarged algebra the elements

$$t^i := \frac{x^i}{r} \quad (36)$$

fulfill the relation (27) characterizing the coordinates of points of the unit sphere  $S^d$ . Choosing  $g(r) = r^{-l}$  in (32-35) we obtain the same relations with  $x^i, X_{l,\pm}^{hk}$  replaced by  $t^i, T_{l,\pm}^{hk} := (t^h \pm i t^k)^l$ . We shall denote by  $Pol_D$  the algebra of complex polynomials in such  $t^i$ , by  $Pol_D^\Lambda$  the subspace of polynomials up to degree  $\Lambda$ , by  $P^\Lambda : Pol_D \rightarrow Pol_D^\Lambda$  the corresponding projection.  $Pol_D$  endowed with the scalar product  $\langle T, T' \rangle := \int_{S^d} d\alpha T^* T'$  is a pre-Hilbert space; its completion is  $\mathcal{L}^2(S^d)$ . We extend  $P^\Lambda$  to all of  $\mathcal{L}^2(S^d)$  by continuity in the norm of the latter. Also  $Pol_D^\Lambda, V_D^l$  are Hilbert subspaces of  $\mathcal{L}^2(S^d)$ .  $Pol_D^\Lambda = W_D^\Lambda r^{-\Lambda} \oplus W_D^{\Lambda-1} r^{1-\Lambda}$  carries a unitary reducible representation of  $O(D)$  [and  $Uso(D)$ ] which splits via  $Pol_D^\Lambda = \bigoplus_{l=0}^\Lambda V_D^l$  into irreps carried by  $V_D^l := \check{V}_D^l / r^l$ . Its dimension is thus

$$\begin{aligned} \dim(Pol_D^\Lambda) &= \sum_{l=0}^\Lambda \dim(\check{V}_D^l) = \sum_{l=0}^\Lambda \dim(W_D^l) - \sum_{l=2}^\Lambda \dim(W_D^{l-2}) \\ &= \dim(W_D^\Lambda) + \dim(W_D^{\Lambda-1}) \\ &= \frac{(D+\Lambda-1)\dots(\Lambda+1)}{(D-1)!} + \frac{(D+\Lambda-2)\dots\Lambda}{(D-1)!} \\ &= \frac{(D+\Lambda-2)\dots(\Lambda+1)}{(D-1)!} (D+2\Lambda-1) =: N \end{aligned} \quad (37)$$

$$\stackrel{(31)}{=} \dim(V_{D+1}^\Lambda). \quad (38)$$

This suggests that  $\mathcal{H}_\Lambda \simeq Pol_D^\Lambda \simeq V_{D+1}^\Lambda$  as  $Uso(D)$  (reducible) representations. We have proved the first isomorphism in section 2 and will prove the second in section 3.2.

the lhs:  $\Delta X_{l,\pm}^{hk} = 0$  and, for all functions  $g(r)$ ,

$$\mathbf{L}^2 g(r) X_{l,\pm}^{hk} = g(r) [\eta(D-2+\eta) - r^2 \Delta] X_{l,\pm}^{hk} = E_l g(r) X_{l,\pm}^{hk}, \quad (33)$$

$$L_{hk} X_{l,\pm}^{hk} g(r) = \pm l X_{l,\pm}^{hk} g(r), \quad L_{hk} X_{l,+}^{hk} X_{m,-}^{hk} g(r) = (l-m) X_{l,+}^{hk} X_{m,-}^{hk} g(r). \quad (34)$$

Denoting by  $\tau = \lceil \frac{D}{2} \rceil$  the rank of  $so(D)$ , as a basis of a Cartan subalgebra of  $so(D)$  one can take any set  $\{H_1 \equiv L_{i_1 i_2}, H_2 \equiv L_{i_3 i_4}, \dots, H_\tau \equiv L_{i_{2\tau-1} i_{2\tau}}\}$ , with  $i_1, i_2, \dots, i_{2\tau} \in \{1, \dots, D\}$  all different from each other. If  $(i_1, i_2) = (h, k)$ , by (33), (34)  $X_{l,+}^{hk}, X_{l,-}^{hk}$  are the corresponding highest, lowest weight vectors, in the sense

$$H_1 X_{l,\pm}^{hk} g(r) = \pm l X_{l,\pm}^{hk} g(r), \quad H_a X_{l,\pm}^{hk} g(r) = 0 \text{ if } a > 1. \quad (35)$$

### 3.1 $O(D)$ -irreps via trace-free completely symmetric projectors

Let  $(\pi, \mathcal{E})$  be the  $D$ -dimensional irreducible unitary representation of  $Uso(D)$  and  $O(D)$ ; the carrier space  $\mathcal{E}$  is isomorphic to  $V_D^1$ . As a vector space  $\mathcal{E} \simeq \mathbb{R}^D$ ; the set of coordinates  $x := (x^1, \dots, x^D) \in \mathbb{R}^D$  can be seen as the set of components of an element of  $\mathcal{E}$  with respect to (w.r.t.) an orthonormal basis. The permutator on  $\mathcal{E}^{\otimes 2} \equiv \mathcal{E} \otimes \mathcal{E}$  is defined via  $\mathbf{P}(u \otimes v) = v \otimes u$  and linearly extended. In all bases it is represented by the  $D^2 \times D^2$  matrix  $\mathbf{P}_{jk}^{hi} = \delta_k^h \delta_j^i$ . The symmetric and antisymmetric projectors  $\mathcal{P}^+, \mathcal{P}^-$  on  $\mathcal{E}^{\otimes 2}$  are obtained as

$$\mathcal{P}^\pm = \frac{1}{2}(\mathbf{1}_{D^2} \pm \mathbf{P}); \quad (39)$$

here and below we denote by  $\mathbf{1}_{D^l}$  the identity operator on  $\mathcal{E}^{\otimes l}$ , which in all bases is represented by the  $D^l \times D^l$  matrix  $\mathbf{1}_{D^l}^{h_1 \dots h_l} := \delta_{i_1}^{h_1} \dots \delta_{i_l}^{h_l}$ . The antisymmetrized tensor product  $\mathcal{P}^- \mathcal{E}^{\otimes 2}$  is an irrep under  $O(D)$ , while the symmetrized one  $\mathcal{P}^+ \mathcal{E}^{\otimes 2}$  contains two irreps: the 1-dim trace one and the trace-free symmetric one. The matrix representation of the 1-dim projector  $\mathcal{P}^t$  on the former is

$$\mathcal{P}_{kl}^{tij} = \frac{1}{D} g^{ij} g_{kl} \quad (40)$$

where the  $D \times D$  metric matrix  $g_{ij}$  (in the chosen basis) is a  $so(D)$ -isotropic symmetric tensor, and  $g^{ij} g_{jh} = \delta_h^i$ , whence  $g^{ij} g_{ij} = D$ . Here we shall use an orthonormal basis of  $\mathcal{E}$ , whence  $g_{ij} = g^{ij} = \delta_{ij}$ , and indices of vector components can be raised or lowered freely, e.g.  $x_i = x^i$ . The  $\frac{1}{2}(D-1)(D+2)$ -dim trace-free symmetric projector  $\mathcal{P}^s$  is given by

$$\mathcal{P}^s := \mathcal{P}^+ - \mathcal{P}^t = \frac{1}{2}(\mathbf{1}_{D^2} + \mathbf{P}) - \mathcal{P}^t. \quad (41)$$

These projectors satisfy the equations

$$\mathcal{P}^\alpha \mathcal{P}^\beta = \mathcal{P}^\alpha \delta^{\alpha\beta}, \quad \sum_\alpha \mathcal{P}^\alpha = \mathbf{1}_{D^2}, \quad (42)$$

where  $\alpha, \beta = -, s, t$ . In the sequel we shall abbreviate  $\mathcal{P} \equiv \mathcal{P}^s$ . This implies in particular  $\mathcal{P}\mathcal{P}' = 0$ , where we have introduced the new projector  $\mathcal{P}' := \mathcal{P}^- + \mathcal{P}^t$ .  $\mathbf{P}, \mathcal{P}^t$  are symmetric matrices, i.e. invariant under transposition  $T$ , and therefore also the other projectors are:

$$\mathbf{P}^T = \mathbf{P}, \quad \mathcal{P}^{\alpha T} = \mathcal{P}^\alpha. \quad (43)$$

Given a (linear) operator  $M$  on  $\mathcal{E}^{\otimes n}$ , for all integers  $l, h$  with  $l > n$ , and  $1 \leq h \leq l+1-n$  we denote by  $M_{h(h+1)\dots(h+n-1)}$  the operator on  $\mathcal{E}^{\otimes l}$  acting as the identity on the first  $h-1$  and the last  $l+1-n-h$  tensor factors, and as  $M$  in the remaining central ones. For instance, if  $M = \mathbf{P}$  and  $l = 3$  we have  $\mathbf{P}_{12} = \mathbf{P} \otimes \mathbf{1}_D$ ,  $\mathbf{P}_{23} = \mathbf{1}_D \otimes \mathbf{P}$ . It is straightforward to check

**Proposition 3.1** *All the projectors  $A = \mathcal{P}^+, \mathcal{P}^-, \mathcal{P}, \mathcal{P}^t, \mathcal{P}'$  fulfill the “braid” relation*

$$A_{12} \mathbf{P}_{23} \mathbf{P}_{12} = \mathbf{P}_{23} \mathbf{P}_{12} A_{23}. \quad (44)$$

Moreover,

$$D \mathcal{P}_{23}^t \mathcal{P}_{12}^t = \mathbf{P}_{12} \mathbf{P}_{23} \mathcal{P}_{12}^t, \quad D \mathbf{P}_{12} \mathcal{P}_{23}^t \mathcal{P}_{12}^t = \mathbf{P}_{23} \mathcal{P}_{12}^t, \quad (45)$$

$$D \mathcal{P}_{12}^t \mathcal{P}_{23}^t = \mathbf{P}_{23} \mathbf{P}_{12} \mathcal{P}_{23}^t, \quad D \mathbf{P}_{23} \mathcal{P}_{12}^t \mathcal{P}_{23}^t = \mathbf{P}_{12} \mathcal{P}_{23}^t, \quad (46)$$

$$D \mathcal{P}_{23}^t \mathcal{P}_{12}^t = \mathcal{P}_{23}^t \mathbf{P}_{12} \mathbf{P}_{23}, \quad D \mathcal{P}_{23}^t \mathcal{P}_{12}^t \mathbf{P}_{23} = \mathcal{P}_{23}^t \mathbf{P}_{12}; \quad (47)$$

Eq. (44-47) hold also for  $l > 3$ , e.g. for all  $2 \leq h \leq l-1$

$$A_{(h-1)h} \mathbf{P}_{h(h+1)} \mathbf{P}_{(h-1)h} = \mathbf{P}_{h(h+1)} \mathbf{P}_{(h-1)h} A_{h(h+1)}. \quad (48)$$

**Proof** Since  $A = \mathbf{1}_{D^2}, \mathbf{P}, \mathcal{P}^t$  fulfill (44), then also  $A = \mathcal{P}^+, \mathcal{P}^-, \mathcal{P}, \mathcal{P}^t, \mathcal{P}'$  do. One can immediately check the first equality in (146) via direct calculation; left multiplying the first by  $\mathbf{P}_{12}$  one obtains the second. Eq. (46) are obtained from (146) exchanging  $1 \leftrightarrow 3$  and using the symmetry of  $\mathbf{P}, \mathcal{P}^t$  under the flip. Eq. (47) are obtained from (46) by transposition.  $\square$

Next, we define and determine the completely symmetric trace-free projector  $\mathcal{P}^l$  on  $\mathcal{E}^{\otimes l}$  generalizing  $\mathcal{P}^2 \equiv \mathcal{P}$  to  $l > 2$ . It projects the tensor product of  $l$  copies of  $\mathcal{E}$  to the carrier space of the  $l$ -fold completely symmetric irrep of  $Uso(D)$ , isomorphic to  $V_D^l$ , therein contained. It is uniquely characterized by the following properties:

$$\begin{aligned} \mathcal{P}^l \mathcal{P}_{n(n+1)}^- &= 0, & \mathcal{P}^l \mathcal{P}_{n(n+1)}^t &= 0, \\ \mathcal{P}_{n(n+1)}^- \mathcal{P}^l &= 0, & \mathcal{P}_{n(n+1)}^t \mathcal{P}^l &= 0, \end{aligned} \quad n = 1, \dots, l-1 \quad (49)$$

$$(\mathcal{P}^l)^2 = \mathcal{P}^l, \quad (50)$$

Consequently, it is also  $\text{tr}_{1\dots l}(\mathcal{P}^l) = \dim(V_D^l)$ , which guarantees that  $\mathcal{P}^l$  acts as the identity (and not as a proper projector) on  $V_D^l$ . The right relations in (49) amount to

$$\mathcal{P}_{j_1 \dots j_l}^{i_1 \dots i_l} \delta^{j_n j_{n+1}} = 0, \quad \delta_{i_n i_{n+1}} \mathcal{P}_{j_1 \dots j_l}^{i_1 \dots i_l} = 0, \quad n = 1, \dots, l-1. \quad (51)$$

Clearly the whole of (49) can be summarized as  $\mathcal{P}^l \mathcal{P}'_{n(n+1)} = 0 = \mathcal{P}'_{n(n+1)} \mathcal{P}^l$ . It is straightforward to prove that the above properties imply also the ones

$$\mathcal{P}^l \mathbf{P}_{n(n+1)} = \mathcal{P}^l, \quad n = 1, \dots, l-1; \quad \mathcal{P}^l \mathcal{P}_{(i+1)\dots(i+h)}^h = \mathcal{P}^l, \quad h < l, \quad 0 \leq i \leq l-h. \quad (52)$$

**Proposition 3.2** *The projector  $\mathcal{P}^{l+1}$  can be expressed as a polynomial in the permutators  $\mathbf{P}_{12}, \dots, \mathbf{P}_{(l-1)l}$  and trace projectors  $\mathcal{P}_{12}^t, \dots, \mathcal{P}_{(l-1)l}^t$  through either recursive relation*

$$\mathcal{P}^{l+1} = \mathcal{P}_{12\dots l}^l M_{l(l+1)} \mathcal{P}_{12\dots l}^l, \quad (53)$$

$$= \mathcal{P}_{2\dots(l+1)}^l M_{12} \mathcal{P}_{2\dots(l+1)}^l, \quad (54)$$

$$\text{where } M \equiv M(l+1) = \frac{1}{l+1} \left[ \mathbf{1}_{D^2} + l \mathbf{P} - \frac{2Dl}{D+2l-2} \mathcal{P}^t \right] \quad (55)$$

As a consequence, the  $\mathcal{P}^l$  are symmetric,  $(\mathcal{P}^l)^T = \mathcal{P}^l$ .

This is the analog of Proposition 1 in [26] for the quantum group  $U_q so(D)$  covariant symmetric projections; the proof is in the appendix. By a straightforward computation one checks that

$$\mathcal{P}_{h j_1 \dots j_l}^{l+1 h i_1 \dots i_l} = \frac{1}{l+1} \left[ D + l - \frac{2l}{D+2l-2} \right] \mathcal{P}_{j_1 \dots j_l}^{l i_1 \dots i_l} \quad (56)$$

( $h$  is summed over). Using (30), (51) one easily shows that the homogeneous polynomials

$$X_l^{i_1 \dots i_l} := \mathcal{P}_{j_1 \dots j_l}^{l i_1 \dots i_l} x^{j_1} \dots x^{j_l} \in \check{V}_D^l \quad (57)$$

are harmonic, i.e. satisfy  $\Delta X_l^{i_1 \dots i_l} = 0$ ; using (28), we find that they are eigenvectors of  $\mathbf{L}^2$ ,

$$\mathbf{L}^2 X_l^{i_1 \dots i_l} = E_l X_l^{i_1 \dots i_l}, \quad (58)$$

with eigenvalues (3). They make up a complete set in  $\check{V}_D^l$ , which can be thus also characterized as the subspace of  $W_D^l$  that is annihilated by  $\Delta$ , whereas  $\Delta \phi \neq 0$  for all  $\phi \in r^2 W_D^{l-2}$ . The  $X_l^{i_1 \dots i_l}$  are not all independent, because they are invariant under permutations of  $(i_1 \dots i_l)$  and by (51) fulfill the linear dependence relations

$$\delta_{i_n i_{n+1}} X_l^{i_1 \dots i_l} = 0, \quad n = 1, \dots, l-1. \quad (59)$$

**Proposition 3.3** *In a compact notation,*

$$\begin{aligned} (\mathcal{P}_{1 \dots l}^l - \mathcal{P}^{l+1}) x_1 \dots x_{l+1} &= \zeta_{l+1} \mathcal{P}_{1 \dots l}^l \mathcal{P}_{l(l+1)}^t x_1 \dots x_{l+1}, \\ (\mathcal{P}_{2 \dots (l+1)}^l - \mathcal{P}^{l+1}) x_1 \dots x_{l+1} &= \zeta_{l+1} \mathcal{P}_{2 \dots (l+1)}^l \mathcal{P}_{12}^t x_1 \dots x_{l+1}, \end{aligned} \quad \zeta_{l+1} = \frac{Dl}{D+2l-2}. \quad (60)$$

The proof is in Appendix 8.2. More explicitly, (60) becomes

$$x^h X_l^{i_1 \dots i_l} - X_{l+1}^{h i_1 \dots i_l} = \frac{\zeta_{l+1}}{D} r^2 \mathcal{P}_{h j_2 \dots j_l}^{l i_1 i_2 \dots i_l} x^{j_2} \dots x^{j_l} = \frac{\zeta_{l+1}}{D} r^2 \mathcal{P}_{h j_2 \dots j_l}^{l i_1 i_2 \dots i_l} X_{l-1}^{j_2 \dots j_l} \quad (61)$$

Contracting the previous relation with  $\delta_{h i_1}$  and using (51) we obtain

$$x^h X_l^{h i_2 \dots i_l} = \frac{\zeta_{l+1}}{D} r^2 \mathcal{P}_{h j_2 \dots j_l}^{l h i_2 \dots i_l} X_{l-1}^{j_2 \dots j_l} \stackrel{(56)}{=} \frac{r^2}{D+2l-2} \left[ D+l-1 - \frac{2l-2}{D+2l-4} \right] X_{l-1}^{i_2 \dots i_l}. \quad (62)$$

In the Appendix we also prove

**Proposition 3.4** *The maps  $L_{hk} : \check{V}_D^l \rightarrow \check{V}_D^l$  explicitly act as follows:*

$$\begin{aligned} i L_{hk} X_l^{i_1 \dots i_l} &= l \frac{\zeta_{l+2}}{\zeta_{l+1}} \left( \mathcal{P}_{k j_1 \dots j_l}^{l+1 h i_1 \dots i_l} - \mathcal{P}_{h j_1 \dots j_l}^{l+1 k i_1 \dots i_l} \right) X_l^{j_1 \dots j_l}, \\ &= l \mathcal{P}_{j_1 \dots j_l}^{l i_1 \dots i_l} \left( \delta^{k j_1} X_l^{h j_2 \dots j_l} - \delta^{h j_1} X_l^{k j_2 \dots j_l} \right). \end{aligned} \quad (63)$$

Dividing (58), (59), (61), (62), (63) by the appropriate powers of  $r$  we find

**Proposition 3.5** *The  $T_l^{i_1 i_2 \dots i_l} := X_l^{i_1 i_2 \dots i_l} / r^l = \mathcal{P}_{j_1 \dots j_l}^{i_1 \dots i_l} t^{j_1} \dots t^{j_l}$  belong to  $V_D^l$ , because*

$$\mathbf{L}^2 T_l^{i_1 \dots i_l} = E_l T_l^{i_1 \dots i_l}; \quad (64)$$

*they make up a complete set  $\mathcal{T}_l$  in it, but not a basis, because they are invariant under permutations of  $(i_1 \dots i_l)$  and by (59) fulfill the linear dependence relations*

$$\delta_{i_n i_{n+1}} T_l^{i_1 \dots i_l} = 0, \quad n = 1, \dots, l-1. \quad (65)$$

*The actions of the operators  $t^h$ ,  $iL_{hk}$  on the  $T_l^{i_1 \dots i_l}$  explicitly read*

$$t^h T_l^{i_1 \dots i_l} = T_{l+1}^{h i_1 \dots i_l} + \frac{l}{D+2l-2} \mathcal{P}_{h j_2 \dots j_l}^{l i_1 i_2 \dots i_l} T_{l-1}^{j_2 \dots j_l} \in V_D^{l+1} \oplus V_D^{l-1}, \quad (66)$$

$$t^i T_l^{i_1 i_2 \dots i_l} = \frac{1}{D+2l-2} \left[ D+l-1 - \frac{2l-2}{D+2l-4} \right] T_{l-1}^{i_2 \dots i_l} \in V_D^{l-1}, \quad (67)$$

$$\begin{aligned} iL_{hk} T_l^{i_1 \dots i_l} &= l \frac{\zeta_{l+2}}{\zeta_{l+1}} \left( \mathcal{P}_{k j_1 \dots j_l}^{l+1 h i_1 \dots i_l} - \mathcal{P}_{h j_1 \dots j_l}^{l+1 k i_1 \dots i_l} \right) T_l^{j_1 \dots j_l}, \\ &= l \mathcal{P}_{j_1 \dots j_l}^{l i_1 \dots i_l} \left( \delta^{k j_1} T_l^{h j_2 \dots j_l} - \delta^{h j_1} T_l^{k j_2 \dots j_l} \right). \end{aligned} \quad (68)$$

For all  $\phi \in \mathcal{H}_s \equiv \mathcal{L}^2(S^d)$  let

$$\phi = \sum_{l=0}^{\infty} \sum_{\mathbf{m} \in l_l} \phi_{\mathbf{m}}^l Y_l^{\mathbf{m}} = \sum_{l=0}^{\infty} \sum_{i_1, \dots, i_l} \phi_{i_1 \dots i_l}^l T_l^{i_1 \dots i_l} \quad (69)$$

be its decompositions in the basis of spherical harmonics and in the complete set  $\mathcal{T} := \bigcup_{l=0}^{\infty} \mathcal{T}_l$ ; here the two sets of coefficients are related by  $\phi_{i_1 \dots i_l}^l = \sum_{\mathbf{m} \in l_l} \phi_{\mathbf{m}}^l A_{i_1 \dots i_l}^{\mathbf{m}}$ , where  $A_{i_1 \dots i_l}^{\mathbf{m}}$  are such that  $Y_l^{\mathbf{m}} = \sum_{i_1 \dots i_l} A_{i_1 \dots i_l}^{\mathbf{m}} T_l^{i_1 \dots i_l}$ . The  $\phi_{i_1 \dots i_l}^l$  are uniquely determined if, as we shall assume, we choose them trace-free and completely symmetric, i.e. fulfilling

$$\phi_{i_1 \dots i_l}^l = \phi_{j_1 \dots j_l}^l \mathcal{P}_{i_1 \dots i_l}^{l j_1 \dots j_l}, \quad (70)$$

whence  $\phi_{i_1 \dots i_l}^l \delta^{i_n i_{n+1}} = 0$  for  $n = 1, \dots, l-1$ . Then (69) can be also written in the form

$$\phi = \sum_{l=0}^{\infty} \sum_{i_1, \dots, i_l} \phi_{i_1 \dots i_l}^l t^{i_1} \dots t^{i_l}. \quad (71)$$

The projection  $P_{\Lambda}$  acts by truncation,  $P_{\Lambda} \phi \equiv \phi_{\Lambda} := \sum_{l=0}^{\Lambda} \sum_{\mathbf{m} \in l_l} \phi_{\mathbf{m}}^l Y_l^{\mathbf{m}} = \sum_{l=0}^{\Lambda} \phi_{i_1 \dots i_l}^l T_l^{i_1 \dots i_l}$  (sum over  $i_1, \dots, i_l$ ). Clearly  $\phi_{\Lambda} \xrightarrow{\Lambda \rightarrow \infty} \phi$  in the  $\mathcal{H}_s$ -norm  $\|\cdot\|$ .

All completely symmetric,  $O(D)$ -isotropic tensors of even rank  $\bar{N}$  are proportional to

$$G_{\bar{N}}^{i_1 \dots i_{\bar{N}}} := \delta^{i_1 i_2} \delta^{i_3 i_4} \dots \delta^{i_{\bar{N}-1} i_{\bar{N}}} + \text{permutations of } (i_1, \dots, i_{\bar{N}}). \quad (72)$$

**Proposition 3.6** The ‘trace’ of  $G_{\bar{N}}$ ,  $tr(G_{\bar{N}}) := \delta_{i_1 i_2} \delta_{i_3 i_4} \dots \delta_{i_{\bar{N}-1} i_{\bar{N}}} G_{\bar{N}}^{i_1 \dots i_{\bar{N}}}$ , is equal to

$$tr(G_{\bar{N}}) = \bar{N}!! D(D+2)\dots(D+\bar{N}-2). \quad (73)$$

The  $O(D)$ -invariant integral over  $S^d$  of the tensor  $H_{\bar{N}}^{i_1 \dots i_{\bar{N}}} := t^{i_1} \dots t^{i_{\bar{N}}}$  is equal to

$$\int_{S^d} d\alpha t^{i_1} \dots t^{i_{\bar{N}}} = C_{\bar{N}} G_{\bar{N}}^{i_1 \dots i_{\bar{N}}}, \quad C_{\bar{N}} = \frac{mis(S^d)}{\bar{N}!! D(D+2)\dots(D+\bar{N}-2)}. \quad (74)$$

In terms of the decompositions (69b)-(70) the scalar product of generic  $\phi, \psi \in \mathcal{H}_s$  is equal to

$$\langle \phi, \psi \rangle = \sum_{l=0}^{\infty} Q_l (\phi_{i_1 \dots i_l}^l)^* \psi_{i_1 \dots i_l}^l, \quad Q_l := C_{2l} 2^l (l!)^2 = \frac{mis(S^d) l! (D-2)!!}{(D+2l-2)!!}. \quad (75)$$

In particular, (75) implies  $\langle T_l^{i_1 \dots i_l}, T_n^{j_1 \dots j_n} \rangle = \delta_{ln} Q_l \mathcal{P}_{i_1 \dots i_l}^{l j_1 \dots j_l}$  and more generally

$$\langle T_l^{i_1 \dots i_l}, \phi \rangle = Q_l \mathcal{P}_{i_1 \dots i_l}^{l j_1 \dots j_l} \phi_{j_1 \dots j_l}^l \stackrel{(70)}{=} Q_l \phi_{i_1 \dots i_l}^l, \quad (76)$$

the second equality holds only if (70) holds. Then, we have also

$$\|\phi\|^2 \equiv \langle \phi, \phi \rangle = \sum_{l=0}^{\infty} Q_l \sum_{i_1, \dots, i_l} |\phi_{i_1 \dots i_l}^l|^2. \quad (77)$$

We now determine the decomposition of  $T_l^{i_1 \dots i_l} T_m^{j_1 \dots j_m}$ .

**Theorem 3.7** The product  $T_l^{i_1 \dots i_l} T_m^{j_1 \dots j_m}$  decomposes as follows into  $V_D^n$  components:

$$T_l^{i_1 \dots i_l} T_m^{j_1 \dots j_m} = \sum_{n \in L^{lm}} S_{k_1 \dots k_n}^{i_1 \dots i_l, j_1 \dots j_m} T_n^{k_1 \dots k_n}, \quad (78)$$

where  $L^{lm} := \{|l-m|, |l-m|+2, \dots, l+m\}$  and, defining  $r := \frac{l+m-n}{2} \in \{0, 1, \dots, m\}$ ,

$$\begin{aligned} S_{k_1 \dots k_n}^{i_1 \dots i_l, j_1 \dots j_m} &= N_n^{lm} V_{k_1 \dots k_n}^{i_1 \dots i_l, j_1 \dots j_m}, & N_n^{lm} &= \frac{(D+2n-2)!! l! m!}{(D+2n+2r-2)!! (l-r)! (m-r)!} \\ V_{k_1 \dots k_n}^{i_1 \dots i_l, j_1 \dots j_m} &= \mathcal{P}_{a_1 \dots a_r c_1 \dots c_{l-r}}^{l i_1 \dots i_l} \mathcal{P}_{a_1 \dots a_r c_{l-r+1} \dots c_n}^{m j_1 \dots j_r j_{r+1} \dots j_m} \mathcal{P}_{c_1 \dots c_n}^{n k_1 \dots k_n}. \end{aligned} \quad (79)$$

The coefficients  $S_{k_1 \dots k_n}^{i_1 \dots i_l, j_1 \dots j_m}$  are the analogs of the Clebsch-Gordon coefficients, which appear in the decomposition of a product of two spherical harmonics into a combination of spherical harmonics, for  $D = 3$ . The first term of the sum (78) is  $T_{l+m}^{i_1 \dots i_l j_1 \dots j_m}$ . This is consistent with the first term in the iterated application of (66). If  $r = m = 1$ , since  $\mathcal{P}_{a_1}^{1 j_1} = \delta_{a_1}^{j_1}$ ,  $n = l-1$ , the result is consistent with the second term in (66):

$$S_{k_1 \dots k_{l-1}}^{i_1 \dots i_l, j_1} = \frac{l}{D+2l-2} \mathcal{P}_{j_1 k_1 \dots k_{l-1}}^{l i_1 \dots i_l}.$$

### 3.2 Embedding in $\mathbb{R}^{D+1}$ , isomorphism $\text{End}(Pol_D^\Lambda) \simeq \pi_{D+1}^\Lambda [Uso(D+1)]$

We naturally embed  $\mathbb{C}[\mathbb{R}^D] \hookrightarrow \mathbb{C}[\mathbb{R}^{\mathbf{D}}]$ , where  $\mathbf{D} \equiv D + 1$ . Henceforth we use real Cartesian coordinates  $(x^i)$  for  $\mathbb{R}^D$  and  $(x^I)$  for  $\mathbb{R}^{\mathbf{D}}$ ;  $h, i, j, k \in \{1, \dots, D\}$ ,  $H, I, J, K \in \{1, \dots, \mathbf{D}\}$ . We naturally embed  $O(D) \hookrightarrow SO(\mathbf{D})$  identifying  $O(D)$  as the subgroup of  $SO(\mathbf{D})$  which is the little group of the  $\mathbf{D}$ -th axis; its Lie algebra, isomorphic to  $so(D)$ , is generated by the  $L_{hk}$ . We shall add  $\mathbf{D}$  as a subscript to distinguish objects in this enlarged dimension from their counterparts in dimension  $D$ , e.g. the distance  $r_{\mathbf{D}}$  from the origin in  $\mathbb{R}^{\mathbf{D}}$ , from its counterpart  $r \equiv r_D$  in  $\mathbb{R}^D$ ,  $\mathcal{P}_{\mathbf{D}}^l$  from  $\mathcal{P}^l \equiv \mathcal{P}_D^l$ , and so on.

We look for the decomposition of each  $\check{V}_{\mathbf{D}}^\Lambda$  into irreps of such a  $Uso(D)$ . Clearly,  $\check{V}_{\mathbf{D}}^0 \simeq \mathbb{C} \simeq V_D^0$ , while  $\check{V}_{\mathbf{D}}^1 = \check{V}_{D,1}^0 \oplus \check{V}_{D,1}^1$ , where  $\check{V}_{D,1}^0 \simeq V_D^0$  is spanned by  $x^{\mathbf{D}}$ , and  $\check{V}_{D,1}^1 \simeq V_D^1$  is spanned by the  $x^i$ . The  $X_{\mathbf{D},2}^{IJ} = x^I x^J - r_{\mathbf{D}}^2 \delta^{IJ} / \mathbf{D}$  span  $\check{V}_{\mathbf{D}}^2$ ; the set of elements

$$X_{\mathbf{D},2}^{\mathbf{D}\mathbf{D}} = x^{\mathbf{D}} x^{\mathbf{D}} - \frac{r_{\mathbf{D}}^2}{\mathbf{D}}, \quad X_{\mathbf{D},2}^{i\mathbf{D}} = x^i x^{\mathbf{D}}, \quad X_2^{ij} = X_{\mathbf{D},2}^{ij} + \frac{\delta^{ij}}{D} X_{\mathbf{D},2}^{\mathbf{D}\mathbf{D}} \quad (80)$$

span carrier spaces  $\check{V}_{D,2}^0, \check{V}_{D,2}^1, \check{V}_{D,2}^2$  respectively isomorphic to  $V_D^0, V_D^1, V_D^2$ , and  $\check{V}_{\mathbf{D}}^2 = \check{V}_{D,2}^2 \oplus \check{V}_{D,2}^1 \oplus \check{V}_{D,2}^0$ . More generally,  $\check{V}_{\mathbf{D}}^\Lambda$  is spanned by the  $X_{\mathbf{D},\Lambda}^{I_1 \dots I_\Lambda}$ , which are homogeneous polynomials of degree  $\Lambda$  in the  $x^I$  obtained by (57) in dimension  $\mathbf{D}$ , i.e.

$$X_{\mathbf{D},l}^{I_1 \dots I_l} := \mathcal{P}_{\mathbf{D},J_1 \dots J_l}^{I_1 \dots I_l} x^{J_1} \dots x^{J_l} \in \check{V}_{\mathbf{D}}^l \quad (81)$$

where the projectors  $\mathcal{P}_{\mathbf{D}}^l$  are constructed as in Proposition 3.2, but with  $D$  replaced by  $\mathbf{D}$ . Any pair of indices  $I_a, I_b$  appears either through the product  $x^{I_a} x^{I_b}$  or through  $r_{\mathbf{D}}^2 \delta^{I_a I_b}$ . If we introduce  $r_{\mathbf{D}}$  as a further generator constrained by the relation  $r_{\mathbf{D}}^2 = x^I x^I$  (sum over  $I$ ), then the  $X_{\mathbf{D},\Lambda}^{I_1 \dots I_\Lambda}$  can be seen also as homogeneous polynomials of degree  $\Lambda$  in  $x^I, r_{\mathbf{D}}$ . Since  $\delta^{j_a \mathbf{D}} = 0$ , in  $X_{\mathbf{D},\Lambda}^{j_1 \dots j_l \mathbf{D} \dots \mathbf{D}}$ : any pair of indices  $\mathbf{D}\mathbf{D}$  appears either through the product  $x^{\mathbf{D}} x^{\mathbf{D}}$  or through  $r_{\mathbf{D}}^2 \delta^{\mathbf{D}\mathbf{D}} = r_{\mathbf{D}}^2$ ; any pair of indices  $j_a, \mathbf{D}$  appears through the product  $x^{j_a} x^{\mathbf{D}}$ ; any pair of indices  $j_a, j_b$  appears either through the product  $x^{j_a} x^{j_b}$  or through  $r_{\mathbf{D}}^2 \delta^{j_a j_b}$ . By property (51), the latter terms completely disappear in any combination  $\mathcal{P}_{j_1 \dots j_l}^{i_1 \dots i_l} X_{\mathbf{D},\Lambda}^{j_1 \dots j_l \mathbf{D} \dots \mathbf{D}}$ ,  $l \in \{2, 3, \dots, \Lambda\}$ . Therefore such a combination can be factorized as follows

$$\check{F}_{\mathbf{D},\Lambda}^{i_1 \dots i_l} := \mathcal{P}_{j_1 \dots j_l}^{i_1 \dots i_l} X_{\mathbf{D},\Lambda}^{j_1 \dots j_l \mathbf{D} \dots \mathbf{D}} = \check{p}_{\Lambda,l} X_l^{i_1 \dots i_l}, \quad (82)$$

where  $\check{p}_{\Lambda,l}$  is a homogeneous polynomial of degree  $\Lambda - l$  in  $x^{\mathbf{D}}, r_{\mathbf{D}}$  of the form

$$\check{p}_{\Lambda,l} = (x^{\mathbf{D}})^{\Lambda-l} + (x^{\mathbf{D}})^{\Lambda-l-2} r_{\mathbf{D}}^2 b_{\Lambda,l+2} + (x^{\mathbf{D}})^{\Lambda-l-4} r_{\mathbf{D}}^4 b_{\Lambda,l+4} + \dots ; \quad (83)$$

the coefficients  $b_{\Lambda,h}$  can be determined from  $\Delta_{\mathbf{D}} \check{F}_{\mathbf{D},\Lambda}^{i_1 \dots i_l} = 0$ , which follows from  $\Delta_{\mathbf{D}} X_{\mathbf{D},\Lambda}^{I_1 \dots I_\Lambda} = 0$ .

**Proposition 3.8**  $\check{V}_{\mathbf{D}}^\Lambda$  decomposes into the following irreducible components of  $Uso(D)$ :

$$\check{V}_{\mathbf{D}}^\Lambda = \bigoplus_{l=0}^{\Lambda} \check{V}_{D,\Lambda}^l, \quad (84)$$



where  $\check{V}_{D,\Lambda}^l \simeq V_D^l$  is spanned by the  $\check{F}_{\mathbf{D},\Lambda}^{i_1 \dots i_l}$ , since the latter are eigenvectors of  $\mathbf{L}^2$ :

$$\mathbf{L}^2 \check{F}_{\mathbf{D},\Lambda}^{i_1 \dots i_l} = E_l \check{F}_{\mathbf{D},\Lambda}^{i_1 \dots i_l}. \quad (85)$$

Denoting by  $[a]$  the integral part of  $a \in \mathbb{R}^+$ , the coefficients of (83) are given by

$$b_{\Lambda, l+2k} = (-)^k \frac{(\Lambda-l)! (2\Lambda-4-2k+\mathbf{D})!!}{(\Lambda-l-2k)! (2k)!! (2\Lambda-4+\mathbf{D})!!}, \quad k = 1, 2, \dots \left[ \frac{\Lambda-l}{2} \right]. \quad (86)$$

The  $\check{F}_{\mathbf{D},\Lambda}^{i_1 \dots i_l}$  transform under  $L_{hk}$  as the  $T_l^{i_1 \dots i_l}$ , and under  $L_{h\mathbf{D}}$  as follows:

$$iL_{h\mathbf{D}} \check{F}_{\mathbf{D},\Lambda}^{i_1 \dots i_l} = (\Lambda-l) \check{F}_{\mathbf{D},\Lambda}^{hi_1 \dots i_l} - \frac{l(\Lambda+l+D-2)}{D+2l-2} \mathcal{P}_{hj_2 \dots j_l}^{li_1 i_2 \dots i_l} \check{F}_{\mathbf{D},\Lambda}^{j_2 \dots j_l}. \quad (87)$$

The proof is in Appendix 8.5. We now determine the decomposition of  $V_{\mathbf{D}}^\Lambda := \check{V}_{\mathbf{D}}^\Lambda / (r_{\mathbf{D}})^\Lambda$  into irreps of  $Uso(D)$ .  $V_{\mathbf{D}}^0 \simeq V_D^0 \simeq \mathbb{C}$  is spanned by 1.  $V_{\mathbf{D}}^1 = V_{D,1}^0 \oplus V_{D,1}^1$ , where  $V_{D,1}^0 \simeq V_D^0$  is spanned by  $t^{\mathbf{D}}$ , and  $V_{D,1}^1 \simeq V_D^1$  is spanned by the  $t^i$  (here  $t^I := x^I / r_{\mathbf{D}}$ ).  $V_{\mathbf{D}}^2$  is spanned by the  $T_{\mathbf{D},2}^{IJ} = t^I t^J - \frac{\delta^{IJ}}{\mathbf{D}}$ ; more explicitly,

$$T_{\mathbf{D},2}^{\mathbf{D}\mathbf{D}} = t^{\mathbf{D}} t^{\mathbf{D}} - \frac{1}{\mathbf{D}} = \frac{D}{\mathbf{D}} - t^h t_h, \quad T_{\mathbf{D},2}^{i\mathbf{D}} = t^i t^{\mathbf{D}}, \quad T_{\mathbf{D},2}^{ij} = t^i t^j - \delta^{ij} \frac{1}{\mathbf{D}};$$

$F_{\mathbf{D},2} := T_{\mathbf{D},2}^{\mathbf{D}\mathbf{D}}$  spans an irrep  $V_{D,2}^0$  isomorphic to  $V_D^0$ , the  $F_{\mathbf{D},2}^i := T_{\mathbf{D},2}^{i\mathbf{D}}$  span an irrep  $V_{D,2}^1$  isomorphic to  $V_D^1$ , while the  $F_{D,2}^{ij} := \mathcal{P}_{hk}^{ij} T_{\mathbf{D},2}^{hk}$  (which fulfill  $F_{D,2}^{ij} \delta_{ij} = 0$ ) span an irrep  $V_{D,2}^2$  isomorphic to  $V_D^2$ . The  $F_{D,2}^{ij}$  can be expressed as combinations of the  $T_{\mathbf{D},2}^{IJ}$ :

$$F_{D,2}^{ij} = T_{\mathbf{D},2}^{ij} + \frac{\delta^{ij}}{D} T_{\mathbf{D},2}^{\mathbf{D}\mathbf{D}} = T_{\mathbf{D},2}^{ij} - \frac{\delta^{ij}}{D} T_{\mathbf{D},2}^{hh} \text{ (sum over } h) = t^i t^j - \frac{\delta^{ij}}{D} t^h t_h$$

This shows the decomposition  $V_{\mathbf{D}}^2 = V_{D,2}^2 \oplus V_{D,2}^1 \oplus V_{D,2}^0 \simeq Pol_D^2$  explicitly. More generally, let

$$F_{\mathbf{D},\Lambda}^{i_1 \dots i_l} := \check{F}_{\mathbf{D},\Lambda}^{i_1 \dots i_l} / (r_{\mathbf{D}})^\Lambda = \mathcal{P}_{j_1 \dots j_l}^{i_1 \dots i_l} T_{\mathbf{D},\Lambda}^{j_1 \dots j_l \mathbf{D} \dots \mathbf{D}} = p_{\Lambda,l} \mathcal{P}_{j_1 \dots j_l}^{i_1 \dots i_l} t^{j_1} \dots t^{j_l} =: p_{\Lambda,l} T_l^{i_1 \dots i_l}, \quad (88)$$

where we have introduced a polynomial  $p_{\Lambda,l}$  of degree  $h = \Lambda - l$  in  $t^{\mathbf{D}}$  (containing only powers of the same parity as  $h$ ) by  $p_{\Lambda,l} := \check{p}_{\Lambda,l}(x^{\mathbf{D}}, r_{\mathbf{D}}) r_{\mathbf{D}}^{l-\Lambda}$ ; more explicitly the latter reads

$$p_{\Lambda,l} = (t^{\mathbf{D}})^{\Lambda-l} + (t^{\mathbf{D}})^{\Lambda-l-2} b_{\Lambda,l+2} + (t^{\mathbf{D}})^{\Lambda-l-4} b_{\Lambda,l+4} + \dots \quad (89)$$

As a direct consequence of Propositions 3.8, dividing all relations by  $(r_{\mathbf{D}})^\Lambda$ , we find

**Corollary 3.9**  $V_{\mathbf{D}}^\Lambda$  decomposes into the following irreducible components of  $Uso(D)$ :

$$V_{\mathbf{D}}^\Lambda = \bigoplus_{l=0}^{\Lambda} V_{D,\Lambda}^l, \quad (90)$$

where  $V_{D,\Lambda}^l \simeq V_D^l$  is spanned by the  $F_{\mathbf{D},\Lambda}^{i_1 \dots i_l}$ . The latter are eigenvectors of  $\mathbf{L}^2$ ,

$$\mathbf{L}^2 F_{\mathbf{D},\Lambda}^{i_1 \dots i_l} = E_l F_{\mathbf{D},\Lambda}^{i_1 \dots i_l}, \quad (91)$$

transform under  $L_{hk}$  as the  $T_l^{i_1 \dots i_l}$ , and under  $L_{h\mathbf{D}}$  as follows:

$$iL_{h\mathbf{D}} F_{\mathbf{D},\Lambda}^{i_1 \dots i_l} = (\Lambda - l) F_{\mathbf{D},\Lambda}^{hi_1 \dots i_l} - \frac{l(\Lambda + l + D - 2)}{D + 2l - 2} \mathcal{P}_{hj_2 \dots j_l}^{i_1 i_2 \dots i_l} F_{\mathbf{D},\Lambda}^{j_2 \dots j_l}. \quad (92)$$

For convenience, we slightly enlarge  $Uso(D)$  by introducing the new generator

$$\lambda = \frac{1}{2} \left[ \sqrt{(D-2)^2 + 4\mathbf{L}^2} - D + 2 \right], \quad (93)$$

which fulfills  $\lambda(\lambda + D - 2) = \mathbf{L}^2$ , so that  $V_D^l$  is a  $\lambda = l$  eigenspace, and  $\lambda F_{\mathbf{D},\Lambda}^{i_1 \dots i_l} = l F_{\mathbf{D},\Lambda}^{i_1 \dots i_l}$ .

**Proposition 3.10** *There exist a  $Uso(D)$ -module isomorphism  $\hat{\chi}_\Lambda : Pol_D^\Lambda \rightarrow V_D^\Lambda$  and a  $Uso(D)$ -equivariant algebra isomorphism  $\hat{\kappa}_\Lambda : End(Pol_D^\Lambda) \rightarrow \pi_D^\Lambda [Uso(\mathbf{D})]$  such that*

$$\hat{\chi}_\Lambda(aT) = \hat{\kappa}_\Lambda(a)\hat{\chi}_\Lambda(T), \quad \forall T \in Pol_D^\Lambda, \quad a \in End(Pol_D^\Lambda). \quad (94)$$

On the  $T_l^{i_1 \dots i_l}$  (spanning  $Pol_D^\Lambda$ ) and on generators  $L_{hi}, P^\Lambda t^i$  of  $End(Pol_D^\Lambda)$  they act by

$$\hat{\chi}_\Lambda(T_l^{i_1 \dots i_l}) := \hat{a}_{\Lambda,l} F_{\mathbf{D},\Lambda}^{i_1 \dots i_l} = \hat{a}_{\Lambda,l} p_{\Lambda,l} T_l^{i_1 \dots i_l}, \quad l = 0, 1, \dots, \Lambda, \quad (95)$$

$$\hat{\kappa}_\Lambda(L_{hi}) := L_{hi}, \quad \hat{\kappa}_\Lambda(P^\Lambda t^i) := \hat{m}_\Lambda^*(\lambda) L_{\mathbf{D}i} \hat{m}_\Lambda(\lambda), \quad (96)$$

where

$$\hat{m}_\Lambda(l) = \sqrt{\frac{\Gamma(\frac{\Lambda+l+D-1}{2}) \Gamma(\frac{\Lambda-l+1}{2})}{2\Gamma(\frac{\Lambda+l+D}{2}) \Gamma(\frac{\Lambda-l}{2} + 1)}}, \quad \hat{a}_{\Lambda,l} = \hat{a}_{\Lambda,0} i^l \sqrt{\frac{\Lambda(\Lambda-1)\dots(\Lambda-l+1)}{(\Lambda+D-1)(\Lambda+D)\dots(\Lambda+l+D-2)}}. \quad (97)$$

The proposition and its proof are obtained from Theorem 5.1 and the associated proof by fixing  $\Lambda$ , taking  $k$  independent of  $\Lambda$  and letting the  $k \rightarrow \infty$ .

## 4 Relations among the $\bar{x}^i, \bar{L}_{hk}$

Since for all fixed  $l = 0, 1, \dots, \Lambda$  the  $T_l^{i_1 i_2 \dots i_l}$ , make up a complete set  $\mathcal{T}_l$  in  $V_D^l$ , then the functions

$$\psi_l^{i_1 i_2 \dots i_l} := T_l^{i_1 i_2 \dots i_l} f_l \quad (98)$$

( $i_h \in \{1, \dots, D\}$  for  $h \in \{1, \dots, l\}$ ) make up a complete set  $\mathcal{S}_{D,\Lambda}^l$  in the eigenspace  $\mathcal{H}_{D,\Lambda}^l$  of  $H, \mathbf{L}^2$ , with eigenvalues  $E_{0,l}, E_l$ . They are not linearly independent, because they are completely symmetric under permutation of the indices and fulfill the relations

$$\delta_{i_n i_{n+1}} \psi_l^{i_1 \dots i_l} = 0, \quad n = 1, \dots, l-1. \quad (99)$$

$\mathcal{S}_{D,\Lambda} := \cup_{l=0}^{\Lambda} \mathcal{S}_{D,\Lambda}^l$  is a complete set in  $\mathcal{H}_{\Lambda}$ . By (98), (68) the  $\bar{L}_{hk}$  act on the  $\psi_l^{i_1 i_2 \dots i_l}$  via

$$i\bar{L}_{hk}\psi_l^{i_1 i_2 \dots i_l} = l\mathcal{P}_{j_1 \dots j_l}^{l i_1 \dots i_l} \left( \delta^{k j_1} \psi_l^{h j_2 \dots j_l} - \delta^{h j_1} \psi_l^{k j_2 \dots j_l} \right). \quad (100)$$

By (26), (66), applying  $P_{\Lambda} = \sum_{l=0}^{\Lambda} P_{\Lambda}^l$  to  $x^i \psi^{i_1 i_2 \dots i_l} = t^i T_l^{i_1 i_2 \dots i_l} r f_l(r)$  we find

$$P_{\Lambda} (x^i \psi_l^{i_1 i_2 \dots i_l}) = c_{l+1} \psi_{l+1}^{i i_1 \dots i_l} + c_l \frac{\zeta_{l+1}}{D} \mathcal{P}_{i j_2 \dots j_l}^{l i_1 i_2 \dots i_l} \psi_{l-1}^{j_2 \dots j_l},$$

$$\text{where } c_l := \begin{cases} \sqrt{1 + \frac{(2D-5)(D-1)}{2k} + \frac{(l-1)(l+D-2)}{k}} & \text{if } 1 \leq l \leq \Lambda, \\ 0 & \text{otherwise,} \end{cases} \quad (101)$$

up to order  $O(k^{-3/2})$ , see appendix 8.6. Hence, at the same order,

$$\bar{x}^i \psi_l^{i_1 i_2 \dots i_l} = c_{l+1} \psi_{l+1}^{i i_1 \dots i_l} + \frac{c_l l}{D+2l-2} \mathcal{P}_{i j_2 \dots j_l}^{l i_1 i_2 \dots i_l} \psi_{l-1}^{j_2 \dots j_l}. \quad (102)$$

The  $O(k^{-3/2})$  corrections depend on the terms proportional to  $(r-1)^k$ ,  $k > 2$ , in the Taylor expansion of  $V$ . These could be set rigorously equal to zero by a suitable choice of  $V$ . Henceforth we adopt (100-102) as exact *definitions* of  $\bar{L}_{hk}, \bar{x}^i$ . In the appendix we prove

**Proposition 4.1** *The  $\bar{x}^i, \bar{L}_{hk}$  are self-adjoint operators generating the  $N^2$ -dimensional  $*$ -algebra  $\mathcal{A}_{\Lambda} := \text{End}(\mathcal{H}_{\Lambda}) \simeq M_N(\mathbb{C})$  of observables on  $\mathcal{H}_{\Lambda}$ ;  $N$  is given by (37). Abbreviating  $\bar{\mathbf{x}}^2 := \bar{x}^i \bar{x}^i$ ,  $\bar{\mathbf{L}}^2 := \bar{L}_{ij} \bar{L}_{ij}/2$ ,  $B := (2D-5)(D-1)/2$ , they fulfill, at leading order in  $1/\sqrt{k}$*

$$[i\bar{L}_{ij}, \bar{x}^h] = \bar{x}^i \delta_j^h - \bar{x}^j \delta_i^h, \quad (103)$$

$$[i\bar{L}_{ij}, i\bar{L}_{hk}] = i(\bar{L}_{ik} \delta_h^j - \bar{L}_{jk} \delta_h^i - \bar{L}_{ih} \delta_k^j + \bar{L}_{jh} \delta_k^i), \quad (104)$$

$$\varepsilon^{i_1 i_2 i_3 \dots i_D} \bar{x}^{i_1} \bar{L}_{i_2 i_3} = 0, \quad (105)$$

$$(\bar{x}^h \pm i\bar{x}^k)^{2\Lambda+1} = 0, \quad (\bar{L}^{hj} + i\bar{L}^{kj})^{2\Lambda+1} = 0, \quad \text{if } h \neq j \neq k \neq h, \quad (106)$$

$$[\bar{x}^i, \bar{x}^j] = \left( -\frac{I}{k} + K P_{\Lambda}^{\Lambda} \right) i\bar{L}_{ij}, \quad K := \frac{1}{k} + \frac{1}{D+2\Lambda-2} \left[ 1 + \frac{B}{k} + \frac{(\Lambda-1)(\Lambda+D-2)}{k} \right], \quad (107)$$

$$\bar{\mathbf{x}}^2 = 1 + \frac{\bar{\mathbf{L}}^2}{k} + \frac{B}{k} - \frac{\Lambda+D-2}{2\Lambda+D-2} \left[ 1 + \frac{B}{k} + \frac{\Lambda(\Lambda+D-1)}{k} \right] P_{\Lambda}^{\Lambda} =: \chi(\bar{\mathbf{L}}^2). \quad (108)$$

This is the analog of Proposition 4.1 in [22]. We obtain a fuzzy sphere choosing  $k$  as a function  $k(\Lambda)$  fulfilling (19), e.g.  $k = \Lambda^2(\Lambda+D-2)^2/4$ ; the commutative limit is  $\Lambda \rightarrow \infty$ . We remark that:

- By (107), (105) can be reformulated also in the form  $\varepsilon^{i_1 i_2 i_3 \dots i_D} \bar{x}^{i_1} \bar{x}^{i_2} \bar{x}^{i_3} = 0$ .
- $\bar{\mathbf{x}}^2$  is not a constant, but by (108), (22) $_{l=\Lambda}$  can be expressed as a polynomial  $\chi$  in  $\bar{\mathbf{L}}^2$  only, with the same eigenspaces  $\mathcal{H}_{\Lambda}^l$ . All its eigenvalues  $r_l^2$ , except  $r_{\Lambda}^2$ , are close to 1, slightly grow with  $l$  and collapse to 1 as  $\Lambda \rightarrow \infty$ . Conversely,  $\bar{\mathbf{L}}^2$  can be expressed as a polynomial  $v$  in  $\bar{\mathbf{x}}^2$ , via  $\bar{\mathbf{L}}^2 = \sum_{l=0}^{\Lambda} E_l P_{\Lambda}^l$  and  $P_{\Lambda}^l = \prod_{n=0, n \neq l}^{\Lambda} \frac{\bar{\mathbf{x}}^2 - r_n^2}{r_l^2 - r_n^2}$ .

- By (107),  $(22)_{l=\Lambda}$  the commutators  $[\bar{x}^i, \bar{x}^j]$  are Snyder-like, i.e. of the form  $\alpha \bar{L}_{ij}$ ; also  $\alpha$  depends only on the  $\bar{L}_{hk}$ , more precisely can be expressed as a polynomial in  $\bar{\mathbf{L}}^2$ .
- Using (103), (104), (107), all polynomials in  $\bar{x}^i, \bar{L}_{hk}$  can be expressed as combinations of monomials in  $\bar{x}^i, \bar{L}_{hk}$  in any prescribed order; the coefficients are complex combinations of 1 and  $P_\Lambda^\Lambda$ , which also can be expressed as a polynomial in  $\bar{\mathbf{L}}^2$  via  $(22)_{l=\Lambda}$ . Hence such ordered monomials make up a basis of the  $N^2$ -dim vector space  $\mathcal{A}_\Lambda$ .
- Actually,  $\bar{x}^i$  generate the  $*$ -algebra  $\mathcal{A}_\Lambda$ , because also the  $\bar{L}_{ij}$  can be expressed as non-ordered polynomials in the  $\bar{x}^i$ : by (107)  $\bar{L}_{ij} = [\bar{x}^j, \bar{x}^i]/\alpha$ , and also  $1/\alpha$ , which depends only on  $P_\Lambda^\Lambda$ , can be expressed itself as a polynomial in  $\bar{\mathbf{x}}^2$ , as shown above.
- Eq. (103-108) are equivariant under the whole group  $O(D)$ , including the inversion  $\bar{x}^i \mapsto -\bar{x}^i$  of one axis, or more (e.g. parity), contrary to Madore's and Hoppe's FS.

The operator norm of  $\bar{\mathbf{x}}^2$  equals its highest eigenvalue,  $\|\bar{\mathbf{x}}^2\|_{op} = r_{\Lambda-1}^2 = 1 + [B + (\Lambda-1)(\Lambda + D-3)]/k$ . For all  $\psi \in \mathcal{H}_\Lambda$  and  $i = 1, \dots, D$ , we find  $\|\bar{x}^i \psi\|^2 = \langle \bar{x}^i \psi, \bar{x}^i \psi \rangle \leq \sum_{i=1}^D \langle \bar{x}^i \psi, \bar{x}^i \psi \rangle = \langle \psi, \bar{\mathbf{x}}^2 \psi \rangle \leq \|\bar{\mathbf{x}}^2\|_{op} \|\psi\|^2$ , whence

$$\|\bar{x}^i\|_{op} \leq \sqrt{1 + \frac{B + (\Lambda-1)(\Lambda + D-3)}{k}} \leq 1 + \epsilon, \quad \epsilon := \frac{B + (\Lambda-1)(\Lambda + D-3)}{2k} \quad (109)$$

## 5 Isomorphisms of $\mathcal{H}_\Lambda, \mathcal{A}_\Lambda$ , and $*$ -automorphisms of $\mathcal{A}_\Lambda$

**Theorem 5.1** *There exist a  $O(D)$ -module isomorphism  $\varkappa_\Lambda : \mathcal{H}_\Lambda \rightarrow V_{\mathbf{D}}^\Lambda$  and a  $O(D)$ -equivariant algebra map  $\kappa_\Lambda : \mathcal{A}_\Lambda \equiv \text{End}(\mathcal{H}_\Lambda) \rightarrow \pi_{\mathbf{D}}^\Lambda[\text{Uso}(\mathbf{D})]$ ,  $\mathbf{D} \equiv D+1$ , such that*

$$\varkappa_\Lambda(a\psi) = \kappa_\Lambda(a)\varkappa_\Lambda(\psi), \quad \forall \psi \in \mathcal{H}_\Lambda, \quad a \in \mathcal{A}_\Lambda. \quad (110)$$

On the  $\psi_l^{i_1 \dots i_l}$  (spanning  $\mathcal{H}_\Lambda$ ) and on generators  $L_{hi}, \bar{x}^i$  of  $\mathcal{A}_\Lambda$  they respectively act as follows:

$$\varkappa_\Lambda(\psi_l^{i_1 \dots i_l}) := a_{\Lambda, l} F_{\mathbf{D}, \Lambda}^{i_1 \dots i_l} = a_{\Lambda, l} p_{\Lambda, l} T_l^{i_1 \dots i_l}, \quad l = 0, 1, \dots, \Lambda, \quad (111)$$

$$\kappa_\Lambda(\bar{L}_{hi}) := \pi_{\mathbf{D}}^\Lambda(L_{hi}), \quad \kappa_\Lambda(\bar{x}^i) := \pi_{\mathbf{D}}^\Lambda[m_\Lambda^*(\lambda) X^i m_\Lambda(\lambda)], \quad (112)$$

where  $X^i := L_{\mathbf{D}i}$ ,  $p_{\Lambda, l} = p_{\Lambda, l}(t^{\mathbf{D}})$  are the polynomials (89), and

$$a_{\Lambda, l} = a_{\Lambda, 0} i^l \sqrt{\frac{\Lambda(\Lambda-1)\dots(\Lambda-l+1)}{(\Lambda+D-1)(\Lambda+D)\dots(\Lambda+l+D-2)}}, \quad (113)$$

$$m_\Lambda(s) = \sqrt{\frac{\Gamma\left(\frac{\Lambda+s+d}{2}\right) \Gamma\left(\frac{\Lambda-s+1}{2}\right) \Gamma\left(\frac{s+1+d/2+iA}{2}\right) \Gamma\left(\frac{s+1+d/2-iA}{2}\right)}{\Gamma\left(\frac{\Lambda+s+D}{2}\right) \Gamma\left(\frac{\Lambda-s}{2} + 1\right) \Gamma\left(\frac{s+d/2+iA}{2}\right) \Gamma\left(\frac{s+d/2-iA}{2}\right) \sqrt{k}}}; \quad (114)$$

here  $A := \sqrt{k + (D-1)(D-3)3/4}$ , and  $\Gamma$  is Euler gamma function.

The proof is in appendix 8.8. The theorem extends Propositions 3.2, 4.2 of [22] to  $d > 2$ . The claims for  $d > 2$  were partially formulated, but not proved, in [23].

As already recalled, the group of  $*$ -automorphisms of  $\mathcal{A}_\Lambda \simeq M_N(\mathbb{C})$  is inner and isomorphic to  $SU(N)$ , i.e. of the type

$$a \mapsto g a g^{-1}, \quad a \in M_N(\mathbb{C}), \quad (115)$$

with  $g$  an unitary  $N \times N$  matrix with unit determinant. We can identify in  $SU(N)$  a subgroup  $\simeq SO(\mathbf{D})$ , acting via the  $N$ -dimensional representation  $(V_{\mathbf{D}}^\Lambda, \pi_{\mathbf{D}}^\Lambda)$ ; namely, it consists of matrices of the form  $g = \pi_{\mathbf{D}}^\Lambda [e^{i\alpha}]$ , where  $\alpha \in so(\mathbf{D})$ . Choosing  $\alpha \in so(D) \subset so(\mathbf{D})$  the automorphism amounts to a  $SO(D) \subset SO(\mathbf{D})$  transformation, i.e. a rotation in the  $x \equiv (x^1, \dots, x^D) \in \mathbb{R}^D$  space.  $O(D) \subset SO(\mathbf{D})$  transformations with determinant  $-1$  in this space keep the same form in the  $\bar{X} \equiv (X^1, \dots, X^D)$  space (where  $X^i \equiv L_{\mathbf{D}i}$ ) and by (112) also in the  $\bar{x} \equiv (\bar{x}^1, \dots, \bar{x}^D)$  space. In particular, those inverting one or more axes of  $\mathbb{R}^D$  (i.e. changing the sign of one or more  $x^i$ , and thus also of  $X^i, \bar{x}^i$ ), e.g. parity, can be also realized as  $SO(\mathbf{D})$  transformations, i.e. rotations in  $\mathbb{R}^D$ . This shows that (112) is equivariant under parity and the whole  $O(D)$ , which plays the role of isometry group of this fuzzy sphere.

## 6 Fuzzy spherical harmonics, and limit $\Lambda \rightarrow \infty$

In this section we suppress Einstein's summation convention over repeated indices. The previous results allow to define  $Uso(D)$ -module Hilbert space isomorphisms

$$\begin{aligned} \sigma_\Lambda : Pol_D^\Lambda = \bigoplus_{l=0}^\Lambda V_D^l &\rightarrow \mathcal{H}_\Lambda = \bigoplus_{l=0}^\Lambda \mathcal{H}_\Lambda^l, \\ \Psi &\Psi \\ T_l^{i_1 \dots i_l}, Y_l^m &\mapsto \psi_l^{i_1 \dots i_l}, \psi_l^m; \end{aligned} \quad (116)$$

here  $\psi_l^m := \sum A_{i_1 \dots i_l}^m \psi_l^{i_1 \dots i_l}$ . The objects at the right of the arrows are our fuzzy analogs of the objects at the left. In the limit  $\Lambda \rightarrow \infty$  the above decomposition of  $\mathcal{H}_\Lambda$  into irreducible components under  $O(D)$  becomes isomorphic to the decomposition of  $Pol_D \simeq \mathcal{H}_s$ .

We define the  $O(D)$ -equivariant embedding  $\mathcal{I} : \mathcal{H}_\Lambda \hookrightarrow \mathcal{H}_s$  by setting  $\mathcal{I}(\psi_l^{i_1 \dots i_l}) := T_l^{i_1 \dots i_l}$  and applying linear extension. Below we drop  $\mathcal{I}$  and identify  $\psi_l^{i_1 \dots i_l} = T_l^{i_1 \dots i_l}$ , or equivalently  $\psi_l^m = Y_l^m$ , as elements of the Hilbert space  $\mathcal{H}_s$ . For all  $\phi \equiv \sum_{l=0}^\infty \phi_{i_1 \dots i_l}^l T_l^{i_1 \dots i_l} \in \mathcal{L}^2(S^2)$  and  $\Lambda \in \mathbb{N}$  let  $\phi_\Lambda := P_\Lambda \phi = \sum_{l=0}^\Lambda \phi_{i_1 \dots i_l}^l T_l^{i_1 \dots i_l}$  be its projection to  $\mathcal{H}_\Lambda$  (or  $\Lambda$ -th truncation). Clearly  $\phi_\Lambda \rightarrow \phi$  in the  $\mathcal{H}_s$ -norm  $\|\cdot\|$ : in this simplified notation,  $\mathcal{H}_\Lambda$  'invades'  $\mathcal{H}_s$  as  $\Lambda \rightarrow \infty$ ;

$\mathcal{I}$  induces the  $O(D)$ -equivariant embedding of operator algebras  $\mathcal{J} : \mathcal{A}_\Lambda \hookrightarrow B(\mathcal{H}_s)$  by setting  $\mathcal{J}(a)\mathcal{I}(\psi) := \mathcal{I}(a\psi)$ ; here  $B(\mathcal{H}_s)$  stands for the  $*$ -algebra of bounded operators on  $\mathcal{H}_s$ . By construction  $\mathcal{A}_\Lambda$  annihilates  $\mathcal{H}_\Lambda^\perp$ . In particular,  $\mathcal{J}(\bar{L}_{hk}) = L_{hk} P^\Lambda$ , and  $\bar{L}_{hk} \phi \xrightarrow{\Lambda \rightarrow \infty} L_{hk} \phi$  for all  $\phi \in D(L_{hk}) \equiv$  the domain of  $L_{hk}$ . More generally,  $f(\bar{L}_{hk}) \rightarrow f(L_{hk})$  strongly on  $D[f(L_{hk})] \subset \mathcal{H}_s$ , for all measurable functions  $f(s)$ .

Continuous functions  $f$  on  $S^d$ , acting as multiplication operators  $f \cdot : \phi \in \mathcal{H}_s \mapsto f\phi \in \mathcal{H}_s$ , make up a subalgebra  $C(S^d)$  of  $B(\mathcal{H}_s)$ . Clearly,  $f$  belongs also to  $\mathcal{H}_s$ . Since  $Pol_D$  is dense in both  $\mathcal{H}_s$ ,  $C(S^d)$ ,  $f_N$  converges to  $f$  as  $N \rightarrow \infty$  in both the  $\mathcal{H}_s$  and the  $C(S^d)$  norm.

We define the  $\Lambda$ -th fuzzy analogs of the  $T_l^{i_1 \dots i_l}$  (seen as an element of  $C(S^d)$ , i.e. acting by multiplication on  $\psi \in \mathcal{H}_s$ ),  $l \leq \Lambda$ , by replacing  $t^i \mapsto \bar{x}^i$  in the definition of  $T_l^{i_1 \dots i_l}$ , i.e. by

$$\widehat{T}_l^{i_1 \dots i_l} := \mathcal{P}_{j_1 \dots j_l}^{l i_1 \dots i_l} \bar{x}^{j_1} \dots \bar{x}^{j_l}, \quad (117)$$

sum over repeated indices (cf. Proposition 3.5); the  $\widehat{T}_l^{i_1 \dots i_l}$  fulfill again (65). Having identified  $\psi_l^{i_1 \dots i_l} \equiv T_l^{i_1 \dots i_l}$ , we rewrite (66), (102) in the form

$$t^h T_l^{i_1 \dots i_l} = T_{l+1}^{h i_1 \dots i_l} + d_l \mathcal{P}_{h j_2 \dots j_l}^{l i_1 i_2 \dots i_l} T_{l-1}^{j_2 \dots j_l}, \quad d_l := \frac{l}{D+2l-2} \quad (118)$$

$$\bar{x}^h T_l^{i_1 i_2 \dots i_l} = c_{l+1} T_{l+1}^{h i_1 \dots i_l} + c_l d_l \mathcal{P}_{h j_2 \dots j_l}^{l i_1 i_2 \dots i_l} T_{l-1}^{j_2 \dots j_l}. \quad (119)$$

Using these formulae in appendix 8.9 we prove the fuzzy version of Proposition 3.7:

**Theorem 6.1** *The action of the  $\widehat{T}_l^{i_1 \dots i_l} = \mathcal{P}_{j_1 \dots j_l}^{l i_1 \dots i_l} \bar{x}^{j_1} \dots \bar{x}^{j_l}$  on  $\mathcal{H}_\Lambda$  is given by*

$$\widehat{T}_l^{i_1 \dots i_l} T_m^{j_1 \dots j_m} = \sum_{n \in L} \widehat{N}_n^{lm} \mathcal{P}_{a_1 \dots a_r c_1 \dots c_{l-r}}^{l i_1 \dots i_l} \mathcal{P}_{a_1 \dots a_r c_{l-r+1} \dots c_n}^{m j_1 \dots j_r j_{r+1} \dots j_m} \mathcal{P}_{c_1 \dots c_n}^{n k_1 \dots k_n} T_n^{k_1 \dots k_n}, \quad (120)$$

with suitable coefficients  $\widehat{N}_n^{lm}$  related to their classical limits  $N_n^{lm} > 0$  of formula (78) by

$$\widehat{N}_n^{lm} = 0 \quad \text{if } l-m > \Lambda, \quad N_n^{lm} \leq \widehat{N}_n^{lm} \leq N_n^{lm} (c_\Lambda)^l \quad \text{otherwise.} \quad (121)$$

As a fuzzy analog of the vector space  $C(S^d)$  we adopt

$$\mathcal{C}_\Lambda := \left\{ \widehat{f}_{2\Lambda} := \sum_{l=0}^{2\Lambda} \sum_{i_1, \dots, i_l} f_{i_1 \dots i_l}^l \widehat{T}_l^{i_1 \dots i_l}, f_{i_1 \dots i_l}^l \in \mathbb{C} \right\} \subset \mathcal{A}_\Lambda \subset B(\mathcal{H}_s); \quad (122)$$

here the highest  $l$  is  $2\Lambda$  because by (121) the  $\widehat{T}_l^{i_1 \dots i_l}$  annihilate  $\mathcal{H}_\Lambda$  if  $l > 2\Lambda$ . By construction,

$$\mathcal{C}_\Lambda = \bigoplus_{l=0}^{2\Lambda} \widehat{V}_D^l, \quad \widehat{V}_D^l := \left\{ \sum_{i_1, \dots, i_l} f_{i_1 \dots i_l}^l \widehat{T}_l^{i_1 \dots i_l}, f_{i_1 \dots i_l}^l \in \mathbb{C} \right\} \quad (123)$$

is the decomposition of  $\mathcal{C}_\Lambda$  into irreducible components under  $O(D)$ .  $\widehat{V}_D^l$  is trace-free for all  $l > 0$ . In the limit  $\Lambda \rightarrow \infty$  (123) becomes the decomposition of  $C(S^d)$ . As a fuzzy analog of  $f \in C(S^d)$  we adopt the sum  $\widehat{f}_{2\Lambda}$  appearing in (122) with the coefficients of the expansion (69) of  $f$  up to  $l = 2\Lambda$ . In appendix 6.2 we prove

**Theorem 6.2** *For all  $f, g \in C(S^d)$  the following strong limits as  $\Lambda \rightarrow \infty$  hold:  $\widehat{f}_{2\Lambda} \rightarrow f \cdot, (\widehat{fg})_{2\Lambda} \rightarrow fg$  and  $\widehat{f}_{2\Lambda} \widehat{g}_{2\Lambda} \rightarrow fg$ .*

The last statement says that the product in  $\mathcal{A}_\Lambda$  of the approximations  $\widehat{f}_{2\Lambda}, \widehat{g}_{2\Lambda}$  goes to the product in  $B(\mathcal{H}_s)$  (the algebra of bounded operators on  $\mathcal{H}_s \equiv \mathcal{L}^2(S^d)$ ) of  $f \cdot, g \cdot$ . We point out that  $\widehat{f}_{2\Lambda}$  does not converge to  $f$  in operator norm, because the operator  $\widehat{f}_{2\Lambda}$  (a polynomial in the  $\bar{x}^i$ ) annihilates  $\mathcal{H}_\Lambda^\perp$  (the orthogonal complement of  $\mathcal{H}_\Lambda^\perp$ ), since so do the  $\bar{x}^i = P^\Lambda x^i \cdot P^\Lambda$ .

Essentially the same claims of this theorem were proved for  $d = 1, 2$  in [22] requiring that  $k(\Lambda)$  diverges much faster than required by (19), and were formulated (without proof) for  $d > 2$  in Theorem 7.1 of [23] with the same strong assumptions on the divergence of  $k(\Lambda)$ .

## 7 Outlook, discussion and conclusions

In this paper we have completed our construction [22, 23] of fuzzy spheres  $S_\Lambda^d$  that be equivariant under the *full* orthogonal group  $O(D)$ ,  $D \equiv d+1$ , for *all*  $d \in \mathbb{N}$ . The construction procedure consists (sections 1, 2) in starting with a quantum particle in  $\mathbb{R}^D$  configuration space subject to a  $O(D)$ -invariant potential  $V(r)$  with a very sharp minimum on the sphere of radius  $r = 1$  and projecting the Hilbert space  $\mathcal{H} = \mathcal{L}^2(\mathbb{R}^D)$  to the subspace  $\mathcal{H}_{\bar{E}}$  with energy below a suitable cutoff  $\bar{E}$ ;  $\bar{E}$  is sufficiently low to exclude all excited radial modes of  $\mathcal{H}$  (this can be considered as a *quantum* version of the constraint  $r = 1$ ), so that on  $\mathcal{H}_{\bar{E}}$  the Hamiltonian essentially reduces to the square angular momentum  $\mathbf{L}^2$  (the Laplacian, i.e. the free Hamiltonian, over the sphere  $S^d$ ). By making both the confining parameter  $k \equiv V''(1)/4$  and  $\bar{E}$  depend on  $\Lambda \in \mathbb{N}$ , and diverge with it, we have obtained a sequence  $\{(\mathcal{H}_\Lambda, \mathcal{A}_\Lambda)\}_{\Lambda \in \mathbb{N}}$  of  $O(D)$ -equivariant approximations of a quantum particle on  $S^d$ .  $\mathcal{H}_\Lambda$  is the  $\Lambda$ -th projected Hilbert space of states and  $\mathcal{A}_\Lambda \equiv \text{End}(\mathcal{H}_\Lambda)$  is the associated  $*$ -algebra of observables. The projected Cartesian coordinates  $\bar{x}^i$  no longer commute (section 4); their commutators  $[\bar{x}^i, \bar{x}^j]$  are of Snyder type, i.e. proportional to the angular momentum components  $L_{ij}$ . Actually, the  $\bar{x}^i$  generate the whole  $\mathcal{A}_\Lambda$ . The square distance  $\bar{x}^2$  from the origin is not identically 1, but a function of  $\mathbf{L}^2$  such that its spectrum is very close to 1 and collapses to 1 as  $\Lambda \rightarrow \infty$ . We have also constructed (section 6) the subspace  $\mathcal{C}_\Lambda \subset \mathcal{A}_\Lambda$  of completely symmetrized trace-free polynomials in the  $\bar{x}^i$ ; this is also spanned by the fuzzy analogs of spherical harmonics (thought as multiplication operators on  $\mathcal{H}_s$ ).  $\mathcal{H}_\Lambda, \mathcal{C}_\Lambda, \mathcal{A}_\Lambda$  carry reducible representations of  $O(D)$ ; as  $\Lambda \rightarrow \infty$  their decompositions into irreps respectively go to the decompositions of  $\mathcal{H}_s \equiv \mathcal{L}^2(S^d)$ , of  $C(S^d)$  (the commutative algebra of continuous functions on  $S^d$  acting as multiplication operators on  $\mathcal{H}_s$ ) and of the algebra  $\mathcal{A}_s$  of observables on  $\mathcal{H}_s$  [see (5), (123)]. Moreover, there are natural embeddings  $\mathcal{H}_\Lambda \hookrightarrow \mathcal{H}_s$ ,  $\mathcal{C}_\Lambda \hookrightarrow C(S^d)$  and  $\mathcal{A}_\Lambda \hookrightarrow \mathcal{A}_s$  such that  $\mathcal{H}_\Lambda \rightarrow \mathcal{H}_s$  in the norm of  $\mathcal{H}_s$ , while  $\mathcal{C}_\Lambda \rightarrow C(S^d)$ ,  $\mathcal{A}_\Lambda \rightarrow \mathcal{A}_s$  strongly as  $\Lambda \rightarrow \infty$  (section 6).

The structure of the pairs  $(\mathcal{H}_\Lambda, \mathcal{A}_\Lambda)$  is made transparent by the discovery (section 5) that these are isomorphic to  $(V_{\mathbf{D}}^\Lambda, \pi_\Lambda[UsO(\mathbf{D})])$ ,  $\mathbf{D} \equiv D+1$ , also as  $O(D)$ -modules;  $\pi_\Lambda$  is the irrep of  $UsO(\mathbf{D})$  on the space  $V_{\mathbf{D}}^\Lambda$  of harmonic polynomials of degree  $\Lambda$  on  $\mathbb{R}^{\mathbf{D}}$ , restricted to  $S^{\mathbf{D}}$ .

We now show that this allows to see  $\{\mathcal{A}_\Lambda\}_{\Lambda \in \mathbb{N}}$  as a fuzzy quantization of a coadjoint orbit of  $O(\mathbf{D})$  that goes to the classical phase space  $T^*S^d$ . We recall that given a Lie group  $G$ , a coadjoint orbit  $\mathcal{O}_\lambda$ , for  $\lambda$  in the dual space  $\mathfrak{g}^*$  of the Lie algebra  $\mathfrak{g}$  of  $G$ , may be defined either extrinsically, as the actual orbit  $\text{Ad}_G^* \lambda$  of the coadjoint action  $\text{Ad}_G^*$  inside  $\mathfrak{g}^*$  passing through  $\lambda$ , or intrinsically as the homogeneous space  $G/G_\lambda$ , where  $G_\lambda$  is the stabilizer of  $\lambda$  with respect to the coadjoint action (this distinction is worth making since the embedding of the orbit may be complicated). Coadjoint orbits are naturally endowed with a symplectic structure arising from the group action. If  $G$  is compact semisimple, identifying  $\mathfrak{g}^*$  with  $\mathfrak{g}$  via the (nondegenerate) Killing form, we can resp. rewrite these definitions in the form

$$\mathcal{O}_\lambda := \{g\lambda g^{-1} \mid g \in G\} \subset \mathfrak{g}^*, \quad \mathcal{O}_\lambda := G/G_\lambda \quad \text{where } G_\lambda := \{g \in G \mid g\lambda g^{-1} = \lambda\}. \quad (124)$$

Clearly,  $G_{\Lambda\lambda} = G_\lambda$  for all  $\Lambda \in \mathbb{C} \setminus \{0\}$ . Denoting as  $\mathcal{H}_\lambda$  the (necessarily finite-dimensional) carrier space of the irrep with highest weight  $\lambda$ , one can regard (see e.g. [27]) the sequence of  $\{\mathcal{A}_\Lambda\}_{\Lambda \in \mathbb{N}}$ , with  $\mathcal{A}_\Lambda := \text{End}(\mathcal{H}_{\Lambda\lambda})$ , as a fuzzy quantization of the symplectic space  $\mathcal{O}_\lambda \simeq G/G_\lambda$ . We recall that the Killing form  $B$  of  $so(\mathbf{D})$  gives  $B(L_{HI}, L_{JK}) =$

$2(\mathbf{D}-2) (\delta_J^H \delta_K^I - \delta_K^H \delta_J^I)$  for all  $H, I, J, K \in \{1, 2, \dots, \mathbf{D} \equiv D+1\}$ . Let  $\sigma := \lfloor \frac{\mathbf{D}}{2} \rfloor = \text{rank of } so(\mathbf{D})$ . As the basis of the Cartan subalgebra  $\mathfrak{h}$  of  $so(\mathbf{D})$  we choose  $\{H_a\}_{a=1}^\sigma$ , where

$$H_\sigma := L_{D\mathbf{D}}, \quad H_{\sigma-1} := L_{(d-1)d}, \quad \dots, \quad H_1 = \begin{cases} L_{12} & \text{if } \mathbf{D} = 2\sigma \\ L_{23} & \text{if } \mathbf{D} = 2\sigma+1 \end{cases} \quad (125)$$

We choose the irrep of  $Uso(\mathbf{D})$  on  $V_{\mathbf{D}}^\Lambda \simeq \mathcal{H}_\Lambda$ ; as the highest weight vector  $\Omega_{\mathbf{D}}^\Lambda \in V_{\mathbf{D}}^\Lambda$  we choose  $\Omega_{\mathbf{D}}^\Lambda := (t^D + it^{\mathbf{D}})^\Lambda$  (for brevity we do not write down the associated partition of roots of  $so(\mathbf{D})$  into positive and negative). The associated weight in the chosen basis, i.e. the joint spectrum of  $H := (H_1, \dots, H_\sigma)$ , is  $\Lambda = (0, \dots, 0, \Lambda)$ . Identifying weights  $\lambda \in \mathfrak{h}^*$  with elements  $H_\lambda \in \mathfrak{h}$  via the Killing form, we find that  $H_\Lambda \propto H_\sigma = L_{D\mathbf{D}}$ . The stabilizer of the latter in  $SO(\mathbf{D})$  is  $SO(2) \times SO(d)$ , where  $SO(2)$ ,  $SO(d)$  have Lie algebra respectively spanned by  $L_{D\mathbf{D}}$  and by the  $L_{ij}$  with  $i, j < D$ . Therefore the corresponding coadjoint orbit  $SO(\mathbf{D})/(SO(2) \times SO(d))$  has dimension

$$\frac{D(D+1)}{2} - 1 - \frac{(D-2)(D-1)}{2} = 2(D-1) = 2d,$$

which is also the dimension of  $T^*S^d$ , the cotangent space of the  $d$ -dimensional sphere  $S^d$  (or phase space over  $S^d$ ). This is consistent with the interpretation of  $\mathcal{A}_\Lambda$  as the algebra of observables (quantized phase space) on the fuzzy sphere. This would have not been the case if we had chosen some other generic irrep of  $Uso(\mathbf{D})$ : the coadjoint orbit would have been some other equivariant bundle over  $S^d$  [27].

For instance, the 4-dimensional fuzzy spheres introduced in [16], as well as the ones of dimension  $d \geq 3$  considered in [17, 18, 19], are based on  $End(V)$ , where  $V$  carries an *irreducible representation* of both  $Spin(D)$  and  $Spin(D+1)$ , and therefore of both  $Uso(D)$  and  $Uso(\mathbf{D})$ . Note: first, this may be only projective as a representation of  $O(D)$ ; second, as  $\Lambda \rightarrow \infty$   $V$  does not go to  $\mathcal{L}^2(S^d)$  as a representation of  $Uso(D)$ , in contrast with our  $\mathcal{H}_\Lambda \simeq V_{\mathbf{D}}^\Lambda$ . The  $X^i := L_{i\mathbf{D}}$  play the role of fuzzy coordinates. As  $\mathbf{x}^2 \equiv X^i X^i$  is central, it can be set  $\mathbf{x}^2 = 1$  identically. The commutation relations are also  $O(D)$ -covariant and Snyder-like, except for the case of the Madore-Hoppe fuzzy sphere [12, 13]. The corresponding coadjoint orbit for  $d = 4$  is the 6-dimensional  $\mathbb{C}P^3$  [20, 21], which can be seen as a  $so(5)$ -equivariant  $S^2$  bundle over  $S^4$  (while [17] does not identify coadjoint orbits for generic  $d$ ).

In [20, 21] the authors consider also constructing a fuzzy 4-sphere  $S_N^4$  through a *reducible* representation of  $Uso(5)$  on a Hilbert space  $V$  obtained decomposing an irrep  $\pi$  of  $Uso(6)$  characterized by a highest weight triple  $(N, 0, n)$  with respect to  $(H'_{\lambda_1}, H'_{\lambda_1}, H'_{\lambda_1})$ , where

$$H'_{\lambda_1} := \frac{i}{2}(L_{34} + L_{12} + L_{56}), \quad H'_{\lambda_2} := iL_{56}, \quad H'_{\lambda_3} := \frac{i}{2}(L_{34} - L_{12} + L_{56}).$$

The  $X^i = L_{i6}$  ( $i = 1, \dots, 5$ ), which make up a basis of the vector space  $so(6) \setminus so(5)$ , still play the role of noncommuting Cartesian coordinates. If  $n = 0$  then  $\mathbf{x}^2 \equiv 1$  ( $V$  carries an irrep of  $O(5)$ ), and one recovers the ( $so(5)$ -equivariant  $S^2$  bundle over the) fuzzy 4-sphere of [16]. If  $n > 0$ , then the  $O(5)$ -scalar  $\mathbf{x}^2 = X^i X^i$  is no longer central, but its spectrum is still very close to 1 provided  $N \gg n$ , because then  $V$  decomposes only in few irreducible  $SO(5)$ -components, all with eigenvalues of  $\mathbf{x}^2$  very close to 1. The associated coadjoint



orbit is 10-dimensional and can be seen as a  $so(5)$ -equivariant  $\mathbb{C}P^2$  bundle over  $\mathbb{C}P^3$ , or a  $so(5)$ -equivariant twisted bundle over either  $S_N^4$  or  $S_n^4$ . On the contrary, with respect to  $(H'_{\lambda_1}, H'_{\lambda_1}, H'_{\lambda_1})$  the highest weight triple of the irrep  $V_6^\Lambda$  considered here is instead  $(\Lambda, \Lambda, \Lambda)$ ; as said,  $\mathbf{x}^2 \equiv \bar{x}^i \bar{x}^i \simeq 1$  is guaranteed by adopting as noncommutative Cartesian coordinates the  $\bar{x}^i = m_\Lambda(L^2) X^i m_\Lambda(L^2)$ , with a suitable function  $m_\Lambda$ , rather than the  $X^i$ , and the associated coadjoint orbit has dimension 8, which is also the dimension of  $T^*S^4$ , as wished.

Many aspects of these new fuzzy spheres deserve investigations: e.g. space uncertainties, optimally localized states and coherent states also for  $d > 2$ , as done in [28, 29, 24] for  $d = 1, 2$ ; a distance between optimally localized states (as done e.g. in [30] for the FS); extending the construction to particles with spin; QFT based on our fuzzy spheres; application of our fuzzy spheres to problems in quantum gravity, or condensed matter physics; etc.

Finally, we point out that a different approach to the construction of noncommutative submanifolds of noncommutative  $\mathbb{R}^D$ , equivariant with respect to a ‘quantum group (twisted Hopf algebra) has been proposed in [32, 31]; it is based on a systematic use of Drinfel’d twists.

## Acknowledgments

We thank F. Pisacane, H. Steinacker for useful discussions in the early stages of the work.

# 8 Appendix

## 8.1 Proof of Proposition 3.2

Our Ansatz is (53) with  $M \equiv M(l+1)$  a  $O(D)$ -invariant matrix to be determined. The most general one is

$$M(l+1) = \alpha_{l+1} (\mathbf{1}_{D^2} + \beta_{l+1} \mathbf{P} + \gamma_{l+1} \mathcal{P}^t) \quad (126)$$

We first determine the coefficients  $\beta_{l+1}, \gamma_{l+1}$  by imposing the conditions (49). By the recursive assumption, only the condition with  $m = l$  is not fulfilled automatically and must be imposed by hand. Actually, it suffices to impose just (49a), due to the symmetry of the Ansatz (53) and of the matrices  $\mathcal{P}^l$  under transposition. Abbreviating  $M' \equiv M(l)$  this amounts to

$$\begin{aligned} 0 &\stackrel{!}{=} \mathcal{P}^{l+1} \mathcal{P}'_{l(l+1)} \stackrel{(53)}{=} \mathcal{P}'_{1\dots l} M_{l(l+1)} \mathcal{P}^l_{1\dots l} \mathcal{P}'_{l(l+1)} \\ &\stackrel{(53)_{l \rightarrow l-1}}{=} \mathcal{P}'_{1\dots l} M_{l(l+1)} \mathcal{P}^{l-1}_{1\dots(l-1)} M'_{(l-1)l} \mathcal{P}^{l-1}_{1\dots(l-1)} \mathcal{P}'_{l(l+1)} \\ &= \mathcal{P}'_{1\dots l} \mathcal{P}^{l-1}_{1\dots(l-1)} M_{l(l+1)} M'_{(l-1)l} \mathcal{P}^{l-1}_{1\dots(l-1)} \mathcal{P}'_{l(l+1)} = \mathcal{P}'_{1\dots l} M_{l(l+1)} M'_{(l-1)l} \mathcal{P}^{l-1}_{1\dots(l-1)} \mathcal{P}'_{l(l+1)} \\ &\propto \mathcal{P}'_{1\dots l} (\mathbf{1}_{D^{l+1}} + \beta_{l+1} \mathbf{P}_{l(l+1)} + \gamma_{l+1} \mathcal{P}^t_{l(l+1)}) (\mathbf{1}_{D^{l+1}} + \beta_l \mathbf{P}_{(l-1)l} + \gamma_l \mathcal{P}^t_{(l-1)l}) \mathcal{P}^{l-1}_{1\dots(l-1)} \mathcal{P}'_{l(l+1)} \\ &= \mathcal{P}'_{1\dots l} \left[ \mathbf{1}_{D^{l+1}} + \beta_{l+1} \mathbf{P}_{l(l+1)} + \gamma_{l+1} \mathcal{P}^t_{l(l+1)} + \beta_l \mathbf{P}_{(l-1)l} + \gamma_l \cancel{\mathcal{P}^t_{(l-1)l}} + \beta_l \beta_{l+1} \mathbf{P}_{l(l+1)} \mathbf{P}_{(l-1)l} \right] \end{aligned}$$

(127) (49) (49) (48), (49)

$$\begin{aligned}
& +\gamma_l\gamma_{l+1}\mathcal{P}_{l(l+1)}^t\mathcal{P}_{(l-1)l}^t+\beta_l\gamma_{l+1}\mathcal{P}_{l(l+1)}^t\mathbf{P}_{(l-1)l}+\gamma_l\beta_{l+1}\mathbf{P}_{l(l+1)}\mathcal{P}_{(l-1)l}^t \Big] \mathcal{P}_{1\dots(l-1)}^{l-1}\mathcal{P}'_{l(l+1)} \\
& \stackrel{(128)}{=} \mathcal{P}_{1\dots l}^l \left\{ \mathbf{1}_{D^{l+1}} + \beta_{l+1} [2\mathcal{P}_{l(l+1)}^t - \mathbf{1}_{D^{l+1}}] + \gamma_{l+1}\mathcal{P}_{l(l+1)}^t + \beta_l\mathbf{1}_{D^{l+1}} + \gamma_l\gamma_{l+1}\mathcal{P}_{l(l+1)}^t\mathcal{P}_{(l-1)l}^t \right. \\
& \quad \left. + \beta_l\gamma_{l+1}D\mathcal{P}_{l(l+1)}^t\mathcal{P}_{(l-1)l}^t \left[ 2\mathcal{P}_{l(l+1)}^t - \mathbf{1}_{D^{l+1}} \right] + \gamma_l\beta_{l+1}D\mathcal{P}_{l(l+1)}^t\mathcal{P}_{(l-1)l}^t \right\} \mathcal{P}_{1\dots(l-1)}^{l-1}\mathcal{P}'_{l(l+1)} \\
& \stackrel{(130)}{=} \mathcal{P}_{1\dots l}^l \left\{ \mathbf{1}_{D^{l+1}} [1 - \beta_{l+1} + \beta_l] + \mathcal{P}_{l(l+1)}^t \left[ 2\beta_{l+1} + \gamma_{l+1} + \beta_l\gamma_{l+1}\frac{2}{D} \right] \right. \\
& \quad \left. + \mathcal{P}_{l(l+1)}^t\mathcal{P}_{(l-1)l}^t [\gamma_l\gamma_{l+1} - \beta_l\gamma_{l+1}D + \gamma_l\beta_{l+1}D] \right\} \mathcal{P}_{1\dots(l-1)}^{l-1}\mathcal{P}'_{l(l+1)}.
\end{aligned}$$

where we have used also the relations

$$\mathcal{P}\mathcal{P}' = \mathcal{P} [\mathcal{P}^t + \mathcal{P}^-] \mathcal{P}' = [\mathcal{P}^t - \mathcal{P}^-] \mathcal{P}' = [2\mathcal{P}^t - \mathbf{1}_{D^2}] \mathcal{P}' \quad (127)$$

$$\begin{aligned}
\mathcal{P}_{l(l+1)}^t\mathbf{P}_{(l-1)l}\mathcal{P}'_{l(l+1)} & \stackrel{(47)}{=} D\mathcal{P}_{l(l+1)}^t\mathcal{P}_{(l-1)l}^t\mathbf{P}_{l(l+1)}\mathcal{P}'_{l(l+1)} \\
& \stackrel{(127)}{=} D\mathcal{P}_{l(l+1)}^t\mathcal{P}_{(l-1)l}^t [2\mathcal{P}_{l(l+1)}^t - \mathbf{1}_{D^{l+1}}] \mathcal{P}'_{l(l+1)}
\end{aligned} \quad (128)$$

$$\mathcal{P}_{1\dots l}^l\mathbf{P}_{l(l+1)}\mathcal{P}_{(l-1)l}^t \stackrel{(146)}{=} D\mathcal{P}_{1\dots l}^l\mathbf{P}_{(l-1)l}\mathcal{P}_{l(l+1)}^t\mathcal{P}_{(l-1)l}^t \stackrel{(49)}{=} D\mathcal{P}_{1\dots l}^l\mathcal{P}_{l(l+1)}^t\mathcal{P}_{(l-1)l}^t \quad (129)$$

$$\mathcal{P}_{l(l+1)}^t\mathcal{P}_{(l-1)l}^t\mathcal{P}_{l(l+1)}^t = \frac{1}{D^2}\mathcal{P}_{l(l+1)}^t. \quad (130)$$

The conditions that the three square brackets vanish

$$\begin{aligned}
1 + \beta_l - \beta_{l+1} &= 0, \\
2\beta_{l+1} + \gamma_{l+1} + \beta_l\gamma_{l+1}\frac{2}{D} &= 0, \\
\gamma_l\gamma_{l+1} - \beta_l\gamma_{l+1}D + \gamma_l\beta_{l+1}D &= 0,
\end{aligned}$$

are recursively solved, starting from  $l = 1$  with initial input  $\beta_1 = 0 = \gamma_1$  (since  $\mathcal{P}^1 = \mathbf{1}_D$ ), by

$$\beta_{l+1} = l, \quad \gamma_{l+1} = -\frac{2Dl}{D+2l-2}.$$

We determine the coefficient  $\alpha_{l+1}$  by imposing the condition (50). This gives

$$\begin{aligned}
0 & \stackrel{!}{=} \mathcal{P}^{l+1} (\mathcal{P}^{l+1} - \mathbf{1}_{D^{l+1}}) \stackrel{(53)}{=} \mathcal{P}^{l+1} (\mathcal{P}_{1\dots l}^l M_{l(l+1)} \mathcal{P}_{1\dots l}^l - \mathbf{1}_{D^{l+1}}) \\
& \stackrel{(52),(55)}{=} \mathcal{P}^{l+1} [\alpha_{l+1} (1 + \beta_{l+1}) \mathcal{P}_{1\dots l}^l - 1] \stackrel{(52)}{=} \mathcal{P}^{l+1} [\alpha_{l+1} (1 + \beta_{l+1}) - 1].
\end{aligned}$$

The condition that the square bracket vanishes is recursively solved, starting from  $l = 0$  with initial input  $\alpha_0 = 1$ , by  $\alpha_{l+1} = 1/(l+1)$ . This makes (126) into (55) [yielding back (41) if  $l = 2$ ]. We have thus proved that the Ansatz (53) fulfills (49-50). Similarly one proves that also the Ansatz (54) does the same job.

## 8.2 Proof of Proposition (3.3)

For  $l = 1$  it is  $\zeta_2 = 1$ ,  $\mathcal{P}^1 = \mathbf{1}_D$ ,  $\mathcal{P}^2 = \mathcal{P}$ , and the claim (60) is true by (41), (6). We now show that (60) <sub>$l \rightarrow l-1$</sub>  implies (60):

$$\begin{aligned}
& (\mathcal{P}_{1\dots l}^l - \mathcal{P}^{l+1}) x_1 \dots x_{l+1} \stackrel{(53)}{=} \mathcal{P}_{1\dots l}^l \left[ \mathbf{1}^{\otimes^{l+1}} - \frac{1}{l+1} \left( \mathbf{1}^{\otimes^{l+1}} + l \mathbf{P}_{l(l+1)} + \gamma_{l+1} \mathcal{P}_{l(l+1)}^t \right) \right] \mathcal{P}_{1\dots l}^l x_1 \dots x_{l+1} \\
&= \frac{1}{l+1} \mathcal{P}_{1\dots l}^l \left[ l \left( \mathbf{1}^{\otimes^{l+1}} - \mathbf{P}_{l(l+1)} \right) - \gamma_{l+1} \mathcal{P}_{l(l+1)}^t \right] \left( \mathcal{P}_{1\dots l}^l - \mathcal{P}_{1\dots(l-1)}^{l-1} + \mathcal{P}_{1\dots(l-1)}^{l-1} \right) x_1 \dots x_{l+1} \\
&\stackrel{(60)_{l \rightarrow l-1}}{=} \frac{1}{l+1} \mathcal{P}_{1\dots l}^l \left[ l \left( \mathbf{1}^{\otimes^{l+1}} - \mathbf{P}_{l(l+1)} \right) - \gamma_{l+1} \mathcal{P}_{l(l+1)}^t \right] \mathcal{P}_{1\dots(l-1)}^{l-1} \left( \mathbf{1}^{\otimes^{l+1}} - \zeta_l \mathcal{P}_{(l-1)l}^t \right) x_1 \dots x_{l+1} \\
&= \frac{1}{l+1} \mathcal{P}_{1\dots l}^l \left[ l \left( \mathbf{1}^{\otimes^{l+1}} - \mathbf{P}_{l(l+1)} \right) - \gamma_{l+1} \mathcal{P}_{l(l+1)}^t \right] \left( \mathbf{1}^{\otimes^{l+1}} - \zeta_l \mathcal{P}_{(l-1)l}^t \right) x_1 \dots x_{l+1} \\
&= \frac{1}{l+1} \mathcal{P}_{1\dots l}^l \left[ \cancel{l \left( \mathbf{1}^{\otimes^{l+1}} - \mathbf{P}_{l(l+1)} \right)}_{x_1 x_2 - \mathbf{P}_{x_1 x_2} = 0} - \gamma_{l+1} \mathcal{P}_{l(l+1)}^t - \zeta_l l \left( \mathbf{1}^{\otimes^{l+1}} - \mathbf{P}_{l(l+1)} \right) \mathcal{P}_{(l-1)l}^t + \zeta_l \gamma_{l+1} \mathcal{P}_{l(l+1)}^t \mathcal{P}_{(l-1)l}^t \right] x_1 \dots x_{l+1} \\
&= \frac{1}{l+1} \mathcal{P}_{1\dots l}^l \left[ -\gamma_{l+1} \mathcal{P}_{l(l+1)}^t - \zeta_l l \left( \cancel{\mathcal{P}_{(l-1)l}^t - \mathbf{P}_{l(l+1)} \mathcal{P}_{(l-1)l}^t} \right) + \zeta_l \gamma_{l+1} \mathcal{P}_{l(l+1)}^t \mathcal{P}_{(l-1)l}^t \mathbf{P}_{l(l+1)} \right] x_1 \dots x_{l+1} \\
&\stackrel{(47)}{=} \frac{1}{l+1} \mathcal{P}_{1\dots l}^l \left[ -\gamma_{l+1} \mathcal{P}_{l(l+1)}^t + \zeta_l l \mathbf{P}_{(l-1)l} \mathbf{P}_{l(l+1)} \mathcal{P}_{(l-1)l}^t + \frac{\zeta_l}{D} \gamma_{l+1} \mathcal{P}_{l(l+1)}^t \mathbf{P}_{(l-1)l} \right] x_1 \dots x_{l+1} \\
&\stackrel{(44)}{=} \frac{1}{l+1} \mathcal{P}_{1\dots l}^l \left[ -\gamma_{l+1} \mathcal{P}_{l(l+1)}^t + \zeta_l l \mathcal{P}_{l(l+1)}^t \mathbf{P}_{(l-1)l} \mathbf{P}_{l(l+1)} + \frac{\zeta_l}{D} \gamma_{l+1} \mathcal{P}_{l(l+1)}^t \mathbf{P}_{(l-1)l} \right] x_1 \dots x_{l+1} \\
&\stackrel{P_{x_1 x_2} = x_1 x_2}{=} \frac{1}{l+1} \mathcal{P}_{1\dots l}^l \left[ -\gamma_{l+1} \mathcal{P}_{l(l+1)}^t + \zeta_l l \mathcal{P}_{l(l+1)}^t + \frac{\zeta_l}{D} \gamma_{l+1} \mathcal{P}_{l(l+1)}^t \right] x_1 \dots x_{l+1} \\
&= \frac{1}{l+1} \left[ -\gamma_{l+1} + \zeta_l l + \frac{\zeta_l}{D} \gamma_{l+1} \right] \mathcal{P}_{1\dots l}^l \mathcal{P}_{l(l+1)}^t x_1 \dots x_{l+1} =: \zeta_{l+1} \mathcal{P}_{1\dots l}^l \mathcal{P}_{l(l+1)}^t x_1 \dots x_{l+1}
\end{aligned}$$

namely the left equality in (60) is fulfilled if the  $\zeta_l$  satisfy the recursion relation

$$\zeta_{l+1} = \frac{1}{(l+1)} \left[ -\gamma_{l+1} + \zeta_l l + \frac{\zeta_l}{D} \gamma_{l+1} \right] = \frac{l}{(l+1)(D+2l-2)} [2D + \zeta_l (D+2l-4)],$$

which setting  $\zeta_2 = 1$  is actually solved by  $\zeta_l = \frac{D(l-1)}{(D+2l-4)} = -\frac{1}{2} \gamma_l$ , as claimed.

## 8.3 Proof of Proposition 3.4

Using Proposition 3.3 we easily find

$$\begin{aligned}
& i L_{hk} X_l^{i_1 \dots i_l} = (x^h \partial^k - x^k \partial^h) \mathcal{P}_{j_1 \dots j_l}^{l i_1 \dots i_l} x^{j_1} \dots x^{j_l} = l (x^h g^{kj_1} - x^k g^{hj_1}) \mathcal{P}_{j_1 \dots j_l}^{l i_1 \dots i_l} x^{j_2} \dots x^{j_l} \quad (131) \\
&\stackrel{(61)}{=} \frac{lD}{\zeta_{l+1} r^2} \left[ x^h \left( x^k X_l^{i_1 \dots i_l} - X_{l+1}^{k i_1 \dots i_l} \right) - x^k \left( x^h X_l^{i_1 \dots i_l} - X_{l+1}^{h i_1 \dots i_l} \right) \right] = \frac{lD}{\zeta_{l+1}} \frac{x^k X_{l+1}^{h i_1 \dots i_l} - x^h X_{l+1}^{k i_1 \dots i_l}}{r^2} \\
&\stackrel{(61)}{=} \frac{lD}{\zeta_{l+1} r^2} \left[ \cancel{X_{l+2}^{k h i_1 \dots i_l}} - \cancel{X_{l+2}^{h k i_1 \dots i_l}} + \frac{\zeta_{l+2}}{D} r^2 \left( g^{kj} \mathcal{P}_{j j_1 \dots j_l}^{l+1 h i_1 \dots i_l} - g^{hj} \mathcal{P}_{j j_1 \dots j_l}^{l+1 k i_1 \dots i_l} \right) X_l^{j_1 \dots j_l} \right] \\
&= \frac{l \zeta_{l+2}}{\zeta_{l+1}} \left( g^{kj} \mathcal{P}_{j j_1 \dots j_l}^{l+1 h i_1 \dots i_l} - g^{hj} \mathcal{P}_{j j_1 \dots j_l}^{l+1 k i_1 \dots i_l} \right) X_l^{j_1 \dots j_l}.
\end{aligned}$$

Alternatively, from (131) we obtain also (63), because

$$\begin{aligned}
& iL_{hk}X_l^{i_1\dots i_l} \stackrel{(52b)}{=} l(x^h g^{kj_1} - x^k g^{hj_1})\mathcal{P}_{j_1\dots j_l}^{l i_1\dots i_l} X_{l-1}^{j_2\dots j_l} \\
& \stackrel{(61)}{=} \mathcal{P}_{j_1\dots j_l}^{l i_1\dots i_l} l g^{kj_1} \left( X_l^{hj_2\dots j_l} + \frac{\zeta_l}{D} r^2 g^{hk_2} \mathcal{P}_{k_2\dots k_l}^{l-1 j_2\dots j_l} X_{l-2}^{k_3\dots k_l} \right) - (h \leftrightarrow k) \\
& = \mathcal{P}_{j_1\dots j_l}^{l i_1\dots i_l} l g^{kj_1} X_l^{hj_2\dots j_l} + l \frac{\zeta_l}{D} r^2 g^{kj_1} g^{hk_2} \mathcal{P}_{j_1 k_2\dots k_l}^{l i_1 i_2\dots i_l} X_{l-2}^{k_3\dots k_l} - (h \leftrightarrow k) \\
& = l \mathcal{P}_{j_1\dots j_l}^{l i_1\dots i_l} \left( g^{kj_1} X_l^{hj_2\dots j_l} - g^{hj_1} X_l^{kj_2\dots j_l} \right).
\end{aligned}$$

Multiplying the previous relations by  $1/r^l$ , which commutes with  $L_{hk}$ , (63) give (68).

## 8.4 Proof of Proposition 3.6 and Theorem 3.7

The right-hand side (rhs) of (72) is the sum of  $\bar{N}!$  terms; in particular,  $G_2^{ij} = \delta^{ij} + \delta^{ji} = 2\delta^{ij}$ .  $G_{\bar{N}}, G_{\bar{N}-2}$  are related by the recursive relation

$$G_{\bar{N}}^{i_1\dots i_{\bar{N}}} = (\delta^{i_1 i_2} + \delta^{i_2 i_1}) G_{\bar{N}-2}^{i_3\dots i_{\bar{N}}} + (\delta^{i_1 i_3} + \delta^{i_3 i_1}) G_{\bar{N}-2}^{i_2 i_4\dots i_{\bar{N}}} + \dots + (\delta^{i_{\bar{N}-1} i_{\bar{N}}} + \delta^{i_{\bar{N}} i_{\bar{N}-1}}) G_{\bar{N}-2}^{i_1\dots i_{\bar{N}-2}}. \quad (132)$$

The rhs is the sum of  $\bar{N}(\bar{N}-1)$  products  $\delta^{\dots} G_{\bar{N}-2}^{\dots}$ . The ‘trace’ of  $G_{\bar{N}}$  equals  $\text{tr}(G_2) = 2D$  for  $\bar{N} = 2$  and by (132) fulfills the recursive relation  $\text{tr}(G_{\bar{N}}) = \bar{N}(D+\bar{N}-2) \text{tr}(G_{\bar{N}-2})$ . In fact, each of the  $\bar{N}$  products  $\delta^{i_1 i_2} G_{\bar{N}-2}^{i_3\dots i_{\bar{N}}}$ ,  $\delta^{i_2 i_1} G_{\bar{N}-2}^{i_3\dots i_{\bar{N}}}$ ,  $\delta^{i_3 i_4} G_{\bar{N}-2}^{i_1 i_2 i_5\dots i_{\bar{N}}}$ ,  $\delta^{i_4 i_3} G_{\bar{N}-2}^{i_1 i_2 i_5\dots i_{\bar{N}}}$ , ...,  $\delta^{i_{\bar{N}} i_{\bar{N}-1}} G_{\bar{N}-2}^{i_1\dots i_{\bar{N}-2}}$  in (132) contributes by  $D \text{tr}(G_{\bar{N}-2})$ , while each of the  $\bar{N}(\bar{N}-2)$  remaining ones contributes by  $\text{tr}(G_{\bar{N}-2})$ . The recursion relation is solved by (73). The integral over  $S^d$  of  $H_{\bar{N}}^{i_1\dots i_{\bar{N}}} := t^{i_1} \dots t^{i_{\bar{N}}}$  is  $O(D)$ -invariant and therefore must be proportional to  $G_{\bar{N}}^{i_1\dots i_{\bar{N}}}$ ; the proportionality coefficient  $C_{\bar{N}}$  is found by consistency contracting both sides with  $\delta_{i_1 i_2} \delta_{i_3 i_4} \dots \delta_{i_{\bar{N}-1} i_{\bar{N}}}$ , and using (27), (73). The scalar product of  $\phi^l, \psi^l \in V_D^l \subset \mathcal{H}_s$  is given by

$$\begin{aligned}
\langle \phi^l, \psi^l \rangle &= \int_{S^d} d\alpha \phi^{l*} \psi^l = (\phi_{i_1\dots i_l}^l)^* \psi_{j_1\dots j_l}^l \int_{S^d} d\alpha t^{i_1} \dots t^{i_l} t^{j_1} \dots t^{j_l} \\
&\stackrel{(74)}{=} C_{2l} (\phi_{i_1\dots i_l}^l)^* \psi_{j_1\dots j_l}^l G_{2l}^{i_1\dots i_l j_1\dots j_l},
\end{aligned}$$

the sum has  $(2l)!$  terms. In fact, all terms where both indices of at least one Kronecker  $\delta$  contained in  $G_{2l}$  are contracted with the two indices of the coefficients  $\phi_{i_1\dots i_l}^l$ , or of the  $\psi_{i_1\dots i_l}^l$ , vanish, by (70). The remaining  $2^l(l!)^2$  terms arise from the  $(l!)^2$  products contained in  $G_{2l}$  of the type  $\delta^{i_{\pi(1)} j_{\pi'(1)}} \dots \delta^{i_{\pi(l)} j_{\pi'(l)}}$ , where  $\pi, \pi'$  are permutations of  $(1, \dots, l)$ , and the other ones which are obtained exchanging the order of the indices  $i_{\pi(h)} j_{\pi'(h)} \mapsto j_{\pi'(h)} i_{\pi(h)}$  in one or more of these Kronecker  $\delta$ 's; they are all equal, again by (70). Hence,

$$\langle \phi^l, \psi^l \rangle = Q_l (\phi_{i_1\dots i_l}^l)^* \psi_{i_1\dots i_l}^l, \quad (133)$$

By the orthogonality  $V_D^l \perp V_D^{l'}$  for  $l \neq l'$  we find that the scalar product of generic  $\phi, \psi \in \mathcal{H}_s$  is given by (75). This concludes the proof of Proposition 3.6.

Applying  $m$  times (66) and absorbing into a suitable combination of  $T_n$ 's ( $n \in L$ ) the  $m$ -degree monomials in  $t^i$  whose combination gives  $T_m^{j_1 \dots j_m}$ , we find that (78) must hold with suitable coefficients  $S_{k_1 \dots k_n}^{i_1 \dots i_l, j_1 \dots j_m} \in \mathbb{R}$ . We determine these coefficients faster using (78) as an Ansatz, making its scalar product with  $T_h^{k_1 \dots k_h}$  and using (76). One finds

$$\begin{aligned} Q_n S_{k_1 \dots k_n}^{i_1 \dots i_l, j_1 \dots j_m} &= \langle T_n^{k_1 \dots k_n}, T_l^{i_1 \dots i_l} T_m^{j_1 \dots j_m} \rangle \\ &= \mathcal{P}_{a_1 \dots a_l}^{l i_1 \dots i_l} \mathcal{P}_{b_1 \dots b_m}^{m j_1 \dots j_m} \mathcal{P}_{c_1 \dots c_n}^{h k_1 \dots k_n} \int_{S^d} d\alpha t^{a_1} \dots t^{a_l} t^{b_1} \dots t^{b_m} t^{c_1} \dots t^{c_n} \\ &= \mathcal{P}_{a_1 \dots a_l}^{l i_1 \dots i_l} \mathcal{P}_{b_1 \dots b_m}^{m j_1 \dots j_m} \mathcal{P}_{c_1 \dots c_n}^{n k_1 \dots k_n} C_{\bar{N}} G_{\bar{N}}^{a_1 \dots a_l b_1 \dots b_m c_1 \dots c_n}, \end{aligned} \quad (134)$$

where  $\bar{N} = l+m+n = n+r =: 2s$  is even. Due to the form of  $G_{\bar{N}}$ , the sum has  $\bar{N}!$  terms, each containing a product of  $s$  Kronecker  $\delta$ 's. All terms where both indices of some Kronecker  $\delta$  contained in  $G_{2s}$  are contracted with two indices of  $\mathcal{P}^l$ , or  $\mathcal{P}^m$ , or  $\mathcal{P}^n$ , vanish, by (51).

As a warm-up, consider first the case  $n = l+m$ . Renaming for convenience  $b_1, \dots, b_m$  as  $a_{l+1}, \dots, a_n$ , the remaining  $2^n (n!)^2$  terms arise from the  $(n!)^2$  products of the type

$$\delta^{c_{\pi(1)} a_{\pi'(1)}} \dots \delta^{c_{\pi(n)} a_{\pi'(n)}}$$

contained in  $G_{\bar{N}}$ , where  $\pi, \pi'$  are permutations of  $(1, \dots, h)$ , and the other ones which are obtained exchanging the order of the indices  $c_{\pi(h)} a_{\pi'(h)} \mapsto a_{\pi'(h)} c_{\pi(h)}$ , in one or more of these  $n$  Kronecker  $\delta$ 's; by the complete symmetry of  $\mathcal{P}^n$ , they are all equal to the term where  $\pi, \pi'$  are the trivial permutations; correspondingly, the product is  $\delta^{c_1 a_1} \dots \delta^{c_n a_n}$ . Hence,

$$Q_n S_{k_1 \dots k_n}^{i_1 \dots i_l, j_1 \dots j_m} = C_{2n} 2^n (n!)^2 \mathcal{P}_{a_1 \dots a_l}^{l i_1 \dots i_l} \mathcal{P}_{a_{l+1} \dots a_n}^{m j_1 \dots j_m} \mathcal{P}_{k_1 \dots k_n}^{n a_1 \dots a_n} = C_{2n} 2^n (n!)^2 \mathcal{P}_{k_1 \dots k_n}^{n i_1 \dots i_l, j_1 \dots j_m}$$

implying  $S_{k_1 \dots k_n}^{i_1 \dots i_l, j_1 \dots j_m} = \mathcal{P}_{k_1 \dots k_n}^{n i_1 \dots i_l, j_1 \dots j_m}$ , i.e. the term of highest rank at the rhs(78) is  $T_n^{i_1 \dots i_l, j_1 \dots j_m}$ . This is consistent with the first term in the (even iterated) application of (66).

For generic  $n \in L$  (134) becomes

$$\begin{aligned} Q_n S_{k_1 \dots k_n}^{i_1 \dots i_l, j_1 \dots j_m} &= C_{2s} F_n^{l,m} \mathcal{P}_{a_1 \dots a_l}^{l i_1 \dots i_l} \mathcal{P}_{b_1 \dots b_m}^{m j_1 \dots j_m} \mathcal{P}_{c_1 \dots c_n}^{n k_1 \dots k_n} \times \\ &\delta^{a_1 b_1} \delta^{a_2 b_2} \dots \delta^{a_r b_r} \delta^{a_{r+1} c_1} \dots \delta^{a_l c_{l-r}} \delta^{b_{r+1} c_{l-r+1}} \dots \delta^{b_m c_n}, \end{aligned} \quad (135)$$

where  $r := \frac{l+m-n}{2}$ ,  $s = \frac{l+m+n}{2}$ ,  $F_n^{l,m} = \frac{2^s s! l! m! (n!)^2}{(l-r)! (m-r)! (s-r)!} = \frac{2^s s! l! m! n!}{(l-r)! (m-r)!}$ . In fact, one of the products of Kronecker  $\delta$ 's contained in  $G_{2s}$  and yielding a nonzero contribution is displayed in the second line. The number  $F_n^{l,m}$  of such products can be determined as follows, starting from the same product with all indices removed: there are  $l$  ways to pick out the first index from the set  $A_l := \{a_1, \dots, a_l\}$ ,  $m$  ways to pick out the second index from the set  $B_m := \{b_1, \dots, b_m\}$ , hence  $lm$  ways to pick out the first index from  $A_l$  and the second from  $B_m$ ; similarly, there are  $lm$  ways to pick out the second index from  $A_l$  and the first from  $B_m$ ; altogether, there are  $2lm$  ways to pick out one of the first two indices from  $A_l$  and the other one from  $B_m$ . After anyone of these choices the corresponding sets of indices  $A_{l-1}$ ,  $B_{m-1}$  will have  $l-1$ ,  $m-1$  indices respectively; therefore there are  $2(l-1)(m-1)$  ways to pick out one among the third, fourth indices from  $A_{l-1}$  and the other from  $B_{m-1}$ . And so on. Therefore there are  $\frac{2^r l! m!}{(l-r)! (m-r)!}$  ways to pick out  $r$  indices appearing in the first  $r$   $\delta$ 's (one for each  $\delta$ ) from  $A_l$  and the  $r$  remaining ones from  $B_m$ . After anyone of these choices the corresponding sets of indices

$A_{l-r}$  and  $B_{m-r}$  will have  $l-r, m-r$  indices respectively, and  $D_n := A_{l-r} \cup B_{m-r}$  will have  $l-r+m-r = n$  indices. Reasoning as in the case  $n = l+m$ , we find that there are  $2^n (n!)^2$  ways to pick out  $n$  indices appearing in the remaining  $n$   $\delta$ 's (one for each  $\delta$ ) from  $D_n$  and the remaining  $h$  ones from  $C_n := \{c_1, \dots, c_n\}$ . Consequently, so far there are  $\frac{2^{n+r} l! m!}{(l-r)! (m-r)!} (n!)^2$  ways to do these operations. The remaining ways are obtained allowing that the  $r$  pair of indices picked one out of  $A_l$  and the other out of  $B_m$  appear not necessarily in the first  $r$   $\delta$ 's, but in any subgroup of  $r$   $\delta$ 's out of the totality of  $s$ ; hence we have to multiply the previous number by the number  $\frac{s!}{(s-r)!}$  of such subgroups, and we finally obtain  $F_n^{l,m} = \frac{s! 2^s l! m! (n!)^2}{(s-r)! (l-r)! (m-r)!}$  (because  $n+r = s$ ), as claimed. Thus (135) implies the following relation, which gives (79):

$$S_{k_1 \dots k_n}^{i_1 \dots i_l, j_1 \dots j_m} = \frac{(D+2n-2)!!}{(D+2s-2)!!} \frac{2^s s! l! m! n!}{n! (2s)!! (l-r)! (m-r)!} \mathcal{P}_{a_1 \dots a_l}^{l i_1 \dots i_l} \mathcal{P}_{b_1 \dots b_m}^{m j_1 \dots j_m} \mathcal{P}_{c_1 \dots c_n}^{n k_1 \dots k_n} \\ \times \delta^{a_1 b_1} \delta^{a_2 b_2} \dots \delta^{a_r b_r} \delta^{a_{r+1} c_1} \dots \delta^{a_l c_{l-r}} \delta^{b_{r+1} c_{l-r+1}} \dots \delta^{b_m c_n}$$

## 8.5 Proof of Proposition 3.8

Since  $L_{hk}$  commute with scalars, and  $[L_{hk}, x^{\mathbf{D}}] = 0$ ,  $\check{p}_{\Lambda, l}$  commutes with all the  $L_{hk}$  and therefore also with  $\mathbf{L}^2$ . Hence, using (58), we find  $\mathbf{L}^2 \check{p}_{\Lambda, l} X_l^{i_1 \dots i_l} = \check{p}_{\Lambda, l} \mathbf{L}^2 X_l^{i_1 \dots i_l} = E_l \check{p}_{\Lambda, l} X_l^{i_1 \dots i_l}$ , i.e (85). To compute the coefficients  $b_{\Lambda, l+2k}$  we preliminarily note that

$$\begin{aligned} \Delta(x^i)^h &= (x^i)^h \Delta + 2h(x^i)^{h-1} \partial^i + h(h-1)(x^i)^{h-2}, \\ \Delta r^h &= r^h \Delta + hr^{h-2}(2\eta + D + h - 2) \\ \Delta_{\mathbf{D}}(x^I)^h &= (x^I)^h \Delta + 2h(x^I)^{h-1} \partial^I + h(h-1)(x^I)^{h-2}, \\ \Delta_{\mathbf{D}} r_{\mathbf{D}}^h &= r_{\mathbf{D}}^h \Delta_{\mathbf{D}} + hr_{\mathbf{D}}^{h-2}(2\bar{\eta} + \mathbf{D} + h - 2) \\ \partial^j \check{p}_{\Lambda, l} &= x^j \left[ 2(x^{\mathbf{D}})^{\Lambda-l-2} b_{\Lambda, l+2} + 4(x^{\mathbf{D}})^{\Lambda-l-4} r_{\mathbf{D}}^2 b_{\Lambda, l+4} + \dots \right] \end{aligned}$$

$$\begin{aligned} \Delta_{\mathbf{D}} \check{p}_{\Lambda, l} &= (\Lambda-l)(\Lambda-l-1) (x^{\mathbf{D}})^{\Lambda-l-2} + [r_{\mathbf{D}}^2 \Delta_{\mathbf{D}} + 2(2\bar{\eta} + \mathbf{D})] (x^{\mathbf{D}})^{\Lambda-l-2} b_{\Lambda, l+2} \\ &+ [r_{\mathbf{D}}^4 \Delta_{\mathbf{D}} + 4r_{\mathbf{D}}^2 (2\bar{\eta} + \mathbf{D} + 2)] (x^{\mathbf{D}})^{\Lambda-l-4} b_{\Lambda, l+4} + \dots \\ &= (\Lambda-l)(\Lambda-l-1) (x^{\mathbf{D}})^{\Lambda-l-2} \\ &+ \left\{ r_{\mathbf{D}}^2 (\Lambda-l-2)(\Lambda-l-3) + 2[2(\Lambda-l-2) + \mathbf{D}] (x^{\mathbf{D}})^2 \right\} (x^{\mathbf{D}})^{\Lambda-l-4} b_{\Lambda, l+2} \\ &+ \left\{ r_{\mathbf{D}}^4 (\Lambda-l-4)(\Lambda-l-5) + 4r_{\mathbf{D}}^2 (x^{\mathbf{D}})^2 [2(\Lambda-l-4) + \mathbf{D} + 2] \right\} (x^{\mathbf{D}})^{\Lambda-l-6} b_{\Lambda, l+4} + \dots \\ &= (x^{\mathbf{D}})^{\Lambda-l-2} \{ (\Lambda-l)(\Lambda-l-1) + 2b_{\Lambda, l+2} [2(\Lambda-l-2) + \mathbf{D}] \} \\ &+ r_{\mathbf{D}}^2 (x^{\mathbf{D}})^{\Lambda-l-4} \{ (\Lambda-l-2)(\Lambda-l-3) b_{\Lambda, l+2} + 4b_{\Lambda, l+4} [2(\Lambda-l-4) + \mathbf{D} + 2] \} \\ &+ r_{\mathbf{D}}^4 (x^{\mathbf{D}})^{\Lambda-l-6} \{ (\Lambda-l-4)(\Lambda-l-5) + 6b_{\Lambda, l+6} [2(\Lambda-l-6) + \mathbf{D} + 4] \} + \dots \end{aligned}$$

We now impose that the  $\check{F}_{\mathbf{D},\Lambda}^{i_1 \dots i_l}$  are harmonic in dimension  $\mathbf{D}$ . By a direct calculation

$$\begin{aligned}
0 &= \Delta_{\mathbf{D}} \check{F}_{\mathbf{D},\Lambda}^{i_1 \dots i_l} = \mathcal{P}_{j_1 \dots j_l}^{i_1 \dots i_l} \{x^{j_1} \dots x^{j_l} \Delta_{\mathbf{D}} + 2lx^{j_1} \dots x^{j_{l-1}} \partial^{j_l}\} \check{p}_{\Lambda,l} = \mathcal{P}_{j_1 \dots j_l}^{i_1 \dots i_l} x^{j_1} \dots x^{j_l} \check{M}_{\Lambda,l+2} \Rightarrow \\
0 &= \check{M}_{\Lambda,l+2} := \Delta_{\mathbf{D}} \check{p}_{\Lambda,l} + 2l \left[ 2(x^{\mathbf{D}})^{\Lambda-l-2} b_{\Lambda,l+2} + 4(x^{\mathbf{D}})^{\Lambda-l-4} r_{\mathbf{D}}^2 b_{\Lambda,l+4} + \dots \right] \\
&= (x^{\mathbf{D}})^{\Lambda-l-2} \{(\Lambda-l)(\Lambda-l-1) + 2b_{\Lambda,l+2} [2l+2(\Lambda-l-2)+\mathbf{D}]\} \\
&+ (x^{\mathbf{D}})^{\Lambda-l-4} r_{\mathbf{D}}^2 \{b_{\Lambda,l+2}(\Lambda-l-2)(\Lambda-l-3) + 4b_{\Lambda,l+4} [2l+2(\Lambda-l-4)+\mathbf{D}+2]\} + \dots \\
&= (x^{\mathbf{D}})^{\Lambda-l-2} \{(\Lambda-l)(\Lambda-l-1) + 2b_{\Lambda,l+2} [2\Lambda-4+\mathbf{D}]\} \\
&+ (x^{\mathbf{D}})^{\Lambda-l-4} r_{\mathbf{D}}^2 \{b_{\Lambda,l+2}(\Lambda-l-2)(\Lambda-l-3) + 4b_{\Lambda,l+4} [2\Lambda-6+\mathbf{D}]\} + \dots;
\end{aligned}$$

the vanishing of the coefficient of each monomial  $(x^{\mathbf{D}})^{\Lambda-l-2-2h} r_{\mathbf{D}}^{2h}$  implies

$$\begin{aligned}
b_{\Lambda,l+2} &= -\frac{(\Lambda-l)(\Lambda-l-1)}{2(2\Lambda-4+\mathbf{D})}, \\
b_{\Lambda,l+4} &= -\frac{(\Lambda-l-2)(\Lambda-l-3)}{4(2\Lambda-6+\mathbf{D})} b_{\Lambda,l+2} = \frac{(\Lambda-l)(\Lambda-l-1)(\Lambda-l-2)(\Lambda-l-3)}{2 \cdot 4(2\Lambda-4+\mathbf{D})(2\Lambda-6+\mathbf{D})}, \\
&\dots;
\end{aligned}$$

namely, more compactly, we obtain (86).

The  $\check{F}_{\mathbf{D},\Lambda}^{i_1 \dots i_l}$  transform as the  $X_l^{i_1 \dots i_l}$  under the action of the  $L_{hk}$ , because the latter commute with  $\check{p}_{\Lambda,l}$ . Using (6), (82), and the fact that  $\partial^{\mathbf{D}}$  annihilates all polynomials in the  $x^i$ , we find

$$\begin{aligned}
iL_{h\mathbf{D}} \check{F}_{\mathbf{D},\Lambda}^{i_1 \dots i_l} &= (x^h \partial^{\mathbf{D}} - x^{\mathbf{D}} \partial^h) \check{p}_{\Lambda,l} X_l^{i_1 \dots i_l} \\
&= \left\{ x^h (x^{\mathbf{D}})^{\Lambda-l-1} (\Lambda-l) + x^h (x^{\mathbf{D}})^{\Lambda-l} \partial^{\mathcal{P}} - (x^{\mathbf{D}})^{\Lambda-l+1} \partial^h \right. \\
&+ b_{\Lambda,l+2} r_{\mathbf{D}}^2 \left[ x^h (x^{\mathbf{D}})^{\Lambda-l-3} (\Lambda-l-2) + x^h (x^{\mathbf{D}})^{\Lambda-l-2} \partial^{\mathcal{P}} - (x^{\mathbf{D}})^{\Lambda-l-1} \partial^h \right] \\
&+ b_{\Lambda,l+4} r_{\mathbf{D}}^4 \left[ x^h (x^{\mathbf{D}})^{\Lambda-l-5} (\Lambda-l-4) + x^h (x^{\mathbf{D}})^{\Lambda-l-4} \partial^{\mathcal{P}} - (x^{\mathbf{D}})^{\Lambda-l-3} \partial^h \right] + \dots \left. \right\} X_l^{i_1 \dots i_l} \\
&= x^h X_l^{i_1 \dots i_l} \check{N}_{\Lambda,l+1} - l x^{\mathbf{D}} \check{p}_{\Lambda,l} \mathcal{P}_{h j_2 \dots j_l}^{i_1 i_2 \dots i_l} X_{l-1}^{j_2 \dots j_l} = X_{l+1}^{h i_1 \dots i_l} \check{N}_{\Lambda,l+1} - \check{Q}_{\Lambda,l-1} \mathcal{P}_{h j_2 \dots j_l}^{i_1 i_2 \dots i_l} X_{l-1}^{j_2 \dots j_l},
\end{aligned}$$

where  $\check{N}_{\Lambda,l+1}$  is the homogeneous polynomials of degree  $l'-1$  in  $x^{\mathbf{D}}, r_{\mathbf{D}}$

$$\begin{aligned}
\check{N}_{\Lambda,l+1} &:= (x^{\mathbf{D}})^{l'-1} l' + b_{\Lambda,l+2} r_{\mathbf{D}}^2 (x^{\mathbf{D}})^{l'-3} (l'-2) + b_{\Lambda,l+4} r_{\mathbf{D}}^4 (x^{\mathbf{D}})^{l'-5} (l'-4) + \dots \\
&= l' \left[ (x^{\mathbf{D}})^{l'-1} - \frac{(l'-1)(l'-2)}{2(2\Lambda-4+\mathbf{D})} r_{\mathbf{D}}^2 (x^{\mathbf{D}})^{l'-3} + \frac{(l'-1) \dots (l'-4)}{8(2\Lambda-4+\mathbf{D})(2\Lambda-6+\mathbf{D})} r_{\mathbf{D}}^4 (x^{\mathbf{D}})^{l'-5} - \dots \right] \\
&= (\Lambda-l) \check{p}_{\Lambda,l+1} \tag{136}
\end{aligned}$$

(here  $l' \equiv \Lambda-l$ ) and  $\check{Q}_{\Lambda,l-1}$  is the homogeneous polynomial of degree  $l'+1$  in  $x^{\mathbf{D}}, r_{\mathbf{D}}$

$$\begin{aligned}
\check{Q}_{\Lambda,l-1} &:= l x^{\mathbf{D}} \check{p}_{\Lambda,l} - \frac{\zeta_{l+1}}{D} r_{\mathbf{D}}^2 \check{N}_{\Lambda,l+1} = l x^{\mathbf{D}} \check{p}_{\Lambda,l} - \frac{l}{D+2l-2} \left[ r_{\mathbf{D}}^2 - (x^{\mathbf{D}})^2 \right] \check{N}_{\Lambda,l+1} \\
&= (x^{\mathbf{D}})^{l'+1} \left[ \frac{ll'}{D+2l-2} + l \right] - r_{\mathbf{D}}^2 (x^{\mathbf{D}})^{l'-1} ll' \left\{ \frac{1}{D+2l-2} \left[ 1 + \frac{(l'-1)(l'-2)}{2(2\Lambda-4+\mathbf{D})} \right] + \frac{l'-1}{2(2\Lambda-4+\mathbf{D})} \right\}
\end{aligned}$$

$$\begin{aligned}
& +r_{\mathbf{D}}^4 (x^{\mathbf{D}})^{l'-3} \frac{l'(l'-1)(l'-2)}{2(2\Lambda-4+\mathbf{D})} \left\{ \frac{1}{D+2l-2} \left[ 1 + \frac{(l'-3)(l'-4)}{2(2\Lambda-6+\mathbf{D})} \right] + \frac{l'-3}{2(2\Lambda-6+\mathbf{D})} \right\} + \dots \\
& = \frac{l(\Lambda+l+D-2)}{D+2l-2} \tilde{p}_{\Lambda, l-1}; \tag{137}
\end{aligned}$$

(equalities (136-137) are proved by direct calculations), whence (87).

## 8.6 Evaluating a class of radial integrals, and proof of (102)

Given a smooth  $h(r)$  not depending on  $k$ , formula (98) of [22] gives the (asymptotic expansion of) the radial integral of its product with  $g_l(r)g_L(r)$  ( $l, L \in \mathbb{N}_0$ ) at lowest order in  $1/k$ :

$$\int_0^\infty g_L(r)g_l(r)h(r)dr = e^{-\frac{\sqrt{k_l k_L}(\tilde{r}_l - \tilde{r}_L)^2}{2(\sqrt{k_l} + \sqrt{k_L})}} \sum_{n=0}^{+\infty} \frac{h^{(2n)}(\hat{r}_{l,L})}{(2n)!! (\sqrt{k_l} + \sqrt{k_L})^n}. \tag{138}$$

$\tilde{r}_l, k_l$  were defined in (16), while  $\hat{r}_{l,L} := \frac{\sqrt{k_l \tilde{r}_l + \sqrt{k_L \tilde{r}_L}}{\sqrt{k_l} + \sqrt{k_L}}$ ,  $h^{(n)}(r) \equiv d^n h/dr^n$ . Up to  $O(k^{-3/2})$  the exponential in (138) is 1, because by explicit computation

$$\begin{aligned}
\sqrt{k_l} &= \sqrt{2k} \left( 1 + \frac{3b(l,D)}{4k} \right) + O(k^{-\frac{3}{2}}), \quad \tilde{r}_l - \tilde{r}_L = \frac{b(l,D)}{3b(l,D)+2k} - \frac{b(L,D)}{3b(L,D)+2k} = O(k^{-1}), \\
\sqrt{k_l \tilde{r}_l} &= \sqrt{2k} \left( 1 + \frac{3b(l,D)}{4k} \right) \left( 1 + \frac{b(l,D)}{2k} \right) + O(k^{-\frac{5}{2}}) = \sqrt{2k} \left( 1 + \frac{5b(l,D)}{4k} \right) + O(k^{-\frac{3}{2}}), \\
\hat{r}_{l,L} &= \frac{2 + \frac{5b(l,D)}{4k} + \frac{5b(L,D)}{4k}}{2 + \frac{3b(l,D)}{4k} + \frac{3b(L,D)}{4k}} + O(k^{-2}) = 1 + \frac{b(l,D)+b(L,D)}{4k} + O(k^{-2}), \quad \frac{\sqrt{k_l k_L}(\tilde{r}_l - \tilde{r}_L)^2}{2(\sqrt{k_l} + \sqrt{k_L})} = O(k^{-\frac{3}{2}}).
\end{aligned}$$

By (26), (66), applying  $P_\Lambda = \sum_{l=0}^\Lambda P_\Lambda^l$  to  $x^i \psi^{i_1 i_2 \dots i_l} = t^i T_l^{i_1 i_2 \dots i_l} r f_l(r)$  we find

$$\begin{aligned}
P_\Lambda (x^i \psi_l^{i_1 i_2 \dots i_l}) &\stackrel{(66)}{=} \tilde{P}_{l+1} [T_{l+1}^{i i_1 \dots i_l} r f_l(r)] + \tilde{P}_{l-1} \left[ \frac{\zeta_{l+1}}{D} g^{ih} \mathcal{P}_{h j_2 \dots j_l}^{l i_1 i_2 \dots i_l} T_{l-1}^{j_2 \dots j_l} r f_l(r) \right] \\
&\stackrel{(26)}{=} c_{l+1} T_{l+1}^{i i_1 \dots i_l} f_{l+1}(r) + c_l \frac{\zeta_{l+1}}{D} \mathcal{P}_{i j_2 \dots j_l}^{l i_1 i_2 \dots i_l} T_{l-1}^{j_2 \dots j_l} f_{l-1}(r) \\
&\stackrel{(98)}{=} c_{l+1} \psi_{l+1}^{i i_1 \dots i_l} + c_l \frac{\zeta_{l+1}}{D} \mathcal{P}_{i j_2 \dots j_l}^{l i_1 i_2 \dots i_l} \psi_{l-1}^{j_2 \dots j_l},
\end{aligned}$$

$$\text{where } c_l := \begin{cases} 0 & \text{if } l = 0, \Lambda, \\ \rho_{l-1,l} := \int_0^\infty r^d dr f_{l-1}^*(r) r f_l(r) = \int_0^\infty dr g_{l-1}(r) g_l(r) r & \text{if } 0 < l < \Lambda. \end{cases}$$

The last integral is of the type (138) with  $L = l-1$  and  $h(r) = r$ . By explicit computation,

$$\begin{aligned}
\rho_{l-1,l} &= 1 + \frac{D^2 - 4D + 3}{k} + \frac{l(l+D-2) + (l+1)(l-1+D)}{2k} + O(k^{-\frac{3}{2}}) \\
&= \sqrt{1 + \frac{(2D-5)(D-1)}{2k} + \frac{(l-1)(l+D-2)}{k}} + O(k^{-\frac{3}{2}}). \tag{139}
\end{aligned}$$

By setting the  $O(k^{-3/2})$  term equal to zero we finally arrive at (101-102).



## 8.7 Proof of Proposition 4.1

The equations in (103), (104), (105) follow from (9), (10), (11) and  $[L_{ij}, P_\Lambda] = 0$ .

By (33-34),  $\psi_{\Lambda, \pm}^{hk} := T_{\Lambda, \pm}^{hk} f_\Lambda(r) \in \mathcal{H}_\Lambda^\Lambda$  are eigenvectors with the highest and lowest eigenvalues,  $\pm\Lambda$ . Given a basis  $\{\psi_m^\alpha\}$  of eigenvectors of  $L_{hk}$ ,  $L_{hk}\psi_m^\alpha = m\psi_m^\alpha$  with  $m \in \{-\Lambda, 1-\Lambda, \dots, \Lambda\}$  ( $\alpha$  are some extra labels), by (32) we find that for all  $m, \alpha$   $(\bar{x}^h + i\bar{x}^k)^j \psi_m^\alpha$  is either zero or an eigenvector of  $L_{hk}$  with eigenvalue  $j+m$ , which must be  $\leq \Lambda$ . Similarly,  $(L_{hj} \pm iL_{kj})^j \psi_m^\alpha$  is either zero or an eigenvector of  $L_{hk}$  with eigenvalue  $j+m$ , which must be  $\leq \Lambda$ . Therefore for  $j > 2\Lambda$  such vectors must be zero, and we obtain (106).

Applying (102) twice we find

$$\begin{aligned}
\bar{x}^i \bar{x}^j \psi_l^{i_1 i_2 \dots i_l} &\stackrel{(102)}{=} \bar{x}^i \left( c_{l+1} \psi_{l+1}^{j i_1 \dots i_l} + c_l \frac{\zeta_{l+1}}{D} g^{jh} \mathcal{P}_{hj_2 \dots j_l}^{l i_1 i_2 \dots i_l} \psi_{l-1}^{j_2 \dots j_l} \right) \\
&\stackrel{(102)}{=} c_l \frac{\zeta_{l+1}}{D} g^{jh} \mathcal{P}_{hj_2 \dots j_l}^{l i_1 i_2 \dots i_l} \left( c_l \psi_l^{i j_2 \dots j_l} + c_{l-1} \frac{\zeta_l}{D} g^{ik} \mathcal{P}_{kk_3 \dots k_l}^{l-1 j_2 \dots j_l} \psi_{l-2}^{k_3 \dots k_l} \right) \\
&\quad + c_{l+1} \left( c_{l+2} \psi_{l+2}^{i j i_1 \dots i_l} + c_{l+1} \frac{\zeta_{l+2}}{D} g^{ih} \mathcal{P}_{hj_1 \dots j_l}^{l+1 j i_1 \dots i_l} \psi_l^{j_1 \dots j_l} \right) \\
&= c_{l+1} c_{l+2} \psi_{l+2}^{i j i_1 \dots i_l} + c_l c_{l-1} \frac{\zeta_{l+1}}{D} \frac{\zeta_l}{D} g^{jh} g^{ik} \mathcal{P}_{hk k_3 \dots k_l}^{l i_1 i_2 i_3 \dots i_l} \psi_{l-2}^{k_3 \dots k_l} \\
&\quad + (c_{l+1})^2 \frac{\zeta_{l+2}}{D} g^{ih} \mathcal{P}_{hj_1 \dots j_l}^{l+1 j i_1 \dots i_l} \psi_l^{j_1 \dots j_l} + (c_l)^2 \frac{\zeta_{l+1}}{D} g^{jh} \mathcal{P}_{hj_2 \dots j_l}^{l i_1 i_2 \dots i_l} \psi_l^{i j_2 \dots j_l} \quad (140)
\end{aligned}$$

Taking the difference of (140) and (140) with  $i, j$  exchanged we find

$$\begin{aligned}
[\bar{x}^i, \bar{x}^j] \psi_l^{i_1 i_2 \dots i_l} &= \bar{x}^i \bar{x}^j \psi_l^{i_1 i_2 \dots i_l} - (i \leftrightarrow j) \\
&\stackrel{(140)}{=} (c_{l+1})^2 \frac{\zeta_{l+2}}{D} (g^{ih} \mathcal{P}_{hj_1 \dots j_l}^{l+1 j i_1 \dots i_l} - g^{jh} \mathcal{P}_{hj_1 \dots j_l}^{l+1 i i_1 \dots i_l}) \psi_l^{j_1 \dots j_l} \\
&\quad + (c_l)^2 \frac{\zeta_{l+1}}{D} (g^{jh} \mathcal{P}_{hj_2 \dots j_l}^{l i_1 i_2 \dots i_l} \psi_l^{i j_2 \dots j_l} - g^{ih} \mathcal{P}_{hj_2 \dots j_l}^{l i_1 i_2 \dots i_l} \psi_l^{j j_2 \dots j_l}) \\
&\stackrel{(63)}{=} \left[ (c_l)^2 \frac{\zeta_{l+1}}{Dl} - (c_{l+1})^2 \frac{\zeta_{l+2}}{Dl} \frac{\zeta_{l+1}}{\zeta_{l+2}} \right] i L^{ij} \psi_l^{i_1 \dots i_l} \\
&\stackrel{(60)}{=} \frac{(c_{l+1})^2 - (c_l)^2}{D+2l-2} i L^{ij} \psi_l^{i_1 \dots i_l} \\
&\stackrel{(139)}{=} i L^{ij} \psi_l^{i_1 \dots i_l} \times \begin{cases} -\frac{1}{k} & \text{if } l < \Lambda \\ \frac{(\rho_\Lambda - 1, \Lambda)^2}{D+2\Lambda-2} & \text{if } l = \Lambda, \end{cases}
\end{aligned}$$

up to order  $O(k^{-3/2})$ , whence (107), using (139). Contracting the indices  $ij$  of (140) we find how  $\bar{x}^2 := \bar{x}^i \bar{x}_i$  acts on the  $\mathcal{H}_\Lambda$  basis elements:

$$\begin{aligned}
\bar{x}^2 \psi_l^{i_1 i_2 \dots i_l} &= g_{ij} \bar{x}^i \bar{x}^j \psi_l^{i_1 i_2 \dots i_l} \\
&\stackrel{(140)}{=} c_{l+1} c_{l+2} g_{ij} \psi_{l+2}^{i j i_1 \dots i_l} + c_l c_{l-1} \frac{\zeta_{l+1}}{D} \frac{\zeta_l}{D} g^{hk} \mathcal{P}_{hk k_3 \dots k_l}^{l i_1 i_2 i_3 \dots i_l} \psi_{l-2}^{k_3 \dots k_l} \\
&\quad + (c_{l+1})^2 \frac{\zeta_{l+2}}{D} \mathcal{P}_{hj_1 \dots j_l}^{l+1 h i_1 \dots i_l} \psi_l^{j_1 \dots j_l} + (c_l)^2 \frac{\zeta_{l+1}}{D} \mathcal{P}_{hj_2 \dots j_l}^{l i_1 i_2 \dots i_l} \psi_l^{h j_2 \dots j_l}
\end{aligned}$$

$$\begin{aligned}
&\stackrel{(56)}{=} (c_{l+1})^2 \frac{\zeta_{l+2}}{D} \frac{1}{l+1} \left[ D + l - \frac{2l}{D+2l-2} \right] \psi_l^{i_1 i_2 \dots i_l} + (c_l)^2 \frac{\zeta_{l+1}}{D} \psi_l^{i_1 i_2 \dots i_l} \\
&\stackrel{(60)}{=} \left\{ (c_{l+1})^2 \left[ 1 - \frac{l}{D+2l-2} \right] + (c_l)^2 \frac{l}{D+2l-2} \right\} \psi_l^{i_1 i_2 \dots i_l} \tag{141}
\end{aligned}$$

up to order  $O(k^{-3/2})$ . We find (108), noting that up to order  $O(k^{-3/2})$  (141) becomes

$$\begin{aligned}
\bar{x}^2 \psi_\Lambda^{i_1 i_2 \dots i_\Lambda} &\stackrel{(101)}{=} \frac{(c_\Lambda)^2 \Lambda}{D+2\Lambda-2} \psi_\Lambda^{i_1 i_2 \dots i_\Lambda} \\
&\stackrel{(139)}{=} \frac{\Lambda}{D+2\Lambda-2} \left\{ 1 + \frac{(2D-5)(D-1)}{2k} + \frac{(\Lambda-1)(\Lambda+D-2)}{k} \right\} \psi_l^{i_1 i_2 \dots i_l} \\
\bar{x}^2 \psi_l^{i_1 i_2 \dots i_l} &= \left\{ (c_{l+1})^2 + \frac{l[(c_l)^2 - (c_{l+1})^2]}{D+2l-2} \right\} \psi_l^{i_1 i_2 \dots i_l} \\
&\stackrel{(139)}{=} \left\{ 1 + \frac{(2D-5)(D-1)}{2k} + \frac{E_l}{k} \right\} \psi_l^{i_1 i_2 \dots i_l}, \quad \text{if } l < \Lambda.
\end{aligned}$$

## 8.8 Proof of Proposition 5.1

We first show that indeed  $\mathcal{A}_\Lambda$  is generated by the  $\bar{L}_{hi}, \bar{x}^i$ . Since the action of  $so(D)$  (which is spanned by the  $\bar{L}_{hi}$ ) is transitive on each irreducible component  $\mathcal{H}_\Lambda^l \simeq V_D^l$  ( $l \in \{0, 1, \dots, \Lambda\}$ ) contained in  $\mathcal{H}_\Lambda$ , it remains to show that some  $\psi_l \in \mathcal{H}_\Lambda^l$  can be mapped into some element of  $\mathcal{H}_\Lambda^m$  for all  $m \neq l$  by applying polynomials in  $\bar{x}^i$ . For all  $\psi_l \in \mathcal{H}_\Lambda^l$  we have  $\bar{x}^i \psi_l \in \mathcal{H}_\Lambda^{l+1} \oplus \mathcal{H}_\Lambda^{l-1}$ ; if  $l = \Lambda$  then  $\bar{x}^i \psi_l \in \mathcal{H}_\Lambda^{\Lambda-1}$  automatically; if  $l < \Lambda$  one can map  $\mathcal{H}_\Lambda^l$  into  $\mathcal{H}_\Lambda^{l-1}$  by the contracted multiplication of (67), while applying  $(\bar{x}^1 + i\bar{x}^2) \cdot$  to  $(t^1 + it^2)^l f_l(r) \in \mathcal{H}_\Lambda^l$  one obtains a vector proportional to  $(t^1 + it^2)^{l+1} f_{l+1}(r) \in \mathcal{H}_\Lambda^{l+1}$ .

The Ansatz (111) with generic coefficients  $a_{\Lambda,l}$  is manifestly  $Uso(D)$ -equivariant, i.e. fulfills (110) for all  $a = L_{hi}$ ; it is also invariant under permutations of  $(i_1 \dots i_l)$  and fulfills relations (99) (both sides give zero when contracted with any  $\delta_{i_a i_b}$ ). More explicitly, (111) read

$$\begin{aligned}
f_0 &\stackrel{\varkappa_\Lambda}{\mapsto} a_{\Lambda,0} F_{\mathbf{D},\Lambda} = a_{\Lambda,0} p_{\Lambda,0} = a_{\Lambda,0} (t^{\mathbf{D}})^\Lambda, \\
t^i f_1 = T_1^i f_1 &\stackrel{\varkappa_\Lambda}{\mapsto} a_{\Lambda,1} F_{\mathbf{D},\Lambda}^i = a_{\Lambda,1} p_{\Lambda,1} T_1^i = a_{\Lambda,1} (t^{\mathbf{D}})^{\Lambda-1} t^i, \\
T_2^{ij} f_2 = \left( t^i t^j - \frac{\delta^{ij}}{D} \right) f_2 &\stackrel{\varkappa_\Lambda}{\mapsto} a_{\Lambda,2} F_{\mathbf{D},\Lambda}^{ij} = a_{\Lambda,2} p_{\Lambda,2} T_2^{ij}, \\
\dots &\stackrel{\varkappa_\Lambda}{\mapsto} \dots
\end{aligned}$$

Similarly, the Ansatz (112) with a generic function  $m_\Lambda(s)$  of a real nonnegative variable  $s$  is manifestly  $Uso(D)$ -equivariant, and by (91), (92) we find

$$m_\Lambda^*(\lambda) L_{\mathbf{D}h} m_\Lambda(\lambda) F_{\mathbf{D},\Lambda}^{i_1 \dots i_l} = i\mu_\Lambda(l)(\Lambda-l) F_{\mathbf{D},\Lambda}^{hi_1 \dots i_l} - i\mu_\Lambda^*(l-1) \frac{l(\Lambda+l+D-2)}{D+2l-2} \mathcal{P}_{hj_2 \dots j_l}^{li_1 i_2 \dots i_l} F_{\mathbf{D},\Lambda}^{j_2 \dots j_l} \tag{142}$$

where we have abbreviated  $\mu_\Lambda(l) \equiv m_\Lambda(l) m_\Lambda^*(l+1)$ . We determine the unknown  $m_\Lambda, a_{\Lambda,l}$  requiring (110) for  $a = \bar{x}^h \cdot$ . Applying  $\varkappa_\Lambda$  to eq. (102) with  $l < \Lambda$ , and imposing (110) we

obtain

$$c_{l+1} a_{\Lambda, l+1} F_{\mathbf{D}, \Lambda}^{hi_1 \dots i_l} + c_l a_{\Lambda, l-1} \frac{l}{D+2l-2} \mathcal{P}_{hj_2 \dots j_l}^{li_1 i_2 \dots i_l} F_{\mathbf{D}, \Lambda}^{j_2 \dots j_l} = m_{\Lambda}^*(\lambda) L_{\mathbf{D}h} m_{\Lambda}(\lambda) a_{\Lambda, l} F_{\mathbf{D}, \Lambda}^{i_1 \dots i_l}$$

$$\stackrel{(142)}{=} ia_{\Lambda, l} \left[ \mu_{\Lambda}(l) (\Lambda - l) F_{\mathbf{D}, \Lambda}^{hi_1 \dots i_l} - \mu_{\Lambda}^*(l-1) \frac{l(\Lambda + l + D - 2)}{D + 2l - 2} \mathcal{P}_{hj_2 \dots j_l}^{li_1 i_2 \dots i_l} F_{\mathbf{D}, \Lambda}^{j_2 \dots j_l} \right],$$

which implies the recursion relations for the coefficients  $a_{\Lambda, l}$

$$c_{l+1} a_{\Lambda, l+1} = i(\Lambda - l) a_{\Lambda, l} \mu_{\Lambda}(l), \quad c_l a_{\Lambda, l-1} = -i(\Lambda + l + D - 2) a_{\Lambda, l} \mu_{\Lambda}^*(l-1). \quad (143)$$

Multiplying (144a) by (144b) <sub>$l \rightarrow l+1$</sub> , i.e. by  $c_{l+1} a_{\Lambda, l} = -i a_{\Lambda, l+1} \mu_{\Lambda}^*(l) (\Lambda + l + D - 1)$ , we find

$$c_{l+1}^2 a_{\Lambda, l} a_{\Lambda, l+1} = a_{\Lambda, l+1} \mu_{\Lambda}^*(l) a_{\Lambda, l} \mu_{\Lambda}(l) (\Lambda - l) (\Lambda + l + D - 1)$$

$$\Rightarrow |\mu_{\Lambda}(l)|^2 (\Lambda - l) (\Lambda + l + D - 1) = c_{l+1}^2,$$

implying

$$m_{\Lambda}^*(l+1) m_{\Lambda}(l) = \mu_{\Lambda}(l) = \frac{\epsilon_{\Lambda, l} c_{l+1}}{\sqrt{(\Lambda - l)(\Lambda + l + D - 1)}}, \quad a_{\Lambda, l+1} = ia_{\Lambda, l} \epsilon_{\Lambda, l} \sqrt{\frac{\Lambda - l}{(\Lambda + l + D - 1)}} \quad (144)$$

where  $\epsilon_{\Lambda, l}$  are arbitrary phase factors. Choosing  $\epsilon_{\Lambda, l} = 1$ ,  $m_{\Lambda}(s)$  positive-definite on  $s \in \mathbb{R}^+$ , and using the property  $\Gamma(z+1) = z\Gamma(z)$  of the Euler gamma function we find that (144) are solved by (114). Applying  $\varkappa_{\Lambda}$  to eq. (102) with  $l = \Lambda$ , and imposing (110) we obtain

$$c_{\Lambda} \frac{\Lambda a_{\Lambda, \Lambda-1}}{D+2\Lambda-2} \mathcal{P}_{hj_2 \dots j_{\Lambda}}^{\Lambda i_1 i_2 \dots i_{\Lambda}} F_{\mathbf{D}, \Lambda}^{j_2 \dots j_{\Lambda}} = m_{\Lambda}^*(\lambda) L_{\mathbf{D}h} m_{\Lambda}(\lambda) a_{\Lambda, \Lambda} F_{\mathbf{D}, \Lambda}^{i_1 \dots i_{\Lambda}}$$

$$\stackrel{(142)}{=} -i a_{\Lambda, \Lambda} \mu_{\Lambda}^*(\Lambda - 1) \frac{\Lambda(2\Lambda + D - 2)}{D + 2\Lambda - 2} \mathcal{P}_{hj_2 \dots j_{\Lambda}}^{\Lambda i_1 i_2 \dots i_{\Lambda}} F_{\mathbf{D}, \Lambda}^{j_2 \dots j_{\Lambda}},$$

which also is satisfied by (114).

## 8.9 Proof of Theorem 6.1

The proof is recursive. By (117)  $\widehat{T}_{l+1}^{h_1 \dots h_{l+1}} = \mathcal{P}^{l+1}_{i_1 \dots i_b}^{h_1 \dots h_{l+1}} \bar{x}^b \widehat{T}_l^{i_1 \dots i_l}$ , which applied to  $T_m^{j_1 \dots j_m}$  gives

$$\widehat{T}_{l+1}^{h_1 \dots h_{l+1}} T_m^{j_1 \dots j_m} = \mathcal{P}^{l+1}_{i_1 \dots i_b}^{h_1 \dots h_{l+1}} \bar{x}^b \widehat{T}_l^{i_1 \dots i_l} T_m^{j_1 \dots j_m} \stackrel{(120)}{=} \sum_{n \in L^{lm}} \mathcal{P}^{l+1}_{i_1 \dots i_b}^{h_1 \dots h_{l+1}} \widehat{N}_n^{lm} V_{b_1 \dots b_n}^{i_1 \dots i_l, j_1 \dots j_m} \bar{x}^b T_n^{b_1 \dots b_n}$$

$$\stackrel{(119)}{=} \sum_{n \in L^{lm}} \widehat{N}_n^{lm} \mathcal{P}^{l+1}_{i_1 \dots i_b}^{h_1 \dots h_{l+1}} \mathcal{P}_{a_1 \dots a_r b_1 \dots b_{l-r}}^{i_1 \dots i_r i_{r+1} \dots i_l} \mathcal{P}_{a_1 \dots a_r b_{l-r+1} \dots b_n}^{j_1 \dots j_r j_{r+1} \dots j_m} (c_{n+1} T_{n+1}^{b_1 \dots b_n} + d_n c_n \mathcal{P}_{bk_2 \dots k_n}^{n b_1 \dots b_n} T_{n-1}^{k_2 \dots k_n})$$

$$= \sum_{n \in L^{lm}} \widehat{N}_n^{lm} \left[ \mathcal{P}_{a_1 \dots a_r b_1 \dots b_{l-r}}^{l+1 h_1 \dots h_r h_{r+1} \dots h_{l+1}} \mathcal{P}_{a_1 \dots a_r b_{l-r+1} \dots b_n}^{j_1 \dots j_r j_{r+1} \dots j_m} (c_{n+1} T_{n+1}^{b_1 \dots b_n} + d_n c_n \mathcal{P}_{bk_2 \dots k_n}^{n b_1 \dots b_n} T_{n-1}^{k_2 \dots k_n}) \right] \quad (145)$$

Now we note that

$$\begin{aligned}
& \mathcal{P}^{l+1 h_1 \dots h_r h_{r+1} \dots h_{l+1}}_{a_1 \dots a_r b_1 \dots b_{l-r} b} \mathcal{P}^{n b_1 b_2 \dots b_n}_{b k_2 \dots k_n} T_{n-1}^{k_2 \dots k_n} = \\
& \mathcal{P}^{l+1 h_1 \dots h_r h_{r+1} \dots h_{l+1}}_{a_1 \dots a_r b_1 \dots b_{l-r} b} \mathcal{P}^{n-1 b_2 b_3 \dots b_n}_{h k_3 \dots k_n} \frac{1}{n} \left[ \delta_b^{b_1} \delta_{k_2}^h + (n-1) \delta_{k_2}^{b_1} \delta_b^h - \frac{2(n-1)}{D+2n-4} \delta^{b_1 h} \delta_{b k_2} \right] T_{n-1}^{k_2 \dots k_n} \\
& = \frac{n-1}{n} \mathcal{P}^{l+1 h_1 \dots h_r h_{r+1} \dots h_{l+1}}_{a_1 \dots a_r b_1 \dots b_{l-r} b} \left[ \mathcal{P}^{n-1 b_2 b_3 \dots b_n}_{b k_3 \dots k_n} T_{n-1}^{b_1 k_3 \dots k_n} - \frac{2}{D+2n-4} \mathcal{P}^{n-1 b_2 b_3 \dots b_n}_{b_1 k_3 \dots k_n} T_{n-1}^{b k_3 \dots k_n} \right] \\
& = \frac{(n-1)(D+2n-6)}{n(D+2n-4)} \mathcal{P}^{l+1 h_1 \dots h_r h_{r+1} \dots h_{l+1}}_{a_1 \dots a_r b_1 \dots b_{l-r} b} \mathcal{P}^{n-1 b_2 b_3 \dots b_n}_{b k_3 \dots k_n} T_{n-1}^{b_1 k_3 \dots k_n},
\end{aligned}$$

for the last equality we have used the symmetry of  $\mathcal{P}^{l+1 h_1 \dots h_r h_{r+1} \dots h_{l+1}}_{a_1 \dots a_r b_1 \dots b_{l-r} b}$  under the exchange  $b \leftrightarrow b_1$ . Iterating the procedure we find

$$\begin{aligned}
& \mathcal{P}^{l+1 h_1 \dots h_r h_{r+1} \dots h_{l+1}}_{a_1 \dots a_r b_1 \dots b_{l-r} b} \mathcal{P}^{n b_1 b_2 \dots b_n}_{b k_2 \dots k_n} T_{n-1}^{k_2 \dots k_n} = \frac{(n-2)(D+2n-8)}{n(D+2n-4)} \mathcal{P}^{l+1 h_1 \dots h_r h_{r+1} \dots h_{l+1}}_{a_1 \dots a_r b_1 \dots b_{l-r} b} \mathcal{P}^{n-2 b_3 b_4 \dots b_n}_{b k_4 \dots k_n} T_{n-1}^{b_1 b_2 k_4 \dots k_n} \\
& = \dots = \frac{(n-l+r)[D+2(n-l+r)-4]}{n(D+2n-4)} \mathcal{P}^{l+1 h_1 \dots h_r h_{r+1} \dots h_{l+1}}_{a_1 \dots a_r b_1 \dots b_{l-r} b} \mathcal{P}^{n-l+r b_{l-r+1} \dots b_n}_{b k_{l-r+2} \dots k_n} T_{n-1}^{b_1 \dots b_{l-r} k_{l-r+2} \dots k_n},
\end{aligned}$$

which replaced in (145) gives

$$\begin{aligned}
& \widehat{T}_{l+1}^{h_1 \dots h_{l+1}} T_m^{j_1 \dots j_m} = \sum_{n \in L^{lm}} \widehat{N}_n^{lm} \left[ \mathcal{P}^{l+1 h_1 \dots h_r h_{r+1} \dots h_{l+1}}_{a_1 \dots a_r b_1 \dots b_{l-r} b} \mathcal{P}^{m j_1 \dots j_r j_{r+1} \dots j_m}_{a_1 \dots a_r b_{l-r+1} \dots b_n} (c_{n+1} T_{n+1}^{b_1 \dots b_n b} \right. \\
& \quad \left. + \frac{(n-l+r)[D+2(n-l+r)-4]}{n(D+2n-4)} \frac{n c_n}{D+2n-2} \mathcal{P}^{n-l+r b_{l-r+1} \dots b_n}_{b k_{l-r+2} \dots k_n} T_{n-1}^{b_1 \dots b_{l-r} k_{l-r+2} \dots k_n} \right) \\
& = \sum_{n \in L^{lm}} \widehat{N}_n^{lm} \left[ c_{n+1} \mathcal{P}^{l+1 h_1 \dots h_r h_{r+1} \dots h_{l+1}}_{a_1 \dots a_r b_1 \dots b_{l-r} b} \mathcal{P}^{m j_1 \dots j_r j_{r+1} \dots j_m}_{a_1 \dots a_r b_{l-r+1} \dots b_n} T_{n+1}^{b_1 \dots b_n b} \right. \\
& \quad \left. + \frac{(n-l+m)[D+n-l+m-4]}{2(D+2n-2)(D+2n-4)} c_n \mathcal{P}^{l+1 h_1 \dots h_r h_{r+1} \dots h_{l+1}}_{a_1 \dots a_r a_{r+1} b_1 \dots b_{l-r}} \mathcal{P}^{m j_1 \dots j_r j_{r+1} \dots j_m}_{a_1 \dots a_r a_{r+1} k_{l-r+2} \dots k_n} T_{n-1}^{b_1 \dots b_{l-r} k_{l-r+2} \dots k_n} \right];
\end{aligned}$$

renaming  $n' \equiv n-1$  in the first sum,  $n' \equiv n+1$  in the second, this becomes

$$\begin{aligned}
& \widehat{T}_{l+1}^{h_1 \dots h_{l+1}} T_m^{j_1 \dots j_m} = \sum_{n' \in L^{lm+1}} \widehat{N}_{n'-1}^{lm} c_{n'} \mathcal{P}^{l+1 h_1 \dots h_r h_{r'+1} \dots h_{l+1}}_{a_1 \dots a_r b b_1 \dots b_{l-r'}} \mathcal{P}^{m j_1 \dots j_r' j_{r'+1} \dots j_m}_{a_1 \dots a_r' b_{l-r'+1} \dots b_{n'-1}} T_{n'}^{b b_1 \dots b_{n'-1}} \\
& \quad + \sum_{n' \in L^{lm-1}} \widehat{N}_{n'+1}^{lm} c_{n'+1} \frac{(n'-l+m+1)[D+n'-l+m-3]}{2(D+2n')(D+2n'-2)} \\
& \quad \mathcal{P}^{l+1 h_1 \dots h_r' \dots h_{l+1}}_{a_1 \dots a_r' b_1 \dots b_{l-r'+1}} \mathcal{P}^{m j_1 \dots j_r' \dots j_m}_{a_1 \dots a_r' k_{l-r'+3} \dots k_{n'+1}} T_{n'}^{b_1 \dots b_{l-r'+1} k_{l-r'+3} \dots k_{n'+1}} \\
& = \sum_{n' \in L^{(l+1)m}} \widehat{N}_{n'}^{(l+1)m} \mathcal{P}^{l+1 h_1 \dots h_r' \dots h_{l+1}}_{a_1 \dots a_r' b_1 \dots b_{l-r'+1}} \mathcal{P}^{m j_1 \dots j_r' j_{r'+1} \dots j_m}_{a_1 \dots a_r' b_{l-r'+2} \dots b_{n'}} T_{n'}^{b_1 \dots b_{n'}}
\end{aligned}$$

where  $r' := \frac{l+1+m-n'}{2}$ , we have used that  $2(n-l+r) = n-l+m$  and we have set

$$\widehat{N}_n^{(l+1)m} = c_n \widehat{N}_{n-1}^{lm} + c_{n+1} \widehat{N}_{n+1}^{lm} \frac{(n-l+m+1)[D+n-l+m-3]}{2(D+2n)(D+2n-2)} \quad (146)$$

for all  $n \in L^{(l+1)m}$ . The relation among  $N_n^{(l+1)m}, N_{n-1}^{lm}, N_{n+1}^{lm}$  is obtained replacing  $c_h \mapsto 1$  and removing  $\hat{\cdot}$ . Since  $c_h \leq c_\Lambda$ , by the induction hypothesis  $(121)_l$  we find  $(121)_{l+1}$ , as claimed:

$$\begin{aligned} \widehat{N}_n^{(l+1)m} &\leq c_\Lambda \left[ \widehat{N}_{n-1}^{lm} + \widehat{N}_{n+1}^{lm} \frac{(n-l+m+1)[D+n-l+m-3]}{2(D+2n)(D+2n-2)} \right] \\ &\leq (c_\Lambda)^{l+1} \left[ N_{n-1}^{lm} + N_{n+1}^{lm} \frac{(n-l+m+1)[D+n-l+m-3]}{2(D+2n)(D+2n-2)} \right] = (c_\Lambda)^{l+1} N_n^{(l+1)m}. \end{aligned}$$

## 8.10 Proof of Theorem 6.2

Setting  $\phi_\Lambda^\perp := \phi - \phi_\Lambda$  we find

$$\begin{aligned} (f - \widehat{f}_{2\Lambda})\phi &= (f - f_{2\Lambda})\phi + (f_{2\Lambda} - \widehat{f}_{2\Lambda})\phi \\ &= (f - f_{2\Lambda})\phi + (f_{2\Lambda} - \widehat{f}_{2\Lambda})\phi_\Lambda^\perp + (f_{2\Lambda} - \widehat{f}_{2\Lambda})\phi_\Lambda, \end{aligned}$$

$$\begin{aligned} \left\| (f - \widehat{f}_{2\Lambda})\phi \right\| &\leq \left\| (f - f_{2\Lambda})\phi \right\| + \left\| (f_{2\Lambda} - \widehat{f}_{2\Lambda})\phi_\Lambda^\perp \right\| + \left\| (f_{2\Lambda} - \widehat{f}_{2\Lambda})\phi_\Lambda \right\| \\ &\leq \|f - f_{2\Lambda}\|_{op} \|\phi\| + \left\| f_{2\Lambda} - \widehat{f}_{2\Lambda} \right\|_{op} \|\phi_\Lambda^\perp\| + \left\| (f_{2\Lambda} - \widehat{f}_{2\Lambda})\phi_\Lambda \right\| \quad (147) \end{aligned}$$

Since  $f \in C(S^d)$ , then  $\|f_{2\Lambda} - f\|_{op} = \|f_{2\Lambda} - f\|_\infty \equiv \sup_{p \in S^d} |f_{2\Lambda}(p) - f(p)|$ , which goes to zero because  $f_{2\Lambda} \rightarrow f$  *uniformly* over  $S^d$  (in fact,  $f_{2\Lambda} - f$  is a continuous function over the compact manifold  $S^d$ , and therefore by Heine-Cantor theorem is also uniformly continuous); therefore the first term at the rhs(147) goes to zero as  $\Lambda \rightarrow \infty$ . As  $\Lambda \rightarrow \infty$  also the second term at the rhs(147) goes to zero because so does  $\|\phi_\Lambda^\perp\| = \|\phi - \phi_\Lambda\|$ , while  $\|f_{2\Lambda} - \widehat{f}_{2\Lambda}\|_{op}$  certainly is bounded (actually one can easily show that this goes to zero as well).

To show that the lhs(147) goes to zero as  $\Lambda \rightarrow \infty$  we now show that the last term at the rhs does, as well. Using the decompositions (69)-(70) for  $\phi_\Lambda, \widehat{f}_{2\Lambda}, f_{2\Lambda}$  we can write

$$\begin{aligned} (\widehat{f}_{2\Lambda} - f_{2\Lambda})\phi_\Lambda &= \sum_{l=0}^{2\Lambda} f_{i_1 \dots i_l}^l (\widehat{T}_l^{i_1 \dots i_l} - T_l^{i_1 \dots i_l}) \sum_{m=0}^{\Lambda} \phi_{j_1 \dots j_m}^m T_m^{j_1 \dots j_m} \\ &\stackrel{(120), (78)}{=} \sum_{l=0}^{2\Lambda} \sum_{m=0}^{\Lambda} f_{i_1 \dots i_l}^l \phi_{j_1 \dots j_m}^m \sum_{n \in L^{lm}} (\widehat{N}_n^{lm} - N_n^{lm}) \mathcal{P}_{a_1 \dots a_r c_1 \dots c_{l-r}}^{m j_1 \dots j_r j_{r+1} \dots j_m} \mathcal{P}_{a_1 \dots a_r c_{l-r+1} \dots c_n}^{n k_1 \dots k_n} T_n^{k_1 \dots k_n} \\ &= \sum_{l=0}^{2\Lambda} \sum_{m=0}^{\Lambda} \sum_{n \in L^{lm}} f_{a_1 \dots a_r c_1 \dots c_{l-r}}^l \phi_{a_1 \dots a_r c_{l-r+1} \dots c_n}^m (\widehat{N}_n^{lm} - N_n^{lm}) T_n^{c_1 \dots c_n} \end{aligned}$$

and taking the square norm in  $\mathcal{H}_s$ ,

$$\begin{aligned} \left\| (\widehat{f}_{2\Lambda} - f_{2\Lambda})\phi_\Lambda \right\|^2 &= \sum_{l, l'=0}^{2\Lambda} \sum_{m, m'=0}^{\Lambda} \sum_{n \in L^{lm}} \sum_{n' \in L^{l'm'}} (\widehat{N}_n^{lm} - N_n^{lm}) (\widehat{N}_{n'}^{l'm'} - N_{n'}^{l'm'}) \\ &\quad (f_{a_1 \dots a_r c_1 \dots c_{l-r}}^l \phi_{a_1 \dots a_r c_{l-r+1} \dots c_n}^m)^* f_{b_1 \dots b_r d_1 \dots d_{l'-r}}^{l'} \phi_{b_1 \dots b_r d_{l'-r+1} \dots d_{n'}}^{m'} \langle T_n^{c_1 \dots c_n}, T_{n'}^{d_1 \dots d_n} \rangle \end{aligned}$$

$$\begin{aligned}
&= \sum_{l,l'=0}^{2\Lambda} \sum_{m,m'=0}^{\Lambda} \sum_{n \in L^{lm}} \sum_{n' \in L^{l'm'}} (\widehat{N}_n^{lm} - N_n^{lm}) (\widehat{N}_{n'}^{l'm'} - N_{n'}^{l'm'}) \\
&\quad (f_{a_1 \dots a_r c_1 \dots c_{l-r}}^l \phi_{a_1 \dots a_r c_{l-r+1} \dots c_n}^m)^* f_{b_1 \dots b_{r'} d_1 \dots d_{l'-r'}}^{l'} \phi_{b_1 \dots b_{r'} d_{l'-r'+1} \dots d_{n'}}^{m'} Q_n \delta_{nm'} \mathcal{P}_{c_1 \dots c_n}^{nd_1 \dots d_n} \\
&= \sum_{l,l'=0}^{2\Lambda} \sum_{m,m'=0}^{\Lambda} \sum_{n \in L^{lm} \cap L^{l'm'}} (\widehat{N}_n^{lm} - N_n^{lm}) (\widehat{N}_n^{l'm'} - N_n^{l'm'}) Q_n \\
&\quad (f_{a_1 \dots a_r c_1 \dots c_{l-r}}^l \phi_{a_1 \dots a_r c_{l-r+1} \dots c_n}^m)^* f_{b_1 \dots b_{r'} c_1 \dots c_{l'-r'}}^{l'} \phi_{b_1 \dots b_{r'} c_{l'-r'+1} \dots c_n}^{m'} \tag{148}
\end{aligned}$$

By (101), (109) it is  $(c_\Lambda)^l \leq (1 + \epsilon)^{l/2} < e^{\epsilon/2} \leq e^{\Lambda\epsilon}$ , because we are using  $l \leq 2\Lambda$ ; hence by Theorem 6.1 we find

$$0 \leq (\widehat{N}_n^{lm} - N_n^{lm}) \leq N_n^{lm} [(c_\Lambda)^l - 1] < N_n^{lm} (e^{\Lambda\epsilon} - 1).$$

Replacing these inequalities in the result above we obtain

$$\begin{aligned}
\left\| (\widehat{f}_{2\Lambda} - f_{2\Lambda}) \phi_\Lambda \right\|^2 &< (e^{\Lambda\epsilon} - 1)^2 \sum_{l,l'=0}^{2\Lambda} \sum_{m,m'=0}^{\Lambda} \sum_{n \in L^{lm} \cap L^{l'm'}} N_n^{lm} N_n^{l'm'} Q_n \\
&\quad (f_{a_1 \dots a_r c_1 \dots c_{l-r}}^l \phi_{a_1 \dots a_r c_{l-r+1} \dots c_n}^m)^* f_{b_1 \dots b_{r'} c_1 \dots c_{l'-r'}}^{l'} \phi_{b_1 \dots b_{r'} c_{l'-r'+1} \dots c_n}^{m'} \\
&= (e^{\Lambda\epsilon} - 1)^2 \|f_{2\Lambda} \phi_\Lambda\|^2. \tag{149}
\end{aligned}$$

The last equality holds because the sum is nothing but  $\|f_{2\Lambda} \phi_\Lambda\|^2$ , i.e. what we would obtain from (148) replacing  $\widehat{N}_n^{lm} \mapsto 0$ , and therefore  $\widehat{f}_{2\Lambda} \mapsto 0$ . Moreover,  $\|f_{2\Lambda} \phi_\Lambda\| \leq \|f_{2\Lambda}\|_{op} \|\phi_\Lambda\|$ ; but both factors have  $\Lambda$ -independent bounds:  $\|\phi_\Lambda\| \leq \|\phi\|$ , while by the triangular inequality,  $\|f_{2\Lambda}\|_{op} \leq \|f\|_{op} + \|f_{2\Lambda} - f\|_{op}$ , and the second term goes to zero as  $\Lambda \rightarrow \infty$ , hence is bounded by some  $\eta \geq 0$ . So we end up with

$$\left\| (\widehat{f}_{2\Lambda} - f_{2\Lambda}) \phi_\Lambda \right\| < (e^{\Lambda\epsilon(\Lambda)} - 1) (\|f\|_{op} + \eta) \|\phi\| \xrightarrow{\Lambda \rightarrow \infty} 0; \tag{150}$$

the limit is zero because by (19)  $k(\Lambda)$  diverges at least as  $\Lambda^4$ , then by (109)  $\Lambda\epsilon(\Lambda)$  goes to zero as  $\Lambda^{-1}$ . Replaced in (147) this yields for all  $f \in C(S^d)$

$$\left\| (f - \widehat{f}_{2\Lambda}) \phi \right\| \xrightarrow{\Lambda \rightarrow \infty} 0 \tag{151}$$

i.e.  $\widehat{f}_{2\Lambda} \rightarrow f \cdot$  strongly, as claimed. Replacing  $f \mapsto fg$ , we find also that  $(\widehat{fg})_{2\Lambda} \rightarrow (fg) \cdot$  strongly for all  $f, g \in C(S^d)$ , as claimed. On the other hand, since  $\|\phi_\Lambda^\perp\| \leq \|\phi\|$  (because  $\phi = \phi_\Lambda + \phi_\Lambda^\perp$  with  $\langle \phi_\Lambda, \phi_\Lambda^\perp \rangle = 0$ ), relations (147) and (147) imply also

$$\left\| (f - \widehat{f}_{2\Lambda}) \phi \right\| \leq \left[ \|f - f_{2\Lambda}\|_{op} + \|f_{2\Lambda} - \widehat{f}_{2\Lambda}\|_{op} + (e^{\Lambda\epsilon(\Lambda)} - 1) (\|f\|_{op} + \eta) \right] \|\phi\| \leq F \|\phi\|,$$

where  $F > 0$  is an upper bound for the expression in the square bracket; hence

$$\|\widehat{f}_{2\Lambda} \phi\| \leq \|(\widehat{f}_{2\Lambda} - f) \phi\| + \|f \phi\| \leq F \|\phi\| + \|f\|_\infty \|\phi\| \leq (F + \|f\|_\infty) \|\phi\|$$

i.e. the operator norms  $\|\hat{f}_{2\Lambda}\|_{op}$  of the  $\hat{f}_{2\Lambda}$  are uniformly bounded:  $\|\hat{f}_{2\Lambda}\|_{op} \leq (F + \|f\|_\infty) \|\phi\|$ . Therefore (151) implies the last claim of the theorem

$$\begin{aligned} \|(fg - \hat{f}_{2\Lambda}\hat{g}_{2\Lambda})\phi\| &\leq \|(f - \hat{f}_{2\Lambda})g\phi\| + \|\hat{f}_{2\Lambda}(g - \hat{g}_{2\Lambda})\phi\| \\ &\leq \|(f - \hat{f}_{2\Lambda})(g\phi)\| + \|\hat{f}_{2\Lambda}\|_{op} \|(g - \hat{g}_{2\Lambda})\phi\| \xrightarrow{\Lambda \rightarrow \infty} 0. \end{aligned} \quad (152)$$

## References

- [1] H. S. Snyder, *Quantized Space-Time*, Phys. Rev. **71** (1947), 38.
- [2] S. Doplicher, K. Fredenhagen, J. E. Roberts, *Spacetime Quantization Induced by Classical Gravity*, Phys. Lett. **B 331** (1994), 39-44; *The quantum structure of spacetime at the Planck scale and quantum fields*, Commun. Math. Phys. **172** (1995), 187-220;
- [3] A. Connes, J. Lott, *Particle models and noncommutative geometry*, Nucl. Phys. (Proc. Suppl.) **B18** (1990), 29-47.
- [4] A.H.Chamseddine, A. Connes, *Noncommutative geometry as a framework for unification of all fundamental interactions including gravity. Part I*, Fortsch. Phys. **58** (2010) 553.
- [5] A. Connes, *Noncommutative geometry*, Academic Press, 1995.
- [6] J. M. Gracia-Bondia, H. Figueroa, J. Varilly, *Elements of Non-commutative geometry*, Birkhauser, 2000.
- [7] G. Landi, Giovanni (1997), *An introduction to noncommutative spaces and their geometries*, Lecture Notes in Physics 51, Springer-Verlag, 1997
- [8] J. Madore, *An introduction to noncommutative differential geometry and its physical applications*, Cambridge University Press, 1999.
- [9] P. Aschieri, H. Steinacker, J. Madore, P. Manousselis, G. Zoupanos *Fuzzy extra dimensions: Dimensional reduction, dynamical generation and renormalizability*, SFIN **A1** (2007) 25-42; and references therein.
- [10] A. Y. Alekseev, A. Recknagel, V. Schomerus, *Non-commutative world-volume geometries: branes on SU(2) and fuzzy spheres*, JHEP **09** (1999) 023.
- [11] Y. Hikida, M. Nozaki and Y. Sugawara, *Formation of spherical d2-brane from multiple D0-branes*, Nucl. Phys. **B617** (2001), 117.
- [12] J. Madore, *Quantum mechanics on a fuzzy sphere*, Journ. Math. Phys. **32** (1991) 332; *The Fuzzy Sphere*, Class. Quantum Grav. **9** (1992), 6947.
- [13] J. Hoppe, *Quantum theory of a massless relativistic surface and a 2-dimensional bound state problem*, PhD thesis MIT1982; B. de Wit, J. Hoppe, H. Nicolai, Nucl. Phys. **B305** (1988), 545.

- [14] H. Grosse, J. Madore, *A noncommutative version of the Schwinger model*, Phys. Lett. **B283** (1992), 218.
- [15] H. Grosse, C. Klimcik, P. Presnajder, *Towards Finite Quantum Field Theory in Non-Commutative Geometry*, Int. J. Theor. Phys. **35** (1996), 231-244.
- [16] H. Grosse, C. Klimcik, P. Presnajder, *On Finite 4D Quantum Field Theory in Non-Commutative Geometry*, Commun. Math. Phys. **180** (1996), 429-438.
- [17] S. Ramgoolam, *On spherical harmonics for fuzzy spheres in diverse dimensions*, Nucl. Phys. **B610** (2001), 461-488; *Higher dimensional geometries related to fuzzy odd-dimensional spheres*, JHEP10(2002)064; and references therein.
- [18] B. P. Dolan, D. O'Connor, *A Fuzzy three sphere and fuzzy tori*, JHEP **0310** (2003) 06.
- [19] B. P. Dolan, D. O'Connor and P. Presnajder, *Fuzzy complex quadrics and spheres*, JHEP **0402** (2004) 055.
- [20] H. Steinacker, *Emergent gravity on covariant quantum spaces in the IKKT model*, J. High Energy Physics 2016: 156.
- [21] M. Sperling, H. Steinacker, *Covariant 4-dimensional fuzzy spheres, matrix models and higher spin*, J. Phys. A: Math. Theor. **50** (2017), 375202.
- [22] G. Fiore, F. Pisacane, *Fuzzy circle and new fuzzy sphere through confining potentials and energy cutoffs*, J. Geom. Phys. **132** (2018), 423-451.
- [23] F. Pisacane,  *$O(D)$ -equivariant fuzzy hyperspheres*, arXiv:2002.01901.
- [24] G. Fiore, F. Pisacane, *Energy cutoff, noncommutativity and fuzziness: the  $O(D)$ -covariant fuzzy spheres*, PoS(CORFU2019)208. <https://pos.sissa.it/376/208>
- [25] G. Fiore, F. Pisacane, *New fuzzy spheres through confining potentials and energy cutoffs*, Proceedings of Science Volume 318, PoS(CORFU2017)184.
- [26] FIORE, G.: *Quantum group covariant (anti)symmetrizers,  $\varepsilon$ -tensors, vielbein, Hodge map and Laplacian*. J. Phys. A: Math. Gen. **37** (2004), 9175-9193.
- [27] E. Hawkins, *Quantization of equivariant vector bundles*, Commun. Math. Phys. **202** (1999), 517-546.
- [28] G. Fiore, F. Pisacane, *The  $x_i$ -eigenvalue problem on some new fuzzy spheres*, J. Phys. A: Math. Theor. **53** (2020), 095201.
- [29] G. Fiore, F. Pisacane, *On localized and coherent states on some new fuzzy spheres*, Lett. Math. Phys. **110** (2020), 1315-1361.
- [30] F. D'Andrea, F. Lizzi, P. Martinetti, *Spectral geometry with a cut-off: Topological and metric aspects*, J. Geom. Phys. **82** (2014), 18-45.



- [31] G. Fiore, D. Franco, T. Weber, *Twisted quadrics and algebraic submanifolds in  $\mathbb{R}^n$* , Math. Phys. Anal. Geom. **23**, 38 (2020)
- [32] G. Fiore, T. Weber, *Twisted submanifolds of  $\mathbb{R}^n$* , Lett. Math. Phys. **111**, 76 (2021)