

# A VARIANT OF SOME CYCLOTOMIC MATRICES INVOLVING TRINOMIAL COEFFICIENTS

YU-BO LI AND NING-LIU WEI

ABSTRACT. In this paper, by using the theory of circulant matrices we study some matrices over finite fields which involve the quadratic character and trinomial coefficients.

## 1. INTRODUCTION

Let  $p$  be an odd prime and let  $(\cdot/p)$  be the Legendre symbol. The study of the matrices involving Legendre symbols can trace back to the works of Lehmer [6] and Carlitz [1]. For example, Carlitz [1] initiated the study of the matrix

$$C_p = \left[ \left( \frac{j-i}{p} \right) \right]_{1 \leq i, j \leq p-1}.$$

Carlitz [1, Thm. 4 (4.9)] proved that the characteristic polynomial of  $C_p$  is

$$f_p(t) = \left( t^2 - (-1)^{\frac{p-1}{2}} p \right)^{\frac{p-3}{2}} \left( t^2 - (-1)^{\frac{p-1}{2}} \right).$$

Later Chapman [2, 3] and Vsemirnov [8, 9] investigated many variants of Carlitz's matrix  $C_p$ . In particular, using sophisticated matrix decompositions Vsemirnov [8, 9] confirmed Chapman's "evil" determinant conjecture which says that

$$\det \left[ \left( \frac{j-i}{p} \right) \right]_{1 \leq i, j \leq \frac{p+1}{2}} = \begin{cases} -a_p & \text{if } p \equiv 1 \pmod{4}, \\ 1 & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

The number  $a_p$  is defined by the following equality

$$\varepsilon_p^{(2 - \frac{2}{p})h_p} = a_p + b_p \sqrt{p}, \quad a_p, b_p \in \mathbb{Q},$$

where  $\varepsilon_p > 1$  and  $h_p$  denote the fundamental unit and class number of the real quadratic field  $\mathbb{Q}(\sqrt{p})$ .

Recently, Sun [7] further studied some variants of Carlitz's matrix  $C_p$ . For example, Sun [7, Thm. 1.2] showed that

$$-\det \left[ \left( \frac{i^2 + j^2}{p} \right) \right]_{1 \leq i, j \leq p-1}$$

---

*Key words and phrases.* Central Trinomial Coefficients, Finite Fields, Determinants.

*2020 Mathematics Subject Classification.* Primary 05A19, 11C20; Secondary 15A18, 15B57, 33B10.

is always a quadratic residue modulo  $p$ . Along this line, for any integers  $c, d$ , the arithmetic properties of the matrix

$$\left[ \left( \frac{i^2 + cij + dj^2}{p} \right) \right]_{1 \leq i, j \leq p-1}$$

were extensively studied. Readers may refer to [5, 7, 10, 11, 13] for details on this topic.

On the other hand, in the same paper Sun posed a conjecture (see [7, Remark 1.3]) which states that

$$2 \det \left[ \frac{1}{i^2 - ij + j^2} \right]_{1 \leq i, j \leq p-1}$$

is a quadratic residue modulo  $p$  whenever  $p \equiv 2 \pmod{3}$  is an odd prime. This conjecture was later proved by Wu, She and Ni [12].

Also, Let

$$D_p = \left[ (i^2 + j^2) \left( \frac{i^2 + j^2}{p} \right) \right]_{1 \leq i, j \leq (p-1)/2}.$$

Recently, Wu, She and Wang [13] proved a conjecture posed by Sun which states that

$$\left( \frac{D_p}{p} \right) = \begin{cases} 1 & \text{if } p \equiv 1 \pmod{4}, \\ (-1)^{\frac{h(-p)-1}{2}} & \text{if } p \equiv 3 \pmod{4}, \end{cases}$$

where  $h(-p)$  is the class number of  $\mathbb{Q}(\sqrt{-p})$ .

Now let  $\mathbb{F}_q = \{0, a_1, \dots, a_{q-1}\}$  be the finite field with  $q$  elements, where  $q$  is an odd prime power. Also, let  $\chi$  be the unique quadratic multiplicative character of  $\mathbb{F}_q$ , i.e.,

$$\chi(x) = \begin{cases} 0 & \text{if } x = 0, \\ 1 & \text{if } x \text{ is a nonzero square,} \\ -1 & \text{otherwise.} \end{cases}$$

Motivated by the above results, in this paper, we shall study the following matrix over  $\mathbb{F}_q$ :

$$S_q := \left[ (a_i^2 + a_i a_j + a_j^2) \chi(a_i^2 + a_i a_j + a_j^2) \right]_{1 \leq i, j \leq q-1}. \quad (1.1)$$

Let  $n$  be a non-negative integer. The central trinomial coefficient  $T_n$  is defined to be the coefficient of  $x^n$  in the polynomial  $(x^2 + x + 1)^n$ . Equivalently,  $T_n$  is the constant term of  $(x + 1 + x^{-1})^n$ . Now we state our main result.

**Theorem 1.1.** *Let  $q$  be an odd prime power. Then there exists an element  $u \in \mathbb{F}_q$  such that*

$$\det S_q = T_{\frac{q+1}{2}} \cdot u^2.$$

As a consequence of Theorem 1.1, we have the following result.

**Corollary 1.1.** *Let  $p$  be an odd prime. Suppose  $p \nmid \det S_p$ . Then*

$$\left( \frac{\det S_p}{p} \right) = \left( \frac{T_{\frac{p+1}{2}}}{p} \right).$$

Next we shall give a necessary and sufficient condition for  $S_q$  to be singular. Let  $n$  be a non-negative integer. Then the trinomial coefficients  $\binom{n}{k}_2$  is defined by

$$\left(x + \frac{1}{x} + 1\right)^n = \sum_{k=-n}^n \binom{n}{k}_2 x^k.$$

Clearly  $\binom{n}{0}_2 = T_n$ . Now we state our last result.

**Theorem 1.2.** *Let  $\mathbb{F}_q$  be the finite field with  $q > 5$  and  $(q, 22) = 1$ . Then*

$$\det S_q = \frac{121}{64} \cdot T_{\frac{q+1}{2}} \cdot \prod_{k=1}^{(p-5)/2} \binom{\frac{q+1}{2}}{k}_2 \in \mathbb{F}_p,$$

where  $p$  is the characteristic of  $\mathbb{F}_q$ . Also,  $S_q$  is a singular matrix over  $\mathbb{F}_q$  if and only if

$$\binom{\frac{q+1}{2}}{k}_2 \equiv 0 \pmod{p}$$

for some  $0 \leq k \leq (q-5)/2$ .

We will prove our main results in Section 2 and Section 3 respectively.

## 2. PROOF OF THEOREM 1.1

We first introduce the definition of the circulant matrices. Let  $R$  be a commutative ring. Let  $m$  be a positive integer and  $t_0, t_1, \dots, t_{m-1} \in R$ . We define the circulant matrix  $C(t_0, \dots, t_{m-1})$  to be an  $m \times m$  matrix whose  $(i-j)$ -entry is  $t_{j-i}$  where the indices are cyclic module  $m$ . Wu [11, Lemma 3.4] obtained the following result.

**Lemma 2.1.** *Let  $R$  be a commutative ring. Let  $m$  be a positive even integer. Let  $t_0, t_1, \dots, t_{m-1} \in R$  such that*

$$t_i = t_{m-i} \text{ for each } 1 \leq i \leq m-1.$$

Then there exists an element  $u \in R$  such that

$$\det C(t_0, \dots, t_{m-1}) = \left(\sum_{i=0}^{m-1} t_i\right) \left(\sum_{i=0}^{m-1} (-1)^i t_i\right) \cdot u^2.$$

We also need the following known result.

**Lemma 2.2.** *Let  $k$  be an integer. Then*

$$\sum_{x \in \mathbb{F}_q \setminus \{0\}} x^k = \begin{cases} -1 & \text{if } p-1 \mid k, \\ 0 & \text{otherwise.} \end{cases}$$

Now we are in a position to prove our main results. For simplicity, the summations  $\sum_{x \in \mathbb{F}_q}$  and  $\sum_{x \in \mathbb{F}_q \setminus \{0\}}$  are abbreviated as  $\sum_x$  and  $\sum_{x \neq 0}$  respectively.

**Proof of Theorem 1.1.** Fix a primitive element  $g$  of  $\mathbb{F}_q$ . Then one can verify that

$$\begin{aligned} \det S_q &= \prod_{i=1}^{q-1} a_i^2 \cdot \det \left[ \left( \left( \frac{a_j}{a_i} \right)^2 + \frac{a_j}{a_i} + 1 \right) \chi \left( \left( \frac{a_j}{a_i} \right)^2 + \frac{a_j}{a_i} + 1 \right) \right]_{1 \leq i, j \leq q-1} \\ &= \det \left[ \frac{1}{g^{j-i}} (g^{2(j-i)} + g^{j-i} + 1)^{\frac{q+1}{2}} \right]_{0 \leq i, j \leq q-2}. \end{aligned}$$

Let  $t_i = g^{-i}(g^{2i} + g^i + 1)^{\frac{q+1}{2}}$  for  $0 \leq i \leq q-2$ . Then

$$\det S_q = \det C(t_0, t_1, \dots, t_{q-2})$$

and  $t_i = t_{q-1-i}$  for  $1 \leq i \leq q-3$ . Applying Lemma 2.1 there is an element  $u \in \mathbb{F}_q$  such that

$$\det S_q = \left( \sum_{i=0}^{q-2} t_i \right) \left( \sum_{i=0}^{q-2} (-1)^i t_i \right) u^2. \quad (2.2)$$

We first evaluate  $\sum_{i=0}^{q-2} t_i$ .

$$\begin{aligned} \sum_{i=0}^{q-2} t_i &= \sum_{x \neq 0} \frac{1}{x} (x^2 + x + 1)^{\frac{q+1}{2}} \\ &= \sum_{x \neq 0} \left( x + \frac{1}{x} + 1 \right) \cdot (x^2 + x + 1)^{\frac{q-1}{2}} \\ &= 2 \sum_{x \neq 0} x \cdot (x^2 + x + 1)^{\frac{q-1}{2}} + \sum_{x \neq 0} (x^2 + x + 1)^{\frac{q-1}{2}} \\ &= -1 + 2 \sum_x x \left( \left( x + \frac{1}{2} \right)^2 + \frac{3}{4} \right)^{\frac{q-1}{2}} + \sum_x \left( \left( x + \frac{1}{2} \right)^2 + \frac{3}{4} \right)^{\frac{q-1}{2}} \\ &= -1 + 2 \sum_x \left( x - \frac{1}{2} \right) \left( x^2 + \frac{3}{4} \right)^{\frac{q-1}{2}} + \sum_x \left( x^2 + \frac{3}{4} \right)^{\frac{q-1}{2}} = -1. \end{aligned}$$

Hence we obtain

$$\sum_{i=0}^{q-2} t_i = -1. \quad (2.3)$$

Next we turn to  $\sum_{i=0}^{q-2} (-1)^i t_i$ .

$$\sum_{i=0}^{q-2} (-1)^i t_i = \sum_{x \neq 0} \frac{1}{x} \cdot (x^2 + x + 1) \cdot \chi \left( \frac{1}{x} \right) \chi (x^2 + x + 1)$$

$$= \sum_{x \neq 0} \left( x + \frac{1}{x} + 1 \right)^{\frac{q+1}{2}} = -T_{\frac{q+1}{2}}.$$

The last equality follows from Lemma 2.2. We therefore obtain

$$\sum_{i=0}^{q-2} (-1)^i t_i = -T_{\frac{q+1}{2}}. \quad (2.4)$$

Combining (2.3) and (2.4) with (2.2), we see that  $\det S_q = T_{\frac{q+1}{2}} \cdot u^2$  for some  $u \in \mathbb{F}_q$ .

This completes the proof.  $\square$

### 3. PROOF OF THEOREM 1.2

We begin with the following known result (see [4, Lemma 10]).

**Lemma 3.3.** *Let  $R$  be a commutative ring and let  $n$  be a positive integer. For any polynomial  $P(T) = p_{n-1}T^{n-1} + \cdots + p_1T + p_0 \in R[T]$  we have*

$$\det [P(X_i Y_j)]_{1 \leq i, j \leq n} = \prod_{i=0}^{n-1} p_i \prod_{1 \leq i < j \leq n} (X_j - X_i)(Y_j - Y_i).$$

We also need the following lemma.

**Lemma 3.4.** *Let  $q$  be an odd prime. Then for any non-zero element  $a \in \mathbb{F}_q$  we have  $(a^2 + a + 1)^{\frac{q+1}{2}} = f(a)$ , where*

$$f(T) = \frac{11}{8} + T + \frac{11}{8}T^2 + \sum_{k=-(q-5)/2}^{(q-5)/2} \binom{\frac{q+1}{2}}{k}_2 T^{k+\frac{q+1}{2}} \quad (3.5)$$

is a polynomial over  $\mathbb{F}_q$ .

*Proof.* As  $a \neq 0$ , we have  $a^{q+k} = a^{k+1}$  for any integer  $k$ . Using this and  $\binom{n}{k}_2 = \binom{n}{-k}_2$  one can verify that  $(a^2 + a + 1)^{\frac{q+1}{2}}$  is equal to

$$\begin{aligned} & \sum_{k=-(q-5)/2}^{(q-5)/2} \binom{\frac{q+1}{2}}{k}_2 a^{k+\frac{q+1}{2}} + \left( \binom{\frac{q+1}{2}}{\frac{q+1}{2}}_2 + \binom{\frac{q+1}{2}}{\frac{q-3}{2}}_2 \right) (1 + a^2) + \left( \binom{\frac{q+1}{2}}{\frac{q-1}{2}}_2 + \binom{\frac{q+1}{2}}{\frac{q-3}{2}}_2 \right) a \\ &= \frac{11}{8} + a + \frac{11}{8}a^2 + \sum_{k=-(q-5)/2}^{(q-5)/2} \binom{\frac{q+1}{2}}{k}_2 a^{k+\frac{q+1}{2}}. \end{aligned}$$

The last equality follows from (below the trinomial coefficient  $\binom{n}{k}_2$  is viewed as an element of  $\mathbb{F}_q$ ).

$$\binom{\frac{q+1}{2}}{\frac{q+1}{2}}_2 = 1, \quad \binom{\frac{q+1}{2}}{\frac{q-1}{2}}_2 = \frac{q+1}{2} = \frac{1}{2}, \quad \binom{\frac{q+1}{2}}{\frac{q-3}{2}}_2 = \frac{1}{2} \cdot \frac{q+1}{2} \cdot \frac{q+3}{2} = \frac{3}{8}.$$

This completes the proof.  $\square$

Now we are in a position to prove our last result.

**Proof of Theorem 1.2.** By Lemma 3.4 one can verify that

$$\begin{aligned} \det S_q &= \prod_{i=1}^{q-1} a_i^{q+1} \cdot \det \left[ \left( \left( \frac{a_j}{a_i} \right)^2 + \frac{a_j}{a_i} + 1 \right)^{\frac{q+1}{2}} \right]_{1 \leq i, j \leq q-1} \\ &= \det \left[ \left( \left( \frac{a_j}{a_i} \right)^2 + \frac{a_j}{a_i} + 1 \right)^{\frac{q+1}{2}} \right]_{1 \leq i, j \leq q-1} \\ &= \det \left[ f \left( \frac{a_j}{a_i} \right) \right]_{1 \leq i, j \leq q-1}, \end{aligned}$$

where  $f$  is defined by (3.5).

Now applying Lemma 3.3 we obtain that  $\det S_q$  is equal to

$$\det \left[ f \left( \frac{a_j}{a_i} \right) \right]_{1 \leq i, j \leq q-1} = \frac{121}{64} \cdot \binom{\frac{q+1}{2}}{0}_2 \cdot \prod_{k=1}^{(p-5)/2} \binom{\frac{q+1}{2}}{k}_2 \cdot \prod_{1 \leq i < j \leq q-1} (a_j - a_i) \left( \frac{1}{a_j} - \frac{1}{a_i} \right). \quad (3.6)$$

By [12, Eq. (3.3)] we further have

$$\prod_{1 \leq i < j \leq q-1} (a_j - a_i) \left( \frac{1}{a_j} - \frac{1}{a_i} \right) = 1. \quad (3.7)$$

Hence by (3.6) and (3.7) we obtain

$$\det S_q = \frac{121}{64} \cdot \binom{\frac{q+1}{2}}{0}_2 \cdot \prod_{k=1}^{(p-5)/2} \binom{\frac{q+1}{2}}{k}_2. \quad (3.8)$$

As  $q > 5$  and  $(q, 22) = 1$ , by (3.8) we see that

$$\det S_q = 0 \Leftrightarrow \binom{\frac{q+1}{2}}{k}_2 \equiv 0 \pmod{p} \text{ for some } 0 \leq k \leq (p-5)/2,$$

where  $p$  is the characteristic of  $\mathbb{F}_q$ . This completes the proof.  $\square$

## REFERENCES

- [1] L. Carlitz, Some cyclotomic matrices, *Acta Arith.* 5 (1959), 293–308.
- [2] R. Chapman, Determinants of Legendre symbol matrices, *Acta Arith.* 115 (2004), 231–244.
- [3] R. Chapman, My evil determinant problem, preprint, December 12, 2012, available at <http://empslocal.ex.ac.uk/people/staff/rjchapma/etc/evildet.pdf>.
- [4] Krattenthaler, C.: Advanced determinant calculus: a complement. *Linear Algebra Appl.* 411 (2005), 68–166
- [5] D. Krachun, F. Petrov, Z.-W. Sun, M. Vsemirnov, On some determinants involving Jacobi symbols, *Finite Fields Appl.* 64 (2020), 101672.
- [6] D. H. Lehmer, On certain character matrices, *Pacific J. Math.* 6 (1956), 491–499.
- [7] Z.-W. Sun, On some determinants with Legendre symbols entries, *Finite Fields Appl.* 56 (2019), 285–307.

- [8] M. Vsemirnov, On the evaluation of R. Chapman’s “evil determinant”, *Linear Algebra Appl.* 436 (2012), 4101–4106.
- [9] M. Vsemirnov, On R. Chapman’s “evil determinant”: case  $p \equiv 1 \pmod{4}$ , *Acta Arith.* 159 (2013), 331–344.
- [10] H.-L. Wu, Determinants concerning Legendre symbols, *C. R. Math. Acad. Sci. Paris* 359 (2021), 651–655.
- [11] H.-L. Wu, Elliptic curves over  $\mathbb{F}_p$  and determinants of Legendre matrices, *Finite Fields Appl.* 76 (2021), 101929.
- [12] H.-L. Wu, Y.-F. She, H.-X. Ni, A conjecture of Zhi-Wei Sun on determinants over finite fields, *Bull. Malays. Math. Sci. Soc.* 45 (2022), 2405–2412.
- [13] H.-L. Wu, Y.-F. She and L.-Y. Wang, Cyclotomic matrices and hypergeometric functions over finite fields, *Finite Fields and Their Applications*, 82 (2022), 102054.

(YU-BO LI) SCHOOL OF SCIENCE, NANJING UNIVERSITY OF POSTS AND TELECOMMUNICATIONS, NANJING 210023, PEOPLE’S REPUBLIC OF CHINA

*Email address:* lybmath2022@163.com

(NING-LIU WEI) SCHOOL OF SCIENCE, NANJING UNIVERSITY OF POSTS AND TELECOMMUNICATIONS, NANJING 210023, PEOPLE’S REPUBLIC OF CHINA

*Email address:* weiningliu6@163.com